

Greater Consideration of Animals Will Enhance Coastal Restoration Outcomes

MICHAEL SIEVERS[✉], CHRISTOPHER J. BROWN[✉], CHRISTINA A. BUELOW, ROBIN HALE, ANDRIA OSTROWSKI, MEGAN I. SAUNDERS, BRIAN R. SILLIMAN, STEPHEN E. SWEARER, MISCHA P. TURSCHWELL, STEPHANIE R. VALDEZ, AND ROD M. CONNOLLY

As efforts to restore coastal habitats accelerate, it is critical that investments are targeted to most effectively mitigate and reverse habitat loss and its impacts on biodiversity. One likely but largely overlooked impediment to effective restoration of habitat-forming organisms is failing to explicitly consider non-habitat-forming animals in restoration planning, implementation, and monitoring. These animals can greatly enhance or degrade ecosystem function, persistence, and resilience. Bivalves, for instance, can reduce sulfide stress in seagrass habitats and increase drought tolerance of saltmarsh vegetation, whereas megaherbivores can detrimentally overgraze seagrass or improve seagrass seed germination, depending on the context. Therefore, understanding when, why, and how to directly manipulate or support animals can enhance coastal restoration outcomes. In support of this expanded restoration approach, we provide a conceptual framework, incorporating lessons from structured decision-making, and describe potential actions that could lead to better restoration outcomes using case studies to illustrate practical approaches.

Keywords: decision science, ecological restoration, rehabilitation, translocation, transplantation

Restoration is a key challenge of the twenty-first century, because ecosystems are being increasingly lost and degraded (McDonald et al. 2016, Gann et al. 2019, Halpern et al. 2019, He and Silliman 2019, Williams et al. 2021). Conservation efforts have traditionally been focused on the protection of intact habitats or the mitigation of stressors, but these approaches have failed at times to reverse widespread trends in ecological degradation (Lotze et al. 2011, Díaz et al. 2019, Griffiths et al. 2020). The restoration of coastal and marine ecosystems is particularly important, because over 775 million people depend on coastal systems; they have a relatively high role in climate mitigation and adaptation, and; have undergone widespread loss (Duarte et al. 2013a, Selig et al. 2019, Dunic et al. 2021, Murray et al. 2022). Restoration is therefore necessary to reverse coastal habitat loss and degradation, enhance biodiversity, and reestablish ecosystem services such as fisheries production, coastline protection, and climate change mitigation (Wood et al. 2019, Abelson et al. 2020, Waltham et al. 2020, Buelow et al. 2022). Coastal restoration efforts are consequently accelerating, supported by international calls to action, including the UN Decade of Ecosystem Restoration and Sustainable Development Goals (Perring et al. 2015, Young and Schwartz 2019, Sheaves et al. 2021). However, despite

some notable exceptions (Saunders et al. 2020), coastal restoration projects tend to be small scale and expensive and have low survival rates of the habitat-forming species (Dale et al. 2014, Bayraktarov et al. 2016, van Katwijk et al. 2016, Cooke et al. 2019).

We posit that an often overlooked but ecologically significant gap in the implementation of restoration is that non-habitat-forming animals are not explicitly and holistically included in restoration planning, implementation, and monitoring (Halpern et al. 2007, Jones and Davidson 2016, Hale et al. 2019). Only 13% of the studies in a review of priorities and motivations of marine coastal restoration research, for instance, measured non-habitat-forming animal responses (Bayraktarov et al. 2020), and only a small proportion of seagrass restoration efforts explicitly manipulate animals (Zhang et al. 2021). In the present article, habitat-forming animals are those that form the structural habitat being restored, such as reef building corals and oysters, whereas non-habitat-forming animals are all other animals. Although the detrimental impacts that animals can have on restored habitats are being considered in some specific restoration efforts (e.g., in those that remove species that predate or graze on transplanted habitat formers), many animals perform a suite of vital functions that are necessary

for ecosystem persistence, enhance ecosystem resilience through assisting disturbance recovery, and drive many of the services that restoration actively seeks to enhance (figure 1; Halpern et al. 2007). Consideration of mutualisms with animals that enhance habitat-forming species growth and survival is particularly important for coastal habitats that are frequently disturbed by natural and anthropogenic sources that make restoration inherently difficult (Lewis and Anderson 2012, Renzi et al. 2019, Gagnon et al. 2020). One of the most well-known mutualisms likely to have significant benefits for coastal restoration initiatives exists between bivalves and coastal vegetation. Bivalves, for example, can reduce sulfide stress in seagrass and mangroves and can facilitate saltmarsh vegetation by providing nutrients and reducing erosion (figure 1; also see Gagnon et al. 2020 and the references within).

Despite hundreds of ecological papers showing that biotic interactions are important to marine foundation species growth and success, we suggest that this knowledge could be better incorporated into coastal restoration planning, implementation, and evaluations of success. We argue that this is true whether animals are a direct goal of restoration, are an impediment to successful restoration, or provide functions that improve outcomes for habitat formers and the ecosystem. Several works describe these interactions within coastal ecosystems, such as mangrove forests (Gedan and Silliman 2009), saltmarshes (Derksen-Hooijberg et al. 2018), seagrass meadows (Valdez et al. 2020), coral reefs (Shaver and Silliman 2017, Seraphim et al. 2020), kelp forests (Eger et al. 2020), and oyster reefs (Reeves et al. 2020). Furthermore, although these general concepts are raised in restoration guidelines (e.g., McDonald et al. 2016, Morris et al. 2020, Eger et al. 2022, Shaver et al. 2022) and although we acknowledge that there are successful restoration projects in which animals have not been explicitly considered, a better understanding of when, why, and how to directly manipulate or support animals in restored habitats could significantly improve outcomes for many coastal restoration initiatives (Derksen-Hooijberg et al. 2018, Renzi et al. 2019, Gagnon et al. 2020).

Restoration objectives and how animals can be incorporated into restoration actions

Coastal restoration efforts can have a range of different objectives that vary both in the importance of animals to meeting them and in the ways in which animals could be manipulated or supported to improve outcomes. For example, animals can be a direct or implicit goal of restoration (e.g., enhancing fisheries or improving habitat for a threatened species; Taylor et al. 2017), can be part of a more holistic goal to restore whole ecosystems (e.g., restoring habitats to reference conditions; McDonald et al. 2016), can provide functions that can benefit alternative goals of ecosystem restoration (Abelson et al. 2016, Gagnon et al. 2020, Valdez et al. 2020), or can be an impediment to restoration goals (e.g., overabundant grazers or invasive species; Morris et al. 2020). There is considerable risk that well-funded

coastal restoration will be attempted globally with limited consideration of animals (Lee et al. 2019), subsequently limiting the success of many projects with a strong or sole habitat-forming species focus.

To encourage and guide scientists and managers to better incorporate animals into coastal restoration planning, implementation, and monitoring, we describe four key contexts in which animals could be manipulated and supported in restored habitats and the actions that can lead to positive outcomes (figure 2). We focus on six coastal marine ecosystems—mangrove forests, saltmarshes, seagrass meadows, coral reefs, macroalgae reefs, and oyster reefs—that have high intrinsic and extrinsic value to society, have unique ecological niches, are under accelerating threats from both land- and sea-based stressors, and are seeing a rapid rise in restoration initiatives (Perillo et al. 2018, Halpern et al. 2019, Bayraktarov et al. 2020). We outline a conceptual framework for an expanded approach that incorporates the benefits of a wider consideration of animals. Notwithstanding differences among restoration objectives, we argue that greater consideration of animals within planning, implementation, and monitoring can have benefits for most coastal restoration initiatives, and we use case studies to illustrate practical approaches (see box 1). Although there are important logistical, financial, legal, permitting, and societal considerations, because these are complex, involved, and require their own treatment to do them justice, we do not cover these in detail in the present article.

Actively add animals to restored habitats (action A). The first action involves situations in which animals could be actively added into restored habitats. The *Field of Dreams* hypothesis, whereby restoring vegetation and physical structure are assumed to lead to animal colonization (i.e., “if you build it, they will come”) and that is sometimes applied to restoration projects, may not come to fruition; animals might not colonize restored habitats because of, for example, dispersal limitation or a lack of suitable source populations (Palmer et al. 1997, Lewis III 2010, Sundermann et al. 2011). In this instance, translocating animals into restored habitats may benefit coastal restoration outcomes, whether the animals are a direct goal (e.g., recovering populations of threatened birds that are incapable of colonizing restored habitats unassisted; figure 2, action 1A) or provide important functions that lead to improved restoration outcomes (e.g., algae herbivory or predation of herbivores that subsequently enhances the survival of the habitat-forming species; figure 2, action 3A; Seddon et al. 2014, Davis et al. 2019). Even when natural colonization is possible, assisting it through translocation may lead to substantial improvements in the recovery and development of habitat formers. For instance, transplanting mussels increased drought tolerance and vegetation growth by upward of 50% in restored saltmarsh (box 1; Angelini et al. 2016, Derksen-Hooijberg et al. 2018), incorporating sponges into coral reef restoration more than doubled successful coral colonization (Biggs 2013), and clam inclusion greatly enhanced seagrass biomass




Figure 1. Conceptual diagram depicting positive (black arrows) and negative (red arrows), indirect (dashed lines) and direct (solid lines) interactions (with respect to effects on the habitat-forming species) between non-habitat-forming animals and coastal habitats. The sets of blue arrows indicate a context dependent pathway of effect. Interactions are categorized as (a) trophic interactions, (b) fear responses, (c) reproduction, (d) environmental alterations, and (e) ecosystem resilience. References: ¹Eger et al. 2020, ²Rotjan and Lewis 2006, ³Seraphim et al. 2020, ⁴Christlanen et al. 2014, ⁵Gangal et al. 2021, ⁶Grabowski 2004, ⁷Tol et al. 2021, ⁸Qiu et al. 2021, ⁹Ellison et al. 1996, ¹⁰van der Heide et al. 2012, ¹¹Angelini et al. 2016, ¹²Hughes et al. 2016, ¹³Foster et al. 2021, ¹⁴Hensel et al. 2021.




Figure 2. Four key contexts (1–4) in which animals could be manipulated and supported within coastal restoration, the actions (A–D) that can lead to positive outcomes, and examples to illustrate links between contexts and actions.

and meadow growth in seeding experiments (box 1; Zhang et al. 2021). Furthermore, adding animals to nursery rearing tanks with habitat formers being cultivated for outplanting can also improve outcomes (e.g., adding algal grazers to enhance reared coral survival; Neil et al. 2021).

Restocking of animals is a common tool in the management of nonmarine aquatic ecosystems, aimed at restoring water quality and vegetation characteristics (e.g., Angeler et al. 2003, Cowx and Gerdeaux 2004). Although less applied in the marine environment, some attempts at restocking have been carried out in marine ecosystems, mainly as a fishery management tool targeted at commercial fish populations (e.g., Lorenzen et al. 2010, Leber 2013). There have also been attempts at restocking invertebrate species such as the grazing gastropod *Trochus* sp. into coral reefs (e.g., Villanueva et al. 2010). Modeling suggests that restocking of grazing fish on coral reefs can facilitate reef recovery and can become profitable within several years (Obolski et al. 2016), and such an approach has been proposed to both significantly benefit the restoration of reef habitats and enhance fisheries stocks (Abelson et al. 2016). Ultimately, when appropriate and feasible, the active addition of animals that are the focus of restoration efforts or that help maintain vital ecological processes can enhance the success of coastal restoration initiatives (Swan et al. 2016).

Actively remove or exclude animals from restored habitats (action B). The second action involves situations in which animals could be actively removed or excluded from restored habitats. Under some circumstances animals can be detrimental to habitat-forming animals and, therefore, restoration outcomes, particularly early on as the habitat-forming species are becoming established (Poore et al. 2012). Targeted animal removal and exclusion can limit detrimental effects, such as impacts from bioturbation from worms (figure 2, action 1B, box 1; Suykerbuyk et al. 2012), overconsumption of planted seagrass by herbivores (figure 2, action 4B; Wendländer et al. 2019), overgrazing of macroalgae by urchins (Eger et al. 2022), and grazing and trampling of saltmarsh by ungulates (Davidson et al. 2017). Small-scale, manipulative experiments show that exclusion of herbivorous urchins and fish (Sharma et al. 2021), and corallivorous snails (Shaver et al. 2018) enhance coastal restoration outcomes via positive effects on habitat-forming species. The optimal intervention will likely depend on whether negative effects are expected or occurring, and the density dependence of those effects. For example, exclusion of urchins may only be necessary at the initial phases of planting macroalgae, until algal density reaches a point where positive density-dependent feedback processes within the population maintain its abundance (Eger et al. 2020). Although in many cases, the complete removal of

Box 1. Case studies of manipulation or support of animals that did or could improve coastal restoration outcomes.
Adding clams during seagrass restoration in the United States (action A)

Zhang and colleagues (2021) planted seagrass seeds within experimental plots (20×20 centimeter plots of *Zostera marina*) and added juvenile quahog clams (*Mercenaria mercenaria*; figure 3a). The addition of ten clams per plot led to significantly increased seagrass shoot length, a 500% expansion in patch size (versus no change in patches without clams), and ten times greater belowground biomass. Aboveground biomass and metrics related to seagrass reproduction—despite being several times higher in patches with clams—were not significantly different to control patches. The most likely casual mechanism was clams enhancing nitrogen availability. Zhang and colleagues (2021) also added harvest-size clams to 50×50 centimeter plots with transplanted adult seagrass (*Zostera marina* and *Halodule wrightii*), but clam addition had no effect.




Figure 3. Images related to the six case studies: (a) clams added to restored seagrass patches, (b) mussels added into transplanted saltmarsh, (c) a crushed shell layer added underneath transplanted seagrass, (d) urchins requiring removal for macroalgae restoration, (e) healthy coral reef sounds played at degraded reefs, and (f) nesting mounds for terrapins added to saltmarsh. See the main text for case study descriptions and references.

Adding mussels during saltmarsh restoration in the United States (action A)

Derksen-Hooijberg and colleagues (2018) cotransplanted ribbed mussels (*Geukensia demissa*) into transplanted saltmarsh plots (*Spartina alterniflora*; 25×15 centimeter, diameter $\times$ depth) to evaluate the strategy for enhancing restoration success (figure 3b). Twenty mussels per plot increased saltmarsh vegetation growth and clonal expansion by 50%, as a result of mussels increasing nutrient levels and reducing sulfide stress. Following experimental vegetation removal that mimicked a disturbance event, vegetation in the plots containing mussels exhibited three times greater survival.

Box 1. Continued.

Excluding lugworms during seagrass restoration in the Netherlands (action B)

Creating physical barriers to separate animals that damage restored vegetation can improve establishment success. For example, Suykerbuyk and colleagues (2012) applied a crushed shell layer to shallow excavations underneath transplanted seagrass (*Zostera noltii*) to reduce interactions between seagrass and bioturbating lugworms (*Arenicola marina*; figure 3c). The shell layer reduced adult lugworm density by over 80%, and therefore reduced the lugworms' negative engineering effects. This was predicted to be the primary driver of the 50%–140% enhancement in seagrass growth.

Removing urchins during macroalgae restoration in Japan (action B)

Herbivorous urchins can dominate rocky reefs and impede the restoration of macroalgal reefs, necessitating their removal (figure 3d; Miller and Shears 2022). Watanuki and colleagues (2010) employed citizen scientists to remove urchins (*Strongylocentrotus nudus*) in order to restore macroalgae beds (*Saccharina japonica* var. *religiosa* and *Undaria pinnatifida*). After 8 months, urchin densities were 0.1, 3.5, and 4.2 individuals per square meter for repeat removal, single removal, or no removal (control), respectively. Removal significantly enhanced kelp colonization and growth, with average standing macroalgae biomass within these treatments of 865, 150, and 0 grams per square meter, respectively.

Supporting colonization of reef fishes using acoustic enrichment in Australia (action C)

Larval fish can use acoustic cues when selecting habitats in which to recruit (Parmentier et al. 2015). Gordon and colleagues (2019) recorded noises from a healthy reef at night and evaluated the effects of playing these recordings on attracting fish to degraded reefs (figure 3e). Fish community development on acoustically enriched coral-rubble patch reefs was significantly enhanced across all major trophic guilds relative to acoustically unmanipulated controls, with doubled overall abundance and 50% greater species richness. Gordon and colleagues (2019) suggested that coupled with active restoration of coral reefs, acoustic attraction methods may expedite recovery processes.

Building nesting mounds with protective boxes for terrapins in the United States (action C)

Animals can be supported in restored habitats by the provision of artificial refuges. Quinn and colleagues (2015) built nesting mounds with protective boxes and electrified wire to reduce nest predation by raccoons and reduce road mortalities for diamond-backed terrapin (*Malaclemys terrapin*; figure 3f). The electrified wire significantly reduced predation rates, and excavated nest boxes afforded high rates of egg survivorship and hatching success, thereby supporting terrapin populations within saltmarsh and adjacent ecosystems. Deploying these within restored saltmarsh could therefore support the survival and reproduction of animals that are functionally important or of conservation concern.

the population (i.e., eradication) can be too costly and even impossible, suppression of the population to a point of "ecological eradication" may be sufficient under certain circumstances (*sensu* Green and Grosholz 2021).

Supporting animals in restored habitats (action C). The third action involves situations in which *supporting* animals (as opposed to actively adding animals; action A) can aid system recovery, enhance ecosystem resilience, and otherwise help meet restoration objectives. Designing restoration programs that identify and support key animal species and their functions can therefore benefit habitat formers and the restored system. Because animals use specific cues when selecting habitats (e.g., host plants, prey species, conspecifics, refuges) and require a suite of resources to persist in that habitat (e.g., sufficient prey resources; Van Dyck 2012), providing species-specific cues and resources can help assist the colonization and persistence of animals that are important for ecosystem function, persistence, and resilience. In addition to designing and modifying structural components of habitats to best support animals, animals can also be encouraged to colonize restored habitats through alternative means, such as with the playback of reef sounds to encourage fish colonization of coral reefs (see box 1; Gordon et al. 2019) or playing conspecific

vocalizations to encourage bird colonization (Jones and Kress 2012). This requires knowledge of animal behavior and habitat requirements and of the species most important to improving restoration outcomes (Hale et al. 2020).

Supporting and attracting animals to benefit coastal restoration is not new; for decades, marine protected areas, for example, have been established in part to increase herbivore and predator abundance, which should, in turn, help with passive restoration of coral reefs (Topor et al. 2019) and kelp forests (Eger et al. 2020). Similarly, sea otter populations in the eastern North Pacific recovered dramatically following various conservation actions implemented decades ago, including restricting harvesting (Bodkin 2015). This recovery and the subsequent decrease in urchin populations has been important for the recovery of kelp at scale and may in fact be the preferred or most feasible action to passively restore habitat formers. Given animals can modulate ecosystem structure and function, similar acknowledgement needs to become the norm within active restorations (where management approaches such as distributing seeds, planting, and constructing habitats are implemented, as opposed to passive approaches that mitigate stressors preventing natural recovery; following Bayraktarov et al. 2016). In fact, where it is feasible, we argue that the explicit support of animals

in restoration initiatives should be the rule rather than the exception. In addition, by taking a more targeted, animal-centric point of view *both* in design and in monitoring responses coastal restoration outcomes could be enhanced, even when the objective is not explicitly related to animals (e.g., blue carbon projects; figure 2, action 3C). One interesting avenue for future work with respect to supporting (or actively adding) animals should be to systematically assess whether a diverse suite of interactions from multiple animal species could best facilitate restoration, rather than focusing only a random or favorite one or two species.

Add habitat-forming species or improve abiotic environment only (action D). The fourth action involves situations in which animals are expected to colonize and persist in restored sites following the restoration of habitat-forming species. There are examples of successful coastal restoration efforts (with respect to creating functioning wildlife habitat) in which animals have not directly been manipulated nor habitats explicitly modified to support specific animal species *per se* (although success may still depend on healthy animal populations being able to colonize). For instance, deploying oyster reefs in Moreton Bay, Australia, led to rapid enhancement of fisheries species (figure 2, action 1D; Gilby et al. 2021); broadcasting seagrass seeds in Chesapeake Bay, in the United States, recovered diverse animal communities (Orth et al. 2020); and transplanting seagrass in California, in the United States, quickly recovered fish populations (Beheshti et al. 2021). Given the likelihood of animals being integral to the long-term health and resilience of restored habitats, ongoing animal monitoring where feasible is recommended and likely beneficial.

Considerations, risks, and challenges. There are a series of important considerations, risks, and challenges when undertaking active interventions to directly manipulate or support animals. For instance, species translocations require sourcing individuals, which can be costly and ethically complex when removing individuals from wild populations (Pettorelli et al. 2018). Notably, many of the examples in this article involve supplementing already existing populations, such as the various bivalve species added to enhance seagrass or saltmarsh restoration (Derksen-Hooijberg et al. 2018, Zhang et al. 2021), with fewer risks relative to introducing new species. There is also a suite of challenges with releasing hatchery-reared animals that may, for example, perform worse in natural environments than wild conspecifics would, in part because of behavioral and cognitive differences (Lorenzen et al. 2013, Abelson et al. 2016, Näslund 2021). In addition, a detailed understanding of the system's ecology and robust predictions of the range of plausible outcomes are needed to help minimize the probability of or to manage unintended consequences (Sarrazin and Barbault 1996, Seddon et al. 2007). Adding or supporting animals can also lead to conflict with humans, such as the recovery of sea otter populations discussed above, which was unpopular in some regions because of otters competing with humans for harvested species such

as clams, crabs, and urchins (Gregar et al. 2020). There are also important ethical considerations with excluding animals. The issue of turtles overgrazing seagrass restoration sites, for example, raises unaddressed questions about the ethics of turtle exclusion, because doing so may affect access to an important food resource for a threatened species and result in animal harm through starvation. There may also be community perception issues around culling animals, particularly if it involves native species, and culling (as per introduction) is likely to trigger a different set of permitting processes from the other restoration activities.

The various considerations, risks, and challenges are highly context specific and, in practice, should be evaluated in detail on a case-by-case basis. Previous works articulate questions that scientists and managers should answer or have some knowledge about prior to directly manipulating or supporting animals in restored habitats, such as those related to selecting source populations, the plausible implications for the wider ecosystem, and various ethical, permitting, and legal considerations (e.g., Sarrazin and Barbault 1996, Seddon et al. 2007, Armstrong and Seddon 2008, Houde et al. 2015, Nogués-Bravo et al. 2016, Berger-Tal et al. 2020). Resources also exist to guide practitioners with respect to translocations and species introductions, including general guidelines (e.g., IUCN 2013) and perspective reports with case studies (e.g., Soorae 2021), and these are also informative when attempting to attract and support animals. Ultimately, however, in many cases there are still gaps in our understanding of the ecology, which will lead to uncertainties in how and whether the manipulation or support of animals will influence the restoration trajectory. This requires ecological research as well as research on the cost-effectiveness of various actions.

Applying structured decision-making to incorporate animals in restoration

We have argued for the benefits of explicit consideration of animals in restoration, and our argument is supported by a strong evidence base. One challenge now will be if, when, and how to scale up experimental restoration that has demonstrated how manipulating and supporting animals can aid restoration. A key impediment to explicitly including animals in restoration at scale will be demonstrating that the benefits exceed the costs (e.g., time, resources, ethical and legal requirements) and the risks (e.g., unintended impacts stemming from a lack of ecological knowledge on key processes, and uncertainty in outcomes). One useful approach for assessing the case for restoration is structured decision-making (SDM), a systematic and transparent approach to natural resource management. SDM is highly amenable to involving stakeholders in decision-making processes and is gaining traction in ecosystem restoration (Guerrero et al. 2017). SDM is based on decision theory and risk analysis and typically has seven key steps, as was articulated for kelp restoration in Gleason and colleagues (2021). We show how the SDM framework can apply to the question of whether to explicitly include

Table 1. Applying a simplified structured decision-making (SDM) process to three hypothetical case studies with varying objective-dependent animal roles.

Key steps	Hypothetical case studies		
1. Problem formulation	1. Restore degraded saltmarsh to reference condition.	2. Enhance fisheries via coral reef restoration.	3. Expand seagrass area for carbon sequestration.
2. Set clear objectives	Reach similar floral and faunal species richness to reference sites within 7 years.	Increase fisheries productivity by 30% within 10 years.	Double carbon stock within 15 years.
3. Identify actions, including parameterization of costs and likelihoods of achieving objectives	A: Translocate supporting animals (e.g., bivalves) B: Plant saltmarsh C: Remove exotic or damaging species (e.g., unguates)	A: Translocate corals B: Breed and release juvenile fisheries species C: Translocate supporting animals (e.g., algae grazing gastropods)	A: Plant seagrass B: Translocate supporting animals (e.g., bivalves, algae grazers) C: Exclude hindering animals (e.g., herbivores)
4. Estimate consequences. This text articulates the prediction of outcomes for one of the actions identified in step 3. A similar approach would be taken for all actions, and the estimates used to inform 5.	Generate predictions for outcomes of actions A, B, and C. For example, using inferences from studies on mussel translocations (e.g., Derksen-Hooijberg et al. 2018) and meta-analyses that compare animal populations between restored and reference ecosystems (e.g., Sievers et al. 2018). For action A, we predict the benefits of translocating mussels at various densities on drought tolerance and growth of saltmarsh vegetation, and relate this to biodiversity benefits. Hypothetical prediction for A: Richness surpasses 80% of reference levels within a 7-year timeframe.	Generate predictions for outcomes of actions A, B, and C. E.g., using numerical fisheries models that predict fish production from coral reef condition (Rogers et al. 2018). For action C, we predict the benefits of translocating algal grazers to the survival and growth of newly transplanted corals, and the subsequent outcomes for fish production. Hypothetical prediction for C: Grazers will enhance coral survival by 30%–50%, leading to a 2–3 tons per hectare increase in fish biomass after 10 years.	Generate predictions for outcomes of actions A, B, and C. E.g., using models that predict CO ₂ capture from restored seagrass extent (e.g., Duarte et al. 2013b). For action C, we predict the exclusion of herbivores promotes seagrass growth (e.g., Burkholder et al. 2013) and, we can link this to predicted carbon sequestration and stocks. Hypothetical prediction for C: Exclusion fences will eliminate grazing by turtles, tripling seagrass biomass, and leading to a doubling of carbon stock within 15 years.
5. Evaluate trade-offs	Evaluate trade-offs across alternative actions from 3 and 4 to determine which one or more best meets that objective (dependent on importance, cost, benefit, degree of certainty, risk, constraints, etc.).		
6. Make decisions	Make decision on the basis of the information gained in steps 3–5, by assessing which options are most likely to achieve the desired goals and objectives set out in steps 1 and 2 within the constraints of the project (budget, time, feasibility, etc.).		
7. Act, monitor and learn	A: Given the ease at which manipulations occur, conduct replicated experiment to examine the effect of mussel addition (and density effects) on marsh growth and survival. Where possible, extend monitoring to other species in the food web. Measure biodiversity across restored and reference sites.	C: Monitor grazers, algal growth, and coral growth and survival. Contrast outcomes with unmanipulated areas. Develop models to identify optimal grazer densities (both densities the system can support, and those that maintain coral survival and growth). Continue to monitor fisheries productivity.	C: Monitor seagrass growth and survival. Once seagrass is established, remove cages to allow potential positive species interactions. Maintain monitoring; if overgrazing continues, refencing may be needed. Quantify carbon stock across natural and restored seagrass meadows.

Note: The SDM approach was based on steps in Gleason and colleagues (2021).

or exclude dependent fauna into restoration of habitat-forming marine and coastal species using three case studies in table 1. This table is intended to be indicative and would require further development to guide restoration science and practice. In reality, applying SDM to restoration requires greater detail, a more holistic understanding of the system, and input from various stakeholders, and ultimately, the best approach may involve a combination of actions (e.g., Gleason et al. 2021).

Conclusions

Ongoing destruction and degradation of coastal habitats and the subsequent loss of service benefits to people have necessitated accelerating restoration efforts. But restoration without animals may not achieve the desired outcome. Although there are many impediments to effective coastal restoration, identifying when, why, and how to directly manipulate or

support animals can lead to substantial improvements in outcomes for habitat-forming species and ecosystem services. By outlining how animals play important roles across different restoration objectives, articulating key contexts in which animals can be explicitly incorporated in coastal restoration, and illustrating these ideas with practical case studies, we hope to encourage scientists and managers to better incorporate animals into coastal restoration planning, implementation, and monitoring.

Acknowledgments

We thank three anonymous reviewers for their insightful comments that greatly improved the manuscript. We acknowledge funding support from The Global Wetlands Project, supported by a charitable organization that neither seeks nor permits publicity for its efforts. MS was funded by Australian Research Council (ARC) Discovery Early Career

Researcher Award no. DE220100079; MIS was supported by a Julius Career Award from CSIRO; CB was funded by ARC Future Fellowship no. FT210100792.

Disclosure statement

The authors declare no conflicts of interest.

References cited

- Abelson A, Obolski U, Regoniel P, Hadany L. 2016. Restocking herbivorous fish populations as a social-ecological restoration tool in coral reefs. *Frontiers in Marine Science* 3: 138.
- Abelson A et al. 2020. Challenges for restoration of coastal marine ecosystems in the Anthropocene. *Frontiers in Marine Science* 7: 544105.
- Angeler DG, Chow-Fraser P, Hanson MA, Sánchez-Carrillo S, Zimmer KD. 2003. Biomanipulation: A useful tool for freshwater wetland mitigation? *Freshwater Biology* 48: 2203–2213.
- Angelini C, Griffin JN, van de Koppel J, Lamers LP, Smolders AJ, Derksen-Hooijberg M, van der Heide T, Silliman BR. 2016. A keystone mutualism underpins resilience of a coastal ecosystem to drought. *Nature Communications* 7: 12473.
- Armstrong DP, Seddon PJ. 2008. Directions in reintroduction biology. *Trends in Ecology and Evolution* 23: 20–25.
- Bayraktarov E, Saunders MI, Abdullah S, Mills M, Behr J, Possingham HP, Mumby PJ, Lovelock CE. 2016. The cost and feasibility of marine coastal restoration. *Ecological Applications* 26: 1055–1074.
- Bayraktarov E, Brisbane S, Hagger V, Smith CS, Wilson KA, Lovelock CE, Gillies C, Steven AD, Saunders MI. 2020. Priorities and motivations of marine coastal restoration research. *Frontiers in Marine Science* 7: 484.
- Beheshti KM, Williams SL, Boyer KE, Endris C, Clemons A, Grimes T, Wasson K, Hughes BB. 2021. Rapid enhancement of multiple ecosystem services following the restoration of a coastal foundation species. *Ecological Applications* 32: e02466.
- Berger-Tal O, Blumstein D, Swaisgood RR. 2020. Conservation translocations: A review of common difficulties and promising directions. *Animal Conservation* 23: 121–131.
- Biggs BC. 2013. Harnessing natural recovery processes to improve restoration outcomes: An experimental assessment of sponge-mediated coral reef restoration. *PLOS ONE* 8: e64945.
- Bodkin JL. 2015. Historic and contemporary status of sea otters in the North Pacific. Pages 43–61 in Larson SE, Bodkin JL, VanBlaricom GR, eds. *Sea Otter Conservation*. Elsevier.
- Buelow CA, Connolly RM, Turschwell MP, Adame MF, Ahmadia GN, Andradi-Brown DA, Bunting P, Canty SW, Dunic JC, Friess DA. 2022. Ambitious global targets for mangrove and seagrass recovery. *Current Biology* 32: 1641–1649.e3.
- Burkholder DA, Heithaus MR, Fourqurean JW, Wirsing A, Dill LM. 2013. Patterns of top-down control in a seagrass ecosystem: Could a roving apex predator induce a behaviour-mediated trophic cascade? *Journal of Animal Ecology* 82: 1192–1202.
- Christianen MJA et al. 2014. Habitat collapse due to overgrazing threatens turtle conservation in marine protected areas. *Proceedings of the Royal Society B* 281: 20132890.
- Cooke SJ, Bennett JR, Jones HP. 2019. We have a long way to go if we want to realize the promise of the "Decade on ecosystem restoration." *Conservation Science and Practice* 1: e129.
- Cowx IG, Gerdeaux D. 2004. The effects of fisheries management practises on freshwater ecosystems. *Fisheries Management and Ecology* 11: 145–151.
- Cunningham F, Young H, Sevilla C, Carrión V, Fessl B. 2011. A trial translocation of the critically endangered mangrove finch: Conservation management to prevent the extinction of Darwin's rarest finch. *Galapagos Report* 2012: 174–179.
- Dale P, Knight J, Dwyer P. 2014. Mangrove rehabilitation: A review focusing on ecological and institutional issues. *Wetlands Ecology and Management* 22: 587–604.
- Davidson KE, Fowler MS, Skov MW, Doerr SH, Beaumont N, Griffin JN. 2017. Livestock grazing alters multiple ecosystem properties and services in salt marshes: A meta-analysis. *Journal of Applied Ecology* 54: 1395–1405.
- Davis RW, Bodkin JL, Coletti HA, Monson DH, Larson SE, Carswell LP, Nichol LM. 2019. Future directions in sea otter research and management. *Frontiers in Marine Science* 5: 510.
- Derksen-Hooijberg M, Angelini C, Lamers LPM, Borst A, Smolders A, Hoogveld JRH, de Paoli H, van de Koppel J, Silliman BR, van der Heide T. 2018. Mutualistic interactions amplify saltmarsh restoration success. *Journal of Applied Ecology* 55: 405–414.
- Díaz SM, Settele J, Brondizio E, Ngo H, Guèze M, Agard J, Arneth A, Balvanera P, Brauman K, Butchart S. 2019. The Global Assessment Report on Biodiversity and Ecosystem Services: Summary for Policy Makers. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
- Duarte CM, Losada IJ, Hendriks IE, Mazarrasa I, Marbà N. 2013. The role of coastal plant communities for climate change mitigation and adaptation. *Nature Climate Change* 3: 961–968.
- Duarte CM, Sintes T, Marbà N. 2013. Assessing the CO₂ capture potential of seagrass restoration projects. *Journal of Applied Ecology* 50: 1341–1349.
- Dunic JC, Brown CJ, Connolly RM, Turschwell MP, Côté IM. 2021. Long-term declines and recovery of meadow area across the world's seagrass bioregions. *Global Change Biology* 27: 4096–4109.
- Eger AM, Marzinelli E, Gribben P, Johnson CR, Layton C, Steinberg PD, Wood G, Silliman BR, Verges A. 2020. Playing to the positives: Using synergies to enhance kelp forest restoration. *Frontiers in Marine Science* 7: 544.
- Eger A, Marzinelli E, Christie H, Fagerli C, Fujita D, Gonzalez A, Hong S, Kim J, Lee L, McHugh T. 2022. Global kelp forest restoration: Past lessons, present status, and future directions. *Biological Reviews* 97: 1449–1475.
- Ellison AM, Farnsworth EJ, Twilley RR. 1996. Facultative mutualism between red mangroves and root-fouling sponges in Belizean mangal. *Ecology* 77: 2431–2444.
- Foster E, Watson J, Lemay MA, Tinker MT, Estes JA, Piercey R, Henson L, Ritland C, Miscampbell A, Nichol L. 2021. Physical disturbance by recovering sea otter populations increases eelgrass genetic diversity. *Science* 374: 333–336.
- Fourqurean JW, Manuel SA, Coates KA, Massey SC, Kenworthy WJ. 2019. Decadal monitoring in Bermuda shows a widespread loss of seagrasses attributable to overgrazing by the green sea turtle *Cheloniemydas*. *Estuaries and Coasts* 42: 1524–1540.
- Gagnon K, Rinde E, Bengil EG, Carugati I, Christianen MJ, Danovaro R, Gambi C, Govers LL, Kipson S, Meysick L. 2020. Facilitating foundation species: The potential for plant-bivalve interactions to improve habitat restoration success. *Journal of Applied Ecology* 57: 1161–1179.
- Gangal M, Gafoor A-B, D'Souza E, Kelkar N, Karkarey R, Marbà N, Arthur R, Alcoverro T. 2021. Sequential overgrazing by green turtles causes archipelago-wide functional extinctions of seagrass meadows. *Biological Conservation* 260: 109195.
- Gann GD, McDonald T, Walder B, Aronson J, Nelson CR, Jonson J, Hallett JG, Eisenberg C, Guariguata MR, Liu J. 2019. International principles and standards for the practice of ecological restoration. *Restoration Ecology* 27: S1–S46.
- Gedan KB, Silliman BR. 2009. Using facilitation theory to enhance mangrove restoration. *AMBIO: A Journal of the Human Environment* 38: 109–109.
- Gilby BL, Olds AD, Brown CJ, Connolly RM, Henderson CJ, Maxwell PS, Schlacher TA. 2021. Applying systematic conservation planning to improve the allocation of restoration actions at multiple spatial scales. *Restoration Ecology* 29: e13403.
- Gleason M, Caselle J, Heady W, Saccomanno V, Zimmerman J, McHugh T. 2021. A Structured Approach for Kelp Restoration and Management Decisions in California. The Nature Conservancy.

- Gordon TA, Radford AN, Davidson IK, Barnes K, McCloskey K, Nedelec SL, Meekan MG, McCormick MI, Simpson SD. 2019. Acoustic enrichment can enhance fish community development on degraded coral reef habitat. *Nature Communications* 10: 5414.
- Grabowski JH. 2004. Habitat complexity disrupts predator-prey interactions but not the trophic cascade on oyster reefs. *Ecology* 85: 995–1004.
- Green SJ, Grosholz ED. 2021. Functional eradication as a framework for invasive species control. *Frontiers in Ecology and the Environment* 19: 98–107.
- Gregg EJ, Christensen V, Nichol L, Martone RG, Markel RW, Watson JC, Harley CD, Pakhomov EA, Shurin JB, Chan KM. 2020. Cascading social-ecological costs and benefits triggered by a recovering keystone predator. *Science* 368: 1243–1247.
- Griffiths LL, Connolly RM, Brown CJ. 2020. Critical gaps in seagrass protection reveal the need to address multiple pressures and cumulative impacts. *Ocean and Coastal Management* 183: 104946.
- Guerrero AM, Shoo L, Iacona G, Standish RJ, Catterall CP, Rumpff L, De Bie K, White Z, Matzek V, Wilson KA. 2017. Using structured decision-making to set restoration objectives when multiple values and preferences exist. *Restoration Ecology* 25: 858–865.
- Hale R, Mac Nally R, Blumstein DT, Swearer SE. 2019. Evaluating where and how habitat restoration is undertaken for animals. *Restoration Ecology* 27: 775–781.
- Hale R, Blumstein DT, Mac Nally R, Swearer SE. 2020. Harnessing knowledge of animal behavior to improve habitat restoration outcomes. *Ecosphere* 11: e03104.
- Halpern BS, Silliman BR, Olden JD, Bruno JP, Bertness MD. 2007. Incorporating positive interactions in aquatic restoration and conservation. *Frontiers in Ecology and the Environment* 5: 153–160.
- Halpern BS, Frazier M, Afflerbach J, Lowndes JS, Micheli F, O'Hara C, Scarborough C, Selkoe KA. 2019. Recent pace of change in human impact on the world's ocean. *Scientific Reports* 9: 11609.
- He Q, Silliman BR. 2019. Climate change, human impacts, and coastal ecosystems in the Anthropocene. *Current Biology* 29: R1021–R1035.
- Hensel MJ, Silliman BR, van de Koppel J, Hensel E, Sharp SJ, Crotty SM, Byrnes JE. 2021. A large invasive consumer reduces coastal ecosystem resilience by disabling positive species interactions. *Nature Communications* 12: 1–10.
- Houde ALS, Garner SR, Neff BD. 2015. Restoring species through reintroductions: Strategies for source population selection. *Restoration Ecology* 23: 746–753.
- Hughes BB, Hammerstrom KK, Grant NE, Hoshijima U, Eby R, Wasson K. 2016. Trophic cascades on the edge: Fostering seagrass resilience via a novel pathway. *Oecologia* 182: 231–241.
- [IUCN] International Union for Conservation of Nature. 2013. Guidelines for Reintroductions and Other Conservation Translocations. IUCN.
- Jones ME, Davidson N. 2016. Applying an animal-centric approach to improve ecological restoration. *Restoration Ecology* 24: 836–842.
- Jones HP, Kress SW. 2012. A review of the world's active seabird restoration projects. *The Journal of Wildlife Management* 76: 2–9.
- Leber KM. 2013. Marine fisheries enhancement, coming of age in the new millennium. Pages 1139–1157 in Christou P, Savin R, Costa-Pierce BA, Misztal I, Whitelaw CBA, eds. *Sustainable Food Production*. Springer.
- Lee SY, Hamilton S, Barbier EB, Primavera J, Lewis RR. 2019. Better restoration policies are needed to conserve mangrove ecosystems. *Nature Ecology and Evolution* 3: 870–872.
- Lewis R. III. 2010. Mangrove field of dreams: If we build it, will they come? *Wetland Science and Practice* 27: 15–18.
- Lewis LS, Anderson TW. 2012. Top-down control of epifauna by fishes enhances seagrass production. *Ecology* 93: 2746–2757.
- Lorenzen K, Leber KM, Blankenship HL. 2010. Responsible approach to marine stock enhancement: An update. *Reviews in Fisheries Science* 18: 189–210.
- Lorenzen K, Agnalt A-L, Blankenship HL, Hines AH, Leber KM, Loneragan NR, Taylor MD. 2013. Evolving context and maturing science: Aquaculture-based enhancement and restoration enter the marine fisheries management toolbox. *Reviews in Fisheries Science* 21: 213–221.
- Lotze HK, Coll M, Magera AM, Ward-Paige C, Airoldi L. 2011. Recovery of marine animal populations and ecosystems. *Trends in Ecology and Evolution* 26: 595–605.
- McDonald T, Gann GD, Jonson J, Dixon KW, Aronson J, Declerck K, Hallett J, Keenleyside K, Nelson C, Walder B. 2016. *International Standards for the Practice of Ecological Restoration*. Society for Ecological Restoration.
- Miller KI, Shears NT. 2022. The efficiency and effectiveness of different sea urchin removal methods for kelp forest restoration. *Restoration Ecology*: e13754.
- Morris RL, Hale R, Strain EM, Reeves SE, Vergés A, Marzinelli EM, Layton C, Shelamoff V, Graham TD, Chevalier M. 2020. Key principles for managing recovery of kelp forests through restoration. *BioScience* 70: 688–698.
- Murray NJ, Worthington TA, Bunting P, Duce S, Hagger V, Lovelock CE, Lucas R, Saunders MI, Sheaves M, Spalding M. 2022. High-resolution mapping of losses and gains of Earth's tidal wetlands. *Science* 376: 744–749.
- Näslund J. 2021. Reared to become wild-like: Addressing behavioral and cognitive deficits in cultured aquatic animals destined for stocking into natural environments: A critical review. *Bulletin of Marine Science* 97: 489–538.
- Neil RC, Humphrey C, Bourne DG, Heyward A. 2021. Co-culture with grazers can improve survival and growth of multiple coral species. *Aquaculture* 544: 737095.
- Nogués-Bravo D, Simberloff D, Rahbek C, Sanders NJ. 2016. Rewilding is the new Pandora's box in conservation. *Current Biology* 26: R87–R91.
- Obolski U, Hadany L, Abelson A. 2016. Potential contribution of fish restocking to the recovery of deteriorated coral reefs: An alternative restoration method? *PeerJ* 4: e1732.
- Orth RJ, Lefcheck JS, McGlathery KS, Aoki L, Luckenbach MW, Moore KA, Oreska MP, Snyder R, Wilcox DJ, Lusk B. 2020. Restoration of seagrass habitat leads to rapid recovery of coastal ecosystem services. *Science Advances* 6: eabc6434.
- Palmer MA, Ambrose RF, Poff NL. 1997. Ecological theory and community restoration ecology. *Restoration Ecology* 5: 291–300.
- Parmentier E, Berten L, Rigo P, Aubrun F, Nedelec S, Simpson SD, Lecchini D. 2015. The influence of various reef sounds on coral-fish larvae behaviour. *Journal of Fish Biology* 86: 1507–1518.
- Perillo G, Wolanski E, Cahoon DR, Hopkinson CS. 2018. *Coastal Wetlands: An Integrated Ecosystem Approach*. Elsevier.
- Perring MP, Standish RJ, Price JN, Craig MD, Erickson TE, Ruthrof KX, Whiteley AS, Valentine LE, Hobbs RJ. 2015. Advances in restoration ecology: Rising to the challenges of the coming decades. *Ecosphere* 6: 1–25.
- Pettorelli N, Barlow J, Stephens PA, Durant SM, Connor B, Schulte to Bühne H, Sandom CJ, Wentworth J, du Toit JT. 2018. Making rewilding fit for policy. *Journal of Applied Ecology* 55: 1114–1125.
- Poore AG, Campbell AH, Coleman RA, Edgar GJ, Jormalainen V, Reynolds PL, Sotka EE, Stachowicz JJ, Taylor RB, Vanderklift MA. 2012. Global patterns in the impact of marine herbivores on benthic primary producers. *Ecology Letters* 15: 912–922.
- Qiu D, Cui B, Ma X, Yan J, Cai Y, Xie T, Gao F, Wang F, Sui H, Bai J. 2021. Reciprocal facilitation between annual plants and burrowing crabs: Implications for the restoration of degraded saltmarshes. *Journal of Ecology* 109: 1828–1841.
- Quinn DP, Kaylor SM, Norton TM, Buhlmann KA. 2015. Nesting mounds with protective boxes and an electric wire as tools to mitigate diamond-backed terrapin (*Malaclemys terrapin*) nest predation. *Herpetological Conservation and Biology* 10: 969–977.
- Reeves SE, Renzi JJ, Fobert EK, Silliman BR, Hancock B, Gillies CL. 2020. Facilitating better outcomes: How positive species interactions can improve oyster reef restoration. *Frontiers in Marine Science* 7: 656.
- Renzi JJ, He Q, Silliman BR. 2019. Harnessing positive species interactions to enhance coastal wetland restoration. *Frontiers in Ecology and Evolution* 7: 131.

- Rogers A, Blanchard JL, Mumby PJ. 2018. Fisheries productivity under progressive coral reef degradation. *Journal of Applied Ecology* 55: 1041–1049.
- Rotjan RD, Lewis SM. 2006. Parrotfish abundance and selective corallivory on a Belizean coral reef. *Journal of Experimental Marine Biology and Ecology* 335: 292–301.
- Sarrazin F, Barbault R. 1996. Reintroduction: Challenges and lessons for basic ecology. *Trends in Ecology and Evolution* 11: 474–478.
- Saunders MI, Doropoulos C, Bayraktarov E, Babcock RC, Gorman D, Eger AM, Vozzo ML, Gillies CL, Vanderklift MA, Steven AD. 2020. Bright spots in coastal marine ecosystem restoration. *Current Biology* 30: R1500–R1510.
- Seddon PJ, Armstrong DP, Maloney RE. 2007. Developing the science of reintroduction biology. *Conservation Biology* 21: 303–312.
- Seddon PJ, Griffiths CJ, Soorae PS, Armstrong DP. 2014. Reversing defaunation: Restoring species in a changing world. *Science* 345: 406–412.
- Selig ER, Hole DG, Allison EH, Arkema KK, McKinnon MC, Chu J, de Sherbinin A, Fisher B, Glew L, Holland MB. 2019. Mapping global human dependence on marine ecosystems. *Conservation Letters* 12: e12617.
- Seraphim MJ, Sloman KA, Alexander ME, Janetski N, Jompa J, Ambo-Rappe R, Snellgrove D, Mars F, Harborne AR. 2020. Interactions between coral restoration and fish assemblages: Implications for reef management. *Journal of Fish Biology* 97: 633–655.
- Sharma R, Swearer SE, Morris RL, Strain EM. 2021. Testing the efficacy of sea urchin exclusion methods for restoring kelp. *Marine Environmental Research* 170: 105439.
- Shaver EC, Silliman BR. 2017. Time to cash in on positive interactions for coral restoration. *PeerJ* 5: e3499.
- Shaver EC, Burkepille DE, Silliman BR. 2018. Local management actions can increase coral resilience to thermally induced bleaching. *Nature Ecology and Evolution* 2: 1075–1079.
- Shaver EC, McLeod E, Hein MY, Palumbi SR, Quigley K, Vardi T, Mumby PJ, Smith D, Montoya-Maya P, Muller EM. 2022. A roadmap to integrating resilience into the practice of coral reef restoration. *Global Change Biology* 28: 4751–4764.
- Sheaves M, Waltham N, Benham C, Bradley M, Mattone C, Diedrich A, Sheaves J, Sheaves A, Hernandez S, Dale P. 2021. Restoration of marine ecosystems: Understanding possible futures for optimal outcomes. *Science of the Total Environment* 796: 148845.
- Sievers M, Hale R, Parris KM, Swearer SE. 2018. Impacts of human-induced environmental change in wetlands on aquatic animals. *Biological Reviews* 93: 529–554.
- Smith NE, Wilcox C, Lessmann JM. 2009. Fiddler crab burrowing affects growth and production of the white mangrove (*Laguncularia racemosa*) in a restored Florida coastal marsh. *Marine Biology* 156: 2255–2266.
- Soorae PS. 2021. Global Conservation Translocation Perspectives: 2021. International Union for Conservation of Nature.
- Sundermann A, Stoll S, Haase P. 2011. River restoration success depends on the species pool of the immediate surroundings. *Ecological Applications* 21: 1962–1971.
- Suykerbuyk W, Bouma TJ, Van Der Heide T, Faust C, Govers LL, Giesen WB, De Jong DJ, Van Katwijk MM. 2012. Suppressing antagonistic bioengineering feedbacks doubles restoration success. *Ecological Applications* 22: 1224–1231.
- Swan KD, McPherson JM, Seddon PJ, Moehrenschrager A. 2016. Managing marine biodiversity: The rising diversity and prevalence of marine conservation translocations. *Conservation Letters* 9: 239–251.
- Taylor MD, Chick RC, Lorenzen K, Agnalt A-L, Leber KM, Blankenship HL, Vander Haegen G, Loneragan NR. 2017. Fisheries enhancement and restoration in a changing world. *Fisheries Research* 186: 407–412.
- Tol SJ, Jarvis JC, York PH, Congdon BC, Coles RG. 2021. Mutualistic relationships in marine angiosperms: Enhanced germination of seeds by mega-herbivores. *Biotropica* 53: 1535–1545.
- Topor ZM, Rasher DB, Duffy JE, Brandl SJ. 2019. Marine protected areas enhance coral reef functioning by promoting fish biodiversity. *Conservation Letters* 12: e12638.
- Valdez SR, Zhang YS, van der Heide T, Vanderklift MA, Tarquinio F, Orth RJ, Silliman BR. 2020. Positive ecological interactions and the success of seagrass restoration. *Frontiers in Marine Science* 7: 91.
- van der Heide T, Govers LL, de Fouw J, Olff H, van der Geest M, van Katwijk MM, Piersma T, van de Koppel J, Silliman BR, Smolders AJ. 2012. A three-stage symbiosis forms the foundation of seagrass ecosystems. *Science* 336: 1432–1434.
- Van Dyck H. 2012. Changing organisms in rapidly changing anthropogenic landscapes: The significance of the “Umwelt”-concept and functional habitat for animal conservation. *Evolutionary Applications* 5: 144–153.
- van Katwijk MM, Thorhaug A, Marbà N, Orth RJ, Duarte CM, Kendrick GA, Althuisen IH, Balestri E, Bernard G, Cambridge ML. 2016. Global analysis of seagrass restoration: The importance of large-scale planting. *Journal of Applied Ecology* 53: 567–578.
- Villanueva RD, Edwards AJ, Bell JD. 2010. Enhancement of grazing gastropod populations as a coral reef restoration tool: Predation effects and related applied implications. *Restoration Ecology* 18: 803–809.
- Waltham NJ, Elliott M, Lee SY, Lovelock C, Duarte CM, Buelow C, Simenstad C, Nagelkerken I, Claassens L, Wen CK. 2020. UN decade on ecosystem restoration 2021–2030: What chance for success in restoring coastal ecosystems? *Frontiers in Marine Science* 7: 71.
- Watanuki A, Aota T, Otsuka E, Kawai T, Iwahashi Y, Kuwahara H, Fujita D. 2010. Restoration of kelp beds on an urchin barren: Removal of sea urchins by citizen divers in southwestern Hokkaido. *Bulletin of Fisheries Research Agency* 32: 83–87.
- Wendländer NS, Lange T, Connolly RM, Kristensen E, Pearson RM, Valdemarsen T, Flindt MR. 2020. Assessing methods for restoring seagrass (*Zostera muelleri*) in Australia's subtropical waters. *Marine and Freshwater Research* 71: 996–1005.
- Williams BA, Watson JE, Beyer HL, Klein CJ, Montgomery J, Runtz RK, Roberson LA, Halpern BS, Grantham HS, Kuempel CD. 2021. The global rarity of intact coastal regions. *Conservation Biology* 36: e13874.
- Wood G, Marzinelli EM, Coleman MA, Campbell AH, Santini NS, Kajlich L, Verdura J, Wodak J, Steinberg PD, Vergés A. 2019. Restoring subtidal marine macrophytes in the Anthropocene: Trajectories and future-proofing. *Marine and Freshwater Research* 70: 936–951.
- Young TP, Schwartz MW. 2019. The decade on ecosystem restoration is an impetus to get it right. *Conservation Science and Practice* 1: e145.
- Zhang YS, Gittman RK, Donaher SE, Trachtenberg SN, Van Der Heide T, Silliman BR. 2021. Inclusion of intra- and interspecific facilitation expands the theoretical framework for seagrass restoration. *Frontiers in Marine Science* 8: 645673.

Michael Sievers (m.sievers@griffith.edu.au), Christopher J. Brown (chris.brown@griffith.edu.au), Christina A. Buelow (c.buelow@griffith.edu.au), Andria Ostrowski (andria.ostrowski@griffithuni.edu.au), Mischa P. Turschwell (m.turschwell@griffith.edu.au), and Rod M. Connolly (r.connolly@griffith.edu.au) are affiliated with the Coastal and Marine Research Centre, the Australian Rivers Institute, and with the School of Environment and Science at Griffith University, in Gold Coast, Queensland, Australia. Robin Hale (rob.hale@delwp.vic.gov.au) is affiliated with the Arthur Rylah Institute for Environmental Research at the Department of Environment, Land, Water, and Planning, in Heidelberg, Victoria, Australia. Megan I. Saunders (megan.saunders@csiro.au) is affiliated with CSIRO Oceans and Atmosphere, in St Lucia, Brisbane, Australia. Brian R. Silliman (brian.silliman@duke.edu) and Stephanie R. Valdez (stephanie.valdez@duke.edu) are affiliated with the Nicholas School of the Environment at Duke University, in Beaufort, North Carolina, in the United States. Stephen E. Swearer (s.swearer@unimelb.edu.au) is affiliated with the National Centre for Coasts and Climate and with the School of Biosciences at the University of Melbourne, in Parkville, Victoria, Australia.