

Mahler Discrete Residues and Summability for Rational Functions

Carlos E. Arreche*

Department of Mathematical Sciences
The University of Texas at Dallas
Richardson, Texas, USA
arreche@utdallas.edu

Yi Zhang†

Department of Foundational Mathematics
Xi'an Jiaotong-Liverpool University
Suzhou, CHINA
Yi.Zhang03@xjtlu.edu.cn

ABSTRACT

We construct Mahler discrete residues for rational functions and show that they comprise a complete obstruction to the Mahler summability problem of deciding whether a given rational function $f(x)$ is of the form $g(x^p) - g(x)$ for some rational function $g(x)$ and an integer $p > 1$. This extends to the Mahler case the analogous notions, properties, and applications of discrete residues (in the shift case) and q -discrete residues (in the q -difference case) developed by Chen and Singer. Along the way we define several additional notions that promise to be useful for addressing related questions involving Mahler difference fields of rational functions, including in particular telescoping problems and problems in the (differential) Galois theory of Mahler difference equations.

CCS CONCEPTS

• Computing methodologies → Algebraic algorithms.

KEYWORDS

Mahler operator, difference fields, difference equations, partial fractions, discrete residues, summability, creative telescoping

ACM Reference Format:

Carlos E. Arreche and Yi Zhang. 2022. Mahler Discrete Residues and Summability for Rational Functions. In *Proceedings of the 2022 International Symposium on Symbolic and Algebraic Computation (ISSAC '22)*, July 4–7, 2022, Villeneuve-d'Ascq, France. ACM, New York, NY, USA, 9 pages. <https://doi.org/10.1145/3476446.3536186>

1 INTRODUCTION

Continuous residues are fundamental and crucial tools in complex analysis, and have extensive and compelling applications in combinatorics [16]. In the last decade, a theory of (q -)discrete residues was proposed in [14] for the study of telescoping problems, which has found essential applications in several other closely related

problems (see [3, 4, 11, 19] for some examples). A theory of residues for skew rational functions was developed in [9], and then extended to Ore polynomials and applied to linearized Reed-Solomon codes in [10]. The elliptic orbit residues defined in [18] have applications in the combinatorial study of walks in the quarter plane. We propose here a theory of Mahler discrete residues aimed at bringing to the Mahler case the successes of these earlier notions of residues.

Let $\mathbb{K}$ be a field of characteristic zero and $\mathbb{K}(x)$ be the field of rational functions in an indeterminate x over $\mathbb{K}$. Fix an integer $p \geq 2$. We study the *Mahler summability problem for rational functions*: given $f(x) \in \mathbb{K}(x)$, decide effectively whether $f(x) = g(x^p) - g(x)$ for some $g(x) \in \mathbb{K}(x)$; if so, we say $f(x)$ is *Mahler summable*.

The motivation to study Mahler difference equations comes from several directions: they find applications in automata theory (automatic sequences), transcendence, and number theory, to name a few. We refer to [15] for more details, and also for an altogether different approach to the Mahler summability problem: the algorithm of [15, §3] computes all the rational solutions to any linear Mahler equation. Thus with this one can decide, in particular, whether any given $f(x) \in \mathbb{K}(x)$ is Mahler summable by computing (or showing non-existence of) a certificate $g(x) \in \mathbb{K}(x)$ such that $f(x) = g(x^p) - g(x)$.

Our goal here is different: we wish to construct a complete obstruction to Mahler summability. Let us elaborate. The image of the $\mathbb{K}$ -linear map $\Delta : g(x) \mapsto g(x^p) - g(x)$ is the kernel of some other $\mathbb{K}$ -linear map (call it ∇) — but what is it? Determining such a ∇ explicitly is algorithmically desirable because it allows to decide the Mahler summability of $f(x) \in \mathbb{K}(x)$ without computing the certificate $g(x) \in \mathbb{K}(x)$, whose computation is often in practice both expensive and not strictly necessary (cf. [5, §1 and Table 1], [6, §1 and Table 1], and [7, §1]). We construct such a ∇ explicitly in Section 4.4, in terms of our new notion of *Mahler discrete residues* for rational functions, and prove in Section 4.3:

MAIN THEOREM. $f(x) \in \mathbb{K}(x)$ is Mahler summable if and only if all of the Mahler discrete residues of $f(x)$ are zero.

The discrete and q -discrete residues developed in [14] comprise complete obstructions to the summability problem of deciding whether $f(x) = g(x+1) - g(x)$ for some $g(x) \in \mathbb{K}(x)$ and the q -summability problem of deciding whether $f(x) = g(qx) - g(x)$ for some $g(x) \in \mathbb{K}(x)$ and $q \in \mathbb{K}$ neither zero nor a root of unity, respectively. This theoretical property of (q -)discrete residues is precisely what enables their applications to the telescoping problems considered in [14] and their indispensable role in the development of the algorithms in [3, 4]. We envision analogous applications of Mahler discrete residues to telescoping problems and in the development of algorithms to compute (differential) Galois groups for Mahler difference equations.

*C.E. Arreche was partially supported by NSF grant CCF-1815108.

†Y. Zhang was supported by XJTLU Research Development Fund No. RDF-20-01-12, the NSFC Young Scientist Fund No. 12101506, and the Natural Science Foundation of the Jiangsu Higher Education Institutions of China No. 21KJB110032.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](https://permissions.acm.org).

ISSAC '22, July 4–7, 2022, Villeneuve-d'Ascq, France

© 2022 Association for Computing Machinery.

ACM ISBN 978-1-4503-8688-3/22/07...\$15.00

<https://doi.org/10.1145/3476446.3536186>

Our strategy is inspired by that of [14] (but see Remark 2.19): we utilize the coefficients in the partial fraction decomposition of $f(x)$ to construct an aspiring certificate $g(x) \in \mathbb{K}(x)$ such that, for

$$\tilde{f}(x) := f(x) + (g(x^p) - g(x)), \quad (1.1)$$

$f(x)$ is Mahler summable if and only if $\tilde{f}(x) = 0$. The Mahler discrete residues of $f(x)$ are (vectors whose components are) the coefficients occurring in the partial fraction decomposition of $\tilde{f}(x)$. This $\tilde{f}(x)$ plays the role of a *Mahler remainder* of $f(x)$, analogous to the remainder of Hermite reduction in the context of integration.

2 PRELIMINARIES

Here we define the notation and conventions used throughout this work, and prove some ancillary results. We fix once and for all an algebraically closed field $\mathbb{K}$ of characteristic zero and an integer $p \geq 2$ (not necessarily prime). We denote by $\mathbb{K}(x)$ the field of rational functions in an indeterminate x over $\mathbb{K}$. We often suppress the functional notation and write simply $f \in \mathbb{K}(x)$ instead of $f(x)$.

Definition 2.1. We denote by $\sigma : \mathbb{K}(x) \rightarrow \mathbb{K}(x)$ the $\mathbb{K}$ -linear endomorphism defined by $\sigma(x) = x^p$, called the *Mahler operator*, so that $\sigma(f(x)) = f(x^p)$ for $f(x) \in \mathbb{K}(x)$. We write $\Delta := \sigma - \text{id}$, so that $\Delta(f(x)) = f(x^p) - f(x)$ for $f(x) \in \mathbb{K}(x)$.

We say that $f \in \mathbb{K}(x)$ is *Mahler summable* if $f = \Delta(g)$ for some $g \in \mathbb{K}(x)$. The *Mahler summability problem for rational functions* is: given $f \in \mathbb{K}(x)$, decide whether f is Mahler summable.

Let $\mathbb{K}^\times = \mathbb{K} \setminus \{0\}$ denote the multiplicative group of $\mathbb{K}$. Let $\mathbb{K}_t^\times$ denote the torsion subgroup of $\mathbb{K}^\times$, i.e., the group of roots of unity in $\mathbb{K}^\times$. For $\zeta \in \mathbb{K}_t^\times$, the *order* of ζ is the smallest $r \in \mathbb{N}$ such that $\zeta^r = 1$. We fix once and for all a *compatible system* of p -power roots of unity $(\zeta_{p^n})_{n \geq 0} \subset \mathbb{K}_t^\times$, that is, each ζ_{p^n} has order p^n and $\zeta_{p^n}^{p^\ell} = \zeta_{p^{n-\ell}}$ for $0 \leq \ell \leq n$. We denote by $\pi_\ell^n : \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Z}/p^\ell\mathbb{Z}$ and by $\pi_n : \mathbb{Z} \rightarrow \mathbb{Z}/p^n\mathbb{Z}$ the canonical projections.

Each $f \in \mathbb{K}(x)$ decomposes uniquely as

$$f = f_L + f_T, \quad \text{where} \quad (2.1)$$

$f_L \in \mathbb{K}[x, x^{-1}]$ is a Laurent polynomial and $f_T = \frac{a}{b}$ for polynomials $a, b \in \mathbb{K}[x]$ such that $b \neq 0$ and, either $a = 0$, or else $\deg(a) < \deg(b)$ and $\gcd(a, b) = 1 = \gcd(x, b)$. The subscript L stands for “Laurent”, whereas the subscript T stands for “Tree” (see Definition 2.6).

LEMMA 2.2. *The decomposition $\mathbb{K}(x) \simeq \mathbb{K}[x, x^{-1}] \oplus \mathbb{K}(x)_T$ given by $f \mapsto f_L \oplus f_T$ as in (2.1) is σ -stable. For $f, g \in \mathbb{K}(x)$, $f = \Delta(g)$ if and only if $f_L = \Delta(g_L)$ and $f_T = \Delta(g_T)$.*

PROOF. We see that $\sigma(f_L) \in \mathbb{K}[x, x^{-1}]$ for any $f_L \in \mathbb{K}[x, x^{-1}]$. By the Euclidean algorithm, $\gcd(\sigma(a), \sigma(b)) = \sigma(\gcd(a, b))$ for any $0 \neq a, b \in \mathbb{K}[x]$. Thus the $\mathbb{K}$ -subspace $\mathbb{K}(x)_T$ is also stabilized by σ . Hence $f = \Delta(g)$ if and only if $f_L = \Delta(g_L)$ and $f_T = \Delta(g_T)$. $\square$

2.1 Mahler trajectories, trees, and cycles

We let $\mathcal{P} := \{p^n \mid n \in \mathbb{Z}_{\geq 0}\}$ denote the multiplicative monoid of non-negative powers of p . Then $\mathcal{P}$ acts on $\mathbb{Z}$ by multiplication, and the set of *maximal trajectories* for this action is

$$\mathbb{Z}/\mathcal{P} := \{\{0\}\} \cup \{\{ip^n \mid n \in \mathbb{Z}_{\geq 0}\} \mid i \in \mathbb{Z} \text{ such that } p \nmid i\}.$$

Remark 2.3. The usage of *trajectory* is perhaps unfamiliar to some readers: it is standard in the context of monoid (and more generally

semigroup) actions, and replaces the more familiar notion of *orbit* for group actions. As in that more familiar setting, the elements $\theta \in \mathbb{Z}/\mathcal{P}$ are pairwise disjoint sets whose union is all of $\mathbb{Z}$.

Definition 2.4. For a maximal trajectory $\theta \in \mathbb{Z}/\mathcal{P}$, we let

$$\mathbb{K}[x, x^{-1}]_\theta := \{\sum_j c_j x^j \in \mathbb{K}[x, x^{-1}] \mid c_j = 0 \text{ for all } j \notin \theta\}, \quad (2.2)$$

and call it the θ -subspace. The θ -component f_θ of $f \in \mathbb{K}(x)$ is the projection of the component f_L of f in (2.1) to $\mathbb{K}[x, x^{-1}]_\theta$ as in (2.2).

LEMMA 2.5. *For $f, g \in \mathbb{K}(x)$, $f_L = \Delta(g_L)$ if and only if $f_\theta = \Delta(g_\theta)$ for every maximal trajectory $\theta \in \mathbb{Z}/\mathcal{P}$.*

PROOF. This follows by observing that the $\mathbb{K}$ -linear decomposition $\mathbb{K}[x, x^{-1}] \simeq \bigoplus_{\theta \in \mathbb{Z}/\mathcal{P}} \mathbb{K}[x, x^{-1}]_\theta$ is σ -stable (cf. [13, §5]). $\square$

Definition 2.6. We denote by $\mathcal{T}_M$ the set of equivalence classes in $\mathbb{K}^\times$ for the equivalence relation $\alpha \sim \gamma \Leftrightarrow \alpha^{p^r} = \gamma^{p^s}$ for some $r, s \in \mathbb{Z}_{\geq 0}$. For $\alpha \in \mathbb{K}^\times$, we denote by $\tau(\alpha) \in \mathcal{T}_M$ the equivalence class of α under $\sim$. The elements $\tau \in \mathcal{T}_M$ are called *Mahler trees*.

Remark 2.7. The usage of *tree* in Definition 2.6 is motivated by the fact that one can define a digraph structure $D(\tau)$ on the vertex set τ with an edge from α to γ whenever $\alpha^p = \gamma$, whose underlying (undirected) graph is connected and acyclic so long as $\tau \cap \mathbb{K}_t^\times = \emptyset$. We find the terminology useful and suggestive also if $\tau \cap \mathbb{K}_t^\times \neq \emptyset$, because even in this exceptional case we do obtain a tree after collapsing the unique cycle in $D(\tau)$ defined below.

Definition 2.8. For a Mahler tree $\tau \in \mathcal{T}_M$, the *Mahler cycle* of τ is

$$C(\tau) := \{\gamma \in \tau \mid \gamma \text{ is a root of unity of order coprime to } p\}.$$

The *cycle length* of τ is defined to be $e(\tau) := |C(\tau)|$.

Example 2.9. (Cf. [15, Figures 4 and 5]). Let us illustrate the definitions of Mahler trees and Mahler cycles with $\mathbb{K} = \mathbb{C}$ and $p = 3$. In this example we write $\zeta_n := e^{\frac{2\pi\sqrt{-1}}{n}} \in \mathbb{C}^\times$, for concreteness.

The vertices in the digraph $D(\tau(2))$ near $\alpha = 2$ are:

$$\begin{array}{c} \{\sqrt[3]{2}, \zeta_3 \sqrt[3]{2}, \zeta_3^2 \sqrt[3]{2}\} \xrightarrow{\quad} \sqrt[3]{2} \\ \{\zeta_9 \sqrt[3]{2}, \zeta_9^4 \sqrt[3]{2}, \zeta_9^7 \sqrt[3]{2}\} \xrightarrow{\quad} \zeta_3 \sqrt[3]{2} \\ \{\zeta_9^2 \sqrt[3]{2}, \zeta_9^5 \sqrt[3]{2}, \zeta_9^8 \sqrt[3]{2}\} \xrightarrow{\quad} \zeta_3^2 \sqrt[3]{2} \end{array} \quad \begin{array}{c} \xrightarrow{\quad} 2 \xrightarrow{\quad} 8 \xrightarrow{\quad} 512 \end{array}$$

For $\alpha = \zeta_4$, we have $C(\tau(\zeta_4)) = \{\zeta_4, \zeta_4^3\}$, so the cycle length $e(\tau(\zeta_4)) = 2$. The vertices in the digraph $D(\tau(\zeta_4))$ near $C(\tau(\zeta_4))$ are:

$$\begin{array}{ccc} \{\zeta_{36}^5, \zeta_{36}^{13}, \zeta_{36}^{25}\} & & \{\zeta_{36}^7, \zeta_{36}^{19}, \zeta_{36}^{31}\} \\ \downarrow \downarrow \downarrow & & \downarrow \downarrow \downarrow \\ \zeta_{12} & & \zeta_{12}^7 \\ & \nearrow \quad \searrow & \\ & \zeta_4 \quad \zeta_4^3 & \\ & \nwarrow \quad \nearrow & \\ \zeta_{12}^5 & & \zeta_{12}^{11} \\ \uparrow \uparrow \uparrow & & \uparrow \uparrow \uparrow \\ \{\zeta_{36}^5, \zeta_{36}^{17}, \zeta_{36}^{29}\} & & \{\zeta_{36}^{11}, \zeta_{36}^{23}, \zeta_{36}^{35}\} \end{array}$$

Remark 2.10. Let us collect some immediate observations about Mahler cycles that we shall use, and refer to, throughout the sequel.

For a Mahler tree $\tau \in \mathcal{T}_M$ it follows from the Definition 2.6 that either $\tau \subset \mathbb{K}_t^\times$ or else $\tau \cap \mathbb{K}_t^\times = \emptyset$. In particular, $C(\tau) = \emptyset \Leftrightarrow e(\tau) = 0$, which occurs precisely when $\tau \not\subset \mathbb{K}_t^\times$ (the *non-torsion case*).

On the other hand, $\mathbb{K}_t^\times$ consists of the pre-periodic points for the action of the monoid $\mathcal{P}$ on $\mathbb{K}^\times$ given by $\alpha \mapsto \alpha^{p^n}$ for $n \in \mathbb{Z}_{\geq 0}$. For $\tau \subset \mathbb{K}_t^\times$ (the *torsion case*), the Mahler cycle $C(\tau)$ is a non-empty set endowed with a simply transitive action of the quotient monoid $\mathcal{P}/\mathcal{P}^e \simeq \mathbb{Z}/e\mathbb{Z}$, where $\mathcal{P}^e := \{p^{ne} \mid n \in \mathbb{Z}\}$, and $e := e(\tau)$. We emphasize that in general $C(\tau)$ is only a set, and not a group. The Mahler tree $\tau(1)$ consists precisely of the roots of unity $\zeta \in \mathbb{K}_t^\times$ whose order r is such that $\gcd(r, p^n) = r$ for some $p^n \in \mathcal{P}$, or equivalently such that every prime factor of r divides p . When $\tau \subset \mathbb{K}_t^\times$ but $\tau \neq \tau(1)$, the cycle length $e(\tau)$ is the order of p in the group of units $(\mathbb{Z}/r\mathbb{Z})^\times$, where $r > 1$ is the common order of the roots of unity $\gamma \in C(\tau)$, and $C(\tau) = \{\gamma^{p^\ell} \mid 0 \leq \ell \leq e-1\}$ for any given $\gamma \in C(\tau)$. We shall abusively write $C(\tau) = \{\gamma^{p^\ell} \mid \ell \in \mathbb{Z}/e\mathbb{Z}\}$.

2.2 Mahler supports and singular supports

Mahler trees allow us to define the following bespoke variants of the singular support $\text{sing}(f)$ of a rational function f (i.e., its set of poles), which are particularly well-suited to the Mahler context.

Definition 2.11. For $f \in \mathbb{K}(x)$, we define $\text{supp}(f) \subset \mathcal{T}_M \cup \{\infty\}$, called the *Mahler support* of f , as follows:

- $\infty \in \text{supp}(f)$ if and only if $f_L \neq 0$; and
- for $\tau \in \mathcal{T}_M$, $\tau \in \text{supp}(f)$ if and only if τ contains a pole of f .

For $\tau \in \mathcal{T}_M$, the *singular support* of f in τ , denoted by $\text{sing}(f, \tau)$, is the (possibly empty) set of poles of f contained in τ .

We omit the straightforward proof of the following lemma.

Lemma 2.12. For $f, g \in \mathbb{K}(x)$ and $0 \neq c \in \mathbb{K}$ we have the following:

- (1) $\text{supp}(f) = \emptyset \Leftrightarrow f = 0$;
- (2) $\text{supp}(\sigma(f)) = \text{supp}(f) = \text{supp}(c \cdot f)$; and
- (3) $\text{supp}(f + g) \subseteq \text{supp}(f) \cup \text{supp}(g)$.

Definition 2.13. For a Mahler tree $\tau \in \mathcal{T}_M$, the τ -subspace is

$$\mathbb{K}(x)_\tau := \{f_T \in \mathbb{K}(x)_T \mid \text{supp}(f_T) \subseteq \{\tau\}\}. \quad (2.3)$$

For $f \in \mathbb{K}(x)$, the τ -component f_τ of f is the projection of the component f_T of f in (2.1) to the τ -subspace $\mathbb{K}(x)_\tau$ in (2.3).

Lemma 2.14. For $f, g \in \mathbb{K}(x)$, $f_T = \Delta(g_T)$ if and only if $f_\tau = \Delta(g_\tau)$ for every Mahler tree $\tau \in \mathcal{T}_M$.

PROOF. It follows from Lemma 2.12 that the $\mathbb{K}$ -linear decomposition $\mathbb{K}(x)_T \simeq \bigoplus_{\tau \in \mathcal{T}_M} \mathbb{K}(x)_\tau$ is σ -stable (cf. [13, §5]). $\square$

2.3 Mahler dispersion

We now define a Mahler variant of the notion of (polar) dispersion used in [14], following the original definitions in [1, 2].

Definition 2.15. For $f \in \mathbb{K}(x)$ and $\tau \in \text{supp}(f)$, the *Mahler dispersion* of f at τ , denoted by $\text{disp}(f, \tau)$, is defined as follows.

If $\tau \in \mathcal{T}_M$, $\text{disp}(f, \tau)$ is the largest $d \in \mathbb{Z}_{\geq 0}$ (if it exists) for which there exists $\alpha \in \text{sing}(f, \tau)$ such that $\alpha^{p^d} \in \text{sing}(f, \tau)$. If there is no such $d \in \mathbb{Z}_{\geq 0}$, then we set $\text{disp}(f, \tau) = \infty$.

If $\tau = \infty$, let us write $f_L = \sum_{i=n}^N c_i x^i \in \mathbb{K}[x, x^{-1}]$ with $c_n c_N \neq 0$.

- If $f_L = c_0 \neq 0$ then we set $\text{disp}(f, \infty) = 0$; otherwise
- $\text{disp}(f, \infty)$ is the largest $d \in \mathbb{Z}_{\geq 0}$ for which there exists an index $i \neq 0$ such that $c_i \neq 0$ and $c_{ip^d} \neq 0$.

For $f \in \mathbb{K}(x)$ and $\tau \in \mathcal{T}_M \cup \{\infty\}$ such that $\tau \notin \text{supp}(f)$, we do not define $\text{disp}(f, \tau)$ at all (cf. [1, 2, 14]).

Similarly as in the shift and q -difference cases (cf. [17, Lemma 6.3] and [14, Lemma 2.4 and Lemma 2.9]), Mahler dispersions will play a crucial role in what follows. As we prove in Corollary 3.2, they already provide a partial obstruction to summability: if $f \in \mathbb{K}(x)$ is Mahler summable then every Mahler dispersion of f is non-zero. Moreover, Mahler dispersions also detect whether f has any “bad” poles (i.e., at roots of unity of order coprime to p) according to:

Lemma 2.16. Let $f \in \mathbb{K}(x)$ and $\tau \in \text{supp}(f)$. Then $\text{disp}(f, \tau) = \infty$ if and only if $\text{sing}(f, \tau) \cap C(\tau) \neq \emptyset$.

PROOF. ($\Rightarrow$). If $\text{disp}(f, \tau) = \infty$, then there exist $\alpha, \gamma \in \text{sing}(f, \tau)$ such that $\alpha^{p^d} = \gamma$ for infinitely many $d \in \mathbb{Z}_{\geq 0}$. Thus, both γ and α are roots of unity. Let r be the order of α . For any $d \in \mathbb{Z}_{\geq 0}$, the order of α^{p^d} is $r_d := r/\gcd(p^d, r)$, and we see that for every sufficiently large d , the order r_d of α^{p^d} is coprime to p , and therefore γ is a root of unity of order coprime to p .

($\Leftarrow$). For $\gamma \in \text{sing}(f, \tau) \cap C(\tau)$ we have $\gamma^{p^{e(\tau) \cdot n}} = \gamma$ for every $n \in \mathbb{Z}_{\geq 0}$ (cf. Remark 2.10), whence $\text{disp}(f, \tau) = \infty$ by Definition 2.15. $\square$

2.4 Mahler coefficients for partial fractions

We now study the effect of the Mahler operator σ on partial fraction decompositions. For $\alpha \in \mathbb{K}^\times$, $m \in \mathbb{N}$, and $1 \leq k \leq m$, we define the *Mahler coefficients* $V_k^m(\alpha) \in \mathbb{K}$ implicitly by

$$\sigma\left(\frac{1}{(x - \alpha^p)^m}\right) = \frac{1}{(x^p - \alpha^p)^m} = \sum_{k=1}^m \sum_{i=0}^{p-1} \frac{V_k^m(\zeta_p^i \alpha)}{(x - \zeta_p^i \alpha)^k}. \quad (2.4)$$

These coefficients are computed explicitly with the following result.

Lemma 2.17. There exist universal coefficients $\mathbb{V}_k^m \in \mathbb{Q}$ such that

$$V_k^m(\alpha) = \mathbb{V}_k^m \cdot \alpha^{k-pm}$$

for every $\alpha \in \mathbb{K}^\times$ and $1 \leq k \leq m$. Moreover, these $\mathbb{V}_k^m$ are the first m Taylor coefficients at $x = 1$ of

$$(x^{p-1} + \dots + x + 1)^{-m} = \sum_{k=1}^m \mathbb{V}_k^m \cdot (x-1)^{m-k} + O((x-1)^m). \quad (2.5)$$

PROOF. We claim that $V_k^m(\alpha) = V_k^m(1) \cdot \alpha^{k-pm}$ for every $\alpha \in \mathbb{K}^\times$. To see this, set $x = \alpha y$ for a new indeterminate y , and note that

$$\begin{aligned} \sum_{k=1}^m \sum_{i=0}^{p-1} \frac{V_k^m(\zeta_p^i \alpha)}{(x - \zeta_p^i \alpha)^k} &= \frac{1}{(x^p - \alpha^p)^m} = \alpha^{-pm} \cdot \frac{1}{(y^p - 1)^m} = \\ &= \alpha^{-pm} \sum_{k=1}^m \sum_{i=0}^{p-1} \frac{V_k^m(\zeta_p^i)}{(y - \zeta_p^i)^k} = \alpha^{-pm} \sum_{k=1}^m \sum_{i=0}^{p-1} \frac{V_k^m(\zeta_p^i) \alpha^k}{(x - \zeta_p^i \alpha)^k}. \end{aligned}$$

It follows that $V_k^m(\zeta_p^i \alpha) = V_k^m(\zeta_p^i) \alpha^{k-pm}$ for $i = 0, \dots, p-1$. In particular for $i = 0$ we obtain $V_k^m(\alpha) = V_k^m(1) \alpha^{k-pm}$, as claimed. Setting $\mathbb{V}_k^m := V_k^m(1)$, we see from (2.4) that $\mathbb{V}_k^m$ is the usual continuous residue of order k of $f(x) := (x^p - 1)^{-m}$. The formula (2.5) follows from [8, Section 2], where it is shown that $\mathbb{V}_k^m = \frac{g^{(m-k)}(1)}{(m-k)!}$, where $g(x) := (x-1)^m f(x) = (x^{p-1} + \dots + 1)^{-m}$. $\square$

The following immediate consequence of Lemma 2.17 is obtained by evaluating (2.5) at $x = 1$.

COROLLARY 2.18. For $\alpha \in \mathbb{K}^\times$, $V_m^m(\alpha) = p^{-m} \alpha^{m-pm}$.

Remark 2.19. We see in (2.4) two phenomena that arise in the Mahler context and have no counterpart in the shift and q -dilation settings considered in [14] — the main inspiration for the present work. Let $f \in \mathbb{K}(x)$ such that $0 \neq f_T$ as in (2.1). Then:

- (1) the number of poles of $\sigma(f_T)$ (counted either with or without multiplicity!) is strictly larger than that of f_T ; and
- (2) the (classical/continuous) higher-order residues of f_T “leak” into the lower-order residues of $\sigma(f_T)$.

These two phenomena are mainly responsible for our need to create new and somewhat intricate bookkeeping devices in the Mahler setting, which were (invisibly) not necessary in the shift and q -dilation settings considered in [14], in order to develop our proposed analogous theory of Mahler discrete residues.

Example 2.20. Let us illustrate the definition of Mahler coefficients with $p = 3$, $m = 2$, and $\alpha^3 = 1$. Then (2.4) becomes

$$\sigma\left(\frac{1}{(x-1)^2}\right) = \frac{1}{(x^3-1)^2} = \sum_{k=1}^2 \sum_{i=0}^2 \frac{\mathbb{V}_k^2 \cdot \zeta_3^{ki}}{(x-\zeta_3^i)^k},$$

because, by Lemma 2.17, $V_k^2(\zeta_3^i) = \mathbb{V}_k^2 \cdot (\zeta_3^i)^{k-6} = \mathbb{V}_k^2 \cdot \zeta_3^{ki}$ for $k = 1, 2$. We find in this case, using (2.5) in Lemma 2.17, that

$$\mathbb{V}_2^2 = (x^2 + x + 1)^{-2} \Big|_{x=1} = \frac{1}{9}; \text{ and } \mathbb{V}_1^2 = ((x^2 + x + 1)^{-2})' \Big|_{x=1} = -\frac{2}{9}.$$

One can verify using a computer algebra system (or by hand!) that the partial fraction decomposition of $9 \cdot (x^3 - 1)^{-2}$ is indeed

$$\frac{1}{(x-1)^2} + \frac{\zeta_3^2}{(x-\zeta_3)^2} + \frac{\zeta_3}{(x-\zeta_3^2)^2} + \frac{-2}{x-1} + \frac{-2\zeta_3}{x-\zeta_3} + \frac{-2\zeta_3^2}{x-\zeta_3^2}.$$

3 MAHLER DISPERSION AND SUMMABILITY

The goal of this section is to prove Corollary 3.2 below. This is an essential ingredient in our proofs, following [14]. The following result is a Mahler analogue of [12, Lemma 2.6].

PROPOSITION 3.1. Let $f, g \in \mathbb{K}(x)$ such that $f = \Delta(g)$. Then $\text{supp}(f) \subseteq \text{supp}(g)$. Moreover, $\text{disp}(f, \tau) = \text{disp}(g, \tau) + 1$ for every $\tau \in \text{supp}(f)$, with the convention that $\infty + 1 = \infty$.

PROOF. By Lemma 2.12, $\tau \in \text{supp}(f) \Rightarrow \tau \in \text{supp}(g)$. We consider separately the two main cases: (1). $\tau = \infty$; and (2). $\tau \in \mathcal{T}_M$.

(1). For $f_L, g_L \in \mathbb{K}[x, x^{-1}]$ as in (2.1), we have $0 \neq f_L = \Delta(g_L)$, since $\infty \in \text{supp}(f)$, and in particular $g_L \notin \mathbb{K}$. Then $f_\theta = \Delta(g_\theta)$ for each $\theta \in \mathbb{Z}/\mathcal{P}$ by Lemma 2.5. Since, for $\theta = \{0\}$, $f_{\{0\}} = \Delta(g_{\{0\}}) = 0$, it follows from Definition 2.15 that

$$\text{disp}(f, \infty) = \max \{ \text{disp}(f_\theta, \infty) \mid \{0\} \neq \theta \in \mathbb{Z}/\mathcal{P}, f_\theta \neq 0 \}.$$

We claim $\text{disp}(\Delta(g_\theta), \infty) = \text{disp}(g_\theta, \infty) + 1$ for every $g_\theta \in \mathbb{K}[x, x^{-1}]_\theta$ with $\{0\} \neq \theta \in \mathbb{Z}/\mathcal{P}$, which will conclude the proof of (1). To prove the claim, let us write $g_\theta = \sum_{j=0}^d c_{ip^j} x^{ip^j}$, where we assume $c_i \neq 0$ and $c_{ip^d} \neq 0$, i.e., $\text{disp}(g_\theta, \infty) = d$. Then

$$\Delta(g_\theta) = c_{ip^d} x^{ip^{d+1}} - c_i x^i + \sum_{j=1}^d (c_{ip^{j-1}} - c_{ip^j}) x^{ip^j},$$

from which it follows that $\text{disp}(\Delta(g_\theta), \infty) = d + 1$, as desired.

(2). By Lemma 2.14, $f_\tau = \Delta(g_\tau)$ for each $\tau \in \mathcal{T}_M$, and in particular for each $\tau \in \text{supp}(f)$. We consider two subcases, depending on whether $\text{disp}(g, \tau)$ is finite or not.

In the first subcase, $\text{disp}(g, \tau) =: d < \infty$. Let $\alpha \in \tau$ be such that α and α^{p^d} are poles of g . Let $\gamma \in \tau$ such that $\gamma^p = \alpha$. Then γ is a pole of $\sigma(g)$ but not of g (by maximality of d), whence γ is a pole of f . On the other hand, $\gamma^{p^{d+1}} = \alpha^{p^d}$ is a pole of g but not of $\sigma(g)$, for if α^{p^d} were a pole of $\sigma(g)$ then $\alpha^{p^{d+1}}$ would be a pole of g , again contradicting maximality of d . Hence $\gamma^{p^{d+1}}$ is a pole of f . Thus $\text{disp}(f, \tau) \geq d + 1$. One can show equality by contradiction: if $\alpha \in \tau$ is a pole of f such that α^{p^s} is also a pole of f for some $s > d + 1$, then each of α and α^{p^s} is either a pole of g or a pole of $\sigma(g)$. This implies (after tedious but straightforward casework) that there exist $0 \leq i, j \leq 1$ such that $\alpha^{p^{s+i}}$ and α^{p^j} are both poles of g , which contradicts the maximality of d since in any case $s + i - j \geq s - 1 > d$. Hence $\text{disp}(f, \tau) = \text{disp}(g, \tau) + 1$ in this first subcase.

In the last remaining subcase where $\text{disp}(g, \tau) = \infty$, there exists $\gamma \in \text{sing}(g, \tau) \cap C(\tau)$ by Lemma 2.16. We claim $\gamma^{p^\ell} \in \text{sing}(f, \tau)$ for some $\ell \in \mathbb{Z}/e\mathbb{Z}$, where $e := e(\tau) \geq 1$ (cf. Remark 2.10, where we discussed the meaning of γ^{p^ℓ} for $\ell \in \mathbb{Z}/e\mathbb{Z}$, rather than $\ell \in \mathbb{Z}_{\geq 0}$). This will imply that $\text{disp}(f, \tau) = \infty = \text{disp}(g, \tau) + 1$, by Lemma 2.16.

Let us prove the claim. Note that the $\mathbb{K}$ -subspace S of $\mathbb{K}(x)_\tau$, consisting of rational functions none of whose poles belongs to $C(\tau)$, or equivalently (by Lemma 2.16), the $\mathbb{K}$ -span of the elements of $\mathbb{K}(x)_\tau$ having finite dispersion, is σ -stable¹. So we may assume

$$g_\tau = \sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{d(k, \ell)}{(x - \gamma^{p^\ell})^k},$$

where $d(k, \ell) \in \mathbb{K}$ such that $d(m, \ell) \neq 0$ for some $\ell \in \mathbb{Z}/e\mathbb{Z}$, without loss of generality, because the other possible poles of g in τ cannot contribute to the possible poles of f in $C(\tau)$ (since S is σ -stable). Then

$$\sigma(g_\tau) = \sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{d(k, \ell)}{(x^{p^e} - \gamma^{p^{\ell e}})^k} = \sum_{i=0}^{p-1} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{V_m^m(\zeta_p^i \gamma^{p^{\ell-1}}) \cdot d(m, \ell)}{(x - \zeta_p^i \gamma^{p^{\ell-1}})^m} \right) + \text{(lower-order terms)},$$

where the V_m^m are as in (2.4), and therefore²

$$f_\tau = \Delta(g_\tau) = \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{V_m^m(\gamma^{p^\ell}) \cdot d(m, \ell + 1) - d(m, \ell)}{(x - \gamma^{p^\ell})^m} + \text{(lower-order terms)} + \text{(elements of } S). \quad (3.1)$$

But the coefficients $V_m^m(\gamma^{p^\ell}) \cdot d(m, \ell + 1) - d(m, \ell)$ cannot be zero for every $\ell \in \mathbb{Z}/e\mathbb{Z}$, for otherwise the computation

$$d(m, \ell) = d(m, \ell) \prod_{j=0}^{e-1} V_m^m(\gamma^{p^j}) = d(m, \ell) \prod_{j=0}^{e-1} \frac{\gamma^{mp^j}}{p^m \gamma^{mp^{j+1}}} = \frac{d(m, \ell)}{p^{em}},$$

where the middle equality is obtained from Corollary 2.18, would imply that $d(m, \ell) = 0$ for every $\ell \in \mathbb{Z}/e\mathbb{Z}$. But this is impossible, concluding the proof of the claim that f has a pole in $C(\tau)$. $\square$

¹If a denominator $b \in \mathbb{K}[x]$ has no roots in $C(\tau)$ then neither does $\sigma(b)$, for if $\gamma \in C(\tau)$ were a root of $\sigma(b)$ then $\gamma^p \in C(\tau)$ would be a root of b .

²See Remark 4.13, where we systematically elaborate on the details of this computation.

COROLLARY 3.2. Suppose that $f \in \mathbb{K}(x)$ is Mahler summable. Then $\text{disp}(f, \tau)$ is non-zero for every $\tau \in \text{supp}(f)$.

PROOF. This is a direct consequence of Proposition 3.1. $\square$

4 MAHLER DISCRETE RESIDUES

In this section we define the Mahler discrete residues of $f \in \mathbb{K}(x)$, in increasing order of complexity: first at infinity, and then at Mahler trees $\tau \in \mathcal{T}_M$, separately in the *non-torsion case* where $\tau \notin \mathbb{K}_t^\times$, and finally in the *torsion case* where $\tau \subset \mathbb{K}_t^\times$ (cf. Remark 2.10).

4.1 Mahler discrete residue at infinity

Here we define the Mahler discrete residue of $f \in \mathbb{K}(x)$ at ∞ in terms of the component $f_L \in \mathbb{K}[x, x^{-1}]$ of f in (2.1), and show that it forms a complete obstruction to the Mahler summability of f_L . The proof of Proposition 4.2 below follows the same strategy as that of [14, Propositions 2.5 and 2.10]: we add to f_L a sequence of Mahler summable elements to eventually obtain a *Mahler remainder* $\tilde{f}_L$ whose apparent dispersion is 0, and then use Corollary 3.2 to conclude that f_L is Mahler summable if and only if this $\tilde{f}_L = 0$.

Definition 4.1. For $f \in \mathbb{K}(x)$, let $f_L = \sum_{j \in \mathbb{Z}} c_j x^j$ with $c_j = 0$ for all but finitely many $j \in \mathbb{Z}$. The *Mahler discrete residue* of f at ∞ is the vector $\text{dres}(f, \infty) := \left(\sum_{j \in \theta} c_j \right)_{\theta \in \mathbb{Z}/\mathcal{P}} \in \bigoplus_{\theta \in \mathbb{Z}/\mathcal{P}} \mathbb{K}$.

PROPOSITION 4.2. For $f \in \mathbb{K}(x)$, the component $f_L \in \mathbb{K}[x, x^{-1}]$ in (2.1) is Mahler summable if and only if $\text{dres}(f, \infty) = 0$.

PROOF. By Lemma 2.5, f_L is Mahler summable if and only if f_θ is Mahler summable for all $\theta \in \mathbb{Z}/\mathcal{P}$. We shall show f_θ is Mahler summable if and only if the θ -component $\text{dres}(f, \infty)_\theta = 0$. We prove this separately in two cases: (1). $\theta = \{0\}$; and (2). $\theta \neq \{0\}$.

(1). For $\theta = \{0\}$, $f_{\{0\}} = c_0 = \text{dres}(f, \infty)_{\{0\}}$ by Definition 4.1. If $\text{dres}(f, \infty)_{\{0\}} = 0$, then $f_{\{0\}} = 0$ is Mahler summable. On the other hand, if $\text{dres}(f, \infty)_{\{0\}} \neq 0$ then $f_{\{0\}} \neq 0$ and $\text{disp}(f_{\{0\}}, \infty) = 0$ by Definition 2.15, so $f_{\{0\}}$ is not Mahler summable by Corollary 3.2.

(2). Suppose $\theta \neq \{0\}$. The claim is trivial in case $f_\theta = 0$; assume $f_\theta \neq 0$. Let us write $f_\theta = \sum_{j \in \theta} c_j x^j \in \mathbb{K}[x, x^{-1}]_\theta$, where: $c_j = 0$ for all but finitely many $j \in \theta$ and $c_j \neq 0$ for at least one $j \in \theta$. Let us write $\theta = \{ip^n \mid n \in \mathbb{Z}_{\geq 0}\}$ for $i \in \theta$ such that $p \nmid i$. Let $h \in \mathbb{Z}_{\geq 0}$ be maximal such that $c_{ip^h} \neq 0$. Let us define recursively: $g_\theta^{(0)} := 0$; and, if $h \geq 1$, then set

$$g_\theta^{(n+1)} := \sum_{k=0}^n \left(\sum_{\ell=0}^k c_{ip^\ell} \right) x^{ip^k} = g_\theta^{(n)} + \left(\sum_{\ell=0}^n c_{ip^\ell} \right) x^{ip^n}$$

for $0 \leq n \leq h-1$. A straightforward induction argument shows:

$$\tilde{f}_\theta^{(n)} := f_\theta + \Delta \left(g_\theta^{(n)} \right) = \sum_{k=n+1}^{h+1} c_{ip^k} x^{ip^k} + \left(\sum_{\ell=0}^n c_{ip^\ell} \right) x^{ip^n} \quad (4.1)$$

for each $0 \leq n \leq h$, whence $\tilde{f}_\theta^{(h)} = (\text{dres}(f, \infty)_\theta) \cdot x^{ip^h}$. The harmless summand for $k = h+1$ in (4.1) is included so that the sum makes sense for $n = h$, but $c_{ip^k} = 0$ for every $k > h$. We see that $\tilde{f}_\theta^{(h)}$ is Mahler summable if and only if f_θ is Mahler summable. In particular, if $\tilde{f}_\theta^{(h)} = 0$ then f_θ is Mahler summable. But if $\tilde{f}_\theta^{(h)} \neq 0$ then $\text{disp}(\tilde{f}_\theta^{(h)}, \infty) = 0$, and by Corollary 3.2 $\tilde{f}_\theta^{(h)}$ is not Mahler summable, so neither is f_θ . Clearly, $\tilde{f}_\theta^{(h)} = 0 \Leftrightarrow \text{dres}(f, \infty)_\theta = 0$. $\square$

Remark 4.3. For $\{0\} \neq \theta \in \mathbb{Z}/\mathcal{P}$ such that $f_\theta \neq 0$, the elements $\tilde{f}_\theta^{(h)}, g_\theta^{(h)} \in \mathbb{K}[x, x^{-1}]_\theta$ constructed in the proof of Proposition 4.2 are the θ -components of the $\tilde{f}, g \in \mathbb{K}(x)$ in (1.1). If $f_\theta = 0$, then we define $\tilde{f}_\theta := 0 =: g_\theta$. In any case, we set $\tilde{f}_{\{0\}} := f_{\{0\}}$ and $g_{\{0\}} := 0$.

4.2 Mahler discrete residues at Mahler trees

Here we define the Mahler discrete residues of $f \in \mathbb{K}(x)$ at a Mahler tree $\tau \in \mathcal{T}_M$, in terms of the partial fraction decomposition of the component $f_\tau \in \mathbb{K}(x)_\tau$ in Definition 2.13, and show they comprise a complete obstruction to the Mahler summability of f_τ . We proceed separately in the *non-torsion case* $\tau \notin \mathbb{K}_t^\times$ and the *torsion case* $\tau \subset \mathbb{K}_t^\times$ (cf. Remark 2.10), depending on which case we represent the poles of f_τ in a particular manner.

LEMMA 4.4. For $f \in \mathbb{K}(x)$ and $\tau \in \text{supp}(f) \cap \mathcal{T}_M$ such that $\tau \notin \mathbb{K}_t^\times$, there exists $\gamma \in \text{sing}(f, \tau)$ and $h \in \mathbb{Z}_{\geq 0}$ such that

$$\text{sing}(f, \tau) \subseteq \beta_h(\gamma) := \{ \zeta_{p^n}^i \gamma^{p^{h-n}} \mid 0 \leq n \leq h; i \in \mathbb{Z}/p^n \mathbb{Z} \}.$$

Moreover, the elements $\zeta_{p^n}^i \gamma^{p^{h-n}} \in \beta_h(\gamma)$ are uniquely determined by $0 \leq n \leq h$ and $i \in \mathbb{Z}/p^n \mathbb{Z}$, relative to the choice of $\gamma \in \text{sing}(f, \tau)$.

PROOF. Note that the set $\beta_h(\gamma)$ (mnemonic: “bouquet” of height h at γ) is precisely the union of the sets of roots of the y -polynomials $y^{p^n} - \gamma^{p^h} = 0$ for all $0 \leq n \leq h$. The elements of $\beta_h(\gamma)$ are uniquely determined by n and i (relative to the choice of γ), because if we had $\zeta_{p^m}^j \gamma^{p^{h-m}} = \zeta_{p^n}^i \gamma^{p^{h-n}}$, then this would force $m = n$, for otherwise $\gamma \in \mathbb{K}_t^\times$ contradicting our assumptions, and then $\zeta_{p^n}^j = \zeta_{p^n}^i$ implies that $j = i$. Let us now show that for any finite set $S \subset \tau$ there exist $\gamma \in S$ and $h \in \mathbb{Z}_{\geq 0}$ such that $S \subseteq \beta_h(\gamma)$. For $\alpha \in S$, let $h(\alpha) \in \mathbb{Z}_{\geq 0}$ be minimal such that $\alpha^{p^{h(\alpha)}} \in \xi^{\mathcal{P}}$ for every $\xi \in S$, where $\xi^{\mathcal{P}} := \{ \xi^{p^t} \mid t \in \mathbb{Z}_{\geq 0} \}$. Choose $\gamma \in S$ such that $h(\gamma) =: h$ is maximal among all elements of S . We claim that $\alpha^{p^{h(\alpha)}} = \gamma^{p^h}$ for every $\alpha \in S$, which will conclude the proof, since $h(\alpha) \leq h$ for every $\alpha \in S$. To prove the claim, note that in any case there exist $t, r \in \mathbb{Z}_{\geq 0}$ such that $\alpha^{p^t} = \gamma^{p^h}$ and $\alpha^{p^{h(\alpha)}} = \gamma^{p^r}$, and the minimality of $h(\alpha)$ and h then imply $t \geq h(\alpha)$ and $r \geq h$. But then

$$\gamma^{p^h} = \alpha^{p^t} = (\alpha^{p^{h(\alpha)}})^{p^{t-h(\alpha)}} = (\gamma^{p^r})^{p^{t-h(\alpha)}} = \gamma^{p^{r+t-h(\alpha)}},$$

and since $\gamma \notin \mathbb{K}_t^\times$ we obtain that $h + h(\alpha) = r + t$, from which it follows that $r = h$ and $t = h(\alpha)$, as claimed. $\square$

LEMMA 4.5. Let $\tau \in \mathcal{T}_M$ with $\tau \subset \mathbb{K}_t^\times$ and $e := e(\tau)$. Choose $\gamma \in C(\tau)$. Then for $\alpha \in \tau$ there are unique $n \in \mathbb{Z}_{\geq 0}$, $i \in \mathbb{Z}/p^n \mathbb{Z}$, and $\ell \in \mathbb{Z}/e\mathbb{Z}$, with either $n = i = 0$ or $p \nmid i$, such that $\alpha = \zeta_{p^n}^i \gamma^{p^{\ell - \pi_e(n)}}$.

PROOF. There exist integers $n, t \in \mathbb{Z}_{\geq 0}$ such that $\alpha^{p^n} = \gamma^{p^t}$, and we may take this n to be as small as possible and replace t with $\ell := \pi_e(t)$. The p^n distinct solutions to $y^{p^n} = \gamma^{p^\ell}$, one of which is $y = \alpha$, are all of the form $\zeta_{p^n}^i \gamma^{p^{\ell - \pi_e(n)}}$ for $i \in \mathbb{Z}/p^n \mathbb{Z}$. Since n is minimal, either $n = i = 0$ or else $1 \leq i \leq p^n - 1$ with $p \nmid i$. $\square$

Definition 4.6. For $f \in \mathbb{K}(x)$ and $\tau \in \text{supp}(f) \cap \mathcal{T}_M$, the *height* of f at τ , denoted by $h(f, \tau)$, is defined as follows.

- If $\tau \notin \mathbb{K}_t^\times$, $h(f, \tau)$ is the smallest $h \in \mathbb{Z}_{\geq 0}$ such that $\text{sing}(f, \tau)$ is contained in $\beta_h(\gamma)$ for some $\gamma \in \text{sing}(f, \tau)$ as in Lemma 4.4.
- If $\tau \subset \mathbb{K}_t^\times$, $h(f, \tau)$ is the smallest $h \in \mathbb{Z}_{\geq 0}$ such that α^{p^h} belongs to $C(\tau)$ for every $\alpha \in \text{sing}(f, \tau)$.

Remark 4.7. Suppose $f \in \mathbb{K}(x)$ and $\tau \in \text{supp}(f) \cap \mathcal{T}_M$. We always have $h(f, \tau) < \infty$. One can show that $h(f, \tau) \geq \text{disp}(f, \tau)$ provided that $\text{disp}(f, \tau) < \infty$, but even then the inequality may be strict.

4.2.1 Mahler discrete residues at Mahler trees: the non-torsion case.

LEMMA 4.8. *Let $f \in \mathbb{K}(x)$ and suppose $\tau \in \text{supp}(f) \cap \mathcal{T}_M$ such that $\tau \notin \mathbb{K}_t^\times$. Then there exists $\gamma \in \text{sing}(f, \tau)$ such that the partial fraction decomposition of f_τ is of the form*

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \left(\sum_{i \in \mathbb{Z}/p^n \mathbb{Z}} \frac{c_\gamma(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k} \right), \quad (4.2)$$

where $m \geq 1$ is the highest order of a pole of f in $\text{sing}(f, \tau)$ and the height $h := h(f, \tau)$ is as in Definition 4.6.

The coefficients $c_\gamma(k, n, i) \in \mathbb{K}$ are uniquely determined by f and the choice of γ , and moreover for any $\gamma, \tilde{\gamma} \in \tau$ as above we have $\tilde{\gamma} = \zeta_{p^h}^j \gamma$ for some $j \in \mathbb{Z}/p^h \mathbb{Z}$, and

$$c_{\tilde{\gamma}}(k, n, i) = c_\gamma(k, n, i + \pi_n^h(j)). \quad (4.3)$$

PROOF. We obtain the existence of $\gamma \in \text{sing}(f, \tau)$ such that $\text{sing}(f, \tau) \subseteq \beta_h(\gamma)$ by Lemma 4.4 and Definition 4.6. The existence and uniqueness of the coefficients $c_\gamma(k, n, i) \in \mathbb{K}$ satisfying (4.2) follows directly from the existence and uniqueness of partial fraction decompositions, since in this case the elements $\zeta_{p^n}^i \gamma^{p^{h-n}} \in \beta_h(\gamma)$ are uniquely determined by n and i (relative to the choice of γ), by Lemma 4.4. For any $\tilde{\gamma} \in \text{sing}(f, \tau)$ such that $\text{sing}(f, \tau) \subseteq \beta_h(\tilde{\gamma})$ we would have $\tilde{\gamma}^{p^{\tilde{n}}} = \gamma^{p^h}$ and $\gamma^{p^n} = \tilde{\gamma}^{p^{\tilde{n}}}$ such that $0 \leq n, \tilde{n} \leq h$, which would force $\tilde{n} = h = n$ since $\gamma \notin \mathbb{K}_t^\times$. Hence $\tilde{\gamma} = \zeta_{p^h}^j \gamma$ for some $j \in \mathbb{Z}/p^h \mathbb{Z}$, and the computation

$$\frac{c_{\tilde{\gamma}}(k, n, i)}{(x - \zeta_{p^n}^i \tilde{\gamma}^{p^{h-n}})^k} = \frac{c_{\tilde{\gamma}}(k, n, i)}{(x - \zeta_{p^n}^{i+j} \gamma^{p^{h-n}})^k} = \frac{c_\gamma(k, n, i+j)}{(x - \zeta_{p^n}^{i+j} \gamma^{p^{h-n}})^k}$$

implies the transformation formula (4.3). $\square$

Remark 4.9. Writing f_τ as in (4.2), let us compute the effect of σ :

$$\begin{aligned} \sigma \left(\sum_{k=1}^m \left(\sum_{i \in \mathbb{Z}/p^n \mathbb{Z}} c_\gamma(k, n, i) \cdot (x - \zeta_{p^n}^i \gamma^{p^{h-n}})^{-k} \right) \right) \\ = \sum_{k=1}^m \left(\sum_{i \in \mathbb{Z}/p^{n+1} \mathbb{Z}} \frac{\sum_{s=k}^m V_k^s \left(\zeta_{p^{n+1}}^i \gamma^{p^{h-(n+1)}} \right) \cdot c_\gamma(s, n, \pi_{n+1}^{n+1}(i))}{(x - \zeta_{p^{n+1}}^i \gamma^{p^{h-(n+1)}})^k} \right) \end{aligned} \quad (4.4)$$

for each $0 \leq n \leq h-1$, where the V_k^s are as in (2.4) for $k \leq s \leq m$.

Definition 4.10. For $f \in \mathbb{K}(x)$ and $\tau \in \mathcal{T}_M$ with $\tau \notin \mathbb{K}_t^\times$, the Mahler discrete residue of f at τ of degree $k \in \mathbb{N}$ is the vector $\text{dres}(f, \tau, k) \in \bigoplus_{\alpha \in \tau} \mathbb{K}$ defined in terms of the $c_\gamma(k, n, i)$ in the partial fraction decomposition of f_τ in Lemma 4.8 as follows.

Set $\text{dres}(f, \tau, k) := \mathbf{0}$ if $\tau \notin \text{supp}(f)$ or if $k > m$. For $\tau \in \text{supp}(f)$ and $\alpha \in \tau$, the component $\text{dres}(f, \tau, k)_\alpha := 0$ whenever $\alpha^{p^h} \neq \gamma^{p^h}$.

For $1 \leq k \leq m$ and $\alpha = \zeta_{p^h}^i \gamma$ with $i \in \mathbb{Z}/p^h \mathbb{Z}$, the component

$$\text{dres}(f, \tau, k)_\alpha := \hat{c}_\gamma(k, h, i);$$

where for $0 \leq n \leq h$ and $i \in \mathbb{Z}/p^n \mathbb{Z}$ we define recursively (in n):

$$\hat{c}_\gamma(k, 0, 0) := c_\gamma(k, 0, 0); \text{ and, for } 1 \leq n \leq h \text{ and } i \in \mathbb{Z}/p^n \mathbb{Z}, \quad (4.5)$$

$$\hat{c}_\gamma(k, n, i) := c_\gamma(k, n, i) + \sum_{s=k}^m V_k^s \left(\zeta_{p^n}^i \gamma^{p^{h-n}} \right) \cdot \hat{c}_\gamma(s, n-1, \pi_{n-1}^n(i))$$

(cf. (4.4)), where the V_k^s are as in (2.4) for $k \leq s \leq m$.

Remark 4.11. Note that the definition of $\text{dres}(f, \tau, k)$ for $\tau \notin \mathbb{K}_t^\times$ given above is independent of the choice of $\gamma \in \text{sing}(f, \tau)$, because for any possibly different $\tilde{\gamma} = \zeta_{p^h}^j \gamma$ with $j \in \mathbb{Z}/p^h \mathbb{Z}$ we obtain $\zeta_{p^h}^i \tilde{\gamma} = \zeta_{p^h}^{i+j} \gamma =: \alpha$. The equality of the expressions

$$\hat{c}_{\tilde{\gamma}}(k, h, i) = \text{dres}(f, \tau, k)_\alpha = \hat{c}_\gamma(k, h, i+j)$$

follows from (4.3), since $\zeta_{p^n}^i \tilde{\gamma}^{p^{h-n}} = \zeta_{p^n}^{i+j} \gamma^{p^{h-n}}$ for all $i \in \mathbb{Z}/p^n \mathbb{Z}$, and therefore $\hat{c}_{\tilde{\gamma}}(k, n, i) = \hat{c}_\gamma(k, n, i + \pi_n^h(j))$ for every $0 \leq n \leq h$.

4.2.2 Mahler discrete residues at Mahler trees: the torsion case.

LEMMA 4.12. *Let $f \in \mathbb{K}(x)$ and suppose $\tau \in \text{supp}(f)$ such that $\tau \subset \mathbb{K}_t^\times$. Then for any $\gamma \in C(\tau)$ the partial fraction decomposition of f_τ is of the form*

$$f_\tau = \sum_{k=1}^m \sum_{n=0}^h \left(\sum'_{i \in \mathbb{Z}/p^n \mathbb{Z}} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{d_\gamma(k, n, i, \ell)}{(x - \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}})^k} \right) \right), \quad (4.6)$$

where: $m \geq 1$ is the highest order of a pole of f in $\text{sing}(f, \tau)$; the height $h := h(f, \tau)$ is as in Definition 4.6; the restricted sum $\sum'$ is taken over $i \in \mathbb{Z}/p^n \mathbb{Z}$ such that $p \nmid i$ whenever $n \neq 0$; and $e := e(\tau) \geq 1$.

The coefficients $d_\gamma(k, n, i, \ell) \in \mathbb{K}$ are uniquely determined by f and γ , and moreover for any $\gamma, \tilde{\gamma} \in C(\tau)$ we have $\tilde{\gamma} = \gamma^{p^j}$ for some $j \in \mathbb{Z}/e\mathbb{Z}$, and

$$d_{\tilde{\gamma}}(k, n, i, \ell) = d_\gamma(k, n, i, \ell + j). \quad (4.7)$$

PROOF. If $\tau \subset \mathbb{K}_t^\times$ then $e \geq 1$ (cf. Remark 2.10). We then have by Lemma 4.5 that for any given choice of $\gamma \in C(\tau)$ the elements $\alpha \in \tau$ can be written uniquely as $\alpha = \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}}$ with $n \in \mathbb{Z}_{\geq 0}$ and $i \in \mathbb{Z}/p^n \mathbb{Z}$ such that either $n = i = 0$ or else $p \nmid i$. The set of elements $\alpha \in \tau$ such that $\alpha^{p^h} \in C(\tau)$ are precisely the $\alpha = \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}}$ with $n \leq h$. It follows that the apparent poles in (4.6) are all distinct and include all the poles of f , and therefore the coefficients $d_\gamma(k, n, i, \ell) \in \mathbb{K}$ are uniquely determined by f and γ . It also follows from Lemma 4.5 that for any other $\tilde{\gamma} \in C(\tau)$ there exists $j \in \mathbb{Z}/e\mathbb{Z}$ such that $\tilde{\gamma} = \gamma^{p^j}$, and therefore the computation

$$\frac{d_{\tilde{\gamma}}(k, n, i, \ell)}{(x - \zeta_{p^n}^i \tilde{\gamma}^{p^{\ell-\pi_e(n)}})^k} = \frac{d_\gamma(k, n, i, \ell)}{(x - \zeta_{p^n}^i \gamma^{p^{\ell+j-\pi_e(n)}})^k} = \frac{d_\gamma(k, n, i, \ell+j)}{(x - \zeta_{p^n}^i \gamma^{p^{\ell+j-\pi_e(n)}})^k}$$

implies the transformation formula (4.7). $\square$

Remark 4.13. Writing f_τ as in (4.6), let us compute the effect of σ :

$$\begin{aligned} \sigma \left(\sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} d_\gamma(k, 0, 0, \ell) \cdot (x - \gamma^{p^\ell})^{-k} \right) \\ = \sum_{k=1}^m \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\sum_{s=k}^m V_k^s \left(\gamma^{p^\ell} \right) \cdot d_\gamma(s, 0, 0, \ell+1)}{(x - \gamma^{p^\ell})^k} \right) + \\ + \sum_{k=1}^m \sum_{i=1}^{p-1} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\sum_{s=k}^m V_k^s \left(\zeta_{p^i}^i \gamma^{p^{\ell-1}} \right) \cdot d_\gamma(s, 0, i, \ell)}{(x - \zeta_{p^i}^i \gamma^{p^{\ell-1}})^k} \right); \end{aligned} \quad (4.8)$$

and for $n \geq 1$ and each $\ell \in \mathbb{Z}/e\mathbb{Z}$ we have

$$\begin{aligned} \sigma \left(\sum_{k=1}^m \sum'_{i \in \mathbb{Z}/p^n \mathbb{Z}} d_\gamma(k, n, i, \ell) \cdot (x - \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}})^{-k} \right) \\ = \sum_{k=1}^m \left(\sum'_{i \in \mathbb{Z}/p^{n+1} \mathbb{Z}} \frac{\sum_{s=k}^m V_k^s \left(\zeta_{p^{n+1}}^i \gamma^{p^{\ell-\pi_e(n+1)}} \right) \cdot d_\gamma(s, n, \pi_{n+1}^{n+1}(i), \ell)}{(x - \zeta_{p^{n+1}}^i \gamma^{p^{\ell-\pi_e(n+1)}})^k} \right); \end{aligned} \quad (4.9)$$

where the V_k^s are as in (2.4) for $k \leq s \leq m$.

The following technical lemma is essential for the definition of Mahler discrete residues in the torsion case. The map $\mathcal{D}_Y^{(m)}$ defined below already appeared (anonymously) in (3.1). It captures the effect of Δ on the (classical/continuous) residues at poles in the Mahler cycle $C(\tau)$ (see Definition 2.8), according to the computation (4.8).

LEMMA 4.14. *Let $\tau \in \mathcal{T}_M$ with $\tau \subset \mathbb{K}_t^\times$ and $e := e(\tau)$. For $\gamma \in C(\tau)$ and $m \in \mathbb{N}$, define $\mathcal{D}_Y^{(m)} : \mathbb{K}^{m \times e} \rightarrow \mathbb{K}^{m \times e}$ by*

$$\mathcal{D}_Y^{(m)} : (c_{k,\ell})_{\substack{1 \leq k \leq m \\ \ell \in \mathbb{Z}/e\mathbb{Z}}} \mapsto (d_{k,\ell})_{\substack{1 \leq k \leq m \\ \ell \in \mathbb{Z}/e\mathbb{Z}}}, \quad (4.10)$$

where $d_{k,\ell} := c_{k,\ell} - \sum_{s=k}^m V_k^s(\gamma^{p^\ell}) \cdot c_{s,\ell+1}$, and where the V_k^s are as in (2.4). Then $\mathcal{D}_Y^{(m)}$ is invertible and has no non-trivial fixed points.

PROOF. Let $\mathbf{0} \neq (c_{k,\ell}) \in \mathbb{K}^{m \times e}$, and write $(d_{k,\ell}) := \mathcal{D}_Y^{(m)}(c_{k,\ell})$. Let $1 \leq r \leq m$ be as large as possible such that $c_{r,\ell} \neq 0$ for some $\ell \in \mathbb{Z}/e\mathbb{Z}$. To see that $(d_{k,\ell}) \neq \mathbf{0}$, note that, for each $\ell \in \mathbb{Z}/e\mathbb{Z}$,

$$d_{r,\ell} = c_{r,\ell} - \sum_{s=r}^m V_r^s(\gamma^{p^\ell}) \cdot c_{s,\ell+1} = c_{r,\ell} - V_r^r(\gamma^{p^\ell}) \cdot c_{r,\ell+1}$$

because $c_{s,\ell+1} = 0$ whenever $s > r$, and we see just as at the end of proof of Proposition 3.1 that the $d_{r,\ell}$ cannot be zero for all $\ell \in \mathbb{Z}/e\mathbb{Z}$ because this would imply that every $c_{r,\ell} = 0$, contradicting our choice of r . Moreover, we also cannot have $d_{k,\ell} = c_{k,\ell}$ for every $1 \leq k \leq m$ and $\ell \in \mathbb{Z}/e\mathbb{Z}$, for this would also imply that $c_{r,\ell} = 0$ for every $\ell \in \mathbb{Z}/e\mathbb{Z}$, again contradicting our choice of r . $\square$

Definition 4.15. With notation as in Lemma 4.14, the inverse of $\mathcal{D}_Y^{(m)}$ is denoted by $\mathcal{L}_Y^{(m)}$.

Definition 4.16. For $f \in \mathbb{K}(x)$ and $\tau \in \mathcal{T}_M$ with $\tau \subset \mathbb{K}_t^\times$, the Mahler discrete residue of f at τ of degree $k \in \mathbb{N}$ is the vector $\text{dres}(f, \tau, k) \in \bigoplus_{\alpha \in \tau} \mathbb{K}$ defined in terms of the $d_Y(k, n, i, \ell)$ in the partial fraction decomposition of f_τ in Lemma 4.12 as follows.

Set $\text{dres}(f, \tau, k) := \mathbf{0}$ if $\tau \notin \text{supp}(f)$ or if $k > m$. For $\tau \in \text{supp}(f)$ and $\alpha \in \tau$, the component $\text{dres}(f, \tau, k)_\alpha := 0$ whenever the smallest integer $r \in \mathbb{Z}_{\geq 0}$ such that $\alpha^{p^r} \in C(\tau)$ is different from h .

If $h = 0$, then for $1 \leq k \leq m$ and $\alpha = \gamma^{p^\ell} \in C(\tau)$ with $\ell \in \mathbb{Z}/e\mathbb{Z}$, the component

$$\text{dres}(f, \tau, k)_{\gamma^{p^\ell}} := d_Y(k, 0, 0, \ell).$$

If $h \geq 1$, then for $1 \leq k \leq m$ and $\alpha = \zeta_{p^h}^i \gamma^{p^{\ell-\pi_e(h)}}$ with $i \in \mathbb{Z}/p^h\mathbb{Z}$ such that $p \nmid i$ and $\ell \in \mathbb{Z}/e\mathbb{Z}$, the component

$$\text{dres}(f, \tau, k)_\alpha := \hat{d}_Y(k, h, i, \ell); \quad \text{where we set}$$

$$\hat{d}_Y(k, 0, 0, \ell) := c_Y(k, \ell), \quad \text{with} \quad (4.11)$$

$$(c_Y(k, \ell))_{\substack{1 \leq k \leq m \\ \ell \in \mathbb{Z}/e\mathbb{Z}}} := \mathcal{L}_Y^{(m)} \left((d_Y(k, 0, 0, \ell))_{\substack{1 \leq k \leq m \\ \ell \in \mathbb{Z}/e\mathbb{Z}}} \right) \quad (4.12)$$

for the linear map $\mathcal{L}_Y^{(m)}$ in Definition 4.15; and for $1 \leq n \leq h$ and $i \in \mathbb{Z}/p^n\mathbb{Z}$ with $p \nmid i$ we define recursively (in n):

$$\hat{d}_Y(k, n, i, \ell) := d_Y(k, n, i, \ell) + \quad (4.13)$$

$$+ \sum_{s=k}^m V_k^s(\zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}}) \cdot \hat{d}_Y(s, n-1, \pi_{n-1}^n(i), \ell),$$

(cf. (4.9)) where the V_k^s are as in (2.4).

Remark 4.17. Note that the definition of $\text{dres}(f, \tau, k)$ for $\tau \subset \mathbb{K}_t^\times$ given above is independent of the choice of $\gamma \in C(\tau)$, because for $\tilde{\gamma} = \gamma^{p^j}$ with $j \in \mathbb{Z}/e\mathbb{Z}$ we have $\zeta_{p^h}^i \tilde{\gamma}^{p^{\ell-\pi_e(h)}} = \zeta_{p^h}^i \gamma^{p^{\ell+j-\pi_e(h)}} =: \alpha$. The equality of the expressions

$$\hat{d}_Y(k, h, i, \ell) = \text{dres}(f, \tau, k)_\alpha = \hat{d}_Y(k, h, i, \ell + j)$$

follows from (4.7), after observing that $\mathcal{D}_{\tilde{\gamma}}^{(m)} \circ \text{cyc}_j = \mathcal{D}_Y^{(m)}$, where $\text{cyc}_j : \mathbb{K}^{m \times e} \rightarrow \mathbb{K}^{m \times e} : (c_{k,\ell}) \mapsto (c_{k,\ell+j})$ for $j \in \mathbb{Z}/e\mathbb{Z}$. It follows that $\text{cyc}_j \circ \mathcal{L}_Y^{(m)} = \mathcal{L}_{\tilde{\gamma}}^{(m)}$ and therefore $\hat{d}_Y(k, n, i, \ell) = \hat{d}_Y(k, n, i, \ell+j)$ for every $0 \leq n \leq h$.

4.3 Proof of the Main Theorem

Our proof of Proposition 4.18 below follows a strategy similar to that of [14, Propositions 2.5 and 2.10]: we add to f_τ a sequence of Mahler summable elements to eventually obtain a Mahler reduction $\tilde{f}_\tau$ whose apparent dispersion is 0, and then use Corollary 3.2 to conclude that f_τ is Mahler summable if and only if this $\tilde{f}_\tau = 0$.

There is a wrinkle: for $\tau \in \text{supp}(f)$ in the torsion case $\tau \subset \mathbb{K}_t^\times$ such that the height $h(f, \tau) = 0$ (see Definition 4.6), $\text{disp}(f, \tau) = \infty$ by Lemma 2.16. Corollary 3.2 remains silent in this case, for which we provide a specialized argument relying on Lemma 4.14.

PROPOSITION 4.18. *For $f \in \mathbb{K}(x)$ and $\tau \in \mathcal{T}_M$, the component f_τ is Mahler summable if and only if $\text{dres}(f, \tau, k) = \mathbf{0}$ for every $k \in \mathbb{N}$.*

PROOF. The Proposition is trivial for $\tau \notin \text{supp}(f) \Leftrightarrow f_\tau = 0$. Assume from now on that $\tau \in \text{supp}(f)$. The proofs in the different cases $\tau \not\subset \mathbb{K}_t^\times$ versus $\tau \subset \mathbb{K}_t^\times$ proceed in parallel below.

Write f_τ as in Lemma 4.8 if $\tau \not\subset \mathbb{K}_t^\times$; and as in Lemma 4.12 if $\tau \subset \mathbb{K}_t^\times$. Let us define recursively: $g_\tau^{(0)} := 0$; and, if $h := h(f, \tau) \geq 1$ as in Definition 4.6, then for $0 \leq n \leq h-1$ set

$$g_\tau^{(n+1)} := g_\tau^{(n)} + \sum_{k=1}^m \left(\sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{\hat{c}_Y(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k} \right)$$

in case $\tau \not\subset \mathbb{K}_t^\times$, with $\hat{c}_Y(k, n, i)$ as in (4.5); and

$$g_\tau^{(n+1)} := g_\tau^{(n)} + \sum_{k=1}^m \left(\sum'_{i \in \mathbb{Z}/p^n\mathbb{Z}} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\hat{d}_Y(k, n, i, \ell)}{(x - \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}})^k} \right) \right)$$

in case $\tau \subset \mathbb{K}_t^\times$, with $\hat{d}_Y(k, n, i, \ell)$ as in (4.11) for $n = 0$ and as in (4.13) for $n \geq 1$. Setting $\tilde{f}_\tau^{(n)} := f_\tau + \Delta(g_\tau^{(n)})$, an induction argument then shows that, for every $0 \leq n \leq h$,

$$\begin{aligned} \tilde{f}_\tau^{(n)} &= \sum_{k=1}^m \sum_{s=n+1}^{h+1} \left(\sum_{i \in \mathbb{Z}/p^s\mathbb{Z}} \frac{c_Y(k, s, i)}{(x - \zeta_{p^s}^i \gamma^{p^{h-s}})^k} \right) + \\ &+ \sum_{k=1}^m \left(\sum_{i \in \mathbb{Z}/p^n\mathbb{Z}} \frac{\hat{c}_Y(k, n, i)}{(x - \zeta_{p^n}^i \gamma^{p^{h-n}})^k} \right) \end{aligned} \quad (4.14)$$

in case $\tau \not\subset \mathbb{K}_t^\times$; and

$$\begin{aligned} \tilde{f}_\tau^{(n)} &= \sum_{k=1}^m \sum_{s=n+1}^{h+1} \left(\sum'_{i \in \mathbb{Z}/p^s\mathbb{Z}} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{d_Y(k, s, i, \ell)}{(x - \zeta_{p^s}^i \gamma^{p^{\ell-\pi_e(s)}})^k} \right) \right) + \\ &+ \sum_{k=1}^m \left(\sum'_{i \in \mathbb{Z}/p^n\mathbb{Z}} \left(\sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\hat{d}_Y(k, n, i, \ell)}{(x - \zeta_{p^n}^i \gamma^{p^{\ell-\pi_e(n)}})^k} \right) \right) \end{aligned} \quad (4.15)$$

in case $\tau \subset \mathbb{K}_t^\times$. The harmless summand for $s = h+1$ in (4.14) and (4.15) is included so that the sums make sense for $n = h$, but we set every $c_Y(k, h+1, i) := 0$ in (4.14) and every $d_Y(k, h+1, i, \ell) := 0$ in (4.15). The induction argument is straightforward, requiring only: the recursive definition of the coefficients $\hat{c}_Y(k, n, i)$ in (4.5), and the

computation (4.4), in case $\tau \notin \mathbb{K}_t^\times$; the recursive definition of the coefficients $\hat{d}_Y(k, n, i, \ell)$ in (4.11) and (4.13), and the computations (4.8) and (4.9), in case $\tau \in \mathbb{K}_t^\times$; and a moderate amount of space and courage to write it down in detail in each case. It then follows from (4.14) and Definition 4.10 (in case $\tau \notin \mathbb{K}_t^\times$), or from (4.15) and Definition 4.16 (in case $\tau \in \mathbb{K}_t^\times$), that

$$\bar{f}_\tau^{(h)} = \sum_{k=1}^m \sum_{\alpha \in \tau} \frac{\text{dres}(f, \tau, k)_\alpha}{(x - \alpha)^k}, \quad (4.16)$$

which holds uniformly in both cases $\tau \notin \mathbb{K}_t^\times$ and $\tau \in \mathbb{K}_t^\times$. Also in both of these cases we see that $\bar{f}_\tau^{(h)} = f_\tau + \Delta(g_\tau^{(h)})$, and therefore f_τ is Mahler summable if and only if $\bar{f}_\tau^{(h)}$ is Mahler summable.

We claim that $\bar{f}_\tau^{(h)}$ is Mahler summable if and only if $\bar{f}_\tau^{(h)} = 0$. This will establish the Proposition, since $\bar{f}_\tau^{(h)} = 0$ if and only if $\text{dres}(f, \tau, k) = 0$ for all $k \in \mathbb{N}$ by (4.16). The non-trivial implication: $\bar{f}_\tau^{(h)} \neq 0 \Rightarrow \bar{f}_\tau^{(h)}$ is not Mahler summable, is proved in two subcases: (1). if either $\tau \notin \mathbb{K}_t^\times$ or $h \neq 0$; and (2). if both $\tau \in \mathbb{K}_t^\times$ and $h = 0$.

(1). In case $\tau \notin \mathbb{K}_t^\times$, by Definition 4.10 $\bar{f}_\tau^{(h)}$ has no poles outside of $\{\zeta_p^i \gamma \mid i \in \mathbb{Z}/p^h\mathbb{Z}\}$. In case $\tau \in \mathbb{K}_t^\times$ and $h \neq 0$, by Definition 4.16 $\bar{f}_\tau^{(h)}$ has no poles outside of $\{\zeta_p^i \gamma^{p^\ell} \mid \ell \in \mathbb{Z}/e\mathbb{Z} \text{ and } i \in (\mathbb{Z}/p^h\mathbb{Z})^\times\}$ (cf. Lemma 4.5). Thus if either $\tau \notin \mathbb{K}_t^\times$ or $h \neq 0$, $\text{disp}(\bar{f}_\tau^{(h)}, \tau) = 0$. By Corollary 3.2, $\bar{f}_\tau^{(h)}$ is not Mahler summable.

(2). Note that $f_\tau = \bar{f}_\tau^{(h)}$ in (4.16) in this last remaining subcase where $h = 0$ and $\tau \in \mathbb{K}_t^\times$, and the Definition 4.16 gives

$$f_\tau = \sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{d_Y(k, \ell)}{(x - \gamma^{p^\ell})^k} = \sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\text{dres}(f, \tau, k)_{\gamma^{p^\ell}}}{(x - \gamma^{p^\ell})^k},$$

where we write $d_Y(k, \ell)$ in lieu of $d_Y(k, 0, 0, \ell)$, to simplify notation. Since $\tau \in \text{supp}(f)$, we must have $\text{dres}(f, \tau, m) \neq 0$. We claim that f_τ cannot be Mahler summable. To prove the claim, let again $g_\tau^{(1)} := \sum_{k=1}^m \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} c(k, \ell) \cdot (x - \gamma^{p^\ell})^{-k}$ despite having $h = 0$, where the $c_Y(k, \ell)$ are as in (4.12). By the computation (4.8) and the Definition 4.15 of the map $\mathcal{L}_Y^{(m)}$,

$$\check{f}_\tau := f_\tau + \Delta(g_\tau^{(1)}) = \sum_{k=1}^m \sum_{i=1}^{p-1} \sum_{\ell \in \mathbb{Z}/e\mathbb{Z}} \frac{\sum_{s=k}^m V_k^s(\zeta_p^i \gamma^{p^{\ell-1}}) \cdot c_Y(k, \ell)}{(x - \zeta_p^i \gamma^{p^{\ell-1}})^k}.$$

Hence f_τ is Mahler summable if and only if $\check{f}_\tau$ is Mahler summable. In particular, if $\check{f}_\tau = 0$, then f_τ is Mahler summable. On the other hand, if $\check{f}_\tau \neq 0$, then $\text{disp}(\check{f}_\tau, \tau) = 0$, in which case $\check{f}_\tau$ cannot be Mahler summable by Corollary 3.2. Hence f_τ is Mahler summable if and only if $\check{f}_\tau = 0$. Let us now show that $f_\tau \neq 0 \Rightarrow \check{f}_\tau \neq 0$.

In any case, the partial fraction coefficients of $\check{f}_\tau$ satisfy

$$\begin{aligned} \sum_{s=k}^m V_k^s(\zeta_p^i \gamma^{p^\ell}) \cdot c_Y(k, \ell + 1) &= \\ &= \zeta_p^{ik} \cdot \sum_{s=k}^m V_k^s(\gamma^{p^\ell}) \cdot c_Y(k, \ell + 1) = \zeta_p^{ik} \cdot (c_Y(k, \ell) - d_Y(k, \ell)) \end{aligned}$$

where the first equality follows from $V_k^s(\zeta_p^i \gamma^{p^\ell}) = \zeta_p^{ik} V_k^s(\gamma^{p^\ell})$ independently of s by Lemma 2.17, and the second equality follows from the Definition 4.15 of $\mathcal{L}_Y^{(m)}$. By Lemma 4.14, since the map $\mathcal{D}_Y^{(m)}$ has no non-trivial fixed points, we cannot have $c_Y(k, \ell) = d_Y(k, \ell)$ for every k and ℓ unless all $d_Y(k, \ell) = 0$. So indeed $f_\tau \neq 0 \Rightarrow \check{f}_\tau \neq 0$. $\square$

Remark 4.19. For $f \in \mathbb{K}(x)$, $\tau \in \text{supp}(f) \cap \mathcal{T}_M$, and $h := h(f, \tau)$ as in Definition 4.6, the $\bar{f}_\tau^{(h)}, g_\tau^{(h)} \in \mathbb{K}(x)_\tau$ constructed in the proof of Proposition 4.18 are the τ -components of the $\bar{f}, g \in \mathbb{K}(x)$ in (1.1).

PROOF OF THE MAIN THEOREM. Let $f \in \mathbb{K}(x)$. By Lemma 2.2, f is Mahler summable if and only if both f_L and f_T are Mahler summable. By Proposition 4.2, f_L is Mahler summable if and only if $\text{dres}(f, \infty) = 0$. By Lemma 2.14, f_T is Mahler summable if and only if f_τ is Mahler summable for all $\tau \in \mathcal{T}_M$. By Proposition 4.18, f_τ is Mahler summable if and only if $\text{dres}(f, \tau, k) = 0$ for all $k \in \mathbb{N}$. $\square$

4.4 Mahler reduction

We can now define the Mahler reduction (1.1): $\bar{f} = f + \Delta(g)$ promised in the introduction for any $f \in \mathbb{K}(x)$, in terms of the decompositions $\bar{f} = \bar{f}_L + \bar{f}_T$ and $g = g_L + g_T$ as in (2.1), by setting

$$\begin{aligned} \bar{f}_L &:= \sum_{\theta \in \mathbb{Z}/\mathcal{P}} \bar{f}_\theta \quad \text{and} \quad g_L := \sum_{\theta \in \mathbb{Z}/\mathcal{P}} g_\theta; \quad \text{and} \\ \bar{f}_T &:= \sum_{\tau \in \text{supp}(f)} \bar{f}_\tau^{(h(f, \tau))} \quad \text{and} \quad g_T := \sum_{\tau \in \text{supp}(f)} g_\tau^{(h(f, \tau))} \end{aligned}$$

as in Remark 4.3 and Remark 4.19. It is clear from the definitions that $c \cdot f = c \cdot \bar{f}$ for $c \in \mathbb{K}$. Setting $\bar{f}_1 + \bar{f}_2 := \bar{f}_1 + \bar{f}_2$ defines a $\mathbb{K}$ -linear structure on $\{\bar{f} \mid f \in \mathbb{K}(x)\}$ such that $\nabla : f \mapsto \bar{f}$ is $\mathbb{K}$ -linear and has the desired property that $\ker(\nabla) = \text{im}(\Delta)$.

5 EXAMPLES

Let us illustrate the Mahler discrete residues at Mahler trees in two small examples, with notation as in Example 2.9. Example 5.1 gives a Mahler summable f in the non-torsion case $\tau \notin \mathbb{K}_t^\times$. Example 5.2 gives a non-Mahler summable f in the torsion case $\tau \in \mathbb{K}_t^\times$.

Example 5.1. Let $\tau = \tau(2)$, and consider the following $f = f_\tau$ with $\text{sing}(f, \tau) = \{2, \sqrt[3]{2}, \zeta_3 \sqrt[3]{2}, \zeta_3^2 \sqrt[3]{2}\}$. By Definition 4.6, $h = h(f, \tau) = 1$.

$$\begin{aligned} f &= \frac{-x^6 + 4x^3 + x^2 - 4x}{(x-2)^2(x^3-2)^2} = \sum_{k=1}^2 \sum_{n=0}^1 \sum_{i=0}^{3^n-1} \frac{c_Y(k, n, i)}{(x - \zeta_{3^n}^i \sqrt[3]{2}^{3^{1-n}})^k} \\ &= \frac{-1}{(x-2)^2} + \frac{1}{18\sqrt[3]{2}} \cdot \sum_{i=0}^2 \frac{\zeta_3^{2i}}{(x - \zeta_3^i \sqrt[3]{2})^2} - \frac{1}{9\sqrt[3]{4}} \cdot \sum_{i=0}^2 \frac{\zeta_3^i}{x - \zeta_3^i \sqrt[3]{2}}, \end{aligned}$$

for $\gamma := \sqrt[3]{2}$ as in Lemma 4.8. By Definition 4.10, for $0 \leq i \leq 2$:

$$\begin{aligned} \text{dres}(f, \tau, 1)_{\zeta_3^i \gamma} &= \frac{-\zeta_3^i}{9\gamma^2} + V_1^2(\zeta_3^i \gamma) \cdot (-1) + V_1^1(\zeta_3^i \gamma) \cdot 0 \\ &= \frac{-\zeta_3^i}{9\gamma^2} - \frac{2}{9} \cdot (\zeta_3^i \gamma)^{-5} = 0; \quad \text{and} \end{aligned}$$

$$\text{dres}(f, \tau, 2)_{\zeta_3^i \gamma} = \frac{\zeta_3^{2i}}{18\gamma} + V_2^2(\zeta_3^i \gamma) \cdot (-1) = \frac{\zeta_3^{2i}}{18\gamma} - 3^{-2}(\zeta_3^i \gamma)^{-4} = 0;$$

by Lemma 2.17 and Example 2.20. Therefore f should be Mahler summable. And indeed, $f = \Delta((x-2)^{-2})$.

Example 5.2. Let $\tau = \tau(\zeta_4)$, and consider the following $f = f_\tau$ with $\text{sing}(f, \tau) = \{\zeta_4^{\pm 1}, \zeta_{12}^{\pm 1}, \zeta_{12}^{\pm 5}\}$. By Definition 4.6, $h = h(f, \tau) = 1$.

$$f = \frac{1}{x^6 + 1} = \frac{1}{6} \left(\frac{\zeta_4^3}{x - \zeta_4} + \frac{\zeta_4}{x - \zeta_4^3} + \frac{\zeta_{12}^7}{x - \zeta_{12}} + \frac{\zeta_{12}}{x - \zeta_{12}^7} + \frac{\zeta_{12}^{11}}{x - \zeta_{12}^5} + \frac{\zeta_{12}^5}{x - \zeta_{12}^{11}} \right).$$

The map $\mathcal{L}_{\zeta_4}^{(1)}$ in Definition 4.15 sends $(\frac{\zeta_4}{6}, \frac{\zeta_4}{6}) \mapsto (\frac{\zeta_4}{4}, \frac{\zeta_4}{4})$. By Definition 4.16, for $1 \leq i \leq 2$; $\ell \geq 1$; with $\alpha_{i, \ell} := \zeta_3^i \zeta_4^{\ell-1} = \zeta_{12}^{4i+3\ell}$:

$$\begin{aligned} \text{dres}(f, \tau, 1)_{\alpha_{i, \ell}} &= \frac{1}{6} \cdot \alpha_{i, \ell+1} + V_1^1(\alpha_{i, \ell}) \cdot \frac{1}{4} \cdot \zeta_4^{3\ell-1} \\ &= \frac{1}{6} \cdot \alpha_{i, \ell+1} + \frac{1}{12} \cdot (\alpha_{i, \ell})^{-2} \cdot \zeta_4^{3\ell-1} = \frac{1}{4} \cdot \alpha_{i, \ell+1} \neq 0, \end{aligned}$$

by Lemma 2.17. Therefore f is not Mahler summable.

Acknowledgements. We thank the anonymous referees for their careful reading and valuable suggestions. We also thank Manuel Kauers for pointing us to relevant literature at a crucial juncture.

REFERENCES

- [1] Sergei A. Abramov. 1971. On the summation of rational functions. *U. S. S. R. Comput. Math. and Math. Phys.* 11, 4 (1971), 324–330. [https://doi.org/10.1016/0041-5553\(71\)90028-0](https://doi.org/10.1016/0041-5553(71)90028-0)
- [2] Sergei A. Abramov. 1974. Solution of linear finite-difference equations with constant coefficients in the field of rational functions. *U. S. S. R. Comput. Math. and Math. Phys.* 14, 4 (1974), 247–251. [https://doi.org/10.1016/0041-5553\(74\)90095-0](https://doi.org/10.1016/0041-5553(74)90095-0)
- [3] Carlos E. Arreche. 2017. Computation of the difference-differential Galois group and differential relations among solutions for a second-order linear difference equation. *Communications in Contemporary Mathematics* 19, 06 (2017), 1650056. <https://doi.org/10.1142/S0219199716500565>
- [4] Carlos E. Arreche and Yi Zhang. 2022. Computing differential Galois groups of second-order linear q-difference equations. *Advances in Applied Mathematics* 132 (2022), 102273. <https://doi.org/10.1016/j.aam.2021.102273>
- [5] Alin Bostan, Shaoshi Chen, Frédéric Chyzak, and Ziming Li. 2010. Complexity of Creative Telescoping for Bivariate Rational Functions. In *Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation* (Munich, Germany) (ISSAC '10). Association for Computing Machinery, New York, NY, USA, 203–210. <https://doi.org/10.1145/1837934.1837975>
- [6] Alin Bostan, Shaoshi Chen, Frédéric Chyzak, Ziming Li, and Guoce Xin. 2013. Hermite Reduction and Creative Telescoping for Hyperexponential Functions. In *Proceedings of the 38th International Symposium on Symbolic and Algebraic Computation* (Boston, Maine, USA) (ISSAC '13). Association for Computing Machinery, New York, NY, USA, 77–84. <https://doi.org/10.1145/2465506.2465946>
- [7] Alin Bostan, Pierre Lairez, and Bruno Salvy. 2013. Creative Telescoping for Rational Functions Using the Griffiths' Dwork Method. In *Proceedings of the 38th International Symposium on Symbolic and Algebraic Computation* (Boston, Maine, USA) (ISSAC '13). Association for Computing Machinery, New York, NY, USA, 93–100. <https://doi.org/10.1145/2465506.2465935>
- [8] Manuel Bronstein. 1992. Formulas for series computations. *Applicable Algebra in Engineering, Communication and Computing* 2, 3 (1992), 195–206. <https://doi.org/10.1007/BF01294333>
- [9] Xavier Caruso. 2021. A theory of residues for skew rational functions. *Journal de l'École polytechnique — Mathématiques* 8 (2021), 1159–1192. <https://doi.org/10.5802/jep.169>
- [10] Xavier Caruso and Amaury Durand. 2021. Duals of linearized Reed-Solomon codes. <https://doi.org/10.48550/ARXIV.2110.12675>
- [11] Shaoshi Chen. 2018. Bivariate Extensions of Abramov's Algorithm for Rational Summation. In *Advances in Computer Algebra*, Carsten Schneider and Eugene Zima (Eds.). Springer International Publishing, Cham, 93–104.
- [12] Shaoshi Chen, Ruyong Feng, Pingchuan Ma, and Michael F. Singer. 2021. Separability Problems in Creative Telescoping. In *Proceedings of the 2021 on International Symposium on Symbolic and Algebraic Computation* (Virtual Event, Russian Federation) (ISSAC '21). Association for Computing Machinery, New York, NY, USA, 83–90. <https://doi.org/10.1145/3452143.3465514>
- [13] Shaoshi Chen, Qing-Hu Hou, George Labahn, and Rong-Hua Wang. 2016. Existence Problem of Telescopers: Beyond the Bivariate Case. In *Proceedings of the ACM on International Symposium on Symbolic and Algebraic Computation* (Waterloo, ON, Canada) (ISSAC '16). Association for Computing Machinery, New York, NY, USA, 167–174. <https://doi.org/10.1145/2930889.2930895>
- [14] Shaoshi Chen and Michael F. Singer. 2012. Residues and Telescopers for Bivariate Rational Functions. *Adv. Appl. Math.* 49, 2 (aug 2012), 111–133. <https://doi.org/10.1016/j.aam.2012.04.003>
- [15] Frédéric Chyzak, Thomas Dreyfus, Philippe Dumas, and Marc Mezzarobba. 2018. Computing solutions of linear Mahler equations. *Math. Comp.* 87, 314 (2018), 2977–3021. <https://doi.org/10.1090/mcom/3359>
- [16] Philippe Flajolet and Robert Sedgewick. 2009. *Analytic Combinatorics*. Cambridge University Press, Cambridge, UK. <https://doi.org/10.1017/CBO9780511801655>
- [17] Charlotte Hardouin and Michael F. Singer. 2008. Differential Galois theory of linear difference equations. *Math. Ann.* 342, 2 (2008), 333–377. <https://doi.org/10.1007/s00208-008-0238-z>
- [18] Charlotte Hardouin and Michael F. Singer. 2021. On differentially algebraic generating series for walks in the quarter plane. *Selecta Mathematica* 27, 5 (2021), 89. <https://doi.org/10.1007/s00029-021-00703-9>
- [19] Qing-Hu Hou and Rong-Hua Wang. 2015. An Algorithm for Deciding the Summability of Bivariate Rational Functions. *Adv. Appl. Math.* 64, C (mar 2015), 31–49. <https://doi.org/10.1016/j.aam.2014.11.002>