



REEFS, CLIMATE CHANGE PAPER

Help Me, Symbionts, You're My Only Hope: Approaches to Accelerate our Understanding of Coral Holobiont Interactions

Colleen B. Bove¹, Maria Valadez Ingersoll and Sarah W. Davies

Department of Biology, Boston University, Boston, MA 02215, USA

¹E-mail: colleenbove@gmail.com

Synopsis Tropical corals construct the three-dimensional framework for one of the most diverse ecosystems on the planet, providing habitat to a plethora of species across taxa. However, these ecosystem engineers are facing unprecedented challenges, such as increasing disease prevalence and marine heatwaves associated with anthropogenic global change. As a result, major declines in coral cover and health are being observed across the world's oceans, often due to the breakdown of coral-associated symbioses. Here, we review the interactions between the major symbiotic partners of the coral holobiont—the cnidarian host, algae in the family Symbiodiniaceae, and the microbiome—that influence trait variation, including the molecular mechanisms that underlie symbiosis and the resulting physiological benefits of different microbial partnerships. In doing so, we highlight the current framework for the formation and maintenance of cnidarian–Symbiodiniaceae symbiosis, and the role that immunity pathways play in this relationship. We emphasize that understanding these complex interactions is challenging when you consider the vast genetic variation of the cnidarian host and algal symbiont, as well as their highly diverse microbiome, which is also an important player in coral holobiont health. Given the complex interactions between and among symbiotic partners, we propose several research directions and approaches focused on symbiosis model systems and emerging technologies that will broaden our understanding of how these partner interactions may facilitate the prediction of coral holobiont phenotype, especially under rapid environmental change.

Introduction

Reef-building corals are the framework engineers for coral reef ecosystems that build one of the most biodiverse habitats in the world (Reaka-Kudla *et al.* 1996), and are economically and ecologically critical (Moberg and Folke 1999; Costanza *et al.* 2014). Dramatic losses in coral cover are being documented globally as anthropogenic greenhouse gas emissions increase and other direct human impacts drive environmental change (De'ath *et al.* 2012; Eddy *et al.* 2021). When seawater temperatures rise or chemistry changes, these shifts can lead to a loss of the coral's symbiotic relationship with its endosymbiotic algae in a process termed coral bleaching (Glynn 1991) and these episodes are increasing in frequency and severity (Sully *et al.* 2019). Predicting coral resilience to changing oceans has become a critical goal of coral reef research, yet, we are just beginning to disentangle the key mechanisms underlying symbiosis establishment, maintenance, and loss,

which are crucial processes in understanding coral resilience.

Understanding symbiosis outcomes under changing oceans is challenging because corals are holobionts—assemblages of the coral host and many other living partners, including millions of single-celled dinoflagellates (family Symbiodiniaceae, (LaJeunesse *et al.* 2018)) living inside coral gastrodermal cells—that together form a discrete ecological unit (Rosenberg *et al.* 2007; Rosenberg and Zilber-Rosenberg 2018). In tropical, oligotrophic waters, these symbiotic algae provide essential organic byproducts to the host from photosynthesis (Muscatine and Cernichiari 1969). Algal symbiont diversity has become better appreciated thanks to molecular genetics approaches (reviewed in Quigley *et al.* 2018; Davies *et al.* 2022) and whole-genome datasets (Dougan *et al.* 2022), which have resulted in taxonomic revisions (LaJeunesse *et al.* 2018, 2021; Nitschke *et al.* 2020; Pochon and LaJeunesse 2021). However, how this

genetic variation predicts functional variation remains an open question. Unique algal isolates can be functionally divergent (e.g., [Parkinson and Baums 2014](#); [Beltrán et al. 2021](#)), which is perhaps predictable given that reefs are highly variable and algae exhibit limited dispersal ([Fitt et al. 1981](#)), hypothetically leading to adaptive divergence in algal communities ([Howells et al. 2011](#)) or long-term acclimatization ([Torda et al. 2017](#)).

Although some coral species exhibit specificity for particular algae ([Thornhill et al. 2014](#); [Hume et al. 2020](#)), others can associate with a diverse array of algae that can potentially shuffle under different environmental conditions ([Baker and Romanski 2007](#)). Hosts can associate with “homologous” (i.e., symbiont strains hosted under ambient conditions) or “heterologous” (i.e., opportunistic strains hosted following disturbance) algal strains ([Davy, Lucas, and Turner 1997](#); [Weis et al. 2001](#)). A host may associate with multiple strains at any given time, the proportion of which can vary under changing environments (see [Reich et al. 2017](#); [Matthews et al. 2018](#)). This potential flexibility is important because not all host–symbiont pairings are equally resilient under stress ([Berkelmans and Van Oppen 2006](#); [Hoadley et al. 2019](#)). For example, whole-genome resequencing of over 250 corals during a bleaching event found that Symbiodiniaceae associations (*Cladocopium* vs. *Durusdinum*) were a stronger predictor of bleaching than host genetic variation ([Fuller et al. 2020](#)). Limited work also suggests that host–symbiont preference could have a host genetic basis ([Quigley et al. 2019](#); [Reich et al. 2021](#)), therefore symbiont-driven thermal tolerance may have higher heritability than expected. Thus, changes in symbiont associations (i.e., symbiont shuffling ([Baker 2003](#))) may enhance resilience to future stress events.

Recently, the roles of other members of the microbiome (e.g., bacteria, viruses, fungi, archaea, etc.) in maintaining holobiont health have become more appreciated ([Pollock et al. 2018](#); [van Oppen and Blackall 2019](#)). The majority of microbiome research has focused on bacteria, which, similar to the algae, provide hosts with essential nutrients ([Robbins et al. 2019](#)), nitrogen cycling ([Rädecker et al. 2015](#)) and carbon cycling ([Kimes et al. 2010](#)). The diversity and community composition of microorganisms within a coral interact to support a healthy holobiont through resource allocation ([Bourne et al. 2016](#)), particularly under stress ([Webster and Reusch 2017](#)). Indeed, corals exhibiting higher microbiome flexibility (i.e., restructuring of microbial community) may possess a unique ecological advantage to better acclimate to environmental change ([Voolstra and Ziegler 2020](#)). These differences between ‘microbiome regulators’ (corals that maintain stable microbiomes; e.g., *Pocillopora verrucosa*) and

‘microbiome conformers’ (corals that modulate microbiomes in response to environment; e.g., *Acropora hemprichii*) may help to predict holobiont phenotypes under stress ([Ziegler et al. 2019](#)).

Clearly, each coral holobiont is a complex metaorganism, and understanding—and ultimately predicting—its response to future change depends on an integrated understanding of all partners and their interactions. Here, we present what is known about the diversity of corals and how their different symbiotic partners interact to influence holobiont physiology and propose a set of research directions that leverage emerging cnidarian model systems to further disentangle how these interactions might influence coral persistence under rapid change.

Host genetic diversity and its influence on holobiont phenotype

Coral host genetic diversity both within and between species is vast, and this diversity can lead to varied responses to environmental stressors. Sequencing technologies for non-model systems have improved our ability to detect this diversity and estimate genetic divergence between hosts across spatial scales, such as depth ([Prada and Hellberg 2013](#); [Serrano et al. 2016](#)) and environmental gradients ([Kenkel and Matz 2016](#)). Similarly, genomic sequencing has facilitated further identification of sibling species (i.e., cryptic species) within previously presumed coral species ([Bickford et al. 2007](#)). Cryptic speciation is a well-known phenomenon in corals ([Knowlton 1993](#); [Prada and Hellberg 2013](#); [Forsman et al. 2020](#); [Fifer et al. 2022](#)); however, recent work suggests that these cryptic lineages may exhibit differential responses to global change ([Gómez-Corrales and Prada 2020](#)) and host unique algal communities ([Rose et al. 2021](#)). While coral genetic diversity may not always directly predict phenotypic diversity, phenotypic variation can be heritable (i.e., narrow-sense heritability, h^2) across various coral traits ([Bairos-Novak et al. 2021](#)), including thermal tolerance ([Dixon et al. 2015](#)), settlement responsiveness ([Meyer et al. 2011](#)), calcification ([Jury et al. 2019](#)), and growth ([Kenkel et al. 2015](#)). This genetic variation highlights the need to include multiple genetic backgrounds in mechanistic studies, as each unique genotype may determine a phenotypic response.

As an additional layer of complexity, corals have a bipartite life cycle that involves planula larval dispersal, followed by recruitment to a substrate, and maturation into a benthic, sedentary adult ([Baird et al. 2009](#)). The selective pressures on planula larvae in the water column are likely very different from those experienced by the recruit or adult coral, which may lead to ontogenetic

shifts in selection pressures, diverse symbiont acquisition strategies (Hartmann et al. 2017; Ali et al. 2019; Epstein et al. 2019), and differences in adaptive potential across life stages (reviewed in Putnam 2021). For example, trait heritability has been shown to vary across coral life stages, with adult and larval stages exhibiting higher heritability of bleaching and growth when compared to the juvenile stage (Bairos-Novak et al. 2021). Such differences may constrain evolution through genetic trade-offs between traits and across life stages. These complex life cycles combined with differences in transmission strategies across hosts and partners make predicting coral performance under changing environments challenging.

The role of the microbiome in coral holobiont phenotype and resilience

Corals are composed of several microhabitats (e.g., tissue, mucus, gastrovascular cavity, and skeleton) hosting different microorganisms (Sweet et al. 2011; Pollock et al. 2018) that support functions within the holobiont. For example, anoxic conditions within coral gastric cavities at night likely support nitrogen fixation by anaerobic microorganisms (Bourne et al. 2016; Bove et al. 2020), which can in turn facilitate photosynthesis of Symbiodiniaceae (Rädecker et al. 2015). However, these compartments are often homogenized when sampling corals (Hughes et al. 2022), potentially concealing more specific microbiome signatures (Pollock et al. 2018; Armitage and Jones 2019; Biagi et al. 2020). Therefore, our understanding of how the spatial arrangement of the microbiome shapes holobiont function remains largely unknown (but see Wooldridge 2010; Morris et al. 2019; Jiang et al. 2021).

The coral microbiome is influenced by many factors, including host developmental stage (Sharp et al. 2012; Damjanovic et al. 2020), spatial scales (e.g., across an individual (Rohwer et al. 2002) or geographic regions (Williams et al. 2022)), and environmental change (van Oppen and Blackall 2019; Voolstra and Ziegler 2020)—especially thermal stress (reviewed in Maire et al. 2022). However, interactions between microbes and other holobiont partners are particularly important for understanding coral resilience. For example, recent work identified relatively stable bacterial assemblages associated with *Durusdinium trenchii* even under thermal stress that may promote thermal resilience (Lawson et al. 2018; Camp et al. 2020). Similar to warming, the effects of ocean acidification on coral microbiomes is species-specific (Meron et al. 2012; Biagi et al. 2020), with changes mostly occurring in the mucus layer (Glasl et al. 2016; Biagi et al. 2020). Because environmental disturbances will continue to occur as global change

persists, a deeper understanding of how multiple stressors interact with these microbial members of the coral holobiont is needed.

While significant advances in our understanding of Symbiodiniaceae–host (see Mieog et al. 2009; Morris et al. 2019) and bacteria–host (see Ainsworth et al. 2010; Webster and Reusch 2017) interactions have been made, the relationships between different members of the microbiome (i.e., Symbiodineaceae and bacteria) remain less studied (see Matthews et al. 2020). Recent characterization of bacterial communities associated with cultured Symbiodiniaceae have showcased high-bacterial diversity across algal strains with three common bacterial groups serving as the core members of Symbiodiniaceae–bacterial assemblages (Lawson et al. 2018). The most abundant core member, *Labrenzia*, produces dimethylsulfoniopropionate (DMSP), which may play a role in the coral stress response (Jones and King 2015). DMSP production has historically been attributed to Symbiodiniaceae; however, if *Labrenzia* are producing DMSP, then they may play a pivotal role in holobiont health. This group was also found to increase in abundance under thermal stress in cultures of *Cladocopium* spp. (Camp et al. 2020), suggesting it may function in the algal stress response as well. These critical Symbiodiniaceae–bacteria interactions are just beginning to be explored and require further elucidation.

Known molecular mechanisms underlying cnidarian–algal symbiosis

Symbiotic cnidarians are faced with the challenge of establishment and maintenance of their algal symbionts while surrounded by a soup of microbes (Hernandez-Agreda et al. 2017). Cnidarians rely on innate immunity to differentiate between beneficial and pathogenic microbes (reviewed by Nyholm and Graf 2012). The primary mechanism for the host to detect friend vs. foe is through Pattern Recognition Receptors (PRRs) that are either secreted by, presented on the surface of, or located entirely within animal cells (reviewed in Akira et al. 2006). Extracellular and transmembrane receptors may be important for symbiont recognition and establishment, while cytoplasmic PRRs may be involved in intracellular symbiont maintenance. PRRs recognize and bind specific patterns on microbial cells and initiate intracellular signaling cascades that modulate the host's immune system (reviewed in Mansfield and Gilmore 2019).

One PRR–microbe interaction implicated in the cnidarian–Symbiodiniaceae symbiosis occurs between host lectin receptors and symbiont glycans (Tortorelli, Rautengarten, et al. 2022). Symbiodiniaceae species have different cell-surface glycan assemblages, which

may facilitate highly specific recognition and establishment (Logan et al. 2010). Phylogenetic analyses have implicated PRRs in symbiosis as demonstrated by an independent expansion of PRRs in symbiotic, but not non-symbiotic cnidarians (Baumgarten et al. 2015; Emery et al. 2021). Cnidarian ficolin-like proteins (CnFLs) are PRRs unique to symbiotic Cnidaria hypothesized to function in highly specific recognition and uptake of compatible symbiont types. In the sea anemone model for symbiosis *Exaiptasia pallida* (Aiptasia), CnFLs transition from upregulated in aposymbiotic individuals to downregulated following establishment of symbiosis with homologous symbionts (Baumgarten et al., 2015).

Symbiosis establishment with (homologous) algal symbionts induces anti-inflammatory immune pathways and reduces pro-inflammatory immune pathways, as measured through Reactive Oxidative Species (ROS) production (reviewed by Weis et al. 2008). Indeed, symbiosis between Aiptasia and homologous symbionts is associated with lower expression of pro-inflammatory gene pathways relative to aposymbiotic Aiptasia (Lehnert et al. 2014). The cytokine Transforming Growth Factor β (TGF β) can suppress the inflammatory response (reviewed in Yoshimura et al. 2010) by downregulating certain cnidarian immune pathways (Fuess et al. 2020). Induction of TGF β and repression of the immune response appear necessary for establishment of homologous symbionts in Aiptasia (Detournay et al. 2012). Nuclear Factor- κ B (NF- κ B) is an evolutionarily conserved transcription factor in the innate immune system activated by PRRs and inhibited by TGF β in Aiptasia (reviewed in Gilmore 2006; Mansfield and Gilmore 2019). NF- κ B is downregulated following establishment of homologous symbionts in adult and larval Aiptasia compared to aposymbiotic larvae and adults (Wolfowicz et al. 2016; Mansfield et al. 2017). Aposymbiotic branches of the facultative coral *Oculina arbuscula* also exhibit enrichment of gene pathways involved in NF- κ B immunity compared to symbiotic branches (Rivera and Davies 2021). Further, field-collected *A. palmata* exhibited upregulation of NF- κ B transcription under short-term thermal stress (DeSalvo et al. 2010).

Interestingly, NF- κ B is not downregulated in adult and larval Aiptasia nor adult *P. damicornis* harboring heterologous symbionts, indicating that the host fails to detect heterologous symbionts through the same mechanism as homologous symbionts (Mansfield et al. 2019). Additionally, Aiptasia populated by heterologous symbionts induce multiple oxidative response pathways and accumulate more ROS and Reactive Nitrogen Species compared to Aiptasia hosting homologous symbionts (Lehnert et al. 2014; Matthews et al. 2017). These data

suggest that hosts associated with heterologous symbionts are chronically responding to oxidative stress even under ambient conditions, but it may also suggest conferred tolerance through antioxidant priming (Matthews et al. 2017). Though hosting heterologous symbiont strains appears to confer maintained immunity, nutritional benefits have been associated with hosting homologous strains (Starzak et al. 2014; Rädecker et al. 2018). Homologous-hosting Aiptasia assimilate more symbiont metabolites than those hosting heterologous symbionts (Matthews et al. 2017, 2018). Symbiosis maintenance involves host control over Symbiodinaceae cell density through regulation of the symbiont's cell cycle and nitrogen cycling in both *O. arbuscula* and Aiptasia (Rivera and Davies 2021; Gorman et al. 2022). It is also noteworthy that heterologous symbiont density is lower (at least in Aiptasia) than homologous symbionts (Matthews et al. 2018; Tsang Min Ching et al. 2022), implicating greater host control over cell proliferation in heterologous partnerships. Further research is required to understand the conference of metabolic benefits by different strains under stress conditions.

In general, this work supports the emerging hypotheses for the molecular mechanisms that might differ between cnidarians harboring homologous and heterologous algal strains, as summarized in Fig. 1. Cnidarians harboring heterologous symbionts are frequently classified as more stress-tolerant than those harboring homologous symbionts because of two hypotheses, which are not mutually exclusive: (1) hosts are able to maintain constitutively higher innate immunity, which allows for immune priming to better quench increased ROS under stress, and (2) lower algal symbiont densities in hosts associated with heterologous symbionts leads to less ROS produced. The role of the molecular pathways explored in this section may be further clarified through targeted knock-out, knock-down, and over-expression experiments. Additionally, the roles of specific localized tissues and cell-types in symbiosis control and nutrient exchange will be resolved with further implementation of emerging technologies (e.g., single-cell RNA-seq (scRNAseq)) across many species, lineages, and treatment conditions as discussed below (see *Leveraging Novel Technologies*).

Proposed paths forward to disentangle the functional consequences of partner interactions

To better understand and predict coral symbiotic partnerships under environmental change, we propose several paths forward that would broaden our understanding of holobiont interactions and resulting phenotypes.




Fig. 1 Proposed mechanism of immunity regulation in three scenarios of cnidarian–algal symbiosis. (A) When a homologous symbiont strain is detected by a host cell, PRRs on the host cell surface recognize and bind glycans (G1) presented on the symbiont cell. This high-affinity binding induces an intracellular signaling cascade that downregulates pathways responsible for inflammation, the reactive oxygen species (ROS) response, and the immune response. This repression may be via the inhibition of NF- κ B—and therefore the inflammatory response and nitric oxide (NO) production—by TGF β . Other glycans (G2) bind CnFLs on host cells, which participate in a negative feedback loop. (B) When a heterologous symbiont strain is detected by a host cell, PRRs on the host cell surface have lower binding affinity for symbiont glycans (G3). Therefore, the intracellular signaling cascade is either induced to a lower degree or fails to be induced (depending on the binding affinity between G3 and PRRs). The reduced signaling cascade (and lower TGF β) means that pro-inflammation, the ROS response, and the immune response (characterized by NF- κ B immunity) are not inhibited. This reduced repression may prime host cells for future stress. It remains unclear whether CnFLs can recognize and bind heterologous symbiont glycans (G4). (C) In aposymbiotic hosts, the intracellular signaling cascade involving TGF β repression of NF- κ B is not induced, so inflammation and the NF- κ B immune response remain high. Additionally, there is a high density of CnFLs presented on the host cell to detect symbionts in the environment. Bolded arrows and text indicate stronger induction/repression of a pathway or response. Dashed arrows indicate transient (e.g. stress) or partial (e.g. inflammatory response) pathway induction/ repression, whereas solid arrows indicate continuous induction/repression.

Increasing the number of host–algal pairings in cnidarian studies

Certain cnidarian model systems have been developed to facilitate the study of the biochemical processes that modulate associations and dissociations of cnidarian hosts with microbial partners. The sea anemone *Aiptasia* has provided a platform for many advances in our molecular understanding of these symbioses because it participates in a facultative symbiosis with certain strains of Symbiodiniaceae, is simple to maintain in laboratories of all scales, can produce larvae naïve to symbionts under inducible conditions, and is easily biochemically manipulatable (Grawunder et al. 2015; Weis et al. 2008).

While the work conducted with *Aiptasia* has significantly shaped our understanding of host–Symbiodiniaceae pairings, it has predominantly been conducted on a single, or very few, *Aiptasia* strains. CC7, the most commonly used *Aiptasia* strain, is male and may have different energetic priorities than female conspecifics. Additionally, significant genetic variation in many traits and different host–symbiont interactions can drive responses to the environment. Therefore, we encourage researchers to expand their work to include additional strains at various developmental stages, to build out strain libraries by pulling genotypes from diverse regions that experience unique environmental conditions, and to make these anemones broadly




Fig. 2 Experimental approaches for future studies on cnidarian holobiont interactions. Focus on increasing host genetic diversity, different monoculture and/or known mixed Symbiodiniaceae strains, and microbial communities of varying complexity. We expect that an individual may exhibit multiple phenotypes (depicted by color of host, represented as a coral and anemone, or culture flask; grey flasks represent failed cultures) depending on associated partners that may result in differing levels of resilience or resistance to stress. Crossing these different pairings under both ambient and stressful conditions may elucidate roles of partners in holobiont responses that help predict holobiont health. The purple arrow depicts future directions once baseline interactions are known across different partner pairings. The coral and anemone shapes were created by G. Puntin.

available to the community (Fig. 2). Many other models for symbiosis are emerging including *Cassiopea xamachana*, which represents a promising model system for some symbiosis questions and demonstrates certain benefits over *Aiptasia*, including the ability to induce settlement and metamorphosis, which has failed in *Aiptasia* (reviewed in Ohdera et al. 2018). The necessary expansion of calcifying coral models will be explored in the subsection *Further develop calcifying coral models*. We recognize that this work involves challenges related to feasibility; however, experiments could be conducted sequentially and collaboration between laboratories is encouraged. We also stress careful documentation of genetic background in husbandry and experiments, and support open-source protocol sharing of animal rearing, spawning, and experimentation, as has been implemented by the *Aiptasia* Symbiosis Resource group (<https://aiptasia-resource.org/>).

This same expansion of genetic backgrounds needs to also be applied to Symbiodiniaceae strains (Fig. 2). Given the diversity of Symbiodiniaceae within and between genera (LaJeunesse et al. 2018; Beltrán et al. 2021), and that each strain can have a unique interaction

with a host genotype (Parkinson and Baums 2014), expansion of strains is critical. Further, it remains important to assess how crossing different host and algal symbiont genetic backgrounds may alter holobiont physiology under both ambient and stressful conditions. Lastly, we recommend consistent and regular genomic screenings of all symbiont libraries and stringent oversight of symbiont culturing to avoid contamination.

Further develop calcifying coral models

While the use of *Aiptasia* (and other emerging models) provides valuable insights into the molecular underpinnings of symbiosis, it remains a solitary polyp that does not calcify like reef-building corals. Given this limitation, other models are emerging to study coral symbiosis, including the facultatively symbiotic species *Astrangia poculata*, which is found along the western Atlantic coastline of the USA (Neff 2020). The use of *A. poculata* may offer a more realistic model for understanding the role of each symbiotic partner (Holcomb et al. 2012; Burmester et al. 2018) given the importance of symbiosis in coral skeletal growth rate (Chalker

and Taylor 1975). *Astrangia poculata* has provided valuable insight into algal symbiosis regulation (Dimond and Carrington 2008), the impact of algal symbionts on microbial community composition (Sharp et al. 2017), how symbiosis may mediate stress in calcifying corals (Burmeister et al. 2017; Wuitchik et al. 2021), and differences in skeletal structure based on symbiotic state (Dellaert et al. 2022). This ability to modulate symbiont state is valuable; however, to date, no studies have reported successfully inoculating *A. poculata* with different (i.e., heterologous) symbiont strains. While other studies have reported acquisition of novel algal symbionts in adult corals (Puntin et al. 2022; Scharfenstein et al. 2022), inoculation of *A. poculata* with heterologous algal strains should remain a priority.

Several other reef building species are emerging as potential models, including *Oculina arbuscula* (Rivera and Davies 2021) and *Galaxea fascicularis* (Puntin et al. 2022), which can also have their symbiotic states experimentally manipulated. In fact, *G. fascicularis* has been successfully bleached and then reinoculated with presumed heterologous strains of Symbiodiniaceae (Puntin et al. 2022), providing a promising system to cross different host genotypes with algal strains as proposed above (Fig. 2). This ability to control and modify symbiosis in combination with previous mechanistic (e.g., ion transport for calcification (Al-Horani et al. 2003, 2005)) and microbiome (Tang et al. 2020; Wepfer et al. 2020) research make it an ideal candidate to explore partner interactions.

Lastly, a wealth of resources have been created for the model tropical reef-building coral *A. millepora*, which was the first to have its transcriptome sequenced (Meyer et al. 2009) and a linkage map developed (Wang et al. 2009). It also has one of the most robust genome assemblies to date (Fuller et al. 2020) and has been used in a plethora of spawning studies. It remains one of the more tractable model systems for genome editing (see *Leveraging novel technologies subsection*; (Cleves et al. 2018)), and robust methods for spawning (both *in situ* and *ex situ*), settlement, and algal symbiosis establishment exist (Craggs et al. 2017, 2020; Pollock et al. 2017).

Conduct research with and without microbiome members

To study specific interactions between the host, algal symbiont, and microbial partners, we suggest employing the use of axenic (i.e., germ-free) and gnotobiotic (i.e., known microbial state) cnidarian host and symbiont models (Fig. 2). The use of axenic and gnotobiotic models is common practice in many systems (especially mammalian models) to assess the functional role that specific host–microbe interactions have on phenotype

(Basic and Bleich 2019) and presents a promising avenue to study partner interactions in symbiotic cnidarians. While gnotobiotic cnidarian models have yet to be successfully produced (Hartman et al. 2022), several strains of axenic Symbiodiniaceae have been developed and have been successfully infected into *Aiptasia* models (Xiang et al. 2013; Matthews et al. 2020). However, additional strains from diverse algal genera would be beneficial along with their associations with specific microbial communities. A protocol for the generation of microbe depleted *Aiptasia* has already been developed (Costa et al. 2021), showcasing progress towards the generation of completely axenic and eventually gnotobiotic *Aiptasia* strains.

While generating axenic and gnotobiotic corals may be in the future, researchers can currently inoculate corals with known beneficial microorganisms for corals (BMC) and evaluate phenotypic outcomes to better understand the microbiome's role in holobiont health (Peixoto et al. 2017; Santoro et al. 2021; Zhang et al. 2021; Li et al. 2022). Inoculating hosts with different microbial communities is widely used across disciplines (Ma et al. 2011; Hansen et al. 2012) to assess a variety of phenotypic and immune responses, and these techniques represent a promising frontier for corals as well. Studying host responses to targeted microorganisms or microbial communities will uncover the role of these associations and how they might be linked with holobiont stress, vulnerability, resistance, and resilience.

Developing high-throughput screening approaches

Development of additional technologies to rapidly assess phenotypic traits is needed and remains a major bottleneck to the types of experiments required to understand complex interactions between multiple symbiotic partners. Recent high-throughput approaches for assessing thermal tolerance at the whole coral level (e.g., coral bleaching automated stress systems (CBASS; (Voolstra et al. 2020)) and single cell levels (Behrendt et al. 2020) have incorporated short-term thermal challenges followed by stress characterization through measurement of a couple physiological variables such as maximum PSII photochemical efficiency (F_v/F_m) and cell density. However, these approaches remain focused on the algal symbiont and development of additional traits is needed. Further developing trait databases, such as the coral trait database (Madin et al. 2016) are urgently needed. While this database hosts data for 56 traits—some of which overlap with the algal symbiont (i.e., gross photosynthesis, mitotic index, symbiont density, algae hosted, algal transmission mode, and

chlorophyll A)—this resource could be complemented by the development of algal and microbiome versions.

Gene expression patterns of each member of the holobiont can also be considered a trait and represents a high-throughput approach; however, linking gene expression to coral phenotypes remains difficult. In the plant community, researchers have better linked gene expression patterns to their trait database (Kattge et al. 2011) by creating a plant gene annotation platform that provides common standards for semantic integration in the form of ontologies (Planteome: Cooper et al. 2018). The development of such a resource for cnidarians would transcend our understanding of gene function. In lieu of developing a novel database, which represents a mammoth task that requires continuous funding to maintain, one approach to link gene expression with cnidarian responses is through meta-analysis. For example, Dixon et al. (2020) re-analyzed 600 coral gene expression profiles (genus *Acropora*) and established two general coral environmental stress responses with contextual annotations for unannotated genes based on their consistent response to stress across independent projects. These sorts of approaches can better link research globally and will help build our understanding of coral diversity in order to better predict responses.

Leveraging novel technologies

Advances in biomedicine have been facilitated by the development of immortal cell lines for molecular manipulation and microscopy analysis. The generation of cnidarian cell lines has been met with limited success (see Rosental et al. 2017; Fricano et al. 2020; Kawamura, Nishitsuji, et al. 2021; Nowotny et al. 2021), although achievements have been made with respect to *in vivo* symbiosis (Kawamura, Sekida, et al. 2021). Functional cell lines of cnidarian models will allow researchers to apply exogenous biochemical stimuli, such as TGF β and NO, which would expand our knowledge of gene function. Current studies using mammalian cell lines expressing cnidarian proteins (i.e., Williams et al. 2018) may be missing key post-translational modifications and/or protein–protein interactions that only occur in native host cells, highlighting the need for further technology development in this area.

ScRNAseq is a developing technology that has been successful in several Cnidaria taxa (Seb  -Pedr  s et al. 2018; Hu et al. 2020; Chari et al. 2021; Levy et al. 2021); however, most studies to date have been conducted under baseline conditions to establish cell-type signatures of gene expression (i.e., cell atlases) through developmental stages. The use of scRNAseq to characterize cell-type gene expression patterns under different stimuli will further unveil the cell types responsible for

organismal responses to various stressors. Additionally, scRNAseq manipulation experiments can help determine the benefit-to-burden balance that hosts must weigh when in symbiosis with homologous vs. heterologous symbiont strains. The development of spatial scRNAseq (reviewed in Longo et al. 2021) or multiplexed error-robust fluorescence *in situ* hybridization (Chen et al. 2015) technologies for other systems can be leveraged in cnidarian research to deepen our understanding of the spatial organization of cell types relative to the organismal body plan. Additionally, proteomic (Oakley et al. 2016; Jaimes-Becerra et al. 2019; Sproles et al. 2019; Tortorelli, Oakley, et al. 2022) and metabolomic studies (Hillyer et al. 2017; Matthews et al. 2017, 2018; Williams et al. 2020) are essential for determining functional patterns of intra- and inter-cellular regulation under different conditions and holobiont partnerships.

Lastly, the development and implementation of gene knockdown (via RNA-interference (Dunn et al. 2007); reviewed in Rentzsch et al. 2019; Quiroga-Artigas et al. 2020) and genome editing (via the CRISPR-Cas9 system; (Cleves et al. 2018)) in Cnidaria has allowed researchers to resolve the functional role of specific genes involved in immunity, development, and more. Indeed, NF-  B has been previously knocked down in the non-symbiotic sea anemone *Nematostella vectensis* via morpholinos, clarifying the protein's role in cnidocyte cell development (Wolenski et al. 2013). We propose that NF-  B, and cnidarian-specific PRRs be knocked down via RNA-interference or knocked out via CRISPR-Cas9 in a symbiotic cnidarian model that can be reared from larvae to adulthood, such as *Cassiopea xamachana*, with the acknowledgement that mosaic phenotypes and inviable phenotypes may be likely outcomes. Additional advances in tractable mutagenesis models in mammal systems have allowed researchers to study the direct link between microbial member networks to host phenotypes (Goodman et al. 2009); we suggest the exploration of implementing similar models in cnidarian systems. Finally, we recommend that transgenic lines be infected with different strains of symbiont and/or microbiome combinations and placed under various challenges to clarify the role of these mechanistic pathways in symbiosis, dysbiosis, and environmental stressors. Combining these data-rich methods in a multi omic context will improve informed diagnostic approaches to coral work *in situ*.

Conclusions

Coral bleaching is perhaps one of the most intriguing processes in nature because it represents the outcome of cellular dysfunction between symbiotic partners that is easily observed by eye, but its influence

spans ecosystem scales. A major goal of coral reef science is to understand and predict these symbiosis outcomes under increasingly changing environments, but this work remains a serious challenge due to the complex interactions between holobiont members, each of which possess immense genetic diversity. While an impressive amount of work has uncovered basic mechanisms involved in cnidarian–algal symbiosis and descriptive studies of microbiome associations are hypothesis generating tools, we propose that mechanistic research would benefit from expanding to use a broader repertoire of genetic diversity and leveraging emerging technologies. To do this, we urge that future work be carried out in collaboration with Indigenous groups throughout the entire research process to center Traditional Ecological Knowledge, without which we cannot properly protect these ecosystems. We are optimistic that the many ways in which these partners interact will offer hope to the coral reef crisis as we are just scratching the surface of understanding how these partners can mix and match, which may offer fuel to the coral holobiont to adapt and acclimate to their rapidly changing environments.

Acknowledgments

We acknowledge that the research proposed in this perspective will be performed on the stolen ancestral land of many Indigenous peoples who historically and currently steward coral reef ecosystems. We also thank Dr Hannah Reich, Dr Daniel Segré, and two anonymous reviewers for providing valuable feedback that improved this perspective. Finally, we thank the previous and ongoing efforts of the many researchers whose work informed the ideas presented in this perspective. The coral and anemone shapes were created by Giulia Puntin and modified for use (https://github.com/sPuntinG/Coral_stuff).

Funding

This work was supported by a Boston University Microbiome Initiative Accelerator grant to CBB. MVI was supported by an National Science Foundation Graduate Research Fellowships Program and National Science Foundation Research Traineeship Division of Graduate Education [grant number 1735087]. Funding for this research came from the National Science Foundation [grant number IOS-SDS1937650 to SWD] and startup funds from Boston University to SWD.

Conflict of interest

The authors declare no conflict of interest.

Data availability

No new data were generated or analyzed in support of this research.

References

Ainsworth TD, Thurber RV, Gates RD. 2010. The future of coral reefs: a microbial perspective. *Trends Ecol Evol* 25:233–40.

Akira S, Uematsu S, Takeuchi O. 2006. Pathogen recognition and innate immunity. *Cell* 124:783–801.

Al-Horani FA, Ferdelman T, Al-Moghrabi SM, de Beer D. 2005. Spatial distribution of calcification and photosynthesis in the scleractinian coral *Galaxea fascicularis*. *Coral Reefs* 24:173–80.

Al-Horani FA, SM Al-Moghrabi, de Beer D. 2003. The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*. *Mar Biol* 142:419–26.

Ali A, Kriefall NG, Emery LE, Kenkel CD, Matz MV, Davies SW. 2019. Recruit symbiosis establishment and Symbiodinaceae composition influenced by adult corals and reef sediment. *Coral Reefs* 38:405–15.

Armitage DW, Jones SE. 2019. How sample heterogeneity can obscure the signal of microbial interactions. *ISME J* 13:2639–46.

Baird AH, Guest JR, Willis BL. 2009. Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. *Annu Rev Ecol Evol Syst* 40:551–71.

Bairois-Novak KR, Hoogenboom MO, van Oppen MJH, Connolly SR. 2021. Coral adaptation to climate change: meta-analysis reveals high heritability across multiple traits. *Global Change Biol* 27:5694–5710.

Baker AC, Romanski AM. 2007. Multiple symbiotic partnerships are common in scleractinian corals, but not in octocorals: Comment on Goulet (2006). *Mar Ecol Prog Ser* 335: 237–42.

Baker AC. 2003. Flexibility and specificity in coral–algal symbiosis: diversity, ecology, and biogeography of *Symbiodinium*. *Annu Rev Ecol Evol Syst* 34:661–89.

Basic M, Bleich A. 2019. Gnotobiotics: past, present and future. *Lab Anim* 53:232–43.

Baumgarten S, Simakov O, Esherick LY, Liew YJ, Lehnert EM, Michell CT, Li Y, Hambleton EA, Guse A, Oates ME, et al. 2015. The genome of *Aiptasia*, a sea anemone model for coral symbiosis. *Proc Natl Acad Sci* 112:11893–8.

Behrendt L, Salek MM, Trampe EL, Fernandez VI, Lee KS, Kühl M, Stocker R. 2020. PhenoChip: a single-cell phenomic platform for high-throughput photophysiological analyses of microalgae. *Sci Adv* 6:1–13.

Beltrán VH, Puill-Stephan E, Howells E, Flores-Moya A, Doblin M, Núñez-Lara E, Escamilla V, López T, van Oppen MJH. 2021. Physiological diversity among sympatric, conspecific endosymbionts of coral (*Cladocopium C1^{acro}*) from the Great Barrier Reef. *Coral Reef* 40: 985–97.

Berkelmans R, Van Oppen MJH. 2006. The role of zooxanthellae in the thermal tolerance of corals: a “nugget of hope” for coral reefs in an era of climate change. *Proc R Soc Lond B Biol Sci* 273:2305–12.

Biagi E, Caroselli E, Barone M, Pezzimenti M, Teixido N, Soverini M, Rampelli S, Turroni S, Gambi MC, Brigidi P, et al. 2020. Patterns in microbiome composition differ with ocean acidification in anatomic compartments of the Mediterranean coral

Astroides calyculus living at CO₂ vents. *Sci Total Environ* 724:138048.

Bickford D, Lohman DJ, Sodhi NS, Ng PKL, Meier R, Winker K, Ingram KK, Das I. 2007. Cryptic species as a window on diversity and conservation. *Trends Ecol Evol* 22: 148–55.

Bourne DG, Morrow KM, Webster NS. 2016. Insights into the coral microbiome: underpinning the health and resilience of reef ecosystems. *Annu Rev Microbiol* 70:317–40.

Bove CB, Whitehead RF, Szmant AM. 2020. Responses of coral gastrovascular cavity pH during light and dark incubations to reduced seawater pH suggest species-specific responses to the effects of ocean acidification on calcification. *Coral Reef* 39:1675–91.

Burmester EM, Breef-Pilz A, Lawrence NF, Kaufman L, Finnerty JR, Rotjan RD. 2018. The impact of autotrophic versus heterotrophic nutritional pathways on colony health and wound recovery in corals. *Ecol and Evol* 8:10805–16.

Burmester EM, Finnerty JR, Kaufman L, Rotjan RD. 2017. Temperature and symbiosis affect lesion recovery in experimentally wounded, facultatively symbiotic temperate corals. *Mar Ecol Prog Ser* 570:87–99.

Camp EF, Kahlke T, Nitschke MR, Varkey D, Fisher NL, Fujise L, Goyen S, Hughes DJ, Lawson CA, Ros M, et al. 2020. Revealing changes in the microbiome of Symbiodiniaceae under thermal stress. *Environ Microbiol* 22:1294–1309.

Chalker BE, Taylor DL. 1975. Light-enhanced calcification, and the role of oxidative phosphorylation in calcification of the coral *Acropora cervicornis*. *Proc R Soc Lond B Biol Sci* 190:323–31.

Chari T, Weissbourd B, Gehring J, Ferraioli A, Leclère L, Herl M, Gao F, Chevalier S, Copley RR, Houlston E, et al. 2021. Whole-animal multiplexed single-cell RNA-seq reveals transcriptional shifts across *Clytia* medusa cell types. *Sci Adv* 7:eaab1683.

Chen KH, Boettiger AN, Moffitt JR, Wang S, Zhuang X. 2015. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* 348:aaa6090.

Cleves PA, Strader ME, Bay LK, Pringle JR, Matz MV. 2018. CRISPR/Cas9-mediated genome editing in a reef-building coral. *Proc Natl Acad Sci* 115:5235–40.

Cooper L, Meier A, Laporte M-A, Elser JL, Mungall C, Sinn BT, Cavaliere D, Carbon S, Dunn NA, Smith B, et al. 2018. The Planteome database: an integrated resource for reference ontologies, plant genomics and phenomics. *Nucleic Acids Res* 46:D1168–80.

Costa RM, Cárdenas A, Loussert-Fonta C, Toullec G, Meibom A, Voolstra CR. 2021. Surface topography, bacterial carrying capacity, and the prospect of microbiome manipulation in the sea anemone coral model *Aiptasia*. *Front Microbiol* 12:637834.

Costanza R, de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I, Farber S, Turner RK. 2014. Changes in the global value of ecosystem services. *Glob Environ Chang* 26:152–8.

Craggs J, Guest J, Davis M, Sweet M. 2020. Completing the life cycle of a broadcast spawning coral in a closed mesocosm. *Invertebr Reprod Dev* 64:244–7.

Craggs J, Guest JR, Davis M, Simmons J, Dashti E, Sweet M. 2017. Inducing broadcast coral spawning ex situ: closed system mesocosm design and husbandry protocol. *Ecol and Evol* 7:11066–78.

Damjanovic K, Menéndez P, Blackall LL, van Oppen MJH. 2020. Early life stages of a common broadcast spawning coral associate with specific bacterial communities despite lack of internalized bacteria. *Microb Ecol* 79:706–19.

Davies S, Gamache MH, Howe-Kerr LI, Kriefall NG, Baker AC, Banaszak AT, Bay LK, Bellantuono AJ, Bhattacharya D, Chan CX, et al. 2022. Building consensus around the assessment and interpretation of Symbiodiniaceae diversity. DOI: 10.20944/preprints202206.0284.v.

Davy SK, Lucas IAN, Turner JR. 1997. Uptake and Persistence of Homologous and Heterologous Zooxanthellae in the Temperate Sea Anemone *Cereus pedunculatus*. *Biol Bull* 192:208–16.

De'ath G, Fabricius KE, Sweatman H, Puotinen M. 2012. The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proc Natl Acad Sci* 109:17995–9.

Dellaert Z, Vargas PA, La Riviere PJ, Roberson LM. 2022. Uncovering the effects of symbiosis and temperature on coral calcification. *Biol Bull* 242:62–73.

DeSalvo MK, Sunagawa S, Voolstra CR, Medina M. 2010. Transcriptomic responses to heat stress and bleaching in the elkhorn coral *Acropora palmata*. *Mar Ecol Prog Ser* 402:97–113.

Detournay O, Schnitzler CE, Poole A, Weis VM. 2012. Regulation of cnidarian-dinoflagellate mutualisms: Evidence that activation of a host TGF β innate immune pathway promotes tolerance of the symbiont. *Dev Comp Immunol* 38:525–37.

Dimond J, Carrington E. 2008. Symbiosis regulation in a facultatively symbiotic temperate coral: zooxanthellae division and expulsion. *Coral Reef* 27:601–4.

Dixon G, Abbott E, Matz M. 2020. Meta-analysis of the coral environmental stress response: *Acropora* corals show opposing responses depending on stress intensity. *Mol Ecol* 29:2855–70.

Dixon GB, Davies SW, Aglyamova GA, Meyer E, Bay LK, Matz MV. 2015. CORAL REEFS. Genomic determinants of coral heat tolerance across latitudes. *Science* 348:1460–62.

Dougan KE, González-Pech RA, Stephens TG, Shah S, Chen Y, Ragan MA, Bhattacharya D, Chan CX. 2022. Genome-powered classification of microbial eukaryotes: focus on coral algal symbionts. *Trends Microbiol* 30:831–40.

Dunn SR, Phillips WS, Green DR, Weis VM. 2007. Knockdown of actin and caspase gene expression by RNA interference in the symbiotic anemone *Aiptasia pallida*. *Biol Bull* 212: 250–58.

Eddy TD, Lam VWY, Reygondeau G, Cisneros-Montemayor AM, Greer K, Palomares MLD, Bruno JF, Ota Y, Cheung WWL. 2021. Global decline in capacity of coral reefs to provide ecosystem services. *One Earth* 4:1278–85.

Emery MA, Dimos BA, Mydlarz LD. 2021. Cnidarian pattern recognition receptor repertoires reflect both phylogeny and life history traits. *Front Immunol* 12:689463.

Epstein HE, Torda G, Munday PL, van Oppen MJH. 2019. Parental and early life stage environments drive establishment of bacterial and dinoflagellate communities in a common coral. *ISME J* 13:1635–8.

Fifer JE, Yasuda N, Yamakita T, Bove CB, Davies SW. 2022. Genetic divergence and range expansion in a western North Pacific coral. *Sci Total Environ* 813:152423.

Fitt WK, Chang SS, Trench RK. 1981. Motility patterns of different strains of the symbiotic dinoflagellate *Symbiodinium* (= *Gymnodinium*) *microadriaticum* (Freudenthal) in culture. *Bull Mar Sci* 31:436–43.

Forsman ZH, Ritson-Williams R, Tisthammer KH, Knapp ISS, Toonen RJ. 2020. Host-symbiont coevolution, cryptic structure, and bleaching susceptibility, in a coral species complex (Scleractinia; Poritidae). *Sci Rep* 10:16995.

Fricano C, Röttinger E, Furla P, Barnay-Verdier S. 2020. Cnidarian cell cryopreservation: a powerful tool for cultivation and functional assays. *Cells* 9:2541.

Fuess LE, Butler CC, Brandt ME, Mydlarz LD. 2020. Investigating the roles of transforming growth factor-beta in immune response of *Orbicella faveolata*, a scleractinian coral. *Dev Comp Immunol* 107:103639.

Fuller ZL, Mocellin VJL, Morris LA, Cantin N, Shepherd J, Sarre L, Peng J, Liao Y, Pickrell J, Andolfatto P, et al. 2020. Population genetics of the coral *Acropora millepora*: toward genomic prediction of bleaching. *Science* 369.6501 1–9.

Gilmore TD. 2006. Introduction to NF-κappaB: players, pathways, perspectives. *Oncogene* 25:6680–84.

Glasl B, Herndl GJ, Frade PR. 2016. The microbiome of coral surface mucus has a key role in mediating holobiont health and survival upon disturbance. *ISME J* 10:2280–92.

Glynn PW. 1991. Coral reef bleaching in the 1980s and possible connections with global warming. *Trends Ecol Evol* 6:175–9.

Gómez-Corrales M, Prada C. 2020. Cryptic lineages respond differently to coral bleaching. *Mol Ecol* 29:4265–73.

Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, Knight R, Gordon JI. 2009. Identifying genetic determinants needed to establish a human gut symbiont in its habitat. *Cell Host Microbe* 6:279–89.

Gorman LM, Konciute MK, Cui G, Oakley CA, Grossman AR, Weis VM, Aranda M, Davy SK. 2022. Symbiosis with dinoflagellates alters cnidarian cell-cycle gene expression. *Cell Microbiol* 2022:1–20.

Grawunder D, Hambleton EA, Bucher M, Wolfowicz I, Bechtoldt N, Guse A. 2015. Induction of gametogenesis in the cnidarian endosymbiosis model *Aiptasia* sp. *Sci Rep* 5:15677.

Hansen CHF, Nielsen DS, Kverka M, Zakostelska Z, Klimesova K, Hudcovic T, Tlaskalova-Hogenova H, Hansen AK. 2012. Patterns of early gut colonization shape future immune responses of the host. *PLoS One* 7:e34043.

Hartman LM, Blackall LL, van Oppen MJH. 2022. Antibiotics reduce bacterial load in, but biofilms hinder its development as a gnotobiotic coral model. *Access Microbiol* 4:000314.

Hartmann AC, Baird AH, Knowlton N, Huang D. 2017. The paradox of environmental symbiont acquisition in obligate mutualisms. *Curr Biol* 27:3711–6.

Hernandez-Agreda A, Gates RD, Ainsworth TD. 2017. Defining the core microbiome in corals' microbial soup. *Trends Microbiol* 25:125–40.

Hillyer KE, Dias DA, Lutz A, Roessner U, Davy SK. 2017. Mapping carbon fate during bleaching in a model cnidarian symbiosis: the application of ¹³C metabolomics. *New Phytol* 214:1551–62.

Hoadley KD, Lewis AM, Wham DC, Pettay DT, Grasso C, Smith R, Kemp DW, LaJeunesse TC, Warner ME. 2019. Host-symbiont combinations dictate the photo-physiological response of reef-building corals to thermal stress. *Sci Rep* 9:9985.

Holcomb M, Cohen AL, McCorkle DC. 2012. An investigation of the calcification response of the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the effects of nutrients, zooxanthellae and gender. *Biogeosciences* 9:29–39.

Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH. 2012. Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat Clim Change* 2:116–20.

Hu M, Zheng X, Fan C-M, Zheng Y. 2020. Lineage dynamics of the endosymbiotic cell type in the soft coral *Xenia*. *Nature* 582:534–8.

Hughes DJ, Raina J-B, Nielsen DA, Suggett DJ, Kühl M. 2022. Disentangling compartment functions in sessile marine invertebrates. *Trends Ecol Evol* 37, 740–48.

Hume BCC, Mejia-Restrepo A, Voolstra CR, Berumen ML. 2020. Fine-scale delineation of Symbiodiniaceae genotypes on a previously bleached central Red Sea reef system demonstrates a prevalence of coral host-specific associations. *Coral Reef* 39:583–601.

Jaimes-Becerra A, Gacesa R, Doonan LB, Hartigan A, Marques AC, Okamura B, Long PF. 2019. "Beyond Primary Sequence"-proteomic data reveal complex toxins in cnidarian venoms. *Integr Comp Biol* 59:777–85.

Jiang J, Wang A, Deng X, Zhou W, Gan Q, Lu Y. 2021. How Symbiodiniaceae meets the challenges of life during coral bleaching. *Coral Reef* 40:1339–53.

Jones GB, King S. 2015. Dimethylsulphoniopropionate (DMSP) as an indicator of bleaching tolerance in scleractinian corals. *J Mar Sci Eng* 3:444–65.

Jury CP, Delano MN, Toonen RJ. 2019. High heritability of coral calcification rates and evolutionary potential under ocean acidification. *Sci Rep* 9:20419.

Kattge J, Díaz S, Lavorel S, Prentice IC, Leadley P, Bönnisch G, Garnier E, Westoby M, Reich PB, Wright IJ, et al. 2011. Establishing sustainable cell lines of a coral, *Acropora tenuis*. *Mar Biotechnol* 23:373–88.

Kattge J, Díaz S, Lavorel S, Prentice IC, Leadley P, Bönnisch G, Garnier E, Westoby M, Reich PB, Wright IJ, et al. 2011. TRY - a global database of plant traits. *Glob Chang Biol* 17:2905–35.

Kawamura K, Sekida S, Nishitsuji K, Shoguchi E, Hisata K, Fujiwara S, Satoh N. 2021. In vitro symbiosis of reef-building coral cells with photosynthetic dinoflagellates. *Front Mar Sci* 8: 900.

Kenkel CD, Matz MV. 2017. Gene expression plasticity as a mechanism of coral adaptation to a variable environment. *Nat Ecol Evol* 1:14.

Kenkel CD, Setta SP, Matz MV. 2015. Heritable differences in fitness-related traits among populations of the mustard hill coral, *Porites astreoides*. *Heredity* 115:509–16.

Kimes NE, Van Nostrand JD, Weil E, Zhou J, Morris PJ. 2010. Microbial functional structure of *Montastraea faveolata*, an important Caribbean reef-building coral, differs between healthy and yellow-band diseased colonies. *Environ Microbiol* 12:541–56.

Knowlton N. 1993. Sibling species in the sea. *Annu Rev Ecol Evol Syst* 24:189–216.

LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018. Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr Biol* 28:2570–80.

LaJeunesse TC, Wiedenmann J, Casado-Amezúa P, D'Ambra I, Turnham KE, Nitschke MR, Oakley CA, Goffredo S, Spano CA, Cubillos VM, et al. 2021. Revival of *Philozoon Geddes* for host-specialized dinoflagellates, "zooxanthellae", in animals from coastal temperate zones of northern and southern hemispheres. *Eur J Phycol* 57:166–180.

Lawson CA, Raina J-B, Kahlke T, Seymour JR, Suggett DJ. 2018. Defining the core microbiome of the symbiotic dinoflagellate. *Environ Microbiol Rep* 10:7–11.

Lehnert EM, Mouchka ME, Burriesci MS, Gallo ND, Schwarz JA, Pringle JR. 2014. Extensive differences in gene expression between symbiotic and aposymbiotic cnidarians. *G3 Genes|Genomes|Genetics* 4:277–95.

Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, Mass T, Sebé-Pedrós A. 2021. A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. *Cell* 184:2973–87.

Li J, Yang Q, Dong J, Sweet M, Zhang Y, Liu C, Zhang Y, Tang X, Zhang W, Zhang S. 2022. Microbiome engineering: a promising approach to improve coral health. *Proc Est Acad Sci Eng*. 1–14. DOI: 10.1016/j.eng.2022.07.010.

Logan DDK, LaFlamme AC, Weis VM, Davy SK. 2010. Flow-cytometric characterization of the cell-surface glycans of symbiotic dinoflagellates (*Symbiodinium* spp.). *J Phycol* 46:525–33.

Longo SK, Guo MG, Ji AL, Khavari PA. 2021. Integrating single-cell and spatial transcriptomics to elucidate intercellular tissue dynamics. *Nat Rev Genet* 22:627–44.

Ma Y, Rajkumar M, Luo Y, Freitas H. 2011. Inoculation of endophytic bacteria on host and non-host plants—effects on plant growth and Ni uptake. *J Hazard Mater* 195:230–37.

Madin JS, Anderson KD, Andreasen MH, Bridge TCL, Cairns SD, Connolly SR, Darling ES, Diaz M, Falster DS, Franklin EC, et al. 2016. The coral trait database, a curated database of trait information for coral species from the global oceans. *Sci Data* 3:160017.

Maire J, Buerger P, Chan WY, Deore P, Dungan AM, Nitschke MR, van Oppen MJH. 2022. Effects of ocean warming on the underexplored members of the coral microbiome. *Integr Comp Biol*. 0 1–10

Mansfield KM, Carter NM, Nguyen L, Cleves PA, Alshanbayeva A, Williams LM, Crowder C, Penrose AR, Finnerty JR, Weis VM, et al. 2017. Transcription factor NF- κ B is modulated by symbiotic status in a sea anemone model of cnidarian bleaching. *Sci Rep* 7:16025.

Mansfield KM, Cleves PA, Van Vlack E, Kriefall NG, Benson BE, Camacho DJ, Hemond O, Pedroza M, Siggers T, Pringle JR, et al. 2019. Varied effects of algal symbionts on transcription factor NF- κ B in a sea anemone and a coral: possible roles in symbiosis and thermotolerance. *bioRxiv*. 1–39

Mansfield KM, Gilmore TD. 2019. Innate immunity and cnidarian–Symbiodiniaceae mutualism. *Dev Comp Immunol* 90:199–209.

Matthews JL, Crowder CM, Oakley CA, Lutz A, Roessner U, Meyer E, Grossman AR, Weis VM, Davy SK. 2017. Optimal nutrient exchange and immune responses operate in partner specificity in the cnidarian–dinoflagellate symbiosis. *Proc Natl Acad Sci* 114:13194–9.

Matthews JL, Oakley CA, Lutz A, Hillyer KE, Roessner U, Grossman AR, Weis VM, Davy SK. 2018. Partner switching and metabolic flux in a model cnidarian–dinoflagellate symbiosis. *Proc Biol Sci* 285.1–10

Matthews JL, Raina J-B, Kahlke T, Seymour JR, van Oppen MJH, Suggett DJ. 2020. Symbiodiniaceae–bacteria interactions: rethinking metabolite exchange in reef-building corals as multi-partner metabolic networks. *Environ Microbiol* 22: 1675–87.

Meron D, Rodolfo-Metalpa R, Cunning R, Baker AC, Fine M, Banin E. 2012. Changes in coral microbial communities in response to a natural pH gradient. *ISME J* 6:1775–85.

Meyer E, Aglyamova GV, Matz MV. 2011. Profiling gene expression responses of coral larvae (*Acropora millepora*) to elevated temperature and settlement inducers using a novel RNA-Seq procedure. *Mol Ecol* 20:3599–3616.

Meyer E, Aglyamova GV, Wang S, Buchanan-Carter J, Abrego D, Colbourne JK, Willis BL, Matz MV. 2009. Sequencing and de novo analysis of a coral larval transcriptome using 454 GSFLx. *BMC Genom* 10:219.

Mieog JC, Olsen JL, Berkelmans R, Bleuler-Martinez SA, Willis BL, van Oppen MJH. 2009. The roles and interactions of symbiont, host and environment in defining coral fitness. *PLoS One* 4:e6364.

Moberg F, Folke C. 1999. Ecological goods and services of coral reef ecosystems. *Ecol Econ* 29:215–33.

Morris LA, Voolstra CR, Quigley KM, Bourne DG. 2019. Nutrient availability and metabolism affect the stability of coral–Symbiodiniaceae symbioses. *Trends Microbiol* 27:678–89.

Muscatine L, Cernichiari E. 1969. Assimilation of photosynthetic products of zooxanthellae by a reef coral. *Biol Bull* 137:506–523.

Neff EP. 2020. The quest for an animal model of coral health and disease. *Lab Anim (NY)* 49:37–41.

Nitschke MR, Craveiro SC, Brandão C, Fidalgo C, Serôdio J, Calado AJ, Frommlet JC. 2020. Description of *Freudenthalidium* gen. nov. and *Halluxium* gen. nov. to formally recognize Clades Fr3 and H as genera in the family Symbiodiniaceae (Dinophyceae). *J Phycol* 56:923–40.

Nowotny JD, Connelly MT, Traylor-Knowles N. 2021. Novel methods to establish whole-body primary cell cultures for the cnidarians *Nematostella vectensis* and *Pocillopora damicornis*. *Sci Rep* 11:4086.

Nyholm SV, Graf J. 2012. Knowing your friends: invertebrate innate immunity fosters beneficial bacterial symbioses. *Nat Rev Microbiol* 10:815–27.

Oakley CA, Ameisemeier MF, Peng L, Weis VM, Grossman AR, Davy SK. 2016. Symbiosis induces widespread changes in the proteome of the model cnidarian *Aiptasia*. *Cell Microbiol* 18:1009–23.

Ohdera AH, Abrams MJ, Ames CL, Baker DM, Suescún-Bolívar LP, Collins AG, Freeman CJ, Gamero-Mora E, Goulet TL, Hofmann DK, et al. 2018. Upside-down but headed in the right direction: review of the highly versatile *Cassiopea xamachana* system. *Front Ecol Evol* 6:1–15

Parkinson JE, Baums IB. 2014. The extended phenotypes of marine symbioses: ecological and evolutionary consequences of intraspecific genetic diversity in coral–algal associations. *Front Microbiol* 5:445.

Peixoto RS, Rosado PM, Leite DC de A, Rosado AS, Bourne DG. 2017. Beneficial microorganisms for corals (BMC): proposed mechanisms for coral health and resilience. *Front Microbiol* 8:341.

Pochon X, LaJeunesse TC. 2021. *Miliolidium* n. gen., a new Symbiodiniacean genus whose members associate with soritid foraminifera or are free-living. *J Eukaryot Microbiol* 68:e12856.

Pollock FJ, Katz SM, van de Water JA JM, Davies SW, Hein M, Torda G, Matz MV, Beltran VH, Buerger P, Puill-Stephan E, et al. 2017. Coral larvae for restoration and research: a large-

scale method for rearing *Acropora millepora* larvae, inducing settlement, and establishing symbiosis. *PeerJ* 5:e3732.

Pollock FJ, McMinds R, Smith S, Bourne DG, Willis BL, Medina M, Thurber RV, Zaneveld JR. 2018. Coral-associated bacteria demonstrate phylosymbiosis and cophylogeny. *Nat Commun* 9:4921.

Prada C, Hellberg ME. 2013. Long prereproductive selection and divergence by depth in a Caribbean candelabrum coral. *Proc Natl Acad Sci* 110:3961–6.

Puntin G, Craggs J, Hayden R, Engelhardt KE, McIlroy S, Sweet M, Baker DM, Ziegler M. 2022. The reef-building coral *Galaxea fascicularis*: a new model system for coral symbiosis. *bioRxiv* 1–33. DOI: 10.1101/2022.06.02.494472.

Putnam HM. 2021. Avenues of reef-building coral acclimatization in response to rapid environmental change. *J Exp Biol* 224:jeb239319.

Quigley KM, Baker AC, Coffroth MA, Willis BL, van Oppen MJH. 2018. Bleaching resistance and the role of algal endosymbionts. In: van Oppen MJH, Lough JM, editors. *Coral bleaching: patterns, processes, causes and consequences*. Cham: Springer International Publishing. p. 111–51.

Quigley KM, Willis BL, Kenkel CD. 2019. Transgenerational inheritance of shuffled symbiont communities in the coral *Montipora digitata*. *Sci Rep* 9:13328.

Quiroga-Artigas G, Duscher A, Lundquist K, Waletich J, Schnitzler CE. 2020. Gene knockdown via electroporation of short hairpin RNAs in embryos of the marine hydroid *Hydractinia symbiolongicarpus*. *Sci Rep* 10:12806.

Rädecker N, Pogoreutz C, Voolstra CR, Wiedenmann J, Wild C. 2015. Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends Microbiol* 23:490–97.

Rädecker N, Raina J-B, Pernice M, Perna G, Guagliardo P, Kilburn MR, Aranda M, Voolstra CR. 2018. Using Aiptasia as a model to study metabolic interactions in cnidarian–Symbiodinium symbioses. *Front Physiol* 9:214.

Reaka-Kudla ML, DE Wilson, Wilson EO. 1996. *Biodiversity II: understanding and protecting our biological resources*. Washington (DC):Joseph Henry Press. p. 527.

Reich HG, Kitchen SA, Stankiewicz KH, Devlin-Durante M, Fogarty ND, Baums IB. 2021. Genomic variation of an endosymbiotic dinoflagellate (Symbiodinium “fitti”) among closely related coral hosts. *Mol Ecol* 30:3500–3514.

Reich HG, Robertson DL, Goodbody-Gringley G. 2017. Do the shuffle: changes in Symbiodinium consortia throughout juvenile coral development. *PLoS One* 12:e0171768.

Rentzsch F, Juliano C, Galliot B. 2019. Modern genomic tools reveal the structural and cellular diversity of cnidarian nervous systems. *Curr Opin Neurobiol* 56:87–96.

Rivera HE, Davies SW. 2021. Symbiosis maintenance in the facultative coral, *Oculina arbuscula*, relies on nitrogen cycling, cell cycle modulation, and immunity. *Sci Rep* 11:21226.

Robbins SJ, Singleton CM, Chan CX, Messer LF, Geers AU, Ying H, Baker A, Bell SC, Morrow KM, Ragan MA, et al. 2019. A genomic view of the reef-building coral *Porites lutea* and its microbial symbionts. *Nat Microbiol* 4:2090–2100.

Rohwer F, Seguritan V, Azam F, Knowlton N. 2002. Diversity and distribution of coral-associated bacteria. *Mar Ecol Prog Ser* 243:1–10.

Rose NH, Bay RA, Morikawa MK, Thomas L, Sheets EA, Palumbi SR. 2021. Genomic analysis of distinct bleaching tolerances among cryptic coral species. *Proc Biol Sci* 288:20210678.

Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. 2007. The role of microorganisms in coral health, disease and evolution. *Nat Rev Microbiol* 5:355–62.

Rosenberg E, Zilber-Rosenberg I. 2018. The hologenome concept of evolution after 10 years. *Microbiome* 6:78.

Rosenthal B, Kozhekbaeva Z, Fernhoff N, Tsai JM, Traylor-Knowles N. 2017. Coral cell separation and isolation by fluorescence-activated cell sorting (FACS). *BMC Cell Biol* 18:30.

Santoro EP, Borges RM, Espinoza JL, Freire M, Messias CSMA, Villela HDM, Pereira LM, Vilela CLS, et al. 2021. Coral microbiome manipulation elicits metabolic and genetic restructuring to mitigate heat stress and evade mortality. *Sci Adv* 7.eabg3088.

Scharfenstein HJ, Chan WY, Buerger P, Humphrey C, van Oppen MJH. 2022. Evidence for de novo acquisition of microalgal symbionts by bleached adult corals. *ISME J* 16:1676–9.

Sebé-Pedrós A, Saudemont B, Chomsky E, Plessier F, Mailhé M-P, Renno J, Loe-Mie Y, Lifshitz A, Mukamel Z, Schmutz S, et al. 2018. Cnidarian cell type diversity and regulation revealed by whole-organism single-cell RNA-Seq. *Cell* 173: 1520–34

Serrano XM, Baums IB, Smith TB, Jones RJ, Shearer TL, Baker AC. 2016. Long distance dispersal and vertical gene flow in the Caribbean brooding coral *Porites astreoides*. *Sci Rep* 6: 21619.

Sharp KH, Distel D, Paul VJ. 2012. Diversity and dynamics of bacterial communities in early life stages of the Caribbean coral *Porites astreoides*. *ISME J* 6:790–801.

Sharp KH, Pratte ZA, Kerwin AH, Rotjan RD, Stewart FJ. 2017. Season, but not symbiont state, drives microbiome structure in the temperate coral *Microbiome* 5:120.

Sproles AE, Oakley CA, Matthews JL, Peng L, Owen JG, Grossman AR, Weis VM, Davy SK. 2019. Proteomics quantifies protein expression changes in a model cnidarian colonised by a thermally tolerant but suboptimal symbiont. *ISME J* 13:2334–45.

Starzak DE, Quinnell RG, Nitschke MR, Davy SK. 2014. The influence of symbiont type on photosynthetic carbon flux in a model cnidarian–dinoflagellate symbiosis. *Mar Biol* 161:711–24.

Sully S, Burkepile DE, Donovan MK, Hodgson G, van Woesik R. 2019. A global analysis of coral bleaching over the past two decades. *Nat Commun* 10:1264.

Sweet MJ, Croquer A, Bythell JC. 2011. Bacterial assemblages differ between compartments within the coral holobiont. *Coral Reef* 30:39–52.

Tang K, Zhan W, Zhou Y, Xu T, Chen X, Wang W, Zeng Z, Wang Y, Wang X. 2020. Antagonism between coral pathogen *Vibrio coralliilyticus* and other bacteria in the gastric cavity of scleractinian coral *Galaxea fascicularis*. *Sci China Earth Sci* 63:157–66.

Thornhill DJ, Lewis AM, Wham DC, LaJeunesse TC. 2014. Host-specialist lineages dominate the adaptive radiation of reef coral endosymbionts. *Evolution* 68:352–67.

Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, Bourne DG, Cantin N, Foret S, Matz M, et al. 2017. Rapid adaptive responses to climate change in corals. *Nat Clim Change* 7:627–36.

Tortorelli G, Oakley CA, Davy SK, van Oppen MJH, McFadden GI. 2022. Cell wall proteomic analysis of the cnidarian ph-

tosymbionts *Breviolum minutum* and *Cladocopium goreau*. *J Eukaryot Microbiol* 69:e12870.

Tortorelli G, Rautengarten C, Bacic A, Segal G, Ebert B, Davy SK, van Oppen MJH, McFadden GI. 2022. Cell surface carbohydrates of symbiotic dinoflagellates and their role in the establishment of cnidarian–dinoflagellate symbiosis. *ISME J* 16:190–99.

Tsang Min Ching SJ, Chan WY, Perez-Gonzalez A, Hillyer KE, Buerger P, van Oppen MJH. 2022. Colonization and metabolic profiles of homologous, heterologous and experimentally evolved algal symbionts in the sea anemone *Exaiptasia diaphana*. *ISME Commun* 2:1–10.

van Oppen MJH, Blackall LL. 2019. Coral microbiome dynamics, functions and design in a changing world. *Nat Rev Microbiol* 17:557–67.

Voolstra CR, Buitrago-López C, Perna G, Cárdenas A, Hume BCC, Rädecker N, Barshis DJ. 2020. Standardized short-term acute heat stress assays resolve historical differences in coral thermotolerance across microhabitat reef sites. *Global Change Biol* 26:4328–43.

Voolstra CR, Ziegler M. 2020. Adapting with microbial help: microbiome flexibility facilitates rapid responses to environmental change. *Bioessays* 42:2000004.

Wang S, Zhang L, Meyer E, Matz MV. 2009. Construction of a high-resolution genetic linkage map and comparative genome analysis for the reef-building coral *Acropora millepora*. *Genome Biol* 10:R126.

Webster NS, Reusch TBH. 2017. Microbial contributions to the persistence of coral reefs. *ISME J* 11:2167–74.

Weis VM, Davy SK, Hoegh-Guldberg O, Rodriguez-Lanetty M, Pringle JR. 2008. Cell biology in model systems as the key to understanding corals. *Trends Ecol Evol* 23:369–76.

Weis VM, Reynolds WS, deBoer MD, Krupp DA. 2001. Host-symbiont specificity during onset of symbiosis between the dinoflagellates *Symbiodinium* spp. and planula larvae of the scleractinian coral *Fungia scutaria*. *Coral Reefs* 20:301–8.

Wepfer PH, Nakajima Y, Hui FKC, Mitarai S, Economo EP. 2020. Metacommunity ecology of *Symbiodiniaceae* hosted by the coral *Galaxea fascicularis*. *Mar Ecol Prog Ser* 633:71–87.

Williams A, Chiles EN, Conetta D, Pathmanathan JS, Cleves PA, Putnam HM, Su X, Bhattacharya D. 2021. Metabolome shift associated with thermal stress in coral holobionts. *Science Advances* 7 1–9

Williams LM, Fuess LE, Brennan JJ, Mansfield KM, Salas-Rodriguez E, Welsh J, Awtry J, Banic S, Chacko C, Chezian A, et al. 2018. A conserved toll-like receptor-to-NF- κ B signaling pathway in the endangered coral *Orbicella faveolata*. *Dev Comp Immunol* 79:128–36.

Williams SD, Klings JG, Zinman S, Clark AS, Bartels E, Villoch Diaz Maurino M, Muller EM. 2022. Geographically driven differences in microbiomes of *Acropora cervicornis* originating from different regions of Florida's Coral Reef. *PeerJ* 10:e13574.

Wolenski FS, Bradham CA, Finnerty JR, Gilmore TD. 2013. NF- κ B is required for cnidocyte development in the sea anemone *Nematostella vectensis*. *Dev Biol* 373:205–15.

Wolfowicz I, Baumgarten S, Voss PA, Hambleton EA, Voolstra CR, Hatta M, Guse A. 2016. *Aiptasia* sp. larvae as a model to reveal mechanisms of symbiont selection in cnidarians. *Sci Rep* 6:32366.

Wooldridge SA. 2010. Is the coral-algae symbiosis really “mutually beneficial” for the partners? *Bioessays* 32:615–25.

Wuitchik DM, Almanzar A, Benson BE, Brennan S, Chavez JD, Liesegang MB, Reavis JL, Reyes CL, Schniedewind MK, Trumble IF, et al. 2021. Characterizing environmental stress responses of aposymbiotic *Astrangia poculata* to divergent thermal challenges. *Mol Ecol* 30:5064–79.

Xiang T, Hambleton EA, DeNofrio JC, Pringle JR, Grossman AR. 2013. Isolation of clonal axenic strains of the symbiotic dinoflagellate *Symbiodinium* and their growth and host specificity(1). *J Phycol* 49:447–58.

Yoshimura A, Wakabayashi Y, Mori T. 2010. Cellular and molecular basis for the regulation of inflammation by TGF-beta. *J Biochem* 147:781–92.

Zhang Y, Yang Q, Ling J, Long L, Huang H, Yin J, Wu M, Tang X, Lin X, Zhang Y, Dong J. 2021. Shifting the microbiome of a coral holobiont and improving host physiology by inoculation with a potentially beneficial bacterial consortium. *BMC Microbiol* 21:130.

Ziegler M, Grupstra CGB, Barreto MM, Eaton M, BaOmar J, Zubier K, Al-Sofyani A, Turki AJ, Ormond R, Voolstra CR. 2019. Coral bacterial community structure responds to environmental change in a host-specific manner. *Nat Commun* 10:3092.