

Multi-experiment Parameter Identifiability of ODEs and Model Theory*

Alexey Ovchinnikov[†], Anand Pillay[‡], Gleb Pogudin[§], and Thomas Scanlon[¶]

Abstract. Structural identifiability is a property of an ODE model with parameters that allows for the parameters to be determined from continuous noise-free data. This is a natural prerequisite for practical identifiability. Conducting multiple independent experiments could make more parameters or functions of parameters identifiable, which is a desirable property to have. How many experiments are sufficient? In the present paper, we provide an algorithm to determine the exact number of experiments for multi-experiment local identifiability and obtain an upper bound that is off at most by one for the number of experiments for multi-experiment global identifiability. Interestingly, the main theoretical ingredient of the algorithm has been discovered and proved using model theory (in the sense of mathematical logic). Based on the insights from the model-theoretic argument, an algebraic proof presented in the paper was obtained. We hope that this unexpected connection will stimulate interactions between applied algebra and model theory, and we provide a short introduction to model theory in the context of parameter identifiability. As another related application of model theory in this area, we construct a nonlinear ODE system with one output such that single-experiment and multiple-experiment identifiability are different for the system. This contrasts with recent results about single-output linear systems. We also present a Monte Carlo randomized version of the algorithm with a polynomial arithmetic complexity. Implementation of the algorithm is provided and its performance is demonstrated on several examples.

Key words. parameter identifiability, multiple experiments, differential algebra, model theory of differential fields

MSC codes. 12H05, 34A55, 92B05, 93C15, 93B25, 93B30, 03C40, 03C45, 03C60

DOI. 10.1137/21M1389845

1. Introduction. Structural identifiability is a property of an ODE system with parameters that allows for the parameters to be uniquely determined (global identifiability) or determined up to finitely many choices (local identifiability) from noiseless data and sufficiently exciting inputs (also known as the persistence of excitation; see [13, 29, 31]). Performing structural identifiability analysis is an important first step in evaluating and, if needed, adjusting the system before a reliable practical parameter identification is performed.

*Received by the editors January 7, 2021; accepted for publication (in revised form) March 25, 2022; published electronically July 28, 2022.

<https://doi.org/10.1137/21M1389845>

Funding: This work was partially supported by NSF grants CCF-1564132 and 1563942, DMS-1760448, 1760413, 1853650, 1665035, 1760212, 1853482, 1800492, and 2054721, by the Paris Ile-de-France Region, and by the Fields Institute for Research in Mathematical Sciences.

[†]Department of Mathematics, CUNY Queens College and Ph.D. Programs in Mathematics and Computer Science, CUNY Graduate Center, New York, NY 11367 USA (aovchinnikov@qc.cuny.edu).

[‡]Department of Mathematics, University of Notre Dame, Notre Dame, IN 46556 USA (Anand.Pillay.3@nd.edu).

[§]Prior addresses: New York University, Courant Institute of Mathematical Sciences; Department of Computer Science, Higher School of Economics (Moscow). Current address: LIX, CNRS, École Polytechnique, Institut Polytechnique de Paris, Palaiseau, France 91120 (gleb.pogudin@polytechnique.edu).

[¶]Department of Mathematics, UC Berkeley, Berkeley, CA 94720-3840 USA (scanlon@math.berkeley.edu).

For an ODE model, some of the parameters or functions of parameters could be non-identifiable from a single experiment, but these parameters could become identifiable if one conducts more than one experiment [27, 17]. Knowing the number of experiments to be conducted to achieve the maximal possible identifiability (that is, if the identifiability did not occur after this many experiments, it will not occur after additional experiments) is important for designing experimental protocols involving several experiments [27, section III.B]. In particular, making this number smaller would allow for less expensive experimental protocols. Also, knowing these bounds, one can use existing software for assessing local [24, 12, 28] or global [9] single-experiment identifiability to check multi-experiment identifiability of parameters (or functions of parameters) of interest. Note that, due to [17, Theorem 19], one can alternatively use software based on input-output equations [23, 16] to find the multi-experiment identifiable functions, but this approach does not determine the number of experiments.

One can find such a number of experiments for the case of local identifiability using the algorithm presented in [27] if all parameters are locally identifiable. In [17, section 4], we gave an algorithm computing, among other things, an upper bound for the number of experiments to achieve the maximal possible global identifiability. The proposed algorithm had two drawbacks: it used the Rosenfeld–Gröbner algorithm for differential elimination, which may be computationally very expensive, and the resulting bound could be arbitrarily far from the exact one.

In this paper, we present an algorithm that computes

- the smallest number of experiments to achieve the maximal possible local identifiability and
- the number of experiments to achieve the maximal possible global identifiability so that this number exceeds the minimal such number by at most one.

We present a randomized Monte Carlo version of this algorithm having polynomial arithmetic complexity (see section 6.1). We have implemented this algorithm in Julia language, demonstrated its performance on several examples, and compared with the algorithm from [17] (see section 6.2).

Our algorithm is based on theoretical properties of multi-experiment identifiability that we establish (summarized in section 3). The process of discovering and establishing these properties originated from model theory (in the sense of mathematical logic), and the paper also includes algebraic proofs of these properties for the reader's convenience. The use of model theory in this area is novel, so we give a brief overview here. Already differential algebra plays a large role in structural identifiability (see, e.g., [13]). Differential algebra and the study of solution sets of ODEs in differential rings and fields are enhanced by model-theoretic perspectives and methods, especially from *stability theory*. In particular, differential fields of definition (of differential ideals) are special cases of *canonical bases* from model theory. Using model-theoretic ideas in a nontrivial way, we will prove new quantitative results on recovering such differential fields of definitions from sufficiently many *independent solutions*. In section 7, we will elaborate on the relationship between the setup of multi-experiment identifiability and that of differential algebra/model theory, where we will give additional references.

We also use model-theoretic tools to construct an example contrasting with recent results about multi-experiment identifiability of linear systems [19]. It follows from [19, Theorem 1]

that single-experiment and multi-experiment identifiability are the same thing for linear ODE systems with one output. In section 4, we show a series of nonlinear systems for which this is not the case.

The paper is organized as follows. In section 2, we give basic definitions from differential algebra and structural identifiability. Section 3 summarizes our main results. Section 4 contains the example of a single-output system for which single-experiment and multi-experiment identifiability do not coincide. In section 5, we give an algebraic proof of Theorem 3.1, which is the main theoretical ingredient of our algorithm. In section 6, we present our algorithm, analyze its complexity, describe our implementation, and demonstrate it on a set of examples (including comparison with the algorithm from [17]). In section 7, we describe the connections between identifiability and model theory and explain the model-theoretic context of the results in this paper. The section is aimed at readers who are interested in knowing why and how model-theoretic methods are useful in or relevant to identifiability problems. Although the reader need not be a specialist in model theory, they should either be acquainted with the basic notions or be willing to follow up the references.

Our implementation together with all examples used in the paper can be found at <https://github.com/pogudingleb/ExperimentsBound>.

2. Preliminaries. We assume that all fields have zero characteristic.

2.1. Differential algebra.

Definition 2.1 (differential rings and fields).

- A *differential ring* $(R, ')$ is a commutative ring with a derivation $' : R \rightarrow R$, that is, a map such that, for all $a, b \in R$, $(a + b)' = a' + b'$ and $(ab)' = a'b + ab'$. A differential ring that is also a field is called a *differential field*.
- For an extension of differential fields $F \subset E$ and elements $a_1, \dots, a_n \in E$, let $F\langle a_1, \dots, a_n \rangle$ denote the smallest differential subfield of E containing F and $a_1, \dots, a_n$.

Definition 2.2 (differential polynomials and differential ideals).

- The *ring of differential polynomials* in the variables $x_1, \dots, x_n$ over a field K is the ring $K[x_j^{(i)} \mid i \geq 0, 1 \leq j \leq n]$ with a derivation defined on the ring by $(x_j^{(i)})' := x_j^{(i+1)}$. This differential ring is denoted by $K\{x_1, \dots, x_n\}$.
- An ideal I of a differential ring $(R, ')$ is called a *differential ideal* if, for all $a \in I$, $a' \in I$. For $F \subset R$, the smallest differential ideal containing set F is denoted by $[F]$.
- For an ideal I and element a in a ring R , we denote $I : a^\infty = \{r \in R \mid \exists \ell : a^\ell r \in I\}$. This set is an ideal in R .

2.2. Identifiability. We will consider an *algebraic differential model*

$$(2.1) \quad \Sigma := \begin{cases} \bar{x}' = \bar{f}(\bar{x}, \bar{\mu}, \bar{u}), \\ \bar{y} = \bar{g}(\bar{x}, \bar{\mu}, \bar{u}), \end{cases}$$

where

- $\bar{f} = (f_1, \dots, f_n)$ and $\bar{g} = (g_1, \dots, g_m)$ are tuples of rational functions over $\mathbb{C}$;
- $\bar{x}$, $\bar{u}$, $\bar{y}$ are state, input, and output variables, respectively;

- $\bar{\mu} = (\mu_1, \dots, \mu_\ell)$ are parameters.

The analytic notion of identifiability [10, Definition 2.5] is equivalent (see [10, Proposition 3.4] and [19, Proposition 4.7]) to the following algebraic definition, which we will use.

We write $\bar{f} = \frac{\bar{F}}{\bar{Q}}$ and $\bar{g} = \frac{\bar{G}}{\bar{Q}}$, where $\bar{F}$ and $\bar{G}$ are tuples of polynomials over $\mathbb{C}(\bar{\mu})$ and $\bar{Q}$ is the common denominator of $\bar{f}$ and $\bar{g}$. Here we consider $\mathbb{C}(\bar{\mu})$ as a differential field of constants. Then we define a differential ideal

$$(2.2) \quad I_\Sigma := [Qx'_1 - F_1, \dots, Qx'_n - F_n, Qy_1 - G_1, \dots, Qy_m - G_m]: Q^\infty \subset \mathbb{C}(\bar{\mu})\{\bar{x}, \bar{y}, \bar{u}\}.$$

Observe that every solution of (2.1) is a solution of I_Σ .

Definition 2.3 (generic solution). A tuple $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ from a differential field $k \supset \mathbb{C}(\bar{\mu})$ is called a *generic solution* of (2.1) if, for every differential polynomial $P \in \mathbb{C}(\bar{\mu})\{\bar{x}, \bar{y}, \bar{u}\}$,

$$P(\bar{x}^*, \bar{y}^*, \bar{u}^*) = 0 \iff P \in I_\Sigma.$$

Remark 2.4. It follows from [10, Lemma 3.2] that I_Σ is a prime differential ideal. Therefore, it has a generic solution.

Definition 2.5 (identifiability: single-experiment). For a model Σ in (2.1), a rational function $h \in \mathbb{C}(\bar{\mu})$ is said to be *globally (resp., locally) single-experiment identifiable* (SE-identifiable) if, for every generic solution $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ of (2.1) considered as a system of differential equations over $\mathbb{C}(\bar{\mu})$, we have

$$h(\bar{\mu}) \in \mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle \quad (\text{resp., } h(\bar{\mu}) \text{ is algebraic over } \mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle).$$

Remark 2.6. The equivalence of Definition 2.5 to the more common analytic definition of identifiability has been established in [10, Proposition 3.4]. For a comparison with other definitions, see [1, section 2.1.1].

Definition 2.7 (identifiability defect). For a model Σ in (2.1), we define the *identifiability defect* as

$$\text{defect}(\Sigma) := \text{trdeg}_{\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{\mu})\langle \bar{y}^*, \bar{u}^* \rangle,$$

where $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ is any generic solution of (2.1) (one can show that the defect does not depend on the choice of the generic solution).

For example, $\text{defect}(\Sigma) = 0$ implies that all the parameters are locally identifiable.

Definition 2.8 (identifiability: multi-experiment [17, Definition 16]).

- For a model Σ and a positive integer r , we define the *r-fold replica* of Σ as

$$\Sigma_r := \begin{cases} \bar{x}'_i = \bar{f}(\bar{x}_i, \bar{\mu}, \bar{u}_i), & i = 1, \dots, r, \\ \bar{y}_i = \bar{g}(\bar{x}_i, \bar{\mu}, \bar{u}_i), & i = 1, \dots, r, \end{cases}$$

where $\bar{x}_1, \dots, \bar{x}_r, \bar{y}_1, \dots, \bar{y}_r, \bar{u}_1, \dots, \bar{u}_r$ are new tuples of indeterminates (note that the vector of parameters is not being replicated).

- For a model Σ , a rational function $h \in \mathbb{C}(\bar{\mu})$ is called *globally (resp., locally) multi-experiment identifiable* (ME-identifiable) if there exists a positive integer r such that $h(\bar{\mu})$ is globally (resp., locally) SE-identifiable in Σ_r .

Notation 2.9. For a differential model Σ , we define $\text{NumExpGlob}(\Sigma)$ (resp., $\text{NumExpLoc}(\Sigma)$) as the smallest integer r such that, for every $h(\bar{\mu}) \in \mathbb{C}(\bar{\mu})$, the following are equivalent:

- h is globally (resp., locally) ME-identifiable for Σ ;
- h is globally (resp., locally) SE-identifiable for Σ_r .

In what follows, we will frequently (but not always) omit writing “global/globally” for brevity.

Remark 2.10. A simple family of linear models Σ_r that reaches arbitrarily high values for the number of experiments (more precisely, growing linearly with the number of state variables, which can be made arbitrarily high in the model), $\text{NumExpLoc}(\Sigma_r) = \text{NumExpGlob}(\Sigma_r) = r$, is [17, Example 30].

3. Main results. Our results consist of a theoretical part and algorithms building upon the theory. The theoretical contribution is summarized in the statements below. In particular, Theorem 3.1 is the basis for an algorithm for computing $\text{NumExpLoc}(\Sigma)$ and obtaining an upper bound for $\text{NumExpGlob}(\Sigma)$ that is off at most by one.

Theorem 3.1. *For every algebraic differential model Σ of the form (2.1), we have the following:*

1. $\text{NumExpLoc}(\Sigma) = \min\{r \mid \text{defect}(\Sigma_r) = \text{defect}(\Sigma_{r+1})\}$.
2. $\text{NumExpLoc}(\Sigma) \leq \text{NumExpGlob}(\Sigma) \leq \text{NumExpLoc}(\Sigma) + 1$.

Note that the theorem does not imply that parameters locally identifiable from r experiments are globally identifiable from $r + 1$ experiments.

Corollary 3.2. *For every algebraic differential model Σ of the form (2.1) with ℓ parameters,*

$$\text{NumExpLoc}(\Sigma) \leq \ell \quad \text{and} \quad \text{NumExpGlob}(\Sigma) \leq \ell + 1.$$

It follows from [19, Theorem 4.2] that, for single-output linear models, identifiable and multi-experiment identifiable functions coincide. Proposition 3.3 indicates that this does not generalize to nonlinear models.

Proposition 3.3. *There is a system of the form (2.1) with a single output such that the fields of identifiable and multi-experiment identifiable functions do not coincide.*

Proofs of Theorem 3.1 and Proposition 3.3 are presented in sections 5 and 4, respectively.

On the algorithmic side, Theorem 3.1 yields a probabilistic algorithm for computing the value $\text{NumExpLoc}(\Sigma)$ and a bound for $\text{NumExpGlob}(\Sigma)$ that is off at most by one with arithmetic complexity being polynomial in the complexity of the system (Proposition 6.5). We implemented this algorithm, and we demonstrate its practical performance and apply it to examples in section 6.

4. Single-output model requiring more than one experiment. In this section, we will prove Proposition 3.3 by showing that the SE-identifiable and ME-identifiable functions do

not coincide for the following model Σ [18, Example 5.8]:

$$(4.1) \quad \begin{cases} x'_1 = 0, \\ x'_2 = x_1x_2 + \mu_1x_1 + \mu_2, \\ y = x_2. \end{cases}$$

We will now give a direct algebraic proof. In section 7.2, we present the model-theoretic argument that has been used to construct this example and can be used to construct more complex ones.

Lemma 4.1. *The field of ME-identifiable functions of (4.1) is $\mathbb{C}(\mu_1, \mu_2)$, but neither μ_1 nor μ_2 is SE-identifiable.*

Proof. We find the field of ME-identifiable functions using [17, Theorem 19]. Differentiating the second equation in (4.1), we get $x''_2 = x_1x'_2$. Using this equation, we can eliminate x_1 from the second equation of (4.1) and obtain

$$(4.2) \quad x_2x''_2 - (x'_2)^2 + \mu_1x''_2 + \mu_2x'_2 = 0.$$

Since x_2 does not satisfy any first-order equation over $\mathbb{C}(\mu_1, \mu_2)$ modulo I_Σ and (4.2) is irreducible, the set consisting of (4.2) is a set of input-output equations for Σ . The coefficients of (4.2) are 1, -1 , μ_1 , and μ_2 . So, according to [17, Definition 15], the coefficients μ_1 and μ_2 are input-output identifiable. The first paragraph of [17, Theorem 19] now implies that the coefficients μ_1 and μ_2 of (4.2) are ME-identifiable.

To prove that μ_1 and μ_2 are not SE-identifiable, consider a generic solution (x_1^*, x_2^*, y^*) of (4.1). Then there is a differential automorphism of $\mathbb{C}(\bar{\mu})\langle x_1^*, x_2^*, y^* \rangle$ defined by

$$\alpha|_{\mathbb{C}\langle x_1^*, x_2^*, y^* \rangle} = \text{id}, \quad \alpha(\mu_1) = \mu_1 + 1, \quad \alpha(\mu_2) = \mu_2 - x_1.$$

Therefore, neither μ_1 nor μ_2 belongs to $\mathbb{C}\langle y^* \rangle$. ■

Remark 4.2. Using [17, Algorithm 1], one can show that the field of SE-identifiable functions of (4.1) is $\mathbb{C}$.

5. Bounding the number of experiments (proof of Theorem 3.1). In this section, we will give an algebraic proof (but with a strong model-theoretic flavor, which we expand in section 7.3) of Theorem 3.1. We start with fixing some notation for the section.

Notation 5.1.

- We use $\bar{*}$ for tuples.
- $\bar{x}, \bar{y}$, and $\bar{z}$ denote tuples of differential indeterminates. Moreover, we will assume that $|\bar{a}| = |\bar{x}|$, $|\bar{b}| = |\bar{y}|$, and $|\bar{c}| = |\bar{z}|$ for tuples $\bar{a}, \bar{b}$, and $\bar{c}$ of elements from some differential field.
- k_0 will be a fixed differential ground field (in applications, $k_0 = \mathbb{Q}, \mathbb{R}, \mathbb{C}$ with zero derivation). We will also consider an extension $K \supset k_0$ such that K is differentially closed and a $|k_0|$ -saturated field. Saturation and differentially closed fields are defined in section 7; in this section, we will use only the following algebraic consequence of these properties [15, Propositions 4.2.13 and 4.3.3 and page 117]: for every differential subfield $k \subset K$ of

cardinality at most $|k_0|$ and every differential automorphism α of k , α can be extended to an endomorphism of K .

- Let E be a field and $\bar{a}$ be a tuple of elements from some extension of E . Then $\text{trdeg}_E \bar{a}$ denotes $\text{trdeg}_E E(\bar{a})$.

Notation 5.2. Let $k \subset K$ be an intermediate differential field (in applications, we will have $k = \mathbb{C}(\bar{\mu})$) and $\bar{a}$ a tuple from K .

- The vanishing ideal of $\bar{a}$ over k is denoted by

$$\mathcal{I}_k(\bar{a}) := \{p \in k\{\bar{x}\} \mid p(\bar{a}) = 0\}.$$

- We denote the differential-algebraic variety of $\bar{a}$ with respect to K defined over k by

$$\mathcal{V}_{K/k}(\bar{a}) = \{\bar{b} \in K^{|\bar{a}|} \mid p(\bar{b}) = 0 \quad \forall p \in \mathcal{I}_k(\bar{a})\} \subset K^{|\bar{a}|}.$$

We consider this as a differential-algebraic variety over K , and it is not necessarily irreducible. For brevity, until the end of the section, by “variety” we will mean “differential-algebraic variety.”

- The field of definition of a (differential) ideal I in $k\{\bar{x}\}$ is the smallest (differential) subfield L of k such that I is generated as an ideal by $I \cap L\{\bar{x}\}$.
- $\text{FD}_k(\bar{a})$ denotes the field generated by k_0 and the field of definition of $\mathcal{I}_k(\bar{a})$.
- Let $\langle \mathcal{I}_k(\bar{a}), \mathcal{I}_k(\bar{b}) \rangle$ denote the ideal in $k\{\bar{x}, \bar{y}\}$ generated by $\mathcal{I}_k(\bar{a}) \subset k\{\bar{x}\}$ and $\mathcal{I}_k(\bar{b}) \subset k\{\bar{y}\}$. For tuples $\bar{a}_1, \dots, \bar{a}_n$, the ideal $\langle \mathcal{I}_k(\bar{a}_1), \dots, \mathcal{I}_k(\bar{a}_n) \rangle$ is defined analogously.
- For tuples $\bar{a}_1$ and $\bar{a}_2$ of the same length, we write $\mathcal{I}_k(\bar{a}_1) \cong \mathcal{I}_k(\bar{a}_2)$ if the ideals $\mathcal{I}_k(\bar{a}_1)$ and $\mathcal{I}_k(\bar{a}_2)$ coincide if being considered in the same ring $k\{\bar{x}\}$.

Lemma 5.3. Let $k \subset K$ be a differential subfield and $\bar{a}$ and $\bar{b}$ be tuples from K such that $\mathcal{I}_k(\bar{a}, \bar{b}) = \langle \mathcal{I}_k(\bar{a}), \mathcal{I}_k(\bar{b}) \rangle$. Then $\mathcal{I}_{k(\bar{b})}(\bar{a})$ is generated by $\mathcal{I}_k(\bar{a})$.

Proof. By clearing denominators, every element $p \in \mathcal{I}_{k(\bar{b})}(\bar{a})$ can be written as $p = q(\bar{x}, \bar{b})/d$, where $d \in k(\bar{b})$ and $q(\bar{x}, \bar{y}) \in \mathcal{I}_k(\bar{a}, \bar{b})$. The differential polynomial $q(\bar{x}, \bar{y})$ can be written as a combination of elements of $\mathcal{I}_k(\bar{a})$ and $\mathcal{I}_k(\bar{b})$. If we plug $\bar{y} = \bar{b}$, the terms from $\mathcal{I}_k(\bar{b})$ will vanish, so $q(\bar{x}, \bar{b})$ can be written as a combination of elements of $\mathcal{I}_k(\bar{a})$. Then the same is true for p . ■

Proposition 5.4 (cf. [17, Theorem 19]—computing an upper bound for N). Let $k \subset K$ be a differential subfield and $\bar{a}_1, \bar{a}_2, \dots$ be tuples of the same length from K with

$$\mathcal{I}_k(\bar{a}_1) \cong \mathcal{I}_k(\bar{a}_2) \cong \dots \quad \text{and} \quad \mathcal{I}_k(\bar{a}_1, \dots, \bar{a}_\ell) = \langle \mathcal{I}_k(\bar{a}_1), \dots, \mathcal{I}_k(\bar{a}_\ell) \rangle \text{ for every } \ell \geq 1.$$

Then there exists N such that (see Notation 5.2)

$$\text{FD}_k(\bar{a}_1) \subset k_0\langle \bar{a}_1, \dots, \bar{a}_N \rangle.$$

Proof. Let L be generated by the field of definition of $J := \mathcal{I}_k(\bar{a}_1)$ and by k_0 . We consider an arbitrary ordering of the monomials of the corresponding differential ring and consider a linear basis of J that is in the reduced row echelon form with respect to this ordering (this construction is described in more detail in the proof of [18, Theorem 4.7]). The elements of this basis form a set of generators $\{f_\lambda\}_{\lambda \in \Lambda}$ of J such that, for every $\lambda \in \Lambda$, the following hold:

- the coefficients of f_λ are in L and at least one of them is 1;
- for every $g \in J \setminus \{f_\lambda\}$, the support of $f_\lambda - g$ is not a proper subset of the support of f_λ .

Then the coefficients of $\{f_\lambda\}_{\lambda \in \Lambda}$ generate L over k_0 . We fix some λ and write $f_\lambda = m_0 + c_1 m_1 + \cdots + c_N m_N$, where $m_1, \dots, m_N$ are differential monomials and $c_1, \dots, c_N \in L$. We will show that $c_1, \dots, c_N \in \mathbb{Q}\langle \bar{a}_1, \dots, \bar{a}_N \rangle$. For every $j \geq 1$, we denote the monomial m_i evaluated at $\bar{a}_j$ by $m_{j,i}$. Then we have a linear system in $c_1, \dots, c_N$

$$\begin{pmatrix} m_{1,1} & \cdots & m_{1,N} \\ \vdots & \ddots & \vdots \\ m_{N,1} & \cdots & m_{N,N} \end{pmatrix} \begin{pmatrix} c_1 \\ \vdots \\ c_N \end{pmatrix} = \begin{pmatrix} -m_{1,0} \\ \vdots \\ -m_{N,0} \end{pmatrix}.$$

We denote the matrix of the above system by S . If $\det S \neq 0$, then, solving the system, we show that $c_1, \dots, c_N \in \mathbb{Q}\langle \bar{a}_1, \dots, \bar{a}_N \rangle$. Assume that $\det S = 0$, and let S_0 be the smallest singular minor of S . Let the index of the first row of S_0 be s , and define $E := k\langle \bar{a}_1, \dots, \bar{a}_{s-1}, \bar{a}_{s+1}, \dots, \bar{a}_N \rangle$. Expanding the equality $\det S_0 = 0$ with respect to the first row, we obtain an element p of $\mathcal{I}_E(\bar{a}_s)$. The minimality of S_0 implies that $p \neq 0$. Since the support of p is a proper subset of the support of f_λ , we conclude that $\mathcal{I}_E(\bar{a}_s)$ is not generated by $\mathcal{I}_k(\bar{a}_s)$. This contradicts Lemma 5.3.

Since L is finitely generated (follows, for example, from [19, Proposition 4.11]), the coefficients of finitely many f_λ 's generate L over k . Taking the maximum of the corresponding N 's will give the desired N and finish the proof. ■

Lemma 5.5. *Let k with $k_0 \subset k \subset K$ be a differential field differentially finitely generated over k_0 and $\bar{a}$ be a tuple from K . Then, for every component C of $\mathcal{V}_{K/k}(\bar{a})$, there exists $\bar{b} \in C$ such that $\mathcal{I}_{k_0}(\bar{a}) \cong \mathcal{I}_{k_0}(\bar{b})$.*

Proof. The ideal $\mathcal{I}_k(\bar{a})$ is prime. Then the Galois group of $k^{\text{alg}} \subset K$ over k acts transitively on the components of $\mathcal{V}_{K/k}(\bar{a})$. Let C_0 be a component containing $\bar{a}$, and let α be an automorphism of k^{alg} over k that maps C_0 to C . By Notation 5.1, since $|k| = |k_0|$, α can be lifted to a differential endomorphism of K which we will denote by α as well. We set $\bar{b} := \alpha(\bar{a})$. Then we have $\mathcal{I}_{k_0}(\bar{a}) \cong \mathcal{I}_{k_0}(\bar{b})$ due to the α -invariance of k_0 . ■

Lemma 5.6. *Let k with $k_0 \subset k \subset K$ be a differential subfield. Let the following hold:*

- $\mathcal{I}_k(\bar{a}, \bar{b}) = \langle \mathcal{I}_k(\bar{a}), \mathcal{I}_k(\bar{b}) \rangle$;
- $\bar{c}$ from k is such that $\text{FD}_k(\bar{a}) \cup \text{FD}_k(\bar{b}) \subset k_0\langle \bar{c} \rangle$ (see Notation 5.2);
- a component $C \subset \mathcal{V}_{K/k_0\langle \bar{a} \rangle}(\bar{c})$ is such that $C \subset \mathcal{V}_{K/k_0\langle \bar{b} \rangle}(\bar{c})$.

Then C is a component of $\mathcal{V}_{K/k_0\langle \bar{a}, \bar{b} \rangle}(\bar{c})$.

Proof. Consider any $p \in \mathcal{I}_{k_0\langle \bar{a}, \bar{b} \rangle}(\bar{c}) \subset k_0\langle \bar{a}, \bar{b} \rangle\{\bar{z}\}$. Let $d \in k_0\langle \bar{a}, \bar{b} \rangle$ be the product of the denominators of the coefficients of p . Then there exists $q \in k_0\{\bar{x}, \bar{y}, \bar{z}\}$ such that $q(\bar{a}, \bar{b}, \bar{z}) = dp$. Since $q(\bar{a}, \bar{b}, \bar{c}) = 0$, we have

$$q(\bar{x}, \bar{y}, \bar{c}) \in \mathcal{I}_{k_0\langle \bar{c} \rangle}(\bar{a}, \bar{b}) = k_0\langle \bar{c} \rangle\{\bar{x}, \bar{y}\} \mathcal{I}_{k_0\langle \bar{c} \rangle}(\bar{a}) + k_0\langle \bar{c} \rangle\{\bar{x}, \bar{y}\} \mathcal{I}_{k_0\langle \bar{c} \rangle}(\bar{b}),$$

where the latter equality follows from $\mathcal{I}_k(\bar{a}, \bar{b}) = \langle \mathcal{I}_k(\bar{a}), \mathcal{I}_k(\bar{b}) \rangle$ and the fact that $k_0\langle \bar{c} \rangle$ contains the fields of definitions of $\mathcal{I}_k(\bar{a})$ and $\mathcal{I}_k(\bar{b})$. By clearing the denominators with respect to $\bar{c}$,

we conclude that there exists $h(\bar{z}) \in k_0\{\bar{z}\}$ such that $h(\bar{c}) \neq 0$ and

$$h(\bar{z})q(\bar{x}, \bar{y}, \bar{z}) \in k_0\{\bar{x}, \bar{y}, \bar{z}\} \mathcal{I}_{k_0}(\bar{a}, \bar{c}) + k_0\{\bar{x}, \bar{y}, \bar{z}\} \mathcal{I}_{k_0}(\bar{b}, \bar{c}).$$

Thus, $h(\bar{z})q(\bar{a}, \bar{b}, \bar{z})$ vanishes on $\mathcal{V}_{K/k_0\langle\bar{a}\rangle}(\bar{c}) \cap \mathcal{V}_{K/k_0\langle\bar{b}\rangle}(\bar{c})$ and, consequently, on C . If $q(\bar{a}, \bar{b}, \bar{z})$ does not vanish on C , then $h(\bar{z})$ does. However, this is impossible due to Lemma 5.5 because $h(\bar{z}) \notin \mathcal{I}_{k_0}(\bar{c})$.

Thus, p vanishes on C , and so $C \subset \mathcal{V}_{K/k_0\langle\bar{a}, \bar{b}\rangle}(\bar{c})$. On the other hand, $\mathcal{V}_{K/k_0\langle\bar{a}, \bar{b}\rangle}(\bar{c}) \subset \mathcal{V}_{K/k_0\langle\bar{a}\rangle}(\bar{c})$, so C is a component of $\mathcal{V}_{K/k_0\langle\bar{a}, \bar{b}\rangle}(\bar{c})$. ■

Lemma 5.7. *Let k with $k_0 \subset k \subset K$ be a differential field differentially finitely generated over k_0 . Consider tuples $\bar{a}_1, \dots, \bar{a}_\ell$ of the same length from K such that*

$$\mathcal{I}_k(\bar{a}_1) \cong \dots \cong \mathcal{I}_k(\bar{a}_\ell) \quad \text{and} \quad \mathcal{I}_k(\bar{a}_1, \dots, \bar{a}_\ell) = \langle \mathcal{I}_k(\bar{a}_1), \dots, \mathcal{I}_k(\bar{a}_\ell) \rangle.$$

Then, for every permutation $\pi \in S_\ell$, there exists an endomorphism α_π of K over k such that $\alpha_\pi(\bar{a}_i) = \bar{a}_{\pi(i)}$ for every $1 \leq i \leq \ell$.

Proof. Consider the ideal $J = \mathcal{I}_k(\bar{a}_1, \dots, \bar{a}_\ell) \subset k\{\bar{x}_1, \dots, \bar{x}_\ell\}$. We fix $\pi \in S_\ell$. Let β_π be the differential k -automorphism of $k\{\bar{x}_1, \dots, \bar{x}_\ell\}$ defined by $\beta_\pi(\bar{x}_i) = \bar{x}_{\pi(i)}$ for every $1 \leq i \leq \ell$. Since the set $\{\mathcal{I}_k(\bar{a}_1), \dots, \mathcal{I}_k(\bar{a}_\ell)\}$, where each $\mathcal{I}_k(\bar{a}_j)$ is considered as a subset of $k\{\bar{x}_1, \dots, \bar{x}_\ell\}$, is β_π -invariant, so is J . Therefore, β_π yields an automorphism, say α_π , of

$$k\{\bar{x}_1, \dots, \bar{x}_\ell\}/J \cong k\{\bar{a}_1, \dots, \bar{a}_\ell\}.$$

α_π can be lifted uniquely to an automorphism of $k\langle\bar{a}_1, \dots, \bar{a}_\ell\rangle$. The resulting automorphism can be lifted to an endomorphism of K by Notation 5.1 since $|k\langle\bar{a}_1, \dots, \bar{a}_\ell\rangle| = |k_0|$. ■

Lemma 5.8. *Let k with $k_0 \subset k \subset K$ be a differential field and $\bar{a}$ be a tuple from K , and let $F := \text{FD}_k(\bar{a})$ (see Notation 5.2) be such that $\text{trdeg}_F k < \infty$. Then $\text{trdeg}_{F\langle\bar{a}\rangle} k = \text{trdeg}_F k$.*

Proof. Let $\alpha_1, \dots, \alpha_N$ be a transcendence basis of k over F . Assume that $\alpha_1, \dots, \alpha_N$ are algebraically dependent over $F\langle\bar{a}\rangle$. Then there exists $P \in F\{\bar{x}\}[y_1, \dots, y_N]$ such that $P(\bar{a}, \alpha_1, \dots, \alpha_N) = 0$ and $P(\bar{a}, y_1, \dots, y_N) \neq 0$. On the other hand, since the field of definition of $\mathcal{I}_k(\bar{a})$ is F , we have $P(\bar{x}, \alpha_1, \dots, \alpha_N) \in k \cdot (\mathcal{I}_k(\bar{a}) \cap F\{\bar{x}\})$. Therefore, since the monomials in $\alpha_1, \dots, \alpha_N$ are F -linearly independent, every coefficient of P as a polynomial in $y_1, \dots, y_N$ vanishes at $\bar{a}$. Thus, $P(\bar{a}, y_1, \dots, y_N) = 0$, a contradiction. ■

Notation 5.9. For an irreducible differential-algebraic variety $X \subset K^n$, let $\text{adim } X$ denote the algebraic dimension, which is the transcendence degree of the algebra of regular functions. The algebraic dimension of an arbitrary differential-algebraic variety is defined as the maximum of the algebraic dimensions of its components.

Proposition 5.10. *Let the following hold:*

- k with $k_0 \subset k \subset K$ is a differential field of finite transcendence degree over k_0 ;
- $\bar{a}_1, \bar{a}_2, \dots$ are tuples of the same length from K such that

$$\mathcal{I}_k(\bar{a}_1) \cong \mathcal{I}_k(\bar{a}_2) \cong \dots \quad \text{and} \quad \mathcal{I}_k(\bar{a}_1, \dots, \bar{a}_\ell) = \langle \mathcal{I}_k(\bar{a}_1), \dots, \mathcal{I}_k(\bar{a}_\ell) \rangle \text{ for every } \ell \geq 1;$$

- r is the smallest integer such that $\text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle} k = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle} k$.

Then the following hold:

- (1) r is the smallest integer such that $\text{FD}_k(\bar{a}_1)$ (see Notation 5.2) is algebraic over $k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle$;
- (2) $\text{FD}_k(\bar{a}_1) \subset k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle$.

Proof. Let $\bar{c}$ be any set of generators of $\text{FD}_k(\bar{a}_1)$. Consider a sequence of varieties (see Notation 5.2)

$$(5.1) \quad X_0 := \mathcal{V}_{K/k_0}(\bar{c}) \supset X_1 := \mathcal{V}_{K/k_0\langle\bar{a}_1\rangle}(\bar{c}) \supset X_2 := \mathcal{V}_{K/k_0\langle\bar{a}_1, \bar{a}_2\rangle}(\bar{c}) \supset \dots$$

Claim. For every $i \geq 0$, if X_i and X_{i+1} have a common component C , then $C = \{\bar{c}\}$.

Let i be such that X_i and X_{i+1} have a common component C . For every $j \geq 1$, we introduce $Y_j := \mathcal{V}_{K/k_0\langle\bar{a}_j\rangle}(\bar{c})$. Since $X_{i+1} \subset Y_{i+1}$, we have $C \subset Y_{i+1}$. We claim that, for every $j \geq i+1$, $C \subset Y_j$. Lemma 5.7 implies that there exists a k -endomorphism α of K such that α leaves $\bar{a}_1, \dots, \bar{a}_i$ invariant and maps $\bar{a}_{i+1}$ to $\bar{a}_j$. Since C is a component of X_i , it is defined over $k_0\langle\bar{a}_1, \dots, \bar{a}_i\rangle^{\text{alg}}$, and therefore C is α -invariant. Thus, by applying α to the inclusion $C \subset Y_{i+1}$, we obtain $C \subset Y_j$.

By applying Lemma 5.6 iteratively to the component C and $\bar{a} = (\bar{a}_1, \dots, \bar{a}_j)$ and $\bar{b} = \bar{a}_{j+1}$ for $j = i+1, i+2, \dots$, we show that C is a component of X_j for every $j \geq i+1$. On the other hand, Proposition 5.4 implies that there exists N such that $X_N = \{\bar{c}\}$. Thus, $C = \{\bar{c}\}$, and the claim is proved.

Since $\bar{a}_1, \bar{a}_2, \dots$ have the same ideals of definition over k and ideals of the form $\mathcal{I}_k(\bar{a}_1, \dots, \bar{a}_s)$ are generated by the ideals of $\bar{a}_i$'s, we have

$$F := \text{FD}_k(\bar{a}_1) = \text{FD}_k(\bar{a}_1, \bar{a}_2) = \text{FD}_k(\bar{a}_1, \bar{a}_2, \bar{a}_3) = \dots$$

Therefore, for every $s \geq 0$,

$$(5.2) \quad \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_s\rangle} k = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_s\rangle} \bar{c} + \text{trdeg}_{k_0\langle\bar{c}, \bar{a}_1, \dots, \bar{a}_s\rangle} k = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_s\rangle} \bar{c} + \text{trdeg}_F k,$$

where the latter equality is due to Lemma 5.8. Thus, r is the smallest integer such that

$$\text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle} \bar{c} = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle} \bar{c}.$$

Then $\text{adim } X_r = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle} \bar{c} = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle} \bar{c} = \text{adim } X_{r+1}$. Therefore, since $X_r \supset X_{r+1}$, the components of X_{r+1} of maximal dimension are components of X_r . By the claim, these components are all equal to $\{\bar{c}\}$. Therefore, every component of X_{r+1} is $\{\bar{c}\}$, and so $X_{r+1} = \{\bar{c}\}$. Thus, $\text{adim } X_r = 0$. The fact that $\text{adim } X_r = 0$ implies the first part of the proposition, and $X_{r+1} = \{\bar{c}\}$ implies the second part of the proposition. ■

Proof of Theorem 3.1. Consider a generic solution

$$(\bar{x}_1^*, \dots, \bar{x}_{\ell+1}^*, \bar{y}_1^*, \dots, \bar{y}_{\ell+1}^*, \bar{u}_1^*, \dots, \bar{u}_{\ell+1}^*)$$

of $\Sigma_{\ell+1}$. We apply Proposition 5.10 with $\bar{a}_i = (\bar{y}_i^*, \bar{u}_i^*)$ for every $1 \leq i \leq \ell+1$, $k_0 = \mathbb{C}$, $k = \mathbb{C}(\bar{\mu})$. Since the sequence $\text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_i\rangle} k$ for $i = 0, \dots, \ell+1$ is nonincreasing, there will be $r \leq \ell$ as in Proposition 5.10. Furthermore, it will be the same as r in the statement of Theorem 3.1. We have that

- $\text{FD}_k(\bar{a}_1)$ is the field of globally ME-identifiable functions (by [17, Theorem 19] or Proposition 5.4) and
- the field of locally ME-identifiable functions is algebraic over $\text{FD}_k(\bar{a}_1)$ because every locally identifiable function is algebraic over the field of globally identifiable functions.

Hence, r being the smallest number such that $\text{FD}_k(\bar{a}_1)$ is algebraic over $k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle$ implies that r is the smallest number such that the field of locally SE-identifiable functions of Σ_r coincides with the field of locally ME-identifiable functions in Σ . Thus, $\text{NumExpLoc}(\Sigma) = r$. Finally, $\text{FD}_k(\bar{a}_1) \subset k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle$ implies that ME-identifiable functions in Σ are SE-identifiable in Σ_{r+1} , so

$$\text{NumExpGlob}(\Sigma) \leq r + 1 = \text{NumExpLoc}(\Sigma) + 1. \quad \blacksquare$$

6. Algorithm, implementation, and examples.

6.1. Algorithm: Theory. Theorem 3.1 implies the correctness of the following algorithm.

Algorithm 6.1 Computing $\text{NumExpLoc}(\Sigma)$ and estimating $\text{NumExpGlob}(\Sigma)$

In: • an algebraic differential model Σ ;

- (optional; for probabilistic version) real number $0 \leq p < 1$;

Out: positive integer r such that $\text{NumExpLoc}(\Sigma) = r$ and $\text{NumExpGlob}(\Sigma) \in \{r, r + 1\}$. In the probabilistic version, this result will be correct with probability at least p .

Set $d_0 = \ell$. For $i = 1, 2, \dots, \ell + 1$, do:

- 1 Using Algorithm 6.2, compute $d_i = \text{defect}(\Sigma_i)$ (see Definition 2.7);
(in the probabilistic version, the input probability for Algorithm 6.2 is $1 - \frac{1-p}{\ell}$)
 - 2 If $d_i = d_{i-1}$, stop and return $i - 1$.
-

Lemma 6.1. Algorithm 6.1 is correct.

Proof. If the outputs of Algorithm 6.2 are correct, the returned result will be correct due to Theorem 3.1. In the probabilistic version, the probability that at least one of the instances of Algorithm 6.2 will return wrong result does not exceed $\ell(1 - (1 - \frac{1-p}{\ell})) = 1 - p$. $\blacksquare$

Our algorithm for computing the identifiability defect will use, as a subroutine, algorithm(s) described in Theorem 6.3 below.

Notation 6.2. We call the *complexity* of a model Σ the maximum of the total number of variables (parameters, states, inputs, and outputs) and the length of a straight-line program (see [2, Chapter 4.1]) computing the numerators and denominators of the right-hand side of Σ . For measuring the complexity of the algorithms in this section, we use the notion of arithmetic complexity, which is the number of arithmetic operations in the ground field; see [30, Chapter 12] for more details.

Theorem 6.3 (see [24]). Consider the following problem:

In: an algebraic differential model Σ without parameters (that is, $\ell = 0$);

Out: $\text{trdeg}_{\mathbb{C}\langle\bar{y}^*, \bar{u}^*\rangle} \mathbb{C}(\bar{x}^*)$, where $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ is any generic solution of Σ .

Then the following hold:

1. *There exists a deterministic algorithm for solving this problem.*
2. *For Σ with coefficients in $\mathbb{Q}$, there exists a probabilistic Monte Carlo algorithm with polynomial arithmetic complexity with respect to the complexity of Σ .*

Proof. The theorem follows from the results from [24] as follows. For part 1, we use [24, Proposition 2.1] together with [24, Remark, p. 742] to show that $\text{trdeg}_{\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{x}^*) = \text{trdeg}_{\mathcal{G}} \mathbb{C}(\bar{x}^*)$, where $\mathcal{G} = \mathbb{C}(\bar{y}^*, \dots, (\bar{y}^*)^{(n+\ell)}, \bar{u}^*, \dots, (\bar{u}^*)^{(n+\ell)})$. Then $\text{trdeg}_{\mathcal{G}} \mathbb{C}(\bar{x}^*)$ can be computed using [24, Corollary 2.1], in which X is $\bar{x}^*$, Y is $\bar{y}^*$, and $\mathcal{G}$ is $\mathcal{G}$.

Part 2 follows from [24, Theorem 1.1] (together with a more precise complexity bound) as follows. The algorithm whose existence is stated in [24, Theorem 1.1] computes the smallest number of nonobservable state variables that are assumed to be known in order to make the system observable. The definition of observability [24, section 2.1] implies that the components of $\bar{x}^*$ corresponding to such a set of state variables is a transcendence basis of $\mathbb{Q}(\bar{x}^*)$ over $\mathbb{Q}\langle \bar{y}^*, \bar{u}^* \rangle$, which is also a transcendence basis of $\mathbb{C}(\bar{x}^*)$ over $\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle$, so the cardinality of this set is the desired transcendence degree. ■

Algorithm 6.2 Computing $\text{defect}(\Sigma)$

In: • an algebraic differential model Σ ;

- (optional; for probabilistic version) real number $0 \leq p < 1$;

Out: $\text{defect}(\Sigma)$. In the probabilistic version, this result is correct with probability at least p .

- 1 Construct two parameter-free algebraic differential models:
 - (a) Σ' obtained from Σ by viewing all parameters as state variables satisfying equations $\mu'_i = 0$ for every $\mu_i \in \bar{\mu}$;
 - (b) Σ'' obtained from Σ' by adding a new output for each state variable corresponding to a parameter of Σ .
 - 2 Run any of the algorithms from Theorem 6.3 on Σ' and Σ'' , denote the results by A and B , respectively (in the probabilistic version, the input probability is $\frac{1+p}{2}$).
 - 3 return $A - B$.
-

Lemma 6.4. *Algorithm 6.2 is correct.*

Proof. Let $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ be a generic solution of (2.1). Then

$$A = \text{trdeg}_{\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{x}^*, \bar{\mu}) \quad \text{and} \quad B = \text{trdeg}_{\mathbb{C}(\bar{\mu})\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{x}^*, \bar{\mu}).$$

Since one can compose a transcendence basis of $(\bar{x}^*, \bar{\mu})$ over $\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle$ by first taking a transcendence basis of $\bar{\mu}$ over $\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle$ which is of cardinality $\text{defect}(\Sigma)$ and then taking a transcendence basis of $\bar{x}^*$ over $\mathbb{C}(\bar{\mu})\langle \bar{y}^*, \bar{u}^* \rangle$, we have

$$\text{defect}(\Sigma) = \text{trdeg}_{\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{\mu}) = \text{trdeg}_{\mathbb{C}\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{x}^*, \bar{\mu}) - \text{trdeg}_{\mathbb{C}(\bar{\mu})\langle \bar{y}^*, \bar{u}^* \rangle} \mathbb{C}(\bar{x}^*, \bar{\mu}) = A - B.$$

Hence, if both A and B have been computed correctly, the returned result is correct. In the probabilistic version, the probability of at least one of them being incorrect does not exceed

$$2 \left(1 - \frac{1+p}{2} \right) = 1 - p. \quad \blacksquare$$

Proposition 6.5. *If Algorithm 6.2 uses the second algorithm from Theorem 6.3, then Algorithm 6.1 is a probabilistic Monte Carlo algorithm of polynomial arithmetic complexity with respect to the complexity of Σ .*

Proof. First we will prove that the arithmetic complexity of Algorithm 6.2 is polynomial. The first and the last steps have polynomial complexity. The fact that the arithmetic complexity of the second step is polynomial follows from Theorem 6.3 and the fact that the complexities of Σ' and Σ'' are polynomial in the complexity of Σ .

Let ℓ be the number of parameters. Since $d_i \leq \ell$ for every $i \geq 0$ except for the last and $d_0 > d_1 > d_2 > \dots$, the counter i in Algorithm 6.1 will not exceed $\ell + 1$. Then there will be at most $\ell + 1$ runs of Algorithm 6.2, and each run will be on a system of complexity at most $\ell + 1$ times the complexity of Σ . Therefore, the total arithmetic complexity will be still polynomial in the complexity of Σ . ■

6.2. Algorithm: Implementation and examples. We implemented the probabilistic version of Algorithm 6.1 for computing the bound from Theorem 3.1 in Julia language using Oscar and Nemo libraries [4] together with a version of the algorithm by Sedoglavic from Theorem 6.3. The code and examples described below are available at <https://github.com/pogudinglab/ExperimentsBound>.

Below we will demonstrate the algorithm and the bound on several examples and compare with the algorithm presented in [17] (see Example 6.13 and Table 2). All of the runtimes reported below have been measured on a laptop with 1.6 GHz processor (Intel Core i5) and 16GB RAM. All of the computations reported below have been performed with the correctness probability of 99% (see the specification of Algorithm 6.1).

Remark 6.6. For some of the examples below, we were able to obtain the exact values of $\text{NumExpGlob}(\Sigma)$ using SIAN [9]. SIAN is software that can check single-experiment identifiability of any fixed function of parameters. We used it as follows:

1. If, for some r , all parameters of Σ_r are globally identifiable, then $\text{NumExpGlob}(\Sigma) \leq r$.
2. If, for some r , the parameter identifiability of Σ_r and Σ_{r+1} are not the same, then $\text{NumExpGlob}(\Sigma) \geq r + 1$.

Example 6.7 (the counterexample from section 4). In section 4, we have shown that $\text{NumExpGlob}(\Sigma) > 1$ for the following system Σ :

$$\begin{cases} x'_1 = 0, \\ x'_2 = x_1x_2 + \mu_1x_1 + \mu_2, \\ y = x_2. \end{cases}$$

Our implementation shows that $\text{NumExpLoc}(\Sigma) = 2$ and $\text{NumExpGlob}(\Sigma) \in \{2, 3\}$. The computation took 0.01 seconds. Using SIAN as described in Remark 6.6, we find that both parameters μ_1 and μ_2 are globally identifiable in Σ_2 . Combining it with $\text{NumExpGlob}(\Sigma) \in \{2, 3\}$ obtained by the algorithm, we conclude that

$$\text{NumExpLoc}(\Sigma) = \text{NumExpGlob}(\Sigma) = 2.$$

The same bound is given by [17, Theorem 21]. The computation took 0.3 seconds.

Example 6.8 (SEIR epidemiological model). Consider the following SEIR model [25, equation (2.2)]:

$$(6.1) \quad \begin{cases} S' = -\beta \frac{SI}{N}, \\ E' = \beta \frac{SI}{N} - \nu E, \\ I' = \nu E - \alpha I, \\ R' = \alpha I, \end{cases}$$

where S , E , I , R are the numbers of individuals susceptible to the infection, exposed, infected, and recovered, respectively, and $N := S + E + I + R$ is the total population which is known. Note that (6.1) implies that $N' = 0$. The output we will consider will be $\gamma I + \delta E$, where γ and δ are constants corresponding to factors such as, for instance, accuracy of the tests for the infection or the percentage of individuals going to a doctor after noticing the symptoms. We will assume that there are several experiments with the same values of $\alpha, \beta, \nu, \delta$ but varying values of γ (e.g., before and after improving the accuracy of the test).

To encode these assumptions into our framework, we will make γ a constant state variable and add an output for it. We will also replace the equation for R from (6.1) with $N' = 0$, as R does not appear in other equations other than inside N . This yields the following model Σ :

$$(6.2) \quad \begin{cases} S' = -\beta \frac{SI}{N}, \\ E' = \beta \frac{SI}{N} - \nu E, \\ I' = \nu E - \alpha I, \\ N' = \gamma' = 0, \\ y_1 = \gamma I + \delta E, \\ y_2 = \gamma, \ y_3 = N. \end{cases}$$

Our implementation shows that $\text{NumExpLoc}(\Sigma) = 1$ and $\text{NumExpGlob}(\Sigma) \in \{1, 2\}$. The computation took 0.05 seconds. Using SIAN as described in Remark 6.6, we find that all the parameters are only locally identifiable from a single experiment but become globally identifiable after 2 experiments. Therefore, $\text{NumExpGlob}(\Sigma) = 2$, so the bound given by the algorithm is exact in this case.

The program for computing a bound for the number of experiments provided in [17] did not finish on this example after 2 hours of computation.

Example 6.9 (linear compartment models with controlled rates). Linear compartment models typically represent a set of compartments in which material is transferred from some compartments to other compartments. It is also allowed to have a leakage of material from some compartments out of the system and input of material into some compartments from outside the system.

Linear compartment models are typically represented as directed graphs with edges labeled by scalar parameters (called rate constants). An example of such a representation is shown in Figure 1. The rules of transforming such a graph into a system of ODEs are the following:

- (compartments) each vertex of the graph corresponds to a state variable (a compartment);

- Downloaded 02/09/23 to 193.55.176.9 . Redistribution subject to SIAM license or copyright; see <https://epubs.siam.org/terms-privacy>



Downloaded 02/09/23 to 193.55.176.9 . Redistribution subject to SIAM license or copyright; see <https://epubs.siam.org/terms-privacy>

Downloaded 02/09/23 to 193.55.176.9 . Redistribution subject to SIAM license or copyright; see <https://epubs.siam.org/terms-privacy>



Downloaded 02/09/23 to 193.55.176.9 . Redistribution subject to SIAM license or copyright; see <https://epubs.siam.org/terms-privacy>

We will consider a modification of these models similar to [27, section III.B]. The modification is motivated by voltage clamp protocols used to identify parameters in ion channel models [5]. Ion channel models are often modeled using Markov models, which are similar to linear compartment models but with parameters depending on input functions (see [27, section III.B] and [22, section 3.2]). In the context of ion channel models, it may be nonrealistic to include a generic time-dependent input into the model. Instead of this, several experiments are performed such that the parameters depend on a constant input that takes different values for different experiments [5, sections 5 and 7]. Such a constant input can be encoded into our framework by adding a new state variable x_0 satisfying $x'_0 = 0$ and a new output equal to x_0 . We will consider the case in which all of the parameters depend linearly on the constant input x_0 ; that is, $a_{ij} = b_{ij} + c_{ij}x_0$ for all i and j , where b_{ij} and c_{ij} are new parameters. A dependence of this form was used for some of the parameters in [3, Supplementary Material] and can be viewed as a linear approximation to the dependencies used in [27, 5]. For example, the cycle model (Figure 2a) with $n = 4$ will be represented as shown in Figure 3 (cf. [27, III.B])




Figure 3. Cyclic model with $n = 4$ compartments with constant input in the reaction rates: graph (left) and ODE model (right)

We have analyzed models from families in Figure 2 with introduced constant input x_0 as described above using our implementation. The resulting values of the bound and the runtimes are summarized in Table 1. The algorithm for computing a bound for the number of experiments described in [17, Remark 22] did not finish on any of the models even for $n = 3$ after 2 hours of computation.

Table 1

Results and runtimes of our implementation on cyclic, catenary, and mammillary models (see Figure 2) with a constant input acting on reaction rates as in Figure 3.

*: for details on SIAN usage in this case, see Remark 6.10

Model	NumExpLoc(Σ)		NumExpGlob(Σ) $\in$		Runtime (sec.)		max n feasible for SIAN*
	$n = 3$	$4 \leq n \leq 15$	$n = 3$	$4 \leq n \leq 15$	$n = 10$	$n = 15$	
Cycle	3	3	{3, 4}	{3, 4}	9.5	41	4
Catenary	4	5	{4, 5}	{5, 6}	45.6	330	3
Mammillary	4	5	{4, 5}	{5, 6}	45.8	320	3

Remark 6.10. For $\text{NumExpGlob}(\Sigma)$, we tried to refine the result to obtain the exact value using SIAN [9] as described in Remark 6.6. The results are the following:

- For the cycle model, we have found that $\text{NumExpGlob}(\Sigma) \geq 4$ for $n = 3, 4$ as described in the second item of Remark 6.6. Combined with the bound given by our implementation, we obtain $\text{NumExpGlob}(\Sigma) = 4$ for $n = 3, 4$, so the bound is exact in this case. Already for $n = 5$, the computation with SIAN did not finish in 10 hours on a server and used more than 20GB of memory.
- For the catenary and mammillary models, a computation with SIAN showed that, for $n = 3$, none of the individual parameters was identifiable after 5 experiments. Therefore, we cannot use SIAN to determine the exact bound in the way we did it for the cycle model. For $n = 4$, the computation with SIAN did not finish in 10 hours on a server and used more than 40GB of memory.

Remark 6.11. For all three series of models, the output of the algorithm stabilizes from $n = 4$. It is natural to conjecture that the result will be the same for all larger values of n . It would be interesting to have a mathematical argument showing this or maybe even a formula for the number of experiments in terms of numerical characteristics of the graph of a model.

Remark 6.12. The same procedure of linearly perturbing the rate constants can be applied to general chemical reaction networks, which yield, in general, highly nonlinear ODEs. In this setup, we also observe that the necessary number of experiments may become larger than 1: for example, for the perturbed version of the phosphorylation model [10, Example 6.1], we get $\text{NumExpLoc}(\Sigma) = 2$ and $\text{NumExpGlob}(\Sigma) \in \{2, 3\}$.

Example 6.13 (examples from [17]). As we mentioned, the algorithm from [17, Remark 22] does not produce any bound for Examples 6.8 and 6.9 in reasonable time. For the sake of comparison, we run our algorithm on the examples collected in [17, section 5]. The comparison is reported in Table 2, in which we also included the above examples for completeness.

Table 2

Comparison of bounds for NumExpGlob and runtimes with [17].

*: obtained by a modification of the method, see [17, section 5.3]; [17, Theorem 21] gives 35

**: our algorithm from the present paper can tackle larger n as well; see Table 1

Model	Approach from [17]		Our algorithm	
	Time (sec.)	Bound	Time (sec.)	Bound
Lotka–Volterra w/control [17, section 5.1]	0.3	1	0.005	2
Slow-fast ambiguity [17, section 5.2]	0.37	2	0.024	2
Lotka–Volterra w/mixture [17, section 5.3]	15	4*	0.01	2
SEIR - prevalence [17, section 5.4]	1	1	0.021	2
SEIR - incidence [17, section 5.4]	340	1	0.032	2
Counterexample from section 4 (Example 6.7)	0.3	2	0.01	3
SEIR w/ mixture (Example 6.8)	> 2 h.	N/A	0.05	2
Cycle for $n = 3^{**}$ (Example 6.9)	> 2 h.	N/A	0.3	4
Catenary for $n = 3^{**}$ (Example 6.9)	> 2 h.	N/A	0.6	5
Mammillary for $n = 3^{**}$ (Example 6.9)	> 2 h.	N/A	0.6	5

7. Model theory and identifiability. The goal of this section is to explain the connections between identifiability and model theory and give an idea how the algebraic arguments from the preceding sections have been inspired and informed by model-theoretic considerations. The section is structured as follows. Section 7.1 introduces some fundamental notions of model theory in the context of differential fields and explains their close relations with the concept of identifiability; the section culminates in the identifiability–model theory dictionary in Table 3. Section 7.2 is about some model theory of differential fields behind (4.1) and Proposition 3.3. Specifically we describe canonical bases of types over constant fields. In section 7.3, we give model-theoretic proofs of the key ingredients of the proof of Theorem 3.1 which are Propositions 5.4 and 5.10. Some of the ideas from these proofs were crucial in the algebraic proof of Theorem 3.1.

Model theory is a kind of abstract algebra, which gives a common environment and common tools for studying algebraic structures such as group, fields, and fields equipped with derivations or automorphisms. Among these tools are *canonical bases* coming from an area of model theory called stability theory and which implicitly play an important role in this paper.

7.1. Setup. We will use basic notions from model theory (such as language, structure, theory, and model). The reader is referred to [15, Chapter 1] for additional details. In this section, we will introduce relevant notions from model theory. We will specialize some of them for simplicity to the case of differential fields and explain their relation to the identifiability problem. Further details can be found in [14] (also [15, 20]). The correspondence between notions from identifiability and model theory is summarized in Table 3 at the end of this section.

One expresses system (2.1) by a formula in the appropriate language (i.e., the conjunction of the system of finitely many equations) as defined below.

Definition 7.1 (extensions of languages). For a language $\mathcal{L}$, a structure M in $\mathcal{L}$, and a subset $A \subseteq M$, let $\mathcal{L}_A$ denote the extension of language $\mathcal{L}$ by adding a constant for each element of A .

The reader should be careful to distinguish “constants” in the sense of constant symbols in logic from constants in the sense of elements of a differential field on which the derivation is zero.

Example 7.2 (language of differential fields). We work in the language of differential fields

$$\mathcal{L}_{DF} := \{+, \cdot, ', 0, 1\}.$$

This language allows one to express differential equations with rational coefficients (not arbitrary complex numbers as in (2.1)), and this is not sufficient to write a system of the form (2.1). However, every equation in (2.1) is a formula in $\mathcal{L}_{DF, \mathbb{C}}$ in variables $\bar{x}, \bar{y}, \bar{u}, \bar{\mu}$.

We will work not in the theory of differential fields but in the theory of differentially closed fields. This ensures that the equations of interest have sufficiently many solutions.

Definition 7.3 (differentially closed fields [15, Definition 4.3.29]). A differential field K is called *differentially closed* if, for all differential polynomials $f, g \in K\{x\} \setminus \{0\}$ with $\text{ord } f > \text{ord } g$ there is $a \in K$ such that $f(a) = 0$ and $g(a) \neq 0$ (for $p \in K \setminus \{0\}$, we define $\text{ord } p := -1$).

These fields share many properties with algebraically closed fields such as the Nullstellensatz: if a system of equations over K has a solution in some extension of K , then it has a solution in K as well [14, Corollary 2.6]. The property of being differentially closed can be written as a list of axioms in $\mathcal{L}_{DF}$. We denote the resulting theory of differentially closed fields by DCF_0 .

Once we have put the system (2.1) into the model-theoretic context, we would like to be able to talk about its solutions and generic solutions. This is done using the language of formulas and types.

Definition 7.4 (types [15, Definition 4.1.1]). Let M be an $\mathcal{L}$ -structure and A a subset of M . Then an n -type over A , relative to the structure M , is a set Φ of formulas in $\mathcal{L}_A$ with free variables $x_1, \dots, x_n$ such that there exists an $\mathcal{L}_A$ -structure N containing M (could be equal to M) such that the following hold:

- all $\mathcal{L}_M$ -sentences true in M are also true in N (such an N is called an *elementary extension*, and we write $M \prec N$);
- there exist $a_1, \dots, a_n \in N$ satisfying all the formulas in Φ .

Such a set or tuple $a_1, \dots, a_n$ is called a *realization of the type*.

Example 7.5 (important classes of types). In this paper, we will encounter mostly types of the following forms:

- Types defined by finitely many formulas (that is, $|\Phi| < \infty$ in Definition 7.4). For example, the system (2.1) (or any other system of differential-algebraic equations) defines such a type over any differential field containing at least one solution of (2.1). Using conjunction, every such type can be defined by a single formula.
- Let M be a model, $A \subset M$ be any subset, and $\bar{a}$ be a tuple from M . Then $\text{tp}_M(\bar{a}/A)$ denotes the set of all formulas in $\mathcal{L}_A$ satisfied by $\bar{a}$ in M . Note that if $M \prec N$, then $\text{tp}_N(\bar{a}/A) = \text{tp}_M(\bar{a}/A)$.
- Let M be an $\mathcal{L}$ -structure, A a subset of M , and p an n -type over A relative to the structure M . Then p is complete if, for every $\phi(\bar{x})$ in $\mathcal{L}_A$, either ϕ or $\neg\phi$ is in p . The complete n -types over A relative to M are precisely of the form $\text{tp}_N(\bar{a}/A)$ for N an elementary extension of M .

For any automorphism f of M , we define a map on the set of complete types over M by applying f to the formulas contained in the types.

Remark 7.6 (types in differentially closed field). The theory DCF_0 admits quantifier elimination [15, Theorem 4.3.32], that is, for every formula ϕ , there is a quantifier-free formula equivalent to ϕ in DCF_0 . Therefore, every type can be defined by a set of quantifier-free formulas.

In particular, if K and L are differentially closed fields and $A, \bar{a} \subset K \cap L$, then the types $\text{tp}_K(\bar{a}/A)$ and $\text{tp}_L(\bar{a}/A)$ are the same. Hence, working in the context of differentially closed fields, we will write simply $\text{tp}(\bar{a}/A)$ without specifying the ambient differentially closed field.

Example 7.7. Consider a generic solution $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ of (2.1) (see Definition 2.3) in a differentially closed field $K \supset \mathbb{C}(\bar{\mu})$. Then we will call $\text{tp}((\bar{x}^*, \bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu}))$ the *type of a generic solution of (2.1)*. This type contains all equations (2.1), but also, for example, any inequation (say, $x'_1 \neq 0$) that is true for at least one solution of (2.1) and thus must be true

for a generic one.

Model theory provides tools to construct large enough differential fields containing realizations of types of generic solutions of all the systems of interest (and in many copies so that we can talk about multiple experiments as well).

Definition 7.8 (saturation [15, Definition 4.3.1]). Let κ be an infinite cardinal. A model M of theory T is called κ -saturated if every complete type Φ such that

$$|\{m \in M \mid m \text{ appears in } \Phi\}| < \kappa$$

has a realization in M . M is called *saturated* if it is $|M|$ -saturated.

Remark 7.9. If M is a model of theory T and $A \subset M$ is a subset and M is saturated with $|M| > \max(|A|, |T|)$, then, for every $\bar{a}, \bar{b}$ in M ,

$\text{tp}_M(\bar{a}/A) = \text{tp}_M(\bar{b}/A) \iff \exists$ automorphism $\alpha: M \rightarrow M$ such that $\alpha(\bar{a}) = \bar{b}$ and $\alpha|_A = \text{id}$ (see [15, Propositions 4.2.13 and 4.3.3]).

Now we define identifiability in the language of model theory.

Definition 7.10 (definability [15, Definition 1.3.1]). A subset $X \subset M^n$ of a structure M in a language $\mathcal{L}$ is called *definable over* a subset $A \subset M$ if there exists a first-order formula $\phi(x_1, \dots, x_n)$ in $\mathcal{L}_A$ such that

$$(a_1, \dots, a_n) \in X \iff \phi(a_1, \dots, a_n) \text{ is true in } M.$$

Example 7.11. Let K be a differential field over a differential subfield k_0 , and let $\bar{a}$ and $\bar{b}$ be tuples of elements of K . It follows from [15, Proposition 1.3.5] and [11, Theorem 2.6] that

$$\bar{a} \text{ definable over } \bar{b} \text{ in } \mathcal{L}_{DF, k_0} \iff \bar{a} \in k_0\langle \bar{b} \rangle$$

(where, for a set A , $\bar{a} \in A$ means $a_i \in A$ for each $i \leq \text{length}(\bar{a})$). Comparing this with Definition 2.5, we see that $h(\bar{\mu}) \in \mathbb{C}(\bar{\mu})$ is *identifiable* if and only if it is $\mathcal{L}_{DF, \mathbb{C}}$ -definable over $(\bar{y}^*, \bar{u}^*)$ for every generic solution $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ of (2.1).

Remark 7.12. For a saturated model M , definability can be restated in terms of automorphisms [15, Proposition 4.3.25]: for $\bar{a}, \bar{b} \in M$, $\bar{a}$ is definable over $\bar{b}$ if and only if

$$\forall \text{ automorphism } \alpha: M \rightarrow M \quad \alpha(\bar{b}) = \bar{b} \implies \alpha(\bar{a}) = \bar{a}.$$

Informally, this can be stated as *if $\bar{b}$ is fixed, then $\bar{a}$ is also fixed*. Syntactically, this is very similar to the analytic definition of identifiability [10, Definition 2.5]. This partially explains why model-theoretic tools were used in proving the equivalence [10, Proposition 3.4] of the analytic definition and Definition 2.5.

In order to define multi-experiment identifiability in model-theoretic terms, we will define the notion of independence.

Definition 7.13 (stationarity, nonforking, and independence). Let k be a differential subfield of a differentially closed field K , and let n be a positive integer. Let $\bar{a}$ be an n -tuple of elements from K , and let $\bar{x}$ denote an n -tuple of differential variables.

- Recall from Notation 5.1 that the vanishing ideal $\mathcal{I}_k(\bar{a})$ of $\bar{a}$ over k is $\{P \in k\{\bar{x}\} : P(\bar{a}) = 0\}$. Note that $\mathcal{I}_k(\bar{a})$ depends only on $p := \text{tp}(\bar{a}/k)$. Moreover, by quantifier elimination of DCF_0 (see Remark 7.6), it also determines $\text{tp}(\bar{a}/k)$.
- Let L be a differential field with $k \subset L \subset K$. We say that $\bar{a}$ is *independent from L over k* if $\mathcal{I}_L(\bar{a})$ is a prime component of $\mathcal{I}_k(\bar{a}) \otimes_k L$. We also express this by saying that $\text{tp}(\bar{a}/L)$ *does not fork over k* , or that $\text{tp}(\bar{a}/L)$ is a *nonforking extension* of $\text{tp}(\bar{a}/k)$.
- We say that $\text{tp}(\bar{a}/k)$ is *stationary* if $\mathcal{I}_k(\bar{a})$ is “absolutely prime,” namely that for each $L \supset k$, $\mathcal{I}_k(\bar{a}) \otimes_k L$ is prime. It is enough to require this for $L = K$.
- If $\bar{b}$ is another finite tuple from K , we say that $\bar{a}$ and $\bar{b}$ are *independent over k* if $\bar{b}$ is independent from $k\langle\bar{a}\rangle$ over k . A sequence of tuples $\bar{a}_1, \bar{a}_2, \dots$ is called *independent over k* if, for every $i \geq 1$, $\bar{a}_{i+1}$ is independent from $k\langle\bar{a}_1, \dots, \bar{a}_i\rangle$ over k .
- In general, given subsets $A \subseteq B$ of K , we say that $\bar{a}$ is *independent from B over A* , or $\text{tp}(\bar{a}/B)$ *does not fork over A* , if $\bar{a}$ is independent from L_2 over L_1 , where L_1 and L_2 are the differential fields generated by A and B , respectively.

Example 7.14 (generic solution of (2.1) is stationary). Let $\bar{a} := (\bar{x}^*, \bar{y}^*, \bar{u}^*)$ be a generic solution of (2.1) (see Definition 2.3). Then, by the definition, $\mathcal{I}_{\mathbb{C}(\bar{\mu})}(\bar{a}) = I_\Sigma$. It follows from [10, Proof of Lemma 3.2] that the ideal I_Σ is prime and remains prime under any field extension. Therefore, $\text{tp}(\bar{a}/\mathbb{C}(\bar{\mu}))$ is stationary.

Remark 7.15 (some properties of independence and forking). In this remark, we use the notation from Definition 7.13.

- (1) One can show that finite tuples $\bar{a}, \bar{b}$ are independent over k if and only if the ideal $\mathcal{I}_k(\bar{a}, \bar{b}) \subseteq k\{\bar{x}, \bar{y}\}$ is a prime component of the ideal I of $k\{\bar{x}, \bar{y}\}$ generated by $\mathcal{I}_k(\bar{a})$ and $\mathcal{I}_k(\bar{b})$. Indeed, $\bar{a}, \bar{b}$ are independent over k if and only if $\mathcal{I}_{k\langle\bar{b}\rangle}(\bar{a})$ is a prime component of $\mathcal{I}_k(\bar{a}) \otimes_k k\langle\bar{b}\rangle$. Let $\varphi : k\{\bar{x}, \bar{y}\} \rightarrow k\langle\bar{b}\rangle\{\bar{x}\}$ be the canonical differential ring homomorphism such that $\varphi(\bar{y}) = \bar{b}$. We have $\ker \varphi = \mathcal{I}_k(\bar{b})$, $\varphi(\mathcal{I}_k(\bar{a}, \bar{b})) \otimes_k k\langle\bar{b}\rangle = \mathcal{I}_{k\langle\bar{b}\rangle}(\bar{a})$, and $\varphi(I) \otimes_k k\langle\bar{b}\rangle = \mathcal{I}_k(\bar{a}) \otimes_k k\langle\bar{b}\rangle$. Hence, since both $\mathcal{I}_k(\bar{a}, \bar{b})$ and I contain $\ker \varphi$, we obtain that $\mathcal{I}_k(\bar{a}, \bar{b})$ is a prime component of I if and only if $\mathcal{I}_{k\langle\bar{b}\rangle}(\bar{a})$ is a prime component of $\mathcal{I}_k(\bar{a}) \otimes_k k\langle\bar{b}\rangle$, which finishes the argument. Moreover, if both $\text{tp}(\bar{a}/k)$ and $\text{tp}(\bar{b}/k)$ are stationary, I is itself prime by the definition of stationarity.
- (2) Since the ideal $\mathcal{I}_k(\bar{a})$ is prime, the ideal $\mathcal{I}_k(\bar{a}) \otimes_k L$ is equidimensional (the transcendence bases corresponding to its prime components are all the same) for every $L \supset k$. Hence, $\mathcal{I}_L(\bar{a})$ is a prime component of $\mathcal{I}_k(\bar{a}) \otimes_k L$ if and only if the transcendence basis corresponding to $\mathcal{I}_L(\bar{a})$ over L is the same as that of $\mathcal{I}_k(\bar{a})$ over k . Therefore, $\bar{a}$ is independent from $L \supset k$ over k if and only if, for every m , we have

$$\text{trdeg}_k k(\bar{a}, \bar{a}', \dots, \bar{a}^{(m)}) = \text{trdeg}_L L(\bar{a}, \bar{a}', \dots, \bar{a}^{(m)}).$$

Together with [21, Proposition 1.16], this implies that the definition of independence in DCF_0 we gave agrees with the general model-theoretic one (as, e.g., in [20, section 2.2, page 28]).

- (3) The definition of stationarity implies that, for every stationary type p over k and every differential field $L \supset k$, there is a unique complete type q that extends p and that does not fork over k . Such a type q will be referred to as *the* nonforking extension of p . Note that the

type q is again stationary. One can show that the converse (the uniqueness of a nonforking extension of p implies the stationarity of p) is also true by using the characterization of independence from the previous item and the fact that, after the extension of scalars, an irreducible variety becomes equidimensional. Indeed, let $p = \text{tp}(\bar{a}/k)$ not be stationary, and so let $L \supset k$ be such that the ideal $\mathcal{I}_k(\bar{a}) \otimes_k L$ is not prime. Let $P_1 = \mathcal{I}_L(\bar{b})$ and $P_2 = \mathcal{I}_L(\bar{c})$ be any two of its prime components. Note that P_1 and P_2 have the same dimension since $\mathcal{I}_k(\bar{a}) \otimes_k L$ is equidimensional. Since P_1 and P_2 are prime components of $P_1 \cap k\{\bar{x}\} \otimes_k L$ and $P_2 \cap k\{\bar{x}\} \otimes_k L$, respectively, the tuples $\bar{b}$ and $\bar{c}$ are each independent from L over k . Thus, $\text{tp}(\bar{b}/L)$ and $\text{tp}(\bar{c}/L)$ are two different extensions of $\text{tp}(\bar{a}/k)$ that do not fork over k .

Example 7.16. Consider the differential field $k = \mathbb{Q}(t)$ with respect to the derivation $\frac{d}{dt}$ and a saturated model $K \supset k$ of DCF_0 . Every formula in the type $p := \text{tp}(t/\mathbb{Q})$ is implied by the single formula $x' = 1$. Then type $q := \text{tp}(t/\mathbb{Q}(t))$ is an extension of p , and it contains a new formula $x = t$, which is not implied by $x' = 1$. We have

$$\text{trdeg}_{\mathbb{Q}} t = 1 \neq 0 = \text{trdeg}_{\mathbb{Q}(t)} t,$$

so the extension of p by q is forking.

Also, from the differential equations theory, we know that the general solution of $x' = 1$ is of the form $x = t + c$, where c is a constant. So we can construct a nonforking extension of p to $\mathbb{Q}(t)$ as $\text{tp}((t+c)/\mathbb{Q}(t))$, where $c \in K$ is a transcendental constant (exists because K is saturated).

Example 7.17 (multi-experiment identifiability via independence). Let $\bar{a} = (\bar{x}_1^*, \bar{y}_1^*, \bar{u}_1^*)$ and $\bar{b} = (\bar{x}_2^*, \bar{y}_2^*, \bar{u}_2^*)$, where $(\bar{x}_1^*, \bar{x}_2^*, \bar{y}_1^*, \bar{y}_2^*, \bar{u}_1^*, \bar{u}_2^*)$ is a generic solution of Σ_2 (see Definition 2.8). By the definition of I_{Σ_2} , it is generated by two copies of I_{Σ} , so $\bar{a}$ and $\bar{b}$ are independent over $\mathbb{C}(\bar{\mu})$. Moreover, $\bar{a}$ and $\bar{b}$ are *independent realizations* of the type of a generic solution of (2.1) (see Example 7.7). Combining this with Example 7.11, we have that $h(\bar{\mu}) \in \mathbb{C}(\bar{\mu})$ is ME-identifiable if and only if it is definable in $\mathcal{L}_{\text{DFC}}$ over some finite number of independent realizations of $\text{tp}((\bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu}))$, where $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ is a generic solution of (2.1).

Finally, it has been shown in [17, Theorem 19] that the field of ME-identifiable functions coincides with the field of definition of the ideal of input-output relations. Any set of generators of the field of definition is called a canonical base in model theory.

Definition 7.18 (canonical base [15, Definition 8.2.2]). Let M be a saturated model of the theory DCF_0 (that is, a large enough differentially closed field; see Definition 7.8), and let p be a complete type over M . Then a set $A \subset M$ is called a *canonical base* of p if and only if

$$\forall \text{ automorphism } \alpha: M \rightarrow M \quad \alpha(p) = p \iff \alpha|_A = \text{id},$$

where the automorphism acts on the type by acting on the formulas defining the type (which are defined over M ; see also Example 7.5). In particular, an automorphism fixes a complete type if it leaves the corresponding set of formulas invariant.

Every canonical base of a complete type p generates the same differential field over k [15, Lemma 8.2.4]. This field will be denoted by $\text{Cb}(p)$ and referred to as *the canonical base* (see [20, page 29]). If k is a differential subfield of a differentially closed K and $\bar{a}$ is a tuple from

K such that $\text{tp}(\bar{a}/k)$ is stationary, then $\text{Cb}(\bar{a}/k)$ denotes the canonical base of the nonforking extension of $\text{tp}(\bar{a}/k)$ to K (see Definition 7.13).

Example 7.19. In the theory of differential fields, the canonical base of stationary $\text{tp}(\bar{a}/k)$ is the field of definition of $\mathcal{I}_k(\bar{a})$ (see Notation 5.2). Therefore, [17, Theorem 19] can be rephrased as follows: *the field of ME-identifiable functions is $\text{Cb}((\bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu}))$, where $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ is any generic solution of (2.1).* This fact sounds natural if one looks at Definition 7.18: the canonical base is fixed if and only if the set of all experimental outcomes for fixed generic parameters is invariant.

We will conclude this subsection by summarizing some properties of forking extensions which will be used in the subsequent proofs.

Remark 7.20 (properties of forking). We fix a differentially closed field $K \supset k_0$, its subsets $A \subseteq B \subseteq C$, and a tuple $\bar{a}$ from K .

- (1) (*transitivity* [20, Proposition 2.20(iii)]) $\text{tp}(\bar{a}/C)$ does not fork over A if and only if it does not fork over B and $\text{tp}(\bar{a}/B)$ does not fork over A .
- (2) (*symmetry* [20, Proposition 2.20(v)]) $\text{tp}(\bar{a}/B)$ does not fork over A if and only if, for every $\bar{b}$ from B , $\text{tp}(\bar{b}/A \cup \bar{a})$ does not fork over A .
- (3) ([20, Proposition 2.20(iv)]) Assume that two distinct types $\text{tp}(\bar{a}/B)$ and $\text{tp}(\bar{b}/B)$ do not fork over A . Assume also that $\text{tp}(\bar{a}/A) = \text{tp}(\bar{b}/A)$. Then there exists an equivalence relation $E(x_1, x_2)$ defined over A with finitely many classes such that, for every $\bar{a}^*$ satisfying $\text{tp}(\bar{a}/B)$ and $\bar{b}^*$ satisfying $\text{tp}(\bar{b}/B)$, we have $\neg E(\bar{a}^*, \bar{b}^*)$. In geometric terms, one can think of E being the relation “belonging to the same component of the variety defined by $\text{tp}(\bar{a}/A)$.”
- (4) (*forking and canonical bases*) Let p be a stationary type over A , and let F be the differential field generated by A . Then $\text{Cb}(p) \subseteq F$ [20, Remark 2.26(i)] and coincides with $\text{Cb}(q)$ whenever q is the nonforking extension of p to a larger set $B \supseteq A$ (this follows from the sentence immediately before [20, Remark 2.26]: if q is a nonforking extension of p , then p and q have a common nonforking extension, namely, q , so that $\text{Cb}(p) = \text{Cb}(q)$). Also $\text{tp}(\bar{a}/B)$, which is a type over B , does not fork over $A \subseteq B$ if and only if $\text{Cb}(\text{tp}(\bar{a}/B))$ is contained in the algebraic closure of F (by [20, Remark 2.26(ii)]).

Table 3
Identifiability: Model theory dictionary.

Identifiability	Model theory
Solution of (2.1)	Realization of the formula (2.1)
Generic solution $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ of (2.1)	Realization of the generic type defined by (2.1) (Example 7.7)
Identifiability (Definition 2.5)	Definability over $(\bar{y}^*, \bar{u}^*)$ (Example 7.11)
Multi-experiment identifiability (Definition 2.8)	Definability over some finite number of independent realizations of $\text{tp}((\bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu}))$ (Example 7.17)
Field of ME-identifiable functions	The canonical base $\text{Cb}((\bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu}))$ (Example 7.19 and Propositions 5.4 and 7.25)

7.2. Single-output model requiring more than one experiment revisited. In this section, we will discuss a model-theoretic construction used to find (4.1). We will work over the field $\mathbb{C}$, that is, in the language $\mathcal{L}_{DF, \mathbb{C}}$. Consider a constant differential field $k = \mathbb{C}(\bar{\mu})$. Let K be a saturated differentially closed field containing k . Let $\mathcal{C}$ denote the constants of K . We will use two technical lemmas.

Lemma 7.21. *For every tuple $\bar{a}$ from K , $\text{Cb}(\bar{a}/\mathcal{C}) = \mathcal{C} \cap \mathbb{C}\langle\bar{a}\rangle$.*

Proof. First we observe that, since $\mathcal{C}$ is algebraically closed [14, Lemma 2.1], the type $\text{tp}(\bar{a}/\mathcal{C})$ is stationary due to [20, Chapter 1, Remark 2.25(i)], so we can use Definition 7.18. Consider any automorphism α of K such that $\alpha(\bar{a}) = \bar{a}$. Since $\mathcal{C}$ is the field of constants of K , we have $\alpha(\mathcal{C}) = \mathcal{C}$. Then α fixes $I_{\mathcal{C}}(\bar{a})$ setwise, so it fixes the nonforking extension of $\text{tp}(\bar{a}/\mathcal{C})$ to K (see Definition 7.13). By Definition 7.18, we conclude that α fixes $\text{Cb}(\bar{a}/\mathcal{C})$. Thus, Remark 7.12 implies that $\text{Cb}(\bar{a}/\mathcal{C}) \subset \mathbb{C}\langle\bar{a}\rangle$.

In the other direction, consider $b \in \mathbb{C}\langle\bar{a}\rangle \cap \mathcal{C}$. There exists a differential rational function f over $\mathbb{C}$ such that $b = f(\bar{a})$. Therefore, the formula $b = f(\bar{x})$ belongs to $\text{tp}(\bar{a}/\mathcal{C})$, so it belongs to its nonforking extension p to K . Then any automorphism α of K fixing p fixes $\bar{b}$. Then Definition 7.18 implies that $b \in \text{Cb}(\bar{a}/\mathcal{C})$. ■

Lemma 7.22. *Let $\bar{a}$ be a tuple from K and $\bar{c}$ be any set of generators of $\text{Cb}(\bar{a}/\mathcal{C})$ as a field over $\mathbb{C}$. Then the following hold:*

1. $\text{Cb}(\bar{a}/k) = \text{Cb}(\bar{c}/k)$;
2. $\mathbb{C}\langle\bar{a}\rangle \cap k = \mathbb{C}\langle\bar{c}\rangle \cap k$.

Proof.

1. Lemma 7.21 implies that $\bar{c}$ is a tuple from $\mathbb{C}\langle\bar{a}\rangle$. Therefore, using Remark 7.12, we obtain $\text{Cb}(\bar{c}/k) \subset \text{Cb}(\bar{a}/k)$.

Consider an automorphism α of K that fixes the nonforking extension p of $\text{tp}(\bar{c}/k)$ to K . Then α fixes $K \otimes_k \mathcal{I}_k(\bar{c})$ setwise, so $\text{tp}(\bar{c}/k) = \text{tp}(\alpha(\bar{c})/k)$. Then Remark 7.9 implies that there exists an automorphism β of K that fixes k and $\beta(\alpha(\bar{c})) = \bar{c}$. Then $\text{tp}(\bar{a}/\mathcal{C}) = \text{tp}(\beta(\alpha(\bar{a}))/\mathcal{C})$, so $\text{tp}(\bar{a}/k) = \text{tp}(\beta(\alpha(\bar{a}))/k)$. Since β^{-1} fixes k , we have

$$\text{tp}(\bar{a}/k) = \text{tp}(\beta(\alpha(\bar{a}))/k) = \text{tp}(\beta^{-1}(\beta(\alpha(\bar{a}))))/k = \text{tp}(\alpha(\bar{a})/k).$$

Therefore, $\mathcal{I}_k(\bar{a}) = \mathcal{I}_k(\alpha(\bar{a}))$, so these types have the same nonforking extensions to K . Hence, α fixes $\text{Cb}(\bar{a}/k)$. Thus, $\text{Cb}(\bar{a}/k) \subset \text{Cb}(\bar{c}/k)$.

2. Using Lemma 7.21, since k is constant, we have

$$\mathbb{C}\langle\bar{a}\rangle \cap k = (\mathbb{C}\langle\bar{a}\rangle \cap \mathcal{C}) \cap k = \mathbb{C}\langle\bar{c}\rangle \cap k. \quad \blacksquare$$

Let $(\bar{x}^*, \bar{y}^*, \bar{u}^*)$ be a generic solution of an algebraic differential model Σ as in (2.1). Then the desired nonequality of the fields of SE- and ME-identifiable functions can be restated, using Examples 7.11 and 7.19, as

$$(7.1) \quad \mathbb{C}\langle\bar{y}^*, \bar{u}^*\rangle \cap \mathbb{C}(\bar{\mu}) \neq \text{Cb}((\bar{y}^*, \bar{u}^*)/\mathbb{C}(\bar{\mu})).$$

We will first construct an example having constant dynamics and satisfying the nonequality (7.1) (as, for example, in [10, Example 2.14]). We then use Lemma 7.22 to “pack” two output variables of the example into a single output variable while preserving (7.1):

(Step 1) Let $\bar{\mu} = (\mu_1, \mu_2)$, and introduce a constant state variable x_1 . We introduce two auxiliary outputs $z_1 = x_1$ and $z_2 = \mu_1 x_1 + \mu_2$. The defining differential ideal of (z_1, z_2) is generated by $z_1', z_2 - \mu_1 z_1 - \mu_2$, so $\text{Cb}((z_1, z_2)/\mathbb{C}(\bar{\mu})) = \mathbb{C}(\bar{\mu})$. On the other hand, since the automorphism of $\mathbb{C}(x_1, \mu_1, \mu_2)$ defined by

$$x_1 \rightarrow x_1, \quad \mu_1 \rightarrow \mu_1 + 1, \quad \mu_2 \rightarrow \mu_2 - x_1$$

fixes z_1 and z_2 but does not fix μ_1 or μ_2 , we conclude that $\mu_1, \mu_2 \notin \mathbb{C}(z_1, z_2)$.

(Step 2) Now we introduce a new state variable x_2 satisfying $x_2' = z_1 x_2 + z_2$ and set the output $y = x_2$. Then we have $\text{Cb}(y/\mathbb{C}) = \mathbb{C}(z_1, z_2)$. Therefore, using Lemma 7.22,

$$\text{Cb}(y/\mathbb{C}(\bar{\mu})) = \text{Cb}((z_1, z_2)/\mathbb{C}(\bar{\mu})) = \mathbb{C}(\bar{\mu}) \neq \mathbb{C}(z_1, z_2) \cap \mathbb{C}(\bar{\mu}) = \mathbb{C}(y) \cap \mathbb{C}(\bar{\mu}).$$

So we get exactly (4.1).

Remark 7.23. Instead of $x_2' = z_1 x_2 + z_2$, we could take any other equation containing z_1 and z_2 among the coefficients, for example, $x_2' = z_1 x_2^2 + 2z_2 x_2 - 3z_1^3$. This would yield an example with the same property (7.1).

Remark 7.24. Another way to obtain an example with the property from Proposition 3.3 is to remove the first output (the equation $y_1 = x_0$) from any of the models in Example 6.9, which can be directly verified using our implementation at <https://github.com/pogudingleb/ExperimentsBound>.

7.3. Model theory way of proving Theorem 3.1. In this section, we will use Notations 5.1 and 5.2. In particular, we work *over a fixed ground differential field* k_0 (that is, in $\mathcal{L}_{DF, k_0}$). Therefore, all fields are assumed to be generated over k_0 and, in particular, whenever we write $\text{tp}(\bar{a}/A)$, this is equivalent to $\text{tp}(\bar{a}/k_0\langle A \rangle)$.

An interested reader could deduce the following proposition from [20, Chapter 1, Lemma 3.19] (see also [15, Exercise 8.4.12]).

Proposition 7.25 (model-theoretic reformulation of Proposition 5.4). *Let $k \subset K$ be a differential subfield, p a stationary type over k , and a sequence $\bar{a}_1, \bar{a}_2, \dots$ of independent realizations of p in K . Then there exists N such that*

$$\text{Cb}(p) \subset k_0\langle \bar{a}_1, \dots, \bar{a}_N \rangle.$$

Lemma 7.26 (cf. Lemma 5.3). *Let $k_0 \subset k \subset K$ be a differential field. Consider tuples $\bar{a}, \bar{b}$ from K such that $\text{tp}(\bar{a}/k)$ is stationary and $\bar{b}$ is independent from $\bar{a}$ over k . Then $\text{Cb}(\bar{a}/k) = \text{Cb}(\bar{a}/k\langle \bar{b} \rangle)$.*

Proof. Let $p_1 := \text{tp}(\bar{a}/k)$ and $p_2 := \text{tp}(\bar{a}/k\langle \bar{b} \rangle)$. Consider a subfield $F \subset k$. Then the following hold:

- the independence of $\bar{a}$ and $\bar{b}$ together with Remark 7.20(2) implies that p_1 does not fork over F if and only if p_2 does not fork over F ;
- since p_1 is the restriction of p_2 to k , p_1 restricted to F is stationary if and only if p_2 restricted to F is stationary.

Applying Remark 7.20(4) twice, with $A = \text{Cb}(p_1)$ and $A = \text{Cb}(p_2)$, we conclude that $\text{Cb}(p_1) = \text{Cb}(p_2)$. ■

Proposition 7.27 (model-theoretic version of Proposition 5.10). *Let there be*

- k with $k_0 \subset k \subset K$ a differential field of finite transcendence degree over k_0 ,
- p a stationary type over k (in particular, p is a complete type),
- a sequence $\bar{a}_1, \bar{a}_2, \dots$ of independent realizations of p in K , and
- r the smallest integer such that

$$\text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle} k = \text{trdeg}_{k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle} k.$$

Then the following hold:

- (1) r is the smallest integer such that $\text{Cb}(p)$ is algebraic over $k_0\langle\bar{a}_1, \dots, \bar{a}_r\rangle$;
- (2) $\text{Cb}(p) \subset k_0\langle\bar{a}_1, \dots, \bar{a}_{r+1}\rangle$.

Proof. Let $\bar{c}$ be any finite tuple of generators of $\text{Cb}(\bar{a}_1/k)$. For every nonnegative integer ℓ , we set $\bar{A}_\ell := (\bar{a}_1, \dots, \bar{a}_\ell)$. We consider the following sequence of types (cf. the varieties X_i in the proof of Proposition 5.10):

$$\text{tp}(\bar{c}/\bar{A}_0) \subset \text{tp}(\bar{c}/\bar{A}_1) \subset \text{tp}(\bar{c}/\bar{A}_2) \subset \dots$$

Claim. $\bar{c}$ is algebraic over $k_0\langle\bar{A}_s\rangle$ if and only if $\text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{c} = \text{trdeg}_{k_0\langle\bar{A}_{s+1}\rangle} \bar{c}$. If the equality of the transcendence degrees does not hold, $\text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{c} > 0$, so $\bar{c}$ is not algebraic over this field.

We will now show the reverse implication. The equality of transcendence degrees implies that $\text{tp}(\bar{c}/k_0\langle\bar{A}_{s+1}\rangle)$ does not fork over $k_0\langle\bar{A}_s\rangle$. The symmetry of forking (Remark 7.20(2)) implies that $\text{tp}(\bar{a}_{s+1}/(\bar{A}_s, \bar{c}))$ does not fork over $\bar{A}_s$. Therefore, by Remark 7.15(2), we have

$$(7.2) \quad \text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{a}_{s+1} = \text{trdeg}_{k_0\langle\bar{c}, \bar{A}_s\rangle} \bar{a}_{s+1}.$$

Since $\bar{a}_1, \dots, \bar{a}_{s+1}$ are independent, $\text{tp}(\bar{a}_{s+1}/k_0\langle\bar{c}, \bar{A}_s\rangle)$ does not fork over $k_0\langle\bar{c}\rangle$. Therefore, Remark 7.20(4) (applying with $A = k_0\langle\bar{c}\rangle$) implies that

$$\text{Cb}(\bar{a}_{s+1}/(\bar{A}_s, \bar{c})) = k_0\langle\bar{c}\rangle.$$

On the other hand, Remark 7.20(4) applied with $A = k_0\langle\bar{A}_s\rangle$ together with (7.2) implies that $\text{Cb}(\bar{a}_{s+1}/(\bar{A}_s, \bar{c}))$ is algebraic over $k_0\langle\bar{A}_s\rangle$. So, the claim is proved.

Fix $s \geq 1$. Since $\bar{c}$ generates $\text{Cb}(\bar{A}_s/k)$, Remark 7.20(4) implies that $\text{tp}(\bar{A}_s/k)$ is a non-forking extension of $\text{tp}(\bar{A}_s/k_0\langle\bar{c}\rangle)$. Let $\bar{d}$ be any finite tuple of generators of k over k_0 . The symmetry of forking (Remark 7.20(2)) implies that $\text{tp}(\bar{d}/k_0\langle\bar{c}, \bar{A}_s\rangle)$ does not fork over $k_0\langle\bar{c}\rangle$. Thus,

$$\text{trdeg}_{k_0\langle\bar{A}_s\rangle} k = \text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{c} + \text{trdeg}_{k_0\langle\bar{c}, \bar{A}_s\rangle} k = \text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{c} + \text{trdeg}_{k_0\langle\bar{c}\rangle} k.$$

Therefore (cf. Lemma 5.8 and (5.2)),

$$\text{trdeg}_{k_0\langle\bar{A}_s\rangle} \bar{c} = \text{trdeg}_{k_0\langle\bar{A}_{s+1}\rangle} \bar{c} \iff \text{trdeg}_{k_0\langle\bar{A}_s\rangle} k = \text{trdeg}_{k_0\langle\bar{A}_{s+1}\rangle} k.$$

Together with the claim, this proves the first statement of the proposition.

Now we prove the second part of the proposition. We will do this by showing that

- (P1) $\text{tp}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle)$ does not fork over $k_0\langle\bar{A}_{r+1}\rangle$ and
 (P2) $\text{tp}(\bar{a}_{r+2}/k_0\langle\bar{A}_{r+1}\rangle)$ is stationary.

If we prove these two statements, then, by Remark 7.20(4), we will have

$$\text{Cb}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle) \subset k_0\langle\bar{A}_{r+1}\rangle.$$

Then Lemma 7.26 will imply that $\text{Cb}(\bar{a}_{r+2}/k) = \text{Cb}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle) \subset k_0\langle\bar{A}_{r+1}\rangle$.

By the first part of the proposition, $\text{Cb}(\bar{a}_{r+2}/k)$ is algebraic over $k_0\langle\bar{A}_r\rangle \subset k_0\langle\bar{A}_{r+1}\rangle$. Lemma 7.26 implies that $\text{Cb}(\bar{a}_{r+2}/k) = \text{Cb}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle)$. Then Remark 7.20(4) applied to type $\text{tp}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle)$ and $A = k_0\langle\bar{A}_{r+1}\rangle$ implies that the type does not fork over A . This proves (P1).

It remains to prove (P2). Assume the contrary, that is, $\text{tp}(\bar{a}_{r+2}/k_0\langle\bar{A}_{r+1}\rangle)$ has at least two nonforking extensions to $k\langle\bar{A}_{r+1}\rangle$. One of them is $p_0 := \text{tp}(\bar{a}_{r+2}/k\langle\bar{A}_{r+1}\rangle)$. Since $\bar{a}_{r+2}$ and $\bar{A}_{r+1}$ are independent over k , p_0 does not fork over k (see Definition 7.13). Since the type $\text{tp}(\bar{a}_{r+2}/k)$ is stationary, Remark 7.15(3) implies that the type p_0 is stationary. We denote an extension different from p_0 by $q_0 := \text{tp}(\bar{b}/k\langle\bar{A}_{r+1}\rangle)$. Since $\text{tp}(\bar{a}_{r+2}/k\langle\bar{A}_r\rangle)$ is stationary (similarly to p_0), the restrictions p_1 and q_1 of p_0 and q_0 , respectively, to $k\langle\bar{A}_r\rangle$ are distinct. Moreover, since p_0 does not fork over $k_0\langle\bar{A}_r\rangle$, the same is true for q_0 . Therefore, Remark 7.20(3) applied to p_0 and q_0 as distinct nonforking extensions of $\text{tp}(\bar{a}_{r+1}/k_0\langle\bar{A}_r\rangle)$ yields a finite equivalence relation E defined over $k_0\langle\bar{A}_r\rangle$ such that $\neg E(\bar{a}_{r+2}, \bar{b})$. Since $\bar{b}$ and $\bar{a}_{r+2}$ are of the same type over $k_0\langle\bar{A}_{r+1}\rangle$, we have $\neg E(\bar{a}_{r+1}, \bar{a}_{r+2})$. Since the type p is stationary and $\bar{a}_{r+1}, \bar{a}_{r+2}, \dots$ are independent over $k\langle\bar{A}_r\rangle$, we have (cf. [20, Lemma 2.28]) that, for every $i \neq j$ such that $i, j > r$,

$$\text{tp}((\bar{a}_i, \bar{a}_j)/k\langle\bar{A}_r\rangle) = \text{tp}((\bar{a}_{r+1}, \bar{a}_{r+2})/k\langle\bar{A}_r\rangle).$$

Therefore, we have $\neg E(\bar{a}_i, \bar{a}_j)$. This contradicts the fact that E defines only finitely many equivalence classes. The contradiction finishes the proof of (P2). ■

Acknowledgments. We are grateful to Julio Banga and Alejandro Villaverde for teaching us about the importance of multi-experiment parameter identifiability at the AIM workshop “Identifiability problems in systems biology,” to Alexandre Sedoglavic for helpful discussions about his algorithm [24], and to the referees for their suggestions.

REFERENCES

- [1] F. ANSTETT-COLLIN, L. DENIS-VIDAL, AND G. MILLÉRIOUX, *A priori identifiability: An overview on definitions and approaches*, Annu. Rev. Control, 50 (2020), pp. 139–149, <https://doi.org/10.1016/j.arcontrol.2020.10.006>.
- [2] P. BÜRGISSE, M. CLAUSEN, AND A. SHOKROLLAHI, *Algebraic Complexity Theory*, Springer-Verlag, Berlin, Heidelberg, 1997, <https://doi.org/10.1007/978-3-662-03338-8>.
- [3] C. E. CLANCY AND Y. RUDY, *Na⁺ channel mutation that causes both Brugada and long-QT syndrome phenotypes*, Circulation, 105 (2002), pp. 1208–1213, <https://doi.org/10.1161/hc1002.105183>.
- [4] C. FIEKER, W. HART, T. HOFMANN, AND F. JOHANSSON, *Nemo/Hecke: Computer algebra and number theory packages for the Julia programming language*, in Proceedings of ISSAC 2017, ACM, New York, 2017, pp. 157–164, <http://doi.acm.org/10.1145/3087604.3087611>.
- [5] M. FINK AND M. NOBLE, *Markov models for ion channels: Versatility versus identifiability and speed*, Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci., 367 (2009), pp. 2161–2179, <http://www.jstor.org/stable/40485576>.

- [6] S. GERBERDING, N. OBATAKE, AND A. SHIU, *Identifiability of linear compartmental models: The effect of moving inputs, outputs, and leaks*, Linear Multilinear Algebra, (2020), <https://doi.org/10.1080/03081087.2020.1812497>.
- [7] E. GROSS, H. HARRINGTON, N. MESHKAT, AND A. SHIU, *Linear compartmental models: Input-output equations and operations that preserve identifiability*, SIAM J. Appl. Math., 79 (2019), pp. 1423–1447, <https://doi.org/10.1137/18m1204826>.
- [8] E. GROSS, N. MESHKAT, AND A. SHIU, *Identifiability of linear compartmental models: The singular locus*, Adv. in Appl. Math., 133 (2022), 102268, <https://doi.org/10.1016/j.aam.2021.102268>.
- [9] H. HONG, A. OVCHINNIKOV, G. POGUDIN, AND C. YAP, *SIAN: Software for structural identifiability analysis of ODE models*, Bioinformatics, 35 (2019), pp. 2873–2874, <https://doi.org/10.1093/bioinformatics/bty1069>.
- [10] H. HONG, A. OVCHINNIKOV, G. POGUDIN, AND C. YAP, *Global identifiability of differential models*, Comm. Pure Appl. Math., 73 (2020), pp. 1831–1879, <https://doi.org/10.1002/cpa.21921>.
- [11] I. KAPLANSKY, *An Introduction to Differential Algebra*, Hermann, Paris, 1957.
- [12] J. KARLSSON, M. ANGUELOVA, AND M. JIRSTRAND, *An efficient method for structural identifiability analysis of large dynamic systems*, IFAC Proc. Vol., 45 (2012), pp. 941–946, <https://doi.org/10.3182/20120711-3-BE-2027.00381>.
- [13] L. LJUNG AND T. GLAD, *On global identifiability for arbitrary model parametrizations*, Automatica, 30 (1994), pp. 265–276, [https://doi.org/10.1016/0005-1098\(94\)90029-9](https://doi.org/10.1016/0005-1098(94)90029-9).
- [14] D. MARKER, *Model theory of differential fields*, in Model Theory of Fields, Springer, Berlin, 1996, pp. 38–113, <http://projecteuclid.org/euclid.lnl/1235423156>.
- [15] D. MARKER, *Model Theory: An Introduction*, Springer, New York, 2002, <https://doi.org/10.1007/b98860>.
- [16] N. MESHKAT, C. KUO, AND J. DiSTEFANO, *On finding and using identifiable parameter combinations in nonlinear dynamic systems biology models and COMBOS: A novel web implementation*, PLoS ONE, 9 (2014), e110261, <https://doi.org/10.1371/journal.pone.0110261>.
- [17] A. OVCHINNIKOV, A. PILLAY, G. POGUDIN, AND T. SCANLON, *Computing all identifiable functions of parameters for ODE models*, Systems Control Lett., 157 (2021), 105030, <https://doi.org/10.1016/j.sysconle.2021.105030>.
- [18] A. OVCHINNIKOV, G. POGUDIN, AND P. THOMPSON, *Parameter identifiability and input-output equations*, Appl. Algebra Eng. Commun. Comput., (2021), <https://doi.org/10.1007/s00200-021-00486-8>.
- [19] A. OVCHINNIKOV, G. POGUDIN, AND P. THOMPSON, *Input-output equations and identifiability of linear ODE models*, IEEE Trans. Automat. Control, (2022), <https://doi.org/10.1109/TAC.2022.3145571>.
- [20] A. PILLAY, *Geometric Stability Theory*, Oxford Logic Guides 32, Clarendon Press, Oxford University, New York, 1996.
- [21] A. PILLAY, *Lecture notes: Applied stability theory*, <https://www3.nd.edu/~apillay/pdf/lecturenotes.applied.pdf>, 2003.
- [22] Y. RUDY AND J. R. SILVA, *Computational biology in the study of cardiac ion channels and cell electrophysiology*, Q. Rev. Biophys., 39 (2006), pp. 57–116, <https://doi.org/10.1017/s0033583506004227>.
- [23] M. SACCOMANI AND L. D’ANGIÓ, *Examples of testing global identifiability with the DAISY software*, IFAC Proc. Vol., 42 (2009), pp. 48–53, <https://doi.org/10.3182/20090706-3-FR-2004.00007>.
- [24] A. SEDOGAVIC, *A probabilistic algorithm to test local algebraic observability in polynomial time*, J. Symb. Comput., 33 (2002), pp. 735–755, <https://doi.org/10.1006/jsc.2002.0532>.
- [25] N. TUNCER AND T. T. LE, *Structural and practical identifiability analysis of outbreak models*, Math. Biosci., 299 (2018), pp. 1–18, <https://doi.org/10.1016/j.mbs.2018.02.004>.
- [26] J. VAN DEN HOF, *Structural identifiability of linear compartmental systems*, IEEE Trans. Automat. Control, 43 (1998), pp. 800–818, <https://doi.org/10.1109/9.679020>.
- [27] A. VILLAYERDE, N. EVANS, M. CHAPPELL, AND J. BANGA, *Input-dependent structural identifiability of nonlinear systems*, IEEE Control Syst. Lett., 3 (2019), pp. 272–277, <https://doi.org/10.1109/LCSYS.2018.2868608>.
- [28] A. F. VILLAYERDE, A. BARREIRO, AND A. PAPACHRISTODOULOU, *Structural identifiability of dynamic systems biology models*, PLoS Comput. Biol., 12 (2016), e1005153, <https://doi.org/10.1371/journal.pcbi.1005153>.
- [29] A. F. VILLAYERDE, N. D. EVANS, M. J. CHAPPELL, AND J. R. BANGA, *Sufficiently exciting inputs for structurally identifiable systems biology models*, IFAC-PapersOnLine, 51 (2018), pp. 16–19, <https://doi.org/10.1016/j.ifacol.2018.08.004>.

- [//doi.org/10.1016/j.ifacol.2018.09.015](https://doi.org/10.1016/j.ifacol.2018.09.015).
- [30] A. WIGDERSON, *Mathematics and Computation*, Princeton University Press, Princeton, NJ, 2019, <https://doi.org/10.1515/9780691192543>.
- [31] X. XIA AND C. MOOG, *Identifiability of nonlinear systems with application to HIV/AIDS models*, IEEE Trans. Automat. Control, 48 (2003), pp. 330–336, <https://doi.org/10.1109/tac.2002.808494>.