Check for updates

Towards climate-smart, three-dimensional protected areas for biodiversity conservation in the high seas

Isaac Brito-Morales ^{[1,2,3,4} ^[2], David S. Schoeman ^{[0,5,6}, Jason D. Everett ^{[0,2,3,7}, Carissa J. Klein^{1,8}, Daniel C. Dunn ^{[0,1,8}, Jorge García Molinos ^{[0,9,10,11}, Michael T. Burrows ^{[0,12}, Kristine Camille V. Buenafe^{1,3}, Rosa Mar Dominguez^{1,8}, Hugh P. Possingham⁸ and Anthony J. Richardson ^{[0,2,3}

Marine species are moving rapidly in response to warming, often in different directions and with variations dependent on location and depth. Given the current impetus to increase the area of protected ocean to 30%, conservation planning must include the 64% of the ocean beyond national jurisdictions, which in turn requires associated design challenges for conventional conservation to be addressed. Here we present a planning approach for the high seas that conserves biodiversity, minimizes exposure to climate change, retains species within reserve boundaries and reduces conflict with fishing. This is developed using data from across four depth domains, considering 12,932 vertebrate, invertebrate and algal species and three climate scenarios. The resultant climate-smart conservation areas cover 6% of the high seas and represent a low-regret option that provides a nucleus for developing a full network of high-seas marine reserves.

uman threats-including fishing and climate change-are impacting marine biodiversity, from the ocean surface to the deepest and most remote places on Earth^{1,2}. Marine protected areas (MPAs) are the most effective and widely used tools in conservation to mitigate threats3. Well-enforced MPAs can protect biodiversity⁴, enhance ecological resilience⁵, enrich local fish stocks⁶ and provide scientific reference sites in modified seascapes7.8. Climate change is considered in the design of some MPA networks, most commonly by using marine species distribution models under different climate scenarios9 and by making management recommendations on how to incorporate climate change into conservation plans^{10,11}. Although most MPA conservation plans include some aspects of both benthic (that is, seafloor) and pelagic (that is, water-column) biodiversity¹², and some include climate change, almost all are in relatively shallow environments with strong mixing where benthic and pelagic environments are considered together. However, in the high seas, we question whether it is appropriate to aggregate biodiversity in the top 200 m of the ocean with the distinct and largely separate biodiversity at 8km depth, especially given that oceanography, biology and climate change impacts vary greatly with depth^{1,13}.

Ensuring that MPAs are climate-smart (that is, considering climate change in their design) is challenging¹⁴. As the climate warms, species are rapidly shifting towards cooler regions^{15–19}, potentially moving beyond MPA boundaries. The retention of species in MPAs is further complicated by the three-dimensional nature of the open ocean. Because rates of ocean warming and spatial temperature gradients change with depth, the speed and the direction of species' movement are also likely to vary across the water column, a phenomenon that will be more pronounced in the future¹³. As MPAs have mostly been designed and declared in coastal and shelf regions without explicit consideration of different depth domains, a new paradigm for the design of MPAs in the open ocean^{20,21} is needed²² one that considers both its three-dimensional nature and the potential consequences of climate change.

Compared with the high seas, national jurisdictions experience intense competition between conservation and human uses, but have higher MPA coverage. For example, while 17% of national waters in exclusive economic zones (EEZs) are in partially protected MPAs, only 1.2% of the high seas is similarly protected, and 0.7% is fully protected²³. This shortfall in the protection of the high seas means that currently only ~7.5% of the ocean is within MPAs²⁴, well short of the goal of protecting 10% of the ocean by 2020 (Aichi Target 11 by the Convention on Biological Diversity). Furthermore, momentum is growing under the Post-2020 Global Biodiversity Framework to increase protection targets to 30% of the ocean by 2030²⁵. Since the inception of protection goals, policymakers have recognized that they would need to use the high seas to meet marine targets, but they have been hindered by fragmented governance regimes and the absence of global mechanisms to implement MPAs in the high seas²⁶. The imminent agreement of a new treaty for the conservation and sustainable use of biodiversity beyond national

¹School of Earth and Environmental Sciences, The University of Queensland, St Lucia, Queensland, Australia. ²Queensland Biosciences Precinct (QBP), Commonwealth Scientific and Industrial Research Organisation (CSIRO) Oceans and Atmosphere, St Lucia, Queensland, Australia. ³School of Mathematics and Physics, The University of Queensland, St Lucia, Queensland, Australia. ⁴Betty and Gordon Moore Center for Science, Conservation International, Arlington, VA, USA. ⁵Global-Change Ecology Research Group, School of Science, Technology and Engineering, University of the Sunshine Coast, Maroochydore, Queensland, Australia. ⁶Centre for African Conservation Ecology, Department of Zoology, Nelson Mandela University, Gqeberha, South Africa. ⁷Centre for Marine Science and Innovation (CMSI), The University of New South Wales, Sydney, New South Wales, Australia. ⁸Centre for Biodiversity and Conservation Science (CBCS), The University of Queensland, St Lucia, Queensland, Australia. ⁹Arctic Research Center, Hokkaido University, Sapporo, Japan. ¹⁰Global Station for Arctic Research, Global Institution for Collaborative Research and Education, Hokkaido University, Sapporo, Japan. ¹¹Graduate School of Environmental Science, Hokkaido University, Sapporo, Japan. ¹²Scottish Association for Marine Science, Oban, UK. ⁵⁵e-mail: i.britomorales@uq.net.au



Fig. 1 The degree of agreement between the climate-smart MPA networks for different planning domains and climate scenarios. The κ index for the relationship between each prioritized climate-smart network design for the four ocean planning domains: epipelagic, mesopelagic, bathyabyssopelagic and the seafloor.

jurisdiction will establish just such a mechanism²⁴, finally providing an opportunity for policymakers to address arguably the biggest gap in biodiversity protection on the planet²². High seas or areas beyond national jurisdiction are thus likely to be a primary focus of marine conservation in the future as global commitments to develop ecologically representative MPA networks increase.

To inform the development of MPAs in the high seas, we present a climate-smart planning approach across different depth domains that prioritizes the conservation of marine biodiversity and excludes fishing²⁷, while minimizing impacts on the fishing industry. Some studies argue that current MPA networks cannot effectively protect biodiversity against climate change^{5,28} because individual MPAs are too small to retain species undergoing range shifts in response to changing environmental conditions. We address this criticism by identifying ecological climate refugia²⁹—areas where biodiversity is likely to be least affected by climate change—and targeting these for protection.

To identify climate refugia, which will have lower exposure to future climate change and in which biodiversity will be more likely to be retained within MPAs, we used two climate-smart metrics. For biodiversity retention, we used climate velocity¹⁵, a generic predictor of the expected speed of species' distribution shifts under climate change^{18,30,31}. Areas with the slowest velocities are likely to see the smallest range shifts and are therefore considered more stable across time. As a metric of relative climate exposure (RCE), we used the temporal rate of temperature change divided by the mean annual temperature range (Methods). This assumes that species exposed to larger annual temperature ranges within any given bioregion (Methods) will exhibit greater thermal plasticity and, along with those exposed to slower warming (that is, where the numerator of the RCE is small), will be more resilient to climate change.

We have focused on temperature because of its fundamental importance as a driver of species' distributions³² and because it is correlated with nutrient availability, dissolved oxygen concentration and pH, thereby also controlling system structure and function³³. These latter correlations, together with our focus on refugia related to temperature, also mean that these areas will probably experience less overall change in terms of interlinked environmental variables than would be experienced in non-refuge areas. Species in climate refugia are thus likely to be more resilient to climate change, irrespective of whether their spatial distribution is towards the centre of the species' range or near their edge. Temperature also has a symmetrical functional relationship with species' biological performance, which is relevant in applications of climate velocity³⁴.

We created spatial plans for three alternative futures based on ocean temperatures from the IPCC Shared Socioeconomic Pathways (SSPs): SSP1-2.6 (an optimistic scenario with peak emissions in 2020), SSP2-4.5 (an intermediate scenario with peak emissions in 2040) and SSP5-8.5 (an unrestrained emission scenario). Both climate-smart metrics were estimated globally across a 0.5°×0.5° grid in four depth domains: three pelagic (epipelagic, mesopelagic and bathyabyssopelagic) and one benthic (seafloor). Each spatial plan included a set proportion of the distribution of 12,932 species of vertebrates (fish, birds, mammals and reptiles), invertebrates (molluscs, arthropods and corals) and macroalgae (green, red and brown) present within each depth zone and bioregion (Extended Data Fig. 1d-g and Supplementary Table 1). It also included 12 geomorphic features recognized for being hotspots of seafloor biodiversity (Methods and Extended Data Fig. 2). Taxa reported as threatened in the IUCN Red List were assigned high protection targets³⁵, whereas those not reported as threatened were assigned targets on the basis of the size of their global distribution (Methods).

ARTICLES



Fig. 2 | Climate-smart networks in the high seas. a-p, Coherence of prioritized networks in the high seas across different combinations of IPCC SSPs (SSP1-2.6, SSP2-4.5 and SSP5-8.5) and low-regret conservation areas, across four depth domains: epipelagic (a-d), mesopelagic (e-h), bathyabyssopelagic (i-l) and the seafloor (m-p). The polygons represent Longhurst provinces for the epipelagic domain, Glasgow provinces for the mesopelagic and bathyabyssopelagic domains, and Global Open Oceans and Deep Seabed provinces for the seafloor domain.

Because high-seas fisheries are a key economic resource³⁶, we minimized the cost to fishing, defined as the product of fish catch and fish value at any point in space (Methods) using the fish catch dataset of Watson³⁷, which includes estimates of illegal, unregulated and unreported catch and discards at sea. We then used integer linear programming to identify priority conservation areas for each depth domain and climate scenario using the R package prioritizr³⁸. Our intent is to answer one pressing conservation question: can we identify climate-smart areas common to all future climate scenarios and across depth domains? These are areas that could form a nucleus around which a network of high-seas MPAs could be built.

When we apply a climate-smart planning approach independently within each depth domain, there is moderate to substantial agreement among configurations of climate-smart MPA networks across emission scenarios in the epipelagic (Cohen's κ coefficient, κ =0.31 to 0.66), the mesopelagic (κ =0.60 to 0.75), the bathyabyssopelagic (κ =0.67 to 0.76) and the seafloor (κ =0.82 to 0.83) (Figs. 1 and 2 and Extended Data Fig. 3). This high degree of spatial correspondence implies that climate-smart conservation networks for particular depth domains could be robust to different climate futures. Given that we do not know the future climate, having substantial agreement among climate scenarios at each depth domain will be key to successful climate-smart conservation and could maximize the protection of biodiversity³⁴.

Running the prioritization analyses without a cost layer had relatively little influence on the spatial pattern of selected planning units (κ =0.91 to 0.95 between the MPA networks with and without a cost layer; Extended Data Figs. 4 and 5). But when estimating the total cost of the scenarios, it was higher for every depth domain and climate scenario (Extended Data Figs. 4 and 5). This suggests that most key conservation areas from a biodiversity perspective are retained when minimizing the cost to fisheries.

Priority areas common across emission scenarios could be considered 'low-regret conservation areas' because they can be conserved independent of the future climate. Low-regret areas for

NATURE CLIMATE CHANGE | www.nature.com/natureclimatechange

Table 1 | Proportion of the high seas (%) covered by all slow conservation feature priorities under different climate scenarios

Depth domain	SSP1- 2.6	SSP2- 4.5	SSP5-8.5	Climate-smart network (all three SSPs)
Epipelagic (0-200 m)	47.8	55.2	56	35.7
Mesopelagic (200–1,000 m)	57.1	57	57.7	47.7
Bathyabyssopelagic (>1,000 m)	48.4	47.4	46.3	39.3
Seafloor	37.6	38	38.2	33.7
Pelagic climate-smart network	-	-	-	12
Pelagic + seafloor climate-smart network	-	-	-	6

conservation include substantial proportions of the mesopelagic (47.7%), bathyabyssopelagic (39.3%), epipelagic (35.7%) and seafloor (33.7%) in the high seas (Fig. 2 and Table 1). These prioritized areas represent biodiversity by depth domain and bioregion that is likely to be both retained and less exposed to warming (Extended Data Fig. 6a). To date, conservation planning in the high seas has largely focused on individual depth domains, usually either the epipelagic³⁹ or the seafloor⁴⁰. In a warming ocean, networks of MPAs that are vertically coherent across depth domains would be the easiest to enforce and implement^{13,34}, but knowing where those areas are located might be the first challenge to address in future marine conservation initiatives.

Identifying vertically coherent low-regret areas for protection is difficult not only because costs differ among depth domains (Extended Data Fig. 1a–d) but also because the threat posed by



Fig. 3 | Low-regret climate-smart networks in the high seas. a-c, Climate-smart prioritization networks as low-regret conservation areas throughout the water column of the ocean for the pelagic domains (epipelagic, mesopelagic and bathyabyssopelagic) and for the pelagic plus the seafloor domains (b). The lateral panels show the latitudinal distribution of the prioritized climate-smart network as a proportion of ocean area for the pelagic domains (**a**) and the pelagic plus the seafloor domains (**b**).

disparate warming futures varies across depths (Extended Data Figs. 7 and 8). For example, due to the flatter spatial temperature gradient exhibited in the mesopelagic layer (Extended Data Fig. 7), its future climate velocity is projected to be faster than in the epipelagic layer, even with strong mitigation of future emissions¹³. This could cause dissociation of bioregions across depths and compromise the functioning, effectiveness and ecosystem services of climate-uninformed MPAs.

This variability in biodiversity, climate metrics and cost among pelagic depth domains (that is, epipelagic, mesopelagic and bathyabyssopelagic) across emission scenarios means that climate-smart MPA network configurations exhibit more spatial agreement between epipelagic and mesopelagic domains ($\kappa = 0.18$ to 0.24 across emission scenarios) and between mesopelagic and bathyabyssopelagic domains ($\kappa = 0.19$ to 0.24) than between epipelagic and bathyabyssopelagic domains ($\kappa = 0.05$ to 0.12) (Fig. 1). The benthic (seafloor) domain exhibits some agreement with the bathyabyssopelagic domain across emission scenarios ($\kappa = 0.24$ to 0.32) but little agreement with the mesopelagic ($\kappa = 0.12$ to 0.14) and the epipelagic domains ($\kappa = 0.04$ to 0.07) (Fig. 1). The spatial agreement (or disagreement) in prioritized networks across depth domains (Fig. 2) highlights not only differences in the cost of fishing with depth (Extended Data Fig. 1a-d) but also the vertical coherence across depths. This vertical coherence, in turn, is the result of vertically structured oceanographic phenomena, the vertical migration of biodiversity and the passive sinking of surface production, particularly through the epipelagic and mesopelagic⁴¹. The declining coherence with greater separation between domains is unsurprising, given that many species are unique to each depth domain⁴² and few vertically migrate far enough to span multiple depth domains. The spatial coherence in the placement of MPAs in the epipelagic and mesopelagic domains is encouraging in terms of establishing vertically coherent climate-smart MPAs in the upper ocean, but the lack of coherence with the bathyabyssopelagic domain and the seafloor suggests that in some cases, separate conservation plans might be needed for upper and lower ocean layers. Vertical zoning of MPAs in the ocean is currently rare⁴³, but examples include the Mexican Caribbean Biosphere Reserve and the Deep Mexican Pacific Biosphere Reserve⁴⁴.

A benefit of targeting ocean protection in areas where responses to climate change are coherent across vertical depth domains and scenarios (Fig. 3 and Extended Data Fig. 3) is that it decreases the risk of decoupling trophic relationships and may allow for the movement of MPA boundaries in response to novel climates^{13,34}. Our results show that 6% of the high seas are such 'low-regret conservation areas' (Fig. 3 and Supplementary Table 2). They represent key areas to conserve pelagic and benthic biodiversity in the high seas now and into the future across depth domains (Extended Data Fig. 6b). Considering only pelagic depth domains (that is, excluding the seafloor), the amount of ocean within low-regret conservation areas expands to 12% (Fig. 3 and Supplementary Table 2). Such low-regret sectoral management measures that operate in parts of the water column (for example, pelagic and benthic) represent 'other effective area-based conservation measures' and could serve as part of a global conservation network⁴³ (Extended Data Fig. 6b).

Biodiversity targets are key in motivating conservation actions, but to date politics and expediency have dominated ecology in the international policy arena⁴⁵, which has dictated how conservation targets are applied in global prioritization exercises^{35,46}. To perform a sensitivity analysis of how representation targets alter the global climate-smart MPA network identified here, we created several alternative climate-smart MPA networks with varying conservation targets for the 12,932 species distributions and the 12 seafloor geomorphic features (Methods). We ran seven additional prioritization analyses with new sets of targets (10% minimum with 10% increments to a maximum of 90%) for every restricted conservation feature (that is, species and geomorphic features in climate refugia in each bioregion) (Extended Data Figs. 9 and 10). As expected, the total area required to achieve these targets in vertically coherent low-regret MPA networks scales positively with target size (Supplementary Table 2), with less area prioritized for targets of 10-30% (targets for wide- and narrow-ranged features; 0.10% of the total high seas) and more area prioritized for targets of 10–90% (3.2% of the total high seas) (Supplementary Table 2). Importantly, despite varying targets for protection, considerable spatial agreement remains among prioritized climate-smart networks across climate scenarios, and with greater spatial agreement in configurations between epipelagic and mesopelagic depth domains than between the epipelagic and either the bathyabyssopelagic or seafloor domains (Extended Data Fig. 9). Higher biodiversity-protection targets in a climate-smart prioritization would increase the size of low-regret conservation areas, not only across different climatic scenarios but also in a coherent way across depth domains. This would provide substantial benefits to the Conservation of Biological Diversity agenda in a warming world.

There are several caveats in our analyses that should be considered. First, while our study focuses on the high seas because they are underprotected, we strongly believe that representative conservation of all coastal and marine habitats is necessary. Protection of the high seas should not replace or undermine efforts to protect critical coastal environments. Second, we focus on placing MPAs in climate refugia, but there are other possible design approaches, including

placing MPAs in areas of high chronic but low acute thermal stress to increase resistance, and potentially providing greater protection for species in areas of rapid change to mitigate non-climate threats¹⁴. Third, by focusing on temperature, we do not consider other climate change impacts that may affect the spatial prioritization. For example, primary productivity influences the community structure of pelagic food webs47, and its magnitude and spatial pattern are likely to change in the future⁴⁸, although uncertainty in this regard is high. Fourth, our climate metrics do not directly link climate warming with a species' thermal preference. For example, a species in a rapidly changing environment may be at relatively low risk if it has a broad thermal safety margin⁴⁹. Nevertheless, by targeting climate refugia²⁹ rather than a broad spectrum of climate change characteristics¹⁴, we focus on areas where the least change is expected, rendering the thermal preferences of resident species less important. Fifth, our approach did not consider the three main aspects of a well-connected marine MPA network in its design⁵⁰: structural connectivity, functional connectivity and climate connectivity. We consider the areas identified to be a nucleus around which to build a larger, broader, more connected MPA network that considers additional criteria. For example, climate-connected MPAs could be designed to enable biodiversity to shift along climate pathways⁵¹, either by moving MPAs with climate change^{52,53} or by designing stepping-stone MPAs³⁴. Sixth, our study assumes that all parts of a species' range are equally important, and meeting a conservation target for a species range will thus not necessarily confer the protection necessary to conserve metapopulation structure or species viability. Seventh, we assume that fishing is the primary driver of cost. In the future, data on the increasing number of mining licences to explore the deep-ocean seafloor could also be included⁵⁴, although they are currently proprietary and unavailable to public scrutiny⁵⁵. International cooperation will be vital in generating open-access information that can be used to minimize political and socio-economic conflicts in the design of open-ocean MPAs. Last, the fisheries cost laver quantifies only current fish catch and prices, both of which could change with future climate, economic and social changes.

Designed and enforced appropriately, a global network of climate-smart MPAs in the high seas will help conserve marine biodiversity in a warming world. An initial priority could be the protection of the low-regret areas across depth domains that we identified (6% of the high seas) because they are vertically coherent, are robust to different climate futures and protect key threatened species. Such protection would increase the coverage of the global MPA network from ~7.5% to ~11.3%. However, because our low-regret climate-smart areas do not consider other ecological criteria, such as spatial connectivity or adequacy²¹, they should be considered a nucleus around which a comprehensive MPA network can be built. In constructing an expanded network, society might need to consider separate reserve networks for different depth domains, if they are to be robust to different climate futures. Since climate-smart low-regret areas are geographically similar between the epipelagic and mesopelagic layers, a second priority could be to focus on the development of sectoral spatial management measures in the upper 1,000 m of the ocean; this could add another 10% of the high seas to the area conserved, mainly representing 'other effective area-based conservation measures'. A greater focus on a separate planning process for the bathyabyssopelagic and the seafloor could consider potential costs beyond deep-sea fishing, including mining and underwater cabling⁵⁵.

As momentum builds towards protection of 30% of the ocean by 2030²⁵, the expansion of the current MPA network is a primary goal of the 2021–2030 Decade of Ocean Science for Sustainable Development. To be effective, conservation and management will need to include the high seas. The climate-smart MPAs proposed here are not meant to be prescriptive but rather to advance the

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at https://doi.org/10.1038/ s41558-022-01323-7.

Received: 13 April 2021; Accepted: 23 February 2022; Published online: 04 April 2022

References

- Levin, L. A. & Le Bris, N. The deep ocean under climate change. Science 350, 766–768 (2015).
- 2. Pecl, G. T. et al. Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science* **355**, eaai9214 (2017).
- 3. Roberts, C. M. et al. Marine reserves can mitigate and promote adaptation to climate change. *Proc. Natl Acad. Sci. USA* **114**, 6167–6175 (2017).
- Davies, T. E., Maxwell, S. M., Kaschner, K., Garilao, C. & Ban, N. C. Large marine protected areas represent biodiversity now and under climate change. *Sci. Rep.* 7, 9569 (2017).
- 5. Bates, A. E. et al. Climate resilience in marine protected areas and the 'protection paradox'. *Biol. Conserv.* **236**, 305–314 (2019).
- Costello, M. J. & Ballantine, B. Biodiversity conservation should focus on no-take marine reserves: 94% of marine protected areas allow fishing. *Trends Ecol. Evol.* 30, 507–509 (2015).
- Ballantine, B. Fifty years on: lessons from marine reserves in New Zealand and principles for a worldwide network. *Biol. Conserv.* 176, 297–307 (2014).
- Lester, S. E. et al. Biological effects within no-take marine reserves: a global synthesis. *Mar. Ecol. Prog. Ser.* 384, 33–46 (2009).
- Jones, K. R., Watson, J. E. M., Possingham, H. P. & Klein, C. J. Incorporating climate change into spatial conservation prioritisation: a review. *Biol. Conserv.* 194, 121–130 (2016).
- Grorud-Colvert, K. et al. The MPA Guide: a framework to achieve global goals for the ocean. *Science* 373, eabf0861 (2021).
- 11. McLeod, E. et al. Integrating climate and ocean change vulnerability into conservation planning. *Coast. Manage.* **40**, 651–672 (2012).
- Magris, R. A. et al. A blueprint for securing Brazil's marine biodiversity and supporting the achievement of global conservation goals. *Divers. Distrib.* 27, 198–215 (2021).
- Brito-Morales, I. et al. Climate velocity reveals increasing exposure of deep-ocean biodiversity to future warming. *Nat. Clim. Change* 10, 576–581 (2020).
- 14. Tittensor, D. P. et al. Integrating climate adaptation and biodiversity conservation in the global ocean. *Sci. Adv.* 5, eaay9969 (2019).
- 15. Burrows, M. T. et al. The pace of shifting climate in marine and terrestrial ecosystems. *Science* **334**, 652–655 (2011).
- Burrows, M. T. et al. Geographical limits to species-range shifts are suggested by climate velocity. *Nature* 507, 492–495 (2014).
- Chaudhary, C., Richardson, A. J., Schoeman, D. S. & Costello, M. J. Global warming is causing a more pronounced dip inmarine species richness around the Equator. *Proc. Natl Acad. Sci. USA* 118, e2015094118 (2021).
- Lenoir, J. et al. Species better track climate warming in the oceans than on land. *Nat. Ecol. Evol.* 4, 1044–1059 (2020).
- Poloczanska, E. S. et al. Global imprint of climate change on marine life. Nat. Clim. Change 3, 919–925 (2013).
- Levin, N., Kark, S. & Danovaro, R. Adding the third dimension to marine conservation. *Conserv. Lett.* 11, e12408 (2018).
- O'Leary, B. C. & Roberts, C. M. Ecological connectivity across ocean depths: implications for protected area design. *Glob. Ecol. Conserv.* 15, e00431 (2018).
- 22. Game, E. T. et al. Pelagic protected areas: the missing dimension in ocean conservation. *Trends Ecol. Evol.* **24**, 360–369 (2009).
- Protected Planet Report 2020 (UNEP-WCMC and IUCN, 2021); https://livereport.protectedplanet.net/
- Wright, G. et al. Marine spatial planning in areas beyond national jurisdiction. *Mar. Policy* 132, 103384 (2021).
- 25. Zero Draft of the Post-2020 Global Biodiversity Framework (Convention on Biological Diversity, 2020).

- Dunn, D. C. et al. The Convention on Biological Diversity's ecologically or biologically significant areas: origins, development, and current status. *Mar. Policy* 49, 137–145 (2014).
- Claudet, J., Loiseau, C., Sostres, M. & Zupan, M. Underprotected marine protected areas in a global biodiversity hotspot. One Earth 2, 380–384 (2020).
- Bruno, J. F. et al. Climate change threatens the world's marine protected areas. Nat. Clim. Change 8, 499–503 (2018).
- Arafeh-Dalmau, N. et al. Incorporating climate velocity into the design of climate-smart networks of marine protected areas. *Methods Ecol. Evol.* 12, 1969–1983 (2021).
- 30. García Molinos, J. et al. Climate velocity and the future global redistribution of marine biodiversity. *Nat. Clim. Change* **6**, 83–88 (2016).
- Pinsky, M. L., Worm, B., Fogarty, M. J., Sarmiento, J. L. & Levin, S. A. Marine taxa track local climate velocities. *Science* 341, 1239–1242 (2013).
- Tittensor, D. P. et al. Global patterns and predictors of marine biodiversity across taxa. *Nature* 466, 1098–1101 (2010).
- Richardson, A. J. In hot water: zooplankton and climate change. ICES J. Mar. Sci. 65, 279–295 (2008).
- Brito-Morales, I. et al. Climate velocity can inform conservation in a warming world. *Trends Ecol. Evol.* 33, 441–457 (2018).
- Jones, K. R. et al. Area requirements to safeguard Earth's marine species. One Earth 2, 188–196 (2020).
- Ortuño Crespo, G. & Dunn, D. C. A review of the impacts of fisheries on open-ocean ecosystems. *ICES J. Mar. Sci.* 74, 2283–2297 (2017).
- 37. Watson, R. A. A database of global marine commercial, small-scale, illegal and unreported fisheries catch 1950–2014. *Sci. Data* **4**, 170039 (2017).
- Hanson, J. O. et al. prioritizr: Systematic Conservation Prioritization in R. R package version 5.0 (2021).
- Visalli, M. E. et al. Data-driven approach for highlighting priority areas for protection in marine areas beyond national jurisdiction. *Mar. Policy* 122, 103927 (2020).
- 40. Dunn, D. C. et al. A strategy for the conservation of biodiversity on mid-ocean ridges from deep-sea mining. *Sci. Adv.* 4, eaar4313 (2018).
- 41. Irigoien, X. et al. Large mesopelagic fishes biomass and trophic efficiency in the open ocean. *Nat. Commun.* 5, 3271 (2014).
- Costello, M. J. & Chaudhary, C. Marine biodiversity, biogeography, deep-sea gradients, and conservation. *Curr. Biol.* 27, R511–R527 (2017).

NATURE CLIMATE CHANGE

- Venegas-Li, R., Levin, N., Possingham, H. & Kark, S. 3D spatial conservation prioritisation: accounting for depth in marine environments. *Methods Ecol. Evol.* 9, 773–784 (2018).
- Menini, E. & Van Dover, C. L. An atlas of protected hydrothermal vents. Mar. Policy 108, 103654 (2019).
- 45. Crespo, G. O. et al. High-seas fish biodiversity is slipping through the governance net. *Nat. Ecol. Evol.* **3**, 1273–1276 (2019).
- Hanson, J. O. et al. Global conservation of species' niches. Nature 580, 232–234 (2020).
- Barton, A. D. et al. The biogeography of marine plankton traits. *Ecol. Lett.* 16, 522–534 (2013).
- Tittensor, D. P. et al. Next-generation ensemble projections reveal higher climate risks for marine ecosystems. *Nat. Clim. Change* 11, 973–981 (2021).
- Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L. & Sunday, J. M. Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature* 569, 108–111 (2019).
- Daigle, R. M. et al. Operationalizing ecological connectivity in spatial conservation planning with Marxan Connect. *Methods Ecol. Evol.* 11, 570–579 (2020).
- Fredston-Hermann, A., Gaines, S. D. & Halpern, B. S. Biogeographic constraints to marine conservation in a changing climate. *Ann. N. Y. Acad. Sci.* 1429, 5–17 (2018).
- Cashion, T. et al. Shifting seas, shifting boundaries: dynamic marine protected area designs for a changing climate. *PLoS ONE* 15, e0241771 (2020).
- Ortuño Crespo, G. et al. Beyond static spatial management: scientific and legal considerations for dynamic management in the high seas. *Mar. Policy* 122, 104102 (2020).
- Levin, L. A., Amon, D. J. & Lily, H. Challenges to the sustainability of deep-seabed mining. *Nat. Sustain.* 3, 784–794 (2020).
- Levin, L. A. et al. Climate change considerations are fundamental to management of deep-sea resource extraction. *Glob. Change Biol.* 26, 4664–4678 (2020).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2022

Methods

Study domain. We used the most recent global dataset of marine regions to define the spatial extent of the high seas (v.11, https://www.marineregions.org/). We defined the high seas as marine areas outside of the EEZs, which represent 95% of habitat on Earth by volume, 64% of the surface of the ocean and 46% of the surface of the Earth²⁴. By excluding EEZs from this study, we omit parts of the ocean that contain most known marine biodiversity and are relatively well protected, and focus instead on areas beyond national jurisdiction, which are likely to be central to efforts to protect 30% of the ocean by 2030. MPAs in the deep ocean also address growing threats. Bottom trawling occurs to depths of 2,200 m (refs. 56,57), and indirect impacts of epipelagic and midwater trawls have been described down to 2,500 m (ref. 58). Beyond fisheries, deep-sea mining is set to begin in 2023 off Nauru, and deepwater cables can also damage habitat on the seafloor at virtually any depth. Here we classified the high seas into three pelagic layers (epipelagic, 0-200 m; mesopelagic, 200-1,000 m; and bathyabyssopelagic, >1,000 m) and one benthic layer (seafloor) using the ETOPO1 bathymetry dataset⁵⁹. We treated each layer as a separate planning domain containing equal-area hexagonal planning units of 2,620 km² (~0.5° at the equator). This yielded a total of 90,065 planning units for the epipelagic domain, 88,528 planning units for the mesopelagic domain and 87,170 planning units for the bathyabyssopelagic domain; there are fewer planning units by depth because of seamounts and underwater mountain ranges. For the seafloor domain, there were 90,065 planning units. Information about climate metrics, species and fishing data were assigned to each depth domain (as described below), each of which could then be prioritized for inclusion in a climate-smart MPA network.

Climate change metrics: sources and processing. Climate change metrics (that is, climate velocity and RCE) were estimated using future ocean temperatures from the Coupled Model Intercomparison Project Phase 6 (Earth System Grid Federation, https://esgf.llnl.gov). For the three pelagic domains, we used sea temperature from a multi-model ensemble mean derived from 11 general circulation models (GCMs), and for the seafloor, we used bottom temperatures from a multi-model ensemble mean derived from 12 GCMs (Supplementary Tables 3 and 4). We used models under three IPCC SSPs⁶⁰: SSP1-2.6, SSP2-4.5 and SSP5-8.5. Pathway SSP1-2.6 represents an optimistic scenario, characterized by a shift to a more sustainable economy and a reduction in inequality, resulting in a peak in radiative forcing of ~3 W m⁻² before 2100. SSP2-4.5 represents an intermediate scenario, with a stabilization of radiative forcing levels at ~4.5 W m⁻² by 2100. SSP5-8.5 is characterized by a continued increase of greenhouse gas emissions resulting from a fossil-fuel-based economy and increased energy demand, with a radiative forcing >8.5 W m⁻² by 2100, rising thereafter.

To construct the multi-model ensemble, we followed published methods¹³. Briefly, for each climate model, we regridded from the original grid to a uniform 0.5° spatial grid using an area-weighted bilinear interpolation⁶¹. We then extracted sea temperature data from the GCMs by depth according to three different ocean layers: epipelagic (0-200 m), mesopelagic (200-1,000 m) and bathyabyssopelagic (>1,000 m)⁶². For each depth layer, we averaged temperatures using a volume-weighting approach, with volumes of 0.5° grid squares in each depth layer. Finally, to avoid artefacts caused by inconsistent numbers of grid cells available by depth for different models, we included only grid cells common to all models within each depth layer¹³. We have assumed that integrating temperatures across broad ocean depths is a reasonable approximation of habitat conditions in the ocean^{34,43,62,63}. Although temperature varies continuously by depth, these ocean layers have been defined on the basis of consistency in their physical, chemical and biological environments63, and these conditions are further integrated by the extensive vertical migration of many marine species⁴¹. Using temperature alone, instead of other climate variables, means that our results reflect only the potential impact of warming on biodiversity. All analysis was undertaken using the software tools Climate Data Operators (version 1.9)64 and R (version 3.5)65 (Supplementary Tables 3 and 4).

Calculation of climate velocity and RCE metrics. Climate change has been considered in previous conservation planning exercises, particularly in coastal regions⁶⁶⁻⁶⁸. Global conservation planning exercises have used sea surface temperature as the metric of climate change^{39,69}, with none using data throughout the water column. Here, to prioritize a climate-smart MPA network, we focused on areas across depth domains (including the seafloor) that accomplish two objectives: (1) high retention of biodiversity and (2) low levels of exposure to future climate warming. We used two metrics of climate change to represent these objectives: climate velocity and RCE. Climate velocity is a metric that gives expectations for species' range shifts under projected future ocean warming^{15,18,30}. It is expected that in areas of slow climate velocity, species' distributions are likely to shift less, promoting their retention within a given area. We estimate local climate velocity at 0.5° resolution for the second half of the century (2050-2100) at each depth domain of the multi-model CMIP6 ensemble. The temporal trend (that is, the numerator of climate velocity) was calculated as the slope of a linear regression of mean annual temperatures (°Cyr⁻¹) for the corresponding climate scenario time period. The spatial gradient (that is, the denominator of climate velocity) was calculated from the vector sum of the latitudinal and longitudinal pairwise

RCE is a metric that we developed to obtain information about the amount of exposure to climate warming that local populations of a species would face relative to its experience of variation in seasonal temperatures. We define exposure to climate change as the presence of a species, ecosystem or habitat at a location where it might be adversely affected by climate change. We calculated RCE as the ratio of the slope of a linear regression of projected mean annual temperatures (°Cyr⁻¹ in 2050–2100) to the current mean seasonal temperature range (°C in 2015–2020):

 $RCE = \frac{Slope of temperature change(^{\circ}C yr^{-1})}{Current seasonal range(^{\circ}C)}$

It is expected that by prioritizing areas with a low climate-exposure index, we will select MPAs that will be more likely to minimize the potential exposure of species to future warming. Although some metrics of exposure have already been incorporated in marine spatial planning^{28,71}, our metric considers not only information on warming but also the local change relative to seasonal variation, and therefore, presumably, the relative vulnerability of resident biodiversity to projected future temperatures. It is important to mention that the RCE metric developed here is only weakly correlated with latitude (r < 0.1 for almost all depth domains and climate scenarios) and shows a similar pattern to climate velocity in terms of greater exposure to warming for species living below the epipelagic layer and under higher emissions (that is, the SSP5-8.5 climate scenario) (Extended Data Fig. 8). For each depth domain, we calculated climate velocity and RCE separately for all three SSPs.

We have chosen to analyse three pelagic depth domains and the seafloor rather than vertical climate velocity for several reasons. First, because species fill their thermal niches72,73, species are likely to move both horizontally and vertically into available habitat. The large horizontal movements (kilometres) are greater than the vertical movements (metres) of species with warming, so our approach of using horizontal climate velocity is best suited to exploring the retention of species within MPA boundaries³⁴. Second, recent empirical evidence⁷⁴ suggests that depth provides refuge from ocean warming only in areas of steep vertical temperature gradients. However, vertical temperature gradients are gentle over most of the open ocean (>500 m depth). Thus, in most of the high seas, vertical shifts would need to be much greater than in coastal areas to avoid climate warming and are less likely74. Such large vertical shifts are rendered less likely because they could dissociate distributions of animals from their food, especially for herbivores, and risk increasing exposure to other climate hazards, such as shoaling oxygen-minimum zones. In the high seas, horizontal climate velocity is likely to be a more useful proxy for community-level responses to climate change than vertical velocity. This is especially pertinent in our analysis because we integrate across large depth domains within which vertical temperature responses might operate. Last, it is not straightforward to assign the relative importance of the horizontal and vertical dimensions of climate velocity in a three-dimensional conservation plan.

Conservation features. To solve the minimum-set conservation prioritization problem, it is necessary to specify a protection target a priori for each conservation feature, which indicates the minimum amount of each feature (that is, species or habitat) to be included within the final prioritized network⁷⁵. Although there are multiple criteria for identifying areas for biodiversity conservation⁷⁶, our conservation features here were marine species distribution maps from AquaMaps⁷⁷ (v.2019) and seafloor geomorphic features⁷⁸ from the Blue Habitats dataset (www.bluehabitats.org).

The AquaMaps dataset predicts marine species distributions using a probability of occurrence (0-1) derived from an environmental niche model based on depth, temperature, salinity and oxygen at 0.5° spatial resolution. It includes 33,518 marine species, 23,700 of which we considered, as their environmental envelopes were generated using at least ten observations¹³. Since published studies have concluded that varying the probability of occurrence to select thresholds in prioritization analysis has little impact in prioritized networks³⁵, we applied the commonly used minimum threshold of 0.5 probability of occurrence^{13,69} to set range maps for our high-seas depth domains. Since most biodiversity in the ocean is located in coastal regions, our selection criteria yielded 12,932 species distribution maps across the depth domains within the high seas (Supplementary Table 1).

We classified the AquaMaps species into those that were pelagic and benthic using information from FishBase⁷⁹ and SeaLifeBase⁸⁰. Pelagic species were those classified as pelagic–neritic, pelagic, bathypelagic, pelagic–oceanic or epipelagic. Benthic species were those that are classified as sessile, demersal, benthic, reef-associated, benthopelagic or bathydemersal. We assigned the pelagic species to each of the three pelagic depth domains on the basis of their depth range from the AquaMaps depth envelopes. This yielded 1,081 conservation features in the epipelagic domain (0–200 m), 1,300 in the mesopelagic domain (200–1,000 m) and 519 in the bathyabyssopelagic domain (>1,000 m; Extended Data Fig. 1e–h and Supplementary Table 1). We also assigned each of the benthic species to the seafloor domain, which yielded 10,860 conservation features.

ARTICLES

In addition to the seafloor conservation features derived from AquaMaps, we included geomorphic features⁷⁸ in the seafloor domain. Geomorphic features are often associated with highly productive zones and are hotspots for biodiversity^{81,82}. We used the 12 categories that best represent biological species-associated information as conservation features for the seafloor domain (Extended Data Fig. 2). These features (basins, bridges, canyons, escarpments, fans, guyots, plateaus, ridges, seamounts, sills, trenches and troughs) bring the total number of conservation features in the seafloor domain to 10,872.

To obtain a better representation of conservation features in the global climate-smart MPA network across latitudes and ocean planning domains, we ensured that each feature is represented in every marine biogeographical provinces that it overlaps with a threshold of 0.5. We used different biogeographical provinces for different depth domains. For the epipelagic domain, we used Longhurst provinces, which are widely recognized by policymakers^{83,84} and which reflect major oceanographic regions and boundaries that are compatible with the current climate-smart approach, although there are other available bioregionalizations⁸⁵. For the mesopelagic and bathyabyssopelagic depth domains, we used the Glasgow provinces⁸⁶, and for the seafloor domain, we used the Global Open Oceans and Deep Seabed provinces⁸⁷. Since most conservation features appear in multiple provinces, this categorization process yielded a total of 66,093 conservation features: 12,791 for the epipelagic domain, 15,141 for the mesopelagic domain, 7,083 for the bathyabyssopelagic domain and 31,078 for the seafloor domain.

Opportunity cost of fishing. When solving the minimum-set problem in spatial prioritization, the objective is to identify areas where MPAs can meet conservation targets at a minimum cost⁷⁵. In our global prioritization analysis, we used the value of fisheries as a cost layer because it is the most common cost layer used in marine planning, as resulting priority areas avoid valuable fishing grounds where possible⁸⁸. Our use of a fisheries cost layer is predicated on the reality that biodiversity conservation in the high seas relies on sectoral implementation of protection measures and their need to balance competing objectives of conservation and fisheries²⁴. However, we have also run a climate-smart prioritization analysis without a cost layer (that is, the base scenario) to focus solely on conservation objectives.

To estimate the fisheries value (US\$) of each planning unit in each depth domain, we used a global database of catch (kg)37 and price (US\$ kg-1)89 for each species caught. The catch data contained fishing records for 1,242 species of fish and invertebrates from different publicly available databases from 1950 to 2014³⁷. This dataset includes landings at 0.5° spatial resolution, is interpolated to account for missing values and includes an estimate of illegal, unreported and unregulated fishing recorded as discards³⁷. For the cost layer, we used a subset of the total catch for the period 2005–2014 in each 0.5° cell (Extended Data Fig. 1a–d) and obtained the mean price of each species (US\$)89 from the Sea Around Us project dataset. Note that these prices are not absolute and are likely to change in the future due to economic factors and fish availability. To obtain a species' price for each depth domain, we categorized species by pelagic and benthic habitats using the same approach as described in 'Conservation features'. For pelagic species, we also used the FishBase database (https://www.fishbase.se/) to classify species by depth range preferences. This process yielded a total of 1,076 prices for species (US\$), 221 prices for the epipelagic layer, 232 prices for the mesopelagic layer, 55 prices for the bathyabyssopelagic layer and 834 prices for benthic species associated with the seafloor. To obtain a final cost layer for each depth layer, we created a cost per square metre for each species and then multiplied it by the number of square metres that each species covers in each pelagic domain. We calculated a total price by adding prices for all species within each 0.5° cell. Finally, we overlapped each cost layer generated with the corresponding depth domain to obtain an area-weighted mean total cost (US\$) for each planning unit (Extended Data Fig. 1a-d). Because fishing is limited in deep ocean layers, we assigned a zero cost in areas or depths where fishing was absent.

Spatial conservation prioritization. We used integer linear programming to find climate-smart MPA networks across depth domains that minimize the overall cost (that is, fishing value in US\$) of the MPA network and achieve the representation targets for each conservation feature. We solved the minimum-set objective⁷⁵ using the R package prioritizr³⁸ and Gurobi optimization software (version 9.0)⁹⁰. We set Gurobi to achieve a solution within 10% of the optimal solution (that is, the solution that achieves the coverage target with the lowest possible cost). This is a relatively arbitrary number that establishes the difference between the upper and lower bounds of the objective function⁹¹. For example, a value of 0.10 results in the optimizer stopping and returning solutions when the difference between the bounds reaches 10% of the upper bound.

Targets for conservation features were generated as follows (Extended Data Fig. 10). First, each conservation feature was intersected with both climate velocity and RCE maps to obtain its corresponding climate change condition in each planning unit. Then, for each conservation feature, we selected only the planning units in which the conservation feature experienced slow climate change. We defined slow climate change as values in the lowest quartile for each of climate velocity and RCE (that is, slow climate velocity and low RCE) within the range of the species under consideration (Extended Data Fig. 10). This process reduced the distribution size of

NATURE CLIMATE CHANGE

each conservation feature (that is, the number of cells represented in the high seas) to one quarter of its initial distribution (Extended Data Fig. 10), yielding 'restricted' conservation features. Next, for each of these restricted conservation features, we assigned targets on the basis of the conservation status reported by the IUCN Red List⁹². For this, we used the R package rredlist⁹³ to obtain the IUCN classification for each conservation feature. For taxa reported as threatened (that is, vulnerable, endangered or critically endangered), every associated restricted conservation feature was assigned a fixed target of 100%35 (that is, 100% of its distribution within the first quartile of our climate-smart metrics for its overall distributional range) (Extended Data Fig. 10). For conservation features not reported as threatened in the IUCN Red List, we assigned relative targets on the basis of the size of their global distribution (that is, the restricted conservation feature) represented in each depth domain, with a minimum of 10% for features with broad ranges and 100% for features with limited ranges (10-100% of its restricted distribution for each conservation feature) (Extended Data Fig. 10 and 'Sensitivity analyses'). We calculated these relative targets using the equation:

$$\text{Target}(\%) = \text{Target}_{\text{max}}(\%) - \frac{\text{PU}_{i}}{\text{PU}_{\text{total}}} \times \left[\text{Target}_{\text{max}}(\%) - \text{Target}_{\text{min}}(\%)\right]$$

where Target_{max} and Target_{min} refer to the maximum and minimum protection targets (%) for restricted conservation feature *i*, PU_i is the number of planning units represented for slow conservation feature *i*, and PU_{total} refers to the total number of planning units for each high-seas depth planning region. By using relative targets instead of fixed targets, we ensured the representation of multiple areas where widely distributed restricted conservation features were conserved, as well as the protection of restricted conservation features within limited distribution ranges³⁵.

We created different prioritization planning scenarios to determine how the incorporation of climate change metrics (that is, slow climate velocity and low RCE) drives the selection of a climate-smart MPA network under alternative climatic futures. We ran three scenarios: one where we included restricted conservation features under SSP1-2.6, one with restricted conservation features under SSP2-4.5 and one with restricted conservation features under SSP5-8.5. The resultant mean and median targets (%) across scenarios and depth domains were 79 ± 12 (n = 12, \pm s.d.) and 74 (70, 82; range as the first and third quartiles), respectively. Each prioritization scenario locked in protection in existing MPAs (data extracted from www.fao.org).

To determine the spatial similarity of selected planning units among prioritization scenarios and depth domains, we calculated the Cohen's κ coefficient⁸⁴. Cohen's κ is a pairwise statistic that indicates the degree of agreement among scenarios, ranging from -1 to 1, where -1 represents complete disagreement, 0 represents agreement due to chance and 1 represents complete agreement⁸⁴.

Sensitivity analyses. The total area protected in a prioritization analysis is highly correlated with the initial target assigned to each conservation feature. To test the sensitivity of our analysis, we tested different targets of protection for the climate-smart prioritization planning approach, with a minimum of 10% and sequentially adding 10% up to a maximum of 90% (10–90% of the restricted distribution for each conservation feature). This gave a total of seven new and different planning scenarios (Supplementary Table 2). For each set of targets (that is, planning scenarios) and for the base scenario, we performed a prioritization analysis using the R package prioritizr and the Gurobi software. As in the main design, we set Gurobi to achieve a solution within 10% of the optimum solution. Given that thousands of marine species were used in this study, a species-by-species and habitat assessment of vulnerabilities and sensitivities to all specific threats was not possible.

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The data used in this study (except the AquaMaps biodiversity and geomorphic features data) are available at Zenodo⁹⁵ under the identifier https://doi.org/10.5281/ zenodo.5912047. The AquaMaps⁷⁷ data are freely available via www.aquamaps.org. The geomorphic features⁷⁸ data are freely available via www.bluehabitats.org.

Code availability

All the scripts are available at Zenodo⁹⁵ under the identifier https://doi.org/10.5281/ zenodo.5912047.

References

- 56. Morato, T., Watson, R., Pitcher, T. J. & Pauly, D. Fishing down the deep. Fish Fish. 7, 24-34 (2006).
- Rogers, A. D. & Gianni, M. Implementation of UNGA Resolutions 61/105 and 64/72 in the Management of Deep-Sea Fisheries on the High Seas (DIANE, 2011).

- Bailey, D. M., Collins, M. A., Gordon, J. D. M., Zuur, A. F. & Priede, I. G. Long-term changes in deep-water fish populations in the Northeast Atlantic: a deeper reaching effect of fisheries? *Proc. R. Soc. B* 276, 1965–1969 (2009).
- NOAA National Geophysical Data Center ETOPO1 1 Arc-Minute Global Relief Model (NOAA National Centers for Environmental Information, 2009).
- 60. O'Neill, B. C. et al. The roads ahead: narratives for Shared Socioeconomic Pathways describing world futures in the 21st century. *Glob. Environ. Change* 42, 169–180 (2017).
- Vrac, M., Stein, M. L., Hayhoe, K. & Liang, X.-Z. A general method for validating statistical downscaling methods under future climate change. *Geophys. Res.* 34, L18701 (2007).
- Rogers, A. D. Environmental change in the deep ocean. Annu. Rev. Environ. Resour. 40, 1–38 (2015).
- 63. Sayre, R. G. et al. A three-dimensional mapping of the ocean based on environmental data. *Oceanography* **30**, 90–103 (2017).
- Schulzweida, U. CDO User Guide (Max Planck Institute for Meteorology, 2019).
- 65. R Core Team *R*: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, 2018).
- 66. Mumby, P. J. et al. Reserve design for uncertain responses of coral reefs to climate change. *Ecol. Lett.* **14**, 132–140 (2011).
- Magris, R. A., Heron, S. F. & Pressey, R. L. Conservation planning for coral reefs accounting for climate warming disturbances. *PLoS ONE* 10, e0140828 (2015).
- Chollett, I., Enríquez, S. & Mumby, P. J. Redefining thermal regimes to design reserves for coral reefs in the face of climate change. *PLoS ONE* 9, e110634 (2014).
- 69. Sala, E. et al. Protecting the global ocean for biodiversity, food and climate. *Nature* **592**, 397–402 (2021).
- García Molinos, J., Schoeman, D. S., Brown, C. J. & Burrows, M. T. VoCC: an R package for calculating the velocity of climate change and related climatic metrics. *Methods Ecol. Evol.* **10**, 2195–2202 (2019).
- Iwamura, T., Wilson, K. A., Venter, O. & Possingham, H. P. A climatic stability approach to prioritizing global conservation investments. *PLoS ONE* 5, e15103 (2010).
- Jorda, G. et al. Ocean warming compresses the three-dimensional habitat of marine life. *Nat. Ecol. Evol.* 4, 109–114 (2020).
- Sunday, J. M., Bates, A. E. & Dulvy, N. K. Thermal tolerance and the global redistribution of animals. *Nat. Clim. Change* 2, 686–690 (2012).
- Burrows, M. T. et al. Ocean community warming responses explained by thermal affinities and temperature gradients. *Nat. Clim. Change* 9, 959–963 (2019).
- 75. Ball, I. R., Possingham, H. P. & Watts, M. in *Spatial Conservation Prioritization: Quantitative Methods and Computational Tools* (eds Moilanen, A. et al.) Ch. 14 (Oxford Univ. Press, 2009).
- Asaad, I., Lundquist, C. J., Erdmann, M. V. & Costello, M. J. Ecological criteria to identify areas for biodiversity conservation. *Biol. Conserv.* 213, 309–316 (2017).
- 77. Kaschner, K. et al. AquaMaps: Predicted Range Maps for Aquatic Species (2019).
- Harris, P. T., Macmillan-Lawler, M., Rupp, J. & Baker, E. K. Geomorphology of the oceans. *Mar. Geol.* 352, 4–24 (2014).
- 79. Froese, R. & Pauly, D. FishBase (2021).
- 80. Palomares, M. L. D. & Pauly, D. SeaLifeBase (2021).
- Morato, T., Hoyle, S. D., Allain, V. & Nicol, S. J. Seamounts are hotspots of pelagic biodiversity in the open ocean. *Proc. Natl Acad. Sci. USA* 107, 9707–9711 (2010).

- Rowden, A. A. et al. A test of the seamount oasis hypothesis: seamounts support higher epibenthic megafaunal biomass than adjacent slopes. *Mar. Ecol.* 31, 95–106 (2010).
- Devred, E., Sathyendranath, S. & Platt, T. Delineation of ecological provinces using ocean colour radiometry. *Mar. Ecol. Prog. Ser.* 346, 1–13 (2007).
- Oliver, M. J. & Irwin, A. J. Objective global ocean biogeographic provinces. Geophys. Res. Lett. 35, L15601 (2008).
- Costello, M. J. et al. Marine biogeographic realms and species endemicity. Nat. Commun. 8, 1057 (2017).
- Sutton, T. T. et al. A global biogeographic classification of the mesopelagic zone. Deep Sea Res. 1 126, 85–102 (2017).
- Global Open Oceans and Deep Seabed (GOODS)—Biogeographic Classification (UNESCO, 2009).
- Ban, N. C. & Klein, C. J. Spatial socioeconomic data as a cost in systematic marine conservation planning. *Conserv. Lett.* 2, 206–215 (2009).
- Tai, T. C., Cashion, T., Lam, V. W. Y., Swartz, W. & Sumaila, U. R. Ex-vessel fish price database: disaggregating prices for low-priced species from reduction fisheries. *Front. Mar. Sci.* 4, 363 (2017).
- 90. Gurobi Optimizer Reference Manual (Gurobi Optimization, 2020).
- Hanson, J. O., Schuster, R., Strimas-Mackey, M. & Bennett, J. R. Optimality in prioritizing conservation projects. *Methods Ecol. Evol.* 10, 1655–1663 (2019).
- 92. IUCN Red List of Threatened Species (IUCN, 2020); https://www.iucnredlist. org/en
- 93. Chamberlain, S. rredlist: 'IUCN' Red List Client. R package version 0.7.0 (2020).
- 94. McHugh, M. L. Interrater reliability: the kappa statistic. *Biochem. Med.* 22, 276–282 (2012).
- Brito-Morales, I. Towards climate-smart, 3-D protected areas for biodiversity conservation in the high seas (v2.0). Zenodo https://doi.org/10.5281/ zenodo.5912047 (2022).

Acknowledgements

I.B.-M. was supported by the Advanced Human Capital Program of the Chilean National Research and Development Agency (ANID Grant No. 72170231). C.J.K. was supported by an ARC Future Fellowship (no. FT200100314). J.D.E. was funded by Australian Research Council Discovery Project No. DP19010229. We thank K. Kaschner, C. Garilao and K. Kesner-Reyes for providing the AquaMaps marine biodiversity data.

Author contributions

I.B.-M., D.S.S. and A.J.R. conceived the research. J.D.E. generated the fishing cost layer. I.B.-M. analysed the data. I.B.-M. wrote the first draft with input from D.S.S., A.J.R., C.J.K. and D.C.D. I.B.-M., D.S.S., A.J.R., C.J.K., D.C.D., J.D.E., J.G.M., M.T.B., K.C.V.B., R.M.D. and H.P.P. contributed equally to the discussion of ideas and analyses, and all authors commented on the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Extended data is available for this paper at https://doi.org/10.1038/s41558-022-01323-7.

Supplementary information The online version contains supplementary material available at https://doi.org/10.1038/s41558-022-01323-7.

Correspondence and requests for materials should be addressed to Isaac Brito-Morales. Peer review information *Nature Climate Change* thanks Rafael Magris, Derek Tittensor and Qianshuo Zhao for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

NATURE CLIMATE CHANGE



Extended Data Fig. 1 Map of monetary value of fishing and biodiversity in the high seas. Opportunity cost of fishing (a, b, c, d) and species richness (number of species) with a probability of occurrence > 0.5 (d, e, f, g) in the high seas at four depth domains. Polygons represent Longhurst provinces for the epipelagic domain (a, e), Glasgow provinces for the mesopelagic (b, f) and bathyabyssopelagic domains (c, g), and the GOOD provinces for seafloor domain (d, h).

ARTICLES



Extended Data Fig. 2 | Geomorphic conservation features in the seafloor domain⁷⁸**.** For each map, green hexagons indicate the presence of each geomorphic feature in each planning unit. Polygons represent the GOODS provinces⁸⁷.

NATURE CLIMATE CHANGE



Extended Data Fig. 3 | **Prioritised climate-smart networks in the high seas.** Prioritised networks for the high seas at three pelagic depth domains and the seafloor, under three IPCC Shared Socioeconomic Pathways (SSP1-2.6, SSP2-4.5 and SSP5-8.5). For each map, green hexagons represent selected planning units while light blue hexagons represent non-selected planning units. Polygons in each map represent Longhurst provinces for the epipelagic domain (a, b, c), Glasgow provinces for the mesopelagic (d, e, f) and bathyabyssopelagic (g, h, i) domains, and the GOODS provinces for seafloor domain (j, k, l).





Extended Data Fig. 4 | The relationship between climate-smart networks with and without the cost layer. Total Opportunity cost among the prioritised base scenario (that is, no cost) and the climate-smart prioritisation scenarios under three IPCC emission pathways (SSP1-2.6, SSP2-4.5 and SSP5-8.5).

Total Opportunity Cost (log₁₀ USD)

ARTICLES

NATURE CLIMATE CHANGE



Extended Data Fig. 5 | Prioritised climate-smart networks in the high seas for the base scenario. Prioritised networks for a base scenario (that is, no cost) for the high seas at three pelagic depth domains and the seafloor, under three IPCC Shared Socioeconomic Pathways (SSP1-2.6, SSP2-4.5 and SSP5-8.5). For each map, green hexagons represent selected planning units while light blue hexagons represent non-selected planning units. Polygons in each map represent Longhurst provinces for the epipelagic domain (a, b, c), Glasgow provinces for the mesopelagic (d, e, f) and bathyabyssopelagic (g, h, i) domains, and the GOODS provinces for seafloor domain (j, k, l).

ARTICLES



Extended Data Fig. 6 | Biodiversity representation for climate-smart networks in the high seas. Average taxonomic group representation (%) in low-regret conservation areas for three pelagic depth domains and the seafloor (a), and throughout the water column for the pelagic domains and pelagic plus the seafloor domain under three IPCC Shared Socioeconomic Pathways: SSP1-2.6, SSP2-4.5, and SSP5-8.5.

NATURE CLIMATE CHANGE



Extended Data Fig. 7 | Climate velocity in the high seas. Climate velocity (km decade⁻¹) in the high seas for projected sea temperatures (2050–2100) at three pelagic depth domains and the seafloor, under three IPCC Shared Socioeconomic Pathways (SSP1-2.6, SSP2-4.5 and SSP5-8.5). Polygons in each map represent Longhurst provinces for the epipelagic domain (a, b, c), Glasgow provinces for the mesopelagic (d, e, f) and bathyabyssopelagic (g, h, i) domains, and the GOODS provinces for the seafloor domain (j, k, l).

ARTICLES



Extended Data Fig. 8 | Relative Climate Exposure (RCE) index in the high seas. RCE index (years) in the high seas for projected sea temperatures (2050–2100) at three pelagic depth domains and the seafloor, under three IPCC Shared Socioeconomic Pathways (SSP1-2.6, SSP2-4.5 and SSP5-8.5). Polygons in each map represent Longhurst provinces for the epipelagic domain (a, b, c), Glasgow provinces for the mesopelagic (d, e, f) and bathyabyssopelagic (g, h, i) domains, and the GOODS provinces for the seafloor domain (j, k, l).

NATURE CLIMATE CHANGE



Extended Data Fig. 9 | The degree of agreement between the climate-smart MPA networks for different sets of conservation targets. The Kappa index for the relationship between each prioritised climate-smart network MPA for different area-based protection targets under four depth domains: Epipelagic, Mesopelagic, Bathyabyssopelagic and the Seafloor. The percentages represent the minimum and maximum targets of protection in each prioritisation analysis.





Extended Data Fig. 10 | Process for setting climate-smart conservation targets. Schematic representation for setting targets for conservation features in the climate-smart prioritisation planning approach.

nature research

Corresponding author(s): Isaac Brito-Morales

Last updated by author(s): Jan 28, 2022

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our <u>Editorial Policies</u> and the <u>Editorial Policy Checklist</u>.

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.				
n/a	Cor	firmed		
	\boxtimes	The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement		
	\boxtimes	A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly		
\boxtimes		The statistical test(s) used AND whether they are one- or two-sided Only common tests should be described solely by name; describe more complex techniques in the Methods section.		
\boxtimes		A description of all covariates tested		
\boxtimes		A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons		
	\boxtimes	A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)		
\boxtimes		For null hypothesis testing, the test statistic (e.g. F, t, r) with confidence intervals, effect sizes, degrees of freedom and P value noted Give P values as exact values whenever suitable.		
\boxtimes		For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings		
\boxtimes		For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes		
\boxtimes		Estimates of effect sizes (e.g. Cohen's d, Pearson's r), indicating how they were calculated		
Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.				

Software and code

Policy information about <u>availability of computer code</u>						
Data collection	No software was used to collect data in this study. All data came from different existing sources listed in the Methods section.					
Data analysis	We used CDO (climate data operators) and R 3.5.0 to analyse the data in this study. Codes and Scripts are available at Zenodo under the identifier https://doi.org/10.5281/zenodo.5510748					

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research guidelines for submitting code & software for further information.

Data

Policy information about availability of data

All manuscripts must include a data availability statement. This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

Data used in this study can be found at Zenodo under the identifier https://doi.org/10.5281/zenodo.5510748

Field-specific reporting

Life sciences

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We collected and used global ocean data on climate models, biodiversity, and fisheries and conducted climate-smart 3D prioritisation analyses to identify a global climate-smart network of marine protected areas across different ocean depths			
Research sample	We used data on: Climate models: Coupled Model Intercomparison Project Phase 6 (Earth System Grid Federation, https://esgf-node.llnl.gov). The ensemble model for each climate scenario and depth domain as been provided in the Zenodo repository. Bathymetry: ETOPO1 Global Relief Model. Included in the Zenodo repository Biodiversity: AquaMaps (standardized distribution maps) and IUCN Red List. These dataset are not provided in the Zenodo repository but can be downloaded from https://www.aquamaps.org/ and https://www.iucn.org/ Marine provinces: Longhurst Provinces from Marine Regions (www.marineregions.org), Glasgow Provinces from Sutton et al. (2017), and the GOODS provinces (Global open oceans and deep seabed: biogeographic classification). These datasets have been provided in the Zenodo repository Fishing data: Sea Around Us and FishBase datasets. Data layers with the costs per fishing are included in the Zenodo repository. Marine Protected Areas from protected planet. Not included in the repository but free available at www.protectedplanet.net Vulnerable Marine Ecosystems from FAO. Not included in the repository but free available at www.fao.org			
Sampling strategy	We used all free available databases that contained global information which allowed us to conduct our 3D climate-smart prioritisation analyses			
Data collection	Data were obtained from online repositories. The previous section describes the sources and availability.			
Timing and spatial scale	We used the most recent databases. Sea temperature from climate models (IPCC Shared Socioeconomic Pathways) range from 2020 to 2100. The previous section describes the sources and availability.			
Data exclusions	No data were excluded from our study			
Reproducibility	N/A - No experiments were performed			
Randomization	N/A - No experiments were performed			
Blinding	N/A - No experiments were performed			
Did the study involve fiel	d work? 🗌 Yes 🔀 No			

Reporting for specific materials, systems and methods

Methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

MRI-based neuroimaging

Materials & experimental systems

n/a	Involved in the study	n/a	Involved in the study
\boxtimes	Antibodies	\boxtimes	ChIP-seq
\boxtimes	Eukaryotic cell lines	\boxtimes	Flow cytometry
\boxtimes	Palaeontology and archaeology	\boxtimes	MRI-based neuroim
\boxtimes	Animals and other organisms		
\boxtimes	Human research participants		
\boxtimes	Clinical data		
\boxtimes	Dual use research of concern		

