

LOCALLY COMPACT SOFIC GROUPS

BY

LEWIS BOWEN*

*Department of Mathematics, University of Texas at Austin
2515 Speedway, Austin, TX 78712, USA
e-mail: lpbowen@math.utexas.edu*

AND

PETER BURTON

*Department of Mathematics, Iowa State University
Carver Hall, 411 Morrill Rd., Ames, IA 50011, USA
e-mail: burtonpeterj@icloud.com*

Dedicated to Benjy Weiss, with utmost admiration

ABSTRACT

We introduce the notion of soficity for locally compact groups and list a number of open problems.

* Supported in part by NSF grant DMS-1900386

Received July 4, 2021 and in revised form October 20, 2021

1. Introduction

A countable discrete group Γ is **sofic** if there exist maps $\sigma_i : \Gamma \rightarrow \text{sym}(V_i)$ (where each V_i is a finite set and $\text{sym}(V_i)$ is its permutation group) satisfying

$$\begin{aligned} 1 &= \lim_{i \rightarrow \infty} |V_i|^{-1} \{v \in V_i : \sigma_i(g)\sigma_i(h)v = \sigma_i(gh)v\} & \forall g, h \in \Gamma, \\ 1 &= \lim_{i \rightarrow \infty} |V_i|^{-1} \{v \in V_i : \sigma_i(g)v \neq v\} & \forall g \in \Gamma \setminus \{1_\Gamma\}. \end{aligned}$$

The first condition ensures that the maps behave asymptotically like homomorphisms and the second condition ensures that, asymptotically, every nontrivial element of Γ behaves like a fixed point-free map on V_i . In this sense, we can think of σ_i as providing a kind of approximation for the left-translation action of Γ on itself.

Sofic groups were defined implicitly by M. Gromov in [Gro99] where he proved they satisfy Gottschalk's surjunctivity conjecture. Benjy Weiss made the subject more accessible by simplifying the proof of Gromov's result in [Wei00] and giving sofic groups their name, which is derived from the Hebrew word *sofi* meaning finite. For an introduction to sofic groups, see [Pes08, PK13, CL15]. At the time of this writing, it is an open problem whether all discrete countable groups are sofic.

Locally compact sofic groups and their entropy theory were introduced by the first author and Sukhpreet Singh in Singh's unpublished 2016 thesis (available upon request to the first author). In this note, we give a new approach to locally compact sofic groups via partial actions and charts. Our main results are informally summarized as follows.

- Theorem 3.2. Every sofic group is unimodular.
- Theorem 3.3. Every unimodular lcsc amenable group is sofic.
- Theorem 3.5. A sequence of local G -spaces is a sofic approximation if and only if the essential injectivity radius of the sequence is infinite.
- Theorem 3.7. The new definition of sofic given in this paper generalizes the previous definitions for discrete countable groups.
- Theorem 3.9. If G admits a sofic lattice subgroup then G is sofic.
- Corollary 3.10. The following groups are sofic: semi-simple Lie groups (e.g., $\text{SL}(n, \mathbb{R})$, $\text{SO}(n, 1)$ etc.), the automorphism group of a regular tree.
- Proposition 3.11. If G is sofic and $H \leq G$ is an open subgroup then H is sofic.

This paper is organized as follows. Fix a locally compact second countable group G . In §2 we introduce local G -spaces as topological spaces with a partial homogeneous action of G . We derive metric and measure-theoretic properties of these spaces. In §3 we define sofic approximations to G as sequences of local G -spaces which, in a sense, approximate the action of G on itself by right-translations. We also prove the main results. The last section gives a series of open problems.

2. Local G -spaces and partial actions

2.1. LOCAL G -SPACES. We use the abbreviation lcsc to mean locally compact second countable. Let G be an lcsc group.

Definition 1: A **partial right-action** of G on a Hausdorff space M is a continuous map $\alpha : \text{dom}(\alpha) \rightarrow M$ where $\text{dom}(\alpha) \subset M \times G$ is open. We require the following axioms hold for all $p \in M$.

Axiom 1. $(p, 1_G) \in \text{dom}(\alpha)$ and $\alpha(p, 1_G) = p$.

Axiom 2. If $(p, g) \in \text{dom}(\alpha)$ then $(\alpha(p, g), g^{-1}) \in \text{dom}(\alpha)$ and $\alpha(\alpha(p, g), g^{-1}) = p$.

Axiom 3. If $(p, g), (\alpha(p, g), h), (p, gh) \in \text{dom}(\alpha)$ then $\alpha(p, gh) = \alpha(\alpha(p, g), h)$.

A partial action α is **homogeneous** if, in addition, it satisfies the following.

Axiom 4. For every $p \in M$ there is an open neighborhood O_p of 1_G in G such that $\{p\} \times O_p \subset \text{dom}(\alpha)$ and the restriction of $\alpha(p, \cdot)$ to $\{p\} \times O_p$ is a homeomorphism onto an open neighborhood of p in M .

Definition 2: A **local G -space** is a pair (M, α) where M is an lcsc space and α is a partial homogeneous right-action.

Notation 1: We will usually denote a local G -space by M (or V), leaving the action α implicit. To simplify notation, we write

$$p.g = \alpha(p, g).$$

If $K \subset M$, we will also write $K.g = \{\alpha(k, g) : k \in K\}$. In particular, $K.g$ is well-defined if and only if $K \times \{g\}$ is in the domain of the action α . Similarly, we write $p.O = \{\alpha(p, g) : g \in O\}$ if $\{p\} \times O \subset \text{dom}(\alpha)$.

Remark 1: By Axiom 3, $p.g_1.g_2 = p.g_1g_2$ when both sides are defined. However this does not imply that $p.g_1.g_2.g_3 = p.g_1g_2g_3$ even when both sides are defined. See Example 3.

LEMMA 2.1: *Let M be a local G -space. Let $g \in G$. Then $\alpha(\cdot, g)$ is injective (where it is defined).*

Proof. Suppose $\alpha(p, g) = \alpha(q, g)$ for some $p, q \in M$. In other words, $p.g = q.g$. By Axiom 2 of Definition 1, $p.g.g^{-1} = p$ and $q.g.g^{-1} = q$ and both are well-defined. Therefore, $p = q$ as required. ■

COROLLARY 2.2: *Let M be a local G -space. Let $K \subset M$ be compact, $g \in G$ and suppose $K.g$ is well-defined. Then the map $\alpha(\cdot, g)$ restricted to K is a homeomorphism onto its image.*

Proof. Because K is compact, it suffices to prove the map is continuous and injective. Continuity follows from joint continuity of α and injectivity follows from the previous lemma. ■

2.2. CHARTS.

Definition 3: Let (M, α) be a local G -space and $p \in M$. A **chart centered at p** is a homeomorphism $f_p : \text{dom}(f_p) \rightarrow \text{rng}(f_p)$ where $\text{dom}(f_p) \subset M$ is an open neighborhood of p , $\text{rng}(f_p)$ is an open neighborhood of the identity in G and $g = f_p(p.g)$ for all $g \in \text{rng}(f_p)$. By Axiom 4 of Definition 1, for every $p \in M$ there exists a chart centered at p .

We will show that the transition functions between two charts are locally given by left-translation in G .

Definition 4: Let $A, B \subset G$ be Borel sets. A map $\phi : A \rightarrow B$ is **locally left-translation** if there exists a decomposition $A = \bigsqcup_{i \in I} A_i$ into relatively open sets and $\{g_i\}_{i \in I} \subset G$ (for some countable index set I) such that $\phi(a) = g_i a$ for all $a \in A_i$. By relatively open we mean A_i is open in A .

PROPOSITION 2.3: *Let (M, α) be a local G -space. Let $p, q \in M$ and let f_p, f_q be charts centered at p, q respectively. Let*

$$\begin{aligned} A &= \{g \in \text{rng}(f_p) : p.g \in \text{dom}(f_p) \cap \text{dom}(f_q)\}, \\ B &= \{g \in \text{rng}(f_q) : q.g \in \text{dom}(f_p) \cap \text{dom}(f_q)\}. \end{aligned}$$

Then there is a map $\tau : A \rightarrow B$ which is locally left-translation such that

$$f_q(r) = \tau(f_p(r))$$

for all $r \in \text{dom}(f_p) \cap \text{dom}(f_q)$. Moreover, τ is bijective and left-Haar-measure-preserving.

Proof. Let $r \in \text{dom}(f_p) \cap \text{dom}(f_q)$. Let $g \in \text{rng}(f_p)$ be such that $p.g = r$. Let $h \in \text{rng}(f_q)$ be such that $q.h = r$. Choose a chart f_r centered at r . After choosing $\text{rng}(f_r)$ smaller if necessary we may assume $g \text{rng}(f_r) \subset \text{rng}(f_p)$ and $h \text{rng}(f_r) \subset \text{rng}(f_q)$. This implies $r.k = p.g.k = p.gk$ for all $k \in \text{rng}(f_r)$. Similarly, $r.k = q.h.k$.

Let $C_r = hg^{-1}$. We claim that $f_q(r.k) = C_r f_p(r.k)$ for all $k \in \text{rng}(f_r)$. This follows from

$$f_q(r.k) = f_q(p.hk) = hk = C_r gk = C_r f_p(p.gk) = C_r f_p(r.k).$$

Since $r.\text{rng}(f_r) = \text{dom}(f_r)$, this implies $f_q(s) = C_r f_p(s)$ for all $s \in \text{dom}(f_r)$.

Because G is lcsc, it follows that there are a countable index set I , and $\{r_i\}_{i \in I} \subset \text{dom}(f_p) \cap \text{dom}(f_q)$ such that

$$\text{dom}(f_p) \cap \text{dom}(f_q) \subset \bigcup_{i \in I} \text{dom}(f_{r_i})$$

where f_{r_i} are as above.

Let $A, B \subset G$ be as in the statement. Define $\tau : A \rightarrow B$ by $\tau(g) = C_{r_i}g$ if $p.g \in \text{dom}(f_{r_i})$. By the previous paragraph

$$\tau(f_p(r)) = f_q(r)$$

for all $r \in \text{dom}(f_p) \cap \text{dom}(f_q)$. In particular, since f_p and f_q are homeomorphisms, τ is well-defined, bijective and continuous. Since $g \mapsto \tau(g)g^{-1}$ is locally constant, τ is locally left-translation and therefore it is left-Haar-measure-preserving. ■

2.2.1. Measures. Here we show that a local G -space admits a canonical measure.

PROPOSITION 2.4 (The canonical measure): *Let (M, α) be a local G -space. Fix a left-Haar measure Haar_G on G . Then there exists a unique Radon measure vol_M on M satisfying the following. If $p \in M$, f_p is a chart centered at p and $K \subset \text{dom}(f_p)$ is Borel, then*

$$(1) \quad \text{vol}_M(K) = \text{Haar}_G(\{g \in \text{rng}(f_p) : p.g \in K\}) = \text{Haar}_G(f_p(K)).$$

Proof. Let $K \subset M$ be Borel. Because M is lcsc there exist a countable index set I , charts $\{f_i\}_{i \in I}$ with f_i centered at $p_i \in M$ such that $K \subset \bigcup_{i \in I} \text{dom}(f_i)$. Therefore, there is a Borel decomposition $K = \bigsqcup_{i \in I} K_i$ with $K_i \subset \text{dom}(f_i)$ for all i . We define

$$\text{vol}_M(K) = \sum_{i \in I} \text{Haar}_G(f_i(K_i)).$$

In order to show this is well-defined, suppose that J is a countable index set, $\{g_j\}_{j \in J}$ are charts, $K = \bigsqcup_{j \in J} L_j$ is a Borel decomposition and $L_j \subset \text{dom}(g_j)$. We must show that

$$\sum_{i \in I} \text{Haar}_G(f_i(K_i)) = \sum_{j \in J} \text{Haar}_G(g_j(L_j)).$$

By countable additivity, it suffices to show that

$$\text{Haar}_G(f_i(K_i \cap L_j)) = \text{Haar}_G(g_j(K_i \cap L_j))$$

for all $i \in I, j \in J$. This follows from Proposition 2.3.

This shows that vol_M is well-defined and satisfies $\text{vol}_M(K) = \text{Haar}_G(f_p(K))$ whenever f is a chart with $K \subset \text{dom}(f)$. Because f_p is a measure-preserving homeomorphism and G is locally compact, it follows that vol_M is a Radon measure. ■

Definition 5: Let $\delta : G \rightarrow \mathbb{R}_{>0}$ be the **modular function**. This means that if $S \subset G$ has a finite Haar measure, then $\text{Haar}_G(Sg) = \delta(g) \text{Haar}_G(S)$ where Haar_G is a left-Haar measure on G . The modular function is a homomorphism; G is **unimodular** if $\delta(g) = 1$ for all $g \in G$. This means that the Haar measure on G is both left and right G -invariant.

LEMMA 2.5 (Locally measure-preserving): *Let (M, α) be a local G -space and suppose $K \times \{g\} \subset \text{dom}(\alpha)$ for some measurable $K \subset M$ and $g \in G$. Then*

$$\text{vol}_M(K.g) = \delta(g) \text{vol}_M(K).$$

Proof. Let $p \in K$. Since $(p, g) \in \text{dom}(\alpha)$ and $\text{dom}(\alpha)$ is open in $M \times G$, there are open neighborhoods U, V of 1_G in G such that $\{p\} \times U \subset \text{dom}(\alpha)$, $\{p\} \times Vg \subset \text{dom}(\alpha)$. By Axiom 4 of Definition 1, there is an open neighborhood O_p of the identity in G such that $\alpha(p, \cdot)$ restricts to a homeomorphism from O_p to an open neighborhood $p.O_p$ of p in M . After intersecting O_p with U and V if necessary, we may assume $O_p \subset U \cap V$.

For every $h \in O_p$, if $p.h \in K$ then $p.h, p.h.g, p.hg$ are all well-defined (and therefore $p.h.g = p.hg$ by Axiom 3 of Definition 1).

Since M is lscs, there exists a countable subset $\{p_i\}_{i \in I} \subset K$ and open neighborhood $O_i = O_{p_i} \subset G$ of the identity as above such that

$$\bigcup_{i \in I} p_i.O_i \supset K.$$

So there is a measurable partition $K = \bigsqcup_{i=1}^{\infty} K_i$ such that for each i , $K_i \subset p_i.O_i$. Because the map $\alpha(\cdot, g) : K \rightarrow M$ is injective (by Lemma 2.1) and

$$\text{vol}_M(K.g) = \sum_{i=1}^{\infty} \text{vol}_M(K_i.g)$$

we may assume without loss of generality that $K \subset p.O_p$ for some $p \in K$.

We claim that $f_p(K.g) = f_p(K)g$ (and both sides are well-defined). To see this, let $k \in K \subset p.O_p$. Then there is $h \in O_p$ such that $k = p.h$. By choice of O_p , $p.h.g = p.hg$ and both are well-defined. Thus

$$f_p(k.g) = f_p(p.hg) = hg = f_p(p.h)g = f_p(k)g.$$

Since $k \in K$ is arbitrary, this proves the claim. So

$$\begin{aligned} \text{Haar}_M(K.g) &= \text{Haar}_G(f_p(K.g)) = \text{Haar}_G(f_p(K)g) \\ &= \delta(g) \text{Haar}_G(f_p(K)) = \delta(g) \text{vol}_M(K). \end{aligned}$$

The first and last equalities hold by definition of $\text{vol}_M(\cdot)$. ■

2.3. METRICS. In this section, we show that, given a left-invariant proper metric d_G on G , there is a canonical induced metric d_M on any local G -space M .

Definition 6: A local G -space M is **transitive** if for every $p, q \in M$ there exist elements $g_1, \dots, g_n \in G$ such that $q = p.g_1 \cdots g_n$. Equivalently, this means there exist charts $f_1, \dots, f_n$ such that $p \in \text{dom}(f_1), q \in \text{dom}(f_n)$ and $\text{dom}(f_i) \cap \text{dom}(f_{i+1}) \neq \emptyset$ for all i .

Definition 7: Let G be an lcsc group with a left-invariant proper metric d_G . Let M be a local G -space. Let $B(\rho) \subset G$ be the open ball of radius ρ centered at 1_G . For $p \in M$, let $\text{injrads}(M, p)$ be the supremum over all $\rho > 0$ such that

- (1) for any $g, h \in G$ with $g, h, gh \in B(\rho)$, $p.g.h = p.gh$ (in particular, both sides are well-defined);
- (2) the restriction of $\alpha(p, \cdot)$ to $B(\rho)$ is a homeomorphism onto its image.

This is the **injectivity radius** at p .

THEOREM 2.6: Let d_G be a left-invariant proper metric on G . Let M be a transitive left- G -space. Then there is a metric d_M on M satisfying the following local condition. For all $p \in M$, if $\rho = \text{injrads}(M, p)$ and $g \in B(\rho)$ then

$$d_M(p, p.g) = d_G(1_G, g).$$

Proof. Let $p, q \in M$. Define

$$d_M(p, q) = \inf \sum_{i=1}^{n-1} d_G(1_G, g_i)$$

where the infimum is over all sequences $g_1, \dots, g_n \in G$ such that $p.g_1 \cdots .g_n = q$.

Let $p \in M$. Let $\rho = \text{inrad}(M, p)$. To finish the proof it suffices to show that if $g \in B(\rho)$ then $d_M(p, p.g) = d_G(1_G, g)$.

So suppose $h_1, \dots, h_n \in G$ and $p.g = p.h_1 \cdots .h_n$. We must show that

$$(2) \quad d_G(1_G, g) \leq \sum_{i=1}^n d_G(1_G, h_i).$$

If for every i , $h_1 \cdots h_i \in B(\rho)$, then (2) is immediate by the triangle inequality. So we assume there is some index m such that $h_1 \cdots h_{m+1} \notin B(\rho)$. We may assume m is the smallest index for which this holds. Thus $h_1 \cdots h_i \in B(\rho)$ for all $i \leq m$ and $h_1 \cdots h_{m+1} \notin B(\rho)$.

Because d_G is left- G -invariant,

$$d_G(1_G, h_1) + d_G(1_G, h_2) = d_G(1_G, h_1) + d_G(h_1, h_1 h_2) \geq d_G(1_G, h_1 h_2).$$

By an inductive argument we obtain

$$\sum_{i=1}^{m+1} d_G(1_G, h_i) \geq d_G(1_G, h_1 \cdots h_{m+1}) \geq \rho > d_G(1_G, g).$$

This proves (2). $\blacksquare$

2.4. EXAMPLES.

Example 1: If $\Gamma < G$ is discrete, then $\Gamma \backslash G$ admits a local G -space structure as follows. Define $\alpha : \Gamma \backslash G \times G \rightarrow \Gamma \backslash G$ by $\alpha(\Gamma g, h) = \Gamma gh$.

Fix $g \in G$ and let $\pi : G \rightarrow \Gamma \backslash G$ be the map $\pi(h) = \Gamma gh$. Since Γ is discrete, π is a covering space map. In particular, it is a local homeomorphism. This implies Axiom 4 of Definition 1. The other Axioms are immediate.

Example 2: Let $M \subset G$ be an open subset. Let

$$\text{dom}(\alpha) = \{(p, g) \in M \times G : pg \in M\}$$

and define $\alpha : \text{dom}(\alpha) \rightarrow M$ by $\alpha(p, g) = pg$. It is immediate that (M, α) is a local G -space. Moreover, vol_M is the restriction of Haar_G to M .

Example 3: In this example, we show that it is possible for $p.g.h.k \neq p.ghk$ even when both sides are well-defined. Let $G = \mathbb{C}$ be the complex plane, as an additive group. Let M be a double cover of $\mathbb{C} \setminus \{0\}$. To be precise, let $M = (0, \infty) \times \mathbb{R}/4\pi\mathbb{Z}$. For $a, b \in \mathbb{R}/4\pi\mathbb{Z}$, define

$$|a - b| = \inf_{n \in \mathbb{Z}} |a' - b' - 4\pi n|$$

where $a', b' \in \mathbb{R}$ satisfy $a' = a \pmod{4\pi\mathbb{Z}}$ and $b' = b \pmod{4\pi\mathbb{Z}}$.

Let

$$\text{dom}(\alpha) = \{(r, \theta), se^{i\phi} - re^{i\theta}\} \in M \times \mathbb{C} : s > 0, |\phi - \theta| < 2\pi/3\}.$$

Define an action $\alpha : \text{dom}(\alpha) \rightarrow M$ by

$$\alpha((r, \theta), z) = (t, \phi)$$

where $te^{i\phi} = re^{i\theta} + z$ and $\phi \in \mathbb{R}/4\pi\mathbb{Z}$ is chosen to minimize $|\theta - \phi|$. Note there are only two different elements $\phi_1, \phi_2 \in \mathbb{R}/4\pi\mathbb{Z}$ that satisfy $te^{i\phi_j} = re^{i\theta} + z$ (for $j = 1, 2$) and $\phi_1 - \phi_2 = 2\pi \pmod{4\pi\mathbb{Z}}$. Because these elements are 2π apart, at most one of them can be within $2\pi/3$ of θ . Moreover, the definition of $\text{dom}(\alpha)$ shows that exactly one of these elements is within $2\pi/3$ of θ . So α is well-defined.

Next we check that (M, α) is a local G -space. Define

$$\tilde{f} : M \rightarrow \mathbb{C}, \quad \tilde{f}(r, \theta) = re^{i\theta}.$$

Then $\tilde{f}$ is a 2-1 covering map of $\mathbb{C} \setminus \{0\}$. Also

$$\tilde{f}(\alpha(p, z)) = \tilde{f}(p) + z.$$

Because $\tilde{f}$ is a local homeomorphism, α satisfies Axiom 4 of Definition 1.

In order to check Axiom 3, suppose $(r, \theta).z_1, (r, \theta).z_1.z_2$ and $(r, \theta).(z_1 + z_2)$ are all well-defined. We must show that $(r, \theta).z_1.z_2 = (r, \theta).(z_1 + z_2)$. Let $(r_1, \theta_1) = (r, \theta).z_1$, $(r_2, \theta_2) = (r, \theta).z_1.z_2$ and $(r_3, \theta_3) = (r, \theta).(z_1 + z_2)$.

The assumption that these are all well-defined implies $|\theta - \theta_1| < 2\pi/3$, $|\theta_1 - \theta_2| < 2\pi/3$ and $|\theta - \theta_3| < 2\pi/3$. The triangle inequality implies

$$|\theta_2 - \theta_3| < 2\pi.$$

On the other hand, the definition of α implies that either $\theta_2 = \theta_3$ or $\theta_2 = \theta_3 + 2\pi \pmod{4\pi}$. So we must have $\theta_2 = \theta_3 \pmod{4\pi}$. Therefore Axiom 3 holds. The other Axioms are immediate.

Next we show the existence of $p \in M$ and $g, h, k \in G$ such that $p.g.h.k \neq p.ghk$ even though both sides are well-defined. Let $p = (1, 0) \in M$. Let $g = i - 1$, $h = -i - 1$, $k = 1 - i$. Then

$$\tilde{f}(p) + g = 1 + (i - 1) = i.$$

So $p.g = (1, \pi/2)$. Also $\tilde{f}(p.g) + h = i + h = -1$. So $p.g.h = (1, \pi)$. Finally,

$$\tilde{f}(p.g.h) + k = -1 + k = -i.$$

So $p.g.h.k = (1, 3\pi/2)$. On the other hand, $g+h+k = -i-1$. So $p.ghk = (1, -\pi/2)$. Because $-\pi/2 \not\equiv 3\pi/2 \pmod{4\pi}$, we have $p.g.h.k \neq p.ghk$.

3. Sofic groups

3.1. DEFINITIONS.

Definition 8: Let $M = (M, \alpha)$ be a local G -space and let $U \subset G$ be open and pre-compact and let $\epsilon > 0$. Let $M[U] = M[\alpha, U]$ be the set of all $p \in M$ such that, if $g, h \in G$ are such that $g, h, gh \in U$, then $p.g.h = p.gh$ (in particular, both sides are well-defined). Moreover, we require that the map $g \mapsto \alpha(p, g)$ is a homeomorphism from U to an open neighborhood of p . We say M is a (U, ϵ) -**sofic approximation to G** if

$$\text{vol}_M(M) < \infty \quad \text{and} \quad \text{vol}_M(M[U]) \geq (1 - \epsilon) \text{vol}_M(M).$$

Definition 9: A **sofic approximation to G** is a sequence $\Sigma = (M_i)_{i=1}^\infty$ where M_i is a (U_i, ϵ_i) -sofic approximation such that the U_i are pre-compact open sets increasing to G and the sequence ϵ_i decreases to 0. We say G is **sofic** if it admits a sofic approximation.

The following lemma will be generally helpful.

LEMMA 3.1: *If $U \subset G$ is a pre-compact open neighborhood of the identity and $g \in U$, then*

$$M[U].g \subset M[U \cap g^{-1}U].$$

Proof. Let $p \in M[U]$, $g \in U$ and $h, k \in G$ be such that $h, k, hk \in U \cap g^{-1}U$. To show $p.g \in M[U \cap g^{-1}U]$, we first show $p.g.h.k = p.g.hk$.

Because $p \in M[U]$, and $g, h, gh \in U$ we have $p.g.h = p.gh$. Because $gh, k, ghk \in U$ we have $p.gh.k = p.ghk$. Combine these equalities to obtain $p.g.h.k = p.g.hk$.

Next define $\beta : U \cap g^{-1}U \rightarrow M$ by $\beta(k) = p.g.k$. We must show β is a homeomorphism onto an open neighborhood of $p.g$. Let $\gamma : U \rightarrow M$ be the map $\gamma(k) = p.k$. This map is a homeomorphism onto an open neighborhood of p since $p \in M[U]$. Moreover, $\beta(k) = p.g.k = p.gk = \gamma(gk)$. So β is the composition of left multiplication by g with γ . This implies it has the claimed properties. ■

3.2. UNIMODULARITY. The goal of this section is to prove:

THEOREM 3.2: *If G is sofic then G is unimodular.*

Proof. Fix a left-Haar measure Haar_G on G . Let $\epsilon > 0$, $g \in G$, and $U \subset G$ be a pre-compact open set containing $\{1_G, g, g^2\}$. Suppose M is a $(U \cup g^{-1}U, \epsilon)$ -sofic approximation.

By Lemma 3.1, $M[g^{-1}U].g^{-1} \subset M[U \cap g^{-1}U]$. Multiply by g on the right to obtain $M[U \cap g^{-1}U].g \supset M[g^{-1}U]$. Thus

$$\begin{aligned} \text{vol}_M(M[U \cap g^{-1}U].g) &\geq \text{vol}_M(M[g^{-1}U]) \geq \text{vol}_M(M[U \cup g^{-1}U]) \\ &\geq (1 - \epsilon) \text{vol}_M(M). \end{aligned}$$

Let $\delta : G \rightarrow \mathbb{R}$ be the modular function. Lemma 2.5 implies

$$\text{vol}_M(M[U \cap g^{-1}U].g) = \delta(g) \text{vol}_M(M[U \cap g^{-1}U]).$$

Combining this with the previous inequality, we obtain

$$\text{vol}_M(M) \geq \text{vol}_M(M[U \cap g^{-1}U]) \geq \delta(g)^{-1}(1 - \epsilon) \text{vol}_M(M).$$

Therefore, $\delta(g) \geq 1 - \epsilon$. Since this is true for every $\epsilon > 0$, $\delta(g) \geq 1$. However, $\delta : G \rightarrow \mathbb{R}_{>0}$ is a homomorphism. So $\delta(g^{-1}) = \delta(g)^{-1}$. Since also $\delta(g^{-1}) \geq 1$, we obtain $\delta(g) = 1$. Because $g \in G$ is arbitrary, G must be unimodular. ■

3.3. AMENABLE GROUPS.

THEOREM 3.3: *Every unimodular amenable lcsc group is sofic.*

Definition 10: A locally compact group G is **amenable** if for every left-Haar measure Haar_G , there exists a sequence $\{F_i\}_{i=1}^\infty$ of measurable sets with finite positive measure such that for every compact $K \subset G$,

$$(3) \quad \lim_{i \rightarrow \infty} \frac{\text{Haar}_G(KF_i)}{\text{Haar}_G(F_i)} = 1.$$

Such a sequence is called a **left-Følner sequence**.

LEMMA 3.4: If G is amenable and unimodular, then there exists a sequence $\{\tilde{F}_i\}_{i=1}^\infty$ of finite measure subsets such that each $\tilde{F}_i$ is pre-compact, open and for every compact $K \subset G$,

$$(4) \quad \lim_{i \rightarrow \infty} \frac{\text{Haar}_G(\{p \in \tilde{F}_i : pK \subset \tilde{F}_i\})}{\text{Haar}_G(\tilde{F}_i)} = 1.$$

Proof. Let $\{F_i\}_{i=1}^\infty$ be a left-Følner sequence. After perturbing slightly, we may assume each F_i is pre-compact. Indeed, since F_i has finite measure, there is a subset $L_i \subset F_i$ which is pre-compact such that $\lim_{i \rightarrow \infty} \frac{\text{Haar}_G(F_i \setminus L_i)}{\text{Haar}_G(F_i)} = 0$. Then $\{L_i\}_{i=1}^\infty$ is left-Følner. So we can replace F_i with L_i .

Let $\{K_i\}_{i=1}^\infty$ be an increasing sequence of pre-compact open subsets with $G = \bigcup_i K_i$. We choose K_i to grow so slowly that

$$(5) \quad \lim_{i \rightarrow \infty} \frac{\text{Haar}_G(K_i F_i)}{\text{Haar}_G(F_i)} = 1.$$

Let $F'_i = K_i F_i$. Then F'_i is pre-compact and open. Because

$$\lim_{i \rightarrow \infty} \frac{\text{Haar}_G(F'_i \triangle F_i)}{\text{Haar}_G(F_i)} = 0,$$

we have that $\{F'_i\}_{i=1}^\infty$ is left-Følner.

For any compact $K \subset G$, there exists i with $K \subset K_i$. Therefore,

$$(6) \quad \begin{aligned} \liminf_{i \rightarrow \infty} \frac{\text{Haar}_G(\{p \in F'_i : Kp \subset F'_i\})}{\text{Haar}_G(F'_i)} &\geq \liminf_{i \rightarrow \infty} \frac{\text{Haar}_G(\{p \in F'_i : K_i p \subset F'_i\})}{\text{Haar}_G(F'_i)} \\ &\geq \liminf_{i \rightarrow \infty} \frac{\text{Haar}_G(F_i)}{\text{Haar}_G(F'_i)} = 1. \end{aligned}$$

Now let $\tilde{F}_i = (F'_i)^{-1}$. Note that $\tilde{F}_i$ is pre-compact and open since F'_i is. Let $K \subset G$ be compact. Suppose $p \in \tilde{F}_i$ satisfies $pK \subset \tilde{F}_i$. Then $K^{-1}p^{-1} \subset F'_i$. The converse is also true. Thus

$$\{p \in \tilde{F}_i : pK \subset \tilde{F}_i\}^{-1} = \{p \in F'_i : K^{-1}p \subset F'_i\}.$$

Because G is unimodular, $\text{Haar}_G(E) = \text{Haar}_G(E^{-1})$ for any measurable $E \subset G$. Thus

$$\liminf_{i \rightarrow \infty} \frac{\text{Haar}_G(\{p \in \tilde{F}_i : pK \subset \tilde{F}_i\})}{\text{Haar}_G(\tilde{F}_i)} = \liminf_{i \rightarrow \infty} \frac{\text{Haar}_G(\{p \in F'_i : K^{-1}p \subset F'_i\})}{\text{Haar}_G(F'_i)} = 1$$

by (6). ■

Proof of Theorem 3.3. Let G be a unimodular amenable lcsc group. Let $\{F_i\}_{i=1}^\infty \subset G$ be a sequence as in Lemma 3.4. As in Example 2 we may regard F_i as a local G -space with vol_{F_i} equal to the restriction of Haar_G to F_i where (as always in this paper) Haar_G is a left-Haar measure. Let $U \subset G$ be pre-compact and $\epsilon > 0$. Then $F_i[U]$ consists of all $p \in F_i$ with $pU \subset F_i$. By Lemma 3.4,

$$\begin{aligned} \text{vol}_{F_i}(F_i[U]) &= \text{Haar}_G(\{p \in F_i : pU \subset F_i\}) \\ &\geq (1 - \epsilon) \text{Haar}_G(F_i) = (1 - \epsilon) \text{vol}_{F_i}(F_i) \end{aligned}$$

for all sufficiently large i . Thus $\{F_i\}_{i=1}^\infty$ is a sofic approximation and G is sofic. ■

3.4. THE METRIC APPROACH TO SOFICITY. For this section, fix an lcsc group G with a left-invariant proper metric d_G .

THEOREM 3.5: *Let $(M_i)_{i=1}^\infty$ be a sequence of local G -spaces. Then the following are equivalent.*

- (1) $\{M_i\}_{i=1}^\infty$ is a sofic approximation to G .
- (2) For every $\rho > 0$

$$\lim_{i \rightarrow \infty} \frac{\text{vol}_{M_i}(\{p \in M_i : \text{injrads}(M_i, p) > \rho\})}{\text{vol}_{M_i}(M_i)} = 1,$$

where injectivity radius is defined in Definition 7.

Proof. Let $U = B(\rho)$, which is the open ball of radius ρ centered at the identity in G . Then $\text{injrads}(M_i, p) \geq \rho$ if and only if $p \in M_i[U]$. So the theorem follows immediately from the definition of sofic approximation. ■

Remark 2: It is possible to define Benjamini–Schramm (BS) convergence for measured metric spaces via the pointed Gromov–Hausdorff–Prokhorov topology [Bow15]. Theorem 3.5 implies that if a sequence of local G -spaces equipped with metrics given by Theorem 2.6 is a sofic approximation to G , then it BS-converges to G . The converse is not true. For example, there are non-isomorphic finitely generated groups G_1, G_2 with Cayley graphs which are isomorphic as unlabeled graphs. Precisely, there are generating sets S_i for G_i ($i = 1, 2$) so that with the corresponding word-metrics there is an isometry $\phi : G_1 \rightarrow G_2$ that maps the identity to the identity and S_1 bijectively onto S_2 .

Suppose $\{M_i\}$ is a sofic approximation to G_1 . Then $\{M_i\}$ BS-converges to G_1 . However, we can think of $\{M_i\}$ as a local G_2 -space in the following way. For $p \in M_i$ and $g \in S_2 \cup \{1_{G_2}\}$, let $p.g$ equal $p.\phi^{-1}(g)$ (if and only if the latter is defined). Because G_2 is finitely generated, this makes each M_i into a local G_2 -space. With this structure, $\{M_i\}$ is not a sofic approximation to G_2 (since G_1 and G_2 are non-isomorphic). But it does BS-converge to G_2 (since G_1 and G_2 are isometric).

3.4.1. Discrete sofic groups. In this section we show that our new definition of sofic agrees with the standard definition if G is a countable discrete group.

Definition 11: Let G be a countable group. Let $\sigma : G \rightarrow \text{sym}(V)$ (where V is a finite set) be a set map. For $U \subset G$, let $V[\sigma, U] \subset V$ be the set of all p such that

- (1) $\sigma(g)\sigma(h)p = \sigma(gh)p$ for all $g, h \in U$;
- (2) $\sigma(g)p \neq \sigma(h)p$ for all $g, h \in U$ with $g \neq h$.

Then σ is a **discrete (U, ϵ) -sofic approximation** if

$$\#V[\sigma, U] \geq (1 - \epsilon)\#V.$$

We call G **sofic as a discrete group** if for every finite $U \subset G$ and $\epsilon > 0$ there exists a (U, ϵ) -sofic approximation to G .

We first show that if G is sofic as a discrete group, then it admits a sofic approximation that is exact with respect to inverses and the identity.

LEMMA 3.6: *If G is sofic as a discrete group, then for every finite $U \subset G$ and $\epsilon > 0$ there exists a discrete (U, ϵ) -sofic approximation $\sigma : G \rightarrow \text{sym}(V)$ such that $\sigma(1_G)$ is the identity and $\sigma(g^{-1}) = \sigma(g)^{-1}$ for all $g \in G$.*

Proof. Without loss of generality, we may assume $U = U^{-1}$ and $1_G \in U$.

Let $\sigma : G \rightarrow \text{sym}(V)$ be a discrete $(U^2, \epsilon|U|^{-1})$ -sofic approximation to G . Let $D \subset G$ be the set of order 2 elements. Let $H \subset G$ be a subset such that for every $g \in G$ there is a unique element in the intersection $H \cap \{g, g^{-1}\}$.

If $v \in V[\sigma, U^2]$ then

$$\sigma(1_G)^2 v = \sigma(1_G) v.$$

Thus $\sigma(1_G)v = v$. If also $g \in D \cap U^2$ is nontrivial then

$$\sigma(g)^2 v = \sigma(g^2)v = \sigma(1_G)v = v, \quad \sigma(g)v \neq \sigma(1_G)v = v.$$

So there exists an element $\sigma'(g) \in \text{sym}(V)$ with order 2 such that $\sigma'(g)v = \sigma(g)v$ for all $v \in V[\sigma, U^2]$. This defines $\sigma'(g)$ for all $g \in D \cap U^2$. Also define

$$\sigma'(g) = \begin{cases} \text{identity,} & g \in \{1_G\} \cup (D \setminus U^2), \\ \sigma(g), & g \in H \setminus (D \cup \{1_G\}), \\ \sigma(g^{-1})^{-1}. & g^{-1} \in H \setminus (D \cup \{1_G\}). \end{cases}$$

This defines σ' on all of G . Note that $\sigma'(1_G)$ is the identity and $\sigma'(g)^{-1} = \sigma'(g^{-1})$ for all $g \in G$. Moreover, $\sigma'(g)v \in \{\sigma(g)v, \sigma(g^{-1})^{-1}v\}$ for all $v \in V[\sigma, U^2]$ and $g \in U^2$.

It now suffices to show σ' is a discrete (U, ϵ) -sofic approximation. To prove this, let

$$W = \{v \in V : \sigma(g)v \in V[\sigma, U^2] \ \forall g \in U\}.$$

We claim that $W \subset V[\sigma', U]$.

To prove this we observe: if $v \in V[\sigma, U^2]$ and $g \in U^2$ then

$$\sigma'(g)v = \sigma(g)v = \sigma(g^{-1})^{-1}v.$$

Indeed

$$\sigma(g^{-1})\sigma(g)v = \sigma(1_G)v = v.$$

Thus $\sigma(g)v = \sigma(g^{-1})^{-1}v$. Since $\sigma'(g)v \in \{\sigma(g)v, \sigma(g^{-1})^{-1}v\}$, it follows that $\sigma'(g)v = \sigma(g)v$. This proves the claim.

Now let $w \in W$ and $g, h \in U$. By definition of W , $\sigma(h)w \in V[\sigma, U^2]$. Since $gh \in U^2$,

$$\sigma'(g)\sigma'(h)w = \sigma'(g)(\sigma(h)w) = \sigma(g)\sigma(h)w = \sigma(gh)w = \sigma'(gh)w.$$

Moreover, if $g \neq h$ then

$$\sigma'(g)w = \sigma(g)w \neq \sigma(h)w = \sigma'(h)w.$$

This shows that $W \subset V[\sigma', U]$ as claimed.

By definition,

$$W = \bigcap_{g \in U} \sigma(g)^{-1}V[\sigma, U^2].$$

Since each $\sigma(g)$ is a permutation and $|V(\sigma, U^2)| \geq (1 - \epsilon|U|^{-1})|V|$, this implies $|W| \geq (1 - \epsilon)|V|$. Thus

$$\#V[\sigma', U] \geq \#W \geq (1 - \epsilon)\#V.$$

This shows that σ' is a discrete (U, ϵ) -sofic approximation. ■

THEOREM 3.7: *Let G be a discrete countable group. Then G is sofic as a discrete group if and only if G is sofic in the sense of Definition 8.*

Proof. Suppose G is sofic as a discrete group. Let d_G be a proper left-invariant metric on G . Recall that $B(\rho)$ denotes the open radius ρ ball centered at the identity in G . It suffices to show that for every radius $\rho > 0$ and $\epsilon > 0$, there is a $(B(\rho), \epsilon)$ -sofic approximation to G (in the sense of Definition 8).

Let $\sigma : G \rightarrow \text{sym}(V)$ be a discrete $(B(2\rho), \epsilon)$ -sofic approximation to G . By Lemma 3.6, we may assume $\sigma(1_G)$ is the identity permutation and $\sigma(g^{-1}) = \sigma(g)^{-1}$ for all $g \in G$.

For $p \in V$, define $\text{injrads}(\sigma, p)$ to be the supremum of $\eta > 0$ such that

- (1) $\sigma(gh)^{-1}p = \sigma(h)^{-1}\sigma(g)^{-1}p$ for all $g, h \in B(\eta)$;
- (2) $\sigma(g)^{-1}p \neq \sigma(h)^{-1}p$ if $g, h \in B(\eta)$ with $g \neq h$.

So $p \in V[\sigma, B(\rho)]$ if and only if $\text{injrads}(\sigma, p) \geq \rho$.

Let $\text{dom}(\alpha)$ be the set of all (p, g) in $V \times G$ such that either $\text{injrads}(\sigma, p) > d_G(1_G, g)$ or $\text{injrads}(\sigma, \sigma(g)^{-1}p) > d_G(1_G, g)$. Define $\alpha : \text{dom}(\alpha) \rightarrow V$ by

$$\alpha(p, g) = \sigma(g)^{-1}p.$$

We claim that (V, α) is a local G -space. It is immediate that Axioms 1, 2 and 4 of Definition 1 hold.

To verify Axiom 3, suppose $(p, g), (\alpha(p, g), h), (p, gh) \in \text{dom}(\alpha)$. Then

$$\begin{aligned}\alpha(\alpha(p, g), h) &= \sigma(h)^{-1}\alpha(p, g) = \sigma(h)^{-1}\sigma(g)^{-1}p, \\ \alpha(p, gh) &= \sigma(gh)^{-1}p.\end{aligned}$$

So we must show that

$$(7) \quad \sigma(h)^{-1}\sigma(g)^{-1}p = \sigma(gh)^{-1}p.$$

Because $(p, g), (\alpha(p, g), h), (p, gh) \in \text{dom}(\alpha)$,

either $\text{injrads}(\sigma, p) > d_G(1_G, g)$ or $\text{injrads}(\sigma, \sigma(g)^{-1}p) > d_G(1_G, g)$,
 either $\text{injrads}(\sigma, \sigma(g)^{-1}p) > d_G(1_G, h)$ or $\text{injrads}(\sigma, \sigma(h)^{-1}\sigma(g)^{-1}p) > d_G(1_G, h)$,
 either $\text{injrads}(\sigma, p) > d_G(1_G, gh)$ or $\text{injrads}(\sigma, \sigma(gh)^{-1}p) > d_G(1_G, gh)$.

Choose $q \in \{p, \sigma(g)^{-1}p, \sigma(h)^{-1}\sigma(g)^{-1}p, \sigma(gh)^{-1}p\}$ to maximize the injectivity radius $\text{injrads}(\sigma, q)$. Note that $\text{injrads}(\sigma, q) > d_G(1_G, f)$ for all $f \in \{g, h, gh\}$.

If $q = p$ then (7) follows by definition of $\text{injr}(\sigma, p)$. If $q = \sigma(g)^{-1}p$ then

$$\sigma(gh)\sigma(h)^{-1}\sigma(g)^{-1}p = \sigma(g)\sigma(g)^{-1}p = p$$

by definition of $\text{injr}(\sigma, \sigma(g)^{-1}p)$ and the assumption $\sigma(h)^{-1} = \sigma(h^{-1})$. This also implies (7) by multiplying both sides by $\sigma(gh)^{-1}$. The other cases are similar. This verifies Axiom 3.

It is immediate that $V[\sigma, B(2\rho)] \subset V[\alpha, B(\rho)]$. So

$$\#V[\alpha, B(\rho)] \geq \#V[\sigma, B(2\rho)] \geq (1 - \epsilon)\#V.$$

This proves (V, α) is $(B(\rho), \epsilon)$ -sofic in the sense of Definition 8. Since ρ, ϵ are arbitrary, this proves G is sofic.

Now suppose G is sofic in the sense of Definition 8. Let $\rho > 0$ be a radius and $\epsilon > 0$. It suffices to show there exists $\sigma : G \rightarrow V$ such that σ is a discrete $(B(\rho), \epsilon)$ -sofic approximation.

By Theorem 3.5, there exists a sofic approximation $M = (M, \alpha)$ to G such that

$$(8) \quad \frac{\text{vol}_M(\{p \in M : \text{injr}(M, p) > 3\rho\})}{\text{vol}_M(M)} > 1 - \epsilon|B(\rho)|^{-1}.$$

Because G is discrete, we choose Haar_G to be counting measure on G . Therefore, vol_M is counting measure on M . In particular, M is finite.

By Lemma 2.1, for $g \in G$, the map $\alpha(\cdot, g)$ is injective on its domain. Therefore, there exists a permutation $\sigma(g^{-1}) \in \text{sym}(M)$ that agrees with $\alpha(\cdot, g)$ on its domain. So there is a map $\sigma : G \rightarrow \text{sym}(M)$ such that $\sigma(g)p = \alpha(p, g^{-1})$ for all $(p, g^{-1}) \in \text{dom}(\alpha)$. We claim that (M, σ) is a $(B(\rho), \epsilon)$ -discrete sofic approximation.

Let D be the set of all $p \in M$ such that $\text{injr}(M, p, g) > 2\rho$ for all $g \in B(\rho)$. By (8), $|D| \geq (1 - \epsilon)|M|$. So it suffices to show that $D \subset M[\sigma, B(\rho)]$.

Let $p \in D$ and $g, h \in B(\rho)$. By the triangle inequality, $gh \in B(2\rho)$. Since $\text{injr}(M, p) > 2\rho$,

$$\sigma(gh)p = \sigma(g)\sigma(h)p.$$

Moreover, if $g \neq h$ then

$$\sigma(g)p \neq \sigma(h)p.$$

These two claims imply $D \subset M[\sigma, B(\rho)]$ and so complete the proof. $\blacksquare$

3.5. STABILITY OF SOFICITY UNDER CONSTRUCTIONS.

3.5.1. *Inducing from a subgroup.* In this section we prove that if G contains a sofic lattice $\Gamma \leq G$ then G is sofic as well. Moreover, if $\Sigma = \{V_i\}_{i \in \mathbb{N}}$ is a sofic approximation to Γ then there is an induced sofic approximation

$$\text{Ind}_\Gamma^G(\Sigma) = \{\text{Ind}_\Gamma^G(V_i)\}_{i \in \mathbb{N}}$$

to G . This is similar to the way that an action or representation of Γ can be induced to G . It depends a priori on a choice of fundamental domain $\Delta \subset G$. We will choose Δ to have some additional properties that will make it easier to prove that the induced map really is a sofic approximation. It seems likely that different fundamental domains lead to essentially the same induced sofic approximation but we make no effort to prove it. The next lemma gives a ‘nice’ fundamental domain.

Definition 12: A **lattice** is a subgroup $\Gamma \leq G$ such that, with the induced topology, Γ is discrete and $\Gamma \backslash G$ has a finite G -invariant Borel measure. A **fundamental domain** for Γ is a Borel set $\Delta \subset G$ such that $\bigsqcup_{g \in \Gamma} g\Delta$ is a partition of G .

Definition 13: Let X be a topological space. A collection $\{Y_i\}_{i \in I}$ of subsets $Y_i \subset X$ is **locally finite** if for every $x \in X$ there exists an open neighborhood O of x in X such that

$$\#\{i \in I : Y_i \cap O \neq \emptyset\} < \infty.$$

This condition implies that for every compact $K \subset X$,

$$\#\{i \in I : Y_i \cap K \neq \emptyset\} < \infty.$$

LEMMA 3.8: *Let $\Gamma \leq G$ be a lattice. Then there exists a fundamental domain Δ for Γ such that the collection $\{g\Delta : g \in \Gamma\}$ is locally finite.*

Proof. Let $\pi : G \rightarrow \Gamma \backslash G$ be the quotient map. Because Γ is discrete, for every $g \in G$ there exists an open pre-compact neighborhood $\tilde{O}$ of g such that the restriction of π to $\tilde{O}$ is injective. So there exists an open cover $\{O_i\}_{i \in I}$ of $\Gamma \backslash G$ such that for each $i \in I$, there is a pre-compact open set $\tilde{O}_i \subset G$ such that π restricted to $\tilde{O}_i$ is a homeomorphism onto O_i . Because G is locally compact, after passing to a sub-cover if necessary, we may assume $\{O_i\}_{i \in I}$ is locally finite. This implies that the cover $\{g\tilde{O}_i\}_{g \in \Gamma, i \in I}$ of G is also locally finite. In fact, because $\tilde{O}_i$ is pre-compact and Γ is discrete, for any compact $K \subset G$

there are only finitely many $g \in \Gamma$ with $g\tilde{O}_i \cap K \neq \emptyset$. On the other hand, there are only finitely many indices $i \in I$ with $\Gamma\tilde{O}_i \cap K \neq \emptyset$ because this condition implies $\Gamma K \cap O_i \neq \emptyset$ and $\{O_i\}_{i \in I}$ is locally finite.

Since G is second countable, the index set I is at most countable, so we may assume that $I \subset \mathbb{N}$. Define

$$\Delta = \bigcup_{i \in I} \tilde{O}_i \setminus \left(\bigcup_{g \in \Gamma} \bigcup_{j < i} g\tilde{O}_j \right).$$

Then Δ is a Borel fundamental domain.

Let $x \in G$. Let $\tilde{O}_x \subset G$ be a pre-compact open neighborhood of x such that the restriction of π to $\tilde{O}_x$ is a homeomorphism onto its image. Because $\{g\tilde{O}_i\}_{g \in \Gamma, i \in I}$ of G is locally finite, there are only finitely many pairs $(g, i) \in \Gamma \times I$ such that $g\tilde{O}_i \cap \tilde{O}_x \neq \emptyset$.

We claim that $\tilde{O}_x$ intersects at most finitely many Γ -translates of Δ . To see this, let $g \in \Gamma$ and suppose $g\Delta \cap \tilde{O}_x \neq \emptyset$. Since $\Delta \subset \bigcup_{i \in I} \tilde{O}_i$, this implies the existence of $i \in I$ with $g\tilde{O}_i \cap \tilde{O}_x \neq \emptyset$. So by the previous paragraph, there are only finitely many $g \in \Gamma$ with $g\Delta \cap \tilde{O}_x \neq \emptyset$. This finishes the lemma. ■

THEOREM 3.9: *Let $\Gamma \leq G$ be a lattice where G is an lcsc group. If Γ is sofic then G is sofic. Moreover, for every sofic approximation $\Sigma = \{V_i\}_{i \in I}$ to Γ there is an induced sofic approximation $\text{Ind}_\Gamma^G(\Sigma) = \{\text{Ind}_\Gamma^G(V_i)\}_{i \in I}$ to G determined only by Σ and a choice of fundamental domain Δ for Γ satisfying Lemma 3.8.*

Proof. Let Δ be a fundamental domain for Γ satisfying Lemma 3.8. Define a section $\sigma : \Gamma \backslash G \rightarrow \Delta$ by $\Gamma\sigma(\Gamma g) = \Gamma g$. This is well-defined because Δ is a fundamental domain. Define $c : \Gamma \backslash G \times G \rightarrow \Gamma$ by

$$c(\Gamma h, g) = \sigma(\Gamma h)g\sigma(\Gamma hg)^{-1}.$$

Then c satisfies the cocycle equation $c(\Gamma h, g)c(\Gamma hg, k) = c(\Gamma h, gk)$ for any g, h, k .

If $K \subset G$ then we write

$$c(\Gamma h, K) = \{c(\Gamma h, k) : k \in K\} \subset \Gamma.$$

We claim that if K is compact then $c(\Gamma h, K)$ is finite. To see this, observe that $c(\Gamma h, k)\Delta \cap \sigma(\Gamma h)K \neq \emptyset$ (for all $k \in K$). In fact,

$$\sigma(\Gamma h)k = c(\Gamma h, k)\sigma(\Gamma hk) \in c(\Gamma h, k)\Delta \cap \sigma(\Gamma h)K.$$

Because the collection $\{g\Delta\}_{g \in \Gamma}$ is locally finite and $\sigma(\Gamma h)K$ is compact, this implies the claim: $c(\Gamma h, K)$ is finite.

Let $\Sigma = \{V_i\}_{i \in I}$ be a sofic approximation to Γ (in the sense of Definition 8). We will denote the partial action of Γ on V_i by $v.g$ for $v \in V_i, g \in \Gamma$ whenever this is well-defined.

For a warm-up exercise, let's handle the special case in which there is a finite-index subgroup $H_i \leq \Gamma$, $V_i = H_i \backslash \Gamma$ and the partial action of Γ on V_i is the usual action by right-translation. In that case, G acts on $V_i \times \Gamma \backslash G$ by

$$(v, \Gamma h).g = (v.c(\Gamma h, g), \Gamma hg).$$

Define a G -equivariant Borel isomorphism

$$\Phi : V_i \times \Gamma \backslash G \rightarrow H_i \backslash G, \quad \Phi(H_i g, \Gamma h) = H_i g \sigma(\Gamma h).$$

The action of G on $V_i \times \Gamma \backslash G$ is not continuous with respect to the product topology. So we re-topologize $V_i \times \Gamma \backslash G$ by pulling back the topology on $H_i \backslash G$.

Because the quotient map $G \rightarrow H_i \backslash G$ is a covering space map, the topology on $H_i \backslash G$ is such that the open sets in $H_i \backslash G$ are images of open sets in G . So the new topology on $V_i \times \Gamma \backslash G$ has a basis of open sets given by sets of the form $p.O$ where $p \in V_i \times \Gamma \backslash G$ and $O \subset G$ is an open neighborhood of the identity.

Now for the general case. For $(v, \Gamma h) \in V_i \times \Gamma \backslash G$ and $g \in G$ we write

$$\alpha'((v, \Gamma h), g) = (v, \Gamma h).g := (v.c(\Gamma h, g), \Gamma hg)$$

whenever this is well-defined. Observe that by Axiom 3 (applied to V_i), if $p \in V_i \times \Gamma \backslash G$ and $g, h \in G$ are such that $p.g, p.g.h$ and $p.gh$ are all well-defined, then

$$p.g.h = p.gh.$$

We say that a subset $O \subset G$ is **good** for a point $p \in V_i \times \Gamma \backslash G$ if

- O is an open neighborhood of the identity,
- for every $h_1, h_2 \in O$, $p.h_1$ and $p.h_1.h_1^{-1}h_2$ are well-defined, and
- the map which sends $g \in O$ to $p.g$ is injective.

Note that if O is good for p , then every open subset of O containing the identity is also good for p . Let $M_i \subset V_i \times \Gamma \backslash G$ be the set of all points p for which there exists a good set $O \subset G$. If $p \in M_i$ and O is good for p then we write

$$p.O = \{p.g : g \in O\}.$$

CLAIM 1: If O is good for p and $g \in O$ then $g^{-1}O$ is good for $p.g$. Moreover,

$$p.O = p.g.g^{-1}O.$$

Proof of Claim 1. Let $h_1, h_2 \in g^{-1}O$. We must show that $p.g.h_1$ and $p.g.h_1.h_1^{-1}h_2$ are well-defined.

Because O is good for p and $g, gh_1 \in O$, it follows that

$$p.g.h_1 = p.g.g^{-1}gh_1$$

is well-defined. Moreover, $p.g$ and $p.gh_1$ are well-defined. Therefore,

$$p.gh_1 = p.g.h_1.$$

So it now suffices to show $p.gh_1.h_1^{-1}h_2$ is well-defined. But this follows from goodness of O because $gh_1, gh_2 \in O$ and $p.gh_1.(gh_1)^{-1}gh_2 = p.gh_1.h_1^{-1}h_2$. This proves the first statement.

Note that if O is good for p and $g, h \in O$ then $p.g.g^{-1}h = p.h$ (because both sides are well-defined). The second statement follows. ■

Claim 1 implies that $p.O \subset M_i$.

CLAIM 2: The collection of subsets of M_i of the form $p.O$ (where $p \in M_i$ and O is good for p) is a base for a topology on M_i .

Proof of Claim 2. It suffices to show that if O is good for p and U is good for q and $r \in p.O \cap q.U$, then there is a good set $W \subset G$ for r such that

$$r.W \subset p.O \cap q.U.$$

Let $r = p.g = q.h$ for some $g \in O, h \in U$. Then $g^{-1}O$ and $h^{-1}U$ are good for r by Claim 1. It follows that $W = g^{-1}O \cap h^{-1}U$ is also good for r . Moreover,

$$r.W = p.g.(g^{-1}O \cap h^{-1}U) \subset p.O.$$

Similarly, $r.W \subset q.U$. ■

From now on, we consider M_i with the topology induced by sets of the form $p.O$ as above. Claim 2 implies that if O is good for p , then the map $g \mapsto p.g$ from O into M_i is a homeomorphism onto an open subset of M_i . In particular, M_i is locally compact.

We claim M_i is second countable. Because G is second countable, there is a collection $\{\Gamma g_j, O_j\}_{j \in J}$ such that

- each $O_j \subset G$ is a pre-compact open neighborhood of the identity,
- the map from O_j to $\Gamma \backslash G$ given by $h \mapsto \Gamma g_j h$ is injective,
- the sets $\{\Gamma g_j O_j\}_{j \in J}$ form a base for the topology on $\Gamma \backslash G$,
- J is countable.

Given $v \in V_i$, $k \in \Gamma$, $j \in J$ define

$$B(v, k, j) = \{(v.kc(\Gamma g_j, s), \Gamma g_j s) : s \in O_j\} \subset V_i \times \Gamma \backslash G$$

whenever this is well-defined. Because V_i, Γ and J are countable, the collection of sets of this form is countable.

Now suppose $p \in M_i$ and $U \subset G$ is good for p . Let $p = (v, \Gamma h)$. Because $\{\Gamma g_j O_j\}_{j \in J}$ is a base and $\Gamma h U$ is open in $\Gamma \backslash G$, there exists $j \in J$ and an open subset $U' \subset U$ with $1_G \in U'$ such that $\Gamma h U' = \Gamma g_j O_j$. Since U' is good for p , $p.U'$ is open in M_i . We claim $p.U' = B(v, k, j)$ for some $k \in \Gamma$. To see this, note that $\Gamma h U' = \Gamma g_j O_j$ implies the existence of $\gamma \in \Gamma$ such that

$$U' = h^{-1} \gamma g_j O_j.$$

If $u \in U'$ then there is an $s \in O_j$ such that $u = h^{-1} \gamma g_j s$. So

$$\begin{aligned} p.u &= (v.c(\Gamma h, u), \Gamma h u) = (v.c(\Gamma h, h^{-1} \gamma g_j s), \Gamma h h^{-1} \gamma g_j s) \\ &= (v.c(\Gamma h, h^{-1} \gamma g_j) c(\Gamma g_j, s), \Gamma g_j s). \end{aligned}$$

Thus $p.U' \subset B(v, k, j)$ where $k = c(\Gamma h, h^{-1} \gamma g_j)$. The reverse inclusion is similar. Thus, the collection of sets of the form $p.U'$ is a countable base for the topology on M_i .

Let $\text{dom}(\alpha)$ be the set of all $(p, g) \in M_i \times G$ such that there is an open neighborhood of (p, g) in $M_i \times G$ on which α' is well-defined. Let α be the restriction of α' to $\text{dom}(\alpha)$.

CLAIM 3: α is continuous. Moreover, if $(q, f) \in \text{dom}(\alpha)$, then there is an open neighborhood W of (q, f) such that $\alpha(W)$ is open in M_i .

Proof of Claim 3. By Claim 2, it suffices to prove: if $p \in M_i$ and $O \subset G$ is good for p then $\alpha^{-1}(p.O)$ is open in $M_i \times G$. So let $(q, f) \in \alpha^{-1}(p.O)$. Let $g \in O$ be such that $p.g = q.f$. By Claim 1, $g^{-1}O$ is good for $p.g = q.f$.

Let $U_1 \subset G$ be a set which is good for q so that

$$(q.U_1 \times \{f\}) \cup (\{q\} \times U_1 f) \subset \text{dom}(\alpha)$$

and $f^{-1}U_1 f \subset g^{-1}O$. We claim that $q.U_1 f \subset p.O$. To see this, let $h \in U_1$. Then $q.hf = q.f.f^{-1}hf$ because both sides are well-defined (by Claim 1 applied to $q.f = p.g$). Since $q.f.f^{-1}hf = p.g.f^{-1}hf \subset p.O$, this proves $q.hf \in p.O$. Since h is arbitrary,

$$q.U_1 f \subset p.O.$$

After choosing U_1 smaller if necessary, we may assume the closure of $q.U_1f$ is contained in $p.O$. Therefore, there is an open neighborhood $U_2 \subset G$ of the identity such that $q.U_1f.U_2 \subset p.O$,

$$(q.U_1 \times fU_2) \cup (\{q\} \times U_1fU_2) \cup (q.U_1f \times U_2) \subset \text{dom}(\alpha)$$

and $f^{-1}U_1fU_2 \subset g^{-1}O$. It follows that

$$W := q.U_1 \times fU_2 \subset M_i \times G$$

is an open neighborhood of (q, f) . Moreover,

$$\alpha(W) = q.U_1.fU_2 = q.U_1fU_2 = q.U_1f.U_2$$

since these are all well-defined. By the assumption on U_2 , this shows that

$$\alpha(W) \subset p.O,$$

which shows that α is continuous.

Note that

$$\alpha(W) = q.U_1fU_2 = q.f.f^{-1}U_1fU_2$$

because both sides are well-defined. Because $f^{-1}U_1fU_2 \subset g^{-1}O$ which is good for $p.g = q.f$, it follows that $\alpha(W)$ is open in M_i . ■

We claim the space M_i with the partial action defined above is a local G -space. Axioms 1, 3 and 4 are immediate. To establish Axiom 2, suppose that $(p, g) \in \text{dom}(\alpha)$. We have to show $(p.g, g^{-1}) \in \text{dom}(\alpha)$. There is an open subset $W \subset \text{dom}(\alpha)$ containing (p, g) . For each $(q, f) \in W$, $q.f.f^{-1}$ is well-defined. Moreover, the set $\{(q.f, f^{-1}) : (q, f) \in W\}$ is an open neighborhood of $(p.g, g^{-1})$. This implies Axiom 2.

Let $U \subset G$ be a precompact open neighborhood of the identity with $U = U^{-1}$ and $\epsilon > 0$. We will show that if i is sufficiently large then M_i is a (U, ϵ) -sofic approximation.

Given $F \subset \Gamma$, let $\Omega(F)$ be the set of all $\Gamma h \in \Gamma \backslash G$ such that, for every $g_1, g_2 \in U^3$ with $g_1g_2 \in U^3$, $c(\Gamma hg_1, g_2) \in F$. Because U is precompact, $c(\Gamma h, U^3)$ is finite for every h . So there exists a finite set $F \subset \Gamma$ such that

$$\text{vol}_{\Gamma \backslash G}(\Omega(F)) > (1 - \epsilon/2) \text{vol}_{\Gamma \backslash G}(\Gamma \backslash G).$$

After choosing F larger if necessary, we may assume $1_G \in F$ and $F = F^{-1}$.

Because Σ is a sofic approximation, there exists I such that $i > I$ implies V_i is an $(F^2, \epsilon/2)$ -sofic approximation to Γ . We claim that if $i > I$ then M_i is a (U, ϵ) -sofic approximation. Because

$$\begin{aligned} \text{vol}_{M_i}(V_i[F^2] \times \Omega(F)) &= \text{vol}_{V_i}(V_i[F^2]) \times \text{vol}_{\Gamma \setminus G}(\Omega(F)) \\ &\geq (1 - \epsilon/2)^2 |V_i| \text{vol}_{\Gamma \setminus G}(\Gamma \setminus G) \\ &\geq (1 - \epsilon/2)^2 \text{vol}_{M_i}(M_i), \end{aligned}$$

it suffices to show that $M_i[U] \supset V_i[F^2] \times \Omega(F)$.

First we claim that if $p = (v, \Gamma h) \in V_i[F^2] \times \Omega(F)$ then $p.U.U^2$ is well-defined. This is equivalent to the well-definedness of $v.c(\Gamma h, g_1).c(\Gamma h g_1, g_2)$ for all $g_1 \in U$ and $g_2 \in U^2$. Because $\Gamma h \in \Omega(F)$, $c(\Gamma h, g_1)$ and $c(\Gamma h, g_1 g_2)$ are in F . By the cocycle equation,

$$c(\Gamma h g_1, g_2) = c(\Gamma h, g_1)^{-1} c(\Gamma h, g_1 g_2) \in F^2.$$

Since $v \in V_i[F^2]$, $v.c(\Gamma h, g_1).c(\Gamma h g_1, g_2)$ is well-defined.

Next we show that U is good for p . The first condition of ‘good’ is trivial and the second condition holds by the paragraph above. To check the third condition, suppose $g_1, g_2 \in U$ and $p.g_1 = p.g_2$. Because $c(\Gamma h, g_i) \in F$ ($i = 1, 2$), and $v \in V_i[F^2]$, the condition $p.g_1 = p.g_2$ implies $c(\Gamma h, g_1) = c(\Gamma h, g_2)$. Equivalently,

$$\sigma(\Gamma h)g_1\sigma(\Gamma h g_1)^{-1} = \sigma(\Gamma h)g_2\sigma(\Gamma h g_2)^{-1}.$$

Because we also have $\Gamma h g_1 = \Gamma h g_2$, it follows that $g_1 = g_2$. This verifies U is good for p . In particular, $p \in M_i$.

Next we claim that for all $g \in U$, $(p, g) \in \text{dom}(\alpha)$. It suffices to show $p.U \times gU$ is an open neighborhood of (p, g) in $\text{dom}(\alpha')$. This is implied by the paragraph above, which shows $p.U.gU$ is well-defined. So $V_i[F^2] \times \Omega(F) \times U \subset \text{dom}(\alpha)$.

Next we show that the map which sends $g \in U$ to $p.g$ is a homeomorphism onto an open subset of M_i . Because U is good for p , this map is well-defined and injective. By Claim 3, it is continuous. Because U is pre-compact, this implies the map is a homeomorphism onto an open subset. Thus $M_i[U] \supset V_i[F] \times \Omega(F)$ as claimed.

Because U, ϵ are arbitrary, this shows $\{M_i\}$ is a sofic approximation to G . ■

COROLLARY 3.10: *The following groups are sofic: semi-simple Lie groups (e.g., $\text{SL}(n, \mathbb{R})$, $\text{SO}(n, 1)$ etc.), the automorphism group of a regular tree.*

Proof. These groups admit residually finite lattices. Since residual finiteness implies soficity, the corollary follows from Theorem 3.9. ■

3.5.2. Restricting to a subgroup. It is well-known that if a countable group Γ is sofic then all of its subgroups are sofic. Indeed, given a sofic approximation $\{\sigma_i: \Gamma \rightarrow \text{sym}(V_i)\}$ one can restrict the maps σ_i to a subgroup Λ to obtain a sofic approximation to Λ . This argument fails in the general setting of locally compact groups because, if $H \leq G$, then the Haar measure on H might be singular to the Haar measure on G . However, the following gives a positive result.

PROPOSITION 3.11: *Let G be a locally compact sofic group. If $H \leq G$ is an open subgroup then H is sofic.*

Proof. Let (M, α) be a local G -space. Define a partial action α_H by

$$\alpha_H: \text{dom}(\alpha_H) \rightarrow M, \quad \alpha_H(p, g) = \alpha(p, g)$$

where

$$\text{dom}(\alpha_H) = \text{dom}(\alpha) \cap (M \times H).$$

We claim that (M, α_H) is a local H -space. To see this, let $p \in M$. By Axiom 4 of Definition 1, there is an open neighborhood O_p of 1_G in G such that the restriction of $\alpha(p, \cdot)$ to O_p is a homeomorphism onto an open neighborhood of p in M . Because H is open, the restriction of $\alpha(p, \cdot)$ to $O_p \cap H$ is also a homeomorphism onto an open neighborhood of p in M . This shows Axiom 4. The other Axioms are immediate.

Let U be an open neighborhood of 1_H in H and $\epsilon > 0$. Because H is open in G , U is also an open neighborhood of 1_G in G . By definition, $M[U, \alpha] = M[U, \alpha_H]$. Because the Haar measure on H equals the Haar measure on G restricted to H , the choice vol_M does not depend on whether we consider M to be a local G -space or a local H -space. So if (M, α) is a (U, ϵ) -sofic approximation to G , then (M, α_H) is also a (U, ϵ) -sofic approximation to H . ■

4. Open problems

4.1. WHICH GROUPS ARE SOFIC?

Problem 1: Are all unimodular lcsc groups sofic? For example, the Neretin group is a unimodular lcsc group without lattices [BCGM12]. Is it sofic?

Problem 2: If G is linear and unimodular then is G sofic? By Mal'cev's Theorem [Mal40] if G is finitely generated and linear then it is residually finite. Because increasing unions of sofic groups are sofic, if G is countable and linear then it is sofic.

Problem 3: If G is connected and sofic then is its universal cover sofic?

Problem 4: Suppose G is a connected unimodular Lie group and let $S \leq G$ be its solvable radical. If G/S is sofic then is G sofic?

Problem 5: Permanence properties for discrete countable sofic groups have been studied in [CHR14, HS18, ABFSG19, AG20] for example. These papers concern graph products, wreath products (restricted and unrestricted) and semi-direct products respectively. Are there analogs of these results for locally compact sofic groups?

Problem 6: Suppose G is a non-unimodular lcsc group and let $\delta : G \rightarrow \mathbb{R}_{>0}$ denote the modular homomorphism. Let $\widehat{G} = \mathbb{R} \rtimes G$ denote the semi-direct product with group law

$$(t, g)(s, h) = (t + \delta(g)s, gh).$$

Then $\widehat{G}$ and $\text{Ker}(\delta)$ are unimodular groups. If $\widehat{G}$ is sofic then is $\text{Ker}(\delta)$ sofic? If $\text{Ker}(\delta)$ is sofic then is $\widehat{G}$ sofic?

4.2. GROUP RINGS.

Problem 7: If G is sofic, then is its group von Neumann algebra Connes-embeddable? Elek and Szabó proved the answer is ‘yes’ in the case of discrete countable groups [ES05].

Problem 8: The algebraic eigenvalue conjecture of J. Dodziuk, P. Linnell, V. Mathai, T. Schick and S. Yates [DLM⁺03] posits that if Γ is a discrete group and $A \in M_n(\mathbb{Z}\Gamma)$ (the ring of $n \times n$ matrices with values in the group ring $\mathbb{Z}\Gamma$) and $l(A)$ is the corresponding operator on $\ell^2\Gamma^{\oplus n}$, then all eigenvalues of $l(A)$ are algebraic integers. This was proven true for sofic groups by A. Thom [Tho08]. Is there an analogous statement for locally compact sofic groups?

Problem 9 (Kaplansky’s direct finiteness conjecture): A ring R is said to be **directly finite** if $xy = 1$ implies $yx = 1$ for all $x, y \in R$. Kaplansky conjectured that if G is a countable group and k is a field, then the group ring kG is directly finite. This is known as Kaplansky’s Direct Finiteness Conjecture. If G is sofic, then as explained in Problem 10 below, it satisfies Gottschalk’s surjunctivity conjecture. This immediately implies kG is directly finite if k is a finite field. The general case follows because all fields are embeddable into ultraproducts of finite fields. See [CL15] for details. Is there an analogous statement in the setting of locally compact groups?

4.3. ACTIONS.

Problem 10 (Gottschalk’s surjunctivity conjecture): Suppose X is a compact Hausdorff space, G is a topological group and $G \times X \rightarrow X$ is a jointly continuous action. This action is said to be **surjunctive** if every continuous injective G -equivariant map $\phi : X \rightarrow X$ is surjective.

Gottschalk conjectured that if G is discrete and A is a finite set, then the full shift $G \curvearrowright A^G$ is surjunctive where

$$A^G = \{x : G \rightarrow A\}$$

has the topology of pointwise convergence and G acts on A^G by

$$(gx)(f) = x(g^{-1}f).$$

M. Gromov proved that sofic groups satisfy Gottschalk’s conjecture [Gro99]. His proof was simplified and made more accessible by B. Weiss [Wei00]. It was re-proven by Kerr and Li using sofic topological entropy [KL11]. Is there an analog of Gottschalk’s conjecture in the locally compact setting?

Problem 11: Sofic approximations have been used to define invariants of actions of discrete countable groups on probability spaces, compact topological spaces and Banach spaces. These invariants include sofic measure entropy [Bow10], topological sofic entropy [KL11], sofic mean dimension [Li13], sofic mean length [LL19] and ℓ^p dimension [Hay14]. These invariants generalize classical invariants of $\mathbb{Z}$ -actions. For an introduction to sofic entropy, see [Bow18]. This motivates the problem: generalize these invariants to actions by locally compact groups. This problem is open except for the fact that Sukhpreet Singh’s thesis generalized some of the foundational results of sofic entropy theory to locally compact groups. He has no plans to publish his thesis, but copies are available upon request to the first author.

4.4. SOFIC APPROXIMATIONS.

Problem 12 (Amenable groups): Elek and Szabó proved a structure theorem for sofic approximations of discrete amenable groups in [ES11]. It states that there is essentially only one sofic approximation to a discrete amenable group (up to asymptotically vanishing perturbations and taking disjoint copies) which is given by a Følner sequence. Is there an analogous statement in the setting of locally compact groups?

Problem 13 (Flexible stability): Let us say that a sofic group G is **flexibly stable** if for every $\delta > 0$ there are pre-compact open $U \subset G$ and $\epsilon > 0$ such that, if M is an (U, ϵ) -sofic approximation to G , then there exist lattice subgroups $\Gamma_1, \dots, \Gamma_k \leq G$, an open subspace $M' \subset M$, an open subspace $X' \subset X$ where

$$X := \bigsqcup_{i=1}^k \Gamma_i \backslash G$$

is the disjoint union and a homeomorphism $\Phi : M' \rightarrow X'$ such that

- $\Phi(p.g) = \Phi(p)g$ whenever both sides are defined;
- $\text{vol}(M') \geq (1 - \delta) \text{vol}(M)$;
- $\text{vol}(X') \geq (1 - \delta) \text{vol}(X)$.

This implies that any sofic approximation to G can, by a small perturbation, be changed into an approximation by a disjoint union of coset spaces. It follows from Benjy Weiss's results in [Wei01] and Elek and Szabó's structure theorem for sofic approximations of amenable groups [ES11] that residually finite amenable discrete groups are flexibly stable. It is a folklore result that free groups are flexibly stable. In recent work, it has been shown that surface groups are flexibly stable [LLM19]. This motivates the following questions: is $\text{PSL}(2, \mathbb{R})$ flexibly stable? $\text{Aut}(T_d)$? $\text{SO}(3, 1)$? $\mathbb{R}^2 \rtimes \text{SL}(2, \mathbb{R})$? Are amenable unimodular groups that admit residually finite lattices flexibly stable? If G is flexibly stable then are all lattice subgroups of G flexibly stable?

Problem 14 (Property (T) and expanders): If M is a complete Riemannian manifold with finite volume, then the **Cheeger constant** of M is

$$h(M) = \inf_{K \subset M} \frac{\text{area}(\partial K)}{\text{vol}(K)}$$

where the infimum is over all compact smooth sub-manifolds $K \subset M$ with $0 < \text{vol}(K) \leq \text{vol}(M)/2$ [Che70].

Suppose G is a connected Lie group with property (T). It is well-known that there is a positive lower bound $\epsilon_0 > 0$ on the Cheeger constants of coset spaces $\Gamma \backslash G$. That is $h(\Gamma \backslash G) \geq \epsilon_0$ for all lattices $\Gamma \leq G$. With this in mind, we conjecture that there exists $\epsilon'_0 > 0$ such that for every $\delta > 0$ there exist a pre-compact open set $U \subset G$ and $\epsilon > 0$, such that if M is any (U, ϵ) -sofic approximation to G , then there exists a smooth submanifold $M' \subset M$ satisfying

- $\text{vol}(M') \geq (1 - \delta) \text{vol}(M)$;
- every connected component of M' has Cheeger constant $\geq \epsilon'_0$.

A similar conjecture for discrete (T) groups by the first author was proven by Gábor Kun [Kun16]. Maybe there is a common generalization to all locally compact (T) groups?

Problem 15: The **sofic dimension** of a countable discrete group measures the growth rate of the number of sofic approximations to the group [DKP14, GP15]. It is a combinatorial version of the free entropy dimension [Voi96]. Moreover, it admits a natural formula with respect to free product with amalgamation over an amenable group. Are there analogs of these results in the locally compact setting?

Problem 16: A countable discrete group G is sofic if and only if it embeds into a metric ultraproduct of finite symmetric groups [ES05, Pes08]. Is there an analogous fact for locally compact sofic groups? Note that $\mathrm{SL}(2, \mathbb{R})$ is sofic (because it admits a residually finite lattice) but it does not continuously embed into a metric ultraproduct of compact groups. This is because $\mathrm{SL}(2, \mathbb{R})$ does not admit a proper bi-invariant metric.

4.5. GROUPOIDS AND MEASURED EQUIVALENCE RELATIONS.

Problem 17: Soficity was generalized to discrete measured equivalence relations and groupoids in [EL10, Pău11, DKP14]. Can this theory be generalized to measured equivalence relations and groupoids with locally compact leaves?

Problem 18: It might be possible to reduce soficity of a non-discrete lcsc group to soficity of a related discrete measured equivalence relation. The latter notion was introduced in [EL10].

It is well-known that a discrete countable group G is sofic if there exists an essentially free action of G on a standard probability space (X, μ) such that the orbit-equivalence relation is sofic. It seems likely that this fact generalizes to locally compact groups as follows.

Suppose G acts on a standard probability space (X, μ) preserving the measure. For simplicity, let us assume the action is essentially free. By [FHM78] there is a complete lacunary section $S \subset X$. This means that S is Borel, GS is conull in X and there is an open neighborhood $U \subset G$ of the identity such that $Ux \cap S = \{x\}$ for all $x \in S$. In particular, if $\mathcal{R}_G^X = \{(x, gx) : x \in X, g \in G\}$ is the orbit-equivalence relation of the G action and $\mathcal{R}_G^S := \mathcal{R}_G \cap (S \times S)$, then $\mathcal{R}_G^S$ is an equivalence relation on S with countable classes.

If G is non-discrete then $\mu(S) = 0$. In spite of this, there is a natural measure, denoted ν , on S which behaves as if it were μ conditioned on S . The measure is defined by

$$\nu(A) = \frac{\mu(VA)}{\text{Haar}_G(V)}$$

where $A \subset S$ is any Borel set and $V \subset G$ is a symmetric open neighborhood of 1_G in G such that $V^2 \subset U$. This does not depend on the choice of V . This is explained in [Avn10] for example.

We conjecture: if there exists a pmp action $G \curvearrowright (X, \mu)$ and a section $S \subset X$ as above such that the discrete measured equivalence relation $(S, \nu, \mathcal{R}_G^S)$ is sofic, then G is sofic. This might give an approach to proving that the group of von Neumann algebras of sofic groups are Connes-embeddable (see [EL10] for related results in the discrete case).

Problem 19: The previous problem gave a sufficient condition for soficity. There is a related equivalent condition. It is well-known that a discrete countable group G is sofic if and only if for every Bernoulli shift action of G the associated measured equivalence relation is sofic. It seems likely that this fact generalizes to locally compact groups as follows.

Consider the Poisson point process on a non-discrete lcsc group G with intensity measure equal to a left-Haar measure on G . Because G is non-discrete, we can consider the law of this process to be a G -invariant probability measure μ on the space Ω of discrete closed subsets of G . Let $\Omega_1 \subset \Omega$ be the set of discrete closed subsets $\omega \subset G$ with $1_G \in \omega$. Even though $\mu(\Omega_1) = 0$, there is a natural probability measure ν on Ω_1 that intuitively represents μ conditioned on Ω_1 (this exists even though Ω_1 is not a lacunary section). Define an equivalence relation $\mathcal{R}$ on Ω_1 by $(\omega_1, \omega_2) \in \mathcal{R} \Leftrightarrow \exists g \in G$ such that $g\omega_1 = \omega_2$. Then $\mathcal{R}$ is discrete and ν -preserving. We conjecture that G is sofic if and only if this measured equivalence relation is sofic.

References

- [ABFSG19] G. Arzhantseva, F. Berlai, M. Finn-Sell and L. Glebsky, *Unrestricted wreath products and sofic groups*, International Journal of Algebra and Computation **29** (2019), 343–355.
- [AG20] G. Arzhantseva and Š. R. Gal, *On approximation properties of semidirect products of groups*, Annales Mathématiques Blaise Pascal **27** (2020), 125–130.

- [Avn10] N. Avni, *Entropy theory for cross-sections*, Geometric and Functional Analysis **19** (2010), 1515–1538.
- [BCGM12] U. Bader, P.-E. Caprace, T. Gelander and S. Mozes, *Simple groups without lattices*, Bulletin of the London Mathematical Society **44** (2012), 55–67.
- [Bow10] L. Bowen, *Measure conjugacy invariants for actions of countable sofic groups*, Journal of the American Mathematical Society **23** (2010), 217–245.
- [Bow15] L. Bowen, *Cheeger constants and L^2 -Betti numbers*, Duke Mathematical Journal **164** (2015), 569–615.
- [Bow18] L. P. Bowen, *A brief introduction of sofic entropy theory*, in *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited Lectures*, World Scientific, Hackensack, NJ, 2018, pp. 1847–1866.
- [Che70] J. Cheeger, *A lower bound for the smallest eigenvalue of the Laplacian*, in *Problems in Analysis (Papers Dedicated to Salomon Bochner, 1969)*, Princeton University Press, Princeton, NJ, 1970, pp. 195–199.
- [CHR14] L. Ciobanu, D. F. Holt and S. Rees, *Sofic groups: graph products and graphs of groups*, Pacific Journal of Mathematics **271** (2014), 53–64.
- [CL15] V. Capraro and M. Lupini, *Introduction to Sofic and Hyperlinear Groups and Connes’ Embedding Conjecture*, Lecture Notes in Mathematics, Vol. 2136, Springer, Cham, 2015.
- [DKP14] K. Dykema, D. Kerr and M. Pichot, *Sofic dimension for discrete measured groupoids*, Transactions of the American Mathematical Society **366** (2014), 707–748.
- [DLM⁺03] J. Dodziuk, P. Linnell, V. Mathai, T. Schick and S. Yates, *Approximating L^2 -invariants and the Atiyah conjecture*, Communications on Pure and Applied Mathematics **56** (2003), 839–873.
- [EL10] G. Elek and G. Lippner, *Sofic equivalence relations*, Journal of Functional Analysis **258** (2010), 1692–1708.
- [ES05] G. Elek and E. Szabó, *Hyperlinearity, essentially free actions and L^2 -invariants. The sofic property*, Mathematische Annalen **332** (2005), 421–441.
- [ES11] G. Elek and E. Szabó, *Sofic representations of amenable groups*, Proceedings of the American Mathematical Society **139** (2011), 4285–4291.
- [FHM78] J. Feldman, P. Hahn and C. C. Moore, *Orbit structure and countable sections for actions of continuous groups*, Advances in Mathematics **28** (1978), 186–230.
- [GP15] R. Graham and M. Pichot, *A free product formula for the sofic dimension*, Canadian Journal of Mathematics **67** (2015), 369–403.
- [Gro99] M. Gromov, *Endomorphisms of symbolic algebraic varieties*, Journal of the European Mathematical Society **1** (1999), 109–197.
- [Hay14] B. Hayes, *An l^p -version of von Neumann dimension for Banach space representations of sofic groups*, Journal of Functional Analysis **266** (2014), 989–1040.
- [HS18] B. Hayes and A. W. Sale, *Metric approximations of wreath products*, Université de Grenoble. Annales de l’Institut Fourier **68** (2018), 423–455.
- [KL11] D. Kerr and H. Li, *Entropy and the variational principle for actions of sofic groups*, Inventiones Mathematicae **186** (2011), 501–558.

- [Kun16] G. Kun, *On sofic approximations of property (T) groups*, <https://arxiv.org/abs/1606.04471>.
- [Li13] H. Li, *Sofic mean dimension*, *Advances in Mathematics* **244** (2013), 570–604.
- [LL19] H. Li and B. Liang, *Sofic mean length*, *Advances in Mathematics* **353** (2019), 802–858.
- [LLM19] N. Lazarovich, A. Levit and Y. Minsky, *Surface groups are flexibly stable*, <https://arxiv.org/abs/1901.07182>.
- [Mal40] A. Malcev, *On isomorphic matrix representations of infinite groups*, *Matematicheskii Sbornik* **8** (1940), 405–422.
- [Pău11] L. Păunescu, *On sofic actions and equivalence relations*, *Journal of Functional Analysis* **261** (2011), 2461–2485.
- [Pes08] V. G. Pestov, *Hyperlinear and sofic groups: a brief guide*, *Bulletin of Symbolic Logic* **14** (2008), 449–480.
- [PK13] V. G. Pestov and A. Kwiatkowska, *An introduction to hyperlinear and sofic group*, in *Appalachian Set Theory 2006–2012*, London Mathematical Society Lecture Note Series, Vol. 406, Cambridge University Press, Cambridge, 2013, pp. 145–185.
- [Tho08] A. Thom, *Sofic groups and Diophantine approximation*, *Communications on Pure and Applied Mathematics* **61** (2008), 1155–1171.
- [Voi96] D. Voiculescu, *The analogues of entropy and of Fisher’s information measure in free probability theory. III. The absence of Cartan subalgebras*, *Geometric and Functional Analysis* **6** (1996), 172–199.
- [Wei00] B. Weiss, *Sofic groups and dynamical systems*, *Sankhyā* **62** (2000), 350–359.
- [Wei01] B. Weiss, *Monotileable amenable groups*, in *Topology, Ergodic Theory, Real Algebraic Geometry*, American Mathematical Society Translations. Series 2, Vol. 202, American Mathematical Society, Providence, RI, 2001, pp. 257–262.