High-Energy and Ultra-High-Energy Neutrinos: A Snowmass White Paper

Editors: Markus Ackermann^a, Mauricio Bustamante^b, Lu Lu^c, Nepomuk Otte^d, Mary Hall Reno^e, Stephanie Wissel^f

Markus Ackermann,¹ Sanjib K. Agarwalla,^{2,3,4} Jaime Alvarez-Muñiz,⁵ Rafael Alves Batista,⁶ Carlos A. Argüelles,⁷ Mauricio Bustamante,⁸ Brian A. Clark,⁹ Austin Cummings,¹⁰ Sudipta Das,^{2,3} Valentin Decoene,¹⁰ Peter B. Denton,¹¹ Damien Dornic,¹² Zhan-Arys Dzhilkibaev,¹³ Yasaman Farzan,¹⁴ Alfonso Garcia,^{6,15} Maria Vittoria Garzelli,¹⁶ Christian Glaser,¹⁷ Aart Heijboer,^{18,19} Jörg R. Hörandel,^{18,20} Giulia Illuminati,^{21,22} Yu Seon Jeong,²³ John L. Kelley,²⁴ Kevin J. Kelly,²⁵ Ali Kheirandish,¹⁰ Spencer R. Klein,^{26,27} John F. Krizmanic,²⁸ Michael J. Larson,²⁹ Lu Lu,²⁴ Kohta Murase,¹⁰ Ashish Narang,² Nepomuk Otte,³⁰ Remy L. Prechelt,³¹ Steven Prohira,^{32,33} Mary Hall Reno,³⁴ Elisa Resconi,³⁵ Marcos Santander,³⁶ Victor B. Valera,⁸ Justin Vandenbroucke,²⁴ Olga Vasil'evna Suvorova,¹³ Lawrence Wiencke,³⁷ Stephanie Wissel,¹⁰ Shigeru Yoshida,³⁸ Tianlu Yuan,²⁴ Enrique Zas,⁵ Pavel Zhelnin,⁷ Bei Zhou³⁹

Endorsers: Luis A. Anchordogui,⁴⁰ Yosuke Ashida,²⁴ Mahdi Bagheri,³⁰ Aswathi Balagopal V.,²⁴ Vedant Basu,²⁴ James Beatty,³² Keith Bechtol,²⁴ Nicole Bell,⁴¹ Abigail Bishop,²⁴ Julia Book,⁷ Anthony Brown,⁴² Alexander Burgman,⁴³ Michael Campana,⁷⁶ Nhan Chau,⁶⁰ Thomas Y. Chen⁴⁴ Alan Coleman,⁴⁵ Amy Connolly,³² Janet M. Conrad,⁴⁶ Pablo Correa,⁴⁷ Cyril Creque-Sarbinowski,³⁹ Austin Cummings,¹⁰ Zachary Curtis-Ginsberg,⁴⁸ Paramita Dasgupta,⁴⁹ Simon De Kockere,⁴⁷ Krijn de Vries,⁴⁷ Cosmin Deaconu,⁴⁸ Abhishek Desai,²⁴ Tyce DeYoung,⁹ Armando di Matteo,⁵⁰ Dominik Elsaesser,⁵¹ Phillip Fürst,⁵² Kwok Lung Fan,²⁹ Anatoli Fedynitch,⁵³ Derek Fox,¹⁰ Erik Ganster,⁵² Martin Ha Minh³⁵ Christian Haack,³⁵ Steffen Hallman,¹ Francis Halzen,²⁴ Andreas Haungs,⁵¹ Aya Ishihara,³⁸ Eleanor Judd,⁵⁴ Timo Karg,¹ Albrecht Karle,²⁴ Teppei Katori,⁵⁵ Alina Kochocki,⁹ Claudio Kopper,⁹ Marek Kowalski,¹ Ilya Kravchenko,⁵⁶ Naoko Kurahashi,⁵⁷ Mathieu Lamoureux,⁵⁸ Hermes León Vargas, ⁵⁹ Massimiliano Lincetto, ⁶⁰ Qinrui Liu, ⁶¹ Jim Madsen, ²⁴ Yuya Makino, ²⁴ Joseph Mammo, ⁵⁶ Zsuzsa Marka⁶² Eric Mayotte³⁷ Kevin Meagher,²⁴ Maximilian Meier,³⁸ Martin Ha Minh,³⁵ Lino Miramonti,^{63,64} Marjon Moulai,²⁴ Katharine Mulrey,²⁰ Marco Muzio,¹⁰ Richard Naab¹ Anna Nelles,¹ William Nichols,²⁴ Alisa Nozdrina,³³ Erin O'Sullivan,¹⁷ Vivian ODell,²⁴ Jesse Osborne,²⁴ Vishvas Pandey,⁶⁵ Ek Narayan Paudel,⁴⁵ Alex Pizzuto,²⁴ Mattias Plum,⁶⁶ Carlos Pobes Aranda,⁶⁷ Lilly Pyras,¹ Christoph Raab,⁶⁸ Zoe Rechav,²⁴ Juan Rojo,^{18,19} Oscar Romero Matamala,³⁰ Marcos Santander,³⁶ Pierpaolo Savina,²⁴ Frank Schroeder,⁴⁵ Lisa Schumacher,³⁵ Sergio Sciutto,⁶⁹ Stephen Sclafani,⁵⁷ Mohammad Ful Hossain Seikh,³⁴ Manuel Silva,²⁴ Rajeev Singh,⁷⁰ Daniel Smith,⁴⁸ Samuel Timothy Spencer,⁷¹ Robert Wayne Springer,⁷² Juliana Stachurska,⁴⁶ Olga Suvorova,⁷³ Ignacio Taboada,³⁰ Simona Toscano,⁴⁹ Matias Tueros,⁶⁹ Jean Pierre Twagirayezu⁹ Nick van Eijndhoven,⁴⁷ Péter Veres,⁷⁴ Abigail Vieregg,⁴⁸ Winnie Wang,²⁴ Nathan Whitehorn,⁹ Walter Winter,¹ Emre Yildizci,²⁴ Shiqi Yu,⁹

 $^{^{\}rm a} markus.ackermann@desy.de$

 $^{^{\}rm b}$ mbustamante@nbi.ku.dk

^clu.lu@icecube.wisc.edu

^dotte@gatech.edu

 $^{^{\}rm e}$ mary-hall-reno@uiowa.edu

^fwissel@psu.edu

- ¹DESY, D-15738 Zeuthen, Germany
- ²Institute of Physics, Sachivalaya Marg, Sainik School Post, Bhubaneswar 751005, India
- ³Homi Bhabha National Institute, Anushakti Nagar, Mumbai 400094, India
- ⁴International Centre for Theoretical Physics, Strada Costiera 11, 34151 Trieste, Italy
- ⁵Instituto Galego de Fisica de Altas Enerxias, Universidade de Santiago de Compostela, Spain
- ⁶Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain
- ⁷Department of Physics & Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA 02138, USA
- ⁸Niels Bohr International Academy, Niels Bohr Institute, University of Copenhagen, Copenhagen 2100, Denmark
- ⁹Dept. of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
- ¹⁰Departments of Physics and Astronomy & Astrophysics, Institute for Gravitation and the Cosmos, Pennsylvania State University, State College, Pennsylvania 16802 USA
- ¹¹High Energy Theory Group, Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA
- ¹²Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France
- ¹³Institute for Nuclear Research, Russian Academy of Sciences, Moscow, Russia
- ¹⁴School of Physics, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran
- ¹⁵Instituto de Física Corpuscular (IFIC), Universitat de València (UV), 46980 Paterna, València, Spain
- ¹⁶II. Institut für Theoretische Physik, Universität Hamburg, D-22761 Hamburg, Germany
- ¹⁷ Uppsala University Department of Physics and Astronomy, Uppsala SE-752 37, Sweden
- ¹⁸Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands
- ¹⁹ University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands
- ²⁰Radboud University, Department of Astrophysics/IMAPP, 6500GL Nijmegen, The Netherlands
- ²¹INFN Sezione di Bologna, Viale Berti-Pichat 6/2, 40127 Bologna, Italy
- ²²Dipartimento di Fisica e Astronomia dell'Università, Viale Berti Pichat 6/2, 40127 Bologna, Italy
- ²³High Energy Physics Center, Chung-Ang University, Seoul 06974, Korea
- ²⁴Department of Physics & Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin–Madison, Madison, WI 53706, USA
- ²⁵ Theoretical Physics Department, CERN, Esplande des Particules, 1211 Geneva 23, Switzerland
- ²⁶Department of Physics, University of California, Berkeley, CA 94720, USA
- ²⁷Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
- ²⁸Laboratory for Astroparticle Physics, NASA/GSFC, Greenbelt, MD 20771, USA
- ²⁹Department of Physics, University of Maryland, College Park, MD 20742, USA
- ³⁰Georgia Institute of Technology, School of Physics & Center for Relativistic Astrophysics, Atlanta, Georgia 30332-0430, USA
- ³¹University of Hawai'i Mānoa, Honolulu, HI 96822, USA
- ³²Department of Physics, Center for Cosmology and AstroParticle Physics, Ohio State University, Columbus, OH 43210
- ³³Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045
- ³⁴Department of Physics and Astronomy, University of Iowa, Iowa City, Iowa 52242 USA
- ³⁵Department of Physics, Technical University of Munich, D-86748 Garching, Germany
- ³⁶Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA
- ³⁷Colorado School of Mines, Golden, CO 80401, USA
- ³⁸International Center for Hadron Astrophysics, Chiba University, Chiba 268-8522, Japan
- ³⁹Department of Physics and Astronomy, Johns Hopkins University, Baltimore, Maryland 21218, USA
- ⁴⁰Department of Physics and Astronomy, Lehman College, City University of New York, Bronx, NY 10468, USA
- ⁴¹School of Physics, The University of Melbourne, Victoria 3010, Australia

- ⁴²Centre for Advanced Instrumentation, Department of Physics, University of Durham, South Road, Durham, DH1 3LE, UK
- ⁴³Lund University, Lund, Sweden
- ⁴⁴Fu Foundation School of Engineering and Applied Science, Columbia University in the City of New York, New York, NY 10027, U.S.A.
- ⁴⁵Bartol Research Institute, Department of Physics and Astronomy, University of Delaware, Newark DE, USA
- ⁴⁶Dept. of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ⁴⁷ Vrije Universiteit Brussel (VUB), Dienst ELEM, B-1050 Brussels, Belgium
- ⁴⁸Dept. of Physics, Enrico Fermi Institue, Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60560, USA
- ⁴⁹ Université Libre de Bruxelles, Science Faculty CP230, B-1050 Brussels, Belgium
- ⁵⁰INFN, Sezione di Torino, Torino, Italy
- ⁵¹Karlsruhe Institute of Technology, Institute for Astroparticle Physics, D-76021 Karlsruhe, Germany
- ⁵²III. Physikalisches Institut B, RWTH Aachen University, 52056 Aachen, Germany
- ⁵³Institute of Physics, Academia Sinica, Taipei, Taiwan
- ⁵⁴Space Sciences Laboratory, University of California, Berkeley, CA, USA
- ⁵⁵Department of Physics, King's College London, WC2R 2LS London, UK
- ⁵⁶Dept. of Physics and Astronomy, University of Nebraska, Lincoln, Nebraska 68588
- ⁵⁷Department of Physics, Drexel University, Philadelphia, Pennsylvania 19104
- ⁵⁸Centre for Cosmology, Particle Physics and Phenomenology CP3, Université catholique de Louvain, Louvain-la-Neuve, Belgium
- ⁵⁹Instituto de Física, Universidad Nacional Autónoma de Mexico, Ciudad de Mexico, Mexico
- ⁶⁰ Fakultät für Physik Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany
- ⁶¹Department of Physics, Engineering Physics and Astronomy, Queen's University, Kingston, Ontario, Canada
- ⁶²Columbia Astrophysics Laboratory, Columbia University in the City of New York, New York, NY 10027, USA
- ⁶³INFN, Sezione di Milano, Milano, Italy
- ⁶⁴Università di Milano, Dipartimento di Fisica, Milano, Italy
- ⁶⁵Department of Physics, University of Florida, Gainesville, FL 32611, USA
- ⁶⁶Physics Department, South Dakota School of Mines and Technology, Rapid City, SD 57701, USA
- ⁶⁷ INMA, CSIC-Universidad de Zaragoza, Pedro Cerbuna 12, 50009 Zaragoza, Spain
- ⁶⁸Institute of Physics of the Czech Academy of Sciences, 182 21 Prague 8, Czechia
- ⁶⁹Departamento de Física, Universidad Nacional de La Plata, C. C.67 1900 La Plata, Argentina
- ⁷⁰Institute of Nuclear Physics Polish Academy of Sciences, PL 31-342 Kraków, Poland
- ⁷¹Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, D-91058 Erlangen, Germany
- ⁷²Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA
- ⁷³Institute for Nuclear Research of Russian Academy of Sciences, Moscow, Russia
- ⁷⁴Center for Space Plasma and Aeronomic Research, University of Alabama in Huntsville, 320 Sparkman Drive, Huntsville, AL 35899, USA

ABSTRACT: Astrophysical neutrinos are excellent probes of astroparticle physics and high-energy physics. With energies far beyond solar, supernovae, atmospheric, and accelerator neutrinos, high-energy and ultrahigh-energy neutrinos probe fundamental physics from the TeV scale to the EeV scale and beyond. They are sensitive to physics both within and beyond the Standard Model through their production mechanisms and in their propagation over cosmological distances. They carry unique information about their extreme non-thermal sources by giving insight into regions that are opaque to electromagnetic radiation. This white paper describes the opportunities astrophysical neutrino observations offer for astrophysics and high-energy physics, today and in coming years.

Executive Summary

With the discovery of high-energy astrophysical neutrinos, we have entered a new era in astroparticle physics. The detection of neutrinos in the TeV–PeV energy range and the expected detection at even higher energies have far-reaching consequences. In particle physics, these neutrinos can address important open questions: What is the nature of dark matter? Do new particles and interactions exist at the highest energies? Are there new fundamental symmetries? Owing to their high energies and cosmological-scale baselines, the highest energy neutrinos are uniquely positioned to address these questions. In astrophysics, they are key to understanding the origin of the highest-energy cosmic rays and gamma rays. We are only seeing the very beginning of what is likely to come.

This white paper describes the rich physics opportunities that high-energy (HE, TeV-100 PeV) and ultrahigh energy (UHE, ≥ 100 PeV) neutrino observations offer for particle physics and astrophysics, potential which has already been acknowledged by international panels like the US Decadal Survey on Astronomy and Astrophysics 2020 (Astro2020) [1] and the European Consortium for Astroparticle Physics (EuCAPT) [2]. The current status and future prospects are:

- Particle Physics [3-5]: In the HE range (TeV-100 PeV), there is an ongoing campaign to test Standard-Model (SM) and beyond-Standard-Model (BSM) neutrino physics: IceCube's discovery of astrophysical high-energy neutrinos has turned theory predictions into data-driven tests. At present, low event rates limit the sensitivity of these tests, but the next generation of neutrino telescopes will overcome this complication. In the UHE (≥ 100 PeV) range, neutrinos have been long-predicted but still await discovery. Thanks to a host of UHE neutrino telescopes currently under planning, they are likely to be detected within the coming years. Detecting UHE neutrinos will push neutrino physics to the forefront of the particle physics energy frontier.
 - High Energy (TeV-100 PeV): We already test the Standard Model with astrophysical neutrinos at TeV-PeV energies and have the sensitivity to search for relatively large BSM effects. Existing studies include the discovery of the Glashow resonance, the first measurement of the HE neutrino-nucleon cross section, searches for neutrinos from dark matter annihilation and decay, and searches for new neutrino self-interactions. With steadily increasing event statistics in this decade and beyond, more subtle effects can be tested.
 - Ultra-High Energy ($\geq 100 \text{ PeV}$): The only way to probe weak-scale physics at center-of-mass energies above 50 TeV is with naturally occurring UHE neutrinos. Because our capacity to probe new physics at these scales will depend on the flux of UHE neutrinos, it should be assessed together with the flux discovery prospects.
 - Neutrino Oscillation Physics: HE and UHE astrophysical neutrinos are sensitive to at least six ($\nu_e \rightarrow \nu_e, \nu_e \rightarrow \nu_\mu, \nu_e \rightarrow \nu_\tau, \nu_\mu \rightarrow \nu_e, \nu_\mu \rightarrow \nu_\mu, \nu_\mu \rightarrow \nu_\tau$) out of the nine neutrino oscillation channels. Furthermore, they are sensitive to oscillations at higher energies and over longer baselines than any other oscillation experiment, allowing for unique searches for BSM oscillations.
 - New Neutrino Interactions: Neutrinos may have BSM interactions, and if the coupling strengths are weak or if heavy mediators mediate the interactions, these interactions may only manifest themselves in the HE and UHE neutrino sector. Possibile scenarios include BSM neutrino interactions with dark matter—including heavy dark matter—and with sterile neutrinos.
 - Synergies with the Neutrino Frontier: Measurements of high-energy prompt atmospheric neutrinos complement existing measurements of heavy-flavor production at LHC experiments and

neutrino experiments at a future Forward Physics Facility [6]. At ultra-high energies, neutrino interactions probe the structure of the proton in kinematic regions that cannot be explored by accelerator experiments.

- Astrophysics [7]: IceCube has discovered a diffuse flux of astrophysical HE (TeV-100 PeV) neutrinos. They have characterized the energy spectrum, distribution of arrival directions, and flavor composition with increasing precision, thanks to steadily improving event statistics, analysis methods, and understanding of detector uncertainties. There is promising observational evidence of the first HE neutrino source and of a connection between electromagnetic and neutrino emission from one flaring blazar event. The next generation of neutrino experiments will firmly establish HE neutrino sources and provide high enough event rates for precision studies. In the UHE range (≥ 100 PeV), we may finally detect longsought UHE neutrinos in the coming years. Detecting them will be a critical step towards finding the origin of ultra-high-energy cosmic rays (UHECRs), the most energetic known particles in the Universe.
 - High Energy (TeV-100 PeV): The discovery of HE neutrinos in the last decade opened a new window into the Universe. We have learned that a large population of dim sources likely explains the bulk of the HE neutrino sources and that their energy density is comparable to that of UHECRs and gamma rays, hinting at the possibility of a common origin. With the improved statistics, sensitivity, and sky coverage offered by upcoming experiments, we can expect to expand our view of the neutrino sky, including firmly establishing neutrino sources. Next-generation telescopes currently under planning or construction will allow detailed studies of HE neutrinos, including their energy spectrum, flavor composition, and the identity of their sources.
 - Ultra-High Energy (≥ 100 PeV): We do not know how the diffuse astrophysical neutrino flux extends to higher energies. Finding out whether it cuts off or not in the 10–100 PeV range is crucial for understanding the physics underlying UHECR accelerators and identifying source classes. Recent advancements in UHECR physics have pushed UHE neutrino flux predictions lower than anticipated: For experiments to probe the entire range of UHE-neutrino models they must target a flux sensitivity of 10^{-10} GeV cm⁻² sr⁻¹ s⁻¹. But current and planned experiments with lower flux sensitivity already constrain properties of UHECRs and their sources even with non-detections of UHE neutrinos.
 - Multi-Messenger Era: Evidence of the first extragalactic HE neutrino source, TXS 0506+056, and the transformational discovery of gravitational-wave sources kicked-off multi-messenger astrophysics. With the advent of more sensitive neutrino experiments and dedicated follow-up facilities, we will likely witness more multi-messenger events at higher statistical significance.

Fully opening the HE and UHE neutrino windows requires a multi-pronged approach that explores a broad range of proposed experimental techniques. With upcoming neutrino telescopes, the goal is to improve the sensitivity ten-fold in the HE range and hundred-fold in the UHE range. In the HE range, we are moving into the high-statistics era, where subtle features may be resolved and multiple sources may be discovered. In the UHE range, neutrino flux predictions and physics tests are well-motivated, but we will likely need new techniques to enable discoveries. The ongoing efforts to develop new instruments will improve sensitivity, increase statistics, and expand the energy range. A broad range of experimental approaches, with complementary capabilities, must be developed and tested in the coming decades. Because it is not clear which techniques will prevail, we advocate for a broad portfolio of experiments in the HE and UHE neutrino sector. Owing to their potential, HE and UHE neutrinos should be at the forefront of the high-energy physics program, as they push the boundaries for the neutrino, cosmic, energy, theory, and instrumentation frontiers.

Contents

1	Introduction and overview		1
	1.1	HE and UHE cosmic neutrinos in particle physics	2
	1.2	HE and UHE cosmic neutrinos in astrophysics	3
	1.3	Detector requirements to achieve the science goals	4
	1.4	Present and future experimental landscape	6
2	Current status and lessons learned		7
	2.1	In particle physics	7
	2.2	In astrophysics	17
3	Goals for the next decade		23
	3.1	In particle physics	23
	3.2	In astrophysics	28
4	Experimental landscape		33
	4.1	Overview	33
	4.2	High-energy range	37
	4.3	Ultra-high-energy range $(> 100 \text{ PeV})$	47

1 Introduction and overview

Cosmic neutrinos are unique probes of extreme environments surrounding the most energetic sources in the Universe and a unique test beam for weak interactions at energies inaccessible through accelerators. Within the last decade, the discovery of high-energy (HE, TeV to 100 PeV) astrophysical neutrinos by IceCube [8] has opened a new window to learn more about cosmic accelerators and neutrino interactions at the highest energies. With next-generation experiments pushing sensitivity and energy reach, we anticipate that the wealth of information will expand dramatically. Ultra-high-energy (UHE, >100 PeV) neutrinos, long-sought but not yet detected, provide the only means of directly investigating processes that occur at energy scales of $EeV (\equiv 10^{18} eV)$ and above in the distant Universe. Discovering them would open new regimes of exploration in high-energy physics, astrophysics, and cosmology. In this white paper, we describe the significant physics opportunities offered by cosmic neutrinos and map out the experimental landscape in the coming decades.

Observations of neutrinos from different sources, across different energies and traveled distances, have led to the fundamental-physics conclusions that neutrinos have mass and mix among flavors. These Nobel-prize winning experimental tests include measurements of neutrinos in the sub-GeV-to-10-PeV energy range from cosmic-ray interactions in the atmosphere [9] and of neutrinos from the Sun [10–15]. Indeed, neutrinos access important questions in the complementary fields of high-energy physics and astrophysics. The wide range of neutrino energies and traveled distances allow us to explore neutrino properties, their interactions, and fundamental symmetries across a wide breadth of parameter space, as shown in Fig. 1. And because they are neutral and weakly interacting, they carry information about the physical conditions at their points of origin; at the highest energies, even from powerful cosmic accelerators at the edge of the observable Universe.

Recently, the discovery of a diffuse flux of HE astrophysical neutrinos, in the TeV–PeV range [8, 16] opened a new view to the Universe. They have made possible the direct measurement of weak interactions in a new energy regime, including the neutrino-nucleon cross section [17–19], inelasticity distribution [19],



Figure 1. Distribution of neutrino sources in energy and distance traveled to the detector, and present and future experiments aimed at detecting them. We focus on high-energy and ultra-high energy neutrinos. Updated from Ref. [4].

and the first Glashow resonance ($\bar{\nu}_e e$) candidate [20, 21]. Ongoing and future observations will refine the measurements of the astrophysical neutrino observables (energy spectrum, flavor composition, distribution of arrival directions and arrival times) and extend them beyond 10 PeV. For particle physics, this means gaining sensitivity to smaller predicted effects and extending the energy scale of fundamental physics that can be tested. For astrophysics, this means probing the most energetic non-thermal sources of the Universe indirectly through the diffuse flux, and directly through the discovery of point sources. Further, neutrinos from transient astrophysical events, detected in spatial or temporal coincidence with cosmic rays and electromagnetic radiation [22–25], will improve our understanding of the extreme physical processes in these environments. UHE neutrinos with energies exceeding 100 PeV, first predicted more than fifty years ago [26] but still undiscovered, are the next frontier in probing fundamental physics and astrophysics at the ultimate neutrino energies.

The preceding decade has ushered in a new era of astroparticle physics, including high-energy neutrino detection. Figure 1 shows that the potential outlined above will be achieved by a rich experimental program of detectors in the next 10–20 years that are presently in different stages of planning, design, and construction. We anticipate that the next decade will result in the construction of multiple high-energy neutrino detectors spanning complementary regions of the sky, with differing sensitivity to different energy ranges between TeV and EeV, and complementary flavor-identification capabilities. While the preceding decade was one of neutrino discovery at high energies, the coming years will be of higher-precision studies at high energies and, plausibly, of discovery at ultra-high energies. These studies are further enhanced by observations with all four messengers – cosmic rays, neutrinos, photons, and gravitational waves. For reviews, see complementary Snowmass Whitepapers on each of these messengers [27–30] and the broad scope of multi-messenger physics [31].

1.1 HE and UHE cosmic neutrinos in particle physics

There is a vast landscape of physics to explore at the highest energies, and high-energy cosmic neutrinos are uniquely well-equipped for the task [4]. Their potential as probes of fundamental physics [3–5, 32, 33] was identified early, but they were only discovered recently, in 2013, when the IceCube Neutrino Observatory observed a diffuse flux of TeV–PeV cosmic neutrinos [8, 34–37]. Since then, there has been a gradual shift of focus from proposing prospective tests of high-energy neutrino physics to performing real, data-driven tests,



Figure 2. Models of new neutrino physics and other new physics classified according to the stage at which they act—at production, during propagation, at detection—and what feature they affect—energy spectrum, arrival directions, flavor composition, arrival times. Figure reproduced from Ref.[5].

of increasing sophistication and based on progressively more and better experimental data. This, paired with a rich present and future experimental program, provides a valuable opportunity to make significant progress.

To seize this opportunity, the neutrino community needs a comprehensive exploration plan that maximizes the potential to probe neutrino properties and the discovery of new physics, identifies target experimental sensitivities, and exploits synergies between TeV–PeV neutrino experiments and their upcoming counterparts at lower and higher energies.

Figure 1 shows why high-energy (TeV–PeV) and ultra-high energy (\geq EeV) neutrinos are incisive probes of new physics. Because they have the highest neutrino energies known—TeV to EeV—they can probe physics at energy scales that are inaccessible to us in the laboratory. Because they travel unscathed for the longest distances—up to a few Gpc, the size of the observable Universe—even tiny effects can accumulate and become observable.

Section 2.1 outlines the current results for UHE neutrino interactions, lepton flavor mixing and on neutrino production and propagation, including standard model and beyond the standard model (BSM) approaches. In Sec. 3.1, we discuss the goals for the future and summarize efforts underway.

1.2 HE and UHE cosmic neutrinos in astrophysics

That neutrinos can emerge from and point back to their sources make them exceptional probes of the extreme environments that produce ultra-high-energy cosmic rays (UHECRs). The luminosity densities of diffuse sub-TeV gamma rays, high-energy neutrinos and UHECRs are comparable, signaling the possible connection between cosmic acceleration and the pp and $p\gamma$ interactions that would produce high-energy neutrinos and photons [38–40]. Measurements of the diffuse high-energy neutrino spectrum and flavor composition, i.e., the relative contribution of ν_e , ν_{μ} , and ν_{τ} in the incoming flux, and a nascent multi-messenger program that includes neutrino telescopes will help identify characteristics of source populations.



Figure 3. Cosmic neutrinos probe fundamental particle physics and astrophysics topics in complementary ways. These physics goals dictate requirements on both physical parameters and experimental design.

The origin and acceleration mechanism of UHECRs, with energies in excess of EeV, remains a fundamental outstanding question in astroparticle physics [41]. A cut-off is observed at energies around 50 EeV [42–46]. This cut-off coincides with the predictions by Greisen, Zatsepin, and Kuzmin [47, 48] for UHE protons interacting with the cosmic microwave background. Alternatively, the cut-off could be a consequence of the maximal energy attainable by cosmic accelerators. In both cases there should be a flux of "cosmogenic neutrinos" produced through interactions of UHECRs with cosmic background photons [26, 49–52] which could reach energies well above ~ 1 EeV. Measurements or constraints on the neutrino energy spectrum would provide much-needed insight into high-energy particle acceleration, the evolution of sources over cosmological length scales, and the mass composition of UHECRs.

UHECRs can also interact with gas or radiation inside the sources themselves to produce UHE neutrinos. Because their interactions are weak, these astrophysical neutrinos can travel long distances across the Universe, undisturbed, and point back to their sources, thus tracing a population of high-energy non-thermal sources over Gpc length scales. Despite many measurements of UHECRs, the sources of UHECRs are still unidentified, though they are believed to be extreme astrophysical environments where charged particles can be accelerated up to 10^{21} eV. To reveal them, complementary information from the neutrinos born in UHECR interactions may be essential. Further, because high-energy neutrinos should be copiously produced in astrophysical environments, they are also rich probes of high-energy astrophysics [7].

The current status and future prospects for diffuse flux and neutrino point source measurements, and of multi-messenger modeling, are described in Sec. 2.2 and Sec. 3.2. The quest for measurements of the cosmogenic neutrino flux and potential BSM sources of neutrinos are also outlined in Secs. 2.2 and 3.2.

1.3 Detector requirements to achieve the science goals

To probe fundamental physics and astrophysics with neutrinos, we can make several measurements with neutrinos. There are four important observables measured by neutrino telescopes that may be used, individually or jointly, to search for new physics and probe astrophysics: the energy spectrum, the distribution of arrival directions, the flavor composition, and the arrival times. Each observable has a standard expectation: respectively, a power law in energy [53], an isotropic diffuse flux [54, 55], a flavor composition at Earth of $\nu_e: \nu_{\mu}: \nu_{\tau} \approx 1: 1: 1$ [53, 56–59], and coincident arrival of neutrinos and other messengers from transient astrophysical sources [60–62]. Indeed, observations of the neutrino characteristics can be made in concert with similar observables with the other messengers — photons, cosmic rays, and gravitational waves. New physics can be imprinted on these observables in several ways as shown in Fig. 2, such that combined information from several experimental parameters can enhance discovery. The flow from physics goals to experimental parameters is shown schematically in Fig. 3.

Fundamentally, large exposures are required to achieve the required sensitivity needed to observe low neutrino fluxes, requiring an order of magnitude improvement in flux sensitivity in the HE range and two orders of magnitude improvement in the UHE range. As the detectors scale to larger volumes, it is critical to maintain low backgrounds to maximize the improvement in flux sensitivity. Energy resolutions of 0.1 in $\log_{10}(E/\text{GeV})$ at the HE scale, and in half-decade energy binning, are sufficient to resolve bumps and dips that may indicate new physics [63–69] while also distinguishing among models of astrophysical neutrino production [53, 70].

Flavor and $\nu/\bar{\nu}$ ratios provide complementary probes of new neutrino physics and neutrino production mechanisms. Because neutrinos are predominately expected from the decays of muons and charged pions, the nominal expectation is that only electron and muon neutrinos are generated at the sources, and that ν and $\bar{\nu}$ are produced in comparable numbers. After leaving the sources, oscillations over cosmological distances are expected to distribute the flux nearly evenly among all flavors by the time the neutrinos reach Earth. In reality, however, different neutrino production channels become accessible at different energies and, as result, the flavor and $\nu/\bar{\nu}$ ratios should vary with energy; see, e.g., Refs. [58, 71–80]. As a result, the expected flavor ratios at Earth might deviate from an equi-flavor composition, and might do so as a function of energy. Thus, we can use the flavor ratios measured at Earth [19, 53, 56, 81], combined with information about the values of the neutrino mixing parameters [82], to infer the flavor ratios at the sources [57, 59, 83]. However, large deviations are possible if there are BSM effects in the oscillations. They can be significantly altered by, for example, Lorentz invariance violation, neutrino decay, and neutrino interactions during propagation over long propagation distances [58, 71, 83–93, 93, 94, 94, 95]. Large event statistics and complementary flavor-specific detection techniques are needed to identify flavor-specific signals and to measure the flavor composition statistically in a sample of collected events. In the TeV–PeV range, water Cherenkov neutrino telescopes are sensitive to all flavors, though with different efficiency. We advocate for the exploration of new techniques to improve flavor separation, like muon and neutron echoes [96]. In the EeV range, some instruments will be sensitive only to certain flavors, while others will be sensitive to all flavors. We advocate for a comprehensive approach that may allow us to combine flavor information from multiple experiments.

Sub-degree pointing resolution is needed to resolve the neutrino sky [97–101] while also reducing the systematic uncertainties on cross section [17, 18, 102–107] and inelasticity measurements [19]. Resolving the neutrino sky will be important to search for BSM physics that causes anisotropies. These can be due to a variety of effects like BSM matter interactions, dark matter clumping [108–114], or Lorentz invariance violation [62, 115].

The flux sensitivity, energy resolution, and pointing resolution are all key parameters for measuring both the neutrino-nucleon cross section [17, 18, 102–107] and inelasticity [19], complementary probes of deep inelastic scattering. Key to both is a large number of events [116]. In the TeV-PeV energy range, we can expect improved precision as instruments gain in sensitivity, but at the EeV energy scale, next-generation detectors will need tens of events to measure the cross section to within an order of magnitude.

Measuring the arrival times of neutrinos is important for both time-domain transient and multi-messenger astrophysics, but also to search for evidence for new physics that would cause photons, neutrinos, and gravitational waves to arrive at Earth at different times [60–62]. Being able to capture the transient behavior of sources requires the ability for instruments to send and respond to real-time alerts. Continuous operation is ideal for detecting transient events and improving overall flux sensitivity.

To make advances, we plan to build on existing, mature experiments while also exploring new technologies. There are several operating and planned experiments, many of which are complementary in terms of their energy range, flavor sensitivity, and sky coverage. We describe them below and in Sec. 4.



Figure 4. Selected present-day measurements of high-energy gamma rays by Fermi-LAT [117], high-energy neutrinos by IceCube [21, 118, 119], and ultra-high-energy cosmic rays by the Pierre Auger Observatory [120] and the Telescope Array (TA) [121]. The three neutrino measurements are independent and utilise different event topology: contained cascades (solid points), partially-contained cascades (light blue diamonds) and through-going tracks (blue crosses). Error bars and upper limits represent 68% confidence intervals. For the track sample only bins in the sensitive energy range are shown while the fit was done from 100 GeV to 100 PeV.

1.4 Present and future experimental landscape

Figure 4 shows the present-day landscape of measurements of high-energy cosmic messengers: gamma rays, neutrinos, and ultra-high-energy cosmic rays. A complete picture of the high-energy Universe is necessarily multi-messenger in nature. Below, we focus on neutrinos, but point out scenarios where gamma rays and cosmic rays offer complementary information.

IceCube, presently the largest neutrino telescope, is an in-ice Cherenkov detector in Antarctica. It instruments 1 km³ of deep underground ice with thousands of photomultipliers that collect the light emitted by particle showers initiated by high-energy neutrino interactions. From the amount of light collected and its spatial and temporal profiles, IceCube infers the energy, flavor, and arrival direction of the neutrinos. Because the bulk of their arrival directions is broadly consistent with an isotropic distribution, the diffuse neutrino flux that IceCube sees is likely of predominant extragalactic origin, though the sources are unknown, save for two promising source associations [22–24]. In particle physics, IceCube has measured the TeV–PeV neutrino-nucleon cross section [17, 18, 122] and inelasticity distribution [19] for the first time, probed charm production in neutrino interactions [19], and seen hints of the first high-energy ν_{τ} [123], and the Glashow resonance (indicating $\bar{\nu}_e$) [20, 21]. ANTARES, a Cherenkov detector in the Mediterranean Sea operating until recently, nears the sensitivity to the IceCube diffuse neutrino flux [16, 124]. Three new telescopes under construction, KM3NeT [124, 125], P-ONE [126], and Baikal-GVD [127], will improve our sensitivity to TeV–PeV neutrinos to the Southern Sky. The next generation of IceCube, IceCube-Gen2 [128], will improve our sensitivity across a broad energy range from the TeV scale to the EeV scale.

In the next 10–20 years, new detectors may improve our sensitivity to neutrino energies above the energy range where that of IceCube becomes too small to detect a significant flux. There are several planned experiments targeting the PeV energy range to determine the high-energy spectrum of the astrophysical flux observed by IceCube. Observation of a spectral cut-off or the continuation of the power-law spectrum would help reveal the sources of these neutrinos while also extending our observations of neutrinos into a new

energy scale. These experiments take different approaches including radar (RET-N [129]), particle showers in a valley (TAMBO [130]), and Earth-skimming tau neutrinos from a mountain (Trinity [131], Ashra NTA [132], CTA [133]), balloons (EUSO-SPB [134]), and satellites (POEMMA [135]).

EeV neutrinos have long been predicted as coming from the interaction of ultra-high energy cosmic rays with cosmic photon backgrounds [26, 47, 48] or photons inside astrophysical sources [136–141], but they have not been discovered yet [142–146]. The flux of these neutrinos is expected to be low [136–141, 147–149]. At even higher energies, ZeV neutrinos might come from cosmic strings [150, 151]. Next-generation, multipurpose EeV–ZeV detectors are being planned or under construction with a variety of detection strategies: in-water and in-ice Cherenkov (IceCube-Gen2 [128]), in-air Cherenkov, fluorescence, and particle showers (EUSO-SPB2 [134], POEMMA [135], AugerPrime [152], GCOS [153]), and radio (GRAND [154], RNO-G [155], PUEO [156], BEACON [157], TAROGE [158], AugerPrime [152], GCOS [153]).

Current and planned detectors are detailed in Sec. 4.

2 Current status and lessons learned

2.1 In particle physics

HE neutrino interactions: With the beam of neutrinos at TeV to PeV energies, we are already exploring tests of neutrino interactions in a new energy regime through their cross sections [17, 18, 122], inelasticity distributions [19], and the Glashow resonance [21, 159]. As statistics grow and new experiments come online, more subtle BSM effects can be explored.

UHE neutrino interactions: The interactions of UHE neutrinos with nucleons have center-of-mass energies of $\sqrt{s} \sim 30$ TeV (vs. ~1 TeV using TeV–PeV neutrinos [17, 18]), providing an excellent opportunity to probe models of the nucleon and nuclear structure [102, 160–162], and new physics in neutrino-nucleon interactions [17, 18, 103, 116, 161–181].

Flavor transitions at the highest energies: Both astrophysical and cosmogenic neutrinos are expected to arrive at Earth with nearly equal fluxes of each of the three flavors, $\nu_e : \nu_\mu : \nu_\tau = 1 : 1 : 1$, as a result of the neutrino production processes at the sources and the values of the oscillation parameters measured in terrestrial experiments. As some experiments have flavor sensitivity (particularly for ν_τ), UHE neutrinos provide a key test for flavor changing processes up to the highest energies [58, 85–94].

Tests of neutrino properties during propagation: Some of the fundamental properties of neutrinos may have an energy dependence that manifests only at the highest energies. Notable examples that have received attention include the breaking of known symmetries or the appearance of new ones [88, 182–193], neutrino self-interactions [64, 65, 67, 91, 194–198], neutrino-dark matter interaction [198–216], and neutrino-dark energy interaction [217, 218].

Below we focus on a selection of tests with high-energy neutrinos that have the potential to yield important new physical insight.

2.1.1 Cross sections

The cross section for neutrino interactions with nucleons is a unique probe of SM and BSM physics. In SM physics, measuring the cross section probes the parton distribution functions (PDFs) indirectly. In BSM physics, measuring the cross section may identify dramatic deviations predicted by various models [151, 165, 167, 180]. Figure 5 shows the neutrino-nucleon interaction cross section, $\sigma_{\nu N}$, measured from GeV to PeV energies, and its projected measurements at hundreds of PeV, compared to a recent SM prediction [161]. In recent years, significant work has gone into predictions for the cross section both within and outside of the SM, motivated by improvements in the determination of PDFs from new collider data and enhanced analysis techniques, and anticipating the advent of next-generation neutrino experiments.



Figure 5. Neutrino-nucleon cross section measurements, compared to deep-inelastic-scattering (DIS) cross section predictions from Ref. [161] (BGR18). In the TeV range, FASER and FASER ν have started measurements [219]. Measurements in the TeV–PeV range are based on IceCube showers [18, 122] and tracks [17]. Projected measurements at energies above 100 PeV [106] are based on 10 years of operation of the radio component of IceCube-Gen2, assuming a resolution in energy of 10% and a resolution in zenith angle of 2°. Since the flux at these energies remains undiscovered, projections for the measurement of the cross section are for different flux predictions. Figure adapted from Ref. [106].

At energies above a few GeV, neutrinos primarily interact with matter via deep inelastic scattering (DIS), where a neutrino exchanges a W (charged-current) or Z (neutral-current) boson with a parton of a nucleon, i.e., a quark or a gluon [220–223]. The differential cross section for this process can be expressed in terms of DIS structure functions, which describe the underlying QCD dynamics of the nuclear medium. The structure functions depend on Bjorken-x—the fraction of nucleon momentum carried by the interacting parton and on Q^2 —the four-momentum transferred to the mediating boson. Structure functions are computed by convoluting PDFs with coefficient functions using perturbation theory. In the last two decades, different groups have studied high-energy neutrino cross sections [102, 160, 161, 224–228] and they have identified several effects that play a significant role in this calculation.

Charm- and top-quark production are important in high-energy neutrino-nucleon interactions. Therefore, heavy-quark mass effects must be calculated to provide an adequate description. At leading order, the slow-rescaling and the modification of light cone momentum fraction in the PDFs must be included [229]. At higher orders, different formalisms have been developed to account for this effect [230–232]. Nevertheless, the contribution of charm or top production can be significantly different depending on the approach [233, 234].

PeV and EeV neutrinos probe the small-x and high- Q^2 region of DIS, a kinematic region with very limited data available from colliders and fixed-target experiments to perform PDF fits [235]. State-of-the-art PDFs have significantly lowered the minimum value of x used to $\mathcal{O}(10^{-8})$ [235–238], but extrapolation must be done if lower values are probed. Uncertainties in the extrapolation of PDFs translate into uncertainties in the predicted cross section, especially at the highest neutrino energies. In addition, the stability of the perturbative expansion is spoiled in the very-small-x region, so resummation corrections must be included in the DGLAP formalism [161, 239–241]. An efficient way to account for resummation and saturation effects is the color dipole approach [225–228]. Depending on the formalism and input PDFs, the predictions can differ by a factor of 3 in the ultra-high-energy regime; see, e.g., Fig. 5 and Fig. 3 in Ref. [18].

High-energy neutrinos are attenuated by interacting with matter during their passage through Earth; the neutrino-nucleon cross section is extracted from measuring the attenuation. The flux is attenuated more strongly the higher the neutrino energy and the longer the path traversed by the neutrino inside the Earth. This feature can be exploited to probe the neutrino cross section at high energies. The first measurement of the neutrino cross section from 6.3 TeV to 980 TeV was done by IceCube using up-going muon neutrinos [17]. Later, two independent measurements were carried out using starting events with energies between 60 TeV and 10 PeV [18, 122]. The results are compatible with the SM predictions described previously. However, it is difficult to draw conclusions beyond that due to the large uncertainties of these measurements.

The neutrino cross section measured in the TeV–PeV region using the astrophysical flux of neutrinos discovered by IceCube has improved our understanding of the cross section since the last Snowmass study a decade ago [242]. We can expect that in the coming decade, improved precision with IceCube, KM3NeT, and Baikal-GVD will reduce the systematic uncertainties, but not necessarily improve the energy reach. The cross section can be measured either by assuming a known flux and inferring the cross section from the observed number of events [104], or by comparing the absorption in the Earth as a function of the observed trajectories that neutrinos travel through [17, 18, 102–107, 122]. The latter requires significant absorption of neutrinos through the Earth and therefore has an energy threshold of 5–10 TeV.

In the PeV range, W-boson production becomes relevant from two processes: electron anti-neutrino scattering on atomic electrons, and neutrino-nucleus interactions in which the hadronic coupling is via a virtual photon. The former produces the distinct Glashow resonance [159], which peaks at 6.3 PeV. The latter can reach up to 5-10% of the DIS cross section in the PeV range [243–247], and it becomes increasingly more important for heavy nuclei as it scales with Z^2 .

IceCube recently reported the detection of a particle shower compatible with it being due to a Glashow resonance [21], with a deposited energy of 6.05 ± 0.72 PeV. The inferred neutrino energy is ~6.3 PeV, after correcting for the invisible energy from particles that do not radiate visible photons. Signatures of low-energy muons were also observed in the event, with energies consistent with expectations of a W^- decay, tagging it as a hadronic cascade. Future analyses with more data could reach a 5σ detection. On the other hand, W-boson production [243, 244, 246, 247] deserves more attention, especially in the near future as much more data is collected. In fact, recent studies have shown that these interactions can play a significant role in the detection of tau neutrinos from cosmic origin [248].

Going beyond standard DIS interactions, it is well known that the quark and gluon PDFs of nucleons bound in nuclei are modified compared to free nucleons. Therefore, high-energy neutrino interactions in ice, water, or rock are modified due to the presence of nuclear effects. The most relevant effect for the calculation of the neutrino DIS is that of shadowing [249], namely, the depletion of nuclear structure functions as compared to their free-nucleon counterparts. In the last years, several collaborations have produced PDF sets in the nuclear sector [250–253]. First calculations of the neutrino cross section using nuclear PDFs find 5–15% suppression in the PeV range, but the uncertainty of the nuclear corrections are still large [161, 234, 254]. Future Electron-Ion Collider (EIC) measurements of nuclear structure functions will reduce the uncertainties in the neutrino cross section due to nuclear corrections [255].

2.1.2 Inelasticity

Inelasticity is the measure of the fraction of neutrino energy transferred to a hadronic target in deep inelastic scattering. The cross section and inelasticity are complementary probes of new physics since a new interaction should have an inelasticity distribution distinct from that expected from conventional charged current deep



Figure 6. Constraints on invisible neutrino decay from a range of experiments. The blue region represents a hint for neutrino decay [268, 269] in IceCube data. Dashed regions represent anticipated sensitivities from future measurements. CMB constraint depends sensitively on the details of the decay; see Ref. [264]. Adapted from Ref. [269].

inelastic scattering, which is modeled well in the Standard Model [256]. Recent results from IceCube made the first measurement of the ν_{μ} inelasticity distribution above 1 TeV by tracking both muons from a chargedcurrent interaction and the shower deposited in the detector [19]. This inelasticity data was also used to constrain the atmospheric $\bar{\nu} : \nu$ ratio. With more data and good atmospheric neutrino rejection, it may be possible to apply this calculation also to astrophysical neutrinos.

2.1.3 Neutrino decay

Because neutrinos have mass, they decay. In the Standard Model, they can decay radiatively, e.g., $\nu_j \rightarrow \nu_i + \gamma$ [257, 258], where ν_j and ν_i are different neutrino mass eigenstates. However, the associated lifetimes are longer than the age of the Universe. Yet, if neutrinos couple to a new light or massless mediator, then the decay rate could be enhanced. This can be tested in a variety of environments. In general, terrestrial and solar constraints are not very strong. The strongest existing constraints come from the cosmic microwave background [259–262], although the bounds may be significantly relaxed [263, 264] and there may even be hints of neutrino decay [260]. In addition, these constraints are somewhat more model-dependent than the others. A measurement of the diffuse supernova neutrino background would provide the next most sensitive probe of neutrino decay [265, 266], followed by a Galactic supernova, although lifetime bounds from SN1987A [267] can be evaded depending on the flavor structure. The existing constraints from terrestrial and astrophysical experiments as well as projected sensitivities are shown in Fig. 6.

After that, the next most relevant constraint comes from high-energy astrophysical neutrinos observed at IceCube [93, 268–273]. Both the fact that neutrinos have been detected as well as the detailed spectral and flavor information have been used to probe neutrino decay. A weak hint for neutrino decay was identified by comparing the spectra of different flavors of neutrinos which makes certain predictions, in particular for the tau neutrino flux [268, 269].

2.1.4 Dark matter

Weakly interacting massive particles (WIMPs) are the primary candidates for particle dark matter (DM). The WIMP hypothesis yields a thermally average cross-section rate, $\langle \sigma v \rangle \simeq 3 \times 10^{-26} \,\mathrm{cm^3 \ s^{-1}}$, which can explain the observed relic abundance after the freeze-out and is independent of the annihilation products. Meanwhile, thermal production of WIMPs in the early Universe implies possible ongoing annihilation of DM to Standard Model (SM) particles. This possibility has facilitated the indirect search for dark matter. There is a distinct possibility that neutrinos might be the principal portal to the dark sector. Such possibility is motivated by the *scotogenic* models where the neutrinos mass is achieved via interaction with DM, see, e.g.,



Figure 7. Constraints on the velocity-averaged annihilation cross section, $\langle \sigma v \rangle$, of supra-TeV dark-matter particles, with mass m_{χ} , into neutrinos. Solid and dashed lines represent, respectively, 90% C.L. existing limits and sensitivities. Figure reproduced from Ref [216]; heart symbols show new constraints introduced therein.

Ref. [274]. Furthermore, the upper limit on the DM annihilation cross-section to neutrinos serves as an upper bound on DM annihilation to SM particles, as the latter is larger [275, 276].

In principle, DM could annihilate to all SM particles. Annihilation to most SM particles results in the production of gamma rays and neutrinos. While the most constraining limits on DM annihilation cross-section for sub-PeV DM are obtained from the absence of signal in multi-wavelength observations, especially from the Milky Way and its satellite galaxies, neutrino observations are shown to be providing stronger probes of very heavy DM with masses > PeV [277]. Moreover, there is a distinct possibility for DM directly annihilating to neutrinos, which makes indirect search via neutrinos important.

Remarkably, for energies ≥ 0.1 MeV, there exists an interrupted coverage of the neutrino flux from the Universe. An extraordinary amount of data has been collected in this range for measuring neutrino fluxes which can be utilized to search for DM annihilation to neutrinos which have been largely used to impose constraints on DM annihilation to neutrinos. While for the low-mass regime ($m_{\rm DM} < \text{TeV}$), the limits are already approaching the thermal relic density values (see [216] for details), the current and upcoming neutrino telescopes aiming at very high energies are opening a new avenue for the indirect search of DM.

Figure 7 shows the upper limits on DM annihilation cross section for DM particles heavier than 1 TeV. Currently, the upper limit on DM annihilation cross section is obtained from the observations of ANTARES and IceCube neutrino observatories. For DM in the mass range of 1–100 TeV, ANTARES dominates the upper limit landscape, thanks to its optimal location to observe the Galactic center. For heavier masses, the best constraints are given by IceCube observation of high-energy cosmic neutrinos, as well as IceCube's search for extremely-high-energy (EHE) neutrinos. The ability of neutrino telescopes to probe DM annihilation will be significantly improved with the upcoming and planned neutrino experiments.



Figure 8. Mean free path of UHE neutrinos in $C\nu B$ field. Only the SM neutrino-neutrino interactions are involved. The sharp resonance structure appears due to the interactions via s-channel of Z^0 boson. The possible BSM interactions involving neutrinophilic mediator are expected to form a similar structure at $E_{\nu} \gtrsim 10^6$ GeV range, which would distort spectral shape of neutrinos. Figure reproduced from Ref. [278].

2.1.5 Secret neutrino interactions

While the SM allows for interactions among neutrinos, these interactions are all highly suppressed by the electroweak scale – any interaction cross section involving only neutrinos scales as the square of the Fermi constant, G_F^2 . It is still unknown whether there are additional BSM interactions solely among neutrinos that are stronger than this, with some effective interaction scale $G_{\text{eff.}} \gg G_F$. In a UV-complete BSM model, this implies the existence of some new electrically-neutral mediator that couples to neutrinos, significantly lighter than the Z^0 boson.

New interactions of this type can significantly modify the character of the HE and UHE neutrino flux arriving at the Earth by the scattering of the HE/UHE neutrinos off of nearly-at-rest cosmic neutrino background (C ν B) neutrinos. The main feature of this effect is the absorption of neutrinos with a characteristic energy $E \approx m_{\phi}^2/(2m_{\nu})$, where m_{ϕ} is the mass of the new neutrinophilic mediator and m_{ν} is the mass of one of the light neutrino eigenstates. The spectral distortion features may appear at PeV-EeV region, depending on the absolute mass of neutrinos. This model is independently motivated by the neutrino mass generation mechanism [195], muon g - 2 anomaly [279], small-scale problems in dark matter substructures [280, 281], and apparent Hubble tension [282–285]. A standard model example of a sharp resonance structure in the mean free path of neutrinos scattering in the C ν B field via Z^0 boson production is shown in Fig. 8.

If we assume $m_{\nu} = 0.1$ eV and $m_{\phi} = 10$ MeV, this predicts absorption of HE, 500 TeV neutrinos. This effect has been explored in, for instance, Refs. [63–65, 67, 68, 194, 195, 197, 198, 286–289]. Beyond the absorption of a characteristic energy of HE/UHE neutrinos, additional effects can be detected, including additional neutrino flux below this energy from a cascade effect [64, 65] and a delayed arrival of post-scattering neutrinos relative to the ones that do not scatter [198].

Constraints on such interactions using current IceCube data have been derived in Refs. [67, 69]. Depending on the flavor structure of the interaction between this new mediator and neutrinos of flavor α and β , these searches can place the most stringent constraints on the interactions for mediator masses between roughly 4 and 40 MeV, even stronger than some laboratory and cosmology constraints [284, 290, 291]. In general, the measurement of astrophysical HE/UHE neutrinos constraints $G_{\text{eff}} \leq 10^{10}G_F$ given current data.



Figure 9. Prompt atmospheric neutrino fluxes for $\nu_{\mu} + \bar{\nu}_{\mu}$ assuming a broken power law (BPL) and the H3a [293] cosmic ray flux. The dashed lines show the contributions to the flux from different collider center-of-mass rapidity ranges for charm production, described in more detail in Ref. [294].

2.1.6 Prompt neutrinos

Prompt neutrinos come from decays of heavy-flavor hadrons, predominantly from charm mesons. In nature, prompt neutrinos can be produced through cosmic-ray interactions with nuclei in the Earth's atmosphere. Due to very short lifetimes of heavy-flavor hadrons, the energy dependence of the flux of prompt atmospheric neutrinos scales roughly according to the cosmic-ray flux, while the energy dependence of conventional atmospheric neutrinos that come from pion and kaon decays falls with an additional power of $1/E_{\nu}$. At energies of $E_{\nu} \gtrsim 100$ TeV-1 PeV, the prompt atmospheric neutrino flux dominates over the conventional neutrino flux and becomes the main component of the flux of atmospheric neutrinos. Meanwhile, high-energy neutrino telescopes such as IceCube and KM3NeT search for astrophysical neutrinos at TeV-10 PeV energies [34-36, 118, 292]. Probing the prompt atmospheric neutrinos is not only interesting for detection itself, but also important in that they are the primary background to astrophysical neutrinos.

Prompt atmospheric neutrinos have not yet been experimentally detected, and the theoretical prediction of their flux has large uncertainties that depend on various factors such as the cosmic-ray spectrum and mass composition, the model for heavy-flavor production, and the parton distribution functions [295–298]. One of most significant contributions to large uncertainties in the prompt neutrino flux originates from the limited knowledge of relevant kinematic regions of heavy-flavor production cross sections in hadron collisions. In evaluating the prompt neutrino flux, models for heavy-flavor production for the charm meson production cross section are compared with the data collected by the LHCb Collaboration [299], which provides data in the most forward region so far, for charm meson rapidities of 2.0 < y < 4.5. To illustrate the charm meson rapidities relevant to the prompt atmospheric neutrino flux as a function of neutrino energy, Fig. 9 shows the prompt atmospheric neutrino fluxes for $\nu_{\mu} + \bar{\nu}_{\mu}$ from the charm produced at different collider rapidity ranges in *pp* collisions [294]. These results are evaluated with one of the modern cosmic-ray spectra analyzed taking into account cosmic-ray composition, H3a [293]. The result with the traditional broken power law (BPL) spectrum is presented for comparison. In the figure, one can see that for the energies of $E_{\nu} \gtrsim 10^5 - 10^6$ GeV, where the prompt neutrinos are most important, the flux depends on charm produced in the equivalent collider rapidity range of $y \gtrsim 4.5$.

As noted, current experimental data for charm production are available for y < 4.5 from LHCb measurements. Recently, two experiments at the LHC, FASER ν [300] and SND@LHC [301], were approved.





Figure 10. Expected measurements of the flavor composition measured at Earth in both 2020 and 2040. Lines show the 99.7% credibility regions of using the astrophysical neutrino flux assuming a composition at Earth of $0.3\nu_e$: $0.36\nu_{\mu}: 0.34\nu_{\tau}$. These can be compared to predictions from different assumed source compositions (π decay, μ -damped π decay, and n decay dominated scenarios), unitarity, and standard oscillations. These predictions will improve over time as oscillation parameters are further refined with new experiments. Figure reproduced from Ref. [83].

Figure 11. Expected flavor ratios observed at Earth under a generic class of new physics assumptions, which can include Lorentz invariance and CPT violation, violation of the equivalence principle, cosmic torsion, and non-standard interactions, among others. The colors represent different assumed flavor compositions at the sources. Figure reproduced from Ref. [90].

They are in preparation to probe more forward regions, $y \gtrsim 7.2$, during Run 3 of the LHC. Already in 2018, a pilot detector for FASER ν collected data from 12.2 fb⁻¹ of pp collisions at the LHC, from which candidate neutrino interaction events were observed [302]. Combined, the Run 3 measurements by FASER ν and SND@LHC will allow for tests of theoretical approaches to forward charm production, for example as reviewed in Refs. [6, 303], to refine theoretical predictions of the prompt atmospheric neutrino flux.

2.1.7 Fundamental symmetries

The Standard Model rests on two fundamental symmetries: the Lorentz and CPT symmetries. According to the CPT theorem, any Lorentz-invariant local quantum field theory with a Hermitian Hamiltonian, like the Standard Model, must respect CPT symmetry. So far, all attempts to find experimental evidence for Lorentz-invariance breaking have failed.

However, because the Standard Model is regarded to be an effective field theory, the Lorentz symmetry upon which it rests might not be truly fundamental, but might be broken at high energies. The full theory of which the Standard Model is an effective one is presently known, but effective-theory extensions of the Standard Model exist that, while not ultraviolet-complete, contain additional Lorentz-violating interactions. The Standard Model Extension (SME) is the best-developed effective field theory of this kind. In it, the coupling strength of the Lorentz-invariance-violating (LIV) interactions, i.e., their Wilson coefficients, have a priori undetermined values. The SME is regularly used to systematically test Lorentz invariance, by constraining the values of the couplings. For neutrinos, the presence of LIV modifies flavor-transition probabilities, possibly significantly. For high-energy neutrinos, the presence of LIV may manifest as deviations in flavor composition of the neutrino flux that reaches Earth, i.e., of the relative amount of ν_e , ν_{μ} , and ν_{τ} in the flux.

In the standard case, i.e., if Lorentz symmetry holds, the Hamiltonian that drives flavor transitions in vacuum, is

$$H_{\rm std} = \frac{1}{2E} U {\rm diag}(0, \Delta {\rm m}_{21}^2, \Delta {\rm m}_{31}^2) {\rm U}^{\dagger} , \qquad (2.1)$$

where E_{ν} is the neutrino energy, $\Delta m_{21}^2 \equiv m_2^2 - m_1^2$ and $\Delta m_{31}^2 \equiv m_3^2 - m_1^2$ are the mass-squared differences between neutrino mass eigenstates, and U is the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix that connects the neutrino mass and flavor bases. The PMNS matrix is conventionally parametrized via four mixing parameters: three mixing angles, θ_{12} , θ_{23} , and θ_{13} , and one CP-violation phase, δ_{CP} , whose values are known from oscillation experiments. For high-energy cosmic neutrinos, the flavor-transition probability derived from the above Hamiltonian oscillates rapidly. Because of limited precision in the knowledge of the distance traveled from its source, and in the measurement of neutrino energy, in practice neutrino telescopes are sensitive only to the average flavor-transition probability. For the transition $\nu_{\alpha} \rightarrow \nu_{\beta}$ ($\alpha, \beta = e, \mu, \tau$), this is

$$P_{\alpha\beta}^{\text{std}} = \sum_{i=1}^{3} |U_{\alpha i}|^2 |U_{\beta i}|^2 , \qquad (2.2)$$

where $U_{\alpha i}$ and $U_{\beta i}$ are components of the PMNS matrix. Thus, the standard average flavor-transition probability is energy-independent, and depends only on the standard mixing parameters.

If neutrinos are produced with a flavor composition of $f_S \equiv (f_{e,S}, f_{\mu,S}, f_{\tau,S})$, then due to oscillations, the flavor composition at Earth will be $f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\beta\alpha}^{\text{std}} f_{\beta,S}$, for $\alpha = e, \mu, \tau$. Figure 10 shows the expected flavor ratios at Earth for three benchmark neutrino production scenarios, and accounting for the uncertainties in the neutrino mixing parameters.

The LIV operators in the SME introduce additional terms to the Hamiltonian, i.e.,

$$H_{\rm LIV} = \sum_{n} \left(\frac{E_{\nu}}{\Lambda_n}\right)^n \tilde{U}_n \operatorname{diag}(O_{n,1}, O_{n,2}, O_{n,3}) \tilde{U}_n^{\dagger} , \qquad (2.3)$$

where $O_{n,i}$ are the eigenstates of the *n*-dimensional Lorentz-violating SME operator, \tilde{U}_n is a PMNS-like mixing matrix, but parametrized by new mixing parameters, ξ_i , and Λ_n is the energy scale characteristic of the *n*-dimensional operator. For instance, n = -1 corresponds to neutrino decay; n = 0, to CPT-odd LIV; and $n \ge 1$ to CPT-even LIV. Equation (2.3) reveals the advantage of using high-energy neutrinos to look for LIV: because $H_{\text{LIV}} \propto E_{\nu}^n$, with $n \ge 0$, its relative importance compared to H_{std} grows with energy. Therefore, even if the values of the LIV parameters are tiny, the LIV effects may become apparent at high energies. Below, we show how this would manifest.

The total Hamiltonian is $H_{\text{tot}} = H_{\text{std}} + H_{\text{LIV}}$. Unlike H_{std} or H_{LIV} by themselves, the total Hamiltonian H_{tot} is no longer diagonalized by the PMNS matrix U or by a group of U_n matrices, but rather by a new matrix, V, that depends on both the standard and LIV mixing parameters. Because of H_{LIV} , the average flavor-transition probability depends not only on the standard mixing parameters, but also on the LIV parameters and the neutrino energy, i.e.,

$$P_{\alpha\beta}^{\rm LIV} = \sum_{i=1}^{3} |V_{\alpha i}|^2 |V_{\beta i}|^2 , \qquad (2.4)$$

where $V_{\alpha i} \equiv V_{\alpha i}(\theta_{12}, \theta_{23}, \theta_{13}, \delta_{CP}, \xi_i, E_{\nu})$ and similarly for $V_{\beta i}$.

Analogously to the standard-oscillation scenario, the flavor composition at Earth with LIV is $f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\beta\alpha}^{\text{LIV}} f_{\beta,\text{S}}$. Figure 11 shows the flavor ratios at Earth in the presence of LIV, for the same benchmark production scenarios. Because the values of the LIV parameters are unknown, when they are allowed to vary large deviations from the standard flavor-composition expectation are possible. Therefore, detecting a

large deviation away from the standard $(f_{e,\oplus}, f_{\mu,\oplus}, f_{\tau,\oplus}) = (1/3, 1/3, 1/3)$ could be interpreted as being due Lorentz-invariance violation.

Another possible test of fundamental symmetries comes from augmenting the Standard Model with a new U(1)' gauge symmetry [304–306]. Such an extension results in a new gauge boson Z' responsible for a new interaction between neutrinos and other elementary massive particles. In the ultra-light limit of the Z'mass, the resulting interaction is long-range in nature, which would affect the propagation of high-energy astrophysical neutrinos. If the Z' exists, neutrinos will in addition interact with matter particles while propagating through the Universe, thus modifying the standard flavor composition f_{\oplus} at Earth [307].

Some well explored BSM extensions are U(1)' models with $L_{\alpha} - L_{\beta}$ ($\alpha, \beta = e, \mu, \tau$) symmetries, where L_{α} is the α -flavor lepton number. For $L_e - L_{\beta}$ ($\beta = \mu, \tau$), neutrinos would interact with electrons via the Z' vector boson subject to the potential $V_{e\beta} = \alpha_{e\beta} \frac{N_e}{R}$, where $\alpha_{e\beta} = g'^2/4\pi$ is the fine structure constant for $L_e - L_{\beta}$ symmetries. However, for $L_{\mu} - L_{\tau}$, since muons and tauons are short-lived and, therefore, rarely encountered by astrophysical neutrinos, the interaction between neutrinos and electrons is only possible via Z - Z' mixing, where Z is the SM-boson of the weak interaction. The effective potential is then given by $V_{\mu\tau} = \pm g'(\xi - s_w \chi) \frac{e}{4s_w c_w} \frac{N_n}{4\pi R}$ where $(\xi - s_w \chi)$ is due to the Z - Z' mixing [308].

In the presence of the new interaction potential, the neutrino propagation Hamiltonian is described by the matrix $H = H_{vac} + V_{mat} + V_{\alpha\beta}$, where H_{vac} is the vacuum neutrino propagation Hamiltonian in the flavor basis, and V_{mat} is the standard neutrino-matter charged-current interaction term. The new interaction matrix $V_{\alpha\beta}$ is the flavor-diagonal matrix having non-zero $\alpha\alpha$ and $\beta\beta$ elements as strength of the respective potential with opposite sign.

Astrophysical neutrinos are ideal probes because they travel cosmological distances over which any minute extension to the SM can potentially accumulate to a measurable effect. Any measured deviation from the standard flavor composition could, therefore, be translated into measurements of the parameters of U(1)' models. Several of the operating neutrino telescopes like IceCube [309], KM3NeT [125], Baikal [310] with sensitivity to >TeV astrophysical neutrinos and their flavors are able to test for f_{\oplus} deviations.

2.1.8 ANITA anomalous events

ANITA has observed a handful of anomalous events; these are events whose observational properties are, at least at first sight, compatible with them being neutrinos. ANITA is a balloon-borne Antarctic UHE particle detector sensitive to the radio emission from neutrino interactions in the ice and air showers that can be generated by both cosmic rays and Earth-skimming neutrinos. The ANITA anomalies come in two types: steeply upcoming air shower and near-horizon air shower events. Air-shower events in ANITA are observed both directly pointed at the payload and reflected off the ice; they can be distinguished with their observed arrival direction and polarity (or phase) inversion due to the reflection off of the ice in the later case. Two events—one each in both the first [311] and third ANITA flights [312]—were observed that were consistent with the geomagnetic signal from an air shower in terms of their polarization and spectrum, but their steep arrival direction (roughly 30° below the horizon) were inconsistent with their observed polarity. In the fourth flight of ANITA, four events near, but below the horizon were observed with an inconsistent polarity indicating that the air shower signal had not been reflected with a significance of $2-3\sigma$ level when considering possible anthropogenic backgrounds and pointing and polarity reconstruction errors.

Several analyses considered the possibility that the steeply upcoming anomalies are due to a possible Earth-skimming tau neutrino origin and find that that they are in strong tension with predictions from Standard Model cross sections and experimental limits at UHE (including the in-ice neutrino channel within the ANITA) [311, 313–319]. This is true for both the diffuse [317, 318] and point-source neutrino hypotheses [318, 320]. Limits from the Pierre Auger Observatory consider the aperture to both generic upgoing air showers [315] and tau lepton induced air showers [314].

The near-horizon ANITA anomalies are more consistent with an Earth-skimming neutrino origin than the steep events in terms of the distributions of their arrival direction and spectral characteristics, but the fluence

implied by the detection of four events in one flight is in strong tension with the Pierre Auger Observatory across all energies and is also in tension with ANITA's own in-ice neutrino limits above $10^{19.3}$ eV [319].

Several explanations have been proposed to explain the steeply upcoming anomalous events. Mundane effects such as transition radiation of cosmic ray air showers piercing the Antarctic ice sheet [321, 322] and subsurface reflections due to anomalous ice features [323] may be possible, although the later is disfavored by the observed waveforms of the events [324]. Beyond-the-Standard Model explanations predict the expected signature of new particles exiting the Earth [325–333]. As of this white paper, the origin of both sets of anomalous events remains unclear; follow-up observations of these unusual events are well-motivated [315].

2.2 In astrophysics

Particle acceleration in extreme environments: The origin of cosmic rays and the associated particle acceleration mechanisms are among the biggest open questions in the astroparticle physics. Possible mechanisms including shock acceleration [334], one-shot/shear acceleration [335, 336], and magnetic reconnection [337]. In most cases, the accelerated cosmic rays will interact with surrounding matter (pp) [338, 339] and light $(p\gamma)$ [49] inside their sources, and produce high-energy neutrinos as a result. Thus, high-energy neutrinos provide a unique probe of acceleration mechanisms in astrophysical environments.

Inside astrophysical sources, pp and $p\gamma$ interactions make charged pions that, upon decaying, make highenergy neutrinos: $\pi^+ \to \mu^+ + \nu_{\mu}$, followed by $\mu^+ \to e^+ + \nu_e + \bar{\nu}_{\mu}$, and the charge-conjugated processes. (The beta-decay of neutrons from in pp and $p\gamma$ interactions makes additional $\bar{\nu}_e$, but those neutrinos have an energy approximately a hundred times smaller than neutrinos from pion decay.) The final-state neutrinos typically carry $\sim 3 - 5\%$ of the parent proton energy, given that energy losses of pions and muons are negligibly small. (The exception is when magnetic fields are intense and synchrotron losses of charged particles are significant; see, e.g., Refs. [340, 341].) Thus, the detection of PeV neutrinos enables us to study the sources of cosmic rays around ~ 100 PeV, i.e., around the second knee of the cosmic-ray spectrum, possibly probing the Galactic-to-extragalactic transition in cosmic rays. Higher-energy neutrinos, especially UHE neutrinos in the EeV range, will be crucial to investigate the accelerators of UHECRs. UHE gamma rays would not be directly observed except for nearby UHECR accelerators within 10–100 Mpc, while UHE neutrinos can be used to study distant sources located in the cosmological distance. Gamma-ray bursts (GRBs), tidal disruption events (TDEs), and blazars are among the candidate accelerators of UHECRs. However, models that may explain the observed neutrinos and UHECRs often require a large baryonic loading, i.e., a large fraction of the available energy imparted to cosmic rays, which may be theoretically challenging.

Neutrinos are also important as a probe of dense environments that are not visible with photons. Cosmic rays can be accelerated in the vicinity of a black hole [342–344], a jet inside a star [345], and dense circumstellar material [346]. There has been significant progress in numerical simulations of particle acceleration in shocks and accretion flows [347–349], and in spark gaps [350, 351]. Non-jetted AGN [352–355], TDEs [356–358], choked jets [359–362], and interacting supernovae [363, 364] are actively discussed as promising targets for high-energy neutrino detection.

Flavor physics at sources: From the full pion decay chain, we nominally expect the flavor composition leaving the neutrino sources to be $f_{e,S} : f_{\mu,S} : f_{\tau,S} = 1 : 2 : 0$, where $f_{\alpha,S}$ is the fraction of $\nu_{\alpha} + \bar{\nu}_{\alpha}$ $(\alpha = e, \mu, \tau)$ at production. However, because neutrinos oscillate, the flavor composition that reaches Earth is different. In particular, for high-energy neutrinos traveling over cosmological-scale distances, the flavortransition probabilities are averaged [365]. As a result, the nominal expectation is for the flavor composition at Earth to be the same for all flavors, i.e., $f_{e,\oplus} : f_{\mu,\oplus} : f_{\tau,\oplus} \approx 1 : 1 : 1$ [85, 366]. There are relatively small variations around this prediction due to uncertainties in the values of the neutrino mixing parameters [58, 83]. However, in highly magnetized sources such as GRBs, pions and muons cool via synchrotron so strongly that the flavor composition at Earth can be modified to $f_{e,\oplus} : f_{\mu,\oplus} : f_{\tau,\oplus} \approx 1 : 1.8 : 1.8$ at high energies [58, 73, 83]. Alternately, in high-gradient sources (more than 1.65 keV cm⁻¹), muons from pion decay will be significantly accelerated before they decay, leading to an equal flux of ν_e and ν_{μ} , and a harder neutrino spectrum [367]. In choked jets associated with supernovae and double neutron star mergers, where the neutrino production region is embedded in the progenitor star and merger ejecta respectively, the flavor ratios can further be affected by matter effects [368–372]. Measuring the flavor composition of high-energy neutrinos is challenging, since signals from ν_e and ν_{τ} are easy to confuse; see, however, Ref. [96]. Nevertheless, IceCube has performed several measurements of the flavor composition [19, 53, 81, 373]; most recently, they have singled out the first ν_{τ} candidates [81]. We can use these measurements, combined with information on the values of the neutrino mixing parameters [82], to infer the flavor composition at the sources and, in turn, constrain their identity [57, 59, 83, 374, 375].

Multi-messenger connection and improved source modeling: Neutrinos are produced by hadronuclear and/or photomeson production processes, in which the neutrino–gamma-ray connection is naturally expected. In particular, the *Fermi* gamma-ray data give us stringent constraints on the candidate sources of IceCube neutrinos. For a simple power-law (SPL) spectrum, the spectral index of the neutrino sources is constrained to be $\gamma_{\text{SPL}} < 2.1-2.2$ [38]. On the other hand, recent shower [118], HESE [70], and track data [119] suggest that the best-fit spectral indices are all softer, and the fact that the all-sky neutrino flux at 10 TeV is larger hints at the existence of hidden neutrino sources that are opaque to GeV–TeV gamma rays [376–378].

Because the energy generation rate densities of UHECRs, TeV–PeV neutrinos, and sub-TeV gamma rays are comparable [40], it is natural that, at least in some cases, they are produced in the same network of processes in the same sources. This can be done by cosmic-ray reservoirs, where cosmic rays confined in magnetized environments produce neutrinos and gamma rays while escaping cosmic rays contribute to the observed UHECR flux [97, 138]. The candidate source classes include galaxy clusters/groups [138, 379– 381] and starburst galaxies [382–386]. In $p\gamma$ scenarios, gamma rays are more likely to be obscured, but the connection between UHECRs and sub-PeV neutrinos is still possible in some source models [387].

Cosmogenic neutrinos: While predictions and modeling of the flux of cosmogenic neutrinos [147–149] have been steadily improving in recent years due to more information from UHECR experiments [44, 45, 388], the uncertainty is still about an order of magnitude. A measurement of the cosmogenic neutrino spectrum will provide valuable inputs to constrain UHECR source properties [389–397].

UHE neutrinos from point sources: Predictions of the flux of cosmogenic neutrinos are rather uncertain due to not only the composition but also the cosmic-ray maximum energy and redshift evolution of the sources. The first detected UHE astrophysical neutrinos may predominantly come from powerful point sources. First, there are source models that predict diffuse neutrino fluxes overwhelming the cosmogenic neutrino flux especially if the UHECR composition is dominated by intermediate or heavy nuclei. Second, the UHECR acceleration requires sufficiently powerful sources that may be rare in the Universe. This would make the source identification easier especially at extremely high energies, where atmospheric backgrounds are negligible [99, 398]. Candidate UHECR accelerators that can also be powerful UHE neutrino emitters include GRBs and hypernovae [399–406], magnetar/pulsar-driven supernovae [136, 407–409], TDEs [410–412], and blazars [137, 140, 413–415].

2.2.1 Diffuse flux of TeV–100 PeV astrophysical neutrinos

There are two ways to disentangle the diffuse flux of astrophysical neutrinos from the atmospheric neutrino flux. The first is based on our knowledge of their distinct energy spectrum. The second is based on the expectation that down-going atmospheric neutrinos can be tagged if accompanied by muons produced in the same air shower [416–418]. Combining both approaches has led to the discovery of an astrophysical flux at the TeV-to-PeV energies with a significance well above 5σ . However, above roughly 10 PeV, the picture remains unclear and likely requires next-generation detectors, possibly in combination [419], for a high-significance measurement of the cosmic neutrino flux. Further exploring this energy regime, as well as refining our current knowledge of the diffuse astrophysical flux will help shed light on source production



Figure 12. Overview of astrophysical diffuse flux measurements to date. The green point and contours shows the most recent best-fit and uncertainties using a sample of predominantly up-going tracks in IceCube with 9.5 years of data taking [119]. It is overlaid with IceCube results using high-energy starting tracks and cascades (red) [70], predominantly contained cascades (blue) [118], and a sample of predominantly starting tracks (orange) [19]. A recent ANTARES result (gray) using both cascades and tracks with 9 years of data is included for comparison [420].

mechanisms. It also opens an avenue to probe fundamental neutrino physics and BSM physics at an energy scale that would be unreachable otherwise.

The global picture of our current knowledge of the diffuse astrophysical neutrino flux is shown in Fig. 12 assuming a single-power-law spectrum. It summarizes spectral constraints derived from the analysis of various IceCube event samples, as well as ANTARES data [420]. Specifically, the IceCube data sets are a sample of high-energy neutrinos which includes both tracks and cascades with interaction vertices within the instrumented volume [70], a sample of up-going tracks (mostly muon neutrinos) [119], a sample of cascade-like events (mostly electron and tau neutrinos) [118], and a sample of tracks that start within the instrumented volume [19]. An apparent slight tension between the different measurements could be due to differences in flavor composition, energy range, the accounting of atmospheric backgrounds and the spectral model used. The combination of individual samples in a multi-sample analysis [421], improved calibration and simulations will lead to improvements of the accuracy of the measurement and a reduction of systematic uncertainties.

The measurement of the flavor composition of astrophysical neutrinos provides important clues about their origin, production mechanisms and physical properties of their sources. The canonical production of neutrinos in astrophysical sources with flavor ratios $f_{e,S} : f_{\mu,S} : f_{\tau,S} = 1 : 2 : 0$ from the decay of charged pions, can be altered by energy losses of pions and muons in the vicinity of the sources, e.g., due to the presence of strong magnetic fields [71, 73–75, 89, 340, 341]. Several measurements of the flavor composition have been performed with IceCube [19, 53, 81, 373]. Figure 13 summarizes the constraints derived. An important milestone was the recent identification of two ν_{τ} candidate events [81]. Current constraints are compatible with several astrophysical production scenarios [57–59, 83, 374, 375, 422–425] and standard neutrino oscillations on their propagation to Earth [59, 83], while high-energy neutrino production from the beta-decay of neutrons is strongly disfavored [59, 83].

The Glashow resonance is an enhancement of the s-channel neutrino charged-lepton cross section over the DIS cross section. Due to the preponderance of electrons in matter, it peaks for electron-antineutrinos at 6.3 PeV in the electron rest frame. Sixty years after its initial proposal [159], IceCube detected an astrophysical neutrino interacting at the resonance energy for the first time [21]. As the resonance occurs on Earth only for electron-antineutrinos, it opens an additional identification channel of both the neutrino flavor



Figure 13. Flavor constraints on the cosmic neutrino flux from various analyses of IceCube data. The constraints from an IceCube analysis identifying first tau neutrino candidates [81] is shown as black contours. Constraints from earlier measurements, a fit encompassing several IceCube data sets [53] and an analysis of the inelasticity distribution of IceCube high-energy events [19] are shown as shaded regions. They are compared to different scenarios of neutrino production in astrophysical sources and the full range of possible flavor compositions assuming Standard Model flavor mixing (gray dotted region). Figure reproduced from Ref. [81].

and charge. As a result, sources of high-energy astrophysical neutrinos can be expected to produce both neutrinos and antineutrinos. Even with only one event detected thus far, the diffuse flux is now expected to extend above the resonance energy, and constraints on cutoff models can be placed. With additional statistics in the future, we should expect more precise constraints on all fronts including the flux, flavor composition, and production mechanisms; see, e.g., Ref. [78, 79, 426].

2.2.2 Diffuse flux of ultra-high energy neutrinos: cosmogenic and from sources

UHECRs with energies reaching ~ 10^{11} GeV are expected to produce UHE neutrinos of energies well above the PeV scale. UHECR nucleons colliding with the cosmic microwave background (CMB) and extragalactic background light (EBL) produce secondary pions via the photomeson production in extragalactic space. The subsequent decays of pions generated by these interactions, also known as the Greisen-Zatsepin-Kuzmin (GZK) process [47, 48], generates the a flux of UHE neutrinos [26]. The GZK process is effective when the energies of UHECR nucleons are higher than ~ 5×10^{10} GeV, because of the threshold effect on the photopion production process. As approximately ~ 5 % of the parent nucleon energy goes into secondary neutrinos, these cosmogenic neutrinos see their main flux around and below ~ 10^9 GeV [51, 148, 389, 391, 427]. In the case of UHECR nuclei with energies below the threshold for photomeson production, beta-decays of unstable cosmic-ray nuclei produced during the photodisintegration chain can also produce cosmogenic neutrinos [428, 429]. In the absence of a significant proton component above ~ 50 EeV, this is the dominant mechanism for cosmogenic neutrino production.

Because the CMB field evolves with redshift following the history of Big Bang cosmology, the neutrino production yield highly depends on the cosmic time at which UHECR nucleons collide with CMB photons. Consequently, the cosmogenic neutrino intensity is a good probe for understanding UHECR source evolution with redshift, i.e., the cosmological evolution of UHECR sources [430]. The present upper limits on the cosmogenic neutrino fluxes obtained by IceCube [144], the Pierre Auger Observatory [431], and ANITA [146] have placed constraints on the source evolution parameter space, putting highly evolved astronomical objects such as FSRQs and GRBs into question as UHECR origins [316].

UHECR sources themselves may yield UHE neutrinos via photonuclear or photohadronic processes, as many of the sources are expected to sit in neighborhood of interstellar gas or photon radiation field [136– 141]. The fact that the UHECR luminosity density is comparable to that of sub-PeV neutrinos measured by IceCube and that of sub-TeV gamma rays measured by *Fermi* may also indicate that the three messenger particles share the same origin [40]. This astroparticle "Grand-Unification" theory has been extensively discussed in the literature. UHECR sources are indeed expected to produce neutrinos and cascaded gamma rays in this scenario, with energies even reaching to 100 PeV and beyond [38, 97, 138, 141]. The generic requirements for being both UHECR accelerators and UHE neutrinos emitters in the unification framework have been extensively discussed for the photohadronic scenario [387].

2.2.3 Neutrino sources

The TeV–PeV astrophysical neutrino flux that IceCube has measured is approximately isotropic, and therefore is likely produced mainly by extragalactic sources [70]. The sources are still almost entirely unresolved and pose a compelling mystery. Two categories of analyses are used to search for the origins of the astrophysical neutrino signal: neutrino-only analyses and multi-messenger analyses. The neutrino-only analyses search for clustering of neutrinos in direction and/or time. They are generally model-independent, but a large trials factor is incurred by searching many directions (and times, for time-dependent studies), which degrades the sensitivity compared to correlation searches with modest numbers of multi-messenger source candidates. Multi-messenger analyses search for correlation between neutrinos and known sources of other messengers, or at least with the directions of those messengers, including electromagnetic radiation, gravitational waves, and cosmic rays. These analyses sometimes search for temporal as well as directional correlation.

Neutrino-only analyses (clustering searches) have established strong upper limits on the flux from pointlike sources across the entire (4π) sky [432]. Dividing the total neutrino flux by these upper limits provides a lower limit on the number of sources (regardless of their distance): there must be at least $O(10^3)$ sources. Although the sources are very likely extragalactic, their distance distribution, and even their distance scale, is unknown.

There is evidence that at least one gamma-ray blazar, TXS 0506+056, contributes to the neutrino signal (and is therefore also a cosmic-ray source) [22, 433]. However, the neutrino sky is remarkably different from the gamma-ray sky. In particular, while gamma rays are an essential component of the evidence that TXS 0506+056 is a neutrino source, the relationship between gamma rays and neutrinos from this and other sources is not simple: the brightest gamma-ray sources are not the brightest neutrino sources [7, 97, 434]. Nevertheless, the neutrino sources could all be GeV and/or TeV gamma-ray sources, but without simple correlation between the neutrino and gamma-ray fluxes: a subset of the ~ 5×10^3 known \gtrsim GeV gamma-ray sources could be sufficient to explain the entire neutrino signal. The lack of simple correlation between neutrino and gamma-ray fluxes how complementary the two messengers are. The lack of correlation could be explained by a combination of (1) the gamma-ray fluxes include leptonic as well as hadronic contributions, (2) the neutrino sources are more distant than the gamma-ray sources, and/or (3) the gamma-ray signals from neutrino sources are attenuated within or near the sources (though in this case gamma rays may be reprocessed to lower energies [376]).

In addition to cross-correlation with gamma-ray sources [435–437], analyses have searched for neutrino emission from known sources across the electromagnetic spectrum, including X-ray, UV/optical/IR, and radio sources. These include blazars as well as the cores of active galactic nuclei (AGN), ultra luminous infrared galaxies (ULIRGs), and radio galaxies [438]. Time-domain (variable or transient) searches have also been performed for neutrino emission from known sources including blazars, gamma-ray bursts (GRBs), fast radio bursts (FRBs), tidal disruption events (TDEs), supernovae, and novae [23, 439–443]. In addition to the electromagnetic spectrum, analyses have searched for coincidence between neutrinos and gravitational waves or cosmic rays [444, 445].

In addition to the predominant extragalactic signal, there could be a significant Galactic neutrino flux [446, 447]. Models predict both a diffuse Galactic signal (from cosmic-ray interactions producing pions) and emission by discrete sources [448, 449]. Discrete source candidates include the highest-energy gamma-ray sources identified by IACTs as well as wide-field instruments including HAWC and LHAASO, including those that have been detected above 100 TeV [450, 451]. As in the case of extragalactic sources, detecting neutrino emission from Galactic sources would identify them as cosmic-ray sources. Detecting diffuse Galactic emission would also provide essential insight into cosmic-ray propagation and interaction. Current upper limits on the diffuse Galactic flux are comparable to model predictions based on the related gamma-ray signal, which is well measured but at lower energies [54, 452, 453].

Although limits on the spatial clustering of the neutrino signal indicate that there must be at least $O(10^3)$ sources, the actual number may be close to this or arbitrarily larger. A more sophisticated analysis than the simple comparison of the total flux to the point-source limits accounts for the source-to-source variation not only in flux but also in redshift and luminosity. Observational constraints can be set in a 2D phase space (source luminosity, source number density) and compared to known source classes characterized with other messengers. As has been established for other messengers and wavebands, there are likely multiple diverse source classes, and they likely vary substantially in contribution to the total flux, such that one or two source classes dominate the total astrophysical neutrino flux.

2.2.4 Multi-messenger modeling of point sources

The coincident observations of a high-energy neutrino event, IceCube-170922A, with X-rays and γ -rays from a blazar TXS 0506+056 [22, 433] demonstrated the power of the multi-messenger approach. Nevertheless, there are still open questions in this case as we still lack a concordance picture of the multi-messenger data. Neutrino production must be accompanied by the production of gamma rays, which, after cascading to lower energies due to interactions on the CMB and EBL, eventually appear in the X-ray and/or gammaray energy range. The resulting flux is predicted to be comparable to the neutrino flux. For the 2017 multi-messenger flare from TXS 0506+056, the electromagnetic spectrum was observed all the way from radio, optical, and X-ray bands to the gamma-ray band, and it showed a valley in X-rays observed by Swift and NuSTAR [454] that is especially constraining. The cascade constraints lead to upper limits on the neutrino flux which are in tension with the observation of IceCube-170922A [454–457]. The situation is more challenging for the 2014-2015 neutrino flare from TXS 0506+056 [458-461], for which various possibilities have been discussed to explain the missing electromagnetic energy [462, 463]. That these results are not sensitive to details of the modeling challenges the single-zone model approach for neutrino and gamma-ray emission. Other coincidences with flares from supermassive black holes have been reported [464-468]. In particular, coincidences with optically-detected tidal disruption events [23] with radio and X-ray counterparts are of special interest [469, 470]. The connection among different messengers remains clouded.

Multi-messenger analyses have begun to help our understanding of the astrophysical environments where cosmic rays are accelerated. The ten-year point-source searches with IceCube have indicated that NGC 1068 is the most significant steady source of neutrinos at a significance of ~ 3σ [432]. Intriguingly, NGC 1068 is known to be not only a type-II Seyfert galaxy but also a starburst galaxy. In the starburst model, NGC 1068 has been considered to be one of the most promising sources [97, 471] in the Northern sky (given that

M82 may not be ideal for confining cosmic rays with ~ 10 - 100 PeV). However, gamma-ray limits provided by MAGIC and HESS observations clearly show that it has to be a hidden neutrino source in the sense that sub-TeV gamma-rays are attenuated or cascaded down to lower energies [472]. High-energy neutrinos may be produced in the vicinity of the supermassive black hole, and particles may be accelerated in hot coronae around the accretion disk [354, 473] or by disk-driven winds [384, 474, 475]. In particular, given that NGC 1068 is among the brightest sources in intrinsic X-rays, the AGN corona model predicts that NGC 1068 is the most promising neutrino source in the Northern sky [354, 476]. The IceCube flux can be explained if a fraction of the accretion power goes to protons, and gamma rays are predicted to appear in the MeV range without violating the observational data by *Fermi*, MAGIC, and HESS.

2.2.5 Multi-messenger alerts

The conclusive identification of astrophysical neutrino sources and their electromagnetic counterparts will not only enable the study of hadronic processes in cosmic-ray accelerators but also determine the baselines over which cosmic neutrinos propagate, a critical ingredient in many of the already-mentioned studies of fundamental physics.

The hunt for neutrino sources is typically performed by searching for an excess of neutrinos in space and time (see Sec. 2.2.3) with respect to the dominant atmospheric background. However, at energies above \sim 100 TeV, the atmospheric background becomes subdominant with respect to the diffuse astrophysical neutrino flux discovered by IceCube [8], which allows to search for sources by correlating in space and time individual neutrino events with a high probability of being astrophysical in origin with other multi-messenger signals. This concept has been realized in current-generation neutrino telescopes such as IceCube and ANTARES, where the detection of single high-energy neutrinos is promptly communicated to the astronomical community so that targeted observations can be collected to identify an electromagnetic (EM) counterpart [477–479]. The blazar TXS 0506+056 was identified as a candidate neutrino using this technique as it was observed in a flaring state in gamma rays in coincidence with the detection of the high-energy neutrino IceCube-170922A [433]. The prompt alert streams maintained by current-generation neutrino observatories have mostly focused on the selection of muon-track events as their positional uncertainty regions (of order 1° radius or better) can be covered by pointed EM telescopes, although more recently particle shower events (or "cascades") have been added which have a higher astrophysical purity given the lower atmospheric backgrounds for this selection although at the cost of a poorer angular resolution (median error radius of $\sim 7^{\circ}$ for IceCube) [480]. These neutrino alerts are currently communicated via prompt Gamma-ray Coordination Network (GCN) notices, circulars and Astronomer's Telegrams and include the information required to facilitate their follow-up, such as sky position (with uncertainty), energy, and false alert rate for each event. The all-sky coverage of neutrino telescopes can also be used to search for correlated emission with multi-messenger alerts of interest, such as gravitational-wave events or the observation of a transient or variable EM sources that are potential neutrino emitters. Real-time programs such as AMON follow up on these external triggers and alert the community if a coincidence of interest is identified [481].

3 Goals for the next decade

3.1 In particle physics

The landscape of new-physics models: New physics may affect various features of high-energy cosmic neutrinos: their energy spectrum, arrival directions, flavor composition, and arrival times, as Fig. 2 illustrates for a representative part of the landscape of BSM models. Some of the outstanding questions to address today and in coming years are: How do neutrino cross sections behave at high energies? [17– 19, 103, 105–107, 116, 122, 161–181, 246, 247, 482, 483] How do flavors mix at high energies? [58, 59, 83, 85– 94, 366, 484–487] What are the fundamental symmetries of Nature? [62, 182–185, 187–192, 256, 488–495] Are neutrinos stable? [91, 268–270, 272, 273, 426, 496, 497] Is there evidence of dark matter in the flux of high-energy neutrinos? [108, 199, 200, 202, 204–208, 212–215, 447, 498–509] Are there hidden interactions with cosmic backgrounds? [63–65, 67–69, 91, 194, 195, 197, 198, 212, 217, 218, 286–289, 307, 510–519] What is the origin of the anomalous EeV events seen by ANITA? [311, 312, 321, 323, 325, 326, 328–333, 520–526] Neutrino telescopes may also probe new neutrino interactions and hypothesized particles, such as magnetic monopoles [527–531], by looking for exotic signatures [3].

Synergies with sub-TeV neutrino experiments: Upcoming sub-TeV experiments will reduce uncertainties that limit the sensitivity of fundamental-physics searches using high-energy cosmic neutrinos. First, the IceCube Upgrade [532], though geared towards low-energy neutrinos, will reduce systematic errors that affect the detection of high-energy neutrinos. Second, oscillation experiments—DUNE [533], JUNO [534], Hyper-Kamiokande [535], the IceCube Upgrade [532], KM3NeT/ORCA [125]—will reduce the uncertainties on lepton mixing parameters [536], permitting more precise tests of new physics via the flavor composition of high-energy cosmic neutrinos [58, 83]. Third, FASER ν will reduce systematic uncertainties in charm production at parton momentum fraction x close to 1 by measuring the high-energy neutrino flux from the LHC in the forward direction [6, 300, 303, 537]. This will improve predictions of the undiscovered background of prompt atmospheric neutrinos that muddle searches of new physics with high-energy cosmic neutrinos.

Given the unique potential and rich experimental outlook of high-energy cosmic neutrinos to extend our view of fundamental physics, this topic should feature prominently in the high-energy physics program, as it pushes the boundaries for the neutrino, cosmic, energy, theory, and instrumentation frontiers. Companion Snowmass Whitepapers study in detail dark matter physics [538–541], Beyond-the-Standard Model physics [542], the Forward Physics Facility [303], prompt neutrinos facilities [543], tau neutrinos [28], multimessenger facilities and capabilities [27, 29–31].

3.1.1 Cross sections

The last decade yielded the first measurements of the deep-inelastic neutrino nucleon interaction (DIS) cross section in the HE range. Future experiments, both at accelerators and with cosmic neutrinos, aim to significantly increase event rates and study neutrino interactions in a new energy region. In particular, both astrophysical and cosmogenic neutrinos provide neutrino beams at energies far beyond the reach of human-made accelerators, which allows us to explore particle physics in the phase space we could not access to otherwise.

Improvements in cross section measurements by neutrino telescopes at the HE scale will largely be due to increased statistics, as future optical experiments increase their sensitivities and exposures. This means that they will not expand their energy reach by a large amount. At the lower energy threshold, systematic uncertainties will dominate how well small absorption features can be resolved and at the higher energy the cross section measurement will be limited by the low cosmic neutrino flux.

To access the higher energies, UHE experiments plan to instrument much larger volumes. Indeed, detector volumes of 100 km³ of Antarctic ice are required to detect a large enough sample of UHE neutrino interactions to probe PDFs with Bjorken-x below 10^{-4} . Other experiments that rely on Earth-skimming tau neutrinos have larger effective volumes and are sensitive to even higher energy neutrinos, in part through neutrino flux attenuation in the Earth. Much progress has been made in recent years in the modeling of neutrino propagation in the Earth [234, 544–546]. These simulation codes will be useful tools to test the standard model and BSM neutrino cross sections inputs.

Beyond DIS cross sections, the first detection of W-boson production via the Glashow resonance [21] is promising. Improved experimental sensitivity in the PeV band with IceCube-Gen2 will further enhance the science gains in this regard. The $\bar{\nu}_e + e$ Glashow resonant production of real W-bosons mainly yields showers in the high-energy neutrino detectors, so Glashow resonant events could be detected via the shower spectrum. Ref. [247] shows that, on top of the shower events from neutrino deep-inelastic scattering with nuclei, W- boson production can be identified with 1σ and 3σ in the conservative and optimistic cases, respectively, with ten years of current IceCube data or with one year of IceCube-Gen2 data.

Through higher-order electroweak processes [234, 243, 244, 246, 247], ν_{μ} induced W-boson production can produce dimuon events [247, 547]. One muon comes from the leptonic vertex while the other from the W muonic decay. In Ref. [547], the authors proposed that dimuons can be detected using the small vertical spacing between the digital optical modules of high-energy neutrino telescopes like IceCube and IceCube-Gen2.

Another source of dimuon events are neutrino charged-current scattering with nuclei that produces a charm hadron. One muon comes from the charged-current scattering, and the second muon comes from the semi-leptonic charm hadron decay. This source of dimuons (DIS dimuons) typically has associated showers. With realistic shower cuts, the DIS dimuons produced in the detector can be almost completely removed, leaving the yield of dimuons from ν_{μ} -induced W-bosons produced in the detector largely background free [547]. IceCube-Gen2 could detect > 6 such W-boson dimuons in ten years [547], and other signatures can be used for detection ν induced W-boson production, including pure electromagnetic showers and showerless single tracks [247]. DIS dimuons event analyses have the potential to measure the strange-quark PDF. In the optimistic case, it is predicted that IceCube can detect $\simeq 130$ DIS dimuons and IceCube-Gen2 can detect $\simeq 620$ DIS dimuons, most from atmospheric neutrinos [547] (see also Ref. [548]).

UHE measurements of the cross section can constrain several proposed models of BSM physics involving the virtual exchange of the Kaluza-Klein graviton [549], microscopic black hole production [172], or neutrinoinduced sphaleron transition [179]. At these energies, again the neutrino-nucleon cross section can be measured by exploiting the attenuation of the flux as neutrinos cross the Earth, particularly because most events occur near the horizon. These experiments can constrain the cross section using the angular distribution of the events if the UHE flux is high, such that more than one event per year is detected [102, 103, 105–107]. Since the event rate of UHE neutrinos by large neutrino telescopes is essentially a product of neutrino flux and the neutrino-nucleon interaction cross section, the upper limit of UHE neutrino fluxes are turned into constraints on these BSM physics, assuming the cosmogenic neutrino flux from proton-dominated UHE-CRs [550]. Any new interactions enhancing the cross section by more than 100 are disfavored under this flux assumption [102].

3.1.2 Inelasticity

Future HE neutrino detectors, particularly optical Cherenkov detectors, are expected to enhance their flux sensitivity – and therefore statistics – by factors of two to four over the current measurements [19, 128]. This will improve precision substantially in that energy range.

At UHE energies, inelasticity measurements are yet to come, but may be possible with future radiodetection experiments [180]. Detectors must be able to detect and separate outgoing leptons and the hadronic cascade produced by neutrino interactions. At energies above about 10^{19-20} eV, the Landau-Pomeranchuk-Migdal (LPM) effect lengthens electromagnetic showers, giving them a longer distance scale than hadronic showers [551]. At still higher energies, the LPM effect will break electromagnetic showers into multiple subshowers, from a single charge-current induced electron. It may be possible to separate the hadronic and electromagnetic cascades on the basis of this length scale, which leads to different radio-emission spectra, allowing for inelasticity measurements in ν_e charged current interactions. Alternately, if multiple subshowers are detected as separate showers, this would also allow for inelasticity measurements. This measurement is most straightforward at extremely high energies (10^{20} eV), promising recent results suggest that hadronic and electromagnetic cascades may be separated on the basis of their different radio-emission spectra [552, 553].

3.1.3 Neutrino decay

Neutrino decay constraints in the future may evolve with measurements of the diffuse supernova neutrino background, cosmological measurements, and possibly with a measurement of the cosmic neutrino back-

ground. At shorter lifetimes, high-energy astrophysical neutrino constraints will continue to improve. These probes also have a benefit over the others in that it is easier to discover neutrino decay given astrophysical uncertainties due to the fact that all three flavors can be differentiated at high energies but not at low energies. The best improvement will come from IceCube augmented by KM3NeT and Baikal-GVD. Experiments sensitive to higher energy neutrinos are less sensitive to neutrino decay since the neutrinos are more boosted and experience less proper time. The increased statistics of events collected by IceCube-Gen2 will help, as well as the presumably increased ν_{τ} data set which is a key data set for neutrino decay.

3.1.4 Dark matter

IceCube and ANTARES have provided the principal constraints on DM annihilation to neutrinos for DM with a mass larger than a TeV. Future experiments will boost the sensitivity of the neutrino telescope in the search for very heavy dark matter. The sensitivity of the future experiments to probe very-heavy DM are shown in Fig. 7. For masses between 1–100 TeV, KM3NeT will provide the leading constraints on the annihilation cross section, closing in on the parameter space near the thermal relic abundance. Above 1 PeV, improved constraints will be provided by IceCube-Gen2, radio arrays like RNO-G and GRAND, and other planned experiments. Experiments operating in the mid-range between 1 PeV to 1 EeV, such as TAMBO, Trinity, or RET-N, can provide a crucial overlap between the PeV scale and the EeV scale experiments. Beyond the EeV scale, POEMMA's fluorescence observation mode could probe DM masses beyond 10¹¹ GeV [554].

Interactions of ultra-light dark matter with neutrinos can also impact the observed flavor ratios. The nearly even expectation for the flavor ratio at the Earth is very robust against varying the condition at the source and even invoking new physics [58]. In fact, Lorentz symmetry implies that neutrinos arrive at the Earth as incoherent combinations of different neutrino mass eigenstates. However, as shown in [5, 90, 514], in the presence of Lorentz violation, the standard expectation of even flavor ratios can be relaxed because the flavor state arriving at the Earth can remain as a coherent combination of the mass eigenstates. In the background of ultralight dark matter, ϕ , neutrinos can obtain a Lorentz violating effective mass of form $(m_{eff})_{\alpha\beta}\nu^{\dagger}_{\alpha}\nu_{\beta}$ originating from vector vector interaction of the following form with ultra light complex dark matter denoted by ϕ [516, 555, 556]

$$\frac{g_{\alpha\beta}}{\Lambda^2}(\phi^*\partial_\mu\phi - \phi\partial_\mu\phi^*)(\bar\nu_\alpha\gamma^\mu\nu_\beta).$$
(3.1)

Then, $(m_{eff})_{\alpha\beta} = (\rho_{DM}/m_{DM})(g_{\alpha\beta}/\Lambda^2)$. In the halo of dark matter where the dark matter density, ρ_{DM} , is relatively large, $(m_{eff})_{\alpha\beta}$ may dominate over $(\Delta m^2/E_{\nu})$. As a result, the energy eigenstates will correspond to the eigenvectors of $(m_{eff})_{\alpha\beta}$ rather than the mass eigenstates in vacuum. Since the DM density variation across the route of neutrinos is smooth, the transition will be adiabatic.

As shown in [516], interaction of form (3.1) can be obtained in a UV complete model based on gauging the $L_{\mu} - L_{\tau}$ symmetry. In this case, $g_{\alpha\beta}$ will be diagonal with a remarkable result that the original flavor ratio at the source located in a dark matter halo will be maintained up to the Earth. That is, if at the source $F_{\nu_{\tau}}^{0} \ll F_{\nu_{\mu}}^{0}, F_{\nu_{e}}^{0}$, the ν_{τ} flux at the Earth will remain negligible. The observation of two ν_{τ} events by IceCube [81] already puts an upper bound on g_{α}/Λ^{2} . However, as discussed in [555], it is still possible that higher energy neutrinos ($E_{\nu} > 10$ PeV) originating from sources located inside DM halos maintain their original flavor ratio. This implies that while detectors designed to detect air showers from Earth-skimming ν_{τ} may report null results, the radio telescopes designed to detect the Askaryan radiation from all neutrino flavors may report a significant flux [555]. Cosmogenic neutrinos, having originated typically outside DM halos will still have the canonical democratic flavor ratios [555].

3.1.5 Secret neutrino interactions

Neutrinos may interact with the cosmic neutrino background through non-standard neutrino interactions [64, 65, 195, 557]. Upcoming experiments will allow for precision measurements of the energy spectrum of highenergy astrophysical neutrinos. The most salient feature imprinted by secret interactions on the high-energy neutrino spectrum are the dips around the resonance energy of the neutrino-neutrino cross section. Thus, previous searches and forecasts have centered predominantly on looking for evidence of spectral dips.

There are two main limitations to these searches; future detectors may try to improve on them. First, in optical Cherenkov neutrino telescopes, the energy E of a neutrino-initiated event is typically reconstructed to within, roughly, 10% in $\log_{10}(E/\text{GeV})$. The limitation stems from uncertainties in the medium properties, in the position of the interaction vertex, and other detector characteristics. The detector energy resolution sets the size of the smallest dip feature that we can expect to detect: if the energy width of the dip is smaller than the energy resolution, the dip will not be detected. Second, when looking for the presence of the dips in the energy spectrum—or other unusual features—searches use samples of contained events, where the neutrino interaction occurs inside the detector volume. In contained events, most or all of the neutrino energy can be inferred from the event energy, and so the shape of the energy spectrum can be inferred with relatively little bias. Unfortunately, contained events are rare; after ten years, IceCube has observed only approximately one hundred of them. Future optical detectors, with better knowledge of the properties of the detector medium, and of larger size, should overcome both of the above limitations.

Further, the couplings to the new mediator need not be flavor-universal: different neutrino flavors experience secret interactions with different strengths. This expands the available parameter space where effects may be present and require searches. Specifically, couplings to ν_{τ} are the least constrained. By including the capability to identify neutrino flavor, future searches could start probing flavor-dependent couplings. However, when doing so, the significant uncertainties in the flavor composition with which neutrinos are produced need to be factored in, and will weaken the probing power. Refs. [69, 91] explored future prospects in detail, highlighting the capability of IceCube-Gen2 to improve constraints on ν_{τ} -coupled mediators by over two orders of magnitude with ten years of data collection, for mediator masses up to 100 MeV. Ultra-high-energy neutrino telescopes could extend the range to tens of GeV [558].

Additionally, measuring neutrinos from identified sources with multi-messenger observations will allow for further constraints on secret neutrino interactions by constraining their mean free path and any possible neutrino echoes or halos [198, 289].

3.1.6 Prompt neutrinos

The extension of IceCube, the proposed IceCube-Gen2, will be able to enhance the detectability of prompt atmospheric neutrinos with higher statistics [128]. It will additionally provide a unique laboratory to study PeV prompt muons as produced in air showers, which are strongly linked to the production of prompt neutrinos using an approach that is complementary to accelerator studies of prompt neutrinos [559]. In the meantime, upcoming forward experiments at the LHC, FASER ν [300] and SND@LHC [301], will measure the prompt neutrinos and reduce the uncertainties in heavy flavor production. In addition, further developing experiments with upgrades of FASER ν and SND@LHC, and additional experiments for the next stage, the so-called Forward Physics Facility (FPF) are under study [6, 303]. The FPF will be designed to cover rapidity ranges beyond those probed by LHCb and by forward experiments deployed during Run 3 of the LHC. The FPF will play a crucial role through measurements of high energy neutrinos from the decays of the charm mesons produced in pp collisions at the LHC at $\sqrt{s} = 14$ TeV. These FPF measurements will then be related to improved theoretical predictions of prompt atmospheric neutrino fluxes [6]. Likewise, measurements of the prompt atmospheric neutrino flux may lead to a better understanding of structure of the proton through the parton distribution functions and, for example, the role of intrinsic charm [560–564].



Figure 14. Neutrino flavor ratio at Earth as a function of long-range potential $V_{\alpha\beta}$ associated with $L_e - L_{\mu}$ (*left*), $L_e - L_{\tau}$ (*middle*), and $L_{\mu} - L_{\tau}$ (*right*) for the three choices of the flavor ratio at the sources $(f_{e,S}:f_{\mu,S}:f_{\tau,S}) = (\frac{1}{3}:\frac{2}{3}:0)$, (0:1:0), and (1:0:0). Equal contribution from neutrino and antineutrino is considered. All the oscillation parameters are varied in 1σ allowed range under the normal mass ordering (NMO) scenario as given in Ref. [565]. Neutrino energy is fixed at $E_{\nu} = 100$ TeV. Dashed blue and dot-dashed red contours respectively show projected IceCube flavor ratios for 8 years runtime and 35 years runtime (15 years IceCube and 10 years IceCube +Gen2). Left and middle ternary diagrams are adapted from Ref. [307].

3.1.7 Fundamental symmetries

 $\mathbf{U}(\mathbf{1})'$ extensions to the Standard Model By measuring the flavor composition of astrophysical neutrinos, next generation flavor-sensitive experiments will significantly improve our sensitivity to possible U(1)' extensions in the Standard Model. Fig. 14, shows the expected neutrino flavor composition at Earth in the SM and in the presence of $L_{\alpha} - L_{\beta}$ interactions [307]. Dashed-blue and dot-dashed-red contours, respectively, show projected IceCube-measured flavor ratios for an 8-year runtime and a 35-year runtime (15 years IceCube and 10 years IceCube +Gen2). Three benchmark scenarios for flux compositions at the source, namely, full pion decay (1:2:0), muon damping (0:1:0), and neutron decay (1:0:0) are considered. The respective flavor composition at Earth is shown by grey blobs in the standard mixing scenario and by yellow-orange strips in the presence of $L_e - L_{\beta}$ long-range force. For all the assumed flavor ratios at source, the flavor ratio at Earth deviates from SM predictions. For $(V_{\alpha\beta} >> (H_{vac})_{ee})$, the original flavor ratio is preserved and not modified during propagation. Therefore, using the IceCube flavor ratio data, it is possible to constrain the mass and the coupling of the new interactions associated with $L_{\alpha} - L_{\beta}$ symmetries. With upcoming experiments like IceCube-Gen2 and the IceCube Upgrade, the flavor composition will be measured with high precision, which will allow for tighter constraints on the coupling and mediator mass of these new interactions.

3.2 In astrophysics

3.2.1 Diffuse flux of TeV–100 PeV astrophysical neutrinos

IceCube has measured, for the first time, the spectrum and flavor composition of cosmic neutrinos over three orders of magnitude in energy (10 TeV - 10 PeV). The apparent isotropic distribution of the observed neutrinos on the sky points to an extragalactic origin. This is also corroborated by the first observational evidence for a neutrino source, the blazar TXS 0506+056.

Over the next decade, IceCube's measurement will be complemented by new neutrino observatories currently under construction in the Northern Hemisphere: the KM3NeT ([125], see also Sec. 4.2.2) neutrino telescope in the Mediterranean Sea, and the Gigaton Volume Detector (GVD) ([566], see also Sec. 4.2.3) in Lake Baikal. The sky coverage of these telescopes is complimentary to that of IceCube, due to their geographic



Figure 15. A comparison of the IceCube measurements of the diffuse astrophysical neutrino flux with projections for the future IceCube-Gen2 detector. The IceCube measurements shown are from high-energy cascades ([118], squares), tracks ([119], circles), a Glashow resonance event ([21], +), and a search for EHE events ([144], line with arrows). The predicted unfolded diffuse spectrum assuming 10 years of IceCube-Gen2 data is shown in blue with error bands representing 68% confidence intervals up to 10 PeV. The upper panel assumes a single power law model with a spectral index of -2.5. The lower panel assumes a two component spectrum, consisting of a power law with an exponential cutoff at approximately 100 TeV, and a component reflecting a model of the $p\gamma$ emission of BLLac which peaks at tens of PeV. Both models are based on fits to the cascade data [118]. With current data from IceCube, it is not statistically significant to reject either hypothesis. IceCube-Gen2 will provide key measurements to reveal spectral features in the astrophysical neutrino component. Above 10 PeV, the expected differential 90% C.L. sensitivities for the combined optical and radio instrumentation for IceCube-Gen2 are shown in solid blue line.

locations. Comparable in size and sensitivity to the latter, they will enhance the precision of the current measurements, and, more importantly, should be able to probe the existence of a Galactic component in the diffuse neutrino flux. The observation of such a diffuse Galactic component, originating from the interactions of TeV-PeV cosmic rays with interstellar gas, and the measurement of its spectrum would be a unique input for understanding the production and propagation of the highest-energy cosmic rays in the Milky Way (e.g., [567]). Joint spectral and flavor measurements combining complementary data from all operating neutrino telescopes will be feasible, and may even be necessary to solve the apparent slight tensions currently seen in IceCube data.

A major step forward in measurement precision for spectrum and flavor composition at TeV and PeV energies will arise from the proposed IceCube-Gen2 observatory ([128], see also section 4.2.5). Figure 15 shows a comparison of current IceCube measurements and projections for 10 years of IceCube-Gen2 data. The IceCube-Gen2 projections [128], with an updated astrophysical flux model and improved characterizations of the surface and radio array efficiencies, give an impression of the expected unprecedented precision and energy coverage of the measurement of the diffuse neutrino spectrum. Together with significantly stronger

constraints of the flavor composition of the cosmic neutrinos (cf. Fig. 14), these data can be used to study the properties of the source populations that dominate the cosmic neutrino flux. Spectral features allow for study of cosmic-ray interaction processes and maximum acceleration energies. Investigations of the resonant production of neutrinos at the Glashow resonance [159] can be used to study the $\nu/\bar{\nu}$ ratio of cosmic neutrinos, and derive constraints on the production environments and targets [79]. Differential measurements of the flavor composition in various energy ranges can probe magnetic field conditions in the neutrino production regions around the sources [128]. For flavor measurements, tau neutrino identification capabilities are of particular importance. See Ref. [28] for a detailed discussion of tau neutrino identification prospects with IceCube-Gen2, KM3NeT, and Baikal-GVD.

3.2.2 Diffuse flux of ultra-high-energy neutrinos: cosmogenic and from sources

The flux of cosmogenic neutrinos is expected to encode complementary and unique information about the nature and flux of UHECR accelerators, including their redshift evolution, their chemical composition, and maximum accelerating energy. The cosmogenic neutrino flux measurements can also be used for probing the Galactic-to-extragalatic transition [389]. Interestingly, there is a "sweet spot" [397] near 1 EeV where the flux of UHE neutrinos is predicted to depend mostly on two parameters: the redshift evolution of sources and the chemical composition of the cosmic rays, in particular the fraction of protons above ~ 40 EeV. For example, a source's redshift evolution is often parameterized as being distributed in z according to $(1+z)^m$; IceCube data already disfavors $m \geq 3.5$ [316] for purely power law sources, constraining otherwise promising sources such as radio-loud FR-II AGN. With regards to composition, there is a long standing discrepancy between two measurements of the UHECR composition, with the Telescope Array experiment favoring a "light" (proton-dominated) composition [568], while the Pierre Auger observatory favors a "heavy" (ironrich) composition [569]. Despite these different interpretations, the results are approximately compatible to each other within uncertainties [570, 571]. These two scenarios have markedly different signatures in the cosmogenic neutrino flux however, with a light proton-dominated UHECR flux giving rise to a much higher cosmogenic neutrino flux than a heavy iron-rich one. The cosmogenic neutrino intensity, therefore, is a completely orthogonal probe of the UHECR composition [390, 397] with different systematic uncertainties.

The IceCube, Auger, and ANITA experiments already constrain the cosmogenic neutrino parameter space, and a major goal for the next generation of observatories is to definitively detect this flux, or improve the constraints by at least two orders of magnitude, reaching a flux sensitivity near 10^{-10} GeV cm⁻² s⁻¹ sr⁻¹ at 1 EeV. Such sensitivity will probe even "pessimistic" cosmogenic models where the UHECR flux is relatively heavy [148, 572], as suggested by the Pierre Auger Observatory. A cosmic-ray composition of even 10-20% protons will lead to a few detected neutrinos per year for typical source evolution models [397]. If the flux is in fact even lighter, as suggested by the Telescope Array, experiments with this level of sensitivity could detect dozens or hundreds of events per year.

3.2.3 Neutrino point sources

IceCube has found the first compelling evidence for an extragalactic neutrino source, and several more hints that point to a rich extragalactic neutrino sky that may be composed of various source populations contributing to the cosmic neutrino flux (cf. Sec.s 2.2.3 and 2.2.4). Ultimately, complementary and more sensitive instruments are needed for revealing the "dominant" population for the observed diffuse neutrino flux (among various source populations) and a comprehensive study of the neutrino sky.

To date, IceCube has found no evidence for Galactic neutrino sources nor, therefore, Galactic cosmic-ray accelerators. Northern Hemisphere observatories (such as the under-construction Baikal-GVD and KM3NeT) will have a much better sensitivity of the inner galactic plane and the Galactic center region, where the non-thermal emission from the Milky Way is brightest and several candidate cosmic-ray accelerators are located, in particular some of the recently established Galactic Pevatrons (sources with observed gamma ray emission at O(100 TeV)) [573–576]. The detection of neutrinos is an unambiguous signature of the acceleration of



Figure 16. Left: Comparison of the effective local density and luminosity density of extragalactic neutrino source populations to the discovery potential of IceCube and IceCube-Gen2 (optical array only). We indicate several candidate populations by the required neutrino luminosity density to account for the full diffuse flux [97] observed by IceCube. The orange band indicates the luminosity / density range that is compatible with the total observed diffuse neutrino flux. The lower (upper) edge of the band assumes rapid (no) redshift evolution. The lines indicate the parameter space (above the lines) for which IceCube and IceCube-Gen2 are able to discover one or more sources of the population. Right: Same comparison for transient neutrino sources parametrized by their local rate density [40].

protons or nuclei. Therefore, the combined data of IceCube and the new generation of Northern Hemisphere detectors may finally resolve the puzzle of where the highest-energy cosmic rays in the Milky Way originate.

An order of magnitude improvement in detector volume, corresponding to a five-fold increase in sensitivity, will be critical for testing the dominant origin of the extragalactic diffuse TeV-PeV neutrinos and investigating the properties of their primary populations with reasonable statistics [97]. Next-generation detectors such as IceCube-Gen2 will be able to identify the brightest sources and transients of populations currently considered as potential cosmic-ray accelerators. This is the case for both cosmic-ray reservoirs that allow the production escape of UHECRs and cosmic-ray accelerators in which cosmic rays are confined and/or depleted [97] (see Fig. 16). It will also enable high-significance observations of the brightest sources and flares similar to the one observed for TXS+0506+056 [128], allowing a precise measurement of the spectral properties of such sources/transients.

The energy generation rate density of 10-100 TeV neutrinos, indicated by the IceCube shower data [118], is surprisingly large. The neutrino flux seems to violate the Waxman-Bahcall bound and the corresponding gamma-ray flux violates the IGRB in the transparent limit [376]. Revealing the sources in the 10 TeV range is important especially if the diffuse flux consists of multiple source populations. This will also enable investigation of particle acceleration and related plasma processes in dense environments that cannot be directly observed by electromagnetic observations, and successful detection will also be useful for probing new physics beyond the Standard Model. The candidate sources include the vicinity of supermassive black holes of AGN [353, 354, 577] and choked jets [359, 361, 362, 372]. For this purpose, neutrino detectors that are sensitive to 10 TeV energies or lower are important. KM3NeT and Baikal-GVD, which has good angular resolutions even for shower topographies, will also help to identify bright sources in the southern sky [476].

For the purpose of identifying UHECR sources, it is important to reveal the origin of neutrinos $\gtrsim 10 - 100$ PeV. Here, novel detector concepts that will be able detect neutrino-induced showers in Earth's natural ice shields or in the atmosphere with effective volumes of hundreds to thousands of km³ (see Sec. 4)


Figure 17. The sensitivity to short (1000 s) burst all-flavor neutrino plus antineutrino spectral fluence for some current and future detectors. The ANTARES, IceCube and Auger limits are 90% confidence level limits in a \pm 500 s window around the gravitational wave event from GW170817 [25]. The dashed blue histogram shows IceCube-Gen2's projected sensitivity for such an event [128], including IceCube-Gen2 radio [578] (shown for a range of declination with cyan). The blue shaded band comes from IceCube's all-sky point-source effective area values tabulated for 2012 with 86 strings [579, 580]. The ANITA-4 NH limit is shown with the solid purple curves, and projected all-flavor sensitivities for PUEO [581] from ν_{τ} -sourced and Askaryan signals are shown with purple and magenta dashed curves, for POEMMA (orange)[582] and for GRAND200K (red) [154] updated according to [583]. The Kimura et al. [584] extended emission short gamma ray burst fluence for on-axis viewing ($\theta = 0^{\circ}$)[584] from 50 Mpc is shown with the solid black curve.

for an overview) will play key roles. Having such large effective volumes, these detectors will be able to identify astrophysical sources and transients that are directly linked to UHECR production in a wide range of scenarios.

3.2.4 Multi-messenger astronomy

The combination of neutrinos with multi-wavelength observations across the electromagnetic spectrum is key for the identification of astrophysical neutrino sources, as well as for the modeling of the physical processes, conditions and environments that enable particle acceleration in such sources (cf. Sec. 2.2.4 and Ref. [31]).

One of the most important results from the multi-messenger observations is that energy generation densities (or luminosity densities) of high-energy neutrinos, sub-TeV gamma rays, and UHECRs are all comparable [38–40]. This may indicate that all three messengers are physically connected. The simplest scenario is that all three messengers come from a single source class. While such a grand-unification scenario is appealing [138], the multi-messenger analysis has indicated other populations, particularly below 100 TeV energies [376]. Even if the physical connection exists, it may be clouded by different source components and/or different emission regions. Indeed, luminosity densities of various sources including AGN and stellar deaths such as supernovae are not much different [40], and dissipation at multi-scales has been observed in these sources. Detailed information from electromagnetic observations from radio, optical, X-ray, and gamma-ray bands will be crucial, and next-generation gamma-ray detectors such as the Cherenkov Telescope Array will provide critical tests for the unification picture.

One of the most successful strategies of multi-messenger astronomy has been the real-time distribution of neutrino alerts (cf. Sec. 2.2.5). Over the coming decade, the quality and quantity of neutrino alerts will improve significantly thanks to the construction and operation of additional gigaton-scale neutrino telescopes such as KM3NeT, Baikal-GVD, and the recently-proposed P-ONE. Their distribution around the world enables a constant monitoring of the entire sky with good sensitivity in the TeV–PeV range. The distribution mechanisms for these alerts will also benefit from current efforts to upgrade the communication infrastructure between astrophysical facilities to prepare for the massive alert rate expected from Rubin observatory and other time-domain instruments.

After completion, the proposed IceCube-Gen2 detector will increase the rate of individual high-energy neutrinos by a factor of five that in combination with improved pointing, and more sensitive follow-up instruments will boost the number of observed correlations with candidate sources [128] by an order of magnitude and enhance their significance. For example, Fig. 17 gives an illustration of the sensitivities to short (1000 s) bursts of neutrinos currently achieved for the gravitational wave event GW170817 from IceCube, ANTARES and Auger [25], and short burst sensitivities of IceCube [579, 580] and ANITA-4 NH, as projected sensitivities from IceCube-Gen2 [128], including IceCube-Gen2 radio [578], PUEO [581], POEMMA [582], and GRAND200K [154].

The construction of neutrino observatories operating in the extremely-high-energy range (beyond PeV) will also provide a new source of neutrino alerts that is yet largely unexplored, but would be of interest given the very low backgrounds expected in this energy range.

4 Experimental landscape

4.1 Overview

In this section, we give a brief overview of the experimental landscape, neutrino detection strategies at the highest energies, and current and planned experiments.

4.1.1 The broad experimental landscape

Because the neutrino-nucleon cross section grows with energy, the Earth is opaque to high-energy neutrinos, reducing the angular coverage for experiments to well below 4π . Charged-current interactions (CC, $\nu_l + N \rightarrow l + X$, where $l = e, \mu, \tau$ and X are final-state hadrons) remove neutrinos from the flux. Neutral-current interactions (NC, $\nu_l + N \rightarrow \nu_l + X$) redistribute high-energy neutrinos to lower energies.

In the TeV–PeV range, the opacity is high [116, 165, 170, 173–176], but some neutrinos still traverse up to thousands of km inside the Earth before interacting [17, 18, 122]. HE neutrinos are detected by instrumenting a cubic-kilometer of water or ice with photon detectors, which capture the optical Cherenkov light from the particle shower following a neutrino interaction. The in-ice optical Cherenkov technique is mature and led to the discovery of HE astrophysical neutrinos. In the near term, we anticipate broader sky coverage as IceCube continues to monitor the Northern sky, while KM3NeT, Baikal-GVD, and P-ONE will open the Southern HE-neutrino sky.

For UHE neutrinos, the interaction length is so short [102, 234, 590, 591] that instruments must either be sensitive to neutrinos interacting in the atmosphere or to neutrinos interacting in rock, ice, or water outside of the detector volume [482, 582, 592]. A variety of suitable UHE-neutrino detection techniques have been developed and are currently being implemented. These techniques include detecting neutrino interactions in dense media (ice or the lunar regolith) and detecting Earth-skimming tau neutrinos from the air showers they generate; we describe them below. While some UHE experiments are sensitive to all neutrino flavors, other are sensitive predominantly or only to tau neutrinos.

The sensitivity of proposed experiments is shown in Fig. 18. In the TeV–PeV energy range, large optical Cherenkov detectors (IceCube [593, 594], KM3NeT [125], Baikal-GVD [127], P-ONE [126]) are expected to survey the full HE neutrino sky with improved statistics in the coming decade. In the 1–100 PeV energy range, IceCube and several proposed experiments (IceCube-Gen2 [128], RET-N [129], TAMBO [130], Trinity [595]) may observe neutrinos from astrophysical sources and study the spectral shape of the diffuse astrophysical neutrinos by extending measurements to higher energies or constraining a spectral cutoff. At even higher energies, we eventually expect a transition from astrophysical neutrinos to cosmogenic neutrinos



Figure 18. The expected differential 90% C.L. sensitivities for a variety of experiments to an all-flavor diffuse neutrino flux computed in decade-wide energy bins and assuming a ten-year integration (unless otherwise noted in the legend). The measurements and sensitivities are compared with a range of cosmogenic neutrino models [141, 397] and astrophysical neutrino models [138, 141, 411]. The blue bordered bands show the astrophysical neutrino flux measured by IceCube using tracks (ν_{μ} [119]) in hatch and using cascade-like events (ν_{e} and ν_{τ} [118]) in solid band. The solid lines show experimental upper limits at higher energies from the Pierre Auger Observatory [431], ARA [585], ARI-ANNA [586], ANITA I-IV [146], and IceCube [587]. The dashed lines show the sensitivities of a selection of proposed experiments currently in various design and prototyping stages (GRAND with 200,000 stations [154], BEACON with 1000 stations [157], TAMBO with 22,000 detectors [130], Trinity with 18 telescopes (updated from [588]), RET-N with 10 stations each with a 100 kW transmitter [129], POEMMA30 [558]) and under construction (RNO-G [155, 589], PUEO [581]). Experiments using the same detection technique are grouped into similar colors (orange, Earth-skimming radio (GRAND, BEACON); dark teal, particle showers (TAMBO); light green, Earth-skimming optical Cherenkov and fluorescence (Trinity, POEMMA30); pink, in-ice radio (ARIANNA, ARA, PUEO, RNO-G, RET-N (radar); blue, in-ice optical Cherenkov (IceCube)). Sensitivity to UHE neutrinos from IceCube-Gen2 (dashed purple) is computed using radio and optical for 10 years (see Fig. 15), and PUEO (dashed orange) uses both in-ice and Earth-skimming radio techniques (shown here for 3 flights, 100 days). Auger (teal) uses particle showers and fluorescence and its upgrade, AugerPrime, will employ radio. The grey downward-pointing arrow is a reminder that experimental sensitivities improve not only as exposure increases with time, but also as new experimental techniques and analysis methods are both demonstrated and scaled to larger detection volumes.

(or a combination of cosmogenic and astrophysical neutrinos). Several experiments are sensitive to these energies (IceCube [593, 594], the Pierre Auger Observatory [431], ARIANNA [596], ARA [585, 597, 598]) or are under construction (RNO-G [155, 589], PUEO [581]). The most constraining limits on UHE neutrinos come from IceCube (in-ice optical), Pierre Auger Observatory (particle showers and fluorescence) [431], and

ANITA experiments (in-ice radio [146]).

Several mid-scale UHE experiments currently operating (ARA [585, 597], ARIANNA [596], TAROGE-M [158]) or under construction (PUEO [581], RNO-G [155, 589], EUSO-SPB2 [599]) have the sensitivity to constrain the proton fraction in UHECR sources [397]. Still other experiments in the design or prototyping phase have promising projected sensitivities (BEACON [157], GRANDProto300 [600], RET [601], POEMMA [558]). The full array of experiments inform our understanding of fundamental physics and cosmic ray sources, but individual instruments also serve as prototypes for larger experiments targeting flux sensitivities consistent with the lowest expectations from UHE neutrino flux models (Trinity-18 [131], BEACON-1K [157], GRAND-200k [154], IceCube-Gen2 [128]) or broad-sky coverage and transient phenomena (POEMMA30 [558]) planned in the coming decades.

4.1.2 Detection strategies

Neutrino detection in ice and in water: Signatures of HE and UHE neutrinos can be detected via optical Cherenkov emission in ice and water. UHE neutrinos can also be detected using their radio signatures.

- Optical detection in ice and water: IceCube [593, 594] and, until very recently, ANTARES [602] search for the optical Cherenkov emission from neutrino-induced showers and tracks from all flavors of neutrinos. Optical Cherenkov experiments are sensitive to HE and UHE neutrinos that interact near or inside the detector volume [316]. Experiments of this type typically look for tracks of secondary leptons produced by charged-current neutrino interactions or cascades produced by neutral-current and charged-current neutrino interactions. Track events have excellent angular resolution due to the long lever arm left by a final-state muon track, while cascade events have superior energy resolution. Typically, astrophysical purity increases with energy as the backgrounds from atmospheric neutrinos and muons fall more steeply with energy than the astrophysical neutrino flux. At UHE energies, these detectors are also sensitive to neutrinos that cascaded from EeV ν_{τ} to PeV energies [603]. In the future, IceCube-Gen2 [128] will have a much improved HE sensitivity and a broader energy coverage. Broader HE sky coverage is achieved by expanding the detector volumes of optical arrays in the Northern Hemisphere, e.g., KM3NeT [125], Baikal-GVD [127], and P-ONE [126].
- Radio detection in ice: In dense media like ice, compact electromagnetic showers generated after UHE neutrino interactions emit coherent Askaryan radiation at the Cherenkov angle. Askaryan radiation—fast, coherent radio-frequency impulses—is due to the excess negative charge in the shower [604]. The long attenuation length at radio frequencies allows the signal to propagate over kilometer-long distances. In-ice radio experiments are sensitive to all three flavors [135, 605], and may have the power to discriminate flavors based on different event topologies [605, 606], such as the stretching of electromagnetic showers due to the Landau-Pomeranchuk-Migdal (LPM) effect [551]. In ARA [598, 607] and ARIANNA [596, 608], radio antennas are buried in the Antarctic ice. RNO-G, based on a similar concept, is a radio detector under construction in Greenland [155]. The experience gained in these experiments will directly feed into the design of the expansive, sparse radio array of IceCube-Gen2 [128]. Radar signals reflecting off in-ice showers is also being explored as a detection method [129, 609, 610].
- Radio detection from the upper atmosphere or in space: ANITA is a balloon-borne radio detector that searches for in-ice UHE neutrino interactions via the Askaryan radio emission that refracts out of the ice [611]. The high altitude provides ANITA with a large effective area at the highest energies [146]. PUEO [156, 581] is an upgrade to ANITA currently under construction. NuMoon searches for radio emission from neutrino interactions in the lunar regolith [612].

Neutrino detection via acoustic signals generated by UHE neutrino interactions in water or solids [613, 614] have also been pursued (see e.g., [615, 616] and references therein).

Air-shower detection techniques of UHE ν_{τ} : The lifetime of the tau and ν_{τ} regeneration [603, 617– 620] allow for unique detection techniques. The CC interaction of a ν_{τ} produces a tau lepton that, at UHE energies, typically decays over a distance of tens to hundreds of kilometers. If the geometry of the experimental setting is right, the neutrino interacts inside the Earth and the tau emerges and decays in the atmosphere. The decay initiates an extensive air shower, which can be detected with particle, air-shower imaging, or radio detectors [621]. But even in cases where a ν_{τ} interacts deep underground and the tau decays before reaching the surface, a new ν_{τ} is produced in the decay which can again interact in the Earth and generate a tau emerging from the surface. This " ν_{τ} regeneration" increases the chances of Earth-skimming ν_{τ} reaching the detector. Although with a lower probability, Earth-emerging muons from tau neutrinos and muon neutrinos (and electron antineutrinos in the PeV energy range due to Glashow Resonance) can undergo catastrophic energy losses in air, initiating deeply penetrating cascades that can increase the sensitivity to neutrinos with energies below 10 PeV for optical experiments [622]. For ν_e , the final-state electrons showers immediately after the CC interaction, i.e., before reaching the surface.) Earth-skimming neutrino fluxes have the added advantage of being unaffected by atmospheric neutrinos and only mildly affected by atmospheric muons [605].

- Air-shower particle detection: The Pierre Auger Observatory (Auger) is a long-running large-scale array of surface water tanks that detect the Cherenkov light from air-shower particles passing through them. Auger is designed to detect UHECRs, but it has been used to search for horizontal showers initiated by UHE neutrinos in the atmosphere [431]. Planned experiments using particle detectors combined with radio detectors include the upgrade to Auger, AugerPrime [152], and GCOS [153]. The Telescope Array (TA) [623] and HAWC [450] experiments have used a similar detection strategy, but have a more limited sensitivity. TAMBO is a planned array of water tanks to be located on one side of an Andean canyon, designed to detect the showers initiated by UHE taus emerging from the opposite side [130].
- Air-shower radio detection: Radio-detection of Earth-skimming tau neutrinos is a promising method due to the long attenuation lengths of radio waves in air. As with in-ice radio detection, sparse arrays can be used to instrument large areas. Radio emission is generated in air showers initiated by tau decays, via the geomagnetic effect, due to charge separation in the magnetic field of Earth as the air showers progress through the atmosphere. Moreover, the narrow Cherenkov cone and fast radio imaging enables sub-degree angular resolution. BEACON [157], in its prototype phase, TAROGE [624], and TAROGE-M [158] are compact antenna arrays in elevated locations that aim to detect UHE ν_{τ} emerging upwards via the radio emission of the air showers that they trigger. ANITA [156] and PUEO [156, 581] are also sensitive to upgoing ν_{τ} , from a higher elevation. GRAND [154] is a planned experiment that will cover large areas with a sparse antenna array to detect the radio emission from air showers triggered by UHE ν_{τ} , cosmic rays, and gamma rays.
- Air-shower imaging: Several air-shower imaging instruments, although optimized for cosmic-ray and gamma-ray detection, have demonstrated that the imaging of air showers via the Cherenkov and fluorescence light radiated by shower particles is a viable detection method of UHE ν_{τ} [133, 625– 627, 627, 628]. Air-shower imaging allows the reconstruction of the air-shower arrival direction with arcminute resolution and the shower energy within a few tens of percent of uncertainty. These excellent reconstruction characteristics are why the very-high-energy gamma-ray and UHECR communities have been using air-shower imaging for quite some time [629]. Two planned instruments optimized for the detection of UHE neutrinos from the ground are Trinity [630] and Ashra NTA [132]. POEMMA is designed to detect the Cherenkov light of UHE ν_{τ} -initiated showers from a satellite. A unique feature of POEMMA is its ability to rapidly reposition to target transient multi-messenger events [558, 582]. EUSO-SPB2 is a telescope mounted on a super-pressure balloon which will fly at high altitudes and serve as a pathfinder for POEMMA [134].



Figure 19. Left: The IceCube detector, showing the three subdetectors described in 4.2.1. The IceCube array is sensitive to interactions above a few hundred GeV. DeepCore lowers the detection threshold to around 5 GeV. The IceTop surface array allows for vetoing of down-going events and for study of atmospheric air showers. Figure reproduced from Ref. [587]. *Right:* The effective area for various charged-current muon neutrino data sets used by IceCube [642], including the "ELowEn" data set for sub-GeV transients [643], the "GRECO Online" data set for sub-TeV transients [642], and the gamma-ray follow-up data set (GFU) used for real-time follow-up analyses [477]. Figure reproduced from Ref. [642].

UHE neutrino detection in lunar regolith: The Moon, with its radio-transparent lunar regolith, provides a 19 million km² target for observing UHE neutrinos with Earth-based radio telescopes [631]. The radio emission generated by showers following a neutrino interaction in the rim of the Moon, are searched for by forming beams on various regions on the moon [632]. The technical requirements on radio telescopes are access to full time-domain voltages of the receiver and multiple beams on- and off-Moon, making not all telescopes suitable for these studies. Past observations have been performed most notably by the Parkes [633] and Westerbork telescopes [634] and the EVLA [635], with past and on-going work being performed with LOFAR [636, 637]. The next step in sensitivity will be obtainable with the Square-Kilometre Array (SKA) [632, 638–640]. While have being able to observe huge target volumes, the technique suffers from an intrinsically high energy threshold beyond 10^{20} eV [641], thus targeting more exotic models beyond the Standard Model. It is complimentary to the energy range of dedicated neutrino experiments and uses existing radio telescope infrastructure without much additional effort.

4.2 High-energy range

4.2.1 IceCube

The IceCube Neutrino Observatory [594], shown in Fig. 19, is the first gigaton neutrino detector. Situated at the geographic South Pole, it consists of the IceTop surface array used for cosmic-ray physics and the IceCube detector, an in-ice Cherenkov detector deployed between 1450 and 2450 meters below the surface of the Antarctic glacier. The IceCube detector contains 5160 Digital Optical Modules (DOMs) arranged in 86 strings. Each DOM contains a single 10-inch-diameter PMT, LEDs used for calibration measurements, and associated electronics housed in a pressure sphere. Seventy-eight strings are arranged in a regular hexagonal grid with strings 125 m apart and DOMs separated by 17 m along each string. Using this configuration, IceCube has measured neutrino fluxes from 100 GeV to several PeV, although its sensitivity to high-energy neutrinos extends beyond 1 EeV [21, 144]. The remaining eight strings are deployed as an infill array at the center of the detector, exploiting the clear ice below 1750 m to lower the IceCube threshold to 10 GeV and to study neutrino oscillations and sub-TeV dark matter. Using specialized studies of detector-wide noise rates, IceCube is also capable of studying nearby MeV scale core-collapse supernova neutrinos.



Figure 20. IceCube expected 90% sensitivity for an E^{-2} (*left*) and E^{-3} (*right*) spectrum [644]. Sensitivities are shown for 10 years of IceCube tracks [432] (grey), IceCube cascades (10 years [644], red; 7 years [480], blue), and for 11 years of ANTARES data [653] (teal). The IceCube track sensitivity dominates the Northern sky while cascade events allow IceCube to remain competitive in the Southern sky for soft spectra. Figures reproduced from Ref. [644]

Able to reconstruct muon tracks in the detector to 0.25° at 1 PeV [432] and shower events to about 7° at 1 PeV [644], IceCube has provided continuous full-sky coverage since detector completion in 2010. The unique location of IceCube at the geographic South Pole results in a clean sample of TeV neutrinos from the Northern sky with world-leading sensitivity to astrophysical neutrino sources, as shown in Fig. 20. These events have been used to test for emission from many source populations, including blazars [645], diffuse Galactic emission [54, 509], gamma ray bursts [442], fast radio bursts [441, 646], and more. IceCube tracks from the Northern sky are also used in broader physics analyses ranging from dark matter searches, flavor physics [81], Lorentz invariance [647], neutrino cross sections [17, 18, 122], and sterile neutrinos [648–651].

IceCube's view of the Southern sky is dominated by muons produced in atmospheric air showers. The harsh cuts necessary to remove atmospheric muons lead to a large reduction in effective area, shown in Fig. 19, limiting the Southern sky sensitivity with tracks. To mitigate the impact of atmospheric backgrounds, the highest-energy tracks, cascades and starting events may be used to perform searches. In 2013, using a sample of these high-energy events starting inside of the detector [35], IceCube reported the first evidence of unresolved flux of astrophysical neutrinos, opening a new window into high-energy physics in the Universe. While no individual sources have yet been identified in the high energy data set [507], characterization of the observed diffuse flux continues with improvements to data selections and analysis methods [53].

IceCube releases information on events likely to be of astrophysical origin to the community in real time. as described in Sec. 2.2.5, allowing partners to search for potential sources of neutrinos within minutes [477]. IceCube also regularly performs follow-up searches for neutrino emission in spatial and temporal coincidence with flares observed with both electromagnetic observatories and gravitational-wave detectors [444]. In 2017, a neutrino released through the IceCube real-time event program, IceCube-170922A, was found to be in coincidence with a gamma-ray flare from blazar TXS 0506+056 [433]. A follow-up analysis unearthed a previously unobserved neutrino emission [22] from the direction of TXS 0506+056. Following the observation of neutrinos from the direction of TXS 0506+056, new analysis tools have been constructed to search the sky for similar emission across 10 years of data [652]. The long operation of IceCube provides more than a decade's worth of data. New data can provide additional sensitivity to TeV physics, but improvements based solely on accumulating data slow the pace of discovery. Instead, significant improvements must come also from progress in selection, reconstruction, and analysis. New studies using starting tracks and shower events reject atmospheric muons better, lowering the energy threshold for observations and dramatically improving the sensitivity in the Southern sky, as shown in Fig. 20, enabling IceCube to continue to perform analyses competitive with planned Northern observatories [644, 654]. Lower-energy data sets targeting astrophysical transients with all-flavor full-sky coverage have recently given new limits on sub-TeV astrophysical phenomena [643, 655] and an expanded data set, shown in Fig. 19, will soon open a new channel for multi-messenger analyses [642]. New event reconstructions relying on updated likelihood constructions [656] or advanced convolutional neural networks [657] show promise in improving the angular and energy resolution necessary for precise characterization of astrophysical fluxes. Recent updates in the point-source likelihood used by IceCube searches aim to improve modeling by utilizing a fuller description of its point spread function [658]. With these updates and future development of machine learning tools, IceCube will continue to be a leader in neutrino astronomy.

4.2.2 KM3NeT

KM3NeT [125] is a multi-purpose cubic-kilometer water Cherenkov neutrino telescope currently being deployed at the bottom of the Mediterranean Sea. It consists of two sub-detectors: ARCA (for Astroparticle Research with Cosmics in the Abyss), located offshore from Capo Passero (Italy) at a depth of 3500 m, and ORCA (for Oscillation Research with Cosmics in the Abyss), placed 2450 m deep offshore from Toulon (France). A KM3NeT string, or Detection Unit (DU), holds 18 Digital Optical Modules (DOMs), each a 17-inch-diameter pressure-resistant glass sphere housing 31 PMTs together with the associated electronics. An array of 115 DUs will constitute a detector building block. Two blocks will be deployed at the ARCA site and one at the ORCA site. Besides the number of building blocks, a different granularity distinguishes ARCA and ORCA. With the DOMs being vertically spaced by 36 m and with a horizontal spacing of 90 m between DUs, ARCA will instrument a total volume of 1 km³ with the primary goal of detecting astrophysical neutrinos in the 100–10⁸ GeV energy range. ORCA, with 9 m (20 m) spacing between DOMs (DUs), is optimized for the study of fundamental neutrino properties using 1–100-GeV atmospheric neutrinos. The combination of these two detectors allows KM3NeT to exploit the full potential of neutrino astronomy, accessing a wide energy range, from MeV-scale core-collapse supernova neutrinos (CCSNs) to astrophysical neutrinos up to several PeV.

ARCA will have an excellent angular resolution: 0.1° at 1 PeV for muon neutrinos, as shown in Fig. 21, and around 1° for showers. It will be a powerful tool for identifying the sources of high-energy cosmic neutrinos. Being located in the Northern Hemisphere, ARCA will have a sensitivity to point-like sources on the Southern sky that will be over one order of magnitude better than the current detectors (see Fig. 21) [659]. Moreover, its geographical location will allow ARCA to test predictions of neutrino production in Galactic sources in the 1–10 TeV energy range, which are based on gamma-ray measurements. As shown in Fig. 21, ARCA will reach the sensitivity to probe the predicted fluxes for several Galactic sources after less than four years of data taking, assuming the observed gamma rays from these sources originate from the interactions of hadrons [659].

The IceCube observation of the diffuse astrophysical flux in 2013 set a milestone in the field of neutrino astronomy. As depicted in Fig. 21, ARCA will quickly confirm the cosmic diffuse neutrino flux with an expected significance of 5σ reached in 1 (0.5) year of operation with one (two) building blocks [125, 660]. It will provide valuable constraints on the neutrino spectrum and flavor composition [659] and its superior angular resolution will enhance the power of multi-messenger follow-up studies for the detected neutrinos, reducing chance coincidences with potential counterpart sources.

ARCA will have a unique sensitivity to detect the diffuse high-energy neutrino emission from the Galactic Plane, especially from the vicinity of the Galactic Center. For certain cosmic-ray propagation models predict-



Figure 21. Top left: ARCA angular resolutions as a function of the neutrino energy for ν_{μ} CC events [659]. Top right: Sensitivity, defined as the median upper limit at 90% CL, of the complete ARCA detector to point-like sources emitting with a E^{-2} spectrum as a function of the source declination after three and seven years of data taking [659]. For comparison, the ANTARES and IceCube sensitivities are also shown. Bottom left: Ratio between the 90% CL sensitivity and the expected neutrino flux normalization for different Galactic sources as a function of the data taking time of the complete ARCA detector [659]. Bottom right: Expected significance in the detection of the cosmic neutrino flux reached by ARCA as a function of the data taking time in either half or full configuration [125, 660].

ing a harder cosmic-ray spectrum in the inner Galaxy, the resulting neutrino flux (the KRA γ models [661]) is expected to be detected with 5 σ significance in four years with the complete detector [660]. Similarly, only three years of data taken by one ARCA building block would be needed to observe the expected neutrino flux from the region of the Fermi Bubbles, in case their gamma-ray emission originates from the decay of pions, and assuming a cutoff in the neutrino energy spectrum at 100 TeV [662].

The world-leading angular resolution, wide energy range, full-sky coverage and 100% duty cycle will make KM3NeT a key player in the field of real-time multi-messenger astronomy, performing fast follow-up-studies of transient phenomena and distributing real-time alerts of interesting neutrino detection [663].

4.2.3 Baikal-GVD

The Baikal Gigaton Volume Detector (Baikal-GVD) [566] is currently the largest operating water Cherenkov neutrino telescope in the Northern Hemisphere, designed to search for high-energy neutrinos of astrophysical origin, whose sources are not yet reliably known. The telescope has a modular structure and consists of clusters. Each cluster is a fully functional detector equipped with detection, triggering, calibration, positioning, and data acquisition systems. The first cluster of Baikal-GVD was deployed on Lake Baikal at a



Figure 22. Top left: Integral distributions of reconstructed events selected by quality cuts versus energy (crosses) and expected events from atmospheric muons (brown histogram) and from a diffuse flux of astrophysical neutrinos (green histogram) [664]. Top right: Accuracy of the angular reconstruction of events in cascade mode for a neutrino energy of 100 TeV. Bottom left: Effective areas of the Baikal-GVD telescope with the 7 clusters deployed as of 2020 in the cascade reconstruction mode for electron neutrino interactions for different ranges of cos(zenith angle) [665]. Bottom right: Preliminary estimates for the single-cluster angular resolution of muon tracks produced in CC neutrino interactions [666].

depth of 1360 meters, 4 km from the shore, starting in 2016. In subsequent seasons, the effective volume of Baikal-GVD was gradually increased by adding one or two functionally independent clusters per season. Further volume expansion to one cubic kilometer by 2025 is ongoing.

Each cluster consists of eight strings 525 m long with optical modules (OM) placed on them to register the Cherenkov radiation in the lake water. The distance between the strings in a cluster is 60 m, where one string is in the center and seven are arranged in a circle around it. The distance between the clusters is 300 m. The OM [667] is a borosilicate glass sphere 42 cm in diameter that houses a PMT of 25.5 cm in diameter. The electronics of the OM are mounted directly on the base of the PMT and include a LED calibration system. Eight GVD clusters have been successfully operating since April 2021 and receiving data from 2304 OMs placed on 64 strings.

The modular structure of the telescope allowed the study of muons and neutrinos in the early stages of detector deployment. Ten (3+7) cascade events have been selected from data collected between 2018 and 2020 with energies above 60 TeV as the first astrophysical neutrino candidates [664]. The energy distributions are shown in the upper left of Fig. 22 for the season 2019-2020 in comparison with expectations from atmospheric muon and diffuse astrophysical neutrino fluxes.

The effective area of the Baikal-GVD telescope with 7 clusters for events from ν_e -induced cascades in the lake water is shown in the lower left of Fig. 22 [665]. At energies above 100 TeV, there is a significant



Figure 23. Left: Design of the Pacific Ocean Neutrino Experiment (P-ONE) showing the strings of photomultipliers, and the light pattern of a 50-TeV neutrino. Right: Charged-current effective areas of $\nu_{\mu} + \bar{\nu}_{\mu}$ for different arrival directions. Figures reproduced from Ref. [126].

decrease in the effective area, due to neutrino absorption in the Earth.

The angular resolution for cascade events is shown in the upper right of Fig. 22. The median value is 2–4 degrees, depending also on neutrino energy and zenith angle. Preliminary estimates of the angular resolution of muon tracks produced in CC neutrino interactions for single GVD cluster are displayed in the bottom right of Fig. 22, along with the its dependence on neutrino energy and the visible track length [666]. An accuracy of better than 1° is achieved for energies above 10 TeV and track lengths greater than 300 m. This precision will allow for significant contributions to multi-messenger searches for neutrino sources via the transmission of neutrino alerts. GVD already participates in follow-up analyses [668]. Presently, the delay in issuing an alert takes on average 3–5 hours [669]. However, a fast mode has been recently implemented [670] that promises low latency alerts on the order of several minutes, which will be useful for multi-messenger alerts.

4.2.4 P-ONE

The vision behind the Pacific Ocean Neutrino Experiment (P-ONE) [126, 671], shown schematically in Fig. 23, is to base a neutrino telescope within a pre-existing large-scale oceanographic infrastructure. As of 2018, the P-ONE collaboration, consisting of researchers in Germany, Canada, and United States, has obtained evidence that the NEPTUNE observatory, operated by Ocean Networks Canada (ONC) since 2009, is an ideal instrumented site to operate a large-volume neutrino telescope.

The 800-km loop of fibre-optic telecommunications cables comprising NEPTUNE provides a high-speed (up to 4 Gbit s⁻¹) and high-power (of 8 kW per node) data link to five nodes that serve as the local hub for the main observations and experiments. A total of 17 primary junction boxes are already wired to the nodes and used to connect hundreds of instruments. The connections are based on underwater mating connectors with field-proven reliability (less than 2% failure in connector pairs deployed over ten years). The ONC success rate of offshore maintenance cruises has been of the order of 95% over several years.



Figure 4.4 Top view of the cecube-cent call detector arrays. Left of the cecube General array, in the negreference "sunflower" design instrumenting). 7.9 km³, Center: the current 2m ever the strings are spaced 240 the low-energy cecube of the focus o detector array (blue), including the DeepCore array with reduced spacing in the center. Above each IceCube string location apettalin Idef opisaidas that be were the source of the source tsphericical on yoto the the idea that a sector of the sec yand Deepe one sitting at a delivered the baseline of ambient bioluminescence and ⁴⁰K and an attenuation length of about 30 meters at 450 nm [672, 673]. The successful pathfinder mission phase triggered a prototype phase in 2021 envisioning the deployment of at least three fully instrumented lines by 2026. The prototyping phase will a characterising ground work and least bill of a power wat 90 Ine multi-cubic kilometer telescope. The scientific goal of P-ONE is to extend IceCube's Gen2, addresses the two main finiting factors in the A second generation observatory, IceCube's Gen2, addresses the two main finiting factors in the the central region of the Galaxy and operate in real time in a complementary way to KM3NeT and GVD. current IceCube instrument: event rate and angular resolution. It does so by improving the event Together, the four telescopes mentioned above can cover the entire sky, optimising possible reciprocal follow-trates and angular resolution by the four telescopes mentioned above can cover the entire sky, optimising possible reciprocal follow-trates and angular interview of the four telescopes mentioned above can cover the entire sky, optimising possible reciprocal follow-trates and angular resolution by the four telescope mentioned above can cover the entire sky. athreeon flogetherathrist results in sensitivity 40 sources five times fainter than visible today. Further, through the addition of an ultra-high-energy radio array, IceCube-Gen2 will expand the accessible

energy range of cosmic peutrinos by several orders of magnitude compared to IceCube.

In-ice optical array (OA): The basic concept for the Gen2 design starts with the idea that the IceCube-Gen2 [128, 674] is a planned next-generation expansion of the IceCube Neutrino Observatory at the primary science goals will target higher energies (from approximately 10 TeV to 10 EeV) than South Pole. IceCube-Gen2 will observe the neutrino sky from TeV to EeV energies and consists of than studes for which IceCube-Gen2 will observe the neutrino sky from TeV to EeV energies and consists of the studes the stude of the rtolienhance the knowledge of the the naturaline medium and of the detector response to pravide

ardireve volution on the near Weal-time multismessenger laterts system for the liow-that legepting high;

tenergy regimes dree the has demonstrated that the diffuse cosmic neutrino flux dominates the

atmosphere backer 2 undiabover 200 the instrument the maps of the construction of the Pole and include the existing IceCube optical array (Fig. 24). DOMs detect the Cherenkov light emitted for hard-spectrum sources comes from events above 10 TeV. Fully exploring these higher energies, from secondary charged particles created in neutrino interactions in the ice and bedrock. The IceCube-Gen2 DOMs will be deployed a requires increasing the active effective area of the detector array. The Gen2 DOMs will be deployed array to cables of surings, with 80 DOMs of sources of string deployed between 1544 n adesize achieves this by neughly doubling the spacing between each string and by deploying more isensitive photom detection modules despendento ithe dee, iresulting the and increase of \$25% in affective rapelutiThesel extensions in the dimensions is concluint the tkn Ewterly gaine drin repetiting "AceCarbe", pattern has menticating the bounder of the standing brand deepi graefinices here down and the stanisht raws of strings). Each IceCube Gen2 DOM has approximately three times the photon collection of the design of the complete array consists of 120 strings (see Frg. 4), where each string is instrumented original IceCube DOM, by employing a multi-photomultiplier-tube design. With optical sensors with similar sensor spacing to that in IceCube. IceCube Gen2 will map the neutrino sky by observing an order of magnitude more neutrino events per year Asl Iplanned, majoritinstrumentation in Gen2 is based on Jerceube feature for the terms of the photomultiplier and the sensor space of the photomultiplier are the photomultiplier and the design.

ferences. A new generation of sensors are being developed for the IceCube Upgrade that have

- 43 -



Figure 25. Left: Earth-skimming technique. The Trinity telescope images the air-shower, which develops in the atmosphere after the interaction of an Earth-skimming tau neutrino inside Earth. *Right:* The integral sensitivity of Trinity to diffuse neutrino fluxes for the prototype telescope in five years and 18 telescopes in ten years, respectively. Updated from Ref. [588].

angular resolution of the optical array for track-like events is approximately 10 arcmin at PeV energies. IceCube-Gen2 can detect neutrino transients from over two times larger distances than IceCube, leading to a factor of 10 source detection volume increase for transient sources. Multi-messenger observations are an ongoing key science goal and will greatly profit from the up to five times higher neutrino alert rate and improved angular resolution when compared to IceCube. IceCube-Gen2's observation of the sky is energy-dependent, shifting from the Northern Hemisphere for energies below 100 TeV to the celestial equator and the Southern Hemisphere for PeV and higher energies.

4.2.6 Trinity

Trinity is a proposed system of 18 air-shower Cherenkov telescopes optimized for detecting Earth-skimming neutrinos with energies between 10 PeV and 1000 PeV [675, 676]. Trinity is an evolution of the original Earth-skimming concept, which proposed monitoring nearby mountains with Cherenkov telescopes [677–679] and has been tested and validated by several groups, for example, by NTA [680] and MAGIC [628].

Trinity's energy range overlaps with IceCube's due to Trinity's PeV threshold. The overlap enables studies of the astrophysical neutrino spectrum in regions of the sky not accessible with in-ice or atmospheric radio experiments (declinations from -75° to 55° [676]). That is because atmospheric radio becomes sensitive at $\sim 10^8$ GeV, and in-ice radio experiments, while sensitive down to $\sim 10^7$ GeV, have a limited sky acceptance due to their locations close to the poles. Trinity closes an important observational gap.

Trinity's telescopes will be located on mountains 2-3 km above the surrounding ground and point at the horizon. At these altitudes, the telescopes can detect air-showers developing as far away as 200 km [630]. The possibility of detecting these very distant showers compensates for the low 20% duty cycle and therefore boosts Trinity's acceptance below 10^8 GeV when compared to other instruments.

Trinity's telescopes are optimized to deliver the best possible detection sensitivity per cost [595]. A key feature of the instrument is the extreme and unique 60-degree wide-field optics (see Fig. 25) [681]. The telescopes use demonstrated technologies lowering costs and improving performance. The mirror technology, for example, has been demonstrated by the Cherenkov Telescope Array Consortium. High efficient, mechanical, and optical robust silicon PMTs populate the focal plane outperforming classical PMTs. The digitizer system has been developed for high-energy physics experiments focusing on low cost and high-channel density. These technologies make it possible to build a high-performance system for a fraction of the cost of a conventional Cherenkov telescope.



Figure 26. Left: The RET-CR concept. The signal from an in-ice transmitter is reflected from the dense in-ice cascade produced by a cosmic ray air shower, and detected by a receiving antenna. Right: The plasma frequency f_p (a proxy for cascade ionization density) versus height for air showers of various energy. Here the 2.4 km surface height is shown at 50 m on the y axis, for clarity. The frequency range of interest to RET-CR is indicated by the vertical red line.

Trinity's sensitivity improves inverse-proportionally with the number of telescopes. One telescope alone can detect astrophysical neutrinos within five years if the spectrum extrapolates from IceCube energies without a cut-off (see Fig. 25).

Trinity is currently in its demonstrator phase [675], which constitutes building a 1 m^2 air-shower Cherenkov telescope that will be deployed on Frisco Peak, UT in 2022. The demonstrator will demonstrate the concept and camera technologies planned for a full Trinity telescope.

4.2.7 RET

The Radar Echo Telescope (RET) is a forthcoming observatory for detecting neutrinos with energies above 10^{16} eV using the radar echo method, where radio waves are reflected from ionization deposits left in the wake of high-energy particle cascades in dense material, such as ice. The concept has recently been demonstrated in the laboratory during experiment T576 at SLAC [129], where a dense beam of high energy electrons was directed into a plastic target, to simulate a high-energy neutrino interaction in ice. The next step, testing the method in the field, is imminent, and will be discussed below.

RET is an umbrella project for two instruments. One, the Radar Echo Telescope for Cosmic Rays (RET-CR) [601] is a prototype system designed to test the radar echo method in nature, shown diagrammatically in Fig. 26. To do this, RET-CR will use the ionized core of a cosmic ray shower—after it has impacted the ice—as a proxy for a neutrino-induced cascade. On a high-elevation ice sheet, 10% or more of the energy of a primary cosmic ray will remain as the cascade impacts the ice. This energy is tightly collimated around the cascade axis, and produces a dense cascade just beneath the surface of the ice. RET-CR will deploy a phased transmitter and several phased receivers (see Refs. [682, 683] for more information on in-ice phased arrays) to detect the ionization left in the wake of these englacial cascades. A detected event rate of roughly 1 event per day with a primary energy at or above approximately 10^{17} eV is expected. RET-CR is under construction at time of writing.

The second system under the RET umbrella is the Radar Echo Telescope for Neutrinos (RET-N), a large-scale, in-ice neutrino detector capable of detecting neutrinos just above the reach of IceCube optical. A RET-N station is comprised of a central transmitter and an array of receivers on baselines of hundreds of meters, all buried ~ 1.5 km deep in the polar ice. RET-N is projected to have good sensitivity down to



TAU AIR-SHOWER MOUNTAIN-BASED OBSERVATORY (TAMBO) - COLCA VALLEY, PERU

Figure 27. Detector concept schematic for TAMBO. A tau neutrino propagates through rock in the mountain. If a charged current interaction occurs, a tau is produced with a range of 50 m–5 km before it decays to produce an air shower. The electromagnetic component of the air shower extends 3–10 km, with a diameter of about 200 m. Such an air shower is detectable by an array of water-Cherenkov tanks, each a m^3 in volume, with 100 m separation. Figure reproduced from Ref. [130]

 10^{16} eV. RET-N is particularly well-suited to flavor studies of UHE neutrinos. Because of the wide angular acceptance of the radar signal, so-called double-bang ν_{τ} events [366] are easily detected—even by a single receiving antenna—leading to an increase in acceptance for such events, providing a promising handle on flavor identification. Furthermore, as the only *active* method for detecting UHE neutrinos, it provides a different method of measuring properties of the cascade, such as direction, energy, and topology. As such, RET-N is complementary to the other radio- and optical-based techniques being explored to bring neutrino astronomy to the UHE regime.

4.2.8 TAMBO

TAMBO (Tau Air-Shower Mountain-Based Observatory) is a proposed detector in the Colca Valley in Peru to measure Earth-skimming astrophysical tau neutrinos in the 1 PeV to 100 PeV energy range (see Fig. 27). Tau neutrinos in that energy range can propagate through Earth with an interaction length of hundreds kilometers [591], which can produce tau particles via charged-current interactions. The produced charged leptons decay within the range of 50 m to 5 km. If the tau neutrino interaction occurs within this distance from the valley, the tau is likely to exit into the air, where it decays producing an air shower with a range of 3–10 km, and a diameter of around 200 m. The air shower can then be detected by water Cherenkov tanks, plastic scintillator panels, or air-Cherenkov telescopes. The preliminary design of TAMBO is based on water Cherenkov detectors, but plastic scintillators are also being studied as a complementary detection medium.

The geographical structure of the Colca valley makes it an ideal place for high-energy tau-neutrino detectors, such as TAMBO. The separation between the slopes of the valley has a length comparable to the decay length scale of a tau. This provides sufficient time for the tau particle to decay to form an extensive air-shower particles, which can eventually be detected by placing a ground array of small water-Cherenkov detectors, each of approximately of volume 1 m³ and separated by ~ 100 m. With this configuration and deploying 22,000 tanks, we expect that TAMBO will have an effective area ten times larger than IceCube's



Figure 28. Schematic of ARIANNA stations at the Ross Ice Shelf. Figure reproduced from Ref. [596]

current tau neutrino effective area at ~ 3 PeV. Compared to ongoing Earth-skimming experiments [684], the deep-valley topography also provides a significant increase to the geometric acceptance compared to a flat ground array. Similarly, the detector location is in the Southern hemisphere, allowing for observation of the Galactic Center and a wide variety of standard astrophysical sources [54, 685] which also holds the prospect of detecting neutrinos from dark matter annihilation [216]. Additionally, the smaller tank-to-tank distance significantly lowers the energy threshold, thereby increasing the rate of detected events. For an $E^{-2.5}$ spectra with the current IceCube best-fit normalization, we expect to detect 7 tau-neutrino events per year over a small background predominantly due to coincident cosmic-ray air showers.

TAMBO will characterize the flux of astrophysical 1–10 PeV neutrinos by measuring the contribution from the tau neutrino component. This would allow a better understanding of high-energy neutrino production [7, 59] and to more precisely test high-energy neutrino physics [5, 58, 90, 93, 686, 687]. Similarly, it would also help determine whether high-energy neutrino sources continue to accelerate particles above 10 PeV, testing some studies [688] that question whether sources of high-energy neutrinos have a cutoff at ~ 6 PeV.

4.3 Ultra-high-energy range (> 100 PeV)

Several experiments search for radio emission from neutrino interactions using embedded radio arrays. Taken together, they serve as complementary test benches for the future radio array of the planned IceCube-Gen2.

4.3.1 ARIANNA

The ARIANNA experiment uses the in-ice radio technique to search for UHE neutrinos [596]. In a uniquely radio quiet area on the Ross Ice Shelf in Antarctica, a hexagonal array of pilot-stations has been taking



Figure 29. Schematic and map ARA stations at the South Pole. Figure reproduced from Ref. [703]

data for several years. In addition, two detector stations have been installed at the South Pole. The station design is shown schematically in Fig. 28. It deployed its first prototype station in 2009 [689] and has been operating a demonstration array since 2014 [596]. Each station uses high gain antennas in both a cosmic ray and veto mode (upward pointing) and in neutrino mode (downward pointing) a few meters below the ice surface. Limits on the UHE neutrino flux from ARIANNA demonstrate the feasibility of the in-ice radio detection technique [690]. The stations provide a continuous test bench for optimization of future larger arrays like IceCube-Gen2 [690, 691] as well as reconstruction and veto methods [552, 692–702].

4.3.2 ARA

The Askaryan Radio Array (ARA) targets the UHE neutrino flux using the in-ice radio technique [597]. An overview of the experiment is provided in Fig. 29. Located near the South Pole in Antarctica, it takes advantage of the deep glacial ice there and the long radio attenuation lengths to search for Askaryan emission from neutrino showers.

In ARA, clusters of radio antennas are deployed 200 m deep below the surface of the ice to access the larger volume of ice visible as the antennas are embedded in denser, glacial ice. Each of five ARA stations uses 16 cylindrical antennas to form an interferometer. The deep antenna deployments at 200 m give the array a wide field of view, stretching from ~ 5° below the horizon to ~ 45° above it. Powered by a DC electric grid, ARA continuously monitors almost one fourth of the sky. One station includes a novel trigger that has been demonstrated to lower the trigger threshold of in-ice Askaryan detectors, thereby lowering the energy threshold and the overall sensitivity of in-ice radio detectors by up to a factor of 2 near 30 PeV [683, 703]. A series of experimental neutrino searches and resultant flux upper limits demonstrate the scalability of the in-ice radio technique [585, 607, 703, 704], and the ability of the detector to both trigger and analyze with high efficiency (> 50%) very low signal to noise ratio data (SNR~ 2), which is essential to achieve the proposed sensitivity of next generation arrays like IceCube-Gen2. ARA anticipates world leading sensitivity to the UHE neutrino flux above 1 EeV by 2023 [585].

Besides searching for neutrinos, ARA has also made a series of measurements which demonstrate the feasibility of the deep technology and the suitability of the South Pole experimental site. Both of these are critical for next generation instruments like IceCube-Gen2. By measuring pulsers deployed on the bottom of IceCube, ARA made the longest horizontal-baseline measurement of the South Pole ice attenuation,



Figure 30. Schematic and map of RNO-G stations being installed at Summit Station in Greenland. Figure reproduced from Ref. [589].

confirming the exceptional clarity (> 1 km attenuation length) of the ice [705]. Using the same pulsers, ARA demonstrated the vertex reconstruction capability of the deep technology, confirming that when a station observes both a direct and reflected pulse, a neutrino vertex can be located to within 1° in zenith and azimuth, and within $\sim 30\%$ in range [706]. This reconstruction capability is essential to the energy resolution of ARA and future radio instruments. ARA data has also revealed potentially novel properties of the Antarctic ice such as birefringence, which mixes the polarization states of a propagating radio signal [706]. Understanding such properties of the medium are critical to neutrino directional reconstruction [707].

4.3.3 RNO-G

The Radio Neutrino Observatory in Greenland (RNO-G) is an in-ice detector that measures neutrinos through the Askaryan emission generated by in-ice showers [155, 589]. The 3 km deep ice sheet above central Greenland with attenuation length of approximately 1 km at the relevant frequencies of 100 MHz to 1 GHz provides a good target material for achieving large effective volumes. The ice will be instrumented with a sparse array of 35 autonomous radio detector stations with a separation of 1.5 km. A schematic and map of the RNO-G stations are shown in Fig. 30. The stations are solar-powered with additional wind generators under development to power the stations during the dark winter months. The stations are connected through an LTE network to Summit station. The first three stations have been installed in 2021, the remaining stations will be installed over the next three years.

Each station is equipped with total of 24 antennas. An interferometric phased array provides a lowthreshold trigger, consisting of 4 bicone antennas installed in close proximity vertically above each other at a depth of approximately 100 m [683]. Additional bicone (vertical signal polarization) and quad-slot (horizontal signal polarization) antennas above the phased array and horizontally displaced on two additional strings provide additional information to reconstruct the properties of the neutrino [708]. LPDA antennas are installed close to the surface providing additional neutrino sensitivity to neutrinos with complementary uncertainties. Each station also comprises three upward facing LPDAs to veto and measure radio emission of air showers which provide in-situ calibration signals [702].



Figure 31. Rendering of the proposed IceCube-Gen2 neutrino observatory combining an optical, surface and radio array.

Due to its relatively low latitude of 72° North, RNO-G observes the majority of the Northern sky within 24 hours adding UHE neutrino information to multi-messenger observations. Its diffuse flux sensitivity is large enough to start probing the parameter space of GZK neutrino production (see Fig. 18). Furthermore, RNO-G will be a technical testbed and pave the way for the much larger radio detector array foreseen for IceCube-Gen2.

4.3.4 IceCube-Gen2 radio

To extend the energy reach to EeV energies, IceCube-Gen2 will include a sparse array of radio detector stations next to its optical component [128] (see Fig. 31). The radio technique allows for a cost-efficient instrumentation of the large volumes required to measure the low flux of UHE neutrinos. The ice at the South Pole provides an optimal target material for radio detection with attenuation lengths of more than 2 km close to the surface where the ice is coldest. The radio array will cover an area of approximately $500 \,\mathrm{km}^2$. The proposed array consists of two types of radio detector stations that measure and reconstruct neutrino properties with complementary uncertainties to maximize the discovery potential by mitigating risks and adding multiple handles for rare background rejection [578]. Hybrid stations combine omni-directional cylindrical antennas (both vertical and horizontal polarization) lowered 150 m below the surface of the ice with high-gain antennas near the surface. The hybrid stations build on the heritage of the deep stations explored by ARA, the shallow stations being explored by ARIANNA, and the hybrid stations currently being deployed for RNO-G. The array design also uses additional shallow-only stations that use LPDA antennas close to the surface with one additional dipole antenna at 15 m to aid event reconstruction. All shallow components are also equipped with upward facing LPDA antennas which provide sensitivity to cosmic rays to veto air-shower induced background [605, 709], as well as to monitor the detector performance (see, e.g., Ref. [702]).

The IceCube-Gen2 radio array will provide sufficient sensitivity to probe GZK neutrino production. The Gen2 sensitivity would reach the current best-fit models to UHECR data measured by the Pierre Auger Observatory, assuming sources identical in UHECR luminosity, spectrum and composition, as well as a



Figure 32. BEACON observes up-going tau neutrinos from a high-elevation mountain. The detection concept is shown on the left while the sensitivity to an all-flavor flux (assuming a 1:1:1 flavor ratio, half-decade energy bins, and a three-year exposure) is shown on the right. The expected 90% confidence upper limit on the differential flux for 100 stations and 1000 stations of both the low- and high-frequency reference designs are compared with models of the cosmogenic flux [147, 391, 710] and upper limits from Auger [431], IceCube [144], and the proposed GRAND-200k experiment [154]. Reproduced from Ref. [157].

rigidity-dependent cut-off and thereby essentially no protons at the highest-energies [148, 149]. In an only slightly more favorable scenario of 10% protons, IceCube-Gen2 will detect at least 2 events per year above \sim 100 PeV [578].

For an unbroken astrophysical neutrino spectrum that follows $E^{-2.28}$, as the one shown in Fig. 18, the radio detector of Gen2 will measure close to ten neutrinos per year where most detected neutrinos will have energies between 10^{17} eV and 10^{18} eV [578]. Due to its location at the South Pole, the instrument continuously observes the same part of the sky with most sensitivity between $\delta \approx -40^{\circ}$ and $\delta \approx 0^{\circ}$ as the Earth is opaque to neutrinos at ultra-high energies. The instantaneous sensitivity will allow to explore neutrino production in transient events such as neutron-star mergers.

The large sensitivity to neutrinos arriving from and slightly below the horizon enables the measurement of the neutrino-nucleon cross section at extremely high energies [102, 103, 105–107]. Also a measurement of the inelasticity seems possible through the detection of high-quality electron neutrino charge-current interaction where the energy of the hadronic shower induced by the breakup of the nucleus can be measured separately from the electromagnetic shower induced by the electron. Furthermore, the production of high-energy muons in air showers with energy above a PeV can be probed through a coincident measurement of the air shower via the in-air radio emission and a muon induced particle shower in the ice via the Askaryan emission.

4.3.5 BEACON

The Beamforming Elevated Array for COsmic Neutrinos (BEACON) is a detector concept that targets Earth-skimming tau neutrinos. BEACON uses the radio technique to search for air showers induced by tau lepton decay, and will be deployed at several sites around the world for a full sky coverage [157]. Because the BEACON concept relies on low power (< 50 W), inexpensive, low channel-count instrumentation, the technique is scalable to many stations that can be deployed at one or at many locations around the world.

BEACON is an efficient design that builds a large neutrino volume by fully exploiting the mountain geometry and by using interferometric phased arrays [157]. The design, shown schematically in Fig. 32, uses the high-elevation of a mountain to monitor a large area of the ground for up-going tau neutrinos fully exploits the topography at a site to maximize the neutrino interaction volume with sparse instrumentation. Stations consist of a clustered phased array used for triggering and antennas placed at longer baselines for

improved pointing resolution (sub-degree). Phased arrays are advantageous in this geometry, because they both improve sensitivity and can be tuned to focus in on the horizon where most of the tau neutrinos are expected. Air showers are expected to be observable up to 100 km away from the detector, making mountains with 2–3 km prominence ideal sites to search for up-going tau neutrinos.

Within 3 years of observations with a full-scale instrument consisting of 1000 stations, BEACON is expected to achieve a sensitivity comparable to pessimistic models of cosmogenic neutrinos, but even with a smaller array of 100 stations, the large effective area will improve upon existing limits within 3 years of observations [157] The sensitivity to a diffuse flux is shown on the right in Fig. 32. The chosen frequency band is 30-80 MHz, but sensitivity studies have shown that a higher frequency band, for instance 200 - 1200 MHz could also be suitable [157]. This leaves the possibility to deploy hybrid frequency-band arrays in order to combine them to improve the pointing and the reconstructions capabilities.

The BEACON experiment is currently in the demonstration phase, with a prototype deployed at the White Mountain Research Station of California. The prototype array consists of 4 dual-polarized electricallyshort-dipole antennas and a two stages of amplification. The trigger system is based on a low-power phased array trigger implemented on an FPGA. While the prototype electronics were originally developed for the ARA experiment [683], custom modular electronics are under design for future stations.

While the prototype is too small to search for tau neutrinos, the technique can be tested thoroughly through searches for cosmic rays at high zenith angles [711]. A cosmic-ray search is underway currently [712], and with the observed flux, the projected neutrino sensitivity will be trained on the measured trigger threshold *in situ*. With this validation of instrument sensitivity, the experiment can be readily expanded to the full array making it competitive to detect UHE neutrinos [148].

4.3.6 GRAND

The Giant Radio Array for Neutrino Detection (GRAND) is a planned observatory for UHE particles, including neutrinos, cosmic rays, and gamma rays. The instrument is a large-scale radio array sensitive to the geomagnetic radio emission air showers in the atmosphere. In particular, GRAND targets UHE neutrinos by searching for very inclined showers, i.e., showers coming from directions close to the horizon, expected from Earth-skimming UHE tau neutrinos [154]. Figure 33 shows the GRAND concept and the differential and integrated neutrino sensitivity limits for several antenna configurations. The design targets diffuse flux sensitivities predicted by pessimistic models of UHE neutrino flux down to $\sim 10^{-10} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$.

To reach this target sensitivity, GRAND will instrument a large 200000 km^2 area with clusters of radio antennas. Each of the 20 clusters will use 10000 radio antennas to monitor the horizon for emerging tau neutrinos. Each antenna operates in the 50–200 MHz band and the local topology will be analyzed to determine the optimal locations of the antennas within the clusters and the array [713] both for improved sensitivity and to enable sub-degree angular resolution [154, 714]. With such precision angular resolution, GRAND will be able to to distinguish between neutrinos and the more prevalent cosmic rays, as the neutrino showers are horizontal or slightly upward-going while cosmic-ray showers are more downward-going. Moreover, GRAND will be an important instrument in transient time-domain radio astronomy, because of its large number of antennas and field of view.

The radio-detection of extensive air showers [715] has been demonstrated by prior experiments and GRAND builds on this heritage. However, scaling to a large array requires a staged approach to further refine the design. The first prototype array, GRANDProto300, will use 300 antennas to validate autonomous radio detection and reconstruction of extensive air showers. This large array will also enable studies of other UHE particles including cosmic rays and photons as well as radio astronomy [600]. The second prototype stage will scale up to 1000 antennas with GRAND10k, building on the design from the 300-antenna array and will explore the feasibility of the radio detection with large-scale and sparse arrays. GRAND10k is expected to reach a diffuse flux sensitivity of 8×10^{-9} GeV cm⁻² s⁻¹ sr⁻¹ after 3 years [154], constraining or discovering UHE neutrinos expected from optimistic models [140, 148]. With this staged approach, the



Figure 33. Left: The GRAND detection principle relies on the observation of EAS induced by both the direct interaction of the cosmic particles in the atmosphere (e.g., cosmic rays and gamma rays), and neutrino underground interaction with subsequent decay of the tau lepton in the atmosphere. The GRAND concept takes advantage of the detector topography in order to increase the detection efficiency [713], and the reconstruction performances [714]. Right: The differential and integrated neutrino sensitivity limits derived from the 10, 000 antennas simulation (GRAND10k, pink area) and its extrapolation to the 20-times larger GRAND array (GRAND200k, maroon line). The gray region represents the expectations for the all-flavor cosmogenic neutrinos flux obtained from the results of the Pierre Auger Observatory [148]. Taken from [154].

design can be refined as the instrument scales up to the full 200000 antennas while simultaneously addressing key science questions in physics and astrophysics.

4.3.7 **POEMMA**

The Probe Of Extreme Multi-Messenger Astrophysics (POEMMA) is a NASA Astrophysics Probe-class, space-based mission designed to perform definitive measurements of Ultra-High Energy Cosmic Rays (UHE-CRs) and searches for cosmic neutrinos. POEMMA was developed as one of the examples of NASA-funded missions to support the definition of the NASA Probe-class, and it is a potential candidate for a future NASA Probe Announcement of Opportunity. POEMMA utilizes the vast amount of atmosphere and Earth viewed using wide Field-of-View (FoV) as neutrino targets and measures the optical signals from the extensive air showers (EAS) from the products of the neutrino interactions [558]. Over a five-year period, POEMMA is anticipated to measure ~ 1,400 UHECRs above 40 EeV and either detect the first neutrino with $E_{\nu} > 20$ PeV for tau neutrinos and $E_{\nu} > 20$ EeV for any flavor of neutrinos or set limits on the diffuse flux and stringent limits on transient source models with significant neutrino flux above 20 PeV.

The POEMMA mission consists of two identical spacecraft that fly in a loose formation at 525 km altitude, 28.5° inclination orbits, separated by a nominal lateral distance of 300 km. Each POEMMA spacecraft contains a Schmidt telescope with 6 m^2 optical collecting area and a 45° full FoV. The focal plane of each telescope is segmented into two sections, each optimized for different EAS optical measurements, and with each section comprised of pixels with 0.086° FoV. The POEMMA Fluorescence Camera occupies 80% of each focal surface and is optimized to record the fluorescence light from EAS initiated by UHECR in the atmosphere. The POEMMA Cherenkov Camera occupies the remaining 20% of the focal surface and is oriented to observe near the Earth's limb, when the POEMMA telescopes are appropriately tilted, to record



Figure 34. Top: POEMMA observing modes: POEMMA-Stereo for UHECRs and UHE ν -induced extensive air showers, and POEMMA-Limb for $\nu_{\tau} \rightarrow \tau$ -sourced up-going extensive air showers. Figure reproduced from Ref. [558]. Bottom left: POEMMA 90% CL per decade, 5-year sensitivity to EAS showers arising from neutrino interactions in the atmosphere and detected via the stereo fluorescence measurements from charged-current and neutral-current interactions from all three neutrino flavors. The solid (dashed) curve is calculated using cross sections from Ref. 591 (Ref. 716). Predictions for strongly coupled string moduli (maroon band) [150] are also shown along with upper limits from ANITA I-IV (blue line) [146] and Westerbork Synthesis Radio Telescope (WSRT; red line with tan band) [717], and the projected sensitivity for LORD (green band) [718]. Bottom right: POEMMA 90% CL per-decade tau neutrino sensitivity assuming the binary neutron star merger (BNS) model of Fang and Metzger [719] for a source at a distance of 10 Mpc. The sky-location sensitivity range for IceCube is shown in the blue band while that for GRAND is in the area defined by the red dash. The black lines show the measured limits to neutrinos from GW170817 set by IceCube and Auger [25]. Figure reproduced from Ref. [558].

the beamed, Cherenkov emission produced by EAS sourced from Earth-skimming neutrino interactions above 20 PeV. The POEMMA telescopes can slew in both azimuth (90° in ~ 8 minutes) and zenith, allowing for unprecedented follow-up on transient astrophysical events by tracking sources as they move across the sky as viewed below the limb of the Earth [582]. The separation of the POEMMA spacecraft can also be decreased to ~25 km to put both telescopes in the upward-moving EAS light pool for each event, thus reducing the neutrino detection energy threshold. The period of the POEMMA spacecraft orbit is 95 minutes. Due to this and the POEMMA telescopes viewing orientation, one of the main advantages of the POEMMA mission is being able to achieve full-sky coverage for both UHECR and UHE neutrino sources. A schematic of the fluorescence telescope configuration viewing down and the Cherenkov telescope viewing towards the limb in the upper panel of Fig. 34 shows POEMMA's two distinct science modes.

The first science operating mode is a precision UHECR and UHE neutrino stereo mode where the telescopes are oriented to co-measure the EAS air fluorescence signal in a common volume corresponding to nearly 10^{13} tons of atmosphere. Due to the high accuracy of the EAS reconstruction from the stereo fluorescence technique when viewing the entire EAS development using the large FoV from low-Earth orbit,

POEMMA can accurately reconstruct the development of the EAS with $\leq 20^{\circ}$ angular resolution, $\leq 20\%$ energy resolution, and $\leq 30 \text{ g/cm}^2 X_{\text{max}}$ resolution [135]. This performance yields excellent sensitivity for all neutrino flavors for UHE EAS that begin deeper in the atmosphere, well-separated from the dominant UHECR flux.

At neutrino energies above EeV, in both charged-current and neutral-current interactions, ~80% of the neutrino energy is carried by the emergent lepton and ~ 20% goes into a hadronic cascade [591]. Thus, the properties of the composite EAS are determined by the emergent UHE lepton for both charged- and neutral-current interactions in the atmosphere. The wide FoV for POEMMA allows the measurement of long, more horizontal EAS generated by neutrino interactions deep in the atmosphere, providing strong rejection of UHECR-induced EAS [135]. Fig. 34 shows the 90% CL per-decade all-flavor limits for 5-years of POEMMA fluorescence UHE measurements assuming two different neutrino cross sections [591, 716] at the highest energies. Comparison to the published limits from ANITA [146] and WSRT [717] observations are shown as well as theoretical predictions of the neutrino flux from strongly coupled string moduli generated early in the formation of the Universe [150].

The second POEMMA science mode has the telescopes pointed to view slightly below the limb of the Earth to be sensitive to the beamed, optical Cherenkov signal sourced by tau-neutrino interactions in the Earth. Tau leptons produced by neutrinos in the Earth can emerge and decay to produce up-moving EAS. This $\nu_{\tau} \rightarrow \tau$ detection channel allows POEMMA to have a sensitivity to energies $E_{\nu} \gtrsim 20$ PeV. POEMMA's FoV constrains viewing to an azimuth band of ~ 30°; however, for an idealized 2π azimuth telescope configuration with POEMMA's optical Cherenkov performance, the limits on the diffuse neutrino flux set by IceCube could be improved by roughly an order of magnitude for $E_{\nu} > 100$ PeV [592, 622]. In principle, POEMMA is also sensitive to Earth-emergent neutrinos through the $\nu_{\mu} \rightarrow \mu$ and $\nu_{\tau} \rightarrow \tau \rightarrow \mu$ interaction channels. These channels improve POEMMA's sensitivity for $E_{\nu} < 10$ PeV due to the relatively long interaction lengths of the muon around 1 PeV, and in the case of the primary ν_{μ} , increased Earthemergence probabilities.

The capability of the POEMMA satellites to quickly slew to the direction of an astrophysical transient neutrino source and follow the source give POEMMA unique observational capabilities using the Cherenkov telescopes. As shown in Fig. 34, it is expected that, compared to ground-based experiments, POEMMA will improve upon the sensitivity to long-burst transient events (duration of 10^{5-6} s from, e.g., binary neutron star mergers and tidal disruption events) for $E_{\nu} > 100$ PeV by nearly an order of magnitude. Additionally, POEMMA will improve on the sensitivity to short-burst (duration of $\sim 10^3$ s from, e.g., short gamma ray bursts with extended emission) by at least an order of magnitude [582], as illustrated in Fig. 17.

4.3.8 EUSO-SPB2

The Extreme Universe Space Observatory aboard a Super Pressure Balloon 2 (EUSO-SPB2) is the follow-up mission to the EUSO-Balloon and EUSO-SPB1 missions, flown in 2014, and 2017, respectively. EUSO-SPB2 will implement the technologies utilized during past EUSO missions in a near-space environment to validate the detection strategy of POEMMA and future space-based observatories by measuring cosmic rays via both fluorescence and optical Cherenkov emission [599, 720].

The EUSO-SPB2 instrument contains two telescopes, each with a different science target: the Fluorescence Telescope (FT) points downward and will measure the microsecond-scale fluorescence tracks of EAS induced by UHECR interactions in the atmosphere, while the Cherenkov Telescope (CT) points towards the limb to measure the nanosecond-scale Cherenkov emission produced by EAS induced by above-the-limb cosmic rays and Earth-skimming neutrinos. The Field of View (FoV) of the EUSO-SPB2 FT is $11^{\circ} \times 35^{\circ}$, while that of the CT is $4^{\circ} \times 12.8^{\circ}$. The EUSO-SPB2 flight train will include an azimuth rotator configured for day and night pointing and an elevation angle tilting mechanism on the CT. EUSO-SPB2 will launch from Wanaka, NZ in Spring of 2023, and aims for a 100-day flight at an altitude of 33 km. In addition to quantifying the night-sky airglow background near the Earth limb for the optimization of future space and near-space missions, the EUSO-SPB2 CT is expected to observe hundreds of direct cosmic rays per hour of operational live time during its flight [721]. These events share many characteristics with the signals generated from neutrino sourced events and are a prime candidate for event reconstruction and evaluation of detector optics, electronics, and triggering algorithms. A larger data set of these above-the-limb events will help to quantify refraction near the Earth limb, and may help evaluate similar events that have been observed by ANITA, and are likely to be observed by PUEO. Due to its relatively short flight duration and small azimuth FoV, EUSO-SPB2 is not competitive with existing ground-based experiments in regards to setting limits on the diffuse neutrino flux [622]. However, by having the ability to slew in azimuth and zenith, EUSO-SPB2 can perform "target of opportunity" observations, where it will search for neutrino sources following alerts of astrophysical transients [722, 723].

4.3.9 ANITA

The Antarctic Impulsive Transient Antenna (ANITA) is a NASA mission that searched for ultra-high energy particles from a long-duration balloon in Antarctica [724]. Originally designed to search for neutrino-induced Askaryan radiation that refracts out of the ice, successive flights and studies demonstrated its capability of detecting geomagnetic radiation from cosmic rays [311, 312, 725–727] and its sensitivity to Earth-skimming tau neutrinos [317, 319]. The upper limits from ANITA's Askaryan channel remain the strongest in the energy range above 30 EeV [146, 611, 728, 729]. In three of the flights, its trigger was sensitive to the geomagnetic radiation from air showers and in each of these flights it reported anomalies that remain outstanding questions in the field [311, 312, 727]. It has additionally set constraints on the search for magnetic monopoles [730] and Lorentz-invariance violation [491].

The great advantage of the sub-orbital platform is that the high float altitude enables the largest instantaneous effective areas in the 10 EeV–ZeV energy range and a unique observing geometry. The ANITA payloads consisted of broadband quad-ridged horn antennas with a maximum bandwidth of 180–1200 MHz with improved trigger sensitivities, lower noise temperatures, and extensions with each subsequent flight. While the trigger used a combinatoric trigger, interferometric analysis of the signal enabled ANITA to reconstruct the radio-signal to within $0.1^{\circ}-0.2^{\circ}$ [731], which translates to an elongated ellipse in the Askaryan channel with a maximum extent of a few degrees and to ~1° for the in-air tau neutrino channel. This beamforming technique is carried forward and implemented at the trigger level for PUEO, dramatically lowering its trigger and energy threshold [581].

4.3.10 PUEO

The Payload for Ultrahigh Energy Observations (PUEO) (Fig. 35) is a long-duration Antarctic balloon experiment designed to have world-leading sensitivity to UHE neutrinos at energies above 1 EeV. The direct successor to the ANITA experiment, it expects more than an order of magnitude more sensitivity than its predecessor below 30 EeV [156, 581]. A multi-purpose instrument for detecting ultra-high-energy particles, PUEO can search for radio emission from neutrinos both from when they interact in the ice and from Earth-skimming tau neutrinos. PUEO is also sensitive to geomagnetic radio emission from UHECRs, including showers that evolve in the stratosphere. The sensitivity of PUEO allows it to probe a host of diffuse cosmogenic and astrophysical neutrino flux models, with multiple neutrinos expected during the nominal 30 day flight under the most optimistic flux models allowed by current limits [581]. PUEO's very large instantaneous aperture makes it well-suited to measuring UHE neutrino fluences from transient astrophysical sources, if they occur in PUEO's field-of-view during the flight.

The main PUEO instrument, seen in Fig. 35, follows the same overall design of ANITA but with significant changes to the internal electronics and improvements to the signal chain to significantly increase the sensitivity in all of its four detection channels (in-ice Askaryan, above-horizon stratospheric UHECR, reflected UHECR, and Earth-skimming τ -induced EAS). Like the ANITA experiment, PUEO will be sensitive



Figure 35. PUEO, the successor to the ANITA experiment. *left* PUEO has two different antenna arrays: the main instrument, comprised of over 100 quad-ridged horn antennas; and a low-frequency instrument, consisting of several channels of sinuous antennas, that drops down below the payload (seen here) and is designed to increase PUEO's sensitivity to extensive air shower-like signals from τ -lepton-decays in the air, and UHECRs. From Ref. [581]. *right* PUEO is designed to search for UHE neutrino interactions in the Antarctic ice via the Askaryan effect, and will have world-leading sensitivity to UHE neutrinos with energies above 1 EeV [156, 581].

to tau neutrinos through the radio emission from both in-ice (Askaryan) events and in-air τ -decay-induced showers.

Unlike ANITA, PUEO is composed of two instruments, each dedicated to different detection channels. The "main instrument" consists of 96 triggered quad-ridged horn antennas (compared to the 48 used in the last flight of ANITA), with an additional ring of downward-canted antennas that increase PUEO's sensitivity to the anomalous steeply upcoming cosmic-ray-like events observed in past ANITA flights [312, 732]. The main instrument targets the 300 MHz to 1200 MHz frequency band, similar to the ANITA instrument, but with a slightly higher low-frequency limit. The multi-channel "low-frequency instrument" targets the 50–300 MHz band designed to detect EAS signals, from UHECRs or τ -lepton decays, extending the frequency range and allowing for an independent measure of polarity for events due to air showers. PUEO will also deploy an improved radio-frequency signal chain and more accurate attitude determination systems, when compared to ANITA-IV.

Despite deploying more than twice the number of antennas as the last flight of ANITA, the biggest improvement in PUEO's sensitivity comes from the real-time interferometric beamforming used in the trigger subsystem. This beamforming trigger computes highly directional beams on the sky, in real-time during the flight, by coherently summing waveforms with different time delays, allowing PUEO to have a 50% trigger efficiency level at a signal-to-noise ratio (SNR) of ~1 [581]. PUEO's beamforming trigger system, with extensive heritage from the phased-array trigger of the Askaryan Radio Array [683], is built on the Xilinx RFSoC (Radio-Frequency System-on-Chip) platform which combines high-bandwidth digitizers, large fieldprogrammable gate-arrays (FPGAs), and digital signal processing (DSP) cores onto a single die.

4.3.11 AugerPrime

With the present Surface Detector (SD) of the Pierre Auger Observatory—an array of ~ 1600 Water-Cherenkov Detectors (WCD) spread over a surface of ~ 3000 km² in an hexagonal grid with 1.5 km separation—UHE neutrinos can be efficiently discriminated in the much larger background of cosmic rayinduced EAS. The neutrino search capabilities are enhanced at EeV energies for EAS developing in the inclined directions with respect to the vertical to ground, at zenith angles $\theta \in (60^{\circ}, 95^{\circ})$, by looking for penetrating showers with a significant amount of electromagnetic component, characteristic of neutrino-induced



Figure 36. *Left:* Picture of an upgraded water-Cherenkov station of AugerPrime. The flat scintillator detector and the SALLA antenna are visible on top of the station, along with the solar panel and the communications Yagi antenna. Source: Pierre Auger Observatory. *Right:* Footprint of a simulated upward-going tau-induced shower produced by a tau neutrino skimming the Earth. The colors indicate the time of arrival of the shower particles at the water-Cherenkov stations from earliest (violet) to latest (red). The black arrow is parallel to the projection of the shower axis on the ground. The radii of the colored stations are proportional to the logarithm of the signal deposited, and the numbers on top are the fraction of the signal due to the electromagnetic component of the shower. The gray dots represent stations of the array without signal. A large fraction of electromagnetic signal in inclined events such as this one, concentrated in the earliest part of the shower, constitute a signature of neutrino-induced showers.

EAS close to the ground [431].

AugerPrime is the upgrade of the Pierre Auger Observatory [152]. It is fully funded, under construction, and foreseen to take data until 2030. A complementary measurement of the shower particles will be provided by an array of plane plastic scintillators (SSD) [733] above the existing WCD (Fig. 36). This allows sampling the shower particles simultaneously with two detectors that have different responses to the muonic and electromagnetic components of the shower, potentially improving the identification of neutrino-induced EAS. The geometrical acceptance of the flat scintillators is reduced for very inclined showers $\theta > 75^{\circ}$ where neutrino identification is currently performed with the highest efficiency [431]. However, for moderate inclination angles $\theta \in (\sim 60^{\circ}, \sim 75^{\circ})$ the expectation is that the SSD can improve neutrino selection in an angular range where the current search algorithms reach efficiencies typically below 75%, and even extend it to $\theta < 60^{\circ}$, where searches for neutrinos are not performed due to the large cosmic-ray background not reducible in an efficient manner with current algorithms.

Besides more than doubling the current statistics, AugerPrime will have superior cosmic-ray composition measurement capabilities compared to the WCD array [733]. Two of the major objectives of AugerPrime are to elucidate the mass composition and the origin of the flux suppression at the highest energies as well as search for a flux contribution of protons up to the highest energies as low as 10%. The proton fraction is known to have a large effect on the intensity of the cosmogenic neutrino flux [397], constituting a decisive



Figure 37. Illustration of total collection area for existing installations and GCOS. From Ref. [153].

ingredient for estimating the physics potential and design of future detectors searching for UHE neutrinos.

An integral part of the AugerPrime detector is the Radio Detector (RD) [734]. Built on the experience gained with the construction and operation of the Auger Engineering Radio Array [735], the goal is to install an antenna on top of each WCD to detect the radio signal induced in EAS by geomagnetic charge separation in the shower [736] and the Askaryan effect [604]. The RD is most efficient for the detection of inclined showers [737], and the radio footprint at ground is known to contain information on X_{max} [738], the depth at which the shower reaches the maximum in its development. The RD will provide a clean measurement of the electromagnetic shower component, ideal to separate neutrino-induced cascades from hadronic showers. These two facts open up the possibility of using the radio detector in the search for penetrating inclined showers induced by UHE neutrinos, complementing and aiding in current searches.

The surface detector stations will be upgraded with faster, better timing accuracy and increased dynamic range electronics [739]. The new electronics has the resources to implement further trigger algorithms on top of the existing ones, targeted to physics tasks such as searches for photons and neutrinos at sub-EeV energies, including the possibility of incorporating the WCD, SSD, and RD to the trigger providing a wealth of information on a single event. This is only starting now to be exploited in Auger [740].

4.3.12 GCOS

The Global Cosmic Ray Observatory (GCOS) is presently in the design phase, with the final detection concept and setup being worked out. More detailed considerations can be found in Ref. [153] and references therein. With the goal of reaching an exposure of at least 2×10^5 km² yr in a period of 10 years and full-sky coverage a set of surface arrays with a total area of about 40000 km² is anticipated as shown in Fig. 37.

GCOS will be designed as a multi-messenger observatory, aiming to register charged cosmic rays, gamma rays, and neutrinos with energies above 10^{19} eV. Different detection concepts are at hand. They need to be optimized to reach the targeted physics case. The design will include a surface detector array with segmented/nested water Cherenkov detectors and radio antennas. Similarly to AugerPrime, as described above, but with a much bigger area/acceptance, the surface detector array of GCOS will be able to search for neutrinos by identifying electron-rich horizontal air showers and distinguish them from hadronic cascades. The radio detector will provide a clean measurement of the electromagnetic component, ideal to identify neutrino-induced cascades. Nested water-Cherenkov detectors will also provide enhanced mass and neutrino sensitivity for horizontal air showers. Identification of the sources of UHE particles will require a good angular resolution. Assuming a detector spacing of the order of 1.6 - 2 km, an angular resolution $< 0.5^{\circ}$ is realistic. GCOS is at present in the phase of developing its science case and studies are being conducted to optimize a suitable detector design. The performance of existing detector systems, in particular AugerPrime, gives concrete and proven design examples to achieve the needed sensitivity for UHE neutrinos. The final design of GCOS will depend on the results that will be obtained with the Auger Observatory and the Telescope Array in the coming decade. New findings will influence the science case and, thus, the design of the observatory. A promising approach towards a full-scale GCOS could be to gradually increase the aperture of the existing arrays. For example, discoveries brought about by increasing the aperture of present-day Auger a few times will improve the understanding about the highest-energy particles and will clarify the neutrino-related design goals for an even larger observatory. Prototype detectors are expected to be built after 2025. The construction of GCOS at multiple sites is expected to start after 2030, with an anticipated operation time of at least twenty years.

Acknowledgements

MA is supported by the Initiative and Networking Fund of the Helmholtz Association, the Deutsches Elektronen Synchrotron (DESY). MB is supported by the VILLUM FONDEN under project no. 29388. LL is supported by the US National Science Foundation (US NSF)-Office of Polar Programs, the US NSF Physics Division through Awards PHY-1913607 and the Wisconsin Alumni Research Foundation. NO is supported by the US NSF through Award 2112769. MHR is supported in part by the US Department of Energy (US DOE) grant DE-SC-0010113. SAW is supported by the US NSF through Awards 2111232 and CAREER 2033500 and NASA through Awards 80NSSC20K1288, 80NSSC20K0925, and 80NSSC20K0775. SKA and AN acknowledge the financial support from the Swarnajayanti Fellowship Research Grant (File no.SB/SJF/2020-21/21) provided by the Science and Engineering Research Board (SERB) and Department of Science and Technology (DST), Govt. of India. SKA would like to thank the United States-India Educational Foundation for providing the financial support through the Fulbright-Nehru Academic and Professional Excellence Fellowship (Award No. 2710/F-N APE/2021). JAM and EZ have received financial support from Xunta de Galicia (Centro singular de investigación de Galicia accreditation 2019-2022), by European Union ERDF, and by the "María de Maeztu" Units of Excellence program MDM-2016-0692 and the Spanish Research State Agency, and from Ministerio de Ciencia e Innovación PID2019-105544GB-I00 and RED2018-102661-T (RENATA). RAB is funded by the "la Caixa" Foundation (ID 100010434) and the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 847648, fellowship code LCF/BQ/PI21/11830030. BAC thanks the National Science Foundation for support through the Astronomy and Astrophysics Postdoctoral Fellowship under Award AST-1903885. PBD acknowledges support from the US DOE under Grant Contract DE-SC0012704. YF would like to acknowledge support from the ICTP through the Associates Programme and from the Simons Foundation through grant number 284558FY19. YSJ is supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government through Ministry of Science and ICT Grant 2021R1A2C1009296. The work of SRK was supported in part by the US NSF under grant number PHY-1307472 and the US DOE under contract number DE-AC-76SF00098. JFK acknowledges support by NASA grant 80NSSC19K0626 at the University of Maryland, Baltimore County and under proposal 17-APRA17-0066 at NASA/GSFC. ER is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) SFB 1258, Cluster of Excellence 2094 ORIGINS, and the European Research Council (ERC) (Grant agreement No. 101053168). ZhD and OVS are supported by the Ministry of Education and Science of the Russian Federation within the financing program of large scientific projects of the "Science" National Project (grant no. 075-15-2020-778). The work of SY is supported by JSPS KAKENHI Grant No. 18H05206. BZ is supported by the Simons Foundation.

References

- Committee for a Decadal Survey on Astronomy and Astrophysics 2020 (Astro2020), Space Studies Board, Board on Physics and Astronomy, Division on Engineering and Physical Sciences, "Pathways to Discovery in Astronomy and Astrophysics for the 2020s." https://www.nap.edu/catalog/26141/ pathways-to-discovery-in-astronomy-and-astrophysics-for-the-2020s.
- R. Alves Batista et al., "EuCAPT White Paper: Opportunities and Challenges for Theoretical Astroparticle Physics in the Next Decade", 2110.10074.
- [3] M. Ahlers, K. Helbing and C. Pérez de los Heros, "Probing Particle Physics with IceCube", <u>Eur. Phys. J. C</u> 78 (2018) 924 [1806.05696].
- [4] M. Ackermann et al., "Fundamental Physics with High-Energy Cosmic Neutrinos", <u>Bull. Am. Astron. Soc.</u> 51 (2019) 215 [1903.04333].
- [5] C. A. Argüelles, M. Bustamante, A. Kheirandish, S. Palomares-Ruiz, J. Salvadó and A. C. Vincent, "Fundamental physics with high-energy cosmic neutrinos today and in the future", <u>PoS</u> ICRC2019 (2020) 849 [1907.08690].
- [6] J. L. Feng et al., "The Forward Physics Facility at the High-Luminosity LHC", in 2022 Snowmass Summer Study, 3, 2022, 2203.05090.
- M. Ackermann et al., "Astrophysics Uniquely Enabled by Observations of High-Energy Cosmic Neutrinos", Bull. Am. Astron. Soc. 51 (2019) 185 [1903.04334].
- [8] ICECUBE Collaboration, "Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector", Science 342 (2013) 1242856 [1311.5238].
- [9] SUPER-KAMIOKANDE Collaboration, "Evidence for oscillation of atmospheric neutrinos", <u>Phys. Rev. Lett.</u> 81 (1998) 1562 [hep-ex/9807003].
- [10] SUPER-KAMIOKANDE Collaboration, "Solar B-8 and hep neutrino measurements from 1258 days of Super-Kamiokande data", Phys. Rev. Lett. 86 (2001) 5651 [hep-ex/0103032].
- [11] SUPER-KAMIOKANDE Collaboration, "Constraints on neutrino oscillations using 1258 days of Super-Kamiokande solar neutrino data", Phys. Rev. Lett. 86 (2001) 5656 [hep-ex/0103033].
- [12] SNO Collaboration, "Measurement of the rate of $\nu_e + d \rightarrow p + p + e^-$ interactions produced by ⁸B solar neutrinos at the Sudbury Neutrino Observatory", Phys. Rev. Lett. 87 (2001) 071301 [nucl-ex/0106015].
- [13] SNO Collaboration, "Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory", Phys. Rev. Lett. 89 (2002) 011301 [nucl-ex/0204008].
- [14] SUPER-KAMIOKANDE Collaboration, "Determination of solar neutrino oscillation parameters using 1496 days of Super-Kamiokande I data", Phys. Lett. B 539 (2002) 179 [hep-ex/0205075].
- [15] SNO Collaboration, "Electron energy spectra, fluxes, and day-night asymmetries of B-8 solar neutrinos from measurements with NaCl dissolved in the heavy-water detector at the Sudbury Neutrino Observatory", <u>Phys.</u> Rev. C 72 (2005) 055502 [nucl-ex/0502021].
- [16] ANTARES Collaboration, "All-flavor Search for a Diffuse Flux of Cosmic Neutrinos with Nine Years of ANTARES Data", Astrophys. J. Lett. 853 (2018) L7 [1711.07212].
- [17] ICECUBE Collaboration, "Measurement of the multi-TeV neutrino cross section with IceCube using Earth absorption", Nature 551 (2017) 596 [1711.08119].
- [18] M. Bustamante and A. Connolly, "Extracting the Energy-Dependent Neutrino-Nucleon Cross Section above 10 TeV Using IceCube Showers", Phys. Rev. Lett. 122 (2019) 041101 [1711.11043].
- [19] ICECUBE Collaboration, "Measurements using the inelasticity distribution of multi-TeV neutrino interactions in IceCube", Phys. Rev. D 99 (2019) 032004 [1808.07629].
- [20] ICECUBE Collaboration, "A novel method of rejecting muon backgrounds for the detection of the highest energy neutrinos", PoS ICRC2019 (2019) 945.

- [21] ICECUBE Collaboration, "Detection of a particle shower at the Glashow resonance with IceCube", <u>Nature</u> **591** (2021) 220 [2110.15051].
- [22] ICECUBE Collaboration, "Neutrino emission from the direction of the blazar TXS 0506+056 prior to the IceCube-170922A alert", Science **361** (2018) 147 [1807.08794].
- [23] R. Stein et al., "A tidal disruption event coincident with a high-energy neutrino", <u>Nature Astron.</u> 5 (2021) 510 [2005.05340].
- [24] P. Giommi, T. Glauch, P. Padovani, E. Resconi, A. Turcati and Y. L. Chang, "Dissecting the regions around IceCube high-energy neutrinos: growing evidence for the blazar connection", <u>Mon. Not. Roy. Astron. Soc.</u> 497 (2020) 865 [2001.09355].
- [25] ANTARES, ICECUBE, PIERRE AUGER, LIGO SCIENTIFIC, VIRGO Collaboration, "Search for High-energy Neutrinos from Binary Neutron Star Merger GW170817 with ANTARES, IceCube, and the Pierre Auger Observatory", Astrophys. J. Lett. 850 (2017) L35 [1710.05839].
- [26] V. Berezinsky and G. Zatsepin, "Cosmic rays at ultrahigh-energies (neutrino?)", Phys. Lett. B 28 (1969) 423.
- [27] A. Coleman et al., "Ultra-High-Energy Cosmic Rays: The Intersection of the Cosmic and Energy Frontiers", 2205.05845.
- [28] R. M. Abraham et al., "Tau Neutrinos in the Next Decade: from GeV to EeV", 2203.05591.
- [29] K. Engel, J. Goodman, P. Huentemeyer, C. Kierans, T. R. Lewis, M. Negro et al., "The Future of Gamma-Ray Experiments in the MeV-EeV Range", in 2022 Snowmass Summer Study, 3, 2022, 2203.07360.
- [30] E. Berti, V. Cardoso, Z. Haiman, D. E. Holz, E. Mottola, S. Mukherjee et al., "Snowmass2021 Cosmic Frontier White Paper: Fundamental Physics and Beyond the Standard Model", in <u>2022 Snowmass Summer Study</u>, 3, 2022, 2203.06240.
- [31] K. Engel, T. Lewis, M. S. Muzio and T. M. Venters, "Advancing the Landscape of Multimessenger Science in the Next Decade", in 2022 Snowmass Summer Study, 3, 2022, 2203.10074.
- [32] L. Anchordoqui and F. Halzen, "IceHEP high energy physics at the south pole", <u>Annals Phys.</u> 321 (2006) 2660 [hep-ph/0510389].
- [33] L. A. Anchordoqui, V. Barger, I. Cholis, H. Goldberg, D. Hooper, A. Kusenko et al., "Cosmic Neutrino Pevatrons: A Brand New Pathway to Astronomy, Astrophysics, and Particle Physics", <u>JHEAp</u> 1-2 (2014) 1 [1312.6587].
- [34] ICECUBE Collaboration, "First observation of PeV-energy neutrinos with IceCube", <u>Phys. Rev. Lett.</u> 111 (2013) 021103 [1304.5356].
- [35] ICECUBE Collaboration, "Observation of High-Energy Astrophysical Neutrinos in Three Years of IceCube Data", Phys. Rev. Lett. **113** (2014) 101101 [1405.5303].
- [36] ICECUBE Collaboration, "Evidence for Astrophysical Muon Neutrinos from the Northern Sky with IceCube", Phys. Rev. Lett. 115 (2015) 081102 [1507.04005].
- [37] ICECUBE Collaboration, "Observation and Characterization of a Cosmic Muon Neutrino Flux from the Northern Hemisphere using six years of IceCube data", Astrophys. J. 833 (2016) 3 [1607.08006].
- [38] K. Murase, M. Ahlers and B. C. Lacki, "Testing the Hadronuclear Origin of PeV Neutrinos Observed with IceCube", Phys. Rev. D 88 (2013) 121301 [1306.3417].
- [39] B. Katz, E. Waxman, T. Thompson and A. Loeb, "The energy production rate density of cosmic rays in the local universe is ~ 10^{44−45} erg Mpc⁻³ yr⁻¹ at all particle energies", 1311.0287.
- [40] K. Murase and M. Fukugita, "Energetics of High-Energy Cosmic Radiations", <u>Phys. Rev. D</u> 99 (2019) 063012 [1806.04194].
- [41] R. Alves Batista et al., "Open Questions in Cosmic-Ray Research at Ultrahigh Energies", Front. Astron. Space Sci. 6 (2019) 23 [1903.06714].
- [42] HIRES Collaboration, "First observation of the Greisen-Zatsepin-Kuzmin suppression", <u>Phys. Rev. Lett.</u> 100 (2008) 101101 [astro-ph/0703099].

- [43] PIERRE AUGER Collaboration, "Observation of the suppression of the flux of cosmic rays above 4×10^{19} eV", Phys. Rev. Lett. **101** (2008) 061101 [0806.4302].
- [44] PIERRE AUGER Collaboration, "Highlights from the Pierre Auger Observatory (ICRC2019)", Pos ICRC2019 (2020) 004 [1909.10791].
- [45] TELESCOPE ARRAY Collaboration, "Highlights from the Telescope Array experiment (ICRC2019)", PoS ICRC2019 (2020) 013.
- [46] PIERRE AUGER Collaboration, "Features of the Energy Spectrum of Cosmic Rays above 2.5×10¹⁸ eV Using the Pierre Auger Observatory", Phys. Rev. Lett. **125** (2020) 121106 [2008.06488].
- [47] K. Greisen, "End to the cosmic ray spectrum?", Phys. Rev. Lett. 16 (1966) 748.
- [48] G. Zatsepin and V. Kuzmin, "Upper limit of the spectrum of cosmic rays", JETP Lett. 4 (1966) 78.
- [49] F. Stecker, "Diffuse Fluxes of Cosmic High-Energy Neutrinos", Astrophys. J. 228 (1979) 919.
- [50] C. T. Hill and D. N. Schramm, "Ultrahigh-Energy Cosmic Ray Neutrinos", Phys. Lett. B 131 (1983) 247.
- [51] S. Yoshida and M. Teshima, "Energy spectrum of ultrahigh-energy cosmic rays with extragalactic origin", Prog. Theor. Phys. 89 (1993) 833.
- [52] R. Engel, D. Seckel and T. Stanev, "Neutrinos from propagation of ultrahigh-energy protons", <u>Phys. Rev. D</u> 64 (2001) 093010 [astro-ph/0101216].
- [53] ICECUBE Collaboration, "A combined maximum-likelihood analysis of the high-energy astrophysical neutrino flux measured with IceCube", Astrophys. J. 809 (2015) 98 [1507.03991].
- [54] ICECUBE Collaboration, "Constraints on Galactic Neutrino Emission with Seven Years of IceCube Data", Astrophys. J. 849 (2017) 67 [1707.03416].
- [55] P. B. Denton, D. Marfatia and T. J. Weiler, "The Galactic Contribution to IceCube's Astrophysical Neutrino Flux", JCAP 08 (2017) 033 [1703.09721].
- [56] ICECUBE Collaboration, "Flavor Ratio of Astrophysical Neutrinos above 35 TeV in IceCube", <u>Phys. Rev. Lett.</u> 114 (2015) 171102 [1502.03376].
- [57] S. Palomares-Ruiz, A. C. Vincent and O. Mena, "Spectral analysis of the high-energy IceCube neutrinos", Phys. Rev. D 91 (2015) 103008 [1502.02649].
- [58] M. Bustamante, J. F. Beacom and W. Winter, "Theoretically palatable flavor combinations of astrophysical neutrinos", Phys. Rev. Lett. 115 (2015) 161302 [1506.02645].
- [59] M. Bustamante and M. Ahlers, "Inferring the flavor of high-energy astrophysical neutrinos at their sources", Phys. Rev. Lett. **122** (2019) 241101 [1901.10087].
- [60] Y. Huang, H. Li and B.-Q. Ma, "Consistent Lorentz violation features from near-TeV IceCube neutrinos", Phys. Rev. D 99 (2019) 123018 [1906.07329].
- [61] U. Jacob and T. Piran, "Neutrinos from gamma-ray bursts as a tool to explore quantum-gravity-induced Lorentz violation", Nature Phys. 3 (2007) 87 [hep-ph/0607145].
- [62] A. Addazi et al., "Quantum gravity phenomenology at the dawn of the multi-messenger era A review", 2111.05659.
- [63] D. Hooper, "Detecting MeV Gauge Bosons with High-Energy Neutrino Telescopes", <u>Phys. Rev. D</u> 75 (2007) 123001 [hep-ph/0701194].
- [64] K. Ioka and K. Murase, "IceCube PeV-EeV neutrinos and secret interactions of neutrinos", <u>PTEP</u> 2014 (2014) 061E01 [1404.2279].
- [65] K. C. Y. Ng and J. F. Beacom, "Cosmic neutrino cascades from secret neutrino interactions", <u>Phys. Rev. D</u> 90 (2014) 065035 [1404.2288].
- [66] M. Dhuria and V. Rentala, "PeV scale Supersymmetry breaking and the IceCube neutrino flux", <u>JHEP</u> 09 (2018) 004 [1712.07138].
- [67] M. Bustamante, C. Rosenstrøm, S. Shalgar and I. Tamborra, "Bounds on secret neutrino interactions from high-energy astrophysical neutrinos", Phys. Rev. D 101 (2020) 123024 [2001.04994].

- [68] C. Creque-Sarbinowski, J. Hyde and M. Kamionkowski, "Resonant neutrino self-interactions", <u>Phys. Rev. D</u> 103 (2021) 023527 [2005.05332].
- [69] I. Esteban, S. Pandey, V. Brdar and J. F. Beacom, "Probing secret interactions of astrophysical neutrinos in the high-statistics era", Phys. Rev. D 104 (2021) 123014 [2107.13568].
- [70] ICECUBE Collaboration, "The IceCube high-energy starting event sample: Description and flux characterization with 7.5 years of data", Phys. Rev. D **104** (2021) 022002 [2011.03545].
- [71] G. Barenboim and C. Quigg, "Neutrino observatories can characterize cosmic sources and neutrino properties", Phys. Rev. D 67 (2003) 073024 [hep-ph/0301220].
- [72] L. A. Anchordoqui, H. Goldberg, F. Halzen and T. J. Weiler, "Galactic point sources of TeV antineutrinos", Phys. Lett. B 593 (2004) 42 [astro-ph/0311002].
- [73] T. Kashti and E. Waxman, "Flavoring astrophysical neutrinos: Flavor ratios depend on energy", Phys.Rev.Lett. 95 (2005) 181101 [astro-ph/0507599].
- [74] M. Kachelriess and R. Tomas, "High energy neutrino yields from astrophysical sources I: Weakly magnetized sources", Phys. Rev. D 74 (2006) 063009 [astro-ph/0606406].
- [75] P. Lipari, M. Lusignoli and D. Meloni, "Flavor Composition and Energy Spectrum of Astrophysical Neutrinos", Phys. Rev. D 75 (2007) 123005 [0704.0718].
- [76] S. Hummer, M. Maltoni, W. Winter and C. Yaguna, "Energy dependent neutrino flavor ratios from cosmic accelerators on the Hillas plot", Astropart. Phys. 34 (2010) 205 [1007.0006].
- [77] S. Hummer, P. Baerwald and W. Winter, "Neutrino Emission from Gamma-Ray Burst Fireballs, Revised", Phys.Rev.Lett. 108 (2012) 231101 [1112.1076].
- [78] V. Barger, L. Fu, J. Learned, D. Marfatia, S. Pakvasa et al., "Glashow resonance as a window into cosmic neutrino sources", 1407.3255.
- [79] D. Biehl, A. Fedynitch, A. Palladino, T. J. Weiler and W. Winter, "Astrophysical Neutrino Production Diagnostics with the Glashow Resonance", JCAP 01 (2017) 033 [1611.07983].
- [80] L. Morejon, A. Fedynitch, D. Boncioli, D. Biehl and W. Winter, "Improved photomeson model for interactions of cosmic ray nuclei", JCAP 11 (2019) 007 [1904.07999].
- [81] ICECUBE Collaboration, "Measurement of Astrophysical Tau Neutrinos in IceCube's High-Energy Starting Events", 2011.03561.
- [82] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, "The fate of hints: updated global analysis of three-flavor neutrino oscillations", JHEP 09 (2020) 178 [2007.14792].
- [83] N. Song, S. W. Li, C. A. Argüelles, M. Bustamante and A. C. Vincent, "The Future of High-Energy Astrophysical Neutrino Flavor Measurements", JCAP 04 (2021) 054 [2012.12893].
- [84] P. Keranen, J. Maalampi, M. Myyrylainen and J. Riittinen, "Effects of sterile neutrinos on the ultrahigh-energy cosmic neutrino flux", Phys. Lett. B 574 (2003) 162 [hep-ph/0307041].
- [85] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa and T. J. Weiler, "Measuring flavor ratios of high-energy astrophysical neutrinos", Phys. Rev. D 68 (2003) 093005 [hep-ph/0307025].
- [86] S. Pakvasa, W. Rodejohann and T. J. Weiler, "Flavor Ratios of Astrophysical Neutrinos: Implications for Precision Measurements", JHEP 02 (2008) 005 [0711.4517].
- [87] M. Bustamante, A. Gago and J. Jones Pérez, "SUSY Renormalization Group Effects in Ultra High Energy Neutrinos", JHEP 05 (2011) 133 [1012.2728].
- [88] M. Bustamante, A. Gago and C. Peña Garay, "Energy-Independent New Physics in the Flavour Ratios of High-Energy Astrophysical Neutrinos", JHEP 04 (2010) 066 [1001.4878].
- [89] P. Mehta and W. Winter, "Interplay of energy dependent astrophysical neutrino flavor ratios and new physics effects", JCAP **03** (2011) 041 [1101.2673].
- [90] C. A. Argüelles, T. Katori and J. Salvadó, "New Physics in Astrophysical Neutrino Flavor", <u>Phys. Rev. Lett.</u> 115 (2015) 161303 [1506.02043].

- [91] I. M. Shoemaker and K. Murase, "Probing BSM Neutrino Physics with Flavor and Spectral Distortions: Prospects for Future High-Energy Neutrino Telescopes", Phys. Rev. D 93 (2016) 085004 [1512.07228].
- [92] M. González-García, M. Maltoni, I. Martínez-Soler and N. Song, "Non-standard neutrino interactions in the Earth and the flavor of astrophysical neutrinos", Astropart. Phys. 84 (2016) 15 [1605.08055].
- [93] R. W. Rasmussen, L. Lechner, M. Ackermann, M. Kowalski and W. Winter, "Astrophysical neutrinos flavored with Beyond the Standard Model physics", Phys. Rev. D 96 (2017) 083018 [1707.07684].
- [94] M. Ahlers, M. Bustamante and S. Mu, "Unitarity Bounds of Astrophysical Neutrinos", <u>Phys. Rev. D</u> 98 (2018) 123023 [1810.00893].
- [95] M. Ahlers, M. Bustamante and N. G. N. Willesen, "Flavors of astrophysical neutrinos with active-sterile mixing", JCAP 07 (2021) 029 [2009.01253].
- [96] S. W. Li, M. Bustamante and J. F. Beacom, "Echo Technique to Distinguish Flavors of Astrophysical Neutrinos", Phys. Rev. Lett. 122 (2019) 151101 [1606.06290].
- [97] K. Murase and E. Waxman, "Constraining High-Energy Cosmic Neutrino Sources: Implications and Prospects", Phys. Rev. D 94 (2016) 103006 [1607.01601].
- [98] M. Ahlers and F. Halzen, "Pinpointing Extragalactic Neutrino Sources in Light of Recent IceCube Observations", Phys.Rev. D90 (2014) 043005 [1406.2160].
- [99] K. Fang, K. Kotera, M. C. Miller, K. Murase and F. Oikonomou, "Identifying Ultrahigh-Energy Cosmic-Ray Accelerators with Future Ultrahigh-Energy Neutrino Detectors", JCAP **12** (2016) 017 [1609.08027].
- [100] I. Bartos, M. Ahrens, C. Finley and S. Marka, "Prospects of Establishing the Origin of Cosmic Neutrinos using Source Catalogs", Phys. Rev. D 96 (2017) 023003 [1611.03861].
- [101] I. Bartos, D. Veske, M. Kowalski, Z. Marka and S. Marka, "The IceCube Pie Chart: Relative Source Contributions to the Cosmic Neutrino Flux", Astrophys. J. 921 (2021) 45 [2105.03792].
- [102] A. Connolly, R. S. Thorne and D. Waters, "Calculation of High Energy Neutrino-Nucleon Cross Sections and Uncertainties Using the MSTW Parton Distribution Functions and Implications for Future Experiments", Phys. Rev. D 83 (2011) 113009 [1102.0691].
- [103] P. B. Denton and Y. Kini, "Ultra-High-Energy Tau Neutrino Cross Sections with GRAND and POEMMA", Phys. Rev. D 102 (2020) 123019 [2007.10334].
- [104] ICECUBE Collaboration, "Measuring total neutrino cross section with IceCube at intermediate energies (~100 GeV to a few TeV)", PoS ICRC2021 (2021) 1132 [2107.09764].
- [105] G.-y. Huang, S. Jana, M. Lindner and W. Rodejohann, "Probing new physics at future tau neutrino telescopes", JCAP 02 (2022) 038 [2112.09476].
- [106] V. B. Valera, M. Bustamante and C. Glaser, "The ultra-high-energy neutrino-nucleon cross section: measurement forecasts for an era of cosmic EeV-neutrino discovery", 2204.04237.
- [107] I. Esteban, S. Prohira and J. F. Beacom, "Detector Requirements for Model-Independent Measurements of Ultrahigh Energy Neutrino Cross Sections", 2205.09763.
- [108] Y. Bai, R. Lu and J. Salvadó, "Geometric Compatibility of IceCube TeV-PeV Neutrino Excess and its Galactic Dark Matter Origin", JHEP 01 (2016) 161 [1311.5864].
- [109] ANTARES Collaboration, "Search for dark matter towards the Galactic Centre with 11 years of ANTARES data", Phys. Lett. B 805 (2020) 135439 [1912.05296].
- [110] ICECUBE, ANTARES Collaboration, "Combined Search for Neutrinos from Dark Matter Annihilation in the Galactic Centre using ANTARES and IceCube", PoS ICRC2019 (2021) 522 [1908.07300].
- [111] ANTARES Collaboration, "Search of Dark Matter Annihilation in the Galactic Centre using the ANTARES Neutrino Telescope", JCAP 10 (2015) 068 [1505.04866].
- [112] M. H. Reno et al., "Neutrino constraints on long-lived heavy dark sector particle decays in the Earth", 2107.01159.
- [113] A. R. Zentner, "High-Energy Neutrinos From Dark Matter Particle Self-Capture Within the Sun", Phys. Rev. D 80 (2009) 063501 [0907.3448].

- [114] ICECUBE Collaboration, "First search for dark matter annihilations in the Earth with the IceCube Detector", Eur. Phys. J. C 77 (2017) 82 [1609.01492].
- [115] ICECUBE Collaboration, "Search for a Lorentz-violating sidereal signal with atmospheric neutrinos in IceCube", Phys. Rev. D 82 (2010) 112003 [1010.4096].
- [116] S. Hussain, D. Marfatia, D. McKay and D. Seckel, "Cross section dependence of event rates at neutrino telescopes", Phys. Rev. Lett. 97 (2006) 161101 [hep-ph/0606246].
- [117] FERMI-LAT Collaboration, "The spectrum of isotropic diffuse gamma-ray emission between 100 MeV and 820 GeV", Astrophys. J. 799 (2015) 86 [1410.3696].
- [118] ICECUBE Collaboration, "Characteristics of the diffuse astrophysical electron and tau neutrino flux with six years of IceCube high energy cascade data", Phys. Rev. Lett. 125 (2020) 121104 [2001.09520].
- [119] ICECUBE Collaboration, "Improved Characterization of the Astrophysical Muon-neutrino Flux with 9.5 Years of IceCube Data", Astrophys. J. 928 (2022) 50 [2111.10299].
- [120] PIERRE AUGER Collaboration, "The energy spectrum of cosmic rays beyond the turn-down around 10¹⁷ eV as measured with the surface detector of the Pierre Auger Observatory", <u>Eur. Phys. J. C</u> 81 (2021) 966 [2109.13400].
- [121] TELESCOPE ARRAY Collaboration, "Energy Spectrum Measured by the Telescope Array", <u>PoS</u> ICRC2019 (2020) 298.
- [122] ICECUBE Collaboration, "Measurement of the high-energy all-flavor neutrino-nucleon cross section with IceCube", Phys. Rev. D 104 (2020) 022001 [2011.03560].
- [123] ICECUBE Collaboration, "First Double Cascade Tau Neutrino Candidates in IceCube and a New Measurement of the Flavor Composition", PoS ICRC2019 (2020) 1015 [1908.05506].
- [124] KM3NET, ANTARES Collaboration, "Status and results from the ANTARES and KM3NeT-ARCA neutrino telescopes", PoS NuFact2019 (2020) 032.
- [125] KM3NET Collaboration, "Letter of intent for KM3NeT 2.0", J. Phys. G 43 (2016) 084001 [1601.07459].
- [126] P-ONE Collaboration, "The Pacific Ocean Neutrino Experiment", Nature Astron. 4 (2020) 913 [2005.09493].
- [127] BAIKAL-GVD Collaboration, "Neutrino Telescope in Lake Baikal: Present and Future", <u>PoS</u> ICRC2019 (2020) 1011 [1908.05427].
- [128] ICECUBE-GEN2 Collaboration, "IceCube-Gen2: the window to the extreme Universe", J. Phys. G 48 (2021) 060501 [2008.04323].
- [129] S. Prohira et al., "Observation of Radar Echoes From High-Energy Particle Cascades", <u>Phys. Rev. Lett.</u> 124 (2020) 091101 [1910.12830].
- [130] TAMBO Collaboration, "An Andean Deep-Valley Detector for High-Energy Tau Neutrinos", in Latin American Strategy Forum for Research Infrastructure, 2, 2020, 2002.06475.
- [131] A. N. Otte, A. M. Brown, A. D. Falcone, M. Mariotti and I. Taboada, "Trinity: An Air-Shower Imaging System for the Detection of Ultrahigh Energy Neutrinos", PoS ICRC2019 (2020) 976 [1907.08732].
- [132] M. Sasaki and G. W.-S. Hou, "Neutrino Telescope Array Letter of Intent: A Large Array of High Resolution Imaging Atmospheric Cherenkov and Fluorescence Detectors for Survey of Air-showers from Cosmic Tau Neutrinos in the PeV-EeV Energy Range", 1408.6244.
- [133] D. F. Fiorillo, G. Miele and O. Pisanti, "Tau Neutrinos with Cherenkov Telescope Array", 2007.13423.
- [134] J. H. Adams et al., "White paper on EUSO-SPB2", 1703.04513.
- [135] L. A. Anchordoqui et al., "Performance and science reach of the Probe of Extreme Multimessenger Astrophysics for ultrahigh-energy particles", Phys. Rev. D 101 (2020) 023012 [1907.03694].
- [136] K. Fang, K. Kotera, K. Murase and A. V. Olinto, "Testing the Newborn Pulsar Origin of Ultrahigh Energy Cosmic Rays with EeV Neutrinos", Phys. Rev. D 90 (2014) 103005 [1311.2044].
- [137] P. Padovani, M. Petropoulou, P. Giommi and E. Resconi, "A simplified view of blazars: the neutrino background", Mon. Not. Roy. Astron. Soc. 452 (2015) 1877 [1506.09135].

- [138] K. Fang and K. Murase, "Linking High-Energy Cosmic Particles by Black Hole Jets Embedded in Large-Scale Structures", Nature Phys. 14 (2018) 396 [1704.00015].
- [139] M. S. Muzio, M. Unger and G. R. Farrar, "Progress towards characterizing ultrahigh energy cosmic ray sources", Phys. Rev. D 100 (2019) 103008 [1906.06233].
- [140] X. Rodrigues, J. Heinze, A. Palladino, A. van Vliet and W. Winter, "Active Galactic Nuclei Jets as the Origin of Ultrahigh-Energy Cosmic Rays and Perspectives for the Detection of Astrophysical Source Neutrinos at EeV Energies", Phys. Rev. Lett. **126** (2021) 191101 [2003.08392].
- [141] M. S. Muzio, G. R. Farrar and M. Unger, "Probing the environments surrounding ultrahigh energy cosmic ray accelerators and their implications for astrophysical neutrinos", Phys. Rev. D 105 (2022) 023022 [2108.05512].
- [142] ARA Collaboration, "Ultra-high energy neutrino search with the Askaryan Radio Array", Pos ICRC2017 (2018) 966.
- [143] PIERRE AUGER Collaboration, "Searches for neutrino fluxes in the EeV regime with the Pierre Auger Observatory", PoS ICRC2017 (2018) 972.
- [144] ICECUBE Collaboration, "Differential limit on the extremely-high-energy cosmic neutrino flux in the presence of astrophysical background from nine years of IceCube data", Phys. Rev. D 98 (2018) 062003 [1807.01820].
- [145] ARIANNA Collaboration, "Performance of the ARIANNA pilot array, and implications for the next generation of UHE neutrino detectors", PoS ICRC2019 (2020) 980.
- [146] ANITA Collaboration, "Constraints on the ultrahigh-energy cosmic neutrino flux from the fourth flight of ANITA", Phys. Rev. D 99 (2019) 122001 [1902.04005].
- [147] A. Romero-Wolf and M. Ave, "Bayesian Inference Constraints on Astrophysical Production of Ultra-high Energy Cosmic Rays and Cosmogenic Neutrino Flux Predictions", JCAP 07 (2018) 025 [1712.07290].
- [148] R. Alves Batista, R. M. de Almeida, B. Lago and K. Kotera, "Cosmogenic photon and neutrino fluxes in the Auger era", JCAP 1901 (2019) 002 [1806.10879].
- [149] J. Heinze, A. Fedynitch, D. Boncioli and W. Winter, "A new view on Auger data and cosmogenic neutrinos in light of different nuclear disintegration and air-shower models", Astrophys. J. 873 (2019) 88 [1901.03338].
- [150] V. Berezinsky, E. Sabancilar and A. Vilenkin, "Extremely High Energy Neutrinos from Cosmic Strings", <u>Phys.</u> Rev. D 84 (2011) 085006 [1108.2509].
- [151] L. A. Anchordoqui, "Ultra-High-Energy Cosmic Rays", Phys. Rept. 801 (2019) 1 [1807.09645].
- [152] PIERRE AUGER Collaboration, "The Pierre Auger Observatory Upgrade Preliminary Design Report", 1604.03637.
- [153] GCOS Collaboration, "GCOS The Global Cosmic Ray Observatory", PoS 395 (2021) 027 [2203.01127].
- [154] GRAND Collaboration, "The Giant Radio Array for Neutrino Detection (GRAND): Science and Design", <u>Sci.</u> China Phys. Mech. Astron. 63 (2020) 219501 [1810.09994].
- [155] RNO-G Collaboration, "Design and Sensitivity of the Radio Neutrino Observatory in Greenland (RNO-G)", JINST 16 (2021) P03025 [2010.12279].
- [156] C. Deaconu et al., "Searches for Ultra-High Energy Neutrinos with ANITA", Pos ICRC2019 (2020) 867 [1908.00923].
- [157] S. Wissel et al., "Prospects for high-elevation radio detection of >100 PeV tau neutrinos", <u>JCAP</u> 11 (2020) 065 [2004.12718].
- [158] J. Nam et al., "High-elevation synoptic radio array for detection of upward moving air-showers, deployed in the Antarctic mountains", PoS ICRC2019 (2020) 967.
- [159] S. L. Glashow, "Resonant Scattering of Antineutrinos", Phys. Rev. 118 (1960) 316.
- [160] A. Cooper-Sarkar, P. Mertsch and S. Sarkar, "The high energy neutrino cross-section in the Standard Model and its uncertainty", JHEP 08 (2011) 042 [1106.3723].
- [161] V. Bertone, R. Gauld and J. Rojo, "Neutrino Telescopes as QCD Microscopes", <u>JHEP</u> 01 (2019) 217 [1808.02034].
- [162] L. A. Anchordoqui, C. García Canal and J. F. Soriano, "Probing strong dynamics with cosmic neutrinos", Phys. Rev. D 100 (2019) 103001 [1902.10134].
- [163] V. Berezinsky and A. Smirnov, "Astrophysical upper bounds on neutrino-nucleon cross-section at energy $E \ge 3 \times 10^{17}$ eV", Phys. Lett. B **48** (1974) 269.
- [164] F. Cornet, J. I. Illana and M. Masip, "TeV strings and the neutrino nucleon cross-section at ultrahigh-energies", Phys. Rev. Lett. 86 (2001) 4235 [hep-ph/0102065].
- [165] A. Kusenko and T. J. Weiler, "Neutrino cross-sections at high-energies and the future observations of ultrahigh-energy cosmic rays", Phys. Rev. Lett. 88 (2002) 161101 [hep-ph/0106071].
- [166] J. Álvarez-Muñiz, F. Halzen, T. Han and D. Hooper, "Phenomenology of high-energy neutrinos in low scale quantum gravity models", Phys. Rev. Lett. 88 (2002) 021301 [hep-ph/0107057].
- [167] L. A. Anchordoqui, J. L. Feng, H. Goldberg and A. D. Shapere, "Black holes from cosmic rays: Probes of extra dimensions and new limits on TeV scale gravity", Phys. Rev. D 65 (2002) 124027 [hep-ph/0112247].
- [168] M. Kowalski, A. Ringwald and H. Tu, "Black holes at neutrino telescopes", <u>Phys. Lett. B</u> 529 (2002) 1 [hep-ph/0201139].
- [169] J. Alvarez-Muñiz, J. L. Feng, F. Halzen, T. Han and D. Hooper, "Detecting microscopic black holes with neutrino telescopes", Phys. Rev. D 65 (2002) 124015 [hep-ph/0202081].
- [170] D. Hooper, "Measuring high-energy neutrino nucleon cross-sections with future neutrino telescopes", <u>Phys.</u> Rev. D 65 (2002) 097303 [hep-ph/0203239].
- [171] J. J. Friess, T. Han and D. Hooper, "TeV string state excitation via high-energy cosmic neutrinos", <u>Phys. Lett.</u> B 547 (2002) 31 [hep-ph/0204112].
- [172] L. A. Anchordoqui, J. L. Feng, H. Goldberg and A. D. Shapere, "Neutrino bounds on astrophysical sources and new physics", Phys. Rev. D 66 (2002) 103002 [hep-ph/0207139].
- [173] L. A. Anchordoqui, J. L. Feng and H. Goldberg, "Particle physics on ice: Constraints on neutrino interactions far above the weak scale", Phys. Rev. Lett. 96 (2006) 021101 [hep-ph/0504228].
- [174] L. A. Anchordoqui, A. M. Cooper-Sarkar, D. Hooper and S. Sarkar, "Probing low-x QCD with cosmic neutrinos at the Pierre Auger Observatory", Phys. Rev. D 74 (2006) 043008 [hep-ph/0605086].
- [175] E. Borriello, A. Cuoco, G. Mangano, G. Miele, S. Pastor, O. Pisanti et al., "Disentangling neutrino-nucleon cross section and high energy neutrino flux with a km³ neutrino telescope", <u>Phys. Rev. D</u> 77 (2008) 045019 [0711.0152].
- [176] S. Hussain, D. Marfatia and D. McKay, "Upward shower rates at neutrino telescopes directly determine the neutrino flux", Phys. Rev. D 77 (2008) 107304 [0711.4374].
- [177] V. Barger and W.-Y. Keung, "Superheavy Particle Origin of IceCube PeV Neutrino Events", <u>Phys. Lett. B</u> 727 (2013) 190 [1305.6907].
- [178] D. Marfatia, D. McKay and T. Weiler, "New physics with ultra-high-energy neutrinos", <u>Phys. Lett. B</u> 748 (2015) 113 [1502.06337].
- [179] J. Ellis, K. Sakurai and M. Spannowsky, "Search for Sphalerons: IceCube vs. LHC", <u>JHEP</u> 05 (2016) 085 [1603.06573].
- [180] S. R. Klein, <u>Probing high-energy interactions of atmospheric and astrophysical neutrinos</u>, p. 75. World Scientific, 2020. 1906.02221. 10.1142/9789813275027_0004.
- [181] K. J. Mack, N. Song and A. C. Vincent, "Signatures of microscopic black holes and extra dimensions at future neutrino telescopes", JHEP 04 (2020) 187 [1912.06656].
- [182] G. Amelino-Camelia, J. R. Ellis, N. Mavromatos, D. V. Nanopoulos and S. Sarkar, "Tests of quantum gravity from observations of gamma-ray bursts", Nature 393 (1998) 763 [astro-ph/9712103].
- [183] D. Hooper, D. Morgan and E. Winstanley, "Lorentz and CPT invariance violation in high-energy neutrinos", Phys. Rev. D 72 (2005) 065009 [hep-ph/0506091].

- [184] M. González-García, F. Halzen and M. Maltoni, "Physics reach of high-energy and high-statistics IceCube atmospheric neutrino data", Phys. Rev. D 71 (2005) 093010 [hep-ph/0502223].
- [185] L. A. Anchordoqui, H. Goldberg, M. González-García, F. Halzen, D. Hooper, S. Sarkar et al., "Probing Planck scale physics with IceCube", Phys. Rev. D 72 (2005) 065019 [hep-ph/0506168].
- [186] J. Bazo, M. Bustamante, A. Gago and O. Miranda, "High energy astrophysical neutrino flux and modified dispersion relations", Int. J. Mod. Phys. A 24 (2009) 5819 [0907.1979].
- [187] A. Kostelecky and M. Mewes, "Neutrinos with Lorentz-violating operators of arbitrary dimension", <u>Phys. Rev.</u> D 85 (2012) 096005 [1112.6395].
- [188] J. S. Diaz, A. Kostelecky and M. Mewes, "Testing Relativity with High-Energy Astrophysical Neutrinos", Phys. Rev. D 89 (2014) 043005 [1308.6344].
- [189] F. W. Stecker, S. T. Scully, S. Liberati and D. Mattingly, "Searching for Traces of Planck-Scale Physics with High Energy Neutrinos", Phys. Rev. D 91 (2015) 045009 [1411.5889].
- [190] F. W. Stecker and S. T. Scully, "Propagation of Superluminal PeV IceCube Neutrinos: A High Energy Spectral Cutoff or New Constraints on Lorentz Invariance Violation", Phys. Rev. D 90 (2014) 043012 [1404.7025].
- [191] G. Tomar, S. Mohanty and S. Pakvasa, "Lorentz Invariance Violation and IceCube Neutrino Events", <u>JHEP</u> 11 (2015) 022 [1507.03193].
- [192] J. Ellis, N. E. Mavromatos, A. S. Sakharov and E. K. Sarkisyan-Grinbaum, "Limits on Neutrino Lorentz Violation from Multimessenger Observations of TXS 0506+056", Phys. Lett. B 789 (2019) 352 [1807.05155].
- [193] R. Laha, "Constraints on neutrino speed, weak equivalence principle violation, Lorentz invariance violation, and dual lensing from the first high-energy astrophysical neutrino source TXS 0506+056", <u>Phys. Rev. D</u> 100 (2019) 103002 [1807.05621].
- [194] J. Lykken, O. Mena and S. Razzaque, "Ultrahigh-energy neutrino flux as a probe of large extra-dimensions", JCAP 12 (2007) 015 [0705.2029].
- [195] K. Blum, A. Hook and K. Murase, "High energy neutrino telescopes as a probe of the neutrino mass mechanism", 1408.3799.
- [196] W. Altmannshofer, C.-Y. Chen, P. Bhupal Dev and A. Soni, "Lepton flavor violating Z' explanation of the muon anomalous magnetic moment", Phys. Lett. B 762 (2016) 389 [1607.06832].
- [197] G. Barenboim, P. B. Denton and I. M. Oldengott, "Constraints on inflation with an extended neutrino sector", Phys. Rev. D 99 (2019) 083515 [1903.02036].
- [198] K. Murase and I. M. Shoemaker, "Neutrino Echoes from Multimessenger Transient Sources", <u>Phys. Rev. Lett.</u> 123 (2019) 241102 [1903.08607].
- [199] B. Feldstein, A. Kusenko, S. Matsumoto and T. T. Yanagida, "Neutrinos at IceCube from Heavy Decaying Dark Matter", Phys. Rev. D 88 (2013) 015004 [1303.7320].
- [200] A. Esmaili and P. D. Serpico, "Are IceCube neutrinos unveiling PeV-scale decaying dark matter?", <u>JCAP</u> 11 (2013) 054 [1308.1105].
- [201] T. Higaki, R. Kitano and R. Sato, "Neutrinoful Universe", JHEP 07 (2014) 044 [1405.0013].
- [202] C. Rott, K. Kohri and S. C. Park, "Superheavy dark matter and IceCube neutrino signals: Bounds on decaying dark matter", Phys. Rev. D 92 (2015) 023529 [1408.4575].
- [203] E. Dudas, Y. Mambrini and K. A. Olive, "Monochromatic neutrinos generated by dark matter and the seesaw mechanism", Phys. Rev. D 91 (2015) 075001 [1412.3459].
- [204] Y. Ema, R. Jinno and T. Moroi, "Cosmic-Ray Neutrinos from the Decay of Long-Lived Particle and the Recent IceCube Result", Phys. Lett. B 733 (2014) 120 [1312.3501].
- [205] J. Zavala, "Galactic PeV neutrinos from dark matter annihilation", Phys. Rev. D 89 (2014) 123516 [1404.2932].
- [206] C. S. Fong, H. Minakata, B. Panes and R. Zukanovich Funchal, "Possible Interpretations of IceCube High-Energy Neutrino Events", JHEP 02 (2015) 189 [1411.5318].

- [207] K. Murase, R. Laha, S. Ando and M. Ahlers, "Testing the Dark Matter Scenario for PeV Neutrinos Observed in IceCube", Phys. Rev. Lett. 115 (2015) 071301 [1503.04663].
- [208] L. A. Anchordoqui, V. Barger, H. Goldberg, X. Huang, D. Marfatia, L. H. M. da Silva et al., "IceCube neutrinos, decaying dark matter, and the Hubble constant", Phys. Rev. D 92 (2015) 061301 [1506.08788].
- [209] S. M. Boucenna, M. Chianese, G. Mangano, G. Miele, S. Morisi, O. Pisanti et al., "Decaying Leptophilic Dark Matter at IceCube", JCAP 12 (2015) 055 [1507.01000].
- [210] T. Cohen, K. Murase, N. L. Rodd, B. R. Safdi and Y. Soreq, "γ-ray Constraints on Decaying Dark Matter and Implications for IceCube", Phys. Rev. Lett. 119 (2017) 021102 [1612.05638].
- [211] P. S. B. Dev, D. Kazanas, R. Mohapatra, V. Teplitz and Y. Zhang, "Heavy right-handed neutrino dark matter and PeV neutrinos at IceCube", JCAP 08 (2016) 034 [1606.04517].
- [212] C. A. Argüelles, A. Kheirandish and A. C. Vincent, "Imaging Galactic Dark Matter with High-Energy Cosmic Neutrinos", Phys. Rev. Lett. 119 (2017) 201801 [1703.00451].
- [213] N. Hiroshima, R. Kitano, K. Kohri and K. Murase, "High-energy neutrinos from multibody decaying dark matter", Phys. Rev. D 97 (2018) 023006 [1705.04419].
- [214] M. Chianese, G. Miele and S. Morisi, "Interpreting IceCube 6-year HESE data as an evidence for hundred TeV decaying Dark Matter", Phys. Lett. B 773 (2017) 591 [1707.05241].
- [215] Y. Sui and P. Bhupal Dev, "A Combined Astrophysical and Dark Matter Interpretation of the IceCube HESE and Throughgoing Muon Events", JCAP 07 (2018) 020 [1804.04919].
- [216] C. A. Argüelles, A. Diaz, A. Kheirandish, A. Olivares-Del-Campo, I. Safa and A. C. Vincent, "Dark Matter Annihilation to Neutrinos: An Updated, Consistent & Compelling Compendium of Constraints", 1912.09486.
- [217] L. Anchordoqui, V. Barger, H. Goldberg and D. Marfatia, "Phase transition in the fine structure constant", Phys. Lett. B 660 (2008) 529 [0711.4055].
- [218] N. Klop and S. Ando, "Effects of a neutrino-dark energy coupling on oscillations of high-energy neutrinos", Phys. Rev. D 97 (2018) 063006 [1712.05413].
- [219] J. Arakawa, J. L. Feng, A. Ismail, F. Kling and M. Waterbury, "Neutrino Detection without Neutrino Detectors: Discovering Collider Neutrinos at FASER with Electronic Signals Only", 2206.09932.
- [220] CTEQ Collaboration, "Handbook of perturbative QCD: Version 1.0", Rev. Mod. Phys. 67 (1995) 157.
- [221] J. M. Conrad, M. H. Shaevitz and T. Bolton, "Precision measurements with high-energy neutrino beams", <u>Rev.</u> Mod. Phys. **70** (1998) 1341 [hep-ex/9707015].
- [222] C. Giunti and C. W. Kim, <u>Fundamentals of Neutrino Physics and Astrophysics</u>. Oxford Scholarship Onine, 2007.
- [223] J. A. Formaggio and G. P. Zeller, "From eV to EeV: Neutrino Cross Sections Across Energy Scales", <u>Rev.</u> Mod. Phys. 84 (2012) 1307 [1305.7513].
- [224] A. Cooper-Sarkar and S. Sarkar, "Predictions for high energy neutrino cross-sections from the ZEUS global PDF fits", JHEP 01 (2008) 075 [0710.5303].
- [225] M. Gluck, P. Jimenez-Delgado and E. Reya, "On the charged current neutrino-nucleon total cross section at high energies", Phys. Rev. D 81 (2010) 097501 [1003.3168].
- [226] V. P. Goncalves and P. Hepp, "A comparative study of the neutrino-nucleon cross section at ultra high energies", Phys. Rev. D 83 (2011) 014014 [1011.2718].
- [227] J. L. Albacete, J. I. Illana and A. Soto-Ontoso, "Neutrino-nucleon cross section at ultrahigh energy and its astrophysical implications", Phys. Rev. D 92 (2015) 014027 [1505.06583].
- [228] C. A. Argüelles, F. Halzen, L. Wille, M. Kroll and M. H. Reno, "High-energy behavior of photon, neutrino, and proton cross sections", Phys. Rev. D 92 (2015) 074040 [1504.06639].
- [229] M. A. G. Aivazis, F. I. Olness and W.-K. Tung, "Leptoproduction of heavy quarks. 1. General formalism and kinematics of charged current and neutral current production processes", <u>Phys. Rev. D</u> 50 (1994) 3085 [hep-ph/9312318].

- [230] M. A. G. Aivazis, J. C. Collins, F. I. Olness and W.-K. Tung, "Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies", <u>Phys. Rev. D</u> 50 (1994) 3102 [hep-ph/9312319].
- [231] R. S. Thorne and R. G. Roberts, "An Ordered analysis of heavy flavor production in deep inelastic scattering", Phys. Rev. D 57 (1998) 6871 [hep-ph/9709442].
- [232] M. Cacciari, M. Greco and P. Nason, "The P(T) spectrum in heavy flavor hadroproduction", <u>JHEP</u> 05 (1998) 007 [hep-ph/9803400].
- [233] Y. S. Jeong and M. H. Reno, "Quark mass effects in high energy neutrino nucleon scattering", <u>Phys. Rev. D</u> 81 (2010) 114012 [1001.4175].
- [234] A. Garcia, R. Gauld, A. Heijboer and J. Rojo, "Complete predictions for high-energy neutrino propagation in matter", JCAP 09 (2020) 025 [2004.04756].
- [235] T.-J. Hou et al., "New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC", Phys. Rev. D 103 (2021) 014013 [1912.10053].
- [236] L. A. Harland-Lang, A. D. Martin, P. Motylinski and R. S. Thorne, "Parton distributions in the LHC era: MMHT 2014 PDFs", Eur. Phys. J. C 75 (2015) 204 [1412.3989].
- [237] NNPDF Collaboration, "Parton distributions from high-precision collider data", <u>Eur. Phys. J. C</u> 77 (2017) 663 [1706.00428].
- [238] JAM Collaboration, "Strange quark suppression from a simultaneous Monte Carlo analysis of parton distributions and fragmentation functions", Phys. Rev. D 101 (2020) 074020 [1905.03788].
- [239] M. Bonvini, S. Marzani and T. Peraro, "Small-x resummation from HELL", <u>Eur. Phys. J. C</u> 76 (2016) 597 [1607.02153].
- [240] M. Bonvini, S. Marzani and C. Muselli, "Towards parton distribution functions with small-x resummation: HELL 2.0", JHEP 12 (2017) 117 [1708.07510].
- [241] R. D. Ball, V. Bertone, M. Bonvini, S. Marzani, J. Rojo and L. Rottoli, "Parton distributions with small-x resummation: evidence for BFKL dynamics in HERA data", Eur. Phys. J. C 78 (2018) 321 [1710.05935].
- [242] S. R. Klein and A. Connolly, "Neutrino Absorption in the Earth, Neutrino Cross-Sections, and New Physics", in Community Summer Study 2013: Snowmass on the Mississippi, 4, 2013, 1304.4891.
- [243] D. Seckel, "Neutrino photon reactions in astrophysics and cosmology", <u>Phys. Rev. Lett.</u> 80 (1998) 900 [hep-ph/9709290].
- [244] I. Alikhanov, "Hidden Glashow resonance in neutrino-nucleus collisions", <u>Phys. Lett. B</u> 756 (2016) 247 [1503.08817].
- [245] R. Gauld, "Precise predictions for multi-TeV and PeV energy neutrino scattering rates", <u>Phys. Rev. D</u> 100 (2019) 091301 [1905.03792].
- [246] B. Zhou and J. F. Beacom, "Neutrino-nucleus cross sections for W-boson and trident production", <u>Phys. Rev.</u> D 101 (2020) 036011 [1910.08090].
- [247] B. Zhou and J. F. Beacom, "W-boson and trident production in TeV-PeV neutrino observatories", <u>Phys. Rev.</u> D 101 (2020) 036010 [1910.10720].
- [248] A. G. Soto, P. Zhelnin, I. Safa and C. A. Argüelles, "Tau Appearance from High-Energy Neutrino Interactions", 2112.06937.
- [249] L. Frankfurt, V. Guzey and M. Strikman, "Leading Twist Nuclear Shadowing Phenomena in Hard Processes with Nuclei", Phys. Rept. 512 (2012) 255 [1106.2091].
- [250] K. Kovarik et al., "nCTEQ15 Global analysis of nuclear parton distributions with uncertainties in the CTEQ framework", Phys. Rev. D 93 (2016) 085037 [1509.00792].
- [251] K. J. Eskola, P. Paakkinen, H. Paukkunen and C. A. Salgado, "EPPS16: Nuclear parton distributions with LHC data", Eur. Phys. J. C 77 (2017) 163 [1612.05741].
- [252] NNPDF Collaboration, "Nuclear parton distributions from lepton-nucleus scattering and the impact of an electron-ion collider", Eur. Phys. J. C 79 (2019) 471 [1904.00018].

- [253] M. Walt, I. Helenius and W. Vogelsang, "Open-source QCD analysis of nuclear parton distribution functions at NLO and NNLO", Phys. Rev. D 100 (2019) 096015 [1908.03355].
- [254] S. R. Klein, S. A. Robertson and R. Vogt, "Nuclear effects in high-energy neutrino interactions", Phys. Rev. C 102 (2020) 015808 [2001.03677].
- [255] R. A. Khalek, J. J. Ethier, E. R. Nocera and J. Rojo, "Self-consistent determination of proton and nuclear PDFs at the Electron Ion Collider", Phys. Rev. D 103 (2021) 096005 [2102.00018].
- [256] L. A. Anchordoqui, C. A. García Canal, H. Goldberg, D. Gomez Dumm and F. Halzen, "Probing leptoquark production at IceCube", Phys. Rev. D 74 (2006) 125021 [hep-ph/0609214].
- [257] S. T. Petcov, "The Processes $\mu \to e + \gamma, \mu \to e + \overline{e}, \nu' \to \nu + \gamma$ in the Weinberg-Salam Model with Neutrino Mixing", Sov. J. Nucl. Phys. **25** (1977) 340.
- [258] W. J. Marciano and A. I. Sanda, "Exotic Decays of the Muon and Heavy Leptons in Gauge Theories", <u>Phys.</u> Lett. B 67 (1977) 303.
- [259] S. Hannestad and G. Raffelt, "Constraining invisible neutrino decays with the cosmic microwave background", Phys. Rev. D 72 (2005) 103514 [hep-ph/0509278].
- [260] M. Escudero and M. Fairbairn, "Cosmological Constraints on Invisible Neutrino Decays Revisited", <u>Phys. Rev.</u> D 100 (2019) 103531 [1907.05425].
- [261] Z. Chacko, A. Dev, P. Du, V. Poulin and Y. Tsai, "Determining the Neutrino Lifetime from Cosmology", <u>Phys.</u> Rev. D 103 (2021) 043519 [2002.08401].
- [262] M. Escudero, J. Lopez-Pavon, N. Rius and S. Sandner, "Relaxing Cosmological Neutrino Mass Bounds with Unstable Neutrinos", JHEP 12 (2020) 119 [2007.04994].
- [263] G. Barenboim, J. Z. Chen, S. Hannestad, I. M. Oldengott, T. Tram and Y. Y. Y. Wong, "Invisible neutrino decay in precision cosmology", JCAP 03 (2021) 087 [2011.01502].
- [264] J. Z. Chen, I. M. Oldengott, G. Pierobon and Y. Y. Wong, "Weaker yet again: mass spectrum-consistent cosmological constraints on the neutrino lifetime", 2203.09075.
- [265] S. Ando, "Decaying neutrinos and implications from the supernova relic neutrino observation", Phys. Lett. B 570 (2003) 11 [hep-ph/0307169].
- [266] G. L. Fogli, E. Lisi, A. Mirizzi and D. Montanino, "Three generation flavor transitions and decays of supernova relic neutrinos", Phys. Rev. D 70 (2004) 013001 [hep-ph/0401227].
- [267] KAMIOKANDE-II Collaboration, "Observation of a Neutrino Burst from the Supernova SN 1987a", <u>Phys. Rev.</u> Lett. 58 (1987) 1490.
- [268] P. B. Denton and I. Tamborra, "Invisible Neutrino Decay Could Resolve IceCube's Track and Cascade Tension", Phys. Rev. Lett. **121** (2018) 121802 [1805.05950].
- [269] A. Abdullahi and P. B. Denton, "Visible Decay of Astrophysical Neutrinos at IceCube", <u>Phys. Rev. D</u> 102 (2020) 023018 [2005.07200].
- [270] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa and T. J. Weiler, "Decay of High-Energy Astrophysical Neutrinos", Phys. Rev. Lett. 90 (2003) 181301 [hep-ph/0211305].
- [271] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa and T. J. Weiler, "Sensitivity to theta(13) and delta in the decaying astrophysical neutrino scenario", Phys. Rev. D 69 (2004) 017303 [hep-ph/0309267].
- [272] P. Baerwald, M. Bustamante and W. Winter, "Neutrino Decays over Cosmological Distances and the Implications for Neutrino Telescopes", JCAP 10 (2012) 020 [1208.4600].
- [273] M. Bustamante, J. F. Beacom and K. Murase, "Testing decay of astrophysical neutrinos with incomplete information", Phys. Rev. D 95 (2017) 063013 [1610.02096].
- [274] C. Boehm, Y. Farzan, T. Hambye, S. Palomares-Ruiz and S. Pascoli, "Is it possible to explain neutrino masses with scalar dark matter?", Phys. Rev. D77 (2008) 043516 [hep-ph/0612228].
- [275] J. F. Beacom, N. F. Bell and G. D. Mack, "General Upper Bound on the Dark Matter Total Annihilation Cross Section", Phys.Rev.Lett. 99 (2007) 231301 [astro-ph/0608090].

- [276] H. Yuksel, S. Horiuchi, J. F. Beacom and S. Ando, "Neutrino Constraints on the Dark Matter Total Annihilation Cross Section", Phys.Rev. D76 (2007) 123506 [0707.0196].
- [277] K. Murase and J. F. Beacom, "Constraining Very Heavy Dark Matter Using Diffuse Backgrounds of Neutrinos and Cascaded Gamma Rays", JCAP 10 (2012) 043 [1206.2595].
- [278] S. Yoshida, H.-y. Dai, C. C. Jui and P. Sommers, "Extremely high-energy neutrinos and their detection", Astrophys. J. 479 (1997) 547 [astro-ph/9608186].
- [279] T. Araki, F. Kaneko, T. Ota, J. Sato and T. Shimomura, "MeV scale leptonic force for cosmic neutrino spectrum and muon anomalous magnetic moment", Phys. Rev. D 93 (2016) 013014 [1508.07471].
- [280] L. G. van den Aarssen, T. Bringmann and C. Pfrommer, "Is dark matter with long-range interactions a solution to all small-scale problems of \Lambda CDM cosmology?", Phys. Rev. Lett. **109** (2012) 231301 [1205.5809].
- [281] S. Tulin and H.-B. Yu, "Dark Matter Self-interactions and Small Scale Structure", Phys. Rept. 730 (2018) 1 [1705.02358].
- [282] F.-Y. Cyr-Racine and K. Sigurdson, "Limits on Neutrino-Neutrino Scattering in the Early Universe", <u>Phys.</u> <u>Rev. D</u> 90 (2014) 123533 [1306.1536].
- [283] C. D. Kreisch, F.-Y. Cyr-Racine and O. Doré, "Neutrino puzzle: Anomalies, interactions, and cosmological tensions", Phys. Rev. D 101 (2020) 123505 [1902.00534].
- [284] N. Blinov, K. J. Kelly, G. Z. Krnjaic and S. D. McDermott, "Constraining the Self-Interacting Neutrino Interpretation of the Hubble Tension", Phys. Rev. Lett. **123** (2019) 191102 [1905.02727].
- [285] J. A. Carpio, K. Murase, I. M. Shoemaker and Z. Tabrizi, "High-energy cosmic neutrinos as a probe of the vector mediator scenario in light of the muon g 2 anomaly and Hubble tension", 2104.15136.
- [286] M. Ibe and K. Kaneta, "Cosmic neutrino background absorption line in the neutrino spectrum at IceCube", Phys. Rev. D 90 (2014) 053011 [1407.2848].
- [287] A. DiFranzo and D. Hooper, "Searching for MeV-Scale Gauge Bosons with IceCube", <u>Phys. Rev. D</u> 92 (2015) 095007 [1507.03015].
- [288] J. F. Cherry, A. Friedland and I. M. Shoemaker, "Short-baseline neutrino oscillations, Planck, and IceCube", 1605.06506.
- [289] K. J. Kelly and P. A. Machado, "Multimessenger Astronomy and New Neutrino Physics", <u>JCAP</u> 10 (2018) 048 [1808.02889].
- [290] J. M. Berryman, A. De Gouvêa, K. J. Kelly and Y. Zhang, "Lepton-Number-Charged Scalars and Neutrino Beamstrahlung", Phys. Rev. D 97 (2018) 075030 [1802.00009].
- [291] K. J. Kelly and Y. Zhang, "Mononeutrino at DUNE: New Signals from Neutrinophilic Thermal Dark Matter", Phys. Rev. D 99 (2019) 055034 [1901.01259].
- [292] KM3NET Collaboration, "Sensitivity of the KM3NeT/ARCA neutrino telescope to point-like neutrino sources", Astropart. Phys. 111 (2019) 100 [1810.08499].
- [293] T. K. Gaisser, "Spectrum of cosmic-ray nucleons, kaon production, and the atmospheric muon charge ratio", Astropart. Phys. 35 (2012) 801 [1111.6675].
- [294] Y. S. Jeong, W. Bai, M. Diwan, M. V. Garzelli, F. K. Kumar and M. H. Reno, "Neutrinos from charm: forward production at the LHC and in the atmosphere", PoS ICRC2021 (2021) 1218 [2107.01178].
- [295] A. Bhattacharya, R. Enberg, M. H. Reno, I. Sarcevic and A. Stasto, "Perturbative charm production and the prompt atmospheric neutrino flux in light of RHIC and LHC", JHEP 06 (2015) 110 [1502.01076].
- [296] R. Gauld, J. Rojo, L. Rottoli, S. Sarkar and J. Talbert, "The prompt atmospheric neutrino flux in the light of LHCb", JHEP 02 (2016) 130 [1511.06346].
- [297] A. Bhattacharya, R. Enberg, Y. S. Jeong, C. S. Kim, M. H. Reno, I. Sarcevic et al., "Prompt atmospheric neutrino fluxes: perturbative QCD models and nuclear effects", JHEP 11 (2016) 167 [1607.00193].
- [298] PROSA Collaboration, "Improved constraints on parton distributions using LHCb, ALICE and HERA heavy-flavour measurements and implications for the predictions for prompt atmospheric-neutrino fluxes", JHEP 04 (2020) 118 [1911.13164].

- [299] LHCB Collaboration, "Measurements of prompt charm production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV", JHEP **03** (2016) 159 [1510.01707].
- [300] FASER Collaboration, "Technical Proposal: FASERnu", 2001.03073.
- [301] SHIP Collaboration, "SND@LHC", 2002.08722.
- [302] FASER Collaboration, "First neutrino interaction candidates at the LHC", <u>Phys. Rev. D</u> 104 (2021) L091101 [2105.06197].
- [303] L. A. Anchordoqui et al., "The Forward Physics Facility: Sites, experiments, and physics potential", <u>Phys.</u> Rept. **968** (2022) 1 [2109.10905].
- [304] R. Foot, X. G. He, H. Lew and R. R. Volkas, "Model for a light Z-prime boson", <u>Phys. Rev. D</u> 50 (1994) 4571 [hep-ph/9401250].
- [305] X.-G. He, G. C. Joshi, H. Lew and R. R. Volkas, "Simplest Z-prime model", Phys. Rev. D 44 (1991) 2118.
- [306] R. Foot, "New Physics From Electric Charge Quantization?", Mod. Phys. Lett. A 6 (1991) 527.
- [307] M. Bustamante and S. K. Agarwalla, "Universe's Worth of Electrons to Probe Long-Range Interactions of High-Energy Astrophysical Neutrinos", Phys. Rev. Lett. **122** (2019) 061103 [1808.02042].
- [308] J. Heeck and W. Rodejohann, "Gauged $L_{\mu} L_{\tau}$ and different Muon Neutrino and Anti-Neutrino Oscillations: MINOS and beyond", J. Phys. G **38** (2011) 085005 [1007.2655].
- [309] ICECUBE Collaboration, "The IceCube Collaboration Contributions to the 37th International Cosmic Ray Conference (ICRC2021)", 2107.06966.
- [310] BAIKAL-GVD Collaboration, "The Baikal-GVD neutrino telescope: First results of multi-messenger study", PoS ICRC2019 (2020) 1013 [1908.05450].
- [311] ANITA Collaboration, "Characteristics of Four Upward-pointing Cosmic-ray-like Events Observed with ANITA", Phys. Rev. Lett. 117 (2016) 071101 [1603.05218].
- [312] ANITA Collaboration, "Observation of an Unusual Upward-going Cosmic-ray-like Event in the Third Flight of ANITA", Phys. Rev. Lett. **121** (2018) 161102 [1803.05088].
- [313] PIERRE AUGER Collaboration, "Improved limit to the diffuse flux of ultrahigh energy neutrinos from the Pierre Auger Observatory", Phys. Rev. D **91** (2015) 092008 [1504.05397].
- [314] PIERRE AUGER Collaboration, "A tau scenario application to a search for upward-going showers with the Fluorescence Detector of the Pierre Auger Observatory", PoS ICRC2021 (2021) 1145.
- [315] PIERRE AUGER Collaboration, "Search for upward-going showers with the Fluorescence Detector of the Pierre Auger Observatory", PoS ICRC2021 (2021) 1140.
- [316] ICECUBE Collaboration, "Constraints on Ultrahigh-Energy Cosmic-Ray Sources from a Search for Neutrinos above 10 PeV with IceCube", Phys. Rev. Lett. 117 (2016) 241101 [1607.05886].
- [317] A. Romero-Wolf et al., "Comprehensive analysis of anomalous ANITA events disfavors a diffuse tau-neutrino flux origin", Phys. Rev. D 99 (2019) 063011 [1811.07261].
- [318] ICECUBE Collaboration, "A search for IceCube events in the direction of ANITA neutrino candidates", Astrophys. J. 892 (2020) 52 [2001.01737].
- [319] ANITA Collaboration, "Analysis of a tau neutrino origin for the near-horizon air shower events observed by the fourth flight of the Antarctic Impulsive Transient Antenna", Phys. Rev. D 105 (2022) 042001 [2112.07069].
- [320] ANITA Collaboration, "Comprehensive estimate of the sensitivity of ANITA to tau neutrinos", PoS ICRC2019 (2021) 1034.
- [321] K. D. de Vries and S. Prohira, "Coherent transition radiation from the geomagnetically-induced current in cosmic-ray air showers: Implications for the anomalous events observed by ANITA", <u>Phys. Rev. Lett.</u> 123 (2019) 091102 [1903.08750].
- [322] P. Motloch, J. Alvarez-Muñiz, P. Privitera and E. Zas, "Can transition radiation explain the ANITA event 3985267?", Phys. Rev. D 95 (2017) 043004 [1606.07059].

- [323] I. M. Shoemaker, A. Kusenko, P. K. Munneke, A. Romero-Wolf, D. M. Schroeder and M. J. Siegert, "Reflections On the Anomalous ANITA Events: The Antarctic Subsurface as a Possible Explanation", <u>Ann.</u> Glaciology (2020) 1 [1905.02846].
- [324] D. Smith et al., "Experimental tests of sub-surface reflectors as an explanation for the ANITA anomalous events", JCAP 04 (2021) 016 [2009.13010].
- [325] B. Chauhan and S. Mohanty, "Leptoquark solution for both the flavor and ANITA anomalies", <u>Phys. Rev. D</u> 99 (2019) 095018 [1812.00919].
- [326] J. H. Collins, P. Bhupal Dev and Y. Sui, "R-parity Violating Supersymmetric Explanation of the Anomalous Events at ANITA", Phys. Rev. D 99 (2019) 043009 [1810.08479].
- [327] D. B. Fox, S. Sigurdsson, S. Shandera, P. Mészáros, K. Murase, M. Mostafá et al., "The ANITA Anomalous Events as Signatures of a Beyond Standard Model Particle, and Supporting Observations from IceCube", 1809.09615.
- [328] A. Esmaili and Y. Farzan, "Explaining the ANITA events by a $L_e L_{\tau}$ gauge model", JCAP 12 (2019) 017 [1909.07995].
- [329] I. Esteban, J. López-Pavón, I. Martínez-Soler and J. Salvadó, "Looking at the axionic dark sector with ANITA", Eur. Phys. J. C 80 (2020) 259 [1905.10372].
- [330] D. Borah, A. Dasgupta, U. K. Dey and G. Tomar, "Connecting ANITA anomalous events to a nonthermal dark matter scenario", Phys. Rev. D 101 (2020) 075039 [1907.02740].
- [331] L. Heurtier, Y. Mambrini and M. Pierre, "Dark matter interpretation of the ANITA anomalous events", <u>Phys.</u> Rev. D 99 (2019) 095014 [1902.04584].
- [332] L. Heurtier, D. Kim, J.-C. Park and S. Shin, "Explaining the ANITA Anomaly with Inelastic Boosted Dark Matter", Phys. Rev. D 100 (2019) 055004 [1905.13223].
- [333] D. Hooper, S. Wegsman, C. Deaconu and A. Vieregg, "Superheavy dark matter and ANITA's anomalous events", Phys. Rev. D 100 (2019) 043019 [1904.12865].
- [334] L. O. Drury, "An introduction to the theory of diffusive shock acceleration of energetic particles in tenuous plasmas", Rept. Prog. Phys. 46 (1983) 973.
- [335] F. M. Rieger, "An Introduction to Particle Acceleration in Shearing Flows", Galaxies 7 (2019) 78 [1909.07237].
- [336] R. Mbarek and D. Caprioli, "Espresso and Stochastic Acceleration of Ultra-high-energy Cosmic Rays in Relativistic Jets", Astrophys. J. 921 (2021) 85 [2105.05262].
- [337] F. Guo, Y.-H. Liu, X. Li, H. Li, W. Daughton and P. Kilian, "Recent progress on particle acceleration and reconnection physics during magnetic reconnection in the magnetically-dominated relativistic regime", <u>Phys.</u> Plasmas 27 (2020) 080501 [2006.15288].
- [338] S. R. Kelner, F. A. Aharonian and V. V. Bugayov, "Energy spectra of gamma-rays, electrons and neutrinos produced at proton-proton interactions in the very high energy regime", <u>Phys. Rev.</u> D74 (2006) 034018 [astro-ph/0606058].
- [339] S. R. Kelner and F. A. Aharonian, "Energy spectra of gamma-rays, electrons and neutrinos produced at interactions of relativistic protons with low energy radiation", Phys. Rev. D78 (2008) 034013 [0803.0688].
- [340] W. Winter, "Photohadronic Origin of the TeV-PeV Neutrinos Observed in IceCube", <u>Phys.Rev.</u> D88 (2013) 083007 [1307.2793].
- [341] M. Bustamante and I. Tamborra, "Using high-energy neutrinos as cosmic magnetometers", <u>Phys. Rev. D</u> 102 (2020) 123008 [2009.01306].
- [342] V. S. Berezinsky, "Proc. of Neutrinos-1977, Elbros.", .
- [343] D. Eichler, "High-energy neutrino astronomy: A probe of galactic nuclei?", Astrophys. J. 232 (1979) 106.
- [344] F. W. Stecker, C. Done, M. H. Salamon and P. Sommers, "High-energy neutrinos from active galactic nuclei", Phys.Rev.Lett. 66 (1991) 2697.
- [345] P. Mészáros and E. Waxman, "TeV neutrinos from successful and choked gamma-ray bursts", <u>Phys.Rev.Lett.</u> 87 (2001) 171102 [astro-ph/0103275].

- [346] K. Murase, T. A. Thompson, B. C. Lacki and J. F. Beacom, "New Class of High-Energy Transients from Crashes of Supernova Ejecta with Massive Circumstellar Material Shells", <u>Phys.Rev.</u> D84 (2011) 043003 [1012.2834].
- [347] M. Hoshino, "Particle Acceleration during Magnetorotational Instability in a Collisionless Accretion Disk", Astrophys. J. 773 (2013) 118 [1306.6720].
- [348] D. Ball, L. Sironi and F. Ozel, "Electron and Proton Acceleration in Trans-Relativistic Magnetic Reconnection: Dependence on Plasma Beta and Magnetization", Astrophys. J. 862 (2018) 80 [1803.05556].
- [349] S. S. Kimura, K. Tomida and K. Murase, "Acceleration and Escape Processes of High-energy Particles in Turbulence inside Hot Accretion Flows", Mon. Not. Roy. Astron. Soc. 485 (2019) 163 [1812.03901].
- [350] A. Levinson and B. Cerutti, "Particle-in-cell simulations of pair discharges in a starved magnetosphere of a Kerr black hole", Astron. Astrophys. 616 (2018) A184 [1803.04427].
- [351] S. Kisaka, A. Levinson and K. Toma, "Comprehensive analysis of magnetospheric gaps around Kerr black holes using 1D GRPIC simulations", Astrophys. J. 902 (2020) 80 [2007.02838].
- [352] S. S. Kimura, K. Murase and K. Toma, "Neutrino and Cosmic-Ray Emission and Cumulative Background from Radiatively Inefficient Accretion Flows in Low-Luminosity Active Galactic Nuclei", <u>Astrophys.J.</u> 806 (2015) 159 [1411.3588].
- [353] O. Kalashev, D. Semikoz and I. Tkachev, "Neutrinos in IceCube from active galactic nuclei", J.Exp.Theor.Phys. 120 (2015) 541.
- [354] K. Murase, S. S. Kimura and P. Meszaros, "Hidden Cores of Active Galactic Nuclei as the Origin of Medium-Energy Neutrinos: Critical Tests with the MeV Gamma-Ray Connection", <u>Phys. Rev. Lett.</u> 125 (2020) 011101 [1904.04226].
- [355] S. S. Kimura, K. Murase and P. Mészáros, "Soft gamma rays from low accreting supermassive black holes and connection to energetic neutrinos", Nature Commun. 12 (2021) 5615 [2005.01934].
- [356] L. Dai and K. Fang, "Can tidal disruption events produce the IceCube neutrinos?", <u>Mon. Not. Roy. Astron.</u> Soc. 469 (2017) 1354 [1612.00011].
- [357] N. Senno, K. Murase and P. Meszaros, "High-energy Neutrino Flares from X-Ray Bright and Dark Tidal Disruption Events", Astrophys. J. 838 (2017) 3 [1612.00918].
- [358] C. Lunardini and W. Winter, "High Energy Neutrinos from the Tidal Disruption of Stars", <u>Phys. Rev. D</u> 95 (2017) 123001 [1612.03160].
- [359] K. Murase and K. Ioka, "TeV-PeV Neutrinos from Low-Power Gamma-Ray Burst Jets inside Stars", Phys.Rev.Lett. 111 (2013) 121102 [1306.2274].
- [360] N. Senno, K. Murase and P. Mészáros, "Choked Jets and Low-Luminosity Gamma-Ray Bursts as Hidden Neutrino Sources", Phys. Rev. D93 (2016) 083003 [1512.08513].
- [361] I. Tamborra and S. Ando, "Inspecting the supernova-gamma-ray-burst connection with high-energy neutrinos", Phys. Rev. D93 (2016) 053010 [1512.01559].
- [362] P. B. Denton and I. Tamborra, "The Bright and Choked Gamma-Ray Burst Contribution to the IceCube and ANTARES Low-Energy Excess", JCAP 04 (2018) 058 [1802.10098].
- [363] K. Murase, "New Prospects for Detecting High-Energy Neutrinos from Nearby Supernovae", <u>Phys. Rev.</u> D97 (2018) 081301 [1705.04750].
- [364] M. Petropoulou, S. Coenders, G. Vasilopoulos, A. Kamble and L. Sironi, "Point-source and diffuse high-energy neutrino emission from Type IIn supernovae", Mon. Not. Roy. Astron. Soc. 470 (2017) 1881 [1705.06752].
- [365] S. Pakvasa, "Neutrino Flavor Goniometry by High Energy Astrophysical Beams", <u>Mod. Phys. Lett. A</u> 23 (2008) 1313 [0803.1701].
- [366] J. G. Learned and S. Pakvasa, "Detecting tau-neutrino oscillations at PeV energies", <u>Astropart. Phys.</u> 3 (1995) 267 [hep-ph/9405296].
- [367] S. R. Klein, R. E. Mikkelsen and J. Becker Tjus, "Muon Acceleration in Cosmic-ray Sources", <u>Astrophys. J.</u> 779 (2013) 106 [1208.2056].

- [368] O. Mena, I. Mocioiu and S. Razzaque, "Oscillation effects on high-energy neutrino fluxes from astrophysical hidden sources", Phys. Rev. D 75 (2007) 063003 [astro-ph/0612325].
- [369] S. Razzaque and A. Y. Smirnov, "Flavor conversion of cosmic neutrinos from hidden jets", JHEP 03 (2010) 031 [0912.4028].
- [370] S. Sahu and B. Zhang, "Effect of Resonant Neutrino Oscillation on TeV Neutrino Flavor Ratio from Choked GRBs", Res. Astron. Astrophys. 10 (2010) 943 [1007.4582].
- [371] D. Xiao and Z. G. Dai, "TeV-PeV Neutrino Oscillation of Low-luminosity Gamma-ray Bursts", <u>Astrophys. J.</u> 805 (2015) 137 [1504.01603].
- [372] J. Carpio and K. Murase, "Oscillation of high-energy neutrinos from choked jets in stellar and merger ejecta", Phys. Rev. D 101 (2020) 123002 [2002.10575].
- [373] ICECUBE Collaboration, "Flavor Ratio of Astrophysical Neutrinos above 35 TeV in IceCube", Phys. Rev. Lett. 114 (2015) 171102 [1502.03376].
- [374] O. Mena, S. Palomares-Ruiz and A. C. Vincent, "Flavor Composition of the High-Energy Neutrino Events in IceCube", Phys. Rev. Lett. 113 (2014) 091103 [1404.0017].
- [375] A. C. Vincent, S. Palomares-Ruiz and O. Mena, "Analysis of the 4-year IceCube high-energy starting events", Phys. Rev. D 94 (2016) 023009 [1605.01556].
- [376] K. Murase, D. Guetta and M. Ahlers, "Hidden Cosmic-Ray Accelerators as an Origin of TeV-PeV Cosmic Neutrinos", Phys. Rev. Lett. 116 (2016) 071101 [1509.00805].
- [377] A. Capanema, A. Esmaili and K. Murase, "New constraints on the origin of medium-energy neutrinos observed by IceCube", Phys. Rev. D 101 (2020) 103012 [2002.07192].
- [378] A. Capanema, A. Esmaili and P. D. Serpico, "Where do IceCube neutrinos come from? Hints from the diffuse gamma-ray flux", JCAP 02 (2021) 037 [2007.07911].
- [379] K. Murase, K. Ioka, S. Nagataki and T. Nakamura, "High-energy cosmic-ray nuclei from high- and low-luminosity gamma-ray bursts and implications for multi-messenger astronomy", <u>Phys. Rev. D</u> 78 (2008) 023005 [0801.2861].
- [380] K. Kotera, D. Allard, K. Murase, J. Aoi, Y. Dubois, T. Pierog et al., "Propagation of ultrahigh energy nuclei in clusters of galaxies: resulting composition and secondary emissions", Astrophys. J. 707 (2009) 370 [0907.2433].
- [381] S. Hussain, R. Alves Batista, E. M. de Gouveia Dal Pino and K. Dolag, "High-energy neutrino production in clusters of galaxies", Mon. Not. Roy. Astron. Soc. 507 (2021) 1762 [2101.07702].
- [382] A. Loeb and E. Waxman, "The Cumulative background of high energy neutrinos from starburst galaxies", JCAP 0605 (2006) 003 [astro-ph/0601695].
- [383] T. A. Thompson, E. Quataert, E. Waxman and A. Loeb, "Assessing The Starburst Contribution to the Gamma-Ray and Neutrino Backgrounds", astro-ph/0608699.
- [384] I. Tamborra, S. Ando and K. Murase, "Star-forming galaxies as the origin of diffuse high-energy backgrounds: Gamma-ray and neutrino connections, and implications for starburst history", <u>JCAP</u> 09 (2014) 043 [1404.1189].
- [385] X.-C. Chang and X.-Y. Wang, "The diffuse gamma-ray flux associated with sub-PeV/PeV neutrinos from starburst galaxies", Astrophys.J. 793 (2014) 131 [1406.1099].
- [386] N. Senno, P. Mészáros, K. Murase, P. Baerwald and M. J. Rees, "Extragalactic star-forming galaxies with hypernovae and supernovae as high-energy neutrino and gamma-ray sources: the case of the 10 TeV neutrino data", Astrophys. J. 806 (2015) 24 [1501.04934].
- [387] S. Yoshida and K. Murase, "Constraining photohadronic scenarios for the unified origin of IceCube neutrinos and ultrahigh-energy cosmic rays", Phys. Rev. D 102 (2020) 083023 [2007.09276].
- [388] PIERRE AUGER, TELESCOPE ARRAY Collaboration, "The energy spectrum of ultra-high energy cosmic rays measured at the Pierre Auger Observatory and at the Telescope Array", Pos ICRC2019 (2020) 234 [2001.08811].

- [389] H. Takami, K. Murase, S. Nagataki and K. Sato, "Cosmogenic neutrinos as a probe of the transition from Galactic to extragalactic cosmic rays", Astropart. Phys. **31** (2009) 201 [0704.0979].
- [390] M. Ahlers, L. A. Anchordoqui and S. Sarkar, "Neutrino diagnostics of ultra-high energy cosmic ray protons", Phys. Rev. D 79 (2009) 083009 [0902.3993].
- [391] K. Kotera, D. Allard and A. Olinto, "Cosmogenic Neutrinos: parameter space and detectability from PeV to ZeV", JCAP 10 (2010) 013 [1009.1382].
- [392] M. Ahlers and F. Halzen, "Minimal Cosmogenic Neutrinos", Phys. Rev. D 86 (2012) 083010 [1208.4181].
- [393] P. Baerwald, M. Bustamante and W. Winter, "Are gamma-ray bursts the sources of ultra-high energy cosmic rays?", Astropart. Phys. 62 (2015) 66 [1401.1820].
- [394] R. Aloisio, D. Boncioli, A. di Matteo, A. Grillo, S. Petrera and F. Salamida, "Cosmogenic neutrinos and ultra-high energy cosmic ray models", JCAP 10 (2015) 006 [1505.04020].
- [395] J. Heinze, D. Boncioli, M. Bustamante and W. Winter, "Cosmogenic Neutrinos Challenge the Cosmic Ray Proton Dip Model", Astrophys. J. 825 (2016) 122 [1512.05988].
- [396] K. Møller, P. B. Denton and I. Tamborra, "Cosmogenic Neutrinos Through the GRAND Lens Unveil the Nature of Cosmic Accelerators", JCAP 05 (2019) 047 [1809.04866].
- [397] A. van Vliet, R. Alves Batista and J. R. Hörandel, "Determining the fraction of cosmic-ray protons at ultrahigh energies with cosmogenic neutrinos", Phys. Rev. D 100 (2019) 021302 [1901.01899].
- [398] D. F. G. Fiorillo, M. Bustamante and V. B. Valera, "Near-future discovery of point sources of ultra-high-energy neutrinos", 2205.15985.
- [399] E. Waxman and J. N. Bahcall, "Neutrino afterglow from gamma-ray bursts: Similar to ~ 10¹⁸ eV", <u>Astrophys.</u> J. 541 (2000) 707 [hep-ph/9909286].
- [400] C. D. Dermer and A. Atoyan, "High energy neutrinos from gamma-ray bursts", <u>Phys. Rev. Lett.</u> 91 (2003) 071102 [astro-ph/0301030].
- [401] K. Murase, "High energy neutrino early afterglows gamma-ray bursts revisited", <u>Phys. Rev. D</u> 76 (2007) 123001 [0707.1140].
- [402] S. Razzaque, "Long-lived PeV-EeV neutrinos from gamma-ray burst blastwave", <u>Phys.Rev.</u> D88 (2013) 103003 [1307.7596].
- [403] J. K. Thomas, R. Moharana and S. Razzaque, "Ultrahigh energy neutrino afterglows of nearby long duration gamma-ray bursts", Phys. Rev. D 96 (2017) 103004 [1710.04024].
- [404] D. Biehl, D. Boncioli, A. Fedynitch and W. Winter, "Cosmic-Ray and Neutrino Emission from Gamma-Ray Bursts with a Nuclear Cascade", Astron. Astrophys. 611 (2018) A101 [1705.08909].
- [405] B. T. Zhang and K. Murase, "Ultrahigh-energy cosmic-ray nuclei and neutrinos from engine-driven supernovae", Phys. Rev. D 100 (2019) 103004 [1812.10289].
- [406] J. Heinze, D. Biehl, A. Fedynitch, D. Boncioli, A. Rudolph and W. Winter, "Systematic parameter space study for the UHECR origin from GRBs in models with multiple internal shocks", <u>Mon. Not. Roy. Astron. Soc.</u> 498 (2020) 5990 [2006.14301].
- [407] K. Murase, P. Mészáros and B. Zhang, "Probing the birth of fast rotating magnetars through high-energy neutrinos", Phys. Rev. D 79 (2009) 103001 [0904.2509].
- [408] K. Fang, B. D. Metzger, K. Murase, I. Bartos and K. Kotera, "Multimessenger Implications of AT2018cow: High-energy Cosmic-Ray and Neutrino Emissions from Magnetar-powered Superluminous Transients", Astrophys. J. 878 (2019) 34 [1812.11673].
- [409] J. A. Carpio, K. Murase, M. H. Reno, I. Sarcevic and A. Stasto, "Charm contribution to ultrahigh-energy neutrinos from newborn magnetars", Phys. Rev. D 102 (2020) 103001 [2007.07945].
- [410] K. Murase, "Astrophysical high-energy neutrinos and gamma-ray bursts", AIP Conf. Proc. 1065 (2008) 201.
- [411] D. Biehl, D. Boncioli, C. Lunardini and W. Winter, "Tidally disrupted stars as a possible origin of both cosmic rays and neutrinos at the highest energies", Sci. Rep. 8 (2018) 10828 [1711.03555].

- [412] C. Guépin, K. Kotera, E. Barausse, K. Fang and K. Murase, "Ultra-High Energy Cosmic Rays and Neutrinos from Tidal Disruptions by Massive Black Holes", Astron. Astrophys. 616 (2018) A179 [1711.11274].
- [413] K. Murase, Y. Inoue and C. D. Dermer, "Diffuse Neutrino Intensity from the Inner Jets of Active Galactic Nuclei: Impacts of External Photon Fields and the Blazar Sequence", <u>Phys. Rev. D</u> 90 (2014) 023007 [1403.4089].
- [414] F. Oikonomou, K. Murase, P. Padovani, E. Resconi and P. Mészáros, "High energy neutrino flux from individual blazar flares", Mon. Not. Roy. Astron. Soc. 489 (2019) 4347 [1906.05302].
- [415] C. Righi, A. Palladino, F. Tavecchio and F. Vissani, "EeV Astrophysical neutrinos from FSRQs?", 2003.08701.
- [416] S. Schonert, T. K. Gaisser, E. Resconi and O. Schulz, "Vetoing atmospheric neutrinos in a high energy neutrino telescope", Phys. Rev. D 79 (2009) 043009 [0812.4308].
- [417] T. K. Gaisser, K. Jero, A. Karle and J. van Santen, "Generalized self-veto probability for atmospheric neutrinos", Phys. Rev. D 90 (2014) 023009 [1405.0525].
- [418] C. A. Argüelles, S. Palomares-Ruiz, A. Schneider, L. Wille and T. Yuan, "Unified atmospheric neutrino passing fractions for large-scale neutrino telescopes", JCAP 07 (2018) 047 [1805.11003].
- [419] L. J. Schumacher, M. Huber, M. Agostini, M. Bustamante, F. Oikonomou and E. Resconi, "PLEnuM: A global and distributed monitoring system of high-energy astrophysical neutrinos", PoS ICRC2021 (2021) 1185.
- [420] ANTARES Collaboration, "Study of the high-energy neutrino diffuse flux with the ANTARES neutrino telescope", PoS ICRC2019 (2020) 891.
- [421] ICECUBE Collaboration, "A Combined Fit of the Diffuse Neutrino Spectrum using IceCube Muon Tracks and Cascades", PoS ICRC2021 (2021) 1129 [2107.10003].
- [422] A. Watanabe, "The spectrum and flavor composition of the astrophysical neutrinos in IceCube", <u>JCAP</u> 08 (2015) 030 [1412.8264].
- [423] A. Palladino, G. Pagliaroli, F. L. Villante and F. Vissani, "What is the Flavor of the Cosmic Neutrinos Seen by IceCube?", Phys. Rev. Lett. 114 (2015) 171101 [1502.02923].
- [424] G. D'Amico, "Flavor and energy inference for the high-energy IceCube neutrinos", <u>Astropart. Phys.</u> **101** (2018) 8 [1712.04979].
- [425] A. Palladino, "The flavor composition of astrophysical neutrinos after 8 years of IceCube: an indication of neutron decay scenario?", Eur. Phys. J. C 79 (2019) 500 [1902.08630].
- [426] M. Bustamante, "New limits on neutrino decay from the Glashow resonance of high-energy cosmic neutrinos", 2004.06844.
- [427] M. Ahlers, L. Anchordoqui, M. Gonzalez-Garcia, F. Halzen and S. Sarkar, "GZK Neutrinos after the Fermi-LAT Diffuse Photon Flux Measurement", Astropart. Phys. 34 (2010) 106 [1005.2620].
- [428] R. Alves Batista, D. Boncioli, A. di Matteo, A. van Vliet and D. Walz, "Effects of uncertainties in simulations of extragalactic UHECR propagation, using CRPropa and SimProp", JCAP 10 (2015) 063 [1508.01824].
- [429] R. Alves Batista, D. Boncioli, A. di Matteo and A. van Vliet, "Secondary neutrino and gamma-ray fluxes from SimProp and CRPropa", JCAP 05 (2019) 006 [1901.01244].
- [430] S. Yoshida and A. Ishihara, "Constraints on the origin of the ultra-high energy cosmic-rays using cosmic diffuse neutrino flux limits: An analytical approach", Phys. Rev. D 85 (2012) 063002 [1202.3522].
- [431] PIERRE AUGER Collaboration, "Probing the origin of ultra-high-energy cosmic rays with neutrinos in the EeV energy range using the Pierre Auger Observatory", JCAP 10 (2019) 022 [1906.07422].
- [432] ICECUBE Collaboration, "Time-Integrated Neutrino Source Searches with 10 Years of IceCube Data", <u>Phys.</u> Rev. Lett. **124** (2020) 051103 [1910.08488].
- [433] ICECUBE, FERMI-LAT, MAGIC, AGILE, ASAS-SN, HAWC, H.E.S.S., INTEGRAL, KANATA, KISO, KAPTEYN, LIVERPOOL TELESCOPE, SUBARU, SWIFT NUSTAR, VERITAS, VLA/17B-403 Collaboration, "Multimessenger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A", Science 361 (2018) eaat1378 [1807.08816].

- [434] A. Silvestri and S. W. Barwick, "Constraints on Extragalactic Point Source Flux from Diffuse Neutrino Limits", Phys.Rev. D81 (2010) 023001 [0908.4266].
- [435] ICECUBE, MAGIC, VERITAS Collaboration, "Very High-Energy Gamma-Ray Follow-Up Program Using Neutrino Triggers from IceCube", JINST 11 (2016) P11009 [1610.01814].
- [436] ICECUBE Collaboration, "Search for PeV Gamma-Ray Emission from the Southern Hemisphere with 5 Years of Data from the IceCube Observatory", Astrophys. J. 891 (2019) 9 [1908.09918].
- [437] AMON TEAM, HAWC, ICECUBE Collaboration, "Multimessenger Gamma-Ray and Neutrino Coincidence Alerts Using HAWC and IceCube Subthreshold Data", Astrophys. J. 906 (2020) 63 [2008.10616].
- [438] ICECUBE Collaboration, "Search for High-energy Neutrinos from Ultraluminous Infrared Galaxies with IceCube", Astrophys. J. 926 (2022) 59 [2107.03149].
- [439] ICECUBE Collaboration, "The contribution of Fermi-2LAC blazars to the diffuse TeV-PeV neutrino flux", Astrophys. J. 835 (2017) 45 [1611.03874].
- [440] FERMI-LAT, ASAS-SN, ICECUBE Collaboration, "Investigation of two Fermi-LAT gamma-ray blazars coincident with high-energy neutrinos detected by IceCube", Astrophys. J. 880 (2019) 880:103 [1901.10806].
- [441] ICECUBE Collaboration, "A Search for MeV to TeV Neutrinos from Fast Radio Bursts with IceCube", Astrophys. J. 890 (2020) 111 [1908.09997].
- [442] ICECUBE Collaboration, "Extending the search for muon neutrinos coincident with gamma-ray bursts in IceCube data", Astrophys. J. 843 (2017) 112 [1702.06868].
- [443] ICECUBE, PTF, SWIFT, PAN-STARRS1 SCIENCE CONSORTIUM Collaboration, "The Detection of a SN IIn in Optical Follow-up Observations of IceCube Neutrino Events", Astrophys. J. 811 (2015) 52 [1506.03115].
- [444] ICECUBE Collaboration, "IceCube Search for Neutrinos Coincident with Compact Binary Mergers from LIGO-Virgo's First Gravitational-wave Transient Catalog", Astrophys. J. Lett. 898 (2020) L10 [2004.02910].
- [445] ANTARES, ICECUBE, AUGER, TELESCOPE ARRAY Collaboration, "Search for Spatial Correlations of Neutrinos with Ultra-High-Energy Cosmic Rays", 2201.07313.
- [446] M. Ahlers and K. Murase, "Probing the Galactic Origin of the IceCube Excess with Gamma-Rays", <u>Phys.Rev.</u> D90 (2014) 023010 [1309.4077].
- [447] M. Ahlers, Y. Bai, V. Barger and R. Lu, "Galactic neutrinos in the TeV to PeV range", <u>Phys. Rev. D</u> 93 (2016) 013009 [1505.03156].
- [448] FERMI Collaboration, "Fermi-lat observations of the diffuse γ -ray emission: Implications for cosmic rays and the interstellar medium", The Astrophysical Journal **750** (2012) 3.
- [449] D. Gaggero, A. Urbano, M. Valli and P. Ullio, "Gamma-ray sky points to radial gradients in cosmic-ray transport", Phys. Rev. D 91 (2015) 083012 [1411.7623].
- [450] H. León Vargas, A. Sandoval, E. Belmont and R. Alfaro, "Capability of the HAWC Gamma-Ray Observatory for the Indirect Detection of Ultrahigh-Energy Neutrinos", Adv. Astron. 2017 (2017) 1932413 [1610.04820].
- [451] LHAASO Collaboration, "Performance of LHAASO-WCDA and Observation of Crab Nebula as a Standard Candle", 2101.03508.
- [452] ANTARES, ICECUBE Collaboration, "Joint Constraints on Galactic Diffuse Neutrino Emission from the ANTARES and IceCube Neutrino Telescopes", Astrophys. J. Lett. 868 (2018) L20 [1808.03531].
- [453] ANTARES, ICECUBE Collaboration, "Combined search for neutrinos from dark matter self-annihilation in the Galactic Center with ANTARES and IceCube", Phys. Rev. D 102 (2020) 082002 [2003.06614].
- [454] A. Keivani et al., "A Multimessenger Picture of the Flaring Blazar TXS 0506+056: implications for High-Energy Neutrino Emission and Cosmic Ray Acceleration", Astrophys. J. 864 (2018) 84 [1807.04537].
- [455] S. Gao, A. Fedynitch, W. Winter and M. Pohl, "Modelling the coincident observation of a high-energy neutrino and a bright blazar flare", Nature Astron. 3 (2019) 88 [1807.04275].
- [456] MAGIC Collaboration, "The blazar TXS 0506+056 associated with a high-energy neutrino: insights into extragalactic jets and cosmic ray acceleration", Astrophys. J. Lett. **863** (2018) L10 [1807.04300].

- [457] M. Cerruti, A. Zech, C. Boisson, G. Emery, S. Inoue and J. P. Lenain, "Leptohadronic single-zone models for the electromagnetic and neutrino emission of TXS 0506+056", <u>Mon. Not. Roy. Astron. Soc.</u> 483 (2019) L12 [1807.04335].
- [458] K. Murase, F. Oikonomou and M. Petropoulou, "Blazar Flares as an Origin of High-Energy Cosmic Neutrinos?", Astrophys. J. 865 (2018) 124 [1807.04748].
- [459] A. Reimer, M. Boettcher and S. Buson, "Cascading Constraints from Neutrino-emitting Blazars: The Case of TXS 0506+056", Astrophys. J. 881 (2019) 46 [1812.05654].
- [460] X. Rodrigues, S. Gao, A. Fedynitch, A. Palladino and W. Winter, "Leptohadronic Blazar Models Applied to the 2014–2015 Flare of TXS 0506+056", Astrophys. J. Lett. 874 (2019) L29 [1812.05939].
- [461] M. Petropoulou et al., "Multi-Epoch Modeling of TXS 0506+056 and Implications for Long-Term High-Energy Neutrino Emission", Astrophys. J. 891 (2020) 115 [1911.04010].
- [462] R. Xue, R.-Y. Liu, M. Petropoulou, F. Oikonomou, Z.-R. Wang, K. Wang et al., "A two-zone model for blazar emission: implications for TXS 0506+056 and the neutrino event IceCube-170922A", <u>Astrophys. J.</u> 886 (2019) 23 [1908.10190].
- [463] B. T. Zhang, M. Petropoulou, K. Murase and F. Oikonomou, "A Neutral Beam Model for High-Energy Neutrino Emission from the Blazar TXS 0506+056", Astrophys. J. 889 (2020) 118 [1910.11464].
- [464] M. Kadler et al., "Coincidence of a high-fluence blazar outburst with a PeV-energy neutrino event", <u>Nature</u> Phys. **12** (2016) 807 [1602.02012].
- [465] P. Giommi, P. Padovani, F. Oikonomou, T. Glauch, S. Paiano and E. Resconi, "3HSP J095507.9+355101: a flaring extreme blazar coincident in space and time with IceCube-200107A", <u>Astron. Astrophys.</u> 640 (2020) L4 [2003.06405].
- [466] M. Petropoulou, F. Oikonomou, A. Mastichiadis, K. Murase, P. Padovani, G. Vasilopoulos et al., "Comprehensive Multimessenger Modeling of the Extreme Blazar 3HSP J095507.9+355101 and Predictions for IceCube", Astrophys. J. 899 (2020) 113 [2005.07218].
- [467] X. Rodrigues, S. Garrappa, S. Gao, V. S. Paliya, A. Franckowiak and W. Winter, "Multiwavelength and Neutrino Emission from Blazar PKS 1502 + 106", Astrophys. J. 912 (2021) 54 [2009.04026].
- [468] F. Oikonomou, M. Petropoulou, K. Murase, A. Tohuvavohu, G. Vasilopoulos, S. Buson et al., "Multi-messenger emission from the parsec-scale jet of the flat-spectrum radio quasar coincident with high-energy neutrino IceCube-190730A", JCAP 10 (2021) 082 [2107.11437].
- [469] K. Murase, S. S. Kimura, B. T. Zhang, F. Oikonomou and M. Petropoulou, "High-Energy Neutrino and Gamma-Ray Emission from Tidal Disruption Events", Astrophys. J. 902 (2020) 108 [2005.08937].
- [470] W. Winter and C. Lunardini, "A concordance scenario for the observed neutrino from a tidal disruption event", Nature Astron. 5 (2021) 5 [2005.06097].
- [471] T. M. Yoast-Hull, J. S. G. III, E. G. Zweibel and J. E. Everett, "Active Galactic Nuclei, Neutrinos, and Interacting Cosmic Rays in NGC 253 and NGC 1068", Astrophys. J. 780 (2014) 137 [1311.5586].
- [472] MAGIC Collaboration, "Constraints on gamma-ray and neutrino emission from NGC 1068 with the MAGIC telescopes", Astrophys. J. 883 (2019) 135 [1906.10954].
- [473] Y. Inoue, D. Khangulyan and A. Doi, "On the Origin of High-energy Neutrinos from NGC 1068: The Role of Nonthermal Coronal Activity", Astrophys. J. Lett. 891 (2020) L33 [1909.02239].
- [474] X. Wang and A. Loeb, "Ultrahigh energy cosmic rays from nonrelativistic quasar outflows", <u>Phys. Rev. D</u> 95 (2017) 063007 [1611.07616].
- [475] R.-Y. Liu, K. Murase, S. Inoue, C. Ge and X.-Y. Wang, "Can winds driven by active galactic nuclei account for the extragalactic gamma-ray and neutrino backgrounds?", Astrophys. J. 858 (2018) 9 [1712.10168].
- [476] A. Kheirandish, K. Murase and S. S. Kimura, "High-energy Neutrinos from Magnetized Coronae of Active Galactic Nuclei and Prospects for Identification of Seyfert Galaxies and Quasars in Neutrino Telescopes", Astrophys. J. 922 (2021) 45 [2102.04475].
- [477] ICECUBE Collaboration, "The IceCube Realtime Alert System", Astropart. Phys. 92 (2017) 30 [1612.06028].

- [478] ICECUBE Collaboration, "The Next Generation of IceCube Real-time Neutrino Alerts", Pos ICRC2019 (2020) 1021 [1908.04884].
- [479] ANTARES Collaboration, "Real-time follow-up of multi-messenger alerts with the ANTARES neutrino telescope", PoS ICRC2019 (2021) 872.
- [480] ICECUBE Collaboration, "Search for Sources of Astrophysical Neutrinos Using Seven Years of IceCube Cascade Events", Astrophys. J. 886 (2019) 12 [1907.06714].
- [481] H. A. Ayala Solares et al., "The Astrophysical Multimessenger Observatory Network (AMON): Performance and science program", Astropart. Phys. 114 (2020) 68 [1903.08714].
- [482] S. Palomares-Ruiz, A. Irimia and T. J. Weiler, "Acceptances for space-based and ground-based fluorescence detectors, and inference of the neutrino-nucleon cross-section above 10¹⁹ eV", <u>Phys. Rev. D</u> 73 (2006) 083003 [astro-ph/0512231].
- [483] P. S. B. Dev, D. K. Ghosh and W. Rodejohann, "R-parity Violating Supersymmetry at IceCube", <u>Phys. Lett.</u> B 762 (2016) 116 [1605.09743].
- [484] J. F. Beacom, N. F. Bell, D. Hooper, J. G. Learned, S. Pakvasa and T. J. Weiler, "PseudoDirac neutrinos: A Challenge for neutrino telescopes", Phys. Rev. Lett. 92 (2004) 011101 [hep-ph/0307151].
- [485] A. Esmaili, "Pseudo-Dirac Neutrino Scenario: Cosmic Neutrinos at Neutrino Telescopes", <u>Phys. Rev. D</u> 81 (2010) 013006 [0909.5410].
- [486] A. Esmaili and Y. Farzan, "Implications of the Pseudo-Dirac Scenario for Ultra High Energy Neutrinos from GRBs", JCAP 12 (2012) 014 [1208.6012].
- [487] Y. Ahn, S. K. Kang and C. Kim, "A Model for Pseudo-Dirac Neutrinos: Leptogenesis and Ultra-High Energy Neutrinos", JHEP 10 (2016) 092 [1602.05276].
- [488] D. Colladay and V. Kostelecky, "Lorentz-violating extension of the Standard Model", <u>Phys. Rev. D</u> 58 (1998) 116002 [hep-ph/9809521].
- [489] V. Kostelecky and M. Mewes, "Lorentz and CPT violation in neutrinos", <u>Phys. Rev. D</u> 69 (2004) 016005 [hep-ph/0309025].
- [490] V. Kostelecky and N. Russell, "Data Tables for Lorentz and CPT Violation", <u>Rev. Mod. Phys.</u> 83 (2011) 11 [0801.0287].
- [491] P. Gorham et al., "Implications of ultra-high energy neutrino flux constraints for Lorentz-invariance violating cosmogenic neutrinos", Phys. Rev. D 86 (2012) 103006 [1207.6425].
- [492] E. Borriello, S. Chakraborty, A. Mirizzi and P. D. Serpico, "Stringent constraint on neutrino Lorentz-invariance violation from the two IceCube PeV neutrinos", Phys. Rev. D 87 (2013) 116009 [1303.5843].
- [493] L. Anchordoqui, V. Barger, H. Goldberg, J. Learned, D. Marfatia, S. Pakvasa et al., "End of the cosmic neutrino energy spectrum", Phys. Lett. B 739 (2014) 99 [1404.0622].
- [494] G. Amelino-Camelia, D. Guetta and T. Piran, "Icecube Neutrinos and Lorentz Invariance Violation", Astrophys. J. 806 (2015) 269.
- [495] J. Liao and D. Marfatia, "IceCube's astrophysical neutrino energy spectrum from CPT violation", <u>Phys. Rev.</u> D 97 (2018) 041302 [1711.09266].
- [496] L. Anchordoqui, T. Han, D. Hooper and S. Sarkar, "Exotic neutrino interactions at the Pierre Auger Observatory", Astropart. Phys. 25 (2006) 14 [hep-ph/0508312].
- [497] G. Pagliaroli, A. Palladino, F. Villante and F. Vissani, "Testing nonradiative neutrino decay scenarios with IceCube data", Phys. Rev. D 92 (2015) 113008 [1506.02624].
- [498] A. Bhattacharya, M. H. Reno and I. Sarcevic, "Reconciling neutrino flux from heavy dark matter decay and recent events at IceCube", JHEP 06 (2014) 110 [1403.1862].
- [499] A. Esmaili, S. K. Kang and P. D. Serpico, "IceCube events and decaying dark matter: hints and constraints", JCAP 12 (2014) 054 [1410.5979].
- [500] C. El Aisati, M. Gustafsson and T. Hambye, "New Search for Monochromatic Neutrinos from Dark Matter Decay", Phys. Rev. D 92 (2015) 123515 [1506.02657].

- [501] S. Troitsky, "Search for Galactic disk and halo components in the arrival directions of high-energy astrophysical neutrinos", JETP Lett. 102 (2015) 785 [1511.01708].
- [502] M. Chianese, G. Miele, S. Morisi and E. Vitagliano, "Low energy IceCube data and a possible Dark Matter related excess", Phys. Lett. B 757 (2016) 251 [1601.02934].
- [503] M. Chianese, G. Miele and S. Morisi, "Dark Matter interpretation of low energy IceCube MESE excess", <u>JCAP</u> 01 (2017) 007 [1610.04612].
- [504] A. Bhattacharya, A. Esmaili, S. Palomares-Ruiz and I. Sarcevic, "Probing decaying heavy dark matter with the 4-year IceCube HESE data", JCAP 07 (2017) 027 [1706.05746].
- [505] ICECUBE Collaboration, "Search for neutrinos from decaying dark matter with IceCube", <u>Eur. Phys. J. C</u> 78 (2018) 831 [1804.03848].
- [506] A. Bhattacharya, A. Esmaili, S. Palomares-Ruiz and I. Sarcevic, "Update on decaying and annihilating heavy dark matter with the 6-year IceCube HESE data", JCAP 05 (2019) 051 [1903.12623].
- [507] ICECUBE Collaboration, "Searches for Connections Between Dark Matter and Neutrinos with the IceCube High-Energy Starting Event Sample", PoS ICRC2019 (2020) 839 [1907.11193].
- [508] M. Chianese, D. F. Fiorillo, G. Miele, S. Morisi and O. Pisanti, "Decaying dark matter at IceCube and its signature on High Energy gamma experiments", JCAP 11 (2019) 046 [1907.11222].
- [509] ANTARES, ICECUBE Collaboration, "Combined search for neutrinos from dark matter self-annihilation in the Galactic Centre with ANTARES and IceCube", 2003.06614.
- [510] T. J. Weiler, "New Physics in Neutrino Telescopes." Presented at TeV Particle Astrophysics (TeVPA) 2006, Madison, WI, USA.
- [511] S. Ando, M. Kamionkowski and I. Mocioiu, "Neutrino Oscillations, Lorentz/CPT Violation, and Dark Energy", Phys. Rev. D 80 (2009) 123522 [0910.4391].
- [512] J. Barranco, O. G. Miranda, C. A. Moura, T. I. Rashba and F. Rossi-Torres, "Confusing the extragalactic neutrino flux limit with a neutrino propagation limit", JCAP 10 (2011) 007 [1012.2476].
- [513] O. Miranda, C. Moura and A. Parada, "Sterile neutrinos, dark matter, and resonant effects in ultra high energy regimes", Phys. Lett. B 744 (2015) 55 [1308.1408].
- [514] P. de Salas, R. Lineros and M. Tórtola, "Neutrino propagation in the galactic dark matter halo", <u>Phys. Rev. D</u> 94 (2016) 123001 [1601.05798].
- [515] M. M. Reynoso and O. A. Sampayo, "Propagation of high-energy neutrinos in a background of ultralight scalar dark matter", Astropart. Phys. 82 (2016) 10 [1605.09671].
- [516] Y. Farzan and S. Palomares-Ruiz, "Flavor of cosmic neutrinos preserved by ultralight dark matter", <u>Phys. Rev.</u> D 99 (2019) 051702 [1810.00892].
- [517] J. Alvey and M. Fairbairn, "Linking Scalar Dark Matter and Neutrino Masses with IceCube 170922A", <u>JCAP</u> 07 (2019) 041 [1902.01450].
- [518] K.-Y. Choi, J. Kim and C. Rott, "Constraining dark matter-neutrino interactions with IceCube-170922A", Phys. Rev. D 99 (2019) 083018 [1903.03302].
- [519] S. Koren, "Neutrino Dark Matter Scattering and Coincident Detections of UHE Neutrinos with EM Sources", JCAP 09 (2019) 013 [1903.05096].
- [520] J. F. Cherry and I. M. Shoemaker, "Sterile neutrino origin for the upward directed cosmic ray showers detected by ANITA", Phys. Rev. D 99 (2019) 063016 [1802.01611].
- [521] L. A. Anchordoqui, V. Barger, J. G. Learned, D. Marfatia and T. J. Weiler, "Upgoing ANITA events as evidence of the CPT symmetric universe", LHEP 1 (2018) 13 [1803.11554].
- [522] G.-y. Huang, "Sterile neutrinos as a possible explanation for the upward air shower events at ANITA", <u>Phys.</u> Rev. D **98** (2018) 043019 [1804.05362].
- [523] L. A. Anchordoqui and I. Antoniadis, "Supersymmetric sphaleron configurations as the origin of the perplexing ANITA events", Phys. Lett. B 790 (2019) 578 [1812.01520].

- [524] J. M. Cline, C. Gross and W. Xue, "Can the ANITA anomalous events be due to new physics?", Phys. Rev. D 100 (2019) 015031 [1904.13396].
- [525] S. Chipman, R. Diesing, M. H. Reno and I. Sarcevic, "Anomalous ANITA air shower events and tau decays", Phys. Rev. D 100 (2019) 063011 [1906.11736].
- [526] M. Abdullah, B. Dutta, S. Ghosh and T. Li, " $(g-2)_{\mu,e}$ and the ANITA anomalous events in a three-loop neutrino mass model", Phys. Rev. D 100 (2019) 115006 [1907.08109].
- [527] P. A. M. Dirac, "Quantised singularities in the electromagnetic field,", Proc. Roy. Soc. Lond. A A133 (1931) 60.
- [528] ICECUBE Collaboration, "Search for Relativistic Magnetic Monopoles with IceCube", <u>Phys. Rev. D</u> 87 (2013) 022001 [1208.4861].
- [529] ICECUBE Collaboration, "Search for non-relativistic Magnetic Monopoles with IceCube", <u>Eur. Phys. J. C</u> 74 (2014) 2938 [1402.3460].
- [530] ICECUBE Collaboration, "Searches for Relativistic Magnetic Monopoles in IceCube", <u>Eur. Phys. J. C</u> 76 (2016) 133 [1511.01350].
- [531] ANTARES Collaboration, "Search for relativistic magnetic monopoles with five years of the ANTARES detector data", JHEP 07 (2017) 054 [1703.00424].
- [532] ICECUBE Collaboration, "The IceCube Upgrade Design and Science Goals", PoS ICRC2019 (2020) 1031 [1908.09441].
- [533] DUNE Collaboration, "Long-baseline neutrino oscillation physics potential of the DUNE experiment", <u>Eur.</u> Phys. J. C 80 (2020) 978 [2006.16043].
- [534] JUNO Collaboration, "Neutrino oscillation studies in JUNO", PoS NuFact2019 (2019) 030.
- [535] HYPER-KAMIOKANDE Collaboration, "Physics potential of Hyper-Kamiokande for neutrino oscillation measurements", PoS NuFact2019 (2019) 040.
- [536] S. A. R. Ellis, K. J. Kelly and S. W. Li, "Current and Future Neutrino Oscillation Constraints on Leptonic Unitarity", JHEP 12 (2020) 068 [2008.01088].
- [537] FASER Collaboration, "Detecting and Studying High-Energy Collider Neutrinos with FASER at the LHC", Eur. Phys. J. C 80 (2020) 61 [1908.02310].
- [538] D. Carney et al., "Snowmass2021 Cosmic Frontier White Paper: Ultraheavy particle dark matter", 2203.06508.
- [539] T. Aramaki et al., "Snowmass2021 Cosmic Frontier: The landscape of cosmic-ray and high-energy photon probes of particle dark matter", 2203.06894.
- [540] E. Berti et al., "Dark Matter In Extreme Astrophysical Environments", in <u>2022 Snowmass Summer Study</u>, 3, 2022, 2203.07984.
- [541] S. Chakrabarti et al., "Snowmass2021 Cosmic Frontier White Paper: Observational Facilities to Study Dark Matter", in 2022 Snowmass Summer Study, 3, 2022, 2203.06200.
- [542] C. A. Argüelles et al., "Snowmass White Paper: Beyond the Standard Model effects on Neutrino Flavor", in 2022 Snowmass Summer Study, 3, 2022, 2203.10811.
- [543] W. Bai, M. V. Diwan, M. V. Garzelli, Y. S. Jeong, K. Kumar and M. H. Reno, "Prompt electron and tau neutrinos and antineutrinos in the forward region at the LHC", JHEAP 34 (2022) 212 [2203.07212].
- [544] V. Niess and O. Martineau-Huynh, "DANTON: a Monte-Carlo sampler of τ from ν_{τ} interacting with the Earth", 1810.01978.
- [545] NUSPACESIM Collaboration, "Monte Carlo simulations of neutrino and charged lepton propagation in the Earth with nuPyProp", PoS ICRC2021 (2021) 1203 [2109.08198].
- [546] I. Safa, J. Lazar, A. Pizzuto, O. Vasquez, C. A. Argüelles and J. Vandenbroucke, "TauRunner: A Public Python Program to Propagate Neutral and Charged Leptons", 2110.14662.
- [547] B. Zhou and J. F. Beacom, "Dimuons in Neutrino Telescopes: New Predictions and First Search in IceCube", 2110.02974.

- [548] C. Sun, F. Zhang, F. Hu, D. Xu and J. Gao, "Search for Charm-quark Production via Dimuons in Neutrino Telescopes", 2201.03580.
- [549] P. Jain, D. W. McKay, S. Panda and J. P. Ralston, "Extra dimensions and strong neutrino nucleon interactions above 10¹⁹-eV: Breaking the GZK barrier", Phys. Lett. B 484 (2000) 267 [hep-ph/0001031].
- [550] S. Yoshida, "Constraints on Neutrino-Nucleon Interactions at energies of 1 EeV with the IceCube Neutrino Observatory", Phys. Rev. D 82 (2010) 103012 [1009.1679].
- [551] L. Gerhardt and S. R. Klein, "Electron and Photon Interactions in the Regime of Strong LPM Suppression", Phys. Rev. D 82 (2010) 074017 [1007.0039].
- [552] S. Stjärnholm, O. Ericsson and C. Glaser, "Neutrino direction and flavor reconstruction from radio detector data using deep convolutional neural networks", PoS ICRC2021 (2021) 1055.
- [553] J. Alvarez-Muniz, R. A. Vazquez and E. Zas, "Characterization of neutrino signals with radiopulses in dense media through the LPM effect", Phys. Rev. D 61 (2000) 023001 [astro-ph/9901278].
- [554] C. Guépin, R. Aloisio, L. A. Anchordoqui, A. Cummings, J. F. Krizmanic, A. V. Olinto et al., "Indirect dark matter searches at ultrahigh energy neutrino detectors", Phys. Rev. D 104 (2021) 083002 [2106.04446].
- [555] Y. Farzan, "On the τ flavor of the cosmic neutrino flux", JHEP 07 (2021) 174 [2105.03272].
- [556] Y. Farzan, "Hide-and-seek with cosmic tau neutrinos", in European Physical Society Conference on High Energy Physics 2021, 10, 2021, 2110.07222.
- [557] E. W. Kolb and M. S. Turner, "Supernova SN 1987a and the Secret Interactions of Neutrinos", <u>Phys. Rev. D</u> 36 (1987) 2895.
- [558] POEMMA Collaboration, "The POEMMA (Probe of Extreme Multi-Messenger Astrophysics) observatory", JCAP 06 (2021) 007 [2012.07945].
- [559] ICECUBE-GEN2 Collaboration, "The Surface Array planned for IceCube-Gen2", Pos ICRC2021 (2021) 407 [2108.00364].
- [560] F. Halzen and L. Wille, "Charm contribution to the atmospheric neutrino flux", <u>Phys. Rev. D</u> 94 (2016) 014014 [1605.01409].
- [561] R. Laha and S. J. Brodsky, "IceCube can constrain the intrinsic charm of the proton", <u>Phys. Rev. D</u> 96 (2017) 123002 [1607.08240].
- [562] A. V. Giannini, V. P. Gonçalves and F. S. Navarra, "Intrinsic charm contribution to the prompt atmospheric neutrino flux", Phys. Rev. D 98 (2018) 014012 [1803.01728].
- [563] A. Bhattacharya and J. R. Cudell, "Forward charm-production models and prompt neutrinos at IceCube", JHEP 11 (2018) 150 [1808.00293].
- [564] V. P. Goncalves, R. Maciula and A. Szczurek, "Impact of intrinsic charm amount in the nucleon and saturation effects on the prompt atmospheric ν_{μ} flux for IceCube", 2103.05503.
- [565] P. F. de Salas, D. V. Forero, C. A. Ternes, M. Tortola and J. W. F. Valle, "Status of neutrino oscillations 2018: 3σ hint for normal mass ordering and improved CP sensitivity", Phys. Lett. B **782** (2018) 633 [1708.01186].
- [566] BAIKAL-GVD Collaboration, "Deep-Water Neutrino Telescope in Lake Baikal", <u>Physics of Atomic Nuclei</u> 84 (2021) 1600.
- [567] S. Gabici, C. Evoli, D. Gaggero, P. Lipari, P. Mertsch, E. Orlando et al., "The origin of Galactic cosmic rays: challenges to the standard paradigm", Int. J. Mod. Phys. D 28 (2019) 1930022 [1903.11584].
- [568] TELESCOPE ARRAY Collaboration, "Depth of Ultra High Energy Cosmic Ray Induced Air Shower Maxima Measured by the Telescope Array Black Rock and Long Ridge FADC Fluorescence Detectors and Surface Array in Hybrid Mode", Astrophys. J. 858 (2018) 76 [1801.09784].
- [569] PIERRE AUGER Collaboration, "Combined fit of spectrum and composition data as measured by the Pierre Auger Observatory", JCAP 04 (2017) 038 [1612.07155].
- [570] PIERRE AUGER, TELESCOPE ARRAY Collaboration, "Testing the agreement between the X_{max} distributions measured by the Pierre Auger and Telescope Array Observatories", PoS ICRC2017 (2018) 522.

- [571] W. Hanlon, "Telescope Array Hybrid Composition and Auger-TA Composition Comparison", <u>EPJ Web Conf.</u> 208 (2019) 02001 [1810.00431].
- [572] K. Murase and J. F. Beacom, "Neutrino Background Flux from Sources of Ultrahigh-Energy Cosmic-Ray Nuclei", Phys.Rev. D81 (2010) 123001 [1003.4959].
- [573] H.E.S.S. Collaboration, "Acceleration of petaelectronvolt protons in the Galactic Centre", <u>Nature</u> **531** (2016) 476 [1603.07730].
- [574] HAWC Collaboration, "Multiple Galactic Sources with Emission Above 56 TeV Detected by HAWC", <u>Phys.</u> Rev. Lett. **124** (2020) 021102 [1909.08609].
- [575] M. Amenomori et al., "First Detection of Photons with Energy Beyond 100 TeV from an Astrophysical Source", Phys. Rev. Lett. **123** (2019) 051101 [1906.05521].
- [576] Z. Cao et al., "Ultrahigh-energy photons up to 1.4 petaelectronvolts from 12 γ -ray Galactic sources", <u>Nature</u> 594 (2021) 33.
- [577] F. W. Stecker, "A note on high energy neutrinos from agn cores", <u>Phys. Rev. D</u> 72 (2005) 107301 [astro-ph/0510537].
- [578] S. Hallmann, B. Clark, C. Glaser and D. Smith for the IceCube-Gen2 Collaboration, "Sensitivity studies for the IceCube-Gen2 radio array", PoS ICRC2021 (2021) 1183 [2107.08910].
- [579] "All-sky point-source IceCube data: years 2010-2012." https: //icecube.wisc.edu/data-releases/2018/10/all-sky-point-source-icecube-data-years-2010-2012/. Accessed: 2021-02-16.
- [580] ICECUBE Collaboration, "All-sky Search for Time-integrated Neutrino Emission from Astrophysical Sources with 7 yr of IceCube Data", Astrophys. J. 835 (2017) 151 [1609.04981].
- [581] PUEO Collaboration, "The Payload for Ultrahigh Energy Observations (PUEO): a white paper", <u>JINST</u> 16 (2021) P08035 [2010.02892].
- [582] T. M. Venters, M. H. Reno, J. F. Krizmanic, L. A. Anchordoqui, C. Guépin and A. V. Olinto, "POEMMA's Target of Opportunity Sensitivity to Cosmic Neutrino Transient Sources", <u>Phys. Rev. D</u> 102 (2020) 123013 [1906.07209].
- [583] K. Kotera, F. Oikonomou et al. personal communication.
- [584] S. S. Kimura, K. Murase, P. Mészáros and K. Kiuchi, "High-Energy Neutrino Emission from Short Gamma-Ray Bursts: Prospects for Coincident Detection with Gravitational Waves", <u>Astrophys. J. Lett.</u> 848 (2017) L4 [1708.07075].
- [585] ARA Collaboration, "Constraints on the diffuse flux of ultrahigh energy neutrinos from four years of Askaryan Radio Array data in two stations", Phys. Rev. D **102** (2020) 043021 [1912.00987].
- [586] A. Anker et al., "A search for cosmogenic neutrinos with the ARIANNA test bed using 4.5 years of data", JCAP **03** (2020) 053 [1909.00840].
- [587] ICECUBE Collaboration, "The IceCube Neutrino Observatory: Instrumentation and Online Systems", <u>JINST</u> 12 (2017) P03012 [1612.05093].
- [588] A. M. Brown, M. Bagheri, M. Doro, E. Gazda, D. Kieda, C. Lin et al., "Trinity: an imaging air Cherenkov telescope to search for Ultra-High-Energy neutrinos", PoS ICRC2021 (2021) 1179.
- [589] RNO-G Collaboration, "The Radio Neutrino Observatory Greenland (RNO-G)", PoS ICRC2021 (2021) 001.
- [590] R. Gandhi, C. Quigg, M. H. Reno and I. Sarcevic, "Ultrahigh-energy neutrino interactions", <u>Astropart. Phys.</u> 5 (1996) 81 [hep-ph/9512364].
- [591] R. Gandhi, C. Quigg, M. H. Reno and I. Sarcevic, "Neutrino interactions at ultrahigh-energies", <u>Phys. Rev. D</u> 58 (1998) 093009 [hep-ph/9807264].
- [592] M. H. Reno, J. F. Krizmanic and T. M. Venters, "Cosmic tau neutrino detection via Cherenkov signals from air showers from Earth-emerging taus", Phys. Rev. D 100 (2019) 063010 [1902.11287].
- [593] F. Halzen and S. R. Klein, "IceCube: An Instrument for Neutrino Astronomy", <u>Rev. Sci. Instrum.</u> 81 (2010) 081101 [1007.1247].

- [594] T. Gaisser and F. Halzen, "IceCube", Ann. Rev. Nucl. Part. Sci. 64 (2014) 101.
- [595] A. N. Otte, A. M. Brown, M. Doro, A. Falcone, J. Holder, E. Judd et al., "Trinity: An Air-Shower Imaging Instrument to detect Ultrahigh Energy Neutrinos", 1907.08727.
- [596] ARIANNA Collaboration, "Targeting ultra-high energy neutrinos with the ARIANNA experiment", <u>Adv.</u> Space Res. **64** (2019) 2595 [1903.01609].
- [597] P. Allison et al., "Design and Initial Performance of the Askaryan Radio Array Prototype EeV Neutrino Detector at the South Pole", Astropart. Phys. 35 (2012) 457 [1105.2854].
- [598] ARA Collaboration, "A low-threshold ultrahigh-energy neutrino search with the Askaryan Radio Array", 2202.07080.
- [599] V. Scotti and G. Osteria, "The EUSO-SPB2 mission", <u>Nuclear Instruments and Methods in Physics Research</u> Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 958 (2020) 162164.
- [600] GRAND Collaboration, "GRANDProto300 experiment: a pathfinder with rich astroparticle and radio-astronomy science case", PoS ICRC2019 (2020) 233 [1909.04893].
- [601] RADAR ECHO TELESCOPE Collaboration, "The Radar Echo Telescope for Cosmic Rays: Pathfinder experiment for a next-generation neutrino observatory", Phys. Rev. D **104** (2021) 102006 [2104.00459].
- [602] ANTARES Collaboration, "ANTARES: the first undersea neutrino telescope", <u>Nucl. Instrum. Meth. A</u> 656 (2011) 11 [1104.1607].
- [603] I. Safa, A. Pizzuto, C. A. Argüelles, F. Halzen, R. Hussain, A. Kheirandish et al., "Observing EeV neutrinos through Earth: GZK and the anomalous ANITA events", JCAP 01 (2020) 012 [1909.10487].
- [604] G. A. Askar'yan, "Excess negative charge of an electron-photon shower and its coherent radio emission", <u>Zh.</u> Eksp. Teor. Fiz. **41** (1961) 616.
- [605] D. García-Fernández, A. Nelles and C. Glaser, "Signatures of secondary leptons in radio-neutrino detectors in ice", Phys. Rev. D 102 (2020) 083011 [2003.13442].
- [606] K.-C. Lai, C.-C. Chen and P. Chen, "The Strategy of Discrimination between Flavors for Detection of Cosmogenic Neutrinos", Nucl. Phys. B Proc. Suppl. 246-247 (2014) 95 [1303.1949].
- [607] ARA Collaboration, "First Constraints on the Ultra-High Energy Neutrino Flux from a Prototype Station of the Askaryan Radio Array", Astropart. Phys. 70 (2015) 62 [1404.5285].
- [608] A. Anker et al., "White Paper: ARIANNA-200 high energy neutrino telescope", 2004.09841.
- [609] K. D. de Vries, K. Hanson and T. Meures, "On the feasibility of RADAR detection of high-energy neutrino-induced showers in ice", Astropart. Phys. 60 (2015) 25 [1312.4331].
- [610] S. Prohira and D. Besson, "Particle-level model for radar based detection of high-energy neutrino cascades", Nucl. Instrum. Meth. A 922 (2019) 161 [1710.02883].
- [611] ANITA Collaboration, "The Antarctic Impulsive Transient Antenna Ultra-high Energy Neutrino Detector Design, Performance, and Sensitivity for 2006-2007 Balloon Flight", Astropart. Phys. 32 (2009) 10 [0812.1920].
- [612] S. Buitink, O. Scholten, J. Bacelar, R. Braun, A. de Bruyn, H. Falcke et al., "Constraints on the flux of Ultra-High Energy neutrinos from WSRT observations", Astron. Astrophys. 521 (2010) A47 [1004.0274].
- [613] G. A. Askaryan and B. Dolgoshein, "Acoustic registration of high-energy neutrinos", JETP Lett. 25 (1977) 213.
- [614] T. Bowen, "Sonic Particle Detection", in <u>International Cosmic Ray Conference</u>, vol. 6 of <u>International Cosmic Ray Conference</u>, p. 277, Jan., 1977.
- [615] R. Nahnhauer, "Acoustic particle detection from early ideas to future benefits", <u>Nucl. Instrum. Meth. A</u> 662 (2012) S20 [1010.3082].
- [616] R. Lahmann, "History of acoustic neutrino detection", EPJ Web Conf. 216 (2019) 01001.
- [617] F. Halzen and D. Saltzberg, "Tau-neutrino appearance with a 1000 megaparsec baseline", <u>Phys. Rev. Lett.</u> 81 (1998) 4305 [hep-ph/9804354].
- [618] S. I. Dutta, M. H. Reno and I. Sarcevic, "Secondary neutrinos from tau neutrino interactions in earth", <u>Phys.</u> Rev. D 66 (2002) 077302 [hep-ph/0207344].

- [619] O. B. Bigas, O. Deligny, K. Payet and V. Van Elewyck, "UHE tau neutrino flux regeneration while skimming the Earth", Phys. Rev. D 78 (2008) 063002 [0806.2126].
- [620] J. Álvarez-Muñiz, W. R. Carvalho, A. L. Cummings, K. Payet, A. Romero-Wolf, H. Schoorlemmer et al., "Comprehensive approach to tau-lepton production by high-energy tau neutrinos propagating through the Earth", Phys. Rev. D 97 (2018) 023021 [1707.00334].
- [621] D. Fargion, "Discovering Ultra High Energy Neutrinos by Horizontal and Upward τ Air-Showers: Evidences in Terrestrial Gamma Flashes?", Astrophys. J. **570** (2002) 909 [astro-ph/0002453].
- [622] A. L. Cummings, R. Aloisio and J. F. Krizmanic, "Modeling of the Tau and Muon Neutrino-induced Optical Cherenkov Signals from Upward-moving Extensive Air Showers", <u>Phys. Rev. D</u> 103 (2021) 043017 [2011.09869].
- [623] TELESCOPE ARRAY Collaboration, "Search for Ultra-High-Energy Neutrinos with the Telescope Array Surface Detector", J. Exp. Theor. Phys. 131 (2020) 255 [1905.03738].
- [624] J. Nam et al., "Design and implementation of the TAROGE experiment", <u>Int. J. Mod. Phys. D</u> 25 (2016) 1645013.
- [625] C. Aramo, A. Insolia, A. Leonardi, G. Miele, L. Perrone, O. Pisanti et al., "Earth-skimming UHE Tau neutrinos at the fluorescence detector of Pierre Auger observatory", Astropart. Phys. 23 (2005) 65 [astro-ph/0407638].
- [626] D. Góra, E. Bernardini and A. Kappes, "Searching for tau neutrinos with Cherenkov telescopes", <u>Astropart.</u> Phys. **61** (2014) 12 [1402.4243].
- [627] D. Góra and E. Bernardini, "Detection of tau neutrinos by Imaging Air Cherenkov Telescopes", <u>Astropart.</u> Phys. 82 (2016) 77 [1606.01676].
- [628] MAGIC Collaboration, "Limits on the flux of tau neutrinos from 1 PeV to 3 EeV with the MAGIC telescopes", Astropart. Phys. 102 (2018) 77 [1805.02750].
- [629] T. C. Weekes et al., "Observation of TeV gamma rays from the Crab nebula using the atmospheric Cerenkov imaging technique", Astrophys. J. 342 (1989) 379.
- [630] A. N. Otte, "Studies of an air-shower imaging system for the detection of ultrahigh-energy neutrinos", <u>Phys.</u> Rev. D 99 (2019) 083012 [1811.09287].
- [631] R. Dagkesamanskii and I. Zheleznykh, "A radio astronomy method of detecting neutrinos and other superhigh-energy elementary particles", Sov. Phys. JETP Lett. 50 (1989) 259.
- [632] J. D. Bray et al., "Lunar detection of ultra-high-energy cosmic rays and neutrinos", in Advancing Astrophysics with the Square Kilometre Array, 8, 2014, 1408.6069.
- [633] J. D. Bray, R. D. Ekers, P. Roberts, J. E. Reynolds, C. W. James, C. J. Phillips et al., "Limit on the ultrahigh-energy neutrino flux from lunar observations with the Parkes radio telescope", <u>Phys. Rev. D</u> 91 (2015) 063002 [1502.03313].
- [634] S. ter Veen, S. Buitink, H. Falcke, C. W. James, M. Mevius, O. Scholten et al., "A new limit on the Ultra-High-Energy Cosmic-Ray flux with the Westerbork Synthesis Radio Telescope", <u>Phys. Rev. D</u> 82 (2010) 103014 [1010.6061].
- [635] T. R. Jaeger, R. L. Mutel and K. G. Gayley, "Project RESUN, a radio EVLA search for UHE neutrinos", Astropart. Phys. 34 (2010) 293 [0910.5949].
- [636] S. Buitink, H. Falcke, C. James, M. Mevius, O. Scholten, K. Singh et al., "Constraints on ultra-high-energy neutrino flux from radio observations of the Moon", Astrophys. Space Sci. Trans. 8 (2012) 29.
- [637] G. K. Krampah et al., "The NuMoon Experiment: Lunar Detection of Cosmic Rays and Neutrinos with LOFAR", PoS ICRC2021 (2021) 1148.
- [638] C. W. James et al., "The lunar Askaryan technique with the Square Kilometre Array", Pos ICRC2015 (2015) 291 [1608.02408].
- [639] J. Bray et al., "Lunar detection of ultra-high-energy cosmic rays and neutrinos with the Square Kilometre Array", PoS AASKA14 (2015) 144.

- [640] C. W. James et al., "Overview of lunar detection of ultra-high energy particles and new plans for the SKA", EPJ Web Conf. 135 (2017) 04001 [1704.05336].
- [641] J. D. Bray, "The sensitivity of past and near-future lunar radio experiments to ultra-high-energy cosmic rays and neutrinos", Astropart. Phys. 77 (2016) 1 [1601.02980].
- [642] ICECUBE Collaboration, "Searching for neutrino transients below 1 TeV with IceCube", Pos ICRC2021 (2021) 1131 [2108.01530].
- [643] ICECUBE Collaboration, "Probing neutrino emission at GeV energies from compact binary mergers with the IceCube Neutrino Observatory", 2105.13160.
- [644] ICECUBE Collaboration, "A Search for Neutrino Sources with Cascade Events in IceCube", Pos ICRC2021 (2021) 1150 [2107.09103].
- [645] ICECUBE Collaboration, "A Search for Time-dependent Astrophysical Neutrino Emission with IceCube Data from 2012 to 2017", Astrophys. J. 911 (2021) 67 [2012.01079].
- [646] ICECUBE Collaboration, "A Search for Neutrino Emission from Fast Radio Bursts with Six Years of IceCube Data", Astrophys. J. 857 (2018) 117 [1712.06277].
- [647] ICECUBE Collaboration, "Neutrino Interferometry for High-Precision Tests of Lorentz Symmetry with IceCube", Nature Phys. 14 (2018) 961 [1709.03434].
- [648] ICECUBE Collaboration, "Search for sterile neutrino mixing using three years of IceCube DeepCore data", Phys. Rev. D 95 (2017) 112002 [1702.05160].
- [649] ICECUBE Collaboration, "Searches for Sterile Neutrinos with the IceCube Detector", Phys. Rev. Lett. 117 (2016) 071801 [1605.01990].
- [650] ICECUBE Collaboration, "eV-Scale Sterile Neutrino Search Using Eight Years of Atmospheric Muon Neutrino Data from the IceCube Neutrino Observatory", Phys. Rev. Lett. **125** (2020) 141801 [2005.12942].
- [651] ICECUBE Collaboration, "Searching for eV-scale sterile neutrinos with eight years of atmospheric neutrinos at the IceCube Neutrino Telescope", Phys. Rev. D 102 (2020) 052009 [2005.12943].
- [652] ICECUBE Collaboration, "Search for Multi-flare Neutrino Emissions in 10 yr of IceCube Data from a Catalog of Sources", Astrophys. J. Lett. 920 (2021) L45 [2109.05818].
- [653] ANTARES Collaboration, "Searches for point-like sources of cosmic neutrinos with 11 years of ANTARES data", PoS ICRC2019 (2020) 920 [1908.08248].
- [654] ICECUBE Collaboration, "Searches for and Characterization of Astrophysical Neutrinos using Starting Track Events in IceCube", PoS ICRC2021 (2021) 1130 [2107.09811].
- [655] ICECUBE Collaboration, "First all-flavor search for transient neutrino emission using 3-years of IceCube DeepCore data", JCAP 01 (2022) 027 [2011.05096].
- [656] ICECUBE Collaboration, "A muon-track reconstruction exploiting stochastic losses for large-scale Cherenkov detectors", JINST 16 (2021) P08034 [2103.16931].
- [657] R. Abbasi et al., "A Convolutional Neural Network based Cascade Reconstruction for the IceCube Neutrino Observatory", JINST 16 (2021) P07041 [2101.11589].
- [658] ICECUBE Collaboration, "A New Search for Neutrino Point Sources with IceCube", Pos ICRC2021 (2021) 1138 [2107.08700].
- [659] KM3NET Collaboration, "Sensitivity estimates for diffuse, point-like and extended neutrino sources with KM3NeT/ARCA", PoS ICRC2021 (2021) 1077.
- [660] KM3NET Collaboration, "The KM3NeT/ARCA detection capability to a diffuse flux of cosmic neutrinos", PoS ICRC2017 (2017) 998.
- [661] D. Gaggero, D. Grasso, A. Marinelli, A. Urbano and M. Valli, "The gamma-ray and neutrino sky: A consistent picture of Fermi-LAT, Milagro, and IceCube results", Astrophys. J. Lett. 815 (2015) L25 [1504.00227].
- [662] KM3NET Collaboration, "Detection Potential of the KM3NeT Detector for High-Energy Neutrinos from the Fermi Bubbles", Astropart. Phys. 42 (2013) 7 [1208.1226].

- [663] KM3NET Collaboration, "Real-time Multi-Messenger Analysis Framework of KM3NeT", Pos ICRC2021 (2021) 941.
- [664] BAIKAL-GVD Collaboration, "Deep underwater Cherenkov detector in Lake Baikal", JETP 161 (2022) 476.
- [665] BAIKAL-GVD Collaboration, "The Baikal-GVD neutrino telescope: search for high-energy cascades", PoS ICRC2021 (2021) 1144 [2108.01894].
- [666] BAIKAL-GVD Collaboration, "Performance of the muon track reconstruction with the Baikal-GVD neutrino telescope", PoS ICRC2021 (2021) 1080 [2108.11217].
- [667] A. D. Avrorin et al., "The optical module of Baikal-GVD", Phys. Part. Nucl. Lett. 13 (2016) 737.
- [668] A. V. Avrorin et al., "High-Energy Neutrino Follow-up at the Baikal-GVD Neutrino Telescope", <u>Astron. Lett.</u> 47 (2021) 94.
- [669] BAIKAL-GVD Collaboration, "Multi-messenger and real-time astrophysics with the Baikal-GVD telescope", PoS ICRC2021 (2021) 946 [2107.14472].
- [670] BAIKAL-GVD Collaboration, "Automatic data processing for Baikal-GVD neutrino observatory", PoS ICRC2021 (2021) 1040 [2107.13939].
- [671] P-ONE Collaboration, "The Pacific Ocean Neutrino Experiment", in 37th International Cosmic Ray Conference, 11, 2021, 2111.13133.
- [672] STRAW Collaboration, "STRAW (STRings for Absorption length in Water): pathfinder for a neutrino telescope in the deep Pacific Ocean", JINST 14 (2019) P02013 [1810.13265].
- [673] N. Bailly et al., "Two-year optical site characterization for the Pacific Ocean Neutrino Experiment (P-ONE) in the Cascadia Basin", Eur. Phys. J. C 81 (2021) 1071 [2108.04961].
- [674] ICECUBE Collaboration, "IceCube-Gen2: A Vision for the Future of Neutrino Astronomy in Antarctica", 1412.5106.
- [675] A. M. Brown, M. Bagheri, M. Doro, E. Gazda, D. Kieda, C. Lin et al., "Trinity: An Imaging Air Cherenkov Telescope to Search for Ultra-High-Energy Neutrinos", in <u>37th International Cosmic Ray Conference</u>, 9, 2021, 2109.03125.
- [676] A. Wang, C. Lin, N. Otte, M. Doro, E. Gazda, I. Taboada et al., "Trinity's Sensitivity to Isotropic and Point-Source Neutrinos", PoS ICRC2021 (2021) 1234 [2108.02751].
- [677] X. Bertou, P. Billoir, O. Deligny, C. Lachaud and A. Letessier-Selvon, "Tau neutrinos in the Auger Observatory: A New window to UHECR sources", Astropart. Phys. 17 (2002) 183 [astro-ph/0104452].
- [678] D. Fargion, P. G. De Sanctis Lucentini and M. De Santis, "Tau air showers from earth", <u>Astrophys. J.</u> 613 (2004) 1285 [hep-ph/0305128].
- [679] J. L. Feng, P. Fisher, F. Wilczek and T. M. Yu, "Observability of earth skimming ultrahigh-energy neutrinos", Phys. Rev. Lett. 88 (2002) 161102 [hep-ph/0105067].
- [680] S. Ogawa and M. Sasaki, "Galactic Bulge VHE Tau-neutrino and Gamma-ray Monitor with Ashra-1 and NTA detectors", PoS ICRC2021 (2021) 970.
- [681] J. Cortina, R. López-Coto and A. Moralejo, "MACHETE: A transit imaging atmospheric Cherenkov telescope to survey half of the very high energy γ-ray sky", Astropart. Phys. 72 (2016) 46 [1507.02532].
- [682] A. G. Vieregg, K. Bechtol and A. Romero-Wolf, "A Technique for Detection of PeV Neutrinos Using a Phased Radio Array", JCAP 02 (2016) 005 [1504.08006].
- [683] P. Allison et al., "Design and performance of an interferometric trigger array for radio detection of high-energy neutrinos", Nucl. Instrum. Meth. A 930 (2019) 112 [1809.04573].
- [684] PIERRE AUGER Collaboration, "Physics and astrophysics of ultra-high energy cosmic rays: recent results from the Pierre Auger Observatory", in <u>70th International conference on Nuclear physics and elementary particle physics:</u> Nuclear physics technologies, 12, 2020, 2012.12943.
- [685] ICECUBE, HAWC Collaboration, "IceCube Search for Galactic Neutrino Sources based on Very High Energy γ -ray Observations", J. Phys. Conf. Ser. **1468** (2020) 012081 [2001.08524].

- [686] L. A. Anchordoqui, H. Goldberg, F. Halzen and T. J. Weiler, "Neutrinos as a diagnostic of high energy astrophysical processes", Phys. Lett. B 621 (2005) 18 [hep-ph/0410003].
- [687] C. A. Argüelles, K. Farrag, T. Katori, R. Khandelwal, S. Mandalia and J. Salvado, "Sterile neutrinos in astrophysical neutrino flavor", JCAP 02 (2020) 015 [1909.05341].
- [688] ICECUBE Collaboration, "The IceCube Neutrino Observatory Contributions to ICRC 2017 Part II: Properties of the Atmospheric and Astrophysical Neutrino Flux", 1710.01191.
- [689] L. Gerhardt, S. Klein, T. Stezelberger, S. Barwick, K. Dookayka, J. Hanson et al., "A prototype station for ARIANNA: a detector for cosmic neutrinos", Nucl. Instrum. Meth. A 624 (2010) 85 [1005.5193].
- [690] ARIANNA Collaboration, "Improving sensitivity of the ARIANNA detector by rejecting thermal noise with deep learning", 2112.01031.
- [691] C. Glaser and S. W. Barwick, "An improved trigger for Askaryan radio detectors", <u>JINST</u> 16 (2021) T05001 [2011.12997].
- [692] S. W. Barwick et al., "Radio detection of air showers with the ARIANNA experiment on the Ross Ice Shelf", Astropart. Phys. 90 (2017) 50 [1612.04473].
- [693] A. Anker et al., "Neutrino vertex reconstruction with in-ice radio detectors using surface reflections and implications for the neutrino energy resolution", JCAP 11 (2019) 030 [1909.02677].
- [694] C. Glaser, A. Nelles, I. Plaisier, C. Welling, S. W. Barwick, D. García-Fernández et al., "NuRadioReco: A reconstruction framework for radio neutrino detectors", Eur. Phys. J. C 79 (2019) 464 [1903.07023].
- [695] C. Welling, C. Glaser and A. Nelles, "Reconstructing the cosmic-ray energy from the radio signal measured in one single station", JCAP 10 (2019) 075 [1905.11185].
- [696] G. Gaswint, "New results on angular reconstruction of deep pulser radio signals", PoS ICRC2019 (2020) 897.
- [697] ARIANNA Collaboration, "Cosmic-ray detection with and novel reconstruction algorithms for the ARIANNA experiment", PoS ICRC2019 (2020) 366.
- [698] ARIANNA Collaboration, "Probing the angular and polarization reconstruction of the ARIANNA detector at the South Pole", JINST 15 (2020) P09039 [2006.03027].
- [699] C. Glaser, S. McAleer, P. Baldi and S. W. Barwick, "Deep learning reconstruction of the neutrino energy with a shallow Askaryan detector", PoS ICRC2021 (2021) 1051.
- [700] G. G. Gaswint, <u>Quantifying the Neutrino Energy and Pointing Resolution of the ARIANNA Detector</u>, Ph.D. thesis, UC, Irvine, 2021. https://escholarship.org/uc/item/lbj9r6rb.
- [701] ARIANNA Collaboration, "Capabilities of ARIANNA: Neutrino Pointing Resolution and Implications for Future Ultra-high Energy Neutrino Astronomy", PoS ICRC2021 (2021) 1151.
- [702] ARIANNA Collaboration, "Measuring the Polarization Reconstruction Resolution of the ARIANNA Neutrino Detector with Cosmic Rays", 2112.01501.
- [703] P. Allison et al., "A low-threshold ultrahigh-energy neutrino search with the Askaryan Radio Array", 2202.07080.
- [704] ARA Collaboration, "Performance of two Askaryan Radio Array stations and first results in the search for ultrahigh energy neutrinos", Phys. Rev. D 93 (2016) 082003 [1507.08991].
- [705] P. Allison et al., "Long-baseline horizontal radio-frequency transmission through polar ice", <u>JCAP</u> **12** (2020) 009 [1908.10689].
- [706] P. Allison et al., "Measurement of the real dielectric permittivity ϵ_r of glacial ice", <u>Astropart. Phys.</u> 108 (2019) 63 [1712.03301].
- [707] A. Connolly, "Impact of biaxial birefringence in polar ice at radio frequencies on signal polarizations in ultra-high energy neutrino detection", 2110.09015.
- [708] J. A. Aguilar et al., "Reconstructing the neutrino energy for in-ice radio detectors: A study for the Radio Neutrino Observatory Greenland (RNO-G)", Eur. Phys. J. C 82 (2022) 147 [2107.02604].

- [709] C. Glaser, D. García-Fernández and A. Nelles, "Prospects for neutrino-flavor physics with in-ice radio detectors", PoS ICRC2021 (2021) 1231.
- [710] K.-H. Kampert and M. Unger, "Measurements of the Cosmic Ray Composition with Air Shower Experiments", Astropart. Phys. 35 (2012) 660 [1201.0018].
- [711] A. Zeolla et al., "Modeling and Validating RF-Only Interferometric Triggering with Cosmic Rays for BEACON", PoS ICRC2021 (2021) 1072.
- [712] BEACON Collaboration, "Searching for RF-Only Triggered Cosmic Ray Events with the High-Elevation BEACON Prototype", PoS ICRC2021 (2021) 1084.
- [713] V. Decoene, N. Renault-Tinacci, O. Martineau-Huynh, D. Charrier, K. Kotera, S. Le Coz et al., "Radio-detection of neutrino-induced air showers: the influence of topography", <u>Nucl. Instrum. Meth. A</u> 986 (2021) 164803 [1903.10466].
- [714] V. Decoene, O. Martineau-Huynh, M. Tueros and S. Chiche, "A reconstruction procedure for very inclined extensive air showers based on radio signals", PoS ICRC2021 (2021) 211 [2107.03206].
- [715] D. Ardouin et al., "First detection of extensive air showers by the TREND self-triggering radio experiment", Astropart. Phys. 34 (2011) 717 [1007.4359].
- [716] M. M. Block, L. Durand and P. Ha, "Connection of the virtual $\gamma^* p$ cross section of ep deep inelastic scattering to real γp scattering, and the implications for νN and ep total cross sections", Phys. Rev. **D89** (2014) 094027 [1404.4530].
- [717] O. Scholten, S. Buitink, J. Bacelar, R. Braun, A. G. de Bruyn, H. Falcke et al., "Improved flux limits for neutrinos with energies above 10²² eV from observations with the Westerbork Synthesis Radio Telescope", Phys. Rev. Lett. 103 (2009) 191301 [0910.4745].
- [718] V. A. Ryabov, V. A. Chechin, G. A. Gusev and K. T. Maung, "Prospects for ultrahigh-energy particle observation based on the lunar orbital LORD space experiment", Adv. Space Res. 58 (2016) 464.
- [719] K. Fang and B. D. Metzger, "High-Energy Neutrinos from Millisecond Magnetars formed from the Merger of Binary Neutron Stars", Astrophys. J. 849 (2017) 153 [1707.04263].
- [720] J. Eser, A. V. Olinto and L. Wiencke, "Science and mission status of EUSO-SPB2", Proceedings of 37th International Cosmic Ray Conference — PoS(ICRC2021) (2021).
- [721] A. Cummings, R. Aloisio, J. Eser and J. Krizmanic, "Modeling the optical cherenkov signals by cosmic ray extensive air showers directly observed from suborbital and orbital altitudes", Physical Review D 104 (2021).
- [722] POEMMA, JEM-EUSO Collaboration, "Astrophysical Implications of Neutrino Target-of-Opportunity Observations with Space-based and Suborbital Optical Cherenkov Detectors", PoS ICRC2021 (2021) 977.
- [723] POEMMA, JEM-EUSO Collaboration, "EAS Optical Cherenkov signatures of tau neutrinos for space and suborbital detectors", PoS ICRC2021 (2021) 1201.
- [724] ANITA Collaboration, "New Limits on the Ultra-high Energy Cosmic Neutrino Flux from the ANITA Experiment", Phys. Rev. Lett. 103 (2009) 051103 [0812.2715].
- [725] ANITA Collaboration, "Observation of Ultra-high-energy Cosmic Rays with the ANITA Balloon-borne Radio Interferometer", Phys. Rev. Lett. 105 (2010) 151101 [1005.0035].
- [726] H. Schoorlemmer et al., "Energy and Flux Measurements of Ultra-High Energy Cosmic Rays Observed During the First ANITA Flight", Astropart. Phys. 77 (2016) 32 [1506.05396].
- [727] ANITA Collaboration, "Ultra-high Energy Air Showers Observed by ANITA-IV", 2008.05690.
- [728] ANITA Collaboration, "Observational Constraints on the Ultra-high Energy Cosmic Neutrino Flux from the Second Flight of the ANITA Experiment", Phys. Rev. D 82 (2010) 022004 [1003.2961].
- [729] ANITA Collaboration, "Constraints on the diffuse high-energy neutrino flux from the third flight of ANITA", Phys. Rev. D 98 (2018) 022001 [1803.02719].
- [730] ANITA-II Collaboration, "Ultra-Relativistic Magnetic Monopole Search with the ANITA-II Balloon-borne Radio Interferometer", Phys. Rev. D 83 (2011) 023513 [1008.1282].

- [731] A. Romero-Wolf et al., "An interferometric analysis method for radio impulses from ultra-high energy particle showers", Astropart. Phys. 60 (2015) 72.
- [732] ANITA Collaboration, "Upward-Pointing Cosmic-Ray-like Events Observed with ANITA", PoS ICRC2017 (2018) 935 [1810.00439].
- [733] PIERRE AUGER Collaboration, "The upgrade of the Pierre Auger Observatory with the Scintillator Surface Detector", PoS ICRC2021 (2021) 251.
- [734] PIERRE AUGER Collaboration, "A Large Radio Detector at the Pierre Auger Observatory Measuring the Properties of Cosmic Rays up to the Highest Energies", PoS ICRC2019 (2021) 395.
- [735] PIERRE AUGER Collaboration, "Antennas for the Detection of Radio Emission Pulses from Cosmic-Ray", JINST 7 (2012) P10011 [1209.3840].
- [736] K. F. Daniel and L. I, "Radiation from cosmic ray air showers", Proc. R. Soc. Lond. 289 (1966) 206.
- [737] PIERRE AUGER Collaboration, "Observation of inclined EeV air showers with the radio detector of the Pierre Auger Observatory", JCAP 10 (2018) 026 [1806.05386].
- [738] PIERRE AUGER Collaboration, "The depth of the shower maximum of air showers measured with AERA", PoS ICRC2021 (2021) 387.
- [739] PIERRE AUGER Collaboration, "New Electronics for the Surface Detectors of the Pierre Auger Observatory", PoS ICRC2019 (2021) 370.
- [740] PIERRE AUGER Collaboration, "Reconstruction of Events Recorded with the Water-Cherenkov and Scintillator Surface Detectors of the Pierre Auger Observatory", PoS ICRC2021 (2021) 218.

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: