

## The magnetic topology of the inverse Evershed flow

A. Prasad<sup>1,2,3</sup>, M. Ranganathan<sup>4,5</sup>, C. Beck<sup>6</sup>, D. P. Choudhary<sup>4</sup>, and Q. Hu<sup>3,7</sup>

- Rosseland Centre for Solar Physics, University of Oslo, Postboks 1029 Blindern, 0315 Oslo, Norway e-mail: avijeet.prasad@astro.uio.no
- <sup>2</sup> Institute of Theoretical Astrophysics, University of Oslo, Postboks 1029 Blindern, 0315 Oslo, Norway
- <sup>3</sup> Center for Space Plasma & Aeronomic Research, The University of Alabama in Huntsville, Huntsville, AL 35899, USA
- <sup>4</sup> Department of Physics & Astronomy, California State University, Northridge, CA 91330-8268, USA
- <sup>5</sup> Institute for Particle and Astrophysics, ETH, 8049 Zürich, Switzerland
- <sup>6</sup> National Solar Observatory (NSO), 3665 Discovery Drive, Boulder, CO 80303, USA
- Department of Space Science, The University of Alabama in Huntsville, Huntsville, AL 35899, USA

Received 5 November 2021 / Accepted 28 February 2022

#### **ABSTRACT**

Context. The inverse Evershed flow (IEF) is a mass motion towards sunspots at chromospheric heights.

Aims. We combined high-resolution observations of NOAA 12418 from the Dunn Solar Telescope and vector magnetic field measurements from the Helioseismic and Magnetic Imager (HMI) to determine the driver of the IEF.

Methods. We derived chromospheric line-of-sight (LOS) velocities from spectra of H and Ca IRI The HMI data were used in a non-force-free magnetic field extrapolation to track closed field lines near the sunspot in the active region. We determined their length and height, located their inner and outer foot points, and derived flow velocities along them.

Results. The magnetic field lines related to the IEF reach on average a height of 3 megameter (Mm) over a length of 13 Mm. The inner (outer) foot points are located at 1.2 (1.9) sunspot radii. The average field strength dierence B between inner and outer foot points is +400 G. The temperature dierence T is anti-correlated with B with an average value of 100 K. The pressure dierence p is dominated by B and is primarily positive with a driving force towards the inner foot points of 1.7 kPa on average. The velocities predicted from p reproduce the LOS velocities of 2–10 km s <sup>1</sup> with a square-root dependence.

Conclusions. We find that the IEF is driven along magnetic field lines connecting network elements with the outer penumbra by a gas pressure dierence that results from a dierence in field strength as predicted by the classical siphon flow scenario.

**Key words.** Sun: chromosphere – Sun: photosphere – sunspots

#### 1. Introduction

The chromospheric inverse Evershed flow (IEF; Evershed 1909) transports material into sunspots along magnetic field lines that connect the boundary of the moat cell with the outer penumbra. These closed loops and the flows along them have a lifetime of a few tens of minutes to more than 1h (Georgakilas & Christopoulou 2003; Beck & Choudhary 2020). The flow velocities of about 2 9 km s <sup>1</sup> (Dialetis et al. 1985; Dere et al. 1990; Beck et al. 2020) carry the material along loops that ascend to a height of a few megameter (Mm) (Maltby 1975; Beck et al. 2020). Flows in dark super-penumbral fibrils were found to be faster than those in bright ones, with fluctuations on timescale of about 25 mins (Georgakilas et al. 2003; Georgakilas & Christopoulou 2003). At the inner foot points in the penumbra, the magnetic field lines return to the photosphere with an average angle of about 65 to the local vertical (Haugen 1969; Beck & Choudhary 2019). As the flow speed at the inner foot point exceeds the photospheric sound speed, the flow terminates in a stationary shock front that heats the lower chromosphere (Thomas & Montesinos 1991; Beck et al. 2014; Choudhary & Beck 2018).

The magnetic and thermodynamic properties and the lifetime of IEF channels comply with a siphon flow scenario (Meyer & Schmidt 1968; Thomas 1988; Montesinos & Thomas 1997), where a stationary flow can be driven along magnetic fields lines (MFLs) that have two foot points (FPs) of dif-

ferent magnetic field strength. If the two FPs have the same total pressure, the dierence in field strength causes a dier-ence in gas pressure as the necessary mechanical driving force of the flow. Observational evidence of siphon flows in the solar atmosphere was reported by, for example, Rueedi et al. (1992), Uitenbroek et al. (2006) and Bethge et al. (2012), but flows along MFLs that connect FPs of unequal field strength can also be driven by other physical processes (Sigwarth et al. 1998). A major problem in diagnosing siphon flows in solar observations is the unequivocal identification of the two FPs — needed in order to determine their field strength —, which requires knowl-edge of the magnetic connectivity.

Narrow filaments in the photosphere and fibrils in the chromosphere are known to trace MFLs connecting the central regions of sunspots with their surroundings (Frazier 1972; Zirin 1972; Schad et al. 2013; Beck & Choudhary 2019), but it is especially dicult to locate the outer FPs for the IEF channels, as their signature in velocity and intensity fades into the background (see, e.g., Beck et al. 2020). A more direct tool to derive the magnetic connectivity of sunspots or whole active regions is magnetic field extrapolation, whereby photospheric (vector) magnetograms are used to infer the coronal magnetic field under certain assumptions. For example, the assumption that the magnetic pressure dominates over the plasma pressure (plasma << 1) allows one to neglect all non-magnetic forces and assume the Lorentz force to be zero. This approach leads to force-free fields, which have been widely used in the solar

community (e.g., Wiegelmann 2008; Wiegelmann & Sakurai 2012). However, it was pointed out by Gary (2001) that the plasma can be of the order of unity in solar photosphere, where the magnetic field measurements are taken, thus emphasizing the need for a dierent approach that incorporates non-force-free eects. One such alternative is the non-force-free-field (NFFF) extrapolation technique (Hu & Dasgupta 2008; Hu et al. 2008, 2010) based on the principle of the minimum energy dissipation rate (Bhattacharyya et al. 2007), where the magnetic field is expressed as the superposition of one potential field and two (constant ) linear force-free fields with distinct parameters. Such NFFF extrapolations have been used in many recent studies to model flaring active regions and coronal iets (Navak et al. 2019; Liu et al. 2020; Yalim et al. 2020; Prasad et al. 2020). Interested readers are referred to Appendix A of Liu et al. (2020), which provides a detailed discussion on the applicability of the NFFF method.

In this paper, we investigate the magnetic field properties of IEF channels using a combination of high-resolution observations of chromospheric spectral lines to trace the flow channels and an NFFF extrapolation to determine the magnetic connectivity. The use of extrapolated field lines allows us to identify the foot points and determine the heights of flow channels that connect the opposite-polarity patches in the magnetograms. As the field gradient and the height of the flow channels play a crucial role in driving the siphon flows, we employ these quantities to study properties such as the strength of mass motions in IEFs. We use the same definition as in Beck et al. (2020), namely that super-penumbral, roughly radially oriented elongated fibrils that connect the penumbra with the super-penumbral boundary and that exhibit a significant flow velocity constitute IEF channels.

Section 2 describes the observations used. Our analysis methods are explained in Sect. 3 and the Appendices A to D. The analysis results are given in Sect. 4. Section 5 discusses the findings, while Sect. 6 provides our conclusions.

## 2. Observations

We observed the decaying active region (AR) NOAA 12418 on 16 September 2015 with the Interferometric Bldimensional Spectrometer (IBIS; Cavallini 2006; Reardon & Cavallini 2008) and the Facility InfraRed Spectropolarimeter (FIRS; Jaeggli et al. 2010) at the Dunn Solar Telescope (DST; Dunn 1969; Dunn & Smartt 1991). The field of view (FOV) covered the isolated leading sunspot of the AR located at about  $x; y = 500^{00}$ ;  $340^{00}$  at a heliocentric angle of 43. The DST observations are described in detail in Beck & Choudhary (2020), and so we only give a short summary here.

IBIS was used to sequentially obtain 400 spectral scans of the two chromospheric spectral lines of H at 656 nm and Ca jį IR at 854.2 nm with about 30 wavelength points each between 14:42 and 15:56 UT. The circular IBIS FOV had a diameter of 95<sup>00</sup> at a spatial sampling of 0:095 pixel <sup>1</sup>. For the current study, we only used the six spectral scans between 14:48 and 15:48 UT that were simultaneous to full-vector obser-vations with the Helioseismic and Magnetic Imager (HMI; Scherrer et al. 2012) at the 12 min cadence of the latter. The IBIS and HMI data were complemented with full-disk images at 171, 304, and 1700 Å from the Atmospheric Imaging Assembly (AIA; Lemen et al. 2012) on board the Solar Dynamics Observa-tory (SDO; Pesnell et al. 2012) and H images from the Global Oscillation Network Group (GONG; Harvey et al. 1996).

Figure 1 shows an overview of the AR in dierent quanti-ties.

The panels in the middle column correspond to the box used

for the magnetic field extrapolation, while the rightmost column corresponds to the IBIS FOV after de-projection. The AR was located in the southeast quadrant of the Sun. The leading positive polarity contained a single, round sunspot, while the trailing negative polarity had decayed to more diuse plage regions. There was no significant amount of magnetic flux of negative polarity to the west of the sunspot.

The H, and AIA 171, and 304 Å images show a large filament of about 200 Mm length towards the east and a smaller filament of 75 Mm length towards the north starting from the outer penumbra of the sunspot. These two filaments are located above polarity inversion lines and are marked with red arrows in Fig. 1. We marked a further two, shorter filaments with blue arrows in Figs. 1 and 4, where the relation to a neutral line is less clear. All of those filaments dier from IEF channels in that they show much stronger absorption, a larger lateral width, an extended velocity signature, and a longer lifetime without any obvious changes over the one-hour duration of the observations (see also Fig. 12 of Beck & Choudhary 2020). They are likely to correspond to the topmost layer of the solar chromosphere that attains enough mass and hence opacity to cause such absorption (Leenaarts & Carlsson 2015).

The AIA 171 Å image shows some closed loops from the sunspot to the trailing plage, but no obvious loops towards the west. The symmetry line of the sunspot with roughly zero line-of-sight (LOS) velocities due to the projection eects on the LOS makes an angle of about 45 going from the northeast to the southwest. The IEF channels are clearly seen in the H line-core intensity and velocity maps, with a roughly even distribution in azimuth around the sunspot. The downflow patches on the center side are more pronounced and isolated than those on the limb side, where the end points of the IEF channels to some extent fall on the neutral line of LOS velocities.

## 3. Data analysis

The majority of the data analysis and the alignment of the high-resolution and full-disk data follow common procedures and are described in detail in Appendices A to D. We only summarize the steps and their outcome here.

From the HMI data, we retrieved the photospheric LOS velocities while compensating the solar rotation across the large FOV (Appendix A.1). Chromospheric LOS velocities were derived from the H and Ca jj IR 854 nm spectra from IBIS using a bi-sector method. The average velocities across the IBIS FOV were set to zero. Both lines were found to yield very similar values and can therefore be used synonymously (Appendix A.2). The HMI continuum intensity was normalized to unity in the quiet Sun and then converted to temperature using the Planck function (Appendix B). The HMI vector magnetograms were extrapolated using the NFFF method to retrieve the chromospheric and coronal magnetic field in a three-dimensional (3D) volume (Appendix C). The ground-based high-resolution and space-based full-disk data were de-projected to correct for the geometrical foreshortening and subsequently aligned to each other with the HMI continuum intensity image as a reference (Appendix D). As an estimate of the true velocity magnitude, the chromospheric LOS velocities were de-projected onto the magnetic field vector in the magnetic field extrapolation at a height of 1 Mm (Appendix D.2) to remove the LOS projection eects assuming field-aligned flows. The data analysis steps that are less common are described in the following two sections.




Fig. 1. Overview figure showing the location of AR NOAA 12418 on 16 September 2015. Left column, top to bottom: Full-disk images in HMI  $I_c$ ,  $B_z$ , GONG H, AIA 304 Å, and AIA 171 Å. Middle column: magnification of the black rectangle (371 Mm 186 Mm) in the left column. The HMI  $I_c$  has been exchanged for the derived temperature instead. Right column: magnification of the black rectangle marked in the middle column (94 Mm 70 Mm). Lower three panels: H line-core intensity and velocity, and the continuum intensity from the IBIS high-resolution spectra. The white dashed line in the H velocity map marks the symmetry line of the sunspot. The GONG H image in the middle column is not de-projected in contrast to all other panels in the right two columns. The red and blue arrows mark examples of H filaments as opposite to IEF channels.

### 3.1. Pressure balance equation

In the simplest approximation of magneto-hydrostatic equilibrium for the ionized plasma of the solar atmosphere in the presence of magnetic fields, the total pressure  $\textbf{p}_{tot}$  is given by the addition of the magnetic pressure  $\textbf{p}_{mag}$  and gas pressure  $\textbf{p}_{gas}$  by (e.g., Steiner et al. 1986; Thomas 1988)

$$p_{tot} = p_{mag} + p_{gas} = \frac{B^2}{2_0} + R.T;$$
 (1)

with the magnetic field B in Tesla T, the magnetic permeability constant  $_0$ = 1:26 10  $^6$  H m  $^1$ , the gas density in kg m  $^3$ , the mean molecular weight of the Sun = 1:3 g mol  $^1$ , the uni-

versal gas constant R =  $8:314 \text{ kg m}^2 \text{ s}^{-2} \text{ K}^{-1} \text{ mol}^{-1}$ , and the gas temperature T in K.

Two points, here represented as "1" and "2", in the solar atmosphere that are connected by MFLs or that are in direct proximity have magnetic and gas pressure dierences p mag and pgas of

$$p_{mag} = \frac{B_{\frac{1}{2}} \frac{1}{2} \frac{2}{and}}{2 \frac{1}{0} p_{gas}} = \frac{R(_{1}T_{1} _{2}T_{2})}{7}.$$
 (2)

with the magnetic field strengths  $B_i$ , temperatures  $T_i$ , and densities with i=1;2.

For spatially separated points, a dierence in gas pressure can drive a mass flow along a connecting field line. For  $B_1 > B_2$ 

Table 1. HSRA values at a few selected heights.

Z	log	Т	gas	р
km	-	K	10 <sup>7</sup> g cm <sup>3</sup>	kPa
77	1.3	9390	3.24	20.01
25	0.3	7140	3.36	15.40
11	0:1	6200	3.06	12.17
23	0:2	6035	2.89	11.20
35	0:3	5890	2.70	10.21
49	0:4	5765	2.50	9.24
63	0:5	5650	2.29	8.31
138	1	5160	1.38	4.56
283	2	4660	0.43	1.27

and total pressure equilibrium,  $p_{gas}^1$  is smaller than  $p_{gas}^2$ , and the flow moves from the location with lower to higher field strength, which is called a siphon flow (Meyer & Schmidt 1968; Cargill & Priest 1980). For  $T_1 < T_2$  at equal density, the flow goes from the point with higher to the one with lower temperature regardless of total pressure equilibrium.

As it is not instantly clear whether or not total pressure equilibrium is valid over large distances, we additionally defined the eective gas pressure dierence  $p_e$  by

$$p_e = \frac{B_1^2 \quad B^2}{2_0} + R(T_2 \quad T_1)$$
 (3)

as the eective driving force of flows along connecting magnetic field lines, assuming  $= -\frac{1}{2}$ .

If the lateral total pressure equilibrium holds over an extended spatial area, Eq. (1) predicts a linear relation between the temperature T and  $B^2$  as

$$T(B^2) = c \frac{1}{2 R^2} B^2;$$
 (4)

where the slope proportional to <sup>1</sup> is the only unknown if T and B are given, while c is proportional to the total pressure.

For a stationary gas flow driven by a pressure dierence p, a first-order estimate of the flow speed v can be derived through the kinetic energy density  $E_{kin} = \frac{1}{2} v^2$  as (Bethge et al. 2012)

$$v(p) = \frac{\frac{s}{2p}}{\frac{2p}{r}}$$
 (5)

while the potential energy E  $_{\rm pot}$  = g h allows one to estimate the maximal height that can be attained through

$$h(p) = \frac{p}{g'} \tag{6}$$

with the solar surface gravity  $g = 273 \,\mathrm{m \, s^{-2}}$ .

Finally, equating kinetic and potential energy density gives the maximal height in dependence of the flow speed as

$$h(v) = \frac{v^2}{2g}. \tag{7}$$

Table 1 lists the values of formation height, optical depth, temperature, gas density, and total pressure in the Harvard Smithsonian Reference Atmosphere (HSRA; Gingerich et al. 1971) over a height range from 100–300 km as an example of typical values of thermodynamic properties.

#### 3.2. Selection of field lines and IEF channels

The top row of Fig. 2 shows an overview of the magnetic field extrapolation results for the whole AR for the first and last magnetograms at 14:48 and 15:58 UT, respectively. The global magnetic topology is described below, but only some MFLs are relevant for the IEF channels. We therefore used two subsets of closed MFLs, defined in an automatic way and the other manually, for closer investigation.

#### 3.2.1. Closed field lines in and near the sunspot

The automatic selection of MFLs relevant for the IEF was based on a 150 150 pixels (54 Mm 54 Mm) square centered on the sunspot as initial seed points (see the black square in the left middle panel of Fig. 2) for the magnetogram at 14:48 UT. We derived the MFLs originating in that area with a simplified approach using the Interactive Data Language. We rejected all MFLs that were open, whose two-dimensional (2D) length in the horizontal plane (see Sect. 4.2.1) was shorter than 5.4 Mm and whose apex height exceeded 7.5 Mm. This provided the spatial positions of 3915 suitable seed points for the VAPOR software (Li et al. 2019), which we then used for all six time steps. The ad hoc limits on the length and height in this step were based on the visibility and appearance of the IEF channels in the Ca ij IR and H lines, that is, the spatial extent of fibrils and the formation heights of the lines.

From the VAPOR output, we then discarded all MFLs where the outer FP was located outside the de-projected aperture stop of IBIS and whose maximal height exceeded 7.5 Mm, but no longer required the minimal 2D length. Table 2 lists the numbers and percentages of the MFLs remaining sequentially with each criterion for the magnetogram at 14:48 UT. This left on average 3158 74 closed MFLs for one time-step with a total of 18948 closed MFLs. This automatically selected sample is labeled the "large sample" in the following. This sample was selected without considering the chromospheric velocities or intensities in the selection process in any way.

Figure 3 shows 1/33 of all the closed MFLs of the first and last magnetogram that were selected by this approach on top of the vertical component of the magnetic field B , the H continuum intensity, and the H LOS velocity. The majority of the closed MFLs extend to the limb side of the sunspot from south to north with a cluster towards the northeast, where the following plage was located, because of the lack of opposite-polarity patches towards the west. MFLs from the umbra and inner penumbra were either open or violated the other two criteria listed above.

#### 3.2.2. Manually selected IEF channels

The large sample of MFLs was complemented by a smaller sample of manually selected IEF channels. To this end, we used the line-core intensity and velocity maps in both H and Ca ii IR to define about 15–25 seed points in each of the six maps and located at the inner end of IEF channels for a total of 123 points. The points were chosen to be at about the end of flow channels or intensity fibrils and about evenly spaced around the sunspot in azimuth. Figure 4 shows these manually selected IEF channels for all six maps, with the MFLs resulting from the seed points. The manually selected sample is a subset of the large sample specifically chosen to catch flow channels and intensity fibrils. The MFLs towards the west are short and connect to moving magnetic features of opposite polarity close to the sunspot, while




Fig. 2. Large-scale magnetic field topology of AR 12418 on 16 September 2015 from NFFF extrapolations. Top row: full domain at 14:48 and 15:48 UT on top of  $B_z$  with open and closed MFLs. Middle (bottom) row: MFLs originating in the sunspot at 14:48 UT on top of  $B_z$  and AIA 1700 Å (AIA 304 Å and AIA 171 Å). The black square in the middle-left panel indicates the area of the initial seed points for the identification of closed MFLs. The FOV of the top row is slightly zoomed out to show the large-scale connectivity.

Table 2. Fraction of MFLs meeting combined selection criteria.

Total	Closed	L > 5:4 Mm	z < 7:5 Mm	Inside IBIS FOV
	7400	4500	2045	2454
22500	7103	4502	3915	3154
100%	32%	20%	17%	14%

most MFLs towards the east do not align with the large and dark intensity filaments marked with arrows in Fig. 4 that stand out prominently, especially in the Ca ii IR line-core images.

The two approaches of automatic and manual selection sample somewhat dierent targets. The automatic sample represents the magnetic topology of closed MFLs in the canopy of the sunspot with a good statistics regardless of the presence of IEF channels. This sample defines the characteristic topology in which IEF channels are found. The manual selection explicitly targets IEFs, that is, flow channels with a clear veloc-

ity signature, but with poorer statistics. Prior studies showed a rather isotropic distribution of IEF channels around sunspots, without a clear dierence between IEFs and their surround-ings in the photospheric magnetic field near their inner end points (Beck & Choudhary 2020). It is not immediately obvious whether the latter is still valid when following the MFLs away from the sunspot and to chromospheric heights. The comparison of the two samples can therefore be used to elucidate whether or not there are special conditions that favor the appearance of IEF channels by looking for eventual dierences in their average properties.

#### 4. Results

## 4.1. Magnetic connectivity throughout the active region

The top row of Fig. 2 shows the MFLs throughout the AR at 14:48 UT and 15:48 UT. The majority of closed MFLs connect




Fig. 3. Examples of automatically selected closed MFLs around the sunspot. Left (right) column: at 14:48 (15:48) UT. Only 1/33 fraction of all the MFLs are drawn here for clarity. The background images show from top to bottom  $B_z$ , the H continuum intensity, and the H LOS velocity.

from the sunspot to the trailing plage region. The MFLs west of the sunspot and east of the plage are primarily open and reach coronal heights. This part of the large-scale topology is reflected by the bright loops that are visible in the AIA 171 Å image in the bottom right panel of Fig. 2. The large filament extending from the sunspot to the east can be traced in the AIA 171 and 304 Å images, but the no MFL passes along its axis. The MFLs near the filament instead form an arcade of loops crossing the polarity inversion line approximately perpendicular to the filament spine. The closed MFLs originating inside or in proximity to the sunspot have an approximately radial orientation away from the sunspot center (Figs. 3 and 4).

The HMI magnetograms showed little to no evolution within the one-hour window of the observations (Figs. 2 and 3), hence the magnetic field extrapolation did not change significantly. The intensity and velocity maps in Fig. 4 show some variation with time; for example the intensity pattern to the west changes completely in the H line-core images, but all dark filaments to the east, north, and southwest and most flow channels persist (see also the temporal average of the same time series in Fig. 12 of Beck & Choudhary 2020).

## 4.2. Properties of closed magnetic field lines in the large sample

## 4.2.1. Topology of closed MFLs: length, apex height, shape, $r_{\text{inner}},$ and $r_{\text{outer}}$

For all closed MFLs, we defined the 3D loop length as the path length along the MFL between its two photospheric FPs, while the 2D loop length was defined as the length of their projection

onto the horizontal plane. The apex height was defined as the maximum height attained along the MFL. The distances of the inner and outer FP from the sunspot center was measured both in absolute units and fractional to the outer penumbral radius of r = 0.16:3 Mm, where the inner FP was defined as the seed point of the MFL inside the square centered on the sunspot.

The top panel of Fig. 5 shows the histogram of the apex height of all closed MFLs in the large sample. The distribution is roughly Gaussian between 0 and 7.5 Mm with a steeper drop o towards greater heights. The average and median values are 2.96 Mm and 2.72 Mm, respectively, far from the rejection threshold of 7.5 Mm in height. Those heights should fall well within the formation height range of the H line, but should exceed that of Ca ji IR. The histograms of the 2D and 3D loop length are fairly similar (bottom panel of Fig. 5) ranging from 0 to about 30 Mm length with average values of about 12 14 6 Mm.

There is a close relation between the 3D loop length and the apex height (Fig. 6), which holds in the same way for the 2D loop length (not shown). The correlation coecient between length and height is about 0.9 with a ratio L:H of about 4:1.

To determine the typical loop shape, we first normalized the 2D length L of each closed MFL to be from 0 to 1 by dividing it by the full length of the MFL. We then calculated the histograms of the height in 20 length bins of 0.05 extent each and sorted the result into a L-height array. Figure 7 shows a slightly smoothed 2D representation of these histograms of the height at each relative length point together with its average value. The closed MFLs form arched loops; these have a larger inclination to the vertical at the inner FPs and are nearly vertical at the outer FPs. The average apex height of 3.65 Mm is attained at a relative length of 0.8 close to the outer FP. Heights below 1 Mm and above 6 Mm are rarely attained (<16%) at any place. The shape roughly matches the parabolic loop of 2.45 Mm height (red line in Fig. 7) that was inferred as best match to the velocity maps of this sunspot in Beck et al. (2020).

Figure 8 shows the histograms of the distance of the inner and outer FPs from the center of the sunspot, while Fig. 9 shows their location on top of maps of photospheric and chromospheric quantities at 14:48 UT. The inner FPs can be found at a radial distance of  $0.6-1.8\,r_0$  with an average value of  $1.19\,r_0$ , slightly outside the outer penumbral boundary. None are found inside the umbra, where the field lines were open. The outer FPs are at  $1.5-2.5\,r_0$  with an average distance of  $1.9\,r_0$  in primarily plage regions of opposite polarity (Fig. 9). Only the short closed MFLs to the southeast (x; y = 20; 20) and southwest (x; y = 60; 20) connect to magnetic elements in the sunspot moat. The locations of the inner FPs coincide to a large extent with the end points of flow fibrils (bottom right panel of Fig. 9).

## 4.2.2. Magnetic field strength

Figure 10 shows the histograms of the magnetic field strength B from the HMI Milne-Eddington inversion at the inner and outer FPs of closed MFLs. The magnetic field strength at the inner FPs has a range of 0–1400 G with an average value of 550 G. The histogram shows a double-peaked distribution with peaks at 250 and 900 G, where the latter peak and the extension to 1400 G corresponds to inner FPs inside the penumbra. The histogram for the outer FPs has a roughly Gaussian distribution from 0 to 400 G with an average value of 180 G. The dierence in field strength between inner and outer FPs B (bottom panel of Fig. 10) ranges from 500 to +1300 G with an average value of +370 G and a similar bimodal shape to that of the inner FPs. About 6% of the




Fig. 4. Magnetic field lines of all manually selected IEF channels. Left to right: at 14:48 to 15:48 UT in steps of 12 min. The background images are from top to bottom B<sub>z</sub>, H and Ca IR continuum intensity, H line-core intensity, H LOS velocity and Ca IR LOS velocity. The white rectangle in the rightmost column indicates the location of the "perfect" IEF channel shown in Fig. 24 below. The red and blue arrows in the fifth column mark H filaments.

MFLs show a negative value of B with higher field strength at the outer FPs.

Figure 11 shows how the field strength along the MFLs varies between the FPs. The field strength B(x; y; z) was determined at the corresponding height of the MFL at each spatial position (x; y) in the horizontal plane, and then plotted against the 2D length L. It drops monotonically from the average value of 450 G with a broad distribution at the inner end up to a relative length of about 0.95. At that length, almost all the MFLs have a similar value of B of 180 G, which represents a small local maximum in B. At the inner and outer FPs, the MFLs are close to the photosphere, while they sample chromospheric layers in between (Fig. 7).


Figure 12 identifies the locations of inner and outer FPs in four bins in B of 400 G width each for the data at 14:48 UT. Most MFLs with B < 0 G belong to comparably short loops, whose inner FPs are in some cases far outside the outer penumbral boundary. For MFLs with increasing values of B; the inner FPs move towards the umbra and the loop height increases, while the outer FP locations do not vary significantly. The MFLs with B > 800 G are found on top of all others (bottom left panel of Fig. 12). The scatter plot in the bottom right panel of Fig. 12 shows a high correlation between the 3D loop length and B, in

line with the displacement of the inner FPs toward the umbra for increasing B, which also increases the length.

#### 4.2.3. Continuum temperature

Figure 13 shows the histograms of the temperatures at the inner and outer FPs, and their dierence T = T  $_{inner}$  T  $_{outer}$ . Similar to the magnetic field strength, the inner FPs exhibit a bimodal distribution with one peak at the quiet sun (QS) temperature of about 5800 K, an extended tail to lower temperatures down to 5300 K, and a smaller peak at 5450 K corresponding to FPs inside the penumbra. The distribution for the outer FPs is Gaussian-shaped around the QS temperature with an average value of 5820 34 K. The temperature dierence T is primar-ily negative from 500 to +100 K with two peaks at 350 and 0 K, again caused by the location of the inner FPs in the QS or the cooler penumbra. About 32% of the closed MFLs show a positive temperature dierence, which could drive a flow oppo-site to the IEF.

The high correlation between T and B of 0.73 in their scatter plot in Fig. 14 confirms a common origin for the behavior of the temperature and field strength dierence, namely the radial gradient in both B and T with distance from the sunspot






Fig. 5. Histograms of the apex height (top panel) and the 3D (blue line) and 2D (red line) loop length (bottom panel) of closed MFLs in the large sample.




Fig. 6. Scatter plot of 3D loop length and apex height for closed MFLs in the large sample.

center and the displacement of inner FPs with higher B towards the umbra. Both B>0 and T<0 create a gas pressure dif-ference in the same direction along a closed MFL, and so their anti-correlation amplifies the potential driving force.




Fig. 7. Loop shape for closed MFLs in the large sample against relative 2D length L. The background image shows the probability distribution of the height with the color bar to the right. The inner (outer) FP is at L = 0 (1). The yellow line indicates the average, while the red line is a parabolic loop with an apex height of 2.45 Mm.




Fig. 8. Histograms of the radial distance of the inner (blue line) and outer FPs (red line) of closed MFLs from the center of the sunspot in the large sample. The green and gray lines show the corresponding histograms for the manually selected sample. The x-axis at the top (bottom) gives the values in Mm (fractional outer penumbral radius  $r_0$ ).

## 4.2.4. Photospheric and chromospheric velocities

For the two chromospheric spectral lines of H and Ca ii IR, we determined the velocity at the inner and outer FPs and the maximal unsigned velocity along each closed MFL. The latter was defined as the maximum velocity encountered along the 2D paths of the MFLs in the horizontal plane, ignoring the height of the loop and any projection eects. The maximal velocity was derived from both the LOS and de-projected velocity maps. We only retrieved the velocity at the outer FPs from the photospheric HMI velocities to check for possible upflows at the outer end, as the velocities at the inner FPs are expected to be strongly contaminated with the regular Evershed flow.

The top two rows of Fig. 15 show the histograms of the three velocities ( $v_{max}$ , v at inner and outer FPs) as derived from the LOS velocities of both chromospheric lines. The maximal flow speeds along closed MFLs range from 0 to 8–10 km s  $^1$  with averages of 2.5 and 3.8 km s  $^1$  for Ca ii IR and H, respectively. All chromospheric velocities at the inner and outer FPs scatter around roughly zero with averages below 0:6 km s  $^1$  apart





Fig. 9. Locations of the inner and outer FPs of closed MFLs at 14:48 UT. The images in the background show (clockwise, starting left top) the H continuum and line-core intensity, the H LOS velocity, and  $B_z$ . Yellow (white) points denote the locations of the inner (outer) FPs.

from the velocity at the inner FPs in H with an average of 1:2 km s <sup>1</sup>, which presumably reflects the larger area coverage of the inner FPs on the limb side with its large-scale blueshift pattern. The maximal velocities in the de-projected velocity maps (bottom left two panels in Fig. 15) largely mirror the same quantity in the LOS velocities, but with an extended range from 0–30 km s <sup>1</sup> and average values of 10–15 km s <sup>1</sup>, which is an increase by a factor of about 5. The photospheric velocity at the outer FPs shows a slight redshift of 0:14 0:33 km s <sup>1</sup> on average, but with a large scatter from 1 to +1 km s <sup>1</sup>. Even with a correction for the convective blueshift of 0:2 km s <sup>1</sup> (see Appendix A.1), no systematic upflows are seen in the photosphere at the outer FPs.

Figure 16 shows scatter plots of the LOS and de-projected velocities against the apex height, B and T. None of the velocities exhibit a strong correlation with the apex height. The majority of the data points cluster around zero on the abscissa in all plots of velocities against B or T (right two columns of Fig. 16). The plots against B clearly reveal that only a small fraction of MFLs have B < 0, while T > 0 happens more frequently in comparison.

All plots of either LOS or de-projected maximal H velocities have upper and lower boundary regions of minimal or maximal velocities that rarely occur. The lower boundary is better defined and indicates the absence of small velocities below <2 km s <sup>1</sup> in the LOS and <4 km s <sup>1</sup> in the de-projected velocities for large values of B > 500 G or T < 200 K. The shape of the distributions matches to first order the predicted squareroot dependence of the flow speed on the magnetic or gas pressure dierences. Using Eq. (5), we calculated the expected velocities for a given value of B and T (solid green and orange lines in Fig. 16). The latter is independent of the gas density, while for the former we used three dierent gas densities in the HSRA model at log = 0:3; 1; and 2. With the close anti-correlation of B and T (Fig. 14), the total driving force is expected to be larger than the individual separate contributions. Both the observed LOS and maximal H velocities exceed a purely thermal driver (right column of Fig. 16). In the case of a driver based on the dierence in magnetic field strength, the predicted velocities in the middle column of Fig. 16 exceed the






Fig. 10. Histograms of the magnetic field strength B (top panel) at the inner (blue line) and outer (red line) FPs, and of their dierence  $B = B_{inner}$  Bouter (bottom panel).




Fig. 11. Field strength for closed MFLs in the large sample against relative 2D length L. The background image shows the probability distribution of B with the color bar to the right. The inner (outer) FP is at L=0 (1). The yellow line indicates the average value.

observed maximal H LOS velocities, but fall into the observed range for the de-projected velocities. Depending on the height in the solar chromosphere, the sound speed as the upper limit of mass flows is 6–20 km s <sup>1</sup>. The highest, but still rather low correlation values of about 0.2 are found for the relation between B or T and the velocities in Ca ii IR at the inner FPs (two bottom-right panels of Fig. 16), where the downflow points of the IEF channels would be expected.




Fig. 12. Properties of MFLs with dierent B. Top left: field lines with B < 0 G (turquoise), 0–400 G (red), 400–800 G (yellow), and >800 G (green) in a top view overlaid on B<sub>2</sub>. Bottom left: same in a side view to highlight the shape. Top right: inner and outer FPs for the same ranges in B. Bottom right: scatter plot of B and 3D loop length. The map in the bottom left panel is rotated relative to the images in the top row for better visibility of the closed MFLs.

## 4.2.5. Pressure balance

Figure 17 shows the histograms of the magnetic, gas, and total pressure at the inner and outer FPs, and their dierences p=p  $p_{inner}$ . For all calculations that include the gas pressure, the photospheric density in the HSRA at log = 0:2 of 2:89 10  $^7$  g cm  $^3$  was used.


The magnetic pressure at the inner FPs ranges from 0 to 8 kPa s with an average of 1.6 kPa. The corresponding values at the outer FPs are much smaller, with 0–0.5 kPa and an average of 0.17 kPa. The resulting dierence p mag is therefore mainly positive, with a similar total range to that of the inner FPs. In the same way as for temperature, the gas pressure at the inner FPs has a bimodal distribution with one peak at about 10.1 kPa corresponding to the penumbra and one at about 10.8 kPa corresponding to the QS, while the histogram for the outer FPs only shows the latter. The gas pressure difference is primarily negative, down to 1 kPa. The total pressure (bottom row of Fig. 17) at the inner FPs with values from 11 to 16 kPa exceeds the one at the outer FPs, which is fairly constant at 10:96 0:3 kPa. This leads to a total pressure imbal-

ance of up to +5 kPa under the assumption of equal gas densities, while 9% of the MFLs have a negative total pressure dierence.

As we cannot simultaneously confirm that the assumption of a global and large-scale total pressure balance holds over the comparably large distances between the inner and outer FPs of closed MFLs and the characteristic values of both B and T would drive a flow in the same direction, we calculated the "eective" pressure dierence following Eq. (3) in addition (Fig. 18). This yields slightly higher pressure dierences from

2 to 8 kPa, but otherwise follows the shape of the corresponding histograms of B and T with only a small fraction of MFLs with p  $_{a}$  < 0.

The scatter plots of p e against maximal de-projected and LOS velocities in Fig. 19 show a similar behavior as before for B and T alone (Fig. 16). The shape of the square-root dependence of the predicted velocities with its lower boundary is matched. The observed LOS velocities fall short of the prediction, while the de-projected velocities span the whole range of predicted velocities when using three dierent gas densities. We only varied the gas density in the application of Eq. (5), but not






Fig. 13. Histograms of the temperature T (top panel) at the inner (blue line) and outer (red line) FPs, and of their dierence  $T = T_{inner}$   $T_{outer}$  (bottom panel).




Fig. 14. Scatter plot of the dierence between inner and outer FPs in temperature  $\mathsf{T}$  and field strength  $\mathsf{B}$ .

in the calculation of p  $_{\rm e}$  that was derived with the HSRA gas density value at log = 0:2.

The question of which gas density is appropriate across a 2D FOV on the solar surface is dicult to answer without additional assumptions. We therefore tried to derive suitable gas densities using the relation between T and B<sup>2</sup> as predicted by Eq. (4) based on the photospheric HMI temperature and magnetic field strength. The left panel of Fig. 20 shows the scatter

plot of T and B<sup>2</sup> for the inner FPs of the large sample, while the right panel shows the same for a square area covering all the umbra, penumbra, and some QS areas outside the outer penumbral boundary. The inner FPs were located close to the outer penumbral boundary, primarily slightly outside the sunspot, with a temperature range of 5250–5950 K. The relation between T and B<sup>2</sup> for the inner FPs has a high correlation of about 0.85. The right panel reveals three dierent regimes in QS, penumbra, and umbra, with a distinct change of behavior at the outer umbral and penumbral boundaries. We therefore fitted three straight lines to the values in the right panel using a spatial mask for the three regimes (see the inset in the right panel of Fig. 20) excluding the areas of transition between them. Table 3 lists the resulting

fit parameters and the derived physical quantities gas density and total pressure  $p_{tot}$ . The correlation values for QS, penumbra, and umbra are somewhat lower at 0.26–0.5. The inferred gas densities of 0:75 4:3 10  $^7\,\mathrm{g}\,\mathrm{cm}^{-3}$  are in the range of the HSRA QS value of 2:89 10  $^7\,\mathrm{g}\,\mathrm{cm}^{-3}$  used in most of the previous calculations, while for the inner FPs the total pressure and the inferred gas density have similar values to the HSRA at log = 0:2 (Table 1). The magnetic pressure contributes more than 50% in the penumbra and completely dominates the total pressure in the umbra. A similar total pressure from the gas pressure alone would only be found for z < 70 km in the HSRA model (Table 1).

### 4.3. Manually selected IEF channels

The manually selected IEF channels are a small subset of the large sample, with seed points for the MFLs chosen based on the appearance of IEF channels in the intensity and velocity maps. All parameters related to the magnetic topology (length, height, locations of FPs) were very similar to the average values of the large sample. We only overplotted the histograms for the locations of the inner and outer FPs in Fig. 8. These reveal that the inner FPs of closed MFLs that explicitly correspond to IEF channels are slightly closer to the penumbra than those of the large sample, with about half of them being inside the sunspot. The histograms of field strength and temperature at the inner and outer FPs in the top two rows of Fig. 21 show that, for those MFLs, the temperature dierence is close to zero (T 20 K) and negligible compared to the field strength difference (B +600 G). The chromospheric maximal LOS flow velocities of 3-4 km s <sup>1</sup> are slightly higher than the averages for the large sample, while the photospheric HMI velocity at the outer FPs again shows no clear indication of blueshifts or upflows.

Figure 22 shows the observed and predicted velocities as a function of p, B; and T for the manually selected IEF chan-nels. In that case, the velocities predicted from the eective pres-sure match the range of observed LOS flow speeds, while they exceed them when only considering the magnetic pressure dif-ference. The temperature dierence is primarily positive, which therefore slightly reduces the eective pressure dierence. The corresponding curve for predicted velocities from T was below the plot range for T > 100 K.

### 4.4. Predicted flow speeds and apex heights

Equation (5) allows us to predict the expected flow velocity given the pressure dierence, where a linear relation between p and v should now hold. The top panel of Fig. 23 tests the prediction for the maximal H LOS velocities in the large sam-ple. The correlation coecient stays comparably low at 0.2, but




Fig. 15. Histograms of chromospheric and photospheric velocities. Top row, left to right: maximal H LOS velocity along field lines, and H LOS velocities at the inner and outer FPs. Middle row: same for Ca iRi and photospheric LOS velocity from HMI at the outer FPs.

the data points are to some extent covered by a one-to-one relation. The prediction of the apex height from Eq. (6) based on the pressure dierence falls short of the apex height measured in the magnetic field extrapolation by a factor of about 50 at a correlation of 0.36 (middle panel of Fig. 23). Apart from being dierent in magnitude, the trend of the data points follows the prediction, with a complete absence of low measured apex heights for low predicted heights. To some extent, this results from the relation between length and apex height (Fig. 6), where longer MFLs have their inner FPs closer to the umbra which causes a larger B (Fig. 12), and hence larger p. A similar mismatch by a fac-tor of about 30 in magnitude is seen for the apex height predicted from the observed LOS velocities (bottom panel of Fig. 23).

Assuming a velocity of the order of the chromospheric sound speed of 10 km s <sup>1</sup> would only correspond to an apex height of 0.18 Mm. A pressure dierence of 5 kPa that is covered by the observed values of p (Fig. 18) would only be able to lift a gas density of 0:006 10 <sup>7</sup> g cm <sup>3</sup> up to the average apex height of 2.96 Mm. This indicates that the Eqs. (6) and (7) related to the apex height do not seem to be strictly valid, as flows on the order of 10 km s <sup>1</sup> are found to appear along MFLs of that height.

### 4.5. The "perfect" IEF channel

To the southeast of the sunspot, one can find several adjacent MFLs (see the square in the rightmost column of Fig. 4) that form an almost "perfect" IEF channel with all theoretically

expected properties: two opposite polarities connected by a low-lying chromospheric loop with blueshifts at the outer and red-shifts at the inner end. At the heliocentric angle of the sunspot, a clear velocity signature of this kind is rare. We use the aver-age properties of these MFLs to summarize our results, while Table 4 additionally lists average values of dierent quantities for the large sample and correlation values between them for completeness.

The left column of Fig. 24 shows a magnification of the 3D view of the MFLs on top of the continuum intensity, B, and the LOS velocity of Ca ji IR, while the right column shows the corresponding magnetic and thermodynamic quantities along their length averaged over about 50 MFLs. The MFLs connect an opposite-polarity patch of magnetic elements in the sunspot moat at about 15 Mm distance from the outer penumbral boundary with the latter (left top two panels of Fig. 24). The inner FPs of the MFLs are slightly inside the penumbra. The MFLs are close to vertical in the magnetic elements at the outer FPs and more inclined in the sunspot penumbra, forming an arched loop with 2 Mm apex height (top right panel of Fig. 24). The field strength of about 700 G at the inner FPs is higher than the 300 G at the outer FPs, which still stand out in B, B, and to their immediate surroundings (middle right panel of Fig. 24). The MFLs show in that case clear upflows of 0.5–1 km s <sup>1</sup> in H and Ca ii IR at the outer and downflows of 1–3 km s <sup>1</sup> at the inner FPs (bottom row of Fig. 24). A slight blueshift is seen in the photospheric HMI LOS velocities.




Fig. 16. Scatter plots of apex height (left column), B (middle column), and T (right column) against flow velocities. Top to bottom: maximal deprojected H velocity, maximal H LOS velocity, and Ca IR LQS velocity at the inner FP. The green lines in the middle column show the predicted velocity from the field strength dierence for HSRA gas densities at (the lowest line to highest) log = 0:3; 1; and 2. The orange lines in the right column show the flow speed predicted by the temperature dierence.

For this "perfect" IEF channel, one therefore finds a picture of mass moving upwards at the outer end of an arched magnetic loop that then streams along it into the sunspot. Both the dierence in field strength B and temperature T would drive a flow in the same direction. For assumed total pressure balance between the inner and outer FPs, the magnetic, thermal, and hydrodynamic topology of these MFLs would all comply with a siphon flow scenario with a mass flow driven primarily by the field strength dierence and the inherent gas pressure dierence it causes.

## 5. Discussion

#### 5.1. Limitations of the current study

The current study suers from a few limitations that for the most part are related to or are intrinsic to the observational data. Our method could benefit from selecting a variety of sunspots to examine MFL connectivity all around sunspots. For the sunspot used, the IEF channels are best seen on the disk center side of the sunspot, where the LOS and the IEF channels are paral-lel (Fig. D.1 and Beck & Choudhary 2019). The magnetic field extrapolation could not provide closed MFLs in that direction due to the lack of opposite-polarity fields to the west. This configuration is similar to that encountered by Kawabata et al. (2020, their Fig. 12). Selecting a leading (trailing) sunspot with trailing (leading) plage to the west (east) of the solar central meridian would improve the situation by likely leading to more

closed MFLs all around the sunspot with our extrapolation technique. A second improvement would be to select a sunspot where the location of the zero line in LOS velocities does not coincide with the inner FPs of the IEF channels on the limb side, which can be achieved by selecting sunspots at small heliocentric angles.

The derivation of the LOS velocities themselves is only possible within limits. As shown in Choudhary & Beck (2018), the IEF is only a satellite component in the spectra close to the inner FPs, which leads to underestimation of the flow velocity by a factor of up to two. The de-projection to the true flow angle and speed was done based on the magnetic field vector at a single height in the NFFF extrapolation, whose accuracy cannot be confirmed with the current data. Additionally, the exact formation heights of the velocities of the chromospheric spectral lines of Ca jj IR at 854 nm and especially H in the canopy of a sunspot o disk center are not well known.

Apart from using the magnetic vector field information as the input for the NFFF extrapolation, the field strength and temperature from the HMI data were directly used. They are to some extent not fully consistent with each other, as their formation heights dier by a few hundred kilometers (Norton et al. 2006). A second limitation is the modulus of the field strength. The HMI magnetograph has only a limited spectral sampling that cannot spectrally resolve the Zeeman splitting for strong magnetic fields. Its spatial sampling of 0.5% prone to unresolved structures within a single pixel in the QS in addition. The onecomponent Milne-Eddington inversion used for the derivation




Fig. 17. Histograms of the magnetic pressure (top row), gas pressure (middle row), and total pressure (bottom row). Left to right: pressures at the inner and outer FPs, and their dierence  $p = p_{inner}$   $p_{outer}$ .

of the vector field from the HMI observations cannot take this latter issue into account. The field strength values are therefore likely to underestimate the true field strength both in the QS and the sunspot; for example the average field strength of 180 G at the outer FPs (Fig. 10) is far from the 1-kG fields expected for magnetic elements in the moat (Beck et al. 2007; Utz et al. 2013). Beck & Choudhary (2019) compared the field strength at the inner FPs between HMI and an inversion of the Fe; lines near 1565 nm and found the HMI values to be about 500 G lower than the average value of 1.3 kG from the Fe lines. Even for fields of 1 kG at the outer FPs, a positive field strength dierence of the same modulus as found here (0.3–0.4 kG) would therefore be maintained (Table 4).

The limitations of the NFFF extrapolation are to some extent caused by the HMI input data. At the photospheric boundary, the values of B are likely inaccurate to some extent and the field inclination in the penumbra corresponds to a weighted average in the case of unresolved magnetic field components with dier-ent inclinations (Bellot Rubio et al. 2004; Beck 2008). The mag-netic connectivity is to a large extent deterministic and can only provide closed MFLs between opposite-polarity FPs, but HMI cannot detect diuse weak magnetic flux below a certain level because of its spatial and spectral resolution. Photospheric highresolution observations of internetwork quiet Sun magnetism reveal a large amount of magnetic flux that is beyond the detection capabilities of the HMI (Lites et al. 2008; Beck & Rezaei 2009; Beck et al. 2017). However, a connection between the outer penumbra and weak magnetic flux favors a siphon flow scenario even more because of the resulting small field strength




Fig. 18. Histogram of the eective pressure dierence pe.

at the outer foot points. The 2D chromospheric flow field of the same sunspot was successfully modeled in Beck et al. (2020) assuming an axisymmetric pattern with IEF channels all around the sunspot.

No magnetic field lines match the strong and large filament to the east, whose negative polarity FP could be at about the center of the FOV of the extrapolation box (bottom left panel with the AIA 304 Å image in Fig. 2). The magnetic field extrapolation here supports a scenario of mass loading on top of an arcade of field lines that cross the neutral line. However, without




Fig. 19. Scatter plots of the eective pressure dierence  $p_e$  against de-projected (left column) and LOS (right column) flow velocities. The green lines show the predicted velocities for HSRA gas densities at log = 0:3; 1; and 2.




Fig. 20. Scatter plots of the relation between T and B<sup>2</sup>. Left panel: for the inner FPs of the large sample. Right panel: For a square covering the whole sunspot and some QS regions in all six time steps. The red lines are straight line fits to all or subsets of data points. The inset in the upper right corner of the right panel shows the corresponding spatial mask for the umbra (purple), penumbra (orange), and QS (red) at 14:48 UT.

Table 3. Line fit and physical parameters derived from the  $\mathsf{T}(\mathsf{B}^2)$  relation.

	Inner FPs	QS	Penumbra	Umbra
Slope	0:0023	0:0014	0:0083	0:0044
Intercept	13.5	8.5	46.3	23.4
Correlation	0.85	0.26	0.33	0.50
$=10^{-7} (g cm^{-3})$	2.69	4.31	0.75	1.42
hTi(K)	5696	5796	5417	3994
hBi (kG)	0.55	0.39	1.26	2.45
p <sub>gas</sub> (kPa)	9.83	15.97	2.60	3.64
p <sub>mag</sub> (kPa)	1.20	0.6	6.27	23.75
p <sub>tot</sub> (kPa)	11.03	16.57	8.87	27.39

introducing an inertial term into it, no extrapolation is able to yield dips in field lines apart from very peculiar configurations in the input magnetograms.

The energy and pressure balance suers from all the short-comings listed above. With the partially unknown formation heights for the LOS velocities, temperature, and field strength, the density that must be attributed to the IEF is not obvious. The IEF has well-defined inner FPs, where it stops at an optical depth of log = 3 z = 400 km (Choudhary & Beck 2018), but the height at which it sets in at the outer FPs is not clear. We plan to investigate the height variation of the IEF using other data sets with more spectral lines that cover a larger range of formation heights (see, e.g., Felipe et al. 2010; Bethge et al. 2012) in the future.




Fig. 21. Histograms of physical properties of the manually selected IEF channels. Top two rows: histograms of B and T at the inner (left column) and outer (middle column) FPs and their dierences B and T (right column). Bottom row, left to right: histograms of maximal chromospheric LOS velocities in H and Ca IR anii photospheric HMI velocity at the outer FPs.

## 5.2. Magnetic properties of the IEF

Thanks to the results of the NFFF extrapolation, we were able to trace individual MFLs related to the IEF. We used two different samples, where the large sample was defined based on the magnetic topology alone, without considering the chromospheric velocities, while the second sample was primarily based on the intensity and velocity signature of IEF channels. Both samples yielded almost identical average values and histograms for the properties of the MFLs and their foot points (Figs. 10, 15, 21 and 22), which indicates that they trace the same structures in the solar atmosphere, namely the IEF channels.

The connectivity found from the magnetic field extrapolation in the current study aligns with the results or assumptions on IEF channels in previous studies. The average distances of inner and outer FPs of 1.2 and 1.9 sunspot radii (Fig. 5) implies that they connect the outer penumbra with the end of the moat cell (Sobotka & Roudier 2007). In Beck et al. (2020), the corresponding values for the same sunspot were 0.98 and  $2 r_0$ , derived or set only using the velocity maps, while peak velocities and intensities for the inner FPs were found at about  $1 r_0$  in Beck & Choudhary (2020).

The average length of closed MFLs of 13 Mm allows one an indirect estimate of the lifetime of IEF channels. Assuming a motion with the chromospheric sound speed of about  $10\,\mathrm{km\,s}^{-1}$ , it takes about 20 min to reach the inner FP from the outer end. Typical lifetimes of IEF channels then should be of the same order when an IEF channel is established, which matches the

lifetimes of 10–60 min found in Beck & Choudhary (2020) from tracing the flow signature of IEF channels with time.

The current average apex height of 3 Mm nicely matches the one of 2.45 Mm inferred for this sunspot based only on the LOS velocity maps in Beck et al. (2020). Aschwanden et al. (2016) found a typical height of up to 4Mm for chromospheric structures such as superpenumbral fibrils, with a similar steep decline of the histogram towards larger heights (Fig. 5 and their Fig. 10). The assumed parabolic loop shape used in Beck et al. (2020), which was previously also used in Maltby (1975), is directly confirmed to first order by the field extrapolation (Fig. 6), which also supports the inclination values for these MFLs being around 30 to the local horizontal (Haugen 1969; Dialetis et al. 1985; Beck & Choudhary 2019; Beck et al. 2020) with a smooth radial variation. The comparably solid ratio between length and height of 4:1 (Fig. 6) might allow one to at least get some estimate of apex heights for closed MFLs in the absence of an extrapolation when the horizontal foot point distance is known.

The average field strengths at the inner and outer FPs of 0.6 kG and 0.2 kG in the HMI data likely underestimate the true values, but the average positive field strength dierence of +0.4 kG appears to be robust in its sign. One of the FPs of the closed MFLs related to the IEF is inside or close to the outer boundary of the penumbra, which implies that nearly any connection will end in an outer FP of lower field strength. The properties of the outer FPs in location, field strength, and inclination (Figs. 7, 9, 11 and 24) match those of magnetic elements in the QS with a higher field strength than their immediate




Fig. 22. Scatter plots of (top to bottom) p, B; and T against the maximum H LOS velocity of manually selected IEF channels. The green lines show the predicted velocities for HSRA gas densities at log = 0:3; 1 and 2.

Fig. 23. Scatter plots of predicted and measured velocities and apex heights. Top to bottom: predicted vs. observed maximal H LOS veloc-ity, measured apex height vs. prediction from  $p_e$ , and measured apex height vs. prediction from the observed maximal H LOS velocity. The green line in the top panel indicates a one-to-one correlation.

surroundings and nearly vertical magnetic field inclinations. The dierence in temperature between the inner and outer FPs seems to play a smaller role than B in most cases, with the caveat that the HMI resolution might prevent us from seeing the true temperatures of the intergranular lanes in which the magnetic elements of the outer foot points are likely to reside.

## 5.3. Pressure balance

The simplified pressure balance of Eq. (2) together with the conversion to the expected flow speed of Eq. (5) correctly predicts the square-root shape of the distribution of the p v scat-ter plots, but the numbers only line up partly with the observed LOS or de-projected velocity values for individual IEF channels

(Fig. 23). The main reasons for this will be the observational limitations discussed in Sect. 5.1. Varying the gas density used in the calculations within reasonable limits ( $z=20\ 200\ km$ , = 0:3 310  $^7$  g cm  $^3$ ) suces to cover the range of observed LOS velocities of 2 10 km s  $^1$ .

For the umbra and the QS with primarily vertical magnetic fields and lower higher order magnetic contributions such as curvature forces to the pressure equation (Steiner et al. 1986; Borrero et al. 2019), the relation between T and B<sup>2</sup> (Fig. 20) might allow one to derive not only average density values, but also spatially resolved density maps when a suited boundary value for the total pressure is prescribed and the Wil-son depression is properly accounted for (Puschmann et al. 2010; Löptien et al. 2020). A combination of the magnetic field




Fig. 24. The "perfect" IEF channel south of the sunspot. Left column: 3D view of the MFLs connecting the sunspot to the opposite-polarity patch on top of the continuum intensity  $I_c$  (top panel),  $B_z$  (middle panel), and the Ca  $_{ii}$  IR LOS velocity (bottom panel). Right column: dierent quantities averaged over the MFLs in the left column as a function of the relative length L. Top panel: temperature (blue line, y-axis at the left) and height (red line, y-axis at the right). Middle panel: LOS magnetic flux  $_{LOS}$  (green line),  $B_z$  (blue line) and field strength B (red line). Bottom panel: LOS velocities of HMI (green line), Ca IR (blue $_i$ tine), and H (red line).

extrapolation results with a thermal inversion of the Caii IR spectra, for example through the Calcium Inversion based on a Spectral Archive code (CAISAR; Beck et al. 2013, 2015), would possibly provide density stratification in a 3D volume.

## 5.4. The driver of the IEF

The IEF channels are well aligned with the magnetic field vector near the inner FPs (Beck & Choudhary 2019). Their visibility in the Ca ji IR data (Figs. 3, 4 and 24) in particular complies with the shape in the magnetic field extrapolation (Fig. 7), which predicts that the IEF channels leave the Ca ji IR formation height of 1–2 Mm around the apex. The primarily radial orientation of the MFLs matches that of the intensity fibrils in H or Ca ji IR in Fig. 3 at most places (see also de la Cruz Rodríguez & Socas-Navarro 2011; Schad et al. 2013; Leenaarts & Carlsson 2015; Kawabata et al. 2020) apart from the dark filaments that supposedly do not correspond to IEF channels. There is therefore no indication that the IEF would not follow the closed MFLs of the NFFF extrapolation wherever the flow cannot be continuously traced in the velocity maps,

or that no IEF channels with similarly shaped closed field lines would exist where the magnetic field extrapolation is unable to yield them. With the positive field strength dierence and the anti-correlated temperature dierence towards the inner FPs, a siphon flow is therefore the most likely explanation for the IEF pattern around sunspots.

The diagnosis of the driver of the IEF in this study was only possible through the combination of high-resolution chromospheric spectra and velocities — or high-resolution observations in general — with a magnetic field extrapolation derived from full-disk data of lower spatial resolution. This highlights the potential of magnetic field extrapolations for the correct interpretation of the physics visible in high-resolution observations at much smaller spatial scales. This approach has a broad range of potential applications that have only partially been explored so far (e.g., Aschwanden et al. 2016; Grant et al. 2018; Yadav et al. 2019; Kawabata et al. 2020; Louis et al. 2021) and could be developed into a standard diagnostic tool for high-resolution observations in the future. The availability of a magnetic field extrapolation also provides additional diagnostic potential through, for instance, the de-projection of LOS

Table 4. Summary of results for the large sample.

Quantity	Apex	2D L	3D L	r <sub>in</sub>	r <sub>out</sub>	B <sub>in</sub>	B <sub>out</sub>	В
units		Mm		r	0		kG	
Χx	3:0 1:8	13 6	14 7	1:2 0:2	1:9 0:2 T	0:6 0:3	0:2 0:1	0:4 0:3
Quantity	Tin	$T_out$		LOS v <sub>H</sub>	Depr v <sub>H</sub>	LOS v <sub>Ca</sub>	Depr v <sub>Ca</sub>	VHMI
Units		kK				km s <sup>1</sup>		
Χx	5:7 0:2	5:8 0:03	0:1 0:2	3:8 1:7	16 8	2:5 1:2	10 5	0:1 0:3
Quantity	p <sup>in</sup> mag	p <sub>mag</sub>	p <sub>mag</sub>	$p_{gas}^{in}$	out	p <sub>gas</sub>	p <sub>e</sub>	p <sub>out</sub> <sup>tot</sup>
kPa	1:6 1:7	0:2 0:3	p <sub>gas</sub>			0:23 0:3	1:67 1:9	10:96 0:3
			1:4 1:6	10:6 0:3	10:8 0:1			
V <sub>H</sub> V <sub>Ca</sub>	L-apex	L B		Correlation v	alues	р v <sub>H</sub>	$T B^2$	h(p)-apex
0.66	0.90	0.56	ВТ	B v <sub>H</sub>	T v <sub>H</sub>	0.17	0.85	0.36
			0.73	0.19	0.17			

A&A, 403, L47

Mechanik. 48. 218

240, 63

velocities to the field direction to obtain true flow speeds. In the other direction, high-resolution observations could help in detailed modeling of for example filaments by determining a corresponding mass density from H spectra to add as a constraint to the extrapolation or the extrapolation results. Combinations of these two dierent approaches have a clear potential to improve future research in solar physics, where the list of techniques applied in the current study is far from exhaustive.

### 6. Conclusions

We find that the inverse Evershed flow of the leading sunspot in AR NOAA 12418 happens along elongated arched loops of about 13 Mm length with an apex height of about 3 Mm. The corresponding closed magnetic field lines connect the outer penumbra with magnetic elements in or at the end of the moat cell. The positive dierence in magnetic field strength of on average +400 G and the negative one in temperature of 100 K both support a siphon flow from the outer foot points towards the sunspot as the driver of the inverse Evershed flow.

Acknowledgements. The Dunn Solar Telescope at Sacramento Peak/NM was operated by the National Solar Observatory (NSO). NSO is operated by the Association of Universities for Research in Astronomy (AURA), Inc. under cooperative agreement with the National Science Foundation (NSF). HMI data are courtesy of NASA/SDO and the HMI science team. IBIS has been designed and constructed by the INAF/Osservatorio Astrofisico di Arcetri with contributions from the Università di Firenze, the Universitàdi Roma Tor Vergata, and upgraded with further contributions from NSO and Queens University Belfast. This work was supported through NSF grants AGS-1413686 and AGS-2050340. Q.H. and A.P. acknowledge partial support of NASA grants 80NSSC17K0016, LWS 80NSSC21K0003 and NSF awards AGS-1650854 and AGS-1954503.This research was also supported by the Research Council of Norway through its Centres of Excellence scheme, project number 262622, as well as through the Synergy Grant No 810218 459 (ERC-2018-SyG) of the European Research Council.

## References

```
Alissandrakis, C. E. 1981, A&A, 100, 197
Aschwanden, M. J., Reardon, K., & Jess, D. B. 2016, ApJ, 826, 61
Beck, C. 2008, A&A, 480, 825
Beck, C., & Rezaei, R. 2009, A&A, 502, 969
Beck, C., & Choudhary, D. P. 2019, ApJ, 874, 6
Beck, C., & Choudhary, D. P. 2020, ApJ, 891, 119
Beck, C., Bellot Rubio, L. R., Schlichenmaier, R., & Sütterlin, P. 2007, A&A, 472, 607
Beck, C., Rezaei, R., & Puschmann, K. G. 2012, A&A, 544, A46
Beck, C., Rezaei, R., & Puschmann, K. G. 2013, A&A, 549, A24
Beck, C., Choudhary, D. P., & Rezaei, R. 2014, ApJ, 788, 183
Beck, C., Choudhary, D. P., Rezaei, R., & Louis, R. E. 2015, ApJ, 798, 100
```

```
Cargill, P. J., & Priest, E. R. 1980, Sol. Phys., 65, 251
Cavallini, F. 2006, Sol. Phys., 236, 415
Chandrasekhar, S., & Kendall, P. C. 1957, ApJ, 126, 457
Choudhary, D. P., & Beck, C. 2018, ApJ, 859, 139
Dasgupta, B., Dasgupta, P., Janaki, M. S., Watanabe, T., & Sato, T. 1998, Phys.
   Rev. Lett., 81, 3144
de la Cruz Rodríguez, J., & Socas-Navarro, H. 2011, A&A, 527, L8
Dere, K. P., Schmieder, B., & Alissandrakis, C. E. 1990, A&A, 233, 207
Dialetis, D., Mein, P., & Alissandrakis, C. E. 1985, A&A, 147, 93
Dunn, R. B. 1969, Sky Teles., 38, 368
Dunn, R. B., & Smartt, R. N. 1991, Adv. Space Res., 11, 139
Evershed, J. 1909, The Observatory, 32, 291
Felipe, T., Khomenko, E., Collados, M., & Beck, C. 2010, ApJ, 722, 131
Frazier, E. N. 1972, Sol. Phys., 24, 98
Gary, G. A. 2001, Sol. Phys., 203, 71
Georgakilas, A. A., & Christopoulou, E. B. 2003, ApJ, 584, 509
Georgakilas, A. A., Christopoulou, E. B., Skodras, A., & Koutchmy, S. 2003,
   A&A, 403, 1123
Gingerich, O., Noyes, R. W., Kalkofen, W., & Cuny, Y. 1971, Sol. Phys., 18, 347
Grant, S. D. T., Jess, D. B., Zaqarashvili, T. V., et al. 2018, Nat. Phys., 14, 480
Harvey, J. W., Hill, F., Hubbard, R. P., et al. 1996, Science, 272, 1284
Haugen, E. 1969, Sol. Phys., 9, 88
Henriques, V. M. J., Nelson, C. J., Rouppe van der Voort, L. H. M., &
   Mathioudakis, M. 2020, A&A, 642, A215
Hu, Q., & Dasgupta, B. 2008, Sol. Phys., 247, 87
Hu, Q., Dasgupta, B., Choudhary, D. P., & Büchner, J. 2008, ApJ, 679, 848
Hu, Q., Dasgupta, B., Derosa, M. L., Büchner, J., & Gary, G. A. 2010, J. Atmos.
   Sol. Terr. Phys., 72, 219
Jaeggli, S. A., Lin, H., Mickey, D. L., et al. 2010, Mem. Soc. Astron. It., 81, 763
Kawabata, Y., Asensio Ramos, A., Inoue, S., & Shimizu, T. 2020, ApJ, 898, 32
Leenaarts, J., Carlsson, M., & Rouppe van der Voort, L. 2015, ApJ, 802, 136
Lemen, J. R., Title, A. M., Akin, D. J., et al. 2012, Sol. Phys., 275, 17
Li, S., Jaroszynski, S., Pearse, S., Orf, L., & Clyne, J. 2019, Atmosphere, 10, 488
Lites, B. W., Kubo, M., Socas-Navarro, H., et al. 2008, ApJ, 672, 1237
Liu, C., Prasad, A., Lee, J., & Wang, H. 2020, ApJ, 899, 34
Löhner-Böttcher, J., Schmidt, W., Schlichenmaier, R., Steinmetz, T., &
   Holzwarth, R. 2019, A&A, 624, A57
Löptien, B., Lagg, A., van Noort, M., & Solanki, S. K. 2020, A&A, 635, A202
Louis, R. E., Prasad, A., Beck, C., Choudhary, D. P., & Yalim, M. S. 2021, A&A,
   652, L4
Mahajan, S. M., & Yoshida, Z. 1998, Phys. Rev. Lett., 81, 4863
Maltby, P. 1975, Sol. Phys., 43, 91
Meyer, F., & Schmidt, H. U. 1968, Zeitschrift Angewandte Mathematik und
```

Beck, C., Fabbian, D., Rezaei, R., & Puschmann, K. G. 2017, ApJ, 842, 37 Beck, C., Choudhary, D. P., & Ranganathan, M. 2020, ApJ, 902, 30

Bellot Rubio, L. R., Balthasar, H., & Collados, M. 2004, A&A, 427, 319

Bobra, M. G., Sun, X., Hoeksema, J. T., et al. 2014, Sol. Phys., 289, 3549 Borrero, J. M., Pastor Yabar, A., Rempel, M., & Ruiz Cobo, B. 2019, A&A, 632,

Bethge, C., Beck, C., Peter, H., & Lagg, A. 2012, A&A, 537, A130

Bhattacharyya, R., & Janaki, M. S. 2004, Phys. Plasmas, 11, 5615

Bellot Rubio, L. R., Balthasar, H., Collados, M., & Schlichenmaier, R. 2003,

Bhattacharyya, R., Janaki, M. S., Dasgupta, B., & Zank, G. P. 2007, Sol. Phys.,

#### A&A 662, A25 (2022)

Sobotka, M., & Roudier, T. 2007, A&A, 472, 277

```
Montesinos, B., & Thomas, J. H. 1997, Nature, 390, 485
Montgomery, D., & Phillips, L. 1988, Phys. Rev. A, 38, 2953
                                                                                  Steiner, O., Pneuman, G. W., & Stenflo, J. O. 1986, A&A, 170, 126
Nayak, S. S., Bhattacharyya, R., Prasad, A., et al. 2019, ApJ, 875, 10
                                                                                  Thomas, J. H. 1988, ApJ, 333, 407
Norton, A. A., Graham, J. P., Ulrich, R. K., et al. 2006, Sol. Phys., 239, 69
                                                                                  Thomas, J. H., & Montesinos, B. 1991, ApJ, 375, 404
                                                                                  Uitenbroek, H., Balasubramaniam, K. S., & Tritschler, A. 2006, ApJ, 645, 776
Pesnell, W. D., Thompson, B. J., & Chamberlin, P. C. 2012, Sol. Phys., 275, 3
Prasad, A., Dissauer, K., Hu, Q., et al. 2020, ApJ, 903, 129
                                                                                  Utz, D., Jurčák, J., Hanslmeier, A., et al. 2013, A&A, 554, A65
Puschmann, K. G., Ruiz Cobo, B., & Martínez Pillet, V. 2010, ApJ, 720,
                                                                                  Valio, A., Spagiari, E., Marengoni, M., & Selhorst, C. L. 2020, Sol. Phys., 295,
Reardon, K. P., & Cavallini, F. 2008, A&A, 481, 897
                                                                                  Wiegelmann, T. 2008, J. Geophys. Res. (Space Phys.), 113, A03S02
                                                                                  Wiegelmann, T., & Sakurai, T. 2012, Liv. Rev. Sol. Phys., 9
Rueedi, I., Solanki, S. K., & Rabin, D. 1992, A&A, 261, L21
Ruiz Cobo, B., & del Toro Iniesta, J. C. 1992, ApJ, 398, 375
                                                                                  Yadav, R., de la Cruz Rodríguez, J., Díaz Baso, C. J., et al. 2019, A&A, 632,
Schad, T. A., Penn, M. J., & Lin, H. 2013, ApJ, 768, 111
Scherrer, P. H., Schou, J., Bush, R. I., et al. 2012, Sol. Phys., 275, 207
                                                                                  Yalim, M. S., Prasad, A., Pogorelov, N. V., Zank, G. P., & Hu, Q. 2020, ApJ,
Schlichenmaier, R., & Schmidt, W. 2000, A&A, 358, 1122
Sigwarth, M., Schmidt, W., & Schuessler, M. 1998, A&A, 339, L53
                                                                                  Zirin, H. 1972, Sol. Phys., 22, 34
```

## Appendix A: Derivation and calibration of line-of-sight velocities

#### A.1. Photospheric velocities from HMI




Fig. A.1. Scatter plot of the LOS velocities of H and Caii IR at the inner FPs of closed MFLs near the outer penumbral boundary. The red line shows a linear regression to the data points. Its slope and intercept, and the linear correlation coecient are given inside the panel. Only every fifth data point is plotted.

The sunspot of AR NOAA 12418 was located at 35.54 eastern longitude. The average LOS velocity caused by solar rotation was 1 km s <sup>1</sup> with an additional variation of 0.4 km s <sup>1</sup> across the box used for the magnetic extrapolation (middle column of Fig. 1), which required the use of a spatially varying correction as well. We determined the contribution from solar rotation by averaging the de-projected (see Appendix D.3 below) HMI velocity maps in the extrapolation box in the x and y directions and fitting straight lines to the average curves. The fitted straight lines were then subtracted from the data. After subtraction of these straight lines, the average HMI velocity in QS regions was found to be approximately zero. Löhner-Böttcher et al. (2019) measured a convective blueshift of about 0:2 km s <sup>1</sup> for the HMI Fe i line at 617.3 nm at = 0:7. We did not apply an additional correction for that, but note that HMI velocities of approximately zero should correspond to small blueshifts of -0.2 km s <sup>1</sup>.

## A.2. Bisector analysis of chromospheric spectra

We used the same bisector analysis as in Beck & Choudhary (2020) and selected the bisector velocity at 93% line depth as LOS velocity of Ca ii IR and H. The average velocity across the whole IBIS FOV was set to zero for each of the six spectral scans, which puts the average umbral velocity at rest as well (Beck et al. 2020; Henriques et al. 2020). The velocities of H and Ca ii IR in and near sunspots are similar (e.g., Beck & Choudhary 2020; Beck et al. 2020), with a correlation coecient above 0.5 and usually higher velocity values in H (Fig. A.1). We primarily show results for H in the following but note that the velocities of the two lines can be used synony-mously. The line-core intensity was defined as the lowest value inside the spectral line in each profile.

# Appendix B: Conversion of HMI intensity to temperature

We used the Planck function to convert the de-projected continuum intensity maps of HMI to temperature (see, e.g., Beck et al.




Fig. B.1. Conversion curves between HMI continuum intensity and temperature as derived from the Planck function (blue line) and using the SIR code (red dashed line). The black dashed vertical lines at I=I(5800) = 0.11 and 1.15 represent the minimum and maximum intensity values in our HMI data.

2012; Valio et al. 2020). Norton et al. (2006) give the formation height of the continuum (line core) of the Fe¡ line at 617 nm as about 20 km (300 km), and so the assumption of an ideal gas in local thermodynamic equilibrium (LTE) with black body radiation is justified. At the heliocentric angle of the observations, the continuum formation height should shift slightly upwards to about 40 km with the optical depth changing from = 0:63 to = 0:63 sin = 0:46 corresponding to log = 0:2 and log = 0:33, respectively.

We calculated the emergent radiation at 617 nm from the Planck function for a temperature range from 3000 to 7000 K and normalized the values by the intensity for 5800 K to obtain a conversion curve from relative intensities to absolute temperatures. The center-to-limb variation (CLV) in the observed intensity maps was removed in a similar way as for the HMI velocities using averages in x and y, but now dividing with the fitted straight lines. After the correction for the CLV, the relative HMI intensities were converted to their corresponding temperatures. With this choice of normalization, the average QS temperature is forced to be 5800 K.

To verify the conversion curve, we generated the same using the Stokes Inversion based on Response functions code (SIR; Ruiz Cobo & del Toro Iniesta 1992), which employs LTE. We synthesized a wavelength window around the Fei line at 617.3 nm that covered continuum wavelengths while modifying the HSRA model with global osets of -2000 to +2000 K at all optical depths. We selected the continuum intensity that corresponds to having a temperature of 5765 K at log = 0:4 in the original unperturbed HSRA model for the normalization in that case. Figure B.1 demonstrates that in the relevant intensity range of the HMI data from 0.11 to 1.15 the dierences between the two approaches are minor, and so the simple conversion using the Planck function can indeed be used, especially as our main focus is on spatial locations with a relative intensity of about unity.

## Appendix C: Non-force-free magnetic field extrapolation

We used the NFFF extrapolation code developed by Hu et al. (2010) to obtain the magnetic field connectivity around the sunspot. The algorithm for this code is based on the principle of

minimum dissipation rate (MDR; Montgomery & Phillips 1988; Dasgupta et al. 1998; Bhattacharyya & Janaki 2004), which originates from a variation approach that allows one to obtain dissipative relaxed states in a two-fluid plasma with an external helicity driving (Bhattacharyya et al. 2007). This therefore makes it suitable for an open system with a flow, as in the solar photosphere.

The MDR approach leads to a set of two decoupled inhomogeneous double-curl Beltrami equations (Mahajan & Yoshida 1998) for the magnetic field B and the fluid vorticity! given by (Bhattacharyya & Janaki 2004; Bhattacharyya et al. 2007)

$$r r B + a_1 r B + b_1 B = r$$
 (C.1)  $r r$   
!  $+ a_2 r ! + b_2 ! = r$ ; (C.2)

where  $a_1$ ;  $a_2$ ;  $b_1$ , and  $b_2$  are constants that depend on the parameters of the system, and and are arbitrary scalar functions that satisfy the Laplace's equation.

For the rest of the description, we focus only on the magnetic field, which is more relevant to the current study. The ambiguity arising from the arbitrary potential can be eliminated by taking the curl of Eq. (C.1), which results in (Hu et al. 2008)

$$r r r B + a_1 r r B + b_1 r B = 0$$
: (C.3)

An exact solution of Eq. (C.3) can be obtained using the linear superposition of three linear force-free fields (LFFFs) arising from the orthogonality of Chandrasekhar-Kendall (CK) eigenfunctions (Chandrasekhar & Kendall 1957). Thus, B is expressed as (Hu et al. 2008):

$$B = B_1 + B_2 + B_3$$
;  $r B_i = {}_{i}B_i$ ; (C.4)

where are distinct constant parameters with i=1;2;3. The equation further requires that one of the is zero. Here we arbitrarily choose = 0, which makes B a potential field. This then implies that a = (+1) and b = -1 Now, we combine Eqs. (C.3) and (C.4) to get

Here V is called the Vandermonde matrix whose elements are of the form  $^{i}$   $^{1}$ j for i; j = 1; 2; 3 (Hu & Dasgupta 2008). Writing the above equation for the z component yields the boundary condition for each LFFF. Assuming a value for the  $_{i}$  parameter, we can then use a standard fast Fourier transform based LFFF solver (Alissandrakis 1981) to obtain the extrapolated field in the full volume.

To obtain an optimal pair of (  $;_1$  ),  $;_2$  we minimize the dier-ence between the observed (B ) and computed (b ) transverse field by defining the following metric (Hu & Dasgupta 2008; Hu et al. 2008):

$$E_n = \int_{i=1}^{XM} jB_{t;i} \quad b_{t;i}j = \int_{i=1}^{M} jB_{t;i}j :$$
(C.6)

Here  $M = N^2$ , is the total number of grids points on the bottom boundary.

One term on the right-hand side of Eq. (C.5) involves the evaluation of a second derivative, (rrB) =  $_z$ r²B, at  $_z$ = 0. This means that we need to provide the boundary conditions at two or more layers. As vector magnetograms are only available at the photospheric boundary, Hu et al. (2010) devised an iter-ative scheme which successively corrects the potential subfield




Fig. D.1. De-projection of LOS velocities onto the magnetic field vector. Top to bottom: observed H LOS velocity,  $\cos(x; y)$ , and de-projected H velocities. No de-projection was applied for all places with  $\cos < 0.1$  and |y| < 0.2 km s<sup>-1</sup> (white areas in the middle panel).

 $B_2$  starting from an initial guess. By setting  $B_2 = 0$ , Eq. (C.5) is first reduced to a second-order matrix equation. This allows us to unambiguously determine the boundary conditions for subfields  $B_1$  and  $B_3$ . If the value of the metric  $E_n$  is above a certain threshold, then a corrector potential field —which is derived from the dierence in the observed and the computed transverse field—is added to B to improve the agreement. For the extrapolations in the present study, the final value of the  $E_n$  was around 0.32, which is similar to those obtained in previous studies (Liu et al. 2020).

## Appendix D: De-projection and alignment of high-resolution and full-disk data

D.1. Spatial de-projection of high-resolution data

Because of the o-center position of the AR, the high-resolution data from the DST required a de-projection to compensate their

geometrical foreshortening. We first determined the center of the sunspot and the orientation of the line that joins the centers of the sunspot and the sun, which made an angle of 35.54 to solar eastwest. We rotated all 2D images by the angle above to have the symmetry line along a row of the data. We then stretched the images by a factor of 1= cos = 1:37 with = 43 being the heliocentric angle of the sunspot in the y-axis and rotated the images back by 35.54 to their original north—south orientation. The resulting images now had 13651000 pixels instead of the initial 10001000 pixels in the uncorrected DST images. The one-hour duration of the observation was too short to cause a significant change in the position of the sunspot on the disk. The same values for the angle and stretching were therefore used for all of the IBIS images recorded during the observation.

### D.2. De-projection of LOS velocities

The observed velocities are the projection of the true velocity vector onto the LOS. The true flow speed can only be determined if the flow angle is known, which cannot be directly derived from the observed spectra, only through additional assump-tions such as axisymmetry (Schlichenmaier & Schmidt 2000) or calculations such as a thermal inversion (Beck & Choudhary 2019). A third possibility is to assume field-aligned flows (e.g., Bellot Rubio et al. 2003; Beck 2008), where ionized and magnetized plasma can only move along MFLs but not perpendicular to them. In the vicinity of a sunspot at chromospheric layers this assumption is largely valid. We thus used the magnetic field extrapolation to retrieve the magnetic field vector B(x; y; z) at a height z = 1 Mm across the FOV. The height corresponds roughly to the formation height of the Ca ji IR line core in which the IEF channels can be seen both in intensity and velocity maps. The scalar product of the LOS vector LOS =  $[\cos \sin; \sin \sin; \cos]$  with = 215:54 and

= 43 with the magnetic field vector gives

LOS 
$$B(x; y) = jBj cos(x; y);$$
 (D.1)

while the de-projected true flow speed is given by  $v_{de\ proj}(x;y) = v_{LOS}(x;y) = \cos(x;y)$ .

Figure D.1 shows an example of the H LOS velocities prior and after the de-projection for one velocity map together with

the corresponding map of cos . On the center side, the LOS is roughly aligned with the magnetic field, and so there is lit-tle change to the flow speed apart from the sign. On the limb side, there is some extended area where the LOS was perpen-dicular to the magnetic field and the values of both the observed LOS velocities and of cos are very small. We decided not to apply the de-projection to all pixels (x;y) with j cos j <0:1 or an unsigned LOS velocity jvj  $<0:2\,km\,s^{-1}$  to avoid introducing spurious high velocities, but set cos to 1 and the LOS veloci-ties to zero at those places. With the lower threshold of 0.1, the deprojection thus increased the LOS flow speeds by a factor of 1–10 depending on the location in the FOV.

#### D.3. De-projection and mapping of full-disk data

For the NFFF extrapolations, we used the "hmi.sharp\_cea\_720s" data series from HMI, which provides the three components of the magnetic field remapped onto a heliographic cylindrical equal-area (CEA) coordinate system centered on an active region cutout (Bobra et al. 2014). The original FOV downloaded from JSOC<sup>1</sup> consisted of 1089541 pixels centered at 192.51 and

15:58 Carrington longitude and latitude, respectively. This was then cropped to 1024512 pixels by shifting the origin to (60,10) pixels to improve the numerical accuracy. The full 3D domain for the extrapolation then consists of 1024 512 512 pixels in the x, y; and z directions, respectively. With the 0 $\alpha$ 5 pixel <sup>1</sup> sampling of HMI, the horizontal extent of the box in x is 371 Mm. All other SDO/AIA filtergrams were also CEA projected and remapped to the same spatial sampling as the magnetic field data with the same FOV.

## D.4. Alignment of high-resolution and full-disk data

The de-projected IBIS data (13651000 pixels) were degraded to the HMI sampling of  $\mathfrak{G}$ :5 pixel  $^1$  (262192 pixels). The resampled IBIS data were then placed into an empty array of 1024512 pixels that matched the size of the FOV used in the extrapolation. The appropriate location for the IBIS images was determined from a visual comparison of the outer penumbral contour line in the HMI and IBIS continuum intensity images. The final images have a common spatial (x; y) coordinate system for all quantities and are aligned to pixel precision at the HMI spatial sampling (rightmost column of Figure 1).

http://jsoc.stanford.edu/ajax/lookdata.html