

EULER–KRONECKER CONSTANTS FOR CYCLOTOMIC FIELDS

LETONG HONG, KEN ONO, AND SHENG TONG ZHANG

ABSTRACT. The Euler-Mascheroni constant $\gamma = 0.5772\dots$ is the $K = \mathbb{Q}$ example of an Euler-Kronecker constant γ_K of a number field K . In this note we consider the size of the $\gamma_q = \gamma_{K_q}$ for cyclotomic fields $K_q := \mathbb{Q}(\zeta_q)$. Assuming the Elliott-Halberstam Conjecture (EH), we prove uniformly in Q that

$$\frac{1}{Q} \sum_{Q < q \leq 2Q} |\gamma_q - \log q| = o(\log Q).$$

In other words, under EH the $\gamma_q / \log q$ in these ranges converge to the one point distribution at 1. This theorem refines and extends a previous result of Ford, Luca, and Moree for prime q . The proof of this result is a straightforward modification of earlier work of Fouvry under the assumption of EH.

1. INTRODUCTION

For a number field K , the *Euler-Kronecker constant* γ_K is given by

$$\gamma_K := \lim_{s \rightarrow 1^+} \left(\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} \right),$$

where $\zeta_K(s)$ is the Dedekind zeta-function for K . The Euler-Mascheroni constant $\gamma = 0.5772\dots$ is the $K = \mathbb{Q}$ case, where $\zeta_{\mathbb{Q}}(s) = \zeta(s)$ is the Riemann zeta-function. We consider the constants $\gamma_q = \gamma_{K_q}$ for cyclotomic fields $K_q := \mathbb{Q}(\zeta_q)$, where $q \in \mathbb{Z}^+$ and ζ_q is a primitive q th root of unity.

The recent interest in the distribution of the γ_q is inspired by work of Ihara [4, 5]. He proposed, for every $\varepsilon > 0$, that there is a $Q(\varepsilon)$ for which

$$(c_1 - \varepsilon) \log q \leq \gamma_q \leq (c_2 + \varepsilon) \log q$$

for every integer $q \geq Q(\varepsilon)$, where $0 < c_1 \leq c_2 < 2$ are absolute constants. This conjecture was disproved by Ford, Luca and Moree in [2] assuming a strong form of the Hardy–Littlewood k -tuple Conjecture. However, assuming the Elliott-Halberstam Conjecture (see [1]), these same authors also proved that the conjecture holds for almost all primes q , with $c_1 = c_2 = 1$. We recall the Elliott-Halberstam Conjecture as formulated in terms of the Von Mangoldt function $\Lambda(n)$, the Chebyshev function $\psi(x)$, and Euler's totient function $\phi(n)$.

Conjecture EH. If we let

$$E(m; x, a) := \sum_{\substack{p \equiv a \pmod{m} \\ p \leq x \text{ prime}}} \Lambda(p) - \frac{\psi(x)}{\varphi(m)},$$

then for every $\varepsilon > 0$ and $A > 0$, we have

$$\sum_{m \leq x^{1-\varepsilon}} \max_{(a,m)=1} |E(m; x, a)| \ll_{A,\varepsilon} \frac{x}{(\log x)^A}.$$

Date: April 19, 2022.

Key words and phrases. Cyclotomic fields; Elliott-Halberstam Conjecture; Euler-Kronecker constants.

2010 *Mathematics Subject Classification.* 11M06, 11N37, 11R18, 11R42, 11Y60.

Assuming EH, Ford, Luca, and Moree proved (see Theorem 6 (i) of [2]), for every $\varepsilon > 0$, that

$$1 - \varepsilon < \frac{\gamma_q}{\log q} < 1 + \varepsilon$$

for almost all primes q (that is, the number of exceptional $q \leq x$ is $o(\pi(x))$ as $x \rightarrow \infty$). Here we extend and refine this result to all integers q .

Theorem 1.1. *Under EH, for $Q \rightarrow +\infty$ we have*

$$\frac{1}{Q} \sum_{Q < q \leq 2Q} |\gamma_q - \log q| = o(\log Q).$$

Remark. Theorem 1.1 shows that EH implies that the distribution of $\gamma_q / \log q$ in $[Q, 2Q]$ converges to the one point distribution supported on 1.

To prove Theorem 1.1, we use of work of Fouvry [3] that allowed him to unconditionally prove that

$$\frac{1}{Q} \sum_{Q < q \leq 2Q} \gamma_q = \log Q + O(\log \log Q).$$

Our conditional result is a point-wise refinement of Fouvry's asymptotic formula under EH.

2. ACKNOWLEDGEMENTS

The authors acknowledge the Thomas Jefferson Fund and the NSF (DMS-2002265 and DMS-2055118) for their generous support. The authors thanks Pieter Moree for helpful discussions regarding his work with Ford and Luca. We thank the referee for suggestions that improved the exposition in this paper.

3. PROOF OF THEOREM 1.1

For brevity, we shall assume that the reader is familiar with Fouvry's paper [3]. The key formula is (see (3) of [3]) the following expression for γ_q in terms of logarithmic derivatives of Dirichlet L -functions:

$$(1) \quad \gamma_q = \gamma + \sum_{1 < q^* | q} \sum_{\chi^* \bmod q^*} \frac{L'(1, \chi^*)}{L(1, \chi^*)}.$$

Here the inner sum runs over the primitive Dirichlet characters χ^* modulo q^* .

We follow the strategy and notation in [3]. However, we replace the sums $\Gamma_i(Q)$ and $\Gamma_{1,j}(Q)$ defined in [3] with the pointwise terms $\gamma_i(q)$ and $\gamma_{1,j}(q)$. Following the approach in [3], which is based on (1), we have

$$\gamma_q = \gamma + A(q) + B(q) - \gamma_2(q) - \gamma_3(q) - (\gamma_{1,1}(q) + \gamma_{1,2}(q) + \gamma_{1,3}(q)),$$

where

$$\begin{aligned} A(q) &= \sum_{q^* | q} \sum_{\chi^* \bmod q^*} \frac{L'}{L}(1, \chi^*) + \Phi_{\chi^*}(x), \\ B(q) &= \sum_{\substack{\chi \bmod q \\ \chi \neq \chi_0}} \Phi_{\chi}(x) - \sum_{q^* | q} \sum_{\chi^* \bmod q^*} \Phi_{\chi^*}(x), \\ \gamma_2(q) &= \frac{1}{x-1} \int_1^x \frac{\varphi(q)\psi(t; q, 1) - \psi(t)}{t} dt, \end{aligned}$$

$$\begin{aligned}
\gamma_3(q) &= \frac{1}{x-1} \int_1^x \sum_{\substack{n \leq t \\ (n,q) \neq 1}} \frac{\Lambda(n)}{n} dt, \\
\gamma_{1,1}(q) &= \frac{1}{x-1} \int_1^x \int_1^{\min(q,t)} \left(\frac{\varphi(q)\psi(u;q,1) - \psi(u)}{u^2} du \right) dt, \\
\gamma_{1,2}(q) &= \frac{1}{x-1} \int_1^x \int_{\min(q,t)}^{\min(x_1,t)} \left(\frac{\varphi(q)\psi(u;q,1) - \psi(u)}{u^2} du \right) dt, \\
\gamma_{1,3}(q) &= \frac{1}{x-1} \int_1^x \int_{\min(x_1,t)}^t \left(\frac{\varphi(q)\psi(u;q,1) - \psi(u)}{u^2} du \right) dt.
\end{aligned}$$

To complete the proof, for $\varepsilon > 0$ we let $x := q^{100}$ and $x_1 := q^{1+\varepsilon}$. Apart from $\gamma_{1,1}(q)$, which gives the $-\log q$ terms in Theorem 1.1, we shall show that these summands are all small.

Estimation of $A(q)$: By Proposition 1 and Remark (i) of [3], we have

$$\sum_{q=Q}^{2Q} |A(q)| = O(Q).$$

Estimation of $B(q)$: For $B(q)$, by equation (26) and Lemma 3 of [3], we simplify

$$\begin{aligned}
B(q) &= -\frac{1}{x-1} \int_1^x \sum_{q^*|q} \sum_{\chi^* \bmod q^*} \sum_{\substack{n \leq t \\ (n,q) > 1}} \frac{\Lambda(n)\chi^*(n)}{n} dt \\
&= -\frac{1}{x-1} \int_1^x \sum_{q^*|q} \sum_{\chi^* \bmod q^*} \sum_{\substack{p^v \leq t \\ p|q}} \frac{\log p \cdot \chi^*(p^v)}{p^v} dt \\
&= -\frac{1}{x-1} \int_1^x \sum_{q^*|q} \sum_{\substack{p^v \leq t \\ p|q}} \sum_{d|(p^v-1, q^*)} \frac{\log p}{p^v} \cdot \varphi(d) \mu\left(\frac{q^*}{d}\right) dt \\
&= -\frac{1}{x-1} \int_1^x \sum_{\substack{p^v \leq t \\ p|q}} \sum_{d|p^v-1} \frac{\log p}{p^v} \cdot \varphi(d) \sum_{\substack{q^*|q \\ d|q^* \\ p \nmid q^*}} \mu\left(\frac{q^*}{d}\right) dt.
\end{aligned}$$

We note that the innermost sum

$$\sum_{\substack{q^*|q \\ d|q^* \\ p \nmid q^*}} \mu\left(\frac{q^*}{d}\right)$$

is always 0 or 1, so we conclude that $B(q) \leq 0$ for any q . Proposition 2 of [3] gives

$$\sum_{q=Q}^{2Q} B(q) = O(Q),$$

and so we have

$$\sum_{q=Q}^{2Q} |B(q)| = O(Q).$$

Estimation of $\gamma_2(q)$: By Lemma 8 of [3], uniformly in Q we have

$$\sum_{q=Q}^{2Q} \psi(u; q, 1) \ll u.$$

Therefore, we have that

$$\sum_{q=Q}^{2Q} |\varphi(q)\psi(t; q, 1) - \psi(t)| = O(Qt),$$

and so we conclude that

$$\sum_{q=Q}^{2Q} |\gamma_2(q)| = O(Q).$$

Estimation of $\gamma_3(q)$: By definition, γ_3 is positive, so by equation (36) of [3], we have

$$\sum_{q=Q}^{2Q} |\gamma_3(q)| = O(Q).$$

Estimation of $\gamma_{1,1}(q)$: Since $\varphi(u; q, 1) = 0$ for $u < q$, we have

$$\gamma_{1,1}(q) = -\frac{1}{x-1} \int_1^x \left(\int_1^{\min(q,t)} \frac{\psi(u)}{u^2} du \right) dt.$$

Dividing both sides of Equation (41) of [3] by Q , we have

$$\gamma_{1,1}(q) = -\log q + O(1).$$

Estimation of $\gamma_{1,2}(q)$: By the same proof as equation (42) of [3], we have

$$\sum_{q=Q}^{2Q} |\gamma_{1,2}(q)| \ll \varepsilon Q \log Q.$$

Summing the above estimates, we conclude unconditionally that

$$\frac{1}{Q} \sum_{q=Q}^{2Q} |\gamma_q - \log q| = \frac{1}{Q} \sum_{q=Q}^{2Q} |\gamma_{1,3}(q)| + O(\varepsilon \log Q).$$

Estimation of $\gamma_{1,3}(q)$: If we assume Conjecture EH holds, then we have (as in Lemma 7 of [3]) that

$$\sum_{\substack{q \leq 2Q \\ (q,a)=1}} \varphi(q) \left| \psi(x; q, a) - \frac{\psi(x)}{\varphi(q)} \right| = O_A(Qx(\log x)^{-A+2}).$$

Therefore, we find that

$$\frac{1}{Q} \sum_{q=Q}^{2Q} |\gamma_{1,3}(q)| = O_{\varepsilon,A}(\log^{-A} Q).$$

By combining these estimates, we obtain the main result

$$\frac{1}{Q} \sum_{q=Q}^{2Q} |\gamma_q - \log q| = o(\log Q),$$

REFERENCES

- [1] P.D.T.A. Elliott and H. Halberstam, *A conjecture in prime number theory*, Symp. Math. 4, 1968–69, 59–71.
- [2] K. Ford, F. Luca, and P. Moree, *Values of the Euler φ -function not divisible by a given odd prime, and the distribution of Euler–Kronecker constants for cyclotomic fields*, Math. Comput. Vol. 83, No. 287 (May 2014), pp. 1447–1476.
- [3] É. Fouvry, *Sum of Euler–Kronecker constants over consecutive cyclotomic fields*, J. Number Theory. (4) 133, 2013, 1346–1361,
- [4] Y. Ihara, *On the Euler–Kronecker constants of global fields and primes with small norms*, in V. Ginzburg, ed., Algebraic Geometry and Number Theory: In Honor of Vladimir Drinfeld’s 50th Birthday, Progress in Mathematics, Vol. 850, Birkhäuser Boston, Cambridge, MA, 2006, 407–451.
- [5] Y. Ihara, *The Euler–Kronecker invariants in various families of global fields*, Proc. of AGCT 2005 (Arithmetic Geometry and Coding Theory 10), Ed. F. Rodier et al., Séminaires et Congrès. 21, 2009, 79–102.

DEPT. OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139
Email address: `clhong@mit.edu`

DEPT. OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904
Email address: `ko5wk@virginia.edu`

DEPT. OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139
Email address: `stzh1555@mit.edu`