

There are no exotic actions of diffeomorphism groups on 1-manifolds

Lei Chen and Kathryn Mann

Abstract. Let M be a manifold and N a 1-dimensional manifold. Assuming that $r \neq \dim M / \mathbb{C} 1$, we show that any nontrivial homomorphism $\varphi: \text{Diff}_c^r(M) \rightarrow \text{Homeo}(N)$ has a standard form: necessarily M is 1-dimensional, and there are countably many embeddings $\iota_i: M \hookrightarrow N$ with disjoint images such that the action of φ is conjugate (via the product of the ι_i) to the diagonal action of $\text{Diff}_c^r(M)$ on $M \times \prod \iota_i(M)$, and trivial elsewhere. This solves a conjecture of Matsumoto. We also show that the groups $\text{Diff}_c^r(M)$ have no countable index subgroups.

1. Introduction

Let $\text{Diff}_c^r(M)$ denote the identity component (in the compact-open C^r topology) of the group of compactly supported C^r diffeomorphisms of a manifold M , for $0 \leq r \leq 1$. These groups are locally contractible, so in fact $\text{Diff}_c^r(M)$ agrees with the group of diffeomorphisms which are isotopic to the identity through a compactly supported isotopy. When we speak of $\text{Diff}_c^r(M)$, we assume that manifolds admit a C^r structure, and a metric structure in the C^0 case, but are otherwise arbitrary. In this paper, we prove the following statement.

Theorem 1.1. Let M be a connected manifold, and $\varphi: \text{Diff}_c^r(M) \rightarrow \text{Homeo}(N)$ is a non-trivial homomorphism, where $N \cong S^1$ or $N \cong \mathbb{R}$, $r \neq \dim M / \mathbb{C} 1$. Then $\dim M \leq 1$ and there are countably many disjoint embeddings $\iota_i: M \hookrightarrow N$ such that $\varphi(g) \cdot \prod \iota_i(M) = \prod \iota_i(g \cdot M)$ and $N \setminus \prod \iota_i(M)$ is globally fixed by the action.

This proves [12, Conjecture 1.3] and generalizes works of Mann [8], Milonin [13], and Matsumoto [12], but with an independent proof. Matsumoto's work [12] proves an analogous result when the target is $\text{Diff}^1(N)$ using rigidity theorems of [3] for solvable affine subgroups of $\text{Diff}^1(\mathbb{R})$. This generalized [8], which proved the result for homomorphisms to $\text{Diff}^2(N)$ using Kopell's lemma. Milonin [13] studies homomorphisms where the source is the group of homeomorphisms of M . Our proof here is comparatively short, and is self-contained modulo the standard but difficult result that $\text{Diff}_c^r(M)$, for $r \neq \dim M / \mathbb{C} 1$, is a simple group, due to Anderson, Mather, and Thurston [1, 10, 11, 18]. Whether simplicity holds for $r \leq \dim M / \mathbb{C} 1$ is an open question; this is responsible for our restrictions on dimension in the statement.

Theorem 1.1 is already known in the case where γ is assumed to be continuous; it is a consequence of the orbit classification theorem of [5], and was likely known to others before. In the case where the target is the group of smooth diffeomorphisms of $\mathbb{N}$, this also follows from work of Hurtado [6] who proves additionally that any such homomorphism is necessarily (weakly) continuous. Here we make no assumptions on continuity, however, our proof suggests that diffeomorphism groups exhibit “automatic continuity”—like properties. Specifically, we show the following small index property.

Theorem 1.2 (The small index property of $\text{Diff}_c^r(M)$). If $r \neq \dim(M) + 1$, then $\text{Diff}_c^r(M)$ has no proper countable index subgroup. Equivalently, $\text{Diff}_c^r(M)$ has no nontrivial homomorphism to the permutation group S_1 .

This is in stark contrast with the case for finite dimensional Lie groups, where we have the following.

Theorem 1.3 (Thomas [17] and Kallman [7]). There is an injective homomorphism

$$SL_n(\mathbb{R}) \hookrightarrow S_1.$$

Thus, one consequence of Theorems 1.2 and 1.3 is that there is no nontrivial homomorphism from $\text{Diff}_c^r(M)$ into a linear group. Of course, this is nearly immediate if one considers only continuous homomorphisms, since $\text{Diff}_c^r(M)$ is infinite dimensional, and one may simply quote the invariance of domain theorem.

If G is a group with a non-open subgroup H of countable index, then the action of G on the coset space G/H gives a discontinuous homomorphism to S_1 . This is one of very few known general recipes for producing discontinuous group homomorphisms (see [16]), so it gives some (weak) evidence that $\text{Diff}_c^r(M)$ might have the automatic continuity property already known to hold for $\text{Homeo}_c(M)$ by [9]. Automatic continuity also holds for homomorphisms between groups of smooth diffeomorphisms by work of Hurtado [6].

Theorem 1.1 also gives new examples of left orderable groups that do not act on the line. It is a well-known fact that any countable group with a left-invariant total order admits a faithful homomorphism to $\text{Homeo}_c(\mathbb{R})$. For $r > 0$, the groups $\text{Diff}_c^r(\mathbb{R}^n)$ for $r > 0$ are known to be left-orderable: the Thurston stability theorem [19] implies that they are locally indicable (any finitely generated subgroup surjects to $\mathbb{Z}$), which implies that they are left-orderable by the Burns-Hale theorem ([4], see also [14, Corollary 2]). Thus, we have the following.

Corollary 1.4. For $r > 0$, the group $\text{Diff}_c^r(\mathbb{R}^n)$ is left-orderable but has no faithful action on the line or the circle.

The proof of Theorem 1.2 uses the idea from the first step of the proof of automatic continuity for homeomorphism groups of [9], following Rosendal [15]. This result is then used to prove Theorem 1.1 by constraining the supports and fixed sets of elements for the action on $\mathbb{N}$. We are then able to use this information to build a map from M to $\mathbb{N}$.

2. Proof of the small index property

In this section, we prove Theorem 1.2. The proof follows a strategy in [9, 15] used in the proof of automatic continuity of $\text{Homeo.M}/$.

Proof. Let M be a manifold and $r \propto \dim.M/C \geq 1$. Let $G \subset \text{Diff}_c^r.M/$, and for an open subset $U \subset M$, denote by G_U the subgroup of $\text{Diff}_c^r.M/$ consisting of maps with compact support contained in U and isotopic to the identity via an isotopy compactly supported in U . Thus, $G_U \cong \text{Diff}_c^r.U/$. (Note that $\text{Diff}_c^r.U/$ is locally contractible, and in particular path connected, for all $0 < r \leq 1$.)

Suppose for contradiction that $H \subset G$ is a countable index subgroup. We will show in Step 1 that there is some ball U in M such that $G_U \subset H$. After this, we will show (Step 2) that H acts transitively on M , thus every $x \in M$ is contained in some open set U_x such that $G_{U_x} \subset H$. The fragmentation property states that $\text{Diff}_c^r.M/$ is generated by the union of such sets G_{U_x} (this is true for any collection of sets U_x which form an open cover of M ; see [2, Chapter 1]), so this is sufficient to prove $H = G$.

Step 1: there is some open ball U in M such that $G_U \subset H$. Let $g_1H; g_2H; \dots$ denote the left cosets of H . Let $B \subset M$ be an open ball, and take a sequence of disjoint balls $B_i \subset B$ such that $\bigcup_{i=1}^{\infty} B_i = B$, with diameter tending to 0 and such that the sequence B_i Hausdorff converges to a point inside B .

We first claim that there exists some $j \in \mathbb{N}$ and a neighborhood U_j of the identity element of G_{B_j} such that the following holds:

- (i) for every $f \in U_j$, there exists $w \in G \setminus G_B$ such that the restriction of w to U_j agrees with f .

Given (i), then we have $w_{\text{id}}^{-1}w \in H \cap g_j^{-1}g_jH \subset H$, and $w_{\text{id}}^{-1}w$ restricts to f on B_j . This shows that every element in U_j agrees with the restriction of an element of H to B_j . Since U_j is an identity neighborhood of G_{B_j} and G_{B_j} is by definition connected, U_j generates G_{B_j} and we conclude that every element of G_{B_j} agrees with the restriction of an element of H to B_j .

We prove this claim by contradiction, using a standard diagonal argument. Inductively, choose neighborhoods U_i of the identity in G_{B_i} so that for any sequence of diffeomorphisms $f_i \in U_i$, the infinite composition $\bigcap_i f_i$ defines an element of G . Supposing that our claim is not true for any U_j , then for each i we can find $f_i \in U_i$ such that there does not exist any $w_i \in G_jH$ supported in B satisfying $w_i|_{B_i} = f_i|_{B_i}$. Let $w \in \bigcap_i U_i f_i$. Then $w \in G_jH$ for some j since $\bigcup_k G_kH = G$. Moreover, the support of w is in B , the restriction of w and that of f on B_j are the same and we have $w \in G_jH$. This is a contradiction and proves the claim.

Now we use a commutator trick. Apply the same argument as above using B_j in place of B . We find a smaller ball $B^0 \subset B_j$ such that every element $f \in G_{B^0}$ agrees with the restriction to B^0 of an element $\psi \in H$, and ψ is supported on B_j . Since $\text{Diff}_c^r.B^0/$ is perfect [1, 10, 11, 18], any element $f \in \text{Diff}_c^r.B^0/$ may be written as a product of commutators $f = \bigcap_{i=1}^k [a_i; b_i]$. The commutator length k of course depends on f , but this is

unimportant to us. We have $\{e_{a_i}; b_i\} \subset \{e_{v_{a_i}}; w_{b_i}\}$ since the supports of v_{a_i} and w_{b_i} intersect only in B , and so $f \in \{e_{v_{a_i}}; w_{b_i}\} \subset H$. This ends the proof of the first step.

Step 2: transitivity. To prove transitivity, let B^0 be the ball from Step 1, and let $x \in B^0$. Suppose that $y \in M$ is some point not in the orbit of x . Let f_t be a flow such that $f_t \cdot y \in B^0$ for all $t \in [0, 2]$. Such a flow can be defined to have support on a neighborhood of a path from x to y . Since B^0 lies in the orbit of x under H , we have that $f_t \in H$ for $t \in [0, 2]$. We know that $H \cap \{f_t \mid t \in \mathbb{R}\}$ is a countable index subgroup of $\{f_t \mid t \in \mathbb{R}\} \cong \mathbb{R}$. Thus, it must intersect every open interval of $\mathbb{R}$; this gives the desired contradiction. As explained above, Steps 1 and 2 together with fragmentation complete the proof of Theorem 1.2. ■

As an immediate consequence, we can conclude that any fixed point free action of such a group on the line or circle is minimal.

Corollary 2.1. With the same restrictions on r as above, if $\text{Diff}_c(M)/\langle f \rangle$ acts on $\mathbb{R}$ or S^1 without global fixed points, then there are no invariant open sets. In particular, every orbit is dense.

Proof. Suppose that the action has an invariant open set. Then $\text{Diff}_c(M)/\langle f \rangle$ permutes the (countably many) connected components of U . The stabilizer of an interval is a countable index subgroup, so, by Theorem 1.2, the permutation action is trivial. Thus each interval is fixed and their endpoints are global fixed points. ■

3. Proof of Theorem 1.1

For the proof of Theorem 1.1, we set the following notation. As in the previous section, we fix some $r \leq \dim(M)/C - 1$ and when $U \subset M$ is an open set we denote by G_U the subgroup of $\text{Diff}_c(M)/\langle f \rangle$ consisting of maps with compact support contained in U and isotopic to the identity via an isotopy compactly supported in U . We additionally use the notation $G^U = \text{Diff}_c(M)/\langle f \rangle$ for the set of elements that pointwise fix U . The open support of a homeomorphism g is the set $\text{Osupp}(g) = \text{supp}(g) \setminus \text{Fix}(g)$; as is standard, the support of g is defined to be the closure of $\text{Osupp}(g)$.

Proof. We will assume that the action on N has no global fixed points, since if the action does have fixed points, then $N \setminus \text{Fix}(f)$ is a union of open intervals, each with a fixed-point free action of $\text{Diff}_c(M)/\langle f \rangle$, so it suffices to understand such actions. In this case, we will show that there is a single homeomorphism $\varphi: M \rightarrow N$ such that the action on N is induced by conjugation by φ .

Lemma 3.1. For any action, if $U \cap V = \emptyset$, then $\text{Osupp}(G_U) \cap \text{Osupp}(G_V) = \emptyset$.

Proof. Since G_U and G_V commute, G_V preserves $\text{Osupp}(G_U)$, permuting its connected components. By Theorem 1.2, this action is trivial. Let I be a connected component of $\text{Osupp}(G_U)$. Suppose that G_V acts nontrivially on I . Since G_V is a simple group, its action on I is faithful. Since G_V is not abelian, Hölder's theorem implies that some nontrivial $g \in G_V$ acts with a fixed point. But then G_U permutes the connected

components of $I \setminus \text{Osupp} \cdot g //$, and this permutation action is trivial. Thus, $\cdot G_U /$ has a fixed point in I , contradicting that $I \setminus \text{Osupp} \cdot G_U //$. ■

We observe the following consequence of the fragmentation property:

Observation 3.2. If $U \setminus W \neq \emptyset$, then G^U and G^W generate $\text{Diff}_c M /$ because $G^U \cap G^W = G^{U \cap W}$ and $G^U \cup G^W = G^{U \cup W}$, and $U \cap W$ and $U \cup W$ cover M . Consequently, our assumption that there are no global fixed points for the action implies that

$$\text{Fix} \cdot G^U / \cap \text{Fix} \cdot G^W / = \emptyset ;$$

Our next goal is to define a map from M to N . For each $x \in M$, pick a neighborhood basis $U_n(x)$ of x so $\bigcap_n U_n(x) = \{x\}$. Let $S_x = \bigcap_n \text{Osupp} \cdot G_{U_n(x)} //$ and let $T_x = \bigcap_n \text{Fix} \cdot G_{U_n(x)} //$. Note that the sets S_x and T_x are independent of the choice of neighborhood basis.

Lemma 3.3. If $x \neq y$, then $S_x \cap S_y = \emptyset$; and $T_x \cap T_y = \emptyset$. Also, S_x and T_x have empty interior.

Proof. The first assertion follows immediately from Lemma 3.1 and the second because $T_x \cap T_y$ would be globally fixed by $\cdot g /$ by our observation above. Furthermore, if $g \cdot x \in D_y$, then $\cdot g / \cdot U_n(x) //$ is a neighborhood basis of y , so we have

$$\begin{aligned} \cdot g / S_x &= \bigcap_n \cdot g / \text{Osupp} \cdot G_{U_n(x)} / \cap D \\ &= \bigcap_n \text{Osupp} \cdot g G_{U_n(x)} g^{-1} \\ &= \bigcap_n \text{Osupp} \cdot G_{g \cdot U_n(x)} / \cap D = S_y ; \end{aligned}$$

Similarly, we have $T_y \subset \cdot g / T_x$. Thus, if some S_x has nonempty interior, disjointness of S_x and S_y would give an uncountable family of disjoint open sets in N , a contradiction. The same applies to the sets T_x . ■

We next prove that these sets, though defined differently, are in fact the same.

Lemma 3.4. For all x , we have $S_x = T_x$.

Proof. Fix x and let $U_n \subset U_n(x)$ be a neighborhood basis of x with the property that $U_n \subset U_{n+1}$ for all n . Thus, by Lemma 3.1, $\cdot G_{U_{n+1}} /$ and $\cdot G_{M \setminus U_n} /$ have disjoint open supports. Since $G^M = \overline{G^{U_n}}$ we conclude that

$$\text{Osupp} \cdot G_{U_{n+1}} / \cap D = \text{Fix} \cdot G_{U_{n+1}} / \cap \text{Fix} \cdot G_{M \setminus U_n} / \cap \text{Fix} \cdot G^{U_n} / ;$$

Also, since U_n and $M \setminus U_{n-1}$ have disjoint closures, Observation 3.2 implies that $\text{Fix} \cdot G^{U_n} / \cap \text{Fix} \cdot G^{M \setminus U_{n-1}} / \cap D = \emptyset$, so

$$\text{Fix} \cdot G^{U_n} / \cap \text{Osupp} \cdot G^{M \setminus U_{n-1}} / \cap \text{Osupp} \cdot G_{U_{n-2}} / ;$$

Combining the two equations above and taking a limit as $n \rightarrow \infty$ shows that $S_x = T_x$, as desired.

Thus $S_x \subset T_x$. For the reverse inclusion, suppose that $z \in T_x \setminus S_x$. Then z

$$\dots \text{Osupp} \cdot G_{U_n} /$$

for some n ; i.e., $z \in \text{Fix} \cdot G_{U_n} /$. Also $z \in \text{Fix} \cdot G^{U_{nc1}} /$ by the definition of T_x . But G_{U_n} and $G^{U_{nc1}}$ together generate $\text{Diff}^f \cdot M /$ (this again is the fragmentation property), so this implies that z is a global fixed point. ■

Lemma 3.5. S_x is nonempty.

Proof. If $N \subset S^1$, this follows immediately since $S_x \subset T_x$ is the intersection of nested, nonempty closed sets. If $N \subset \mathbb{R}$, the same is true provided that $\text{Fix} \cdot G^{U_n \cdot x} /$ (or equivalently $\text{Osupp} \cdot G_{U_n \cdot x} /$) does not leave every compact set as $n \rightarrow \infty$. Note that this holds for some x if and only if it holds for all x because $g/S_x \subset S_y$ when $g \cdot x / \subset y$.

We proceed by contradiction. Suppose that, for each $x \in M$, as $n \rightarrow \infty$ we have that $\text{Osupp} \cdot G_{U_n \cdot x} /$ does leave every compact set. Fixing some compact $K \subset \mathbb{R}$, this means that for each $x \in M$ there is a neighborhood $U \cdot x /$ of x such that

$$\text{Osupp} \cdot G_{U \cdot x} / \cap K \subset \emptyset ;$$

Let $\mathcal{O}$ denote the open cover formed by such sets $U \cdot x /$. By fragmentation, $\text{Diff}_c \cdot M /$ is generated by the subgroups $G \cdot U \cdot x /$. Thus, $\text{Osupp} \cdot \text{Diff}_c \cdot M / \cap K \subset \emptyset$, contradicting the fact that M has no global fixed points. ■

Construction of φ . To finish the proof, we wish to show that S_x is a singleton, and the assignment $\varphi : S_x \rightarrow M$ is a homeomorphism conjugating φ with the standard action of $\text{Diff}_c \cdot M /$ on M . We will actually show first that $x \mapsto S_x$ is a local homeomorphism, use this to conclude that S_x is discrete, and proceed from there.

Step 1: definition of φ locally. Let $I \subset \mathbb{R}$; a, b be a connected component of $N \cap S_x$, chosen so that $a \neq b$ if $N \subset \mathbb{R}$. If $N \subset S^1$ and S_x is a singleton, it is possible that both “endpoints” of this interval agree. For simplicity, we treat the case where $a \neq b$; the case $a = b$ on the circle can be handled with exactly the same strategy and in fact the argument simplifies quite a bit since S_x is already a singleton.

Fix a neighborhood basis $U_n \subset U_{nc1}$ of x . For $n \in \mathbb{N}$, denote by O_n the connected component of $\text{Osupp} \cdot G_{U_n} /$ that contains a . Since $\bigcap_k O_k \subset S_x$ and it contains a , and since $a, b \in N \cap S_x$, we can conclude that for all k sufficiently large a is the right-most point of $S_x \cap O_k$.

Fix such a k . We will show that, for $y \in U_k$, the set $S_y \cap O_k$ also has a rightmost point. This allows us to define a map from U_k to O_k , sending y to this rightmost point, which we will then show to be the desired local homeomorphism. First, to see that $S_y \cap O_k$ has a rightmost point, take some $g \in G_{U_k}$ with $g \cdot x / \subset y$. Thus $g/S_x \subset S_y$. Since g fixes endpoints of O_k by definition, we know that $g \cdot a / \in S_y$ and it is the rightmost point of $S_y \cap O_k$. This proves our claim.

Define $\varphi : U_k \rightarrow O_k$ by setting $\varphi(y)$ to be the rightmost point of $S_y \cap O_k$. An equivalent definition of φ is that $\varphi(y) = D \cdot g \cdot a /$, where g is any diffeomorphism in G_{U_k} such

that $g.x / D y$. Our argument above shows that this is independent of choice of g . Furthermore, if we repeat the definition using U_{kC1} instead of U_k , the map we will obtain is simply the restriction of π to U_{kC1} .

Step 2: local continuity of π on U_k . We first show that π is continuous at x . Suppose that $x_n \rightarrow x$ is a convergent sequence. Passing to a subsequence and reindexing if needed, we may assume that $x_n \in U_n$ and that our index set starts at k . Then we may take $g \in G_{U_n}$ so that $g.x / D x_n$, so $x_n / D .g/.a/$. Since the sequence of connected components of $O\text{supp}..G_{U_n} //$ containing x converges to x , we get that $x_n / \rightarrow a$.

To show that π is continuous on U_k , let $x^0 \in U_k$, and take a sequence $x_n^0 \rightarrow x^0$ in U_k . There exists $g \in G_{U_k}$ such that $g.x / D x^0$ and $g^{-1}.x_n^0 /$ is a sequence converging to x . It follows from continuity at x that $.g^{-1}.x_n^0 /_n$ converges to $.x /$. By definition,

$$.g / g^{-1}.x_n^0 / D .x_n^0 /;$$

so we conclude that $.x_n^0 /_n$ converges to $.x^0 /$.

Note also that π is injective by Lemma 3.3. Thus, by invariance of domain, we conclude that M is one-dimensional so equal to $\mathbb{R}$ or S^1 , and π gives a homeomorphism from U_k onto an open interval A containing a in N . In particular, this shows that a is an isolated point of S_x .

Step 3: extension of π globally. The last step is to show that π extends to a globally defined homeomorphism $M \rightarrow N$; to do this we actually work with the inverse of π . First, note that the orbit of A under $.G /$ is an open, $.G /$ -invariant set, so by Corollary 2.1, $.G /.A / D N$.

This topological transitivity implies that, for all x , every point of S_x is an isolated point, i.e., S_x is discrete. Extend π^{-1} to a map $\tilde{\pi}$ defined on N by setting $\tilde{\pi}.S_x / D x$. The work in step 2 and the fact that $.g /.S_x / D S_{g.x /}$ implies that $\tilde{\pi}$ is a local homeomorphism, hence a covering map, and is equivariant with respect to the actions of $\text{Diff}_c^r.M /$ by its standard action on M and by π on N . If $M \cong \mathbb{R}$, we immediately conclude that $N \cong \mathbb{R}$, and $\tilde{\pi}$ conjugates π to the standard action of $\text{Diff}_c^r.\mathbb{R} /$.

If $M \cong S^1$, we can also conclude that $N \cong S^1$ because $\text{Diff}_c^r.S^1 /$ contains torsion, so cannot faithfully act on $\mathbb{R}$. Thus, $\bigcup_x S_x \rightarrow N$ is a finite cover, and $\tilde{\pi}$ is a lift of the standard action of $\text{Diff}_c.S^1 /$ on S^1 . Identifying the rotation subgroup $SO(2) /$ with S^1 , and considering $.SO(2) //$ which is a continuous lift, covering space theory tells us the degree of the cover must be 1. Alternatively, one can derive a contradiction by looking at the action of finite order elements: an order two rotation lifted to a degree d cover will have order $2d$. ■

Acknowledgments. The authors would like to thank an anonymous referee for the help with the writing.

Funding. Research of the first author supported by NSF 2005409. Research of the second author supported by DMS-1844516 and a Sloan fellowship.

References

- [1] R. D. Anderson, The algebraic simplicity of certain groups of homeomorphisms. *Amer. J. Math.* 80 (1958), 955–963 Zbl [0090.38802](#) MR [98145](#)
- [2] A. Banyaga, The structure of classical diffeomorphism groups. *Math. Appl.* 400, Kluwer Academic Publishers Group, Dordrecht, 1997 Zbl [0874.58005](#) MR [1445290](#)
- [3] C. Bonatti, I. Monteverde, A. Navas, and C. Rivas, Rigidity for C^1 actions on the interval arising from hyperbolicity I: solvable groups. *Math. Z.* 286 (2017), no. 3–4, 919–949 Zbl [1433.37030](#) MR [3671566](#)
- [4] R. G. Burns and V. W. D. Hale, A note on group rings of certain torsion-free groups. *Canad. Math. Bull.* 15 (1972), 441–445 Zbl [0244.16006](#) MR [310046](#)
- [5] L. Chen and K. Mann, Structure theorems for actions of homeomorphism groups. 2019, arXiv:[1902.05117](#)
- [6] S. Hurtado, Continuity of discrete homomorphisms of diffeomorphism groups. *Geom. Topol.* 19 (2015), no. 4, 2117–2154 Zbl [1322.57026](#) MR [3375524](#)
- [7] R. R. Kallman, Every reasonably sized matrix group is a subgroup of S_1 . *Fund. Math.* 164 (2000), no. 1, 35–40 Zbl [0967.20002](#) MR [1784652](#)
- [8] K. Mann, Homomorphisms between diffeomorphism groups. *Ergodic Theory Dynam. Systems* 35 (2015), no. 1, 192–214 Zbl [1311.57047](#) MR [3294298](#)
- [9] K. Mann, Automatic continuity for homeomorphism groups and applications. With an appendix by Frédéric Le Roux and Mann. *Geom. Topol.* 20 (2016), no. 5, 3033–3056 Zbl [1362.57044](#) MR [3556355](#)
- [10] J. N. Mather, Commutators of diffeomorphisms. *Comment. Math. Helv.* 49 (1974), 512–528 Zbl [0289.57014](#) MR [356129](#)
- [11] J. N. Mather, Commutators of diffeomorphisms. II. *Comment. Math. Helv.* 50 (1975), 33–40 Zbl [0299.58008](#) MR [375382](#)
- [12] S. Matsumoto, Actions of groups of diffeomorphisms on one-manifolds by C^1 diffeomorphisms. In *Geometry, dynamics, and foliations 2013*, pp. 441–451, *Adv. Stud. Pure Math.* 72, Math. Soc. Japan, Tokyo, 2017 Zbl [1388.57027](#) MR [3726723](#)
- [13] E. Militon, Actions of groups of homeomorphisms on one-manifolds. *Groups Geom. Dyn.* 10 (2016), no. 1, 45–63 Zbl [1373.57053](#) MR [3460330](#)
- [14] D. Rolfsen, A topological view of ordered groups. In *Knots in Poland III. Part III*, pp. 357–369, *Banach Center Publ.* 103, Polish Acad. Sci. Inst. Math., Warsaw, 2014 Zbl [1316.06020](#) MR [3363818](#)
- [15] C. Rosendal, Automatic continuity in homeomorphism groups of compact 2-manifolds. *Israel J. Math.* 166 (2008), 349–367 Zbl [1155.54025](#) MR [2430439](#)
- [16] C. Rosendal, Automatic continuity of group homomorphisms. *Bull. Symbolic Logic* 15 (2009), no. 2, 184–214 Zbl [1173.03037](#) MR [2535429](#)
- [17] S. Thomas, Infinite products of finite simple groups. II. *J. Group Theory* 2 (1999), no. 4, 401–434 Zbl [0938.20025](#) MR [1718758](#)
- [18] W. Thurston, Foliations and groups of diffeomorphisms. *Bull. Amer. Math. Soc.* 80 (1974), 304–307 Zbl [0295.57014](#) MR [339267](#)
- [19] W. P. Thurston, A generalization of the Reeb stability theorem. *Topology* 13 (1974), 347–352 Zbl [0305.57025](#) MR [356087](#)

Received 21 July 2020.

Lei Chen

Department of Mathematics, University of Maryland, College Park, MD 20742, USA;
chenlei@umd.edu

Kathryn Mann

Department of Mathematics, Cornell University, Ithaca, NY 14853, USA;
k.mann@cornell.edu