



COUNTEREXAMPLES TO THE COLORFUL TVERBERG CONJECTURE FOR HYPERPLANES

J. P. CARVALHO^{1,†} and P. SOBERÓN^{2,*,‡}

¹Haverford College, Haverford, PA 19041, USA
e-mail: jdecarvalh@haverford.edu

²Baruch College, City University of New York, New York, NY 10010, USA
e-mail: pablo.soberon-bravo@baruch.cuny.edu

(Received August 26, 2021; revised February 22, 2022; accepted May 7, 2022)

Abstract. Karasev [16] conjectured that for every set of r blue lines, r green lines, and r red lines in the plane, there exists a partition of them into r colorful triples whose induced triangles intersect. We disprove this conjecture for every r and extend the counterexamples to higher dimensions.

1. Introduction

Tverberg's theorem is a central result in combinatorial geometry. It states that *any set of $(r-1)(d+1)+1$ points in $\mathbb{R}^d$ can be partitioned into r sets whose convex hulls intersect*. It was proven by Helge Tverberg [28] and its generalizations and extensions have sparked significant interest in the area [7,11,20].

One of the most intriguing open questions regarding Tverberg's theorem is the colorful version. This was conjectured by Bárány and Larman.

CONJECTURE 1.1 (Bárány, Larman [4]). *Let r, d be positive integers. Given $d+1$ sets $X_1, \dots, X_{d+1}$, each consisting of r points of $\mathbb{R}^d$, there exists a partition of their union into r sets $A_1, \dots, A_r$ such that each A_j has exactly one point of each X_i and the convex hulls of $A_1, \dots, A_r$ intersect.*

* Corresponding author.

This project was done as part of the 2021 New York Discrete Math REU, funded by NSF grant DMS 2051026.

[†] Carvalho's research was supported by the KINSC Summer Scholar program by Haverford College.

[‡] Soberón's research is supported by NSF grant DMS 2054419 and a PSC-CUNY TRADB52 award.

Key words and phrases: Tverberg's theorem, hyperplane arrangement, combinatorial geometry.

Mathematics Subject Classification: 52A37, 52C35.

The conjecture above has been confirmed when $r + 1$ is prime [9,10], or when $d = 2$ [4]. If each X_i is allowed to have more than r points, there are plenty of positive results (see, e.g., [8,30] and the references therein).

One popular way to find variations of Tverberg's theorem is to replace the convex hull operator. The topological versions of Tverberg's theorem are the most notorious examples, in which one wants to show that any continuous mapping from an $(r - 1)(d + 1)$ -dimensional simplex to $\mathbb{R}^d$ has points covered by the image of r pairwise disjoint faces [6,24,29]. In this case, one must either introduce conditions on the parameters, such as being prime powers, or use a larger number of points [3,13,14,21]. Determining the validity of the topological Tverberg theorem for non-prime powers was a crucial problem in topological combinatorics (see [22]) and it was a surprise to the community the prime power conditions were necessary. Results whose proofs use similar methods from equivariant topology always bring the question of whether such conditions are needed. For instance see the progress of the Nandakumar–Ramana-Rao problem in measure partitions [1,12,18,26], or the cake-cutting partition results when one removes the “hungry guest” condition [2,23].

There are other ways to replace the convex hull in Tverberg's theorem. A recent example is if we consider for any two points the ball centered at their midpoint that contains both points in its boundary [15,27]. Huemer et al. showed that the natural extension of the colorful Tverberg theorem is valid in this setting in the plane. Another way to replace the convex hull is with Tverberg's theorem for hyperplanes.

In the hyperplane version, we are given sets of hyperplanes in $\mathbb{R}^d$ in general position instead of points. We say that a family of hyperplanes in $\mathbb{R}^d$ is in general position if the normal vectors of any d hyperplanes are linearly independent and no $d + 1$ hyperplanes have a point in common. With these conditions, any $d + 1$ hyperplanes uniquely define a simplex.

THEOREM 1.2 (Karasev [16]). *Let r be a prime power and d be a positive integer. Any family of $r(d + 1)$ hyperplanes in $\mathbb{R}^d$ in general position can be split into r sets of $d + 1$ hyperplanes each so that their induced simplices all intersect.*

The case $d = 2$ was proved earlier by Rousseeuw and Hubert without conditions on r [25]. Karasev also proved further extensions of the theorem above [17]. Bárány and Pach proved similar extensions of other results in combinatorial geometry for hyperplanes [5]. Karasev [16] conjectured that a colorful version of the Tverberg-type theorems for hyperplanes should hold similar to the Bárány–Larman conjecture.

CONJECTURE 1.3 (Karasev [16]). *Given r blue, r red, and r green lines in the plane in general position, it is possible to split them into r colorful triples whose induced triangles intersect.*

If r is a prime power, Karasev proved a relaxed version of the conjecture above if each family has $2r - 1$ lines but we only look for r colorful triples whose triangles intersect. This conjecture was previously known to be false when $r = 2m$ for odd m [19]. As Karasev's proofs of Theorem 1.2 and the cases for the colorful version use strong topological techniques, it is of interest to know if his conjecture remains valid under some conditions of the parameters.

In this note, we disprove Conjecture 1.3 for every r . Our examples work in higher dimensions, so we also disprove the natural extension to $\mathbb{R}^d$. In other words, Theorem 1.2 cannot be made colorful as in the Bárány–Larman conjecture. An interesting point is the structure of our counterexamples, which are based in hypercubes with two particular cross-sections, equipped with a doubling argument. We found it surprising that the dual of Lovász's parametrization [4] (an octahedron with opposite vertices of the same color) for the proof of the colorful Radon theorem helps one find the counterexamples for the dual of the problem.

THEOREM 1.4. *Let $r, d \geq 2$ be positive integers. There exists a family of $d + 1$ sets $X_1, \dots, X_{d+1}$, each consisting of r hyperplanes in $\mathbb{R}^d$, so that their union is in general position and the following statement holds. For any partition of their union into r sets $A_1, \dots, A_r$ so that each A_j has a hyperplane of each X_i , there are two sets $A_j, A_{j'}$ whose induced simplices do not intersect.*

In Section 3 we discuss the implications of our examples to the topological methods used by Karasev. Since his proof of Theorem 1.2 used the Borsuk–Ulam type result that combines Brouwer's fixed point theorem with the standard configuration space for Tverberg's theorem, no such extension can be done with the configuration space for the known cases of the colorful Tverberg theorem. We also hope that the doubling argument used in the examples can be useful in other results in combinatorial geometry.

2. Counterexamples

We first construct the counterexample in dimension two, and extend our constructions in higher dimensional spaces. Given $d + 1$ sets of hyperplanes in $\mathbb{R}^d$, we say that they are in *weak general position* if every time we pick d hyperplanes of different sets, their normal vectors are linearly independent. This guarantees that their intersection is a single point.

Our constructions are made for sets in weak general position, and a small perturbation argument gives Theorem 1.4. We consider each set X_i as a color class. Given $d + 1$ sets $X_1, \dots, X_{d+1}$ of r hyperplanes each in $\mathbb{R}^d$, we say that a colorful partition is a partition of their union into r sets $A_1, \dots, A_r$, each with one element of each X_i . For a colorful $(d + 1)$ -tuple A in weak




Figure 1: An illustration of the example for $d = 2$

general position we denote by $\Delta(A)$ the simplex generated by A . If the hyperplanes of A are concurrent, then $\Delta(A)$ is their point of intersection.

THEOREM 2.1. *Let $r \geq 2$ be an integer. Consider the set X_1 (red) of $r - 1$ copies of the line $x = 1$ and one copy of the line $x = -1$, the set X_2 (green) of $r - 1$ copies of the line $y = -1$ and one copy of $y = 1$, and the set X_3 (blue) of $r - 1$ copies of the line $x = -y$ and one copy of the line $x = y$. Then, every colorful partition $A_1, \dots, A_r$ of X_1, X_2, X_3 has two sets $A_j, A_{j'}$ so that $\Delta(A_j) \cap \Delta(A_{j'}) = \emptyset$.*

The construction can be seen in Figure 1.

PROOF. To make a colorful partition of the three colors we first make a colorful partition of red and green and then assign blue lines to each part. Choosing a pair of red and green lines is equivalent to choosing a vertex of the square $[-1, 1]^2$. The number of times a vertex can be chosen is bounded by the number of green lines containing it and by the number of red lines containing it.

If the point $(-1, -1)$ is chosen, then all the other $r - 1$ vertices chosen have to be $(1, 1)$. If the line associated with the point $(-1, -1)$ is $x = y$, then all the other points are associated with the line $x = -y$. So the first simplex is just the point $(-1, -1)$ that is not contained in the simplex defined by $(x = 1, y = 1, x = -y)$. If the blue line associated with $(-1, -1)$ is $x = -y$, then one of the copies of $(1, 1)$ is associated with $x = y$. That last simplex is simply the point $(1, 1)$, which is not contained in the triangle spanned by $(x = -1, y = -1, x = -y)$.

If, on the other hand, the point $(-1, 1)$ is chosen, then the point $(-1, -1)$ cannot be (they share the same red line). Similarly, the point $(1, 1)$ can be chosen at most $r - 2$ times (they share the same green line). Therefore the points in the partition have to be $(-1, 1)$, $(1, -1)$, and $r - 2$ copies of $(1, 1)$. One of $(-1, 1)$ or $(1, -1)$ will have to be associated with the line $x = -y$, since there is only one copy of the line $x = y$. This means that one of the

simplices will be a point, say $(-1, 1)$. But then one of the other simplices is either the point $(1, -1)$ or defined by this point and the line $x = y$, and in either case these two simplices are disjoint. The case analysis is analogous if $(1, -1)$ is chosen. $\square$

Now we extend this construction to $\mathbb{R}^d$. For point $x \in \mathbb{R}^d$ we denote its coordinates by $x = (x_1, \dots, x_d)$.

THEOREM 2.2. *Let $d \geq 3$, $r \geq 2$ be integers, and $\varepsilon > 0$ be a real number smaller than $\frac{1}{2\sqrt{2}(d-2)}$. Consider the sets $X_1, \dots, X_{d+1}$ of hyperplanes in $\mathbb{R}^d$ defined as follows. For $i = 1, \dots, d$, the set X_i is made by $r - 1$ copies of the hyperplane $x_i = 1$ and one copy of the hyperplane $x_i = -1$. The set X_{d+1} is made by $r - 1$ copies of the hyperplane $x_1 + x_2 + \varepsilon(x_3 + \dots + x_d) = 0$ and one copy of the hyperplane $x_1 - x_2 + \varepsilon(x_3 + \dots + x_d) = 0$. Then, every colorful partition $A_1, \dots, A_r$ of $X_1, \dots, X_{d+1}$ has two sets $A_j, A_{j'}$ so that $\Delta(A_j) \cap \Delta(A_{j'}) = \emptyset$.*

PROOF. Let H be the two-dimensional plane spanned by e_1, e_2 . Let $\Pi : \mathbb{R}^d \rightarrow H$ be the orthogonal projection. Notice that every colorful partition of $X_1, X_2, \dots, X_{d+1}$ can be made by first making a colorful partition of X_1, X_2, X_{d+1} and then assign each of the hyperplanes in X_i to different sets in the partition. The colorful partition of X_1, X_2, X_{d+1} intersected with H is essentially the same construction as in Theorem 2.1. We will show that there are two sets $A_j, A_{j'}$ such that $\Pi(\Delta(A_j)) \cap \Pi(\Delta(A_{j'})) = \emptyset$, which would prove the theorem.

Let us compute the first two coordinates of every vertex of A_j . We denote by $\lambda_1, \lambda_2, \lambda_3$ the elements of $\{-1, 1\}$ so that the following holds. The colorful $(d + 1)$ -tuple A_j has the hyperplane $x_1 = \lambda_1$ from X_1 , the hyperplane $x_2 = \lambda_2$ from X_2 , and the hyperplane $x_1 - \lambda_3 x_2 + \varepsilon(x_3 + \dots + x_d)$ from X_{d+1} . The $(d + 1)$ -tuple A_j also has some hyperplane from each of $X_3, \dots, X_d$.

To compute the first two coordinates of a vertex of $\Delta(A_j)$, we have to remove one of the $d + 1$ hyperplanes of A_j and find the intersection of the other d . If neither the hyperplanes from X_1 nor X_2 are removed, the first two coordinates from the intersection are (λ_1, λ_2) . If the hyperplane $x_1 = \lambda_1$ is removed, then the second coordinate is λ_2 . We also have $x_1 - \lambda_3 x_2 + \varepsilon(x_3 + \dots + x_d) = 0$. Therefore,

$$|x_1 - \lambda_3 \lambda_2| = \varepsilon |x_3 + \dots + x_d| \leq \varepsilon(d - 2).$$

In the last case, if the hyperplane $x_2 = \lambda_2$ is removed, the first coordinate is λ_1 . We also have $x_1 - \lambda_3 x_2 + \varepsilon(x_3 + \dots + x_d) = 0$, so $\lambda_3 x_1 - x_2 + \lambda_3 \varepsilon(x_3 + \dots + x_d) = 0$, as $(\lambda_3)^2 = 1$. Therefore,

$$|x_2 - \lambda_3 \lambda_1| = \varepsilon |\lambda_3| \cdot |x_3 + \dots + x_d| \leq \varepsilon(d - 2).$$

Therefore, the vertices of $\Pi(\Delta(A_j))$ are very close to the vertices of the triangle in the corresponding partition of X_1, X_2, X_{d+1} in Theorem 2.1. Since in Theorem 2.1 we could always find two (possibly degenerate) disjoint triangles at distance greater than or equal to $\frac{1}{\sqrt{2}}$, then as long as $\varepsilon < \frac{1}{2\sqrt{2}(d-2)}$ we will have the same conclusion. $\square$

Finally, we make the approximation argument explicit to prove Theorem 1.4.

PROOF OF THEOREM 1.4. We take the construction from Theorem 2.2 and denote it Λ . For each colorful partition P of Λ , there exist two colorful triples A_1, A_2 so that $\Delta(A_1)$ and $\Delta(A_2)$ are strictly separated by a hyperplane H . Let

$$\delta(P) = \min\{\text{dist}(\Delta(A_1), H), \text{dist}(\Delta(A_2), H)\}, \quad \delta = \min_P \delta(P),$$

where the minimum defining δ ranges over all colorful partitions of Λ . We apply a small enough perturbation to Λ so that for every colorful d -tuple X of hyperplanes, the distance $\bigcap X$ moves is less than δ . This guarantees that the hyperplanes that separate colorful simplices still work. $\square$

3. Additional remarks

Karasev's proof of Theorem 1.2 relies on a clever geometric idea. Given a set of $r(d+1)$ hyperplanes in $\mathbb{R}^d$ in general position, and an additional point p , we can consider S the set of projections of p onto each hyperplane, and include p . This gives us a set of $r(d+1) + 1$ points in $\mathbb{R}^d$, so Tverberg's theorem can be applied. We obtain a partition of the point set into $r+1$ sets whose convex hulls intersect. Karasev showed that if in this partition one of the parts is simply $\{p\}$ and the rest are $(d+1)$ -tuples, then the partition they induce on the hyperplanes is the one we seek. Moreover, for every $(d+1)$ -tuple A_j of hyperplanes we would have $p \in \Delta(A_j)$. Then, the proof boils down to showing that for some point p a partition as mentioned works.

At first glance, it might seem tempting to apply a similar idea by replacing Tverberg's theorem by the optimal colorful Tverberg theorem by Blagojević, Matschke, and Ziegler [9,10], as the number of points matches Tverberg's theorem. One interpretation of the counterexamples found in this note is that controlling the size of the sets in an optimal colorful partition is significantly harder than for Tverberg's theorem, even with a similar degree of freedom in the configuration of points. This is a bit surprising since the projection of p onto a fixed hyperplane is an affine map, so the reason why Karasev's conjecture fails is not topological in essence.

References

- [1] A. Akopyan, S. Avvakumov and R. Karasev, Convex fair partitions into an arbitrary number of pieces, arXiv, fmath.MG (2018), 12 pp.
- [2] S. Avvakumov and R. Karasev, Envy-free division using mapping degree, *Mathematika*, **67** (2021), 36–53.
- [3] S. Avvakumov, R. Karasev and A. Skopenkov, Stronger counterexamples to the topological Tverberg conjecture, arXiv:1908.08731, (2019).
- [4] I. Bárány and D. G. Larman, A colored version of Tverberg’s theorem, *J. London Math. Soc.*, **45** (1992), 314–320.
- [5] I. Bárány and J. Pach, Homogeneous selections from hyperplanes, *J. Combin. Theory Ser. B*, **104** (2014), 81–87.
- [6] I. Bárány, S. B. Shlosman and A. Szücs, On a topological generalization of a theorem of Tverberg, *J. London Math. Soc.*, **2** (1981), 158–164.
- [7] I. Bárány and P. Soberón, Tverberg’s theorem is 50 years old: A survey, *Bull. Amer. Math. Soc.*, **55** (2018), 459–492.
- [8] P. V. M. Blagojević, F. Frick and G. M. Ziegler, Tverberg plus constraints, *Bull. London Math. Soc.*, **46** (2014), 953–967.
- [9] P. V. M. Blagojević, B. Matschke and G. M. Ziegler, Optimal bounds for a colorful Tverberg-Vrećica type problem, *Adv. in Math.*, **226** (2011), 5198–5215.
- [10] P. V. M. Blagojević, B. Matschke and G. M. Ziegler, Optimal bounds for the colored Tverberg problem, *J. European Math. Soc.*, **17** (2015), 739–754.
- [11] P. V. M. Blagojević and G. M. Ziegler, Beyond the Borsuk–Ulam theorem: the topological Tverberg story, in: *A Journey Through Discrete Mathematics*, Springer (Cham, 2017). 273–341.
- [12] P. V. M. Blagojević and G. M. Ziegler, Convex equipartitions via equivariant obstruction theory, *Israel J. Math.*, **200** (2014), 49–77.
- [13] F. Frick, Counterexamples to the topological Tverberg conjecture, *Oberwolfach Reports*, **12** (2015), 318–312.
- [14] F. Frick and P. Soberón, The topological Tverberg problem beyond prime powers, arXiv, math.CO (2020).
- [15] C. Huemer, P. Pérez-Lantero, C. Seara and R. I. Silveira, Matching points with disks with a common intersection, *Discrete Math.*, **342** (2019), 1885–1893.
- [16] R. N. Karasev, Dual theorems on central points and their generalizations, *Sbornik: Math.*, **199** (2008), 1459–1479.
- [17] R. N. Karasev, Tverberg-Type Theorems for intersecting by rays, *Discrete & Comput. Geom.*, **45** (2011), 340–347.
- [18] R. N. Karasev, A. Hubard and B. Aronov, Convex equipartitions: the spicy chicken theorem, *Geom. Dedicata*, **170** (2014), 263–279.
- [19] S. Lee and K. Yoo, On a conjecture of Karasev, *Comput. Geom.*, **75** (2018), 1–10.
- [20] J. A. D. Loera, X. Goaoc, F. Meunier and N. H. Mustafa, The discrete yet ubiquitous theorems of Carathéodory, Helly, Sperner, Tucker, and Tverberg, *Bull. Amer. Math. Soc.*, **56** (2019), 1–97.
- [21] I. Mabillard and U. Wagner, Eliminating Tverberg points. I. An analogue of the Whitney trick, in: *Proc. 30th Annual Symp. Comput. Geom. (SOCG)*, ACM (Kyoto, 2014), pp. 171–180.
- [22] J. Matoušek, *Using the Borsuk–Ulam Theorem: Lectures on Topological Methods in Combinatorics and Geometry*, Springer (Berlin, Heidelberg, 2003).
- [23] F. Meunier and S. Zerbib, Envy-free cake division without assuming the players prefer nonempty pieces, *Israel J. Math.*, **234** (2019), 907–925.
- [24] M. Özaydin, Equivariant maps for the symmetric group (1987), unpublished preprint, <https://minds.wisconsin.edu/bitstream/handle/1793/63829/Ozaydin.pdf>.

- [25] P. J. Rousseeuw and M. Hubert, Depth in an arrangement of hyperplanes, *Discrete Comput. Geom.*, **22** (1999), 167–176.
- [26] P. Soberón, Balanced convex partitions of measures in R^d , *Mathematika*, **58** (2012), 71–76.
- [27] P. Soberón and Y. Tang, Tverberg’s theorem, disks, and Hamiltonian cycles, arXiv:2011.12218 (2020).
- [28] H. Tverberg, A generalization of Radon’s theorem, *J. London Math. Soc.*, **41** (1966), 123–128.
- [29] A. Y. Volovikov, On a topological generalization of the Tverberg theorem, *Math. Notes*, **59** (1996), 324–326.
- [30] R. T. Živaljević and S. T. Vrećica, The colored Tverberg’s problem and complexes of injective functions, *J. Combin. Theory, Series A*, **61** (1992), 309–318.