Ecological and Evolutionary Origin of *Costus flammulus* (Costaceae): A New Species from the Montane Cloud Forests of the Volcanic Cordilleras in Northern Costa Rica

Pedro Juárez, 1,3 Oscar M. Vargas, 2 and Kathleen M. Kay1

¹Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, California 95060, USA; pjuarez1@ucsc.edu; kmkay@ucsc.edu

²Department of Biological Sciences, California Polytechnic State University, Humboldt, Arcata, California 95521, USA; ov20@humboldt.edu

³Author for correspondence (pjuarez1@ucsc.edu)

Communicating Editor: James I. Cohen

Abstract—Costus flammulus is a new herbaceous species endemic to montane cloud forests of the volcanic cordilleras in northern Costa Rica. Costus flammulus has been mistaken for C. wilsonii, but phylogenetic evidence demonstrates that it is closely related to the widespread lowland species C. pulverulentus. Here, we use an integrated framework of species concepts to evaluate whether C. flammulus and C. pulverulentus are distinct species. First, we re-evaluate prior phylogenetic analyses to assess whether C. flammulus bifurcated from or budded off from within C. pulverulentus and whether C. flammulus is monophyletic. We then compare phenotypic traits to determine which diagnostic vegetative and inflorescence traits can be used to identify species in herbarium specimens and examine whether floral traits may confer floral isolation. We compare pollinator assemblages to examine whether pollinator specificity may contribute to reproductive isolation. Finally, we model species distributions and climatic niche overlap to assess ecogeographic isolation. We found that C. flammulus is a monophyletic species phenotypically, ecologically, and geographically distinct from C. pulverulentus and may have speciated as a peripheral isolate at the high elevation range edge of C. pulverulentus. Several lines of evidence, such as C. pulverulentus paraphyly, range size asymmetry, and C. flammulus' nested distribution and vegetative traits, suggest that C. flammulus budded off from a C. pulverulentus-like progenitor species, evolving to tolerate a colder and more seasonal montane environment.

Keywords—Budding speciation, ecogeographic isolation, floral isolation, integrative taxonomy, Neotropical flora, species concepts.

Costus flammulus K.M.Kay & P.Juárez (Costaceae) is a newly proposed plant species restricted to the montane cloud forests of the volcanic cordilleras in northern Costa Rica (Fig. 1). It is a perennial hummingbird-pollinated herb found in forest understory and rocky ravines. This new taxon (hereafter C. flammulus) was first identified by phylogenetic analyses of the Neotropical Costus L. (Kay et al. 2005) and its phylogenetic placement was confirmed with the recent 756 gene phylogenomic analyses of Vargas et al. (2020). Costus flammulus is closely related to the widespread species C. pulverulentus C.Presl (Fig. 1), and diverged ~714,000 yr ago (95% confidence interval: 180,000–1,420,000 yr; Vargas et al. 2020). Vargas et al. (2020) suggested that C. flammulus may be phylogenetically nested within a paraphyletic C. pulverulentus and geographically nested within its range.

Previously, *C. flammulus* had been included in *C. wilsonii* Maas (e.g. Stratton 1989; Haber 2000; Maas and Maas-van de Kamer 2003), apparently due to its similar phenotype and habitat (see Notes). Both species are hummingbird-pollinated understory herbs restricted to montane forests with similar foliage and inflorescence characteristics. However, phylogenetic evidence by Kay et al. (2005) and Vargas et al. (2020) show that *C. flammulus* and *C. wilsonii* are independent evolutionary lineages that may have converged on a similar phenotype. Their most recent common ancestor lived ~3 MYA in the Pliocene when *Costus* first diverged in the Neotropics (Vargas et al. 2020); thus, *C. flammulus* and *C. wilsonii* are not closely related within Neotropical *Costus*.

To understand the ecology, delimitation, and evolutionary origin of *C. flammulus*, we investigate its phylogenetic position, phenotype, distribution, ecological niche, and reproductive isolation from *C. pulverulentus*. Phylogenetic analysis is indispensable for assessing monophyly, the primary criterion for defining species and delimiting taxa under the Phylogenetic Species Concept (reviewed in Baum and Smith 2013).

Phylogenetic evidence also may be the most compelling for analyzing modes of speciation, such as budding speciation or splitting cladogenesis (Foote 1996; Sianta and Kay 2022). In plants with low dispersal and strong local adaptation among populations, budding speciation is hypothesized to be a particularly common mode of speciation (Grossenbacher et al. 2014). Budding speciation involves the isolation of marginal populations at the periphery of a species range (Crawford 2010) and initially results in a monophyletic 'derivative' species phylogenetically nested within a paraphyletic 'progenitor' species (Baldwin 2005; Sianta and Kay 2022). Phenotypic distinguishability is essential for species classification and identification (Zapata and Jiménez 2012), and phenotypic comparisons can suggest potential forms of reproductive isolation, which are key criteria for species delimitation under the Biological Species Concept (Mayr 1942). For instance, floral divergence between closely related species may indicate divergence in pollination and/or mating systems that may contribute to reproductive isolation (Van der Niet et al. 2014) and can be investigated with field studies. Moreover, comparing geographic distributions and ecological niches of closely related species can suggest the geographic mode of speciation and whether ecogeographic isolation may be important (Raxworthy et al. 2007; Rissler and Apodaca 2007). This is relevant because divergent adaptation to local environmental conditions often may be the primary source of divergence (Mayr 1947; Futuyma 1998; Hendry 2017). An integrative approach to species delimitation is useful for understanding the roles of ecological divergence and reproductive isolation in speciation (De Queiroz 1998, 2007; Dayrat 2005).

Here, we combine a traditional protologue with an analysis of ecological divergence and putative reproductive isolating mechanisms related to ecogeographic and floral isolation to compare *C. flammulus* with its putative progenitor or sister species, *C. pulverulentus* (Fig. 1). First, we reevaluate the phylogenetic evidence from Vargas et al. (2020) to assess whether




Fig. 1. Costus flammulus and C. pulverulentus. A. Map of occurrences and the 80% probability of occurrence for each species derived from species distribution models (SDM) based on bioclimatic data. White outline shows southern Nicaragua, Costa Rica, and Northwestern Panama. B. Pruned phylogram inferred from the concatenated matrix of 756 genes for Costus (Vargas et al. 2020). Numbers to the right of nodes indicate phylogenetic support, left: ultrafast bootstrap, right: Shimodaira-Hasegawa approximate ratio; nodes without values are fully supported at 100. C–D. Photographs of C. flammulus (C) and C. pulverulentus (D). Photographs by P. Juárez (C); and K. M. Kay (D).

C. flammulus bifurcated or budded off from C. pulverulentus and whether it is monophyletic. We then compare phenotypic traits to determine which diagnostic vegetative and inflorescence traits can be used to identify species in herbarium specimens and examine floral traits that may contribute to reproductive isolation. We compare pollinator assemblages to examine to what extent pollinator specificity may contribute to reproductive isolation. Finally, we model species distributions and climatic niche overlap to assess ecogeographic isolation.

MATERIALS AND METHODS

Species Concepts and Species Delimitation—We use an integrated framework of species concepts to assess whether C. flammulus and C. pulverulentus are two phylogenetically, morphologically, and ecologically distinct species. First, we use the Phylogenetic Species Concept framework to delimit C. flammulus (Donoghue 1985; Mishler 1985). Second, we use the traditional Phenetic Species Concept framework to study phenotypic trait divergence and identify diagnostic phenotypic traits and classify species based on overall similarity (Michener 1970; Sokal and Crovello 1970; Sneath and Sokal 1973). Third, we use the Ecological Species Concept framework to determine whether C. flammulus and C. pulverulentus are two ecologically distinct species that occupy different climate and pollinator niches (Van Valen 1976; Andersson 1990). Finally, we use the Biological Species Concept framework to study putative reproductive isolating barriers of C. flammulus, such as ecogeographic and floral isolation (Mayr 1942; Dobzhansky 1970; Coyne and Orr 2004).

Study System—Neotropical Costus comprises approximately 60 named species found in habitats from sea level to montane forests throughout tropical Central and South America (Maas 1972, 1977; Maas and Maas-van de Kamer 2003, 2014; Kay et al. 2005; Vargas et al. 2020). Costus are perennial herbs found in forest edges, forest gaps, forest understory, ravines, and riparian habitats. Neotropical Costus is a recent and rapid species radiation since dispersing from Africa approximately 3 MYA. Neotropical Costus species are primarily pollinated by euglossine bees (tribe Euglossini) or hermit hummingbirds (subfamily Phaethornithinae) (Kay and Schemske 2003; Kay 2006; Kay and Grossenbacher 2022) and there have been many evolutionary transitions from bee to hummingbird pollination (Kay and Grossenbacher 2022). Prior studies on Neotropical Costus suggest that ecogeographic and floral isolation play an important role in segregating species (Kay and Schemske 2003; Kay 2006; Chen 2013; Chen and Schemske 2015). Ecogeographic isolation is geographic isolation resulting from genetically based ecological differences between species (Sobel et al. 2010; Sobel 2014). Floral isolation is reproductive isolation mediated by floral morphology and pollinator behavior (Grant 1994; Kay and Sargent 2009; Schiestl and Schlüter 2009).

Specimen Sampling—We studied herbarium specimens from Costa Rica determined as Costus, C. pulverulentus, and C. wilsonii of the following herbaria: CR, HLDG, MO, and USJ. Herbarium acronyms cited in this paper follow Thiers (2022). All specimens we identified as C. flammulus were used to extract occurrence data. We selected one representative C. pulverulentus specimen from every district where the species occurs in Costa Rica to measure vegetative and inflorescence traits and compare elevation (Supplemental Table S1; Juárez et al. 2023). For the elevation comparison, we excluded those herbarium specimens for which elevation was reported as a range. We also visited the type locality of C. flammulus (Estación Biologica Monteverde, Quebrada Máquina, 1,524 m, 10°19′03′′N, 84°48′27′′W) in August 2021 to record pollinator visitation and to collect

and measure flowers for floral trait comparisons. *Costus pulverulentus* flowers were collected and measured from several Costa Rica and Panama locations (Table S2, Juárez et al. 2023).

Phylogenetic Analyses—The concatenated analysis of 756 nuclear genes of Vargas et al. (2020) showed that two samples of C. flammulus (C. sp. nov. 18020, C. sp. nov. 18049) appear monophyletic (support: 100% bootstrap, 161 gene topologies) and nested within C. pulverulentus (support: 91% bootstrap, 1 gene topology) (Fig. 1). Precisely, C. flammulus appears to be sister to a clade comprising C. pulverulentus samples from Costa Rica, Panama, and South America (hereafter widespread); with samples of C. pulverulentus from Mexico (hereafter Mexican) sister to a clade containing C. flammulus and widespread C. pulverulentus (support: 78% bootstrap, 1 gene topology). This topological pattern suggests that C. flammulus may be a derivative from a C. pulverulentus-like progenitor species (Crawford 2010; Sianta and Kay 2022). Here, we further investigate gene tree support for different phylogenetic placements of C. flammulus and C. pulverulentus by calculating the support per gene of alternative topologies in the Vargas et al. (2020) results. We asked how many gene trees recover these alternative topologies: a) C. flammulus and C. pulverulentus as comprising a single clade (regardless of the monophyly of each species), b) C. pulverulentus as monophyletic, c) C. flammulus as monophyletic, d) C. flammulus and widespread C. pulverulentus as a single clade, e) C. flammulus sister to widespread C. pulverulentus (as shown in the concatenated phylogeny), f) C. flammulus and Mexican C. pulverulentus comprising a single clade, and g) C. flammulus sister to Mexican C. pulverulentus. We calculated the raw number of gene trees that support each alternative topology, and their percentages relative to the number of gene trees that contain any phylogenetic signal for the position of C. flammulus and C. pulverulentus. Trees with strong phylogenetic signal are defined as those in which any sample belonging to the focal species forms a wellsupported clade (≥ 90 bootstrap support) with any other sample, as calculated by Vargas et al. (2020). We used a custom R script to analyze trees and calculate their gene tree support using the packages phytools (Revell 2012) and MonoPhy (Schwery and O'Meara 2016).

Phenotypic Trait Measurements and Comparisons—We measured five vegetative and five inflorescence traits (Table 1; Fig. 2) on herbarium specimens of C. flammulus (n=21) and C. pulverulentus (n=28) to determine which diagnostic traits can be used to separate the two species. Vegetative traits include stem diameter (mm), ligule length (mm), petiole length (mm), and leaf length and width (cm). Inflorescence traits include inflorescence length and width (cm), bract length and width (cm), and nectarial callus length (mm). The nectarial callus is located on top of the bracts (Maas 1972) and commonly produces extrafloral nectar that attracts ants that deter florivores and fruit predators (Schemske 1982), although in some species the callus is vestigial.

To determine floral divergence that may contribute to floral isolation, we measured eight floral traits (Table 1; Fig. 2) on fresh flowers of *C. flammulus* (n = 7) and *C. pulverulentus* (n = 17). Floral traits include corolla

length (cm), corolla tube length (cm), corolla dorsal lobe length (cm), labellum length (cm), petaloid stamen length (cm), petaloid stamen exsertion length (mm), anther length (mm), and style length (cm). We separately compared herbarium specimens and floral trait measurements with multiple two-tailed t tests and adjusting p values within data sets with false discovery rate (FDR) controlling procedures (Benjamini and Hochberg 1995) using the R package Stats (R Core Team 2021).

We used two separate principal component analyses (PCA) to visualize phenotypic distributions for each dataset using the R package Stats; one for vegetative and inflorescence traits measured on herbarium specimens and the other for floral traits measured on fresh flowers. The PCAs allowed us to reduce data complexity and graphically represent differences between the two species (Ringnér 2008). To plot PCAs, we used the R packages ggfortify (Tang et al. 2016) and ggplot2 (Wickham 2009).

Pollinator Assemblage Sampling—We recorded pollinator visitation to six *C. flammulus* individuals at the type locality for seven days in August 2021 to determine pollinator identities and visitation rates. We recorded visitation with video cameras from 6:00 AM to 2:00 PM, when pollinators start activity to about when daily pollinator activity decreases, and flowers wilt. We used Canon PowerShot SX420 IS and Canon PowerShot SX540 HS digital cameras with auto-controlled aperture settings to record visitation. We used the Canon Hack Development Kit (CHDK, https://chdk.fandom.com) to automate motion detection features. We recorded 15-sec videos with the motion detection script MDFB2013 setting the trigger threshold to 5 and burst to 15-sec. Pollinator visitation to *C. pulverulentus* was reported previously (Kay and Schemske 2003; Kay 2006).

Species Distribution Models—To compare suitable habitats, niches, and geographic ranges, we built maximum entropy species distribution models (SDMs) for C. flammulus and C. pulverulentus with climate data following the methods in Goff et al. (2021). We obtained occurrence data of both species from herbarium databases (CR, https://biodiversidad. museocostarica.go.cr; MO, https://tropicos.org) and field observations by K. M. Kay and P. Juárez. We excluded occurrences with duplicated collection localities and uncertain taxonomic identifications. To avoid potential misidentifications, we considered only occurrences with identifications made by one of four taxonomic experts: P. Maas, D. Skinner, K. M. Kay, and P. Juárez. We used 21 specimens for C. flammulus from Costa Rica identified by P. Juárez and 593 for C. pulverulentus (Table S3, Juárez et al. 2023). The specimen number disparity was expected since C. flammulus is only known from a few locations on volcanic cordilleras in northern Costa Rica (Fig. 1; see Distribution and Habitat), whereas C. pulverulentus is a widespread species ranging from Mexico to southern Ecuador (Maas 1972).

We used occurrence data for 29 Neotropical flowering plant genera (Table S4, Juárez et al. 2023) as pseudo-absence data in the SDMs to represent the range of environmental conditions available to *C. flammulus* and *C. pulverulentus* and to correct for geographic variation in plant collecting

Table 1. Phenotypic and elevational comparisons between *C. flammulus* and *C. pulverulentus*. Abbreviations is for abbreviations in Fig. 2. ns $p \ge 0.05$, $p \le 0.05$, $p \le 0.05$, $p \le 0.00$, $p \ge 0.00$, $p \ge 0.00$.

		Mean ± 95% CI (n)		
Trait	Abbreviations	C. flammulus	C. pulverulentus	p
Stem diameter (mm)	Stm	$6.79 \pm 0.90 (21)$	8.96 ± 1.13 (28)	*
Ligule length (mm)	Lg	5.90 ± 0.73 (21)	$7.32 \pm 1.67 (28)$	ns
Petiole length (mm)	P	$5.09 \pm 0.77 (21)$	$5.17 \pm 0.61 (28)$	ns
Leaf blade length (cm)	Lvsl	$14.67 \pm 1.08 (21)$	$18.23 \pm 1.78 (28)$	*
Leaf blade width (cm)	Lvsw	4.30 ± 0.45 (21)	$5.34 \pm 0.47 (28)$	*
Inflorescence length (cm)	Infl	$6.44 \pm 0.80 (21)$	9.78 ± 1.16 (28)	***
Inflorescence width (cm)	Infw	3.01 ± 0.22 (21)	3.69 ± 0.82 (28)	ns
Bract length (cm)	Bl	1.76 ± 0.13 (21)	2.56 ± 0.21 (28)	***
Bract width (cm)	Bw	1.56 ± 0.13 (21)	1.87 ± 0.14 (28)	*
Nectarial callus length (mm)	Cal	$6.14 \pm 0.80 (21)$	$8.82 \pm 1.12 (28)$	**
Corolla length (cm)	С	3.44 ± 0.27 (7)	$6.19 \pm 0.27 (17)$	***
Corolla tube length (cm)	Ct	1.11 ± 0.21 (7)	$2.06 \pm 0.26 (17)$	***
Corolla dorsal lobe length (cm)	Cdl	$3.47 \pm 0.50 (7)$	$4.20 \pm 0.24 (17)$	*
Labellum length (cm)	L	3.22 ± 0.35 (7)	$4.76 \pm 0.23 (17)$	***
Petaloid stamen length (cm)	S	3.52 ± 0.21 (7)	$6.37 \pm 0.26 (17)$	***
Petaloid stamen exsertion (mm)	Se	$5 \pm 0 (7)$	$16.27 \pm 2.30 (17)$	***
Anther length (mm)	A	$7 \pm 0 \ (7)$	8.32 ± 0.73 (17)	***
Style length (cm)	St	2.91 ± 0.21 (7)	$5.87 \pm 0.21 (17)$	***
Elevation (m)	-	$1310.7 \pm 688.2 (14)$	$443.7 \pm 1045.6 (25)$	***




Fig. 2. Phenotypic comparisons between Costus flammulus and C. pulverulentus. A. Scatterplot from principal component analysis of the vegetative and inflorescence traits measured in herbarium specimens; PC1 and PC2 explain 41.04% and 16.52% of the total phenotypic variance, respectively. Costus flammulus vegetative and inflorescence traits appear as a nested subset of C. pulverulentus. B. Diagram of the vegetative and inflorescence traits measured on herbarium specimens. Image based on Haber, Ivey & Zuchowski 10798 (CR, MO). C. Scatterplot from principal component analysis of the floral traits measured on fresh flowers; PC1 and PC2 explain 75.82% and 13.14% of the total phenotypic variance, respectively. Costus flammulus and C. pulverulentus mainly differentiate along PC1, with high values corresponding to all floral traits measured. D. Diagram of the floral traits measured on fresh flowers. Photographs by M. Morello (B); and C. Girvin (D).

TABLE 2. Number of gene trees that support specific relationships in *C. flammulus* and *C. pulverulentus*. Percentage is calculated over the 443 gene trees that contain phylogenetic signal for the focal species.

Relationship	Number of trees	Percentage
C. flammulus and C. pulverulentus are a clade	6	1.4
C. pulverulentus is monophyletic	20	4.6
C. flammulus is monophyletic	243	54.9
C. flammulus and Widespread C. pulverulentus are a clade	8	1.8
C. flammulus is sister to Widespread C. pulverulentus	2	0.5
C. flammulus and Mexican C. pulverulentus are a clade	3	0.7
C. flammulus is sister to Mexican C. pulverulentus	3	0.7

effort and accessibility. Using pseudo-absence data that is similarly spatially biased to the actual occurrence data of *Costus* generates less-biased models than when using randomly selected pseudo-absence data (Phillips et al. 2009). We bounded the pseudo-absence points for the coordinates of the terrestrial political borders of Costa Rica (08° to 11° latitude and -86° to -82° longitude) and subsampled them so that there were five pseudo-absence points per 10-min resolution grid cell, totaling 20,424 points (Fig. S1, Juárez et al. 2023). We limited the analysis to the boundaries of Costa Rica, even though the actual *C. pulverulentus* distribution range extends beyond these because we wanted to compare SDMs and ecological niches from co-occurring populations in the same geographical area.

We used 19 BioClim variables at 0.5-min resolution from WorldClim (https://www.worldclim.org) as explanatory variables for the SDMs. To build SDMs, we combined the climate data for *C. flammulus* and *C. pulverulentus* and pseudo-absence data using the default MaxEnt settings with the R package dismo (Hijmans et al. 2017). We used 80% of the occurrence data for each *Costus* species to train the model and 20% to test its performance. We assessed models with the area under the receiver operating characteristic (ROC) curves (AUC) and qualitatively compared the consistency of the percentage contribution of variables to the models.

Climatic Niche Comparisons—We made a multivariate comparison of the climatic niche by circumscribing each Costus species' occurrencebased niche relative to the available niche space established by pseudoabsence data using the R package ecospat (Broennimann et al. 2012). We extracted values for the climatic variables from each spatial cell at 0.5-min resolution included in the geographic extent of the pseudoabsence occurrence points used for the SDMs. We used PCA of the 19 Bio-Clim variables to visualize axes of niche overlap, projecting C. flammulus and C. pulverulentus into an available niche space and tested for niche differentiation. We generated 100 × 100 PCA unit grid cells and used the Costus species' occurrence data to project the density of each species into an environmental space. This projection applies a kernel density function to determine the density of occurrences in each cell in the environmental space for each species and lessen sampling strategy bias effects when calculating niche differences (Broennimann et al. 2012). We calculated niche overlap as Schoener's ${\it D}$ (Schoener 1970) between ${\it C. flammulus}$ and ${\it C. pul-}$ verulentus and tested for niche equivalency and similarity (Warren et al. 2008) using ecospat. Niche similarity indicates whether two species occupy a similar subset of the total available environment, whereas niche equivalency indicates whether the niches of the two species are statistically distinguishable (Warren et al. 2008; Broennimann et al. 2012).

RESULTS

Phylogenetics—Only 443 genes, out of the 756 genes, have some phylogenetic signal for the position of *C. flammulus*. The monophyly of *C. flammulus* is supported by 243 genes (54.9%) (Table 2). Our support per gene analysis regarding the position of *C. flammulus* relative to *C. pulverulentus* is inconclusive. Twenty genes support the monophyly of all *C. pulverulentus* samples. Only two genes recover *C. flammulus* as sister to widespread *C. pulverulentus*, while three genes show *C. flammulus* as sister to Mexican *C. pulverulentus*. Six genes support the position of *C. flammulus* nested in *C. pulverulentus* and eight genes support *C. flammulus* forming a clade with widespread *C. pulverulentus* (regardless of species-level monophyly). Despite the uncertainty shown by the gene support

analysis, our results support that *C. flammulus* samples form a monophyletic clade, possibly nested within *C. pulverulentus*. The concatenated analysis in Vargas et al. (2020), in which *C. flammulus* is nested within *C. pulverulentus* and reciprocally monophyletic with widespread *C. pulverulentus*, may be our best hypothesis, but we only find strong support for the monophyly of *C. flammulus*.

Phenotypic Trait Comparisons—The phenotypic trait comparisons reveal consistent floral trait differences between C. flammulus and C. pulverulentus, but fewer differences in vegetative and inflorescence traits (Table 1; Fig. 2). Costus flammulus has significantly narrower stems and smaller leaves than C. pulverulentus. Inflorescence length, bract size, and nectarial callus are significantly different, with C. flammulus having shorter inflorescences and smaller bracts than *C. pulverulentus*. The nectarial callus is also shorter in *C. flammulus*, and according to field observations is not functional and does not attract ants, unlike the active extrafloral nectaries of *C. pulverulentus* (Schemske 1982). All floral traits differ significantly; C. flammulus has smaller flowers than C. pulverulentus. The PCAs demonstrate that C. flammulus vegetative and inflorescence traits are a nested subset of C. pulverulentus; conversely, floral traits form clearly separate phenotypic clusters (Fig. 2).

Pollinator Assemblage Overlap—We did not detect an overlap in pollinator assemblage between *C. flammulus* and *C. pulverulentus*. Costus flammulus was almost exclusively visited by the green hermit (*Phaethornis guy* Lesson) (0.250 visits/hour; 14 legitimate visits) during observations in the type locality; with an additional visit by the black-crested coquette (*Lophornis helenae* Delattre) (0.018 visits/hour; 1 legitimate visit), and several nectar robbing approaches by the white-throated mountaingem (*Lampornis castaneoventris* Gould). Kay and Schemske (2003) and Kay (2006) previously recorded pollinator assemblages for several *Costus* species and determined that *C. pulverulentus* is solely visited by the long-billed hermit (*P. longirostris* Delattre) at all their study sites in Costa Rica and Panama.

Species Distribution Models—The SDMs for *C. flammulus* predicted almost no probability of environmental suitability for *C. pulverulentus* within the range of *C. flammulus* and vice versa (Fig. 1), with the predicted areas of the probability of occurrence closely overlapping the actual species occurrences. The SDMs predict that *C. flammulus* is restricted to the volcanic cordilleras in northern Costa Rica, whereas *C. pulverulentus* occurs primarily in the wet lowlands of Costa Rica, especially in the northern Caribbean plains and southern Pacific region. The SDMs built using 19 BioClim as explanatory variables performed well, as indicated by the area under the receiver operating characteristic curve (AUC of ROC) scores, which are typical measures of SDM performance.

The AUC of ROC was 0.945 for *C. flammulus* and 0.86 for *C. pulverulentus*.

Climatic Niche Overlap—Costus flammulus and C. pulverulentus occupy significantly different climatic niches (Fig. 3). Within the available climatic niche determined by the pseudoabsence occurrence data, C. flammulus and C. pulverulentus differentiate along a combination of PC1 and PC2, with C. flammulus occurring in colder climates within the drier and more seasonal part of C. pulverulentus' precipitation range (Fig. 3). Costus pulverulentus occupies a broader climatic niche space, as expected given its widespread distribution, and there is little climate niche overlap between the species. The D metric of niche overlap for C. flammulus and C. pulverulentus was 0.0038, which is significantly lower than when aggregated occurrences were randomly assigned to species (ecospat test of niche equivalency; p = 0.0099), indicating that the two species occupy different climatic niches (Fig. 3; Warren et al. 2008). The ecospat test for niche similarity shows that the species' niches are more similar than expected when sampling the total available environment at random (p = 0.4545), as expected for closely related taxa (Warren et al. 2008).

DISCUSSION

We describe *C. flammulus*, a new herbaceous species endemic to the montane cloud forests of the volcanic cordilleras in northern Costa Rica. *Costus flammulus* is a monophyletic species that is phenotypically, ecologically, reproductively, and geographically distinct from its closest relative, *C. pulverulentus*. It may have speciated as a peripheral isolate at the high elevation range edge of the widespread, lowland *C. pulverulentus*. *Costus flammulus* can be distinguished from *C. pulverulentus* by its smaller leaves, inflorescences, and flowers and distribution at higher elevations in montane cloud forests.

Our integrative approach allows us to circumscribe C. flammulus under multiple species concepts. (a) Under the Phenetic Species Concept framework (Michener 1970; Sokal and Crovello 1970; Sneath and Sokal 1973), C. flammulus is a phenotypically distinct species from C. pulverulentus, forming a divergent phenotypic cluster in a floral phenotypic space (Fig. 2). Additionally, our phenotypic comparisons show that C. flammulus and C. pulverulentus significantly differ in leaf, inflorescence, and flower size (Table 1). (b) Under the Ecological Species Concept framework (Van Valen 1976; Andersson 1990), C. flammulus is an ecologically distinct species from C. pulverulentus because it occupies fundamentally different pollinator and climate niches. Our pollinator observations show that C. flammulus and C. pulverulentus are consistently visited by different species of hermit hummingbirds with no pollinator assemblage overlap, although that might be because their specific hummingbird pollinators, the green hermit, and the long-billed hermit, also segregate geographically. Our SDMs and ecological niche comparisons show that C. flammulus and C. pulverulentus niches have diverged and that there is little climatic niche overlap. (c) Under the Biological Species Concept framework (Mayr 1942; Dobzhansky 1970; Coyne and Orr 2004), C. flammulus is a reproductively isolated species from C. pulverulentus because their effective ecogeographic isolation is nearly complete, such that their populations are unlikely to experience gene flow, and their floral trait divergence could mediate floral isolation if they were to come into geographic contact. (d) Under the Phylogenetic Species Concept framework (Donoghue 1985; Mishler 1985), our conclusions are less certain. Our re-evaluation of Vargas et al.'s (2020) phylogenetic analysis shows strong evidence that *C. flammulus* is a monophyletic species, and possibly nested within *C. pulverulentus*, although its exact position is uncertain, and we do not have strong support for reciprocal monophyly of *C. flammulus* and any subset of *C. pulverulentus* samples.

Costus flammulus' phylogenetic position may make C. pulverulentus paraphyletic. This topological pattern is congruent with budding speciation theory (Crawford 2010), in which C. pulverulentus and C. flammulus may be a progenitorderivative species pair, i.e. C. flammulus may be a derivative from a C. pulverulentus-like progenitor species (see below). If true, according to the Phylogenetic Species Concept, C. pulverulentus may be considered two separate species, with a possible division between widespread and Mexican samples. Under this scenario, one of these two C. pulverulentus phylogenetic entities would need a new name, which would require a detailed examination of the type specimen of C. pulverulentus to determine whether it belongs to the Mexican or widespread C. pulverulentus. Although Mexico is the cited type locality in the original protologue (Presl 1827), there is uncertainty about the location. The C. pulverulentus type specimen was collected in 1790, possibly in Nicaragua, based on the documented itinerary of the collector (D. Skinner pers. comm.). Further investigation of *C. pulverulentus* taxonomy should involve phylogenetic analysis of samples from the putative type locality and could benefit from more extensive genetic data to better resolve relationships. Given our current data, splitting up *C. pulverulentus* is premature.

Phenotypic comparisons detected significant floral divergence with a reduction in floral size in *C. flammulus*, whereas no such pattern was found for vegetative and inflorescence traits. This suggests little diversification of vegetative and inflorescence traits but an evolutionary force acting on flowers. The maintenance of vegetative trait similarity accompanied by floral shifts may be expected between recently diverged species, as observed in other groups (Grant 1949; Van der Niet and Johnson 2009; Grossenbacher and Whittall 2011; Pessoa et al. 2020). This pattern may reflect recent shared ancestry, pollination or mating system divergence, and the establishment of prezygotic reproductive isolation mechanisms.

The quantified floral divergence suggests that floral isolation could be important in segregating C. flammulus and C. pulverulentus if they were to establish in sympatry. Floral divergence can mediate isolation between closely related species (Grant 1949, 1993, 1994; Fulton and Hodges 1999; Ramsey et al. 2003; Kay 2006; Schiestl and Schlüter 2009). Although we do not have direct measures of floral isolation, we can speculate based on our knowledge of isolating barriers between other species pairs. For instance, the reduction of stamen exsertion beyond the labellum could be particularly relevant for floral isolation. Our pollinator observations show that C. flammulus may place its pollen on the distal third of the hummingbirds' beak, whereas C. pulverulentus places its pollen on the forehead (Kay 2006; Kay and Surget-Groba 2022). Studies estimating reproductive barriers between sympatric C. pulverulentus and C. scaber Ruiz & Pav. have shown that divergence in stamen exsertion mediates mechanical isolation by placing pollen in different sites of the hummingbird; a pollen placement shift from the hummingbird beak to forehead produces nearly complete isolation (Kay 2006). Differences




Fig. 3. Climate niches of C. flammulus and C. pulverulentus. A, B. 100×100 gridded environmental spaces representing the climate environment available to Costus flammulus and C. pulverulentus along these PC axes. Gray shading indicates the density of occurrences within a cell and dotted and solid lines encompass 50% and 100% of the available environment, respectively. Overall, C. flammulus inhabits a colder and seasonal habitat, whereas C. pulverulentus occupies warmer and very humid environments. The niches of Costus flammulus and C. pulverulentus are significantly different from each other. C–D. Histograms showing the observed niche overlap D between the species' ranges (red bars with a diamond) and simulated niche overlaps (grey bars) on which test of niche equivalency (C) and niche similarity (D) are calculated from D0 iterations. C1. Biplot from the principal components analysis of the bioclimatic environmental space defined by pseudo-absence occurrence data used in the species distribution models. Correlated bioclimatic variables are circled and jointly labeled.

in pollen placement sites are important in isolating species in other Neotropical groups as well, such as *Burmeistera* (Muchhala and Thomson 2012), *Heliconia* (Stiles 1975, 1979; Kress 1983), and Orchidaceae (reviewed in Schiestl and Schlüter 2009). Thus, divergent floral traits like stamen exsertion could play a major role in isolating these species by differential pollen placement. However, floral isolation studies are needed to confirm this hypothesis.

Our pollinator assemblage observations show that C. flammulus and C. pulverulentus occupy non-overlapping pollinator niches. Costus flammulus is visited by montane hummingbirds, primarily by the green hermit, whereas C. pulverulentus is exclusively visited by the long-billed hermit (Kay and Schemske 2003; Kay 2006). It is well known that pollinator specificity contributes to adaptive divergence and reproductive isolation (Van der Niet et al. 2014). As such, pollinators select on floral traits, and plant populations adapt to pollinators' preferences, behavior, and morphology to maximize visitation and efficient pollen transfer (Grant and Grant 1965; Stebbins 1970; Fenster et al. 2004; Johnson 2006; Schiestl and Johnson 2013). In this way, geographic variation in pollinator-mediated selection can drive floral divergence between plant populations. However, C. flammulus speciation may not be related to pollinator specialization, but rather the differences in pollinator identity may be a byproduct of elevational divergence. The green hermit, common at medium to high elevations, and the long-billed hermit, abundant in lowlands and rare above medium elevations (Garrigues and Dean 2014), are ecological equivalents, both being medium-sized forest understory hermit hummingbirds with similar behavior (Stiles 1983; Skutch and Stiles 1989). Thus, it seems likely that C. flammulus' pollination niche shift is confounded with climatic adaptation rather than local pollinator availability. It has been hypothesized that climate-mediated divergence in the pollination environment may drive the adoption of a novel pollinator niche (Grant and Grant 1965; Stebbins 1970; Phillips et al. 2020). To better understand the role of pollinator specificity in C. flammulus speciation, it is necessary to carry out reciprocal translocation experiments to tease apart the relative contribution of climatic and pollinator niches and their interactions (Miller et al. 2014; Van der Niet et al. 2014; Phillips et al. 2020).

We found almost no geographic and climatic niche overlap, suggesting that strong ecogeographic isolation is the main barrier segregating *C. flammulus* and *C. pulverulentus*. Ecogeographic differentiation is thought to evolve in the early stages of speciation; thus, studying ecogeographic isolation is crucial to understanding incipient species divergence (Schemske 2000, 2010; Sobel et al. 2010). It has been demonstrated that ecogeographic differentiation and geographic isolation are drivers of speciation in several Neotropical plant groups (Kay 2006; Nakazato et al. 2010; Castillo-Cardenas et al. 2015; Chen and Schemske 2015; Cuevas et al. 2018; Maciel et al. 2020; Misiewicz et al. 2020; Vargas et al. 2020). Non-overlapping ranges and climatic niches suggest that *C. flammulus* and *C. pulverulentus* differ significantly in the niches they can occupy.

Together, our results are consistent with the hypothesis of *C. flammulus* having budded off from a *C. pulverulentus*-like progenitor species. Budding, or progenitor-derivative, speciation involves an initially small colonizing population becoming reproductively isolated from a larger-ranged species, generally in a marginal habitat at the range edge (Mayr 1954; Grant 1981). Budding speciation is recognized as an important

phenomenon in plants (Richerson and Lum 1980; Gottlieb 2004; Crawford 2010; Anacker and Strauss 2014; Grossenbacher et al. 2014; Sianta and Kay 2022). However, in proportion to its importance, the role of budding speciation in the Neotropical flora is still poorly understood (but see Vargas et al. 2020). Whether *C. flammulus* is derived from a *C. pulverulentus*-like progenitor or not is still uncertain and could benefit from detailed population genomic studies of its demographic and speciation history. Nevertheless, several lines of evidence, such as possible *C. pulverulentus* paraphyly, range size asymmetry, and *C. flammulus*' nested distribution of vegetative traits, suggest that *C. flammulus* budded off from a *C. pulverulentus*-like progenitor species (Crawford 2010; Sianta and Kay 2022), evolving to tolerate a colder and more seasonal montane environment.

TAXONOMIC TREATMENT

Costus flammulus K.M.Kay & P.Juárez, sp. nov. Type: Costa Rica. Puntarenas, Monteverde. E. B. Monteverde, Quebrada Máquina (10°19′03′′N, 84°48′27′′W), 1524 m, 18 Aug 2022 (fl), P. Juárez & C. Girvin 1620 (holotype: USJ!, isotypes CR!, MO!).

Closely related and vegetatively similar to *C. pulverulentus*, it differs in the smaller stem diameter (6.79 mm vs. 8.96 mm in diam), the smaller leaves $(14.67 \times 4.30 \text{ cm vs. } 18.23 \times 5.34 \text{ cm})$, the shorter inflorescences (6.44 vs. 9.78 cm long), the smaller bracts $(1.76 \times 1.56 \text{ cm vs. } 2.56 \times 1.87 \text{ cm})$, the shorter nectarial callus (non-functional, 6.14 mm vs. functional, 8.82 mm long), the smaller flowers (Table 1), the antrorse, red and distally yellow corolla lobes (vs. retrorse, red corolla lobes), the yellow to red, antrorse, shortly 5-lobed labellum (vs. yellow, 5-lobed, lobes retrorse), the petaloid stamen shortly exserted by 5 mm (vs. petaloid stamen exserted by 16.27 mm), and geographic distribution (montane cloud forest of the Tilarán and Guanacaste cordilleras in Costa Rica, at elevations of 1000–1650 m vs. lowland wet forests from Mexico to southern Ecuador, at elevations of 0–1400 m). The measurement contrasts are means. See Table 1 for detailed comparisons.

Perennial herb, (0.5) 1.5 (3) m tall, hermaphroditic, geophyte. Rhizomes reddish, sympodial, every element ending in a shoot. Stems 6.7-9 mm in diam, orthotropic, green to reddish near the base, terete, glabrous, with alternate-spiraled phyllotaxy. Sheaths 6.7–9 mm in diam, green, glabrous. Ligules 5-5.7 (8) mm long, green, truncate, or slightly twolobed, glabrous, margins sometimes dilacerating into fibers. Leaves $18.5-24\times4-7$ cm; petioles 5-5.7 (8) mm long; blade elliptic-obovate, base rounded to cordate, apex acuminate, margins undulate, adaxial surface glabrous, abaxial surface slightly glaucous, chartaceous, venation parallel. Inflorescences strobilaceous-capitate spike, terminal, (4.6) 7.9–13.7 \times (2) 3.2–5.8 cm, cylindrical-ovoid. Bracts $1.4-2.3 \times 1-2.2$ cm, red to distally yellow, ovate-triangular, apex acute, margins glabrescent, sometimes dilacerating into fibers, glabrous, coriaceous, subtending 1 flower per bract; callus 4-7 mm long, yellow, linear, non-functional. Bracteole 1 (1.5) cm long, red, glabrous. Flowers hermaphroditic, sympetalous, zygomorphic; calyx connate, 4.2 mm long, red, truncate or slightly 3-lobed, glabrous; corolla connate, 2.9–4.1 cm long, proximally white to red to distally yellow, venation red, 3-lobed, lobes antrorse, proximally tubular; tube (0.7) 1.2 cm long, white,




Fig. 4. Costus flammulus K. M. Kay & P. Juárez. Line illustration based on P. Juárez & C. Girvin 1620 (USJ, CR, MO) and live plants from the type locality. A. Stem with terminal spike with flower. B. Flower, side view. C. Dorsal corolla lobe, abaxial view. D. Lateral corolla lobe, abaxial view. E. Lateral corolla lobe, adaxial view. F. Labellum, adaxial view. G. Petaloid stamen, style, and stigma. H. Detail of bilamellated stigma. I. Ovary and calyx. J. Capsule with persistent calyx. K. Floral diagram of Costus. Drawn by P. Juárez.



Fig. 5. *Costus flammulus*. A. Terminal spike with flower. B. Stem, petiole, and ligules. Note ligules' margins sometimes dilacerating into fibers. C. *Phaethornis guy* visiting *C. flammulus*' flower. D. Young capsule subtended by bracteole. Note that bracts have been removed. E. Leaves, abaxial view. Note undulate blade margins. F. *Costus wilsonii*, a species with which *C. flammulus* has been commonly mistaken in herbaria. Photographs by P. Juárez (A, C, F); J. E. Jiménez (B, E); and C. Girvin (D).

glabrous, throat adaxially covered with yellow hairs; **dorsal lobe** 2.5– 4.1×0.7 –1.2 cm, obovate, apex mucronate, margin slightly involute; **lateral lobes** 2.4– 3.7×0.6 –0.9 cm, elliptic, apex obtuse, margin involute; **labellum** 3– 3.7×1.4 –1.8 cm, yellow to red, spatulate, apex irregular, antrorse, shortly 5-lobed, lobes < 1 mm long, lateral lobes involute, forming a tube of ca. 4.5 mm in diameter; **stamen** 3.1– 3.8×0.6 –0.8 cm, yellow to reddish, petaloid, narrowly elliptic, apex obtuse, exserted, ca. 5 mm exceeding the labellum, anthers 7 mm long, thecae open in longitudinal slits, pollen white; **ovary** trilocular, 3.4×4.2 mm, white, globose, trigonous, many ovules; **style** 2.7–3.3 cm long, whitish, filiform; **stigma** ca. 3 mm wide, whitish, bilamellated, margin ciliate, appressed to anthers. **Capsules** 1– 1.3×0.7 –0.9 cm, white, obovoid, glabrous, calyx persistent. **Seeds** ca. 4 mm wide, black. Figures 4, 5.

Distribution and Habitat—Costus flammulus is endemic to the primary and secondary montane cloud forests, at elevations of 1000-1650 m, on the Tilarán and Guanacaste volcanic cordilleras in northern Costa Rica (Fig. 1). The montane cloud forests of the northern volcanic cordilleras are characterized by high edaphic, microclimatic, and topographic diversity, as well as by high inter-annual precipitation variation (Lawton et al. 2016). Costus flammulus has been collected in several relevant protected areas, both state and private, such as the Guanacaste, Miravalles-Jorge Manuel Dengo, and Tenorio Volcano national parks, and the Alberto Brenes and Monteverde Cloud Forest biological reserves. Locally, C. flammulus is frequently found near streams, in rocky ravines, or in the understory. In the Tilarán Cordillera, C. flammulus co-occurs with the distantly related C. montanus Maas, which likely hybridize due to microhabitat and pollinator assemblage overlap (P. Juárez pers. obs.). See Table S3 for detailed information on C. flammulus occurrences.

Phenology—*Costus flammulus* flowers in the dry and rainy seasons. It has been recorded flowering in February, April, July, August, and October, and fruiting in October and November.

Etymology—The specific epithet is a diminutive of the Latin *flamma* or *flammae*, meaning "flame" (Gledhill 2008), referring to the flower's resemblance to a little flame.

Notes—Costus flammulus has been mistakenly included in C. wilsonii (e.g. Stratton 1989; Haber 2000; Maas and Maas-van de Kamer 2003). Costus flammulus vegetative morphology and habitat affinity resemble that of C. wilsonii; thus, distinguishing herbarium specimens could be difficult without flowers. Besides, Costus flowers are delicate and generally not preserved well in herbarium specimens. Presumably, C. flammulus and C. wilsonii have been confused because both species are hummingbird-pollinated herbs from montane forests with truncated or slightly two-lobed ligules and undulate leaf margins.

Costus flammulus and C. wilsonii can be differentiated by their dried foliage color, ligule margin, and distribution range in herbarium specimens. Costus flammulus foliage usually dries yellowish, whereas C. wilsonii foliage dries grayish. Costus flammulus ligule margins sometimes dilacerate into fibers (Fig. 5), whereas C. wilsonii does not. Costus flammulus is endemic to the Tilarán and Guanacaste cordilleras in northern Costa Rica. Costus wilsonii occurs in the Talamanca Cordillera, Fila Cruces, and the most southern volcanic peaks of the Central Cordillera, ranging from the southern half of Costa Rica to the north of Panama.

Although *C. flammulus* and *C. wilsonii* herbarium specimens may be confused because of their vegetative resemblance and habitat preferences, their flowers are distinguishably different (Fig. 5). *Costus wilsonii* flowers are yellow and differ from the rest of hummingbird-pollinated *Costus* by their expanded labellum

with red nectar guides on the limb. *Costus wilsonii* is the only hummingbird-pollinated *Costus* with nectar guides on the labellum, which potentially is a retained trait related to ancestral euglossine pollination. *Costus wilsonii* is the sister to *C. laevis* (Kay 2006; Vargas et al. 2020), a euglossine-pollinated widespread species inhabiting lowland wet forests of Central and South America.

Paratypes—Costa Rica. —ALAJUELA: Balsa. 10 km N of San Ramón, 1000 m, 13 Sep. 1974 (fl) P. J. M. Mass & R. W. Lent 1218 (MO!); San Carlos. Monteverde Reserve, El Valle trail, near continental divide with Atlantic exposure, lower montane rain forest (10°20'N, 84°50'W), 1600 m, 26 Oct. 1985 (fl) W. Haber & E. Bello 3161 (MO!); Peñas Blancas. 1200 m, 14 Sep. 1985 (fl) W. Haber & E. Bello 2725 (MO!); San Ramón. Cerro San Isidro, 1260 m, 25 Mar. 1982 (fl), A. Carvajal 191 (MO!); Angeles. A la vera del río S. Lorenzo, Faldas de la Fila Volcán Muerto, 1000-1100 m, 31 July 1982 (fl), J. Gómez-Laurito 8865 (CR!); Reserva de San Ramón, Estación Sn Lorencito, 700-1000 m, 02 July 1986 (fl), A. Chacón, I. Chacón & V. Mora 1981 (CR!); San Lorenzo. 12 km NNW of San Ramón by road on way to San Lorenzo (10°10'N, 84°29'W), 1100 m, 25 April 1983 (fl), R. Liesner & E. Judiewicz 14993 (MO!); Piedades Norte. Cuenca del Barranca, Sector Burio. Cerro Azahar (10°08'24"N, 84°33'40''W), 1400 m, 04 Oct. 1997 (fl), B. Hammel & H. Kennedy 21075 (CR!); Cuenca del Barranca, Sector Burio. Cerro Azahar (10°08'24''N, 84°33'40''W), 1400 m, 04 Oct. 1997 (fl), B. Hammel & H. Kennedy 21078 (CR!, MO!). —GUANA-CASTE: Abangares, La Cruz de Abangares, 1400 m, 15 July 1985 (fl) W. Haber & E. Bello 2027 (MO, digital image); Bagaces, La Fortuna. Parque Nacional Volcán Miravalles, Cabro Muco, sobre sendero camino hacia la catarata, bosque secundario (10°43'27.10"N, 85°08'38.62"W), 1246 m, 14 April 2022 (fl) J. E. Jiménez & M. F. Cordero 6527 (USJ!); La Cruz. SW slope of Volcán Orosí (10°58′00′′N, 85°28′30′′W), 1100–1400 m, 22 Aug. 2007 (fl) M. H. Grayum & A. Rojas 12835 (MO!); Liberia, Ángeles. SW (Pacific) slope of Cerro Cacao (10°55′53′′N, 85°27′35′′W), 1300–1450 m, 10 Aug. 2007 (fl) M. H. Grayum & D. García 12580 (MO!); Tilarán, Quebrada Grande. Cuenca del San Carlos, Reserva Biológica Bosque Nuboso, alrededores del laboratorio A. Skutch (10°20'40.67''N, 84°49'14''W), 1540 m, 29 Aug. 2011 (fl), D. Santamaría 8888 (CR!); Tierras Morenas. Z. P. Tenorio, Cabec. Río San Lorenzo (10°36'40''N, 84°59'45"W), 1050 m, 23 Aug. 1994 (fl), G. Rodriguez 307 (CR!, MO!); Tronadora. San Gerardo Biological Station. Road to Río Caño Negro, secondary forest (10°22'00''N, 84°47'00''W), 1100 m, 1 Apr. 1995 (fl), D. Penneys 294 (CR!, MO!). —Puntarenas: Puntarenas, Monteverde. Monteverde Cloud Forest Reserve, 1540-1620 m, 12 Aug. 1982 (fl), P. Feinsinger et al. 409 (MO!); Santa Elena de Monteverde, Tilarán road, c. 1 km from S. Elena. 1300 m, 17 Feb. 1984 (fl), R. Khan, M. Tebbs & A. R. Vickery 1091 (CR!); R. B. Bosque Eterno de los Niños, Cuencas del Lagarto y Guacimal, sendero Chomogo y el Roble (10°18′00′′N, 84°48′00′′W), 1600 m, 19 Oct. 1997 (fl), A. Soto 73 (CR!, MO!); Cloud forest and swamp forest (premontane and lower montane rain forest formations) in and around the Monteverde Nature reserve, mostly on the Pacific watershed (10°18'N, 84°47'W), 1450-1650 m, 31 Oct. & 2 Nov. 1975 (fl) W. Burger & R. Baker 9720 (MO!); Quebrada Máquina, Pacific slope (10°18′N, 84°48′W), 1400 m, 9 Aug. 1991 (fl), W. Haber, C. Ivey & W. Zuchowski 10798 (CR!, MO!).

ACKNOWLEDGMENTS

We thank M. Blanco (USJ), S. Lobo, J. Sánchez, and A. Quesada (CR), R. Quiros (HLDG), and M. Morello (MO) for helping in herbaria and providing specimen images. We thank C. Girvin for assistance during fieldwork and for providing valuable comments on the manuscript, and D. Tenorio for measuring phenotypic traits on herbarium specimens. We thank the Estación Biológica Monteverde for facilitating fieldwork. All field research was conducted with appropriate permits in Costa Rica (M-P-SINAC-PNI-ACAT-031-2021, and CONAGEBIO R-056-2019-OT and R-058-2019-OT). This research was funded by the National Science Foundation Dimensions of Biodiversity grant DEB-1737878 to K. M. Kay.

AUTHOR CONTRIBUTIONS

PJ and KMK did the fieldwork. PJ analyzed all data, except for OV who did the phylogenetic analysis. PJ prepared the figures and tables. PJ wrote the manuscript with edits of KMK and OV.

LITERATURE CITED

Anacker, B. L. and S. Y. Strauss. 2014. The geography and ecology of plant speciation: range overlap and niche divergence in sister species. *Proceedings. Biological Sciences* 281: 20132980.

- Andersson, L. 1990. The driving force: Species concepts and ecology. *Taxon* 39: 375–382.
- Baldwin, B. G. 2005. Origin of the serpentine-endemic herb *Layla discoidea* from the widespread *L. landulosa* (Compositae). *Evolution* 59: 2473–2479.
- Baum, D. A. and S. D. Smith. 2013. Tree Thinking. An Introduction to Phylogenetic Biology. New York: W. H. Freeman & Company.
- Benjamini, Y. and Y. Hochberg. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. Journal of the Royal Statistical Society. Series B. Methodological 57: 289–300.
- Broennimann, O., M. C. Fitzpatrick, P. B. Pearman, B. Petitpierre, L. Pellissier, N. G. Yoccoz, W. Thuiller, M. J. Fortin, C. Randin, N. E. Zimmermann, C. H. Graham, and A. Guisan. 2012. Measuring ecological niche overlap from occurrence and spatial environmental data. *Global Ecology and Biogeography* 21: 481–497.
- Castillo-Cardenas, M. F., J. A. Ramirez-Śilva, O. Sanjur, and N. Toro-Perea. 2015. Evidence of incipient speciation in the Neotropical mangrove Pelliciera rhizophorae (Tetrameristaceae) as revealed by molecular, morphological, physiological and climatic characteristics. Botanical Journal of the Linnean Society 179: 499–510.
- Chen, G. F. 2013. Sexual isolation in two bee-pollinated Costus (Costaceae). Plant Reproduction 26: 3–16.
- Chen, G. F. and D. W. Schemske. 2015. Ecological differentiation and local adaptation in two sister species of Neotropical Costus (Costaceae). Ecology 96: 440–449.
- Coyne, J. A. and H. A. Orr. 2004. Speciation. Sunderland: Sinauer Associates. Crawford, D. J. 2010. Progenitor-derivative species pairs and plant speciation. Taxon 59: 1413–1423.
- Cuevas, E., J. Espino, and I. Marques. 2018. Reproductive isolation between Salvia elegans and S. fulgens, two hummingbird-pollinated sympatric sages. Plant Biology 20: 1075–1082.
- Dayrat, B. 2005. Towards integrative taxonomy. *Biological Journal of the Linnean Society. Linnean Society of London* 85: 407–417.
- De Queiroz, K. 1998. The general lineage concept of species, species criteria, and the process f speciation. Pp. 57–75 in *Endless Forms: Species and Speciation*, eds. D. J. Howard and S. H. Berlocher. New York: Oxford University Press.
- De Queiroz, K. 2007. Species concepts and species delimitation. *Systematic Biology* 56: 879–886.
- Dobzhansky, T. 1970. Genetics of the Evolutionary Process. New York: Columbia University Press.
- Donoghue, M. J. 1985. A critique of the biological species concept and recommendations for a phylogenetic alternative. *The Bryologist* 88: 72–181.
- Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004. Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* 35: 375–403.
- Foote, M. 1996. On the probability of ancestors in the fossil record. *Paleobiology* 22: 141–151.
- Fulton, M. and S. A. Hodges. 1999. Floral isolation between Aquilegia formosa and Aquilegia pubescens. Proceedings of the Royal Society of London. Series B, Biological Sciences 266: 2247–2252.
- Futuyma, D. J. 1998. Evolutionary Biology. Sunderland: Sinauer Associates. Garrigues, R. and R. Dean. 2014. The Birds of Costa Rica: A Field Guide. New York: Cornell University Press.
- Gledhill, D. 2008. The Names of Plants. Cambridge: Cambridge University Press.
- Grant, V. 1949. Pollination systems as isolating mechanisms in angiosperms. Evolution 3: 82–97.
- Grant, V. 1981. Plant Speciation. New York: Columbia University Press.
- Grant, V. 1993. Origin of floral isolation between ornithophilous and sphingophilous plant species. Proceedings of the National Academy of Sciences USA 90: 7729–7733.
- Grant, V. 1994. Modes and origins of mechanical and ethological isolation in angiosperms. Proceedings of the National Academy of Sciences USA 91: 3–10.
- Grant, V. and K. A. Grant. 1965. Flower Pollination in the Phlox Family. New York: Columbia University Press.
- Grossenbacher, D. L. and J. B. Whittall. 2011. Increased floral divergence in sympatric monkeyflowers. *Evolution* 65: 2712–2718.
- Grossenbacher, D. L., S. D. Veloz, and J. P. Sexton. 2014. Niche and range size patterns suggest that speciation begins in small, ecologically diverged populations in North American monkeyflowers (*Mimulus* spp.). Evolution 68: 1270–1280.

- Goff, K. A., C. Martinez Del Rio, and K. M. Kay. 2021. A greenhouse experiment partially supports inferences of ecogeographic isolation from niche models of *Clarkia* sister species. *American Journal of Botany* 108: 2002–2014.
- Gottlieb, L. D. 2004. Rethinking classic examples of recent speciation in plants. *The New Phytologist* 161: 71–82.
- Haber, W. A. 2000. Plants and vegetation. Pp. 39–94 in Monteverde: Ecology and Conservation of a Tropical Cloud Forest, eds. N. M. Nadkarni and N. T. Wheelwright. New York: Oxford University Press.
- Hendry, A. P. 2017. Eco-Evolutionary Dynamics. Princeton, New Jersey: Princeton University Press.
- Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith. 2017. dismo: Species distribution modeling. R package version 1.1–4.
- Johnson, S. D. 2006. Pollinator-driven speciation in plants. Pp. 295–310 in The Ecology and Evolution of Flowers, eds. L. D. Harder and S. C. H. Barrett. Oxford: Oxford University Press.
- Juárez, P., O. M. Vargas, and K. M. Kay. 2023. Data from: Ecological and evolutionary origin of Costus flammulus (Costaceae): A new species from the montane cloud forests of the volcanic cordilleras in northern Costa Rica. Dryad Digital Repository. https://datadryad.org/ stash/share/ZzsH-Cqz62XBtMm737eN7FnXIkNuH-A0AVmCX6KU.
- Kay, K. M. 2006. Reproductive isolation between two closely related hummingbird pollinated neotropical gingers. Evolution 60: 538–552.
- Kay, K. M. and D. L. Grossenbacher. 2022. Evolutionary convergence on hummingbird pollination in Neotropical Costus provides insight into the causes of pollinator shifts. The New Phytologist 236: 1572–1583.
- Kay, K. M. and R. D. Sargent. 2009. The role of animal pollination in plant speciation: Integrating ecology, geography, and genetics. *Annual Review of Ecology, Evolution, and Systematics* 40: 637–656.
- Kay, K. M. and D. W. Schemske. 2003. Pollinator assemblages and visitation rates for 11 species of Neotropical Costus (Costaceae). Biotropica 35: 198–207.
- Kay, K. M. and Y. Surget-Groba. 2022. The genetic basis of floral mechanical isolation between two hummingbird-pollinated Neotropical understorey herbs. *Molecular Ecology* 31: 4351–4363.
- Kay, K. M., P. A. Reeves, R. G. Olmstead, and D. W. Schemske. 2005. Rapid speciation and the evolution of hummingbird pollination in neotropical *Costus* subgenus *Costus* (Costaceae): Evidence from nrDNA ITS and ETS sequences. *American Journal of Botany* 92: 1899–1910.
- Kress, W. J. 1983. Crossability barriers in Neotropical Heliconia. Annals of Botany 52: 131–147.
- Lawton, R. O., M. F. Lawton, R. M. Lawton, and J. D. Daniels. 2016. The montane cloud forests of the volcanic cordilleras. Pp. 415–450 in Costa Rican Ecosystems, eds. M. Kappelle, T. E. Lovejoy, and R. G. Lobo. Chicago: University of Chicago Press.
- Maas, P. J. M. 1972. Costoideae (Zingiberaceae). Flora Neotropica 8: 1–139.
- Maas, P. J. M. 1977. *Renealmia* (Zingiberaceae-Zingiberoideae) Costoideae (Additions) (Zingiberaceae). *Flora Neotropica* 18: 1–218.
- Maas, P. J. M. and H. Maas-van de Kamer. 2003. Costaceae. Pp. 413–423 in *Manual de Plantas de Costa Rica*, vol. II, eds. B. E. Hammel, M. H. Grayum, C. Herrera, and N. Zamora. St. Louis: Missouri Botanical Garden Press.
- Maciel, J. R., G. Zizka, and M. Alves. 2020. Differential geographical and ecological dynamics allow diversification of morphologically convergent giant bromeliads in the Atlantic Forest. *Journal of Biogeography* 47: 2684–2697.
- Mayr, E. 1942. Systematics and the Origin of Species. New York: Columbia University Press.
- Mayr, E. 1947. Ecological factors in speciation. Evolution 4: 263-288.
- Mayr, E. 1954. Change of genetic environment and evolution. Pp. 157–180 in *Evolution as a Process*, eds. J. Huxley, A. C. Hardy, and E. B. Ford. New York: Collier Books.
- Michener, C. D. 1970. Diverse approaches to systematics. *Evolutionary Biology* 4: 1–38.
- Miller, T. J., R. A. Raguso, and K. M. Kay. 2014. Novel adaptation to hawk-moth pollinators in *Clarkia* reduces efficiency, not attraction of diurnal visitors. *Annals of Botany* 113: 317–329.
- Mishler, B. D. 1985. The morphological, developmental, and phylogenetic basis of species concepts in bryophytes. *The Bryologist* 88: 207–214.
- Misiewicz, T. M., T. S. Simmons, and P. V. Fine. 2020. The contribution of multiple barriers to reproduction between edaphically divergent lineages in the Amazonian tree *Protium subserratum* (Burseraceae). *Ecology and Evolution* 10: 6646–6663.

- Muchhala, N. and J. D. Thomson. 2012. Interspecific competition in pollination systems: Costs to male fitness via pollen misplacement. Functional Ecology 26: 476–482.
- Nakazato, T., D. L. Warren, and L. C. Moyle. 2010. Ecological and geographic modes of species divergence in wild tomatoes. *American Journal of Botany* 97: 680–693.
- Pessoa, E., M. Sader, A. Pedrosa-Harand, and M. Alves. 2020. A natural hybrid, an autopolyploid, or a new species? An integrative case study of a distinctive Costus species (Costaceae) from the Atlantic Forest of Brazil. Systematics and Biodiversity 18: 42–56.
- Phillips, R. D., R. Peakall, T. Van der Niet, and S. D. Johnson. 2020. Niche perspectives on plant–pollinator interactions. *Trends in Plant Science* 25: 779–793.
- Phillips, S. J., M. Dudík, J. Elith, C. H. Graham, A. Lehmann, J. Leathwick, and S. Ferrier. 2009. Sample selection bias and presence-only distribution models: Implications for background and pseudo-absence data. *Ecological Applications* 19: 181–197.
- Presl, K. B. 1827. Reliquiae Haenkeanae 1.
- R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ramsey, J., H. D. Bradshaw Jr., and D. W. Schemske. 2003. Components of reproductive isolation between the monkeyflowers *Minulus lewisii* and *M. cardinalis* (Phrymaceae). *Evolution* 57: 1520–1534.
- Raxworthy, C. J., C. M. Ingram, N. Rabibisoa, and R. G. Pearson. 2007. Applications of ecological niche modeling for species delimitation: A review and empirical evaluation using day geckos (*Phelsuma*) from Madagascar. Systematic Biology 56: 907–923.
- Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other things). Methods in Ecology and Evolution 3: 217–223.
- Richerson, P. J. and K. Lum. 1980. Patterns of plant species diversity in California: Relation to weather and topography. American Naturalist 116: 504–536.
- Ringnér, M. 2008. What is principal component analysis? Nature Biotechnology 26: 303–304.
- Rissler, L. J. and J. J. Apodaca. 2007. Adding more ecology into species delimitation: Ecological niche models and phylogeography help define cryptic species in the black salamander (*Aneides flavipunctatus*). Systematic Biology 56: 924–942.
- Schemske, D. W. 1982. Ecological correlates of a neotropical mutualism: Ant assemblages at Costus extrafloral nectaries. Ecology 63: 932–941.
- Schemske, D. W. 2000. Understanding the origin of species. Evolution 54: 1069–1073.
- Schemske, D. W. 2010. Adaptation and the origin of species. American Naturalist 176: S4–S25.
- Schiestl, F. P. and S. D. Johnson. 2013. Pollinator-mediated evolution of floral signals. *Trends in Ecology & Evolution* 28: 307–315.
- Schiestl, F. P. and P. M. Schlüter. 2009. Floral isolation, specialized pollination, and pollinator behavior in orchids. Annual Review of Entomology 54: 425–446.
- Schoener, T. W. 1970. Nonsynchronous spatial overlap of lizards in patchy habitats. *Ecology* 51: 408–418.
- Schwery, O. and B. O'Meara. 2016. MonoPhy: A simple R package to find and visualize monophyly issues. *PeerJ. Computer Science* 2: e56.

- Sianta, S. A. and K. M. Kay. 2022. Phylogenomic analysis does not support a classic but controversial hypothesis of progenitor-derivative origins for the serpentine endemic *Clarkia franciscana*. Evolution 76: 1246–1259.
- Skutch, A. F. and F. G. Stiles. 1989. A Guide to the Birds of Costa Rica. Utica: Cornell University Press.
- Sneath, P. H. A. and R. R. Sokal. 1973. Numerical Taxonomy: The Principles and Practice of Numerical Classification. San Francisco: W. H. Freeman & Company.
- Sobel, J. M. 2014. Ecogeographic isolation and speciation in the genus *Mimulus. American Naturalist* 184: 565–579.
- Sobel, J. M., G. F. Chen, L. R. Watt, and D. W. Schemske. 2010. The biology of speciation. Evolution 64: 295–315.
- Sokal, R. R. and T. J. Crovello. 1970. The biological species concept: A critical evaluation. *American Naturalist* 104: 127–153.
- Stebbins, G. L. 1970. Adaptive radiation of reproductive characteristics in angiosperms. I. Pollination mechanisms. Annual Review of Ecology and Systematics 1: 307–326.
- Stiles, F. G. 1975. Ecology, flowering phenology, and hummingbird pollination of some Costa Rican Heliconia species. Ecology 56: 285–301.
- Stiles, F. G. 1979. Notes on the natural history of *Heliconia* (Musaceae) in Costa Rica. *Brenesia* 15: 151–180.
- Stiles, F. G. 1983. Birds. Pp. 597–599 in Costa Rican Natural History, ed. D. H. Janzen. Chicago: University of Chicago Press.
- Stratton, D. A. 1989. Longevity of individual flowers in a Costa Rican cloud forest: Ecological correlates and phylogenetic constraints. *Bio-tropica* 21: 308–318.
- Tang, Y., M. Horikoshi, and W. Li. 2016. ggfortify: Unified interface to visualize statistical results of popular R packages. *The R Journal* 8: 474–485.
- Thiers, B. 2022 [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. http://sweetgum.nybg.org/science/ih/ (last accessed June 2022).
- Van der Niet, T. and S. D. Johnson. 2009. Patterns of plant speciation in the Cape floristic region. Molecular Phylogenetics and Evolution 51: 85–93
- Van der Niet, T., R. Peakall, and S. D. Johnson. 2014. Pollinator-driven ecological speciation in plants: New evidence and future perspectives. Annals of Botany 113: 199–212.
- Van Valen, L. 1976. Ecological species, multispecies, and oaks. Taxon 25: 233–239.
- Vargas, O. M., B. Goldston, D. L. Grossenbacher, and K. M. Kay. 2020. Patterns of speciation are similar across mountainous and lowland regions for a Neotropical plant radiation (Costaceae: Costus). Evolution 74: 2644–2661.
- Warren, D. L., R. E. Glor, and M. Turelli. 2008. Environmental niche equivalency versus conservatism: Quantitative approaches to niche evolution. Evolution 62: 2868–2883.
- Wickham, W. 2009. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag.
- Zapata, F. and I. Jiménez. 2012. Species delimitation: Inferring gaps in morphology across geography. Systematic Biology 61: 179–194.