

Cross-Selective Aza-Pinacol Coupling via Atom Transfer Catalysis

Sean M. Rafferty,[†] Joy E. Rutherford,[†] Lumin Zhang, Lu Wang, and David A. Nagib*

Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH 43210, United States.

Supporting Information Placeholder

ABSTRACT: A cross-selective aza-pinacol coupling of aldehydes and imines has been developed to afford valuable β -amino alcohols. This strategy enables chemo-selective conversion of aliphatic aldehydes to ketyl radicals – in the presence of more easily reduced imines and other functional groups. Upon carbonyl-specific activation by AcI, a photoinitiated Mn catalyst selectively reduces the resulting α -oxy iodide by an atom transfer mechanism. The ensuing ketyl radical selectively couples to imines – precluding homo-dimerization by a classical reductive approach. In this first example of reductive, ketyl coupling by atom transfer catalysis, Zn serves as a terminal reductant to facilitate Mn catalyst turnover. This new strategy also enables ketyl radical couplings to alkenes, alkynes, aldehydes, propellanes, and chiral imines.


Ketyl radicals provide versatility to the synthetically valuable carbonyl group by reversing its reactivity from $2e^-$ electrophile to $1e^-$ nucleophile.¹ This polarity reversal strategy enables streamlined access to useful motifs such as 1,2-diols via pinacol coupling of carbonyls.²⁻⁴ Similarly, β -amino alcohols – a privileged structure found in nature, catalysis, and medicine⁵⁻⁶ – can be accessed by aza-pinacol coupling of an aldehyde and imine.⁷ This convergent C-C coupling provides synthetic modularity and avoids regioselectivity issues associated with typical routes to β -amino alcohols via alkenes or epoxides.⁵⁻⁶ Nonetheless, underlying thermodynamic and chemoselectivity challenges remain for direct aza-pinacol coupling (Figure 1). Notably, the generation of ketyl radical anions requires strong reductants (e.g. Na, Mg, Ti, Sm) with highly negative redox potentials necessary to reduce carbonyls ($E_{\text{red}} > -1.9$ V vs SCE)⁸⁻¹² – especially aliphatic aldehydes (-2.9 V).⁸ Moreover, since imines are more easily reduced (Ph: -1.9 V; Ms: -1.5 V),⁹⁻¹¹ it is difficult to chemoselectively generate ketyl radicals in their presence. Instead, single-electron transfer (SET) reduction of an imine – and subsequent homo-dimerization of the resulting α -amino radical – is typically observed (Figure 1a).^{1,13}

Recent strategies by Yoon, Knowles, and others to access ketyl radicals by milder reductants, such as Ru and Ir photocatalysts, entail Brønsted or Lewis acid-activation of carbonyls to lower their reduction potential.¹⁴⁻²¹ Nonetheless, such key advances remain subject to thermodynamic favorability for reducing imines versus aldehydes. To address this ongoing challenge of cross-selectivity,¹ aza-pinacol couplings typically entail either (1) intramolecularity, or (2) slow addition of an α -amino radical precursor to super-stoichiometric quantities of reductant (e.g. Sm) and the less reducible, carbonyl partner.²²⁻²⁶ Other recent approaches to bypass this challenge entail metal-catalyzed hydro/silyl functionalization²⁷⁻²⁹ or reductive couplings,³⁰⁻³¹ alkyl-Cr addition to carbonyls,³²⁻³⁴ and Sn-mediated electrochemistry.³⁵

To solve the aza-pinacol coupling challenge, we proposed an alternate strategy to harness ketyl radicals by an atom transfer mechanism.³⁶ In this approach (Figure 1b), an aldehyde is first combined with AcI *in situ* to form an α -oxy iodide, which is nearly 2 V easier

to reduce than the parent carbonyl. This improved radical precursor contains a weak C-I bond (58 kcal/mol) that may also be cleaved by iodine atom abstraction to chemoselectively form ketyl radicals of aliphatic aldehydes in the presence of more easily reduced imines. Whereas we have shown *redox-neutral* addition of ketyl radicals to alkynes afford Z-vinyl iodides,³⁶ we reasoned a suitable, terminal H-donor may now also afford *reductive* reactivity – in the form of an aza-pinacol coupling.

a. Challenge: Cross-selectivity in aza-pinacol coupling



b. Strategy: Chemoselective ketyl radical generation via atom transfer




Figure 1. Design of cross-selective strategy for aza-pinacol coupling.

To test our hypothesis (Figure 2), pentanal **1** and imine **2a** were irradiated with visible light in the presence of a photocatalyst (1% Ir(ppy)₂(dtbbpy)PF₆) and terminal reductant (3 equiv *n*Bu₃N). As expected from previous reports,¹³ this SET strategy does not provide aza-pinacol adduct **3**, but instead affords only imine-derived homo-dimer **4a** (68%). Conversely, to probe our atom transfer strategy, AcI and pentanal **1** were first combined in CH₂Cl₂ at 0 °C for 15 minutes. The *in situ* generated α -oxy iodide **1a** was then

subjected to imine **2b** and photocatalytic conditions. To our delight, a complete switch in reactivity was observed, wherein *cross-selective* aza-pinacol adduct **3** is now obtained exclusively (76%) – without homo-dimer **4** (0%). Desiring an imine with a more easily removed protecting group, we replaced N-aryl imines **2a-b** with sulfonimine **2c** (R = Ms). However, this more easily reduced imine (Ms: –1.5 V vs Ph: –1.9 V)⁹⁻¹¹ falls within the redox window of the Ir photocatalyst (–1.5 V)³⁷⁻³⁸ and thus yields exclusive dimerization (64% **4c**). To solve this problem, we pivoted from an SET reduction mechanism to atom transfer activation by a Mn₂(CO)₁₀ catalyst.³⁹⁻⁴¹ In this case, iodide **1a** and sulfonimine **2c** selectively afford cross-coupled adduct **3c** (56%) without any dimer **4** (0%).




Figure 2. Development of cross-selective aza-pinacol coupling.

To further improve this atom transfer-enabled, aza-pinacol coupling, several reaction parameters were examined (Table 1). First, including a co-reductant (e.g. Hantzsch ester) affords cross-coupled adduct **3** in >20:1 chemoselectivity over dimer **4** – for three classes of imines (R = Ph, Ts, Ms, entries 1-3). Although excluding the Hantzsch reductant diminishes efficiency (10% vs 56%, entries 3-4), replacing with Zn recovers reactivity (40%, entry 5) and affords a simpler purification (separation of Zn salts versus co-eluting Hantzsch pyridine). A switch from irradiation by blue LED to a broader spectrum white CFL further improves reactivity (70%, entry 6). And the hindered base, Cy₂NMe, proves superior to either *i*Pr₂NEt or KOAc (90%, entries 6-8) – suggesting its likely role as H-atom donor to the transient N-radical. Finally, increased catalyst loading (15% vs 5%) affords >99% yield within two hours (entry 9), and 1:1 stoichiometry (i.e. without excess aldehyde) also provides efficient cross-selective reactivity (entry 10).

The scope and generality of this mild, cross-selective aza-pinacol coupling was then explored using a wide variety of imines and aldehydes (Table 2). For example, several *ortho*-, *meta*-, and *para*-substituted aryl aldimines were shown to be effective imine partners (**5-14**). Notably, an exceptionally broad range of electronically diverse substituents were tolerated – spanning Hammett constants (σ_p) of –0.3 (OMe) to +0.5 (CF₃) as well as a wide redox window,⁹ including imines that may otherwise afford homodimerization via

Table 1. Reaction optimization

entry	R	light	reductant	base	yield 3	3 : 4
1	Ph	blue LED	Hantzsch	<i>i</i> Pr ₂ NEt	28%	>20:1
2	Ts	blue LED	Hantzsch	<i>i</i> Pr ₂ NEt	50%	>20:1
3	Ms	blue LED	Hantzsch	<i>i</i> Pr ₂ NEt	56%	>20:1
4	Ms	blue LED	none	<i>i</i> Pr ₂ NEt	10%	>20:1
5	Ms	blue LED	Zn	<i>i</i> Pr ₂ NEt	40%	>20:1
6	Ms	white CFL	Zn	<i>i</i> Pr ₂ NEt	70%	>20:1
7	Ms	white CFL	Zn	KOAc	29%	>20:1
8	Ms	white CFL	Zn	Cy ₂ NMe	90%	>20:1
9 ^a	Ms	white CFL	Zn	Cy ₂ NMe	99%	>20:1
10 ^b	Ms	white CFL	Zn	Cy ₂ NMe	70%	>20:1


Conditions: 0.2 mmol imine, pentanal (3 eq), AcI (2.6 eq), 5% Mn₂(CO)₁₀, reductant (2 eq), base (5 eq), CH₂Cl₂ (1 mL), 24 h. ^a15% Mn₂(CO)₁₀, 2 h. ^baldehyde (1 eq).

an SET manifold. Steric variation similarly does not inhibit efficiency, although *ortho* substituents strongly influence diastereoselectivity with *o*-Me groups affording β -amino alcohols in up to 9:1 dr (**11**, **15**). Bis-substitution is well-tolerated for both sterically (**15**) and electronically exaggerated cases, such as bis-CF₃ arene (**16**) and bis-F pyridine (**17**) – again without dimerization. Lastly, a pair of mechanistic probes were investigated, containing weak, benzylic C-H bonds that may facilitate H-atom transfer (HAT) to the N-radical intermediate.^{42,43} However, since no remote functionalization products were observed (by either inter- or intra-molecular HAT), including by the strong H-atom donor, fluorene, (**18-19**), we conclude N-radical termination occurs more rapidly.

To critically probe the chemoselectivity of this cross-selective coupling, we designed a series of experiments examining imines with acid-labile or easily reducible groups that would not be tolerated under typical SmI₂, photocatalytic, or other highly reducing conditions. In each of these cases, aza-pinacol coupling was the exclusive product observed. For example, benzofuran (**20**), acetal (**21**), aryl nitrile (**22**), and aryl iodides (**23-25**) all remain intact. Most notably, benzophenone (**26**; –1.3 V)⁴⁴ is unperturbed despite its less negative potential than the imine (–1.5 V) – or the aldehyde (–2.9 V) that is chemoselectively converted to a ketyl radical.


Next, a range of aliphatic aldehydes were investigated to determine the generality of β -amino alcohols accessible by this ketyl coupling. Steric effects appear minimal as both smaller acetaldehyde (**27-28**) and larger isobutyraldehyde (**29**) provide similar efficiency, with the imine partner having a stronger stereoselectivity influence (**28**; 10:1 dr with *o*-tolyl imine). As further chemoselectivity probes, reductively prone functionality was also preserved on the aldehyde component, including esters, nitriles, and alkyl halides (**30-33**). Synthetically useful heteroatom substitution is also tolerated, such as ethers, imides, and amides (**34-36**), including within a natural product derivative of deoxycholic acid (**38**). Conversely, both the aldehyde and imine partners may also be entirely aliphatic (**18**, **37**).⁴⁵

Table 2. Scope, generality, and chemoselectivity of catalytic, atom transfer, cross-aza-pinacol coupling



^a0.2 mmol imine, aldehyde (3 eq), AcI (2.6 eq), 5% Mn₂(CO)₁₀, Zn (2 eq), Cy₂NMe (5 eq), CH₂Cl₂, (1 mL), 2 x 23W white CFL, 24 h. ^b10% Mn₂(CO)₁₀. ^c*i*Pr₂NEt instead of Cy₂NMe. ^dwith Hantzsch ester (0.75 eq). See SI for full details.

Given the strong influence of the imine on controlling diastereoselectivity, we sought to access a single enantiomer of β -amino alcohols by use of a chiral auxiliary (**Eq 1**).⁴⁶⁻⁴⁷ To this end, we found S-aryl sulfinimines enable more efficient radical-coupling than the S-*t*Bu analog (<10%). Moreover, 2,4-di-F-phenyl sulfinyl aldimine (**38**) affords superior stereoselectivity (18:3:3:1 dr **39**; 18:1 dr at C-N stereocenter) vs typical variants: mesityl (7:1 dr) or tolyl (8:1 dr). Oxidation of the auxiliary by *m*CPBA then affords protected sulfonamide **40** in >98% ee.



We next sought to examine if other classes of *reductive* ketyl radical couplings could also be enabled by these mild atom transfer conditions (Figure 3). To complement our previous, *redox-neutral* ketyl couplings with alkynes (yielding *Z*-vinyl iodides),³⁶ we were pleased to find inclusion of Zn as a terminal reductant instead affords *E*-allyl acetate **41**. Notably, these mild conditions do not over-reduce the acrylate product. Separately, addition to acrylates yield γ -acetoxy ester **42** via a ketyl coupling that was not observed under redox-neutral conditions. Lastly, pinacol cross-coupling of pentanal with more easily reduced glyoxylate (−1.4 V)⁴⁸ affords α -hydroxy- β -acetoxy ester **43**. This unique *cross*-selectivity of our atom transfer strategy is highlighted by divergent reactivity compared to SET-mediated photocatalysis,⁴⁹ which exclusively yields homo-dimeric tartrate **44**. Finally, given significant recent interest in the medicinal utility of bicyclo(1.1.1)pentane bioisosteres,⁵⁰⁻⁵¹ we

tested [1.1.1]propellane as a ketyl coupling partner via both reductive and Zn-free conditions. Interestingly, both pathways yield redox-neutral adduct **45** exclusively – indicating the rapid rate of iodine atom trapping by the bicyclopentyl radical and stability of the resulting alkyl iodide in this first ketyl-propellane coupling.




Figure 3. Additional classes of ketyl couplings enabled by atom-transfer catalysis.

Mechanistic studies were conducted to elucidate the nature of the key reductive turnover step in this Mn-catalyzed transformation. Unlike our previous *redox-neutral* studies, wherein radical rebound abstracts $I^\bullet$ from $[Mn]-I$,³⁶ catalyst turnover in this *reductive* process necessitates that a mild reductant liberates I^- to regenerate the 17-electron $[Mn^\bullet]$ complex. Mindful of redox data indicating Zn may not be strong enough to reduce $[Mn]-I$, we were especially curious about this selective reduction event. To investigate the role of

Zn in catalyst turnover, we first employed the proposed intermediate, $(CO)_5MnI$, as a catalyst instead of $Mn_2(CO)_{10}$ (Figure 4a). Interestingly, under photolytic conditions in the absence of Zn, we observed Cy_2NMe can activate $[Mn]-I$ to afford 25% yield, likely via an electron-donor-acceptor (EDA) complex. However, addition of both Zn and amine significantly improves efficiency (66% yield) – suggesting a productive interplay of all three components. This hypothesis was further supported by UV-vis analysis (Figure 4b). Background absorption spectra were obtained for $Mn_2(CO)_{10}$ ($\lambda_{max} = 343$ nm, yellow line) and $(CO)_5MnI$ ($\lambda_{max} = 300$ nm, orange line). Notably, the diagnostic absorption at 343 nm was not observed for combination of $(CO)_5MnI$ and Zn alone (grey line). However, when all three components were combined ($[Mn]-I$, Zn, Cy_2NMe), the signal was restored (blue line). Further evidence that an EDA complex promotes this reduction is a concentration-dependent loss of the $[Mn]-I$ signal with increasing $[Cy_2NMe]$, along with formation of a new 250 nm peak. Together, these experiments illustrate the combined action of both Zn and amine in reducing $[Mn]-I$ to $[Mn^\bullet]$ and enabling catalyst turnover.

To further elucidate the role of amine in enabling reduction, CV analysis confirms that Zn (-1.0 V) reduction of either $(CO)_5MnI$ ($E_{p/2} = -1.1$ V) or $Mn_2(CO)_{10}$ ($E_{p/2} = -1.2$ V) is thermodynamically disfavored (Figure 4c). Conversely, a 100mV anodic shift in reduction potential was observed upon addition of Cy_2NMe to $(CO)_5MnI$ ($E_{p/2} = -1.0$ V), indicating the $[Mn]-I:NR_3$ EDA complex facilitates milder reductive turnover of the Mn catalyst. Intriguingly, these results suggest photoinitiation may not be required, which we have validated by a pair of additional experiments conducted in the absence of light (Figure 4d). First, since $[Mn^\bullet]$ generation from $Mn_2(CO)_{10}$ is dependent on visible light homolysis of the Mn-Mn bond, minimal product was observed (10%) without this initiation. However, when proposed intermediate $[Mn]-I$, Zn, and amine were combined to initiate generation of the active $[Mn^\bullet]$ species, reactivity was observed in the dark (51%). This data suggests photolysis




Figure 4. Mechanistic experiments probing Mn catalyst turnover.

is required only for initiation and when $[\text{Mn}^\bullet]$ dimerization ($k = 9.5 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$)⁵² occasionally outcompetes ketyl radical generation by iodine abstraction.

Merging the insights from these experiments, a proposed mechanism is illustrated in Figure 5. Upon selective *in situ* activation of aldehyde **1** by AcI, this least reducible component is transformed into the most easily reduced, α -oxy iodide **1a** (−2.9 V to −1.1 V). To initiate catalysis, visible light homolysis of the weak Mn–Mn bond of $\text{Mn}_2(\text{CO})_{10}$ provides $(\text{CO})_5\text{Mn}^\bullet$ (**A**). Ketyl radical generation is chemoselectively accomplished by iodine atom transfer of the weak C–I bond (58 kcal/mol)³⁶ by **A** to form $\text{Mn}(\text{CO})_5\text{I}$ (**B**). The resulting α -acyloxy radical **C** selectively cross-couples to imine **2** (or other π -acceptors, such as alkynes, alkenes, aldehydes, and propellane) to form N-centered radical **D** (or corresponding C-radical). This electron-deficient amidyl radical is then terminated via either: (i) SET reduction by Zn and R_3N (both are necessary), or (ii) HAT from the polarity-matched α -amino C–H of Cy_2NMe .^{53,54} This final step occurs more rapidly than intramolecular HAT from even weak, benzylic C–H bonds – to afford aza-pinacol adduct **3**. Finally, catalyst turnover of $(\text{CO})_5\text{MnI}$ (**B**) is mediated by R_3N (via EDA complex **E**), whose ground state reduction potential is lowered by 100 mV (−1.1 V to −1.0 V) – allowing Zn to serve as a mild, chemoselective reductant. The regenerated catalyst **A** may propagate non-photolytic catalytic cycles until dimerization to $\text{Mn}_2(\text{CO})_{10}$ necessitates visible light homolysis to reinitiate the catalytic cycle.




Figure 5. Proposed mechanism of Mn-catalyzed aza-pinacol coupling.

In summary, a cross-selective aza-pinacol coupling has been developed for the synthesis of β -amino alcohols. An atom transfer catalytic approach enables chemoselective reduction of aldehydes to ketyl radicals in the presence of more easily reduced imines. This atypical selectivity facilitates cross-coupling reactivity and precludes homodimerization of the imine partner, as might otherwise be observed under typical reductive manifolds. A wide range of β -amino alcohols are accessed, including valuable chemoselectivity probes containing acid- and reductant-sensitive groups that would not otherwise be tolerated by Sm, Na, or strong photoreductants.

The synthetic utility of this strategy is further illustrated by ketyl couplings of alkynes, alkenes, aldehydes, [1.1.1]propellane, and stereoselective aza-pinacol coupling with chiral sulfinimine. This represents the first *reductive* version of our Mn-catalyzed, atom transfer strategy for accessing ketyl radical reactivity. Mechanistic studies illustrate the shared role of Zn and amine in Mn catalyst turnover and the rapid rates of termination of the radical relay. We envision this strategy will enable further discovery of non-classical ketyl radical couplings via atom transfer mechanisms.

ASSOCIATED CONTENT

The Supporting Information is available free of charge on the ACS Publications website at DOI:

Experimental procedures and characterization for all new compounds (PDF)

^1H and ^{13}C NMR spectral data (PDF)

AUTHOR INFORMATION

Corresponding Author

nagib.1@osu.edu

ORCID

David A. Nagib: 0000-0002-2275-6381

†Authors contributed equally

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank the National Science Foundation (CAREER 1654656) and National Institutes of Health (NIH R35 GM119812) for financial support. JER is supported by an NIH Fellowship (F31).

REFERENCES

- (1) Streuff, J. The Electron-Way: Metal-Catalyzed Reductive Umpolung Reactions of Saturated and α,β -Unsaturated Carbonyl Derivatives. *Synthesis* **2013**, 45, 281–307.
- (2) Procter, D. J.; Flowers, R. A.; Skrydstrup, T. *Organic Synthesis Using Samarium Diodide*; Royal Society of Chemistry: Cambridge, **2009**. <https://doi.org/10.1039/9781849730754>.
- (3) Kagan, H. B. Twenty-Five Years of Organic Chemistry with Diiodosamarium: An Overview. *Tetrahedron* **2003**, 59, 10351–10372.
- (4) Molander, G. A.; Harris, C. R. Sequencing Reactions with Samarium(II) Iodide. *Chem. Rev.* **1996**, 96, 307–338.
- (5) Ager, D. J.; Prakash, I.; Schaad, D. R. 1,2-Amino Alcohols and Their Heterocyclic Derivatives as Chiral Auxiliaries in Asymmetric Synthesis. *Chem. Rev.* **1996**, 96, 835–875.
- (6) Bergmeier, S. C. The Synthesis of Vicinal Amino Alcohols. *Tetrahedron* **2000**, 56, 2561–2576.
- (7) Burchak, O. N.; Py, S. Reductive Cross-Coupling Reactions (RCCR) between CN and CO for β -Amino Alcohol Synthesis. *Tetrahedron* **2009**, 65, 7333–7356.
- (8) (a) Maekawa, H.; Yamamoto, Y.; Shimada, H.; Yonemura, K.; Nishiguchi, I. Mg-Promoted Mixed Pinacol Coupling. *Tetrahedron Lett.* **2004**, 45, 3869–3872.
- (9) Roth, H. G.; Romero, N. A.; Nicewicz, D. A. Experimental and Calculated Electrochemical Potentials of Common Organic Molecules for Applications to Single-Electron Redox Chemistry. *Synlett* **2016**, 27, 714–723.

- (10) Uraguchi, D.; Kinoshita, N.; Kizu, T.; Ooi, T. Synergistic Catalysis of Ionic Brønsted Acid and Photosensitizer for a Redox Neutral Asymmetric α -Coupling of N-Arylaminoethanes with Aldimines. *J. Am. Chem. Soc.* **2015**, *137*, 13768–13771.
- (11) Zhu, X.-Q.; Liu, Q.-Y.; Chen, Q.; Mei, L.-R. Hydride, Hydrogen, Proton, and Electron Affinities of Imines and Their Reaction Intermediates in Acetonitrile and Construction of Thermodynamic Characteristic Graphs (TCGs) of Imines as a “Molecule ID Card.” *J. Org. Chem.* **2010**, *75*, 789–808.
- (12) All Potentials Reported vs SCE (Saturated Calomel Electrode).
- (13) Nakajima, M.; Fava, E.; Loescher, S.; Jiang, Z.; Rueping, M. Photoredox-Catalyzed Reductive Coupling of Aldehydes, Ketones, and Imines with Visible Light. *Angew. Chem. Int. Ed.* **2015**, *54*, 8828–8832.
- (14) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. Efficient Visible Light Photocatalysis of [2+2] Enone Cycloadditions. *J. Am. Chem. Soc.* **2008**, *130*, 12886–12887.
- (15) Du, J.; Espelt, L. R.; Guzei, I. A.; Yoon, T. P. Photocatalytic Reductive Cyclizations of Enones: Divergent Reactivity of Photogenerated Radical and Radical Anion Intermediates. *Chem. Sci.* **2011**, *2*, 2115.
- (16) Tarantino, K. T.; Liu, P.; Knowles, R. R. Catalytic Ketyl-Olefin Cyclizations Enabled by Proton-Coupled Electron Transfer. *J. Am. Chem. Soc.* **2013**, *135*, 10022–10025.
- (17) Rono, L. J.; Yayla, H. G.; Wang, D. Y.; Armstrong, M. F.; Knowles, R. R. Enantioselective Photoredox Catalysis Enabled by Proton-Coupled Electron Transfer: Development of an Asymmetric Aza-Pinacol Cyclization. *J. Am. Chem. Soc.* **2013**, *135*, 17735–17738.
- (18) Amador, A. G.; Sherbrook, E. M.; Yoon, T. P. Enantioselective Photocatalytic [3 + 2] Cycloadditions of Aryl Cyclopropyl Ketones. *J. Am. Chem. Soc.* **2016**, *138*, 4722–4725.
- (19) Lee, K. N.; Lei, Z.; Ngai, M.-Y. β -Selective Reductive Coupling of Alkenylpyridines with Aldehydes and Imines via Synergistic Lewis Acid/Photoredox Catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 5003–5006.
- (20) Seo, H.; Jamison, T. F. Catalytic Generation and Use of Ketyl Radical from Unactivated Aliphatic Carbonyl Compounds. *Org. Lett.* **2019**, *21*, 10159–10163.
- (21) Review: Xia, Q.; Dong, J.; Song, H.; Wang, Q. Visible-Light Photocatalysis of the Ketyl Radical Coupling Reaction. *Chem. Eur. J.* **2019**, *25*, 2949–2961.
- (22) Roskamp, E. J.; Pedersen, S. F. The First Practical Niobium(III) Reagent in Organic Synthesis. A Convenient Route to 2-Amino Alcohols via the Coupling of Imines with Aldehydes or Ketones Promoted by NbCl₅(DME). *J. Am. Chem. Soc.* **1987**, *109*, 6551–6553.
- (23) Machrouhi, F.; Namy, J.-L. Samarium Diiodide/Nickel Diiodide an Efficient System for Homo and Heterocoupling Reactions of Imines. *Tetrahedron Lett.* **1999**, *40*, 1315–1318.
- (24) Taniguchi, N.; Uemura, M. Asymmetric Synthesis of β -Amino Alcohols by Cross-Pinacol Coupling of Planar Chiral Ferrocene-carboxaldehydes with Imines [2]. *J. Am. Chem. Soc.* **2000**, *122*, 8301–8302.
- (25) Zhong, Y. W.; Dong, Y. Z.; Fang, K.; Izumi, K.; Xu, M. H.; Lin, G. Q. A Highly Efficient and Direct Approach for Synthesis of Enantiopure β -Amino Alcohols by Reductive Cross-Coupling of Chiral N-Tert-Butanesulfinyl Imines with Aldehydes. *J. Am. Chem. Soc.* **2005**, *127*, 11956–11957.
- (26) Lin, X.; Bentley, P. A.; Xie, H. Auxiliary Strategies for the Preparation of β -Amino Alcohols with Reductive Cross-Coupling and a Synthesis of (–)-Cytozone. *Tetrahedron Lett.* **2005**, *46*, 7849–7852.
- (27) Shi, S. L.; Wong, Z. L.; Buchwald, S. L. Copper-Catalysed Enantioselective Stereodivergent Synthesis of Amino Alcohols. *Nature* **2016**, *532*, 353–356.
- (28) Takeda, M.; Mitsui, A.; Nagao, K.; Ohmiya, H. Reductive Coupling between Aromatic Aldehydes and Ketones or Imines by Copper Catalysis. *J. Am. Chem. Soc.* **2019**, *141*, 3664–3669.
- (29) Yang, Z. P.; Fu, G. C. Convergent Catalytic Asymmetric Synthesis of Esters of Chiral Dialkyl Carbinols. *J. Am. Chem. Soc.* **2020**, *142*, 5870–5875.
- (30) Nguyen, K. D.; Park, B. Y.; Luong, T.; Sato, H.; Garza, V. J.; Krische, M. J. Metal-Catalyzed Reductive Coupling of Olefin-Derived Nucleophiles: Reinventing Carbonyl Addition. *Science* **2016**, *354*, aah5133.
- (31) Wang, H.; Lu, G.; Sormunen, G. J.; Malik, H. A.; Liu, P.; Montgomery, J. NHC Ligands Tailored for Simultaneous Regio- and Enantiocontrol in Nickel-Catalyzed Reductive Couplings. *J. Am. Chem. Soc.* **2017**, *139*, 9317–9324.
- (32) Schwarz, J. L.; Schäfers, F.; Tlahuext-Aca, A.; Lückemeier, L.; Glorius, F. Diastereoselective Allylation of Aldehydes by Dual Photoredox and Chromium Catalysis. *J. Am. Chem. Soc.* **2018**, *140*, 12705–12709.
- (33) Schwarz, J. L.; Kleinmans, R.; Paulisch, T. O.; Glorius, F. 1,2-Amino Alcohols via Cr/Photoredox Dual-Catalyzed Addition of α -Amino Carbanion Equivalents to Carbonyls. *J. Am. Chem. Soc.* **2020**, *142*, 2168–2174.
- (34) Tanabe, S.; Mitsunuma, H.; Kanai, M. Catalytic Allylation of Aldehydes Using Unactivated Alkenes. *J. Am. Chem. Soc.* **2020**, *142*, 12374–12381.
- (35) Hu, P.; Peters, B. K.; Malapit, C. A.; Vantourout, J. C.; Wang, P.; Li, J.; Mele, L.; Echeverria, P.-G.; Minter, S. D.; Baran, P. S. Electoreductive Olefin–Ketone Coupling. *J. Am. Chem. Soc.* **2020**, *142*, 20979–20986.
- (36) Wang, L.; Lear, J. M.; Rafferty, S. M.; Fosu, S. C.; Nagib, D. A. Ketyl Radical Reactivity via Atom Transfer Catalysis. *Science* **2018**, *362*, 225–229.
- (37) Slinker, J. D.; Gorodetsky, A. A.; Lowry, M. S.; Wang, J.; Parker, S.; Rohl, R.; Bernhard, S.; Malliaras, G. G. Efficient Yellow Electroluminescence from a Single Layer of a Cyclometalated Iridium Complex. *J. Am. Chem. Soc.* **2004**, *126*, 2763–2767.
- (38) Nagib, D. A.; Scott, M. E.; MacMillan, D. W. C. Enantioselective α -Trifluoromethylation of Aldehydes via Photoredox Organocatalysis. *J. Am. Chem. Soc.* **2009**, *131*, 10875–10877.
- (39) Koumura, K.; Satoh, K.; Kamigaito, M. Manganese-Based Controlled/Living Radical Polymerization of Vinyl Acetate, Methyl Acrylate, and Styrene: Highly Active, Versatile, and Photoresponsive Systems. *Macromolecules* **2008**, *41*, 7359–7367.
- (40) McMahon, C. M.; Renn, M. S.; Alexanian, E. J. Manganese-Catalyzed Carboacylations of Alkenes with Alkyl Iodides. *Org. Lett.* **2016**, *18*, 4148–4150.
- (41) Nuhant, P.; Oderinde, M. S.; Genovino, J.; Juneau, A.; Gagné, Y.; Allais, C.; Chinigo, G. M.; Choi, C.; Sach, N. W.; Bernier, L.; Fobian, Y. M.; Bundesmann, M. W.; Khunte, B.; Frenette, M.; Fadéy, O. O. Visible-Light-Initiated Manganese Catalysis for C–H Alkylation of Heteroarenes: Applications and Mechanistic Studies. *Angew. Chem. Int. Ed.* **2017**, *56*, 15309–15313.
- (42) Stateman, L. M.; Nakafuku, K. M.; Nagib, D. A. Remote C–H Functionalization via Selective Hydrogen Atom Transfer. *Synthesis* **2018**, *50*, 1569–1586.
- (43) Liu, R. Z.; Li, J.; Sun, J.; Liu, X. G.; Qu, S.; Li, P.; Zhang, B. Generation and Reactivity of Amidyl Radicals: Manganese-Mediated Atom-Transfer Reaction. *Angew. Chem. Int. Ed.* **2020**, *59*, 4428–4433.
- (44) Tsierekos, N. G. Investigation of the Electrochemical Reduction of Benzophenone in Aprotic Solvents Using the Method of Cyclic Voltammetry. *J. Solution Chem.* **2007**, *36*, 1301–1310.
- (45) α,β -unsaturated aldehydes do not work in this strategy (acrolein undergoes 1,4-addition by AcI; benzaldehyde-derived radicals are reduced before cross-coupling). However, these aldehydes have lower reduction potentials and may react via SET pathways instead (see Ref 21).

- (46) Friestad, G. K.; Qin, J. Intermolecular Alkyl Radical Addition to Chiral N-Acylhydrazones Mediated by Manganese Carbonyl. *J. Am. Chem. Soc.* **2001**, *123*, 9922–9923.
- (47) Friestad, G. K. Radical Additions to Chiral Hydrazones: Stereoselectivity and Functional Group Compatibility. In *Topics in Current Chemistry*; Springer, Berlin, Heidelberg, 2012; Vol. 320, pp 61–91.
- (48) Meyer, M. L.; DeAngelis, T. P.; Heineman, W. R. Mercury-Gold Minigrid Optically Transparent Thin-Layer Electrode. *Anal. Chem.* **1977**, *49*, 602–606.
- (49) Zhu, J.; Yuan, Y.; Wang, S.; Yao, Z.-J. Synthesis of 2,3-Dialkylated Tartaric Acid Esters via Visible Light Photoredox-Catalyzed Reductive Dimerization of α -Ketoesters. *ACS Omega* **2017**, *2*, 4665–4677.
- (50) Kanazawa, J.; Uchiyama, M. Recent Advances in the Synthetic Chemistry of Bicyclo[1.1.1]Pentane. *Synlett* **2019**, *30*, 1–11.
- (51) Nugent, J.; Arroniz, C.; Shire, B. R.; Sterling, A. J.; Pickford, H. D.; Wong, M. L. J.; Mansfield, S. J.; Caputo, D. F. J.; Owen, B.; Mousseau, J. J.; Duarte, F.; Anderson, E. A. A General Route to Bicyclo[1.1.1]Pentanes through Photoredox Catalysis. *ACS Catal.* **2019**, *9*, 9568–9574.
- (52) Wegman, R. W.; Olsen, R. J.; Gard, D. R.; Faulkner, L. R.; Brown, T. L. Flash Photolysis Study of the Metal-Metal Bond Homolysis in Dimanganese Decacarbonyl and Dirhenium Decacarbonyl. *J. Am. Chem. Soc.* **1981**, *103*, 6089–6092.
- (53) While an α -amino C-H is not required (Me₄piperidine: 52%), it improves efficiency (Me₃piperidine: 71%; Cy₂NMe: 90%), likely via termination by HAT to the N-radical. The resulting α -amino radical may then be reduced by Zn (regenerating Cy₂NMe). Although turnover by I-abstraction is also possible, Cy₂NH (by iminium formation/hydrolysis) is not observed.
- (54) Constantin, T.; Zanini, M.; Regni, A.; Sheikh, N. S.; Juliá, F.; Leonori, D. Aminoalkyl Radicals as Halogen-Atom Transfer Agents for Activation of Alkyl and Aryl Halides. *Science* **2020**, *367*, 1021–1026.

Table of Contents

