

Complex sexual-social environments produce high boldness and low aggression behavioral syndromes

Philip S. Queller^{1*}, Yasmin Shirali^{1,2}, Kelly J. Wallace^{1,3}, Ross S. DeAngelis¹, Vural Yurt^{1,4}, Luke P. Reding¹, Molly E. Cummings^{1*}

1: Department of Integrative Biology, The University of Texas at Austin, TX USA

2: Texas Department of State Health Services, Austin, TX USA

3: Department of Psychology, Emory University, Atlanta, GA USA

4: HAS University of Applied Sciences, 's-Hertogenbosch, the Netherlands

* Correspondence:

pqueller@utexas.edu

mcummings@austin.utexas.edu

Keywords: social complexity, behavioral development, animal personality, coping styles, mating systems

Abstract

Evidence of animal personality and behavioral syndromes is widespread across animals, yet the development of these traits remains poorly understood. Previous research has shown that exposure to predators, heterospecifics, and urbanized environments can influence personality and behavioral syndromes. Yet, to date, the influence of early social experiences with conspecifics on the development of adult behavioral traits is far less known. We use swordtail fish (*Xiphophorus nigrensis*), a species with three genetically-determined male mating strategies (courtship display, coercion, or mixed strategy) to assess how different early-life social experiences shape adult behavioral development. We raised female swordtails from birth to adulthood in density-controlled sexual-social treatments that varied in the presence of the type of male mating tactics (coercers only, displayers only, coercers and displayers, and mixed-strategists only). At adulthood, we tested females' boldness, shyness, aggression, sociality, and activity. We found that the number of different mating strategies females were raised with (social complexity) shaped behavioral development more than any individual mating strategy. Females reared in complex environments with two male mating tactics were bolder, less shy, and less aggressive than females reared with a single male mating tactic (either courtship only or coercion only). Complex sexual-social environments produced females with behavioral syndromes (correlations between aggression and activity, shyness and aggression, and social interaction and activity), whereas simple environments did not. Importantly, the characteristics of these socially-induced behavioral syndromes differ from those driven by predation, but converge on characteristics emerging from animals found in urban environments. Our findings suggest that complexity of the sexual-social environment shapes the development of personality and behavioral syndromes to facilitate social information gathering. Furthermore, our research highlights the previously overlooked influence of sexual selection as a significant contributing factor to diverse behavioral development.

Introduction

Early social experiences are known to have powerful effects on adult physiology (stress-response, Laubach et al. 2021; Murthy and Gould 2018) and behavior (e.g. mate preferences, ten Cate et al. 2006), and are expected to be important factors shaping a multitude of adult behaviors (Stamps and Groothuis 2010a; Stamps and Groothuis 2010b; Taborsky 2016; Cabrera et al. 2021). Researchers have manipulated early-life experience with variation in food availability (Groothuis et al. 2005; Edenbrow and Croft 2013), immune stress (Butler et al. 2012; DiRienzo et al. 2015), predation (Bell and Sih 2007; Edenbrow and Croft 2013; Niemelä et al. 2012; Castellano and Friard 2021), heterospecific exposure (Delclos et al. 2020), social isolation (Liedtke et al. 2015; Naguib et al. 2011), and physical enrichment (Liedtke et al. 2015; Xu et al. 2020) —all showing that these developmental experiences can shape traits like boldness, activity, aggression, and exploration. These traits are often studied in the context of animal personality (i.e. consistent inter-individual variation in behavior (Stamps and Groothuis 2010a; Stamps and Groothuis 2010b; Bergmüller and Taborsky 2010)) and behavioral syndromes (i.e. correlated behaviors across time and context (Sih et al. 2004a; Sih et al. 2004b)). Yet, there are few empirical studies that investigate how early-life social experiences affect the development of personality and behavioral syndromes in adults (Stamps and Groothuis 2010a; Stamps and Groothuis 2010b; Taborsky 2016; Cabrera et al. 2021).

Surprisingly, one of the untapped areas of research into personality and behavioral syndromes is how they are shaped by social interactions between the sexes. Sexual behavior is known to lead to selection for different behavioral traits. For instance, in mating systems that rely on male coercion, females tend to adopt behavioral strategies to evade sexual harassment (Dadda 2015; Pilastro et al. 2003; Darden and Croft 2008; Rowe et al. 1994). Meanwhile, in mating systems that rely on male display or courtship, females adopt cooperative behaviors (e.g. visiting leks (Uy et al. 2001; Rosenthal 2017; DuVal 2013); solicitation displays (Searcy and Capp 1997)). Differences in mating systems and specific male mating tactics are expected to lead to changes in non-reproductive female behaviors such as shoaling (Magurran and Macias Garcia 2000) as well as cognition (Cummings 2018). However, whether experience with different mating tactics influences the development of female personality or behavioral syndromes has not yet been investigated.

In addition, the complexity of the social environment itself may shape personality and behavioral syndromes. Social complexity has been hypothesized to drive the evolution of cognition (Dunbar 1998) and communication (Freeberg et al. 2012), and correlates with brain evolution in primates (Dunbar 2009), birds (Burish et al. 2004), and ants (Kamhi et al. 2016). How might complexity relate to personality and behavioral syndromes? It has been posited that in complex physical environments (Bengston et al. 2014) constraints on how individuals behave towards the complex environment have pleiotropic effects on other types of behaviors. Less is known about how social complexity affects these behavioral traits. At this point, we know that group-housed animals develop different behaviors as adults than those reared in isolation (Ballen et al. 2014; Liedtke et al. 2015). However, it may not simply be the presence or number of conspecifics that alters behavioral development, but the diversity of interactions individuals have with different types of conspecifics (Hobson et al. 2019).

Species with complex mating systems and alternative reproductive tactics allow us to begin to explore how specific social interactions alter behavioral development. Here, we manipulate lifetime social experiences (from birth to adulthood) of female El Abra swordtails (*Xiphophorus nigrensis*) to test how exposure to specific mating tactics (e.g. coercion, courtship display, or some combination of the two) as well as differing degrees of social complexity affect the development of personality and behavioral syndromes. Males of this species exhibit 1 of 3 genetically determined sexual phenotypes: large males are ornamented and court females with swimming displays, small males are drab and coerce females by chasing after them, and intermediate-sized males are moderately ornamented and exhibit a mixed strategy that includes both courtship and coercion (Ryan and Causey 1989).

We raised female swordtails from birth to adulthood (generally > 1 year) in one of five different social-rearing environments that maintained similar social densities but varied the combination of different male phenotypes in order to isolate the role of specific male tactics and the number of male tactics that females were exposed to in shaping female personality and behavioral syndromes. We ask whether females raised in different sexual-social environments (i) differ in their activity, sociability, boldness, shyness, and aggression; (ii) whether these behaviors are repeatable (form personality traits); and (iii) whether they correlate with one another (form a behavioral syndrome). Furthermore, our design allows us to compare the effects of (iv) specific sexual-social experiences (courtship only vs coercion only experiences), and (v) simple relative to complex mating environments (e.g. treatments with a single mating tactic vs treatments with two mating tactics) on the development of personality and behavioral syndromes.

Methods

Sexual-social rearing treatments

We raised female *Xiphophorus nigrensis* from birth to adulthood (1-1.5 years) in five socially controlled rearing treatments followed by behavioral testing in early adulthood. Experimental females were first introduced into experimental 30 gallon aquaria as fry (<10 mm) from broods produced in community tanks with individual broods split across all treatments to control for genetic effects. The fifteen experimental aquaria (5 treatments x 3 replicates each) were initially stocked with 30-40 fry of unknown sex along with 4 adult model females (identified with elastomer tags) and 4 adult males (see below for more details). Fish were fed Cargill and TetraMin flakes once daily and were supplemented with brine shrimp periodically. Tanks were enriched with plastic plants, flower pots, and gravel, and were illuminated with full spectrum aquarium lights. Experimental tanks were visually inspected 3 times a week to quantify juvenile densities and remove developing males. Developing males can be identified by a thickening of the anal fin as it develops into a gonopodium. Juvenile densities increased over time due to additional brood production by adult model females; and fluctuations in tank densities over time were due to changes in births, deaths, and removal of developing males. To minimize the variation in social densities across treatments and replicates over time, fry would be added to or removed from replicate tanks to maintain similar densities. Adult male stimuli were also moved between replicate tanks within a social treatment every 3 months to ensure developing females experienced multiple individual males of a given phenotype.

Our experimental females were reared in one of five social rearing treatments that varied by the type of male sexual behavior females experienced. The rearing treatments varied by the different types of stimulus males present: large courtship displaying males only (D), small coercive males only (C), mixed strategy intermediate-sized males only (M), combination of small coercive and large displaying males (C&D), and females only (F). To control for adult density, each tank had 8 stimulus adults including four male types described above (D, C or M males) and four adult model females. The C&D group contained two small, coercive males and two large, displaying males. The M, D, and C groups each had 4 males of their respective phenotype (see Figure 1a-d); and the F group had 8 adult model females. Adult stimulus (or model) females were selected from a community tank where they had experience with all types of *X. nigrensis* males and served as social learning models for developing females. We selected large adult females (usually > 30mm) to serve as models in our treatment tanks as avoidance of small males and preference for courting males increases with female size which is a proxy for age (Rios-Cardenas et al. 2007; Wong et al. 2011), with preference for associating with large courting males existing even in the absence of coercive males (Reding and Cummings 2017). Adult model females were dorsally tagged with white elastomer markings to differentiate them from the developing experimental females. It takes approximately one year for *X. nigrensis* females to reach sexual maturity, and sexually mature females are distinguishable from immature females by a melanized brood patch on the ventral area near the gonopore. We maintained experimental females in treatment tanks for 15-18 months in order to characterize developmental effects from the combined influences of social learning (from observing model females) and their own direct sexual-social experiences as adults. As mate choice copying has been documented in other poeciliids (Dugatkin and Godin 1993; Briggs et al. 1996), we expect social learning of mate preferences to strongly shape developing female preferences in adulthood (Davies et al. 2020; Dugatkin 1996). We conducted multiple rounds of this social rearing experiment from mid-2016 to 2019. Juvenile females that were introduced into experimental tanks in mid-2016 were tested as adults in 2018, and juvenile females introduced into social treatments in 2017-2018 were tested as adults in 2019.

To quantify the sexual-social interactions females from different social treatments experienced during their developmental period, we recorded weekly 15 min video recordings of all social rearing tanks between March 2016 to June 2017 for a total of 149 videos. From each video, we selected three 10 second video clips (from 1, 6, and 11 min. timestamps) that were then hand-scored by 7 undergraduate researchers naïve to the studies objectives (for a total of 411 scored video clips). Observers scored rates of coercive chases and courtship displays oriented toward adult females in each treatment group, along with average number of female foraging bouts and activity levels (time spent moving) across the rearing treatments. As expected in a system with genotypically-determined alternative male reproductive strategies, we were able to verify that females experienced different rates of coercion and courtship attempts across the different treatments (Figure 1e,f). However, we were not able to quantify adult female responses to these attempts (e.g. successful copulations) due to constraints of the video-recordings (see Supplemental videos 1-5 for representative videos of our social treatment tanks). Nonetheless, an extensive literature on *Xiphophorus* female preferences for large displaying males and aversion of small coercive males leads us to reasonably expect female responses in our experiment to be consistent with patterns of past research (Ryan and Causey 1989; Cummings and Mollaghan 2006; Rios-Cardenas et al 2007; Wong et al. 2011; Reding and Cummings 2017).

174 Behavioral Assessment

175 After experimental females reached adulthood, we evaluated their shyness, boldness, aggression,
176 sociability, and activity levels by testing them in 3 assays (scototaxis, mirror aggression, and
177 shoaling assays described below, see Table 1 for information on which behaviors came from
178 which assays). In total, we tested 60 experimental females with sample sizes from each treatment
179 including: D, display-only (n = 15); C, coercive-only (n = 9); C&D, coercive and displayers (n =
180 20); M, mixed-strategy (n = 10); and F female-only (n = 6). For information on sample sizes
181 from replicate tanks see Table S1 (Supplementary Information). We used the initial trials from
182 all experimental females to evaluate how treatment influenced specific behaviors. To assess
183 consistency of these behavioral traits (e.g. personality), we measured repeatability of these
184 behaviors from experimental fish tested twice on each of the three assays across a 17 day period
185 with an interval of 14 days between scototaxis trials, 13 days between mirror/aggression trials,
186 and 16 days between sociability trials. After removal from treatment tanks for testing, focal
187 females were individually-housed in adjacent 25 by 24 by 30 cm aquaria that allowed fish to see
188 and interact with the neighboring fish but remain separate for continuous identification of focal
189 fish. The visual social interaction was intended to reduce any stress associated with social
190 isolation. We ran repeat assays for 60 fish with initial sociability assay run on Day 1 while the
191 initial mirror and scototaxis trials run on Day 2 with a minimum resting interval of 2 hours
192 between their 1st (aggression) and 2nd (scototaxis) assay of the day (see Table S2 in
193 Supplementary Information for complete schedule). For 18 fish, we ran females through the
194 three assays only once (with 12 following the same initial assay schedule as above, and 6 fish
195 having longer intervals between trials: Day 1 sociability, Day 2 aggression; Day 4 scototaxis).
196 Due to technical problems with videos, along with some mortality of our experimental fish (n =
197 4), our final sample sizes for initial behavioral trials included sociability (n = 54), scototaxis (n
198 = 58), and aggression (n = 54); and final sample sizes for repeat trials included sociability (n =
199 41), scototaxis (n = 41), and aggression (n = 26). All videos were recorded using Nikon D3300
200 cameras with 18-55mm lenses.

201 Scototaxis

202 We measured shyness and boldness in females using a scototaxis assay (Maximino et al. 2010).
203 Our scototaxis tank involved a 25 x 50 cm aquarium with one half lined with white felt and the
204 other half lined with black felt (Figure 2a). The black side is a darker environment and can serve
205 as a refuge, while the white side is more exposed and activities in this region have been
206 associated with boldness (Maximino et al. 2010; Ramsey et al. 2014; Etheredge et al. 2018). We
207 evaluated ‘shyness’ as time spent in the edges of the black half (black thigmotaxis) as it
208 represents the most sheltered portion of the tank, and we characterized ‘boldness’ as the time
209 spent in the white center because it represented the most exposed zone (Figure 2a). To start the
210 trial, the tank is filled with 10 cm of water and the overhead fluorescent lights turned on. The fish
211 is then introduced to the center of the arena and constrained to a 4cm space by white barrier on
212 the white side and a black barrier on the black side. After a 5 minute habituation period, the
213 barriers are removed and the fish’s behavior was recorded for 10 minutes. Full water changes
214 were conducted between experimental fish to control for accumulation of cortisol in the water.

215 To quantify boldness and shyness in this assay, we divided the tank into 4 zones: black center,
216 white center, black thigmotaxis (2cm nearest the walls on the black side, see red dashed area in

Figure 2a), and white thigmotaxis (2cm nearest the walls on the white side, see blue dashed area in Figure 2a). We superimposed grids onto scototaxis videos using Microsoft Powerpoint and iMovie to create the 4 zones and time spent in each zone was scored using cowlog (“CowLog – Cross-Platform Application for Coding Behaviours from Video” 2016) by a trained undergraduate (RB). Video names were blinded such that the scorer was uninformed which treatment the subject female was raised within.

Mirror assay

We tested females’ aggression towards their mirror image following a similar protocol to other poeciliid studies (Franck et al. 1985; Wilson et al. 2011; Moretz and Morris 2003). Three sides of a 28 cm x 9 cm tank were lined with grey felt and a mirror was secured to the unfelted end (Figure 2b), with a video camera recordings collected directly above the tank. A focal female was introduced to a PVC tube on the opposite end as the mirror for a 5 minute habituation period. Afterwards, the PVC tube is removed and the fish allowed to explore the tank for 5 min with an opaque barrier covering the mirror end (movement control). The fish is then gently constrained back into the PVC tube for 1 min, during which the opaque barrier is removed and the mirror exposed. After 1 min, the camera is turned on and the trial recorded for 5 minutes. All water is changed between fish to control for accumulation of cortisol. A single trained undergraduate (NJ) scored all videos for aggressive behavior towards the mirror by counting the total number of bites focal fish directed toward their mirror image. Video names were blinded such that the scorer was uninformed which treatment the subject female was raised within.

Shoaling assay

We measured two behaviors from the shoaling assay: activity and sociability. Activity was measured in the shoaling assay because it was the least stressful assay and movements in less risky environments are considered to be better measures of general activity that are unconfounded by stress (Brown et al. 2007; Etheredge et al 2018). The assay was conducted in a rectangular glass aquarium measuring 120 cm in length and 30 cm in width and filled with water to a height of 18 cm (Figure 2c). The walls of the aquarium were lined with blue felt to reduce stress and maximize video tracking success. The felt also prevents the fish from becoming distracted by its reflection in the glass. A clear plexiglass partition was inserted 24 cm from the wall on each side of the tank (lengthwise), creating a 72 cm arena between the barriers. The tank was illuminated by four fluorescent aquarium bulbs covered with teal and dark blue filters. A shoal group, consisting of 5 females was placed on the side of the glass partition opposite of the focal fish and allowed 30 minutes to acclimate prior to testing. On the other side of the tank, an artificial plant was inserted on the same side as the focal fish, and the glass partition covered with brown felt opposite the plant. Between tests, the shoal side and plant side were switched to prevent side biases.

To begin each assay, the focal fish was placed in a 10cm diameter PVC habituation tube for 5-minutes, and then released and recorded for 10-minutes. Following this initial observation period, black felted plastic dividers were used to corral the focal fish back into the center of the arena. The fish was then held between the dividers (separated by 10cm) for an additional 5-minute habituation period. During this time the shoal of five females was moved to the opposite side of the aquarium and vice versa with the plant. The black felted dividers were then removed,

and video recordings conducted again for 10-minutes. Focal fish were then subsequently returned to their home aquaria.

We analyzed the video recordings using EthovisionXT15 (Noldus) behavioral tracking software to quantify time spent 2 cm adjacent to the plexiglass partition on the shoal side (see red dashed line in Figure 2c). We quantified female behaviors during the middle 5-minute portion for each 10 min observation period for a total of 10 min of behavioral tracking per subject (5 min from each observation period). Tracking began approximately 2 minutes after the start of each observation period to allow fish time to adjust to the new environment following removal of the habituation tube or black felted dividers. For each observation period, we collected total time spent in the social interaction zone as well as total distance moved as a measure of activity.

Statistics

All analyses were conducted in R version 4.1.2. To test for personality, we used Spearman's Rank correlation to test for repeatability between first and second measures of each behavior across the females that experienced repeat scototaxis, aggression and shoaling assays (Nakagawa and Schielzeth 2010). To test for effects of rearing environment on behavioral development, we used the initial measurements for all subject females' behavioral data as input into a Principal Components Analysis (PCA). To understand how rearing environment affected the mean values of each behavior, we also used ANOVA, using social treatment as the explanatory variable and behavioral response as the response variable. To evaluate whether mating system complexity drove differences in behavioral development, we combined measures from simple mating environments (treatments with only 1 male tactic: C and D) and combined measures from complex mating environments (treatments with 2 male tactics: C&D and M). We used generalized linear mixed effect models (GLMMs) with complexity as a fixed effect and treatment nested within complexity as a random effect to control for the non-independence between observations from treatment within each level of complexity. We also included replicate tank ID as a random effect to control for tank effects. Our complexity GLMM model did not converge for shyness and activity data, so for these analyses we used a non-parametric Kruskal-Wallis test. To evaluate behavioral syndromes (correlations between different behavior measures), we used bivariate correlation matrices as well as PCA coupled with ANOVA to test whether differences in loadings on each of the first two axes of variation differed across social complexity levels.

Results

Sexual-social rearing conditions

Video recordings collected during the developmental period from the experimental tanks revealed that females from different rearing environments experienced different levels of courtship (Figure 1e, ANOVA: $F = 61.2$, $p < 0.001$) and coercion (Figure 1f, ANOVA: $F = 10.5$, $p < 0.001$). Tukey post hoc analysis reveals that females raised in the D, M, and C&D social treatments experienced higher courtship than females raised in C and F treatments. Conversely, females raised in C, M, and C&D treatments experienced more coercion than females raised in D and F treatments, as expected. We also ran these analyses without the F treatment to make sure the absence of courtship and coercion in these females did not skew our results. Removal of this

treatment confirmed that the male-exposed rearing environments differed in the amount of courtship (ANOVA: $F = 28.8$, $p < 0.001$) and coercion (ANOVA: $F = 5.91$, $p < 0.001$). Tukey post hoc analysis without the F-females reveals females raised in D, C&D, and M treatments experienced more courtship than females reared in C treatments, with no difference in courtship rates between D, C&D, and M females. Conversely, females raised in C treatments experienced more coercion than females in M or D, but not C&D treatments (See Figure S1). Measurements of adult female models in each of the social treatments (while swimming freely within these experimental tanks) exhibited no significant difference in general activity (e.g. time spent moving, Figure 1g, $p = 0.2397$) or foraging rates (Figure 1h, $p = 0.1523$).

Density measurements ($n = 1184$ tri-weekly counts over 18 months across replicate tanks) of juveniles in the rearing tanks showed that females reared in different sexual-social treatments experienced similar levels of social densities across their 13-15 month developmental period (Figure 1i). To assess whether treatments differed in juvenile density, we used linear mixed effect models using lme4 package in R with treatment as a fixed effect. Across all treatments (C,D,C&D,M and F groups), there was a borderline statistically significant effect of treatment on juvenile densities ($X^2 = 9.03$, $df = 4$, $p = 0.060$ across all treatments), though this effect was mainly due to the female-only group (F, see Figure 1i, $X^2 = 0.39$, $df = 3$, $p = 0.94$ for only male-exposed treatments). Multiple regression analysis revealed no significant difference in slopes between treatments (p value of slope estimated differences for each treatment relative to the C social rearing group: $p = 0.727$ (C&D), $p = 0.601$ (D), $p = 0.668$ (M), and $p = 0.246$ (F)).

Personality

Following the developmental rearing period (13-15 months), experimental females were assessed in repeat trials of individual aggression, scototaxis and shoaling assays that revealed significant correlations across trials for some behavior metrics. We found that bites ($\rho = 0.37$, $p = 0.03$), shyness ($\rho = 0.40$, $p = 0.008$), boldness ($\rho = 0.45$, $p = 0.003$), and activity (distance traveled in shoaling assay, $\rho = 0.58$, $p < 0.0001$, Table 1) exhibited significant correlations across repeat trials. However, we found no statistically significant correlations between scored social interaction time between repeat shoaling assay trials (Table 1).

Sexual-social rearing environment shaping behavioral traits

Given the female-only social treatment had the smallest sample size ($n = 6$) and a marginally significant trend for lower densities (Figure 1i), we excluded the female-only group from subsequent analyses. Restricting our analyses to male-exposed rearing treatments (D, C, C&D, and M) enables us to compare how exposure to diverse mating types during development influences female behavioral development, while developing a balanced comparison between groups that involved a single mating tactic (C, D treatments) and groups that involved two mating tactics (C&D, M treatments). Comparisons including the F group can be found in Figure S2 in Supplementary Information.

Boldness & Shyness

The sexual-social rearing environment had significant effects on boldness (proportion of time spent in white center, Figure 2d, ANOVA: $F = 4.22$ $p = 0.01$) and shyness (proportion of time

spent in black thigmotaxis zone; Figure 2f, ANOVA: $F = 2.90$, $p = 0.04$). Tukey post hoc analysis shows that females raised in the C&D treatment were bolder than females in the C ($p = 0.04$) and D ($p = 0.02$) treatments. In addition, C&D females were less shy than D females ($p = 0.05$). When pooling treatment groups based on the number of mating tactics females were exposed to, we found that females reared with two mating tactics (complex environments: C&D and M treatments) were significantly bolder (Figure 2e: GLMM: $X^2 = 7.89$, $p = 0.005$) and less shy (Figure 2g: Kruskal-Wallis: $X^2 = 7.09$, $p < 0.008$) than females raised with only a single mating tactic (pooled C and D treatments).

Aggression

We found that sexual-social rearing experience affected intrasexual aggression (number of bites performed at mirror, Figure 2h, ANOVA: $F = 3.07$, $p = 0.04$). Tukey post hoc analysis shows females experienced with courtship displaying male types only (D treatment) showed significantly higher number of bites than C&D females ($p = 0.03$). Overall, females from multiple male tactic environments (complex environments) performed fewer bites than females from single male tactic environments (simple environments; Figure 2i, GLMM: $X^2 = 8.41$, $p = 0.004$).

Activity

There were no significant differences across treatments in the distance moved during the sociability assay (Figure 2j, ANOVA: $F = 0.55$, $p = 0.65$), nor were there differences between simple and complex groups (Figure 2k, Kruskal-Wallis: $X^2 = 1.02$, $p = 0.31$).

Sociability

There were no significant differences across treatments in the amount of social interaction (Figure 2l, ANOVA: $F = 1.38$, $p = 0.26$), nor were there differences between simple and complex groups (Figure 2m, GLMM: $X^2 = 0.05$, $p = 0.82$).

Sexual-social-rearing environment shaping behavioral syndromes

When we explored individual bi-variate correlations across the five behavioral metrics, we found several significant correlations affected by social rearing conditions (Figure 3). The one consistent trend shared across all treatment groups was a negative relationship between shyness and boldness (e.g. spearman correlation $\rho = -0.86$, $p = 0.003$ for C females (Figure 3b); $\rho = -0.7$, $p = 0.031$ for M females (Figure 3d); with similarly non-significant trends in D ($p = 0.08$, Figure 3a) and C&D ($p = 0.13$, Figure 3c) females). This correlation is expected because these traits represent opposing ends of the boldness/shyness spectrum, but we interpret this correlation with caution due to the non-independence of these measures that come from the same assay. However, we observed variation in correlations between behaviors measured in different contexts based on sexual-social rearing environment. Among females reared with displaying courters (D females), we found no significant correlations between behavioral traits across contexts. Females raised with coercive males only (C) had a positive trend between social interaction and aggression (Figure 3b, $\rho = 0.74$, $p = 0.058$). Meanwhile, females raised with both types of phenotypes (C&D) revealed a positive correlation between aggression and shyness (Figure 3c, $\rho = 0.48$, $p = 0.039$). And females raised with mixed strategy males (M females)

and both of these phenotypes (C&D females) exhibited a positive correlation between activity and social interaction (Figure 3d, $\rho = 0.78$, $p = 0.012$; Figure 3c, $\rho = 0.56$ $p = 0.038$).

More broadly, we found that females reared in environments with two mating tactics showed multiple significant correlations between behaviors whereas females reared in simple environments with a single mating tactic largely did not. Pooling females from simple sexual-social environments (C and D treatments) showed the expected negative correlation between boldness and shyness (Figure 3e, $\rho = -0.66$, $p = 0.001$). Meanwhile, pooling females from complex sexual-social environments with two mating tactics (C&D and M treatments) showed positive correlations between aggression and activity (Figure 3f, $\rho = 0.52$ $p = 0.019$), aggression and shyness ($\rho = 0.51$ $p = 0.009$), social interaction and activity ($\rho = 0.62$ $p = 0.001$), along with a near-significant negative correlation between boldness and aggression ($\rho = -0.38$, $p = 0.065$) and the expected negative correlation between shyness and boldness ($\rho = -0.47$, $p = 0.01$).

Principal Components Analysis (PCA)

Incorporating all behavior metrics as inputs into a PCA, we found that the number of mating tactics females were exposed to during development significantly predicted individual variation along the primary axis in this multidimensional reduction (Figure 4a). The first two principal components explain 66.79% of the multivariate variation (PC1 = 42.86 %, PC2 = 23.93%). PC1 primarily captures variation associated with boldness, shyness and aggression while PC2 captures variation associated with social interaction and activity (Figure 4b). Females from complex sexual-social environments (two mating tactics) had significantly higher PC1 scores than females from simple, single mating tactic environments (Figure 4c, ANOVA: $F = 14.32$, $p = 0.0006$). There was also a non-significant trend for females from complex environments to score higher on PC2 than those from simple environments (Figure 4d, ANOVA: $F = 2.689$, $p = 0.0835$).

Discussion

In this study, we successfully created discrete sexual-social experiences that females encountered for the entirety of their developmental period (from parturition to adulthood) by manipulating female exposure to different combinations of alternative male reproductive phenotypes. Importantly, females raised in these different social groups experienced similar densities (Figure 1i), enabling us to examine the effect of male mating tactic and sexual-social complexity on female development independent of social density effects.

Personality

We found that aggression, activity, boldness, and shyness were stable over repeat measurements (Table 1). These findings demonstrate that *X. nigrensis* possess similarly stable personality traits as have been reported in other poeciliids. For instance, aggression (Wilson et al. 2011), boldness (Boulton et al. 2014), activity (Cote et al. 2010; Biro and Adriaenssens 2013; Blake and Gabor 2014), and exploration (Blake and Gabor 2014; Heinen-Kay et al. 2016) have been established as personality traits in other swordtail and mosquitofish species. Variation in some of these personality traits such as boldness and aggression has previously been shown to correlate with

421 predation (Blake and Gabor 2014; Heinen-Kay et al. 2016; Castellano and Friard 2021;
422 Huntingford 1976; Dingemanse et al. 2007; Bell 2004), but this is among the first to explore how
423 specific social experiences shape the development of these traits.

424 Sexual-social rearing environment shaping personality

425 We show that social experience during ontogeny leads to long lasting and repeatable changes in
426 personality during adulthood, adding to a growing body of literature on the ontogeny of
427 personality (Stamps and Groothuis 2010a; Stamps and Groothuis 2010b; Hedrick 2017; Cabrera
428 et al. 2021). Importantly, we quantified behavior in the rearing environment (Figure 1) as well as
429 during adulthood allowing a direct comparison of what type or combination of early-life
430 experiences affect adult personality (Taborsky 2016). Why would developmental exposure to
431 both courtship and coercion lead to bolder and less aggressive females? One possible
432 interpretation is that responding to multiple male sexual behaviors is more cognitively
433 challenging than responding to males that predictably employ the same tactic. In a complex
434 sexual-social environment, an increase in boldness and a reduction in aggression may increase
435 females' exposure to important social information and thereby allow females to appropriately
436 respond. Boldness may be required to explore the sexual-social environment, while too much
437 intrasexual aggression may detract attention away from male behavior and thereby decrease
438 information gathering related to male conspecifics. Our results are consistent with findings in
439 other taxa where social isolation leads to lower exploration in individuals relative to individuals
440 raised with others (Liedtke et al. 2015). Indeed, boldness has been shown to positively correlate
441 with a number of cognitive traits such as spatial learning in lizards (Carazo et al. 2014) and
442 learning speed in chickadees (Guillette et al. 2009). Access to information likely explains
443 differences in aggression in dominance hierarchies (Hobson 2020), and our results suggest that
444 aggression may be modulated to increase social information gathering when social contexts are
445 complex. In this study, experience with multiple male mating behaviors leads to bolder and less
446 aggressive females, which may lead to increased information acquisition.

447 Sexual-social rearing environment shaping behavioral syndromes

448 Why might behavioral syndromes emerge from a complex, but not simple, mating environment
449 during development? According to the social niche specialization hypotheses, behavioral
450 syndromes may emerge from animals optimizing their behavior in a socioecological context
451 through state-behavior feedbacks (Bergmüller and Taborsky 2010; Sih et al. 2015; Sih 2011).
452 Perhaps sexual-social complexity that reflects the natural social conditions of *X. nigrensis*
453 requires some constraint on the development of female behavior. If, for example, an increase in
454 female boldness in response to social complexity aids in information processing and improved
455 social decision making, behaviors such as aggression might be constrained to lower levels to
456 facilitate this relationship. In support of this view, social complexity is associated with increased
457 social competence across taxa (Taborsky and Oliveira 2012). Female behavior from a simple
458 social environment may be less constrained, with reduced pressure to form behavioral
459 syndromes. In addition, research on pace-of-life syndromes suggests ecological conditions drive
460 differences in resource allocation to survival and reproduction that may also affect personality
461 and behavioral syndromes (Polverino et al. 2018). In this experiment, social complexity may
462 have shaped female behavior through its effect on pace-of-life syndromes. Physical complexity
463 in the rearing environment has been shown to affect behavioral syndromes (Liedtke et al. 2015;

Xu et al. 2020), and here we show that sexual-social complexity is also an important factor in shaping behavioral syndromes.

Previous research in sticklebacks (Dingemanse et al. 2007; Bell and Sih 2007) has shown that exposure to predators leads to the formation of behavioral syndromes, whereas predator-naïve sticklebacks lack behavioral syndromes. Here we find that exposure to multiple mating tactics leads to the formation of female behavioral syndromes, which are absent in females from environments with lower mating tactic complexity. Our results suggest that sexual selection pressures might be just as powerful as natural selection pressures at forming behavioral syndromes. Interestingly, the correlations we report are in opposite directions as what is commonly reported in studies where predation pressure is implicated in behavioral syndrome formation (Huntingford 1976; Bell 2004; Dingemanse et al. 2007). Whereas many predation studies report a positive correlation between boldness, aggression and activity in focal prey taxa, we found negative correlations between activity and aggression and between aggression and boldness in females raised with multiple male mating tactics. This unusual behavioral syndrome in females reared in complex mating environments may reflect a constraint females face in the more complex social environment that link together sets of behaviors possibly involved in social information gathering. Interestingly, this same pattern of low aggression linked with high boldness is found among animals experiencing urbanization (eg. northern cardinals (Huang et al. 2020), lizards (Batabyal and Thaker 2019) and several other taxa (Sadoul et al. 2021)). Researchers have suggested that boldness may emerge in the more complex urban environments as a result of habituation to humans (Samia et al. 2015); and that lower aggression may facilitate greater cognitive flexibility required in these more complex urban environments (Sadoul et al. 2021). The convergence of the same behavioral syndrome traits emerging from socially complex conspecific environments as those that emerge from exposure to humans may suggest that complexity in a number of different forms can lead to selection for a specific suite of behavioral traits.

An alternative, non-mutually exclusive, explanation for how the social environment affects the development of behavioral syndromes is that diverse social experiences represent different social stressors that produce different coping styles. Coping styles refers to variation in how individuals respond behaviorally and physiologically to stressful events with ‘proactive’ animals tending to be more bold and aggressive, and ‘reactive’ animals tending to be the opposite (Koolhaas et al. 1999; Bensch et al. 2017). The environment is expected to shape behavioral syndromes through shaping the stress axes in early development (Killen et al. 2013; Careau et al. 2014; Guenther et al. 2014; Farine et al. 2015; Sih 2011). For example, in the cavy (*Cavia aperea*), variation in birth season (associated with different stressors) leads to behavioral syndrome differences in adults (Guenther et al. 2014). In crickets, agonistic experience during development shapes individual differences in approach-avoidance behaviors, aggressiveness, and motility, which the authors argue supports a proactive and reactive coping style syndromes in that species (Balsam and Stevenson 2021). The social environment can represent different levels of stress, and certainly variation in the type of mating interactions is likely to be a large source of stress variation. Further research may be able to determine if these differences in behavioral syndromes correlate with differences in social information processing and/or differences in stress physiology (coping styles).

Conclusions

In summary, we found that sexual-social experiences during development affect the emergence of behavioral syndromes. We were able to manipulate the type of social interactions females experienced across development while keeping social group size the same—allowing us to differentiate the type of social interactions from the quantity. Our results have broad implications on the power of sexual selection on the evolution and maintenance of behavioral syndromes and personality. If female experience with single or multiple male mating tactics in the laboratory has the power to shape behavioral syndromes, then we should also see this pattern in the wild. Comparative work should examine whether species with single mating tactics repeatedly diverge in personality and behavioral syndromes from taxa with multiple alternative male mating tactics; and whether species with multiple mating tactics exhibit stronger evidence for behavioral syndromes (and whether a boldness-low aggression syndrome is the most dominant). Moreover, future studies should consider multiple sources of environmental variation on behavioral syndrome development including both sexual selection (mating system, social complexity) and natural selection (predation, habitat complexity) pressures.

Acknowledgments

We are grateful to the Brackenridge Field Laboratory at UT Austin which provided a semi-natural outdoor setting for our seed populations of *X. nigrensis*. We heartily thank all of the undergraduate students that helped maintain social environments, conduct density counts, and score data including Rob Bailey, Niveya James, Kathryn Kaihlanen, Camille Akins, Amogh Kashyap, Adam Redmer, Jessika McFarland, Caleb Fleischer; and thank the National Science Foundation for funding this research (Grant NSF-IOS-1911826 to MEC).

References

- Ballen, C., Shine, R., & Olsson, M. (2014). Effects of early social isolation on the behaviour and performance of juvenile lizards, *Chamaeleo calyptratus*. *Animal Behaviour*, 88, 1-6.
- Balsam, J. S., & Stevenson, P. A. (2021). Agonistic experience during development establishes inter-individual differences in approach-avoidance behaviour of crickets. *Scientific Reports*, 11(1), 1-13.
- Batabyal, A., & Thaker, M. (2019). Social coping styles of lizards are reactive and not proactive in urban areas. *General and comparative endocrinology*, 270, 67-74.
- Bell, A. M., & Sih, A. (2007). Exposure to predation generates personality in threespined sticklebacks (*Gasterosteus aculeatus*). *Ecology letters*, 10(9), 828-834.
- Bell, A. M. (2005). Differences between individuals and populations of threespined stickleback. *J. Evol. Biol*, 18, 464-473.
- Bengston, S. E., Pruitt, J. N., & Riechert, S. E. (2014). Differences in environmental enrichment generate contrasting behavioural syndromes in a basal spider lineage. *Animal Behaviour*, 93, 105-110.

- Bensky, M. K., Paitz, R., Pereira, L., & Bell, A. M. (2017). Testing the predictions of coping styles theory in threespined sticklebacks. *Behavioural Processes*, 136, 1-10.
- Bergmüller, R., & Taborsky, M. (2010). Animal personality due to social niche specialisation. *Trends in Ecology & Evolution*, 25(9), 504-511.
- Biro, P. A., & Adriaenssens, B. (2013). Predictability as a personality trait: consistent differences in intraindividual behavioral variation. *The American Naturalist*, 182(5), 621-629.
- Blake, C. A., & Gabor, C. R. (2014). Effect of prey personality depends on predator species. *Behavioral Ecology*, 25(4), 871-877.
- Boulton, K., Grimmer, A. J., Rosenthal, G. G., Walling, C. A., & Wilson, A. J. (2014). How stable are personalities? A multivariate view of behavioural variation over long and short timescales in the sheephead swordtail, *Xiphophorus birchmanni*. *Behavioral Ecology and Sociobiology*, 68(5), 791-803.
- Briggs, S. E., Godin, J. G. J., & Dugatkin, L. A. (1996). Mate-choice copying under predation risk in the Trinigadian guppy (*Poecilia reticulata*). *Behavioral Ecology*, 7(2), 151-157.
- Burish, M. J., Kueh, H. Y., & Wang, S. S. H. (2004). Brain architecture and social complexity in modern and ancient birds. *Brain, Behavior and Evolution*, 63(2), 107-124.
- Butler, M. W., Toomey, M. B., McGraw, K. J., & Rowe, M. (2012). Ontogenetic immune challenges shape adult personality in mallard ducks. *Proceedings of the Royal Society B: Biological Sciences*, 279(1727), 326-333.
- Cabrera, D., Nilsson, J. R., & Griffen, B. D. (2021). The development of animal personality across ontogeny: a cross-species review. *Animal Behaviour*, 173, 137-144.
- Carazo, P., Noble, D. W., Chandrasoma, D., & Whiting, M. J. (2014). Sex and boldness explain individual differences in spatial learning in a lizard. *Proceedings of the Royal Society B: Biological Sciences*, 281(1782), 20133275.
- Careau, V., Buttemer, W. A., & Buchanan, K. L. (2014). Early-developmental stress, repeatability, and canalization in a suite of physiological and behavioral traits in female zebra finches. *Integrative and Comparative Biology*, 54(4), 539-554.

- Castellano, S., & Friard, O. (2021). Environmental effects on the ontogenesis of tadpole personality. *Animal Behaviour*, 175, 153-161.
- ten Cate, C., Verzijden, M. N., & Etman, E. (2006). Sexual imprinting can induce sexual preferences for exaggerated parental traits. *Current Biology*, 16(11), 1128-1132.
- Cote, J., Fogarty, S., Weinersmith, K., Brodin, T., & Sih, A. (2010). Personality traits and dispersal tendency in the invasive mosquitofish (*Gambusia affinis*). *Proceedings of the Royal Society B: Biological Sciences*, 277(1687), 1571-1579.
- Pastell, M. (2016). CowLog-cross-platform application for coding behaviours from video.
- Cummings, M., & Mollaghan, D. (2006). Repeatability and consistency of female preference behaviours in a northern swordtail, *Xiphophorus nigrensis*. *Animal Behaviour*, 72(1), 217-224.
- Cummings, M. E. (2018). Sexual conflict and sexually dimorphic cognition—reviewing their relationship in poeciliid fishes. *Behavioral Ecology and Sociobiology*, 72(4), 1-13.
- Dadda, M. (2015). Female social response to male sexual harassment in poeciliid fish: a comparison of six species. *Frontiers in Psychology*, 6, 1453.
- Darden, S. K., & Croft, D. P. (2008). Male harassment drives females to alter habitat use and leads to segregation of the sexes. *Biology Letters*, 4(5), 449-451.
- Davies, A. D., Lewis, Z., & Dougherty, L. R. (2020). A meta-analysis of factors influencing the strength of mate-choice copying in animals. *Behavioral Ecology*, 31(6), 1279-1290.
- Delclos, P. J., Forero, S. A., & Rosenthal, G. G. (2020). Divergent neurogenomic responses shape social learning of both personality and mate preference. *Journal of Experimental Biology*, 223(6), jeb220707.
- Dingemanse, N. J., Wright, J., Kazem, A. J., Thomas, D. K., Hickling, R., & Dawnay, N. (2007). Behavioural syndromes differ predictably between 12 populations of three-spined stickleback. *Journal of Animal Ecology*, 1128-1138.
- DiRienzo, N., Niemelä, P. T., Skog, A., Vainikka, A., & Kortet, R. (2015). Juvenile pathogen exposure affects the presence of personality in adult field crickets. *Frontiers in Ecology and Evolution*, 3, 36.

- Dugatkin, L. A., & Godin, J. G. J. (1993). Female mate copying in the guppy (*Poecilia reticulata*): age-dependent effects. *Behavioral Ecology*, 4(4), 289-292.
- Dugatkin, L. A. (1996). Copying and mate choice. *Social learning in animals: The roots of culture*, 85-105.
- Dunbar, R. I. (1998). The social brain hypothesis. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews*, 6(5), 178-190.
- DuVal, E. H. (2013). Female mate fidelity in a lek mating system and its implications for the evolution of cooperative lekking behavior. *The American Naturalist*, 181(2), 213-222.
- Edenbrow, M., & Croft, D. P. (2013). Environmental and genetic effects shape the development of personality traits in the mangrove killifish *Kryptolebias marmoratus*. *Oikos*, 122(5), 667-681.
- Etheredge, R. I., Avenas, C., Armstrong, M. J., & Cummings, M. E. (2018). Sex-specific cognitive-behavioural profiles emerging from individual variation in numerosity discrimination in *Gambusia affinis*. *Animal Cognition*, 21(1), 37-53.
- Farine, D. R., Spencer, K. A., & Boogert, N. J. (2015). Early-life stress triggers juvenile zebra finches to switch social learning strategies. *Current Biology*, 25(16), 2184-2188.
- Franck, D., Hannes, R. P., Lanffermann, H., & Ribowski, A. (1985). Effects of social isolation on aggressiveness in fish with special reference to the swordtail (*Xiphophorus helleri*). *Behavioural processes*, 10(4), 415-427.
- Overgaard, M., & Sandberg, K. (2012). Kinds of access: different methods for report reveal different kinds of metacognitive access. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1594), 1287-1296.
- Groothuis, T. G., Koolhaas, J. M., Drent, P. J., & Carere, C. (2005). Epigenetic effects on personality traits: early food provisioning and sibling competition. *Behaviour*, 142(9-10), 1329-1355.
- Günther, A., Finkemeier, M. A., & Trillmich, F. (2014). The ontogeny of personality in the wild guinea pig. *Animal Behaviour*, 90, 131-139.
- Guenther, A., Brust, V., Dersen, M., & Trillmich, F. (2014). Learning and personality types are related in caviar (*Cavia aperea*). *Journal of Comparative Psychology*, 128(1), 74.

- Guillette, L. M., Reddon, A. R., Hurd, P. L., & Sturdy, C. B. (2009). Exploration of a novel space is associated with individual differences in learning speed in black-capped chickadees, *Poecile atricapillus*. *Behavioural Processes*, 82(3), 265-270.
- Hedrick, A. V. (2017). The development of animal personality. *Frontiers in Ecology and Evolution*, 5, 14.
- Heinen-Kay, J. L., Schmidt, D. A., Stafford, A. T., Costa, M. T., Peterson, M. N., Kern, E. M., & Langerhans, R. B. (2016). Predicting multifarious behavioural divergence in the wild. *Animal behaviour*, 121, 3-10.
- Hobson, E. A. (2020). Differences in social information are critical to understanding aggressive behavior in animal dominance hierarchies. *Current Opinion in Psychology*, 33, 209-215.
- Hobson, E. A., Ferdinand, V., Kolchinsky, A., & Garland, J. (2019). Rethinking animal social complexity measures with the help of complex systems concepts. *Animal Behaviour*, 155, 287-296.
- Huang, P., St. Mary, C. M., & Kimball, R. T. (2020). Habitat urbanization and stress response are primary predictors of personality variation in northern cardinals (*Cardinalis cardinalis*). *Journal of Urban Ecology*, 6(1), juaa015.
- Huntingford, F. A. (1976). The relationship between anti-predator behaviour and aggression among conspecifics in the three-spined stickleback, *Gasterosteus aculeatus*. *Animal Behaviour*, 24(2), 245-260.
- Kamhi, J. F., Gronenberg, W., Robson, S. K., & Traniello, J. F. (2016). Social complexity influences brain investment and neural operation costs in ants. *Proceedings of the Royal Society B: Biological Sciences*, 283(1841), 20161949.
- Killen, S. S., Marras, S., Metcalfe, N. B., McKenzie, D. J., & Domenici, P. (2013). Environmental stressors alter relationships between physiology and behaviour. *Trends in Ecology & Evolution*, 28(11), 651-658.
- Koolhaas, J. M., Korte, S. M., De Boer, S. F., Van Der Vegt, B. J., Van Reenen, C. G., Hopster, H., De Jong, I.C., Ruis, M.A.W., & Blokhuis, H. J. (1999). Coping styles in animals: current status in behavior and stress-physiology. *Neuroscience & Biobehavioral Reviews*, 23(7), 925-935.

- Laubach, Z. M., Greenberg, J. R., Turner, J. W., Montgomery, T. M., Pioon, M. O., Sawdy, M. A., ... & Perng, W. (2021). Early-life social experience affects offspring DNA methylation and later life stress phenotype. *Nature Communications*, 12(1), 1-15.
- Liedtke, J., Redekop, D., Schneider, J. M., & Schuett, W. (2015). Early environmental conditions shape personality types in a jumping spider. *Frontiers in Ecology and Evolution*, 3, 134.
- Loffredo, C. A., & Borgia, G. (1986). Male courtship vocalizations as cues for mate choice in the satin bowerbird (*Ptilonorhynchus violaceus*). *The Auk*, 103(1), 189-195.
- Magurran, A. E., & Garcia, C. M. (2000). Sex differences in behaviour as an indirect consequence of mating system. *Journal of Fish Biology*, 57(4), 839-857.
- Maximino, C., Marques de Brito, T., Dias, C. A. G. D. M., Gouveia, A., & Morato, S. (2010). Scototaxis as anxiety-like behavior in fish. *Nature Protocols*, 5(2), 209-216.
- Moretz, J. A., & Morris, M. R. (2003). Evolutionarily labile responses to a signal of aggressive intent. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1530), 2271-2277.
- Murthy, S., & Gould, E. (2018). Early life stress in rodents: animal models of illness or resilience? *Frontiers in Behavioral Neuroscience*, 12, 157.
- Naguib, M., Flörcke, C., & van Oers, K. (2011). Effects of social conditions during early development on stress response and personality traits in great tits (*Parus major*). *Developmental Psychobiology*, 53(6), 592-600.
- Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists. *Biological Reviews*, 85(4), 935-956.
- Niemelä, P. T., DiRienzo, N., & Hedrick, A. V. (2012). Predator-induced changes in the boldness of naïve field crickets, *Gryllus integer*, depends on behavioural type. *Animal Behaviour*, 84(1), 129-135.
- Pilastro, A., Benetton, S., & Bisazza, A. (2003). Female aggregation and male competition reduce costs of sexual harassment in the mosquitofish *Gambusia holbrooki*. *Animal Behaviour*, 65(6), 1161-1167.
- Polverino, G., Santostefano, F., Díaz-Gil, C., & Mehner, T. (2018). Ecological conditions drive pace-of-life syndromes by shaping relationships between life history, physiology and behaviour in two populations of Eastern mosquitofish. *Scientific Reports*, 8(1), 1-10.

- Ramsey, M. E., Vu, W., & Cummings, M. E. (2014). Testing synaptic plasticity in dynamic mate choice decisions: N-methyl d-aspartate receptor blockade disrupts female preference. *Proceedings of the Royal Society B: Biological Sciences*, 281(1785), 20140047.
- Reding, L., & Cummings, M. E. (2017). Context-dependent preferences vary by multicomponent signals in a swordtail. *Animal Behaviour*, 129, 237-247.
- Rios-Cardenas, O., Tudor, M. S., & Morris, M. R. (2007). Female preference variation has implications for the maintenance of an alternative mating strategy in a swordtail fish. *Animal Behaviour*, 74(3), 633-640.
- Rosenthal, G. G. (2017). *Mate choice: the evolution of sexual decision making from microbes to humans*. Princeton University Press.
- Rowe, L., Arnqvist, G., Sih, A., & Krupa, J. J. (1994). Sexual conflict and the evolutionary ecology of mating patterns: water striders as a model system. *Trends in Ecology & Evolution*, 9(8), 289-293.
- Ryan, M. J., & Causey, B. A. (1989). "Alternative" mating behavior in the swordtails *Xiphophorus nigrensis* and *Xiphophorus pygmaeus* (Pisces: Poeciliidae). *Behavioral Ecology and Sociobiology*, 24(6), 341-348.
- Sadoul, B., Blumstein, D. T., Alfonso, S., & Geffroy, B. (2021). Human protection drives the emergence of a new coping style in animals. *PLoS Biology*, 19(4), e3001186.
- Samia, D. S., Nakagawa, S., Nomura, F., Rangel, T. F., & Blumstein, D. T. (2015). Increased tolerance to humans among disturbed wildlife. *Nature Communications*, 6(1), 1-8.
- Searcy, W. A., & Capp, M. S. (1997). Estradiol dosage and the solicitation display assay in red-winged blackbirds. *The Condor*, 99(3), 826-828.
- Sih, A. (2011). Effects of early stress on behavioral syndromes: an integrated adaptive perspective. *Neuroscience & Biobehavioral Reviews*, 35(7), 1452-1465.
- Sih, A., Bell, A., & Johnson, J. C. (2004a). Behavioral syndromes: an ecological and evolutionary overview. *Trends in ecology & evolution*, 19(7), 372-378.
- Sih, A., Bell, A. M., Johnson, J. C., & Ziemba, R. E. (2004b). Behavioral syndromes: an integrative overview. *The Quarterly Review of Biology*, 79(3), 241-277.

- Sih, A., Mathot, K. J., Moirón, M., Montiglio, P. O., Wolf, M., & Dingemanse, N. J. (2015).
Animal personality and state–behaviour feedbacks: a review and guide for empiricists.
Trends in ecology & evolution, 30(1), 50-60.
- Stamps, J. A., & Groothuis, T. G. (2010a). Developmental perspectives on personality:
implications for ecological and evolutionary studies of individual differences.
Philosophical Transactions of the Royal Society B: Biological Sciences, 365(1560),
4029-4041.
- Stamps, J., & Groothuis, T. G. (2010b). The development of animal personality: relevance,
concepts and perspectives. *Biological Reviews*, 85(2), 301-325.
- Taborsky, B. (2016). Opening the black box of developmental experiments: behavioural
mechanisms underlying long-term effects of early social experience. *Ethology*, 122(4),
267-283.
- Taborsky, B., & Oliveira, R. F. (2012). Social competence: an evolutionary approach. *Trends in
ecology & evolution*, 27(12), 679-688.
- Wilson, A. J., de Boer, M., Arnott, G., & Grimmer, A. (2011). Integrating personality research
and animal contest theory: aggressiveness in the green swordtail *Xiphophorus helleri*.
PLoS One, 6(11), e28024.
- Wong, R. Y., So, P., & Cummings, M. E. (2011). How female size and male displays influence
mate preference in a swordtail. *Animal Behaviour*, 82(4), 691-697.
- Xu, W., Yao, Q., Zhang, W., Zhang, F., Li, H., Xu, R., Li, C., & Zhang, B. (2021).
Environmental complexity during early life shapes average behavior in adulthood.
Behavioral Ecology, 32(1), 105-113.

Figures/Tables:

Assay (sample size)	Metric	rho	<i>p</i> -value
Mirror (<i>N</i> =26)	Aggression (bites at mirror)	0.369	0.0348
Scototaxis (<i>N</i> =41)	Shyness (time in black thigmotaxis)	0.408	0.0086
Scototaxis (<i>N</i> =41)	Boldness (time in white center)	0.449	0.0032
Shoaling (<i>N</i> =41)	Activity (distance moved)	0.579	<0.001
Shoaling (<i>N</i> =41)	Social interaction (time < 2 cm from shoal)	0.111	0.518

Table 1: Spearman's rank correlation between the two repeat measurements for each behavior from each assay.




Figure 1: Social-rearing environment conditions

Schematic of different social rearing environments: D, displayers only (a); C, coercers only (b); C&D, coercers and displayers together (c); M, mixed strategy males (d); and F, female only environments (not shown). Females from different rearing environments experienced different rates of male courtship (e) and coercion (f) while exhibiting no differences in their activity (time spent moving) (g) or foraging (h) rates. Box and whisker plots show median (center line) and interquartile range. Letters demonstrate similarity from Tukey post hoc analysis. Data were from 411 video clips (10 secs each) collected from the treatment tanks during the developmental rearing period (> 1 year) prior to behavioral testing of the experimental females at adulthood (number of videos per treatment shown above box and whisker plots). Weekly density measurements across replicate tanks for each treatment (i) reveal that females in the F (female only treatment) showed a near-significant difference in juvenile densities over time relative to the male-exposed treatments ($p = 0.06$). Excluding the female-only treatment group revealed that females from male-exposure treatments (D, C, C&D, and M) experienced similar density levels over time ($p = 0.94$). Bold colored lines represent the average densities across replicate tanks within a social treatment: (D) blue, (C) red, (C&D) green, (M) yellow, (F) grey, while faint lines show density counts from individual replicate tanks (represented as lighter colors than the treatment average line).




Figure 2: Behavioral effects of social rearing environment on adult behavior

At adulthood, females were tested in (a) scototaxis, (b) aggression, and (c) shoaling assays. We measured (d) boldness (white center, proportion of time in white center—denoted by dashed blue line), and (f) shyness (black thigmotaxis, proportion of time on black side within 2cm of wall—denoted by dashed red line) in the scototaxis assay. We assessed (h) aggression as the number of bites at the mirror; and activity (j) as the total movement (pixels traversed on video screen):

arbitrary units) during the shoaling assay; and social interaction (I) as the proportion of time individuals spent in the social interaction zone (2cm from barrier, see red line in (c)). We used ANOVA to test for differences across groups for each behavioral metric with associated p value shown for each test, and significant differences between pair wise comparisons using Tukey-post hoc test denoted with different letters above each treatment group. We further examined behavioral developmental differences by social complexity (e,g,i,j,m) by using GLMM (and Kruskal-Wallis for non-normal data) between ‘simple’ (pooled rearing environments with a single male mating tactic (C, D) shown in purple) and ‘complex’ (pooled rearing environments with two male mating tactics (C&D, M) shown in orange).




Figure 3: Heatmap of behavioral correlations by treatment and complexity
Spearman's rank correlation of behaviors by treatment (a-d) and sexual-social complexity (e,f) with negative correlations shown in cooler colors and positive correlations in warmer colors. Significant spearman's rank correlations ($p < 0.05$) are denoted with bold font. A significant or

near-significant trend for a negative relationship was observed between shyness and boldness which is expected given the non-independence of these two behavioral characteristics (measured within the same assay). Evidence for behavioral syndromes (correlations of behaviors measured between contexts, *sensu* Sih et al. 2004b) were found only in complex sexual-social rearing environments (f, aggression and shyness ($p = 0.009$), aggression and activity ($p = 0.019$), activity and social interaction ($p = 0.001$), and near-significant boldness and aggression ($p = 0.065$)) but not in simple sexual-social rearing environments (e).




Figure 4: Multivariate analysis of behavioral traits by social complexity

Principal component analysis (PCA) including all 5 behavioral metrics revealed females reared in simple (purple) and complex (orange) mating environments show behavioral differences in multidimensional space. (a) The first two principal components explain 66.79% of the multivariate variation, with PC1 explaining 42.9% of the variance. Points are colored by treatment and ellipses represent 95% confidence intervals. (b) Vector-plot of PC loadings reveal that boldness, shyness and aggression load most prominently on PC1, whereas activity and shoaling tendencies load more strongly onto PC2. (c) ANOVA reveals females from complex environments (C&D and M groups) score higher on PC1 than females from simple environments (C and D groups, $p = 0.0006$); with (d) non-significant trend for complex-reared females to have higher PC2 scores than simple-reared females ($p = 0.08$).