

BOUNDS ON ORTHOGONAL POLYNOMIALS AND SEPARATION OF THEIR ZEROS

¹ELI LEVIN AND ²D. S. LUBINSKY

ABSTRACT. Let $\{p_n\}$ denote the orthonormal polynomials associated with a measure μ with compact support on the real line. Let μ be regular in the sense of Stahl, Totik, and Ullmann, and I be a subinterval of the support in which μ is absolutely continuous, while μ' is positive and continuous there. We show that boundedness of the $\{p_n\}$ in that subinterval is closely related to the spacing of zeros of p_n and p_{n-1} in that interval. One ingredient is proving that "local limits" imply universality limits.

Research supported by NSF grant DMS1800251

1. RESULTS

Let μ be a finite positive Borel measure with compact support, which we denote by $\text{supp}[\mu]$. Then we may define orthonormal polynomials

$$p_n(x) = \gamma_n x^n + \dots, \gamma_n > 0,$$

$n = 0, 1, 2, \dots$ satisfying the orthonormality conditions

$$\int p_n p_m d\mu = \delta_{mn}.$$

The zeros of p_n are real and simple. We list them in decreasing order:

$$x_{1n} > x_{2n} > \dots > x_{n-1,n} > x_{nn}.$$

They interlace the zeros y_{jn} of p'_n :

$$p'_n(y_{jn}) = 0 \text{ and } y_{jn} \in (x_{j+1,n}, x_{jn}), 1 \leq j \leq n-1.$$

It is a classic result that the zeros of p_n and p_{n-1} also interlace. The three term recurrence relation has the form

$$(x - b_n) p_n(x) = a_{n+1} p_{n+1}(x) + a_n p_{n-1}(x),$$

where for $n \geq 1$,

$$a_n = \frac{\gamma_{n-1}}{\gamma_n} = \int x p_{n-1}(x) p_n(x) d\mu(x); \quad b_n = \int x p_n^2(x) d\mu(x).$$

Uniform boundedness of orthonormal polynomials is a long studied topic. For example, given an interval I , one asks whether

$$\sup_{n \geq 1} \|p_n\|_{L^\infty(I)} < \infty.$$

There is an extensive literature on this fundamental question - see for example [1], [2], [3], [4], [12]. In this paper, we establish a connection to the distance between zeros of p_n and p_{n-1} .

The results require more terminology: we let $\text{dist}(a, \mathbb{Z})$ denote the distance from a real number a to the integers. We say that μ is *regular* (in the sense of Stahl, Totik, and Ullmann) if for every sequence of non-zero polynomials $\{P_n\}$ with degree P_n at most n ,

$$\limsup_{n \rightarrow \infty} \left(\frac{|P_n(x)|}{(\int |P_n|^2 d\mu)^{1/2}} \right)^{1/n} \leq 1$$

for quasi-every $x \in \text{supp}[\mu]$ (that is except in a set of logarithmic capacity 0). If the support consists of finitely many intervals, and $\mu' > 0$ a.e. in each subinterval, then μ is regular, though much less is required [15]. An equivalent formulation involves the leading coefficients $\{\gamma_n\}$ of the orthonormal polynomials for μ :

$$\lim_{n \rightarrow \infty} \gamma_n^{1/n} = \frac{1}{\text{cap}(\text{supp}[\mu])},$$

where cap denotes logarithmic capacity.

Recall that the equilibrium measure for the compact set $\text{supp}[\mu]$ is the probability measure that minimizes the energy integral

$$\int \int \log \frac{1}{|x - y|} d\nu(x) d\nu(y)$$

amongst all probability measures ν supported on $\text{supp}[\mu]$. If I is an interval contained in $\text{supp}[\mu]$, then the equilibrium measure is absolutely continuous in I , and moreover its density, which we denote throughout by ω , is positive and continuous in the interior I° of I [13, p.216, Thm. IV.2.5]. Given sequences $\{x_n\}, \{y_n\}$ of non-0 real numbers, we write

$$x_n \sim y_n$$

if there exists $C > 1$ such that for $n \geq 1$,

$$C^{-1} \leq x_n/y_n < C.$$

Similar notation is used for functions and sequences of functions.

Our main result is

Theorem 1.1

Let μ be a regular measure on $\mathbb{R}$ with compact support. Let I be a closed subinterval of the support and assume that in some open interval containing I , μ is absolutely continuous, while μ' is positive and continuous. Let ω be the density of the equilibrium measure for the support of μ . Let $A > 0$. The following are equivalent:

(a) There exists $C > 0$ such that for $n \geq 1$ and $x_{jn} \in I$,

$$(1.1) \quad \text{dist}(n\omega(x_{jn})(x_{jn} - x_{j,n-1}), \mathbb{Z}) \geq C.$$

(b) There exists $C > 0$ such that for $n \geq 1$ and $y_{jn} \in I$,

$$(1.2) \quad \text{dist}(n\omega(y_{jn})(y_{jn} - y_{j,n-1}), \mathbb{Z}) \geq C.$$

(c) Uniformly for $n \geq 1$ and $x \in I$,

$$(1.3) \quad \|p_{n-1}\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \|p_n\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \sim 1.$$

(d) There exists $C > 0$ such that for $n \geq 1$ and $x \in I$,

$$(1.4) \quad \|p_{n-1}\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \|p_n\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \leq C.$$

Moreover, under any of (a), (b), (c), (d), we have

$$(1.5) \quad \sup_{n \geq 1} \sup_{x \in I} |x - b_n|^{1/2} p_n(x) < \infty.$$

Remarks

(a) The main idea behind the proof is that universality limits and "local" limits give

$$|p_{n-1}(y_{j,n-1}) p_n(y_{jn})| |\sin[\pi n \omega(y_{jn})(y_{jn} - y_{j,n-1})] + o(1)| \sim 1,$$

uniformly in j, n , while p_n has a local extremum at y_{jn} .

(b) We could replace $x_{j,n-1} - x_{jn}$ in (1.1) by $x_{j,n-1} - x_{j,n+k}$, for any fixed integer k (see Lemma 4.1).

(b) Under additional assumptions, involving the spacing of zeros of p_n and p_{n-2} , we can remove the factor $|x - b_n|^{1/2}$ in (1.5):

Theorem 1.2

Let μ be a regular measure on $\mathbb{R}$ with compact support. Let I be a closed subinterval of the support and assume that in some open interval containing I , μ is absolutely continuous, while μ' is positive and continuous. Let ω be the density of the equilibrium measure for the support of μ . Let $A > 0$. Assume that (1.1) holds in I . The following are equivalent:

(a) There exist $C_1 > 0$ such that for $n \geq 1$ and $x_{jn} \in I$,

$$(1.6) \quad |n(x_{jn} - x_{j-1,n-2})| \geq C_1 |x_{jn} - b_{n-1}|.$$

(b) *Uniformly for $x \in I$ and $n \geq 1$,*

$$(1.7) \quad \|p_n\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \sim 1.$$

(c)

$$(1.8) \quad \sup_{n \geq 1} \|p_n\|_{L_\infty(I)} < \infty.$$

Remark

We note that because of the interlacing, both x_{jn} and $x_{j-1, n-2}$ belong to the interval $(x_{j, n-1}, x_{j-1, n-1})$.

Two important ingredients in our proofs are universality and local limits. The so-called universality limit involves the reproducing kernel

$$(1.9) \quad K_n(x, y) = \sum_{k=0}^{n-1} p_k(x) p_k(y) = \frac{\gamma_{n-1}}{\gamma_n} \frac{p_n(x) p_{n-1}(y) - p_{n-1}(x) p_n(y)}{x - y}.$$

For x in the interior of $\text{supp}[\mu]$ (the "bulk" of the support), at least when $\mu'(x)$ is finite and positive, the universality limit typically takes the form [6], [8], [14], [17]

$$(1.10) \quad \lim_{n \rightarrow \infty} \frac{K_n\left(x + \frac{a}{\mu'(x)K_n(x, x)}, x + \frac{b}{\mu'(x)K_n(x, x)}\right)}{K_n(x, x)} = \mathbb{S}(a - b),$$

uniformly for a, b in compact subsets of $\mathbb{C}$. Here $\mathbb{S}$ is the sinc kernel,

$$\mathbb{S}(a) = \frac{\sin \pi a}{\pi a}.$$

Universality limits holds far more generally than pointwise asymptotics for orthonormal polynomials, that at one stage were used to prove them. In a series of recent papers [7], [9], [10], [11], it was shown that one can go in the other direction, namely from universality limits, to "local ratio limits" for orthogonal polynomials.

Under fairly general conditions on μ , the *Christoffel function* $K_n(x, x)$ admits the asymptotic [16]

$$\lim_{n \rightarrow \infty} \frac{1}{n} K_n(x, x) \mu'(x) = \omega(x)$$

for x in the interior of the support of μ . This allows us to reformulate the universality limit (1.10) as

$$(1.11) \quad \lim_{n \rightarrow \infty} \frac{K_n\left(x + \frac{a}{n\omega(x)}, x + \frac{b}{n\omega(x)}\right) \mu'(x)}{n\omega(x)} = \mathbb{S}(a - b),$$

uniformly for a, b in compact subsets of $\mathbb{C}$.

Using this universality limit, we proved in [10]:

Theorem A

Assume that μ is a regular measure with compact support. Let I be a closed subinterval of the support in which μ is absolutely continuous, and μ' is positive and continuous. Let J be a compact subset of the interior I° of I . Then

$$(1.12) \quad \lim_{n \rightarrow \infty} \frac{p_n \left(y_{jn} + \frac{z}{n\omega(y_{jn})} \right)}{p_n(y_{jn})} = \cos \pi z$$

uniformly for $y_{jn} \in J$ and z in compact subsets of $\mathbb{C}$.

A secondary goal of this paper is to prove a converse of Theorem A, namely to show that local limits such as (1.12) imply a universality limit like (1.11). For measures on the unit circle this was undertaken in [11] - however the results necessarily take a quite different form.

Theorem 1.3

Let μ be a measure with compact support. Assume that we are given a bounded sequence of real numbers $\{\xi_n\}$ such that

$$(1.13) \quad \sup_{n \geq 1} n |\xi_n - \xi_{n-1}| < \infty,$$

and a sequence $\{\tau_n\}$ of positive numbers with $\tau_n \sim 1$ such that

$$(1.14) \quad \lim_{n \rightarrow \infty} \frac{\tau_n}{\tau_{n-1}} = 1$$

and uniformly for z in compact subsets of $\mathbb{C}$,

$$(1.15) \quad \lim_{n \rightarrow \infty} \frac{p_n \left(\xi_n + \frac{\tau_n}{n} z \right)}{p_n(\xi_n)} = \cos \pi z.$$

Let $A > 0$. Then uniformly for a, b in compact subsets of $\mathbb{C}$, and x_n such that

$$(1.16) \quad |x_n - \xi_n| \leq \frac{A}{n}$$

we have

$$(1.17) \quad \frac{K_n \left(x_n + \frac{\tau_n}{n} a, x_n + \frac{\tau_n}{n} b \right)}{K_n(x_n, x_n)} = \mathbb{S}(a - b) + o \left(\frac{\frac{\gamma_{n-1}}{\gamma_n} n |p_{n-1}(\xi_{n-1}) p_n(\xi_n)|}{K_n(x_n, x_n)} \right).$$

Moreover,

$$(1.18) \quad \frac{K_n \left(x_n + \frac{\tau_n}{n} a, x_n + \frac{\tau_n}{n} b \right)}{K_n(x_n, x_n)} = \mathbb{S}(a - b) + o(1),$$

provided either

$$(1.19) \quad \liminf_{n \rightarrow \infty} \operatorname{dist} \left(\frac{n}{\tau_n} (\xi_n - \xi_{n-1}), \mathbb{Z} \right) > 0$$

or

$$(1.20) \quad \sup_{n \geq 1} \frac{\frac{\gamma_{n-1}}{\gamma_n} n |p_{n-1}(\xi_{n-1}) p_n(\xi_n)|}{K_n(x_n, x_n)} < \infty.$$

We prove Theorem 1.3 in the next section and Theorem 1.1 in Section 3. Theorem 1.2 is proved in Section 4. In the sequel $C, C_1, C_2, \dots$ denote constants independent of n, x, θ . The same symbol does not necessarily denote the same constant in different occurrences.

2. PROOF OF THEOREM 1.3

Throughout this section, we assume the hypotheses of Theorem 1.3. Write for $n \geq 1$ and $m = n - 1, n$,

$$(2.1) \quad x_n = \xi_m + \Delta_{n,m} \frac{\tau_m}{m}$$

and

$$(2.2) \quad \chi_n = \left(\frac{\tau_n}{n} \right) / \left(\frac{\tau_{n-1}}{n-1} \right).$$

Recall from (1.14) that $\chi_n \rightarrow 1$ as $n \rightarrow \infty$. Note too that in view of (1.13), (1.14), (1.16), $\{\Delta_{n,n}\}$ and $\{\Delta_{n,n-1}\}$ are bounded sequences. We start with:

Lemma 2.1

(a) *Uniformly for z in compact subsets of $\mathbb{C}$,*

$$(2.3) \quad \lim_{n \rightarrow \infty} \frac{\tau_n p'_n \left(\xi_n + \frac{\tau_n}{n} z \right)}{n p_n(\xi_n)} = -\pi \sin \pi z.$$

(b) *Uniformly for a, b in compact subsets of $\mathbb{C}$,*

$$\begin{aligned} & \left(p_n \left(x_n + \frac{\tau_n}{n} a \right) - p_n \left(x_n + \frac{\tau_n}{n} b \right) \right) / p_n(\xi_n) \\ &= -\pi \int_b^a \sin \pi (\Delta_{n,n} + t) dt + o(|a - b|). \end{aligned}$$

(c) Moreover,

$$\begin{aligned} & \left(p_{n-1} \left(x_n + \frac{\tau_n}{n} a \right) - p_{n-1} \left(x_n + \frac{\tau_n}{n} b \right) \right) / p_{n-1} (\xi_{n-1}) \\ &= -\pi \int_b^a \sin \pi (\Delta_{n,n-1} + t) dt + o(|a - b|). \end{aligned}$$

Proof

(a) As the asymptotic (1.15) holds uniformly for z in compact subsets of the plane, we can differentiate it to obtain (2.3).

(b) Now

$$\begin{aligned} & \left(p_n \left(x_n + \frac{\tau_n}{n} a \right) - p_n \left(x_n + \frac{\tau_n}{n} b \right) \right) / p_n (\xi_n) \\ &= \int_b^a p'_n \left(x_n + \frac{\tau_n}{n} t \right) \frac{\tau_n}{n} dt / p_n (\xi_n). \end{aligned}$$

Note that this is meaningful even for complex a, b , with the integral being taken over the directed line segment from b to a . Using (2.1) and (2.3), we continue this as

$$\begin{aligned} & \int_b^a \frac{p'_n \left(\xi_n + \frac{\tau_n}{n} (\Delta_{n,n} + t) \right) \frac{\tau_n}{n}}{p_n (\xi_n)} dt \\ &= \int_b^a (-\pi \sin \pi (\Delta_{n,n} + t) + o(1)) dt \\ &= -\pi \int_b^a \sin \pi (\Delta_{n,n} + t) dt + o(|a - b|). \end{aligned}$$

(c) Using (2.2),

$$\begin{aligned} & \left(p_{n-1} \left(x_n + \frac{\tau_n}{n} a \right) - p_{n-1} \left(x_n + \frac{\tau_n}{n} b \right) \right) / p_{n-1} (\xi_{n-1}) \\ &= \int_b^a p'_{n-1} \left(x_n + \frac{\tau_n}{n} t \right) \frac{\tau_n}{n} dt / p_{n-1} (\xi_{n-1}) \\ &= \int_b^a \frac{p'_{n-1} \left(\xi_{n-1} + \frac{\tau_{n-1}}{n-1} (\Delta_{n,n-1} + \chi_n t) \right) \frac{\tau_{n-1}}{n-1}}{p_{n-1} (\xi_{n-1})} \chi_n dt \\ &= \int_b^a (-\pi \sin (\pi (\Delta_{n,n-1} + \chi_n t)) + o(1)) dt \\ &= -\pi \int_b^a \sin \pi (\Delta_{n,n-1} + t) dt + o(|a - b|). \end{aligned}$$

■

Proof of Theorem 1.3

We apply (1.15) and (b), (c) of Lemma 2.1. Now if $a \neq b$,

$$\begin{aligned}
& \frac{\tau_n}{np_{n-1}(\xi_{n-1})p_n(\xi_n)} K_n \left(x_n + \frac{\tau_n}{n}a, x_n + \frac{\tau_n}{n}b \right) \\
&= \frac{\gamma_{n-1} \left[p_n \left(x_n + \frac{\tau_n}{n}a \right) - p_n \left(x_n + \frac{\tau_n}{n}b \right) \right] p_{n-1} \left(x_n + \frac{\tau_n}{n}b \right)}{\gamma_n (a-b)p_n(\xi_n)p_{n-1}(\xi_{n-1})} \\
& \quad + \frac{\gamma_{n-1} p_n \left(x_n + \frac{\tau_n}{n}b \right) \left[p_{n-1} \left(x_n + \frac{\tau_n}{n}b \right) - p_{n-1} \left(x_n + \frac{\tau_n}{n}a \right) \right]}{\gamma_n (a-b)p_n(\xi_n)p_{n-1}(\xi_{n-1})} \\
&= \frac{\gamma_{n-1}}{\gamma_n} \left[\frac{-\pi}{a-b} \int_b^a \sin \pi (\Delta_{n,n} + t) dt + o(1) \right] [\cos \pi (\Delta_{n,n-1} + b\chi_n) + o(1)] \\
& \quad + \frac{\gamma_{n-1}}{\gamma_n} [\cos \pi (\Delta_{n,n} + b) + o(1)] \left[\frac{\pi}{a-b} \int_b^a \sin \pi (\Delta_{n,n-1} + t) dt + o(1) \right]
\end{aligned}$$

by (1.15) and (b), (c) of Lemma 2.1. Note that because of the uniformity of the limits, this holds in a confluent form even if $a = b$. We continue this, using $\chi_n = 1 + o(1)$, as

$$\begin{aligned}
&= \frac{\gamma_{n-1}}{\gamma_n} \frac{\pi}{a-b} \int_a^b [\sin \pi (\Delta_{n,n} + t) \cos \pi (\Delta_{n,n-1} + b) - \cos \pi (\Delta_{n,n} + b) \sin \pi (\Delta_{n,n-1} + t)] dt \\
& \quad + o \left(\frac{\gamma_{n-1}}{\gamma_n} \right).
\end{aligned}$$

(2.4)

Next, we expand the integrand using double angle formulae, in a straightforward but tedious fashion:

$$\begin{aligned}
& \sin \pi (\Delta_{n,n} + t) \cos \pi (\Delta_{n,n-1} + b) - \cos \pi (\Delta_{n,n} + b) \sin \pi (\Delta_{n,n-1} + t) \\
&= [\sin \pi \Delta_{n,n} \cos \pi t + \cos \pi \Delta_{n,n} \sin \pi t] [\cos \pi \Delta_{n,n-1} \cos \pi b - \sin \pi \Delta_{n,n-1} \sin \pi b] \\
& \quad - [\cos \pi \Delta_{n,n} \cos \pi b - \sin \pi \Delta_{n,n} \sin \pi b] [\sin \pi \Delta_{n,n-1} \cos \pi t + \cos \pi \Delta_{n,n-1} \sin \pi t] \\
&= \cos \pi t \cos \pi b \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}) + \sin \pi t \sin \pi b \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}) \\
&= \cos (\pi (t - b)) \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}).
\end{aligned}$$

We can then continue (2.4) as

$$\begin{aligned}
& \frac{\gamma_{n-1}}{\gamma_n} \frac{\pi}{a-b} \int_a^b [\cos (\pi (t - b)) \sin \pi (\Delta_{n,n} - \Delta_{n,n-1})] dt + o \left(\frac{\gamma_{n-1}}{\gamma_n} \right) \\
&= \frac{\gamma_{n-1}}{\gamma_n} \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}) \frac{1}{a-b} (-\sin \pi (a - b)) + o \left(\frac{\gamma_{n-1}}{\gamma_n} \right) \\
&= -\pi \frac{\gamma_{n-1}}{\gamma_n} \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}) \mathbb{S}(a - b) + o \left(\frac{\gamma_{n-1}}{\gamma_n} \right).
\end{aligned}$$

In summary, uniformly for a, b in compact subsets of the plane,

$$(2.5) \quad \frac{\tau_n}{np_{n-1}(\xi_{n-1})p_n(\xi_n)} K_n\left(x_n + \frac{\tau_n}{n}a, x_n + \frac{\tau_n}{n}b\right) \\ = -\pi \frac{\gamma_{n-1}}{\gamma_n} \sin \pi (\Delta_{n,n} - \Delta_{n,n-1}) \mathbb{S}(a-b) + o\left(\frac{\gamma_{n-1}}{\gamma_n}\right).$$

Next observe from (2.1), (1.13), and (1.14), that

$$x_n = \xi_n + \Delta_{n,n} \frac{\tau_n}{n} = \xi_{n-1} + \Delta_{n,n-1} \frac{\tau_n}{n} + o\left(\frac{1}{n}\right) \\ \Rightarrow \frac{\tau_n}{n} [\Delta_{n,n} - \Delta_{n,n-1}] = \xi_{n-1} - \xi_n + o\left(\frac{1}{n}\right).$$

As τ_n is bounded below, this allows us to reformulate (2.5) as

$$\frac{\tau_n}{n} K_n\left(x_n + \frac{\tau_n}{n}a, x_n + \frac{\tau_n}{n}b\right) \\ = -\pi \frac{\gamma_{n-1}}{\gamma_n} p_{n-1}(\xi_{n-1}) p_n(\xi_n) \left\{ \sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right] \mathbb{S}(a-b) + o(1) \right\}. \quad (2.6)$$

In particular, setting $a = b = 0$,

$$\frac{\tau_n}{n} K_n(x_n, x_n) \\ = -\pi \frac{\gamma_{n-1}}{\gamma_n} p_{n-1}(\xi_{n-1}) p_n(\xi_n) \left\{ \sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right] + o(1) \right\}, \quad (2.7)$$

so that (2.6) can be recast as

$$\frac{\tau_n}{n} K_n\left(x_n + \frac{\tau_n}{n}a, x_n + \frac{\tau_n}{n}b\right) \\ = \frac{\tau_n}{n} K_n(x_n, x_n) \mathbb{S}(a-b) + o\left(\frac{\gamma_{n-1}}{\gamma_n} |p_{n-1}(\xi_{n-1}) p_n(\xi_n)|\right),$$

giving (1.17). If (1.19) holds, then $\sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right]$ is bounded away from 0, so we can reformulate (2.6) as

$$\frac{\tau_n}{n} K_n\left(x_n + \frac{\tau_n}{n}a, x_n + \frac{\tau_n}{n}b\right) \\ = -\pi \frac{\gamma_{n-1}}{\gamma_n} p_{n-1}(\xi_{n-1}) p_n(\xi_n) \sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right] \{ \mathbb{S}(a-b) + o(1) \}$$

and (2.7) as

$$\begin{aligned} & \frac{\tau_n}{n} K_n(x_n, x_n) \\ &= -\pi \frac{\gamma_{n-1}}{\gamma_n} p_{n-1}(\xi_{n-1}) p_n(\xi_n) \sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right] \{1 + o(1)\}. \end{aligned}$$

Together these give (1.18). Finally if (1.20) holds, then we see from (2.6) that necessarily $\sin \left[\pi \frac{n}{\tau_n} (\xi_{n-1} - \xi_n) \right]$ is bounded away from 0 and again (1.18) follows. ■

3. PROOF OF THEOREM 1.1

Recall that y_{jn} is the zero of p'_n in $(x_{j+1,n}, x_{jn})$. We begin with:

Lemma 3.1

Let μ be a regular measure on $\mathbb{R}$ with compact support. Let I be a closed subinterval of the support and assume that in some open interval containing I , μ is absolutely continuous, while μ' is positive and continuous.

(a) Uniformly for $y_{jn} \in I$,

$$(3.1) \quad \lim_{n \rightarrow \infty} n(x_{jn} - y_{jn}) \omega(x_{jn}) = \frac{1}{2} = \lim_{n \rightarrow \infty} n(y_{jn} - x_{j+1,n}) \omega(x_{jn});$$

$$(3.2) \quad \lim_{n \rightarrow \infty} n(x_{jn} - x_{j+1,n}) \omega(x_{jn}) = 1;$$

$$(3.3) \quad \lim_{n \rightarrow \infty} n(y_{jn} - y_{j+1,n}) \omega(x_{jn}) = 1.$$

(b) Uniformly for $y_{jn} \in I$,

$$(3.4) \quad \frac{\gamma_{n-1}}{\gamma_n} |p_{n-1}(y_{j,n-1}) p_n(y_{jn})| |\sin [\pi n \omega(y_{jn}) (y_{j,n-1} - y_{jn})] + o(1)| \sim 1.$$

(c) Fix $A > 0$. Uniformly for $n \geq 1$ and $x \in I$,

$$(3.5) \quad \|p_n\|_{L_\infty[x - \frac{A}{n}, x + \frac{A}{n}]} \sim |p_n(y_{jn})|,$$

where $y_{jn} \in [x - \frac{A}{n}, x + \frac{A}{n}]$ or is the closest zero of p'_n to this interval.

Proof

(a) First note that uniformly for $y_{jn} \in I$ and z in compact subsets of $\mathbb{C}$,

$$(3.6) \quad \lim_{n \rightarrow \infty} \frac{p_n\left(y_{jn} + \frac{z}{n\omega(y_{jn})}\right)}{p_n(y_{jn})} = \cos \pi z.$$

This was proved in [10] and is Theorem A above. Next, Theorem 2.1 in [17] shows that (3.2) holds uniformly for $x_{jn} \in I$. In particular $x_{jn} - x_{j+1,n} = O\left(\frac{1}{n}\right)$ uniformly for $x_{jn} \in I$. Write

$$x_{jn} = y_{jn} + \frac{z_n}{n\omega(y_{jn})},$$

so that $z_n > 0$ and $z_n = O(1)$. From (3.6), we have

$$0 = \frac{p_n(x_{jn})}{p_n(y_{jn})} = \cos \pi z_n + o(1)$$

so necessarily for some non-negative integer j_n ,

$$z_{jn} = j_n + \frac{1}{2} + o(1).$$

If $j_n \geq 1$ for infinitely many n , then Hurwitz' Theorem shows that there would be other zeros of p_n between x_{jn} and y_{jn} , which contradicts that $y_{jn} \in (x_{j+1,n}, x_{jn})$. So $j_n = 0$ for n large enough, which gives the first limit in (3.1). Note too that $\omega(x_{jn})/\omega(y_{jn}) = 1 + o(1)$ by continuity of ω . The second is similar. Both (3.2) and (3.3) follow from (3.1), though as noted, (3.2) appears in [17].

(b) Because of (3.6), we can apply Theorem 1.3 and results from its proof. In that theorem, we set $x_n = y_{jn}$, $\tau_n = \frac{1}{\omega(y_{jn})}$; $\xi_n = y_{jn}$; so that $\xi_{n-1} = y_{j,n-1}$. Note that (1.13), (1.14), (1.16) are satisfied because of the spacing estimates in Lemma 3.1, and the continuity of ω . From (2.7),

$$\begin{aligned} & \frac{1}{n\omega(y_{jn})} K_n(y_{jn}, y_{jn}) \\ &= -\pi \frac{\gamma_{n-1}}{\gamma_n} p_{n-1}(y_{j,n-1}) p_n(y_{jn}) \{ \sin [\pi n \omega(y_{jn}) (y_{j,n-1} - y_{jn})] + o(1) \}. \end{aligned}$$

(3.7)

Next, Theorem 2.2 in [17] establishes that uniformly for $t \in I$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} K_n(t, t) \mu'(t) = \omega(t).$$

Since ω is positive and continuous in I as is μ' , we then obtain (3.4) from (3.7).

(c) This follows directly from the limit in (3.6) and the fact that $|p_n(y_{jn})|$ is the maximum of $|p_n|$ in $[x_{j+1,n}, x_{jn}]$. ■

Proof that Theorem 1.1(a) $\Leftrightarrow$ (b)

This follows directly from Lemma 3.1(a). ■

Proof that Theorem 1.1(b) $\Rightarrow$ (c).

First note that as $\text{supp}[\mu]$ is compact [5, p. 41],

$$(3.8) \quad \frac{\gamma_{n-1}}{\gamma_n} \leq C.$$

Our hypothesis (1.2), as well as (3.4) give that uniformly for $y_{jn} \in I$,

$$(3.9) \quad \frac{\gamma_{n-1}}{\gamma_n} |p_{n-1}(y_{j,n-1}) p_n(y_{jn})| \sim 1.$$

Then (3.5) gives uniformly for $x \in I$,

$$(3.10) \quad \frac{\gamma_{n-1}}{\gamma_n} \|p_{n-1}\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \|p_n\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \sim 1.$$

Let $I_{jn} = [y_{j+1,n}, y_{jn}]$ for all j, n . We similarly obtain from (3.6) and (3.9) and our spacing that

$$\frac{\gamma_{n-1}}{\gamma_n} \left(\int_{I_{j,n-1}} p_{n-1}^2 d\mu \right)^{1/2} \left(\int_{I_{jn}} p_n^2 d\mu \right)^{1/2} \geq \frac{C}{n}.$$

Here we are also using that μ' is positive and continuous in I . Adding over $y_{jn} \in I$, and using that there are necessarily $\geq Cn$ such y_{jn} , because of the spacing, we obtain

$$\frac{\gamma_{n-1}}{\gamma_n} \sum_{y_{jn} \in I} \left(\int_{I_{j,n-1}} p_{n-1}^2 d\mu \right)^{1/2} \left(\int_{I_{jn}} p_n^2 d\mu \right)^{1/2} \geq C.$$

Cauchy-Schwarz' inequality gives

$$\frac{\gamma_{n-1}}{\gamma_n} \left(\int p_{n-1}^2 d\mu \int p_n^2 d\mu \right)^{1/2} \geq C$$

so that

$$\frac{\gamma_{n-1}}{\gamma_n} \geq C.$$

Together with (3.8), this gives

$$(3.11) \quad a_n = \frac{\gamma_{n-1}}{\gamma_n} \sim 1.$$

So from (3.10), uniformly in $x \in I$,

$$(3.12) \quad \|p_{n-1}\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \|p_n\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \sim 1.$$

■

Proof that Theorem 1.1(c) $\Rightarrow$ (d).

This is immediate. ■

Proof that Theorem 1.1(d) $\Rightarrow$ (b).

From (3.4), (3.11), and our assumed bound (1.4),

$$|\sin [\pi n \omega (y_{jn}) (y_{j,n-1} - y_{jn})] + o(1)| \geq C.$$

This yields

$$\text{dist} (n \omega (y_{jn}) (y_{jn} - y_{j,n-1}), \mathbb{Z}) \geq C.$$

■

Proof of the bound (1.5)

From the recurrence relation and (3.11),

$$\begin{aligned} & \| (x - b_n) p_n \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \| p_n \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \\ & \leq C \left(\| p_{n+1} \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \| p_n \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} + \| p_{n-1} \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \| p_n \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \right) \\ & \leq C, \end{aligned}$$

by (1.4). Then also uniformly in $x \in I$,

$$\| (x - b_n) p_n^2 \|_{L_\infty [x - \frac{A}{n}, x + \frac{A}{n}]} \leq C$$

and we obtain (1.5). ■

4. PROOF OF THEOREM 1.2

We begin with:

Lemma 4.1

Let μ be a regular measure on $\mathbb{R}$ with compact support. Let I be a closed subinterval of the support and assume that in some open interval containing I , μ is absolutely continuous, while μ' is positive and continuous. Assume (1.1). Let $A > 0$.

(a) Let $L \geq 1$. There exists n_0 such that uniformly for $n \geq n_0$, for $x_{jn} \in I$ and $|k - j| \leq L$,

$$(4.1) \quad \text{dist} (n \omega (x_{jn}) (x_{k,n-1} - x_{jn}), \mathbb{Z}) \geq C.$$

(b) Let

$$(4.2) \quad \delta_{jn} := n \omega (x_{jn}) (x_{jn} - x_{j-1,n-2}).$$

There exist $n_0, \eta_0 > 0$ such that uniformly for $n \geq n_0$, and for $x_{jn} \in I$,

$$(4.3) \quad |\delta_{jn}| \leq 1 - \eta_0.$$

(c) There exist $n_0, C_1 > 0$ such that uniformly for $n \geq n_0$ and for $x_{jn} \in I$, we have

$$(4.4) \quad |x_{jn} - b_{n-1}| \sim \| p_{n-2} \|_{L_\infty [x_{jn} - \frac{A}{n}, x_{jn} + \frac{A}{n}]}^2 |\delta_{jn}|.$$

Here if $x_{jn} - b_{n-1} = 0$, both sides are 0.

Proof

(a) Using the spacing (3.2),

$$\begin{aligned} & \text{dist}(n\omega(x_{jn})(x_{k,n-1} - x_{jn}), \mathbb{Z}) \\ &= \text{dist}(n\omega(x_{jn})(x_{j,n-1} - x_{jn}), \mathbb{Z}) + o(1) \end{aligned}$$

so (1.1) gives the result.

(b) The interlacing of the zeros of successive orthogonal polynomials shows that both x_{jn} and $x_{j-1,n-2}$ lie in the interval $(x_{j,n-1}, x_{j-1,n-1})$. Even more, the bounds given in (a) show that for n large enough, both x_{jn} and $x_{j-1,n-2}$ lie in the interval $\left(x_{j,n-1} + \frac{C_1}{n\omega(x_{jn})}, x_{j-1,n-1} - \frac{C_1}{n\omega(x_{jn})}\right)$ for some $C_1 > 0$. Then

$$\begin{aligned} |\delta_{jn}| &= |n\omega(x_{jn})(x_{jn} - x_{j-1,n-2})| \\ &\leq n\omega(x_{jn})(x_{j,n-1} - x_{j+1,n-1}) - 2C_1 = 1 - 2C_1 + o(1), \end{aligned}$$

by (3.2).

(c) From the recurrence relation,

$$(4.5) \quad (x_{jn} - b_{n-1})p_{n-1}(x_{jn}) = a_{n-1}p_{n-2}(x_{jn}).$$

We now examine the behavior of the left and right-hand side as $n \rightarrow \infty$. By (3.1) to (3.3), the local asymptotic (3.6), and the fact that $x_{jn} - y_{j,n-1} = O\left(\frac{1}{n}\right)$,

$$\begin{aligned} \frac{p_{n-1}(x_{jn})}{p_{n-1}(y_{j,n-1})} &= \cos \pi(n\omega(y_{j,n-1})(x_{jn} - y_{j,n-1})) + o(1) \\ &= \cos \pi(n\omega(y_{j,n-1})(x_{jn} - x_{j,n-1} + x_{j,n-1} - y_{j,n-1})) + o(1) \\ &= \cos \pi\left(n\omega(y_{j,n-1})(x_{jn} - x_{j,n-1}) + \frac{1}{2}\right) + o(1) \\ &= -\sin \pi(n\omega(y_{j,n-1})(x_{jn} - x_{j,n-1})) + o(1) \end{aligned}$$

so using our original condition (1.1), we obtain for some threshold n_0 that is independent of j , and for $n \geq n_0$,

$$(4.6) \quad |p_{n-1}(x_{jn})| \sim |p_{n-1}(y_{j,n-1})|.$$

Next, in analyzing the term on the right in (4.5), we use the differentiated form of (3.6): uniformly for $y_{jn} \in I$ and z in compact subsets of $\mathbb{C}$,

$$(4.7) \quad \lim_{n \rightarrow \infty} \frac{p'_n\left(y_{jn} + \frac{z}{n\omega(y_{jn})}\right)}{n\omega(y_{jn})p_n(y_{jn})} = -\pi \sin \pi z.$$

Then noting that we can replace n by $n \pm 2$ in the term involving z , we see that

$$\begin{aligned} \frac{p_{n-2}(x_{jn})}{p_{n-2}(y_{j-1,n-2})} &= \int_{(x_{j-1,n-2}-y_{j-1,n-2})n\omega(y_{j-1,n-2})}^{(x_{jn}-y_{j-1,n-2})n\omega(y_{j-1,n-2})} \frac{p'_{n-2}\left(y_{j-1,n-2} + \frac{t}{n\omega(y_{j-1,n-2})}\right)}{n\omega(y_{j-1,n-2})p_{n-2}(y_{j-1,n-2})} dt \\ &= \int_{(x_{j-1,n-2}-y_{j-1,n-2})n\omega(y_{j-1,n-2})}^{(x_{jn}-y_{j-1,n-2})n\omega(y_{j-1,n-2})} (-\pi \sin \pi t + o(1)) dt. \end{aligned}$$

Here the lower limit of integration is

$$(x_{j-1,n-2} - y_{j-1,n-2}) n\omega(y_{j-1,n-2}) = \frac{1}{2} + o(1),$$

(by (3.1)), so we can continue the above as

$$\begin{aligned} \frac{p_{n-2}(x_{jn})}{p_{n-2}(y_{j-1,n-2})} &= \int_0^{(x_{jn}-x_{j-1,n-2})n\omega(y_{j-1,n-2})} \left(-\pi \sin \left(\pi \left(t + \frac{1}{2} \right) \right) + o(1) \right) dt + o(\delta_{jn}) \\ &= \int_0^{(x_{jn}-x_{j-1,n-2})n\omega(y_{j-1,n-2})} (-\pi \cos \pi t + o(1)) dt + o(\delta_{jn}) \\ &= -\sin \pi \delta_{jn} + o(\delta_{jn}). \end{aligned}$$

Here we are also using that $\omega(y_{j-1,n-2})/\omega(x_{jn}) \rightarrow 1$ as $n \rightarrow \infty$ by continuity of ω in the interior of I . Next, from (b), $|\delta_{jn}| \leq 1 - \varepsilon$, so $|\sin \pi \delta_{jn}| \sim |\delta_{jn}|$ and we can continue this as

$$\frac{p_{n-2}(x_{jn})}{p_{n-2}(y_{j-1,n-2})} = -(\sin \pi \delta_{jn})(1 + o(1)).$$

It is possible here that $\delta_{jn} = 0$, but in such a case both sides are 0. Combining this with (4.5), (4.6) and (3.11) gives uniformly in j and n , for $n \geq n_0$,

$$|x_{jn} - b_{n-1}| |p_{n-1}(y_{j,n-1})| \sim |p_{n-2}(y_{j-1,n-2})| |\sin \pi \delta_{jn}| \sim |p_{n-2}(y_{j-1,n-2})| |\delta_{jn}|.$$

Here by our local limits and (1.3),

$$|p_{n-1}(y_{j,n-1})| = \|p_{n-1}\|_{L_\infty[x_{j+1,n-1}, x_{j,n-1}]} \sim \|p_{n-2}\|_{L_\infty[x_{jn}-\frac{A}{n}, x_{jn}+\frac{A}{n}]}^{-1}.$$

A related assertion holds for $p_{n-2}(y_{j-1,n-2})$. We deduce that

$$|x_{jn} - b_{n-1}| \sim \|p_{n-2}\|_{L_\infty[x_{jn}-\frac{A}{n}, x_{jn}+\frac{A}{n}]}^2 |\delta_{jn}|.$$

Again if $x_{jn} = b_{n-1}$, $\delta_{jn} = 0$. ■

Proof that Theorem 1.2(a) $\Leftrightarrow$ (c)

If first (1.6) holds, then $|\delta_{jn}| \geq C|x_{jn} - b_{n-1}|$ and (4.4) gives

$$C|\delta_{jn}| \geq \|p_{n-2}\|_{L_\infty[x_{jn}-\frac{A}{n}, x_{jn}+\frac{A}{n}]}^2 |\delta_{jn}|,$$

which forces

$$\|p_{n-2}\|_{L_\infty[x_{jn}-\frac{A}{n}, x_{jn}+\frac{A}{n}]}^2 \leq C_1,$$

uniformly in $x_{jn} \in I$, provided no $\delta_{jn} = 0$. Since $\delta_{jn} = 0$ can occur for at most one j , namely when $x_{jn} = b_{n-1}$ (as follows from the recurrence relation), that exceptional interval can be covered by others with A large enough. So we have (1.8).

Conversely, suppose we have (1.8). Then from (4.4),

$$|x_{jn} - b_{n-1}| \leq C |\delta_{jn}|,$$

so that we have (1.6). ■

Proof that Theorem 1.2(b) $\Leftrightarrow$ (c)

It is immediate that (b) $\Rightarrow$ (c). For the converse we note that if (c) holds, then from Theorem 1.1(c),

$$\|p_{n-1}\|_{L_\infty[x-\frac{A}{n}, x+\frac{A}{n}]} \geq C$$

uniformly for $x \in I$. This together with (1.8), gives (1.7). ■

REFERENCES

- [1] M.U. Ambroladze, *On the possible rate of growth of polynomials that are orthogonal with a continuous positive weight*, Math. USSR-Sb. 72 (1992), 311–331.
- [2] A. Aptekarev, S. Denisov, D. Tulyakov, *On a problem by Steklov*, J. Amer. Math. Soc. 29 (2016), 1117–1165.
- [3] S. Denisov, *On the size of the polynomials orthonormal on the unit circle with respect to a measure which is a sum of the Lebesgue measure and P point masses*, Proc. Amer. Math. Soc. 144 (2016), 1029–1039.
- [4] S. Denisov, K. Rush, *Orthogonal polynomials on the circle for the weight w satisfying conditions $w, w^{-1} \in BMO$* , Constr. Approx. 46 (2017), 285–303.
- [5] G. Freud, *Orthogonal Polynomials*, Pergamon Press/ Akademiai Kiado, Budapest, 1971.
- [6] Eli Levin and D.S. Lubinsky, *Universality Limits in the Bulk for Varying Measures*, Advances in Mathematics, 219(2008), 743–779.
- [7] Eli Levin and D.S. Lubinsky, *Local Limits for Orthogonal Polynomials for Varying Weights via Universality*, J. Approx. Theory, 254(2020), 105394.
- [8] D.S. Lubinsky, *A New Approach to Universality Limits involving Orthogonal Polynomials*, Annals of Mathematics, 170(2009), 915–939.
- [9] D.S. Lubinsky, *Pointwise Asymptotics for Orthonormal Polynomials at the Endpoints of the Interval via Universality*, International Maths Research Notices, <https://doi.org/10.1093/imrn/rny042>.
- [10] D.S. Lubinsky, *Local Asymptotics for Orthonormal Polynomials in the Interior of the Support via Universality*, Proc. Amer. Math. Soc., 147(2019), 3877–3886.
- [11] D.S. Lubinsky, *Local Asymptotics for Orthonormal Polynomials on the Unit Circle via Universality*, J. d' Analyse. Math., 141(2020), 285–304.

- [12] E.A. Rakhmanov, *Estimates of the growth of orthogonal polynomials whose weight is bounded away from zero*, (Russian) Mat. Sb. (N.S.) 114(156) (1981), 269–298.
- [13] E.B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer, New York, 1997.
- [14] B. Simon, *Orthogonal Polynomials on the Unit Circle*, Parts 1 and 2, American Mathematical Society, Providence, 2005.
- [15] H. Stahl and V. Totik, *General Orthogonal Polynomials*, Cambridge University Press, Cambridge, 1992.
- [16] V. Totik, *Asymptotics for Christoffel Functions for General Measures on the Real Line*, J. d'Analyse Math., 81(2000), 283–303.
- [17] V. Totik, *Universality and fine zero spacing on general sets*, Arkiv för Matematik, 47(2009), 361–391.

¹MATHEMATICS DEPARTMENT, THE OPEN UNIVERSITY OF ISRAEL, P.O. BOX 808, RAANANA 43107, ISRAEL;, ²SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA., LUBINSKY@MATH.GATECH.EDU