

Communications in Contemporary Mathematics
(2021) 2150083 (47 pages)
© World Scientific Publishing Company
DOI: 10.1142/S0219199721500838



**Bernstein–Sato functional equations,
V-filtrations, and multiplier ideals of direct summands**

Josep Àlvarez Montaner

*Departament de Matemàtiques
and Institut de Matemàtiques de la UPC-BarcelonaTech (IMTech)
Universitat Politècnica de Catalunya, Barcelona 08028, Spain
josep.alvarez@upc.edu*

Daniel J. Hernández

*Department of Mathematics
University of Kansas, Lawrence, KS 66045, USA
hernandez@ku.edu*

Jack Jeffries

*University of Nebraska-Lincoln, Lincoln, NE 68502, USA
jack.jeffries@unl.edu*

Luis Núñez-Betancourt

*Centro de Investigación en Matemáticas
Guanajuato, Gto., México
luisnub@cimat.mx*

Pedro Teixeira

*Department of Mathematics, Knox College
Galesburg, IL 61401, USA
pteixeir@knox.edu*

Emily E. Witt*

*Department of Mathematics, University of Kansas
Lawrence, KS 66045, USA
witt@ku.edu*

Received 25 April 2020

Accepted 23 July 2021

Published

*Corresponding author.

J. Álvarez Montaner et al.

This paper investigates the existence and properties of a Bernstein–Sato functional equation in nonregular settings. In particular, we construct D -modules in which such formal equations can be studied. The existence of the Bernstein–Sato polynomial for a direct summand of a polynomial over a field is proved in this context. It is observed that this polynomial can have zero as a root, or even positive roots. Moreover, a theory of V -filtrations is introduced for nonregular rings, and the existence of these objects is established for what we call *differentially extensible summands*. This family of rings includes toric, determinantal, and other invariant rings. This new theory is applied to the study of multiplier ideals and Hodge ideals of singular varieties. Finally, we extend known relations among the objects of interest in the smooth case to the setting of singular direct summands of polynomial rings.

Keywords: D -module; Bernstein–Sato polynomial; direct summand; V -filtration; ring of invariants; multiplier ideal.

Mathematics Subject classification 2020: 14F10, 13N10, 13A35, 16S32, 13D45, 14B05, 14M25, 13A50

Contents

1. Introduction	2
2. The Bernstein–Sato Functional Equation	8
2.1. Rings of differential operators	8
2.2. The Bernstein–Sato functional equation for polynomial and power series rings	9
2.3. The Bernstein–Sato functional equation for nonregular rings	12
2.4. Some existence results and examples of Bernstein–Sato polynomials	16
3. Differential Direct Summands and Differential Extensibility	18
3.1. Differential extensibility	18
3.2. Differential direct summands and differentially extensible summands	20
3.3. Bernstein–Sato functional equation for direct summands	24
4. V -Filtrations	27
4.1. V -filtrations for nonregular rings	28
4.2. V -filtrations for differentially extensible summands	31
4.3. Hodge ideals	34
5. Multiplier Ideals	36
5.1. Test ideals and Cartier extensibility	37
5.2. Multiplier ideals and reduction modulo p	39
References	43

1. Introduction

The theory of D -modules — modules over rings of differential operators — on a smooth analytic or complex algebraic variety has been an active research area over the last 40 years. It serves as a powerful tool in solving problems arising in a wide range of mathematical disciplines, from analysis to algebraic geometry, the topology

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

of varieties, representation theory, and commutative algebra. This paper centers around two major fundamental constructions in D -module theory: *Bernstein–Sato polynomials* and *V -filtrations*.

The Bernstein–Sato polynomial of a holomorphic or regular function f over $\mathbb{C}$ is the monic polynomial $b(s)$ in $\mathbb{C}[s]$ of least degree for which there exists a polynomial differential operator $\delta(s)$ in the indeterminate s that satisfies the functional equation

$$\delta(s) \cdot f^{s+1} = b(s)f^s.$$

This object originated in independent constructions by Bernstein [3], to establish meromorphic extensions of distributions, and by Sato, as the b -function in the theory of prehomogeneous vector spaces [69, 70].

The existence of a nonzero polynomial satisfying the functional equation above was proved by Bernstein over polynomial rings, and extended by Björk [6, 7] to power series rings. Kashiwara [39] proved that the roots of the Bernstein–Sato polynomial for holomorphic functions are negative rational numbers, extending a result of Malgrange [44, 45] for functions with isolated singularities. Indeed, Malgrange exhibited a relation between these roots and the eigenvalues of the monodromy of the Milnor fiber, and the rationality of the roots means that the monodromy is quasi-unipotent.

Since its inception, the Bernstein–Sato polynomial has found broad applications in the study of singularities. For instance, the roots of the Bernstein–Sato polynomial are related to the jumping numbers of multiplier ideals [18, 26, 41], to spectral numbers [14, 28, 67, 68], and to poles of zeta functions [24]. The Bernstein–Sato polynomial also plays a key role in understanding algorithmic aspects of local cohomology modules and de Rham cohomology [61–63, 83, 84].

Bernstein–Sato polynomials are closely related to the notion of a *V -filtration* on a D -module along an element f , first introduced by Malgrange and Kashiwara [40, 46]. This is a decreasing filtration on a D -module indexed by the rational numbers, and satisfying several conditions; see Definitions 4.1 and 4.3. In fact, a V -filtration can be described using a relative version of the Bernstein–Sato polynomial [49, 65]. These filtrations were originally introduced to relate Deligne’s nearby and vanishing cycles [22] to their corresponding D -modules via the Riemann–Hilbert correspondence.

In the most general form, V -filtrations are known to exist for regular holonomic D -modules with quasi-unipotent monodromy. This condition is satisfied, for example, by the coordinate ring of a smooth variety. Quite nicely, the D -module-theoretic notion of V -filtration in this case is essentially equivalent to the algebro-geometric notion of *multiplier ideals* [18].

Both the notion of the Bernstein–Sato polynomial and the V -filtration have been generalized to the setting of nonprincipal ideals by Budur *et al.* [17]. The roots of the Bernstein–Sato polynomial in this context are still negative rational numbers, and the V -filtration of the coordinate ring along the ideal essentially coincides with the filtration of multiplier ideals of the ideal.

J. Álvarez Montaner et al.

On varieties that are not smooth, there is a natural notion of a ring of differential operators, as defined by Grothendieck [27], but does not necessarily have the same favorable ring-theoretic properties. To start with, they are generally not generated by homotheties (maps of the form $s \mapsto rs$ for fixed r) and derivations, and a full description of these rings is only known in special cases. Moreover, rings of differential operators are in general neither left- nor right-Noetherian [4], and a theory of holonomic D -modules is no longer available in this case. For many applications, the theory of D -modules over singular varieties can be approached by Kashiwara's equivalence, in which a singular variety is embedded into a smooth one [38], so that one can consider the subcategory of D -modules over the smooth variety supported on the singular subvariety. However, this approach is not always satisfactory, as the coordinate ring of the singular variety is not a D -module here.

Nonetheless, multiplier ideals make sense for all normal varieties [21], and one might hope to develop appropriate Bernstein–Sato and V -filtration theories compatible with the multiplier ideal theory, in a way analogous to the smooth case, at least for a reasonable class of singularities.

The first major advance in this direction was due to Huneke *et al.* [1]. They proved that every element f (or more generally, ideal) in a direct summand of a polynomial or formal power series ring admits a Bernstein–Sato polynomial in a weaker sense, in which the functional equation $\delta(t) \bullet f^{t+1} = b(t)f^t$ is satisfied for all *integer* values t . Moreover, they showed that the Bernstein–Sato polynomial of f considered as an element of the direct summand divides the Bernstein–Sato polynomial of f considered as an element of the polynomial ring, but that they are not equal in general. In fact, under the extra condition that every differential operator of the direct summand extends to the polynomial ring, equality always holds [13, Theorem 6.11].

In the pursuit of a theory of Bernstein–Sato polynomials in nonregular rings, previous results [1] are not fully satisfactory, in the sense that they do not realize the functional equation as a *formal equality* in an appropriate D -module. Moreover, the lack of a theory of V -filtrations obstructs our understanding of the relationship between Bernstein–Sato polynomials and multiplier ideals in nonregular contexts.

This paper develops a full theory of Bernstein–Sato polynomials and V -filtrations for a large class of direct summands of polynomial rings. Namely, we concentrate on direct summands satisfying the aforementioned condition on the extensibility of differential operators, a class of rings that includes rings of invariants of finite groups, toric rings, and determinantal rings.

Section 2 builds the foundation for a full theory of Bernstein–Sato polynomials in nonregular rings. Given a field $\mathbb{K}$ of characteristic zero, consider a $\mathbb{K}$ -algebra A , and its ring of $\mathbb{K}$ -linear differential operators $D_{A|\mathbb{K}}$. For simplicity, we highlight our main results for Bernstein–Sato polynomials of a *single element* $f \in A$ here, but all results extend to those associated to a sequence of elements $\underline{f} = f_1, \dots, f_\ell \in A$, and to the relative version of the Bernstein–Sato polynomial.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

The functional equation

$$\delta(s) \bullet f f^s = b(s) f^s$$

should be understood formally as an equality in

$$M^A[\mathbf{f}^s] := A_f[s] \mathbf{f}^s,$$

which is the free rank-one $A_f[s]$ -module generated by the formal symbol $\mathbf{f}^s$. That said, the specialized version of the Bernstein–Sato functional equation for direct summands [1], which reads as $\delta(t) \bullet f^{t+1} = b(t) f^t$ for all $t \in \mathbb{Z}$, is only a family of equalities in the localization A_f . The obstruction to the formal version in this context is the existence of a $D_{A|\mathbb{K}}[s]$ -module structure on $M^A[\mathbf{f}^s]$, an issue that was not previously addressed. The major goal of Sec. 2 is to determine such a structure.

Theorem A (Theorem 2.12). *Suppose that A is either finitely generated over $\mathbb{K}$, or local and complete. Then there exists a unique $D_{A|\mathbb{K}}[s]$ -module structure on $M^A[\mathbf{f}^s]$ that is compatible with the $D_{A|\mathbb{K}}$ -module structure on A_f after specialization.*

With this structure in hand, Proposition 2.14 (see also Corollary 2.16) establishes that the formal and the specialized versions of the Bernstein–Sato polynomial coincide. That is, the equality $\delta(s) \bullet f f^s = b(s) f^s$ holds in $M^A[\mathbf{f}^s]$ if and only if $\delta(t) \bullet f^{t+1} = b(t) f^t$ for all $t \in \mathbb{Z}$. In particular, one deduces the existence of a formal Bernstein–Sato functional equation for direct summands of polynomial rings (see Theorem 2.17).

Theorem 2.12 opens the door to the study of the Bernstein–Sato polynomial for a large class of $\mathbb{K}$ -algebras. To this purpose, necessary and sufficient conditions for its existence are presented in Proposition 2.18, and examples are presented in which the roots of the Bernstein–Sato polynomial include zero, or even positive rational numbers.

Section 3 introduces the class of rings for which we can develop the theory of V -filtrations. When A is a direct summand of a ring T , any differential operator on T yields a differential operator on A after composing with the splitting. On the other hand, we have the notion of *differentially extensible* rings [13] in which we are provided an “opposite” process, since in these rings, every differential operator on A can be extended to a differential operator on T . This work combines both properties in the class of *direct summands of polynomial rings that are differentially extensible*, which contains many rings of interest, including:

- many cases of rings of invariants of finite groups (see Example 3.3),
- toric rings (see Example 3.4, Lemma 3.5 and Remark 3.6), and
- determinantal rings (see Example 3.7).

When we turn our attention to modules, we consider a slight generalization of the category of differential direct summands [1] that allows us to consider $D_{A|\mathbb{K}}[s]$ -modules. Combining this with differential extensibility builds the notion

J. Álvarez Montaner et al.

of *differentially extensible summand* (see Definition 3.16 for details). In particular, we show in Theorem 3.21 that the $D_{A|\mathbb{K}}[s]$ -module $M^A[\mathbf{f}^s]$ is a differential direct summand of the $D_{T|\mathbb{K}}[s]$ -module $M^T[\mathbf{f}^s]$ and it is also a differentially extensible summand (Theorem 3.22).

This fact yields a direct proof of the existence of the formal Bernstein–Sato polynomial for direct summands of a polynomial ring. More importantly, through a slight generalization of a recent result of Mustață [59], we see that the Bernstein–Sato polynomial of a sequence of elements only depends on the ideal generated by these elements, thus extending previous results [17] to this nonregular context.

In Sec. 4, we develop the theory of V -filtrations for nonregular rings. We follow the same ideas used in the smooth case, to axiomatically define the notion of a V -filtration along an ideal of a Noetherian $\mathbb{K}$ -algebra. As in the smooth case, we begin by assuming that the ideal defines a smooth subvariety, and it is generated by a collection of variables (see Definitions 4.1 and 4.3 for details).

A main result of this paper is the existence of V -filtrations for differentially extensible summands.

Theorem B (Corollary 4.11). *Let R be a polynomial ring over a field $\mathbb{K}$ of characteristic zero. Let A be a $\mathbb{K}$ -subalgebra of R such that A is a direct summand of R , and for which the inclusion $A \subseteq R$ is differentially extensible. Let I be an ideal of A , and M a $D_{A|\mathbb{K}}$ -module that is a differentially extensible summand of a regular holonomic $D_{R|\mathbb{K}}$ -module N that has quasi-unipotent monodromy. Then M admits a V -filtration along I .*

This result applies, in particular, when M is A itself, or any local cohomology module of A .

As a consequence, we are able to define Hodge ideals in this singular ambient setting. We also extend known relations of Hodge ideals to minimal exponents in Sec. 4.3.

Corollary C (Corollary 4.14). *Suppose that $\mathbb{K}$ has characteristic zero and R be a polynomial ring over $\mathbb{K}$. Let A be a $\mathbb{K}$ -subalgebra that is a direct summand of R , for which the inclusion $A \subseteq R$ is differentially extensible. Then for all $f \in A$ and $\lambda \in \mathbb{Q}_{\geq 0}$, $I_k^A(f^\lambda)$ exists. Furthermore,*

$$I_k^A(f^\lambda) = I_k^R(f^\lambda) \cap A.$$

We also use this new theory of V -filtrations in Sec. 5 to study multiplier ideals in differentially extensible summands. In particular, we extend results previously obtained only for smooth varieties [17, 18].

Theorem D (Theorem 5.3). *Let R be a polynomial ring over $\mathbb{K}$, and let $A \subseteq R$ be a differentially extensible inclusion of finitely generated $\mathbb{K}$ -algebras, such that A is a direct summand of R . For every ideal I of A , and real number $\lambda \geq 0$, the*

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

following ideals coincide:

- (1) $\mathcal{J}_R((IR)^\lambda) \cap A$,
- (2) $\bigcup_{\alpha > \lambda} V^\alpha A$, where the V -filtration is taken along I , and
- (3) $\{g \in A : \gamma > \lambda \text{ if } b_{I,g}^A(-\gamma) = 0\}$.

We study whether the equality $\mathcal{J}_A(I^\lambda) = \mathcal{J}_R((IR)^\lambda) \cap A$ holds, and appeal to positive characteristic methods to address this question. It is known that via reduction to positive characteristic, multiplier ideals reduce to test ideals when A has KLT singularities [19]. We replace the notion of differential extensibility with the parallel notion of *Cartier extensibility* (see Definition 5.7) for rings of positive characteristic, and obtain the following comparison results.

Theorem E (Proposition 5.9, Theorem 5.15).

- (1) Let $A \subseteq R$ be an extension of rings of positive characteristic, such that R is a regular F -finite domain and A is a Cartier extensible direct summand of R . Then for every ideal I of A , and every real number $\lambda \geq 0$,

$$\tau_A(I^\lambda) = \tau_R((IR)^\lambda) \cap A.$$

- (2) Let R be a polynomial ring over a field $\mathbb{K}$ of characteristic zero, and let $A \subseteq R$ be a differentially extensible inclusion, such that A is a direct summand of R and that A is finitely generated over $\mathbb{K}$ with KLT singularities. Suppose that the reduction modulo p is also Cartier extensible for each prime $p \gg 0$. Then for every ideal I of A and every real number $\lambda \geq 0$,

$$\mathcal{J}_A(I^\lambda) = \mathcal{J}_R((IR)^\lambda) \cap A.$$

- (3) Under the hypotheses of part 2 above, one has equalities

$$\mathcal{J}_A(I^\lambda) = \bigcup_{\alpha > \lambda} V^\alpha A = \{g \in A : \gamma > \lambda \text{ if } b_{I,g}^A(-\gamma) = 0\},$$

where the V -filtration is taken along I .

In particular, this partially answers an open question regarding jumping numbers [1, Question 4.15] for the rings considered here. Moreover, in Theorem 5.19, we relate jumping numbers with the roots of the Bernstein–Sato polynomials, thus extending results in smooth varieties [17, 18, 26] to this nonregular setting.

The hypothesis of Cartier extensibility after reduction modulo a prime may not be necessary, as Example 5.16 shows. Finally, we point out that a similar comparison result [25, Theorem 1.1] was shown for V -filtrations and multiplier ideals in the case of an inclusion of a smooth divisor in a smooth variety. There are also analogous results to Proposition 5.9 for test ideals in special cases [12, 73].

Setup 1.1. Throughout the paper, we generally use the following notation.

- $\mathbb{K}$ is a field,

J. Álvarez Montaner et al.

- A and T denote Noetherian commutative rings containing $\mathbb{K}$, and
- R is a polynomial ring over $\mathbb{K}$ in indeterminates $x_1, \dots, x_d$.

For the most part, the statements of lemmas, propositions, and theorems are self-contained, and do not rely on the conventions above, besides the fact that $\mathbb{K}$ *always denotes a field*.

2. The Bernstein–Sato Functional Equation

2.1. Rings of differential operators

We start this section recalling some notions from the theory of rings of differential operators, as introduced by Grothendieck [27, §16.8].

A *differential operator of order zero* on a ring T is defined by the multiplication by an element $s \in T$. Given an integer $m \geq 1$, a *differential operator of order at most m* is an additive map $\delta \in \text{Hom}_{\mathbb{Z}}(T, T)$ such that the commutator $[\delta, s] = \delta s - s\delta$ is a differential operator of order at most $m - 1$ for each $s \in T$. The set consisting of all differential operators of order at most m is denoted by D_T^m .

Sums and compositions of differential operators are themselves differential operators, hence the differential operators form a subring D_T of $\text{Hom}_{\mathbb{Z}}(T, T)$ that admits a filtration

$$D_T^0 \subseteq D_T^1 \subseteq \dots \subseteq \bigcup_{m \in \mathbb{N}} D_T^m = D_T.$$

We also consider the subring $D_{T|\mathbb{K}} \subseteq D_T$ of $\mathbb{K}$ -linear differential operators.

For a polynomial or power series ring R in the variables $x_1, \dots, x_d$ over a field $\mathbb{K}$, one has equalities [27, §16.11.2]

$$D_{R|\mathbb{K}}^n = R \left\langle \frac{\partial^{a_1}}{a_1!} \dots \frac{\partial^{a_d}}{a_d!} : a_1 + \dots + a_d \leq n \right\rangle,$$

where the operator $\frac{\partial^{a_i}}{a_i!}$ is characterized by rule

$$\frac{\partial^{a_i}}{a_i!} (x_1^{b_1} \dots x_d^{b_d}) = \binom{b_i}{a_i} x_1^{b_1} \dots x_i^{b_i - a_i} \dots x_d^{b_d}.$$

If $\mathbb{K}$ has characteristic zero, this agrees with the familiar description

$$D_{R|\mathbb{K}}^n = R \langle \partial_{x_1}^{a_1} \dots \partial_{x_d}^{a_d} : a_1 + \dots + a_d \leq n \rangle,$$

where $\partial_{x_i}^{a_i} = a_i! \frac{\partial^{a_i}}{a_i!}$ is the usual iterated partial derivative operator.

Let R be a polynomial ring over a field $\mathbb{K}$, and $A = R/I$ for some ideal I of R . The ring of $\mathbb{K}$ -linear differential operators of A has been described in terms of the $\mathbb{K}$ -linear differential operators in R [48, Theorem 15.5.13] (see also [51, 52]). Namely, we have

$$D_{A|\mathbb{K}} \cong \frac{D_{R|\mathbb{K}}(-\log I)}{ID_{R|\mathbb{K}}}, \quad (2.1)$$

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

where $D_{R|\mathbb{K}}(-\log I) := \{\delta \in D_{R|\mathbb{K}} : \delta \bullet I \subseteq I\}$ and the map in the “left” direction corresponds to restriction. The same results also hold when R is a formal power series ring. We point out that the order of the differential operators is preserved; that is, if $D_{R|\mathbb{K}}^n(-\log I) := \{\delta \in D_{R|\mathbb{K}}^n : \delta \bullet I \subseteq I\}$, then we have

$$D_{A|\mathbb{K}}^n \cong \frac{D_{R|\mathbb{K}}^n(-\log I)}{ID_{R|\mathbb{K}}^n}. \quad (2.2)$$

When $\mathbb{K}$ has characteristic $p > 0$, every additive map is $\mathbb{Z}/p\mathbb{Z}$ -linear, and thus $D_T = D_{T|(\mathbb{Z}/p\mathbb{Z})}$. Moreover, let $T^{p^e} \subseteq T$ be the subring consisting of the p^e th powers of all elements of T , and set $D_T^{(e)} := \text{Hom}_{T^{p^e}}(T, T)$. In this case,

$$D_{T|\mathbb{K}} \subseteq D_T \subseteq \bigcup_{e \in \mathbb{N}} D_T^{(e)}.$$

We have $D_{T|\mathbb{K}} = D_T$ whenever $\mathbb{K}$ is a perfect field. On the other hand $D_T = \bigcup_{e \in \mathbb{N}} D_T^{(e)}$ when T is F -finite, that is, when T is finitely generated as a T^{p^e} -module for some (equivalently, all) $e > 0$; see [78, Theorem 2.7; 85, Theorem 1.4.9].

The ring structure and module theory of rings of differential operators is well understood when T is either a polynomial ring or a formal power series ring over a field, but much less is known when T is not a regular ring. As usual, a module over a ring of differential operators will mean a left module.

2.2. The Bernstein–Sato functional equation for polynomial and power series rings

In this section, we retain Setup 1.1, with the additional assumption that $\mathbb{K}$ has characteristic zero.

Consider $D_{R|\mathbb{K}}[s]$, the polynomial ring in a new variable s over the ring of $\mathbb{K}$ -linear differential operators $D_{R|\mathbb{K}}$. For every $f \in R$, there exist $\delta(s) \in D_{R|\mathbb{K}}[s]$ and a nonzero polynomial $b(s) \in \mathbb{K}[s]$ such that the following *functional equation* is satisfied:

$$\delta(s) \bullet f f^s = b(s) f^s. \quad (2.3)$$

We note that this equation can be interpreted in two different senses. First, we may view it as a family of equations inside the module R_f , indexed by $s \in \mathbb{Z}$; that is, one has $\delta(t) \bullet f^{t+1} = b(t) f^t$ in R_f for all $t \in \mathbb{Z}$. We temporarily say that the polynomial $b(s)$ satisfies the *specialized* functional equation in this case. Later on, in Corollary 2.16, we prove that this interpretation is equivalent to the *a priori* stronger interpretation of (2.3) as an equality in a D -module

$$M^R[f^s] := R_f[s] f^s,$$

which is the free rank-one $R_f[s]$ -module generated by the formal symbol f^s . We say that the polynomial $b(s)$ satisfies the functional equation *formally* in this case.

J. Álvarez Montaner et al.

A key point is that $M^R[\underline{f}^s]$ has a $D_{R[\mathbb{K}[s]]}$ -module structure given by the action of the partial derivatives as follows: for $h \in R_f[s]$, we have

$$\partial_{x_r} \bullet h \underline{f}^s = \left(\frac{\partial h}{\partial x_r} + shf^{-1} \frac{\partial f}{\partial x_r} \right) \underline{f}^s.$$

Henceforth in this section, we consider the functional equation formally. The collection of all polynomials $b(s)$ satisfying a functional equation as in (2.3), for some $\delta(s) \in D_{R[\mathbb{K}[s]]}$, is an ideal in $\mathbb{K}[s]$, and the unique monic generator of this ideal is known as the *Bernstein–Sato polynomial* associated to f ; we denote this polynomial as $b_f^R(s)$. Alternatively, the Bernstein–Sato polynomial $b_f^R(s)$ is the monic polynomial of smallest degree among those polynomials $b(s)$ such that $b(s)\underline{f}^s$ lies in the $D_{R[\mathbb{K}[s]]}$ -submodule of $M^R[\underline{f}^s]$ generated by $\underline{f}^s$.

The existence of $b_f^R(s)$ in the case that R is a polynomial ring is due to Bernstein [3], and this polynomial coincides with the *b-function* in the theory of prehomogeneous vector spaces developed by Sato [69, 70]. Björk established the existence of $b_f^R(s)$ when R is a power series ring [6, 7]. A fundamental property of the Bernstein–Sato polynomial is that its roots are rational numbers, and consequently $b_f^R(s) \in \mathbb{Q}[s]$, in the case that R is either a polynomial ring or a ring of convergent power series. This was proved by Malgrange for elements f such that R/fR has an isolated singularity [44, 45], and by Kashiwara in general [39].

Budur *et al.* extended the notion of a Bernstein–Sato polynomial to be associated to an *ideal* in a polynomial ring R [17], and we work in this more general framework. We point out that recent work of Mustață [59] implies that this construction is also valid for formal power series rings.

Definition 2.1 (The module $M^R[\underline{f}^s]$). Fix $\underline{f} = f_1, \dots, f_\ell \in R$. Now, given indeterminates $\underline{s} = s_1, \dots, s_\ell$, we define the free rank-one $R_{f_1 \dots f_\ell}[\underline{s}]$ -module

$$M^R[\underline{f}^s] := R_{f_1 \dots f_\ell}[\underline{s}] \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell},$$

where $\underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}$ is a formal symbol for the generator.

The $R_{f_1 \dots f_\ell}[\underline{s}]$ -module $M^R[\underline{f}^s]$ has a natural $D_{R[\mathbb{K}[\underline{s}]]}$ -module structure (and so also a $D_{R[\mathbb{K}]}$ -module structure), defined as follows: for $h \in R_{f_1 \dots f_\ell}$, set

$$\partial_{x_r} \bullet h \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell} = \left(\frac{\partial h}{\partial x_r} + h \sum_{i=1}^{\ell} s_i f_i^{-1} \frac{\partial f_i}{\partial x_r} \right) \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}. \quad (2.4)$$

Since $D_{R[\mathbb{K}[\underline{s}]]}$ is generated by R , $s_1, \dots, s_\ell$, and the partial derivatives $\partial_{x_1}, \dots, \partial_{x_\ell}$, this gives a recipe for an action by any element of $D_{R[\mathbb{K}[\underline{s}]]}$.

Definition 2.2 (The exponent specialization map φ_t on $M^R[\underline{f}^s]$). The *exponent specialization map* on $M^R[\underline{f}^s]$ associated to $t = (t_1, \dots, t_\ell) \in \mathbb{Z}^\ell$ is the $R_{f_1 \dots f_\ell}$ -linear map

$$\varphi_t: M^R[\underline{f}^s] \rightarrow R_{f_1 \dots f_\ell}$$

given by $\varphi_t(\underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}) = \underline{f}_1^{t_1} \cdots \underline{f}_\ell^{t_\ell}$, and $\varphi_t(s_i) = t_i$.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

For each $t \in \mathbb{Z}^\ell$, $\delta \in D_{R[\mathbb{K}][s]}$, and $v \in M^R[\underline{f}^s]$ we have

$$\varphi_t(\delta \bullet v) = \delta(t) \bullet \varphi_t(v), \quad (2.5)$$

showing, in particular, that φ_t is $D_{R[\mathbb{K}]}$ -linear. Indeed, the verification of this equality reduces to the case of $\delta = \partial_{x_r}$, in which case it follows from (2.4).

Definition 2.3 (Bernstein–Sato functional equation for polynomial rings).

Fix follow this style globally $\underline{f} = f_1, \dots, f_\ell \in R$. For an integer $m \geq 0$, let $\binom{s_i}{m}$ denote $s_i(s_i - 1) \cdots (s_i - m + 1)/m! \in \mathbb{K}[s]$. There exists a nonzero polynomial $b(s) \in \mathbb{Q}[s]$ for which $b(s_1 + \cdots + s_\ell) \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}$ is in the $D_{R[\mathbb{K}][s]}$ -submodule of $M^R[\underline{f}^s]$ generated by the elements

$$\prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \cdots f_\ell^{c_\ell} \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell},$$

where $c = (c_1, \dots, c_\ell)$ runs over the elements of $\mathbb{Z}^\ell$ for which $|c| := c_1 + \cdots + c_\ell = 1$, and $\text{nsupp}(c) := \{i : c_i < 0\}$. Equivalently, there exists $\delta_c \in D_{R[\mathbb{K}][s]}$, with $\delta_c = 0$ for all but finitely many c , for which the following *functional equation* holds:

$$\begin{aligned} \sum_{\substack{c \in \mathbb{Z}^\ell \\ |c|=1}} \delta_c \bullet \prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \cdots f_\ell^{c_\ell} \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell} \\ = b(s_1 + \cdots + s_\ell) \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}. \end{aligned} \quad (2.6)$$

Definition 2.4 (Bernstein–Sato polynomial associated to an ideal in a polynomial ring). Let I denote the ideal generated by $\underline{f} = f_1, \dots, f_\ell \in R$. The *Bernstein–Sato polynomial* $b_I^R(s)$ is the unique monic polynomial $b(s) \in \mathbb{Q}[s]$ of smallest degree satisfying the functional equation (2.6), for some $\delta_c \in D_{R[\mathbb{K}][s]}$.

The Bernstein–Sato polynomial of I does not depend on the choice of generators of I [17, Theorem 2.5]. For a principal ideal $I = \langle f \rangle$, $b_I^R(s)$ agrees with $b_f^R(s)$, and in general, like $b_f^R(s)$, all roots of $b_I^R(s)$ are negative rational numbers [17].

Example 2.5. Let $R = \mathbb{C}[x, y, z]$, $f_1 = xy$, and $f_2 = xz$. There is a functional equation for $\underline{f} = f_1, f_2$ involving the generators corresponding to the vectors $c = (1, 0)$ and $(0, 1)$. Namely,

$$\partial_x \partial_y \bullet f_1 \underline{f}_1^{s_1} \underline{f}_2^{s_2} + \partial_x \partial_z \bullet f_2 \underline{f}_1^{s_1} \underline{f}_2^{s_2} = (s_1 + s_2 + 1)(s_1 + s_2 + 2) \underline{f}_1^{s_1} \underline{f}_2^{s_2},$$

so the Bernstein–Sato polynomial $b_{\langle xy, xz \rangle}^R$ divides $(s+1)(s+2)$. Using the methods from [17, Sec. 4.3], one can show that equality holds here.

Example 2.6. Let $R = \mathbb{C}[x, y, z]$, $f_1 = xz^2$, and $f_2 = yz^3$. There is a functional equation for $\underline{f} = f_1, f_2$ using the vectors $c = (0, 1)$, $(1, 0)$, $(2, -1)$, and $(3, -2)$:

$$\begin{aligned} \delta_1 \bullet f_2 \underline{f}_1^{s_1} \underline{f}_2^{s_2} + \delta_2 \bullet f_1 \underline{f}_1^{s_1} \underline{f}_2^{s_2} + \delta_3 \bullet \binom{s_2}{1} f_1^2 f_2^{-1} \underline{f}_1^{s_1} \underline{f}_2^{s_2} + \delta_4 \bullet \binom{s_2}{2} f_1^3 f_2^{-2} \underline{f}_1^{s_1} \underline{f}_2^{s_2} \\ = b(s_1 + s_2) \underline{f}_1^{s_1} \underline{f}_2^{s_2}, \end{aligned}$$

J. Álvarez Montaner et al.

where

$$\begin{aligned}
 b(s) &= (s+1)^2(s+2) \left(s + \frac{1}{2}\right) \left(s + \frac{2}{3}\right) \left(s + \frac{4}{3}\right), \\
 \delta_1 &= \frac{1}{2592}(-66 - 66s_1 + 31s_2 + 79s_1s_2 + 96s_2^2) \partial_y \partial_z^3, \\
 \delta_2 &= \frac{1}{2592}(1350 + 3300s_1 + 2592s_1^2 + 648s_1^3 + 3315s_2 + 5128s_1s_2 \\
 &\quad + 1944s_1^2s_2 + 2684s_2^2 + 2114s_1s_2^2 + 915s_2^3) \partial_x \partial_z^2, \\
 \delta_3 &= \frac{1}{1296}(-156 - 132s_1 + 59s_2 + 158s_1s_2 + 192s_2^2) y \partial_x^2 \partial_z, \\
 \delta_4 &= \frac{1}{108}(3 - s_2) y^2 \partial_x^3.
 \end{aligned}$$

Using the methods from [17, Sec. 4.3], one can show that $b(s)$ is the Bernstein–Sato polynomial of $\underline{f} = f_1, f_2$.

The following generalization of Definition 2.4 is useful for applications to birational geometry.

Definition 2.7 (Relative Bernstein–Sato polynomial in a polynomial ring). Fix an ideal I generated by $f_1, \dots, f_\ell \in R$, and an element $g \in R$. The *Bernstein–Sato polynomial of I relative to g* , denoted $b_{I,g}^R(s)$, is the unique monic polynomial $b(s)$ of the smallest degree satisfying the equation obtained from (2.6), after replacing $\underline{f}^{s_1} \cdots \underline{f}^{s_\ell}$ with $g \underline{f}^{s_1} \cdots \underline{f}^{s_\ell}$ on both sides.

As with Definition 2.4, it is a fact that this notion is well-defined (see [16, Remark 2.3; 17, Sec. 2.10]). Moreover, if $g = 1$, then $b_{I,g}^R(s) = b_I^R(s)$.

2.3. The Bernstein–Sato functional equation for nonregular rings

We now proceed to define a module analogous to the one considered in Definition 2.1 and functional equations as considered in Definition 2.4 for more general rings. Again, we may consider functional equations either in the *specialized* or *formal* sense, as in the previous section. The papers [1] and [35] study Bernstein–Sato polynomials in the specialized sense; that is, the monic polynomial of smallest degree among those polynomials $b(s)$ for which there exists $\delta(s) \in D_{R|\mathbb{K}}[s]$ such that $\delta(t) \bullet f^{t+1} = b(t)f^t$ for all $t \in \mathbb{Z}$. Here, we develop a framework for formal Bernstein–Sato polynomials. The following object plays a central role in this theory.

Definition 2.8 ($M^A[\underline{f}^s]$ as an $A_{f_1 \dots f_\ell}[s]$ -module). Let A be a $\mathbb{K}$ -algebra. Given $\underline{f} = f_1, \dots, f_\ell \in A$, we define

$$M^A[\underline{f}^s] := A_{f_1 \dots f_\ell}[s] \underline{f}_1^{s_1} \cdots \underline{f}_\ell^{s_\ell}$$

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

as an $A_{f_1 \dots f_\ell}[s]$ -module. Moreover, for $t = (t_1, \dots, t_\ell) \in \mathbb{Z}^\ell$, we define the exponent specialization map $\varphi_t: M^A[\underline{f}^s] \rightarrow A_{f_1 \dots f_\ell}$ in the manner directly analogous to that defined in Definition 2.2.

We want to endow $M^A[\underline{f}^s]$ with a $D_{A[\mathbb{K}]}[s]$ -module structure that is compatible with the $D_{A[\mathbb{K}]}$ -module structure on $A_{f_1 \dots f_\ell}$ via the exponent specialization maps. In general, this allows for at most one such structure, as Theorem 2.12 shows. The following lemma is a key observation in establishing this. Although this is likely a well-known result, we include its simple proof, for its centrality in our constructions, and for lack of an appropriate reference.

Lemma 2.9. *Let T be an algebra over a field of characteristic zero, and let $h \in T[s_1, \dots, s_\ell]$. Suppose h has degree at most m in each variable. Then h vanishes on the set $\mathcal{A} = \{0, 1, \dots, m\}^\ell$ if and only if $h = 0$.*

Proof. We prove the “only if” implication, starting with polynomials in one variable. The result is clear for $m = 0$, so suppose m is positive and the result holds for one-variable polynomials of degree strictly less than m . Suppose $h \in T[s]$ has degree at most m and vanishes on $\{0, \dots, m\}$. Then $h = (s - m)g$, for some $g \in T[s]$ of degree strictly less than m . Since $h(a) = 0$ and $a - m$ is a unit in T for each $a \in \{0, \dots, m - 1\}$, it follows that $g(a) = 0$ for all such a . By our induction hypothesis, $g = 0$, and consequently $h = 0$.

Now suppose the result holds for polynomials in $\ell - 1$ variables, and suppose $h \in T[s_1, \dots, s_\ell]$ is nonzero and has degree at most m in each variable. Without loss of generality, we assume that m is positive and the indeterminate s_ℓ effectively appears in h , and write

$$h = h_0 + h_1 s_\ell + \dots + h_n s_\ell^n,$$

with $h_i \in T[s_1, \dots, s_{\ell-1}]$, for $i = 0, \dots, n$, and $h_n \neq 0$. By our induction hypothesis, there exists $a_1, \dots, a_{\ell-1} \in \{0, \dots, m\}$ such that $h_n(a_1, \dots, a_{\ell-1}) \neq 0$. We conclude that $h(a_1, \dots, a_{\ell-1}, s_\ell)$ is a nonzero polynomial, and there exists $a_\ell \in \{0, \dots, m\}$ such that $h(a_1, \dots, a_{\ell-1}, a_\ell) \neq 0$. $\square$

Corollary 2.10. *Under the assumptions of Lemma 2.9, the polynomial h has coefficients in $\mathbb{Q}[h(\mathcal{A})] \subseteq \mathbb{Q}[h(\mathbb{Z}^\ell)]$.*

Proof. This follows from the identity

$$h = \sum_{a \in \mathcal{A}} h(a) \prod_{i=1}^{\ell} \prod_{\substack{0 \leq j \leq m \\ j \neq a_i}} \frac{s_i - j}{a_i - j},$$

which is obtained by applying Lemma 2.9 to the difference between the left- and the right-hand sides. $\square$

J. Álvarez Montaner et al.

The following technical lemma deals with the behavior of differential operators with respect to localization. We remark that, given $\delta \in D_{T|\mathbb{K}}$ and $g \in T$, there is a unique $\delta_g \in D_{T_g|\mathbb{K}}$ that extends δ [47, Theorem 2.2.10].

Lemma 2.11. *Given a $\mathbb{K}$ -algebra T , fix $g \in T$ and an ideal I of T . Then the image of any element of $D_{T|\mathbb{K}}(-\log I)$ in $D_{T_g|\mathbb{K}}$ under the localization map lies in $D_{T_g|\mathbb{K}}(-\log IT_g)$.*

Proof. First observe that if $h \in T$ and $\delta \in D_{T|\mathbb{K}}^n(-\log I)$ for some $n \geq 1$, then for all $a \in I$, $[\delta, h] \bullet a = \delta \bullet (ha) - h\delta \bullet a$ is again in I . Thus, $[\delta, h] \in D_{T|\mathbb{K}}^{n-1}(-\log I)$.

For elements of $D_{T|\mathbb{K}}(-\log I)$ of order zero, the statement clearly holds. Proceeding by induction on the order of a differential operator, fix $n \geq 1$ and $\delta \in D_{T|\mathbb{K}}^n(-\log I)$, and suppose that the claim holds for all operators in $D_{T|\mathbb{K}}^{n-1}(-\log I)$. Since $\delta \circ g^t = g^t \circ \delta + [\delta, g^t]$, it follows that for $a \in I$ and $t \geq 0$,

$$\begin{aligned} \delta \bullet a &= \delta \bullet \left(g^t \cdot \frac{a}{g^t} \right) = g^t \delta \bullet \frac{a}{g^t} + [\delta, g^t] \bullet \frac{a}{g^t}, \\ \text{so } \delta \bullet \frac{a}{g^t} &= \frac{\delta \bullet a - [\delta, g^t] \bullet \frac{a}{g^t}}{g^t}. \end{aligned}$$

Note that $[\delta, g^t]$ is in $D_{T|\mathbb{K}}^{n-1}(-\log I)$ by our initial observation, so that it is also in $D_{T_g|\mathbb{K}}(-\log IT_g)$ by the inductive hypothesis. Finally, since $\delta \bullet a \in I$ by our choice of δ , we conclude that $\delta \bullet \frac{a}{g^t} \in IT_g$. $\square$

Theorem 2.12. *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero. Suppose that either A is finitely generated over $\mathbb{K}$, or that A is local and complete. Given $\underline{f} = f_1, \dots, f_\ell \in A$, there is a unique $D_{A|\mathbb{K}}[\underline{s}]$ -module structure on $M^A[\underline{f}^{\underline{s}}]$ with the following property: for all $v \in M^A[\underline{f}^{\underline{s}}]$, $\delta \in D_{A|\mathbb{K}}[\underline{s}]$, and $t \in \mathbb{Z}^\ell$*

$$\varphi_t(\delta \bullet v) = \delta(t) \bullet \varphi_t(v).$$

Proof. First write $A = R/I$ for some polynomial or power series ring R over $\mathbb{K}$, and I an ideal in R . We claim that if $\delta \in D_{R|\mathbb{K}}(-\log I)[\underline{s}]$, then

$$\delta \bullet IM^R[\underline{f}^{\underline{s}}] \subseteq IM^R[\underline{f}^{\underline{s}}]. \quad (2.7)$$

Toward (2.7), fix $a \in IR_{f_1 \dots f_\ell}[\underline{s}]$, so that $a \underline{f}^{\underline{s}} \in IM^R[\underline{f}^{\underline{s}}]$. Observe that for $t \in \mathbb{Z}^\ell$, $\delta(t) \in D_{R|\mathbb{K}}(-\log I)$ and $a(t) \in IR_{f_1 \dots f_\ell}$, so that $\delta(t) \in D_{R_{f_1 \dots f_\ell}|\mathbb{K}}(-\log IR_{f_1 \dots f_\ell})$ by Lemma 2.11. Therefore, $\varphi_t(\delta \bullet a \underline{f}^{\underline{s}}) = \delta(t) \bullet a(t) f_1^{t_1} \dots f_\ell^{t_\ell}$ is in $IR_{f_1 \dots f_\ell}$. Writing $\delta \bullet a \underline{f}^{\underline{s}} = b \underline{f}^{\underline{s}}$ for some $b \in R_{f_1 \dots f_\ell}[\underline{s}]$, consider the image of b in the polynomial ring over the $\mathbb{K}$ -algebra $A_{f_1 \dots f_\ell} = (R/I)_{f_1 \dots f_\ell}$. As a polynomial over this ring, our work thus far shows that b takes the value zero for all $t \in \mathbb{Z}^\ell$, and hence is the zero polynomial by Lemma 2.9. That is, $\delta \bullet a \underline{f}^{\underline{s}} \in IR_{f_1 \dots f_\ell}[\underline{s}] \underline{f}^{\underline{s}}$, proving (2.7).

Note also that if $\delta \in ID_{R|\mathbb{K}}$, then $\delta \bullet M^R[\underline{f}^{\underline{s}}] \subseteq IM^R[\underline{f}^{\underline{s}}]$. Now, using (2.1), we obtain a $D_{A|\mathbb{K}}[\underline{s}]$ -module structure on $M^{\bar{R}}[\underline{f}^{\underline{s}}]/IM^R[\underline{f}^{\underline{s}}] \cong M^A[\underline{f}^{\underline{s}}]$ that is compatible with the specialization maps.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

To see the uniqueness of this structure, let $\star$ and $\bullet$ denote two $D_{A|\mathbb{K}}[s]$ -actions that satisfy the given property. Fix v and δ as in the statement, and then fix $a, b \in A_{f_1 \dots f_\ell}[s]$ for which

$$\delta \star v = a f_1^{s_1} \dots f_\ell^{s_\ell} \quad \text{and} \quad \delta \bullet v = b f_1^{s_1} \dots f_\ell^{s_\ell}.$$

Since, by our hypothesis,

$$a(t) f_1^{t_1} \dots f_\ell^{t_\ell} = \varphi_t(\delta \star v) = \delta(t) \bullet \varphi_t(v) = \varphi_t(\delta \bullet v) = b(t) f_1^{t_1} \dots f_\ell^{t_\ell}$$

for every $t = (t_1, \dots, t_\ell) \in \mathbb{Z}^\ell$, we have that $a(t) = b(t)$ for every $t \in \mathbb{Z}^\ell$. Lemma 2.9 then allows us to conclude that $a = b$, and the claim follows. $\square$

Convention 2.13. When referring to a $D_{A|\mathbb{K}}[s]$ -module structure on $M^A[\underline{f}^s]$, we use the unique $D_{A|\mathbb{K}}[s]$ -module structure compatible with specialization described in Theorem 2.12.

From Theorem 2.12, we obtain a suitable D -module in which to judge the existence of a Bernstein–Sato polynomial in the formal sense. We show now that the notions of formal and specialized Bernstein–Sato polynomials are equivalent.

Proposition 2.14. *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero. Suppose that A is either finitely generated over $\mathbb{K}$, or that A is local and complete. Then for $f \in A$, the equality $\delta \bullet f f^s = b(s) f^s$ holds in $M^A[\underline{f}^s]$ if and only if $\delta(t) \bullet f^{t+1} = b(t) f^t$ for all $t \in \mathbb{Z}$.*

Proof. If $\delta \bullet f f^s = b(s) f^s$ holds in $M^A[\underline{f}^s]$, we can apply the map φ_t to both sides to obtain $\delta(t) \bullet f^{t+1} = b(t) f^t$ for any t . Conversely, suppose that $\delta(t) \bullet f^{t+1} = b(t) f^t$ for all t . We then have $\varphi_t(\delta \bullet f f^s) = \varphi_t(b(s) f^s)$ for all t . Writing $(\delta f - b(s)) \bullet f^s = a f^s$ for some $a \in A_f[s]$, we have $a(t) = \varphi_t(a) = 0$ for all t , so $a = 0$ by Lemma 2.9. It follows that $\delta \bullet f f^s = b(s) f^s$ in $M^A[\underline{f}^s]$. $\square$

An analogous (omitted) proof also holds for the general case we present in the following.

Proposition 2.15. *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero. Suppose that A is either finitely generated over $\mathbb{K}$, or that A is local and complete. Fix $\underline{f} = f_1, \dots, f_\ell \in A$. The equality*

$$\sum_{\substack{c \in \mathbb{Z}^\ell \\ |c|=1}} \delta_c \bullet \prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \dots f_\ell^{c_\ell} f_1^{s_1} \dots f_\ell^{s_\ell} = b(s_1 + \dots + s_\ell) f_1^{s_1} \dots f_\ell^{s_\ell}$$

holds in $M^A[\underline{f}^s]$ if and only if

$$\sum_{\substack{c \in \mathbb{Z}^\ell \\ |c|=1}} \delta_c(t) \bullet \prod_{i \in \text{nsupp}(c)} \binom{t_i}{-c_i} f_1^{t_1+c_1} \dots f_\ell^{t_\ell+c_\ell} = b(t_1 + \dots + t_\ell) f_1^{t_1} \dots f_\ell^{t_\ell}$$

for all $t = (t_1, \dots, t_\ell) \in \mathbb{Z}^\ell$.

J. Álvarez Montaner et al.

Corollary 2.16. *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero. Suppose that A is either finitely generated over $\mathbb{K}$, or that A is local and complete. Fix $\underline{f} = f_1, \dots, f_\ell \in A$. Then there exists a formal Bernstein–Sato polynomial for $\underline{f}$ if and only if there exists a specialized Bernstein–Sato polynomial for $\underline{f}$, and in this case, these two polynomials agree.*

The existence of the specialized Bernstein–Sato polynomial for direct summands of polynomial rings was recently proven [1]. Thus, we can extend their result to the formal setting.

Theorem 2.17 ([1, Theorem 3.14]). *Let A be a finitely generated $\mathbb{K}$ -subalgebra of a polynomial ring R , such that A is a direct summand of R . Then the Bernstein–Sato polynomial $b_{\underline{f}}^A(s)$ exists for any $\underline{f} = f_1, \dots, f_\ell \in A$. Moreover $b_{\underline{f}}^A(s)$ divides $b_{\underline{f}}^R(s)$.*

The question whether the Bernstein–Sato polynomial $b_{\underline{f}}^A(s)$ only depends on the ideal generated by $\underline{f}$ was not considered in the initial work on direct summands [1]. We address this issue in Sec. 3.3.

2.4. Some existence results and examples of Bernstein–Sato polynomials

A sufficient condition for the existence of Bernstein–Sato polynomials in the polynomial ring case is the fact that $M^R[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is a D -module of finite length. This result was extended to the case of differentiably admissible $\mathbb{K}$ -algebras [50, 60]. In general, for the case of $\mathbb{K}$ -algebras, we provide necessary and sufficient conditions for the existence of Bernstein–Sato polynomials.

Proposition 2.18. *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero. Suppose that A is either finitely generated over $\mathbb{K}$, or that A is local and complete. Fix an element $f \in A$. Then, the following are equivalent:*

- (1) *There exists a Bernstein–Sato polynomial for f .*
- (2) *$M^A[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is a finitely-generated $D_{A|\mathbb{K}}[s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ -module.*
- (3) *$M^A[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is generated by $\mathbf{f}^s$ as a $D_{A|\mathbb{K}}[s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ -module.*

Proof. We observe first that for $\delta(s) \in D_{A|\mathbb{K}}[s]$ and $b(s) \in \mathbb{K}[s]$ and for any integer j , the equality $\delta(s) \cdot f \mathbf{f}^s = b(s) \mathbf{f}^s$ holds in $M^A[\mathbf{f}^s]$ if and only if $\delta(s+j) \cdot f^{j+1} \mathbf{f}^s = b(s+j) f^j \mathbf{f}^s$; this is a straightforward application of Lemma 2.9, as in the results of the previous section.

Now, assume that $M^A[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is finitely generated over $D_{A|\mathbb{K}}[s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$, and take generators $a_1 f^{j_1} \mathbf{f}^s, \dots, a_n f^{j_n} \mathbf{f}^s$ with $a_i \in A$ and $j_i \in \mathbb{Z}$. We can replace each j_i by $j = \min\{j_i\}$. Then, there exist operators $\delta_1, \dots, \delta_n \in D_{A|\mathbb{K}}(s)$ such that $\sum_i \delta_i \cdot (a_i f^j \mathbf{f}^s) = f^{j-1} \mathbf{f}^s$. We can rewrite this as $(\sum_i \delta_i a_i) \cdot f^j \mathbf{f}^s = f^{j-1} \mathbf{f}^s$. The element $\sum_i \delta_i a_i$ can be written as a quotient of an element $\delta(s) \in D_{A|\mathbb{K}}[s]$ and an

element $b(s) \in \mathbb{K}[s]$. We then have the equality $\delta(s) \bullet f^j \mathbf{f}^s = b(s) f^{j-1} \mathbf{f}^s$ in $M^A[\mathbf{f}^s]$. By our observation above, there exists a Bernstein–Sato polynomial for f .

On the other hand, if there exists a Bernstein–Sato polynomial $b(s)$ for f , then $\delta(s) \bullet f \mathbf{f}^s = b(s) \mathbf{f}^s$, for some $\delta(s) \in D_{A|\mathbb{K}[s]}$, and by the observation above, $\delta(s+j) \bullet f^{j+1} \mathbf{f}^s = b(s+j) f^j \mathbf{f}^s$ for any integer j . We then have $\frac{\delta(s+j)}{b(s+j)} \bullet f^{j+1} \mathbf{f}^s = f^j \mathbf{f}^s$ for any integer j . Since $M^A[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is generated as an A -module by $\{f^j \mathbf{f}^s : j \in \mathbb{Z}\}$, we conclude that $M^A[\mathbf{f}^s] \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$ is generated by $\mathbf{f}^s$ as a module over $D_{A|\mathbb{K}[s]} \otimes_{\mathbb{K}[s]} \mathbb{K}(s)$. $\square$

In the following, we show that for nonregular rings we have examples where the roots of the Bernstein–Sato polynomial may be zero or positive rational numbers.

Example 2.19. Let $A = \mathbb{C}[x, y]/\langle xy \rangle$ and $f = x$. The operator $\delta = x\partial_x^2 \in D_{\mathbb{C}[x, y]|\mathbb{C}}$ is easily seen to stabilize the ideal $\langle xy \rangle$, and hence descends to a differential operator on A . The equation $\delta \bullet x^{t+1} = (t+1)tx^t$ holds for all nonnegative integers $t \in \mathbb{N}$, and thus, by Proposition 2.14, the functional equation $\delta \bullet x \mathbf{x}^s = (s+1)s \mathbf{x}^s$ holds in $M^A[\mathbf{x}^s]$. Thus, $b_x^A(s)$ divides $s(s+1)$.

In fact, the equality $b_x^A(s) = s(s+1)$ holds here. To see this, it suffices to show that if $\delta_{-1} \bullet x^0 = c_{-1}x^{-1}$ and $\delta_0 \bullet x^1 = c_0x^0$ with $\delta_{-1}, \delta_0 \in D_{A|\mathbb{C}}$ and $c_{-1}, c_0 \in \mathbb{C}$, then $c_{-1} = c_0 = 0$. Observe that A is a $D_{A|\mathbb{C}}$ -module, that $\langle x \rangle$ and $\langle y \rangle$ are $D_{A|\mathbb{C}}$ -submodules of A [13, Lemma 3.3], and thus the sum $\langle x, y \rangle \subseteq A$ is a $D_{A|\mathbb{C}}$ -submodule. The claim then follows.

Example 2.20. Let $A = \mathbb{C}[x^2, x^3]$ and $f = x^2$. Consider the endomorphism of A given by

$$\delta = (x\partial_x - 1) \circ \partial_x^2 \circ (x\partial_x - 1)^{-1},$$

where $(x\partial_x - 1)^{-1}$ is the inverse function of $x\partial_x - 1$ on A (i.e. the linear operator sending x^n to $\frac{1}{n-1}x^n$ for all $n \neq 1$). One can verify by the definition that $\delta \in D_{A|\mathbb{C}}^2$; alternatively, see [77, 79]. The equation $\delta \bullet x^{2(t+1)} = (2t+2)(2t-1)x^{2t}$ holds for all nonnegative integers $t \in \mathbb{N}$, and thus, by Proposition 2.14, the functional equation

$$\delta \bullet x^2 (\mathbf{x}^2)^s = (2s+2)(2s-1)(\mathbf{x}^2)^s$$

holds in $M^A[(\mathbf{x}^2)^s]$. Thus, $b_{x^2}^A(s)$ divides $(s - \frac{1}{2})(s+1)$.

In fact, the equality $b_{x^2}^A(s) = (s - \frac{1}{2})(s+1)$ holds here. By the same argument as in the previous example, $s = -1$ is a root of $b_{x^2}^A(s)$. By a specific description of $D_{A|\mathbb{C}}$ [77, 79], every differential operator of degree -2 on A can be written as $(x\partial_x - 1) \circ \partial_x^2 \circ \Upsilon \circ (x\partial_x - 1)^{-1}$, for some $\Upsilon \in \mathbb{C}[x\partial_x]$. Since $M^A[(\mathbf{x}^2)^s]$ is a graded module we can decompose the functional equation as a sum of homogeneous pieces, and thus there exists a functional equation of the form

$$P(s) \bullet x^2 (\mathbf{x}^2)^s = b_{x^2}^A(s) (\mathbf{x}^2)^s,$$

where $b_{x^2}^A(s)$ is the Bernstein–Sato polynomial, and $P(s)$ is a polynomial expression in s with coefficients consisting of differential operator of degree -2 on A . Using

J. Álvarez Montaner et al.

the description of such operators given above, it is clear that $s = \frac{1}{2}$ must be a root of $b_{x^2}^A(s)$.

Finally we remark that we can find rings where the Bernstein–Sato polynomial does not exist. The ring $A = \mathbb{C}[x, y, z]/\langle x^3 + y^3 + z^3 \rangle$ is such an example (see [1, Example 3.19] for details).

3. Differential Direct Summands and Differential Extensibility

3.1. Differential extensibility

In this section, we introduce the class of rings at the focal point of this paper, the *differentially extensible* subrings of polynomial rings. This notion was formalized to study *differential signature* [13], but was implicitly used earlier by Levasseur and Stafford [43] and Schwarz [71].

Definition 3.1 (Differential extensibility [13, Definition 6.1]). Consider a $\mathbb{K}$ -algebra homomorphism $\varphi: A \rightarrow T$.

- (1) If $\delta \in D_{A|\mathbb{K}}$, we say that a differential operator $\tilde{\delta} \in D_{T|\mathbb{K}}$ *extends* δ (or call $\tilde{\delta}$ an *extension* of δ) if $\varphi \circ \delta = \tilde{\delta} \circ \varphi$; i.e. the following diagram commutes.

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & T \\ \delta \downarrow & & \downarrow \tilde{\delta} \\ A & \xrightarrow{\varphi} & T \end{array}$$

If φ is simply an inclusion map, this is equivalent to the condition that $\tilde{\delta}|_A = \delta$.

- (2) The map φ is *differentially extensible* if for every $\delta \in D_{A|\mathbb{K}}$, there exists $\tilde{\delta} \in D_{T|\mathbb{K}}$ that extends δ .
- (3) The map φ is *differentially extensible with respect to the order filtration*, or *order-differentially extensible*, if for every nonnegative integer n , if $\delta \in D_{A|\mathbb{K}}^n$, then there exists $\tilde{\delta} \in D_{T|\mathbb{K}}^n$ that extends δ .
- (4) If $\mathbb{K}$ has positive characteristic, we say that φ is *differentially extensible with respect to the level filtration*, or *level-differentially extensible*, if for every non-negative integer e , given $\delta \in D_{A|\mathbb{K}}^{(e)}$, there exists $\tilde{\delta} \in D_{T|\mathbb{K}}^{(e)}$ that extends δ .

Remark 3.2. If $\varphi: A \rightarrow T$ is as above, and $\underline{s} = s_1, \dots, s_\ell$ are indeterminates, we say that an element $\tilde{\delta} \in D_{T|\mathbb{K}}[\underline{s}]$ *extends* an element $\delta \in D_{A|\mathbb{K}}[\underline{s}]$ if

$$(\varphi \otimes_{\mathbb{K}} \mathbb{K}[\underline{s}]) \circ \delta = \tilde{\delta} \circ (\varphi \otimes_{\mathbb{K}} \mathbb{K}[\underline{s}]).$$

We observe that an element $\tilde{\delta} \in D_{T|\mathbb{K}}[\underline{s}]$ extends $\delta \in D_{A|\mathbb{K}}[\underline{s}]$ if and only if each of its coefficients is an extension of the corresponding coefficient of δ .

Several maps of interest are differentially extensible.

Example 3.3 (37; 13, Proposition 6.4, Theorem 7.1). Let R be a polynomial ring over $\mathbb{K}$, and let G be a finite group acting by degree-preserving automorphisms

on R such that

- (1) $|G|$ is a unit in $\mathbb{K}$, and
- (2) No element of G fixes a hyperplane in $[R]_1$.

Then the inclusion $R^G \subseteq R$ is order-differentially extensible. Moreover, if $\mathbb{K}$ has positive characteristic, then the inclusion is also level-differentially extensible.

Example 3.4 (35, Proof of Theorem 3.4, 53; 13, Lemma 7.3). Let A be a pointed normal affine semigroup ring over a field $\mathbb{K}$ of characteristic zero. Then there is an embedding of A into a polynomial ring R as a monomial subalgebra, such that A is a direct summand of R , and for which the embedding is order-differentially extensible.

Lemma 3.5. *Suppose that $\mathbb{K}$ has characteristic $p > 0$, and d is a positive integer. Let $V \subseteq \mathbb{R}^d$ be a rational linear subspace, and $L \subseteq \mathbb{Z}^d$ be a lattice of finite index. Suppose that p does not divide the index $[\mathbb{Z}^d : L]$, and that L surjects onto $\mathbb{Z}$ via each coordinate map. Then the inclusion of normal semigroup rings $\mathbb{K}[\mathbb{N}^d \cap V \cap L] \subseteq \mathbb{K}[\mathbb{N}^d]$ is level-differentially extensible.*

Proof. We briefly describe our plan: First, we show that $\mathbb{K}[\mathbb{N}^d \cap V \cap L] \subseteq \mathbb{K}[\mathbb{N}^d \cap L]$ and $\mathbb{K}[\mathbb{N}^d \cap L] \subseteq \mathbb{K}[\mathbb{N}^d]$ are each level-differentially extensible inclusions. From here, it follows that the composition is also level-differentially extensible.

It can be shown that the inclusion $\mathbb{K}[\mathbb{N}^d \cap V \cap L] \subseteq \mathbb{K}[\mathbb{N}^d \cap L]$ is level-differentially extensible by the same argument used to compute differential signature [13, Proposition 6.6]. By our assumptions on L , $\mathbb{K}[\mathbb{N}^d \cap L] \subseteq \mathbb{K}[\mathbb{N}^d]$ is the inclusion of a ring of invariants of a finite diagonal abelian group. Any diagonal element g that fixes a hyperplane in the space of one-forms must fix every variable except one, say x_1 , and multiply that element by a t th root of unity. The invariants of g must then be contained in the subring $\mathbb{K}[x_1^t, x_2, \dots, x_d]$, contradicting the assumption that L surjects onto $\mathbb{Z}$ via the first coordinate map. Thus, the inclusion $\mathbb{K}[\mathbb{N}^d \cap L] \subseteq \mathbb{K}[\mathbb{N}^d]$ is level-differentially extensible by Example 3.3. $\square$

Remark 3.6. An embedding defined by Hsiao and Matusevich [35, Notation 1.2] satisfies the hypotheses of the previous lemma for all but finitely many characteristics p , and as a consequence, every normal affine semigroup ring of characteristic zero can be realized as a direct summand of a polynomial ring by a map that is order-differentially extensible, and such that the reduction modulo p of this map for all but finitely many p is level-differentially extensible (see Sec. 5 for details).

Example 3.7 ([43]). Suppose that $\mathbb{K}$ has characteristic zero. The generic determinantal, symmetric determinantal, and skew-symmetric determinantal rings are order-differentially extensible summands of polynomial rings.

In the lemmas that follow, we focus on studying whether a differential operator can be extended to the ring of differential operators of a localization.

J. Álvarez Montaner et al.

Remark 3.8. Let A be an algebra over $\mathbb{K}$. Then for every element f of A , the localization map $A \rightarrow A_f$ is differentially extensible with respect to the order filtration. Moreover, as we mentioned in the paragraph preceding Lemma 2.11, given $\delta \in D_{A|\mathbb{K}}$, there is a unique $\delta_f \in D_{A_f|\mathbb{K}}$ that extends δ [47, Theorem 2.2.10].

We remark that the usual $D_{A|\mathbb{K}}$ -module structure on A_f (see, e.g., [1, p. 303]) agrees with the rule $\delta \bullet v = \delta_f \bullet v$, where the right-hand side is the tautological $D_{A_f|\mathbb{K}}$ -action.

Notation 3.9. Take A and f as in Remark 3.8. Given $\delta \in D_{A|\mathbb{K}}$, we retain the notation from the remark, using δ_f to denote the unique extension of δ in $D_{A_f|\mathbb{K}}$ corresponding to the localization map $A \rightarrow A_f$.

Lemma 3.10. Let $\varphi: A \rightarrow T$ be a $\mathbb{K}$ -algebra homomorphism, and consider differential operators $\delta, \eta \in D_{A|\mathbb{K}}$. If $\tilde{\delta}$ and $\tilde{\eta}$ are differential operators in $D_{T|\mathbb{K}}$ that extend δ and η , respectively, then $[\tilde{\delta}, \tilde{\eta}] \in D_{T|\mathbb{K}}$ extends $[\delta, \eta] \in D_{A|\mathbb{K}}$.

Proof. Since $\varphi \circ \delta = \tilde{\delta} \circ \varphi$ and $\varphi \circ \eta = \tilde{\eta} \circ \varphi$, it is a brief and routine computation to verify that $\varphi \circ [\delta, \eta] = [\tilde{\delta}, \tilde{\eta}] \circ \varphi$. $\square$

Lemma 3.11. Let $A \subseteq T$ be $\mathbb{K}$ -algebras. For $f \in A$, if $\tilde{\delta} \in D_{T|\mathbb{K}}$ extends $\delta \in D_{A|\mathbb{K}}$, then $\tilde{\delta}_f$ extends δ_f .

Proof. We proceed by induction on the order, n , of δ . If $n = 0$, then δ is given by multiplication by an element of A , and an extension is given by multiplication by the same element.

Assume the claim for operators with order less than n , and consider $\delta \in D_{A|\mathbb{K}}$ with order n . For a nonnegative integer t , set $\rho = [\delta, f^t] \in D_{A|\mathbb{K}}$, and note that its order is strictly less than n . Let $\tilde{\rho} = [\tilde{\delta}, f^t] \in D_{T|\mathbb{K}}$, which by Lemma 3.10 is an extension of ρ . Then for any $h \in A$, it is straightforward to check that

$$\delta_f \bullet \frac{h}{f^t} = \frac{\delta \bullet h - \rho_f \bullet \frac{h}{f^t}}{f^t} \quad \text{and} \quad \tilde{\delta}_f \bullet \frac{h}{f^t} = \frac{\tilde{\delta} \bullet h - \tilde{\rho}_f \bullet \frac{h}{f^t}}{f^t}.$$

By our inductive hypothesis, $\rho_f \bullet \frac{h}{f^t} = \tilde{\rho}_f \bullet \frac{h}{f^t}$, while $\delta \bullet h = \tilde{\delta} \bullet h$ by assumption; thus, we conclude that $\delta_f \bullet \frac{h}{f^t} = \tilde{\delta}_f \bullet \frac{h}{f^t}$. $\square$

Applying Lemma 3.11 to the zero differential operator, we obtain the following.

Corollary 3.12. Let $A \subseteq T$ be $\mathbb{K}$ -algebras. If $\delta \in D_{T|\mathbb{K}}$ vanishes on A , then for every $f \in A$, $\delta_f \in D_{T_f|\mathbb{K}}$ vanishes on A_f .

3.2. Differential direct summands and differentially extensible summands

Let $A \subseteq T$ be an extension of Noetherian $\mathbb{K}$ -algebras such that A is a direct summand of T , with A -linear splitting $\beta: T \rightarrow A$. A theory of *differential direct summands* was previously developed to study the structure of D -modules over A [1].

The definition that we give here is slightly more general than the original, since we also consider $D_{A|\mathbb{K}}[s]$ -modules, though the idea is the same.

Definition 3.13 (Differential direct summand [1, Definition 3.2]). Let $A \subseteq T$ be an inclusion of $\mathbb{K}$ -algebras with A -linear splitting $\beta: T \rightarrow A$. Let $\underline{s}$ be a sequence of indeterminates (which may be empty). Recall that, for $\zeta \in D_{T|\mathbb{K}}$, the map $\beta \circ \zeta|_A: A \rightarrow A$ is an element of $D_{A|\mathbb{K}}$. By abuse of notation, for $\delta \in D_{T|\mathbb{K}}[s]$, we write $\beta \circ \delta|_A$ for the element of $D_{A|\mathbb{K}}[s]$ obtained from δ by applying $\beta \circ -|_A$ coefficientwise as a polynomial in $\underline{s}$.

We say that a $D_{A|\mathbb{K}}[s]$ -module M is a *differential direct summand* of a $D_{T|\mathbb{K}}[s]$ -module N if $M \subseteq N$ and there exists an A -linear splitting $\Theta: N \rightarrow M$, called a *differential splitting*, such that

$$\Theta(\delta \bullet v) = (\beta \circ \delta|_A) \bullet v$$

for every $\delta \in D_{T|\mathbb{K}}[s]$ and $v \in M$, where the action on the left-hand side is the $D_{T|\mathbb{K}}[s]$ -action, considering v as an element of N , and the action on the right-hand side is the $D_{A|\mathbb{K}}[s]$ -action.

Definition 3.14 (Morphism of differential direct summands [1, Definition 3.5]). Let $A \subseteq T$ be $\mathbb{K}$ -algebras such that A is a direct summand of T . Fix $D_{A|\mathbb{K}}[s]$ -modules M_1 and M_2 that are differential direct summands of $D_{T|\mathbb{K}}[s]$ -modules N_1 and N_2 , respectively, with differential splittings $\Theta_1: N_1 \rightarrow M_1$ and $\Theta_2: N_2 \rightarrow M_2$. We call $\phi: N_1 \rightarrow N_2$ a *morphism of differential direct summands* if $\phi \in \text{Hom}_{D_{T|\mathbb{K}}[s]}(N_1, N_2)$, $\phi(M_1) \subseteq M_2$, $\phi|_{M_1} \in \text{Hom}_{D_{A|\mathbb{K}}[s]}(M_1, M_2)$, and the following diagram commutes:

$$\begin{array}{ccccc} M_1 & \xrightarrow{\subseteq} & N_1 & \xrightarrow{\Theta_1} & M_1 \\ \downarrow \phi|_{M_1} & & \downarrow \phi & & \downarrow \phi|_{M_1} \\ M_2 & \xrightarrow{\subseteq} & N_2 & \xrightarrow{\Theta_2} & M_2 \end{array}$$

For simplicity of notation, we often write ϕ instead of $\phi|_{M_1}$.

Further, a complex $M_\bullet$ of $D_{A|\mathbb{K}}[s]$ -modules is called a *differential direct summand* of a complex $N_\bullet$ of $D_{T|\mathbb{K}}[s]$ -modules if each M_i is a differential direct summand of N_i , and each differential is a morphism of differential direct summands.

Remark 3.15. It is known that the property of being a differential direct summand is preserved under localization, taking kernels, and taking cokernels [1, Proposition 3.6, Lemma 3.7].

We now move on to develop the notion of *differential extension* in this context, which is an essential ingredient in the study of V -filtrations and multiplier ideals in later sections.

Definition 3.16 (Differentially extensible summand). Let $A \subseteq T$ be a differentially extensible inclusion of $\mathbb{K}$ -algebras, making A a direct summand of T . We

J. Álvarez Montaner et al.

say that a $D_{A|\mathbb{K}}[s]$ -module M is a *differentially extensible summand* of a $D_{T|\mathbb{K}}[s]$ -module N (or that $M \subseteq N$ is a *differentially extensible summand*) if the following conditions hold:

- (1) M is a differential direct summand of N .
- (2) For any $\delta \in D_{A|\mathbb{K}}[s]$ and any $\tilde{\delta} \in D_{T|\mathbb{K}}[s]$ that extends δ , one has $\tilde{\delta} \bullet m = \delta \bullet m$ for all $m \in M$.

Now we study the properties of differentially extensible summands. In particular, we see that the operations of localization, taking kernels, and taking cokernels all preserve the property of being a differentially extensible summand. This is important, due to our desire to work with local cohomology, which we use to investigate V -filtrations.

Lemma 3.17. *Let $A \subseteq T$ be a differentially extensible inclusion of $\mathbb{K}$ -algebras, with A a direct summand of T . If a $D_{A|\mathbb{K}}$ -module M is a differentially extensible summand of a $D_{T|\mathbb{K}}$ -module N , then for any $f \in A$, the localization M_f is a differentially extensible summand of N_f . Furthermore, the localization map is a morphism of differential direct summands.*

Proof. It has already been established that M_f is a differential direct summand of N_f [1, Proposition 3.6]. Given an extension $\tilde{\delta} \in D_{T|\mathbb{K}}$ of $\delta \in D_{A|\mathbb{K}}$, we have that $\tilde{\delta} \bullet \frac{v}{f^t} = \delta \bullet \frac{v}{f^t}$ for all $v \in M$, and all integers $t \geq 0$, by an argument directly analogous to the proof of Lemma 3.11, but with $v \in M$ replacing $h \in A$. $\square$

Lemma 3.18. *Let $A \subseteq T$ be a differentially extensible inclusion of $\mathbb{K}$ -algebras, with A a direct summand of T . Consider $D_{A|\mathbb{K}}$ -modules M_1 and M_2 that are differentially extensible summands of $D_{T|\mathbb{K}}$ -modules N_1 and N_2 . If $\phi: N_1 \rightarrow N_2$ is a morphism of differential direct summands, then $\text{Ker}(\phi|_{M_1}) \subseteq \text{Ker}(\phi)$, $\text{Im}(\phi|_{M_1}) \subseteq \text{Im}(\phi)$, and $\text{Coker}(\phi|_{M_1}) \subseteq \text{Coker}(\phi)$ are also differentially extensible summands.*

Proof. It is known that $\text{Ker}(\phi|_{M_1}) \subseteq \text{Ker}(\phi)$, $\text{Im}(\phi|_{M_1}) \subseteq \text{Im}(\phi)$, and $\text{Coker}(\phi|_{M_1}) \subseteq \text{Coker}(\phi)$, and that, moreover, each is a differential direct summand [1, Lemma 3.7].

Take an extension $\tilde{\delta} \in D_{T|\mathbb{K}}$ of $\delta \in D_{A|\mathbb{K}}$. Since $\tilde{\delta} \bullet v = \delta \bullet v$ for all $v \in M_1$, this holds for all $v \in \text{ker}(\phi|_{M_1}) \subseteq M_1$, and $\tilde{\delta} \bullet u = \delta \bullet u$ for all $u \in \text{Coker}(\phi|_{M_1})$. Moreover, since $\delta \bullet w = \tilde{\delta} \bullet w$ for all $w \in M_2$, the same holds for all $w \in \text{Im}(\phi|_{M_1}) \subseteq M_2$. $\square$

Proposition 3.19. *Let $A \subseteq T$ be a differentially extensible inclusion of $\mathbb{K}$ -algebras, with A a direct summand of T . Let M be a $D_{A|\mathbb{K}}$ -module, and N a $D_{T|\mathbb{K}}$ -module, with M a differentially extensible summand of N . Then for any ideal I of A , and any integer i , $H_I^i(M)$ is a differentially extensible summand of $H_{IT}^i(N)$.*

Proof. Fix generators $f = f_1, \dots, f_\ell \in A$ for I . The Čech-like complex $\check{C}^\bullet(f; M)$ defining the local cohomology modules of M with support in I is a differentially

extensible summand of the complex $\check{C}^\bullet(f; N)$ defining the local cohomology of N with support in IT : It is apparent that $\check{C}^j(f; M) \subseteq \check{C}^j(f; N)$ for all integers j , and each is a differentially extensible summand by Lemma 3.17; moreover, the fact that each differential is a morphism of differentially extensible summand holds by this same lemma, since each localization is such a morphism. Therefore, by Lemma 3.18, the inclusion of cokernels $H_I^i(M) \subseteq H_{IT}^i(N)$ is a differentially extensible summand. $\square$

We now study the notions of differentiable direct summands and differentially extensible summands in the context of the modules $M^A[\underline{f}^s]$ introduced in Sec. 2.3.

Setup 3.20. In the remainder of this section, $\mathbb{K}$ is a field of characteristic zero, and $A \subseteq T$ are Noetherian $\mathbb{K}$ -algebras such that A is a direct summand of T . We fix elements $\underline{f} = f_1, \dots, f_\ell \in A$, and assume that $M^A[\underline{f}^s]$ and $M^T[\underline{f}^s]$ have D -module structures compatible with specialization, as in Theorem 2.12. Recall that such a structure, if it exists, is unique, by the argument used in the proof of Theorem 2.12; moreover, that structure does exist for $\mathbb{K}$ -algebras that are either finitely generated or local and complete.

Theorem 3.21. *In the context of Setup 3.20, if $\beta: T \rightarrow A$ is an A -linear splitting, the $D_{A|\mathbb{K}}[s]$ -module $M^A[\underline{f}^s]$ is a differential direct summand of the $D_{T|\mathbb{K}}[s]$ -module $M^T[\underline{f}^s]$, with differential splitting*

$$\Theta = M^A[\underline{f}^s] \otimes_A \beta: M^T[\underline{f}^s] \rightarrow M^A[\underline{f}^s].$$

Proof. Given $\eta \in D_{T|\mathbb{K}}[s]$ we consider the differential operator $\delta := \beta \circ \eta|_A \in D_{A|\mathbb{K}}[s]$. Then, for any $v \in M^A[\underline{f}^s]$ we have to check that $\delta \bullet v = \Theta(\eta \bullet v)$. By Lemma 2.9, it suffices to show that $\varphi_t(\Theta(\eta \bullet v)) = \varphi_t(\delta \bullet v)$ for all $t \in \mathbb{Z}^\ell$. Indeed, for each such t we have

$$\begin{aligned} \varphi_t(\Theta(\eta \bullet v)) &= \beta_{f_1 \dots f_\ell}(\varphi_t(\eta \bullet v)) = (\beta \circ \eta(t)|_A) \bullet \varphi_t(v) \\ &= \delta(t) \bullet \varphi_t(v) = \varphi_t(\delta \bullet v), \end{aligned}$$

where we have used, in turn, the definition of Θ , the fact that $A_{f_1 \dots f_\ell}$ is a differential direct summand of $T_{f_1 \dots f_\ell}$ with differential splitting $\beta_{f_1 \dots f_\ell}$ [1, Proposition 3.6] and the definition of δ . $\square$

Theorem 3.22. *Under Setup 3.20, if $A \subseteq T$ is a differentially extensible inclusion, then the $D_{A|\mathbb{K}}[s]$ -module $M^A[\underline{f}^s]$ is a differentially extensible summand of the $D_{T|\mathbb{K}}[s]$ -module $M^T[\underline{f}^s]$.*

Proof. We proved in Theorem 3.21 that $M^A[\underline{f}^s]$ is a differential direct summand of $M^T[\underline{f}^s]$. To prove the second condition of differential extensibility, fix $v \in M^A[\underline{f}^s]$ and $\delta \in D_{A|\mathbb{K}}[s]$, and suppose $\tilde{\delta} \in D_{T|\mathbb{K}}[s]$ extends δ . Write $\delta \bullet v = g \underline{f}_1^{s_1} \dots \underline{f}_\ell^{s_\ell}$ and $\tilde{\delta} \bullet v = h \underline{f}_1^{s_1} \dots \underline{f}_\ell^{s_\ell}$, with $g \in A_{f_1 \dots f_\ell}[s]$ and $h \in T_{f_1 \dots f_\ell}[s]$. Since the D -module

J. Álvarez Montaner et al.

structures on $M^A[\underline{f}^s]$ and $M^T[\underline{f}^s]$ are compatible with specialization, for each $t \in \mathbb{Z}^\ell$ we have

$$\begin{aligned} g(t)f_1^{t_1} \cdots f_\ell^{t_\ell} &= \varphi_t(\delta \bullet v) = \delta(t) \bullet \varphi_t(v) = \tilde{\delta}(t) \bullet \varphi_t(v) \\ &= \varphi_t(\tilde{\delta} \bullet v) = h(t)f_1^{t_1} \cdots f_\ell^{t_\ell}, \end{aligned}$$

where the central equality follows from Lemma 3.11. This shows that $g(t) = h(t)$, for all $t \in \mathbb{Z}^\ell$, so $g = h$ by Lemma 2.9, and $\delta \bullet v = \tilde{\delta} \bullet v$. $\square$

3.3. Bernstein–Sato functional equation for direct summands

As we have done in Theorem 2.17, the existence of the formal Bernstein–Sato polynomial for direct summands can be proved by invoking the existence of the specialized version [1]. However we can provide a direct proof using the D -module structure of the module $M^A[\underline{f}^s]$ described in Theorem 2.12 and following the ideas used for the specialized version [1]. For completeness we work out the details in this section in full generality. Moreover, we address the issue of whether the Bernstein–Sato polynomial associated to a sequence of elements only depends on the ideal generated by these elements.

From now on we consider the following setup but we point out that all the results also hold for direct summands of formal power series rings.

Setup 3.23. In the remainder of this section, $\mathbb{K}$ is a field of characteristic zero, R is a polynomial ring over $\mathbb{K}$, and A is a $\mathbb{K}$ -subalgebra of R , such that A is a direct summand of R and that A is finitely generated over $\mathbb{K}$. Moreover, following Convention 2.13, given $\underline{f} = f_1, \dots, f_\ell \in A$, when referring to the $D_{A|\mathbb{K}}[\underline{s}]$ -module structure on $M^A[\underline{f}^s]$, we use the structure described in Theorem 2.12.

For the sake of clarity, we first present the more familiar case of principal ideals.

Theorem 3.24. *Under Setup 3.23, given $f, g \in A$, there exists $\delta \in D_{A|\mathbb{K}}[s]$, and a nonzero polynomial $b(s) \in \mathbb{Q}[s]$, such that the functional equation*

$$\delta \bullet fgf^s = b(s)gf^s$$

holds in $M^A[\underline{f}^s]$.

Proof. There exist $\eta \in D_{R|\mathbb{K}}[s]$ and a nonzero polynomial $b(s) \in \mathbb{Q}[s]$ such that

$$\eta \bullet fgf^s = b(s)gf^s$$

in $M^R[\underline{f}^s]$, as in Definition 2.7. By applying the splitting $\Theta: M^R[\underline{f}^s] \rightarrow M^A[\underline{f}^s]$ induced by the splitting β making A a direct summand of R , we obtain that

$$b(s)gf^s = \Theta(b(s)gf^s) = \Theta(\eta \bullet fgf^s) = (\beta \circ \eta|_A) \bullet fgf^s$$

in $M^A[\underline{f}^s]$. Taking $\delta = \beta \circ \eta|_A \in D_{A|\mathbb{K}}[s]$, we obtain the desired equation. $\square$

Example 3.25. Let $A = \frac{\mathbb{C}[u,v,w]}{\langle u^3-vw \rangle}$. We determine a functional equation for the element $uv \in A$. To do this, we realize $A \subseteq R = \mathbb{C}[x, y]$ via the inclusion given by $f(u) = xy$, $f(v) = x^3$, $f(w) = y^3$. Observe that the image of A under f is the ring of polynomials invariant under the action of the cyclic group G generated by g , where $g(x) = e^{2\pi i/3}x$ and $g(y) = e^{-2\pi i/3}y$. Since g does not fix any one-forms in R , the inclusion map f is differentially extensible by Example 3.3. From here we find a functional equation for $f(uv) = x^4y$ in R , namely

$$\frac{1}{4^4} \partial_x^4 \partial_y \bullet x^4 y (x^4 y)^s = (s+1)^2 \left(s + \frac{3}{4}\right) \left(s + \frac{1}{2}\right) \left(s + \frac{1}{4}\right) (x^4 y)^s.$$

The existence of such an operator comes from the standard theory of Bernstein–Sato polynomials. We then observe that the differential operator $\frac{1}{4^4} \partial_x^4 \partial_y \in D_{R|\mathbb{C}}[s]$ sends G -invariant polynomials to G -invariant polynomials, and is thus a differential operator on A . This yields the functional equation we seek.

Thanks to Theorem 3.24, the following definition is well founded for a direct summand of a polynomial ring.

Definition 3.26 (Relative Bernstein–Sato polynomial of an element of a direct summand of a polynomial ring). Given $A \subseteq R$ and $f, g \in A$ as in Theorem 3.24, we call the monic polynomial of smallest degree satisfying the conclusion of this theorem the *Bernstein–Sato polynomial of f relative to g* , which we denote $b_{f,g}^A(s)$. If $g = 1$, we call this polynomial the *Bernstein–Sato polynomial of f* , and denote it $b_f^A(s)$.

We now present the Bernstein–Sato polynomial $b_{f,g}^A(s)$ associated to a set of nonzero generators $\underline{f} = f_1, \dots, f_\ell$ of an ideal I of A , relative to an element $g \in A$.

Theorem 3.27. *Under Setup 3.23, given $\underline{f} = f_1, \dots, f_\ell \in A$ and $g \in A$, there exists a nonzero polynomial $b(s) \in \mathbb{Q}[s]$ such that*

$$b(s_1 + \dots + s_\ell) g \mathbf{f}_1^{s_1} \dots \mathbf{f}_\ell^{s_\ell}$$

is in the $D_{A|\mathbb{K}}[s]$ -submodule of $M^A[\underline{\mathbf{f}}^s]$ generated by all

$$\prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \dots f_\ell^{c_\ell} g \mathbf{f}_1^{s_1} \dots \mathbf{f}_\ell^{s_\ell},$$

where $c = (c_1, \dots, c_\ell)$ runs over the elements of $\mathbb{Z}^\ell$ such that $|c| = c_1 + \dots + c_\ell = 1$, and $\text{nsupp}(c) = \{i : c_i < 0\}$.

Proof. Let N denote the $D_{A|\mathbb{K}}[s]$ -submodule of $M^A[\underline{\mathbf{f}}^s]$ described in the statement of the theorem. By the existence results given for Bernstein–Sato polynomials for varieties in smooth varieties [17, Sec. 2.10] (see also [16, Definition 2.2]), there exists a nonzero polynomial $b(s) \in \mathbb{Q}[s]$ such that $b(s_1 + \dots + s_\ell) g \mathbf{f}_1^{s_1} \dots \mathbf{f}_\ell^{s_\ell}$ is in the

J. Álvarez Montaner et al.

$D_{R|\mathbb{K}}[s]$ -submodule of $M^R[\underline{f}^s]$ generated by all

$$\prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \cdots f_\ell^{c_\ell} g f_1^{s_1} \cdots f_\ell^{s_\ell},$$

where $c = (c_1, \dots, c_\ell)$ runs over the elements of $\mathbb{Z}^\ell$ such that $|c| = 1$. Now, applying the splitting $\Theta: M^R[\underline{f}^s] \rightarrow M^A[\underline{f}^s]$, we obtain that

$$b(s_1 + \cdots + s_\ell) g f_1^{s_1} \cdots f_\ell^{s_\ell} \in N. \quad \square$$

As in the case of a single element f , we have justified the following definition in the case of a direct summand of a polynomial ring over a field of characteristic zero.

Definition 3.28 (Relative Bernstein–Sato polynomial of a sequence of elements in a direct summand of a polynomial ring). We call the unique monic polynomial $b(s) \in \mathbb{Q}[s]$ of smallest degree satisfying the conclusion of Theorem 3.27 the *Bernstein–Sato polynomial of $\underline{f}$ relative to g* , and we denote it $b_{\underline{f},g}^A(s)$. If $g = 1$, we call this the *Bernstein–Sato polynomial of $\underline{f}$* , and we denote it $b_{\underline{f}}^A(s)$.

It follows from the proof of Theorem 3.27 that, for a direct summand $A \subseteq R$, the Bernstein–Sato polynomial $b_{\underline{f},g}^A(s)$ divides $b_{\underline{f},g}^R(s)$ (see [1, Theorem 3.14] for the specialized version). Under the extra condition that $A \subseteq R$ is differentially extensible, these two polynomials coincide as it was shown for the specialized version [13, Theorem 6.11].

Theorem 3.29. *Consider Setup 3.23 under the extra condition that the inclusion $A \subseteq R$ is differentially extensible. Given $\underline{f} = f_1, \dots, f_\ell \in A$ and $g \in A$, we have that $b_{\underline{f},g}^A(s) = b_{\underline{f},g}^R(s)$.*

Proof. From the proof of Theorem 3.27, we obtain that $b_{\underline{f},g}^A(s)$ divides $b_{\underline{f},g}^R(s)$. We now prove that $b_{\underline{f},g}^R(s)$ divides $b_{\underline{f},g}^A(s)$. Choose $\delta_c \in D_{A|\mathbb{K}}[s]$ for which

$$\begin{aligned} & b_{\underline{f},g}^A(s_1 + \cdots + s_\ell) g f_1^{s_1} \cdots f_\ell^{s_\ell} \\ &= \sum_{\substack{c \in \mathbb{Z}^\ell \\ |c|=1}} \delta_c \cdot \prod_{i \in \text{nsupp}(c)} \binom{s_i}{-c_i} f_1^{c_1} \cdots f_\ell^{c_\ell} g f_1^{s_1} \cdots f_\ell^{s_\ell}. \end{aligned}$$

The above equation remains true after replacing each $\delta_c \in D_{A|\mathbb{K}}[s]$ with an extension $\tilde{\delta}_c \in D_{R|\mathbb{K}}[s]$, justifying our claim. $\square$

Using a mild generalization of a recent result by Mustață [59] we can prove that the Bernstein–Sato polynomial does not depend on the choice of generators for an ideal $I \subseteq A$, justifying the notation $b_{I,g}^A(s)$, or $b_I^A(s)$ if $g = 1$.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

Theorem 3.30 (Musta ). *Let A be an algebra over a field $\mathbb{K}$ of characteristic zero, that is finitely generated over $\mathbb{K}$. Assume that A is a direct summand of a polynomial ring over $\mathbb{K}$. Given nonzero $\underline{f} = f_1, \dots, f_\ell \in A$ and $g \in A$, set $h = y_1 f_1 + \dots + y_\ell f_\ell \in A[y_1, \dots, y_\ell]$, where $y_1, \dots, y_\ell$ are indeterminates. Then*

$$b_{\underline{f},g}^A(s) = \frac{b_{h,g}^{A[y_1, \dots, y_\ell]}(s)}{(s+1)}.$$

Furthermore, $b_{\underline{f},g}^A(s)$ only depends on the ideal generated by $\underline{f}$.

The proof of this result is essentially the same as the one presented in work previously mentioned [59, Theorem 1.1 and Remark 2.1], so we omit it. We only note that the hypothesis that A (and hence also $A[y_1, \dots, y_\ell]$) is a direct summand of a polynomial ring guarantees the existence of the relative Bernstein–Sato polynomial $b_{h,g}^{A[y_1, \dots, y_\ell]}(s)$, and that $(s+1)$ is a factor of this polynomial, since h involves some variables that g does not.

Theorems 3.27 and 3.30 allow us to define the (relative) Bernstein–Sato polynomial of an ideal in a direct summand of a polynomial ring.

Definition 3.31 (Relative Bernstein–Sato polynomial in a direct summand of a polynomial ring). Let A be an algebra over a field $\mathbb{K}$ of characteristic zero that is finitely generated over $\mathbb{K}$. Assume that A is a direct summand of a polynomial ring over $\mathbb{K}$. Let I and g be an ideal and an element of A . Given generators $\underline{f} = f_1, \dots, f_\ell$ for I , we call the polynomial $b_{\underline{f},g}^A \in \mathbb{Q}[s]$ the *Bernstein–Sato polynomial of I relative to g* , and denote it $b_{I,g}^A(s)$. If $g = 1$, we call this the *Bernstein–Sato polynomial of I* , and denote it $b_I^A(s)$.

4. V -Filtrations

The theory of the V -filtration, originally defined by Malgrange [46] and Kashiwara [40], is a central object in the study of D -modules over a smooth variety. Our goal in this section is to extend this theory to the setting of a differentially extensible summand of a polynomial ring. In the following section, we apply this newly-developed theory to the study of multiplier ideals.

Suppose that $\mathbb{K}$ has characteristic zero. Let $R = \mathbb{K}[\underline{x}] = \mathbb{K}[x_1, \dots, x_d]$ be a polynomial ring over $\mathbb{K}$ and set $X = \text{Spec}(R)$. Let $I \subseteq R$ be an ideal generated by $\underline{f} = f_1, \dots, f_\ell \in R$. We may assume that I defines a smooth subvariety, by considering the graph embedding $i_{\underline{f}}: X \rightarrow X \times \mathbb{A}^\ell$ mapping $x \mapsto (x, f_1(x), \dots, f_\ell(x))$, and then replacing X by $X \times \mathbb{A}^\ell$ and each f_i by the projection onto the i th coordinate of $\mathbb{A}^\ell$. Therefore, if $\underline{t} = t_1, \dots, t_\ell$ is a system of coordinates of $\mathbb{A}^\ell$, we may simply assume that I is $\langle \underline{t} \rangle = \langle t_1, \dots, t_\ell \rangle \subseteq R[t_1, \dots, t_\ell]$.

The V -filtration along $\langle \underline{t} \rangle$ on the ring of $\mathbb{K}$ -linear differential operators $D_{R[\underline{t}]|\mathbb{K}}$ is obtained by assigning degree 0 to the elements of $D_{R|\mathbb{K}}$, degree 1 to the indeterminates $t_1, \dots, t_\ell$, and degree -1 to the partial derivatives $\partial_{t_1}, \dots, \partial_{t_\ell}$. Equivalently,

J. Álvarez Montaner et al.

a differential operator δ lies in $V^i D_{R[t]|\mathbb{K}}$ if and only if $\delta \bullet \langle t \rangle^j \subseteq \langle t \rangle^{j+i}$ for all $j \in \mathbb{Z}$, where we use the convention that $\langle t \rangle^j = R[t]$ for negative exponents j .

For a given $D_{R|\mathbb{K}}$ -module N , one considers its direct image with respect to the graph embedding

$$i_{f*}(N) := N \otimes_{\mathbb{K}} \mathbb{K}[\partial_{t_1}, \dots, \partial_{t_\ell}],$$

which is a $D_{R[t]|\mathbb{K}}$ -module isomorphic to the local cohomology module

$$H_{\langle t-f \rangle}^\ell(N[t]),$$

where $\langle t-f \rangle$ denotes the ideal $\langle t_1 - f_1, \dots, t_\ell - f_\ell \rangle$ of $R[t]$. If N is a *regular holonomic* $D_{R|\mathbb{K}}$ -module with *quasi-unipotent monodromy*, then there exists a V -filtration along $\langle t \rangle$ on $i_{f*}(N)$, which is a decreasing filtration $V_{\langle t \rangle}^\alpha i_{f*}(N)$ indexed over the rational numbers satisfying certain conditions (see Definition 4.3). The existence of such a filtration is a result of Malgrange [46] and Kashiwara [40], and is essentially equivalent to the existence of relative Bernstein–Sato polynomials in a generalized sense. We do not go into detail on the definition of regular holonomic modules with quasi-unipotent monodromy, so we encourage the interested reader to take a look at [46] and [40]; we only point out that the ring R itself, and the local cohomology modules of the form $H_I^i(R)$, satisfy this property.

Finally, one defines the V -filtration along f of the $D_{R|\mathbb{K}}$ -module N as

$$V_{\langle f \rangle}^\alpha N := V_{\langle t \rangle}^\alpha i_{f*}(N) \cap N.$$

It is known that the V -filtration only depends on the ideal generated by f , and not on the choice of generators [17, Proposition 2.6].

4.1. V -filtrations for nonregular rings

In order to develop a theory of V -filtrations for a not-necessarily-regular ring, we follow the same ideas as in the smooth case and define them using a set of axioms.

Definition 4.1 (*V -filtration on $D_{T[t]|\mathbb{K}}$*). Suppose that $\mathbb{K}$ has characteristic zero. Let T be a Noetherian commutative $\mathbb{K}$ -algebra, and let $T[t] = T[t_1, \dots, t_\ell]$ be a polynomial ring over T . The V -filtration along the ideal $\langle t \rangle = \langle t_1, \dots, t_\ell \rangle$ on the ring of differential operators $D_{T[t]|\mathbb{K}}$ is the filtration indexed by integers $i \in \mathbb{Z}$ defined by

$$V_{\langle t \rangle}^i D_{T[t]|\mathbb{K}} = \{ \delta \in D_{T[t]|\mathbb{K}} : \delta \bullet \langle t \rangle^j \subseteq \langle t \rangle^{j+i} \text{ for all } j \in \mathbb{Z} \},$$

where we use the convention that $\langle t \rangle^j = T[t]$ for integers $j \leq 0$.

Remark 4.2. As a graded $D_{T|\mathbb{K}}$ -module, we have the following description:

$$V_{\langle t \rangle}^i D_{T[t]|\mathbb{K}} = \bigoplus_{\substack{a, b \in \mathbb{N}^\ell \\ |a| - |b| \geq i}} D_{T|\mathbb{K}} \cdot t_1^{a_1} \cdots t_\ell^{a_\ell} \partial_{t_1}^{b_1} \cdots \partial_{t_\ell}^{b_\ell},$$

where, given $u = (u_1, \dots, u_\ell) \in \mathbb{N}^\ell$, we denote $|u| = u_1 + \dots + u_\ell$.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

The V -filtration along the ideal $\langle t \rangle = \langle t_1, \dots, t_\ell \rangle$ on a $D_{T[t]|\mathbb{K}}$ -module M is defined axiomatically as follows.

Definition 4.3 (*V -filtration on a $D_{T[t]|\mathbb{K}}$ -module along indeterminates*). Let M be a $D_{T[t]|\mathbb{K}}$ -module. A V -filtration on M along the ideal $\langle t \rangle = \langle t_1, \dots, t_\ell \rangle$ is a decreasing filtration $\{V_{\langle t \rangle}^\alpha M\}_\alpha$ on M , indexed by $\alpha \in \mathbb{Q}$, satisfying the following conditions.

- (1) For all $\alpha \in \mathbb{Q}$, $V_{\langle t \rangle}^\alpha M$ is a Noetherian $V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}$ -submodule of M .
- (2) The union of the $V_{\langle t \rangle}^\alpha M$, over all $\alpha \in \mathbb{Q}$, is M .
- (3) The filtration is left-semicontinuous and discrete. More precisely, there exists some $m \in \mathbb{N}$ such that for every $n \in \mathbb{Z}$ and all $\alpha \in (\frac{n-1}{m}, \frac{n}{m}]$, $V_{\langle t \rangle}^\alpha M = V_{\langle t \rangle}^{n/m} M$.
- (4) For all $\alpha \in \mathbb{Q}$ and all $1 \leq i \leq \ell$,

$$t_i \bullet V_{\langle t \rangle}^\alpha M \subseteq V_{\langle t \rangle}^{\alpha+1} M \quad \text{and} \quad \partial_{t_i} \bullet V_{\langle t \rangle}^\alpha M \subseteq V_{\langle t \rangle}^{\alpha-1} M,$$

i.e. the filtration is compatible with the V -filtration on $D_{T[t]|\mathbb{K}}$.

- (5) For all $\alpha \gg 0$, $\sum_{i=1}^\ell (t_i \bullet V_{\langle t \rangle}^\alpha M) = V_{\langle t \rangle}^{\alpha+1} M$.
- (6) For all $\alpha \in \mathbb{Q}$,

$$\sum_{i=1}^\ell \partial_{t_i} t_i - \alpha$$

acts nilpotently on $V_{\langle t \rangle}^\alpha M / (\bigcup_{\gamma > \alpha} V_{\langle t \rangle}^\gamma M)$.

As in the case of polynomial rings, the V -filtration is unique in this broader context, provided it exists. The proof of this fact is analogous to that of the original result in the case that T is a polynomial ring over $\mathbb{K}$ (see [15, Proposition 1.3]), so we omit it.

Proposition 4.4. *Let M be a finitely generated $D_{T[t]|\mathbb{K}}$ -module. If a V -filtration on M along $\langle t \rangle$ exists, then it is unique.*

Our goal is to prove the existence of the V -filtration for differentially extensible summands of polynomial rings. For this reason, we introduce certain auxiliary Rees algebras that lead to an equivalent way of defining V -filtrations.

Definition 4.5. Suppose that $\mathbb{K}$ has characteristic zero. Let T be a Noetherian commutative $\mathbb{K}$ -algebra, and let $T[t] = T[t_1, \dots, t_\ell]$ be a polynomial ring over T . Given an auxiliary indeterminate w , we define the following Rees algebras:

$$\begin{aligned} \mathfrak{V}_{\langle t \rangle} D_{T[t]|\mathbb{K}} &:= \bigoplus_{i \in \mathbb{Z}} (V_{\langle t \rangle}^i D_{T[t]|\mathbb{K}}) w^i \\ &= D_{T|\mathbb{K}}[t_1 w, \dots, t_\ell w, \partial_{t_1} w^{-1}, \dots, \partial_{t_\ell} w^{-1}] \\ \mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} &:= \bigoplus_{i \geq 0} (V_{\langle t \rangle}^i D_{T[t]|\mathbb{K}}) w^i \oplus \bigoplus_{i < 0} (V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}) w^i \\ &= D_{T|\mathbb{K}}[w^{-1}, t_i w, \partial_{t_i} t_j \mid 1 \leq i, j \leq \ell] \end{aligned}$$

J. Álvarez Montaner et al.

Moreover, given a $D_{T[t]|\mathbb{K}}$ -module M , for $\alpha \in \mathbb{Q}$,

$$\mathfrak{V}_{\langle t \rangle}^\alpha M := \bigoplus_{i \in \mathbb{Z}} \left(V_{\langle t \rangle}^{\alpha+i} M \right) w^i.$$

In the polynomial ring case, these algebras are Noetherian.

Lemma 4.6. *Suppose that $\mathbb{K}$ has characteristic zero. Let $R = \mathbb{K}[x] = \mathbb{K}[x_1, \dots, x_d]$, and consider the polynomial ring $R[t] = R[t_1, \dots, t_\ell]$ over R . Then $V_{\langle t \rangle}^0 D_{R[t]|\mathbb{K}}$, $\mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}}$, and $\mathfrak{V}_{\langle t \rangle}^+ D_{R[t]|\mathbb{K}}$ are Noetherian rings.*

Proof. We aim to show that each ring admits a filtration for which the associated graded ring is commutative and Noetherian. Let $\mathcal{F}$ denote the order filtration on $D_{R[t]|\mathbb{K}}$, so that

$$\mathrm{gr}_{\mathcal{F}}(D_{R[t]|\mathbb{K}}) \cong \mathbb{K}[\underline{x}, \underline{y}, \underline{t}, \underline{u}],$$

where $\underline{y} = y_1, \dots, y_d$ and $\underline{u} = u_1, \dots, u_\ell$ correspond to the images of the partial derivatives $\partial_{x_1}, \dots, \partial_{x_d}$ and $\partial_{t_1}, \dots, \partial_{t_\ell}$. Let $\mathcal{G}_n = (V_{\langle t \rangle}^0 D_{R[t]|\mathbb{K}}) \cap \mathcal{F}_n$, and consider the ring inclusion

$$\varphi: \mathrm{gr}_{\mathcal{G}}(V_{\langle t \rangle}^0 D_{R[t]|\mathbb{K}}) \rightarrow \mathrm{gr}_{\mathcal{F}}(D_{R[t]|\mathbb{K}}).$$

Set $a = (a_1, \dots, a_\ell)$ and $b = (b_1, \dots, b_\ell)$. We have that

$$\begin{aligned} \mathrm{gr}_{\mathcal{G}}(V_{\langle t \rangle}^0 D_{R[t]|\mathbb{K}}) &\cong \mathrm{Im}(\varphi) \cong \mathbb{K}[\underline{x}, \underline{y}, t_1^{a_1} \dots t_\ell^{a_\ell} u_1^{b_1} \dots u_\ell^{b_\ell}]_{\{|a| \geq |b|\}} \\ &\cong \mathbb{K}[\underline{x}, \underline{y}, \underline{t}, t_i u_j]_{\{i, j\}}, \end{aligned}$$

which is a finitely generated $\mathbb{K}$ -algebra, and so, a Noetherian ring. We conclude that $V_{\langle t \rangle}^0 D_{R[t]|\mathbb{K}}$ is left- and right-Noetherian since $\mathcal{F}_0 \cong R[t]$.

Now, consider the subring $\mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}}$ of $D_{R[t]|\mathbb{K}}[w]$, which is $\mathbb{Z}$ -graded as a polynomial ring in w . We refine this to a $\mathbb{Z} \times \mathbb{N}$ grading by taking the filtration

$$\mathcal{G}'_j = \{\delta \in \mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}} : \delta \in \mathcal{F}_j \otimes_{D_{R[t]|\mathbb{K}}} D_{R[t]|\mathbb{K}}[w]\}.$$

We obtain an inclusion $\varphi: \mathrm{gr}_{\mathcal{G}'}(\mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}}) \rightarrow \mathrm{gr}_{\mathcal{F}}(D_{R[t]|\mathbb{K}})[w]$. With the same coordinates that we used for $\mathrm{gr}_{\mathcal{F}}(D_{R[t]|\mathbb{K}})$ above

$$\begin{aligned} \mathrm{gr}_{\mathcal{G}'}(\mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}}) &\cong \mathrm{Im}(\varphi) \cong \mathbb{K}[\underline{x}, \underline{y}, t_1^{a_1} \dots t_\ell^{a_\ell} u_1^{b_1} \dots u_\ell^{b_\ell} w^j]_{\{|a| \geq |b|+j\}} \\ &\cong \mathbb{K}[\underline{x}, \underline{y}, t_1 w, \dots, t_\ell w, u_1 w^{-1}, \dots, u_\ell w^{-1}], \end{aligned}$$

which is again a finitely generated $\mathbb{K}$ -algebra, and hence Noetherian. We conclude that $\mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}}$ is left- and right-Noetherian.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

Finally, consider $\mathfrak{V}_{\langle t \rangle}^+ D_{R[t]|\mathbb{K}} \subseteq \mathfrak{V}_{\langle t \rangle} D_{R[t]|\mathbb{K}} \subseteq D_{R[t]|\mathbb{K}}[w]$. We have that

$$\begin{aligned} \mathrm{gr}_{G'}(\mathfrak{V}^+ D_{R[t]|\mathbb{K}}) \\ &\cong \mathbb{K}[x, y, t_1^{a_1} \cdots t_\ell^{a_\ell} u_1^{b_1} \cdots u_\ell^{b_\ell} w^j]_{\{|a| \geq |b| + \max\{j, 0\}\}} \\ &\cong \mathbb{K}[x, y, t_1 w, \dots, t_\ell w, w^{-1}, t_i u_j]_{\{i, j\}}, \end{aligned}$$

and by the same argument above, we are done. $\square$

Using the algebras introduced in Definition 4.5, we can give another characterization of the stabilization conditions appearing in the definition of the V -filtration.

Lemma 4.7. *Suppose that $\mathbb{K}$ has characteristic zero. Let T be a commutative Noetherian $\mathbb{K}$ -algebra, and let $T[t] = T[t_1, \dots, t_\ell]$ be a polynomial ring over T . Given a $D_{T[t]|\mathbb{K}}$ -module M , a decreasing filtration $\{V_{\langle t \rangle}^\alpha M\}_\alpha$ on M indexed by $\alpha \in \mathbb{Q}$ satisfies conditions (1) and (5) of Definition 4.3 if*

- (a) $\mathfrak{V}_{\langle t \rangle}^\alpha M$ is a Noetherian $\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}}$ -module for all $\alpha \in \mathbb{Q} \cap [0, 1)$.

Proof. We first show that (a) implies (1). Let $\{M_j\}$ an ascending chain of $V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}$ -submodules of $V_{\langle t \rangle}^\alpha M$. We consider the induced chain $\{\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} M_j\}$. By (a), there exists an integer a such that for every $j > a$, we have $\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} M_j = \mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} M_a$. By the grading, we obtain that $M_j = M_a$ as $V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}$ -modules. We conclude that $V_{\langle t \rangle}^\alpha M$ is a Noetherian left $V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}$ -submodule of M .

We now show that (a) implies (5). Let $\mathfrak{V}_{\langle t \rangle}^\alpha M = \sum_{i=1}^s \mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} m_i w^{d_i}$ as above. For $d > \max\{d_1, \dots, d_s\}$, using the previous equation, we obtain that

$$\begin{aligned} V_{\langle t \rangle}^{\alpha+d+1} M &= \sum V_{\langle t \rangle}^{d-d_i+1} D_{T[t]|\mathbb{K}} m_i \\ &= \sum \langle t \rangle \bullet V_{\langle t \rangle}^{d-d_i} D_{T[t]|\mathbb{K}} m_i = \langle t \rangle \bullet V_{\langle t \rangle}^{\alpha+d} M. \end{aligned} \quad \square$$

4.2. V -filtrations for differentially extensible summands

The main result of this section is the existence of V -filtrations for differentially extensible summands.

Theorem 4.8. *Let T be a commutative Noetherian ring containing a field $\mathbb{K}$ of characteristic zero. Let A be a $\mathbb{K}$ -subalgebra of T such that A is a direct summand of T , and for which the inclusion $A \subseteq T$ is differentially extensible. Let M be a $D_{A[t]|\mathbb{K}}$ -module that is a differentially extensible summand of a $D_{T[t]|\mathbb{K}}$ -module N . If $V_{\langle t \rangle}^\alpha N$ is a V -filtration on N along the ideal $\langle t \rangle = \langle t_1, \dots, t_\ell \rangle$ over $T[t]$, then $W_{\langle t \rangle}^\alpha M := V_{\langle t \rangle}^\alpha N \cap M$ is a V -filtration on M along $\langle t \rangle$ over $A[t]$.*

Proof. We show that $W_{\langle t \rangle}^\alpha M$ is a V -filtration for M by verifying Conditions (2), (3), (4), and (6) of Definition 4.3, and Condition (a) of Lemma 4.7.

J. Álvarez Montaner et al.

The fact that Condition (2) holds follows from the fact that $V^\alpha N$ is exhaustive.

To see that Condition (3) holds, observe that since $V_{\langle t \rangle}^\alpha N$ is a V -filtration for N , there exists some $m \in \mathbb{N}$ for which $V_{\langle t \rangle}^\alpha N$ is constant on each interval of the form $(\frac{n-1}{m}, \frac{n}{m}]$, where $n \in \mathbb{Z}$. Consequently, $W_{\langle t \rangle}^\alpha M = V_{\langle t \rangle}^\alpha N \cap M$ is constant on each such interval.

Toward verifying Condition (4), we first show that $W_{\langle t \rangle}^\alpha M$ is a $V_{\langle t \rangle}^0 D_{A[t]|\mathbb{K}}$ -submodule of M . Let

$$\delta = \sum_{|a| \geq |b|} \delta_{a,b} t^a \partial^b \in V_{\langle t \rangle}^0 D_{A[t]|\mathbb{K}},$$

where $\delta_{a,b} \in D_{A|\mathbb{K}}$, $t^a = t_1^{a_1} \cdots t_\ell^{a_\ell}$, $\partial^b = \partial_{t_1}^{b_1} \cdots \partial_{t_\ell}^{b_\ell}$, and the sum is taken over finitely many $a, b \in \mathbb{N}^\ell$. For each $\delta_{a,b}$, there exists $\tilde{\delta}_{a,b} \in D_{T|\mathbb{K}}$ such that $\tilde{\delta}_{a,b}|_A = \delta_{a,b}$; set

$$\tilde{\delta} = \sum_{|a| \geq |b|} \tilde{\delta}_{a,b} t^a \partial^b.$$

Given $w \in W_{\langle t \rangle}^\alpha M$, we have $\delta \bullet w = \tilde{\delta} \bullet w$, as M is a differentially extensible summand of N . Since $\tilde{\delta} \in V_{\langle t \rangle}^0 D_{T[t]|\mathbb{K}}$, we obtain that $\tilde{\delta} \bullet w \in V_{\langle t \rangle}^\alpha N$, hence $\delta \bullet w \in W_{\langle t \rangle}^\alpha M$.

To conclude that Condition (4) holds, it suffices to show that $t^a \partial^b \bullet W_{\langle t \rangle}^\alpha M \subseteq W_{\langle t \rangle}^{\alpha+|a|-|b|} M$, but this follows from the string of containments

$$\begin{aligned} t^a \partial^b \bullet W_{\langle t \rangle}^\alpha M &= t^a \partial^b \bullet (V_{\langle t \rangle}^\alpha N \cap M) \subseteq (t^a \partial^b \bullet V_{\langle t \rangle}^\alpha N) \cap M \\ &\subseteq (V_{\langle t \rangle}^{\alpha+|a|-|b|} N) \cap M = W_{\langle t \rangle}^{\alpha+|a|-|b|} M. \end{aligned}$$

From the definition, it is clear that Condition (6) holds, since

$$\frac{W_{\langle t \rangle}^\alpha M}{\bigcup_{\beta > \alpha} W_{\langle t \rangle}^\beta M} = \frac{V_{\langle t \rangle}^\alpha N \cap M}{\bigcup_{\beta > \alpha} V_{\langle t \rangle}^\beta N \cap M} \hookrightarrow \frac{V_{\langle t \rangle}^\alpha N}{\bigcup_{\beta > \alpha} V_{\langle t \rangle}^\beta N},$$

and $\sum_{i=1}^r \partial_{t_i} t_i - \alpha$ acts nilpotently on the module on the right-hand side.

Finally, to verify that Condition (a) holds, we show that the following stronger condition holds: $\mathfrak{W}_{\langle t \rangle}^\alpha M$ is a left-Noetherian $\mathfrak{V}^+ D_{A[t]|\mathbb{K}}$ -module. It suffices to show that if $M_1, M_2 \subseteq \mathfrak{W}_{\langle t \rangle}^\alpha M$ are $\mathfrak{V}_{\langle t \rangle}^\alpha D_{A[t]|\mathbb{K}}$ -modules such that

$$(\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}})M_1 = (\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}})M_2,$$

then $M_1 = M_2$, because $\mathfrak{W}_{\langle t \rangle}^\alpha N$ is a left-Noetherian $\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}}$ -module. We denote by $\Theta: N \rightarrow M$ a differential splitting. Let $w \in M_1$. Since $\mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} M_1 = \mathfrak{V}_{\langle t \rangle}^+ D_{T[t]|\mathbb{K}} M_2$, there exists $\xi_1, \dots, \xi_s \in D_{T|\mathbb{K}}$ and $w_1, \dots, w_s \in M_2$ such that $w = \xi_1 \bullet w_1 + \cdots + \xi_s \bullet w_s$. We apply Θ to both sides to obtain

$$\begin{aligned} w' &= \Theta(w') = \Theta(\xi_1 \bullet w_1 + \cdots + \xi_s \bullet w_s) \\ &= \Theta(\xi_1 \bullet w_1) + \cdots + \Theta(\xi_s \bullet w_s) = \tilde{\xi}_1 \bullet w_1 + \cdots + \tilde{\xi}_s \bullet w_s, \end{aligned}$$

where the last equality follows from the fact that Θ is a differential splitting compatible with β . Thus, $w' \in \mathfrak{V}_{\langle \underline{t} \rangle}^+ D_{A[\underline{t}]|\mathbb{K}} M_2 = M_2$. Hence, $M_1 \subseteq M_2$. Similarly, one can show that $M_2 \subseteq M_1$. $\square$

Corollary 4.9. *Suppose that $\mathbb{K}$ has characteristic zero. Let R be a polynomial ring over $\mathbb{K}$, and let A be a $\mathbb{K}$ -subalgebra that is a direct summand of R , and such that the inclusion $A \subseteq R$ is differentially extensible. Let M be a $D_{A[\underline{t}]|\mathbb{K}}$ -module that is a differentially extensible summand of a regular holonomic $D_{R[\underline{t}]|\mathbb{K}}$ -module N with quasi-unipotent monodromy. Then there exists a V -filtration on M along the ideal $\langle \underline{t} \rangle = \langle t_1, \dots, t_\ell \rangle$.*

Finally, we aim to define a V -filtration on a $D_{A|\mathbb{K}}$ -module M along $\underline{f} = f_1, \dots, f_\ell \in A$, where M is a differential direct summand of a $D_{R|\mathbb{K}}$ -module N . The direct image of N under the graph embedding $i_{\underline{f}}$ is the local cohomology module $H_{\langle \underline{t} - \underline{f} \rangle}^\ell(N[\underline{t}])$, where $\langle \underline{t} - \underline{f} \rangle = \langle t_1 - f_1, \dots, t_\ell - f_\ell \rangle$. This $D_{R[\underline{t}]|\mathbb{K}}$ -module has $H_{\langle \underline{t} - \underline{f} \rangle}^\ell(M[\underline{t}])$ as a differential direct summand by Proposition 3.19, and we use this fact in defining our desired V -filtration. In Theorem 4.11, we show that the V -filtration depends on the ideal generated by $\underline{f}$, but not on the particular choice of its generators.

Definition 4.10 (*V -filtration on a $D_{A|\mathbb{K}}$ -module along $\underline{f}$*). Suppose that $\mathbb{K}$ has characteristic zero. Let A be a commutative Noetherian $\mathbb{K}$ -algebra. Given indeterminates $\underline{t} = t_1, \dots, t_\ell$, and $\underline{f} = f_1, \dots, f_\ell \in A$, consider the ideal $\langle \underline{t} - \underline{f} \rangle$ of the polynomial ring $A[\underline{t}]$ generated by $t_1 - f_1, \dots, t_\ell - f_\ell$. For a $D_{A|\mathbb{K}}$ -module M , let M' denote the $D_{A[\underline{t}]|\mathbb{K}}$ -module $H_{\langle \underline{t} - \underline{f} \rangle}^\ell(M[\underline{t}])$, and identify M with the isomorphic module $0 :_{M'} \langle \underline{t} - \underline{f} \rangle \subseteq M'$. Suppose that M' admits a V -filtration along $\langle \underline{t} \rangle$ over $A[\underline{t}]$. Then the V -filtration on M along $\underline{f}$ is defined, for $\alpha \in \mathbb{Q}$, as

$$V_{\langle \underline{f} \rangle}^\alpha M := V_{\langle \underline{t} \rangle}^\alpha M' \cap M = (0 :_{V_{\langle \underline{t} \rangle}^\alpha M'} \langle \underline{t} - \underline{f} \rangle).$$

Using the proof of the previous theorem, we show that a V -filtration over A along $\underline{f}$ only depends on the ideal $I = \langle f_1, \dots, f_\ell \rangle$ and not on the generators chosen.

Corollary 4.11. *Suppose that $\mathbb{K}$ has characteristic zero. Let R be a polynomial ring over $\mathbb{K}$, and let A be a $\mathbb{K}$ -subalgebra that is a direct summand of R , for which the inclusion $A \subseteq R$ is differentially extensible. Let M be a $D_{A[\underline{t}]|\mathbb{K}}$ -module that is a differentially extensible summand of a regular holonomic $D_{R[\underline{t}]|\mathbb{K}}$ -module N with quasi-unipotent monodromy. Suppose that $\underline{f} = f_1, \dots, f_\ell$ and $\underline{g} = g_1, \dots, g_m$ both generate an ideal $I \subseteq A$. Set $M' = H_{\langle \underline{t} - \underline{f} \rangle}^\ell(M[\underline{t}])$ and $M'' = H_{\langle \underline{u} - \underline{g} \rangle}^m(M[\underline{u}])$, for another sequence of indeterminates $\underline{u} = u_1, \dots, u_m$. We identify M with both $0 :_{M'} \langle \underline{t} - \underline{f} \rangle$ and $0 :_{M''} \langle \underline{u} - \underline{g} \rangle$. Then $V_{\langle \underline{t} \rangle}^\alpha M' \cap M = V_{\langle \underline{u} \rangle}^\alpha M'' \cap M$. In particular, the V -filtration of M along $\underline{f}$ depends only on the ideal I .*

Proof. Let $N' = H_{\langle \underline{t} - \underline{f} \rangle}^\ell(N[\underline{t}])$ and $N'' = H_{\langle \underline{u} - \underline{g} \rangle}^m(N[\underline{u}])$. We observe that M' is a differentially extensible summand of N' , and M'' is a differentially extensible

J. Álvarez Montaner et al.

summand of N'' by Proposition 3.19. Since both N' and N'' have V -filtrations (along $\underline{t}$ and $\underline{u}$), so do M' and M'' by Theorem 4.8. We have that N is identified with $0 :_{N'} \langle \underline{t} - \underline{f} \rangle$ and $0 :_{N''} \langle \underline{u} - \underline{g} \rangle$. Then $V_{\langle \underline{t} \rangle}^\alpha N' \cap N = V_{\langle \underline{u} \rangle}^\alpha N'' \cap N$ [17, Proposition 2.6], and thus we have the equalities

$$\begin{aligned} V_{\langle \underline{t} \rangle}^\alpha M' \cap M &= (V_{\langle \underline{t} \rangle}^\alpha N' \cap M') \cap M = V_{\langle \underline{t} \rangle}^\alpha N' \cap M' \cap N \cap M \\ &= (V_{\langle \underline{t} \rangle}^\alpha N' \cap N) \cap M' \cap M = (V_{\langle \underline{u} \rangle}^\alpha N'' \cap N) \cap M'' \cap M \\ &= (V_{\langle \underline{u} \rangle}^\alpha N'' \cap M'') \cap M = V_{\langle \underline{u} \rangle}^\alpha M'' \cap M. \end{aligned} \quad \square$$

4.3. Hodge ideals

Given $f \in R = \mathbb{C}[x_1, \dots, x_d]$, the $D_{R|\mathbb{C}}$ -module of regular functions on $X = \text{Spec}(R)$ with poles along the divisor $Y = \text{div}(f)$ is given by $\mathcal{O}_X(*Y) = R_f$. This module is equipped with the Hodge filtration $\mathcal{F}_\bullet(\mathcal{O}_X(*Y))$ [56, 66], which is an invariant measuring the singularities of Y . Mustařă and Popa [56] defined a sequence of ideals, the *Hodge ideals* of f , that encode data of the Hodge filtration. Later on, they extended this notion to $\mathbb{Q}$ -divisors [57, 58]. In this framework, they considered a filtration $\mathcal{F}_\bullet \mathcal{O}_X(*Y) \cdot f^{1-\lambda}$ on the free $D_{R|\mathbb{C}}$ -module of rank one $\mathcal{O}_X(*Y) \cdot f^{1-\lambda} = R_f \cdot f^{1-\lambda}$, where $\lambda \in \mathbb{Q}_{\geq 0}$. If we denote the support of Y as $Z = Y_{\text{red}}$, then the sequence of *Hodge ideals* of f^λ , which we denote as $I_k(f^\lambda)$, $k \geq 0$, is defined by the equations:

$$\mathcal{F}_k \mathcal{O}_X(*Y) \cdot f^{1-\lambda} = I_k(\lambda Y) \otimes \mathcal{O}_X(kZ + Y) \cdot f^{1-\lambda}.$$

We point out that Hodge ideals can be also defined for ideals of R , and $\lambda \in (0, 1] \cap \mathbb{Q}$ [54]. This description boils down to the computation of Hodge ideals of $\mathbb{Q}$ -divisors.

An interesting feature is that Hodge ideals of $\mathbb{Q}$ -divisors are characterized completely in terms of V -filtrations. For simplicity we will just consider the case where the divisor is reduced. Set $Q_j(x) := x(x+1) \cdots (x+j-1) \in \mathbb{Z}[x]$.

Theorem 4.12 ([58]). *Let Y be a reduced divisor defined by $f \in R$. Then for all $\lambda \in \mathbb{Q}_{\geq 0}$ and $k \geq 0$, we have*

$$I_k(f^\lambda) = \left\{ g = \sum_{j=0}^k Q_j(\lambda) f^{k-j} h_j \in R \mid \sum_{j=0}^k h_j \partial_t^j \frac{1}{f-t} \in V_{\langle \underline{t} \rangle}^\lambda H_{\langle f-t \rangle}^1(R[t]) \right\}.$$

We refer to the work of Mustařă and Popa [55–58] for properties of these ideals.

Theorem 4.12 allows us to define Hodge ideals whenever V -filtrations are defined.

Definition 4.13. Suppose that $\mathbb{K}$ has characteristic zero. Let T be a $\mathbb{K}$ -algebra that admits a V -filtration along a reduced $f \in T$. For any $\lambda \in \mathbb{Q}_{\geq 0}$, and every $k \geq 0$, the k th Hodge ideal of f^λ is defined by

$$I_k^T(f^\lambda) = \left\{ g = \sum_{j=0}^k Q_j(\lambda) f^{k-j} h_j \in T \mid \sum_{j=0}^k h_j \partial_t^j \frac{1}{f-t} \in V_{\langle \underline{t} \rangle}^\lambda H_{\langle f-t \rangle}^1(T[t]) \right\}.$$

If the ring T is clear from the context, we write $I_k(f^\lambda)$ instead of $I_k^T(f^\lambda)$.

From our results on V -filtrations, we can show that Hodge ideals exist for extensible direct summands of polynomial rings.

Corollary 4.14. *Suppose that $\mathbb{K}$ has characteristic zero. Let R be a polynomial ring over $\mathbb{K}$, and let A be a $\mathbb{K}$ -subalgebra that is a direct summand of R , for which the inclusion $A \subseteq R$ is differentially extensible. Then $I_k^A(f^\lambda)$ exists for all $f \in A$ and $\lambda \in \mathbb{Q}_{\geq 0}$. Furthermore,*

$$I_k^A(f^\lambda) = I_k^R(f^\lambda) \cap A. \quad (4.1)$$

Proof. The existence of Hodge ideals follows from Corollary 4.9. We prove (4.1) by showing both containments. First take $g \in I_k^A(f^\lambda)$. There exists $h_0, \dots, h_k \in A \subseteq R$ such that $g = \sum_{j=0}^k Q_j(\lambda) f^{k-j} h_j$ and

$$\sum_{j=0}^k h_j \partial_t^j \frac{1}{f-t} \in V_{\langle t \rangle}^\lambda H_{\langle f-t \rangle}^1(A[t]) \subseteq V_{\langle t \rangle}^\lambda H_{\langle f-t \rangle}^1(R[t])$$

so that $g \in I_k^R(f^\lambda)$.

On the other hand, for $g \in I_k^R(f^\lambda) \cap A$, there exist $h_0, \dots, h_k \in R$ such that $g = \sum_{j=0}^k Q_j(\lambda) f^{k-j} h_j$ and

$$\sum_{j=0}^k h_j \partial_t^j \frac{1}{f-t} \in V_{\langle t \rangle}^\lambda H_{\langle f-t \rangle}^1(A[t]) \subseteq V_{\langle t \rangle}^\lambda H_{\langle f-t \rangle}^1(R[t]).$$

Let $\beta: R \rightarrow A$ a splitting, and $\theta: H_{\langle f-t \rangle}^1(R[t]) \rightarrow H_{\langle f-t \rangle}^1(A[t])$ a differentiable splitting. Then

$$\begin{aligned} \theta \left(\sum_{j=0}^k h_j \partial_t^j \frac{1}{f-t} \right) &= \sum_{j=0}^k \theta \left(h_j \partial_t^j \frac{1}{f-t} \right) \\ &= \sum_{j=0}^k (\beta \circ h_j \partial_t^j) \frac{1}{f-t} \\ &= \sum_{j=0}^k (\beta h_j) \partial_t^j \frac{1}{f-t}. \end{aligned}$$

Since $\beta(h_0), \dots, \beta(h_k) \in A$, we conclude that $g \in I_k^A(f^\lambda)$. $\square$

We now recall a definition related to the log canonical threshold.

Definition 4.15. Suppose that $\mathbb{K}$ has characteristic zero. Let T be a $\mathbb{K}$ -algebra such that $b_f^T(s)$ exists for every $f \in T$. The *minimal exponent* of f , $\tilde{\alpha}_f$, is the negative of the largest root of $b_f(s)/(s+1)$.

J. Álvarez Montaner et al.

As a consequence of Corollary 4.14, we can relate Hodge ideals and minimal exponent.

Theorem 4.16. *Suppose that $\mathbb{K}$ has characteristic zero. Let R be a polynomial ring over $\mathbb{K}$, and let A be a $\mathbb{K}$ -subalgebra that is a direct summand of R , for which the inclusion $A \subseteq R$ is differentially extensible. Given $\lambda \in (0, 1] \cap \mathbb{Q}$, $I_k^A(f^\lambda) = 0$ if and only if $k \leq \tilde{\alpha}_f - \lambda$.*

Proof. Since this equivalence is known for polynomial rings [58, Theorem C], the result follows from Theorem 3.29 and 4.14. $\square$

5. Multiplier Ideals

Given a field $\mathbb{K}$ of characteristic zero, let R be a polynomial ring over $\mathbb{K}$ and $X = \text{Spec}(R)$. To any ideal $I \subseteq R$, we associate a family of *multiplier ideals* $\mathcal{J}_R(I^\lambda)$ parametrized by nonnegative real numbers λ . There exists a discrete sequence of rational numbers

$$0 = \lambda_0 < \lambda_1 < \lambda_2 < \cdots$$

called the *jumping numbers* of I , for which $\mathcal{J}_R(I^\lambda)$ is constant for $\lambda_i \leq \lambda < \lambda_{i+1}$, and $\mathcal{J}_R(I^{\lambda_i}) \subsetneq \mathcal{J}_R(I^{\lambda_{i+1}})$. The first nonzero jumping number, λ_1 , is known as the *log canonical threshold* of I , denoted $\text{lct}_R(I)$.

We begin this section by recalling the definition of the multiplier ideals in this setting, and we refer to [42] for details, and any unexplained terminology.

Definition 5.1 (Multiplier ideals over a polynomial ring). Let R be a polynomial ring over a field $\mathbb{K}$ of characteristic zero, let I be an ideal of R , and let $X = \text{Spec}(R)$. Let $\pi: Y \rightarrow X$ be a log resolution of I such that $I\mathcal{O}_Y = \mathcal{O}_Y(-F_\pi)$, where $F_\pi = \sum r_i E_i$, and $K_{Y/X} = \sum b_i E_i$ is the relative canonical divisor. The *multiplier ideal* of I with exponent $\lambda \in \mathbb{R}_{\geq 0}$ is defined by

$$\begin{aligned} \mathcal{J}_R(I^\lambda) &= \pi_* \mathcal{O}_Y(\lceil K_{Y/X} - \lambda F_\pi \rceil) \\ &= \{h \in R : \text{ord}_{E_i}(\pi^* h) \geq \lfloor \lambda r_i - b_i \rfloor \ \forall i\}. \end{aligned}$$

In our setting, there is a characterization of multiplier ideals in terms of the V -filtration that, roughly speaking, says that the chain of multiplier ideals is essentially the V -filtration of R along the ideal I [17, 18]. In fact, this relation has been used to develop algorithms to compute multiplier ideals, based on existent algorithms for computing the Bernstein–Sato polynomial that use Gröbner bases over the ring of differential operators [2, 74].

Theorem 5.2 ([17, Theorem 1, Corollary 2]). *Let R be a polynomial ring over a field $\mathbb{K}$ of characteristic zero, and let I be an ideal of R . Given a real number $\lambda \geq 0$, the following ideals coincide:*

- (1) $\mathcal{J}_R(I^\lambda)$,

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

- (2) $\bigcup_{\alpha > \lambda} V^\alpha R$, where the V -filtration is taken along I , and
- (3) $\{g \in R : \gamma > \lambda \text{ if } b_{I,g}^R(-\gamma) = 0\}$.

The definition of the multiplier ideals can be easily extended to the case that X is $\mathbb{Q}$ -Gorenstein, where it is possible to construct the relative canonical divisor. A version of multiplier ideals in normal varieties was defined by de Fernex and Hacon [21], and later extended by Chieochio *et al.* to include a boundary divisor term [19].

Our aim in this section is to study multiplier ideals for direct summands of coordinate rings of smooth varieties, and to extend Theorem 5.2 to this setting. Our most general result in this direction is the following.

Theorem 5.3. *Let R be a polynomial ring over a field $\mathbb{K}$ of characteristic zero, and let $A \subseteq R$ be a differentially extensible inclusion of finitely generated $\mathbb{K}$ -algebras, such that A is a direct summand of R . Then for every ideal I of A , and every real number $\lambda \geq 0$, the following ideals coincide:*

- (1) $\mathcal{J}_R((IR)^\lambda) \cap A$,
- (2) $\bigcup_{\alpha > \lambda} V^\alpha A$, where the V -filtration is taken along I , and
- (3) $\{g \in A : \gamma > \lambda \text{ if } b_{I,g}^A(-\gamma) = 0\}$.

Proof. From the equality $b_{I,g}^A(s) = b_{IR,g}^R(s)$ given in Theorem 3.29, we deduce

$$\begin{aligned} \{g \in A : \gamma > \lambda \text{ if } b_{I,g}^A(-\gamma) = 0\} \\ = \{g \in R : \gamma > \lambda \text{ if } b_{IR,g}^R(-\gamma) = 0\} \cap A. \end{aligned}$$

Moreover, $\bigcup_{\alpha > \lambda} V^\alpha A$ is the intersection $(\bigcup_{\alpha > \lambda} V^\alpha R) \cap A$, where the V -filtrations are taken along I in A , and along IR in R . The claim then follows from Theorem 5.2. $\square$

Of course, it would be desirable if $\mathcal{J}_R((IR)^\lambda) \cap A$ were equal to $\mathcal{J}_A(I^\lambda)$, and we show this, under certain conditions, in Theorem 5.15. To prove this result, we make a detour through positive characteristic methods, relying on the well-known relation between multiplier ideals and test ideals [29, 30, 76].

5.1. Test ideals and Cartier extensibility

Throughout this section, T denotes a commutative Noetherian ring of prime characteristic p . The e th iteration of the Frobenius endomorphism F^e on T , given by $r \mapsto r^{p^e}$, yields a new T -module structure on T by restriction of scalars. We denote this module action on T by $F_*^e T$, so that for $r \in T$ and $F_*^e x \in F_*^e T$,

$$r \cdot F_*^e x = F_*^e(r^{p^e} x) \in F_*^e T.$$

We say that T is F -finite if $F_*^e T$ is a finitely generated T -module for some (equivalently, all) $e > 0$.

J. Álvarez Montaner et al.

The iterated Frobenius endomorphism can be used to classify singularities.

Definition 5.4. Let T be an F -finite Noetherian domain of prime characteristic.

- (1) We call T *F-pure* if the inclusion map $T \rightarrow F_*^e T$ splits for some (equivalently, all) $e > 0$.
- (2) We call T *strongly F-regular* if for every $r \in T$, there exists $e \in \mathbb{N}$ such that the T -linear map $T \rightarrow F_*^e T$ sending 1 to $F_*^e r$ splits.

A regular ring is strongly F -regular, and a direct summand of a strongly F -regular ring is itself strongly F -regular [31, Theorem 3.1].

Definition 5.5. Let T be an F -finite Noetherian domain of prime characteristic p . An additive map $\psi: T \rightarrow T$ is a p^{-e} -linear map if $\psi(r^{p^e} f) = r\psi(f)$. Such a map is also referred to as a *Cartier operator*. The set of all the p^{-e} -linear maps on T is denoted $\mathcal{C}_T^e$; in the notation of Definition 5.5, $\mathcal{C}_T^e \cong \text{Hom}_T(F_*^e T, T)$.

Test ideals first appeared in the theory of tight closure developed by Hochster and Huneke [32, 33]. Hara and Yoshida subsequently defined test ideals associated to pairs (T, I^λ) , where I is an ideal of a regular ring T , and λ is a nonnegative real parameter [30]. A tight closure-free version of test ideals was developed by Blickle *et al.* using Cartier operators over regular rings [10, 11]. This new approach was extended to nonregular rings [8, 9, 72]. For strongly F -regular rings, we have the following description (see [82]).

Definition 5.6. Let T be an F -finite Noetherian domain of prime characteristic p that is strongly F -regular. The *test ideal* of an ideal I of T with respect to a real parameter $\lambda \geq 0$ is defined by

$$\tau_T(I^\lambda) = \bigcup_{e>0} \left(\sum_{\psi \in \mathcal{C}_T^e} \psi(I^{\lceil p^e \lambda \rceil}) \right).$$

Cartier operators are closely related to differential operators, so it is natural to consider our condition of differentiable extensibility in this context, as well.

Definition 5.7 (Cartier extensibility). Consider a ring homomorphism $\varphi: A \rightarrow T$, and fix a Cartier operator $\psi \in \mathcal{C}_A^e$. We say that

- (1) A Cartier operator $\tilde{\psi} \in \mathcal{C}_T^e$ *extends* ψ if $\varphi \circ \psi = \tilde{\psi} \circ \varphi$; i.e. the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & T \\ \psi \downarrow & & \downarrow \tilde{\psi} \\ A & \xrightarrow{\varphi} & T \end{array}$$

If φ is simply an inclusion, then this is equivalent to the condition that $\tilde{\psi}|_A = \psi$.

- (2) The map φ is *Cartier extensible* if for every $\psi \in \mathcal{C}_A^e$, there exists $\tilde{\psi} \in \mathcal{C}_T^e$ that extends ψ .

Lemma 5.8. *An inclusion $A \subseteq T$ of F -finite F -pure domains of prime characteristic $p > 0$ that is differentially extensible with respect to the level filtration is also Cartier extensible.*

Proof. Let $i_T: T \rightarrow F_*^e T$ be the inclusion map, and $\pi_T: F_*^e T \rightarrow T$ a T -linear splitting. Likewise, let $i_A: A \rightarrow F_*^e A$ be the inclusion map, and π_A an A -splitting. Given an A -linear map $\psi: F_*^e A \rightarrow A$, we obtain an A -linear map $i_A \circ \psi$ on $F_*^e A$. Note that $\psi = \pi_A \circ (i_A \circ \psi)$. By assumption, there is a T -linear map β on $F_*^e T$ such that $\beta|_{F_*^e A} = i_A \circ \psi$. Then since $\text{im}(\beta|_{F_*^e A}) \subseteq A$ and $(\pi_T)|_A$ is the identity map, $\pi_T \circ \beta \in \mathcal{C}_T^e$ extends ψ . $\square$

Proposition 5.9. *Let R be a regular F -finite domain of prime characteristic $p > 0$. Let $A \subseteq R$ be an extension of Noetherian rings, such that A is a Cartier extensible direct summand of R . Then for every ideal I of A , and every real number $\lambda \geq 0$.*

$$\tau_A(I^\lambda) = \tau_R((IR)^\lambda) \cap A.$$

Proof. Since A is a direct summand of R , $\tau_R((IR)^\lambda) \cap A \subseteq \tau_A(I^\lambda)$ [1, Proposition 4.9], and we focus on the other containment. For any $f \in \tau_A(I^\lambda)$, there exists $e \in \mathbb{N}$ for which $f \in \mathcal{C}_A^e I^{[p^e \lambda]}$. Then there exist $g_1, \dots, g_\ell \in I^{[p^e \lambda]}$ and $\psi_1, \dots, \psi_\ell \in \mathcal{C}_A^e$ such that $f = \psi_1 g_1 + \dots + \psi_\ell g_\ell$. Given extensions $\tilde{\psi}_i \in \mathcal{C}_R^e$ of ψ_i ,

$$f = \psi_1 g_1 + \dots + \psi_\ell g_\ell = \tilde{\psi}_1 g_1 + \dots + \tilde{\psi}_\ell g_\ell \in \mathcal{C}_R^e I^{[p^e \lambda]} \subseteq \tau_R((IR)^\lambda)$$

so that $\tau_A(I^\lambda) \subseteq \tau_R((IR)^\lambda) \cap A$. $\square$

5.2. Multiplier ideals and reduction modulo p

The relationship between multiplier ideals and test ideals using reduction to prime characteristic p was originally observed in work of Smith [75, 76] and Hara [29]. For the case of pairs, we refer to work of Hara and Yoshida [30], to that of Takagi [80] for $\mathbb{Q}$ -Gorenstein varieties, and to that of de Fernex *et al.* for the extension to the numerically $\mathbb{Q}$ -Gorenstein case [20]. Herein, we call upon a result of Chiechio *et al.* that states that multiplier ideals reduce to test ideals via reduction to prime characteristic for a large class of rings [19].

Before explaining the result in [19], we recall the basics on reduction to prime characteristic. For more details, we refer to Hochster and Huneke’s notes on tight closure in characteristic zero [34].

Definition 5.10 (Model for an ideal in a finitely generated $\mathbb{K}$ -algebra).

Suppose that A is a finitely generated $\mathbb{K}$ -algebra, where $\mathbb{K}$ has characteristic zero, and fix an ideal I of A . Fix a finitely generated $\mathbb{Z}$ -subalgebra B of $\mathbb{K}$, and a B -subalgebra A_B of A of finite type over B , that satisfy the following conditions: A_B is flat over B , $A_B \otimes_B \mathbb{K} \cong A$, and $I_B A = I$, where $I_B = I \cap A_B \subseteq A_B$. For every closed point $s \in \text{Spec}(B)$, we set $A_s = A_B \otimes_B \kappa(s)$ and $I_s = I_B A_s \subseteq A_s$, where $\kappa(s)$ denotes the residue field of s . We say that (B, A_B, I_B) is a *model* for $(\mathbb{K}, A, I)$.

J. Álvarez Montaner et al.

If, additionally, R is another finitely generated $\mathbb{K}$ -algebra, and $\varphi: A \rightarrow R$ is a $\mathbb{K}$ -algebra homomorphism, then we say that $(B, A_B, I_B, R_B, \varphi_B)$ is a *model* for $(\mathbb{K}, A, I, R, \varphi)$ if (B, A_B, I_B) is a model for $(\mathbb{K}, A, I)$, $(B, R_B, 0)$ is a model for $(\mathbb{K}, R, 0)$, and $\varphi_B: A_B \rightarrow R_B$ satisfies $\varphi = \varphi_B \otimes_B \mathbb{K}$. We write $\varphi_s = \varphi_B \otimes_B \kappa(s)$ for $s \in \text{Spec}(B)$ closed.

Remark 5.11. Note that if A is a subalgebra of a polynomial ring R over a field $\mathbb{K}$ of characteristic zero, and A is generated by polynomials with integer coefficients, then one can take $B = \mathbb{Z}$ in a model for A , and $A_{\mathbb{Z}}$ the $\mathbb{Z}$ -algebra with the same generators. In this case, a condition on A_s , for all s in a dense open subset of $\text{Spec}(\mathbb{Z})$ is equivalent to this condition on $A_{\mathbb{Z}}/pA_{\mathbb{Z}}$, for all $p \gg 0$. We write A_p to denote $A_{\mathbb{Z}}/pA_{\mathbb{Z}}$ in this case.

Definition 5.12 (Anticanonical cover). Let A be a normal ring that is either local and complete, or $\mathbb{N}$ -graded and finitely generated over its zeroth graded component. The *anticanonical cover* of A is the symbolic Rees algebra

$$\mathcal{R}(-\omega_A) = \bigoplus_{n \geq 0} \omega_A^{(-n)},$$

where ω_A is the canonical module of A .

By recent results in the minimal model program [5, Corollary 1.1.9], we know that for varieties with KLT singularities, $\mathcal{R}(-\omega_A)$ is finitely generated.

Theorem 5.13 ([19, Theorem D]). *Suppose that A is a finitely generated $\mathbb{K}$ -algebra over an algebraically closed field of characteristic zero such that the anticanonical cover $\mathcal{R}(-\omega_A)$ is finitely generated. Let I be an ideal of A , and fix $\lambda \in \mathbb{R}_{\geq 0}$. Let (B, A_B, I_B) be a model for $(\mathbb{K}, A, I)$. Then*

$$[\mathcal{J}_A(I^\lambda)]_s = \tau_{A_s}(I_s^\lambda)$$

for every closed point s in a dense open subset of $\text{Spec}(B)$.

Consider the following setup.

Setup 5.14. Let R be a polynomial ring with coefficients in an algebraically closed field $\mathbb{K}$ of characteristic zero. Suppose that $A \subseteq R$ is differentially extensible, that the inclusion map φ makes A a direct summand of R , and that A is finitely generated over $\mathbb{K}$ with KLT singularities. Moreover, let I be an ideal of A , and let $(B, A_B, I_B, R_B, \varphi_B)$ be a model for $(\mathbb{K}, A, I, R, \varphi)$. Suppose that for all s in a dense open subset U of $\text{Spec}(B)$, A_s is a direct summand of R_s and the inclusion φ_s of A_s into R_s is also Cartier extensible.

In this setting, we fully understand the relationship between the multiplier ideals $\mathcal{J}_A(I^\lambda)$ and $\mathcal{J}_R(IR)$.

Theorem 5.15. *In the context of Setup 5.14, for every real number $\lambda \geq 0$,*

$$\mathcal{J}_A(I^\lambda) = \mathcal{J}_R((IR)^\lambda) \cap A.$$

Proof. Take $f \in \mathcal{J}_A(I^\lambda)$, and for $s \in \text{Spec}(B)$, let f_s denote the image of f in A_s . Then $f_s \in [\mathcal{J}_A(I^\lambda)]_s$ for every s , so $f_s \in \tau_{A_s}(I_s^\lambda)$ for $s \in U$ by Theorem 5.13. Thus, $f_s \in \tau_{R_s}((IR_s)^\lambda) \cap A_s$ by Proposition 5.9. In particular, $f_s \in \tau_{R_s}((IR_s)^\lambda)$, so that $f_s \in [\mathcal{J}_R((IR)^\lambda)]_s$ for $s \in U$, and $f \in \mathcal{J}_R((IR)^\lambda) \cap A$.

Now take $f \in \mathcal{J}_R((IR)^\lambda) \cap A$, so that $f_s \in [\mathcal{J}_R(I^\lambda)]_s \cap A_s$ for every s . Then $f_s \in \tau_{R_s}((IR_s)^\lambda) \cap A_s = \tau_{A_s}(I_s^\lambda)$ for all s in the dense open set described in Theorem 5.13 and Proposition 5.9. Then $f_s \in [\mathcal{J}_A(I^\lambda)]_s$ for such s , and so $f \in \mathcal{J}_A(I^\lambda)$. $\square$

The hypotheses that the inclusion $A \subseteq R$ is differentially extensible, and split as A -modules, are essential for Theorem 5.15, as the following example shows.

Example 5.16. Given $A = \mathbb{C}[x^2, y^2] \subseteq R = \mathbb{C}[x, y]$, and $I = \langle x^2, y^2 \rangle \subseteq A$, then $\mathcal{J}_A(I) = A$, whereas $\mathcal{J}_R(IR) = \langle x, y \rangle$. In particular, $\mathcal{J}_R(IR) \cap A \neq \mathcal{J}_A(I)$.

The following example indicates that the assumption in Theorem 5.15 that, after reduction to prime characteristic, $A_p \subseteq R_p$ is a direct summand that is also Cartier extensible for $p \gg 0$, might not be necessary.

Example 5.17. Let X, Y, Z be 3×3 , 3×2 , and 2×3 matrices of indeterminates, respectively. Let $T = \mathbb{C}[X]$, $A = T/\langle \det(X) \rangle$, and $R = \mathbb{C}[Y, Z]$. Then the $\mathbb{C}$ -algebra map sending x_{ij} in A to the (i, j) -entry of the matrix product YZ realizes A as an extensible direct summand of R [43, IV 1.9]. Let $I \subseteq A$ be the ideal generated by the 2×2 minors of X .

Using the methods of Johnson [36, Chaps. 4 and 5] along with techniques of Takagi [81, Theorem 3.1], one can compute $\mathcal{J}_A(I^\lambda) = \mathfrak{m}^{\lfloor 2\lambda \rfloor - 5} \cap I^{\lfloor \lambda \rfloor - 1}$, where $\mathfrak{m}$ is the ideal generated by the entries of X . We have that $\mathcal{J}_R(IR^\lambda) = (I_1(Y)I_1(Z))^{\lfloor 2\lambda \rfloor - 5} \cap (I_2(Y)I_2(Z))^{\lfloor \lambda \rfloor - 1}$ using [36, Theorem 5.4; 23, Variant 2.5], where $I_i(-)$ denotes the ideal of $i \times i$ -minors of a matrix. It is then easy to verify that $\mathcal{J}_R(IR^\lambda) \cap A = \mathcal{J}_A(I^\lambda)$.

However, for p prime, the inclusion $A_p \subseteq R_p$ is not a direct summand. Indeed, suppose it were. Then there would be an injection $H_{\mathfrak{m}_p}^8(A_p) \hookrightarrow H_{\mathfrak{m}_p R_p}^8(R_p)$. However, $H_{\mathfrak{m}_p}^8(A_p) \neq 0$ by Grothendieck nonvanishing, and since $\text{pd}_{R_p}(\mathfrak{m}_p R_p) = 7$, we know that $H_{\mathfrak{m}_p R_p}^8(R_p) = 0$ by Peskine–Szpiro vanishing [64, Proposition 4.1], yielding a contradiction.

We note that we do not know whether the inclusions $A_p \subseteq R_p$ are Cartier extensible or differentially extensible.

Motivated by Theorems 5.3 and 5.15 and Example 5.17, we wonder whether it is possible to drop the condition of Cartier extensibility after reduction to prime characteristic. Namely, we pose the following question:

Question 5.18. Let A be a differentially extensible summand of a polynomial ring over a field of characteristic zero. Given an ideal I of A , and a real number $\lambda \geq 0$,

J. Álvarez Montaner et al.

do the following ideals coincide?

- (1) $\mathcal{J}_A(I^\lambda)$,
- (2) $\bigcup_{\alpha > \lambda} V^\alpha A$, where the V -filtration is taken along I , and
- (3) $\{g \in A : \gamma > \lambda \text{ if } b_{I,g}^A(-\gamma) = 0\}$.

The following is an extension to our setting of results for smooth varieties that relate the jumping numbers of multiplier ideals with the roots of Bernstein–Sato polynomials [17, 26].

Theorem 5.19. *Under Setup 5.14, the log canonical threshold of I in A is the smallest root α_I of $b_I(-s)$, and every jumping number of I in $[\alpha_I, \alpha_I + 1)$ is a root of $b_I(-s)$.*

Proof. Since $\mathcal{J}_A(I^\lambda) = \mathcal{J}_R((IR)^\lambda) \cap A$, we have that $\mathcal{J}_A(I^\lambda) = A$ if and only if $\mathcal{J}_R((IR)^\lambda) = R$. As a consequence, $\text{lct}_A(I) = \text{lct}_R(IR)$. Since $b_I^A(-s) = b_{IR}^R(-s)$ by Theorem 3.29, the first claim follows. Furthermore, every jumping number of I in A is also a jumping number of IR in R . Then the second claim follows again from Theorem 3.29. $\square$

The above results apply to pointed normal affine toric varieties. Recall that by Example 3.4, Lemma 3.5, and Remark 3.6, there is an embedding of A into a polynomial ring R over $\mathbb{K}$ such that, $A \subseteq R$ is order-differentially extensible, and $A_p \subseteq R_p$ is level-differentially extensible for $p \gg 0$. The ring R in the following theorem is chosen in this way.

Corollary 5.20. *Let I be an ideal of A , the coordinate ring of a pointed normal affine toric variety over a field $\mathbb{K}$ of characteristic zero. Then there exists an embedding of A into a polynomial ring R over $\mathbb{K}$ such that $\mathcal{J}_A(I^\lambda) = \mathcal{J}_R((IR)^\lambda) \cap A$ for all real numbers $\lambda \geq 0$.*

Proof. With R chosen as in the discussion before the statement, $A_p \subseteq R_p$ is Cartier extensible for $p \gg 0$ by Lemma 5.8, so we can apply Theorem 5.15. $\square$

The following result is already established for monomial ideals I [35].

Corollary 5.21. *Let $I \subseteq A$ be an ideal in an affine normal toric ring over an algebraically closed field $\mathbb{K}$ of characteristic zero. Then the log canonical threshold of I in A coincides with the smallest root α_I of the Bernstein–Sato polynomial $b_I(-s)$, and any jumping numbers of I in $[\alpha_I, \alpha_I + 1)$ are roots of $b_I(-s)$.*

We obtain similar results for rings of invariants under the action of a finite group.

Corollary 5.22. *Let $\mathbb{K}$ be a field of characteristic zero, and G a finite group acting linearly on a polynomial ring R over $\mathbb{K}$. Suppose that G contains no element that*

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

fixes a hyperplane in the space of one-forms $[R]_1$. Let $A = R^G$ denote the ring of invariants. Then the log canonical threshold of I in A coincides with the smallest root α_I of $b_I(-s)$, and every jumping number of I in $[\alpha_I, \alpha_I + 1)$ is a root of $b_I(-s)$.

In the polynomial ring setting, we can understand Hodge ideals as a generalization of multiplier ideals. Indeed,

$$I_0^R(f^\lambda) = \mathcal{J}_R(f^{\lambda-\varepsilon})$$

for $\varepsilon > 0$ small enough [57, Proposition 9.1]. In the case of ideals $I \subseteq R$, the same statement holds when $\lambda \in (0, 1] \cap \mathbb{Q}$ [54]. As a consequence of Theorem 5.15, we have the same property for extensible direct summands of polynomial rings.

Proposition 5.23. *In the context of Setup 5.14, for every $\lambda \in \mathbb{Q}_{\geq 0}$ and $\varepsilon > 0$ small enough, we have*

$$I_0^A(f^\lambda) = \mathcal{J}_A(f^{\lambda-\varepsilon}).$$

Proof. Using Corollary 4.14 and Theorem 5.15 we have

$$I_0^A(f^\lambda) = I_0^R(f^\lambda) \cap A = \mathcal{J}_R(f^{\lambda-\varepsilon}) \cap A = \mathcal{J}_A(f^{\lambda-\varepsilon}). \quad \square$$

Acknowledgments

This project was initiated and developed during an American Institute of Mathematics Structured Quartet Research Ensemble (SQuaRE) titled *Roots of Bernstein–Sato polynomials*. We are grateful for AIM’s hospitality, and for their support of this project. We thank Devlin Mallory for helping us compute Example 5.17, and Eamon Quinlan for a careful reading of the paper. We also thank Nero Budur, Manuel González, Craig Huneke, Mircea Mustață, Luis Narváez-Macarro, Claude Sabbah, and Karl Schwede for helpful conversations and correspondences. In particular, we thank Manuel González for bringing our attention to Hodge ideals, and for discussions regarding Sec. 4.3. We thank the referee for the helpful comments. Josep Àlvarez Montaner is partially supported by grants PID2019-103849GB-I00 (AEI/10.13039/501100011033) and 2017SGR-932 (AGAUR). Daniel J. Hernández is partially supported by the NSF Grants DMS-1304250 and DMS-1600702. Jack Jeffries is partially supported by the NSF Grant DMS-1606353. Luis Núñez-Betancourt is partially supported by the NSF CAREER Award DMS 194561, CONACYT Grant 284598, and Cátedras Marcos Moshinsky. Emily E. Witt is partially supported by the NSF Grant DMS-1623035 and Simons Foundation Grant 582574.

References

- [1] J. Àlvarez Montaner, C. Huneke and L. Núñez-Betancourt, D -modules, Bernstein–Sato polynomials and F -invariants of direct summands, *Adv. Math.* **321** (2017) 298–325.

J. Álvarez Montaner et al.

- [2] C. Berkesch and A. Leykin, Algorithms for Bernstein–Sato polynomials and multiplier ideals, in *ISSAC 2010 — Proc. 2010 Int. Symp. Symbolic and Algebraic Computation* (ACM, New York, 2010) pp. 99–106.
- [3] J. N. Bernšteĭn, Analytic continuation of generalized functions with respect to a parameter, *Funktsional. Anal. i Prilozhen.* **6**(4) (1972) 26–40.
- [4] J. N. Bernšteĭn, I. M. Gel’fand and S. I. Gel’fand, Differential operators on a cubic cone, *Uspekhi Mat. Nauk* **27**(1(163)) (1972) 185–190.
- [5] C. Birkar, P. Cascini, C. D. Hacon and J. McKernan, Existence of minimal models for varieties of log general type, *J. Amer. Math. Soc.* **23**(2) (2010) 405–468.
- [6] J.-E. Björk, Dimensions of modules over algebras of differential operators, in *Fonctions Analytiques de Plusieurs Variables et Analyse complexe*, “Agora Mathematica” Vol. 1, Colloques internationaux du Centre national de la Recherche scientifique, Vol. 208 (Gauthier-Villars, Paris, 1972/1974), pp. 6–11.
- [7] J.-E. Björk, *Rings of Differential Operators*, North-Holland Mathematical Library, Amsterdam–New York, Vol. 21 (1979).
- [8] M. Blickle, Test ideals via algebras of p^{-e} -linear maps, *J. Algebraic Geom.* **22**(1) (2013) 49–83.
- [9] M. Blickle and G. Bockle, Cartier modules: Finiteness results, *J. Reine Angew. Math.* **661** (2011) 85–123.
- [10] M. Blickle, M. Mustață and K. E. Smith, Discreteness and rationality of F -thresholds, *Michigan Math. J.* **57** (2008) 43–61.
- [11] M. Blickle, M. Mustață and K. E. Smith, F -thresholds of hypersurfaces, *Trans. Amer. Math. Soc.* **361**(12) (2009) 6549–6565.
- [12] A. Bravo and K. E. Smith, Behavior of test ideals under smooth and étale homomorphisms, *J. Algebra* **247**(1) (2002) 78–94.
- [13] H. Brenner, J. Jeffries and L. Núñez-Betancourt, Quantifying singularities with differential operators, *Adv. Math.* **358** (2019), Article ID: 106843, 89pp.
- [14] N. Budur, On Hodge spectrum and multiplier ideals, *Math. Ann.* **327**(2) (2003) 257–270.
- [15] N. Budur, On the V -filtration of D -modules, in *Geometric Methods in Algebra and Number Theory, Progress in Mathematics* Vol. 235 (Birkhäuser, Boston, MA, 2005) pp. 59–70.
- [16] N. Budur, Bernstein–Sato polynomials, Lecture notes for the summer school, Multiplier Ideals, Test Ideals, and Bernstein–Sato Polynomials, at UPC Barcelona, available at <https://perswww.kuleuven.be/u0089821/Barcelona/BarcelonaNotes.pdf> (2015).
- [17] N. Budur, M. Mustață and M. Saito, Bernstein–Sato polynomials of arbitrary varieties, *Compos. Math.* **142**(3) (2006) 779–797.
- [18] N. Budur and M. Saito, Multiplier ideals, V -filtration, and spectrum, *J. Algebr. Geom.* **14**(2) (2005) 269–282.
- [19] A. Chiechio, F. Enescu, L. E. Miller and K. Schwede, Test ideals in rings with finitely generated anti-canonical algebras, *J. Inst. Math. Jussieu* **17**(1) (2018) 171–206.
- [20] T. de Fernex, R. Docampo, S. Takagi and K. Tucker, Comparing multiplier ideals to test ideals on numerically $\mathbb{Q}$ -Gorenstein varieties, *Bull. Lond. Math. Soc.* **47**(2) (2015) 359–369.
- [21] T. de Fernex and C. D. Hacon, Singularities on normal varieties, *Compos. Math.* **145**(2) (2009) 393–414.
- [22] P. Deligne and N. Katz, *Groupes de Monodromie en Géométrie Algébrique II*, Lecture Notes in Mathematics, Vol. 340 (Springer-Verlag, Berlin–New York, 1973) (Séminaire de Géométrie Algébrique du Bois-Marie SGA 7 II, 1967–1969).

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

- [23] J.-P. Demailly, L. Ein and R. Lazarsfeld, A subadditivity property of multiplier ideals, *Michigan Math. J.* **48** (2000) 137–156.
- [24] J. Denef and F. Loeser, Caractéristiques d’Euler-Poincaré, fonctions zêta locales et modifications analytiques, *J. Amer. Math. Soc.* **5**(4) (1992) 705–720.
- [25] A. Dimca, P. Maisonobe, M. Saito and T. Torrelli, Multiplier ideals, V -filtrations and transversal sections, *Math. Ann.* **336**(4) (2006) 901–924.
- [26] L. Ein, R. Lazarsfeld, K. E. Smith and D. Varolin, Jumping coefficients of multiplier ideals, *Duke Math. J.* **123**(3) (2004) 469–506.
- [27] A. Grothendieck, Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas (Quatrième partie), *Inst. Hautes Études Sci. Publ. Math.*, No. 32 (1967) 361.
- [28] A. G. Guimarães and A. Hefez, Bernstein–Sato polynomials and spectral numbers, *Ann. Inst. Fourier (Grenoble)* **57**(6) (2007) 2031–2040.
- [29] N. Hara, Geometric interpretation of tight closure and test ideals, *Trans. Amer. Math. Soc.* **353**(5) (2001) 1885–1906.
- [30] N. Hara and K.-I. Yoshida, A generalization of tight closure and multiplier ideals, *Trans. Amer. Math. Soc.* **355**(8) (2003) 3143–3174.
- [31] M. Hochster and C. Huneke, Tight closure and strong F -regularity, *Mém. Soc. Math. France (N.S.)*, No. 38 (1989) 119–133; *Colloque en l’honneur de Pierre Samuel* (Orsay, 1987).
- [32] M. Hochster and C. Huneke, Tight closure, invariant theory, and the Briançon–Skoda theorem, *J. Amer. Math. Soc.* **3**(1) (1990) 31–116.
- [33] M. Hochster and C. Huneke, F -regularity, test elements, and smooth base change, *Trans. Amer. Math. Soc.* **346**(1) (1994) 1–62.
- [34] M. Hochster and C. Huneke, Tight closure in equal characteristic zero, Unpublished, available at <http://www.math.lsa.umich.edu/~hochster/tcz.pdf> (1999).
- [35] J.-C. Hsiao and L. F. Matusevich, Bernstein–Sato polynomials on normal toric varieties, *Michigan Math. J.* **67**(1) (2018) 117–132.
- [36] A. A. Johnson, Multiplier ideals of determinantal ideals, PhD thesis, University of Michigan, ProQuest LLC, Ann Arbor, MI (2003), p. 83.
- [37] J.-M. Kantor, Formes et opérateurs différentiels sur les espaces analytiques complexes, *Bull. Soc. Math. France Mém.*, No. 53 (1977) 5–80.
- [38] M. Kashiwara, Algebraic study of systems of linear partial differential equations, Master’s thesis, University of Tokyo (1971).
- [39] M. Kashiwara, B -functions and holonomic systems. Rationality of roots of B -functions, *Invent. Math.* **38**(1) (1976/77) 33–53.
- [40] M. Kashiwara, Vanishing cycle sheaves and holonomic systems of differential equations, in *Algebraic Geometry*, Lecture Notes in Mathematics, Vol. 1016 (Springer, Berlin, Tokyo/Kyoto, 1982/1983), pp. 134–142.
- [41] J. Kollár, Singularities of pairs, in *Algebraic geometry — Santa Cruz 1995*, *Proc. Symp. Pure Math.* Vol. 62 (American Mathematical Society, Providence, RI, 1997), pp. 221–287.
- [42] R. Lazarsfeld, *Positivity in Algebraic Geometry II* (Springer-Verlag, Berlin, 2004).
- [43] T. Lvasseur and J. T. Stafford, Rings of differential operators on classical rings of invariants, *Mem. Amer. Math. Soc.* **81**(412) (1989) vi+117.
- [44] B. Malgrange, Sur les polynômes de I. N. Bernstein, in *Séminaire Goulaouic-Schwartz 1973–1974*, Équations aux dérivées partielles et analyse fonctionnelle, *Exp. No. 20* (Centre de Mathematics, École Polytechnique, Paris, 1974), p. 10.
- [45] B. Malgrange, Le polynôme de Bernstein d’une singularité isolée, in *Fourier Integral Operators and Partial Differential Equations*, Lecture Notes in Mathematics,

J. Álvarez Montaner et al.

- Vol. 459 (Colloquium International, University, Nice, 1974) (Springer, Berlin, 1975), pp. 98–119.
- [46] B. Malgrange, Polynômes de Bernstein–Sato et cohomologie évanescence, in *Analysis and Topology on Singular Spaces, II, III* (Luminy, 1981), *Astérisque* Vol. 101 (Société mathématique, France, Paris, 1983), pp. 243–267.
 - [47] G. Masson, Rings of differential operators and étale homomorphisms, PhD thesis, Massachusetts Institute of Technology, ProQuest LLC, Ann Arbor, MI (1991) .
 - [48] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian Rings*, Pure and Applied Mathematics (John Wiley & Sons, Chichester, 1987). With the cooperation of L. W. Small, A Wiley-Interscience Publication.
 - [49] Z. Mebkhout, *Le formalisme des six opérations de Grothendieck pour les $\mathcal{D}_X$ -modules cohérents*, Travaux en Cours, Vol. 35 (Hermann, Paris, 1989). With supplementary material by the author and L. Narváez Macarro.
 - [50] Z. Mebkhout and L. Narváez-Macarro, La théorie du polynôme de Bernstein-Sato pour les algèbres de Tate et de Dwork-Monsky-Washnitzer, *Ann. Sci. École Norm. Sup. (4)* **24**(2) (1991) 227–256.
 - [51] D. Milićić, Lectures on algebraic theory of D -modules, University of Utah course notes, available at <https://www.math.utah.edu/~milicic/Eprints/dmodules.pdf> (1986).
 - [52] L. N. Moncada Morales, Rings of differential operators, Master’s thesis, Centro de Investigación en Matemáticas, Guanajuato, Mexico (2018).
 - [53] I. M. Musson, Rings of differential operators on invariant rings of tori, *Trans. Amer. Math. Soc.* **303**(2) (1987) 805–827.
 - [54] M. Mustața and M. Popa, Hodge ideals and minimal exponents of ideals, *Rev. Roumaine Math. Pures Appl.* **65**(3) (2020) 327–354.
 - [55] M. Mustața and M. Popa, Restriction, subadditivity, and semicontinuity theorems for Hodge ideals, *Int. Math. Res. Not. IMRN*, No. 11 (2018) 3587–3605.
 - [56] M. Mustața and M. Popa, Hodge ideals, *Mem. Amer. Math. Soc.* **262**, No. 1268, (2019) v+80.
 - [57] M. Mustața and M. Popa, Hodge ideals for $\mathbf{Q}$ -divisors: Birational approach, *J. Éc. Polytech. Math.* **6** (2019) 283–328.
 - [58] M. Mustața and M. Popa, Hodge ideals for $\mathbf{Q}$ -divisors, V -filtration, and minimal exponent, *Forum Math. Sigma* **8** (2020), Paper No: e19, 41pp.
 - [59] M. Mustața, Bernstein–sato polynomials for general ideals vs. principal ideals, preprint (2019), arXiv:1906.03086 [math.AG].
 - [60] L. Núñez-Betancourt, On certain rings of differentiable type and finiteness properties of local cohomology, *J. Algebra* **379** (2013) 1–10.
 - [61] T. Oaku, An algorithm of computing b -functions, *Duke Math. J.* **87**(1) (1997) 115–132.
 - [62] T. Oaku and N. Takayama, Algorithms for D -modules — restriction, tensor product, localization, and local cohomology groups, *J. Pure Appl. Algebra* **156**(2–3) (2001) 267–308.
 - [63] T. Oaku, N. Takayama and U. Walther, A localization algorithm for D -modules, *J. Symbolic Comput.* **29**(4–5) (2000) 721–728.
 - [64] C. Peskine and L. Szpiro, Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck, *Inst. Hautes Études Sci. Publ. Math.*, No. 42 (1973) 47–119.
 - [65] C. Sabbah, D -modules et cycles évanescents (d’après B. Malgrange et M. Kashiwara), in *Géométrie algébrique et applications, III*, Travaux en Cours, Vol. 24 (La Rábida, 1984) (Hermann, Paris, 1987), pp. 53–98.

Bernstein–Sato functional equations, V -filtrations, and multiplier ideals of direct summands

- [66] M. Saito, Mixed hodge modules, *Publ. Res. Inst. Math. Sci.* **26**(2) (1990) 221–333.
- [67] M. Saito, On b -function, spectrum and rational singularity, *Math. Ann.* **295**(1) (1993) 51–74.
- [68] M. Saito, Multiplier ideals, b -function, and spectrum of a hypersurface singularity, *Compos. Math.* **143**(4) (2007) 1050–1068.
- [69] M. Sato, Theory of prehomogeneous vector spaces (algebraic part) — the English translation of Sato’s lecture from Shintani’s note, *Nagoya Math. J.* **120** (1990) 1–34. Notes by Takuro Shintani, Translated from the Japanese by Masakazu Muro.
- [70] M. Sato, M. Kashiwara, T. Kimura and T. Oshima, Microlocal analysis of prehomogeneous vector spaces, *Invent. Math.* **62**(1) (1980/81) 117–179.
- [71] G. W. Schwarz, Lifting differential operators from orbit spaces, *Ann. Sci. École Norm. Sup. (4)* **28**(3) (1995) 253–305.
- [72] K. Schwede, Test ideals in non- $\mathbb{Q}$ -Gorenstein rings, *Trans. Amer. Math. Soc.* **363**(11) (2011) 5925–5941.
- [73] K. Schwede and K. Tucker, On the behavior of test ideals under finite morphisms, *J. Algebraic Geom.* **23**(3) (2014) 399–443.
- [74] T. Shibuta, Algorithms for computing multiplier ideals, *J. Pure Appl. Algebra* **215**(12) (2011) 2829–2842.
- [75] K. E. Smith, F -rational rings have rational singularities, *Amer. J. Math.* **119**(1) (1997) 159–180.
- [76] K. E. Smith, The multiplier ideal is a universal test ideal, *Comm. Algebra* **28**(12) (2000) 5915–5929.
- [77] S. P. Smith, An example of a ring Morita equivalent to the Weyl algebra A_1 , *J. Algebra* **73**(2) (1981) 552–555.
- [78] S. P. Smith, The global homological dimension of the ring of differential operators on a nonsingular variety over a field of positive characteristic, *J. Algebra* **107**(1) (1987) 98–105.
- [79] S. P. Smith and J. T. Stafford, Differential operators on an affine curve, *Proc. London Math. Soc. (3)* **56**(2) (1988) 229–259.
- [80] S. Takagi, An interpretation of multiplier ideals via tight closure, *J. Algebraic Geom.* **13**(2) (2004) 393–415.
- [81] S. Takagi, Adjoint ideals along closed subvarieties of higher codimension, *J. Reine Angew. Math.* **641** (2010) 145–162.
- [82] S. Takagi and R. Takahashi, D -modules over rings with finite F -representation type, *Math. Res. Lett.* **15**(3) (2008) 563–581.
- [83] U. Walther, Algorithmic computation of local cohomology modules and the local cohomological dimension of algebraic varieties, *J. Pure Appl. Algebra* **139**(1–3) (1999) 303–321.
- [84] U. Walther, Algorithmic computation of de Rham cohomology of complements of complex affine varieties, *J. Symbolic Comput.* **29**(4–5) (2000) 795–839.
- [85] A. Yekutieli, An explicit construction of the Grothendieck residue complex, *Astérisque* (208) (1992) 127. With an appendix by Pramathanath Sastry.