

A NOTE ON THE DISSIPATION FOR THE GENERAL MUSKAT PROBLEM

BY

SUSANNA V. HAZIOT (*Department of Mathematics, Brown University, Providence, RI 02912*)

AND

BENOÎT PAUSADER (*Department of Mathematics, Brown University, Providence, RI 02912*)

This paper is dedicated to Professor C. Dafermos, with love and admiration.

Abstract. We consider the dissipation of the Muskat problem and we give an elementary proof of a surprising inequality of Constantin-Cordoba-Gancedo-Strain [J. Eur. Math. Soc. (JEMS) 15 (2013), pp. 201–227 and Amer. J. Math. 138 (2016), pp. 1455–1494] which holds in greater generality.

1. Introduction.

1.1. *The Muskat problem.* The general Muskat problem describes the dynamics of two immiscible fluids in a porous medium with different densities $\rho^\pm$ and viscosities $\mu^\pm$. Let us denote the interface between the two fluids by Σ and assume that it is the graph of a time-dependent function $\eta(x, t)$, i.e.

$$\Sigma_t = \{(x, \eta(x, t)) : x \in \mathbb{R}^d\}.$$

The associated time-dependent fluid domains are then given by

$$\begin{aligned}\Omega_t^+ &= \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \eta(x, t) < y < \underline{b}^+(x)\}, \\ \Omega_t^- &= \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \underline{b}^-(x) < y < \eta(x, t)\},\end{aligned}\tag{1.1}$$

where $\underline{b}^\pm$ are the parametrizations of the rigid boundaries

$$\Gamma^\pm = \{(x, \underline{b}^\pm(x)) : x \in \mathbb{R}^d\}.$$

Received October 17, 2022.

2020 *Mathematics Subject Classification.* Primary 35R35, Secondary 35Q35.

Key words and phrases. Muskat, Hele-Shaw, free boundary problems.

This material is based upon work supported by the National Science Foundation under Grant No. DMS-1929284 while the authors were in residence at the Institute for Computational and Experimental Research in Mathematics in Providence, RI, during the program “Hamiltonian Methods in Dispersive and Wave Evolution Equations”. The work of the first author was funded by the National Science Foundation through the award DMS-2102961. The work of the second author was partially supported by a Simons fellowship and by NSF Grant No. DMS-2154162.

Email address: susanna_haziot@brown.edu

Email address: benoit_pausader@brown.edu

The incompressible fluid velocity $u^\pm$ in each region is governed by Darcy's law:

$$\mu^\pm u^\pm + \nabla_{x,y} p^\pm = -(0, \rho^\pm), \quad \operatorname{div}_{x,y} u^\pm = 0 \quad \text{in } \Omega_t^\pm. \quad (1.2)$$

At the interface Σ_t , the normal velocity is continuous, the jump¹ of the pressure is related to the surface tension coefficient $\sigma \geq 0$ and the interface moves with the fluid: for $\nu = \frac{1}{\sqrt{1+|\nabla\eta|^2}}(-\nabla\eta, 1)$ the upward pointing unit normal to Σ_t ,

$$u^+ \cdot \nu = u^- \cdot \nu, \quad \llbracket p \rrbracket = \sigma \kappa(x) \quad \text{on } \Sigma_t, \quad \partial_t \eta = \sqrt{1+|\nabla\eta|^2} u^- \cdot \nu|_{\Sigma_t}, \quad (1.3)$$

where the mean curvature is given by

$$\kappa(x) := -\operatorname{div} \left(\frac{\nabla\eta}{\sqrt{1+|\nabla\eta|^2}} \right).$$

Finally, at the two rigid boundaries, the no-penetration boundary conditions are imposed:

$$u^\pm \cdot \nu^\pm = 0 \quad \text{on } \Gamma^\pm, \quad (1.4)$$

where $\nu^\pm = \pm \frac{1}{\sqrt{1+|\nabla b^\pm|^2}}(-\nabla b^\pm, 1)$ denotes the outward pointing unit normal to $\Gamma^\pm$. In case one or both of $\Gamma^\pm$ is empty (infinite depth), (1.4) is then replaced by the vanishing of ∇u at infinity.

As in [8, 12], we shall refer to the system (1.1)-(1.4) as the two-phase Muskat problem. When the top phase corresponds to vacuum, i.e. $\mu^+ = \rho^+ = 0$, the two-phase Muskat problem reduces to the one-phase Muskat problem and (1.3) becomes

$$p^- = \sigma \kappa(x) \quad \text{on } \Sigma_t, \quad \partial_t \eta = \sqrt{1+|\nabla\eta|^2} u^- \cdot \nu|_{\Sigma_t}.$$

In the following, we will make the following assumptions which are enough to use the trace theory developed in [10] (see also [12, Section 3 and Appendix A] for the notations and slight expansions):

(H1) either $\Gamma^\pm = \emptyset$ or $b^\pm \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ and $\operatorname{dist}(\Gamma^\pm, \Sigma) > 0$.

(H2) $\eta \in L^1 \cap \dot{W}^{1,\infty}(\mathbb{R}^d)$. In particular, $\eta \in H^{\frac{1}{2}} \subset \tilde{H}_\Theta^{\frac{1}{2}}$, the space of trace functions used in [12].

1.2. *A variational formulation for the velocities.* Starting from (1.2) and taking a divergence we get, with the incompressibility condition in (1.2), that u is related to the gradient of a harmonic function:

$$\begin{aligned} u^\pm &= -\frac{1}{\mu^\pm} \nabla q^\pm, \quad \Delta q^\pm = 0, \quad q^\pm = p + \rho^\pm y, \quad \text{in } \Omega^\pm, \\ \frac{1}{\mu^+} \partial_\nu q^+ &= \frac{1}{\mu^-} \partial_\nu q^-, \quad \llbracket q^\pm \rrbracket = \sigma \kappa(x) + \llbracket \rho \rrbracket \eta(x), \quad \text{on } \Sigma_t, \\ \partial_\nu q^\pm &= 0 \quad \text{on } \Gamma^\pm. \end{aligned} \quad (1.5)$$

We can introduce the kinetic energy

$$\mathcal{E}(t) := \mu \int_\Omega |u(x, t)|^2 dx = \mu^+ \int_{\Omega^+} |u^+(x, t)|^2 dx + \mu^- \int_{\Omega^-} |u^-(x, t)|^2 dx$$

¹Here and in the following, we denote the jump of a quantity at the interface to be

$$\llbracket f \rrbracket = f^- - f^+.$$

and using (1.5) and (1.3), we also obtain the energy dissipation equality

$$\frac{d}{dt} \left(\frac{\llbracket \rho \rrbracket}{2} \int_{\mathbb{R}^d} \eta^2(x, t) dx + \sigma \text{rArea}(\Sigma_t) \right) = -\mathcal{E}(t),$$

where we define the renormalized area to be

$$\text{rArea}(\Sigma_t) := \int_{\mathbb{R}^d} \left\{ \sqrt{1 + |\nabla \eta(x)|^2} - 1 \right\} dx.$$

The monotonicity of the L^2 norm is a classical important result for the Muskat equation [5] (see also [1] for a list of other monotone quantities). Using e.g. [12] and the notations therein, without surface tension, in infinite depth, and assuming a Rayleigh-Taylor condition, the energy dissipation can be recasted in the form

$$-\frac{d}{dt} \frac{\llbracket \rho \rrbracket}{2} \int_{\mathbb{R}^d} \eta^2(x, t) dx = \frac{1}{\mu^-} \int_{\mathbb{R}^d} \eta \cdot G^-(\eta) f^- dx = \frac{1}{\mu^+ + \mu^-} \|\nabla|\cdot|^{\frac{1}{2}} \eta\|_{L^2}^2 + O(\eta^3),$$

$$|O(\eta^3)| \lesssim F(\|\eta\|_{H^{2d+1}}) \|\nabla \eta\|_{L^\infty} \|\eta\|_{H^{\frac{1}{2}}}^2$$

which would suggest that the dissipation rate for the Muskat equation may control derivatives of the solutions and be a useful quantity in the analysis. However, the surprising result of [5, Section 2] (later expanded to $3d$ in [6]) shows that, at least in certain settings (no surface tension, absence of boundary, equal viscosities), a much simpler (and weaker) lower bound, in fact, holds:

$$0 \leq \mathcal{E}(t) \leq C \|\eta\|_{L^1(\mathbb{R}^d)}. \quad (1.6)$$

The purpose of this note is to give an elementary, variational, proof of (1.6) which extends this inequality to various settings.

THEOREM 1.1. Assume **(H1)** and **(H2)**, $\sigma \geq 0$, $\llbracket \rho \rrbracket > 0$ and $\min\{\mu^-, \mu^+\} > 0$, then there holds that

$$0 \leq \mathcal{E}(t) < \llbracket \rho \rrbracket^2 [\mu^+ \|\eta_-(t)\|_{L^1(\mathbb{R}^d)} + \mu^- \|\eta_+(t)\|_{L^1(\mathbb{R}^d)}]. \quad (1.7)$$

Moreover, the constants are optimal.

In the inequality above and in the rest of the paper, we define the positive and negative parts of a function with lower subscripts: $\eta_- := \min\{0, \eta\}$, $\eta_+ := \max\{\eta, 0\}$.

REMARK 1.2.

- (1) Although the constants in (1.7) are optimal, the inequality is never saturated, unless $\eta \equiv 0$.
- (2) A simple variation of the analysis allows to consider piecewise smooth interfaces and thus allows corners, which is significant in view of [9].
- (3) Straightforward modifications allow to consider other domains than $\mathbb{R}^d$, e.g. $\mathbb{T}^d$.

We finish with some open questions:

- (1) It would be interesting to understand the minimal analytic setting under which Theorem 1.1 holds. In particular, whether in the infinite depth setting, one can extend the result to $\eta \in L^1 \cap \dot{H}^{\frac{3}{2}}$, which seems to be optimal in view of recent advances in local well-posedness theory in [2, 3]. In case $\Gamma^\pm \neq \emptyset$, it would be interesting to know the largest space allowable for η and $\underline{b}^\pm$ in order for the boundary problem to be well-defined.

- (2) In case $x \in \mathbb{R}$ and η has different limits at $\pm\infty$ (as e.g. [7]), it seems that the Dirichlet energy is unbounded for infinite bottom, but can be finite in case $\|\underline{b}^+ - \underline{b}^-\|_{L^\infty} < \infty$. In this case, it would be interesting to investigate whether there is an analogue of Theorem 1.1 involving some renormalization of η .
- (3) In a similar spirit, it would be interesting to extend the above results to the case when the interface is not a graph, especially since such interfaces can dynamically form [4], and whether there is a connection with the problematic local well-posedness theory.
- (4) It would be great to clarify what is true and what is not in the one-fluid case, and whether the two-fluid problem is “more (or less) stable” in a sense to be made precise. We refer to [11] which may suggest that stronger bounds than (1.6) hold in the one-fluid case.

2. A minimization problem.

2.1. *The minimization problem.* All of our considerations are instantaneous; from now on, we will fix an interface $\eta \in L^1 \cap \dot{W}^{1,\infty}(\mathbb{R}^d)$. We would like to express $\mathcal{E}$ as the solution of the following minimization problem:

$$\mathfrak{m} := \min\{\mathbb{E}[f^+, f^-], (f^+, f^-) \in \mathcal{A}\}, \quad (2.1)$$

where the set of admissible pairs is given by

$$\mathcal{A} := \{(f^+, f^-) \in \dot{H}^1(\Omega^+) \times \dot{H}^1(\Omega^-) : \llbracket f \rrbracket = \eta(x) \llbracket \rho \rrbracket\}$$

and the energy is given by

$$\mathbb{E}[f^+, f^-] := \mu^+ \int_{\Omega^+} |\nabla_{x,y} f^+(x, y, t)|^2 dx dy + \mu^- \int_{\Omega^-} |\nabla_{x,y} f^-(x, y, t)|^2 dx dy.$$

LEMMA 2.1. Assume **(H1)** and **(H2)**. The affine set $\mathcal{A}$ is well-defined, non-empty, and every bounded sequence in $\mathcal{A}$ for the natural semi-norm has a subsequence that converges weakly.

Proof. Let $u(x, t) = (e^{t|\nabla|}\eta)(x)$ be the harmonic extension of η to the lower half-plane. Define

$$f_0^+(x, y) = 0, \quad \text{and} \quad f_0^-(x, y) = \llbracket \rho \rrbracket u(x, y - \eta(x)) \chi(r^{-1}(y - \eta(x))) \mathbf{1}_{\Omega^-}(x, y),$$

where $0 < r < \text{dist}(\Sigma, \Gamma^-)$ and $\chi \in C_c^\infty(-1, 1)$ is such that $\chi \equiv 1$ in a neighborhood of 0. We have that $(f_0^+, f_0^-) \in \mathcal{A}$, which is then non-empty. It is proved in [12, Proposition 3.3.] that $\dot{H}^1(\Omega^\pm)$ is a Hilbert space. Using [10, Theorem 5.1] or [12, Theorem A.1.], we see that $Tr^\pm : f^\pm \mapsto f^\pm(x, \eta(x))$ is well-defined and continuous from $\dot{H}^1(\Omega^\pm)$ into $\tilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d)$. Thus we see that $\mathcal{A} \subset \dot{H}^1(\Omega^+) \times \dot{H}^1(\Omega^-)$ is a closed affine subset of a Hilbert space. $\square$

LEMMA 2.2. Assuming **(H1)** and **(H2)**, there exists a unique minimizer of (2.1), (q^+, q^-) , and in addition, $q^\pm$ satisfies (1.5).

Proof. We refer to [12, Section 3.1.] for similar computations. Since $\mathcal{A}$ is affine and $\mathbb{E}$ is strictly convex, uniqueness is direct. Considering a subsequence and using Lemma 2.1, we

obtain the existence. Harmonicity inside the domain follows by considering variations supported inside $\Omega^\pm$, and the Neumann conditions at $\Gamma^\pm$ and at Σ follow similarly by considering perturbations supported near the top and bottom, and the interface, respectively. $\square$

2.2. Comparison principle and proof of the main theorem.

Proof of Theorem 1.1. Once we have established that the velocity field u comes from a minimizer, it suffices to find competitors in $\mathcal{A}$. A natural example is given by

$$f^+(x, y) := \llbracket \rho \rrbracket \max\{-y, 0\}, \quad f^-(x, y) := \llbracket \rho \rrbracket \max\{0, y\}. \quad (2.2)$$

The competitor (f^+, f^-) is admissible and we conclude that

$$\mathcal{E} < \mathbb{E}[f^+, f^-] = \llbracket \rho \rrbracket^2 [\mu^+ \|\eta_-\|_{L^1} + \mu^- \|\eta_+\|_{L^1}]$$

which gives (1.7). The inequality above is strict by uniqueness of the minimizer since (f^+, f^-) are not harmonic. The optimality of the bound (1.7) follows from Lemma 2.3. $\square$

2.3. *Extension to the one-fluid case.* In the one-fluid setting, the same argument gives a variational interpretation for $\mathcal{E}$. However, in this case, the competitor (2.2) only works for a wave of elevation: $\eta \geq 0$. In particular, this gives

$$\mathcal{E}_{1f} < \llbracket \rho \rrbracket^2 \mu \int_{\mathbb{R}^d} |\eta| dx.$$

In fact, this can also be obtained from the observation that² [12, Lemma 4.2.] $G(\eta)\eta < 1$.

2.4. *Optimality.* One may wonder whether (1.7) gives a bound which is optimal in any way. Simple scaling arguments show that it can be saturated, at least up to a multiplicative constant.

LEMMA 2.3. Assume **(H1)** and **(H2)**. There exists a sequence of smooth, compactly supported functions η_n for which the inequality in (1.7) is saturated in the sense that:

$$\mathcal{E}[\eta_n]/(\mu^+ \|(\eta_n)_-\|_{L^1} + \mu^- \|(\eta_n)_+\|_{L^1}) \rightarrow \llbracket \rho \rrbracket^2.$$

REMARK 2.4. It is interesting to note that one can construct scenarios for which $\mathcal{E} < \infty$ even though $\|\eta\|_{L^1} = \infty$.

Proof of Lemma 2.3. We will prove this in the case of a surface of elevation: $\eta \geq 0$. The case $\eta \leq 0$ follows similarly. In this case, we will only consider one domain and we define the Dirichlet energy of a function η on a domain Ω to be $\mathcal{D}_\Omega(\eta) := \int_\Omega |\nabla \eta|^2 dx dy$.

We first study the model case of a step function $\eta = \sigma := H \cdot \mathbf{1}_{\{0 \leq x \leq L\}}$, for some constant height H , when the computations can be carried out explicitly and extend by continuity of the Dirichlet energy under deformation of the domain. The arguments are elementary but we give the details for the sake of completeness. We consider the rectangle $R := \Omega^- \cap \{y \geq 0\} = \{0 \leq x \leq L, 0 \leq y \leq H\}$, and we also partition the boundary into $\partial R = \mathcal{T} \cup \mathcal{B}$ with “free” boundary at the bottom $\mathcal{B} := \{0 \leq x \leq L, y = 0\}$. Here $\mathcal{T}$ denotes the union of the top with the two vertical sides of the rectangle.

²We thank Huy Nguyen for this observation.

Let u_R be the harmonic function satisfying the Dirichlet boundary condition $u_R = y$ on $\mathcal{T}$ and the Neumann condition $\partial_y u_R = 0$ on $\mathcal{B}$. By standard arguments, u_R is the minimizer of the Dirichlet energy with Dirichlet condition on $\mathcal{T}$:

$$\mathcal{D}_R(u_R) := \min\{\mathcal{D}_R(g) : g(x, y) = y \text{ on } \mathcal{T}\}$$

and since f^- is admissible, we see that

$$\mathcal{D}_R(u_R) \leq \mathcal{D}_R(f^-) := \iint_R |\nabla f^-(x, y)|^2 dx dy \leq \iint_{\Omega^-} |\nabla f^-(x, y)|^2 dx dy.$$

The energy of u_R can be computed: letting $v = y - u_R$, we see that it satisfies 0 Dirichlet boundary condition on top and sides and the Neumann condition $\partial_y v = 1$ at the bottom. We can then expand into Fourier series to get

$$v(x, y) := \sum_{n \geq 1} a_n \sin\left(\frac{\pi n x}{L}\right) \sinh\left(\frac{n\pi}{L}(y - H)\right), \quad 1 = \sum_{n \geq 1} \frac{n\pi}{L} \cosh\left(\frac{n\pi}{L}H\right) a_n \sin\left(\frac{\pi n x}{L}\right).$$

The Fourier coefficients can easily be computed to be

$$a_n = \frac{2(1 + (-1)^n)}{n\pi} \frac{L}{n\pi} \frac{1}{\cosh\left(\frac{n\pi}{L}H\right)}$$

and we conclude that

$$\begin{aligned} \|\nabla_{x,y} v\|_{L^2(R)}^2 &= \sum_{n \geq 1} \frac{n^2 \pi^2}{4L} H \frac{\sinh(nx)}{nx} a_n^2 \\ &= (LH) \sum_{n \geq 1} \frac{1}{n^2 \pi^2} \frac{\left(\frac{\sinh(2nx)}{2nx}\right)}{\cosh(2nx) + 1} \end{aligned}$$

for $x = 2\pi H/L$. Notice that as $\frac{H}{L} \rightarrow \infty$, $\|\nabla_{x,y} v\|_{L^2(R)}^2 \rightarrow 0$. As a result, we see that

$$\begin{aligned} \mathcal{E} &> \mu^- \|\rho\|^2 \|\nabla_{x,y} u\|_{L^2(R)}^2 = \mu^- \|\rho\|^2 \|\nabla_{x,y} (y - v)\|_{L^2(R)}^2 \geq \mu^- \|\rho\|^2 LH (1 + o_{H/L \rightarrow \infty}(1)) \\ &> \mu^- \|\rho\|^2 \|\eta_+\|_{L^1} (1 + o(1)). \end{aligned}$$

It remains to modify η to have a smooth function. This can be done by choosing η as a smooth function bounded by two step functions of same height but slightly different support: $\sigma_1 \leq \eta \leq \sigma_2$ with $\|\sigma_2 - \sigma_1\|_{L^1} \ll \|\sigma_1\|_{L^1}$. We let R_1 be the rectangle associated to σ_1 and R_2 the rectangle associated to σ_2 and $\Omega_+^- := \Omega^- \cap \{y \geq 0\}$ so that $R_1 \subset \Omega_+^- \subset R_2$ and $\text{Area}(R_2 \setminus R_1) \ll \text{Area}(\Omega_+^-)$. We call u_{R_1} , u_{R_2} the minimizers of the Dirichlet energy as before, and given a function u defined on one of the above domains, we define u^{ext} its extension to a bigger domain by y (this is a Lipschitz function defined in a bounded domain). Then, we see that

$$\begin{aligned} \mathcal{D}_{R_2}(u_{R_2}) &< \mathcal{D}_{R_2}((f^-)^{ext}) = \mathcal{D}_{\Omega_+^-}(\eta) + \|\sigma_2 - \eta\|_{L^1}, \\ \mathcal{D}_{\Omega_+^-}(\eta) &< \mathcal{D}_{\Omega_+^-}(u_{R_1}^{ext}) = \mathcal{D}_{R_1}(u_{R_1}) + \|\eta - \sigma_1\|_{L^1}, \end{aligned}$$

and in particular

$$|\mathcal{D}_{\Omega_+^-}(\eta) - \|\eta\|_{L^1}| \leq 3\|\sigma_2 - \sigma_1\|_{L^1} \ll \|\eta\|_{L^1}.$$

This finishes the proof of Lemma 2.3. □

Acknowledgments. The authors would like to thank Prof. J. Gomez-Serrano for many stimulating discussions and constructive comments. They would like to thank Prof. I. Tice for interesting discussions and pointing out an omission in a first draft.

REFERENCES

- [1] T. Alazard, N. Meunier, and D. Smets, *Lyapunov functions, identities and the Cauchy problem for the Hele-Shaw equation*, Comm. Math. Phys. **377** (2020), no. 2, 1421–1459, DOI 10.1007/s00220-020-03761-w. MR4115021
- [2] T. Alazard and Q.-H. Nguyen, *Endpoint sobolev theory for the Muskat equation*, Arxiv preprint, arXiv:2010.06915, 2020.
- [3] T. Alazard and Q.-H. Nguyen, *On the Cauchy problem for the Muskat equation. II: Critical initial data*, Ann. PDE **7** (2021), no. 1, Paper No. 7, 25, DOI 10.1007/s40818-021-00099-x. MR4242131
- [4] Á. Castro, D. Córdoba, C. Fefferman, and F. Gancedo, *Breakdown of smoothness for the Muskat problem*, Arch. Ration. Mech. Anal. **208** (2013), no. 3, 805–909, DOI 10.1007/s00205-013-0616-x. MR3048596
- [5] P. Constantin, D. Córdoba, F. Gancedo, and R. M. Strain, *On the global existence for the Muskat problem*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 1, 201–227, DOI 10.4171/JEMS/360. MR2998834
- [6] P. Constantin, D. Córdoba, F. Gancedo, L. Rodríguez-Piazza, and R. M. Strain, *On the Muskat problem: global in time results in 2D and 3D*, Amer. J. Math. **138** (2016), no. 6, 1455–1494, DOI 10.1353/ajm.2016.0044. MR3595492
- [7] F. Deng, Z. Lei, and F. Lin, *On the two-dimensional Muskat problem with monotone large initial data*, Comm. Pure Appl. Math. **70** (2017), no. 6, 1115–1145, DOI 10.1002/cpa.21669. MR3639321
- [8] P. T. Flynn and H. Q. Nguyen, *The vanishing surface tension limit of the Muskat problem*, Comm. Math. Phys. **382** (2021), no. 2, 1205–1241, DOI 10.1007/s00220-021-03980-9. MR4227171
- [9] E. García-Juárez, J. Gómez-Serrano, H. Q. Nguyen, and B. Pausader, *Self-similar solutions for the Muskat equation*, Adv. Math. **399** (2022), Paper No. 108294, 30, DOI 10.1016/j.aim.2022.108294. MR4385135
- [10] G. Leoni and I. Tice, *Traces for homogeneous Sobolev spaces in infinite strip-like domains*, J. Funct. Anal. **277** (2019), no. 7, 2288–2380, DOI 10.1016/j.jfa.2019.01.005. MR3989147
- [11] H. Q. Nguyen, *Coercivity of the Dirichlet-to-Neumann operator and applications to the Muskat problem*, Preprint, arXiv:2206.02321.
- [12] H. Q. Nguyen and B. Pausader, *A paradiifferential approach for well-posedness of the Muskat problem*, Arch. Ration. Mech. Anal. **237** (2020), no. 1, 35–100, DOI 10.1007/s00205-020-01494-7. MR4090462