

Traffic noise inhibits inhibitory control in wild-caught songbirds

Christopher N Templeton^{1,2}, Amber O'Connor^{1,3}, Sarah Strack^{1,4}, Franco Meraz¹, Katri Herranen¹

¹Department of Biology, Pacific University, Forest Grove, OR, 97116, USA

²Corresponding author and lead contact: templeton@pacificu.edu

³Current address: Department of Surgery, Oregon Health Science University, Portland, OR, 97239, USA

⁴Current address: Carlson College of Veterinary Medicine, Oregon State University, Corvallis, OR, 97331, USA

Summary

Anthropogenic noise is ubiquitous across environments and can have negative effects on animals, ranging from physiology to community structure. Recent work with captive-bred zebra finches demonstrated that traffic noise also affects cognitive performance. We examined whether these results extend to animals that have experienced noise in the wild. We collected black-capped chickadees from areas frequently exposed to road traffic noise and tested them on a detour reaching task, a commonly used measure of inhibitory control. Those chickadees exposed to traffic noise playback had much lower performance on the task than control birds, indicating that noise negatively impacts inhibitory control. These data corroborate previously findings in lab-reared zebra finches. Further, these results suggest that prior experience with traffic noise is not sufficient for animals to habituate to noise and overcome its negative effects on cognitive performance. Instead, noise-induced cognitive effects might have broad impacts on animal species living in noise-polluted habitats.

Keywords

Animal cognition, anthropogenic noise, black-capped chickadee, detour reaching, traffic noise, urban ecology

Introduction

Rapid human population growth rates have driven a dramatic increase in anthropogenic noise pollution, which can have a variety of negative impacts on wildlife ^{1,2}. Sound can directly impact animal anatomy and physiology, for example high intensity sound waves can physically damage animal auditory systems ³. Sound can also indirectly impact animals, by influencing their behavior, and while these effects can be less obvious, they can still have significant impacts on individuals, populations, and communities. For example, because many animals use sound to communicate information, anthropogenic noise has been widely shown to negatively impact animal behavior by masking acoustic signals related to sexual reproduction ⁴, begging calls ⁵, and alarm calls ^{6,7}. Noise pollution modifies animal stress response, gut microbiome, reproductive success, habitat use, and predator-prey interactions, suggesting that the acoustic environment can influence both individual fitness and population structures ^{1,8-11}

In addition to the ecological effects of noise pollution, noise can also impact cognitive processing. These impacts were first observed in humans, with several studies suggesting that individuals in noisy environments, for example children attending school near airports, tended to perform poorer on cognitive assessments ^{12,13}. While this topic has been less studied in animal systems, there is

recent evidence that noise also impacts cognitive performance in animals. Prolonged playback of high-amplitude, white noise produced oxidative tissue damage in the brains of mice, in turn reducing their cognitive performance¹⁴. Chronic noise exposure also causes stress and sleep disturbances, in mice and rats respectively, both of which in turn impact cognitive processes^{15,16}. Experimental playback of noise during development resulted in decreased song learning accuracy in zebra finches (*Taeniopygia guttata*), with further impacts on immune function¹⁷ and smaller brain regions related to vocal learning¹⁸. In addition to damaging brains' anatomy and physiology, noise could impact cognitive performance by distracting animals from other tasks^{19,20} or increasing vigilance²¹⁻²³. Thus, anthropogenic noise pollution could have cognitive impacts in animals similar to those observed in humans.

Recent research indicates that anthropogenic noise pollution may have substantial negative impacts on avian cognitive processes. Osbrink et al.²⁴ played road traffic noise at realistic amplitudes to captive-bred zebra finches and demonstrated that just hearing traffic noise significantly reduced cognitive performance on a variety of different foraging tasks. While zebra finches are an exceptionally useful study system that has been used to address many questions related to perception and learning²⁵, they are also domesticated lab animals that have been in captivity for over a hundred generations²⁶ and it is therefore important to determine how generalizable previous results are to other species^{27,28}. Further, although the Osbrink et al.²⁴ findings suggest the potential for widespread impacts of noise on cognitive processes in animals, we know little about how these results actually translate to animals living in the real world, where noise exposure can vary across both space and time.

To begin examining the extent to which anthropogenic noise impacts cognitive processing in other animal species, we expand this line of research by focusing on another small songbird, the black-capped chickadee (*Poecile atricapillus*). Chickadees are an ideal study system for this type of research for a number of reasons. First, they are common and widely-distributed throughout much of North America, where they frequently inhabit both urban areas and roadside habitats as well as more natural habitats²⁹. Next, cognitive behavior of chickadees, and closely related species, has been heavily researched, both in the laboratory and in natural environments³⁰⁻³³. Last, it has been suggested that their cognitive ability and cognitive flexibility have allowed chickadees to succeed in novel environments, including urban and urbanizing areas³⁴. Previous work has shown that chickadee communication is negatively impacted by noise³⁵⁻³⁷, but the impact of this noise on their cognition has not been studied.

Here we use wild-caught chickadees to test whether the negative impacts of traffic noise on cognitive processing observed in zebra finches²⁴ are generalizable to other songbirds. We presented chickadees with a detour reaching task that has been broadly used across human and animal systems to investigate inhibitory control, the ability to inhibit prepotent but ultimately counterproductive behavior³⁸. Inhibitory control is related to brain size, correlates positively with behavioral flexibility, and underlies other types of learning³⁹⁻⁴². This measure, therefore has been suggested to provide a useful window into problem solving, planning, and flexibility^{39,41}, important traits for adapting to changing environments, such as those presented in human-dominated urban environments^{43,44}. We collected chickadees from locations near moderately busy highways to ensure that they had prior experience with traffic noise in the wild and tested them on a detour reaching task in the lab under experimentally manipulated levels of noise. If the patterns observed in Osbrink et al.²⁴ are generalizable to other songbirds, we predicted that experimentally induced traffic noise exposure would also decrease cognitive performance observed in chickadees. Further, the magnitude of difference between control birds and noise-exposed birds compared with previous work on captive bred zebra finches should provide insights into the ability of animals to

habituate to noise pollution with prior experience, potentially mediating its negative cognitive effects. If prior experience with traffic noise allows chickadees to overcome the negative effects of noise pollution, then we predict that the difference in cognitive performance between control and noise-exposed birds should be reduced compared with that observed in zebra finches. If habituation to prior noise pollution does not enable animals to overcome the negative cognitive effects of noise, then we expected to find patterns similar to those previously observed in captive-bred zebra finches.

Results

The presence of experimental traffic noise significantly decreased the performance of birds on the detour reaching task ($t_{15} = 5.85$, $p < 0.0001$, Cohen's $d = 2.83$). Those birds performing the task in the presence of traffic noise were more than twice as likely to fail a given trial (mean $\pm$ SEM: $84.20 \pm 5.8\%$ failed trials compared with $36.5 \pm 5.7\%$ failed for control birds; Figure 1), by pecking at the clear cylinder before detouring to the open end. Control birds rapidly learned to inhibit their pecking response and detour around the clear cylinder, beginning to consistently solve the task after just a few trials (mean $\pm$ SEM: 4.5 ± 0.68 trials to the first of three successes in a row). In contrast, while birds exposed to experimental traffic noise did solve the task, none of the individuals managed to achieve the criteria of three successful trials in a row.

In contrast to trial performance, participation rates did not differ significantly between treatments ($t_{15} = 1.47$, $p = 0.172$, Cohen's $d = 0.682$). Participation rates were generally high for all birds, with control birds participating in $96.25 \pm 2.63\%$ of trials compared and noise-exposed birds participating in $83.33 \pm 8.33\%$ of trials.

Discussion

Chickadees exposed to experimental traffic noise had reduced performance on a detour reaching task. Compared with individuals tested under quiet control conditions, those birds exposed to noise were less than half as likely to correctly solve the task. In contrast, participation rates, a likely correlate of neophobia and stress level, did not significantly vary between treatments. These results indicate that the negative impacts of traffic noise on cognitive processing previously observed in zebra finches extend to detour reaching performance in wild-caught chickadees.

Given that performance on detour reaching has been shown to be a reliable indicator of the behavioral flexibility and problem-solving capabilities^{42,45–48}, our results suggest that anthropogenic noise could negatively affect black-capped chickadees inhabiting noise polluted environments in the wild. Previous work has demonstrated that chickadees change their behavior in the presence of noise, altering vocalizations³⁷, vigilance and foraging⁴⁹, and use of alarm signals^{35,50,51}. Closely related mountain chickadees (*P. gambeli*) shift home ranges away from noise sources when exposed to traffic noise in the wild⁵². Despite these effects on behavior, chickadees and other Parids have relatively highly-developed cognitive abilities^{32,53–55}, which have helped them survive in a variety of human-dominated landscapes, such as urban areas and those near roads²⁹. These environments typically provide novel food sources, such as bird feeders, roadkill carrion, or garbage containers, and the ability of chickadees to effectively exploit these novel problems has helped them succeed in these environments. Yet, our results demonstrating diminished performance on detour reaching suggests that some of the very cognitive traits that help chickadees and other common urban species^{44,56,57} adapt to human-dominated environments could be negatively affected by anthropogenic noise. Does noise pollution counteract these species' cognitive adaptations or have disproportionate effects on more cognitively-advanced species? Better

understanding how noise impacts detour reaching could provide a useful window into the cognitive flexibility necessary for chickadees and other species to overcome the challenges posed by urban areas and other human-dominated environments⁵⁸⁻⁶⁰.

The specific mechanisms driving noise-induced cognitive inhibition are not yet clear. Noise pollution is known to induce a stress response in animals⁶¹⁻⁶⁴ and is associated with increased vigilance^{2,21} and it is possible that these could negatively impact cognitive performance despite likely being adaptive responses. Further, a change in foraging behavior, such as avoiding foraging tactics that pose higher risk of predation, could be adaptive under noisy conditions. While we have not measured stress levels or vigilance behavior in this study directly, it seems likely that these would impact the degree to which animals participate in trials in addition to cognitive performance. Yet, we observed no significant effect of noise pollution on participation rates between experimental and control birds. This lack of difference in participation suggests that the impact of noise pollution on cognitive performance was not primarily driven by stress, increased vigilance, or other adaptive changes in behavior under noise. While the mechanisms in which noise pollution diminishes cognitive performance are not yet clear, another reasonable hypothesis would be that noise reduces attention to the task by acting as a distractor¹⁹. Future work should further examine the mechanisms underlying noise-induced decreases in cognitive performance.

Our results suggest the potential for the cognitive effects of noise to translate to negative impacts on fitness of animals living in environments polluted with traffic noise. While there are many types of cognitive function potentially relevant to individual success⁶⁵, detour reaching could provide a useful window into the cognitive flexibility necessary to overcome new challenges in human-dominated environments. As detour reaching tasks assess the ability of an individual to inhibit their first instinct and instead find alternative solutions to a problem, the decrease in inhibitory control that our subjects demonstrated in the presence of traffic noise implies that traffic noise interferes with black-capped chickadees' ability to quickly and efficiently solve problems. While the link between an individual's cognitive ability and its fitness has been little tested, there is some evidence that cognitive ability predicts fitness. For example, Ashton et al.⁶⁶ demonstrated that cognitive performance of female Australian magpies (*Cracticus tibicen dorsalis*) on a similar detour reaching and other cognitive tasks predicted reproductive success. Cole et al.³³ showed that performance on novel problem-solving tasks also correlated with reproductive success in great tits (*Parus major*). Although the extent to which individual cognitive abilities impact fitness are still under debate⁶⁵, it seems likely that a significant reduction in cognitive performance due to noise pollution could affect an animal's fitness, regardless of its initial cognitive abilities. In a natural environment polluted by traffic noise, the extra time and energy necessary to solve problems could be fatal, especially to small avian species like black-capped chickadees that have high metabolic rates⁶⁷. In addition to foraging-related cognition, it is likely that noise pollution could also impact other aspects of cognitive behavior in these and other animals. For example, song learning is generally thought to be a different cognitive module from the sorts of foraging tasks described here⁶⁸, but is also known to be impacted by anthropogenic noise pollution^{18,69}. Similarly, learning to discriminate among auditory stimuli, such as contact calls⁷⁰ or sexual signals⁷¹ are further examples of the potential for noise to disrupt a variety of cognitive processes in animals.

The findings with chickadees extend previous research demonstrating that lab-reared zebra finches have diminished cognitive performance when exposed to traffic noise²⁴. In addition to demonstrating that these noise-induced cognitive effects could be generalizable to other songbird species, comparing the magnitude of the effect could shed light on the ability of wild birds to adapt to noise pollution through habituation (Figure 2). Surprisingly, both chickadees and zebra finches

participating in the detour reaching task under quiet control conditions had nearly identical performance levels, with failure rates of $36.5 \pm 5.7\%$ (chickadees) and $35.0 \pm 5.8\%$ (zebra finches). This suggests that on average, individuals of both species would exhibit similar levels of inhibitory control in the absence of noise pollution. However, chickadees exposed to noise pollution had much lower levels of performance ($84.2 \pm 5.8\%$ failure rate) compared with zebra finches ($53.4 \pm 5.3\%$ failure rate). Chickadee participation rates (control: $96 \pm 3\%$ and noise: $88 \pm 8\%$ of trials) were similar to those previously observed in zebra finches (control: $96 \pm 4\%$ and noise: $92 \pm 4\%$; Osbrink et al. ²⁴), suggesting that noise had similar and relatively minor effects on neophobia for both species, despite differentially impacting cognitive performance between species. In addition to species differences, the birds used in these two studies had differences in their developmental history, with finches reared in the lab environment and chickadees reared in the wild and brought into the lab for testing. While chickadees were habituated to the lab environment before the experiment began and we observed no difference in performance of control birds compared with zebra finches, it could be that the testing environment was more novel for chickadees, heightening their vigilance levels so that the noise playback had a stronger impact. Another difference in rearing environment was the presence of noise pollution. Finches raised in the lab had little experience with traffic noise, though the chickadees were all collected from locations in which they would have previously heard traffic noise in the wild. The findings that noise playback had similar or even greater impact on chickadees that had previous experience with traffic noise suggests that any habituation to noise pollution developed through previous experience does not alleviate the effects of noise on animal cognition, at least in the way it was tested in this experiment. It will be important for future work to compare responses of chickadees from quiet environments to further understand the effects of potential habituation to noise pollution in this species. Quinn et al. ²¹ demonstrated a similar lack of habituation to repeated noise playback in relationship to vigilance behavior, suggesting that animals might not be able to tune out noise pollution even with previous experience.

In addition to the prior experience and species of birds, other factors could potentially help explain the variation in effect size observed between the present data and the previous study ²⁴. Chickadees used in this experiment were not provided with prior experience with detour reaching tasks following a protocol similar to Ashton et al. ⁶⁶, whereas Osbrink et al. ²⁴ provided zebra finches with prior exposure to an opaque version of the task ³⁸. Because chickadees are less neophobic, we elected to avoid the training phase, as it has been suggested that this prior exposure can make detour reaching results more challenging to interpret because success with the clear cylinder could result from inhibitory control or from a holdover of a previously-learned rule from the prior training ⁶⁶. Chickadees were given ten trials with the detour reaching task, compared with four trials (after training) in Osbrink et al. ²⁴, which provided more opportunities for chickadees to learn about the cylinder and helped counteract for the lack of previous experience. Further, if this difference in prior exposure was the major factor determining the outcome of the task, then we would have expected to observe species differences in the control group in addition to the noise group, which were not detected. It therefore seems likely that the differences in neophobia and prior training likely counterbalance each other between the two species, making the data reasonably comparable between experiments. Future work examining the degree to which different species respond to noise pollution would further our understanding of the implications of noise-induced cognitive decline for different species inhabiting anthropogenic landscapes.

Limitations of the study

While this study has now shown that the effects of traffic noise exposure on cognitive performance extend beyond lab-reared birds, further research would be beneficial to better understand how

noise pollution might affect cognitive behavior of wild animals. First, only one type of cognitive task—inhibitory control—was employed in this study. Inhibitory control is a good first measure because it is correlated with brain size³⁸, is related to planning, problem-solving, and judgement³⁹, and has been proposed to be a foundational ability that underlies many other types of cognition⁴¹. However, the repeatability of this task is currently debated in the literature^{72–75}, so future work demonstrating repeatability of these scores for chickadees would be useful. Future research could also benefit from applying a battery of different tasks that assess different aspects of cognition, especially those cognitive components that are particularly relevant to the species of interest^{65,76}. For highly social animals with cognitive adaptations for caching, some of the most relevant tasks for chickadees might focus on assessing social learning and spatial memory^{54,77,78}. Future work examining the impact of traffic noise should ensure that the tasks are ecologically relevant to the biology of the particular study species being used^{65,76}. In addition to measuring performance on only one task, we only assessed birds with similar levels of prior exposure to traffic noise. All birds were collected from locations where they had previously been exposed to intermediate levels of traffic noise. While these results suggest that this level of previous exposure was not sufficient to habituate to the cognitive effects of traffic noise, it is possible that the effects we observed would vary for birds that had experienced different traffic noise regimes during development. For example, chickadees living in more highly urbanized environments are more tolerant of noise playback at feeding stations⁷⁹. It is also possible that animals habituate to very specific ambient noise profiles in their environments and if this is the case, the birds' previous experience with noise would not necessarily generalize to help them overcome the experimental noise conditions used in this study. Future work will examine birds originating from different noise environments, including habitats with both noisier and more acoustically pristine conditions, will better determine how noise regimes affect the cognitive impacts of noise pollution.

Acknowledgements

We thank May Murakami-Smith and Carrie Nunnenkamp for assistance with data collection, Ali Osbrink and Lauren Chan for helpful discussions, and Ursula Valdez and Matt Chemielski for assistance capturing birds. Funding was provided by NSF (IOS-2207395), the MJ Murdock Charitable Trust, and Pacific University.

Author contributions

CNT: project conception and development, data collection, data archiving, data analysis, manuscript writing, funding acquisition, and supervision. AOC: Project development, data collection, data archiving, animal care, manuscript writing. SS: Project development, experimental design, data collection, data archiving, animal care, manuscript writing. FM: Project development, experimental design, data collection, data archiving, animal care, manuscript writing. KH: data collection, data archiving, animal care.

Declaration of interests

The authors declare no competing interests

Inclusion and diversity

We support inclusive, diverse, and equitable conduct of research

Figure Legends

Figure 1. Traffic noise negatively impacts inhibitory control in black-capped chickadees. A) Birds hearing traffic noise playback were more than twice as likely to fail trials on a detour reaching task, although B) their participation rate did not significantly differ from control birds. Data are represented as mean $\pm$ SEM, with individual data points depicted for each bird.

Figure 2. Wild-caught chickadee detour reaching performance compared with that of lab-reared zebra finches. Despite control birds of each species having similar failure rates, chickadees exposed to traffic noise playback had higher failure rates than those observed in zebra finches. Data are represented as mean $\pm$ SEM percent of trials. Zebra finch data adapted from Osbrink et al. ²⁴.

STAR Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Chris Templeton (templeton@pacificu.edu)

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Study Animals and Housing

Wild black-capped chickadees were captured from locations on and within 3km of the Pacific University campus in Forest Grove, Oregon USA (45.5204N, -123.1093W). All birds were collected in locations in which they would have been previously exposed to traffic noise from adjacent highways (80-300m from collection sites) and other roads that feature frequent (average daily vehicles traveling in each direction: 735 ± 354 ; ⁸⁰) medium speed traffic from vehicles ranging from passenger cars to heavy logging trucks. While we did not collect specific data on the specific traffic noise profiles of these sites, they were generally similar to the playback stimuli (described below). A total of 17 individuals from four different winter flocks were captured using mistnets between February 2019 and March 2020. We were not able to accurately determine sex of the subjects because chickadees are sexually monomorphic ²⁹, though like previous researchers ⁵⁴, we had no reason to believe that males and females would perform differently on this task and did not deem it worth conducting the invasive tests necessary to determine sex. Exact ages were also unknown, but all birds were independent adults at the time of capture.

Birds were housed with their flock mates in a covered outdoor aviary on the Pacific University campus. The aviary was made from a wood frame with aviary netting and measured 2.3 x 3.8 x 2.2m high. The aviary contained numerous plants, perches, a roosting box, and fresh water provided from a pool with a fountain. Chickadees were given *ad lib* access to sunflower seeds, suet, and peanuts, with live mealworms also provided twice a day. Each bird was allowed to habituate to

captivity for at least one week prior to testing and birds were kept in the aviary except when they were actively being tested.

Ethical Statement

All work described in this paper was approved by Pacific University IACUC (910252) and conforms to ABS/ASAB guidelines for treatment of animals in behavioral research.

METHOD DETAILS

Experimental Preparation

To measure inhibitory control, we constructed a detour reach task that was based on the task used in numerous other studies with birds and other animals³⁸. The task consisted of a clear 11.5cm long x 6.3cm diameter tube made from thin plastic that was glued to a 13cm x 18cm wooden board.

We randomly split birds into two groups, those exposed to experimental traffic noise, and control birds receiving no noise playback. To ensure that we could compare our data directly with those data collected on zebra finches, we followed the detour reaching procedure established in Osbrink et al.²⁴ and used the same noise playback stimuli found in that study. The chickadees in the traffic noise group received playback of road noise, which featured four exemplars of mostly continuous noise recorded from a moderately busy highway. To increase external validity, we used playback recordings that were derived from natural traffic recordings and presented at realistic sound levels^{17,24,81}. Recording details and complete noise profile information is included in Templeton et al.⁶, but briefly, recordings were made approximately 10 m from a rural two-lane highway in Germany with moderate levels of traffic and road noise using a Sennheiser ME66/K6 microphone and Marantz PMD660 digital recorder. We edited the recordings to create 30s playback files that included mostly constant noise using the 'Mix Paste' and 'Crossfade' functions in Adobe AUDITION 3.0 (Adobe Inc., San Jose, CA, USA). Traffic noise was broadcast from a Pignose 7-100 speaker (Pignose Amps, Las Vegas NV USA) located 1m from the focal bird's cage that was calibrated with a Cel-246 SPL meter (L_{AFmax} setting, maximum sound pressure level, frequency weighting: A, time weighting: F; 20 μ PA reference value) so that the noise amplitude at the location of the bird was 70dB (to realistically simulate cars passing about 30m from the bird;⁶. The speaker was also present for the control birds to control for any visual effect of the speaker, but no sound was played during these trials (ambient testing room amplitude ~30dB).

Testing Procedure

Individuals were captured from the aviary and placed in a testing cage (45 × 76 × 45 cm) located in an adjacent room that was acoustically and visually isolated from the other birds in the aviary. Each bird was given half an hour to habituate to the test cage with ad lib access to food and water. After habituation, food was removed for an additional 45 minute period to help increase motivation during the following experimental trials.

Prior to the trial, the bird was sequestered in one side of the cage using an opaque divider, so that the task apparatus could be placed on the other side of the cage out of view. Also out of view of the chickadee, mealworm halves were placed inside the clear tube of the detour reaching task at $\frac{1}{4}$, $\frac{1}{2}$, and $\frac{3}{4}$ of the length of the tube, so that a chickadee needed to reach its head into either end of the tube to access a food reward and each bird could obtain up to 1.5 mealworms per successful trial. The tube was positioned in the side of the cage that did not contain the chickadee, oriented so that the side of the tube, not the openings, faced towards the chickadee when the divider was removed. We conducted 10 trials per chickadee, each lasting as long as it took for the subject to interact with

the apparatus but not exceeding 5 minutes. Between each trial, there was a 5 minute break where the bird was again sequestered to one side and we restocked the mealworms in the task before the next trial. For the experimental subjects, traffic noise was played during the foraging trials, but not during breaks, following Osbrink et al. ²⁴.

Trials were scored as successful if the bird detoured around to the open side of the tube without first attempting to obtain the mealworm through the side of the tube. If birds interacted with the side of the tube (e.g., pecking, flapping, scratching with feet, etc.) in an attempt to obtain the mealworm, we considered the trial a fail. If the subject made no attempt to obtain the mealworm during the 5 minute period, this trial did not count as a pass or fail and non-participation was excluded from the detour reaching score.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical Analysis

To determine whether traffic noise impacted subjects' inhibitory control, we compared the percent of trial successes and failures between the noise and control groups using an independent samples t-test. We tested a total of 8 birds in the control treatment and 9 birds in the noise treatment (n=17 birds). In addition to examining the percent of successful trials for each bird, we measured the number of trials required to become proficient with the task, as evidenced by successfully detouring on three trials in a row ⁶⁶. In addition to comparing the detour reaching scores, we assessed participation rates of each group to ensure that any apparent differences in cognitive performance were not due to lack of participation. Again, we used a t-test to compare participation rates, but because the two groups did not have equal variances (Levene's test, $F=8.56$, $p=0.01$), we used a Welch's t-test which does not assume equal variances. All statistics were computed in SPSS v. 28.0 (IBM Corp, Armonk, NY) and all tests report two-tailed p values.

References

1. Kight, C.R., and Swaddle, J.P. (2011). How and why environmental noise impacts animals: an integrative, mechanistic review. *Ecol. Lett.* 14, 1052–1061. 10.1111/j.1461-0248.2011.01664.x.
2. Shannon, G., McKenna, M.F., Angeloni, L.M., Crooks, K.R., Fristrup, K.M., Brown, E., Warner, K.A., Nelson, M.D., White, C., Briggs, J., et al. (2016). A synthesis of two decades of research documenting the effects of noise on wildlife. *Biol. Rev.* 91, 982–1005. 10.1111/brv.12207.
3. Liberman, M.C., and Beil, D.G. (1979). Hair cell condition and auditory nerve response in normal and noise-damaged cochleas. *Acta Otolaryngol.* 88, 161–176. 10.3109/00016487909137156.
4. Brumm, H., and Slabbekoorn, H. (2005). Acoustic Communication in Noise. *Adv. Study Behav.* 35, 151–209. 10.1016/S0065-3454(05)35004-2.
5. McIntyre, E., Leonard, M.L., and Horn, A.G. (2013). Ambient noise and parental communication of predation risk in tree swallows, *Tachycineta bicolor*. *Anim. Behav.* 87, 85–89.
6. Templeton, C.N., Zollinger, S.A., and Brumm, H. (2016). Traffic noise drowns out great tit alarm calls. *Curr. Biol.* 26, R1173--R1174. 10.1016/j.cub.2016.09.058.
7. Zhou, Y., Radford, A.N., and Magrath, R.D. (2019). Why does noise reduce response to alarm calls? Experimental assessment of masking, distraction and greater vigilance in wild birds.


- 445 Funct. Ecol. 33, 1280–1289. 10.1111/1365-2435.13333.
- 446 8. Barber, J.R., Burdett, C.L., Reed, S.E., Warner, K.A., Formichella, C., Crooks, K.R., Theobald,
447 D.M., and Fristrup, K.M. (2011). Anthropogenic noise exposure in protected natural areas:
448 Estimating the scale of ecological consequences. *Landsc. Ecol.* 26, 1281–1295.
449 10.1007/s10980-011-9646-7.
- 450 9. Benítez-López, A., Alkemade, R., and Verweij, P.A. (2010). The impacts of roads and other
451 infrastructure on mammal and bird populations: A meta-analysis. *Biol. Conserv.* 143, 1307–
452 1316. 10.1016/j.biocon.2010.02.009.
- 453 10. Fahrig, L., and Rytwinski, T. (2009). Effects of roads on animal abundance: An empirical
454 review and synthesis. *Ecol. Soc.* 14. 10.5751/ES-02815-140121.
- 455 11. Injaian, A.S., Poon, L.Y., and Patricelli, G.L. (2018). Effects of experimental anthropogenic
456 noise on avian settlement patterns and reproductive success. *Behav. Ecol.* 29, 1181–1189–
457 1181–1189.
- 458 12. Clark, C., and Paunovic, K. (2018). WHO environmental noise guidelines for the european
459 region: A systematic review on environmental noise and cognition. *Int. J. Environ. Res. Public*
460 *Health* 15, 2400. 10.3390/ijerph15112400.
- 461 13. Stansfeld, S.A., Berglund, B., Clark, C., Lopez-Barrio, I., Fischer, P., Öhrström, E., Haines, M.M.,
462 Head, J., Hygge, S., Van Kamp, I., et al. (2005). Aircraft and road traffic noise and children's
463 cognition and health: A cross-national study. *Lancet* 365, 1942–1949. 10.1016/S0140-
464 6736(05)66660-3.
- 465 14. Cheng, L., Wang, S.-H.H., Chen, Q.-C.C., and Liao, X.-M.M. (2011). Moderate noise induced
466 cognition impairment of mice and its underlying mechanisms. *Physiol. Behav.* 104, 981–988.
467 10.1016/j.physbeh.2011.06.018.
- 468 15. Jafari, Z., Kolb, B.E., and Mohajerani, M.H. (2018). Chronic traffic noise stress accelerates
469 brain impairment and cognitive decline in mice. *Exp. Neurol.* 308, 1–12.
470 10.1016/j.expneurol.2018.06.011.
- 471 16. Rabat, A., Bouyer, J.J., George, O., Le Moal, M., and Mayo, W. (2006). Chronic exposure of rats
472 to noise: Relationship between long-term memory deficits and slow wave sleep
473 disturbances. *Behav. Brain Res.* 171, 303–312. 10.1016/j.bbr.2006.04.007.
- 474 17. Brumm, H., Goymann, W., Derégnaucourt, S., Geberzahn, N., and Zollinger, S.A. (2021). Traffic
475 noise disrupts vocal development and suppresses immune function. *Sci. Adv.* 7.
476 10.1126/sciadv.abe2405.
- 477 18. Potvin, D.A., Curcio, M.T., Swaddle, J.P., and MacDougall-Shackleton, S.A. (2016).
478 Experimental exposure to urban and pink noise affects brain development and song learning
479 in zebra finches (*Taenopygia guttata*). *PeerJ* 2016, 1–21. 10.7717/peerj.2287.
- 480 19. Chan, A.A.Y.-H., Giraldo-Perez, P., Smith, S., and Blumstein, D.T. (2010). Anthropogenic noise
481 affects risk assessment and attention: the distracted prey hypothesis. *Biol. Lett.* 6, 458–461.
482 10.1098/rsbl.2009.1081.
- 483 20. Morris-Drake, A., Kern, J.M., and Radford, A.N. (2016). Cross-modal impacts of anthropogenic
484 noise on information use. *Curr. Biol.* 26, R911–R912. 10.1016/j.cub.2016.08.064.
- 485 21. Quinn, J.L., Whittingham, M.J., Butler, S.J., and Cresswell, W. (2006). Noise, predation risk
486 compensation and vigilance in the chaffinch *Fringilla coelebs*. *J. Avian Biol.* 37, 601–608.
487 10.1111/j.2006.0908-8857.03781.x.
- 488 22. Meillere, A., Brischoux, F., and Angelier, F. (2015). Impact of chronic noise exposure on
489 antipredator behavior: An experiment in breeding house sparrows. *Behav. Ecol.* 26, 569–
490 577. 10.1093/beheco/aru232.
- 491 23. Klett-Mingo, J.I., Pavón, I., and Gil, D. (2016). Great tits, *Parus major*, increase vigilance time
492 and reduce feeding effort during peaks of aircraft noise. *Anim. Behav.* 115, 29–34.
493 10.1016/j.anbehav.2016.02.021.
- 494 24. Osbrink, A., Meatte, M.A., Tran, A., Herranen, K.K., Meek, L., Murakami-Smith, M., Ito, J.,

- Bhadra, S., Nunnenkamp, C., and Templeton, C.N. (2021). Traffic noise inhibits cognitive performance in a songbird. *Proc. R. Soc. B* 288, 20202851.
25. Hauber, M.E., Louder, M.I., and Griffith, S.C. (2021). The natural history of model organisms neurogenomic insights into the behavioral and vocal development of the zebra finch. *Elife* 10, 1–19. 10.7554/eLife.61849.
26. Zann, R.A. (1996). *The zebra finch: a synthesis of field and laboratory studies* (Oxford University Press).
27. Farrar, B.G., Voudouris, K., and Clayton, N.S. (2021). Replications, comparisons, sampling and the problem of representativeness in animal cognition research. *Anim. Behav. Cogn.* 8, 273–295. 10.26451/abc.08.02.14.2021.
28. Webster, M.M., and Rutz, C. (2020). How STRANGE are your study animals? *Nature* 582, 337–340. 10.1038/d41586-020-01751-5.
29. Foote, J.R., Mennill, D.J., Ratcliffe, L.M., and Smith, S.M. (2020). Black-capped Chickadee (*Poecile atricapillus*). In *Birds of the World*, A. F. Poole, ed. (Cornell Lab of Ornithology). <https://doi.org/10.2173/bow.bkcchi.01>.
30. Aplin, L.M., Farine, D.R., Mann, R.P., Sheldon, B.C., and B, P.R.S. (2014). Individual-level personality influences social foraging and collective behaviour in wild birds. *Proc. R. Soc. B* 281, 20131016.
31. Healy, S.D., and Krebs, J.R. (1996). Food storing and the hippocampus in Paridae. *Brain. Behav. Evol.* 47, 195–199.
32. Guillette, L.M., Hahn, A.H., Hoeschele, M., Przyslupski, A.M., and Sturdy, C.B. (2015). Individual differences in learning speed, performance accuracy and exploratory behaviour in black-capped chickadees. *Anim. Cogn.* 18, 165–178. 10.1007/s10071-014-0787-3.
33. Cole, E.F., Morand-Ferron, J., Hinks, A.E., and Quinn, J.L. (2012). Cognitive ability influences reproductive life history variation in the wild. *Curr. Biol.* 22, 1808–1812. 10.1016/j.cub.2012.07.051.
34. Kozlovsky, D.Y., Weissgerber, E.A., and Pravosudov, V. V. (2017). What makes specialized food-caching mountain chickadees successful city slickers? *Proc. R. Soc. B Biol. Sci.* 284, 25–27. 10.1098/rspb.2016.2613.
35. Damsky, J., and Gall, M.D. (2016). Anthropogenic noise reduces approach of Black-capped Chickadee (*Poecile atricapillus*) and Tufted Titmouse (*Baeolophus bicolor*) to Tufted Titmouse mobbing calls. *Condor* 119, 26–33. 10.1650/CONDOR-16-146.1.
36. Billings, A.C. (2018). The low-frequency acoustic structure of mobbing calls differs across habitat types in three passerine families. *Anim. Behav.* 138, 39–49. 10.1016/j.anbehav.2018.02.001.
37. Lazerte, S.E., Slabbekoorn, H., and Otter, K.A. (2016). Learning to cope: Vocal adjustment to urban noise is correlated with prior experience in black-capped chickadees. *Proc. R. Soc. B Biol. Sci.* 283, 20161058. 10.1098/rspb.2016.1058.
38. MacLean, E.L., Hare, B., Nunn, C.L., Addessi, E., Amici, F., Anderson, R.C., Aureli, F., Baker, J.M., Bania, A.E., Barnard, A.M., et al. (2014). The evolution of self-control. *Proc. Natl. Acad. Sci.* 111, E2140–E2148. 10.1073/pnas.1323533111.
39. Neill, W.T., Valdes, L.A., and Terry, K.M. (1995). Selective attention and the inhibitory control of cognition. *Interf. Inhib. Cogn.*, 207–261. 10.1016/B978-012208930-5/50008-8.
40. Diamond, A. (1990). Developmental Time Course in Human Infants and Infant Monkeys, and the Neural Bases of, Inhibitory Control in Reaching. *Ann. N. Y. Acad. Sci.* 608, 637–676. 10.1111/j.1749-6632.1990.tb48913.x.
41. Shadmehr, R., and Holcomb, H.H. (1999). Inhibitory control of competing motor memories. *Exp. Brain Res.* 126, 235–251. 10.1007/s002210050733.
42. Amici, F., Aureli, F., and Call, J. (2008). Fission-fusion dynamics, behavioral flexibility, and inhibitory control in primates. *Curr. Biol.* 18, 1415–1419. 10.1016/j.cub.2008.08.020.


43. Lee, V.E., and Thornton, A. (2021). Animal cognition in an urbanised world. *Front. Ecol. Evol.* 9, 1–20. 10.3389/fevo.2021.633947.
44. Griffin, A.S., Tebbich, S., and Bugnyar, T. (2017). Animal cognition in a human-dominated world. *Anim. Cogn.* 20, 1–6. 10.1007/s10071-016-1051-9.
45. Shaw, R.C., Boogert, N.J., Clayton, N.S., and Burns, K.C. (2015). Wild psychometrics: Evidence for “general” cognitive performance in wild New Zealand robins, *Petroica longipes*. *Anim. Behav.* 109, 101–111. 10.1016/j.anbehav.2015.08.001.
46. Tangney, J.P., Baumeister, R.F., and Boone, A.L. (2004). High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *J. Pers.* 72, 271–324. 10.1111/j.0022-3506.2004.00263.x.
47. Moffitt, T.E., Arseneault, L., Belsky, D., Dickson, N., Hancox, R.J., Harrington, H.L., Houts, R., Poulton, R., Roberts, B.W., Ross, S., et al. (2011). A gradient of childhood self-control predicts health, wealth, and public safety. *Proc. Natl. Acad. Sci. U. S. A.* 108, 2693–2698. 10.1073/pnas.1010076108.
48. Boogert, N.J., Anderson, R.C., Peters, S., Searcy, W.A., and Nowicki, S. (2011). Song repertoire size in male song sparrows correlates with detour reaching, but not with other cognitive measures. *Anim. Behav.* 81, 1209–1216. 10.1016/j.anbehav.2011.03.004.
49. Campbell, K.A., Proppe, D.S., Congdon, J. V, Scully, E.N., Miscler, S.K., and Sturdy, C.B. (2019). The effects of anthropogenic noise on feeding behaviour in black-capped chickadees (*Poecile atricapillus*). *Behav. Processes* 158, 53–58. 10.1016/j.beproc.2018.10.015.
50. Grace, M.K., and Anderson, R.C. (2015). “D” notes of Carolina chickadee calls in response to traffic noise. *Behav. Ecol. Sociobiol.* 69, 253–263. 10.1007/s00265-014-1838-0.
51. Grade, A.M., and Sieving, K.E. (2016). When the birds go unheard: Highway noise disrupts information transfer between bird species. *Biol. Lett.* 12, 2016–2019. 10.1098/rsbl.2016.0113.
52. McClure, C.J.W., Ware, H.E., Carlisle, J., Kaltenecker, G., and Barber, J.R. (2013). An experimental investigation into the effects of traffic noise on distributions of birds: avoiding the phantom road. *Proc. R. Soc. B Biol. Sci.* 280, 20132290. 10.1098/rspb.2013.2290.
53. Sonnenberg, B.R., Branch, C.L., Pitera, A.M., Bridge, E., and Pravosudov, V. V (2019). Natural selection and spatial cognition in wild food- caching mountain chickadees. *Curr. Biol.*, 1–7. 10.1016/j.cub.2019.01.006.
54. Feeney, M.C., Roberts, W.A., and Sherry, D.F. (2009). Memory for what, where, and when in the black-capped chickadee (*Poecile atricapillus*). *Anim. Cogn.* 12, 767–777. 10.1007/s10071-009-0236-x.
55. Aplin, L.M., Farine, D.R., Cockburn, A., Thornton, A., and Sheldon, B.C. (2014). Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature*, 1–4. 10.1038/nature13998.
56. Sol, D., Lapedra, O., and González-Lagos, C. (2013). Behavioural adjustments for a life in the city. *Anim. Behav.* 85, 1101–1112. <https://doi.org/10.1016/j.anbehav.2013.01.023>.
57. Maklakov, A.A., Immler, S., Gonzalez-Voyer, A., Rönn, J., and Kolm, N. (2011). Brains and the city: big-brained passerine birds succeed in urban environments. *Biol. Lett.* 7, 730–732. 10.1098/rsbl.2011.0341.
58. Clucas, B., and Marzluff, J.M. (2013). Coupled relationships between humans and other organisms in urban areas. In *Urban ecology: patterns, processes, and applications*, J. Niemelä, ed. (Oxford University Press), pp. 135–147. 10.1093/acprof:oso/9780199563562.003.0017.
59. Bonier, F., Martin, P.R., and Wingfield, J.C. (2007). Urban birds have broader environmental tolerance. *Biol. Lett.* 3, 670–673. 10.1098/rsbl.2007.0349.
60. Lowry, H., Lill, A., and Wong, B.B.M. (2013). Behavioural responses of wildlife to urban environments. *Biol. Rev.* 88, 537–549. 10.1111/brv.12012.
61. Kleist, N.J., Guralnick, R.P., Cruz, A., Lowry, C.A., and Francis, C.D. (2018). Chronic

- anthropogenic noise disrupts glucocorticoid signaling and has multiple effects on fitness in an avian community. *Proc. Natl. Acad. Sci.* *115*, E648–E657. 10.1073/pnas.1709200115.
62. Mills, S.C., Beldade, R., Henry, L., Laverty, D., Nedelec, S.L., Simpson, S.D., and Radford, A.N. (2020). Hormonal and behavioural effects of motorboat noise on wild coral reef fish. *Environ. Pollut.* *262*, 114250. <https://doi.org/10.1016/j.envpol.2020.114250>.
 63. Wright, A.J., Soto, N.A., Baldwin, A.L., Bateson, M., Beale, C.M., Clark, C., Deak, T., Edwards, E.F., Fernández, A., and Godinho, A. (2007). Anthropogenic noise as a stressor in animals: a multidisciplinary perspective. *Int. J. Comp. Psychol.* *20*.
 64. Zollinger, S.A., Dorado-Correa, A., Goymann, W., Forstmeier, W., Knief, U., Bastidas-Urrutia, A.M., and Brumm, H. (2019). Traffic noise exposure depresses plasma corticosterone and delays offspring growth in breeding zebra finches. *Conserv. Physiol.* *7*, 1–15. 10.1093/conphys/coz056.
 65. Morand-Ferron, J., Cole, E.F., and Quinn, J.L. (2016). Studying the evolutionary ecology of cognition in the wild: A review of practical and conceptual challenges. *Biol. Rev.* *91*, 367–389. 10.1111/brv.12174.
 66. Ashton, B.J., Ridley, A.R., Edwards, E.K., and Thornton, A. (2018). Cognitive performance is linked to group size and affects fitness in Australian magpies. *Nature* *554*, 364–367. 10.1038/nature25503.
 67. Olson, J.R., Cooper, S.J., Swanson, D.L., Braun, M.J., and Williams, J.B. (2010). The relationship of metabolic performance and distribution in black-capped and carolina chickadees. *Physiol. Biochem. Zool.* *83*, 263–275. 10.1086/648395.
 68. Searcy, W.A., and Nowicki, S. (2019). Birdsong learning, avian cognition and the evolution of language. *Anim. Behav.* *151*, 217–227. 10.1016/j.anbehav.2019.01.015.
 69. Moseley, D.L., Derryberry, G.E., Phillips, J.N., Danner, J.E., Danner, R.M., Luther, D.A., and Derryberry, E.P. (2018). Acoustic adaptation to city noise through vocal learning by a songbird. *Proc. R. Soc. B Biol. Sci.* *285*. 10.1098/rspb.2018.1356.
 70. Horn, A.G., Aikens, M., Jamieson, E., Kingdon, K., and Leonard, M.L. (2020). Effect of noise on development of call discrimination by nestling tree swallows, *Tachycineta bicolor*. *Anim. Behav.* *164*, 143–148. <https://doi.org/10.1016/j.anbehav.2020.04.008>.
 71. Montenegro, C., Service, W.D., Scully, E.N., Mischler, S.K., Sahu, P.K., Benowicz, T.J., Fox, K.V.R., and Sturdy, C.B. (2021). The impact of anthropogenic noise on individual identification via female song in Black-capped chickadees (*Parus atricapillus*). *Sci. Rep.* *11*, 17530. 10.1038/s41598-021-96504-3.
 72. Van Horik, J.O., Langley, E.J.G., Whiteside, M.A., Laker, P.R., Beardsworth, C.E., and Madden, J.R. (2018). Do detour tasks provide accurate assays of inhibitory control? *Proc. R. Soc. B Biol. Sci.* *285*. 10.1098/rspb.2018.0150.
 73. Ashton, B.J., Thornton, A., Cauchoix, M., and Ridley, A.R. (2022). Long-Term repeatability of cognitive performance. *R. Soc. Open Sci.* *9*. 10.1098/rsos.220069.
 74. Davidson, G.L., Reichert, M.S., Coomes, J.R., Kulahci, I.G., de la Hera, I., and Quinn, J.L. (2022). Inhibitory control performance is repeatable over time and across contexts in a wild bird population. *Anim. Behav.* *187*, 305–318. 10.1016/j.anbehav.2022.02.007.
 75. Soha, J.A., Peters, S., Anderson, R.C., Searcy, W.A., and Nowicki, S. (2019). Performance on tests of cognitive ability is not repeatable across years in a songbird. *Anim. Behav.* *158*, 281–288. 10.1016/j.anbehav.2019.09.020.
 76. Pritchard, D.J., Hurly, T.A., Tello-Ramos, M.C., and Healy, S.D. (2016). Why study cognition in the wild (and how to test it)? *J. Exp. Anal. Behav.* *105*, 41–55. 10.1002/jeab.195.
 77. Heinen, V.K., Pitera, A.M., Sonnenberg, B.R., Benedict, L.M., Bridge, E.S., Farine, D.R., and Pravosudov, V. V. (2021). Food discovery is associated with different reliance on social learning and lower cognitive flexibility across environments in a food-caching bird. *Proc. R. Soc. B Biol. Sci.* *288*. 10.1098/rspb.2020.2843.

- 645 78. Aplin, L.M., Sheldon, B.C., and Morand-Ferron, J. (2013). Milk bottles revisited: Social learning
646 and individual variation in the blue tit, *Cyanistes caeruleus*. *Anim. Behav.* 85, 1225–1232.
647 10.1016/j.anbehav.2013.03.009.
- 648 79. Van Donselaar, J.L., Atma, J.L., Kruijf, Z.A., LaCroix, H.N., and Proppe, D.S. (2018). Urbanization
649 alters fear behavior in black-capped chickadees. *Urban Ecosyst.* 21, 1043–1051.
650 10.1007/s11252-018-0783-5.
- 651 80. Oregon Traffic Monitoring System (2022). Traffic Count Database System. accessed 27
652 December 2022. <https://ordot.public.ms2soft.com/tcds/>
- 653 81. Liu, Y., Zollinger, S.A., and Brumm, H. (2020). Chronic exposure to urban noise during the
654 vocal learning period does not lead to increased song frequencies in zebra finches. *Behav.*
655 *Ecol. Sociobiol.* 75, 3. 10.1007/s00265-020-02942-w.
656
657



659 Figure 1. Traffic noise negatively impacts inhibitory control in black-capped chickadees. A) Birds
660 hearing traffic noise playback were more than twice as likely to fail trials on a detour reaching task,
661 although B) their participation rate did not significantly differ from control birds. Data are
662 represented as mean $\pm$ SEM, with individual data points depicted for each bird.



667 Figure 2. Wild-caught chickadee detour reaching performance compared with that of lab-reared
668 zebra finches. Despite control birds of each species having similar failure rates, chickadees exposed
669 to traffic noise playback had higher failure rates than those observed in zebra finches. Data are
670 represented as mean $\pm$ SEM percent of trials. Zebra finch data adapted from Osbrink et al. ²⁴.