

Ryugu's nucleosynthetic heritage from the outskirts of the Solar System

Research article to be submitted to *Science Advances*

Abstract: 145 words

Main text: 2332 words

Tables: 1

Figures: 4

Materials and Methods: 732 words

References: 65

References in Supplementary Materials: 21

Supplementary Tables: 2

Supplementary Figures: 4

Timo Hopp^{1*†}, Nicolas Dauphas¹, Yoshinari Abe², Jérôme Aléon³, Conel M. O'D. Alexander⁴, Sachiko Amari^{5,6}, Yuri Amelin⁷, Ken-ichi Bajo⁸, Martin Bizzarro⁹, Audrey Bouvier¹⁰, Richard W. Carlson⁴, Marc Chaussidon¹¹, Byeon-Gak Choi¹², Andrew M. Davis¹, Tommaso Di Rocco¹³, Wataru Fujiya¹⁴, Ryota Fukai¹⁵, Ikshu Gautam¹⁶, Makiko K. Haba¹⁶, Yuki Hibiya¹⁷, Hiroshi Hidaka¹⁸, Hisashi Homma¹⁹, Peter Hoppe²⁰, Gary R. Huss²¹, Kiyohiro Ichida²², Tsuyoshi Iizuka²³, Trevor R. Ireland²⁴, Akira Ishikawa¹⁶, Motoo Ito²⁵, Shoichi Itoh²⁶, Noriyuki Kawasaki⁸, Noriko T. Kita²⁷, Kouki Kitajima²⁷, Thorsten Kleine²⁸, Shintaro Komatani²², Alexander N. Krot²¹, Ming-Chang Liu^{29,30}, Yuki Masuda¹⁶, Kevin D. McKeegan²⁹, Mayu Morita²², Kazuko Motomura³¹, Frédéric Moynier¹¹, Izumi Nakai³¹, Kazuhide Nagashima²¹, David Nesvorný³², Ann Nguyen³³, Larry Nittler^{4,34}, Morihiko Onose²², Andreas Pack¹³, Changkun Park³⁵, Laurette Piani³⁶, Liping Qin³⁷, Sara S. Russell³⁸, Naoya Sakamoto³⁹, Maria Schönbächler⁴⁰, Lauren Tafla²⁹, Haolan Tang²⁹, Kentaro Terada⁴¹, Yasuko Terada⁴², Tomohiro Usui¹⁵, Sohei Wada⁸, Meenakshi Wadhwa³⁴, Richard J. Walker⁴³, Katsuyuki Yamashita⁴⁴, Qing-Zhu Yin⁴⁵, Tetsuya Yokoyama¹⁶, Shigekazu Yoneda⁴⁶, Edward D. Young²⁹, Hiroharu Yui⁴⁷, Ai-Cheng Zhang⁴⁸, Tomoki Nakamura⁴⁹, Hiroshi Naraoka⁵⁰, Takaaki Noguchi²⁵, Ryuji Okazaki⁵⁰, Kanako Sakamoto¹⁵, Hikaru Yabuta⁵¹, Masanao Abe¹⁵, Akiko Miyazaki¹⁵, Aiko Nakato¹⁵, Masahiro Nishimura¹⁵, Tatsuaki Okada¹⁵, Toru Yada¹⁵, Kasumi Yogata¹⁵, Satoru Nakazawa¹⁵, Takanao Saiki¹⁵, Satoshi Tanaka¹⁵, Fuyuto Terui⁵², Yuichi Tsuda¹⁵, Sei-ichiro Watanabe¹⁸, Makoto Yoshikawa¹⁵, Shogo Tachibana⁵³, Hisayoshi Yurimoto⁸

Affiliations:

¹Origins Laboratory, Department of the Geophysical Sciences and Enrico Fermi Institute, The University of Chicago, 5734 South Ellis Avenue, Chicago 60637, USA.

²Graduate School of Engineering Materials Science and Engineering, Tokyo Denki University; Tokyo 120-8551, Japan.

³Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Sorbonne Université, Museum National d'Histoire Naturelle, CNRS UMR 7590, IRD; 75005 Paris, France.

⁴Earth and Planets Laboratory, Carnegie Institution for Science; Washington, DC, 20015, USA.

⁵McDonnell Center for the Space Sciences and Physics Department, Washington University; St. Louis, MO 63130, USA.

⁶Geochemical Research Center, The University of Tokyo, Tokyo, 113-0033, Japan

⁷Guangzhou Institute of Geochemistry, Chinese Academy of Sciences; Guangzhou, GD 510640, China.

⁸Natural History Sciences, IIL, Hokkaido University; Sapporo 001-0021, Japan.

⁹Centre for Star and Planet Formation, GLOBE Institute, University of Copenhagen; Copenhagen, K 1350, Denmark.

¹⁰Bayerisches Geoinstitut, Universität Bayreuth; Bayreuth 95447, Germany.

¹¹Université Paris Cité, Institut de physique du globe de Paris, CNRS; 75005 Paris, France

¹²Department of Department of Earth Science Education, Seoul National University; Seoul 08826, Republic of Korea.

¹³Faculty of Geosciences and Geography, University of Göttingen; Göttingen, D-37077, Germany.

¹⁴Faculty of Science, Ibaraki University; Mito 310-8512, Japan.

¹⁵ISAS/JSEC, JAXA; Sagamihara 252-5210, Japan.

¹⁶Department of Earth and Planetary Sciences, Tokyo Institute of Technology; Tokyo 152-8551, Japan.

¹⁷General Systems Studies, The University of Tokyo; Tokyo 153-0041, Japan.

¹⁸Earth and Planetary Sciences, Nagoya University; Nagoya 464-8601, Japan.

¹⁹Osaka Application Laboratory, SBUWDX, Rigaku Corporation; Osaka 569-1146, Japan.

²⁰Max Planck Institute for Chemistry; Mainz 55128, Germany.

²¹Hawai‘i Institute of Geophysics and Planetology, University of Hawai‘i at Mānoa; Honolulu, HI 96822, USA.

²²Analytical Technology, Horiba Techno Service Co., Ltd.; Kyoto 601-8125, Japan.

²³Earth and Planetary Science, The University of Tokyo; Tokyo 113-0033, Japan.

²⁴School of Earth and Environmental Sciences, The University of Queensland; St Lucia QLD 4072, Australia.

²⁵Kochi Institute for Core Sample Research, JAMSTEC; Kochi 783-8502, Japan.

²⁶Earth and Planetary Sciences, Kyoto University; Kyoto 606-8502, Japan.

²⁷Geoscience, University of Wisconsin- Madison; Madison, WI 53706, USA.

²⁸Max Planck Institute for Solar System Research; 37077 Göttingen, Germany.

²⁹Earth, Planetary, and Space Sciences, UCLA; Los Angeles, CA 90095, USA.

³⁰Lawrence Livermore National Laboratory, Livermore, CA 94550, USA.

- ³¹Thermal Analysis, Rigaku Corporation; Tokyo 196-8666, Japan.
- ³²Department of Space Studies, Southwest Research Institute, Boulder, CO 80302, USA.
- ³³Astromaterials Research and Exploration Science, NASA Johnson Space Center; Houston, TX 77058, USA.
- ³⁴School of Earth and Space Exploration, Arizona State University; Tempe, AZ 85281, USA.
- ³⁵Earth-System Sciences, Korea Polar Research Institute; Incheon 21990, Korea.
- ³⁶Centre de Recherches Pétrographiques et Géochimiques, CNRS - Université de Lorraine; 54500 Nancy, France.
- ³⁷University of Science and Technology of China, School of Earth and Space Sciences; Anhui 230026, China.
- ³⁸Department of Earth Sciences, Natural History Museum; London, SW7 5BD, UK.
- ³⁹IIL, Hokkaido University; Sapporo 001-0021, Japan.
- ⁴⁰Institute for Geochemistry and Petrology, Department of Earth Sciences, ETH Zurich, Zurich, Switzerland.
- ⁴¹Earth and Space Science, Osaka University; Osaka 560-0043, Japan.
- ⁴²Spectroscopy and Imaging, Japan Synchrotron Radiation Research Institute; Hyogo 679-5198 Japan.
- ⁴³Geology, University of Maryland, College Park, MD 20742, USA.
- ⁴⁴Graduate School of Natural Science and Technology, Okayama University; Okayama 700-8530, Japan.
- ⁴⁵Earth and Planetary Sciences, University of California; Davis, CA 95616, USA.
- ⁴⁶Science and Engineering, National Museum of Nature and Science; Tsukuba 305-0005, Japan.
- ⁴⁷Chemistry, Tokyo University of Science; Tokyo 162-8601, Japan.
- ⁴⁸School of Earth Sciences and Engineering, Nanjing University; Nanjing 210023, China.
- ⁴⁹Department of Earth Science, Tohoku University; Sendai, 980-8578, Japan.
- ⁵⁰Department of Earth and Planetary Sciences, Kyushu University; Fukuoka 819-0395, Japan.
- ⁵¹Earth and Planetary Systems Science Program, Hiroshima University; Higashi-Hiroshima, 739-8526, Japan.
- ⁵²Kanagawa Institute of Technology; Atsugi 243-0292, Japan.
- ⁵³UTokyo Organization for Planetary and Space Science, University of Tokyo; Tokyo 113-0033, Japan.

* Corresponding author. Email: hopp@mps.mpg.de

† Now at the Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany.

Teaser: (113/125 characters) **Asteroid Ryugu and Ivuna-type carbonaceous meteorites may have originated from the outskirts of the Solar System.**

Abstract: (145/150 words)

Little is known about the origin of the spectral diversity of asteroids and what it says about conditions in the protoplanetary disk. Here we show that samples returned from Cb-type asteroid Ryugu have Fe isotopic anomalies **indistinguishable from Ivuna-type (CI)** chondrites, which are distinct from all other carbonaceous chondrites. Iron isotopes, therefore, demonstrate that Ryugu and CI chondrites formed in a reservoir that was **different** from the source regions of other carbonaceous asteroids. **Growth and migration of the giant planets destabilized nearby planetesimals and ejected some inwards to be implanted into the Main Belt. In this framework,** most carbonaceous chondrites may have originated from regions around the birthplaces of Jupiter and Saturn, while the distinct isotopic composition of CI chondrites and Ryugu may reflect their formation further away in the disk, owing their presence in the inner Solar System to excitation by Uranus and Neptune.

Main Text:

Introduction

Main Belt asteroids show great compositional diversity (1), ranging from metallic objects that are remnants of differentiated planetesimals (2) to carbon-rich objects with comet-like dust-ejection activity (3). The original formation locations of these diverse objects are unknown. Meteorites are remnants of planetesimals and protoplanets that formed at various heliocentric distances within the first few million years after the birth of the Solar System. They, therefore, provide invaluable insights into the early evolution of the Solar System and the building blocks of the terrestrial planets. Most meteorites are fragments of Main Belt asteroids, but direct asteroid-meteorite links are scarce (1). Establishing such links is important as it provides clues on the relationship and formation locations of meteorite parent bodies, asteroids, and other small bodies in the Solar System. Cb-type asteroid (162173) Ryugu is a near-Earth object (NEO) that most likely originated from the inner Main Belt (4). Chemical and mineralogical analyses of Ryugu samples returned to Earth by JAXA's Hayabusa2 mission (5) show that they share chemical and mineralogical characteristics with Ivuna-type carbonaceous (CI) chondrites (6). The latter is the only group of meteorites containing most non-volatile elements in proportions nearly equal to those measured in

the solar photosphere (7). The physical and chemical similarities of Ryugu and CI chondrites are, however, not diagnostic of a shared heritage because the low-temperature conditions required to explain their solar-like chemical compositions could have been widespread in the outer Solar System disk.

To better constrain Ryugu's nucleosynthetic heritage, we measured the Fe isotopic compositions of four Ryugu samples collected during the first and second touchdown (5), 11 different carbonaceous chondrites from five different groups (CI, CM, CV, CO, CR), and two ungrouped chondrites. Materials formed in the Solar System display variations in the isotopic compositions of some elements that stem from the heterogeneous distribution and processing of highly anomalous presolar materials in the protosolar nebula (8–11). Such anomalies cannot easily be modified by physicochemical processes on planetary bodies and, therefore, provide lasting isotopic fingerprints of the regions where planetary bodies formed (12–14). For several elements, meteorites display a dichotomy in their isotopic anomalies between non-carbonaceous (NC) and carbonaceous (CC) meteorite groups (13). The origin of this dichotomy could have involved the physical separation between inner and outer Solar System reservoirs by Jupiter (13, 15, 16) and planetesimal formation at distinct locations in an evolving protoplanetary disk (17–21). Although isotopic analyses of Ti and Cr show that Ryugu's building blocks formed in the CC reservoir and support a possible kinship between Ryugu and CI chondrites (6), an unambiguous genetic link to a specific carbonaceous chondrite group could not be established because the Ti and Cr isotopic anomalies of Ryugu overlap with several carbonaceous chondrites and achondrites (Fig. 1A). Carbonaceous chondrites also display mass-independent variations in O isotopes that correlate with Ti and Cr isotopic anomalies (13). Ryugu and CI chondrites have similar O isotopic anomalies and represent an endmember composition for the CC cluster in O-Cr and O-Ti spaces (Fig. 1B,C) (6). Those results support the view that CI chondrites and Ryugu formed within the CC reservoir, but because of correlations between O, Cr, and Ti anomalies (Fig. 1), there is significant redundancy in evidence presented thus far and further work is needed to better understand the isotopic architecture of the outer Solar System.

The Fe isotopic composition of CI chondrites (22) is clearly distinct from all other carbonaceous meteorites (22–24), where all CC meteorites except CI chondrites show significant excesses in ⁵⁴Fe. **Iron isotopes, therefore, provide a diagnostic tool to evaluate if Ryugu has the same distinct nucleosynthetic heritage as CI chondrites.**

Results

Ryugu samples and all carbonaceous chondrites display limited mass-dependent isotopic variation relative to the terrestrial standard IRMM-524a (Table S1), meaning that all isotopic anomalies reported here are real and not artifacts from the internal normalization scheme used to correct natural and instrumental isotopic fractionation (14). Mass-independent Fe isotopic analyses confirm that most carbonaceous chondrites display variable excesses of $\sim +15$ to $\sim +40$ in $\mu^{54}\text{Fe}$ but no resolvable variations in $\mu^{58}\text{Fe}$ (see Table 1 for a definition of the μ -notation) (Table 1; Fig. 2). By contrast, all three CI chondrites analyzed in this study are clearly distinct from other carbonaceous chondrites, defining an average $\mu^{54}\text{Fe} = +3 \pm 2$, consistent with previous measurements (Fig. 2; Fig. S1) (22). The four Ryugu samples (A0106 and A0106-A0107 collected during the first touchdown from the surface; C0107 and C0108 collected during the second touchdown and possibly sampling material from the subsurface) define an average $\mu^{54}\text{Fe}$ value of $+1 \pm 4$, which is indistinguishable from the composition of CI chondrites but distinct from all other carbonaceous chondrites (Fig. 2). The $\mu^{54}\text{Fe}$ difference of Ryugu and CI chondrites compared to all other carbonaceous chondrites cannot be due to prolonged exposure to cosmic rays in space because (i) CI chondrites have low cosmic ray exposure ages (25) and (ii) cosmogenic effects would induce a positive shift in $\mu^{54}\text{Fe}$ that would correlate with a negative shift in $\mu^{58}\text{Fe}$ (24, 26), which is not observed (Table 1; Fig. S2). Thus, the distinct $\mu^{54}\text{Fe}$ values of CI chondrites and Ryugu represent the nucleosynthetic heritage of their formation reservoir in the protosolar nebula. Examination of $\mu^{54}\text{Fe}$ - $\mu^{50}\text{Ti}$ (Fig. 3) and $\mu^{54}\text{Fe}$ - $\mu^{54}\text{Cr}$ (Fig. S3) isotopic anomalies show that Ryugu and CI chondrites form a compositional cluster that is distinct from the NC and CC fields defined by other meteorites. Thus, whereas the isotopic anomalies of Ti, Cr, and O in CI chondrites (10, 11, 27–31) and Ryugu (6) tie them to the CC reservoir, the Fe isotopic data reveal that CI chondrites and Ryugu formed in a reservoir that is rarely sampled by meteorites.

Discussion

Recent models tie the distinct isotopic characteristics of meteorites to planetesimal formation at different locations and/or at different time in an isotopically heterogeneous protoplanetary disk (9, 15, 17, 18, 20, 21). The main driver of the isotopic heterogeneity could be a change in the composition of infalling material during collapse of the parental molecular cloud core of the Solar System (14, 32, 33), or alternatively, unmixing of presolar carriers by disk processes (9, 10, 19).

Irrespective of the origin of the heterogeneity, it requires the existence of large-scale isotopic heterogeneities throughout the disk. Interestingly, the Fe isotopic compositions of CI chondrites and Ryugu are similar to Earth's mantle and NC meteorites, **while** isotopic anomalies **in other elements** are similar to CC meteorites and significantly different compared to inner Solar System material (Fig. 1; Fig. 3). This suggests that CI chondrites and Ryugu derive from an isotopic reservoir that has a distinctive nucleosynthetic heritage from NC and CC meteorites. **While** the Fe isotopic characteristics **of** CI chondrites could be explained by an inner Solar System origin, the isotopic anomalies of Ti, Cr, and Mo clearly tie CI chondrites to **the** CC reservoir and the outer Solar System (10–12, 16, 27). More importantly, the unfractionated chemical compositions and volatile element-rich nature of CI chondrites and Ryugu support the view that they formed **beyond** the snow line, mostly from material that experienced minimal thermal processing. **The heavy hydrogen and nitrogen isotopic compositions of Ryugu are also consistent with an outer Solar System origin (34).** The findings that CI chondrites and Ryugu share the same nucleosynthetic heritage (Fig. 3) and have close mineralogical, chemical, and isotopic characteristics (6, 34), therefore, suggest that these objects formed contemporaneously and were co-located in the same outer Solar System reservoir. It is even possible, although not required by the data, that CI chondrites and the NEO Ryugu originally derived from the same precursor object, which was fragmentated by collision during its residence in the inner Main Belt (4, 35).

The distinct isotopic heritage of CI chondrites and Ryugu is unlikely to reflect a temporal change in the isotopic composition of the CC reservoir because (i) CI chondrites and Ryugu are distinct from other carbonaceous chondrites and intermediate compositions are missing (Fig. 2), and (ii) CI chondrites have similar inferred accretion ages **as** other carbonaceous chondrites **(~2.5 to 4 Myr after condensation of refractory inclusions)** (36). Thus, the **distinctive** isotopic heritage of CI chondrites and Ryugu is most likely caused by spatial separation of their source region from the CC reservoir, **or as discussed below**, bias in the implantation of planetesimals from distinct heliocentric distances into the Main Belt.

Cb-type asteroids like Ryugu represent ~10–20% of all C-type asteroids (37, 38), suggesting that a substantial portion of Main Belt asteroids formed in the same outer Solar System reservoir where CI chondrites and Ryugu formed. Planetesimals from a large range of heliocentric distances could have been implanted into the inner Solar System during the growth and migration of the giant planets within the protoplanetary disk, before dissipation of nebular gas (39). Planetesimals from

the outer Solar System could have also been implanted during subsequent instabilities in the orbits of Uranus and Neptune after dissipation of nebular gas (40–43), but the efficiency of this process is low and early implantation of C-type asteroids in the Main Belt while nebular gas was still present is more likely. The planetesimals scattered by the growth and migration of the giant planets were originally on highly eccentric orbits but they experienced a strong headwind from nebular gas that circularized their orbits, leading to their trapping in the Main Belt. Simulations of the growth and migration of the giant planets show that most outer Solar System planetesimals implanted in the Main Belt originated from the formation region of the gas giant planets Jupiter and Saturn (*e.g.*, ~4-12 AU), but some could have come from further away in the formation region of ice giant planets Uranus and Neptune (*e.g.*, 13-25 AU) (39). The dynamical process of orbital excitation and circularization introduces strong biases in the original orbital radius of the planetesimals that are eventually implanted in the Main Belt (Fig. S4). In that context, most CC meteorites could have come from the birth region of Jupiter and Saturn, while the **distinctive** Fe isotopic heritage (Fig. 3) and primitive chemical characteristics of CI chondrites and Ryugu (6) could be explained if they were implanted into the Main Belt from a reservoir that was located further outside, possibly in the vicinity of the birthplaces of Uranus and Neptune (Fig. 4; Fig. S4). If correct, CI chondrites and Ryugu would possibly share a common heritage with Oort cloud comets (44, 45).

A common source region for the parent bodies of CI chondrites/Ryugu and Oort cloud comets would need to be reconciled with their present-day distinct chemical and physical characteristics (6, 34, 46). **Deuterium/hydrogen (D/H) ratios of water in carbonaceous chondrites (47) and Ryugu (34) are lower than those of Oort cloud comets and overlap partially with Jupiter-family comets (48). Simulations of ice transport in the nebula predict a spatially and temporally complex evolution of water D/H ratio in the nebula (49). Furthermore, the water D/H ratio of active comets and asteroids might have been modified by water sublimation (50, 51) and water-rock reactions (52) during their lifetimes. Therefore, the present-day water D/H ratio of ice in comets and rock-bound water in carbonaceous chondrites provide little insights into formation locations.** CI chondrites and Ryugu show evidence for extensive aqueous alteration as late as ~5 Myr after the birth of the Solar System (6, 53, 54). Water responsible for this aqueous alteration would have presumably been accreted as ice, with melting caused by decay of ²⁶Al. Such melting could have been hampered in ice-rich comets if much of radioactive heat

from ^{26}Al was consumed by ice sublimation rather than melting (55). While most dust grains captured in the coma of comet 81P/Wild2 were anhydrous (46), Berger et al. (56) found evidence for low-temperature aqueous activity in 81P/Wild2 under conditions akin to those inferred for CI chondrites. The Deep Impact mission also found signatures of carbonates, phyllosilicates, sulfides, water gas, and ice in the ejecta of comet 9P/Tempel, which is consistent with extensive aqueous alteration (57). While these observations support the presence of aqueous activity on extant comets, a comparison between icy planetesimals that were scattered inwards and outwards by the growth and migration of the giant planets is difficult because they would have experienced very different thermal histories. Indeed, the planetesimals scattered inwards would have been put on eccentric orbits with low perihelion, well **inside** the snow line, where ice could have been sublimated and the more volatile compounds could have been lost. For a ~ 100 km planetesimal, the timescale for damping eccentricity through gas interaction is on the order of several tens of thousands of years, which is in the order of the expected lifetime of short-period comets (58). Thus, the rock/ice ratio of ice-rich planetesimals formed around Uranus and Neptune may have increased substantially by the time these planetesimals were implanted in the Main Belt. Such processing could also have affected other characteristics of the icy planetesimals, notably their inventories of organics, mineralogical compositions, and physical properties. Further transformations would have taken place due to collisions during residence in the Main Belt (35, 59). Consequently, although the Ryugu asteroid and Oort cloud comets may have been born in the same region of the protoplanetary disk, they would have rapidly diverged in their chemical evolution after being scattered **inwards** and outwards by the ice giant planets.

In our model, Cb-type asteroids formed in a reservoir that was located at the outskirts of the planetary accretion region and were possibly implanted into the Main Belt due to excitation by Uranus and Neptune (Fig. 4). The outer extent of this **isotopic reservoir** is unknown. The main source region of Kuiper Belt objects (KBOs) was likely the trans-Neptunian disk (>20 – 25 AU) (60). These KBOs are out of reach for sample return missions but possible trans-Neptunian objects 203 Pompeja and 269 Justitia were recently found in the Main Belt (61). Measuring the Fe and Ti isotopic compositions of these objects would provide important new insights into the isotopic architecture of the early Solar System and help evaluate the extent of the CI reservoir.

Materials and Methods

Samples, preparation, and chemical purification. The Hayabusa2 spacecraft returned a total of ~5 g of material from Cb-type (162173) asteroid Ryugu. Surface samples were collected in Chamber A (~3 g) during the first touchdown and subsurface samples in Chamber C (~2 g) during the second touchdown (5, 62). Two subsamples from Chamber A (A0106, A0106-A0107) and two from Chamber C (C0107, C0108) were digested for isotopic analysis (Table 1). Sample masses were 14.2, 23.88, 14.20, 12.90, and 22.24 mg, respectively. For samples A0106 and C0107 soluble organic matter was separated prior to digestion by acids. Sample A0108 was analyzed by XRF before digestion. Approximately 20–25 mg of six carbonaceous chondrite powders (Table 1: Orgueil-4, Alais, Murchison-2, Allende-3, Tagish Lake-2, and Tarda) were digested and processed together with the Hayabusa2 samples. Sample digestion for these samples was conducted at Tokyo Tech. Powder aliquots were digested using mixtures of HF-HNO₃-HCl-H₂O₂ on hot plate and under ultrasonic agitation. Approximately 80% of the solutions was taken for sequential separation of several elements for isotopic analysis. We measured 20 additional carbonaceous chondrite samples and three terrestrial geostandards (Table 1) to provide some context for interpreting Ryugu's results. Some samples were digested for this study while others are elution cuts from previous studies focused on elements other than Fe. The masses digested, original masses homogenized, and details on the processing history for each sample are summarized in Table S2. The first step in the chemical processing of Fe in the four Ryugu samples and six accompanying carbonaceous chondrites was conducted at Tokyo Tech and involved: (i) separation of major elements, including Fe, from Zn, Pb, and highly siderophile elements using anion exchange chromatography (AG-1X8), (ii) Separation of Fe and U from remaining major elements using AG-1X8, and (iii) Separation of U from Fe using Eichrom UTEVA resin. A 20% aliquot of the Fe solution was then purified from remaining traces of Cr, Ni, Co, and Cu at the University of Chicago using an established protocol (24). Approximately 0.5-1 mg of Fe was loaded in 0.25 ml 10 M HCl onto 10.5 cm long PFA columns (0.62 cm inner diameter) filled with 3 ml pre-cleaned AG1-X8 (200-400 mesh) anion resin. Matrix elements were eluted in 5 ml 10 M HCl. Other possible contaminants (*e.g.*, Cu, Cr) were eluted from the resin using 30 ml 4 M HCl. Iron was eluted using 9 ml 0.4 M HCl. The samples not previously processed at Tokyo Institute of Technology were purified using the same procedure, but the elution was repeated using new resin. The overall Fe yield is >99% and the procedural blank is negligible (~70 ng) compared to the amount of Fe

purified for each sample (0.5-1 mg Fe). Interfering elements Cr (^{54}Cr on ^{54}Fe) and Ni (^{58}Ni on ^{58}Fe) were present at low enough levels ($\text{Cr/Fe} \leq 9 \times 10^{-6}$ and $\text{Ni/Fe} \leq 2 \times 10^{-5}$) to not affect the accuracy of the analyses (24).

Iron isotopic measurements. High precision Fe isotopic compositions were measured following the protocol used for analysis of Fe isotopic anomalies in iron meteorites (24). Measurements were conducted with a Thermo Scientific Neptune multicollector inductively coupled plasma mass spectrometer (MC-ICP-MS) at the **Origins Laboratory of the** University of Chicago. Measurements were made on the flat-topped peak shoulder in medium-resolution (MR) mode (63). Ion beams of $^{54}\text{Fe}^+$, $^{56}\text{Fe}^+$, $^{57}\text{Fe}^+$, and $^{58}\text{Fe}^+$ were analyzed statically on Faraday collectors. All isotopes were measured using $10^{11} \Omega$ amplifiers, except for high abundance $^{56}\text{Fe}^+$, which was measured using a $10^{10} \Omega$ amplifier. Isobaric interferences from $^{54}\text{Cr}^+$ and $^{58}\text{Ni}^+$ were determined simultaneously by monitoring $^{53}\text{Cr}^+$ and $^{60}\text{Ni}^+$ using $10^{12} \Omega$ amplifiers. The purified Fe solutions ($10 \mu\text{g/g}$ in 0.3 M HNO_3) were introduced into the MC-ICP-MS using an ESI PFA nebulizer with an uptake rate of $\sim 100 \mu\text{l/min}$ combined with a cyclonic glass spray chamber. **Iron isotopic composition was measured at a typical** $^{56}\text{Fe}^+$ ion signal intensity of 1.3 nA on a $10^{10} \Omega$ amplifier. Each measurement consisted of 50 cycles of 8.369 s each. Sample analyses were bracketed by measurements of IRMM-524a in a **standard-sample-standard scheme**. On peak zero intensities from a blank solution measured at the start of each **sequence** were subtracted from measurements. A washout time of 210 s was used between each analysis. The Fe concentrations of the samples and standards were matched to within $\leq 2 \%$, which is required for accurate and precise Fe isotopic analysis (24).

References

1. F. E. DeMeo, B. J. Burt, M. Marsset, D. Polishook, T. H. Burbine, B. Carry, R. P. Binzel, P. Vernazza, V. Reddy, M. Tang, C. A. Thomas, A. S. Rivkin, N. A. Moskovitz, S. M. Slivan, S. J. Bus, Connecting asteroids and meteorites with visible and near-infrared spectroscopy. *Icarus*. **380**, 114971 (2022).
2. L. T. Elkins-Tanton, E. Asphaug, J. F. Bell, C. J. Bierson, B. G. Bills, W. F. Bottke, S. W. Courville, S. D. Dobb, I. Jun, D. J. Lawrence, S. Marchi, T. J. McCoy, J. M. G. Merayo, R. Oran, J. G. O'Rourke, R. S. Park, P. N. Peplowski, T. H. Prettyman, C. A. Raymond, B. P. Weiss, M. A. Wieczorek, M. T. Zuber, Distinguishing the origin of asteroid (16) Psyche. *Space Sci Rev.* **218**, 17 (2022).
3. D. S. Laretta, C. W. Hergenrother, S. R. Chesley, J. M. Leonard, J. Y. Pelgrift, C. D. Adam, M. al Asad, P. G. Antreasian, R. L. Ballouz, K. J. Becker, C. A. Bennett, B. J. Bos, W. F. Bottke, M. Brozović, H. Campins, H. C. Connolly, M. G. Daly, A. B. Davis, J. de León, D. N. DellaGiustina, C. Y. Drouet d'Aubigny, J. P. Dworkin, J. P. Emery, D. Farnocchia, D. P. Glavin, D. R. Golish, C. M. Hartzell, R. A. Jacobson, E. R. Jawin, P. Jenniskens, J. N. Kidd, E. J. Lessac-Chenen, J. Y. Li, G. Libourel, J. Licandro, A. J. Liounis, C. K. Maleszewski, C. Manzoni, B. May, L. K. McCarthy, J. W. McMahon, P. Michel, J. L.

- Molaro, M. C. Moreau, D. S. Nelson, W. M. Owen, B. Rizk, H. L. Roper, B. Rozitis, E. M. Sahr, D. J. Scheeres, J. A. Seabrook, S. H. Selznick, Y. Takahashi, F. Thuillet, P. Tricarico, D. Vokrouhlický, C. W. V. Wolner, Episodes of particle ejection from the surface of the active asteroid (101955) Bennu. *Science*. **366**, eaay3544 (2019).
4. H. Campins, J. de León, A. Morbidelli, J. Licandro, J. Gayon-Markt, M. Delbo, P. Michel, The origin of asteroid 162173 (1999 JU3). *Astronomical Journal*. **146** (2013).
5. S. Tachibana, H. Sawada, R. Okazaki, Y. Takano, K. Sakamoto, Y. N. Miura, C. Okamoto, H. Yano, S. Yamanouchi, P. Michel, Y. Zhang, S. Schwartz, F. Thuillet, H. Yurimoto, T. Nakamura, T. Noguchi, H. Yabuta, H. Naraoka, A. Tsuchiyama, N. Imae, K. Kurosawa, A. M. Nakamura, K. Ogawa, S. Sugita, T. Morota, R. Honda, S. Kameda, E. Tatsumi, Y. Cho, K. Yoshioka, Y. Yokota, M. Hayakawa, M. Matsuoka, N. Sakatani, M. Yamada, T. Kouyama, H. Suzuki, C. Honda, T. Yoshimitsu, T. Kubota, H. Demura, T. Yada, M. Nishimura, K. Yogata, A. Nakato, M. Yoshitake, A. I. Suzuki, S. Furuya, K. Hatakeda, A. Miyazaki, K. Kumagai, T. Okada, M. Abe, T. Usui, T. R. Ireland, M. Fujimoto, T. Yamada, M. Arakawa, H. C. Connolly Jr, A. Fujii, S. Hasegawa, N. Hirata, C. Hirose, S. Hosoda, Y. Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, S. Kikuchi, K. Kitazato, D. S. Lauretta, G. Libourel, B. Marty, K. Matsumoto, T. Michikami, Y. Mimasu, A. Miura, O. Mori, K. Nakamura-Messenger, N. Namiki, A. N. Nguyen, L. R. Nittler, H. Noda, R. Noguchi, N. Ogawa, G. Ono, M. Ozaki, H. Senshu, T. Shimada, Y. Shimaki, K. Shirai, S. Soldini, T. Takahashi, Y. Takei, H. Takeuchi, R. Tsukizaki, K. Wada, Y. Yamamoto, K. Yoshikawa, K. Yumoto, M. E. Zolensky, S. Nakazawa, F. Terui, S. Tanaka, T. Saiki, M. Yoshikawa, S. Watanabe, Y. Tsuda, Pebbles and sand on asteroid (162173) Ryugu: In situ observation and particles returned to Earth. *Science*. **375**, 1011–1016 (2022).
6. T. Yokoyama, K. Nagashima, I. Nakai, E. D. Young, Y. Abe, J. Aléon, C. M. O’R. D. Alexander, S. Amari, Y. Amelin, K.-I. Bajo, M. Bizzarro, A. Bouvier, R. W. Carlson, M. Chaussidon, B.-G. Choi, N. Dauphas, A. M. Davis, T. di Rocco, W. Fujiya, R. Fukai, I. Gautam, M. K. Haba, Y. Hibiya, H. Hidaka, H. Homma, P. Hoppe, G. R. Huss, K. Ichida, T. Iizuka, T. R. Ireland, A. Ishikawa, M. Ito, S. Itoh, N. Kawasaki, N. T. Kita, K. Kitajima, T. Kleine, S. Komatani, A. N. Krot, M.-C. Liu, Y. Masuda, K. D. Mckeegan, M. Morita, K. Motomura, F. Moynier, A. Nguyen, L. Nittler, M. Onose, A. Pack, C. Park, L. Piani, L. Qin, S. S. Russell, N. Sakamoto, M. Schönbächler, L. Tafla, H. Tang, K. Terada, Y. Terada, T. Usui, S. Wada, M. Wadhwa, R. J. Walker, K. Yamashita, Q.-Z. Yin, S. Yoneda, H. Yui, A.-C. Zhang, H. C. Connolly, H. Sawada, H. Senshu, Y. Shimaki, K. Shirai, S. Sugita, Y. Takei, H. Takeuchi, S. Tanaka, E. Tatsumi, F. Terui, Y. Tsuda, R. Tsukizaki, K. Wada, S.-I. Watanabe, M. Yamada, T. Yamada, Y. Yamamoto, H. Yano, Y. Yokota, K. Yoshihara, M. Yoshikawa, K. Yoshikawa, S. Furuya, K. Hatakeda, T. Hayashi, Y. Hitomi, K. Kumagai, A. Miyazaki, A. Nakato, M. Nishimura, H. Soejima, A. Suzuki, T. Yada, D. Yamamoto, K. Yogata, M. Yoshitake, S. Tachibana, H. Yurimoto, The first returned samples from a C-type asteroid show kinship to the chemically most primitive meteorites. *Science*. **first release**, 1–8 (2022).
7. K. Lodders, Relative Atomic Solar System Abundances , Mass Fractions , and Atomic Masses of the Elements and Their Isotopes , Composition of the Solar Photosphere , and Compositions of the Major Chondritic Meteorite Groups. *Space Sci Rev*. **217**, 44 (2021).
8. N. Dauphas, B. Marty, L. Reisberg, Molybdenum evidence for inherited planetary scale isotope heterogeneity of the protosolar nebula. *Astrophys J*. **565**, 640–644 (2002).
9. M. Ek, A. C. Hunt, M. Lugaro, M. Schönbächler, The origin of s-process isotope heterogeneity in the solar protoplanetary disk. *Nat Astron*. **4**, 273–281 (2020).
10. A. Trinquier, T. Elliott, D. Ulfbeck, C. Coath, A. N. Krot, M. Bizzarro, Origin of nucleosynthetic isotope heterogeneity in the solar protoplanetary disk. *Science*. **324**, 374–376 (2009).
11. A. Trinquier, J.-L. Birck, C. J. Allègre, Widespread ⁵⁴Cr heterogeneity in the inner Solar System. *Astrophysical Journal*. **655**, 1179–1185 (2007).
12. C. Burkhardt, T. Kleine, F. Oberli, A. Pack, B. Bourdon, R. Wieler, Molybdenum isotope anomalies in meteorites: Constraints on solar nebula evolution and origin of the Earth. *Earth Planet Sci Lett*. **312**, 390–400 (2011).

13. P. H. Warren, Stable-isotopic anomalies and the accretionary assemblage of the Earth and Mars: A subordinate role for carbonaceous chondrites. *Earth Planet Sci Lett.* **311**, 93–100 (2011).
14. N. Dauphas, E. A. Schauble, Mass fractionation laws, mass-independent effects, and isotopic anomalies. *Annu Rev Earth Planet Sci.* **44**, 709–783 (2016).
15. T. S. Kruijer, C. Burkhardt, G. Budde, T. Kleine, Age of Jupiter inferred from the distinct genetics and formation times of meteorites. *Proc Natl Acad Sci U S A.* **114**, 6712–6716 (2017).
16. G. Budde, C. Burkhardt, G. A. Brennecka, M. Fischer-Gödde, T. S. Kruijer, T. Kleine, Molybdenum isotopic evidence for the origin of chondrules and a distinct genetic heritage of carbonaceous and non-carbonaceous meteorites. *Earth Planet Sci Lett.* **454**, 293–303 (2016).
17. T. Lichtenberg, J. Drazkowska, M. Schönbächler, G. J. Golabek, T. O. Hands, Bifurcation of planetary building blocks during Solar System formation. *Science.* **371**, 365–370 (2021).
18. A. Morbidelli, K. Baillié, K. Batygin, S. Charnoz, T. Guillot, D. C. Rubie, T. Kleine, Contemporary formation of early Solar System planetesimals at two distinct radial locations. *Nat Astron.* **6**, 72–79 (2022).
19. M. Schiller, M. Bizzarro, V. A. Fernandes, Isotopic evolution of the protoplanetary disk and the building blocks of Earth and the Moon. *Nature.* **555**, 501–510 (2018).
20. B. Liu, A. Johansen, M. Lambrechts, M. Bizzarro, T. Haugbølle, Natural separation of two primordial planetary reservoirs in an expanding solar protoplanetary disk. *Sci. Adv.* **8**, eabm3045 (2022).
21. A. Izidoro, R. Dasgupta, S. N. Raymond, R. Deienno, B. Bitsch, A. Isella, Planetesimal rings as the cause of the Solar System’s planetary architecture. *Nat Astron.* **6**, 357–366 (2022).
22. M. Schiller, M. Bizzarro, J. Siebert, Iron isotope evidence for very rapid accretion and differentiation of the proto-Earth. *Sci. Adv.* **6**, eaay7604 (2020).
23. D. L. Cook, B. S. Meyer, M. Schönbächler, Iron and nickel isotopes in IID and IVB iron meteorites : Evidence for admixture of an SN II component and implications for the initial abundance of ^{60}Fe . *Astrophys J.* **917**, 59 (2021).
24. T. Hopp, N. Dauphas, F. Spitzer, C. Burkhardt, T. Kleine, Earth’s accretion inferred from iron isotopic anomalies of supernova nuclear statistical equilibrium origin. *Earth Planet Sci Lett.* **577**, 117245 (2022).
25. O. Eugster, Cosmic-ray exposure ages of meteorites and lunar rocks and their significance. *Chem. Erde.* **63**, 3–30 (2003).
26. D. L. Cook, I. Leya, M. Schönbächler, Galactic cosmic ray effects on iron and nickel isotopes in iron meteorites. *Meteorit Planet Sci.* **55**, 2758–2771 (2020).
27. I. Leya, M. Schönbächler, U. Wiechert, U. Krähenbühl, A. N. Halliday, Titanium isotopes and the radial heterogeneity of the solar system. *Earth Planet Sci Lett.* **266**, 233–244 (2008).
28. R. N. Clayton, T. K. Mayeda, Oxygen isotope studies of carbonaceous chondrites. *Geochim Cosmochim Acta.* **63**, 2089–2104 (1999).
29. A. Shukolyukov, G. W. Lugmair, Manganese-chromium isotope systematics of carbonaceous chondrites. *Earth Planet Sci Lett.* **250**, 200–213 (2006).
30. L. Qin, C. M. O. D. Alexander, R. W. Carlson, M. F. Horan, T. Yokoyama, Contributors to chromium isotope variation of meteorites. *Geochim Cosmochim Acta.* **74**, 1122–1145 (2010).
31. J. Zhang, N. Dauphas, A. M. Davis, I. Leya, A. Fedkin, The proto-Earth as a significant source of lunar material. *Nat Geosci.* **5**, 251–255 (2012).
32. J. A. M. Nanne, F. Nimmo, J. N. Cuzzi, T. Kleine, Origin of the non-carbonaceous–carbonaceous meteorite dichotomy. *Earth Planet Sci Lett.* **511**, 44–54 (2019).
33. C. Burkhardt, N. Dauphas, U. Hans, B. Bourdon, T. Kleine, Elemental and isotopic variability in solar system materials by mixing and processing of primordial disk reservoirs. *Geochim Cosmochim Acta.* **261**, 145–170 (2019).
34. M. Ito, N. Tomioka, M. Uesugi, A. Yamaguchi, N. Shirai, T. Ohigashi, M. C. Liu, R. C. Greenwood, M. Kimura, N. Imae, K. Uesugi, A. Nakato, K. Yogata, H. Yuzawa, Y. Kodama, A. Tsuchiyama, M. Yasutake, R. Findlay, I. A. Franchi, J. A. Malley, K. A. McCain, N. Matsuda, K. D. McKeegan, K. Hirahara, A. Takeuchi, S. Sekimoto, I. Sakurai,

- I. Okada, Y. Karouji, M. Arakawa, A. Fujii, M. Fujimoto, M. Hayakawa, N. Hirata, N. Hirata, R. Honda, C. Honda, S. Hosoda, Y. ichi Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, K. Kawahara, S. Kikuchi, K. Kitazato, K. Matsumoto, M. Matsuoka, T. Michikami, Y. Mimasu, A. Miura, O. Mori, T. Morota, S. Nakazawa, N. Namiki, H. Noda, R. Noguchi, N. Ogawa, K. Ogawa, T. Okada, C. Okamoto, G. Ono, M. Ozaki, T. Saiki, N. Sakatani, H. Sawada, H. Senshu, Y. Shimaki, K. Shirai, S. Sugita, Y. Takei, H. Takeuchi, S. Tanaka, E. Tatsumi, F. Terui, R. Tsukizaki, K. Wada, M. Yamada, T. Yamada, Y. Yamamoto, H. Yano, Y. Yokota, K. Yoshihara, M. Yoshikawa, K. Yoshikawa, R. Fukai, S. Furuya, K. Hatakeda, T. Hayashi, Y. Hitomi, K. Kumagai, A. Miyazaki, M. Nishimura, H. Soejima, A. Iwamae, D. Yamamoto, M. Yoshitake, T. Yada, M. Abe, T. Usui, S. ichiro Watanabe, Y. Tsuda, A pristine record of outer Solar System materials from asteroid Ryugu's returned sample. *Nat Astron* (2022), doi:<https://doi.org/10.1038/s41550-022-01745-5>.
35. W. F. Bottke, M. Brož, D. P. O'Brien, A. C. Bagatin, A. Morbidelli, S. Marchi, "The collisional evolution of the main asteroid belt" in *Asteroids IV* (University of Arizona Press, 2015), pp. 701–724.
36. N. Sugiura, W. Fujiya, Correlated accretion ages and $\epsilon^{54}\text{Cr}$ of meteorite parent bodies and the evolution of the solar nebula. *Meteorit Planet Sci.* **49**, 772–787 (2014).
37. A. S. Rivkin, The fraction of hydrated C-complex asteroids in the asteroid belt from SDSS data. *Icarus.* **221**, 744–752 (2012).
38. F. E. DeMeo, R. P. Binzel, S. M. Slivan, S. J. Bus, An extension of the Bus asteroid taxonomy into the near-infrared. *Icarus.* **202**, 160–180 (2009).
39. S. N. Raymond, A. Izidoro, Origin of water in the inner Solar System: planetesimals scattered inward during Jupiter and Saturn's rapid gas accretion. *Icarus.* **297**, 134–148 (2017).
40. H. F. Levison, W. F. Bottke, M. Gounelle, A. Morbidelli, D. Nesvorný, K. Tsiganis, Contamination of the asteroid belt by primordial trans-Neptunian objects. *Nature.* **460**, 364–366 (2009).
41. B. Liu, S. N. Raymond, S. A. Jacobson, Early Solar System instability triggered by dispersal of the gaseous disk. *Nature.* **604**, 643–646 (2022).
42. D. Vokrouhlický, W. F. Bottke, D. Nesvorný, Capture of trans-Neptunian planetesimals in the main asteroid belt. *Astron J.* **152**, 39 (2016).
43. R. de S. Ribeiro, A. Morbidelli, S. N. Raymond, A. Izidoro, R. Gomes, E. Vieira Neto, Dynamical evidence for an early giant planet instability. *Icarus.* **339**, 113605 (2020).
44. J. A. Fernández, W.-H. Ip, Dynamical Evolution of a Cometary Swarm in the Outer Planetary Region. *Icarus.* **47**, 470–479 (1981).
45. M. Duncan, T. Quinn, S. Tremaine, The Formation and Extent of the Solar System Comet Cloud. *Astronomical Journal.* **94**, 1330–1339 (1987).
46. D. Brownlee, P. Tsou, J. Aléon, C. M. O. D. Alexander, T. Araki, S. Bajt, G. A. Baratta, R. Bastien, P. Bland, P. Bleuet, J. Borg, J. P. Bradley, A. Brearley, F. Brenker, S. Brennan, J. C. Bridges, N. D. Browning, J. R. Brucato, E. Bullock, M. J. Burchell, H. Busemann, A. Butterworth, M. Chaussidon, A. Cheuvront, M. Chi, M. J. Cintala, B. C. Clark, S. J. Clemett, G. Cody, L. Colangeli, G. Cooper, P. Cordier, C. Daghlán, Z. Dai, Z. Djouadi, G. Dominguez, T. Duxbury, J. P. Dworkin, D. S. Ebel, T. E. Economou, S. Fakra, S. A. J. Fairey, S. Fallon, G. Ferrini, T. Ferroir, H. Fleckenstein, C. Floss, G. Flynn, I. A. Franchi, M. Fries, Z. Gainsforth, J. Gallien, M. Genge, M. K. Gilles, P. Gillet, J. Gilmour, D. P. Glavin, M. Gounelle, M. M. Grady, G. A. Graham, P. G. Grant, S. F. Green, F. Grossemy, L. Grossman, J. N. Grossman, Y. Guan, K. Hagiya, R. Harvey, P. Heck, G. F. Herzog, P. Hoppe, F. Hörz, J. Huth, I. D. Hutcheon, K. Ignatyev, H. Ishii, M. Ito, D. Jacob, C. Jacobsen, S. Jacobsen, S. Jones, D. Joswiak, A. Jurewicz, A. T. Kearsley, L. P. Keller, H. Khodja, A. David Kilcoyne, J. Kissel, A. Krot, F. Langenhorst, A. Lanzirotti, L. Le, L. A. Leshin, J. Leitner, L. Lemelle, H. Leroux, M.-C. Liu, K. Luening, I. Lyon, G. MacPherson, M. A. Marcus, K. Marhas, B. Marty, G. Matrajt, K. McKeegan, D. A. Papanastassiou, R. Palma, M. E. Palumbo, R. O. Pepin, D. Perkins, M. Perronnet, P. Pianetta, W. Rao, F. J. M. Rietmeijer, F. Robert, D. Rost, A. Rotundi, R. Ryan, S. A. Sandford, C. S. Schwandt, T. H. See, D. Schlutter, J. Sheffield-Parker, A. Simionovici, S. Simon, I. Sitnitsky, C. J. Snead, M. K. Spencer, F. J. Stadermann, A. Steele, T. Stephan, R. Stroud, J. Susini, S. R. Sutton, Y.

- Suzuki, M. Taheri, S. Taylor, N. Teslich, K. Tomeoka, N. Tomioka, A. Toppani, J. M. Trigo-Rodríguez, D. Troadec, A. Tsuchiyama, A. J. Tuzzolino, T. Tyliczszak, K. Uesugi, M. Velbel, J. Vellenga, E. Vicenzi, L. Vincze, J. Warren, I. Weber, M. Weisberg, A. J. Westphal, S. Wirick, D. Wooden, B. Wopenka, P. Wozniakiewicz, I. Wright, H. Yabuta, H. Yano, E. D. Young, R. N. Zare, T. Zega, K. Ziegler, L. Zimmerman, E. Zinner, M. Zolensky, Comet 81P/Wild2 under a microscope. *Science* (1979). **314**, 1711–1716 (2006).
47. C. M. O. D. Alexander, R. Bowden, M. L. Fogel, K. T. Howard, C. D. K. Herd, L. R. Nittler, The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets. *Science*. **337**, 721–723 (2012).
48. D. R. Müller, K. Altwegg, J. J. Berthelier, M. Combi, J. de Keyser, S. A. Fuselier, N. Hänni, B. Pestoni, M. Rubin, I. R. H. G. Schroeder, S. F. Wampfler, High D/H ratios in water and alkanes in comet 67P/Churyumov-Gerasimenko measured with Rosetta/ROSINA DFMS. *Astron Astrophys.* **662**, A69 (2022).
49. L. Yang, F. J. Ciesla, C. M. O. D. Alexander, The D/H ratio of water in the solar nebula during its formation and evolution. *Icarus*. **226**, 256–267 (2013).
50. D. C. Lis, D. Bockelée-Morvan, R. Güsten, N. Biver, J. Stutzki, Y. Delorme, C. Durán, H. Wiesemeyer, Y. Okada, Terrestrial deuterium-to-hydrogen ratio in water in hyperactive comets. *Astron Astrophys.* **625**, L5 (2019).
51. R. H. Brown, D. S. Lauretta, B. Schmidt, J. Moores, "Experimental and theoretical simulations of ice sublimation with implications for the chemical, isotopic, and physical evolution of icy objects" in *Planetary and Space Science* (2012), vol. 60, pp. 166–180.
52. C. M. O. D. Alexander, S. D. Newsome, M. L. Fogel, L. R. Nittler, H. Busemann, G. D. Cody, Deuterium enrichments in chondritic macromolecular material-Implications for the origin and evolution of organics, water and asteroids. *Geochim Cosmochim Acta*. **74**, 4417–4437 (2010).
53. A. J. Brearley, "The action of water" in *Meteorites and the Early Solar System*, D. S. Lauretta, H. Y. McSween, Eds. (University of Arizona Press, II., 2006), pp. 587–624.
54. E. Nakamura, K. Kobayashi, R. Tanaka, T. Kunihiro, H. Kitagawa, C. Potiszil, T. Ota, C. Sakaguchi, M. Yamanaka, D. M. Ratnayake, H. Tripathi, R. Kumar, M. L. Avramescu, H. Tsuchida, Y. Yachi, H. Miura, M. Abe, R. Fukai, S. Furuya, K. Hatakeda, T. Hayashi, Y. Hitomi, K. Kumagai, A. Miyazaki, A. Nakato, M. Nishimura, T. Okada, H. Soejima, S. Sugita, A. Suzuki, T. Usui, T. Yada, D. Yamamoto, K. Yogata, M. Yoshitake, M. Arakawa, A. Fujii, M. Hayakawa, N. Hirata, N. Hirata, R. Honda, C. Honda, S. Hosoda, Y. I. Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, K. Kawahara, S. Kikuchi, K. Kitazato, K. Matsumoto, M. Matsuoka, T. Michikami, Y. Mimasu, A. Miura, T. Morota, S. Nakazawa, N. Namiki, H. Noda, R. Noguchi, N. Ogawa, K. Ogawa, C. Okamoto, G. Ono, M. Ozaki, T. Saiki, N. Sakatani, H. Sawada, H. Senshu, Y. Shimaki, K. Shirai, Y. Takei, H. Takeuchi, S. Tanaka, E. Tatsumi, F. Terui, R. Tsukizaki, K. Wada, M. Yamada, T. Yamada, Y. Yamamoto, H. Yano, Y. Yokota, K. Yoshihara, M. Yoshikawa, K. Yoshikawa, M. Fujimoto, S. I. Watanabe, Y. Tsuda, On the origin and evolution of the asteroid Ryugu: A comprehensive geochemical perspective. *Proc Jpn Acad Ser B Phys Biol Sci*. **98**, 227–282 (2022).
55. M. Gounelle, A. Morbidelli, P. A. Bland, E. D. Young, M. Sephton, "Meteorites from the outer solar system?" in *The Solar System Beyond Neptune*, M. A. Barucci, H. Boehnhardt, D. P. Cruikshank, A. Morbidelli, Eds. (Univ Arizona Press, 2008), pp. 525–541.
56. E. L. Berger, T. J. Zega, L. P. Keller, D. S. Lauretta, Evidence for aqueous activity on comet 81P/Wild 2 from sulfide mineral assemblages in Stardust samples and CI chondrites. *Geochim Cosmochim Acta*. **75**, 3501–3513 (2011).
57. C. M. Lisse, J. Vancleve, A. C. Adams, M. F. A'hearn, Y. R. Fernández, T. L. Farnham, L. Armus, C. J. Grillmair, J. Ingalls, M. J. S. Belton, O. Groussin, L. A. Mcfadden, K. J. Meech, P. H. Schultz, B. C. Clark, L. M. Feaga, J. M. Sunshine, Spitzer spectral observations of the deep impact ejecta. *Science* (1979). **313**, 635–640 (2006).
58. I. Ferrin, C. Gil, The aging of comets Halley and Encke. *Astron Astrophys.* **194**, 288–296 (1988).
59. T. Okada, T. Fukuhara, S. Tanaka, M. Taguchi, T. Arai, H. Senshu, N. Sakatani, Y. Shimaki, H. Demura, Y. Ogawa, K. Suko, T. Sekiguchi, T. Kouyama, J. Takita, T.

- Matsunaga, T. Imamura, T. Wada, S. Hasegawa, J. Helbert, T. G. Müller, A. Hagermann, J. Biele, M. Grott, M. Hamm, M. Delbo, N. Hirata, N. Hirata, Y. Yamamoto, S. Sugita, N. Namiki, K. Kitazato, M. Arakawa, S. Tachibana, H. Ikeda, M. Ishiguro, K. Wada, C. Honda, R. Honda, Y. Ishihara, K. Matsumoto, M. Matsuoka, T. Michikami, A. Miura, T. Morota, H. Noda, R. Noguchi, K. Ogawa, K. Shirai, E. Tatsumi, H. Yabuta, Y. Yokota, M. Yamada, M. Abe, M. Hayakawa, T. Iwata, M. Ozaki, H. Yano, S. Hosoda, O. Mori, H. Sawada, T. Shimada, H. Takeuchi, R. Tsukizaki, A. Fujii, C. Hirose, S. Kikuchi, Y. Mimasu, N. Ogawa, G. Ono, T. Takahashi, Y. Takei, T. Yamaguchi, K. Yoshikawa, F. Terui, T. Saiki, S. Nakazawa, M. Yoshikawa, S. Watanabe, Y. Tsuda, Highly porous nature of a primitive asteroid revealed by thermal imaging. *Nature*. **579**, 518–522 (2020).
60. D. Nesvorný, Dynamical evolution of the Early Solar System. *Annu. Rev. Astron. Astrophys.* **56**, 137–174 (2018).
 61. S. Hasegawa, M. Marsset, F. E. DeMeo, S. J. Bus, J. Geem, M. Ishiguro, M. Im, D. Kuroda, P. Vernazza, Discovery of two TNO-like bodies in the asteroid belt. *Astrophys J Lett.* **916**, L6 (2021).
 62. T. Yada, M. Abe, T. Okada, A. Nakato, K. Yogata, A. Miyazaki, K. Hatakeda, K. Kumagai, M. Nishimura, Y. Hitomi, H. Soejima, M. Yoshitake, A. Iwamae, S. Furuya, M. Uesugi, Y. Karouji, T. Usui, T. Hayashi, D. Yamamoto, R. Fukai, S. Sugita, Y. Cho, K. Yumoto, Y. Yabe, J. P. Bibring, C. Pilorget, V. Hamm, R. Brunetto, L. Riu, L. Louri, D. Loizeau, G. Lequertier, A. Moussi-Soffys, S. Tachibana, H. Sawada, R. Okazaki, Y. Takano, K. Sakamoto, Y. N. Miura, H. Yano, T. R. Ireland, T. Yamada, M. Fujimoto, K. Kitazato, N. Namiki, M. Arakawa, N. Hirata, H. Yurimoto, T. Nakamura, T. Noguchi, H. Yabuta, H. Naraoka, M. Ito, E. Nakamura, K. Uesugi, K. Kobayashi, T. Michikami, H. Kikuchi, N. Hirata, Y. Ishihara, K. Matsumoto, H. Noda, R. Noguchi, Y. Shimaki, K. Shirai, K. Ogawa, K. Wada, H. Senshu, Y. Yamamoto, T. Morota, R. Honda, C. Honda, Y. Yokota, M. Matsuoka, N. Sakatani, E. Tatsumi, A. Miura, M. Yamada, A. Fujii, C. Hirose, S. Hosoda, H. Ikeda, T. Iwata, S. Kikuchi, Y. Mimasu, O. Mori, N. Ogawa, G. Ono, T. Shimada, S. Soldini, T. Takahashi, Y. Takei, H. Takeuchi, R. Tsukizaki, K. Yoshikawa, F. Terui, S. Nakazawa, S. Tanaka, T. Saiki, M. Yoshikawa, S. ichiro Watanabe, Y. Tsuda, Preliminary analysis of the Hayabusa2 samples returned from C-type asteroid Ryugu. *Nat Astron.* **6**, 214–220 (2022).
 63. S. Weyer, J. B. Schwieters, High precision Fe isotope measurements with high mass resolution MC-ICPMS. *Int J Mass Spectrom.* **226**, 355–368 (2003).
 64. C. Burkhardt, F. Spitzer, A. Morbidelli, G. Budde, J. H. Render, T. S. Kruijer, T. Kleine, Terrestrial planet formation from lost inner solar system material. *Sci. Adv.* **7**, 7601 (2021).
 65. N. Dauphas, The isotopic nature of the Earth’s accreting material through time. *Nature*. **541**, 521–524 (2017).
 66. C. Burkhardt, N. Dauphas, H. Tang, M. Fischer-Gödde, L. Qin, J. H. Chen, S. S. Rout, A. Pack, P. R. Heck, D. A. Papanastassiou, In search of the Earth-forming reservoir: Mineralogical, chemical, and isotopic characterizations of the ungrouped achondrite NWA 5363/NWA 5400 and selected chondrites. *Meteorit Planet Sci.* **52**, 807–826 (2017).
 67. M. E. Sanborn, J. Wimpenny, C. D. Williams, A. Yamakawa, Y. Amelin, A. J. Irving, Q. Z. Yin, Carbonaceous achondrites Northwest Africa 6704/6693: Milestones for early Solar System chronology and genealogy. *Geochim Cosmochim Acta.* **245**, 577–596 (2019).
 68. S. Gerber, C. Burkhardt, G. Budde, K. Metzler, T. Kleine, Mixing and transport of dust in the early solar nebula as inferred from titanium isotope variations among chondrules. *Astrophys J.* **841**, L17 (2017).
 69. Z. A. Torrano, D. L. Schrader, J. Davidson, R. C. Greenwood, D. R. Dunlap, M. Wadhwa, The relationship between CM and CO chondrites: Insights from combined analyses of titanium, chromium, and oxygen isotopes in CM, CO, and ungrouped chondrites. *Geochim Cosmochim Acta.* **301**, 70–90 (2021).
 70. C. D. Williams, M. E. Sanborn, C. Defouilloy, Q.-Z. Yin, N. T. Kita, D. S. Ebel, A. Yamakawa, K. Yamashita, K. performed research, K. analyzed data, Chondrules reveal large-scale outward transport of inner Solar System materials in the protoplanetary disk. *National Academy of Sciences of the United States of America.* **117**, 23426–23435 (2020).

71. A. Anand, J. Pape, M. Wille, K. Mezger, B. Hofmann, Early differentiation of magmatic iron meteorite parent bodies from Mn-Cr chronometry. *Geochem Perspect Lett.* **20**, 6–10 (2021).
72. C. S. Dey, Q.-Z. Yin, M. Zolensky, "Exploring the planetary genealogy of Tarda - A unique new carbonaceous chondrite" in *52nd Lunar and Planetary Science Conference* (2021), p. 2548.
73. B. Mougél, F. Moynier, C. Göpel, Chromium isotopic homogeneity between the Moon, the Earth, and enstatite chondrites. *Earth Planet Sci Lett.* **481**, 1–8 (2018).
74. C. Göpel, J. L. Birck, A. Galy, J. A. Barrat, B. Zanda, Mn-Cr systematics in primitive meteorites: Insights from mineral separation and partial dissolution. *Geochim Cosmochim Acta.* **156**, 1–24 (2015).
75. K. K. Larsen, A. Trinquier, C. Paton, M. Schiller, D. Wielandt, M. A. Ivanova, J. N. Connelly, Å. Nordlund, A. N. Krot, M. Bizzarro, Evidence for magnesium isotope heterogeneity in the solar protoplanetary disk. *Astrophysical Journal Letters.* **735**, L37 (2011).
76. E. M. M. E. van Kooten, D. Wielandt, M. Schiller, K. Nagashima, A. Thomen, K. K. Larsen, M. B. Olsen, Å. Nordlund, A. N. Krot, M. Bizzarro, Isotopic evidence for primordial molecular cloud material in metal-rich carbonaceous chondrites. *Proc Natl Acad Sci U S A.* **113**, 2011–2016 (2016).
77. K. Zhu, F. Moynier, M. Schiller, D. Wielandt, K. K. Larsen, E. M. M. E. van Kooten, J.-A. Barrat, M. Bizzarro, Chromium Isotopic Constraints on the Origin of the Ureilite Parent Body. *Astrophys J.* **888**, 126 (2020).
78. M. Petit, J. L. Birck, T. H. Luu, M. Gounelle, The chromium isotopic composition of the ungrouped carbonaceous chondrite Tagish lake. *Astrophysical Journal.* **736**, 23 (2011).
79. K. Zhu, J. Liu, F. Moynier, L. Qin, C. M. O. Alexander, Y. He, Chromium Isotopic Evidence for an Early Formation of Chondrules from the Ornans CO Chondrite. *Astrophys J.* **873**, 82 (2019).
80. L. Qin, D. Rumble, C. M. O. D. Alexander, R. W. Carlson, P. Jenniskens, M. H. Shaddad, The chromium isotopic composition of Almahata Sitta. *Meteorit Planet Sci.* **45**, 1771–1777 (2010).
81. M. Schiller, E. van Kooten, J. C. Holst, M. B. Olsen, M. Bizzarro, Precise measurement of chromium isotopes by MC-ICPMS. *J Anal At Spectrom.* **29**, 1406–1416 (2014).
82. K. Zhu, F. Moynier, M. Schiller, C. M. O. D. Alexander, J. A. Barrat, A. Bischoff, M. Bizzarro, Mass-independent and mass-dependent Cr isotopic composition of the Rumuruti (R) chondrites: Implications for their origin and planet formation. *Geochim Cosmochim Acta.* **293**, 598–609 (2021).
83. J. M. Schneider, C. Burkhardt, Y. Marrocchi, G. A. Brennecka, T. Kleine, Early evolution of the solar accretion disk inferred from Cr-Ti-O isotopes in individual chondrules. *Earth Planet Sci Lett.* **551** (2020).
84. E. van Kooten, L. Cavalcante, D. Wielandt, M. Bizzarro, The role of Bells in the continuous accretion between the CM and CR chondrite reservoirs. *Meteorit Planet Sci.* **55**, 575–590 (2020).
85. A. Yamakawa, K. Yamashita, A. Makishima, E. Nakamura, Chromium isotope systematics of achondrites: Chronology and isotopic heterogeneity of the inner solar system bodies. *Astrophysical Journal.* **720**, 150–154 (2010).
86. D. L. Cook, B. S. Meyer, M. Schönbächler, Iron and Nickel Isotopes in IID and IVB Iron Meteorites: Evidence for Admixture of an SN II Component and Implications for the Initial Abundance of ^{60}Fe . *Astrophys J.* **917**, 59 (2021).

Acknowledgments: Hayabusa2 was developed and built under the leadership of Japan Aerospace Exploration Agency (JAXA), with contributions from the German Aerospace Center (DLR) and the Centre National d'Études Spatiales (CNES), and in collaboration with NASA, and other universities, institutes, and companies in Japan. The curation system was developed by JAXA in collaboration with companies in Japan. We thank the Smithsonian National Museum of Natural History, the Robert A. Pritzker Center for Meteoritics at the Field Museum, the Muséum National d'Histoire Naturelle, and the Institut für Planetologie in Münster for providing carbonaceous chondrites samples for this study. Nicole X. Nie, and Reika Yokochi are thanked for discussions.

Funding: This work was supported by NASA grants NNX17AE86G (LARS), 80NSSC17K0744 (HW), 000306-002 (HW), 80NSSC21K0380 (EW), 0995GXB174 (EW), NSF grant EAR-2001098 (CSEDI), and funding from DOE to N.D.; and Kaken-hi grants to S.T., T. Y. and H. Y.

Author contributions: H.Y. and T.Y. coordinated the isotopic analyses of the samples among members of the Hayabusa2-initial-analysis chemistry team. T.H. and T.Y. processed the samples and separated Fe from the matrix. T.H. measured the Fe isotopic composition. T.H. and N.D. wrote the paper, with contributions from all co-authors.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: The authors declare that data supporting the findings of this study are available within the paper and its Supplementary Materials. Data of Hayabusa2 sample and other data from the mission are available at the DARTS archive at www.darts.isas.jaxa.jp/curation/hayabusa2 and www.darts.isas.jaxa.jp/planet/project/hayabusa2/.

Tables

Table 1. Iron isotopic compositions of Ryugu samples and carbonaceous chondrites. Calculated mass-independent ($\mu^{54}\text{Fe}$, $\mu^{58}\text{Fe}$) and mass-dependent ($\delta^{56}\text{Fe}$) Fe isotopic compositions in Ryugu samples, carbonaceous chondrites, and geostandards. Uncertainties of individual samples are 95% confidence intervals of the mean of N standard-sample-standard bracketing analyses. Uncertainties on group averages are the 95% confidence intervals of the mean. The isotopic anomalies are calculated by internal normalization to $^{57}\text{Fe}/^{56}\text{Fe}=0.023095$ and expressed in μ -notation (22, 24) defined as the parts-per-million deviation of the internally normalized $^{54}\text{Fe}/^{56}\text{Fe}$ ratio in the sample relative to bracketing measurements of IRMM-524a. The mass-dependent Fe isotopic variations are calculated by sample-standard bracketing and given as δ -notation defined **parts-per-thousand** deviation of the $^{56}\text{Fe}/^{54}\text{Fe}$ ratio of the sample relative to the IRMM-524a standard solution.

Sample	N	$\mu^{54}\text{Fe}$	$\mu^{58}\text{Fe}$	$\delta^{56}\text{Fe}$
<i>Ryugu</i>				
A0106	30	-2 $\pm$ 5	14 $\pm$ 14	-0.01 $\pm$ 0.02
A0106-A0107	30	2 $\pm$ 5	10 $\pm$ 10	-0.01 $\pm$ 0.02
C0107	25	4 $\pm$ 4	15 $\pm$ 17	-0.02 $\pm$ 0.02
C0108	25	2 $\pm$ 5	11 $\pm$ 7	0.01 $\pm$ 0.02
Average Ryugu		1$\pm$4	13$\pm$4	-0.01$\pm$0.02
<i>CI chondrites</i>				
Orgueil-1	30	2 $\pm$ 5	12 $\pm$ 8	-0.02 $\pm$ 0.02
Orgueil-2	30	3 $\pm$ 5	2 $\pm$ 11	0.01 $\pm$ 0.02
Orgueil-3	14	4 $\pm$ 9	3 $\pm$ 12	0.00 $\pm$ 0.02
Orgueil-4	30	3 $\pm$ 5	12 $\pm$ 8	0.06 $\pm$ 0.02
Ivuna	30	1 $\pm$ 4	16 $\pm$ 11	0.00 $\pm$ 0.02
Alais	28	6 $\pm$ 6	17 $\pm$ 12	-0.02 $\pm$ 0.02
Average CI		3$\pm$2	10$\pm$6	0.00$\pm$0.02
<i>CM chondrites</i>				
Murchison-1	14	23 $\pm$ 5	6 $\pm$ 7	0.00 $\pm$ 0.02
Murchison-2	15	18 $\pm$ 6	2 $\pm$ 9	0.01 $\pm$ 0.02
Mighei	14	27 $\pm$ 8	7 $\pm$ 9	-0.01 $\pm$ 0.02
Average CM		24$\pm$7	6$\pm$11	0.00$\pm$0.02
<i>CV chondrites</i>				
Allende-1	15	29 $\pm$ 4	2 $\pm$ 9	-0.02 $\pm$ 0.02
Allende-2	15	32 $\pm$ 6	2 $\pm$ 9	-0.05 $\pm$ 0.02
Allende-3	15	21 $\pm$ 7	3 $\pm$ 12	0.08 $\pm$ 0.02
Vigarano	15	29 $\pm$ 5	-7 $\pm$ 13	0.02 $\pm$ 0.02
Average CV		28$\pm$6	0$\pm$6	0.01$\pm$0.08
<i>CO chondrites</i>				
Ormans	15	21 $\pm$ 5	0 $\pm$ 8	-0.03 $\pm$ 0.02
Lance	15	18 $\pm$ 5	1 $\pm$ 10	0.01 $\pm$ 0.02
<i>CR chondrites</i>				
Acfer 139	14	38 $\pm$ 8	9 $\pm$ 11	-0.11 $\pm$ 0.02
GRA 06100	15	39 $\pm$ 5	9 $\pm$ 6	0.00 $\pm$ 0.02
<i>Ungrouped chondrites</i>				
Tagish Lake-1	15	18 $\pm$ 7	2 $\pm$ 14	-0.03 $\pm$ 0.02
Tagish Lake-2	15	15 $\pm$ 6	2 $\pm$ 22	0.05 $\pm$ 0.02
Tarda	30	24 $\pm$ 8	3 $\pm$ 12	-0.01 $\pm$ 0.02
<i>Geostandards</i>				
BHVO-2-1	15	1 $\pm$ 7	-9 $\pm$ 8	0.09 $\pm$ 0.02
BHVO-2-2	15	1 $\pm$ 7	-1 $\pm$ 11	0.06 $\pm$ 0.02
AGV-2	30	2 $\pm$ 4	1 $\pm$ 6	0.08 $\pm$ 0.02
Average Geostandards		1$\pm$1	-3$\pm$14	0.09$\pm$0.06

Figure captions




Fig. 1. Previously published isotopic anomalies of Ti, Cr, and O in Ryugu and other Solar System materials. Plots of $\mu^{50}\text{Ti}$ vs. $\mu^{54}\text{Cr}$ (A), $\mu^{50}\text{Ti}$ vs. $\Delta^{17}\text{O}$ (B), and $\mu^{54}\text{Cr}$ vs. $\Delta^{17}\text{O}$ (C). In these diagrams, Ryugu and CI represent an endmember to the CC array. Data for Ryugu are from Yokoyama et al. (6). The average Ti and Cr isotopic compositions of non-carbonaceous (NC) and carbonaceous (CC) meteorite groups and Earth's mantle are from the data compilation of Burkhardt et al. (64) and O isotopic data from the compilation of Dauphas (65). Uncertainties for individual groups are the 95% confidence interval of the mean. If not visible, error bars are smaller than symbols.




Fig. 2. Fe isotopic anomalies of returned samples from Cb-type asteroid (162173) Ryugu and carbonaceous chondrites (Table 1). Ryugu samples and CI chondrites have identical $\mu^{54}\text{Fe}$ values, which is distinct from all other carbonaceous chondrite groups (CM, CV, CO, CR, Ung=ungrouped). The Open triangle is the CI chondrite average from Schiller et al. (22).




Fig. 3. Isotopic anomalies of Fe and Ti in Solar System materials. Ryugu samples and CI chondrites have identical $\mu^{54}\text{Fe}$ and $\mu^{50}\text{Ti}$ values that are distinct from other meteorites. Red circles correspond to non-carbonaceous (NC) chondrite groups (E=enstatite; R=rumuruti; OC=ordinary chondrites), red diamonds to NC achondrites (Ure=ureilites; Dio=diogenites), and blue circles to carbonaceous (CC) chondrite groups (TL=Tagish Lake). The green square is Earth's mantle. The average composition of Ryugu and Ivuna-type carbonaceous chondrites (CI) are shown as triangles. Average compositions of meteorite groups and Earth's mantle are calculated using data from this study (Table 1) and the data compilation of Burkhardt et al. (64) (Table S1). **If not visible, error** bars are smaller than symbols.




Fig. 4. Schematic of the possible source region of Cb-type asteroids and CI chondrites. Planetesimals formed in different regions of the protoplanetary disk. Volatile-poor planetesimals (red circles) formed in the inner region, while volatile-rich planetesimals (blue circles) formed beyond Jupiter's orbit. The growth and migration of the gas and ice giant planets implanted some of the planetesimals into the Main Belt (small arrows), while the majority of planetesimals were transported outwards or ejected from the disk (large arrows) (39). A plausible explanation for the different Fe nucleosynthetic heritage and primitive chemical composition of CI chondrites and Ryugu is that they were implanted in the Main Belt by excitation from Uranus and Neptune (filled **bright** blue circles), while other CCs formed in more internal regions near Jupiter and Saturn (filled dark blue circles) (Fig. S4). The icy planetesimals that were formed around Uranus and Neptune and were ejected outwards went to populate the Oort cloud (44, 45). CI chondrites and Ryugu may thus share some parentage with long-period comets. Such a scenario could explain the trichotomy between NC, CC, and CI for nucleosynthetic anomalies (Fig. 3).

Supplementary Materials for

Ryugu's nucleosynthetic heritage from the outskirts of the Solar System

Timo Hopp*, Nicolas Dauphas, Yoshinari Abe, Jérôme Aléon, Conel M. O'D. Alexander, Sachiko Amari, Yuri Amelin, Ken-ichi Bajo, Martin Bizzarro, Audrey Bouvier, Richard W. Carlson, Marc Chaussidon, Byeon-Gak Choi, Andrew M. Davis, Tommaso Di Rocco, Wataru Fujiya, Ryota Fukai, Ikshu Gautam, Makiko K. Haba, Yuki Hibiya, Hiroshi Hidaka, Hisashi Homma, Peter Hoppe, Gary R. Huss, Kiyohiro Ichida, Tsuyoshi Iizuka, Trevor R. Ireland, Akira Ishikawa, Motoo Ito, Shoichi Itoh, Noriyuki Kawasaki, Noriko T. Kita, Kouki Kitajima, Thorsten Kleine, Shintaro Komatani, Alexander N. Krot, Ming-Chang Liu, Yuki Masuda, Kevin D. McKeegan, Mayu Morita, Kazuko Motomura, Frédéric Moynier, Izumi Nakai, Kazuhide Nagashima, David Nesvorný, Ann Nguyen, Larry Nittler, Morihiko Onose, Andreas Pack, Changkun Park, Laurette Piani, Liping Qin, Sara S. Russell, Naoya Sakamoto, Maria Schönbächler, Lauren Tafla, Haolan Tang, Kentaro Terada, Yasuko Terada, Tomohiro Usui, Sohei Wada, Meenakshi Wadhwa, Richard J. Walker, Katsuyuki Yamashita, Qing-Zhu Yin, Tetsuya Yokoyama, Shigekazu Yoneda, Edward D. Young, Hiroharu Yui, Ai-Cheng Zhang, Tomoki Nakamura, Hiroshi Naraoka, Takaaki Noguchi, Ryuji Okazaki, Kanako Sakamoto, Hikaru Yabuta, Masanao Abe, Akiko Miyazaki, Aiko Nakato, Masahiro Nishimura, Tatsuaki Okada, Toru Yada, Kasumi Yogata, Satoru Nakazawa, Takanao Saiki, Satoshi Tanaka, Fuyuto Terui, Yuichi Tsuda, Sei-ichiro Watanabe, Makoto Yoshikawa, Shogo Tachibana, Hisayoshi Yurimoto

*Corresponding author. Email: hopp@mps.mpg.de

This PDF file includes:

Figs. S1 to S4
Tables S1 and S2




Fig. S1. Comparison of $\mu^{54}\text{Fe}$ values of carbonaceous chondrites determined in this study with literature data of Schiller et al. (22). The data from this study agree with previous studies. Uncertainties of individual samples are 95% confidence intervals calculated from replicate analysis (N=14-30).




Fig. S2. Average nucleosynthetic isotopic anomalies of carbonaceous chondrites and Ryugu samples normalized to $^{57}\text{Fe}/^{56}\text{Fe}$. All carbonaceous chondrites except CI chondrites and Ryugu display resolvable $\mu^{54}\text{Fe}$ (dark blue lines). The $\mu^{58}\text{Fe}$ values of most carbonaceous chondrites are not resolvable from the terrestrial standard. While the calculated averages of CI chondrites and Ryugu display excesses in $\mu^{58}\text{Fe}$ resolvable from the standard, these are not resolvable from the other carbonaceous chondrites. Comparison with expected anomalies produced by sole variations in ^{54}Fe , ^{56}Fe , and ^{57}Fe (black lines) reveals that the observed $\mu^{54}\text{Fe}$ anomalies are best explained by ^{54}Fe variations (24). Exposure to galactic cosmic rays induces positive shifts in $\mu^{54}\text{Fe}$ that correlate with negative shifts in $\mu^{58}\text{Fe}$ (24, 26). If Fe isotopic compositions of CI chondrites and Ryugu had been affected by exposure to galactic cosmic rays, their primary Fe isotopic compositions would be even more distinct from other carbonaceous chondrites because the pre-exposure $\mu^{54}\text{Fe}$ and $\mu^{58}\text{Fe}$ values would shift towards more negative and positive values, respectively.




Fig. S3. Isotopic anomalies of Fe and Cr in Ryugu and other Solar System materials. The average $\mu^{54}\text{Fe}$ and $\mu^{54}\text{Cr}$ data of Ryugu samples are from this study and Yokoyama et al. (6). The average compositions of non-carbonaceous (NC) and carbonaceous (CC) meteorite groups and Earth's mantle (BSE) are given in the data compilation Table S1 and are calculated using data from this study and the data compilation of Burkhardt et al. (64). Uncertainties for individual groups are the 95% confidence interval of the mean. E-enstatite; R-rumuruti; OC-ordinary chondrites; Ure=ureilites; Dio=diogenites. Error bars are smaller than symbols if not visible.




Fig. S4. Example of possible source regions for CC and CI asteroids in the Main Belt [modified from Fig. 12 of (39)]. This figure depicts the fractions of asteroids from different starting orbital radii that are implanted into the Main Belt in a simulation involving the growth and migration of the gas and ice giant planets. In that simulation, Jupiter, Saturn, Uranus, and Neptune start with initial orbital radii of 10, 15, 20, and 25 AU and grow and migrate inwards on a timescale of ~ 2.5 Myr (39). As shown, the processes of planetesimal excitation by the giant planets and interaction with nebular gas leads to a far-reaching and uneven sampling of planetesimals. We speculate that CC chondrites could have formed in the birth region of Jupiter and Saturn while CI chondrites and Ryugu derived from planetesimals that could have been implanted into the Main Belt by the growth and migration of Uranus and Neptune, explaining their distinct isotopic and chemical heritage. **While the fraction of planetesimals implanted into the Main Belt from the outskirts of the Solar System (e.g., CI) is small ($\sim 1\%$ of the planetesimals located in these regions are implanted into the Main Belt), the total number of planetesimals from these regions implanted into the Main Belt can be significant because of more total mass. This could explain that based on spectral observations up to 20% of carbonaceous (C-type) asteroids are of Cb-type.**

Table S1: Average O, Ti, Cr, and Fe isotopic anomalies of Ryugu, meteorites, and Earth's mantle. Averages adopted from the data compilation of Dauphas (2017) (65) and Burkhardt et al. (64) and updated with more recently published data.

Samples	Reservoir	$\Delta^{17}\text{O}$	$\pm$	95% CI	$\mu^{50}\text{Ti}$	$\pm$	95% CI	$\mu^{54}\text{Cr}$	$\pm$	95% CI	$\mu^{54}\text{Fe}$	$\pm$	95% CI	$\mu^{58}\text{Fe}$	$\pm$	95% CI
Hayabusa2																
Ryugu A	CC		$\pm$		163	$\pm$	20	135	$\pm$	21	0	$\pm$	4	12	$\pm$	4
Ryugu C	CC		$\pm$		202	$\pm$	19	125	$\pm$	10	3	$\pm$	3	13	$\pm$	4
Ryugu Mean	CC	0.61	$\pm$	0.08	183	$\pm$	27	127	$\pm$	18	1	$\pm$	4	13	$\pm$	4
Chondrites																
CI	CC	0.39	$\pm$	0.10	189	$\pm$	15	155	$\pm$	10	2	$\pm$	3	10	$\pm$	7
CM	CC	-2.92	$\pm$	0.44	301	$\pm$	10	101	$\pm$	13	23	$\pm$	6	5	$\pm$	6
CV	CC	3.62	$\pm$	0.48	345	$\pm$	19	94	$\pm$	8	27	$\pm$	6	0	$\pm$	7
CO	CC	-4.32	$\pm$	0.26	377	$\pm$	99	77	$\pm$	33	19	$\pm$	5	1	$\pm$	2
CK	CC	-4.47	$\pm$	0.21	342	$\pm$	105	48	$\pm$	42	26	$\pm$	6		$\pm$	
CR	CC	-1.48	$\pm$	0.55	251	$\pm$	45	128	$\pm$	7	33	$\pm$	7		$\pm$	
CH	CC	-1.55		0.27				137	$\pm$	29	16	$\pm$	7	9	$\pm$	0
CB	CC				204		7	120		9						
Tagish Lake	CC	-0.91	$\pm$	0.53	276	$\pm$	26	133	$\pm$	26	16	$\pm$	5	2	$\pm$	12
Tarda	CC							122	$\pm$	10	24	$\pm$	8	3	$\pm$	12
EC	NC	-0.02	$\pm$	0.05	-20	$\pm$	8	3	$\pm$	3	6	$\pm$	1			
OC	NC	0.98	$\pm$	0.48	-66	$\pm$	6	-37	$\pm$	6	11	$\pm$	3			
R	NC	2.64	$\pm$	0.15		$\pm$		-7	$\pm$	3	6	$\pm$	1			
Achondrites/ Iron meteorites																
Diogenites	NC	-0.29	$\pm$	0.08	-123	$\pm$	5	-69	$\pm$	8	14	$\pm$	3			
Ureilites	NC	-1.16	$\pm$	0.15	-200	$\pm$	32	-90	$\pm$	4	12	$\pm$	2			
IIAB	NC							-83		17	16	$\pm$	4			
IIIAB	NC							-78		6	11	$\pm$	6			
IVA	NC							-47		6	8	$\pm$	4			
Earth's mantle																
		0	$\pm$	0	-2	$\pm$	3	10	$\pm$	13	0	$\pm$	1	1	$\pm$	5

Data sources: O – Data compilation of (65), (6); Ti – (6, 10, 66–70); Cr – (6, 11, 29, 30, 66, 67, 69–85); Fe – (22, 24, 26, 86).

Table S2: Fe isotopic data of samples investigated in this study. Calculated mass-independent ($\mu^{54}\text{Fe}$, $\mu^{58}\text{Fe}$) and mass-dependent ($\delta^{5X}\text{Fe}$) Fe isotopic compositions in Ryugu samples, carbonaceous chondrites, and geostandards. Uncertainties of individual samples are 95% confidence intervals of the mean of standard-sample-standard bracketing analyses (N). The chemical procedure used at Tokyo Institute of Technology is described in the Methods and Yokoyama et al. (6). Samples processed at the University of Chicago (UofC) were digestion aliquots taken after digestion with HF-HNO₃-HCl-HClO₄ mixtures. Matrix aliquots containing the bulk Fe of samples processed at the Institut für Planetologie Münster (IfP) were digested HF-HNO₃-HCl-HClO₄ mixtures and then separated from Sr using cation exchange resin (AG50W-X8). *Samples for which soluble organic matter was extracted prior digestion. †Sample analyzed with XRF before digestion.

Sample	Mass digested (g)	Mass homogenized (g)	Processed?	N	$\mu^{54}\text{Fe}$	$\mu^{58}\text{Fe}$	$\delta^{56}\text{Fe}$	$\delta^{57}\text{Fe}$	$\delta^{58}\text{Fe}$
<i>Ryugu</i>									
A0106*	0.0146	0.0175	Fe, U (Tokyo Tech)	30	-2±5	14±14	-0.01±0.02	-0.01±0.02	0.00±0.05
A0106-A0107	0.0239	0.0289	Fe, U (Tokyo Tech)	30	2±5	10±10	-0.01±0.02	-0.02±0.03	-0.08±0.05
C0107*	0.0128	0.0174	Fe, U (Tokyo Tech)	25	4±4	15±17	-0.02±0.02	-0.03±0.03	-0.02±0.05
C0108†	0.0222	0.0333	Fe, U (Tokyo Tech)	25	2±5	11±7	0.01±0.02	0.02±0.03	0.03±0.05
<i>CI</i>									
Orgueil-1	0.401	1.12	Sr matrix cut (IfP)	30	2±5	12±8	-0.02±0.02	-0.02±0.03	-0.02±0.05
Orgueil-2	0.012	~0.1	Fe (UofC)	30	3±5	2±11	0.01±0.02	0.02±0.03	0.02±0.05
Orgueil-3	0.010	~0.1	Fe (UofC)	14	4±9	3±12	0.00±0.02	0.00±0.03	0.00±0.05
Orgueil-4	0.020	0.050	Fe, U (Tokyo Tech)	30	3±5	12±8	0.06±0.02	0.09±0.03	0.13±0.05
Ivuna	0.099	~0.1	Fe (UofC)	30	1±4	16±11	0.00±0.02	0.00±0.03	0.01±0.05
Alais	0.022	0.051	Fe, U (Tokyo Tech)	28	6±6	17±12	-0.02±0.02	-0.02±0.03	-0.01±0.05
<i>CM</i>									
Murchison-1	0.010	~0.3	Fe (UofC)	14	23±5	6±7	0.00±0.02	0.01±0.03	0.03±0.05
Murchison-2	0.025	1.65	Fe, U (Tokyo Tech)	15	18±6	2±9	0.01±0.02	0.03±0.03	0.05±0.05
Mighei	0.011	~0.1	Fe (UofC)	14	27±8	7±9	-0.01±0.02	0.00±0.03	0.01±0.05
<i>CV</i>									
Allende-1	0.012	~4000 (USNM)	Fe (UofC)	15	29±4	2±9	-0.02±0.02	-0.02±0.03	-0.01±0.05
Allende-2	0.514	~100 g (MS-A)	Sr matrix cut (IfP)	15	32±6	2±9	-0.05±0.02	-0.06±0.03	-0.07±0.05
Allende-3	0.025	~4000 (USNM)	Fe, U (Tokyo Tech)	15	21±7	3±12	0.08±0.02	0.13±0.03	0.18±0.05
Vigarano	0.014	~0.3	Fe (UofC)	15	29±5	-7±13	0.02±0.02	0.05±0.03	0.07±0.05
<i>CO</i>									
Ornans	0.041	~0.3	Fe (UofC)	15	21±5	0±8	-0.03±0.02	-0.03±0.03	-0.04±0.05
Lance	0.092	~0.5	Fe (UofC)	15	18±5	1±10	0.01±0.02	0.03±0.03	0.04±0.05
<i>CR</i>									
Acfer 139	0.525	0.525	Mo matrix cut (IfP)	14	38±8	9±11	-0.11±0.02	-0.15±0.03	-0.17±0.05
GRA 06100	0.281	0.281	Sr matrix cut (IfP)	15	39±5	9±6	0.00±0.02	0.02±0.03	0.04±0.05
<i>Ungrouped</i>									
Tagish Lake-1	0.486	1.5	Sr matrix cut (IfP)	15	18±7	2±14	-0.03±0.02	-0.04±0.03	-0.04±0.05
Tagish Lake-2	0.025	1.06	Fe, U (Tokyo Tech)	15	15±6	2±22	0.05±0.02	0.08±0.03	0.11±0.05
Tarda	0.025	0.212	Fe, U (Tokyo Tech)	30	24±8	3±12	-0.01±0.02	0.00±0.03	0.01±0.05
<i>Geostandard</i>									
BHVO-2-1	0.012	-	Fe (UofC)	15	1±7	-9±8	0.09±0.02	0.14±0.03	0.18±0.05
BHVO-2-2	0.010	-	Fe (UofC)	15	1±7	-1±11	0.06±0.02	0.10±0.03	0.13±0.05
AGV-2	0.012	-	Fe (UofC)	30	2±4	1±6	0.08±0.02	0.13±0.03	0.16±0.05