



Cite this article: Walmsley SF, Boutin S, Dantzer B, Lane JE, Coltman DW, McAdam AG. 2023 Benefits of living closer to kin vary by genealogical relationship in a territorial mammal. *Proc. R. Soc. B* **290**: 20221569. <https://doi.org/10.1098/rspb.2022.1569>

Received: 10 August 2021

Accepted: 5 December 2022

Subject Category:

Behaviour

Subject Areas:

behaviour, ecology, evolution

Keywords:

cooperation, kinship, reproductive success, survival, social evolution, familiarity

Author for correspondence:

Sam F. Walmsley

e-mail: sam.walmsley@dal.ca

[†]Present address: Department of Biology, Dalhousie University, 1355 Oxford Street, Halifax, Nova Scotia, Canada B3H 4J1.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6350103>.

Benefits of living closer to kin vary by genealogical relationship in a territorial mammal

Sam F. Walmsley^{1,†}, Stan Boutin², Ben Dantzer^{3,4}, Jeffrey E. Lane⁵, David W. Coltman^{2,6} and Andrew G. McAdam¹

¹Department of Ecology and Evolutionary Biology, University of Colorado, 1900 Pleasant Street, Boulder, CO 80309, USA

²Department of Biological Sciences, University of Alberta, 11455 Saskatchewan Drive, Edmonton, Alberta, Canada T6G 2E9

³Department of Psychology and ⁴Department of Ecology and Evolutionary Biology, University of Michigan, 500 South State Street, Ann Arbor, MI 48109, USA

⁵Department of Biology, University of Saskatchewan, 112 Science Place, Saskatoon, Canada S7N 5E2

⁶Biology Department, Western University, 1151 Richmond Street, London, Ontario, Canada N6A 5B7

SFW, 0000-0001-8577-6884; BD, 0000-0002-3058-265X; JEL, 0000-0003-0849-7181

While cooperative interactions among kin are a key building block in the societies of group-living species, their importance for species with more variable social environments is unclear. North American red squirrels (*Tamiasciurus hudsonicus*) defend individual territories in dynamic neighbourhoods and are known to benefit from living among familiar conspecifics, but not relatives. However, kin-directed behaviours may be restricted to specific genealogical relationships or strongly mediated by geographical distance, masking their influence at broader scales. Using distance between territories as a proxy for the ability of individuals to interact, we estimated the influence of primary kin (parents, offspring, siblings) on the annual survival and reproductive success of red squirrels. This approach revealed associations between fitness and access to kin, but only for certain genealogical relationships and fitness components. For example, females had enhanced annual survival when living closer to their daughters, though the reverse was not true. Most surprising was the finding that males had higher annual reproductive success when living closer to their father, suggesting possible recognition and cooperation among fathers and sons. Together, these findings point to unexpected nuance in the fitness consequences of kinship dynamics for a species that is territorial and largely solitary.

1. Introduction

How individuals disperse, settle, and die has important consequences for the evolution of cooperation, constraining the mechanisms by which affiliative relationships can form [1,2]. For example, in most group-living mammals, female-biased philopatry results in social units composed of matrilineal relatives [3]. Here, kin selection provides the opportunity for the development of stable and cooperative groups, bolstered by the additional benefits of inclusive fitness [3,4]. While the challenges of navigating group dynamics can be significant, members often rely on predictable dominance and kinship-based hierarchies to structure their interactions. For example, spotted hyenas (*Crocuta crocuta*) are born into matrilineal clans and inherit the social rank and network of their mother, a key determinant of lifelong fitness [5–7]. Less understood is whether kinship shapes cooperative relationships in species where access to kin is more variable [8–10], especially those lacking obvious social units, such as solitary species [11].

Non-traditional cooperative relationships among kin may be particularly difficult to study if they rely on more cryptic forms of interaction. For example, kin-directed behaviours may simply involve increased spatial tolerance (e.g. Galápagos sea lions, *Zalophus wollebaeki* [12]; Atlantic salmon, *Salmo salar* [13]),

or subtle reductions in antagonism (e.g. grey squirrels, *Sciurus carolinensis* [14]). An important follow-up question is whether these subtler processes should translate into meaningful effects on individual variation in survival and reproductive success. Quantifying the fitness consequences of these less-conspicuous interactions is a crucial step towards understanding whether the benefits of living with kin are mirrored in species with highly variable social environments.

Evidence from group-living species suggests that variable access to kin can result in a compensatory investment in mutualistic relationships with non-relatives. Vampire bats (*Desmodus rotundus*) rely on regular food sharing, which typically occurs among close kin [15]. However, to mitigate the risk of resting at a different roost from these preferred partners, individuals 'hedge their bets' by simultaneously developing affiliative relationships with unrelated conspecifics. Similar strategies are found in baboons: female yellow baboons (*Papio cynocephalus*) preferentially associate with their mothers or daughters but rely on bonds with non-kin when necessary [16], and female Chacma baboons (*Papio ursinus*) increase investment in new grooming relationships when a preferred kin partner dies [17]. These findings demonstrate that cooperation with unrelated individuals can buffer individuals from the risks of temporarily or permanently losing access to a close relative. In other words, despite lacking the advantage of inclusive fitness benefits, group-living animals may choose to invest in mutualistic bonds with non-kin if it leads to a more predictable social environment [18,19]. So, while kin selection is typically considered the lower-threshold pathway to prosocial bonds [20], mammals living in less-structured social environments may instead prioritize mutualistic bonds with potentially unrelated individuals.

North American red squirrels (*Tamiasciurus hudsonicus*; hereafter red squirrels) defend exclusive territories containing a central hoard of stored food resources (midden) from nearby conspecifics. Populations are characterized by fluctuating membership, driven by an influx of juveniles in the summertime [21], particularly during episodic resource pulses. Red squirrels sometimes establish territories near their parents [21,22], creating some spatial scaffolding for kin cooperation. However, mean relatedness among neighbours remains low [23], and future access to kin depends almost entirely on where an individual can establish a territory. Compared to many group-living mammals, red squirrels cannot count on living near relatives as adults. So, are red squirrels exploiting this variation in access to kin, or does the unpredictability of the social environment lead to a prioritization of mutualistic bonds with available neighbours, regardless of relatedness?

Siracusa *et al.* found that familiarity, defined as the amount of time two individuals lived within 130 m of one another, enhanced the survival and reproductive success of red squirrels, whereas the mean relatedness of neighbouring conspecifics had no effect [23]. Though consistent with the hypothesis that animals benefit by social bet-hedging when access to kin is unpredictable [15], this finding is surprising given evidence of rare but seemingly cooperative kin-biased behaviour. Related red squirrels sometimes nest communally for warmth [24], adopt orphaned juveniles [25], and even bequeath entire territories to their offspring [21]. Possible explanations for this discord could be that benefits of kin-directed behaviour only occur at finer spatial scales [23], or perhaps, that kin benefits are limited to specific genealogical

relationships (e.g. mother-daughter) [26,27], especially if kin recognition and/or affiliation are contingent on early associations in the natal nest [28], which may be the case for red squirrels [29].

Using a long-term study of wild red squirrels in Yukon, Canada, we ask whether kinship plays a role in structuring fitness-enhancing interactions in a social environment where access to kin is highly variable across individuals and over time. Red squirrels risk intrusions from neighbours when away from their territory [30], probably making it costly to interact with individuals that are far away. Following studies on kin benefits in humans [31], we aimed to exploit the spatial limitations of any affiliative behaviours, hypothesizing that the influence of kin on annualized survival and reproductive success would be mediated by geographical distance. Given the possibility that beneficial kin-directed behaviours vary by genealogical relationship [26,32], we focused our analysis on primary kin (mothers, fathers, sons, daughters, full siblings), estimating separate influences for each kinship class. We predicted that any influences of kin on annual survival or annual reproductive success (ARS) would dissipate as a function of distance. Also, while we were interested in possible interactions among fathers and offspring, we expected that benefits were more likely to occur among kin that shared a natal nest (e.g. mother–offspring dyads).

2. Material and methods

(a) Study site and long-term data collection

All data used in this study were collected as part of the Kluane Red Squirrel Project (KRSP), a long-term research project investigating populations of wild red squirrels living in southwestern Yukon, Canada (61° N, 138° W). Since 1987, KRSP has tagged, enumerated, and monitored individual red squirrels throughout their lives using alphanumeric ear tags (Monel #1; National Band and Tag, Newport, KY, USA). Red squirrel territories are typically centred on caches (middens) containing hoarded white spruce cones (*Picea glauca*). White spruce trees undergo masting events every 4–7 years, resulting in resource pulses for red squirrels.

Our analysis focused on two study areas ('Kloo' and 'Sulphur'). Both study areas were staked at 30 m intervals, with locations identified to tenths of an interval, meaning that any spatial position could be specified within approximately 3 m. Both male and female red squirrels are exclusively territorial, and typically defend the same territory year-round for life, allowing us to reliably locate individuals and map out the distances between them. Territory sizes vary considerably in this population [33], but are estimated to be 0.34 ha on average [34], corresponding to a radius of approximately 33 m for a hypothetical circular territory. Territory ownership was determined during a census each May through repeated live-trapping at each midden or nest on the study area (Tomahawk Live Trap Co., Tomahawk, WI, USA), and by behavioural observations (principally, evidence that an individual consistently used its territorial 'rattle' vocalization at a given location [30]). Pregnant females are regularly monitored in the spring so that juveniles can be tagged and linked to their maternal pedigree. Previous work has confirmed that these linkages are unlikely to be biased by rare adoption events [25]. Starting in 2003, all censused individuals have also been genotyped at 16 microsatellite loci. These data have been used to supplement maternal linkages with paternities (99% confidence, CERVUS 3.0 [35]), resulting in a comprehensive, long-term pedigree including nearly every individual living on

each study area [36]. For additional information on the study site and long-term data collection protocols, see [21,37,38].

(b) Study subjects

For all known red squirrels living on the Kloo or Sulphur study areas between 2003 and 2014, we identified pairwise relationships among primary kin (mother, father, son, daughter, full sibling) using the multigenerational pedigree and the *pedantics* package [39]. Full sibling relationships were not partitioned by sex owing to a lack of data: high rates of both multiple paternity [40], and juvenile mortality meant that it was relatively rare for full siblings to persist in the population (median two sibling dyads per study area per year, range 0–7). Given our interest in estimating annual survival, we only included individuals with known birth years and excluded any squirrels that were known to have died of unnatural causes, as these deaths were expected to be unrelated to influences of the social environment. Geographical distance between kin territories was measured as the linear distance between the centre of the primary midden (cone cache) of each squirrel, which is typically situated at the centre of the territory. Red squirrel middens are easily identifiable as piles of spruce cone scales and spines, often containing tunnels where cones are stored. We included territory locations of several individuals who lacked full, established middens but were known to defend a nest in the study area ($n = 4$). For individuals without any primary kin in the study area in a given year, we calculated their distance to the territory of a randomly selected individual, coded as 'non-kin'. Note that this category could include more distantly related kin, but no full siblings, parents, or offspring. These 'non-kin' observations served as a reference category, anchoring estimated influences of primary kin in subsequent models to a meaningful baseline. Following Siracusa *et al.* [23], we also calculated familiarity for each observation as the number of days that the dyad was known to live within 130 m of one another. This measure was mean-centred and scaled to have a standard deviation of 1 prior to further analysis. The distance threshold of 130 m corresponds to the estimated range at which red squirrel territorial vocalizations are heard, i.e. the acoustic social neighbourhood [41], and also approximates the spatial scale at which red squirrels respond to variation in density [42].

Prior to exploring fitness effects, we used a generalized linear mixed model (GLMM) with Gamma family (log link) to test whether males and females differed in distance to their nearest primary kin. To avoid repeated inclusions of the same observation, we randomly sampled a single distance measure per dyad for this model. We also included a random effect to account for repeated measurements of an individual across multiple dyads.

(c) Estimating influences of kin on survival

We measured annual survival as the detection of an individual on or after 1 January of the following calendar year (note however that trapping typically began each year at approximately 1 March). Because red squirrels rarely change territory locations [43], individuals were assumed to have died if they disappeared from the study area and were never seen or trapped again. Previous work on this population has estimated the probability of re-detecting living adults to be 100% [44].

Prior to analysis, we subset the data to only include observations of the *nearest* parent, offspring, or full sibling for each "focal" squirrel in each year, resulting in a dataset with distances to different types of kin. This allowed us to directly test whether influences on fitness differed by kin relationship, while also focusing on the smallest distance values, expecting that any behaviourally mediated influences should be most apparent when squirrels were very close to one another.

First, we used generalized additive models (GAMs) to investigate possible nonlinearities in the relationships between fitness and

distance to specific types of primary kin. This was necessary given that influences of other squirrels could be negligible at very large distances, in which case a linear fit would not be justifiable for the full spatial extent of the data. We fitted GAMs with a binomial family and log link to relate annualized survival as the response variable. We used a basis dimension of 3 for each GAM, based on examination of the estimated degrees of freedom after fitting [45]. Crucially, we fitted an interaction between relationship type and linear distance. This allowed us to test whether the fitness-distance relationships were significantly different between different types of primary kin, compared to the reference category (non-kin). We also included mast year (yes/no) as a predictor, expecting that resource pulses could be a common cause influencing survival, and potentially, access to primary kin (i.e. environmental co-variance). Lastly, we included quadratic effects of age given the expectation that the probability of annual survival should decrease for both juveniles and senescent adults [23].

These preliminary GAMs indicated potentially nonlinear relationships between fitness and distance to specific types of primary kin. However, relationships within the previously defined 'social neighbourhood' in this population [23,29], i.e. up to a maximum distance between territories of 130 m, tended to be linear, with apparent changes in the relationship typically occurring at or beyond this point. Accordingly, we fitted GLMMs with linear effects of distance to the restricted, 'social neighbourhood' datasets, only including distance records less than or equal to 130 m. These models were fitted with binomial family and a logit link function and used the same response and predictor structure as above, explaining survival as a function of mast year, age (linear and quadratic terms) and an interaction between genealogical relationship and geographical distance. Here, we also included dyadic measures of familiarity (following the method for calculating familiarity from Siracusa *et al.* [23]) so that benefits of kin were estimated over and above any influences of familiarity-driven relationships. Though these within-neighbourhood (i.e. <130 m) influences of kin were most relevant for our research question, we also provide additional GLMMs in the electronic supplementary material that include distances spanning the full range of each study area as well as a quadratic effect of distance to account for apparent nonlinear associations at this scale.

All GLMMs were fitted with crossed random effects to account for the inclusion of multiple measures of the focal squirrel (focal identity (ID)) across years and of possibly bi-directional records of the same dyad of squirrels (dyad ID). We also included a random effect for year of the study period to account for any annual variation in fitness measures not captured by the mast-year fixed effect.

(d) Estimating influences of kin on reproductive success

Following previous work on this population [23], we calculated separate metrics of ARS for males and females to reflect differences in parental care. For females, we defined ARS as the number of pups that successfully recruited into the population (i.e., were confirmed alive for at least 200 days), excluding any non-breeding individuals. As with measures of annual survival, we first used GAMs with a Poisson family and log link to visualize any nonlinear influences of each type of primary kin on ARS. To test the sensitivity of our results to this exclusion, we also fitted identical GAMs including non-breeding females ($n = 181$), attributing them ARS values of 0. Because male red squirrels provide no obvious paternal care, ARS for males was measured as the number of pups sired per year.

As with the annual survival models, we identified possible nonlinearities across the full spatial extent of each study area, but linear relationships within the social neighbourhood. Here we used GLMMs with the Poisson family and log link function to explain reproductive success as a function of distance to primary




Figure 1. Spatial overview of kin relationships on the 'Kloo' study area (see the electronic supplementary material, figure S2 for overview of 'Sulphur' area). Lines linking certain individuals represent relationships among primary kin, with colours distinguishing dyads as a sibling pair or by the sex of the parent. Grey circles show the locations of middens or territory centres for all territories occupied each year. Mast years (resource pulses) occurred in 2005, 2010 and 2014.

kin, fitted with an interaction between distance and genealogical relationship. We also included dyadic familiarity, mast year, linear and quadratic effects of age, and crossed random intercepts representing variation across individuals, dyads and years.

The model with male reproductive success as the response variable showed moderate overdispersion, so was re-fitted using the negative binomial distribution. For all GLMMs, measures of distance were \log_{10} -transformed, reducing the use of very large distance values. No additional mean-centring or scaling was applied, so that main effect coefficients correspond to benefits of each type of primary kin having a separate territory at very close distances (1 m). Note though that such close distances do not occur biologically: in our study, the closest distance between two squirrels with distinct territories was 9.5 m.

Prior to analysing any data, we developed and publicly uploaded an analysis plan to avoid inadvertently hypothesizing after results were known (HARKing) [46]. This plan is available at <https://github.com/KluaneRedSquirrelProject/Analysis-Plans>. Some changes to our approach were made during analysis, such as the inclusion of dyadic familiarity and year-specific random effects. All analyses were conducted in R v. 4.0.5 [47]. GAMs were fitted using the *mgcv* package while GLMMs were fitted using *glmmTMB* [48,49]. For each GLMM, we used *DHARMa* to assess model diagnostics [50], and *MuMIn* to calculate conditional and marginal R^2 [51,52]. Detailed sample sizes including the number of observations, focal individuals, and unique dyads are available in the electronic supplementary material, tables S1–S9. Coding scripts are available at github.com/KluaneRedSquirrelProject/Walmsley-Projects.

3. Results

In total, our main analysis included observations of ARS and overwinter survival from 679 individuals between 2003 and 2014. Compared to females, male squirrels generally lived

farther from their nearest primary kin (95% confidence interval (CI) of predicted mean for males: 124–160 m, and for females: 90–111 m; $\beta = 0.34 \pm 0.08$ s.e., $p < 0.001$; electronic supplementary material, table S1 and figure S1). As expected, the individuals in our study had highly variable access to primary kin. In any given year, it was common for both female and male red squirrels not to have any close relatives in their social neighbourhood (mean 0.56 primary kin within 130 m, range 0–6). These spatial associations were also transient, with kin dyads remaining in the same social environment for just 1.4 years on average (range 1–4; for 130 m neighbourhood). Access to kin was not much improved when considering the larger spatial scale of each study area (approx. 40 ha), with each squirrel having a mean of 0.83 primary kin in the study area (range 0–7) in a given year, and with overlap among kin lasting 1.5 years on average (range 1–4). The number of kin relationships appeared to vary in tandem with changes in population density attributable to resource pulses (figure 1; electronic supplementary material, figure S2). Specifically, both density and the number of kin relationships tended to increase immediately after mast years, gradually declining thereafter.

(a) Influences of kin on survival

For females, annual survival was strongly associated with access to daughters. At the scale of the social neighbourhood, females whose nearest primary kin was a daughter were more likely to survive when that daughter was closer, as evidenced by the significantly more negative distance-survival relationship for daughters compared to the reference category, 'non-kin' ($\beta = -3.78 \pm 1.47$ s.e., $p = 0.010$; electronic supplementary material, table S2). Crucially, this effect waned near the threshold of the social neighbourhood (130 m), meaning that the survival benefit of having a




Figure 2. Lines represent predicted probabilities of female survival, estimated with GAMs while controlling for the influences of age (linear and quadratic) and resource pulses. Here, probabilities are shown for a focal individual of age 2 years in a non-mast year. Points represent raw data, so are not an exact reflection of the partial smooth. Shaded bands show 95% confidence intervals. The dotted vertical line corresponds to 130 m, a commonly used radius of red squirrels' social neighbourhoods. 'Non-kin' records are made up of individuals with no primary kin in the study area, for which distances were calculated to a randomly chosen individual.

daughter as nearest kin occurred only at short distances (e.g. less than approximately 40 m; figure 2). This effect was estimated while controlling for a quadratic effect of the age of the focal individual (i.e. in the present example, the mother), which, as expected, was also a significant predictor of survival ($\beta^2 = -0.10 \pm 0.04$ s.e., $p = 0.015$). The additional GLMM including distance measures greater than 130 m did not identify a significant effect of the proximity of daughters on female survival (electronic supplementary material, table S3), further supporting the idea that daughter-induced benefits are only present within the finer spatial scale of the social neighbourhood.

Adult males were less likely to survive during mast years compared to non-mast years ($\beta = -0.69 \pm 0.33$ s.e., $p = 0.036$). However, there were no significant differences in the distance-survival relationship for any types of primary kin, compared to the reference effect calculated for non-kin conspecifics (electronic supplementary material, table S4) indicating no effect of access to kin on male survival. Results were similar when including the entire study area, however the quadratic component of the daughter-distance interaction was statistically significant and negative ($\beta = -30.31 \pm 11.84$ s.e., $p = 0.010$; electronic supplementary material, table S5), suggesting that male survival increases with distance to daughters up to approximately 130 m, then decreases (electronic supplementary material, figure S5). For models of both male and female survival, effects of kin were identified while conditioning on familiarity between the focal and target

individuals, which itself had little influence on annual survival (electronic supplementary material, tables S2 and S4).

(b) Influences of kin on reproductive success

For females, ARS was strongly influenced by masting events, with breeding females producing approximately three times as many recruiting offspring during mast years versus non-mast years, on average ($\beta = 1.16 \pm 0.18$ s.e., $p < 0.001$). However, reproductive success was not associated with distance to any type of primary kin, relative to non-kin (electronic supplementary material, table S6 and figure S3). These results were similar for models including distances to kin across the entire study area (electronic supplementary material, table S7). The additional analysis including non-breeding individuals (attributing them zero pups recruited) revealed the same pattern whereby we found no evidence that distance to kin affected female reproductive success (electronic supplementary material, figure S4), so we proceeded to interpret values from GLMMs using ARS of breeders alone. As with female survival, we found little influence of dyadic familiarity on fitness ($\beta = 0.04 \pm 0.06$ s.e., $p = 0.471$).

By contrast, we found clear links between ARS and distance to kin for males. Outputs from the GAMs pointed to the possibility of sex-specific trends: males sharing their social neighbourhood with a father or son seemed to sire more pups when living closer to that individual (figure 3). Oppositely, distance to a mother or daughter's territory




Figure 3. Lines represent predicted probabilities of male annual reproductive success, estimated with GAMs, controlling for the influences of age (linear and quadratic) and resource pulses, with shaded bands representing 95% confidence intervals. Here, probabilities are shown for a focal individual of age 2 years in a non-mast year. Points represent raw data, so are not an exact reflection of the partial smooth. The dotted vertical line corresponds to 130 m, a commonly used radius of red squirrels' social neighbourhoods. Non-kin records are made up of individuals with no primary kin in the study area, and distances were calculated relative to a randomly chosen individual.

seemed to be positively associated with male reproductive success (i.e. implying a relative cost of living closer to female kin; figure 3). This general pattern was supported by the parametric outputs from the GLMM, though only effects of fathers and mothers were statistically significant in the social neighbourhood analysis. For males whose nearest primary kin was their father, those living closer to their father sired more pups, reflected in the more negative distance-ARS relationship for fathers compared to non-kin ($\beta = -5.07 \pm 2.26$ s.e., $p = 0.025$). It is worth noting that this effect was detected with a smaller sample size than some other kin-specific effects, as it was less common for males to have a father as their nearest primary kin than a mother or daughter, for example (figure 3). Oppositely, the GLMM suggested that among males whose nearest primary kin was their mother, those living *farther* from their mother were more successful ($\beta = 2.79 \pm 1.42$ s.e., $p = 0.049$), though this effect was smaller in magnitude than the benefits of access to fathers. As before, these effects were estimated while conditioning on mastng events and age (linear and quadratic terms), all of which also had statistically significant effects on male ARS (electronic supplementary material, table S8). Here we also detected a positive effect of dyadic familiarity on male reproductive success ($\beta = 0.12 \pm 0.05$ s.e., $p = 0.022$). Though distance to daughters did not significantly influence male ARS at the scale of the social neighbourhood, the effect of

daughters emerged as a significant influence on the number of pups sired in the model based on the entire study area (electronic supplementary material, table S9). This suggests that any influence of daughters on male reproductive success (negative but non-significant at the scale of the social neighbourhood; electronic supplementary material, table S8) probably changes after a certain distance threshold, and that nearby daughters may negatively impact their father's reproductive success (figure 3). Finally, we detected a significant effect of distance to non-kin (electronic supplementary material, table S9), suggesting that when considering a large spatial scale, males may sire more pups at higher densities overall, possibly a result of an increased availability of females. However, a similar effect was not detected when considering the social environment alone (i.e. the 130 m social neighbourhood; electronic supplementary material, table S8).

4. Discussion

In this study, we used the spatial distance between territories of solitary red squirrels to test whether access to kin influenced individual fitness. We found that red squirrels had improved survival and reproductive success when living near primary kin, though these effects were surprisingly nuanced, hinging

on genealogical relationship, sex and fitness measure. The dilution of fitness consequences from nearby kin across relatively short distances (0–130 m) is consistent with the hypothesis that these benefits are caused by kin-specific behavioural interactions, though we cannot rule out the possibility that other (unknown) factors also play a role. Together, our findings suggest complex kinship effects on survival and reproductive success in a solitary mammal, highlighting the value of multigenerational datasets for uncovering otherwise cryptic social dynamics.

The increased survival of mothers living closer to daughters converges with similar findings from other mammals, where philopatry facilitates further cooperative or tolerative interactions [11,53]. This includes a growing literature documenting female-biased kin benefits in less social mammalian species. For example, Eurasian lynx (*Lynx lynx*) mothers and daughters appear to share space when resources are patchy [28]. It is worth noting the magnitude of the effect of proximity to daughters in red squirrels. For example, among females whose nearest primary kin was a daughter, having that daughter 30 m away rather than 130 m away corresponded to a change in the mother's predicted probability of overwinter survival from approximately 37% to 82% (calculated for a 2-year old female in a non-mast year with average dyadic familiarity; electronic supplementary material, table S2). Interestingly, daughters did not appear to receive similar benefits from access to their mother. We also note the lack of any influences of nearby kin on ARS in females. Social thermoregulation is a possible mechanism driving these kin-based increases in survival, which is known to occur among females whose territories are close together (mean of 43.4 ± 18.0 m s.d. apart) [24]. However, we would generally expect the benefits of social thermoregulation to be reciprocal, i.e. also increase survival for daughters living near mothers [54]. Previous work has shown that territorial intrusions (possibly to pilfer cones) sometimes involve mothers invading offspring territories, and again is more likely to occur when distances between territories are small [29]. Thus, a seemingly more likely explanation for these unidirectional benefits is that mothers increase their survival by maintaining access to the resources of nearby daughters. It is important to note though that generally, pilfered cones appear to make up a very small proportion of an individual's food stores in this population [55]. An additional explanation for these patterns is that generally successful mothers are both more likely to recruit daughters nearby and to survive through the winter. However, the benefits of access to daughters were detected within the sample of mothers who had a daughter within 130 m—that daughters benefited mothers only when living very close makes it less likely that correlations between reproductive output and survival are the major driver of these patterns.

For male red squirrels, annual survival was not associated with access to kin, but was instead linked to environmental variation (i.e. masting events). This converges with evidence from a separate study which identified a similar, though non-significant, trend of reduced survival during mast years in this population [56]. One possibility is that this reduction in overwinter survival for males may be a cost of intense investment in reproduction: males expand their territory sizes nearly three-fold during mast years [33], possibly a result of increased efforts to find and mate with receptive females.

Much more surprising was the finding that male red squirrels had increased reproductive success when living closer to

their fathers. A typical male living 50 m from its father was predicted to sire two pups on average, but no pups if that father lived 130 m away (calculated for 2-year old during non-mast year with average dyadic familiarity). This was unexpected for two reasons: first, while red squirrels appear to discriminate kin and non-kin based on rattle vocalizations [57], it remains unclear whether this recognition stems from shared early life (natal) experience [58], which would limit recognition to maternal kin. This points to the possibility of another mechanism by which fathers and offspring can recognize one another, such as phenotypic matching [58,59], thought to facilitate kin recognition in other species (e.g. cichlids (e.g. *Pelvicachromis taeniatus*) [60]; Siberian jays (*Perisoreus infaustus*) [61]). Second, male red squirrels have been found to have enhanced lifetime reproductive success as immigrants to a new population [62], congruent with the expectation that they are avoiding competition with related males. In a similar vein, evidence in this study of reduced male reproductive success when closer to female kin (figure 3), could reflect the costs of reduced access to unrelated mates. However, there appears to be little inbreeding avoidance in this population: offspring from related parents are common ($r > 0.125$ for 19% of mates) and do not face reductions in birth mass, growth, or survival [63], suggesting that other factors may explain this pattern.

In some species, males form reproductive alliances to cooperate in securing and defending mating opportunities with females, leading to improved reproductive success [64–67]. Alliance members are often related, leading to indirect benefits for individuals that do not enjoy increased mating opportunities themselves. Much rarer associations between fathers and sons have been linked to reproductive benefits in red howler monkeys (*Alouatta seniculus*) [68] and yellow baboons (*Papio cynocephalus*) [69], for example. And while male-male alliances are typically found in group-living species, it has been suggested that rudimentary aspects of coalitionary behaviour may be exhibited in otherwise solitary males [70]. So, could the observed ARS benefits observed among father-son dyads in red squirrels be evidence of cryptic coalitions? Theory suggests that male reproductive coalitions are more likely to form when (i) females are receptive for short periods of time, and (ii) cannot be monopolized by a single mate [70,71]. Both are true of the 'scramble competition' mating system of red squirrels, where oestrus lasts just a single day, during which an oestrous female mates with an average of seven males [36,63]. However, male red squirrels are not known to engage in any observable affiliative interactions, making even basic aspects of coalitionary behaviour unlikely. Perhaps more likely is that fathers and sons are simply less likely to disrupt one another's copulations during mating chases. Hypothetically, related males living together could also benefit from socially transmitted information about the locations of receptive females. One avenue for future work would be to investigate where proximity between fathers and sons predicts co-parentage of offspring in the same litter (above and beyond any influences of distance to the mother), expected if male kin are somehow facilitating mating opportunities. Alternatively, increased tolerance among fathers and sons could reduce the need for defence and thereby allow males to exert more energy searching for mates.

More generally, these findings place an even greater emphasis on territory acquisition as a determinant of fitness for red squirrels [72], by increasing the stakes of co-locating with certain types of kin. Red squirrels are one of very few

species known to exhibit bequeathal, where an adult female undergoes breeding dispersal thus allowing an offspring (with a slight bias for daughters over sons) to inherit her territory and resource cache [21,73]. Such an extreme display of nepotism is striking, given the critical importance of territory ownership for survival. Interestingly then, previous work in this population has not identified a survival cost for bequeathing mothers [74]. Theory suggests that if a mother can secure a new territory near their previous site, increased access to offspring might lead to greater downstream fitness benefits via kin cooperation [75]. Consistent with this hypothesis, our findings suggest that bequeathal may provide additional, direct benefits to mothers in addition to enhancements in offspring survival [74], at least when the inheritor is a daughter. Compared to females, male red squirrels tend to live farther from primary kin, with no obvious pattern of father-son dyads living closer together than father–daughter dyads, for example (see the electronic supplementary material, figure S1). Thus we find no obvious indication of kin-biased dispersal, whereby males might be expected to preferentially acquire territories near male kin [67,76], though a thorough test of this possibility may be a worthwhile direction for future work.

Using distance among kin as a proxy for their ability to interact allowed us to directly measure fitness consequences without a clear sense of the specific behaviours that are important [31]. Beyond age and resource pulses (which were controlled for in our models), we are unaware of any likely confounds of proximity among kin and individual fitness in the population. Moreover, the importance of small differences in distance to kin (i.e. within the 130 m social neighbourhood) and the variation in effects across genealogical relationship are difficult to explain without distance-constrained social behaviours. However, future work is necessary to understand the contribution of behavioural interactions or other processes in causing these patterns.

Consistent with previous findings [23], male squirrels also had enhanced reproductive success when living near familiar individuals. Unlike previous work, however, we did not find significant effects of familiarity on survival for either males or females. We expect that this difference is a result of the dyadic approach in the present study—in other words, the benefits of familiarity may be small when considering a single target individual. This distinction highlights the separate spatial and social scales at which familiarity and kinship operate to influence red squirrel fitness. Whereas the benefits of familiarity emerge primarily as a property of a wider socio-spatial environment (the red squirrel neighbourhood) [23], the benefits of kinship occur via close proximity to a small set of specific individuals. Crucially, access to primary kin may not necessarily translate into variation in local genetic relatedness, raising questions about the use of local relatedness as a measure of possible kin-directed effects on fitness (reviewed in [9]).

The idiosyncratic nature of our results suggests that consequences of access to kin may be overlooked in systems where associations are not parsed into specific genealogical relationships. It also suggests that squirrels require kin recognition to maintain these relationships, contrary to many species where spatial heuristics (e.g. ‘cooperate with those around you’) are sufficient [77,78]. In tandem with previous studies, our work contributes to an increasingly complex depiction of sociality in a mostly solitary species. Though further work is needed to understand the specific mechanisms causing these benefits, it appears that kinship can have multifaceted and impactful contributions to fitness in solitary mammals.

Ethics. This research was approved by the University of Guelph Animal Care Committee (AUP number 1807).

Data accessibility. The data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.vt4b8gtwf> [79]. Analysis scripts can be found at <https://github.com/KluaneRedSquirrelProject/Walmsley-Projects>.

The data are provided in the electronic supplementary material [80].

Authors' contributions. S.F.W.: conceptualization, formal analysis, funding acquisition, investigation, methodology, visualization, writing—original draft, writing—review and editing; S.B.: data curation, funding acquisition, resources, writing—review and editing; B.D.: data curation, funding acquisition, resources, writing—review and editing; J.E.L.: data curation, funding acquisition, resources, writing—review and editing; D.W.C.: data curation, funding acquisition, resources, writing—review and editing; A.G.M.: conceptualization, data curation, funding acquisition, investigation, methodology, project administration, supervision, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This work was supported by Fulbright Canada (Special Foundation Fellowship to S.F.W.), the USA National Science Foundation (grant no. DEB-0515849 to A.G.M, IOS-1110436 to A.G.M. and B.D., IOS-1749627 to B.D.), the Natural Sciences and Engineering Research Council of Canada (Discovery Grants RGPIN-371579-2009 and RGPIN-2015-04707 to A.G.M, RGPIN-2020-06781 to J.E.L.; RGPIN-2018-04354 to D.W.C.; RGPIN-2014-05874 to S.B.) and the Ontario Ministry of Research and Innovation (grant no. ER08-05-119 to A.G.M.).

Acknowledgements. We thank Agnes MacDonald and her family for long-term access to her trapline, and the Champagne and Aishihik First Nations for allowing us to conduct our work within their traditional territory. We are grateful to each of the field technicians who have contributed to the long-term Kluane Red Squirrel Project database over the years. We thank Erin Siracusa and April Martinig for providing helpful advice regarding analysis and interpretation. We also thank members of the McAdam laboratory including Shelby Bohn, David Delaney, Simon Denomme-Brown, Alex Hare, Katie Kariatsumari and Quinn Webber for valuable and supportive commentary on an earlier version of this work. Lastly, we thank three anonymous reviewers for helpful comments that improved this work.

References

1. Silk JB. 2007 The adaptive value of sociality in mammalian groups. *Phil. Trans. R. Soc. B* **362**, 539–559. (doi:10.1098/rstb.2006.1994)
2. Mitani JC, Watts DP, Pepper JW, Merriwether DA. 2002 Demographic and social constraints on male chimpanzee behaviour. *Anim. Behav.* **64**, 727–737. (doi:10.1006/anbe.2002.4014)
3. Clutton-Brock TH, Lukas D. 2012 The evolution of social philopatry and dispersal in female mammals. *Mol. Ecol.* **21**, 472–492. (doi:10.1111/j.1365-294X.2011.05232.x)
4. Thompson ME. 2019 How can non-human primates inform evolutionary perspectives on female-biased kinship in humans? *Phil. Trans. R. Soc. B* **374**, 20180074. (doi:10.1098/rstb.2018.0074)
5. Holekamp KE, Smale L. 1991 Dominance acquisition during mammalian social development: the

'inheritance' of maternal rank. *Integr. Comp. Biol.* **31**, 306–317. (doi:10.1093/icb/31.2.306)

6. Turner JW, Robitaille AL, Bills PS, Holekamp KE. 2021 Early-life relationships matter: social position during early life predicts fitness among female spotted hyenas. *J. Anim. Ecol.* **90**, 183–196. (doi:10.1111/1365-2656.13282)

7. Ilany A, Holekamp KE, Akçay E. 2021 Rank-dependent social inheritance determines social network structure in spotted hyenas. *Science* **373**, 348–352. (doi:10.1126/science.abc1966)

8. Shah SS, Rubenstein DR. 2021 Prenatal environmental variation, alternative reproductive tactics and the formation of a mixed-kin cooperative society. *bioRxiv* (doi:10.1101/2021.02.10.430612)

9. Croft DP, Weiss MN, Nielsen MLK, Grimes C, Cant MA, Ellis S, Franks DW, Johnstone RA. 2021 Kinship dynamics: patterns and consequences of changes in local relatedness. *Proc. R. Soc. B* **288**, 20211129. (doi:10.1098/rspb.2021.1129)

10. Cantor M *et al.* 2021 The importance of individual-to-society feedbacks in animal ecology and evolution. *J. Anim. Ecol.* **90**, 27–44. (doi:10.1111/1365-2656.13336)

11. Holekamp KE, Sawdy MA. 2019 The evolution of matrilineal social systems in fissiped carnivores. *Phil. Trans. R. Soc. B* **374**, 20180065. (doi:10.1098/rstb.2018.0065)

12. Wolf JBW, Trillmich F. 2008 Kin in space: social viscosity in a spatially and genetically substructured network. *Proc. R. Soc. B* **275**, 2063–2069. (doi:10.1098/rspb.2008.0356)

13. Griffiths SW, Armstrong JD. 2002 Kin-biased territory overlap and food sharing among Atlantic salmon juveniles. *J. Anim. Ecol.* **71**, 480–486. (doi:10.1046/j.1365-2656.2002.00614.x)

14. Koprowski JL. 1993 The role of kinship in field interactions among juvenile gray squirrels (*Sciurus carolinensis*). *Can. J. Zool.* **71**, 224–226. (doi:10.1139/z93-031)

15. Carter GG, Farine DR, Wilkinson GS. 2017 Social bet-hedging in vampire bats. *Biol. Lett.* **13**, 1–4. (doi:10.1098/rsbl.2017.0112)

16. Silk JB, Altmann J, Alberts SC. 2006 Social relationships among adult female baboons (*Papio cynocephalus*) I. Variation in the strength of social bonds. *Behav. Ecol. Sociobiol.* **61**, 183–195. (doi:10.1007/s00265-006-0249-2)

17. Engh AL, Beehner JC, Bergman TJ, Whitten PL, Hoffmeier RR, Seyfarth RM, Cheney DL. 2006 Behavioural and hormonal responses to predation in female chacma baboons (*Papio hamadryas ursinus*). *Proc. R. Soc. B* **273**, 707–712. (doi:10.1098/rspb.2005.3378)

18. Chakrabarti S, Kolipakam V, Bump JK, Jhala YV. 2020 The role of kinship and demography in shaping cooperation amongst male lions. *Sci. Rep.* **10**, 1–13. (doi:10.1038/s41598-020-74247-x)

19. Moor DD, Roos C, Ostner J, Schülke O. 2020 Bonds of bros and brothers: kinship and social bonding in postdispersal male macaques. *Mol. Ecol.* **29**, 3346–3360. (doi:10.1111/mec.15560)

20. West SA, Cooper GA, Ghoul MB, Griffin AS. 2021 Ten recent insights for our understanding of cooperation. *Nat. Ecol. Evol.* **5**, 419–430. (doi:10.1038/s41559-020-01384-x)

21. Bertiaux D, Boutin S. 2000 Breeding dispersal in female North American red squirrels. *Ecology* **81**, 1311–1326. (doi:10.1890/0012-9658(2000)081[1311:BDIFNA]2.0.CO;2)

22. Larsen KW, Boutin S. 1994 Movements, survival, and settlement of red squirrel (*Tamiasciurus hudsonicus*). *Ecology* **75**, 214–223. (doi:10.2307/1939395)

23. Siracusa E, Boutin SA, Dantzer B, Lane JE, Coltman DW, McAdam AG. 2021 Familiar neighbors, but not relatives, enhance fitness in a territorial mammal. *Curr. Biol.* **31**, 438–445. (doi:10.1016/j.cub.2020.10.072)

24. Williams CT, Gorrell JC, Lane JE, McAdam AG, Humphries MM, Boutin S. 2013 Communal nesting in an 'asocial' mammal: social thermoregulation among spatially dispersed kin. *Behav. Ecol. Sociobiol.* **67**, 757–763. (doi:10.1007/s00265-013-1499-4)

25. Gorrell JC, McAdam AG, Coltman DW, Humphries MM, Boutin S. 2010 Adopting kin enhances inclusive fitness in asocial red squirrels. *Nat. Commun.* **1**, 22. (doi:10.1038/ncomms1022)

26. Wright BM, Stredulinsky EH, Ellis GM, Ford JKB. 2016 Kin-directed food sharing promotes lifetime natal philopatry of both sexes in a population of fish-eating killer whales, *Orcinus orca*. *Anim. Behav.* **115**, 81–95. (doi:10.1016/j.anbehav.2016.02.025)

27. Emigh H, Truax J, Highfill L, Vonk J. 2020 Not by the same token: a female orangutan (*Pongo pygmaeus*) is selectively prosocial. *Primates* **61**, 237–247. (doi:10.1007/s10329-019-00780-7)

28. Aronsson M, Åkesson M, Low M, Persson J, Andrén H. 2020 Resource dispersion and relatedness interact to explain space use in a solitary predator. *Oikos* **129**, 1174–1184. (doi:10.1111/oik.07258)

29. Siracusa E, Boutin S, Humphries MM, Gorrell JC, Coltman DW, Dantzer B, Lane JE, McAdam AG. 2017 Familiarity with neighbours affects intrusion risk in territorial red squirrels. *Anim. Behav.* **133**, 11–20. (doi:10.1016/j.anbehav.2017.08.024)

30. Siracusa E, Morandini M, Boutin S, Humphries MM, Dantzer B, Lane JE, McAdam AG. 2017 Red squirrel territorial vocalizations deter intrusions by conspecific rivals. *Behaviour* **154**, 1259–1273. (doi:10.1163/1568539X-00003467)

31. Engelhardt SC, Bergeron P, Gagnon A, Dillon L, Pelletier F. 2019 Using geographic distance as a potential proxy for help in the assessment of the grandmother hypothesis. *Curr. Biol.* **29**, 651–656.e3. (doi:10.1016/j.cub.2019.01.027)

32. Ellis S, Franks DW, Weiss MN, Cant MA, Domenici P, Balcomb KC, Ellifrit DK, Croft DP. 2021 Mixture models as a method for comparative sociality: social networks and demographic change in resident killer whales. *Behav. Ecol. Sociobiol.* **75**, 438–445. (doi:10.1007/s00265-021-03006-3)

33. Webber QMR, Dantzer B, Lane JE, Boutin S, McAdam AG. 2022 Density-dependent plasticity in territoriality revealed using social network analysis. *J. Anim. Ecol.* (doi:10.1111/1365-2656.13846)

34. LaMontagne JM, Williams CT, Donald JL, Humphries MM, McAdam AG, Boutin S. 2013 Linking intraspecific variation in territory size, cone supply, and survival of North American red squirrels. *J. Mammal.* **94**, 1048–1058. (doi:10.1644/12-MAMM-A-245.1)

35. Kalinowski ST, Taper ML, Marshall TC. 2007 Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.* **16**, 1099–1106. (doi:10.1111/j.1365-294X.2007.03089.x)

36. Lane JE, Boutin S, Gunn MR, Coltman DW. 2009 Sexually selected behaviour: red squirrel males search for reproductive success. *J. Anim. Ecol.* **78**, 296–304. (doi:10.1111/j.1365-2656.2008.01502.x)

37. McAdam AG, Boutin S, Sykes AK, Humphries MM. 2007 Life histories of female red squirrels and their contributions to population growth and lifetime fitness. *Ecoscience* **14**, 362. (doi:10.2980/1195-6860(2007)14[362:LHFRS]2.0.CO;2)

38. Dantzer B, McAdam AG, Humphries MM, Lane JE, Boutin S. 2020 Decoupling the effects of food and density on life-history plasticity of wild animals using field experiments: insights from the steward who sits in the shadow of its tail, the North American red squirrel. *J. Anim. Ecol.* **89**, 2397–2414. (doi:10.1111/1365-2656.13341)

39. Morrissey MB, Wilson AJ. 2010 PEDANTICS: an R package for pedigree-based genetic simulation and pedigree manipulation, characterization and viewing. *Mol. Ecol. Resour.* **10**, 711–719. (doi:10.1111/j.1755-0998.2009.02817.x)

40. Lane JE, Boutin S, Gunn MR, Slate J, Coltman DW. 2008 Female multiple mating and paternity in free-ranging North American red squirrels. *Anim. Behav.* **75**, 1927–1937. (doi:10.1016/j.anbehav.2007.10.038)

41. Smith CC. 1978 Structure and function of the vocalizations of tree squirrels (*Tamiasciurus*). *J. Mammal.* **59**, 793–808. (doi:10.2307/1380144)

42. Dantzer B, Boutin S, Humphries MM, McAdam AG. 2012 Behavioral responses of territorial red squirrels to natural and experimental variation in population density. *Behav. Ecol. Sociobiol.* **66**, 865–878. (doi:10.1007/s00265-012-1335-2)

43. Larsen KW, Boutin S. 1995 Exploring territory quality in the North American red squirrel through removal experiments. *Can. J. Zool.* **73**, 1115–1122. (doi:10.1139/z95-133)

44. Descamps S, Boutin S, McAdam AG, Berteaux D, Gaillard JM. 2009 Survival costs of reproduction vary with age in North American red squirrels. *Proc. R. Soc. B* **276**, 1129–1135. (doi:10.1098/rspb.2008.1401)

45. Wood SN. 2017 *Generalized additive models: an introduction with R*, 2nd edn. CRC/Taylor & Francis, New York, New York, USA.

46. Kerr NL. 1998 HARKing: hypothesizing after the results are known. *Pers. Soc. Psychol. Rev.* **2**, 196–217. (doi:10.1207/s15327957pspr0203_4)

47. R Core Development Team. 2021 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.

48. Wood SN. 2011 Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J. R. Stat. Soc. B* **73**, 3–36. (doi:10.1111/j.1467-9868.2010.00749.x)

49. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Mächler M, Bolker BM. 2017 glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R J.* **9**, 378–400. (doi:10.32614/rj-2017-066)

50. Hartig F. 2018 DHARMA: residual diagnostics for hierarchical (multi-level/mixed) regression models. See <https://CRAN.R-project.org/package=DHARMA>.

51. Barton K, Barton MK. 2020 MuMln: Multi-Model Inference. R package version 1.43.17. Version 1, 18. See <https://CRAN.R-project.org/package=MuMln>.

52. Nakagawa S, Johnson PCD, Schielzeth H. 2017 The coefficient of determination R^2 and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *J. R. Soc. Interface* **14**, 20170213. (doi:10.1098/rsif.2017.0213)

53. Ashbury AM, Willems EP, Suci S, Atmoko U, Saputra F, Schaik CPV. 2020 Home range establishment and the mechanisms of philopatry among female Bornean orangutans (*Pongo pygmaeus wurmbii*) at Tuanan. *Behav. Ecol. Sociobiol.* **74**, 1–21.

54. Lutermann H, Schmelting B, Radespiel U. 2006 The role of survival for the evolution of female philopatry in a solitary forager, the grey mouse lemur (*Microcebus murinus*). *Proc. R. Soc. B* **273**, 2527–2533. (doi:10.1098/rspb.2006.3603)

55. Donald JL, Boutin S. 2011 Intraspecific cache pilferage by larder-hoarding red squirrels (*Tamiasciurus hudsonicus*). *J. Mammal.* **92**, 1013–1020. (doi:10.1644/10-MAMM-A-340.1)

56. McAdam AG, Webber QMR, Dantzer B, Lane JE, Boutin S. 2022 Social effects on annual fitness in red squirrels. *J. Hered.* **113**, 69–78. (doi:10.1093/jhered/esab051)

57. Wilson DR, Goble AR, Boutin S, Humphries MM, Coltman DW, Gorrell JC, Shonfield J, McAdam AG. 2015 Red squirrels use territorial vocalizations for kin discrimination. *Anim. Behav.* **107**, 79–85. (doi:10.1016/j.anbehav.2015.06.011)

58. Shonfield J, Gorrell JC, Coltman DW, Boutin S, Humphries MM, Wilson DR, McAdam AG. 2017 Using playback of territorial calls to investigate mechanisms of kin discrimination in red squirrels. *Behav. Ecol.* **28**, 382–390. (doi:10.1093/beheco/awr165)

59. Widdig A. 2007 Paternal kin discrimination: the evidence and likely mechanisms. *Biol. Rev.* **82**, 319–334. (doi:10.1111/j.1469-185X.2007.00011.x)

60. Hesse S, Bakker TCM, Baldauf SA, Thünken T. 2012 Kin recognition by phenotype matching is family- rather than self-referential in juvenile cichlid fish. *Anim. Behav.* **84**, 451–457. (doi:10.1016/j.anbehav.2012.05.021)

61. Griesser M, Halvarsson P, Drobniak SM, Vilà C. 2015 Fine-scale kin recognition in the absence of social familiarity in the Siberian jay, a monogamous bird species. *Mol. Ecol.* **24**, 5726–5738. (doi:10.1111/mec.13420)

62. Martinig AR, McAdam AG, Dantzer B, Lane JE, Coltman DW, Boutin S. 2020 The new kid on the block: immigrant males win big whereas females pay fitness cost after dispersal. *Ecol. Lett.* **23**, 430–438. (doi:10.1111/ele.13436)

63. Lane JE, Boutin S, Gunn MR, Slate J, Coltman DW. 2007 Genetic relatedness of mates does not predict patterns of parentage in North American red squirrels. *Anim. Behav.* **74**, 611–619. (doi:10.1016/j.anbehav.2006.12.017)

64. Parsons KM, Durban JW, Claridge DE, Balcomb KC, Noble LR, Thompson PM. 2003 Kinship as a basis for alliance formation between male bottlenose dolphins, *Tursiops truncatus*, in the Bahamas. *Anim. Behav.* **66**, 185–194. (doi:10.1006/anbe.2003.2186)

65. Wiszniewski J, Corrigan S, Beheregaray LB, Mo LM. 2012 Male reproductive success increases with alliance size in Indo-Pacific bottlenose dolphins (*Tursiops aduncus*). *J. Anim. Ecol.* **81**, 423–431. (doi:10.1111/j.1365-2656.2011.01910.x)

66. Ryder TB, Blake JG, Parker PG, Loiselle BA. 2011 The composition, stability, and kinship of reproductive coalitions in a lekking bird. *Behav. Ecol.* **22**, 282–290. (doi:10.1093/beheco/qrq213)

67. Williams DA, Rabenold KN. 2005 Male-biased dispersal, female philopatry, and routes to fitness in a social corvid. *J. Anim. Ecol.* **74**, 150–159. (doi:10.1111/j.1365-2656.2004.00907.x)

68. Pope TR. 1990 The reproductive consequences of male cooperation in the red howler monkey: paternity exclusion in multi-male and single-male troops using genetic markers. *Behav. Ecol. Sociobiol.* **27**, 439–446. (doi:10.1007/BF00164071)

69. Charpentier MJE, Van Horn RC, Altmann J, Alberts SC. 2008 Paternal effects on offspring fitness in a multimale primate society. *Proc. Natl Acad. Sci. USA* **105**, 1988–1992. (doi:10.1073/pnas.0711219105)

70. Olson LE, Blumstein DT. 2009 A trait-based approach to understand the evolution of complex coalitions in male mammals. *Behav. Ecol.* **20**, 624–632. (doi:10.1093/beheco/arp040)

71. Teichroeb JA, Wikberg EC, Ting N, Sicotte P. 2014 Factors influencing male affiliation and coalitions in a species with male dispersal and intense male-male competition, *Colobus vellerosus*. *Behaviour* **151**, 1045–1066. (doi:10.1163/1568539X-00003089)

72. Hendrix JG, Fisher DN, Martinig AR, Boutin S, Dantzer B, Lane JE, McAdam AG. 2020 Territory acquisition mediates the influence of predators and climate on juvenile red squirrel survival. *J. Anim. Ecol.* **89**, 1408–1418. (doi:10.1111/1365-2656.13209)

73. Fisher DN, Boutin S, Dantzer B, Humphries MM, Lane JE, McAdam AG. 2017 Multilevel and sex-specific selection on competitive traits in North American red squirrels. *Evolution* **71**, 1841–1854. (doi:10.1111/evo.13270)

74. Lane JE, McAdam AG, Charmantier A, Humphries MM, Coltman DW, Fletcher Q, Gorrell JC, Boutin S. 2015 Post-weaning parental care increases fitness but is not heritable in North American red squirrels. *J. Evol. Biol.* **28**, 1203–1212. (doi:10.1111/jeb.12633)

75. Clarke PMR, McElreath MB, Barrett BJ, Mabry KE, McElreath R. 2018 The evolution of bequeathal in stable habitats. *Ecol. Evol.* **8**, 10594–10607. (doi:10.1002/ece3.4549)

76. Pollack L, Rubenstein DR. 2015 The fitness consequences of kin-biased dispersal in a cooperatively breeding bird. *Biol. Lett.* **11**, 2–5. (doi:10.1098/rsbl.2015.0336)

77. Langer P, Nowak MA, Hauert C. 2008 Spatial invasion of cooperation. *J. Theor. Biol.* **250**, 634–641. (doi:10.1016/j.jtbi.2007.11.002)

78. Rand DG, Nowak MA. 2013 Human cooperation. *Trends Cogn. Sci.* **17**, 413–425. (doi:10.1016/j.tics.2013.06.003)

79. Walmsley SF, Boutin S, Dantzer B, Lane JE, Coltman DW, McAdam AG. 2023 Data from: Benefits of living closer to kin vary by genealogical relationship in a territorial mammal. Dryad Digital Repository. (doi:10.5061/dryad.vt4b8gtwf)

80. Walmsley SF, Boutin S, Dantzer B, Lane JE, Coltman DW, McAdam AG. 2023 Benefits of living closer to kin vary by genealogical relationship in a territorial mammal. Figshare. (doi:10.6084/m9.figshare.c.6350103)