

**Seven replacement names for bees in the genus  
Protandrena Cockerell, 1896 (Hymenoptera: Andrenidae:  
Panurginae)**

Authors: Bossert, Silas, and Neff, John L.

Source: The Pan-Pacific Entomologist, 98(3) : 226-229

Published By: Pacific Coast Entomological Society

URL: <https://doi.org/10.3956/2022-98.3.226>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Scientific Note

**Seven replacement names for bees in the genus *Protandrena*  
Cockerell, 1896 (Hymenoptera: Andrenidae: Panurginae)**

The Protandrenini is a New World tribe of 436 species of andrenid bees (Ascher & Pickering 2022, Bossert et al. 2022). The North American protandrenines form a single clade (Bossert et al. 2022, Ramos et al. 2022) of 186 species (Ascher & Pickering 2022). Over the past 170 years, the North American protandrenine bees have been placed in many different genera, including *Panurginus* Nylander, 1848 and *Psaenythia* Gerstacker, 1868 (Michener 1944). More recently they have been considered to consist of between two (Ascher & Pickering 2022, Michener 1944, Michener, 1951, Mitchell 1960), three (Michener 2000, 2007), five (Ramos et al. 2022), six (Michener et al. 1994, Timberlake 1973), or seven (Hurd 1973) genera, sometimes including South American taxa (Michener 1944, 2000, 2007; Mitchell 1960). A recent molecular analysis (Bossert et al. 2022) has shown that none of these classifications adequately reflects the phylogeny of the group, all being paraphyletic.

Pending further research, we believe the best solution for now is to recognize a single genus (*Protandrena*) with five subgenera: *Anthemurgus* Robertson, 1902, *Heterosarus* Robertson, 1918, *Metapsaenythia* Timberlake, 1969, *Pseudopanurgus* Cockerell, 1897, *Pterosarus* Timberlake, 1967 and *Protandrena* (s. str.). A possible taxonomic alternative would be to recognize *Pseudopanurgus* along with *Anthemurgus* and the currently recognized subgenera of *Protandrena* (*sensu* Michener 2007) as genera. Unfortunately, however, it is not at all obvious that either *Heterosarus* (see Ramos 2011) or *Pterosarus* (see Ramos 2011, Timberlake 1967) is monophyletic. Far more extensive sampling is necessary to resolve the relationships of the North American protandrenines. In the absence of such a study, the one genus solution is most adequate to retain genera that correspond to natural groups. This results in seven cases of secondary homonymy for which we provide replacement names below. Only one of these species is relatively common, the rest being known from only one sex, often just the holotype.

## SYSTEMATICS

Tribe Protandrenini Robertson, 1904

Genus *Protandrena* Cockerell, 1896

Subgenus *Heterosarus* Robertson, 1918

*Protandrena* (*Heterosarus*) *chamaesycephila*, **nom. nov.**

*Heterosarus euphorbiae* Timberlake, 1975:24, *nomen praeoccupatum* (junior secondary homonym), *nec* *Psaenythia euphorbiae* Timberlake, 1955 = *Protandrena* (*Protandrena*) *euphorbiae* (Timberlake, 1955).

**Etymology:** The name refers to *Chamaesyce*, a subgenus of the huge genus *Euphorbia* L. (Euphorbiaceae) whose species are the primary pollen hosts of this oligolectic bee.

**Comments:** This is a common bee known from both males and females and occurring from southern Arizona to Chiapas, Mexico (Timberlake 1975).

Subgenus *Protandrena*

*Protandrena (Protandrena) impunctata*, **nom. nov.**

*Protandrena pernitens* Timberlake, 1976:193, *nomen praeoccupatum* (junior secondary homonym), nec *Heterosarus pernitens* Timberlake, 1975 = *Protandrena (Heterosarus) pernitens* (Timberlake, 1975).

Etymology: The name refers to the nearly impunctate male metasoma.

Comments: Known only from males from Morelos and Oaxaca, Mexico (Timberlake 1976).

*Protandrena (Protandrena) zacateca*, **nom. nov.**

*Protandrena fasciata* Timberlake, 1976:164 *nomen praeoccupatum* (junior secondary homonym), nec *Pseudopanurgus fasciatus* Timberlake, 1973 = *Protandrena (Pseudopanurgus) fasciata* (Timberlake, 1975)

Etymology: The name refers to the locale of the type female.

Comments: Known only from the type female from Zacatecas, Mexico (Timberlake 1976).

*Protandrena (Protandrena) pseudomexicanorum*, **nom. nov.**

*Protandrena durangoensis* Timberlake, 1976:159, *nomen praeoccupatum* (junior secondary homonym), nec *Heterosarus durangoensis* Timberlake, 1975 = *Protandrena (Heterosarus) durangoensis* (Timberlake, 1975)

Etymology: The name follows the suggestion of Timberlake (1976) that the species is very closely related to *P. (Protandrena) mexicanorum* (Cockerell).

Comments: Known only from the type female from Durango, Mexico (Timberlake 1976).

*Protandrena (Protandrena) rozeni*, **nom. nov.**

*Protandrena lateralis* Timberlake, 1976: 177, *nomen praeoccupatum* (junior secondary homonym), nec *Pseudopanurgus lateralis* Timberlake, 1975 = *Protandrena (Protandrena) lateralis* (Timberlake, 1975)

Etymology: The species is named for Jerome Rozen, Jr., whose numerous studies have greatly expanded what we know of the nests and larval morphology of the North American Protandrenini (Rozen 1967, 1970, 1989).

Comments: Known only from the type male from Durango, Mexico (Timberlake 1976).

*Protandrena (Protandrena) santacruzana*, **nom. nov.**

*Protandrena amplipennis* Timberlake, 1976:146, *nomen praeoccupatum* (junior secondary homonym), nec *Heterosarus amplipennis* Timberlake, 1975 = *Protandrena (Heterosarus) amplipennis* (Timberlake, 1975)

Etymology: The species is named for Santa Cruz County, Arizona, the locale of the type female.

Comments: Known only from the type female (Timberlake 1976).

*Protandrena (Protandrena) tauformis*, **nom. nov.**

*Protandrena semilevis* Timberlake, 1976:203, *nomen praeoccupatum* (junior secondary homonym), nec *Pseudopanurgus semilevis* Timberlake, 1973 = *Protandrena (Pseudopanurgus) semilevis* (Timberlake, 1973)

Etymology: The name refers to a t-shaped mark on the female clypeus.

Comments: Known from a series of females from Guerrero, Morelos and Oaxaca, Mexico (Timberlake 1976).

*Protandrena (Protandrena) maurula* (Cockerell, 1896)

*Protandrena texana* Timberlake 1976: 40, **syn. nov.**, nec *Pseudopanurgus texanus* Timberlake, 1973 = *Protandrena (Pseudopanurgus) texana* (Timberlake, 1973).

Comments: Only males have been described for *P. (Protandrena) texana* while only females have been described for *P. (Protandrena) maurula*. We have seen a series of male and female *Protandrena* collected by Sam Droege at one site in Kent County, Texas on 22 April 2016. The females of this series are identical with the description of *P. (Protandrena) maurula* while the males agree with *P. (Protandrena) texana* supporting the synonymy of *P. (Protandrena) texana* under *P. (Protandrena) maurula*.

*Acknowledgments.* We thank two anonymous reviewers for catching problems in an early version of the manuscript.

Silas Bossert, *Department of Entomology, Washington State University, Pullman, Washington 99164, U.S.A., e-mail: silas.bossert@wsu.edu; and John L. Neff (corresponding author), Central Texas Melittological Institute, 7307 Running Rope, Austin, Texas 78731, U.S.A., e-mail: jlnatctmi@yahoo.com.*

#### LITERATURE CITED

- Ascher, J. S. & J. Pickering. 2022. Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). Available from <http://www.discoverlife.org/mp/20q?guide=Apoidea> species (accessed 10 July 2022).
- Bossert, S., T. J. Wood, S. Patiny, D. Michez, E. A. B. Almeida, R. L. Minckley, L. Packer, J. L. Neff, R. S. Copeland, J. Straka, A. Pauly, T. L. Griswold, S. G. Brady, B. N. Danforth & E. A. Murray. 2022. Phylogeny, biogeography, and rapid diversification of the mining bee family Andrenidae. *Systematic Entomology* 47:283–302 <https://doi.org/10.1111/syen.12530>
- Hurd, P. D., Jr. 1979. Superfamily Apoidea, pp 1043–1255. In: K. V. Krombein, D. R. Smith & B. D. Burks (Eds.), *Hymenoptera in America North of Mexico—Synoptic Catalog*. U.S. Department of Agriculture Monograph No. 2, Washington, D.C., xvi + 2209 pp.
- Michener, C. D. 1944. Comparative external morphology, phylogeny, and a classification of the bees (Hymenoptera). *Bulletin of the American Museum of Natural History* 82(6):151–326.
- Michener, C. D. 1951. Superfamily Apoidea, pp 1741–2209. In: C. F. W. Muesebeck & K. V. Krombein (Eds.), *Catalog of Hymenoptera in America North of Mexico, Vol. 2*. Smithsonian Institution Press, Washington, D.C., 1420 pp.
- Michener, C. D. 2000. *The Bees of the World*. The Johns Hopkins University Press, Baltimore, xiv + 913 pp.
- Michener, C. D. 2007. *The Bees of the World, Second Edition*. The Johns Hopkins University Press, Baltimore, xvi + 953 pp.
- Mitchell, T. B. 1960. Bees of the Eastern United States, Volume I. *North Carolina Agricultural Experiment Station Technical Bulletin* 141:1–538.
- Michener, C. D., T. J. McGinley & B. N. Danforth. 1994. *The Bee Genera of North and Central America (Hymenoptera: Apoidea)*. Smithsonian Institution Press, Washington, D.C., viii + 209 pp.
- Ramos, K. D. S. 2011. Relações filogenéticas entre as abelhas da subfamília Andreninae como enfoque nas tribos Calliopsini, Protandrenini e Protomelitturgini (Hymenoptera, Apidae). Ph. D. dissertation, Departamento de Zoologia, Programa de Pós-Graduação em Entomologia, Universidade Federal do Paraná, Curitiba, Brasil, 156 pp.
- Ramos, K. D. S., A. C. Martins & G. A. R. Melo. 2022. Evolution of andrenine bees reveals a long and complex history of faunal interchanges through the Americas during the Mesozoic and Cenozoic. *Molecular Phylogenetics and Evolution* 172(2):107484 <https://doi.org/10.1111/syen.107484>
- Rozen, J. G., Jr. 1967. Review of the biology of panurgine bees, with observations on North American forms (Hymenoptera: Andrenidae). *American Museum Novitates* 2297:1–44.

- Rozen, J. G., Jr. 1970. Biology and immature stages of the panurgine bee genera *Hypomacrotera* and *Psaenythia* (Hymenoptera, Apoidea). *American Museum Novitates* 2416:1–16.
- Rozen, J. G., Jr. 1989. Life history studies of the “primitive” panurgine bees (Hymenoptera: Andrenidae: Panurginae). *American Museum Novitates* 2962:1–27.
- Timberlake, P. H. 1973. Revision of the genus *Pseudopanurgus* of North America (Hymenoptera, Apoidea). *University of California Publications in Entomology* 72:1–58.
- Timberlake, P. H. 1975. The North American species of *Heterosarus* Robertson (Hymenoptera, Apoidea). *University of California Publications in Entomology* 77:1–56.
- Timberlake, P. H. 1976. Revision of the North American bees of the genus *Protandrena* Cockerell (Hymenoptera: Apoidea). *Transactions of the American Entomological Society* 102(2): 133–227.
- Timberlake, P. H. 1977. Descriptions of two new species of *Heterosarus* Robertson (Hymenoptera, Apoidea). *The Pan-Pacific Entomologist* 53 (1):56–59.

*Received 23 May 2022; accepted 14 July 2022. Publication date 16 September 2022*

*Subject editor Michael C. Orr*