

Title: H atom collisions with a semiconductor efficiently promote electrons to the conduction band

Author list: Kerstin Krüger¹, Yingqi Wang², Sophia Tödter¹, Felix Debbeler¹, Anna Matveenko¹, Nils Hertl³, Xueyao Zhou⁴, Bin Jiang⁴, Hua Guo², Alec M. Wodtke^{1,3,5}, Oliver Bünermann^{1,5,*}

Affiliations:

¹Institute of Physical Chemistry, Georg-August University, Göttingen, Germany

²Department of Chemistry and Chemical Biology, University of New Mexico, New Mexico, USA

³Department of Dynamics at Surfaces, Max-Planck-Institute for Multidisciplinary Sciences, Göttingen, Germany

⁴Hefei National Laboratory for Physical Science at the Microscale, Department of Chemical Physics, University of Science and Technology of China, Hefei, Anhui, China

⁵International Center of Advanced Studies of Energy Conversion, Georg-August University, Göttingen, Germany

*Corresponding author. Email: oliver.buenermann@chemie.uni-goettingen.de

Abstract

The Born-Oppenheimer Approximation (BOA) is the keystone of modern computational chemistry and there is wide interest in understanding under what conditions it remains valid. H atom scattering from insulator, semi-metal and metal surfaces has helped provide such information. For insulators, the BOA is adequate. For metals, the BOA fails but not severely. Here, we present H atom scattering from a semiconductor – Ge(111)c(2×8). Experiments reveal bimodal energy-loss distributions reflecting two channels. Molecular dynamics (MD) trajectories within the BOA reproduce one channel quantitatively. The second channel transfers much more energy and is absent in MD simulations. It grows with H atom incidence energy and exhibits an energy-loss onset equal to the Ge surface bandgap. This leads us to conclude that H atom collisions at the surface of a semiconductor are capable of promoting electrons from the valence to the conduction band with high efficiency. Our current understanding fails to explain these observations.

Main text

Atoms and molecules colliding at solid surfaces create time-varying electric fields that, due to their finite masses and associated low speeds, represent frequencies typically $\leq 10^{13}$ Hz, whereas much lighter electrons in solids oscillate at frequencies one to two orders of magnitude higher than this. This separation of time-scales is used to justify the Born-Oppenheimer approximation (BOA)¹, the bedrock of computational surface chemistry², where electronic quantum states rapidly adjust to the motion of nuclei. Inelastic H atom surface scattering experiments have provided excellent benchmarks against which theoretical methods can and have been tested and proven³. Using this approach, the BOA has been shown to be justified for H atom scattering from Xe, where molecular dynamics simulations using a full-dimensional potential energy surface quantitatively reproduced energy-losses measured in high-resolution scattering experiments⁴. The validity of the BOA in this case is not surprising since the lowest energy electronic excitations in Xe exceeded the energies of that work. Similar energy-loss measurements from experiments scattering H and D from the semi-metal graphene, where low energy electron-hole pair (EHP) excitations are possible, also showed no signs of BOA failure⁵⁻⁷. Despite these successes, there are reasons to question the BOA's validity^{8,9}. For example, energetic H atoms colliding at metal surfaces always excite EHPs^{10,11}; albeit, theory could successfully treat this with a weak-coupling "electronic friction" approximation^{12,13} suggesting BOA failure is not severe and can be accounted for in a perturbative fashion.

Experiments with semiconductors present an opportunity to make predictions from our current understanding about a fundamentally different class of solids. This is true if semiconductors behave in some hybrid fashion, reflecting some intermediate to insulators and metals. But let us consider semiconductors from the point of view of another kind of time varying electric field. We know visible light with electric fields oscillating at $\sim 10^{14-15}$ Hz efficiently excites electrons from the valence band (VB) to the conduction band (CB), forming the basis for a large fraction of optical science and technology. This raises the question: if collisions of atoms and molecules with semiconductors could produce time varying electric fields oscillating at similar frequencies, would they not also excite VB electrons to the CB and might this not provide important new avenues of research with promise of new technology? If we were to adopt the physical picture derived from our study of metals, where electronic friction describes BOA failure, the answer to this question would certainly be "no" or more precisely "only weakly" as electronic friction theories lead to hot EHP distributions that still favor low energy excitation near the Fermi level¹². Unfortunately, scattering experiments with semiconductors that test the validity of the BOA are rare. Transient currents were observed

when Xe atoms with energies between 3 and 10 eV were scattered from surfaces of semiconductors¹⁴⁻¹⁶; however, this resulted from the creation of a local hot spot where initial phonon excitation subsequently transferred energy to EHPs. While these experiments provide us clear evidence of BOA failure in a semiconductor, we can gain only little insight into the dynamics of the atom-surface collision. In fact, an electronically adiabatic model could describe the energy-loss of scattered Xe atoms.

In this work, we produce H atoms whose speeds are high enough to test the limits of the BOA directly by investigating the characteristics of their collisions with a semiconductor surface. The measured H atom energy-loss spectra and angular distributions reveal the excitations appearing in the solid on the sub-ps time scale. We find that, not only is VB-CB excitation possible, at sufficiently high energies it dominates the energy transfer dynamics, showing that new physical mechanisms are at play. Specifically, we present translational energy-loss measurements on energetic H atoms scattered from a reconstructed Ge(111)*c*(2×8) surface along with first principles electronically adiabatic molecular dynamics (MD) simulations, performed with a newly developed high-dimensional neural network potential energy surface (NN-PES). When incidence energies are below the bandgap, only one scattering channel arises with small energy-losses nearly identical to those seen in the MD simulations. These exhibit collision dynamics similar to those seen in H scattering from Xe. Surprisingly, at higher incidence energies, a second channel appears whose energy-loss onset is coincident with the semiconductor bandgap. This channel is absent in the MD simulations with and without electronic friction. The importance of this channel increases rapidly with H atom velocity — a signature of BOA failure — and accounts for ~90% probability at the highest H atom incidence energies of this work.

Results

Fig. 1 shows experimental translational energy-loss distributions for H atoms scattered from Ge(111)*c*(2×8)¹⁷ at incidence energies E_i above and below the 0.49 eV surface bandgap¹⁸. We note that the given value for the surface band gap was determined at a surface temperature of 30 K. However, a similar value is expected at room temperature since the reconstruction of the surface is unchanged. Also shown are the predictions of the electronically adiabatic MD trajectory calculations. Below the bandgap (Fig. 1A) only a single feature appears in the energy-loss distribution. The MD simulations reproduce the experimental result extremely well. Molecular dynamics with electronic friction¹⁹ at the level of local density friction approximation (LDFA)²⁰ fail to describe the energy loss distributions—see Fig. E1. Analysis of adiabatic MD trajectories shows that H atoms interact with the Ge surface for only a few fs

and that energy exchange is limited. Figs. 1B-D show energy-loss distributions for three values of E_i larger than the surface bandgap—in all three cases, the distributions are bimodal and the MD trajectories reproduce only the feature seen at low values of energy-loss. Hereafter, we refer to this feature as the adiabatic channel. The second feature appearing at higher energy-losses is absent in the adiabatic MD simulations, strongly suggesting that this channel involves conversion of H atom translational energy to electronic excitation of the Ge solid. This idea is further supported by the observation that the energy-loss onset of this feature is coincident (within experimental uncertainty) with the Ge surface bandgap of 0.49 eV at all values of E_i . Furthermore, as expected for a channel involving BOA failure, this channel is strongly promoted by incidence translational energy, becoming about 90% of the observed scattering at the highest value of $E_i = 6.17$ eV. For these reasons, we assign the high energy-loss feature to an electronically non-adiabatic process where the collision of the H atom at the surface promotes an electron above the bandgap of the Ge surface. We refer to this mechanism hereafter as the VB-CB channel.

Fig. 2 shows differential properties from both experiment and theory for H atoms incident at three angles ϑ_i and at $E_i = 0.99$ eV. Here, polar plots display the final translational energy E_f as a function of final scattering angle ϑ_f . A white dashed line shows the expected minimal energy loss for excitation of an electron across the surface band gap, which demarcates the adiabatic from the VB-CB channel. Experiment shows that the VB-CB channel exhibits a much narrower angular distribution (Table 1) than the adiabatic channel at all three incidence angles. The MD simulations yield similar differential scattering maps as seen in experiment for the adiabatic channel only. The energy loss agrees with experiment and even the experimentally observed dependency of the angular distribution on ϑ_i is reproduced. The VB-CB channel is absent in the MD simulations.

Fig. 3 shows polar plot representations similar to Fig. 2 emphasizing the incidence energy dependence of the scattering. As before, the experimental results show bimodal scattering distributions with two well-resolved channels separated in energy space by the bandgap energy, marked as a white dashed line. The angular distributions of both channels broaden between $E_i = 0.99$ eV and 1.92 eV; but the VB-CB channel broadens significantly more as it is narrower at $E_i = 0.99$ eV (Table 1). The adiabatic MD simulations (Fig. 3C & 3D) reproduce this effect for the adiabatic channel.

The average energy losses derived from the experiments are summarized in Table 2. Notice that for the adiabatic channel, the average energy transferred to the surface $\langle E_i - E_f \rangle$ is a small and nearly constant fraction ($10 \pm 5\%$) of E_i . The VB-CB channel behaves differently—the fraction of incidence energy transferred to the solid goes up dramatically as E_i is reduced. This is an influence of the surface bandgap, where the absolute of energy lost must exceed 0.49 eV, regardless of E_i . Hence at lower values of E_i , the fractional energy loss must sharply increase. Also notice that the average energy lost decreases only slightly with increasing ϑ_i for both channels.

Discussion

We start by highlighting some of the key observations just presented and their implications. First, Fig. 2 shows clearly that the most probable value of ϑ_f depends on the chosen value of ϑ_i , proving the scattered atoms did not thermalize with the solid—thermalization occurs on the ps timescale. Thus, we conclude that the scattered atoms in both channels experience a sub-picosecond interaction time with the surface. Second, there is evidence of sticking, despite the fact that integrated scattering probabilities like sticking probabilities cannot be easily obtained from in-plane differential scattering measurements, since the fraction of incident atoms that scatter out of the detection plane may also depend on incidence conditions and branching channel. We can nevertheless integrate the observed scattering at flux over E_f and ϑ_f . These integrals scaled to the experimentally observed adiabatic channel at $E_i = 0.99$ eV and $\vartheta_i = 45^\circ$ appear as bold numbers next to each differential scattering diagram in Figs. 2 & 3. They are given as ratios that report the relative contributions of the two scattering channels. There is an overall loss of signal between $E_i = 1.92$ eV and 0.99 eV. If we were to assume the out-of-plane scattering fraction were independent of E_i , we would conclude that the sticking probability decreases with increasing incidence energy. A similar trend is seen in the MD simulations. Also notice that the branching ratios shown in Figs. 3A & B are consistent with those of Figs. 1B & C, which represent the branching between the two scattering channels detected at $\vartheta_f = 45^\circ$ only. This agreement suggests that the branching seen in Fig. 1C ($E_i = 1.92$ eV) is representative of other scattering angles.

The major outcome of this work is the observation that an H atom scattering from a semiconductor may experience one or the other of two types of interactions, either a mechanical interaction well described within the BOA or a strong non-adiabatic interaction capable of promoting an electron to energies above the bandgap. We emphasize that while there are similarities with past work, the behavior seen here is qualitatively different from previous observations involving insulators, metals or semi-metals. For example, the adiabatic channel

seen in Figs. 1-3 exhibits marked similarities to H atom scattering from insulating Xe; however, that system exhibited no BOA failure whatsoever. Conversely, H scattering trajectories describing collisions with metals simultaneously excite both phonons and EHPs¹⁰⁻¹³, the two excitations being inextricably linked to one another. The question remains, what gives rise to the branching between the two channels in the H/Ge system?

The fact that H scattering from Ge exhibits branching behavior between two distinct dynamical channels is consistent with a two-state picture. We envision that the H atom proceeds initially along the ground electronic state until it encounters a seam of crossing associated with a short lived electronically excited state. (Note that the word state is used here loosely as many electronic states are involved in the VB and CB of the system.) We assume that this state rapidly decays into unoccupied electronic states within the CB. At low incidence energies, reaching the seam of crossing requires specific approach, but at higher energies, other regions of the seam become accessible with reduced steric restrictions.

Evidence supporting this picture can be found in observations of this work, especially Fig. 2. Notice that the VB-CB channel exhibits a narrow angular distribution, peaking near the specular scattering angle (arrows in Fig. 2). This shows that there is no preference for loss of incidence energy parallel or perpendicular to the surface when inducing electronic excitation. A narrow angular distribution is typical of scattering influenced by directional forces associated with atomic orbitals with preferred orientations, consistent with the suggested mechanism of a curve crossing, where H atom collisions must occur at specific surface sites (Ge atoms) and with specific approach geometries. Fig. 3 shows that at higher energy these steric restrictions appear to be less severe; consequently, the VB-CB scattering angular distribution broadens.

Contrasting this behavior, the adiabatic channel exhibits a markedly broader angular distribution even at low incidence energy. This indicates a large corrugation of the potential energy surface experienced by the atoms passing through the adiabatic channel. Despite the many final scattering angles, the energy loss follows a hard-sphere line-of-centers (LOC) binary collision model (black dashed line). This indicates that the H atom scattered through the adiabatic channel is experiencing binary collisions with many impact parameters, not surprising due to the complex surface structure of Ge(111)c(2×8) surface if the H atoms scattering through the adiabatic channel sample a large fraction of the surface unit cell.

Bimodal energy-loss distributions may be produced without electronic excitation. For example, H scattering from a graphene layer involves trajectories that either fail or succeed in surmounting a chemisorption barrier⁵⁻⁷. H atoms reflected from the barrier experience weak van der Waals interactions with little energy transferred, while H atoms surmounting the barrier

couple strongly to in-plane phonons of the graphene layer ⁵. In contrast to this behavior, the electronically adiabatic MD simulations carried out in this work show no sign of bimodal distributions, consistent with the absence of a chemisorption barrier in the H/Ge system. The combined strength of the experimental and theoretical results supports our assignment of an electronically adiabatic and a non-adiabatic channel.

While it is common knowledge that absorption of photons in the bulk of a semiconductor excites electrons from the VB to the CB, this work shows that a colliding atom may efficiently promote electrons in a similar way in a purely surface specific process. The probability to convert translational energy of the H atom to electronic excitation of the solid dramatically increases with incidence energy, as does the average excitation energy. The large excitation probability as well as the large energy-loss is inconsistent with electronic friction theories; hence, this work stands as a challenge for new theories of electronically non-adiabatic surface chemistry. We hasten to add that the designation of this behavior as VB-CB represents a simplified viewpoint. The precise nature of the excited electronic states involved is still unknown – transient surface localized excitations (even plasmons) might be important. Nevertheless, the observation that electronic excitation dominates the dynamics in collisions of a simple atom with a semiconductor opens new horizons for research into non-adiabatic effects in surface chemistry and chemical sensors.

Acknowledgments:

We thank Daniel Auerbach and Alexander Kandratsenka for helpful discussions. A.M.W. thanks the Max Planck Society for the advancement of science. O.B. and A.M.W. acknowledge support from the Deutsche Forschungsgemeinschaft (DFG) under SFB 1073 (217133147), project A04 and from the DFG, the Ministerium für Wissenschaft und Kultur, Niedersachsen and the Volkswagenstiftung under grant no. INST 186/902-1. A.M.W. and H.G. acknowledge support from the Alexander von Humboldt Foundation. H.G. acknowledges support from the National Science Foundation under grant No. CHE-1951328. B.J. acknowledges support from the National Natural Science Foundation of China under grant no. 22073089 and from the K. C. Wong Education Foundation under grant no. GJTD-2020-15.

Author Contributions:

A.M.W. and O.B. conceived the project. O.B. supervised the experiment, H.G. and B.J. the theoretical simulations. K.K. performed the experiments, S.T., F.D. and A.M. assisted in part

of the experiments. Y.W., X.Z., B.J. and H.G. constructed the NN-PES. Y.W. performed the MD simulations. K.K. and Y.W. analyzed the data. K.K. and O.B. wrote the manuscript with feedback from A.M.W., Y.W., N.H., B.J. and H.G.

Competing Interests: The authors declare no competing interests.

Tables:

Table 1: Angular full width at half maximum for the experimental angular distributions of this work.

E_i	VB – CB			Adiabatic		
	$\vartheta_i = 30^\circ$	$\vartheta_i = 45^\circ$	$\vartheta_i = 60^\circ$	$\vartheta_i = 30^\circ$	$\vartheta_i = 45^\circ$	$\vartheta_i = 60^\circ$
0.99 eV	24°	31°	24°	> 56°	44°	34°
1.92 eV	–	> 70°	–	–	> 73°	–

Table 2: Average energy-loss in experimentally obtained specular ($\vartheta_i = \vartheta_f$) H atom scattering. Values in parentheses were computed from adiabatic MD trajectories.

	ϑ_i	VB – CB		Adiabatic	
		$\langle E_i - E_f \rangle$	$\frac{\langle E_i - E_f \rangle}{E_i} \times 100$	$\langle E_i - E_f \rangle$	$\frac{\langle E_i - E_f \rangle}{E_i} \times 100$
0.37 eV	45°	–	–	0.05 eV	14% (13%)
0.99 eV	30°	0.75 eV	75%	0.15 eV	15% (17%)
	45°	0.71 eV	72%	0.13 eV	13% (12%)
	60°	0.69 eV	70%	0.10 eV	10% (8.1%)
1.92 eV	45°	1.12 eV	58%	0.20 eV	10% (14%)
6.17 eV	45°	2.28 eV	37%	0.32 eV	5.2% (7.7%)

Figure Legends/Captions (for main text figures):

Fig. 1. Translational energy-loss distributions for H atoms scattered from Ge(111)c(2×8). The incident H atoms travel along the $[\bar{1}10]$ surface direction, while the polar incidence and scattering angles ϑ_i and ϑ_f , respectively, were both 45° with respect to the surface normal. The surface temperature T_s was 300 K. Experimental data (+) and the results of adiabatic molecular dynamics simulations (solid lines) for four H atom translational incidence energies are shown: $E_i = 0.37$ eV (A), 0.99 eV (B), 1.92 eV (C), and 6.17 eV (D). The band gap of the surface is 0.49 eV and indicated by the vertical dashed line. The experimentally obtained ratio of the adiabatic to the VB-CB channel appears in each panel. All experimental curves are normalized to the peak intensity. The MD curves are scaled to fit the adiabatic channel.

Fig. 2. Incidence angle dependence of H atoms scattered from Ge(111)c(2×8). Energy resolved angular distributions derived from in-plane scattering flux are shown for three incidence angles, $\vartheta_i = 30^\circ, 45^\circ, 60^\circ$ and an incidence translational energy $E_i = 0.99$ eV. The surface temperature was $T_s = 300$ K. Experimental results (A to C) are compared to MD simulations (D to F). The adiabatic and the VB-CB channels both exhibit maximum scattering flux near the specular scattering angle (arrows). The MD simulations reproduce the behavior of the adiabatic channel only. To construct the experimental plots, data was recorded in 5° increments from $\vartheta_f = 0^\circ$ to 75° . All six polar plots are normalized to the incident H atom flux. Bold numbers show the ratios of the experimentally observed scattering channels with respect to the adiabatic channel for an incidence angle of $\vartheta_i = 45^\circ$, the left one corresponds to the VB-CB channel, the right one to the adiabatic channel. The MD simulations are scaled to experiment such that at an incidence angle of $\vartheta_i = 45^\circ$, the integrated adiabatic channels are equal in both. The black dashed lines represent the final energy predicted by a line-of-centers binary collision model: $E_f =$

$E_i \{1 - \cos^2[(\vartheta_i + \vartheta_f)/2] \times [1 - (m_H - m_{Ge})^2/(m_H + m_{Ge})^2]\}$. The white dashed lines indicates the surface band gap of 0.49 eV.

Fig. 3. Incidence energy dependence of H atoms scattered from Ge(111)c(2×8). Energy resolved angular distributions derived from in-plane scattering flux are shown for two incidence translational energies $E_i = 0.99$ eV (A & C) and 1.92 eV (B & D). The surface temperature was $T_S = 300$ K and the incidence angle is $\vartheta_i = 45^\circ$. Experimental results (A & B) are compared to MD simulations (C & D). The MD simulations reproduce the behavior of the adiabatic channel only. To construct the experimental plots, data was recorded in 5° increments from $\vartheta_f = 0^\circ$ to 75° . All four polar plots are normalized to the incident H atom flux. Bold numbers show the ratios of the experimentally observed scattering channels with respect to the adiabatic channel for an incidence energy of $E_i = 0.99$ eV, the left one corresponds to the VB-CB channel, the right one to the adiabatic channel. The MD simulations are scaled to experiment such that at an incidence energy of $E_i = 0.99$ eV, the integrated adiabatic channels are equal in both. The black dashed lines represent the final energy predicted by a line-of-centers binary collision model: $E_f = E_i \{1 - \cos^2[(\vartheta_i + \vartheta_f)/2] \times [1 - (m_H - m_{Ge})^2/(m_H + m_{Ge})^2]\}$. The white dashed lines indicates the surface band gap of 0.49 eV.

References

- 1 Born, M. & Oppenheimer, R. Zur Quantentheorie der Molekeln. *Ann. Phys* **84**, 0457-0484, (1927).
- 2 Tully, J. C. Perspective on "Zur Quantentheorie der Molekeln" - Born M, Oppenheimer R (1927) *Ann Phys* **84** : 457. *Theor. Chem. Acc.* **103**, 173-176, (2000).
- 3 Bünermann, O., Kandratsenka, A. & Wodtke, A. M. Inelastic Scattering of H Atoms from Surfaces. *J. Phys. Chem. A* **125**, 3059-3076, (2021).
- 4 Hertl, N., Kandratsenka, A., Bünermann, O. & Wodtke, A. M. Multibounce and Subsurface Scattering of H Atoms Colliding with a van der Waals Solid. *J. Phys. Chem. A* **125**, 5745-5752, (2021).
- 5 Jiang, H. Y. *et al.* Imaging covalent bond formation by H atom scattering from graphene. *Science* **364**, 379-382, (2019).
- 6 Jiang, H. Y. *et al.* Small Nuclear Quantum Effects in Scattering of H and D from Graphene. *J. Phys. Chem. Lett.* **12**, 1991-1996, (2021).
- 7 Wille, S. *et al.* An experimentally validated neural-network potential energy surface for H-atom on free-standing graphene in full dimensionality. *Phys. Chem. Chem. Phys.* **22**, 26113-26120, (2020).
- 8 Tully, J. C. Chemical dynamics at metal surfaces. *Annu. Rev. Phys. Chem.* **51**, 153-178, (2000).
- 9 Wodtke, A. M., Tully, J. C. & Auerbach, D. J. Electronically non-adiabatic interactions of molecules at metal surfaces: Can we trust the Born-Oppenheimer approximation for surface chemistry? *Int. Rev. Phys. Chem.* **23**, 513-539, (2004).
- 10 Bünermann, O. *et al.* Electron-hole pair excitation determines the mechanism of hydrogen atom adsorption. *Science* **350**, 1346-1349, (2015).
- 11 Jiang, H. Y., Dorenkamp, Y., Krüger, K. & Bünermann, O. Inelastic H and D atom scattering from Au(111) as benchmark for theory. *J. Chem. Phys.* **150**, 184105, (2019).
- 12 Kandratsenka, A. *et al.* Unified description of H-atom-induced chemicurrents and inelastic scattering. *PNAS* **115**, 680-684, (2018).
- 13 Dorenkamp, Y. *et al.* Hydrogen collisions with transition metal surfaces: Universal electronically nonadiabatic adsorption. *J. Chem. Phys.* **148**, 034706, (2018).
- 14 Amirav, A. & Cardillo, M. J. Electron-Hole Pair Creation by Atomic Scattering at Surfaces. *Phys. Rev. Lett.* **57**, 2299-2302, (1986).
- 15 Amirav, A. *et al.* Electron-Hole Pair Creation at a Ge(100) Surface by Ground-State Neutral Xe Atoms. *J. Appl. Phys.* **59**, 2213-2215, (1986).
- 16 Weiss, P. S., Amirav, A., Trevor, P. L. & Cardillo, M. J. Hyperthermal Gas-Surface Scattering. *J. Vac. Sci. Technol., A* **6**, 889-894, (1988).
- 17 Chadi, D. J. & Chiang, C. New c -2x8 Unit-Cell for the Ge(111) Surface. *Phys. Rev. B* **23**, 1843-1846, (1981).
- 18 Feenstra, R. M., Lee, J. Y., Kang, M. H., Meyer, G. & Rieder, K. H. Band gap of the Ge(111)c(2x8) surface by scanning tunneling spectroscopy. *Phys. Rev. B* **73**, 035310, (2006).
- 19 Head-Gordon, M. & Tully, J. C. Molecular Dynamics with Electronic Frictions. *J. Chem. Phys.* **103**, 10137-10145, (1995).

- 20 Juaristi, J. I., Alducin, M., Muino, R. D., Busnengo, H. F. & Salin, A. Role of electron-hole pair excitations in the dissociative adsorption of diatomic molecules on metal surfaces. *Phys. Rev. Lett.* **100**, 116102, (2008).

Methods

The experimental setup is described in detail in refs. ^{3,21}. Briefly, UV ($\lambda_{\text{photolysis}} = 248.35 \text{ nm}$) or VUV ($\lambda_{\text{photolysis}} = 121.4 \text{ nm}$) photodissociation of a supersonic molecular beam of hydrogen iodide produces a H atom beam with translational energies of $E_i = 0.37, 0.99, 1.92$ or 6.17 eV that then passes two differential pumping chambers to enter an UHV scattering chamber before colliding with a Germanium crystal. The Ge sample is held on a 5-axis manipulator allowing the variation of the polar incidence angle ϑ_i with respect to the surface normal. The scattered H atoms are excited to a long-lived Rydberg state just below the ionization limit ²² and fly 250 mm before they are field-ionized and detected by a multichannel plate (MCP) assembly. A multichannel scaler records the arrival time to obtain the time-of-flight (TOF) distributions, which we convert to energy spectra applying the appropriate Jacobians. The detector is rotatable in the plane defined by the incident H atom beam and the surface normal allowing TOF distributions to be obtained at various final scattering angles ϑ_f . The used Ge crystal is undoped with a purity of 99.999%. The Ge(111) surface was cleaned with cycles of Ar^+ ion sputtering and annealing to $\sim 670^\circ\text{C}$. Auger electron spectroscopy (AES) and low-energy electron diffraction (LEED) validated the cleanliness and $c(2\times 8)$ structure of the surface.

To perform theoretical simulations, a neural-network potential (NN-PES) was constructed for the $\text{H@Ge}(111)c(2\times 8)$ system and molecular dynamics (MD) simulations were performed. Data for the NN fitting were obtained with spin-polarized DFT calculations, carried out with the Vienna Ab initio Simulation Package (VASP) ^{23,24} with the frozen-core all-electron projector-augmented wave (PAW) method ^{25,26}. The electronic wave function was expanded using plane waves with an energy cutoff of 250 eV. The electron exchange-correlation energies were described by the Perdew–Burke–Ernzerhof (PBE) functional within the generalized gradient approximation (GGA) ²⁷. The reconstructed $\text{Ge}(111)c(2\times 8)$ surface was modeled by repeated slabs separated by a vacuum space of 16 Å in z direction. Each slab contains eight atomic layers, with four additional Ge adatoms added on top of the first layer. The Ge atoms in the bottom layer not seen by the scattering H atoms in the MD simulations were capped by Ge-H bonds. The Ge adatoms and top six layers were allowed to move while the remaining atoms were fixed throughout the calculations and there are thus a total of 101 movable atoms in the unit cell. The Brillouin zone was sampled with a $3\times 1\times 1$ k -point grid. AIMD trajectories were used to provide training data for the NN fitting. The AIMD trajectories employed initial positions of the H atom randomly sampled 6 Å above the

surface. About 100 AIMD trajectories were run for an H atom incidence energy of 0.99 eV and 1.92 eV each, with an incidence angle of 45°, and a surface temperature of 300 K, providing ~150,000 points. Additional single point DFT calculations were performed to augment the AIMD points. The data set was culled using an Euclidean distance of 0.3 Å to remove points that were too close to one another. About 26,000 points (including both energy and gradient) were finally selected to fit a 303-dimensional PES using an embedded atom neural network (EANN) approach²⁸. The EANN PES obtained in this way was thoroughly tested, giving a root mean square error (RMSE) of about 80 meV/cell (or 0.8 meV/atom). MD trajectories were calculated with a modified Venus program²⁹. The timesteps were chosen separately for each incidence energy, 0.1, 0.05, 0.03 and 0.01 fs for 0.37, 0.99, 1.92 and 6.17 eV, respectively.

To study possible non-adiabatic effects, an electron friction model was applied^{19,30}. The electronic friction coefficient was calculated based on the local-density friction approximation (LDFA)^{20,31}. The electron density of the Ge(111)c(2x8) surface was obtained from about 100 configurations at 300 K. To obtain an analytical expression for the electron density the data was again fitted with the EANN method.

Data Availability

Source data are provided with this paper. Data are also available from the corresponding author upon reasonable request.

Code Availability

The EANN code of B.J. is available at <https://github.com/zhangylch/REANN>. The VENUS code is available at <https://www.depts.ttu.edu/chemistry/Venus/index.php>

Methods-only references

- 3 Bünemann, O., Kandratenka, A. & Wodtke, A. M. Inelastic Scattering of H Atoms from Surfaces. *J. Phys. Chem. A* **125**, 3059-3076, (2021).
- 19 Head-Gordon, M. & Tully, J. C. Molecular Dynamics with Electronic Frictions. *J. Chem. Phys.* **103**, 10137-10145, (1995).
- 20 Juaristi, J. I., Alducin, M., Muino, R. D., Busnengo, H. F. & Salin, A. Role of electron-hole pair excitations in the dissociative adsorption of diatomic molecules on metal surfaces. *Phys. Rev. Lett.* **100**, 116102, (2008).
- 21 Bünemann, O., Jiang, H. Y., Dorenkamp, Y., Auerbach, D. J. & Wodtke, A. M. An ultrahigh vacuum apparatus for H atom scattering from surfaces. *Rev. Sci. Instrum.* **89**, 094101, (2018).
- 22 Schnieder, L., Seekampprahn, K., Liedeker, F., Steuwe, H. & Welge, K. H. Hydrogen-Exchange Reaction H + D₂ in Crossed Beams. *Faraday Discuss.* **91**, 259-269, (1991).

- 23 Kresse, G. & Furthmüller, J. Efficient iterative schemes for ab initio total-energy calculations using a plane-wave basis set. *Phys. Rev. B* **54**, 11169-11186, (1996).
- 24 Kresse, G. & Furthmüller, J. Efficiency of ab-initio total energy calculations for metals and semiconductors using a plane-wave basis set. *Comput. Mater. Sci.* **6**, 15-50, (1996).
- 5 25 Blöchl, P. E. Projector Augmented-Wave Method. *Phys. Rev. B* **50**, 17953-17979, (1994).
- 26 Kresse, G. & Joubert, D. From ultrasoft pseudopotentials to the projector augmented-wave method. *Phys. Rev. B* **59**, 1758-1775, (1999).
- 27 Perdew, J. P., Burke, K. & Ernzerhof, M. Generalized gradient approximation made simple. *Phys. Rev. Lett.* **77**, 3865-3868, (1996).
- 10 28 Zhang, Y. L., Hu, C. & Jiang, B. Embedded Atom Neural Network Potentials: Efficient and Accurate Machine Learning with a Physically Inspired Representation. *J. Phys. Chem. Lett.* **10**, 4962-4967, (2019).
- 29 Hase, W. L. *et al.* VENUS96: A general chemical dynamics computer program. *QCPE Bulletin* **16**, 671, (1996).
- 30 Li, Y. G. & Wahnström, G. Nonadiabatic Effects in Hydrogen Diffusion in Metals. *Phys. Rev. Lett.* **68**, 3444-3447, (1992).
- 15 31 Puska, M. J. & Nieminen, R. M. Atoms Embedded in an Electron-Gas - Phase-Shifts and Cross-Sections. *Phys. Rev. B* **27**, 6121-6128, (1983).