

Invariant Measures for Horospherical Actions and Anosov Groups

Minju Lee¹ and Hee Oh^{1,2,*}

¹Mathematics Department, Yale University, New Haven, CT 06520, USA
and ²Korea Institute for Advanced Study, Seoul, 02455, Republic of Korea

**Correspondence to be sent to: e-mail: hee.oh@yale.edu*

Let Γ be a Zariski dense Anosov subgroup of a connected semisimple real algebraic group G . For a maximal horospherical subgroup N of G , we show that the space of all non-trivial NM -invariant ergodic and A -quasi-invariant Radon measures on $\Gamma \backslash G$, up to proportionality, is homeomorphic to $\mathbb{R}^{\text{rank } G-1}$, where A is a maximal real split torus and M is a maximal compact subgroup that normalizes N . One of the main ingredients is to establish the NM -ergodicity of all Burger–Roblin measures.

1 Introduction

Let G be a connected semisimple real algebraic group and $\Gamma < G$ a Zariski dense discrete subgroup. A subgroup N of G is called *horospherical* if there exists a diagonalizable element $a \in G$ such that

$$N = \{g \in G : a^k g a^{-k} \rightarrow \infty \quad \text{as } k \rightarrow +\infty\},$$

or equivalently, N is the unipotent radical of a parabolic subgroup of G . We assume that N is a maximal horospherical subgroup, which exists uniquely up to conjugation. We are interested in the measure rigidity property of the N -action on the homogeneous space $\Gamma \backslash G$. When Γ is a lattice, that is, when $\Gamma \backslash G$ has finite volume, the well-known

Received June 2, 2021; Revised May 23, 2022; Accepted September 2, 2022

measure rigidity theorems of Furstenberg [17], Veech [57], and Dani [14] give a complete classification of Radon measures (=locally finite Borel measures) invariant by N . This rigidity phenomenon extends to any unipotent subgroup action by the celebrated theorem of Ratner in [45].

When G has rank one and Γ is geometrically finite, the horospherical subgroup action on $\Gamma \backslash G$ is known to be essentially uniquely ergodic; there exists a unique non-trivial invariant ergodic Radon measure on $\Gamma \backslash G$, called the Burger–Roblin measure ([10], [46], [59]). When Γ is geometrically infinite, there may be a continuous family of horospherically invariant ergodic measures as first discovered by Babillot and Ledrappier ([3], [4]). For a certain class of geometrically infinite groups, a complete classification of horospherically invariant ergodic measures has been obtained; see [50], [51], [32], [36], [28], [29], etc. We refer to a recent article by Landesberg and Lindenstrauss [28] for a more precise description on the rank one case.

When G has rank at least 2 and Γ has infinite co-volume in G , very little is known about invariant measures. In this paper, we focus on a special class of discrete subgroups, called *Anosov subgroups*. In the rank one case, this class coincides with the class of convex cocompact subgroups, and hence the class of Anosov subgroups can be considered as a generalization of convex cocompact subgroups of rank one groups to higher rank. The works of Burger [11] and Quint [40] on a higher rank version of the Patterson–Sullivan theory supply a continuous family of maximal horospherically invariant Burger–Roblin measures, as was introduced in [15]. We show that all of these Burger–Roblin measures are ergodic for maximal horospherical foliations, and classify all ergodic *non-trivial* Radon measures for maximal horospherical foliations, which are also quasi-invariant under Weyl chamber flow. In particular, we establish a homeomorphism between the space of these measures and the interior of the projective limit cone of Γ , which is again homeomorphic to $\mathbb{R}^{\text{rank } G-1}$.

In order to formulate our main result precisely, we begin with the definition of an Anosov subgroup of G . Let P be the normalizer of N , that is, a minimal parabolic subgroup of G and $\mathcal{F} := G/P$ the Furstenberg boundary. We denote by $\mathcal{F}^{(2)}$ the unique open G -orbit in $\mathcal{F} \times \mathcal{F}$. A Zariski dense discrete subgroup $\Gamma < G$ is called an *Anosov subgroup* (with respect to P) if it is a finitely generated word hyperbolic group which admits a Γ -equivariant embedding ζ of the Gromov boundary $\partial\Gamma$ into \mathcal{F} such that $(\zeta(x), \zeta(y)) \in \mathcal{F}^{(2)}$ for all $x \neq y$ in $\partial\Gamma$.

First introduced by Labourie [27] as the images of Hitchin representations of surface groups ([23], [18]), this definition is due to Guichard and Wienhard [22], who showed that Anosov subgroups (more precisely, Anosov representations) form an open

subset in the representation variety $\text{Hom}(\Gamma, G)$. The class of Anosov groups includes Schottky subgroups [41], and hence any Zariski dense discrete subgroup of G contains an Anosov subgroup ([6], [42]). We also refer to the work of Kapovich, Leeb, and Porti [25] for other equivalent characterizations of Anosov groups, as well as to the excellent survey articles by Kassel [26] and Wienhard [58] on higher Teichmüller theory.

We let $P = NMA$ be the Langlands decomposition of P , so that N is the unipotent radical of P , A is a maximal real split torus of G , and M is a compact subgroup that commutes with A . Note that any maximal horospherical subgroup arises in this way, that is, as the unipotent radical of a minimal parabolic subgroup.

The limit set Λ of Γ is the unique minimal Γ -invariant closed subset of \mathcal{F} . Hence the following set

$$\mathcal{E} := \{[g] \in \Gamma \backslash G : gP \in \Lambda\}$$

is the unique minimal P -invariant closed subset of $\Gamma \backslash G$. We call a P -quasi-invariant measure on $\Gamma \backslash G$ *non-trivial* if its support is contained in \mathcal{E} .

Theorem 1.1. For any Anosov subgroup $\Gamma < G$, the space \mathcal{Q}_Γ of all non-trivial NM -invariant ergodic and A -quasi-invariant Radon measures on $\Gamma \backslash G$, up to constant multiples, is homeomorphic to $\mathbb{R}^{\text{rank } G-1}$.

In order to describe the explicit homeomorphism, we need to define Burger-Roblin measures on \mathcal{E} . Denote by \mathfrak{a} the Lie algebra of A and fix a positive Weyl chamber $\mathfrak{a}^+ \subset \mathfrak{a}$ so that $\log N$ is the sum of positive root subspaces. Fix a maximal compact subgroup K of G so that the Cartan decomposition $G = K(\exp \mathfrak{a}^+)K$ holds. Let $\mu : G \rightarrow \mathfrak{a}^+$ denote the Cartan projection map (Def. 2.2). We denote by $\mathcal{L}_\Gamma \subset \mathfrak{a}^+$ the limit cone of Γ , which is the asymptotic cone of $\mu(\Gamma)$ (Def. 2.16). Let $\psi_\Gamma : \mathfrak{a} \rightarrow \mathbb{R} \cup \{-\infty\}$ denote the growth indicator function of Γ (Def. 2.17).

For Anosov subgroups, the following two spaces are homeomorphic to each other:

$$D_\Gamma^* := \{\psi \in \mathfrak{a}^* : \psi \geq \psi_\Gamma, \psi(v) = \psi_\Gamma(v) \text{ for some } v \in \text{int } \mathcal{L}_\Gamma\} \simeq \text{int}(\mathbb{P}\mathcal{L}_\Gamma)$$

where $\text{int}(\mathbb{P}\mathcal{L}_\Gamma)$ denotes the interior of the projective limit cone $\mathbb{P}\mathcal{L}_\Gamma$ (Proposition 4.4). Since $\text{int}(\mathcal{L}_\Gamma)$ is a non-empty open convex cone of \mathfrak{a}^+ [6, Thm. 1.2], it follows that D_Γ^* is

homeomorphic to $\mathbb{R}^{\text{rank } G-1}$. We remark that D_Γ^* is in fact a closed analytic submanifold of \mathfrak{a}^* [38, Prop. 4.11].

For a linear form $\psi \in \mathfrak{a}^*$, a Borel probability measure ν on the limit set Λ is called a (Γ, ψ) -Patterson–Sullivan measure if for all $\gamma \in \Gamma$ and $\xi \in \mathcal{F}$,

$$\frac{d\gamma_\star \nu}{d\nu}(\xi) = e^{\psi(\beta_\xi(o, \gamma o))} \quad (1.2)$$

where $o = [K] \in G/K$ and $\beta : \mathcal{F} \times G/K \times G/K \rightarrow \mathfrak{a}$ denotes the \mathfrak{a} -valued Busemann function (Def. 3.2). Quint constructed a (Γ, ψ) -Patterson–Sullivan measure for each $\psi \in D_\Gamma^*$ [40]; for Γ Anosov, this measure exists uniquely (hence Γ -ergodic), which we denote by ν_ψ (see Theorem 4.3 and references therein).

In the rest of the introduction, we let $\Gamma < G$ be an Anosov subgroup. By a Patterson–Sullivan measure on Λ , we mean a (Γ, ψ) -Patterson–Sullivan measure on Λ for some $\psi \in \mathfrak{a}^*$. We show:

Theorem 1.3. The map $\psi \mapsto \nu_\psi$ is a homeomorphism between D_Γ^* and the space of all Patterson–Sullivan measures on Λ . Moreover, Patterson–Sullivan measures are pairwise mutually singular.

When Γ is the fundamental group of a closed negatively curved manifold, the above theorem also follows from [33].

We also denote by ν_ψ the M -invariant lift of ν_ψ on $\mathcal{F} \simeq K/M$ to K by abuse of notation. The Burger–Roblin measure m_ψ^{BR} on $\Gamma \backslash G$ is induced from the following Γ -invariant measure $\tilde{m}_\psi^{\text{BR}}$ on G : for $g = k(\exp b)n \in KAN$,

$$d\tilde{m}_\psi^{\text{BR}}(g) = e^{\psi(b)} dn db d\nu_\psi(k) \quad (1.4)$$

where dn and db are Lebesgue measures on N and \mathfrak{a} , respectively.

The following is a more elaborate version of Theorem 1.1:

Theorem 1.5 (Classification). The map $\psi \mapsto [m_\psi^{\text{BR}}]$ defines a homeomorphism between D_Γ^* and \mathcal{Q}_Γ .

While the P -ergodicity of m_ψ^{BR} follows from the Γ -ergodicity of ν_ψ , establishing the ergodicity of m_ψ^{BR} , and hence the well-definedness of the above map, is the most significant part of Theorem 1.5:

Theorem 1.6 (Ergodicity). For each $\psi \in D_\Gamma^*$, m_ψ^{BR} is NM -ergodic.

Since \mathcal{E} is a second countable topological space, Theorem 1.6 implies:

Corollary 1.7. For m_ψ^{BR} almost all $x \in \mathcal{E}$,

$$\overline{xNM} = \mathcal{E}.$$

A Radon measure m on $\Gamma \backslash G$ is called P -semi-invariant if there exists a character $\chi : P \rightarrow \mathbb{R}^*$ such that $p_*m = \chi(p)m$ for all $p \in P$. Note that any P -semi-invariant Radon measure is necessarily NM -invariant since NM is unimodular. We show that any P -semi-invariant Radon measure on \mathcal{E} is of the form m_ψ^{BR} for some $\psi \in D_\Gamma^*$ (Proposition 10.25). Hence Theorem 1.6 implies:

Corollary 1.8. The space of all P -semi-invariant Radon measures on \mathcal{E} coincides with \mathcal{Q}_Γ , up to constant multiples.

Discussion on the proof of Theorem 1.6.

Fix $\psi \in D_\Gamma^*$. Defining a Γ -invariant Radon measure $\widehat{\nu}_\psi$ on $\mathcal{H} := G/NM \simeq \mathcal{F} \times \mathfrak{a}$ by

$$d\widehat{\nu}_\psi(gP, b) = e^{\psi(b)} d\nu_\psi(gP) db,$$

the standard duality theorem implies that the NM -ergodicity of m_ψ^{BR} is equivalent to the Γ -ergodicity of $\widehat{\nu}_\psi$.

Generalizing the observation of Schmidt [52] (also see [46]) to a higher rank situation, the Γ -ergodicity of $\widehat{\nu}_\psi$ follows if the closed subgroup, say $E_{\nu_\psi} = E_{\nu_\psi}(\Gamma)$, consisting of all ν_ψ -essential values is equal to \mathfrak{a} (Proposition 9.2):

Definition 1.9. An element $v \in \mathfrak{a}$ is called a (Γ, ν_ψ) -essential value, if for any $\varepsilon > 0$ and any Borel set $B \subset \mathcal{F}$ with $\nu_\psi(B) > 0$, there exists $\gamma \in \Gamma$ such that

$$B \cap \gamma B \cap \{\xi \in \mathcal{F} : \|\beta_\xi(o, \gamma o) - v\| < \varepsilon\}$$

has a positive ν_ψ -measure.

Recalling that the Jordan projection $\lambda(\Gamma)$ of Γ generates a dense subgroup of \mathfrak{a} [6], the following is the main ingredient of our proof of Theorem 1.6:

Proposition 1.10. For each $\psi \in D_\Gamma^*$, there exists a finite subset $F_\psi \subset \lambda(\Gamma)$ such that

$$\lambda(\Gamma) - F_\psi \subset E_{\nu_\psi}(\Gamma).$$

In particular, $E_{\nu_\psi}(\Gamma) = \mathfrak{a}$.

See Proposition 10.2 for a more general version stated for any Zariski dense normal subgroup of Γ .

Among other things, the following three key properties of Anosov groups play important roles in the proof of Proposition 1.10:

- (1) (Antipodality) $\Lambda \times \Lambda - \{(\xi, \xi)\} \subset \mathcal{F}^{(2)}$;
- (2) (Regularity) If $\gamma_i \rightarrow \infty$ in Γ , then $\alpha(\mu(\gamma_i)) \rightarrow \infty$ for each simple root α of $\text{Lie}(G)$ with respect to \mathfrak{a}^+ ;
- (3) (Morse property) There exists a constant $D > 0$ such that any discrete geodesic ray $[e, x)$ in Γ tending to $x \in \partial\Gamma$ is contained in the D -neighborhood of some gA^+ in G where $g \in G$ satisfies $gP = \zeta(x)$.

(1) is a part of the definition of an Anosov subgroup. (2) follows from the fact that $\mathcal{L}_\Gamma \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ ([41], [48], [8]) in view of Lemma 7.2. (3) is proved in [25] (see also Proposition 5.12).

Many aspects of our proof of Proposition 1.10 can be simplified for a special class of $\psi \in D_\Gamma^*$ with certain strong positivity property (cf. Lemma 5.1); however, as our eventual goal is the classification theorem as stated in Theorem 1.1, we need to address all $\psi \in D_\Gamma^*$, which makes the proof much more intricate and requires the full force of the Anosov property of Γ .

Fix $\gamma_0 \in \Gamma$. We aim to show that $\lambda(\gamma_0) \in E_{\nu_\psi}(\Gamma)$. More precisely, for any $\varepsilon > 0$ and any Borel subset $B \subset \mathcal{F}$ with $\nu_\psi(B) > 0$, there exists $\gamma \in \Gamma$ such that

$$\nu_\psi(B \cap \gamma \gamma_0 \gamma^{-1} B \cap \{\xi \in \mathcal{F} : \|\beta_\xi(o, \gamma \gamma_0 \gamma^{-1} o) - \lambda(\gamma_0)\| < \varepsilon\}) > 0. \quad (1.11)$$

For $p \in G/K$, we define

$$d_{\psi, p}(\xi_1, \xi_2) = e^{-[\xi_1, \xi_2]_{\psi, p}}$$

for any $\xi_1 \neq \xi_2$ in Λ , where $[\cdot, \cdot]_{\psi, p}$ denotes the ψ -Gromov product based at p (Def. 6.1). Its well-definedness is due to the antipodality (1). In the rank one case, this is simply the restriction of the classical visual metric to the limit set Λ . In general, it is not even

symmetric but we show that any sufficiently small power of $d_{\psi,p}$ is comparable to some genuine metric on Λ :

Theorem 1.12. For all sufficiently small $s > 0$, there exist a metric d_s on Λ and $C_s > 0$ such that for all $\xi_1 \neq \xi_2$ in Λ ,

$$C_s^{-1} d_s(\xi_1, \xi_2) \leq d_{\psi,p}(\xi_1, \xi_2)^s \leq C_s d_s(\xi_1, \xi_2).$$

Remark 1.13. In the process of proving this theorem, we also show that the Gromov product on $\partial\Gamma$ and the ψ -Gromov product $[\cdot, \cdot]_{\psi,p}$ are equivalent to each other (see Theorem 6.13).

As a consequence of Theorem 1.12, $d_{\psi,p}$ can be used to define virtual balls with respect to which Vitali type covering lemma can be applied (Lemma 6.12). Let $\xi_0 \in \mathcal{F}$ denote the attracting fixed point of γ_0 and consider the family

$$D(\gamma\xi_0, r) := \mathbb{B}_p(\gamma\xi_0, \frac{1}{3}e^{-\psi(\underline{a}(\gamma^{-1}p, p) + i\underline{a}(\gamma^{-1}p, p))}r), \quad \gamma \in \Gamma, r > 0$$

where $\underline{a}(q, p)$ denotes the \mathfrak{a}^+ -valued distance from q to p (Def. 2.4). We then show that for all sufficiently small $r > 0$, there are infinitely many $D(\gamma\xi_0, r)$ satisfying (1.11) (Lemma 10.12). The key ingredient in this step is the following:

Lemma 1.14. There exists $C = C(\psi, p) > 0$ such that for all $\gamma \in \Gamma$ and $\xi \in \Lambda$,

$$-\psi(\underline{a}(p, \gamma p)) - C \leq \psi(\beta_\xi(\gamma p, p)) \leq \psi(\underline{a}(\gamma p, p)) + C.$$

In the rank one case, a stronger statement $-d(p, q) \leq \beta_\xi(q, p) \leq d(p, q)$ holds for all $q, p \in G/K$ and $\xi \in \mathcal{F}$, which generalizes to strongly positive linear forms (Lemma 5.1). For a general $\psi \in D_\Gamma^*$, our proof of Lemma 1.14 is based on the property that the orbit map $\gamma \mapsto \gamma(o)$ sends a shadow in the word hyperbolic group Γ to a shadow in the symmetric space G/K (Proposition 5.14), as well as the following lemma, which is of independent interest: we denote by $|\cdot|$ the word length on Γ with respect to a fixed finite symmetric generating subset.

Lemma 1.15. There exists $R > 0$ such that for any $\gamma_1, \gamma_2 \in \Gamma$ with $|\gamma_1 \gamma_2| = |\gamma_1| + |\gamma_2|$, we have

$$\|\mu(\gamma_1 \gamma_2) - \mu(\gamma_1) - \mu(\gamma_2)\| < R.$$

We emphasize that this lemma does not follow from the property of Anosov groups that $(\Gamma, |\cdot|) \rightarrow G$ is a quasi-isometric embedding [22, Thm. 1.7], due to the non-trivial multiplicative constant.

To establish (1.11), we approximate a general Borel subset $B \subset \mathcal{F}$ by some $D(\gamma \xi_0, r)$ satisfying (1.11). In this step, we prove the following higher rank generalization of Tukia's theorem [56, Thm. 4A] (see also [34], [1], [35]):

Theorem 1.16. For any Patterson–Sullivan measure ν on Λ , the set of Myrberg limit points (Def. 8.1) has full ν -measure.

It follows that for the *AM*-invariant Bowen–Margulis–Sullivan measure m_ψ^{BMS} on $\Gamma \backslash G$, almost all points have dense $A^+ M$ orbits (Corollary 8.12). Using the property that virtual balls $\mathbb{B}_p(\gamma \xi_0, r)$ satisfy a covering lemma (Lemma 6.12) that is a consequence of Theorem 1.12, we show that ν_ψ -almost all Myrberg limit points satisfy the Lebesgue density type statement for the family $\{D(\gamma \xi_0, r) : \gamma \in \Gamma, r > 0\}$ (Proposition 10.17). By Theorem 1.16, this gives a desired approximation of B by some $D(\gamma \xi_0, r)$ satisfying (1.11).

We finally remark that in our subsequent work [30], we present refined versions of Theorems 1.1 and 1.6, building on the main results of this paper.

Organization: In section 2, we go over basic definitions and properties of Zariski dense discrete subgroups of G . In section 3, we discuss the notion of \mathfrak{a} -valued Gromov product and define the generalized BMS measures for a pair of (Γ, ψ) -conformal densities on \mathcal{F} . From section 4, we assume that Γ is Anosov. In section 4, we observe that the BMS measure m_ψ^{BMS} is *AM*-ergodic for each $\psi \in D_\Gamma^*$. Sections 5 and 6 are devoted to proving Lemma 1.14 and Theorem 1.12, respectively. In section 7, we prove that the space of PS-measures on Λ is homeomorphic to D_Γ^* , which is the first part of Theorem 1.3. In section 8, we show that the set of Myrberg limit points of Γ has full measure for any PS-measure on Λ . In section 9, we discuss the relation between the set of essential values of ν_ψ and the *NM*-ergodicity of m_ψ^{BR} . In the final section 10, we prove Theorems 1.6, 1.5, and the second part of Theorem 1.3.

Added to proof: It was recently shown that for any $\psi \in D_\Gamma^*$, any ψ -conformal measure is necessarily supported on Λ , and hence there exists a unique (Γ, ψ) -conformal

measure on \mathcal{F} (not only on Λ), first in [16] for ranks at most 3 and in [31] for general ranks. As a consequence, the space of all Γ -Patterson–Sullivan measures on Λ in Theorem 1.3 is equal to the space of all (Γ, ψ) -conformal measures on \mathcal{F} , $\psi \in D_\Gamma^*$.

2 Limit Set and Limit Cone

Let G be a connected, semisimple real algebraic group with finite center, and $\Gamma < G$ be a Zariski dense discrete subgroup. We fix, once and for all, a Cartan involution θ of the Lie algebra \mathfrak{g} of G , and decompose \mathfrak{g} as $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where \mathfrak{k} and \mathfrak{p} are the $+1$ and -1 eigenspaces of θ , respectively. We denote by K the maximal compact subgroup of G with Lie algebra \mathfrak{k} , and by $X = G/K$ the associated symmetric space. We also choose a maximal abelian subalgebra \mathfrak{a} of \mathfrak{p} . Choosing a closed positive Weyl chamber \mathfrak{a}^+ of \mathfrak{a} , let $A := \exp \mathfrak{a}$ and $A^+ = \exp \mathfrak{a}^+$. The centralizer of A in K is denoted by M , and we set N to be the contracting horospherical subgroup: for $a \in \text{int } A^+$, $N = \{g \in G : a^{-n}ga^n \rightarrow e \text{ as } n \rightarrow +\infty\}$. Note that $\log N$ is the sum of all positive root subspaces for our choice of A^+ . Similarly, we also consider the expanding horospherical subgroup N^+ : for $a \in \text{int } A^+$, $N^+ = \{g \in G : a^n ga^{-n} \rightarrow e \text{ as } n \rightarrow +\infty\}$. We set

$$P^+ = MAN^+, \quad \text{and} \quad P = P^- = MAN;$$

they are minimal parabolic subgroups of G that are opposite to each other. The quotient $\mathcal{F} = G/P$ is known as the Furstenberg boundary of G and is isomorphic to K/M .

Let $N_K(\mathfrak{a})$ be the normalizer of \mathfrak{a} in K . Let $\mathcal{W} := N_K(\mathfrak{a})/M$ denote the Weyl group. Fixing a left G -invariant and right K -invariant Riemannian metric on G induces a \mathcal{W} -invariant inner product on \mathfrak{a} , which we denote by $\langle \cdot, \cdot \rangle$. The identity coset $[e]$ in G/K is denoted by o .

Denote by $w_0 \in \mathcal{W}$ the unique element in \mathcal{W} such that $\text{Ad}_{w_0} \mathfrak{a}^+ = -\mathfrak{a}^+$; it is the longest Weyl element. Note that $w_0 P w_0^{-1} = P^+$.

Definition 2.1 (Visual map). For each $g \in G$, we define

$$g^+ := gP \in G/P \quad \text{and} \quad g^- := gw_0P \in G/P.$$

For all $g \in G$ and $m \in M$, observe that $g^\pm = (gm)^\pm = g(e^\pm)$. Let $\mathcal{F}^{(2)}$ denote the unique open G -orbit in $\mathcal{F} \times \mathcal{F}$:

$$\mathcal{F}^{(2)} = G(e^+, e^-) = \{(g^+, g^-) \in \mathcal{F} \times \mathcal{F} : g \in G\}.$$

Note that the stabilizer of (e^+, e^-) is the intersection $P^- \cap P^+ = MA$.

We say that $\xi, \eta \in \mathcal{F}$ are in general position if $(\xi, \eta) \in \mathcal{F}^{(2)}$. The Bruhat decomposition says that G is the disjoint union $\cup_{w \in \mathcal{W}} NwP^+$, and NP^+ is Zariski open and dense in G . Hence $(\xi, \eta) \notin \mathcal{F}^{(2)}$ if and only if $(\xi, \eta) \in G(e^+, we^-)$ for some $w \in \mathcal{W} - \{e\}$.

Cartan projection and \mathfrak{a}^+ -valued distance

Definition 2.2 (Cartan projection). For each $g \in G$, there exists a unique element $\mu(g) \in \mathfrak{a}^+$, called the Cartan projection of g , such that

$$g \in K \exp(\mu(g))K.$$

When $\mu(g) \in \text{int } \mathfrak{a}^+$ and $g = k_1 \exp(\mu(g))k_2$, k_1, k_2 are determined uniquely up to mod M , more precisely, if $g = k'_1 \exp(\mu(g))k'_2$, then for some $m \in M$, $k_1 = k'_1 m$ and $k_2 = m^{-1} k'_2$. We write

$$\kappa_1(g) := [k_1] \in K/M \quad \text{and} \quad \kappa_2(g) := [k_2] \in M \setminus K.$$

Lemma 2.3. [6, Lem. 4.6] For any compact subset $L \subset G$, there exists a compact subset $Q = Q(L) \subset \mathfrak{a}$ such that for all $g \in G$,

$$\mu(LgL) \subset \mu(g) + Q.$$

Definition 2.4 (\mathfrak{a}^+ -valued distance). We define $\underline{a} : X \times X \rightarrow \mathfrak{a}^+$ by

$$\underline{a}(p, q) := \mu(g^{-1}h)$$

where $p = g(o)$ and $q = h(o)$.

Accumulation of points of X on \mathcal{F}

Let Π denote the set of all simple roots of \mathfrak{g} with respect to \mathfrak{a}^+ .

Definition 2.5. We write that

- (1) $v_i \rightarrow \infty$ regularly in \mathfrak{a}^+ if $\alpha(v_i) \rightarrow \infty$ as $i \rightarrow \infty$ for all $\alpha \in \Pi$;
- (2) $a_i \rightarrow \infty$ regularly in A^+ if $\log a_i \rightarrow \infty$ regularly in \mathfrak{a}^+ ;
- (3) $g_i \rightarrow \infty$ regularly in G if $\mu(g_i) \rightarrow \infty$ regularly in \mathfrak{a}^+ .

If $a_i \rightarrow \infty$ regularly in A^+ , then for all $n \in N^+$,

$$\lim_{i \rightarrow \infty} a_i n a_i^{-1} = e$$

uniformly on compact subsets of N .

Definition 2.6. We call Γ *regular* if for any sequence $\gamma_i \in \Gamma$ going to ∞ in G , $\gamma_i \rightarrow \infty$ regularly in G .

Lemma 2.7. If the closure of $\{(\xi_i, e^-) : i = 1, 2, \dots\}$ is contained in $\mathcal{F}^{(2)}$, then $a_i \xi_i \rightarrow e^+$ for any sequence $a_i \rightarrow \infty$ regularly in A^+ .

Proof. The hypothesis implies that $\xi_i = n_i e^+$ for a bounded sequence $n_i \in N^+$. Hence $a_i \xi_i = a_i n_i e^+ = (a_i n_i a_i^{-1}) e^+ \rightarrow e^+$ as $a_i \rightarrow \infty$ regularly in A^+ . ■

Definition 2.8.

- (1) A sequence $g_i \in G$ is said to converge to $\xi \in \mathcal{F}$, if $g_i \rightarrow \infty$ regularly in G and $\lim_{i \rightarrow \infty} \kappa_1(g_i)^+ = \xi$.
- (2) A sequence $p_i = g_i(o) \in X$ is said to converge to $\xi \in \mathcal{F}$ if g_i does.

Lemma 2.9. Consider a sequence $g_i = k_i a_i h_i^{-1}$ where $k_i \in K$, $a_i \in A^+$, $h_i \in G$ satisfy that $k_i^+ \rightarrow k_0^+$ in K , $h_i \rightarrow h_0$ in G , and $a_i \rightarrow \infty$ regularly in A^+ . Then for any $\xi \in \mathcal{F}$ in general position with h_0^- , we have

$$\lim_{i \rightarrow \infty} g_i \xi = k_0^+.$$

Proof. As $(\xi, h_0^-) \in \mathcal{F}^{(2)}$, we have $(h_0^{-1} \xi, e^-) \in \mathcal{F}^{(2)}$. Since $\mathcal{F}^{(2)}$ is open and $h_i^{-1} \xi \rightarrow h_0^{-1} \xi$, we have $(h_i^{-1} \xi, e^-) \in \mathcal{F}^{(2)}$ for all large i . By Lemma 2.7, $a_i h_i^{-1} \xi \rightarrow e^+$ as $i \rightarrow \infty$. Therefore, $\lim_{i \rightarrow \infty} g_i \xi = \lim_{i \rightarrow \infty} k_i (a_i h_i^{-1} \xi) = k_0^+$. ■

Lemma 2.10. If $g_i \in G$ converges to $\xi \in \mathcal{F}$ and $p_i \in X$ is bounded, then $\lim_{i \rightarrow \infty} g_i p_i = \xi$.

Proof. Write $g_i = k_i a_i \ell_i^{-1} \in KA^+K$. The hypothesis implies that $a_i \rightarrow \infty$ regularly in A^+ and $k_i^+ \rightarrow \xi$ as $i \rightarrow \infty$. Let $k_0 \in K$ be such that $k_0^+ = \xi$, and $g'_i \in G$ be such that $g'_i(o) = p_i$. Write $g_i g'_i = k'_i a'_i (\ell'_i)^{-1} \in KA^+K$. We need to show that $\lim_{i \rightarrow \infty} k'_i = k_0^+$. As $k_i^+ \rightarrow k_0^+$, it suffices to show that any limit of the sequence $k_i^{-1} k'_i$ belongs to $M = \text{Stab}_K e^+$.

Set $q_i := k_i^{-1}k'_i$. Let q be a limit of the sequence q_i . By passing to a subsequence, we may suppose $q_i \rightarrow q \in K$. Since $d(o, p_i) = d(g_i o, g_i p_i) = d(a_i o, q_i a'_i o)$, the sequence $h_i^{-1} := a_i^{-1} q_i a'_i$ is bounded. Passing to a subsequence, assume that h_i converges to some $h_0 \in G$ as $i \rightarrow \infty$. Choose $\eta \in \mathcal{F}$ that is in general position with both h_0^- and e^- . Then $\lim_{i \rightarrow \infty} a_i h_i^{-1} \eta = e^+$ and $\lim_{i \rightarrow \infty} q_i a'_i \eta = q^+$ by Lemma 2.9. Since $a_i h_i^{-1} \eta = q_i a'_i \eta$, we get $e^+ = q^+ = q(e^+)$. This implies $q \in \text{Stab}_K e^+ = M$. \blacksquare

Lemma 2.11. If $g_i \rightarrow g$ in G and $a_i \rightarrow \infty$ regularly in A^+ , then for any $p \in X$, $\lim_{i \rightarrow \infty} g_i a_i(p) = g^+$ and $\lim_{i \rightarrow \infty} g_i a_i^{-1}(p) = g^-$.

Proof. By Lemma 2.10, it suffices to consider the case when $p = o$. Write $g_i a_i = k_i b_i \ell_i^{-1} \in KA^+K$. As the sequence g_i is bounded, it follows from Lemma 2.3 that $b_i \rightarrow \infty$ regularly in A^+ . In order to show that $g_i a_i(o) \rightarrow g^+$, it suffices to show that if $k_i \rightarrow k_0$, then $k_0^+ = g^+$. By passing to a subsequence, we may assume that $\ell_i \rightarrow \ell_0$ in K . Choose $\xi \in \mathcal{F}$ that is in general position with both ℓ_0^- and e^- . Then $g_i a_i \xi \rightarrow k_0^+$ by Lemma 2.9. On the other hand, as $(\xi, e^-) \in \mathcal{F}^{(2)}$, $g_i a_i \xi \rightarrow g^+$ by Lemma 2.7. Hence $g^+ = k_0^+$, proving the first claim. Now the second claim follows since $g_i a_i^{-1} = g_i w_0 b_i w_0^{-1}$ for some $b_i \in A^+$, and $g_i w_0 b_i w_0^{-1}(o) = g_i w_0 b_i(o) \rightarrow (gw_0)^+ = g^-$. \blacksquare

Limit set and Limit cone

Denote by m_o the K -invariant probability measure on $\mathcal{F} \simeq K/M$.

Definition 2.12 (Limit set). The limit set Λ of Γ is defined as the set of all points $\xi \in \mathcal{F}$ such that the Dirac measure δ_ξ is a limit point of $\{\gamma_* m_o : \gamma \in \Gamma\}$ in the space of Borel probability measures on \mathcal{F} .

Benoist showed that Λ is the unique minimal Γ -invariant closed subset of \mathcal{F} . Moreover, Λ is Zariski dense in \mathcal{F} ([6, Section 3.6], see also [15, Lem. 2.10] for a stronger statement).

Lemma 2.13. For any $p \in X$, we have

$$\Lambda = \left\{ \lim_{i \rightarrow \infty} \gamma_i p \in \mathcal{F} : \gamma_i \in \Gamma \right\}.$$

Proof. Let $(\gamma_i)_* m_o \rightarrow \delta_\xi$, and write $\gamma_i = k_i a_i \ell_i^{-1} \in KA^+K$. Suppose $k_i \rightarrow k$. Then $(a_i)_* m_o \rightarrow \delta_{k^{-1}\xi}$. It follows that $a_i \rightarrow \infty$ regularly in A^+ and $k^{-1}\xi = e^+$, that is, $\xi = k^+$.

Hence $\gamma_i \rightarrow \xi$ and hence $\gamma_i(p) \rightarrow \xi$ by Lemma 2.10. This proves the inclusion \subset . If $\gamma_i p \rightarrow \xi$ and $\gamma_i = k_i a_i \ell_i^{-1} \in KA^+K$, then $a_i \rightarrow \infty$ regularly and $k_i^+ \rightarrow \xi$. Since $(a_i)_* m_o$ converges to δ_{e^+} , we have $(\gamma_i)_* m_o \rightarrow \delta_\xi$. This proves the other inclusion. ■

Any element $g \in G$ can be written as the commuting product $g_h g_e g_u$, where g_h , g_e , and g_u are unique elements that are conjugate to elements of A^+ , K , and N , respectively. When g_h is conjugate to an element of $\text{int } A^+$, g is called *loxodromic*; in such a case, $g_u = e$. If a loxodromic element $g \in G$ satisfies $\varphi^{-1} g_h \varphi \in \text{int } A^+$ for $\varphi \in G$, then

$$Y_g := \varphi^+ \quad (2.14)$$

is called the attracting fixed point of g . We then have $Y_{g^{-1}} = \varphi^-$.

Lemma 2.15. [6, Lem. 3.6] The set

$$\{(Y_\gamma, Y_{\gamma^{-1}}) \in \Lambda \times \Lambda : \gamma \text{ is a loxodromic element of } \Gamma\}$$

is dense in $\Lambda \times \Lambda$.

The Jordan projection of g is defined as $\lambda(g) \in \mathfrak{a}^+$, where $\exp \lambda(g)$ is the element of A^+ conjugate to g_h .

Definition 2.16 (Limit cone). The *limit cone* $\mathcal{L}_\Gamma \subset \mathfrak{a}^+$ of Γ is defined as the smallest closed cone containing the *Jordan projection* $\lambda(\Gamma)$. Alternatively, it can be defined as the asymptotic cone of $\mu(\Gamma)$ [6, Thm. 1.2].

The limit cone \mathcal{L}_Γ is a convex subset of \mathfrak{a}^+ with non-empty interior [6, Thm. 1.2].

Definition 2.17 (Growth indicator function). The growth indicator function $\psi_\Gamma : \mathfrak{a}^+ \rightarrow \mathbb{R} \cup \{-\infty\}$ is defined as a homogeneous function, that is, $\psi_\Gamma(tu) = t\psi_\Gamma(u)$, such that for any unit vector $u \in \mathfrak{a}^+$,

$$\psi_\Gamma(u) := \inf_{\substack{\text{open cones } \mathcal{C} \subset \mathfrak{a}^+ \\ u \in \mathcal{C}}} \tau_{\mathcal{C}}$$

where $\tau_{\mathcal{C}}$ is the abscissa of convergence of the series $\sum_{\gamma \in \Gamma, \mu(\gamma) \in \mathcal{C}} e^{-t\|\mu(\gamma)\|}$.

We may consider ψ_Γ as a function on \mathfrak{a} by setting $\psi_\Gamma = -\infty$ outside of \mathfrak{a}^+ . Quint showed the following:

Theorem 2.18. [39, Thm. IV.2.2] The growth indicator function ψ_Γ is concave, is upper-semicontinuous, and satisfies

$$\mathcal{L}_\Gamma = \{u \in \mathfrak{a}^+ : \psi_\Gamma(u) > -\infty\}.$$

Moreover, ψ_Γ is non-negative on \mathcal{L}_Γ and positive on $\text{int } \mathcal{L}_\Gamma$.

3 \mathfrak{a} -Valued Gromov Product and Generalized BMS Measures

Iwasawa cocycle and \mathfrak{a} -valued Busemann function

The Iwasawa decomposition says that the product map $K \times A \times N \rightarrow G$ is a diffeomorphism.

Definition 3.1. The Iwasawa cocycle $\sigma : G \times \mathcal{F} \rightarrow \mathfrak{a}$ is defined as follows: for $(g, \xi) \in G \times \mathcal{F}$, $\sigma(g, \xi) \in \mathfrak{a}$ is the unique element satisfying

$$gk \in K \exp(\sigma(g, \xi))N \tag{3.2}$$

where $k \in K$ is such that $\xi = k^+$.

It satisfies the cocycle relation

$$\sigma(g_1 g_2, \xi) = \sigma(g_1, g_2 \xi) + \sigma(g_2, \xi)$$

for all $g_1, g_2 \in G$ and $\xi \in \mathcal{F}$.

Definition 3.2. The \mathfrak{a} -valued Busemann function $\beta : \mathcal{F} \times X \times X \rightarrow \mathfrak{a}$ is defined as follows: for $\xi \in \mathcal{F}$ and $g(o), h(o) \in X$,

$$\beta_\xi(g(o), h(o)) := \sigma(g^{-1}, \xi) - \sigma(h^{-1}, \xi).$$

Observe that the Busemann function is continuous in all three variables. To ease the notation, we will write $\beta_\xi(g, h) = \beta_\xi(g(o), h(o))$. We can check that for all $g, h, q \in G$

and $\xi \in \mathcal{F}$,

$$\begin{aligned}\beta_\xi(g, h) + \beta_\xi(h, q) &= \beta_\xi(g, q), \\ \beta_{g\xi}(gh, gq) &= \beta_\xi(h, q), \text{ and} \\ \beta_\xi(g, e) &= \sigma(g^{-1}, \xi).\end{aligned}\tag{3.4}$$

Geometrically, if $\xi = k^+ \in \mathcal{F}$ for $k \in K$, then for any unit vector $u \in \mathfrak{a}^+$,

$$\langle \beta_\xi(g, h), u \rangle = \lim_{t \rightarrow +\infty} d(g(o), \xi_t) - d(h(o), \xi_t)$$

where $\xi_t = k \exp(tu)o \in X$.

Lemma 3.5. For any loxodromic element $g \in G$ and $p \in X$,

$$\beta_{Y_g}(p, gp) = \lambda(g) \quad \text{and} \quad \beta_{Y_{g^{-1}}}(p, gp) = -\lambda(g^{-1}).$$

Proof. Let $\varphi \in G$ be so that $g = \varphi a m \varphi^{-1}$ for some $a \in A^+$ and $m \in M$. If $p = h(o)$ for $h \in G$, then, since g^{-1} fixes $\varphi^+ = Y_g$,

$$\beta_{Y_g}(p, gp) = \beta_{\varphi^+}(ho, gho) = \sigma(h^{-1}, \varphi^+) - \sigma(h^{-1}g^{-1}, \varphi^+) = -\sigma(g^{-1}, \varphi^+).$$

Writing $\varphi = kb$ with $k \in K$ and $b \in P$, we have

$$g^{-1}k = \varphi(am)^{-1}\varphi^{-1}k = kb(am)^{-1}b^{-1} \in Ka^{-1}N.$$

This gives $\sigma(g^{-1}, \varphi^+) = \log a^{-1} = -\lambda(g)$, and hence the first identity. The second identity follows from the first, by replacing g with g^{-1} . ■

\mathfrak{a} -valued Gromov product

Definition 3.6 (Opposition involution). The involution $i : \mathfrak{a} \rightarrow \mathfrak{a}$ defined by

$$i(u) = -\text{Ad}_{w_0}(u)$$

is called the opposition involution; it preserves \mathfrak{a}^+ . Note that for all $g \in G$,

$$\lambda(g^{-1}) = i(\lambda(g)) \quad \text{and} \quad \mu(g^{-1}) = i(\mu(g)).$$

It follows that

$$i(\mathcal{L}_\Gamma) = \mathcal{L}_\Gamma \quad \text{and} \quad \psi_\Gamma \circ i = \psi_\Gamma. \quad (3.7)$$

Definition 3.8. We define the \mathfrak{a} -valued Gromov product on $\mathcal{F}^{(2)}$ as follows: for $(\xi, \eta) \in \mathcal{F}^{(2)}$,

$$\mathcal{G}(\xi, \eta) := \beta_{g^+}(e, g) + i \beta_{g^-}(e, g)$$

where $g \in G$ satisfies $g^+ = \xi$ and $g^- = \eta$.

The definition does not depend on the choice of a representative of $[g] \in G/AM$. For all $h \in G$ and $(x, y) \in \mathcal{F}^{(2)}$, we have the following identity:

$$\mathcal{G}(hx, hy) - \mathcal{G}(x, y) = \sigma(h, x) + i \sigma(h, y). \quad (3.9)$$

As $\mathcal{G}(y, x) = i \mathcal{G}(x, y)$, the Gromov product is not symmetric in general.

Lemma 3.10. [55] There exists a family of irreducible representations (ρ_α, V_α) , $\alpha \in \Pi$, of G so that

- (1) the highest weight χ_α of ρ_α is a positive integral multiple of the fundamental weight ϖ_α corresponding to α ;
- (2) the highest weight space of ρ_α is one dimensional.

For $\alpha \in \Pi$, denote by V_α^+ the highest weight space of ρ_α , and by V_α^- its unique complementary A -invariant subspace in V_α . We have $\rho_\alpha(P)V_\alpha^+ = V_\alpha^+$, and hence the map $g \mapsto (\rho_\alpha(g)V_\alpha^+)_{\alpha \in \Pi}$ factors through a proper immersion

$$\mathcal{F} = G/P \rightarrow \prod_{\alpha \in \Pi} \mathbb{P}(V_\alpha).$$

Let $\langle \cdot, \cdot \rangle_\alpha$ be a K -invariant inner product on V_α with respect to which A is symmetric; then V_α^+ and V_α^- are orthogonal to each other. We denote by $\|\cdot\|_\alpha$ the norm on V_α induced by $\langle \cdot, \cdot \rangle_\alpha$. For $\varphi \in V_\alpha^*$, $\|\varphi\|_\alpha$ means the operator norm of φ . We also use the notation $\|\cdot\|_\alpha$ for a bi- $\rho_\alpha(K)$ -invariant norm on $\mathrm{GL}(V_\alpha)$.

Lemma 3.11. For all $\alpha \in \Pi$ and $g \in G$,

$$\chi_\alpha(\mathcal{G}(g^+, g^-)) = -\log \frac{|\varphi(v)|}{\|\varphi\|_\alpha \|v\|_\alpha} \quad (3.12)$$

where $v \in gV_\alpha^+$ and $\varphi \in V_\alpha^*$ is such that $\ker \varphi = gV_\alpha^<$.

Proof. If we define $\mathcal{G}'(g^+, g^-)$ to be the unique element of \mathfrak{a} satisfying (3.12), it is shown in [47, Lem 4.12] that \mathcal{G}' satisfies (3.9). Hence for all $h \in G$,

$$\mathcal{G}'(h^+, h^-) - \mathcal{G}'(e^+, e^-) = \mathcal{G}(h^+, h^-) - \mathcal{G}(e^+, e^-).$$

We claim that $\mathcal{G}'(e^+, e^-) = 0$; to check this, take φ to be the projection $V \rightarrow V_\alpha^+$ parallel to $V_\alpha^<$. Since V_α^+ and $V_\alpha^<$ are orthogonal, it follows that $\|\varphi\|_\alpha = 1$. Now for $v \in V_\alpha^+$, we have

$$\frac{|\varphi(v)|}{\|\varphi\|_\alpha \|v\|_\alpha} = \frac{\|v\|_\alpha}{\|v\|_\alpha} = 1.$$

Since $\mathcal{G}(e^+, e^-) = 0$, we conclude $\mathcal{G} = \mathcal{G}'$ on $\mathcal{F}^{(2)}$. ■

Remark 3.13. In view of this lemma, our definition of Gromov product differs by $-i$ from the one given in [47].

Patterson–Sullivan measures on Λ

Definition 3.14 (Conformal measures). Given a closed subgroup $\Gamma < G$ and $\psi \in \mathfrak{a}^*$, a Borel probability measure ν on \mathcal{F} is called a (Γ, ψ) -conformal measure if, for any $\gamma \in \Gamma$ and $\xi \in \mathcal{F}$,

$$\frac{d\gamma_* \nu}{d\nu}(\xi) = e^{\psi(\beta_\xi(e, \gamma))} \quad (3.15)$$

where $\gamma_* \nu(Q) = \nu(\gamma^{-1} Q)$ for any Borel subset $Q \subset \mathcal{F}$.

If 2ρ denotes the sum of all positive roots of G with respect to \mathfrak{a}^+ , then it is a standard fact that a $(G, 2\rho)$ -conformal measure is precisely the unique K -invariant probability measure m_o on \mathcal{F} (cf. [44, Prop. 3.3]).

Fix a Zariski dense discrete subgroup $\Gamma < G$ in the rest of this section.

Definition 3.16 (Patterson–Sullivan measures). For $\psi \in \mathfrak{a}^*$, a (Γ, ψ) -conformal measure supported on the limit set Λ will be called a (Γ, ψ) -PS measure. By a PS measure on Λ , we mean a (Γ, ψ) -PS measure for some $\psi \in \mathfrak{a}^*$.

Set

$$D_\Gamma := \{\psi \in \mathfrak{a}^* : \psi \geq \psi_\Gamma\}.$$

The following collection of linear forms is of particular importance:

$$D_\Gamma^* := \{\psi \in D_\Gamma : \psi(u) = \psi_\Gamma(u) \text{ for some } u \in \mathcal{L}_\Gamma \cap \text{int } \mathfrak{a}^+\}. \quad (3.17)$$

By (3.7), $\psi \circ i \in D_\Gamma^*$ for all $\psi \in D_\Gamma^*$. The concavity of ψ_Γ and the non-emptiness of $\text{int } \mathcal{L}_\Gamma$ imply that D_Γ^* is non-empty by the Hahn–Banach theorem. When $\psi(u) = \psi_\Gamma(u)$, we say ψ is tangent to ψ_Γ at u .

Generalizing the work of Patterson and Sullivan ([37], [53]), Quint [40] constructed a (Γ, ψ) -PS measure for every $\psi \in D_\Gamma^*$.

Generalized BMS-measure m_{ν_1, ν_2}

Given a pair of Γ -conformal measures on \mathcal{F} , we now define an *MA*-semi invariant measure on $\Gamma \backslash G$, which we call a generalized BMS-measure.

Definition 3.18 (Hopf parametrization). The map

$$gM \rightarrow (g^+, g^-, b = \beta_{g^+}(e, g))$$

gives a homeomorphism between G/M and $\mathcal{F}^{(2)} \times \mathfrak{a}$, which is called the Hopf parametrization of G/M .

Fixing a pair of Γ -conformal measures $\nu_{\psi_1}, \nu_{\psi_2}$ on \mathcal{F} for a pair of linear forms $\psi_1, \psi_2 \in \mathfrak{a}^*$, we define a Radon measure $\tilde{m}_{\nu_{\psi_1}, \nu_{\psi_2}}$ on G/M as follows: for $g = (g^+, g^-, b) \in \mathcal{F}^{(2)} \times \mathfrak{a}$,

$$d\tilde{m}_{\nu_{\psi_1}, \nu_{\psi_2}}(g) = e^{\psi_1(\beta_{g^+}(e, g)) + \psi_2(\beta_{g^-}(e, g))} d\nu_{\psi_1}(g^+) d\nu_{\psi_2}(g^-) db, \quad (3.19)$$

where $db = d\ell(b)$ is the Lebesgue measure on \mathfrak{a} . This measure is left Γ -invariant, and hence induces a measure on $\Gamma \backslash G/M$. We denote by $m_{\nu_{\psi_1}, \nu_{\psi_2}}$ its *M*-invariant lift to $\Gamma \backslash G$.

It is A -semi-invariant as

$$a_* m_{v_{\psi_1}, v_{\psi_2}} = e^{(\psi_2 \circ i - \psi_1)(\log a)} m_{v_{\psi_1}, v_{\psi_2}} \quad (3.20)$$

for all $a \in A$ [15, Lem. 3.6].

BMS-measures: $m_{v_\psi, v_{\psi \circ i}}^{\text{BMS}}$. Let $\psi \in \mathfrak{a}^*$ and let v_ψ and $v_{\psi \circ i}$ be respectively (Γ, ψ) and $(\Gamma, \psi \circ i)$ -PS measures. We set

$$m_{v_\psi}^{\text{BMS}} := m_{v_\psi, v_{\psi \circ i}} \quad (3.21)$$

and call it the Bowen–Margulis–Sullivan measure associated to $(v_\psi, v_{\psi \circ i})$. It is right MA -invariant and its support is given by

$$\Omega := \{x \in \Gamma \backslash G : x^\pm \in \Lambda\};$$

since Λ is Γ -invariant, the condition $x^\pm \in \Lambda$ is well-defined. Note that for $[g] \in G/M$,

$$dm_{v_\psi}^{\text{BMS}}[g] = e^{\psi(\mathcal{G}(g^+, g^-))} d\nu_\psi(g^+) d\nu_{\psi \circ i}(g^-) db. \quad (3.22)$$

N -invariant BR-measures: $m_{v_\psi}^{\text{BR}}$. We set

$$m_{v_\psi}^{\text{BR}} := m_{v_\psi, m_o} \quad (3.23)$$

and call it the N -invariant Burger–Roblin measure associated to v_ψ . See [15, Section 3] for the equivalence of this definition with the one given in (1.4). The support of $m_{v_\psi}^{\text{BR}}$ is given by

$$\mathcal{E} := \{x \in \Gamma \backslash G : x^+ \in \Lambda\}.$$

4 Anosov Groups and AM-Ergodicity of BMS Measures

Let Γ be a Zariski dense discrete subgroup of G , and set $\Lambda^{(2)} := (\Lambda \times \Lambda) \cap \mathcal{F}^{(2)}$.

Definition 4.1. We say that $\Gamma < G$ is Anosov, if it is a finitely generated word hyperbolic group admitting a Γ -equivariant homeomorphism $\zeta : \partial\Gamma \rightarrow \Lambda$ such that $(\zeta(x), \zeta(y)) \in \Lambda^{(2)}$ for all $x \neq y \in \partial\Gamma$, where $\partial\Gamma$ denotes the Gromov boundary of Γ .

Such ζ is Hölder continuous and exists uniquely ([27, Prop. 3.2] and [8, Lem. 2.5]). We call it the limit map of Γ . We note that the antipodal property of Λ follows directly:

$$\Lambda \times \Lambda - \{(\xi, \xi)\} = \Lambda^{(2)}. \quad (4.2)$$

In the literature, this definition is referred to as P -Anosov for a minimal parabolic subgroup P of G . See [22], [21], and [25] for equivalent characterizations of Anosov subgroups.

In the rest of this section, let Γ be an Anosov subgroup of G .

The following theorem was proved by Quint [41, Prop. 3.2 and Thm. 4.7] for Schottky groups and by Sambarino [48, Coro. 3.12, 3.13 and 4.9] and by [15, Thm. 7.9] for general Anosov subgroups in view of the results in [13] (see also [15, Remark 7.10]):

Theorem 4.3.

- (1) $\mathcal{L}_\Gamma \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ and every non-trivial element of Γ is loxodromic ([27], [22]).
- (2) ψ_Γ is strictly concave and analytic on $\text{int } \mathcal{L}_\Gamma$.
- (3) $D_\Gamma^* = \{\psi \in D_\Gamma : \psi(u) = \psi_\Gamma(u) \text{ for some } u \in \text{int } \mathcal{L}_\Gamma\}$.
- (4) For any $\psi \in D_\Gamma^*$, $\psi > 0$ on $\mathcal{L}_\Gamma - \{0\}$.
- (5) For any $\psi \in D_\Gamma^*$, there exists a unique (Γ, ψ) -PS measure, say ν_ψ , on \mathcal{F} . In particular, ν_ψ is Γ -ergodic.

(1) and (3) imply that if $\psi \in D_\Gamma$ is tangent to ψ_Γ at some $u \in \mathcal{L}_\Gamma - \{0\}$, then $u \in \text{int } \mathcal{L}_\Gamma$. Note also that (1) implies that any Anosov subgroup is regular as defined in Def. 2.6. For $u \in \mathcal{L}_\Gamma$, we denote by $D_u \psi_\Gamma$ the directional derivative of ψ_Γ at u , whenever it exists.

Proposition 4.4. For each unit vector $u \in \text{int } \mathcal{L}_\Gamma$, $\psi_u := D_u \psi_\Gamma \in D_\Gamma^*$ and $D_u \psi_\Gamma(u) = \psi_\Gamma(u)$. Moreover, the map $u \mapsto \psi_u$ induces a homeomorphism between the set of unit vectors of $\text{int } \mathcal{L}_\Gamma$ ($\simeq \text{int } \mathbb{P}\mathcal{L}_\Gamma$) and D_Γ^* . Hence $D_\Gamma^* \simeq \mathbb{R}^{\text{rank } G-1}$.

Proof. See ([48, Thm. A], [15, Lem. 2.23]) for the first claim. The fact that $\psi_u \in D_\Gamma^*$ (and hence the well-definedness) and surjectivity of the map $u \mapsto D_u \psi_\Gamma$ follows from it, and the injectivity follows from the strict concavity of ψ_Γ as in Theorem 4.3(2). Continuity follows from the analyticity of ψ_Γ on $\text{int } \mathcal{L}_\Gamma$. We claim that if $D_{u_i} \psi_\Gamma \rightarrow D_u \psi_\Gamma$ for some unit vectors $u_i, u \in \text{int } \mathcal{L}_\Gamma$, then $u_i \rightarrow u$. Let $v \in \mathcal{L}_\Gamma$ be a limit of the sequence u_i . By passing to a subsequence, assume $u_i \rightarrow v$. By the upper-semi continuity of ψ_Γ

(Theorem 2.18), we have

$$\psi_\Gamma(v) \geq \limsup_{i \rightarrow \infty} \psi_\Gamma(u_i).$$

Since $\psi_\Gamma(u_i) = D_{u_i}\psi_\Gamma(u_i)$ and $D_{u_i}\psi_\Gamma \rightarrow D_u\psi_\Gamma$, we get $\psi_\Gamma(v) \geq D_u\psi_\Gamma(v)$. Since $D_u\psi_\Gamma \in D_\Gamma^*$, we have $\psi_\Gamma(v) = D_u\psi_\Gamma(v)$. It follows from Theorem 4.3(1) and (3) that $v \in \text{int } \mathcal{L}_\Gamma$. Since $\psi_\Gamma(u) = D_u\psi_\Gamma(u)$, the strict concavity of ψ_Γ on $\text{int } \mathcal{L}_\Gamma$ implies that $u = v$, establishing the homeomorphism. Since $\text{int}(\mathcal{L}_\Gamma)$ is a non-empty open convex cone of \mathfrak{a}^+ , $\text{int}(\mathbb{P}\mathcal{L}_\Gamma) \simeq \mathbb{P}\text{int}(\mathcal{L}_\Gamma)$ is homeomorphic to $\mathbb{R}^{\text{rank } G-1}$. \blacksquare

We denote by $\nabla\psi_\Gamma$ the gradient of ψ_Γ so that $D_u\psi_\Gamma(v) = \langle \nabla\psi_\Gamma(u), v \rangle$ for $u \in \text{int } \mathcal{L}_\Gamma$ and $v \in \mathfrak{a}$. Set

$$O_\Gamma := \{\nabla\psi_\Gamma(u) \in \mathfrak{a} : u \in \text{int } \mathcal{L}_\Gamma\}, \quad (4.5)$$

which is an open convex cone of $\mathfrak{a} - \{0\}$. By Proposition 4.4, the map $w \mapsto \langle w, \cdot \rangle$ gives a homeomorphism between $\{\nabla\psi_\Gamma(u) \in \mathfrak{a} : u \in \text{int } \mathcal{L}_\Gamma, \|u\| = 1\}$ and D_Γ^* , and hence a homeomorphism

$$O_\Gamma \simeq \mathbb{R}_+ D_\Gamma^*.$$

Quint showed that there exists a unique unit vector, say $u_\Gamma \in \text{int } \mathfrak{a}^+$, such that $\psi_\Gamma(u_\Gamma) = \max_{\|u\|=1} \psi_\Gamma(u)$. The vector u_Γ is called the direction of maximal growth of Γ . If we set $\delta_\Gamma := \psi_\Gamma(u_\Gamma)$, then $\nabla\psi_\Gamma(u_\Gamma) = \delta_\Gamma u_\Gamma$ and

$$\delta_\Gamma = \limsup_{T \rightarrow \infty} \frac{\log \#\{\gamma \in \Gamma : \|\mu(\gamma)\| < T\}}{T}.$$

Consider the following dual cone to \mathcal{L}_Γ :

$$\mathcal{L}_\Gamma^* := \{w \in \mathfrak{a} : \langle w, v \rangle \geq 0 \text{ for all } v \in \mathcal{L}_\Gamma\}.$$

Note that $\text{int } \mathcal{L}_\Gamma^* = \{w \in \mathfrak{a} : \langle w, v \rangle > 0 \text{ for all non-zero } v \in \mathcal{L}_\Gamma\}$.

Lemma 4.6. We have $O_\Gamma = \text{int } \mathcal{L}_\Gamma^*$. In particular,

$$\mathfrak{a}^+ - \{0\} \subset O_\Gamma \subset \mathfrak{a} - \{0\}.$$

Proof. If $w \in O_\Gamma$, then $\langle w, \cdot \rangle \in \mathbb{R}_+ D_\Gamma^*$, and hence by Theorem 4.3 (3), $\langle w, v \rangle > 0$ for all $v \in \mathcal{L}_\Gamma - \{0\}$. Hence $O_\Gamma \subset \text{int } \mathcal{L}_\Gamma^*$. Now suppose $w \in \text{int } \mathcal{L}_\Gamma^*$. Setting $\psi(v) := \langle w, v \rangle$, we claim that $\psi \in \mathbb{R}_+ D_\Gamma^*$; this implies $\text{int } \mathcal{L}_\Gamma^* \subset O_\Gamma$. Since $w \in \text{int } \mathcal{L}_\Gamma^*$, $\psi > 0$ on $\mathcal{L}_\Gamma - \{0\}$ and hence $c := \max_{\|v\|=1, v \in \mathcal{L}_\Gamma} \psi(v) > 0$. Since $\delta_\Gamma c^{-1} \psi \geq \psi_\Gamma$ on \mathcal{L}_Γ , and hence on \mathfrak{a} , it follows that for some $\varepsilon > 0$, $\varepsilon \psi \in D_\Gamma^*$, that is, $\psi \in \mathbb{R}_+ D_\Gamma^*$.

Since $\mathcal{L}_\Gamma - \{0\} \subset \text{int } \mathfrak{a}^+$ and the angle between any two walls of \mathfrak{a}^+ is at most $\pi/2$, the second claim follows. \blacksquare

AM-ergodicity of m_ψ^{BMS}

We fix $\psi \in D_\Gamma^*$ and set

$$v := v_\psi \quad \text{and} \quad m_\psi^{\text{BMS}} := m_{v_\psi, v_{\psi \circ i}}^{\text{BMS}}. \quad (4.7)$$

The composition $c := \psi \circ \sigma : \Gamma \times \Lambda \rightarrow \mathbb{R}$ is a Hölder cocycle satisfying $c(\gamma, y_\gamma) = \psi(\lambda(\gamma)) > 0$ for all non-trivial $\gamma \in \Gamma$.

Consider the action of Γ on $\Lambda^{(2)} \times \mathbb{R}$ given as follows: for $\gamma \in \Gamma$ and $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$,

$$\gamma \cdot (\xi, \eta, t) = (\gamma \xi, \gamma \eta, t + c(\gamma, \xi)).$$

The \mathbb{R} -action on $\Lambda^{(2)} \times \mathbb{R}$ defined by

$$\tau_s(\xi, \eta, t) = (\xi, \eta, t + s)$$

will be called translation flow.

The following is proved in [49, Thm. 3.2] when Γ is the fundamental group of a closed negatively curved manifold, and can be extended for general Anosov groups, using ingredients from [8]. The sketch of the proof can be found in [12, Appendix A].

Theorem 4.8. The action of Γ on $\Lambda^{(2)} \times \mathbb{R}$ is proper and cocompact, and the measure $\tilde{m}_\psi(\xi, \eta, t) = e^{\psi(\mathcal{G}(\xi, \eta))} d\nu_\psi(\xi) \otimes d\nu_{\psi \circ i}(\eta) \otimes dt$ induces the measure of maximal entropy, say m_ψ , for $\{\tau_s : s \in \mathbb{R}\}$ on $\Gamma \backslash \Lambda^{(2)} \times \mathbb{R}$. In particular, m_ψ is $\{\tau_s : s \in \mathbb{R}\}$ -ergodic.

In terms of the Hopf parametrization, Γ acts on $\Lambda^{(2)} \times \mathfrak{a} = \text{supp } \tilde{m}_\psi^{\text{BMS}}$ as follows: for $\gamma \in \Gamma$ and $(\xi, \eta, v) \in \Lambda^{(2)} \times \mathfrak{a}$,

$$\gamma \cdot (\xi, \eta, v) = (\gamma \xi, \gamma \eta, v + \sigma(\gamma, \xi)).$$

Corollary 4.9. For any $\psi \in D_\Gamma^*$, the *AM*-action on $(\Gamma \backslash G, m_\psi^{\text{BMS}})$ is ergodic and if $\text{rank } G \geq 2$, $|m_\psi^{\text{BMS}}| = \infty$.

Proof. The $\{\tau_s : s \in \mathbb{R}\}$ -ergodicity of m_ψ is equivalent to ergodicity of $(\Lambda^{(2)}, \Gamma, \nu_\psi \otimes \nu_{\psi \circ i}|_{\Lambda^{(2)}})$, which is again equivalent to the *AM*-ergodicity of m_ψ^{BMS} . Consider the projection map $\pi : \Gamma \backslash \Lambda^{(2)} \times \mathfrak{a} \rightarrow \Gamma \backslash \Lambda^{(2)} \times \mathbb{R}$ induced by the Γ -equivariant map $\Lambda^{(2)} \times \mathfrak{a} \rightarrow \Lambda^{(2)} \times \mathbb{R}$ given by $(\xi, \eta, v) \mapsto (\xi, \eta, \psi(v))$. Then π is a principal $\ker \psi$ -bundle, which is trivial as $\ker \psi$ is a vector group. It follows that there exists a $\ker \psi$ -equivariant homeomorphism between $\Gamma \backslash \Lambda^{(2)} \times \mathfrak{a}$ and $(\Gamma \backslash \Lambda^{(2)} \times \mathbb{R}) \times \ker \psi$. Therefore, m_ψ^{BMS} disintegrates over the measure m_ψ with conditional measure being the Lebesgue measure on $\ker \psi \simeq \mathbb{R}^{\text{rank } G-1}$ so that $m_\psi^{\text{BMS}} \simeq m_\psi \otimes \text{Leb}_{\ker \psi}$ (cf. [47, Prop. 3.5]). This gives the infinitude of $|m_\psi^{\text{BMS}}|$ when G has rank at least 2. ■

5 Comparing \mathfrak{a} -Valued Busemann Functions and Distances via ψ

When G has rank one, for any $p, q \in X$, the maximum and minimum of Busemann function $\beta_\xi(p, q), \xi \in \mathcal{F}$ are always achieved as $\pm d(p, q)$. A higher rank generalization of this fact can be stated as follows.

Lemma 5.1. Let $\psi \in \mathfrak{a}^*$ be strongly positive, in the sense that ψ is a non-negative linear combination of fundamental weights ϖ_α , $\alpha \in \Pi$. Then for any $p, q \in X$ and $\xi \in \mathcal{F}$, we have

$$-\psi(\underline{a}(q, p)) \leq \psi(\beta_\xi(p, q)) \leq \psi(\underline{a}(p, q)). \quad (5.2)$$

Proof. We use notations introduced in Lemma 3.10. Since ϖ_α is a positive multiple of χ_α , it suffices to prove the claim when $\psi = \chi_\alpha$ for $\alpha \in \Pi$.

Write $q = go$, and $p = hq$ for some $g, h \in G$. Note that

$$\chi_\alpha(\underline{a}(p, q)) = \chi_\alpha(\mu(g^{-1}h^{-1}g)) = \log \|\rho_\alpha(g^{-1}h^{-1}g)\|_\alpha.$$

Write $g^{-1}\xi = k^+$ for some $k \in K$ and $g^{-1}h^{-1}gk = k' an \in KAN$. Then

$$\beta_\xi(p, q) = \sigma(g^{-1}h^{-1}g, k^+) = \log a.$$

Hence for a unit vector $v \in V_\alpha$,

$$\chi_\alpha(\beta_\xi(p, q)) = \log \|\rho_\alpha(g^{-1}h^{-1}g)\rho_\alpha(k)v\| \leq \log \|\rho_\alpha(g^{-1}h^{-1}g)\|_\alpha = \chi_\alpha(\underline{a}(p, q)).$$

Since $\|\rho_\alpha(g^{-1})\|^{-1} \leq \|\rho_\alpha(g)v\|$ and $\chi_\alpha(\underline{a}(q, p)) = \log \|\rho_\alpha(g^{-1}hg)\|_\alpha$, we also get

$$\chi_\alpha(\beta_\xi(p, q)) \geq \log \|\rho_\alpha(g^{-1}hg)\|_\alpha^{-1} = -\chi_\alpha(\underline{a}(q, p)).$$

■

There are $\psi \in D_\Gamma^*$ that are not strongly positive (see Lemma 4.6). We establish the following modification for Anosov groups, which is the main goal of this section:

Theorem 5.3. Let $\Gamma < G$ be Anosov. For any $\psi \in D_\Gamma^*$ and $p \in X$, there exists $C = C(\psi, p) > 0$ such that for all $\gamma \in \Gamma$ and $\xi \in \Lambda$,

$$-\psi(\underline{a}(p, \gamma p)) - C \leq \psi(\beta_\xi(\gamma p, p)) \leq \psi(\underline{a}(\gamma p, p)) + C.$$

We begin by noting that $\psi(\underline{a}(\gamma p, p))$ is always positive possibly except for finitely many γ 's:

Lemma 5.4. Let $\psi \in D_\Gamma^*$ and $p \in X$. For any sequence $\gamma_i \rightarrow \infty$ in Γ , $\psi(\underline{a}(\gamma_i p, p)) \rightarrow +\infty$.

Proof. By Lemma 2.3, it suffices to check that $\psi(\mu(\gamma_i)) \rightarrow +\infty$ as $i \rightarrow \infty$. Setting $t_i := \|\mu(\gamma_i)\|^{-1}$, passing to a subsequence, we may assume that $t_i \mu(\gamma_i)$ converges to some unit vector $u \in \mathfrak{a}$. Since \mathcal{L}_Γ is the asymptotic cone of $\mu(\Gamma)$, we have $u \in \mathcal{L}_\Gamma$. Hence, we have $\psi(u) > 0$ by Lemma 4.3. Since $\psi(t_i \mu(\gamma_i)) \rightarrow \psi(u)$ and $\psi(\mu(\gamma_i)) = t_i^{-1} \psi(t_i \mu(\gamma_i))$, we have $\psi(\mu(\gamma_i)) \rightarrow +\infty$. ■

The following is the main ingredient of the proof of Theorem 5.3:

Proposition 5.5. For $p \in X$, there exists $C = C(p) > 0$ such that for each $(\gamma, \xi) \in \Gamma \times \Lambda$, we can find $\gamma_1 = \gamma_1(\xi), \gamma_2 = \gamma_2(\xi) \in \Gamma$ satisfying

- (1) $\gamma = \gamma_1 \gamma_2$ and $|\gamma| = |\gamma_1| + |\gamma_2|$;

- (2) $\|\beta_\xi(\gamma p, p) + \mu(\gamma_1) - \mu(\gamma_2^{-1})\| \leq C;$
- (3) $\|\underline{a}(\gamma p, p) - \mu(\gamma_1^{-1}) - \mu(\gamma_2^{-1})\| \leq C.$

Proof of Theorem 5.3 using Proposition 5.5: For $\gamma \in \Gamma$ and $\xi \in \Lambda$, choose $\gamma_1, \gamma_2 \in \Gamma$ as in Proposition 5.5. Then

$$\begin{aligned}\psi(\beta_\xi(\gamma p, p)) &\leq \psi(\mu(\gamma_2^{-1}) - \mu(\gamma_1)) + C\|\psi\| \\ &\leq \psi(\mu(\gamma_2^{-1}) + \mu(\gamma_1^{-1})) + C\|\psi\| \\ &\leq \psi(\underline{a}(\gamma p, p)) + 2C\|\psi\|,\end{aligned}$$

where the second inequality is valid because $\psi(\mu(\gamma_1^{\pm 1})) \geq 0$. Similarly, we get

$$\begin{aligned}\psi(\beta_\xi(\gamma p, p)) &\geq \psi(\mu(\gamma_2^{-1}) - \mu(\gamma_1)) - C\|\psi\| \\ &\geq -\psi(\mu(\gamma_2) + \mu(\gamma_1)) - C\|\psi\|.\end{aligned}$$

Since $i\mu(g^{-1}) = \mu(g)$, $i\underline{a}(p, q) = \underline{a}(q, p)$ and the norm is i -invariant, we get $\psi(\beta_\xi(\gamma p, p)) \geq \underline{\psi}(\underline{a}(p, \gamma p)) - 2C\|\psi\|$. ■

The rest of this section is devoted to a proof of Proposition 5.5 in which shadows of \mathcal{F} and $\partial\Gamma$ as well as their relationship play important roles.

Shadows in \mathcal{F}

Let $q \in X$ and $r > 0$. The shadows of the ball $B(q, r)$ viewed from $p \in X$ and $\xi \in \mathcal{F}$ are respectively defined as

$$O_r(p, q) := \{gk^+ \in \mathcal{F} : k \in K, gk \text{ int } A^+ o \cap B(q, r) \neq \emptyset\}$$

where $g \in G$ satisfies $p = g(o)$, and

$$O_r(\xi, q) := \{h^+ \in \mathcal{F} : h^- = \xi, ho \in B(q, r)\}.$$

The following two lemmas 5.6 and 5.7 are proved for $G = \text{SL}_n(\mathbb{R})$ in [54]. For $r > 0$, we set $A_r = \{a \in A : \|\log a\| \leq r\}$, and $A_r^+ = A_r \cap A^+$.

Lemma 5.6. If a sequence $q_i \in X$ converges to $\xi \in \mathcal{F}$, then for any $r > 0$, $q \in X$ and $\varepsilon > 0$, we have, for all sufficiently large i ,

$$O_{r-\varepsilon}(q_i, q) \subset O_r(\xi, q) \subset O_{r+\varepsilon}(q_i, q).$$

Proof. Since $O_r(\xi, g(o)) = gO_r(g^{-1}\xi, o) = kgO_r(e^+, o)$ for $k \in K$ with $k^+ = g^{-1}\xi$, it suffices to consider the case when $q = o$ and $\xi = e^+$. Since $q_i \rightarrow e^+$, we have $q_i = k_i a_i o$ for some $k_i \rightarrow e$ in K and $a_i \rightarrow \infty$ regularly in A^+ . As $k_i \rightarrow e$, $O_{r-\varepsilon/2}(e^+, o) \subset k_i^{-1}O_r(e^+, o) \subset O_{r+\varepsilon/2}(e^+, o)$ for all sufficiently large i . Therefore, we may assume without loss of generality that $q_i = a_i$. Let $\xi \in O_r(e^+, o)$, that is, $\xi = h^+$ for some $h \in Pw_0 \cap KA_r^+K$. Note that the sequence $a_i^{-1}h w_0^{-1}a_i$ is bounded as $a_i \in A^+$ and $h w_0^{-1} \in P$. If we write $a_i^{-1}h = \tilde{k}_i \tilde{a}_i \tilde{n}_i \in KAN$, then the following gives the KAN^+ -decomposition of $a_i^{-1}h w_0^{-1}a_i$:

$$a_i^{-1}h w_0^{-1}a_i = \tilde{k}_i \tilde{a}_i \tilde{n}_i w_0^{-1}a_i = (\tilde{k}_i w_0^{-1})(w_0 \tilde{a}_i w_0^{-1}a_i)(a_i^{-1}w_0 \tilde{n}_i w_0^{-1}a_i) \in KAN^+.$$

As the product map $K \times A \times N^+ \rightarrow G$ is a diffeomorphism, there exists $R > 1$ such that $\{w_0 \tilde{a}_i w_0^{-1}a_i \in A : i \in \mathbb{N}\} \subset A_R$. Moreover, as the sequence $a_i^{-1}w_0 \tilde{n}_i w_0^{-1}a_i$ must be bounded while $a_i \rightarrow \infty$ regularly in A^+ , it follows that $\tilde{n}_i \rightarrow e$ as $i \rightarrow \infty$. We now claim that for all sufficiently large $i \gg 1$,

$$a_i \tilde{k}_i \text{int } A^+ \cap KA_{r+\varepsilon}^+K \neq \emptyset$$

and hence $\xi = h^+ = a_i \tilde{k}_i^+ \in O_{r+\varepsilon}(a_i o, o)$. Set $b_i = w_0 a_i^{-1} w_0^{-1}$. Then

$$a_i \tilde{k}_i b_i = a_i \tilde{k}_i (w_0 a_i^{-1} w_0^{-1}) = h \tilde{n}_i^{-1} (\tilde{a}_i^{-1} w_0 a_i^{-1} w_0^{-1}).$$

Since $\tilde{n}_i \rightarrow e$ as $i \rightarrow \infty$ and $\tilde{a}_i^{-1} w_0 a_i^{-1} w_0^{-1} \in A_R$, we can find $\tilde{b}_i \in b_i A_R$ such that $a_i \tilde{k}_i \tilde{b}_i \rightarrow h$ as $i \rightarrow \infty$. On the other hand, by the strong wavefront lemma [20, Thm. 2.1], there exists a neighborhood \mathcal{O} of e in G such that $KA_r^+ K \mathcal{O} \subset KA_{r+\varepsilon}^+ K$. Since $h \in KA_r^+ K$ and $\tilde{b}_i \in \text{int } A^+$ for all large $i \gg 1$, we obtain that $a_i \tilde{k}_i \tilde{b}_i \in KA_{r+\varepsilon}^+ K$ for all sufficiently large i , proving the inclusion on the right hand side.

Now, in order to show $O_{r-\varepsilon}(a_i, o) \subset O_r(e^+, o)$, let $\eta_i \in O_{r-\varepsilon}(a_i o, o)$ be arbitrary. Since $O_{r-\varepsilon}(a_i o, o) = a_i O_{r-\varepsilon}(o, a_i^{-1} o)$, there exists $\tilde{k}_i \in K$ and $b_i \in \text{int } A^+$ such that $\tilde{k}_i^+ \in O_{r-\varepsilon}(o, a_i^{-1} o)$, $\eta_i = a_i \tilde{k}_i^+$ and $a_i \tilde{k}_i b_i \in KA_{r-\varepsilon}^+ K$. Since $a_i^{-1} o \rightarrow e^-$, it follows that $\tilde{k}_i^+ \rightarrow e^-$ as $i \rightarrow \infty$. Hence, k_i^- is in general position with e^- for all large i and hence $(a_i \tilde{k}_i b_i)^- = a_i \tilde{k}_i^- \rightarrow e^+$ as $i \rightarrow \infty$. For all large i , we have $n_i \in N$ such that $(a_i \tilde{k}_i b_i n_i)^- = e^+$ and

$d(n_i o, o) < \varepsilon$. Set $h_i := a_i \tilde{k}_i b_i n_i$. It follows that $h_i^+ = \eta_i$ and $h_i^- = e^+$. Since

$$\begin{aligned} d(h_i o, o) &= d(a_i \tilde{k}_i b_i n_i o, o) \leq d(a_i \tilde{k}_i b_i n_i o, a_i \tilde{k}_i b_i o) + d(a_i \tilde{k}_i b_i o, o) \\ &= d(n_i o, o) + d(a_i \tilde{k}_i b_i o, o) < \varepsilon + (r - \varepsilon) = r, \end{aligned}$$

we have $\eta_i = h_i^+ \in O_r(e^+, o)$ as desired. This finishes the proof. \blacksquare

The following is an analogue of Sullivan's shadow lemma:

Lemma 5.7. There exists $\kappa > 0$ such that for any $p, q \in X$ and $r > 0$, we have

$$\sup_{\xi \in O_r(p, q)} \|\beta_\xi(p, q) - \underline{a}(p, q)\| \leq \kappa r.$$

We will prove this lemma using the following:

Lemma 5.8.

(1) There exists $c_1 > 1$ such that for all $r \geq 0$,

$$KA_r^+ K \subset KA_{c_1 r} N.$$

(2) There exists $c_2 > 1$ such that for all $g \in G$ and $r \geq 0$, we have

$$\mu(gKA_r^+ K) \subset \mu(g) + \log A_{c_2 r}.$$

Proof. We use notations introduced in Lemma 3.10. Since $\chi_\alpha, \alpha \in \Pi$, form a dual basis of \mathfrak{a}^* , $\|\cdot\|_* := \sum_{\alpha \in \Pi} |\chi_\alpha(\cdot)|$ defines a norm on \mathfrak{a} . Let $k \in K$ and $a \in A_r^+$ be arbitrary. Write $ak = k'bn \in KAN$. Let $\alpha \in \Pi$. For $v_\alpha \in V_\alpha^+ \setminus \{0\}$, we have

$$\|\rho_\alpha(ak)v_\alpha\|_\alpha = \|\rho_\alpha(k'bn)v_\alpha\|_\alpha = \|\rho_\alpha(b)v_\alpha\|_\alpha = e^{\chi_\alpha(\log b)} \|v_\alpha\|_\alpha.$$

On the other hand, we have

$$e^{-\chi_\alpha(\log a)} \|v_\alpha\|_\alpha \leq \|\rho_\alpha(ak)v_\alpha\|_\alpha \leq e^{\chi_\alpha(\log a)} \|v_\alpha\|_\alpha.$$

Hence $|\chi_\alpha(\log b)| \leq |\chi_\alpha(\log a)|$ for all $\alpha \in \Pi$; so $\|\log b\|_* \leq \|\log a\|_*$. Since $\|\cdot\|_*$ and $\|\cdot\|$ are comparable, the first claim follows.

Note that $e^{\chi_\alpha(\mu(g))} = \|\rho_\alpha(g)\|_\alpha$ for $g \in G$. For all $k \in K$ and $a \in A^+$,

$$\|\rho_\alpha(gka)\|_\alpha \leq \|\rho_\alpha(g)\|_\alpha \|\rho_\alpha(k)\|_\alpha \|\rho_\alpha(a)\|_\alpha = e^{\chi_\alpha(\log a)} \|\rho_\alpha(g)\|_\alpha,$$

and similarly,

$$\|\rho_\alpha(g)\|_\alpha \leq \|\rho_\alpha(gka)\|_\alpha \|\rho_\alpha(k^{-1})\|_\alpha \|\rho_\alpha(a^{-1})\|_\alpha = e^{\chi_\alpha(\log a)} \|\rho_\alpha(gka)\|_\alpha.$$

It follows that $-\chi_\alpha(\log a) \leq \chi_\alpha(\mu(gka) - \mu(g)) \leq \chi_\alpha(\log a)$. This means that $\|\mu(gka) - \mu(g)\|_* \leq \|\log a\|_*$, finishing the proof. \blacksquare

Since $i(\mu(g)) = \mu(g^{-1})$, Lemma 5.8(2) easily implies that for all $g \in G$ and $r \geq 0$, we have

$$\mu(A_r^+ K g K A_r^+) \subset \mu(g) + \log A_{2c_2 r}. \quad (5.9)$$

Proof of Lemma 5.7. It suffices to prove the claim for $p = o$ and $q = a^{-1}o$ for $a \in A^+$. Let $\xi = k^+ \in O_r(o, a^{-1}o)$ for $k \in K$. Then there exists $b \in \text{int } A^+$ such that $d(kbo, a^{-1}o) < r$. Hence $akb \in KA_r^+ K$. Now note that $ak \in K e^{-\beta_\xi(o, a^{-1}o)} N$ by the definition of $\beta_\xi(o, a^{-1}o)$ and hence

$$akb \in K b e^{-\beta_\xi(o, a^{-1}o)} N \cap KA_r^+ K.$$

By Lemma 5.8(1), $be^{-\beta_\xi(o, a^{-1}o)} \in A_{c_1 r}$. On the other hand, $b \in K a^{-1} K A_r^+ K$, and hence $\log b \in \mu(a^{-1} K A_r^+ K)$. By Lemma 5.8(2), $b \in a^{-1} A_{c_2 r}$. Since $\underline{a}(o, a^{-1}o) = \log(a^{-1})$, the lemma is now proved. \blacksquare

Lemma 5.7 implies Theorem 5.3 for those $\xi \in O_r(\gamma p, p)$. In order to control the value of $\beta_\xi(\gamma p, p)$ when $\xi \notin O_r(\gamma p, p)$, we use the Anosov property of Γ . Let us recall some basic terminologies for hyperbolic groups for which we refer to [9] and [24].

Discrete geodesics

Let Γ be a finitely generated word hyperbolic group. We fix a finite symmetric generating subset S of Γ once and for all. Let $|\cdot| : \Gamma \rightarrow \mathbb{N} \cup \{0\}$ denote the word length associated to S . We denote by d_w the associated left-invariant word metric, that is, $d_w(\gamma_1, \gamma_2) := |\gamma_1^{-1} \gamma_2|$ for $\gamma_1, \gamma_2 \in \Gamma$.

A finite sequence $(\gamma_0, \dots, \gamma_n)$ of elements of Γ will be called a finite path if $\gamma_i^{-1}\gamma_{i+1} \in S$ for all i . Such a path will be called a geodesic segment if $|\gamma_0^{-1}\gamma_n| = n$. Infinite and bi-infinite paths can be defined analogously. They will be called geodesic rays and geodesic lines, respectively, if all of their finite subpaths are geodesic segments.

Let $\partial\Gamma$ denote the Gromov boundary of Γ , that is, $\partial\Gamma$ is the set of equivalence classes of geodesic rays, where two rays are equivalent to each other if and only if their Hausdorff distance is finite. For a geodesic ray $(\gamma_0, \gamma_1, \dots)$, we use the notation $[\gamma_0, \gamma_1, \dots]$ for its equivalence class in $\partial\Gamma$.

Let $(\cdot|\cdot)$ denote the Gromov product in the hyperbolic space Γ based at $e \in \Gamma$: $(\gamma_1|\gamma_2) := \frac{1}{2}(d_w(\gamma_1, e) + d_w(\gamma_2, e) - d_w(\gamma_1, \gamma_2))$. This extends to $\partial\Gamma$: for $x, y \in \partial\Gamma$, $(x|y) := \sup \liminf_{i,j \rightarrow \infty} (\gamma_i|\gamma'_j)$ where the supremum is taken over all sequences γ_i and γ'_j such that $x = \lim \gamma_i$ and $y = \lim \gamma'_j$. The union $\Gamma \cup \partial\Gamma$ is a compact space with the topology given as follows: a sequence $\gamma_i \in \Gamma$ converges to $x \in \partial\Gamma$ if and only if $\lim_{i \rightarrow \infty} (\gamma_i|v_i) = \infty$ for any geodesic ray (e, v_1, v_2, \dots) representing x .

For any $x, y \in \Gamma \cup \partial\Gamma$, there exists a discrete geodesic starting from x and ending at y , which may not be unique. By $[x, y]$, we mean one of those geodesics and by $[x, y)$ we mean $[x, y] - \{y\}$.

A geodesic triangle is a union of three geodesics, pairwise sharing a common endpoint in $\Gamma \cup \partial\Gamma$. Since Γ is hyperbolic, there exists $\delta = \delta(\Gamma, S) > 0$ such that for any geodesic triangle Δ , we can find a point on each edge of Δ so that the set of these triples has diameter less than δ .

Shadows in $\partial\Gamma$

For $R > 0$ and $\gamma_1, \gamma_2 \in \Gamma$, the shadow of the ball $B_R(\gamma_2)$ viewed from γ_1 is given by

$$O_R(\gamma_1, \gamma_2) = \{x \in \partial\Gamma : [\gamma_1, x] \cap B_R(\gamma_2) \neq \emptyset \text{ for some geodesic ray } [\gamma_1, x]\}.$$

Shadows satisfy the equivariance property: for any $\gamma, \gamma_1, \gamma_2 \in \Gamma$ and $R > 0$,

$$\gamma O_R(\gamma_1, \gamma_2) = O_R(\gamma\gamma_1, \gamma\gamma_2). \quad (5.10)$$

Lemma 5.10. There exist $R_0 > 1$ and $N_0 > 0$ such that the following holds: if $\gamma_1, \gamma_2 \in \Gamma$ with $|\gamma_1|, |\gamma_2| \geq N_0$ satisfies $|\gamma_1\gamma_2| = |\gamma_1| + |\gamma_2|$, then for all $R \geq R_0$,

$$O_R(\gamma_1\gamma_2, e) \cap O_R(\gamma_1\gamma_2, \gamma_1) \cap O_R(\gamma_1, e) \neq \emptyset.$$

Proof. Since $|\gamma_1\gamma_2| = |\gamma_1| + |\gamma_2|$, there exists a geodesic segment $[\gamma_1\gamma_2, e]$ passing through γ_1 , say $\alpha = (\gamma_1\gamma_2, \dots, \gamma_1, \dots, e)$. Since Γ is word hyperbolic, there exists $C > 0$ such that α lies in the C -neighborhood of some geodesic line, say $(\dots, u_{-1}, u_0, u_1, \dots)$. Set $N_0 := 4C$. Choose u_m, u_n , and u_ℓ to be elements closest to $\gamma_1\gamma_2, \gamma_1$, and e , respectively.

We claim that $|m - \ell| \geq \max(|m - n|, |n - \ell|)$. By the triangle inequality,

$$|n - \ell| = d_w(u_n, u_\ell) \leq d_w(\gamma_1, e) + 2C = |\gamma_1| + 2C;$$

$$|m - n| = d_w(u_m, u_n) \leq d_w(\gamma_1\gamma_2, \gamma_1) + 2C = |\gamma_2| + 2C.$$

Since $|\gamma_1\gamma_2| = |\gamma_1| + |\gamma_2|$ and $|\gamma_1\gamma_2| \leq d_w(u_m, u_\ell) + 2C = |m - \ell| + 2C$, it follows that

$$\begin{aligned} |\gamma_2| - 2C &\geq \max(|\gamma_1|, |\gamma_2|) - 2C + N_0 \\ &= \max(|\gamma_1|, |\gamma_2|) + 2C \\ &\geq \max(|n - \ell|, |m - n|). \end{aligned}$$

This proves the claim. ■

Now possibly after flipping the geodesic, we may assume that $m \leq \ell$. Then the claim implies that $\ell - m = |m - n| + |n - \ell|$ and hence $m \leq n \leq \ell$. Set $x := [u_0, u_1, u_2, \dots] \in \partial\Gamma$. Choose geodesic rays $[\gamma_1\gamma_2, x)$ and $[\gamma_1, x)$. Since the Hausdorff distance between $[\gamma_1\gamma_2, x)$ and the ray (u_m, u_{m+1}, \dots) is at most $d_w(\gamma_1\gamma_2, u_m) + \delta \leq C + \delta$, it follows that there exist $v_1, v_2 \in \Gamma$ lying on $[\gamma_1\gamma_2, x]$ such that $d_w(u_n, v_1) < C + \delta$ and $d_w(u_\ell, v_2) < C + \delta$. Since the Hausdorff distance between $[\gamma_1, x)$ and the ray (u_n, u_{n+1}, \dots) is at most $d_w(\gamma_1, u_n) + \delta < C + \delta$, there exists $v_3 \in \Gamma$ lying on $[\gamma_1, x)$ such that $d_w(u_\ell, v_3) < C + \delta$. These altogether imply that

$$x \in O_{2C+\delta}(\gamma_1\gamma_2, e) \cap O_{2C+\delta}(\gamma_1\gamma_2, \gamma_1) \cap O_{2C+\delta}(\gamma_1, e).$$

In the rest of this section, we assume that Γ is an Anosov subgroup of G . The following Morse property of Kapovich–Leeb–Porti [25, Prop. 5.16] says that a discrete geodesic line (resp. ray) of Γ is contained in a uniform neighborhood of some A -orbit (resp. A^+ -orbit) in X .

Proposition 5.12 (Morse property). For any Anosov subgroup $\Gamma < G$, there exists $R_1 > 0$ such that

(1) If $(\cdots, \gamma_{-1}, \gamma_0, \gamma_1, \cdots)$ is a geodesic line in (Γ, d_w) , then

$$\sup_{k \in \mathbb{Z}} d(\gamma_k o, gA o) \leq R_1$$

for any $g \in G$ such that $g^+ = \zeta([\gamma_0, \gamma_1, \cdots]), g^- = \zeta([\gamma_0, \gamma_{-1}, \cdots])$.

(2) If $(\gamma_0, \gamma_1, \cdots)$ is a geodesic ray in (Γ, d_w) , then

$$\sup_{k \in \mathbb{N}} d(\gamma_k o, gA^+ o) \leq R_1,$$

where $g \in \gamma_0 K$ is the unique element satisfying $g^+ = \zeta([\gamma_0, \gamma_1, \cdots])$.

Using this proposition, we will show that shadows in the Gromov boundary $\partial\Gamma$ are mapped to shadows in the Furstenberg boundary \mathcal{F} by the limit map $\zeta : \partial\Gamma \rightarrow \Lambda$ (Proposition 5.14). We will need the following lemma:

Lemma 5.13. There exists $C > 0$ such that for all $\gamma \in \Gamma$, $\|\mu(\gamma)\| \leq C|\gamma|$. In particular, $d(o, \gamma o) \leq Cd_w(e, \gamma)$.

Proof. We use notations from Lemma 3.10. Consider the norm $\|\cdot\|_* := \sum_{\alpha \in \Pi} |\chi_\alpha(\cdot)|$ on \mathfrak{a} . Let $\gamma \in \Gamma$ be arbitrary, and write $\gamma = s_1 \cdots s_\ell$ with $s_i \in S$ and $\ell = |\gamma|$. Since $\chi_\alpha(\mu(g)) = \log \|\rho_\alpha(g)\|_\alpha$ for all $g \in G$ and $\|\rho_\alpha(s_1 \cdots s_\ell)\|_\alpha \leq \|\rho_\alpha(s_1)\|_\alpha \cdots \|\rho_\alpha(s_\ell)\|_\alpha$, it follows that for each $\alpha \in \Pi$,

$$\chi_\alpha(\mu(s_1 \cdots s_\ell)) \leq \chi_\alpha(\mu(s_1)) + \cdots + \chi_\alpha(\mu(s_\ell)).$$

Noting that χ_α is positive on \mathfrak{a}^+ , we have

$$\begin{aligned} \|\mu(\gamma)\|_* &= \sum_{\alpha \in \Pi} |\chi_\alpha(\mu(\gamma))| = \sum_{\alpha \in \Pi} \chi_\alpha(\mu(\gamma)) \\ &\leq \sum_{\alpha \in \Pi} (\chi_\alpha(\mu(s_1)) + \cdots + \chi_\alpha(\mu(s_\ell))) \leq C|\gamma|, \end{aligned}$$

where $C := \max \{ \sum_{\alpha \in \Pi} \chi_\alpha(\mu(s)) : s \in S \}$. Since $\|\cdot\|$ and $\|\cdot\|_*$ are comparable, this finishes the proof. ■

Proposition 5.14 (Shadows go to shadows). There exists $c > 0$ such that for all $R > 1$ and $\gamma, \gamma' \in \Gamma$,

$$\zeta(O_R(\gamma', \gamma)) \subset O_{cR}(\gamma' o, \gamma o).$$

Proof. By (5.10), it suffices to consider the case $\gamma' = e$. Let $x \in O_R(e, \gamma)$. By the definition of $O_R(e, \gamma)$, there exists a geodesic ray $(\gamma'_0 = e, \gamma'_1, \gamma'_2, \dots)$ representing x such that $d_w(\gamma'_m, \gamma) < R$ for some $m \in \mathbb{N}$. Let $R_1 > 0$ be the constant from Proposition 5.12, and $k \in K$ be an element such that $k^+ = \zeta([e, \gamma'_1, \gamma'_2, \dots])$. Then by Proposition 5.12(2), there exists $a \in A^+$ such that $d(\gamma'_m o, kao) \leq R_1$. By Lemma 5.13, we have

$$d(\gamma o, \gamma'_m o) = \|\mu(\gamma^{-1} \gamma'_m)\| < C d_w(\gamma, \gamma'_m) < CR.$$

Therefore,

$$d(\gamma o, kao) \leq d(\gamma o, \gamma'_m o) + d(\gamma'_m o, kao) \leq CR + R_1.$$

This implies that $\zeta(x) \in O_{CR+R_1}(o, \gamma o)$. Since $R > 1$, the conclusion follows by setting $c := C + R_1$. \blacksquare

Corollary 5.15. There exists $R_2 > 0$ such that for all $\gamma_1, \gamma_2 \in \Gamma$ with $|\gamma_1 \gamma_2| = |\gamma_1| + |\gamma_2|$, we have

$$\|\mu(\gamma_1 \gamma_2) - \mu(\gamma_1) - \mu(\gamma_2)\| \leq R_2.$$

Proof. Let N_0 and R_0 be given by Lemma 5.10. If one of $|\gamma_1|, |\gamma_2|$ is less than N_0 , then the claim holds by Lemma 2.3. Now assume that $|\gamma_1|, |\gamma_2| \geq N_0$. Then by Lemma 5.10 and Proposition 5.14, we can choose

$$\xi \in O_{cR_0}(\gamma_1 \gamma_2 o, o) \cap O_{cR_0}(\gamma_1 \gamma_2 o, \gamma_1 o) \cap O_{cR_0}(\gamma_1 o, o),$$

where c is as in Proposition 5.14. By Lemma 5.7 and the cocycle identity

$$\beta_\xi(\gamma_1 \gamma_2 o, o) = \beta_\xi(\gamma_1 \gamma_2 o, \gamma_1 o) + \beta_\xi(\gamma_1 o, o),$$

we have

$$\|\underline{a}(\gamma_1 \gamma_2 o, o) - \underline{a}(\gamma_1 o, o) - \underline{a}(\gamma_2 o, o)\| \leq 3\kappa c R_0.$$

Since $\underline{a}(go, o) = i\mu(g)$ for all $g \in G$ and i preserves $\|\cdot\|$,

$$\|\mu(\gamma_1\gamma_2) - \mu(\gamma_1) - \mu(\gamma_2)\| \leq 3\kappa cR_0.$$

■

Proof of Proposition 5.5. We may assume that $p = o$ by Lemma 2.3. Let $\gamma \in \Gamma$ and $\xi \in \Lambda$ be arbitrary. If $\gamma = \gamma_1\gamma_2$, we have

$$\beta_\xi(\gamma o, o) = \beta_\xi(\gamma o, \gamma_1 o) - \beta_\xi(o, \gamma_1 o).$$

We claim that we can find $\gamma_1, \gamma_2 \in \Gamma$ so that $\gamma = \gamma_1\gamma_2$, $|\gamma| = |\gamma_1| + |\gamma_2|$, and

$$\xi \in O_{c(\delta+1)}(\gamma o, \gamma_1 o) \cap O_{c(\delta+1)}(o, \gamma_1 o) \quad (5.16)$$

where $c > 0$ is as in Proposition 5.14.

If $\xi \in O_{c(\delta+1)}(o, \gamma o)$, then we may simply set $\gamma_1 = \gamma$ and $\gamma_2 = e$. In general, we find γ_1 as follows. Consider a geodesic triangle Δ whose vertices are $e, \gamma \in \Gamma$, and $\zeta^{-1}(\xi) \in \partial\Gamma$. Since Γ is hyperbolic, we can find three points on Δ , one on each edge, whose diameter is less than δ . Let $\gamma_1 \in \Gamma$ be the point on the geodesic segment joining e and γ , and set $\gamma_2 := \gamma_1^{-1}\gamma$. We then have $|\gamma| = |\gamma_1| + |\gamma_2|$, and $\zeta^{-1}(\xi) \in O_\delta(\gamma, \gamma_1) \cap O_\delta(e, \gamma_1)$. Now the claim follows from Proposition 5.14.

Therefore, by Lemma 5.7,

$$\max(\|\beta_\xi(\gamma o, \gamma_1 o) - \mu(\gamma_2^{-1})\|, \|\beta_\xi(o, \gamma_1 o) - \mu(\gamma_1)\|) \leq \kappa c(\delta + 1)$$

and hence

$$\|\beta_\xi(\gamma o, o) + \mu(\gamma_1) - \mu(\gamma_2^{-1})\| \leq 2\kappa c(\delta + 1).$$

Since $|\gamma| = |\gamma_1| + |\gamma_2|$ and S is symmetric, we have $|\gamma^{-1}| = |\gamma_1^{-1}| + |\gamma_2^{-1}|$. As $\underline{a}(\gamma o, o) = \mu(\gamma^{-1})$, we have, by Corollary 5.15,

$$\|\underline{a}(\gamma o, o) - \mu(\gamma_1^{-1}) - \mu(\gamma_2^{-1})\| \leq R_2.$$

Hence it suffices to set $C := \max(2\kappa c(\delta + 1), R_2)$. ■

6 Virtual Visual Metrics via ψ -Gromov Product

In this section, we let $\Gamma < G$ be an Anosov subgroup, and fix $\psi \in D_\Gamma^*$. The main aim here is to show that exponentiating the following ψ -Gromov product defines a virtual visual metric on Λ up to a small power.

Definition 6.1. The ψ -Gromov product based at o is a function $\mathcal{F}^{(2)} \rightarrow \mathbb{R}$ defined as follows: for any $(\xi_1, \xi_2) \in \mathcal{F}^{(2)}$,

$$[\xi_1, \xi_2]_{\psi, o} := \psi(\mathcal{G}(\xi_1, \xi_2)),$$

where \mathcal{G} is the \mathfrak{a} -valued Gromov product defined in Definition 3.8. For $p = g(o) \in X$, we set

$$[\xi_1, \xi_2]_{\psi, p} := [g^{-1}\xi_1, g^{-1}\xi_2]_{\psi, o}.$$

For simplicity, we set $[\xi_1, \xi_2]_p := [\xi_1, \xi_2]_{\psi, p}$.

Define $d_p = d_{\psi, p} : \mathcal{F}^{(2)} \rightarrow \mathbb{R}_{\geq 0}$ by

$$d_p(\xi_1, \xi_2) = e^{-[\xi_1, \xi_2]_p}. \quad (6.2)$$

It follows from (3.9) that for all $g \in G$, $p \in X$, and $(\xi_1, \xi_2) \in \mathcal{F}^{(2)}$, we have

$$d_{gp}(\xi_1, \xi_2) = e^{-\psi(\beta_{\xi_1}(gp, p) + i\beta_{\xi_2}(gp, p))} d_p(\xi_1, \xi_2) = d_p(g^{-1}\xi_1, g^{-1}\xi_2). \quad (6.3)$$

We set $[\xi, \xi]_p = +\infty$ and $d_p(\xi, \xi) = 0$ for all $\xi \in \mathcal{F}$. By the antipodal property (4.2), $[\cdot, \cdot]_p$ and d_p are defined on all of $\Lambda \times \Lambda$. The following is the main theorem of this section:

Theorem 6.4. Fix $p \in X$. For all sufficiently small $\varepsilon > 0$, there exist a metric $d_\varepsilon = d_\varepsilon(p)$ on Λ and a constant $C_\varepsilon = C_\varepsilon(p) > 0$ such that for all $\xi_1, \xi_2 \in \Lambda$,

$$C_\varepsilon^{-1} d_{\psi, p}(\xi_1, \xi_2)^\varepsilon \leq d_\varepsilon(\xi_1, \xi_2) \leq C_\varepsilon d_{\psi, p}(\xi_1, \xi_2)^\varepsilon.$$

This is an analogue of [9, Part III, Prop. 3.21] for Gromov hyperbolic spaces.

Weak ultrametric inequality

A well-known construction [19, Section 7.3] shows the existence of a metric in Theorem 6.4, provided there exists $C > 0$ such that for all $\xi_1, \xi_2, \xi_3 \in \Lambda$, we have

- (1) (weak symmetry) $d_p(\xi_1, \xi_2) \leq e^C d_p(\xi_2, \xi_1)$;
- (2) (weak ultrametric inequality) $d_p(\xi_1, \xi_3) \leq e^C \max(d_p(\xi_1, \xi_2), d_p(\xi_2, \xi_3))$.

Hence Theorem 6.4 follows from the following proposition:

Proposition 6.5. There exists $C = C(p) > 0$ such that for all $\xi_1, \xi_2, \xi_3 \in \Lambda$, we have

$$\begin{aligned} [\xi_1, \xi_2]_p &\geq [\xi_2, \xi_1]_p - C; \\ [\xi_1, \xi_3]_p &\geq \min([\xi_1, \xi_2]_p, [\xi_2, \xi_3]_p) - C. \end{aligned}$$

In the case of $X = \mathbb{H}^2$, the classical Gromov product satisfies that there exists a uniform constant $C > 0$ such that for any $x, y \in \partial\mathbb{H}^2$,

$$|\mathcal{G}(x, y) - 2d(o, z)| \leq C$$

where z is the unique projection of o to the geodesic connecting x and y . In the following lemma 6.6, we establish the analogous property for α -valued Gromov products on $\Lambda \times \Lambda$ using the Morse property of Anosov groups.

For $\gamma \in \Gamma$ and any geodesic segment α in Γ , we define the set of projections of γ to α by

$$\pi_\alpha(\gamma) := \{\gamma' \in \alpha : d_w(\gamma, \gamma') = d_w(\gamma, \alpha)\}.$$

Since Γ is hyperbolic, the diameter of $\pi_\alpha(\gamma)$ is less than 4δ .

Lemma 6.6. There exists $C_1 > 0$ such that for any $x \neq y$ in $\partial\Gamma$ and $\gamma \in \pi_{[x,y]}(e)$, we have

$$\|\mathcal{G}(\zeta(x), \zeta(y)) - (\mu(\gamma) + i\mu(\gamma))\| \leq C_1.$$

In particular, \mathcal{G} is almost symmetric on Λ : for any $\xi_1 \neq \xi_2 \in \Lambda$,

$$\|\mathcal{G}(\xi_1, \xi_2) - \mathcal{G}(\xi_2, \xi_1)\| \leq 2C_1.$$

Proof. Let $\alpha := (u_0 = e, u_1, u_2, \dots)$ and $\alpha' := (v_0 = e, v_1, v_2, \dots)$ be geodesic representatives of x and y , respectively. Let $\gamma \in \pi_{[x,y]}(e)$ be arbitrary, and $f, g, h \in G$ be elements satisfying the following:

- $f(o) = o$ and $f^+ = \zeta(x)$;
- $g(o) = o$ and $g^+ = \zeta(y)$;
- $h^+ = \zeta(x)$ and $h^- = \zeta(y)$.

Applying Proposition 5.12(1) to the geodesic line $[x, y]$, we have

$$d(ho, \gamma o) < R_1$$

after replacing h with some element of hA . Hence by Lemma 2.3, there exists $C' = C'(R_1) > 0$ such that

$$\|\mu(h) - \mu(\gamma)\| \leq C'. \quad (6.7)$$

Noting

$$\mathcal{G}(\zeta(x), \zeta(y)) = \mathcal{G}(h^+, h^-) = \beta_{h^+}(o, ho) + i\beta_{h^-}(o, ho),$$

it is now sufficient to show that for some uniform constant $C_1 > 0$,

$$\|\beta_{h^+}(o, ho) - \mu(h)\| \leq C_1 \quad \text{and} \quad \|\beta_{h^-}(o, ho) - \mu(h)\| \leq C_1.$$

By Lemma 5.7, this claim follows if we show

$$h^+, h^- \in O_R(o, ho) \quad (6.8)$$

for some uniform constant $R > 0$.

Since Γ is hyperbolic, the diameter of the set $\pi_{\alpha'}(x) \cup \pi_{\alpha}(y) \cup \pi_{[x,y]}(e)$ is at most $C\delta$ for some uniform constant $C > 1$. In particular, we can find $k, \ell \in \mathbb{N}$ such that the set $\{u_k, v_\ell, \gamma\}$ has diameter less than $C\delta$. Applying Proposition 5.12(2) to the geodesic ray α , we find $a_1 \in A^+$ such that

$$d(fa_1 o, u_k o) < R_1.$$

Since $d_w(u_k, \gamma) = |u_k^{-1}\gamma| \leq C\delta$, we have

$$d(u_k o, \gamma o) = \|\mu(u_k^{-1}\gamma)\| \leq \sup\{\|\mu(\gamma')\| : |\gamma'| \leq C\delta\}.$$

Therefore,

$$\begin{aligned} d(fa_1 o, ho) &\leq d(fa_1 o, u_k o) + d(u_k o, \gamma o) + d(\gamma o, ho) \\ &\leq 2R_1 + \sup\{\|\mu(\gamma')\| : |\gamma'| \leq C\delta\}. \end{aligned}$$

Setting $R := 2R_1 + \sup\{\|\mu(\gamma')\| : |\gamma'| \leq C\delta\}$, it follows that $h^+ = f^+ \in O_R(o, ho)$. Similar argument shows that $h^- = g^+ \in O_R(o, ho)$. This proves (6.8). \blacksquare

Lemma 6.9. For any compact subset $C \subset X$, the set $\{\beta_\xi(p, o) : \xi \in \mathcal{F}, p \in C\}$ is bounded.

Proof. This follows from Lemma 5.1 by setting $\psi = \sum_{\alpha \in \Pi} \varpi_\alpha$. \blacksquare

Proof of Proposition 6.5. Observe that the identity (6.3) gives that for any $\xi_1 \neq \xi_2 \in \Lambda$,

$$[\xi_1, \xi_2]_p - [\xi_1, \xi_2]_o = \psi(\beta_{\xi_1}(p, o) + i\beta_{\xi_2}(p, o)).$$

Now Lemma 6.9 shows the existence of $C = C(p, \psi) > 0$ such that $|[\xi_1, \xi_2]_p - [\xi_1, \xi_2]_o| \leq C$. Therefore it suffices to show the claim for $p = o$. The first inequality is an immediate consequence of Lemma 6.6 with $C > 2C_1\|\psi\|$.

To show the second inequality, let $C_1 > 0$ be a constant from Lemma 6.6 so that we have

$$[\xi_1, \xi_3]_o \geq \psi(\mu(\gamma_2) + i\mu(\gamma_2)) - C_1\|\psi\|. \quad (6.9)$$

Set $x_i := \xi^{-1}(\xi_i) \in \partial\Gamma$ for $i = 1, 2, 3$. For each i , we fix a geodesic line $[x_i, x_{i+1}]$ joining x_i and x_{i+1} , and choose $\gamma_{i+2} \in \pi_{[x_i, x_{i+1}]}(e)$, where all the indices are to be interpreted mod 3. By the hyperbolicity of Γ , for some uniform constant $C > 0$, there exists $1 \leq i \leq 3$ such that $d_w(\gamma_i, \gamma_{i+1}) < C\delta$ and for some $\gamma' \in [e, \gamma_{i+2}]$, the diameter of $\{\gamma', \gamma_i, \gamma_{i+1}\}$ is at most $C\delta$.

We first consider the case when $i = 1$. Since

$$d(\gamma_1 o, \gamma_2 o) \leq d_w(\gamma_1, \gamma_2) \max_{s \in S} d(o, s) < C\delta \max_{s \in S} d(o, s),$$

it follows from Lemma 2.3 that for some uniform $C_2 > 0$,

$$\|\mu(\gamma_1) - \mu(\gamma_2)\| \leq C_2.$$

In view of (6.9), we now obtain

$$\begin{aligned} [\xi_1, \xi_3]_o &\geq \psi(\mu(\gamma_1) + i\mu(\gamma_1)) - C_1\|\psi\| - 2C_2 \\ &\geq [\xi_2, \xi_3]_o - 2C_1\|\psi\| - 2C_2 \text{ by Lemma 6.6} \\ &\geq \min([\xi_1, \xi_2]_o, [\xi_2, \xi_3]_o) - 2C_1\|\psi\| - 2C_2. \end{aligned}$$

The case $i = 2$ can be handled similarly by interchanging the roles of γ_2 and γ_3 . Finally, in the case when $i = 3$, let R_2 be as in Corollary 5.15. Since $(e, \dots, \gamma', \dots, \gamma_2)$ is a geodesic, we have by Corollary 5.15 that

$$\|\mu(\gamma_2) - \mu(\gamma') - \mu(\gamma'^{-1}\gamma_2)\| \leq R_2.$$

By (6.9) and the fact $\psi(\mu((\gamma'^{-1}\gamma_2)^{\pm 1})) \geq 0$, we deduce

$$\begin{aligned} [\xi_1, \xi_3]_o &\geq \psi(\mu(\gamma') + i\mu(\gamma')) - C_1\|\psi\| - 2R_2\|\psi\| \\ &\geq \psi(\mu(\gamma_1) + i\mu(\gamma_1)) - (C_1 + 2C_2 + 2R_2)\|\psi\|, \end{aligned}$$

as the diameter of $\{\gamma', \gamma_1, \gamma_3\}$ is less than δ . The rest of the proof is similar to the case $i = 1$. ■

Covering lemma

Using Theorem 6.4, we obtain:

Lemma 6.11 (Triangle inequality). There exists $N = N(\psi, p) \geq 1$ such that for any $\xi_1, \xi_2, \xi_3 \in \Lambda$,

$$d_p(\xi_1, \xi_3) \leq N(d_p(\xi_1, \xi_2) + d_p(\xi_2, \xi_3)).$$

In particular, $d_p(\xi_1, \xi_2) \leq Nd_p(\xi_2, \xi_1)$. Moreover, $N(\psi, p)$ can be taken uniformly for all p in a fixed compact subset of X .

Proof. Choose $\varepsilon > 0$ sufficiently small so that Theorem 6.4 holds, and set $d := d_\varepsilon$, $C := C_\varepsilon$. We then have

$$d_p(\xi_1, \xi_3)^\varepsilon \leq Cd(\xi_1, \xi_3) \leq C(d(\xi_1, \xi_2) + d(\xi_2, \xi_3)) \leq C^2(d_p(\xi_1, \xi_2)^\varepsilon + d_p(\xi_2, \xi_3)^\varepsilon).$$

Since $(a^\varepsilon + b^\varepsilon)^{1/\varepsilon} \leq \alpha(a + b)$ for all $a, b \geq 0$ for some uniform constant $\alpha = \alpha(\varepsilon) > 0$, it suffices to take the $1/\varepsilon$ power in each side of the above. Now the second part follows from (6.3) and Lemma 6.9. \blacksquare

For $\xi \in \Lambda$ and $r > 0$, set

$$\mathbb{B}_p(\xi, r) := \{\eta \in \Lambda : d_{\psi, p}(\xi, \eta) < r\}.$$

Lemma 6.12 (Covering lemma). There exists $N_0(\psi, p) \geq 1$ satisfying the following: for any finite collection $\mathbb{B}_p(\xi_1, r_1), \dots, \mathbb{B}_p(\xi_n, r_n)$ with $\xi_i \in \Lambda$ and $r_i > 0$, there exists a disjoint subcollection $\mathbb{B}_p(\xi_{i_1}, r_{i_1}), \dots, \mathbb{B}_p(\xi_{i_\ell}, r_{i_\ell})$ such that

$$\mathbb{B}_p(\xi_1, r_1) \cup \dots \cup \mathbb{B}_p(\xi_n, r_n) \subset \mathbb{B}_p(\xi_{i_1}, 3N_0 r_{i_1}) \cup \dots \cup \mathbb{B}_p(\xi_{i_\ell}, 3N_0 r_{i_\ell}).$$

Moreover, $N_0(\psi, p)$ can be taken uniformly for all p in a fixed compact subset of X .

Proof. Let $N = N(\psi, p)$ be as given by Lemma 6.11. For simplicity, set $B_i := \mathbb{B}_p(\xi_i, r_i)$. We may assume $r_1 \geq \dots \geq r_n$ without loss of generality and define inductively

$$i_1 = 1, i_{j+1} = \min\{i > i_j : B_i \cap (B_{i_1} \cup \dots \cup B_{i_j}) = \emptyset\},$$

as long as possible, to obtain a maximal disjoint subcollection $\{B_{i_1}, \dots, B_{i_\ell}\}$. Let $\xi \in B_j$ for some $1 \leq j \leq n$. Then there exists $1 \leq k \leq \ell$ such that $B_j \cap B_{i_k} \neq \emptyset$ and $r_{i_k} \geq r_j$. Choose $\eta \in B_j \cap B_{i_k}$. Then by Lemma 6.11, we have $d_p(\eta, \xi) \leq N(d_p(\eta, \xi_j) + d_p(\xi_j, \xi)) < 2N^2 r_j \leq 2N^2 r_{i_k}$ and $d_p(\xi_{i_k}, \eta) < r_{i_k}$. Hence

$$d_p(\xi_{i_k}, \xi) \leq N(d_p(\xi_{i_k}, \eta) + d_p(\eta, \xi)) < 3N^3 r_{i_k}.$$

Hence it suffices to set $N_0 := N^3$. \blacksquare

Comparing Gromov products

Although we will not be using it in the rest of the paper, we record the following theorem, which is of independent interest:

Theorem 6.13. For any $\psi \in D_\Gamma^*$, there exist $c_1 = c_1(\psi) \geq 1, c_2 = c_2(\psi) > 0$ such that for all $x \neq y \in \partial\Gamma$,

$$c_1^{-1}(x|y) - c_2 \leq \psi(\mathcal{G}(\zeta(x), \zeta(y))) \leq c_1(x|y) + c_2.$$

Note that if $\gamma \in \pi_{[x,y]}(e)$ for $x \neq y$ in $\partial\Gamma$, then $|(x|y) - |\gamma|| \leq C$ for some uniform constant $C > 0$ (cf. [9]). Given this fact, Theorem 6.13 follows immediately from Lemma 6.6 and the following lemma:

Lemma 6.14. For any $\psi \in D_\Gamma^*$, there exist constants $C_\psi, c_\psi > 0$ such that for all $\gamma \in \Gamma$,

$$C_\psi^{-1}|\gamma| - c_\psi \leq \psi(\mu(\gamma)) \leq C_\psi|\gamma|.$$

Proof. Since $\psi > 0$ on \mathcal{L}_Γ , we have

$$0 < d := \min_{\|u\|=1, u \in \mathcal{L}_\Gamma} \psi(u) \leq D := \max_{\|u\|=1, u \in \mathcal{L}_\Gamma} \psi(u) < \infty.$$

Hence $d\|\mu(\gamma)\| \leq \psi(\mu(\gamma)) \leq D\|\mu(\gamma)\|$ for all $\gamma \in \Gamma$. So the upper bound follows from Lemma 5.13, and the lower bound follows from the well-known property of Anosov groups that for some uniform $C > 0$, $C^{-1}|\gamma| - C \leq \|\mu(\gamma)\|$ for all $\gamma \in \Gamma$ [22]. \blacksquare

7 Conical points, Divergence Type, and Classification of PS Measures

In this section, we show that for Anosov groups, the space of all PS-measures on Λ is homeomorphic to D_Γ^* .

Conical limit points

For a discrete subgroup $\Gamma < G$ and $x \in \Gamma \backslash G$, we mean by $\limsup xA^+M$ the set of all limit points $\lim_{i \rightarrow \infty} xa_i m_i$ where $a_i \rightarrow \infty$ in A^+ and $m_i \in M$.

Definition 7.1 (Conical limit points). We call $\xi \in \mathcal{F}$ a conical limit point of Γ if $\limsup \Gamma g A^+ M \neq \emptyset$ for some $g \in G$ with $g^+ = \xi$. Equivalently, $\xi \in \mathcal{F}$ is conical if there

exists $R > 0$ such that $\xi \in O_R(o, \gamma_i o)$ for some sequence $\gamma_i \rightarrow \infty$ in Γ . We denote by Λ_c the set of all conical limit points of Γ .

Lemma 7.2. Let Γ be a Zariski dense discrete subgroup and $\mathfrak{c} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ be a closed convex cone whose interior contains $\mathcal{L}_\Gamma - \{0\}$. If $\gamma_i g_i a_i$ is a bounded sequence where $g_i \in G$ is bounded, $\gamma_i \in \Gamma$ and $a_i \rightarrow \infty$ in A^+ , then

$$\log a_i \in \mathfrak{c} \quad \text{for all sufficiently large } i. \quad (7.1)$$

In particular, for any $x \in \Gamma \backslash G$, $\limsup x A^+ M$ coincides with the set

$$\{\lim_{i \rightarrow \infty} x a_i m_i : m_i \in M, \log a_i \rightarrow \infty \text{ in } \mathfrak{c}\}.$$

Proof. As g_i and $\gamma_i g_i a_i$ are bounded sequences, the sequence $\mu(\gamma_i^{-1}) - \log a_i$ is also bounded by Lemma 2.3. Hence $\lim_{i \rightarrow \infty} \frac{\log a_i}{\|\log a_i\|}$ belongs to the asymptotic cone of $\mu(\Gamma)$, which is equal to \mathcal{L}_Γ [6, Thm. 1.2]. Since $\mathcal{L}_\Gamma - \{0\} \subset \text{int } \mathfrak{c}$, it follows that $\log a_i \in \mathfrak{c}$ for all large \blacksquare

We note that for $\mathfrak{c} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ as above, there exists a constant $s = s(\mathfrak{c}) > 0$ such that for any sequence $v_i \rightarrow \infty$ in \mathfrak{c} ,

$$\min_{\alpha \in \Pi} \liminf_{i \rightarrow \infty} \alpha\left(\frac{v_i}{\|v_i\|}\right) \geq s.$$

We deduce from Proposition 5.12: recall $\mathcal{E} = \{[g] \in \Gamma \backslash G : g^+ \in \Lambda\}$.

Proposition 7.4. For Γ Anosov, there exist a compact subset \mathcal{Q} of \mathcal{E} and a closed convex cone $\mathfrak{c} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ such that for any $x \in \mathcal{E}$, there exists $\log a_i \rightarrow \infty$ in \mathfrak{c} such that

$$x a_i \in \mathcal{Q} \quad \text{for all } i \geq 1.$$

In particular, $\Lambda = \Lambda_c$.

Proof. For Γ Anosov, we have $\mathcal{L}_\Gamma - \{0\} \subset \text{int } \mathfrak{a}^+$ by Theorem 4.3. Hence we can find a closed convex cone $\mathfrak{c} \subset \text{int } \mathfrak{a}^+ \cup \{0\}$ such that $\mathcal{L}_\Gamma \subset \text{int } \mathfrak{c} \cup \{0\}$. We first check that $\Lambda_c \subset \Lambda$. Let $g^+ \in \Lambda_c$ for some $g \in G$. Then there exists $\gamma_i \in \Gamma$ and $a_i m_i \rightarrow \infty$ in $A^+ M$ such that $\gamma_i g a_i m_i$ is bounded. By Lemma 7.2, it follows that $\log a_i \rightarrow \infty$ in \mathfrak{c} . In particular, $a_i \rightarrow \infty$ regularly in A^+ . Hence by Lemma 2.11, $g a_i o \rightarrow g^+$ as $i \rightarrow \infty$. Since $d(g a_i o, \gamma_i^{-1} o)$ is bounded, $\gamma_i^{-1} o \rightarrow g^+$ as $i \rightarrow \infty$. By Lemma 2.13, $g^+ \in \Lambda$.

Let $g^+ = \xi \in \Lambda$ and $z \in \partial\Gamma$ be such that $\xi = \zeta(z)$. Choose a geodesic ray $r = (\gamma_0 = e, \gamma_1, \gamma_2, \dots)$ representing z . Note that if $g^+ = h^+$, then for any sequence $a_i \rightarrow \infty$ in A^+ , there exists $b_i \in A^+$ such that $d(ga_i o, hb_i o) \leq 1$ for all sufficiently large i . Hence we may assume that $g \in K$ by replacing g by an element of gP . By Proposition 5.12, $\gamma_i o$ is contained in the R_1 -neighborhood of $ga^+ o$, with R_1 given therein. Hence for some $a_i \rightarrow \infty$ in A^+ , $\Gamma \backslash \Gamma g a_i \in \mathcal{Q}$ where $\mathcal{Q} = \Gamma \backslash \Gamma \{h \in G : d(o, ho) \leq R_1\} \cap \mathcal{E}$. Hence $g^+ \in \Lambda_c$. Moreover, by Lemma 7.2, $\log a_i \in \mathfrak{c}$ for all sufficiently large i . This finishes the proof. \blacksquare

Classification of PS measures on Λ

Lemma 7.5. Let $\psi_i \in \mathfrak{a}^*$ and ν_{ψ_i} be a (Γ, ψ_i) -PS measure for $i = 1, 2$. If $\nu_{\psi_1} = \nu_{\psi_2}$, then $\psi_1 = \psi_2$.

Proof. Suppose that $\nu_{\psi_1} = \nu_{\psi_2}$. Then for all $\gamma \in \Gamma$ and $\xi \in \Lambda$, we have

$$\psi_1(\beta_\xi(e, \gamma)) = \psi_2(\beta_\xi(e, \gamma)).$$

By setting $\xi = \gamma_\gamma$, we obtain $\lambda(\gamma) \in \ker(\psi_1 - \psi_2)$ for all $\gamma \in \Gamma$, by Lemma 3.5. Hence $\mathcal{L}_\Gamma \subset \ker(\psi_1 - \psi_2)$. Since \mathcal{L}_Γ has nonempty interior [6, Thm. 1.2], this implies that $\psi_1 = \psi_2$. \blacksquare

Remark 7.6. When Γ is an Anosov subgroup, ν_{ψ_1} and ν_{ψ_2} are even mutually singular to each other whenever $\psi_1 \neq \psi_2$ (See Theorem 10.20 below).

We denote by \mathcal{S}_Γ the space of all PS measures on Λ . Recall that for $\psi \in D_\Gamma^*$, Quint constructed a (Γ, ψ) -PS measure on Λ [40]. In the Anosov case, such a measure is unique, which we denote by ν_ψ . By Lemma 7.5, the map $\psi \mapsto \nu_\psi$ from D_Γ^* to \mathcal{S}_Γ is injective.

Theorem 7.7. For $\Gamma < G$ Anosov, the map $\psi \mapsto \nu_\psi$ is a homeomorphism between D_Γ^* and \mathcal{S}_Γ .

In the rank one case, there exists a unique Patterson–Sullivan measure on Λ and its dimension is given by the critical exponent of Γ . The above theorem generalizes such phenomenon.

To prove that the map $\psi \mapsto \nu_\psi$ is surjective, we need the following shadow lemma. It was first presented in [2, Thm. 3.3] and then in [40, Thm. 8.2] in slightly different forms.

Lemma 7.8 (Size of shadow). Let $\Gamma < G$ be a Zariski dense discrete subgroup. For $\psi \in \mathfrak{a}^*$, let ν_ψ be a (Γ, ψ) -conformal measure on \mathcal{F} . Then

- (1) for some $R = R(\nu_\psi) > 0$, we have $c := \inf_{\gamma \in \Gamma} \nu_\psi(O_R(\gamma o, o)) > 0$;
- (2) for all $r \geq R$ and for all $\gamma \in \Gamma$,

$$c \cdot e^{-\|\psi\|\kappa r} e^{-\psi(\mu(\gamma))} \leq \nu_\psi(O_r(o, \gamma o)) \leq e^{\|\psi\|\kappa r} e^{-\psi(\mu(\gamma))},$$

where $\kappa > 0$ is a constant given in Lemma 5.7. In particular, if $\Lambda = \Lambda_c$ in addition, then ν_ψ is atom-free on Λ .

Proof. Suppose that there exist sequences $R_i \rightarrow \infty$ and $\gamma_i \in \Gamma$ such that for all $i \geq 1$, $\nu_\psi(O_{R_i}(\gamma_i^{-1} o, o)) < 1/i$. Write $\gamma_i = k_i a_i \ell_i \in KA^+K$ with $k_i, \ell_i \in K$ and $a_i \in A^+$. Passing to a subsequence, we may assume that $\ell_i \rightarrow \ell_0$ as $i \rightarrow \infty$.

We claim that

$$\limsup O_{R_i}(a_i^{-1} o, o) \supset N^+ e^+. \quad (7.9)$$

Fix an arbitrary $h \in N^+$ and $a_i h = k_i b_i n_i \in KAN$ be the Iwasawa decomposition of $a_i h$. Then the Iwasawa decomposition of $a_i h a_i^{-1}$ is given by

$$a_i h a_i^{-1} = k_i (b_i a_i^{-1}) (a_i n_i a_i^{-1}) \in KAN.$$

Since $a_i h a_i^{-1}$ is uniformly bounded, both $b_i a_i^{-1}$ and $a_i n_i a_i^{-1}$ are also uniformly bounded for all i . It follows that the sequence $n_i \in N$ is uniformly bounded as well. To prove the claim, we observe that for all $i \geq 1$,

$$\begin{aligned} h^+ &\in O_{R_i}(a_i^{-1} o, o) \\ \Leftrightarrow a_i h^+ &\in O_{R_i}(o, a_i o) \\ \Leftrightarrow k_i^+ &\in O_{R_i}(o, a_i o) \\ \Leftrightarrow k_i A^+ o &\cap B(a_i o, R_i) \neq \emptyset \\ \Leftrightarrow a_i^{-1} k_i A^+ o &\cap B(o, R_i) \neq \emptyset \\ \Leftrightarrow h n_i^{-1} b_i^{-1} A^+ o &\cap B(o, R_i) \neq \emptyset. \end{aligned}$$

On the other hand, by the uniform boundedness of n_i and $b_i^{-1} a_i$ and since $R_i \rightarrow \infty$, we have $h n_i^{-1} (b_i^{-1} a_i) o \in B(o, R_i)$ for all sufficiently large i . Hence the claim (7.9) follows.

Since $O_{R_i}(\gamma_i^{-1}o, o) = \ell_i^{-1}O_{R_i}(a_i^{-1}o, o)$, the hypothesis $\nu_\psi(O_{R_i}(\gamma_i^{-1}o, o)) < 1/i$ now implies that $\nu_\psi(\ell_0^{-1}N^+e^+) = 0$ by Claim (7.9). Since N^+e^+ is Zariski open in \mathcal{F} , this contradicts the fact that $\Lambda \subset \text{supp } \nu_\psi$ is Zariski dense in \mathcal{F} . This proves the claim (1).

Now let $\gamma \in \Gamma$ and $r > R$ be arbitrary. By Lemma 5.7, for all $\xi \in O_r(\gamma^{-1}o, o)$, we have

$$\|\beta_\xi(\gamma^{-1}o, o) - \mu(\gamma)\| \leq \kappa r.$$

Since

$$\nu_\psi(O_r(o, \gamma o)) = \int_{O_r(\gamma^{-1}o, o)} e^{-\psi(\beta_\xi(\gamma^{-1}o, o))} d\nu_\psi(\xi),$$

(2) then follows from (1).

Suppose that $\Lambda = \Lambda_c$. Then for any $\xi \in \Lambda$, there exist $r > 0$ and a sequence $\gamma_i \rightarrow \infty$ in Γ such that $\xi \in \bigcap_i O_r(o, \gamma_i o)$. Since $\nu_\psi(\xi) \leq \nu_\psi(O_r(o, \gamma_i o)) \leq Ce^{-\psi(\mu(\gamma_i))}$ and $\psi(\mu(\gamma_i)) \rightarrow +\infty$ as $i \rightarrow \infty$, $\nu_\psi(\xi) = 0$. Hence the second claim follows. \blacksquare

Lemma 7.10. [39, Lem. III.1.3] Let $\theta : \mathfrak{a} \rightarrow \mathbb{R}$ be a continuous function satisfying $\theta(tu) = t\theta(u)$ for all $t \geq 0$ and $u \in \mathfrak{a}$. If $\theta(u) > \psi_\Gamma(u)$ for all $u \in \mathfrak{a} - \{0\}$, then

$$\sum_{\gamma \in \Gamma} e^{-\theta(\mu(\gamma))} < \infty.$$

If there exists $u \in \mathfrak{a}$ such that $\theta(u) < \psi_\Gamma(u)$, then $\sum_{\gamma \in \Gamma} e^{-\theta(\mu(\gamma))} = \infty$.

Lemma 7.11. Suppose that Γ is Zariski dense. Let $\psi \in \mathfrak{a}^*$. If there exists a (Γ, ψ) -conformal measure ν_ψ such that $\nu_\psi(\Lambda_c) > 0$, then

$$\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} = \infty.$$

Moreover, $\psi(v) = \psi_\Gamma(v)$ for some non-zero $v \in \mathcal{L}_\Gamma$.

Proof. Note that Λ_c is an increasing union $\bigcup_{N=1}^{\infty} \Lambda_N$, where

$$\Lambda_N := \{\xi \in \Lambda : \text{there exists } \gamma_i \rightarrow \infty \text{ in } \Gamma \text{ such that } \xi \in O_N(o, \gamma_i o)\}.$$

Hence $\nu_\psi(\Lambda_{N_0}) > 0$ for some $N_0 \geq 1$. Fix $N \geq \max\{R(\nu), N_0\}$, and set $C' := e^{\|\psi\|\kappa N}$ where $R(\nu)$ is as in Lemma 7.8. Observe that for any $m \geq 1$,

$$\Lambda_N \subset \bigcup_{\gamma \in \Gamma, d(o, \gamma o) > m} O_N(o, \gamma o).$$

Hence

$$0 < \nu_\psi(\Lambda_N) \leq \sum_{d(o, \gamma o) > m} \nu_\psi(O_N(o, \gamma o)) \leq C' \sum_{d(o, \gamma o) > m} e^{-\psi(\mu(\gamma))}.$$

Since $m > 1$ is arbitrary, the first claim follows.

We note that $\psi \geq \psi_\Gamma$ by [40, Thm. 8.1]. If $\psi(u) > \psi_\Gamma(u)$ for all $u \in \mathcal{L}_\Gamma - \{0\}$, and hence for all $u \in \mathfrak{a} - \{0\}$, then Lemma 7.10 implies $\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} < \infty$. This is a contradiction by the first claim. \blacksquare

When Γ is Anosov, $\Lambda = \Lambda_c$ and hence by Theorem 4.3(5),

Corollary 7.12. If Γ is Anosov, then $\sum_{\gamma \in \Gamma} e^{-\psi(\mu(\gamma))} = \infty$ for any $\psi \in D_\Gamma^*$.

Proof of Theorem 7.7. In order to prove surjectivity, suppose that there exists a (Γ, ψ) -PS measure, say ν_ψ , for $\psi \in \mathfrak{a}^*$. By Lemma 7.11, $\psi(v) = \psi_\Gamma(v)$ for some non-zero $v \in \mathcal{L}_\Gamma$. By Theorem 4.3(1), it follows that $\psi \in D_\Gamma^*$, proving surjectivity.

If $\psi_i \rightarrow \psi$ in D_Γ^* , then any weak-limit of ν_{ψ_i} is a (Γ, ψ) -PS measure. By the uniqueness of (Γ, ψ) -conformal measure, ν_{ψ_i} converges to ν_ψ as $i \rightarrow \infty$. Hence the map $\psi \mapsto \nu_\psi$ is continuous. Now suppose $\nu_{\psi_i} \rightarrow \nu_\psi$ where $\psi_i, \psi \in D_\Gamma^*$. Since the closed cone generated by $\mu(\Gamma)$ is equal to \mathcal{L}_Γ that has non-empty interior, we can find $\gamma_1, \dots, \gamma_k \in \Gamma$ such that $\mu(\gamma_i)$'s form a basis of \mathfrak{a} . For each γ_ℓ and $r > 0$, we have $\nu_{\psi_i}(O_r(o, \gamma_\ell o)) \rightarrow \nu_\psi(O_r(o, \gamma_\ell o))$. Hence $\{(\psi_i - \psi)(\mu(\gamma_\ell)) : i = 1, 2, \dots\}$ is bounded by Lemma 7.8. It follows that $\{\psi_i : i = 1, 2, \dots\}$ is a relatively compact subset of \mathfrak{a}^* . Suppose that $\phi \in \mathfrak{a}^*$ is a limit of $\{\psi_i\}$. By passing to a subsequence, assume that $\psi_i \rightarrow \phi \in \mathfrak{a}^*$. Since $\nu_{\psi_i} \rightarrow \nu_\psi$, it follows that ν_ψ is a (Γ, ϕ) -PS measure. Since $\psi \mapsto \nu_\psi$ is a bijection between D_Γ^* and \mathcal{S}_Γ , we have $\phi \in D_\Gamma^*$ and $\nu_\phi = \nu_\psi$. By Lemma 7.5, we have $\phi = \psi$. Since every limit of the sequence ψ_i is ψ , it follows that ψ_i converges to ψ as $i \rightarrow \infty$. This finishes the proof. \blacksquare

Critical exponents. Recall the definition of O_Γ from (4.5). For each unit vector $w \in O_\Gamma$, consider the Poincare series

$$\mathcal{P}_w(s, p) := \sum_{\gamma \in \Gamma} e^{-s\langle w, \underline{a}(p, \gamma p) \rangle}.$$

Define the critical exponent δ_w to be the abscissa of convergence of $\mathcal{P}_w(s, p)$, which is independent of $p \in G/K$:

$$\delta_w := \inf\{s \in \mathbb{R} : \mathcal{P}_w(s, p) < \infty\}. \quad (7.13)$$

Corollary 7.14. Let $w \in O_\Gamma$ be a unit vector.

(1) For $w = \frac{\nabla \psi_\Gamma(u)}{\|\nabla \psi_\Gamma(u)\|} \in O_\Gamma$ for $u \in \text{int } \mathcal{L}_\Gamma$ with $\|u\| = 1$, we have

$$\delta_w = \|\nabla \psi_\Gamma(u)\|.$$

In particular, $w \mapsto \delta_w$ is analytic on $\{w \in O_\Gamma : \|w\| = 1\}$.

(2) For any $p \in G/K$, $\mathcal{P}_w(\delta_w, p) = \infty$.

Proof. (1) follows from Lemmas 7.10 and 7.11 together with Theorem 4.3. (2) is a direct consequence of Lemma 7.11. ■

8 Myrberg Limit Points of Anosov Groups

In this section, we discuss the notion of Myrberg limit points. We show that for Anosov groups, the set of Myrberg limit points has full measure for any PS measure on Λ . In the rank one case, this was proved by Tukia [56, Thm. 4A]. Let $\Gamma < G$ be a Zariski dense discrete subgroup.

Definition 8.1 (Myrberg points). Let $p \in X$. We call a point $\xi_0 \in \Lambda$ a Myrberg limit point for Γ if, for any $\xi \neq \eta$ in Λ , there exists a sequence $\gamma_i \in \Gamma$ such that $\gamma_i p \rightarrow \xi$ and $\gamma_i \xi_0 \rightarrow \eta$ as $i \rightarrow \infty$.

Note that this definition is independent of the choice of $p \in X$ by Lemma 2.10. We denote by $\Lambda_M \subset \Lambda$ the set of all Myrberg limit points for Γ .

When G is of rank one, a Myrberg limit point $\xi \in \Lambda$ is characterized by the property that any geodesic ray toward ξ is dense in the space of all geodesics connecting limit points. The following proposition generalizes this to a general Anosov subgroup.

Proposition 8.2. Let Γ be Anosov. We have $\xi_0 \in \Lambda_M$ if and only if for any $g \in G$ with $g^+ = \xi_0$,

$$\limsup \Gamma \backslash \Gamma g A^+ M = \Omega.$$

Let $\Gamma < G$ be an Anosov subgroup for the rest of this section.

Lemma 8.3. Let $b_i \in A$ be a sequence tending to ∞ such that $w^{-1}b_i^{-1}w \in A^+$ for some $w \in \mathcal{W}$. If $\gamma_i g b_i \rightarrow h$ for some $h, g \in G$ and $\gamma_i \in \Gamma$, then $\lim_{i \rightarrow \infty} \gamma_i g o = h w^+ \in \Lambda$. In particular, if $b_i \in A^+$, then $\lim_{i \rightarrow \infty} \gamma_i g o = h^-$.

Proof. Let $c_i := h^{-1} \gamma_i g b_i$ and $a_i := w^{-1} b_i^{-1} w \in A^+$. Then $g w = \gamma_i^{-1} h c_i w a_i$. Hence by Lemma 7.2, $a_i \rightarrow \infty$ regularly in A^+ . Lemma 2.11 implies that $h c_i w a_i(o) \rightarrow h w^+$. Since $\gamma_i g w = h c_i w a_i$, we have $\gamma_i g w(o) = \gamma_i g o \rightarrow h w^+$. This proves the first claim by Lemma 2.13. If $b_i \in A^+$, then $w_0^{-1} b_i^{-1} w_0 \in A^+$. Since $w_0^+ = e^-$, the last claim follows. ■

The following is proved in [25, Coro. 5.8]:

Theorem 8.4 (The limit map as a continuous extension of the orbit map). For any $p \in X$, the map $\Gamma \cup \partial\Gamma \rightarrow X \cup \mathcal{F}$ given by $\gamma \mapsto \gamma p$ for $\gamma \in \Gamma$ and $x \mapsto \zeta(x)$ for $x \in \partial\Gamma$ is continuous.

We need the following basic fact about word hyperbolic groups.

Lemma 8.5. Let $x \neq y$ in $\partial\Gamma$. If $\gamma_i \in \Gamma$ is an infinite sequence such that $(\gamma_i x, \gamma_i y) \rightarrow (x', y') \in \partial\Gamma \times \partial\Gamma$, then γ_i converges to either x' or y' .

Proof. Choose a geodesic line $[x, y]$, and its representative $(\dots, u_2, u_1, u_0 = v_0, v_1, v_2, \dots)$. Note that $x = [u_0, u_1, u_2, \dots]$ and $y = [v_0, v_1, v_2, \dots]$. It suffices to show that $\gamma_i u_0$ converges to either x' or y' . Suppose not. Then by passing to a subsequence, we have $\gamma_i u_0 \rightarrow z'$ where $z' \notin \{x', y'\}$. Since $(z'|x'), (z'|y') < \infty$, there exists a subsequence n_k such that $\sup_k (\gamma_k u_0 | \gamma_k u_{n_k}) + (\gamma_k u_0 | \gamma_k v_{n_k}) < \infty$. Let $\mathcal{L}_k^- := [\gamma_k u_0, \gamma_k u_{n_k}]$ and $\mathcal{L}_k^+ := [\gamma_k u_0, \gamma_k v_{n_k}]$, so that $\sup_k d_w(e, \mathcal{L}_k^\pm) < \infty$. The thin triangle property of the hyperbolic group Γ implies that if the projection of e to the geodesic segment $\mathcal{L}_k^- \cup \mathcal{L}_k^+$ lies in \mathcal{L}_k^\pm , then $d_w(e, \gamma_k u_0)$ is equal to $d_w(e, \mathcal{L}_k^\mp)$ up to a uniform additive constant. And hence $d_w(e, \gamma_k u_0)$ is uniformly bounded, which is a contradiction as $\gamma_k \rightarrow \infty$ as $k \rightarrow \infty$. ■

The following is immediate from Theorem 8.4 and Lemma 8.5:

Corollary 8.6. Let $\gamma_i \in \Gamma$ be an infinite sequence such that $(\gamma_i \xi, \gamma_i \eta) \rightarrow (\xi', \eta')$ in $\Lambda^{(2)}$ as $i \rightarrow \infty$. Then for any $p \in X$, $\gamma_i p$ converges to either ξ' or η' .

Lemma 8.7. Let $g \in G$ be such that $g^\pm \in \Lambda$. If $\lim_{i \rightarrow \infty} \gamma_i g^\pm = \xi$ for some infinite sequence $\gamma_i \in \Gamma$, then $\lim_{i \rightarrow \infty} \gamma_i g o = \xi$.

Proof. Set $x^\pm := \zeta^{-1}(g^\pm)$ and $y = \zeta^{-1}(\xi)$. Since $\zeta : \partial\Gamma \rightarrow \Lambda$ is a homeomorphism, we have $\gamma_i x^\pm \rightarrow y$ as $i \rightarrow \infty$. By Lemma 8.5, we have $\gamma_i \rightarrow y$ as $i \rightarrow \infty$. By Theorem 8.4, we get $\lim_{i \rightarrow \infty} \gamma_i o = \xi$. By Lemma 2.10, $\lim_{i \rightarrow \infty} \gamma_i g o = \xi$ as desired. \blacksquare

Since the fibers of the visual map $g \mapsto g^+$ are P -orbits, the following lemma is an easy consequence of the regularity lemma 7.2.

Lemma 8.8. If $g, h \in G$ satisfy $g^+ = h^+$, then

$$\limsup \Gamma g A^+ M = \limsup \Gamma h A^+ M.$$

Proof of Proposition 8.2. Set $\tilde{\Omega} := \{g \in G : g^\pm \in \Lambda\}$. Suppose $\xi_0 \in \Lambda_M$ and $g^+ = \xi_0$. We claim that $\Gamma g A^+ M = \tilde{\Omega}$. By Lemma 8.8, we may assume that $g^- \in \Lambda$. Let $h \in \tilde{\Omega}$. As $\xi_0 \in \Lambda_M$, there exists $\gamma_i \in \Gamma$ such that $\gamma_i g^+ \rightarrow h^+$ and $\gamma_i g o \rightarrow h^-$. By Lemma 8.5, by passing to a subsequence, $\gamma_i g^-$ converges to h^- . Therefore, $\gamma_i g A M \rightarrow h A M$ in G/AM ; there exists $b_i m_i \in AM$ such that $\gamma_i g b_i m_i \rightarrow h$. We claim that $b_i \in A^+$ for all large i . If not, by passing to a subsequence, we have m_i^{-1} converges to some $m_0 \in M$ and there exists $w \in \mathcal{W} - \{e\}$ such that $a_i := w^{-1} b_i w \in A^+$. Then $\gamma_i g w a_i \rightarrow h m_0 w$. By Lemma 8.3, $\gamma_i g o \rightarrow h m_0 w^-$, and hence $h m_0 w^- = h^-$. It follows that $w = e$, yielding a contradiction. Therefore, $h \in \limsup \Gamma g A^+ M$, proving the claim.

Now suppose that $\limsup \Gamma g A^+ M = \tilde{\Omega}$. We claim that $g^+ \in \Lambda_M$. Let $\xi \neq \xi'$ in Λ , and let $h \in G$ be such that $h^+ = \xi$ and $h^- = \xi'$. By the hypothesis and Lemma 7.2, there exist $\gamma_i \in \Gamma$, $m_i \in M$ and $a_i \rightarrow \infty$ regularly in A^+ such that $\gamma_i g a_i m_i \rightarrow h$ in G . Then $\gamma_i g^+ \rightarrow h^+ = \xi$. By Lemma 8.3, $\gamma_i g o \rightarrow h^- = \xi'$. Hence $g^+ \in \Lambda_M$. This finishes the proof of Proposition 8.2. \blacksquare

Lemma 8.9. Let $\psi \in D_\Gamma^*$, and $(\xi, \xi'), (\eta_1, \eta_2) \in \Lambda^{(2)}$. If $\gamma_i \in \Gamma$ and $t_i \rightarrow +\infty$ are such that

$$\lim_{i \rightarrow \infty} (\gamma_i \xi, \gamma_i \xi', t_i + \psi(\beta_{\gamma_i \xi}(o, \gamma_i o))) = (\eta_1, \eta_2, 0),$$

then $\lim_{i \rightarrow \infty} \gamma_i(o) = \eta_2$.

Proof. Write $x = \zeta^{-1}(\xi)$, $x' = \zeta^{-1}(\xi')$, $y_1 = \zeta^{-1}(\eta_1)$, $y_2 = \zeta^{-1}(\eta_2)$, and choose $u \in [x, x']$. Since the triangle $[\gamma_i x, \gamma_i x'] \cup [\gamma_i u, \gamma_i x] \cup [\gamma_i u, \gamma_i x']$ is δ -thin, it follows that for all i , either $\gamma_i x \in O_\delta(u, \gamma_i u)$ or $\gamma_i x' \in O_\delta(u, \gamma_i u)$. We claim the latter holds for all large i .

Suppose not. Then by passing to a subsequence, we may assume that $\gamma_i x \in O_\delta(u, \gamma_i u)$ for all i . Then by Proposition 5.14 and Lemma 5.7, there exists a uniform constant $c > 0$ such that $\gamma_i \xi \in O_{c(\delta+1)}(uo, \gamma_i uo)$ and

$$|\psi(\beta_{\gamma_i \xi}(uo, \gamma_i uo)) - \psi(\mu(\gamma_i))| < \|\psi\| \kappa c(\delta + 1).$$

Since $\psi(\mu(\gamma_i)) \rightarrow +\infty$ as $i \rightarrow \infty$ by Lemma 5.4, and $\psi(\beta_{\gamma_i \xi}(uo, \gamma_i uo))$ and $\psi(\beta_{\gamma_i \xi}(o, \gamma_i o))$ are uniformly close to each other, $\psi(\beta_{\gamma_i \xi}(o, \gamma_i o)) \rightarrow +\infty$. This contradicts the hypothesis that the sequence $t_i + \psi(\beta_{\gamma_i \xi}(o, \gamma_i o))$ converges to a finite number as $i \rightarrow \infty$. It follows that for all sufficiently large i ,

$$\gamma_i x' \in O_\delta(u, \gamma_i u). \quad (8.10)$$

On the other hand, $\gamma_i u \rightarrow y_\ell$ for some $\ell \in \{1, 2\}$ by Lemma 8.5. Since $\gamma_i x' \rightarrow y_2$ and $O_\delta(u, \gamma_i u)$ converges to y_ℓ , (8.10) implies that $\gamma_i u \rightarrow y_2$. Therefore, $\gamma_i o \rightarrow \eta_2$ by Lemma 8.4. \blacksquare

Theorem 8.11. For any PS-measure ν on Λ , $\nu(\Lambda_M) = 1$.

Proof. By Theorem 7.7, $\nu = \nu_\psi$ for some $\psi \in D_\Gamma^*$. Let m_ψ be the $\mathbb{R} := \{\tau_s : s \in \mathbb{R}\}$ -ergodic finite measure on $\Gamma \backslash \Lambda^{(2)} \times \mathbb{R}$ in Theorem 4.8. Let $Z_\psi \subset \Gamma \backslash \Lambda^{(2)} \times \mathbb{R}$ denote the set of elements with dense \mathbb{R}_+ -orbits, and \tilde{Z}_ψ be its lift in $\Lambda^{(2)} \times \mathbb{R}$. By the Birkhoff ergodic theorem, Z_ψ has full m_ψ -measure, and hence $\nu(\pi(\tilde{Z}_\psi)) = \nu(\Lambda)$ where $\pi : \Lambda^{(2)} \times \mathbb{R} \rightarrow \Lambda$ denotes the projection map $\pi(\xi, \eta, t) = \xi$. It is now sufficient to prove that $\pi(\tilde{Z}_\psi) \subset \Lambda_M$.

Let $\xi \in \pi(\tilde{Z}_\psi)$ and $(\eta_1, \eta_2) \in \Lambda^{(2)}$ be arbitrary. We need to show that there exists $\gamma_i \in \Gamma$ such that $\gamma_i \xi \rightarrow \eta_1$ and $\gamma_i o \rightarrow \eta_2$ as $i \rightarrow \infty$. Choose $(\xi, \xi', 0) \in \tilde{Z}_\psi$. By definition, we can find $\gamma_i \in \Gamma$ and $t_i \rightarrow +\infty$ such that the sequence

$$\gamma_i(\tau_{t_i}(\xi, \xi', 0)) = \gamma_i(\xi, \xi', t_i) = (\gamma_i \xi, \gamma_i \xi', t_i + \psi(\beta_{\gamma_i \xi}(o, \gamma_i o)))$$

converges to $(\eta_1, \eta_2, 0)$. Since $\gamma_i o \rightarrow \eta_2$ by Lemma 8.9, this finishes the proof. \blacksquare

In the rank one case, the BMS measure is finite, and $A = \{a_t\}$ is the union of $A^+ = \{a_t : t \geq 0\}$ and $A^- = \{a_t : t \leq 0\}$. The AM -ergodicity of the BMS measure implies

that for almost all $x \in \Gamma \backslash G$, $xA^\pm M$ is dense in $\Omega = \{x \in \Gamma \backslash G : x^\pm \in \Lambda\}$. In general, $A = \cup_{w \in \mathcal{W}} wA^+w^{-1}$, and we have the following corollary of Theorem 8.11:

Corollary 8.12. Let $\psi \in D_\Gamma^*$. For m_ψ^{BMS} -almost all $x \in \Omega$, each xwA^+M and xw_0A^+M is dense in Ω .

Proof. Note that for $x = \Gamma g \in \Omega$, xwA^+M is dense in Ω if and only if $gw^+ \in \Lambda_M$ by Proposition 8.2. For $w = e$ (resp. $w = w_0$), the claim follows as $\nu_\psi(\Lambda_M) = 1$ (resp. $\nu_{\psi \circ i}(\Lambda_M) = 1$) by Theorem 8.11. \blacksquare

We also observe:

Lemma 8.13. For any $x \in \mathcal{E}$ and $w \in \mathcal{W} - \{e, w_0\}$, the map $A^+M \rightarrow xwA^+M$ is proper.

Proof. Note that if $(g^+, gw^+) \in \mathcal{F}^{(2)}$ for $g \in G$ and $w \in \mathcal{W}$, then $w = w_0$. Choose $g \in G$ so that $\Gamma g = x \in \mathcal{E}$. Since $g^+ \in \Lambda$ and $\Lambda \times \Lambda - \{(\xi, \xi)\} \subset \mathcal{F}^{(2)}$ by the antipodality, $gw^+ \in \Lambda$ can happen only for $w \in \{e, w_0\}$. Suppose for some $\gamma_i \in \Gamma$ and $a_i \rightarrow \infty$ in A^+ , $\gamma_i g w a_i$ converges to some $h \in G$ as $i \rightarrow \infty$. This means that $d(g w a_i, \gamma_i^{-1} h) \rightarrow 0$ as $i \rightarrow \infty$, and hence $gw^+ \in \Lambda$. Hence, for each $w \in \mathcal{W} - \{e, w_0\}$, $\limsup xwA^+M = \emptyset$, proving the claim. \blacksquare

9 Criterion for Ergodicity via Essential Values

In this section, let $\Gamma < G$ be a Zariski dense discrete subgroup, and let ν_ψ be a (Γ, ψ) -conformal measure on \mathcal{F} for $\psi \in \mathfrak{a}^*$. Consider the action of G on $\mathcal{F} \times \mathfrak{a}$ by

$$g(\xi, v) = (g\xi, v + \beta_\xi(g^{-1}, e)).$$

Then the map $g \mapsto (g^+, b := \beta_{g^+}(e, g))$ induces a G -equivariant homeomorphism $G/NM \cong \mathcal{F} \times \mathfrak{a}$. Using this homeomorphism, we define a Γ -invariant Radon measure $\widehat{\nu}_\psi$ on $G/NM \cong \mathcal{F} \times \mathfrak{a}$ by

$$d\widehat{\nu}_\psi(gNM) = d\nu_\psi(g^+)e^{\psi(b)}db.$$

Since $dm_\psi^{\text{BR}} = d\widehat{\nu}_\psi dm dn$, the NM -ergodicity of m_ψ^{BMS} is equivalent to the Γ -ergodicity of $\widehat{\nu}_\psi$. For simplicity, we set $\nu := \nu_\psi$ and $\widehat{\nu} := \widehat{\nu}_\psi$ for the rest of the section. Schmidt gave a characterization of Γ -ergodicity of $\widehat{\nu}$ using the notion of ν -essential values in the rank one case ([52], see also [46, Prop. 2.1]).

Definition 9.1. An element $v \in \mathfrak{a}$ is called a (ν, Γ) -essential value, if for any Borel set $B \subset \mathcal{F}$ with $\nu(B) > 0$ and any $\varepsilon > 0$, there exists $\gamma \in \Gamma$ such that

$$\nu \left(B \cap \gamma^{-1}B \cap \{\xi \in \mathcal{F} : \|\beta_\xi(\gamma^{-1}o, o) - v\| < \varepsilon\} \right) > 0.$$

Let $E_\nu = E_\nu(\Gamma)$ denote the set of all (ν, Γ) -essential values in \mathfrak{a} . It is easy to see that E_ν is a closed subgroup of \mathfrak{a} . The main goal of this section is to prove the following criterion of Γ -ergodicity of $\widehat{\nu}$, which can be considered as a higher rank version of [46, Prop. 2.1].

Proposition 9.2. $(G/NM, \Gamma, \widehat{\nu})$ is ergodic if and only if $(G/P, \Gamma, \nu)$ is ergodic and $E_\nu(\Gamma) = \mathfrak{a}$.

Fixing ν , we set $E := E_\nu(\Gamma)$ in the rest of this section. Our proof of Proposition 9.2 is an easy adaptation of the proof of [46, Prop. 2.1] to a higher rank case. We begin with the following lemma.

Lemma 9.3. Let $h : G/NM = \mathcal{F} \times \mathfrak{a} \rightarrow [0, 1]$ be a Γ -invariant Borel function such that for each $\xi \in \mathcal{F}$, $h(\xi, \cdot)$ is a C -Lipschitz function on \mathfrak{a} for some $C > 0$ independent of ξ . Then for each $\log a \in E$, $h(xa) = h(x)$ for $\widehat{\nu}$ -a.e. $x \in G/NM$.

Proof. Suppose that $\widehat{\nu}\{x \in G/NM : h(x) \neq h(xa)\} > 0$ for some $\log a \in E$. We will then find a subset $\mathcal{A}^* = \mathcal{A}^*(a) \subset G/NM$ with $\widehat{\nu}(\mathcal{A}^*) > 0$ and $\gamma \in \Gamma$ such that $h(\gamma^{-1}x) \neq h(x)$ for all $x \in \mathcal{A}^*$; this contradicts the Γ -invariance of h .

By replacing h with $-h$ if necessary, we may assume that $\widehat{\nu}\{x \in G/NM : h(x) < h(xa)\} > 0$. Hence there exist $r, \varepsilon > 0$ such that

$$Q_a := \{x \in G/NM : h(x) < r - C\varepsilon < r + C\varepsilon < h(xa)\}$$

has a positive $\widehat{\nu}$ -measure. Now we can choose a ball $\mathcal{O} = B_{\mathfrak{a}}(v_0, \varepsilon/2) \subset \mathfrak{a}$ such that

$$\widehat{\nu}((\mathcal{F} \times \mathcal{O}) \cap Q_a) > 0.$$

Set $F_a := \{\xi \in \mathcal{F} : (\{\xi\} \times \mathcal{O}) \cap Q_a \neq \emptyset\}$. We claim that

$$\text{if } (\xi, w) \in F_a \times \mathcal{O}, \text{ then } h(\xi, w + \log a) > r > h(\xi, w). \quad (9.4)$$

Note that there exists $v \in \mathfrak{a}$ with $\|v\| < \varepsilon$ such that $(\xi, w + v) \in Q_a$ and hence

$$|h(\xi, w)| \leq |h(\xi, w) - h(\xi, w + v)| + |h(\xi, w + v)| < C\|v\| + (r - C\varepsilon) \leq r.$$

Similarly,

$$\begin{aligned} |h(\xi, w + \log a)| &\geq |h(\xi, w + v + \log a)| - |h(\xi, w + \log a) - h(\xi, w + v + \log a)| \\ &> (r + C\varepsilon) - C\|v\| > r, \end{aligned}$$

which verifies the claim (9.4).

Since $-\log a \in E$ and $\nu(F_a) > 0$, there exists $\gamma \in \Gamma$ such that

$$\mathcal{A} := F_a \cap \gamma F_a \cap \{\xi \in G/P : \|\beta_\xi(o, \gamma o) + \log a\| < \varepsilon/2\}$$

has a positive ν -measure. For $\xi \in \mathcal{A}$, set

$$\mathcal{O}_\xi := \{w \in \mathcal{O} : w - (\beta_\xi(o, \gamma o) + \log a) \in \mathcal{O}\}.$$

Since $\|\beta_\xi(o, \gamma o) + \log a\| < \varepsilon/2$, and \mathcal{O} is a Euclidean ball of diameter ε , there is a uniform positive lower bound for the volume of \mathcal{O}_ξ . It follows that

$$\mathcal{A}^* := \bigcup_{\xi \in \mathcal{A}} \{\xi\} \times \mathcal{O}_\xi$$

has positive $\widehat{\nu}$ -measure. We now claim that $h \circ \gamma^{-1} > h$ on \mathcal{A}^* .

Let $(\xi, w) \in \mathcal{A}^*$. Since $(\xi, w) \in F_a \times \mathcal{O}$, (9.4) implies that $h(\xi, w) < r$.

Write $\gamma^{-1}(\xi, w) = (\gamma^{-1}\xi, w - (\beta_\xi(o, \gamma o) + \log a) + \log a)$. Since $(\gamma^{-1}\xi, w - (\beta_\xi(o, \gamma o) + \log a)) \in F_a \times \mathcal{O}$, (9.4) says that

$$h(\gamma^{-1}(\xi, w)) > r;$$

this proves the claim. ■

Proof of Proposition 9.2. Assume that $(G/NM, \Gamma, \widehat{\nu})$ is ergodic. Let $\pi : G/NM \rightarrow G/P$ denote the projection map. Since $\pi_* \widehat{\nu}$ is absolutely continuous with respect to ν , it follows that $(G/P, \Gamma, \nu)$ is ergodic.

To show $E = \mathfrak{a}$, fix an arbitrary Borel set $B \subset G/P$ of positive ν -measure. For any $w \in \mathfrak{a}$ and $\varepsilon > 0$, we define

$$\mathcal{B}_{w,\varepsilon} := \{(\xi, v) \in G/P \times \mathfrak{a} : \xi \in B, \|v - w\| < \varepsilon\} \subset G/NM.$$

Observe that

$$\widehat{\nu}(\mathcal{B}_{0,\varepsilon}) = \int_{G/P} \int_{\mathfrak{a}} \mathbf{1}_{\mathcal{B}_{0,\varepsilon}}(\xi, b) e^{\psi(b)} db d\nu(\xi) \geq \text{Vol}(\mathcal{B}_{\mathfrak{a}}(0, \varepsilon)) e^{-\|\psi\|\varepsilon} \nu(B) > 0.$$

Hence it follows from the ergodicity of $(G/NM, \Gamma, \widehat{\nu})$ that $\widehat{\nu}(G/NM - \Gamma \mathcal{B}_{0,\varepsilon}) = 0$. In particular, there exists $\gamma \in \Gamma$ such that $\widehat{\nu}(\mathcal{B}_{w,\varepsilon} \cap \gamma \mathcal{B}_{0,\varepsilon}) > 0$. Finally, note that if $(\xi, v) \in \mathcal{B}_{w,\varepsilon} \cap \gamma \mathcal{B}_{0,\varepsilon}$, then $\xi \in B \cap \gamma B$, and

$$\|\beta_{\xi}(e, \gamma) - w\| \leq \|\beta_{\xi}(e, \gamma) - v\| + \|v - w\| \leq \varepsilon + \varepsilon = 2\varepsilon.$$

This, together with the fact $\pi_* \widehat{\nu} \ll \nu$, implies that

$$\nu(B \cap \gamma B \cap \{\xi \in G/P : \|\beta_{\xi}(e, \gamma) - w\| \leq 2\varepsilon\}) > 0,$$

which finishes the proof of (\Rightarrow) .

We now assume that $(G/P, \Gamma, \nu)$ is ergodic and $E = \mathfrak{a}$. Let $h : G/NM \rightarrow [0, 1]$ be a Γ -invariant Borel function. We need to show that h is constant $\widehat{\nu}$ -a.e. Identifying $\mathfrak{a} \simeq \mathbb{R}^r$ with $r = \text{rank } G$, for each $\tau = (\tau_1, \dots, \tau_r) \in \mathfrak{a}$, we define a Γ -invariant Borel function $h_{\tau} : G/NM \rightarrow \mathbb{R}$ as follows:

$$h_{\tau}(x) = \int_0^{\tau_1} \cdots \int_0^{\tau_r} h(x \exp(t_1, \dots, t_r)) dt_r \cdots dt_1.$$

Note that h_{τ} satisfies the hypothesis of Lemma 9.3. Hence by the hypothesis $E_{\nu} = \mathfrak{a}$, for each $a \in A$, $h_{\tau}(x) = h_{\tau}(xa)$ for $\widehat{\nu}$ -a.e. $x \in G/NM$.

Let $\{a_n : n \in \mathbb{N}\}$ be a countable dense subset of A . Then there exists Ω_n of full $\widehat{\nu}$ -measure such that for all $x \in \Omega_n$, $h_{\tau}(x) = h_{\tau}(xa_n)$. Set $\Omega := \cap_{n=1}^{\infty} \Omega_n$. Then for all $x \in \Omega$, we have $h_{\tau}(x) = h_{\tau}(xa)$ for all $a \in A$, as $h_{\tau}(\xi, \cdot)$ is continuous on \mathfrak{a} . Now h_{τ} is a Γ -invariant function on G/NM , which is also A -invariant $\widehat{\nu}$ -a.e.

Since $(G/P, \Gamma, \nu)$ is ergodic, there exists $c(\tau) \in \mathbb{R}$ such that $h_{\tau} = c(\tau)$ $\widehat{\nu}$ -a.e. on G/NM .

Next, fix $1 \leq i \leq r$ and $\tau_1, \dots, \tau_{i-1}, \tau_{i+1}, \dots, \tau_r \geq 0$, and define

$$f(t) := (\tau_1, \dots, \tau_{i-1}, t, \tau_{i+1}, \dots, \tau_r) \in \mathfrak{a}.$$

Then $t \mapsto c(f(t))$ is linear; indeed, by definition, we have

$$h_{f(t+s)} = h_{f(t)} + h_{f(s)} \circ \exp(te_i)$$

for all $t, s \geq 0$ and hence $c(f(t+s)) = c(f(t)) + c(f(s))$. We conclude $c(\tau) = \kappa \tau_1 \cdots \tau_r$, for some $\kappa \in \mathbb{R}$.

Hence for each $\tau \in \mathfrak{a}$, $h_\tau = \kappa \tau_1 \cdots \tau_r$ $\widehat{\nu}$ -a.e. Since $|h_{\tau+\sigma} - h_\tau| \leq 2^r \|\sigma\| \|\tau\|^{r-1}$ and hence $\tau \rightarrow h_\tau$ is continuous, using a countable dense subset of \mathfrak{a} , we conclude there exists a subset Ω of full $\widehat{\nu}$ -measure such that

$$h_\tau(x) = \kappa \tau_1 \cdots \tau_r \quad \text{for all } x \in \Omega \text{ and } \tau \in \mathfrak{a}.$$

By restricting h_τ to each fiber of $\pi : G/NM \rightarrow G/P$, and applying the Lebesgue differentiation theorem, we conclude that $\frac{1}{\tau_1 \cdots \tau_r} h_\tau(x) \rightarrow h(x)$ as $\tau \rightarrow 0$ for $\widehat{\nu}$ -a.e. x . Consequently, $h = \kappa$ $\widehat{\nu}$ -a.e., finishing the proof. \blacksquare

10 Ergodicity of m_ψ^{BR} and Classification

Let $\Gamma < G$ be an Anosov subgroup. Recall the NM -invariant BR measure m_ψ^{BR} defined in (3.23). We prove the following theorem in this section:

Theorem 10.1. For each $\psi \in D_\Gamma^*$, m_ψ^{BR} is NM -ergodic.

Recall the definition of $\widehat{\nu}_\psi$ and ν_ψ from section 9. Since $(\mathcal{F}, \Gamma, \nu_\psi)$ is ergodic by Theorem 4.3, the following proposition implies that $(G/NM, \Gamma, \widehat{\nu}_\psi)$, and hence $(\Gamma \backslash G, NM, m_\psi^{\text{BR}})$, is ergodic by Proposition 9.2.

Proposition 10.2. Let Γ_0 be a Zariski dense normal subgroup of Γ . For any $\psi \in D_\Gamma^*$, we have $E_{\nu_\psi}(\Gamma_0) = \mathfrak{a}$. In particular, $E_{\nu_\psi}(\Gamma) = \mathfrak{a}$.

Most of the section is devoted to the proof of Proposition 10.2. We fix a Zariski dense normal subgroup Γ_0 of Γ .

Lemma 10.3. For any finite subset $S_0 \subset \lambda(\Gamma_0)$, the subgroup generated by $\lambda(\Gamma_0) - S_0$ is dense in \mathfrak{a} .

Proof. Let F denote the closure of the subgroup generated by $\lambda(\Gamma_0) - S_0$. Suppose that $F \neq \mathfrak{a}$. Identifying $\mathfrak{a} = \mathbb{R}^r$, since F is infinite, there exist $1 \leq k < r$ and $0 \leq m \leq r$ such that $F = \sum_{i=1}^k \mathbb{R}v_i + \sum_{i=1}^m \mathbb{Z}w_i$ where v_i, w_i are linearly independent vectors. For each $s = \lambda(\gamma) \in S_0$, $\lambda(\gamma^n) = n\lambda(\gamma) \rightarrow \infty$ as γ is loxodromic. Hence, there exists $n_s \in \mathbb{N}$ so that $n_s\lambda(\gamma) \in F$. Setting $N := \prod_{s \in S_0} n_s$, we have $S_0 \subset \sum_{i=1}^k \mathbb{R}v_i + N^{-1} \sum_{i=1}^m \mathbb{Z}w_i$.

Therefore, the closure of the subgroup generated by $F \cup S_0$ is contained in $\sum_{i=1}^k \mathbb{R}v_i + N^{-1} \sum_{i=1}^m \mathbb{Z}w_i$. Since $\lambda(\Gamma_0) \subset \sum_{i=1}^k \mathbb{R}v_i + N^{-1} \sum_{i=1}^m \mathbb{Z}w_i$ and $\lambda(\Gamma_0)$ generates a dense subgroup of \mathfrak{a} [7], it follows that $k = \dim \mathfrak{a}$, yielding a contradiction. ■

Proposition 10.4. For any $\psi \in D_\Gamma^*$ and $C > 0$, the set $\{\lambda(\gamma) \in \mathfrak{a}^+ : \gamma \in \Gamma_0, \psi(\lambda(\gamma)) \geq C\}$ generates a dense subgroup of \mathfrak{a} .

Proof. Theorem 3.2 in [49] extends to general Anosov subgroups (see also [12, Thm. A.2-(2)]), and hence the cocycle $c = \psi \circ \sigma$ has a finite exponential growth rate. In particular,

$$\#\{\lambda(\gamma) : \gamma \in \Gamma, \psi(\lambda(\gamma)) < C\} \leq \#[\gamma] \in [\Gamma] : \psi(\lambda(\gamma)) < C\} < \infty \quad (10.5)$$

where $[\Gamma]$ denotes the set of conjugacy classes in Γ . Hence $\#\{\lambda(\gamma) : \gamma \in \Gamma_0, \psi(\lambda(\gamma)) < C\} < \infty$ and the claim follows from Lemma 10.3. ■

Lemma 10.6. There exists a compact subset $\mathcal{C} \subset G$ such that for any $\xi \in \Lambda$, there exists $g \in \mathcal{C}$ such that $g^+ = \xi$ and $g^- \in \Lambda$.

Proof. In the Gromov hyperbolic space Γ , there exists a finite subset $F \subset \Gamma$ such that for any $x \in \partial\Gamma$, there exists $y \in \partial\Gamma$ such that $[x, y] \cap F \neq \emptyset$. It suffices to choose a compact subset $\mathcal{C} \subset G$ such that $\mathcal{C}(o)$ contains the R_1 -neighborhood of $F(o)$ with R_1 given in Proposition 5.12. ■

We set

$$N_0 := \max_{p \in \mathcal{C}(o)} N_0(\psi, p) < \infty$$

with $N_0(\psi, p)$ and \mathcal{C} given by Lemmas 6.12 and 10.6, respectively.

In view of Proposition 10.4, Proposition 10.2 is an immediate consequence of the following:

Proposition 10.7. For any $\gamma_0 \in \Gamma_0$ with $\psi(\lambda(\gamma_0)) \geq 1 + \log 3N_0$,

$$\lambda(\gamma_0) \in E_{\nu_\psi}(\Gamma_0).$$

Essential values of ν_ψ

Most of this section is devoted to the proof of this proposition. We fix $\gamma_0 \in \Gamma_0$ with

$$\psi(\lambda(\gamma_0)) \geq \log 3N_0 + 1.$$

Since $\psi > 0$ on $\lambda(\Gamma) - \{0\}$ by Theorem 4.3(4), we have

$$\psi(i\lambda(\gamma_0)) + \psi(\lambda(\gamma_0)) > \log 3N_0 + 1. \quad (10.8)$$

Definition of $\mathcal{B}_R(\gamma_0, \varepsilon)$

Let $0 < \varepsilon < \|\psi\|^{-1}$ be an arbitrary number. We fix $g \in \mathcal{C}$ such that $g^+ = y_{\gamma_0}$ and $g^- \in \Lambda$, given by Lemma 10.6. Set $p := go \in \mathcal{C}(o)$, $\xi_0 := y_{\gamma_0}$, and $\eta := g^-$.

For $\xi \in \Lambda$ and $r > 0$, set

$$\mathbb{B}_p(\xi, r) := \{\eta \in \Lambda : d_{\psi, p}(\xi, \eta) < r\}$$

where $d_{\psi, p}$ is the virtual visual metric defined in section 6.

For each $\gamma \in \Gamma$, define $r_p(\gamma) > 0$ to be the supremum $r \geq 0$ such that

$$\max_{\xi \in \mathbb{B}_p(\gamma \xi_0, 3N_0 r)} \|\beta_\xi(p, \gamma \gamma_0^{\pm 1} \gamma^{-1} p) \mp \lambda(\gamma_0)\| < \varepsilon. \quad (10.9)$$

For each $R > 0$, we define the family of virtual-balls as follows:

$$\mathcal{B}_R(\gamma_0, \varepsilon) = \{\mathbb{B}_p(\gamma \xi_0, r) : \gamma \in \Gamma, 0 < r < \min(R, r_p(\gamma))\}.$$

Equivalently, $\mathbb{B}_p(\gamma \xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$ if and only if $r < R$ and one has $\|\beta_\eta(p, \gamma \gamma_0 \gamma^{-1} p) - \lambda(\gamma_0)\| \leq \varepsilon$ for all $\eta \in \mathbb{B}_p(\gamma \xi_0, r)$.

Let $C = C(\psi, p) > 0$ be as in Theorem 5.3. Since $\xi_0 \in O_{\varepsilon/(8\kappa)}(\eta, p)$ where $\kappa > 0$ is as in Lemma 5.7, we can choose $0 < s = s(\gamma_0) < R$ small enough such that

$$\mathbb{B}_p(\xi_0, e^{\psi(\lambda(\gamma_0) + i\lambda(\gamma_0)) + \frac{1}{2}\|\psi\|\varepsilon + 2C}s) \subset O_{\varepsilon/(8\kappa)}(\eta, p); \quad (10.10)$$

$$\sup_{x \in \mathbb{B}_p(\xi_0, e^{2C}s)} \|\beta_x(p, \gamma_0^{\pm 1}p) \mp \lambda(\gamma_0)\| < \varepsilon/4. \quad (10.11)$$

For each $\gamma \in \Gamma$ and $r > 0$, set

$$D(\gamma\xi_0, r) := \mathbb{B}_p(\gamma\xi_0, \frac{1}{3N_0}e^{-\psi(\underline{a}(\gamma^{-1}p, p) + i\underline{a}(\gamma^{-1}p, p))}r).$$

Lemma 10.12. Fix $R > 0$. If $\xi \in \Lambda$ and $\gamma_i \in \Gamma$ is a sequence such that $\gamma_i^{-1}p \rightarrow \eta$ and $\gamma_i^{-1}\xi \rightarrow \xi_0$ as $i \rightarrow \infty$, then for any $0 < r \leq s(\gamma_0)$, there exists $i_0 = i_0(r) > 0$ such that for all $i \geq i_0$,

$$D(\gamma_i\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon) \quad \text{and} \quad \xi \in D(\gamma_i\xi_0, r).$$

In particular, for any $R > 0$,

$$\Lambda_M \subset \bigcup_{D \in \mathcal{B}_R(\gamma_0, \varepsilon)} D.$$

Proof. Set $\Gamma_p := \{\gamma \in \Gamma : \psi(\underline{a}(\gamma^{-1}p, p) + i\underline{a}(\gamma^{-1}p, p)) > 0\}$; note that $\Gamma - \Gamma_p$ is a finite subset by Lemma 5.4. Hence we may assume that for all i , $\gamma_i \in \Gamma_p$. Since $\gamma_i^{-1}p \rightarrow \eta$ as $i \rightarrow \infty$, we may assume by Lemma 5.6 that for all i ,

$$O_{\varepsilon/(8\kappa)}(\eta, p) \subset O_{\varepsilon/(4\kappa)}(\gamma_i^{-1}p, p). \quad (10.13)$$

To prove that $D(\gamma_i\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$, we need to check that

$$\max_{\xi' \in \mathbb{B}_p(\gamma_i\xi_0, 3N_0s_i)} \|\beta_{\xi'}(p, \gamma_i\gamma_0^{\pm 1}\gamma_i^{-1}p) \mp \lambda(\gamma_0)\| < \varepsilon,$$

where $s_i = \frac{1}{3N_0}e^{-\psi(\underline{a}(\gamma_i^{-1}p, p) + i\underline{a}(\gamma_i^{-1}p, p))}r$. Let $\xi' \in \mathbb{B}_p(\gamma_i\xi_0, 3N_0s_i)$. We only prove that $\|\beta_{\xi'}(p, \gamma_i\gamma_0\gamma_i^{-1}p) - \lambda(\gamma_0)\| < \varepsilon$, as the other case can be treated similarly. First, observe

that

$$\begin{aligned}
d_p(\xi_0, \gamma_i^{-1}\xi') &= d_p(\gamma_i\xi_0, \xi') e^{\psi(\beta_{\xi_0}(\gamma_i^{-1}p, p) + i\beta_{\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p))} \\
&\leq e^{-\psi(\underline{a}(\gamma_i^{-1}p, p) + i\underline{a}(\gamma_i^{-1}p, p)) + \psi(\beta_{\xi_0}(\gamma_i^{-1}p, p) + i\beta_{\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p))} r \\
&\leq e^{2C} r \text{ by Theorem 5.3.}
\end{aligned} \tag{10.14}$$

Since $r \leq s(\gamma_0)$, this implies that

$$\|\beta_{\gamma_i^{-1}\xi'}(p, \gamma_0 p) - \lambda(\gamma_0)\| < \varepsilon/4.$$

Hence, by (6.3), we have

$$\begin{aligned}
d_p(\xi_0, \gamma_0^{-1}\gamma_i^{-1}\xi') &= e^{-\psi(\beta_{\xi_0}(\gamma_0 p, p) + i\beta_{\gamma_i^{-1}\xi'}(\gamma_0 p, p))} d_p(\xi_0, \gamma_i^{-1}\xi') \\
&\leq e^{\psi(\lambda(\gamma_0) + i\lambda(\gamma_0)) + \frac{1}{2}\|\psi\|\varepsilon + 2C} r.
\end{aligned} \tag{10.15}$$

Since $r \leq s(\gamma_0)$, it follows from (10.14), (10.15), and (10.10) that both $\gamma_i^{-1}\xi'$ and $\gamma_0^{-1}\gamma_i^{-1}\xi'$ belong to $\mathcal{O}_{\varepsilon/(8\kappa)}(\eta, p)$. Since, $\gamma_i^{-1}\xi', \gamma_0^{-1}\gamma_i^{-1}\xi' \in O_{\varepsilon/(4\kappa)}(\gamma_i^{-1}p, p)$ by (10.13), it follows from Lemma 5.7 that

$$\|\beta_{\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p)\| < 2\kappa(\varepsilon/4\kappa) = \varepsilon/2.$$

Now we have

$$\begin{aligned}
&\|\beta_{\xi'}(p, \gamma_i\gamma_0\gamma_i^{-1}p) - \lambda(\gamma_0)\| \\
&\leq \|\beta_{\xi'}(\gamma_i p, \gamma_i\gamma_0 p) - \lambda(\gamma_0)\| + \|\beta_{\xi'}(p, \gamma_i p) - \beta_{\xi'}(\gamma_i\gamma_0\gamma_i^{-1}p, \gamma_i\gamma_0 p)\| \\
&= \|\beta_{\gamma_i^{-1}\xi'}(p, \gamma_0 p) - \lambda(\gamma_0)\| + \|\beta_{\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\xi'}(\gamma_i^{-1}p, p)\| \\
&\leq \varepsilon/4 + \varepsilon/2 < \varepsilon,
\end{aligned}$$

which verifies that $D(\gamma_i\xi_0, r)$ belongs to the family $\mathcal{B}_R(\gamma_0, \varepsilon)$.

We now check that $\xi \in D(\gamma_i\xi_0, r)$. Since $\gamma_i^{-1}\xi \rightarrow \xi_0$, we may assume that for all i ,

$$d_p(\xi_0, \gamma_i^{-1}\xi) < \frac{1}{3N_0} e^{-\|\psi\|\varepsilon} r. \tag{10.16}$$

Since $r \leq s(\gamma_0)$, (10.10), (10.13), and (10.16) imply that $\gamma_i^{-1}\xi \in \mathcal{O}_{\varepsilon/(4\kappa)}(\gamma_i^{-1}p, p)$. Since $\xi_0 \in \mathcal{O}_{\varepsilon/(4\kappa)}(\gamma_i^{-1}p, p)$ as well, we have

$$\|\beta_{\gamma_i^{-1}\xi}(\gamma_i^{-1}p, p) - \underline{a}(\gamma_i^{-1}p, p)\| \leq \varepsilon/4 \text{ and } \|\beta_{\xi_0}(\gamma_i^{-1}p, p) - \underline{a}(\gamma_i^{-1}p, p)\| \leq \varepsilon/4,$$

by Lemma 5.7. Note that

$$\begin{aligned} d_p(\gamma_i\xi_0, \xi) &= d_{\gamma_i^{-1}p}(\xi_0, \gamma_i^{-1}\xi) \\ &= e^{-\psi(\beta_{\xi_0}(\gamma_i^{-1}p, p) + i\beta_{\gamma_i^{-1}\xi}(\gamma_i^{-1}p, p))} d_p(\xi_0, \gamma_i^{-1}\xi) \\ &\leq e^{-\psi(\underline{a}(\gamma_i^{-1}p, p) + i\underline{a}(\gamma_i^{-1}p, p) + \frac{1}{2}\|\psi\|\varepsilon)} d_p(\xi_0, \gamma_i^{-1}\xi) \\ &\leq \frac{1}{3N_0} e^{-\psi(\underline{a}(\gamma_i^{-1}p, p) + i\underline{a}(\gamma_i^{-1}p, p))} r \text{ by 10.16.} \end{aligned}$$

This proves that $\xi \in D(\gamma_i\xi_0, r)$. ■

Consider the following measure $\nu_p = \nu_{\psi, p}$ on Λ :

$$d\nu_p(\xi) = e^{\psi(\beta_{\xi}(o, p))} d\nu_{\psi}(\xi).$$

Proposition 10.17. Let $B \subset \mathcal{F}$ be a Borel subset with $\nu_p(B) > 0$. Then for ν_p -a.e. $\xi \in B$,

$$\lim_{R \rightarrow 0} \sup_{\xi \in D, D \in \mathcal{B}_R(\gamma_0, \varepsilon)} \frac{\nu_p(B \cap D)}{\nu_p(D)} = 1.$$

Proof. For a given Borel function $h : \mathcal{F} \rightarrow \mathbb{R}$, we define $h^* : \mathcal{F} \rightarrow \mathbb{R}$ by

$$h^*(\xi) = \lim_{R \rightarrow 0} \sup_{\xi \in D, D \in \mathcal{B}_R(\gamma_0, \varepsilon)} \frac{1}{\nu_p(D)} \int_D h \, d\nu_p.$$

By Lemma 10.12, h^* is well defined on Λ_M . Since Λ_M has a full ν_p measure by Theorem 8.11, h^* is defined ν_p -a.e. on \mathcal{F} . We will prove that $h = h^*$, ν_p -a.e.; by taking $h = \mathbf{1}_B$, the conclusion of the lemma will follow. Note that $h = h^*$ when h is continuous. To deal with the general case, we proceed as follows.

Step 1: For all $\alpha > 0$,

$$\nu_p(\{h^* > \alpha\}) \leq \frac{e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon}}{\alpha} \int_{\mathcal{F}} |h| \, d\nu_p.$$

Letting Q be an arbitrary compact subset of $\{\xi : h^*(\xi) > \alpha\}$, it suffices to show that

$$\nu_p(Q) \leq \frac{e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon}}{\alpha} \int_{\mathcal{F}} |h| \, d\nu_p.$$

Fix $R > 0$. By definition, for each $x \in Q$, there exists $D_x \in \mathcal{B}_R(\gamma_0, \varepsilon)$ containing x such that

$$\frac{1}{\nu_p(D_x)} \int_{D_x} h \, d\nu_p > \alpha.$$

Since K is compact, there exists a finite subcover of $\{D_x : x \in Q\}$, say $D_i = \mathbb{B}_p(\gamma_i \xi_0, s_i)$ ($i = 1, \dots, n$) where $\gamma_i \in \Gamma$ and $s_i = \frac{1}{3N_0} e^{-\psi(\underline{a}(\gamma_i^{-1} p, p) + i \underline{a}(\gamma_i^{-1} p, p))} r_i$ for some $0 < r_i < R$.

For brevity, we will write $3N_0 D_i := \mathbb{B}_p(\gamma_i \xi_0, 3N_0 s_i)$. By Lemma 6.12, there exists a disjoint subcollection $\{D_{i_1}, \dots, D_{i_\ell}\}$ such that

$$\bigcup_{k=1}^n D_k \subset \bigcup_{j=1}^\ell 3N_0 D_{i_j}.$$

Now we claim that $3N_0 D_{i_j} \subset \gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} D_{i_j}$: note that for $\xi \in 3N_0 D_{i_j}$,

$$\begin{aligned} d_p(\gamma_{i_j} \xi_0, \gamma_{i_j} \gamma_0 \gamma_{i_j}^{-1} \xi) &= d_{\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} p}(\gamma_{i_j} \xi_0, \xi) \\ &= e^{-\psi(\beta_{\gamma_{i_j} \xi_0}(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} p, p) + i \beta_\xi(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} p, p))} d_p(\gamma_{i_j} \xi_0, \xi) \\ &\leq 3N_0 e^{-\psi(\lambda(\gamma_0) + i \lambda(\gamma_0)) + \|\psi\|\varepsilon} s_{i_j} < s_{i_j}, \end{aligned}$$

by (6.3), (5.2), (10.9), and (10.8). Hence

$$\begin{aligned} \nu_p(3N_0 D_{i_j}) &\leq \nu_p(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} D_{i_j}) \\ &= \int_{D_{i_j}} e^{\psi(\beta_\xi(e, \gamma_{i_j} \gamma_0 \gamma_{i_j}^{-1}))} \, d\nu_p(\xi) \\ &\leq e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon} \nu_p(D_{i_j}), \end{aligned}$$

where the last inequality follows from (10.9). Therefore,

$$\begin{aligned} \nu_p(Q) &\leq \sum_{j=1}^{\ell} \nu_p(3N_0 D_{ij}) \leq \sum_{j=1}^{\ell} e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon} \nu_p(D_{ij}) \\ &\leq \frac{e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon}}{\alpha} \sum_{j=1}^{\ell} \int_{D_{ij}} h \, d\nu_p \leq \frac{e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon}}{\alpha} \int_{\mathcal{F}} |h| \, d\nu_p, \end{aligned}$$

which was to be proved.

Step 2: $h(\xi) = h^*(\xi)$ for ν_p -a.e ξ .

We first prove that $h(\xi) \leq h^*(\xi)$ for ν_p -a.e ξ . Let $\alpha > 0$ be arbitrary. It suffices to show that $\nu_p(\{\xi : h(\xi) - h^*(\xi) > \alpha\}) = 0$. Let h_n be a continuous function converging to h in $L^1(\nu_p)$. Note that $h_n^* = h_n$ and

$$\begin{aligned} \nu_p(\{\xi : h(\xi) - h^*(\xi) > \alpha\}) &\leq \nu_p(\{\xi : h(\xi) - h_n(\xi) > \alpha/2\}) + \nu_p(\{\xi : h_n^*(\xi) - h^*(\xi) > \alpha/2\}) \\ &\leq \frac{2}{\alpha} \|h - h_n\|_1 + \frac{2}{\alpha} e^{\psi(\lambda(\gamma_0)) + \|\psi\|\varepsilon} \|h - h_n\|_1. \end{aligned}$$

Taking $n \rightarrow \infty$, we get

$$\nu_p(\{\xi : h(\xi) - h^*(\xi) > \alpha\}) = 0.$$

As $\alpha > 0$ is arbitrary, it follows that $h \leq h^*$, ν_p -a.e. A similar argument shows that $h^* \leq h$, ν_p -a.e. ■

Proof of Proposition 10.7. It is easy to check that $E_\nu(\Gamma_0) = E_{\nu_p}(\Gamma_0)$. Hence it suffices to show $\lambda(\gamma_0) \in E_{\nu_p}(\Gamma_0)$. Let $B \subset \mathcal{F}$ be a Borel subset with $\nu_p(B) > 0$ and $\varepsilon > 0$. By Proposition 10.17, there exists $D = \mathbb{B}_p(\gamma\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$ for $\gamma \in \Gamma$ and $r > 0$ such that

$$\nu_p(D \cap B) > (1 + e^{-\psi(\lambda(\gamma_0)) - \|\psi\|\varepsilon})^{-1} \nu_p(D). \quad (10.18)$$
■

Since $r < r_p(\gamma)$, we have

$$\begin{aligned} D &\subset \{\xi : \|\beta_\xi(p, \gamma\gamma_0^\pm\gamma^{-1}p) \mp \lambda(\gamma_0)\| \leq \varepsilon\} \\ &\subset \{\xi : |\psi(\beta_\xi(p, \gamma\gamma_0^\pm\gamma^{-1}p)) \mp \psi(\lambda(\gamma_0))| \leq \|\psi\|\varepsilon\}. \end{aligned}$$

We note that $\gamma\gamma_0\gamma^{-1}D \subset D$: if $\xi \in D$, by (6.3),

$$\begin{aligned} d_p(\gamma\xi_0, \gamma\gamma_0\gamma^{-1}\xi) &= d_{\gamma\gamma_0^{-1}\gamma^{-1}p}(\gamma\xi_0, \xi) \\ &= e^{\psi(\beta_{\gamma\xi_0}(p, \gamma\gamma_0^{-1}\gamma^{-1}p) + i\beta_{\xi}(p, \gamma\gamma_0^{-1}\gamma^{-1}p))} d_p(\gamma\xi_0, \xi) \\ &\leq e^{-\psi(\lambda(\gamma_0) + i\lambda(\gamma_0)) + \|\psi\|\varepsilon} r < r. \end{aligned}$$

Since

$$B \cap \gamma\gamma_0\gamma^{-1}B \cap \{\xi : \|\beta_{\xi}(p, \gamma\gamma_0\gamma^{-1}p) - \lambda(\gamma_0)\| < \varepsilon\} \supset (D \cap B) \cap \gamma\gamma_0\gamma^{-1}(D \cap B),$$

it suffices to prove that $(D \cap B) \cap \gamma\gamma_0\gamma^{-1}(D \cap B)$ has a positive ν_p -measure. Note that

$$\begin{aligned} \nu_p(\gamma\gamma_0\gamma^{-1}(D \cap B)) &= \int_{D \cap B} e^{\psi(\beta_{\xi}(p, \gamma\gamma_0^{-1}\gamma^{-1}p))} d\nu_p(\xi) \\ &\geq e^{-\psi(\lambda(\gamma_0)) - \|\psi\|\varepsilon} \nu_p(D \cap B). \end{aligned}$$

Hence by (10.18),

$$\nu_p(D \cap B) + \nu_p(\gamma\gamma_0\gamma^{-1}(D \cap B)) > (1 + e^{-\psi(\lambda(\gamma_0)) - \|\psi\|\varepsilon}) \nu_p(D \cap B) > \nu_p(D).$$

Since both $D \cap B$ and $\gamma\gamma_0\gamma^{-1}(D \cap B)$ are contained in D , this implies that their intersection has a positive ν_p -measure. Since $\gamma\gamma_0\gamma^{-1} \in \Gamma_0$, it follows that $\lambda(\gamma_0) \in E_{\nu_p}(\Gamma_0)$.

In view of Proposition 9.2, we obtain the following corollary:

Corollary 10.19. Let Γ_0 be a Zariski dense normal subgroup of an Anosov subgroup $\Gamma < G$. Let $\psi \in D_{\Gamma}^*$. If ν_{ψ} is Γ_0 -ergodic, then m_{ψ}^{BR} , considered as a measure on $\Gamma_0 \backslash G$, is NM-ergodic.

Patterson–Sullivan measures are mutually singular

Theorem 10.20. Let $\Gamma < G$ be an Anosov subgroup. Then $\{\nu_{\psi} : \psi \in D_{\Gamma}^*\}$ are pairwise mutually singular.

Proof. Since $\Gamma < G$ is Anosov, the family $\{\nu_{\psi} : \psi \in D_{\Gamma}^*\}$ consists of Γ -ergodic measures (see the remark following Theorem 4.3). Hence any ν_{ψ_1} and ν_{ψ_2} in this family are either mutually singular or absolutely continuous to each other. Now the claim follows from Lemma 10.21 below, in view of Proposition 10.2. ■

Lemma 10.21. For $i = 1, 2$, let ν_{ψ_i} be a (Γ, ψ_i) -PS measure for some $\psi_i \in \mathfrak{a}^*$. If $E_{\nu_{\psi_2}} = \mathfrak{a}$ and $\nu_{\psi_1} \ll \nu_{\psi_2}$, then $\psi_1 = \psi_2$.

Proof. Suppose that $\nu_{\psi_1} \ll \nu_{\psi_2}$ and that $\psi_1 \neq \psi_2$. Consider the Radon–Nikodym derivative $f := \frac{d\nu_{\psi_1}}{d\nu_{\psi_2}} \in L^1(\Lambda, \nu_{\psi_2})$. Note that there exists a ν_{ψ_2} -conull set $E \subset \Lambda$ such that for all $\xi \in E$ and $\gamma \in \Gamma$, we have

$$f(\gamma^{-1}\xi) = e^{(\psi_1 - \psi_2)(\beta_\xi(e, \gamma))}f(\xi). \quad (10.22)$$

If f were continuous, then $f \neq 0$ and by applying $\xi = y_\gamma$ in the above, we get $\psi_1(\lambda(\gamma)) = \psi_2(\lambda(\gamma))$ for all $\gamma \in \Gamma$. Since $\lambda(\Gamma)$ generates a dense subgroup of \mathfrak{a} , it follows that $\psi_1 = \psi_2$.

In general, we use the hypothesis $E_{\nu_{\psi_2}} = \mathfrak{a}$. Choose $0 < r_1 < r_2$ such that

$$B := \{\xi \in \Lambda : r_1 < f(\xi) < r_2\}$$

has a positive ν_{ψ_2} -measure. Since $\psi_1 \neq \psi_2$, we can choose $w \in \mathfrak{a}$ such that

$$e^{(\psi_1 - \psi_2)(w)} > \frac{2r_2}{r_1}. \quad (10.23)$$

Choose $\varepsilon > 0$ such that $e^{\|\psi_1 - \psi_2\|\varepsilon} < 2$. Since $\nu_{\psi_2}(B) > 0$ and $E_{\nu_{\psi_2}} = \mathfrak{a}$, there exists $\gamma \in \Gamma$ such that

$$B' := B \cap \gamma B \cap \{\xi \in \Lambda : \|\beta_\xi(e, \gamma) - w\| < \varepsilon\}$$

has a positive ν_{ψ_2} -measure. Now note that

$$\begin{aligned} \int_{B'} f(\gamma^{-1}\xi) d\nu_{\psi_2}(\xi) &> e^{(\psi_1 - \psi_2)(w) - \|\psi_1 - \psi_2\|\varepsilon} \int_{B'} f(\xi) d\nu_{\psi_2}(\xi) \\ &> \frac{r_2}{r_1} \int_{B'} f(\xi) d\nu_{\psi_2}(\xi) \end{aligned}$$

by (10.22), (10.23), and the choice of ε . In particular,

$$\nu_{\psi_2} \left\{ \xi \in B' : f(\gamma^{-1}\xi) > \frac{r_2}{r_1} f(\xi) \right\} > 0.$$

It follows that there exists $\xi \in B' \cap E$ such that

$$f(\gamma^{-1}\xi) > \frac{r_2}{r_1}f(\xi). \quad (10.24)$$

On the other hand, for $\xi \in B'$, both ξ and $\gamma^{-1}\xi$ belong to B . Hence, by definition of B , for all $\xi \in B'$, we have

$$f(\gamma^{-1}\xi) < \frac{r_2}{r_1}f(\xi).$$

This is a contradiction to (10.24). ■

P-semi-invariant measures

In this section, we establish that *P*-semi-invariant Radon measures supported in $\mathcal{E} = \{x \in \Gamma \backslash G : x^+ \in \Lambda\}$, up to constant multiples, are parametrized by D_Γ^* .

If μ is *P*-semi-invariant, then there exists a linear form $\chi_\mu \in \mathfrak{a}^*$ such that for all $a \in A$,

$$a_*\mu = e^{-\chi_\mu(\log a)}\mu.$$

We set $\psi_\mu := \chi_\mu + 2\rho \in \mathfrak{a}^*$. The first part of the following proposition is known in the rank one case (see e.g., [3], [10], and [28]), and the proof can be easily adapted to the higher rank case.

Proposition 10.25. For any Zariski dense discrete subgroup $\Gamma < G$, any *P*-semi-invariant Radon measure μ on $\Gamma \backslash G$ is proportional to $m_{v_{\psi_\mu}, m_0}$ where v_{ψ_μ} is a (Γ, ψ_μ) -conformal measure and $\psi_\mu \in D_\Gamma$. Moreover, if μ is supported on \mathcal{E} , then μ is proportional to $m_{\psi_\mu}^{\text{BR}}$. If Γ is Anosov, we also have $\psi_\mu \in D_\Gamma^*$.

Proof. For simplicity, set $\chi = \chi_\mu$ and $\psi = \psi_\mu$. Let $\tilde{\mu}$ be the Γ -invariant lift of μ to G and $\pi : G \rightarrow G/P$ be the projection. Choose a section $c : G/P \rightarrow K$ so that $\pi \circ c = \text{id}$ and consider the measurable isomorphism

$$\begin{aligned} G/P \times M \times A \times N &\rightarrow G \\ (\xi, m, a, n) &\rightarrow c(\xi)man. \end{aligned}$$

Let dm , dn , and da be the Haar measures on M , N , and A . As $\tilde{\mu}$ is a P -semi-invariant Radon measure, there exist $\tilde{\chi} \in \mathfrak{a}^*$ and a Radon measure ν on G/P such that

$$d\tilde{\mu}(c(\xi)man) = e^{\tilde{\chi}(\log a)} dn da dm d\nu(\xi).$$

Without loss of generality, we may assume that $|\nu| = 1$. Because $d\tilde{\mu}(\cdot a) = e^{\chi(\log a)} d\tilde{\mu}(\cdot)$, we have

$$\chi = \tilde{\chi} - 2\rho, \text{ or equivalently, } \tilde{\chi} = \psi.$$

Note that G is measurably isomorphic to the product $G/P \times P$ and the left Γ -action with respect to these coordinates is given by $\gamma \cdot (\xi, p) = (\gamma \cdot \xi, \Phi(\gamma, \xi)p)$ for some P -valued cocycle $\Phi : \Gamma \times G/P \rightarrow P$ where $\gamma \in \Gamma$ and $(\xi, p) \in G/P \times P$. One can check that

$$\Phi(\gamma, \xi) = m(\gamma, \xi) \exp(\beta_\xi(\gamma^{-1}, e)) n(\gamma, \xi)$$

for some $m(\gamma, \xi) \in M$ and $n(\gamma, \xi) \in N$. Hence, for $p = man$, the MAN -coordinates for $\Phi(\gamma, \xi)p$ are given by

$$\Phi(\gamma, \xi)p = (m(\gamma, \xi)m) (\exp(\beta_\xi(\gamma^{-1}, e))a) ((ma)^{-1}n(\gamma, \xi)man). \quad (10.26)$$

Since $\tilde{\mu}$ is left- Γ -invariant, we have for any $f \in C_c(G)$ and any $\gamma \in \Gamma$,

$$\begin{aligned} \int_G f(g) d\tilde{\mu}(g) &= \int_G f(g) d(\gamma_* \tilde{\mu})(g) \\ &= \int_{G/P} \int_P f((\gamma \xi, \Phi(\gamma, \xi)p) e^{\psi(\log a)} dn da dm d\nu(\xi) \\ &= \int_{G/P} \int_P f(\xi, p) e^{\psi(\log a - \beta_{\gamma^{-1}\xi}(\gamma^{-1}, e))} dn da dm d(\gamma_* \nu)(\xi), \end{aligned}$$

where in the last equality, we have used (10.26) and the change of variables $a' = a \exp(\beta_\xi(e, \gamma^{-1}))$. On the other hand, we have

$$\int_G f(g) d\tilde{\mu}(g) = \int_{G/P} \int_P f(\xi, p) e^{\psi(\log a)} dn da dm d\nu(\xi).$$

By comparing these two identities, we get that for any $\gamma \in \Gamma$,

$$d(\gamma_* \nu)(\xi) = e^{\psi(\beta_\xi(e, \gamma))} d\nu(\xi),$$

that is, ν is a (Γ, ψ) -conformal measure. By [40, Thm. 8.1], $\psi \in D_\Gamma$.

Finally, recall that for all $g \in G$ and $\phi \in C_c(G)$,

$$\int_N \phi(gn) dn = \int_{G/P} \phi(gn) e^{2\rho(\beta_{gn^-}(e, gn))} dm_o(gn^-).$$

For $g = c(\xi)man \in KAN$, we have $\beta_{g^+}(e, g) = \log a$ and $g^+ = \xi$. Hence, for any $f \in C_c(G)$,

$$\begin{aligned} \int_G f(g) d\tilde{\mu}(g) &= \int_{G/P} \int_P f(c(\xi)man) e^{\psi(\log a)} dn da dm d\nu(\xi) \\ &= \int_{G/M} \int_M f(g) e^{2\rho(\beta_{g^-}(e, g))} e^{\psi(\beta_{g^+}(e, g))} dm da dm_o(g^-) d\nu(g^+) \\ &= \tilde{m}_{\nu, m_o}(f). \end{aligned}$$

Therefore, $\tilde{\mu}(f) = \tilde{m}_{\nu, m_o}(f)$.

Now, if μ is supported on \mathcal{E} , then ν is supported on Λ . Hence ν is a (Γ, ψ) -PS measure; so $\mu = m_\psi^{\text{BR}}$. When Γ is Anosov, $\psi \in D_\Gamma^*$ by Theorem 7.7. \blacksquare

Let \mathcal{P}_Γ be the space of all P -semi-invariant Radon measures on \mathcal{E} up to proportionality. Let \mathcal{Q}_Γ be the space of all NM -invariant, ergodic, and A -quasi-invariant Radon measures supported on \mathcal{E} up to proportionality.

Theorem 10.27. Let $\Gamma < G$ be an Anosov subgroup. We have $\mathcal{P}_\Gamma = \mathcal{Q}_\Gamma$ and the map $D_\Gamma^* \rightarrow \mathcal{Q}_\Gamma$ given by $\psi \mapsto [m_\psi^{\text{BR}}]$ is a homeomorphism between D_Γ^* and \mathcal{Q}_Γ . In particular, \mathcal{Q}_Γ is homeomorphic to $\mathbb{R}^{\text{rank } G-1}$.

Proof. For $\mu \in \mathcal{Q}_\Gamma$ and $a \in A$, $a_*\mu$ and μ are equivalent to each other, and by the NM -ergodicity of μ , the Radon–Nikodym derivative $da_*\mu/d\mu$ is constant, say $\chi(a)$. Now the function $a \mapsto \chi(a)$ gives the semi-invariance of μ by A and hence by P . This implies $\mathcal{Q}_\Gamma \subset \mathcal{P}_\Gamma$. The other direction $\mathcal{P}_\Gamma \subset \mathcal{Q}_\Gamma$ follows from Proposition 10.25 and Theorem 10.1.

Let $\mathcal{Q}_\Gamma^\blacklozenge$ be the space of all NM -ergodic A -quasi-invariant Radon measures supported on $\{x \in \Gamma \backslash G : x^+ \in \Lambda\}$, so that $\mathcal{Q}_\Gamma = \mathcal{Q}_\Gamma^\blacklozenge / \sim$. Set $\iota(\psi) = m_\psi^{\text{BR}}$ for $\psi \in D_\Gamma^*$. Since m_ψ^{BR} is NM -ergodic by Theorem 10.1, the map $\iota : D_\Gamma^* \rightarrow \mathcal{Q}_\Gamma^\blacklozenge$ is well defined and injective by Lemma 10.21. By Proposition 10.25, $\iota(D_\Gamma^*)$ contains precisely one representative of each class in \mathcal{Q}_Γ . Hence it suffices to show that the map ι gives a homeomorphism between D_Γ^* and its image $\iota(D_\Gamma^*)$. Continuity of ι follows from Theorem 7.7. Now, suppose that $m_{\psi_i}^{\text{BR}} \rightarrow m_\psi^{\text{BR}}$ for some sequence $\psi_i, \psi \in D_\Gamma^*$. Then the

A -semi-invariance of the BR-measures given by (3.20) and the convergence $a_*m_{\psi_i}^{\text{BR}} \rightarrow a_*m_{\psi}^{\text{BR}}$ implies that $\lim_{i \rightarrow \infty} e^{(2\rho - \psi_i)(\log a)} m_{\psi_i}^{\text{BR}}(f) = e^{(2\rho - \psi)(\log a)} m_{\psi}^{\text{BR}}(f)$ for all $f \in C_c(\Gamma \backslash G)$. Since $\lim_{i \rightarrow \infty} m_{\psi_i}^{\text{BR}}(f) = m_{\psi}^{\text{BR}}(f)$, we get $\lim_{i \rightarrow \infty} e^{(2\rho - \psi_i)(\log a)} = e^{(2\rho - \psi)(\log a)}$ for all $a \in A$. Hence $\psi_i \rightarrow \psi$. This proves that D_{Γ}^* and \mathcal{Q}_{Γ} are homeomorphic to each other. The last claim follows from Proposition 4.4. \blacksquare

Funding

This work was supported in part by NSF grants DMS-1900101 [to H.O.].

References

- [1] Agard, S. "A geometric proof of Mostow's rigidity theorem for groups of divergence type." *Acta Math.* 151 (1983): 231–52. <https://doi.org/10.1007/BF02393208>.
- [2] Albuquerque, P. "Patterson–Sullivan theory in higher rank symmetric spaces." *Geom. Funct. Anal.* 9 (1999): 1–28. <https://doi.org/10.1007/s000390050079>.
- [3] Babillot, M. "On the Classification of Invariant Measures for Horosphere Foliations on Nilpotent Covers of Negatively Curved Manifolds." *Random Walks and Geometry*, 319–35. Walter de Gruyter, Berlin, 2004, <https://doi.org/10.1515/9783110198089.2.319>.
- [4] Babillot, M. and F. Ledrappier. "Geodesic Paths and Horocycle Flow on Abelian Covers." *Lie Groups and Ergodic Theory (Mumbai, 1996)*. TIFR Stud. Math., vol. 14. 1–32.
- [5] Benoist, Y. "Actions propres sur les espaces homogenes réductifs." *Ann. Math.* (2) 144 (1996): 315–47.
- [6] Benoist, Y. "Propriétés asymptotiques des groupes linéaires." *Geom. Funct. Anal.* 7, no. 1 (1997): 1–47. <https://doi.org/10.1007/PL00001613>.
- [7] Benoist, Y. "Propriétés asymptotiques des groupes linéaires. II." *Adv. Stud. Pure Math.* 26. Tokyo, Japan: Math. Soc., 2000.
- [8] Bridgeman, M., R. Canary, F. Labourie, and A. Sambarino. "The pressure metric for Anosov representations." *Geom. Funct. Anal.* 25 (2015): 1089–179. <https://doi.org/10.1007/s00039-015-0333-8>.
- [9] Bridson, M. and A. Haefliger. "Metric Spaces of Non-Positive Curvature." *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 319. Springer, Berlin, 1999, <https://doi.org/10.1007/978-3-662-12494-9>.
- [10] Burger, M. "Horocycle flow on geometrically finite surfaces." *Duke Math. J.* 61, no. 3 (1990): 779–803. <https://doi.org/10.1215/S0012-7094-90-06129-0>.
- [11] Burger, M. "Intersection, the Manhattan curve and Patterson–Sullivan theory in rank 2." *Int. Math. Res. Not. IMRN* 7 (1993): 217–25. <https://doi.org/10.1155/S1073792893000236>.
- [12] Carvajales, L. "Growth of quadratic forms under Anosov subgroups." *IMRN*. Preprint arXiv:2004.05903,.
- [13] Chow, M. and P. Sarkar. "Local mixing of one-parameter diagonal flows on Anosov homogeneous spaces." Preprint arXiv:2105.11377.

- [14] Dani, S. G. "Invariant measures and minimal sets of horospherical flows." *Invent. Math.* 64, no. 2 (1981): 357–85. <https://doi.org/10.1007/BF01389173>.
- [15] Edwards, S., M. Lee, and H. Oh. "Anosov groups: local mixing, counting, and equidistribution." *Geom. Topol.* Preprint arXiv:2003.14277.
- [16] Edwards, S., M. Lee, and H. Oh. "Uniqueness of conformal measures and local mixng for Anosov groups." *Michigan Math. J. (Prasad volume)* 72 (2022): 243–59.
- [17] Furstenberg, H. "The Unique Ergodicity of the Horocycle Flow." *Recent Advances in Topological Dynamics (Proc. Conf. Yale U. 1972 in honor of Hedlund)*. Lecture Notes in Math., vol 318. Berlin: Springer, 1973.
- [18] Fock, V. and A. Goncharov. "Moduli spaces of local systems and higher Teichmuller theory." *Publ. Math. Inst. Hautes Études Sci.* 103 (2006): 1–211. <https://doi.org/10.1007/s10240-006-0039-4>.
- [19] Ghys, E. and P. de la Harpe. *Sur les Groupes Hyperboliques d'apr s Mikhael Gromov*. Boston, MA: Birkh user, 1990, <https://doi.org/10.1007/978-1-4684-9167-8>.
- [20] Gorodnik, A. and H. Oh. "Orbits of discrete subgroups on a symmetric space and the Furstenberg boundary." *Duke Math. J.* 139 (2007): 483–525. <https://doi.org/10.1215/S0012-7094-07-13933-4>.
- [21] Gueritaud, F., O. Guichard, F. Kassel, and A. Wienhard. "Anosov representations and proper actions." *Geom. Topol.* 21, no. 1 (2017): 485–584. <https://doi.org/10.2140/gt.2017.21.485>.
- [22] Guichard, O. and A. Wienhard. "Anosov representations: domains of discontinuity and applications." *Invent. Math.* 190, no. 2 (2012): 357–438. <https://doi.org/10.1007/s00222-012-0382-7>.
- [23] Hitchin, N. J. "Lie groups and Teichmuller space." *Topology* 31, no. 3 (1992): 449–73. [https://doi.org/10.1016/0040-9383\(92\)90044-I](https://doi.org/10.1016/0040-9383(92)90044-I).
- [24] Kapovich, I. and N. Benakli. "Boundaries of Hyperbolic Groups." *Combinatorial and Geometric Group Theory (New York, 2000/Hoboken, NJ, 2001)*, 39–93. Contemp. Math., 296. Amer. Math. Soc. Colloq. Publ., Providence, RI. <https://doi.org/10.1090/conm/296/05068>
- [25] Kapovich, M., B. Leeb, and J. Porti. "Anosov subgroups: dynamical and geometric characterizations." *Eur. J. Math.* 3, no. 4 (2017): 808–98. <https://doi.org/10.1007/s40879-017-0192-y>.
- [26] Kassel, F. "Geometric Structures and Representations of Discrete Groups." *Proceedings of the International Congress of Mathematicians—Rio de Janeiro*, vol. II, 1115–51. 2018.
- [27] Labourie, F. "Anosov flows, surface groups and curves in projective space." *Invent. Math.* 165, no. 1 (2006): 51–114. <https://doi.org/10.1007/s00222-005-0487-3>.
- [28] Landesberg, O. and E. Lindenstrauss. "On Radon measures invariant under horospherical flows on geometrically infinite quotients." *IMRN* (2019): Preprint, arXiv:1910.08956.
- [29] Landesberg, O. "Horospherically invariant measures and finitely generated Kleinian groups." *J. Modern Dyn.* 17 (2021): 337–52. <https://doi.org/10.3934/jmd.2021012>.
- [30] Lee, M. and H. Oh. "Ergodic decompositions of geometric measures on Anosov homogeneous spaces." *Israel J. Math.* Preprint, arXiv:2010.11337.

- [31] Lee, M. and H. Oh. "Dichotomy and measures on limit sets of Anosov groups." Preprint, arXiv:2203.06794.
- [32] Ledrappier, F. "Invariant measures for the stable foliation on negatively curved periodic manifolds." *Univ. Grenoble. Ann. Inst. Fourier. Univ. Grenoble I* 58 (2008): 85–105. <https://doi.org/10.5802/aif.2345>.
- [33] Ledrappier, F. "Structure au Bord des Variétés à Courbure Négative." *Séminaire de Théorie Spectrale et Géométrie, No. 13*. Année 1994–1995, 97–122.
- [34] Myrberg, P. J. "Ein approximationssatz für die Fuchschen Gruppen." *Acta Math.* 57 (1931): 389–409. <https://doi.org/10.1007/BF02403050>.
- [35] Nakanishi, T. and P. J. On. "Myrberg's approximation theorem for some Kleinian groups." *J. Math. Kyoto Univ.* 25 (1985): 405–19. <https://doi.org/10.1215/kjm/1250521061>.
- [36] Oh, H. and W. Pan. "Local mixing and invariant measures for horospherical subgroups on abelian covers." *Int. Math. Res. Not. IMRN.* 2019 (2019): 6036–88. <https://doi.org/10.1093/imrn/rnx292>.
- [37] Patterson, S. J. "The limit set of a Fuchsian group." *Acta Math.* 136 (1976): 241–73. <https://doi.org/10.1007/BF02392046>.
- [38] Potrie, R. and A. Sambarino. "Eigenvalues and entropy of a Hitchin representation." *Invent. Math.* 209, no. 3 (2017): 885–925. <https://doi.org/10.1007/s00222-017-0721-9>.
- [39] Quint, J.-F. "Divergence exponentielle des sous-groupes discrets en rang supérieur." *Comment. Math. Helv.* 77, no. 3 (2002): 563–608. <https://doi.org/10.1007/s00014-002-8352-0>.
- [40] Quint, J.-F. "Mesures de Patterson–Sullivan en rang supérieur." *Geom. Funct. Anal.* 12, no. 4 (2002): 776–809. <https://doi.org/10.1007/s00039-002-8266-4>.
- [41] Quint, J.-F. "L'indicateur de croissance des groupes de Schottky." *Ergodic Theory Dynam. Systems* 23, no. 1 (2003): 249–72. <https://doi.org/10.1017/S0143385702001268>.
- [42] Quint, J.-F. "Groupes de Schottky et comptage." *Univ. Grenoble. Ann. Inst. Fourier. Univ. Grenoble I* 55 (2005): 373–429. <https://doi.org/10.5802/aif.2102>.
- [43] Quint, J.-F. "Propriété de Kazhdan et sous-groupes discrets de covolume infini." *Trav. Math.* 14 (2003): 143–51. Univ. Luxemb., Luxembourg.
- [44] Quint, J.-F. *An Overview of Patterson–Sullivan Theory*. Lecture Notes available at math.u-bordeaux.fr/jquint/.
- [45] Ratner, M. "On Raghunathan's measure conjecture." *Ann. Math. (2)* 134 (1991): 545–607.
- [46] Roblin, T. "Ergodicité et équidistribution en courbure négative." *Mém. Soc. Math. Fr. (N.S.)* 95 (2003): vi+96.
- [47] Sambarino, A. "The orbital counting problem for Anosov representations." *Ann. Inst. Fourier (Grenoble) (Grenoble)* 65, no. 4 (2015): 1755–97. <https://doi.org/10.5802/aif.2973>.
- [48] Sambarino, A. "Hyperconvex representations and exponential growth." *Ergodic Theory Dynam. Systems* 34, no. 3 (2014): 986–1010. <https://doi.org/10.1017/etds.2012.170>.
- [49] Sambarino, A. "Quantitative properties of convex representations." *Comment. Math. Helv.* 89, no. 2 (2014): 443–88. <https://doi.org/10.4171/CMH/324>.
- [50] Sarig, O. "Invariant Radon measures for horocycle flows on abelian covers." *Invent. Math.* 157 (2004): 519–51. <https://doi.org/10.1007/s00222-004-0357-4>.

- [51] Sarig, O. "The horocycle flow and the Laplacian on hyperbolic surfaces of infinite genus." *Geom. Funct. Anal.* 19 (2010): 1757–812. <https://doi.org/10.1007/s00039-010-0048-9>.
- [52] Schmidt, K. "Cocycles on ergodic transformation groups." *Macmillan Lectures in Mathematics*, vol. 1. 202. Delhi: Macmillan Company of India, Ltd, 1977.
- [53] Sullivan, D. "The density at infinity of a discrete group of hyperbolic motions." *Publ. Math. Inst. Hautes Études Sci.* 50 (1979): 171–202. <https://doi.org/10.1007/BF02684773>.
- [54] Thirion, X. "Sous-groupes discrets de $SL(d, \mathbb{R})$ et équidistribution dans les espaces symétriques." PhD thesis, Université de Tours, 2007.
- [55] Tits, J. "Représentations linéaires irréductibles d'un groupe réductif sur un corps quelconque." *J. Reine Angew. Math.* 1971 (1971): 196–220. <https://doi.org/10.1515/crll.1971.247.196>.
- [56] Tukia, P. "The Poincaré series and the conformal measure of conical and Myrberg limit points." *J. Anal. Math.* 62 (1994): 241–59. <https://doi.org/10.1007/BF02835956>.
- [57] Veech, W. "Unique ergodicity of horospherical flows." *Am. J. Math.* 99 (1977): 827–59. <https://doi.org/10.2307/2373868>.
- [58] Wienhard, A. "An Invitation to Higher Teichmüller Theory." *Proceedings of the International Congress of Mathematicians—Rio de Janeiro*, vol. II, 1013–39. 2018.
- [59] Winter, D. "Mixing of frame flow for rank one locally symmetric spaces and measure classification." *Israel J. Math.* 210, no. 1 (2015): 467–507.