



Studies

SPECIAL ISSUE: The Role of Seed Dispersal in Plant Populations: Perspectives and Advances in a Changing World**Low-intensity logging and hunting have long-term effects on seed dispersal but not fecundity in Afrotropical forests**Chase L. Nuñez^{1,2*}, James S. Clark^{1,2}, Connie J. Clark² and John R. Poulsen^{1,2}¹University Program in Ecology, Duke University, Durham, NC 27708, USA²Nicholas School of the Environment, Duke University, Durham, NC 27708, USA**Received:** 30 August 2018 **Editorial decision:** 30 November 2018 **Accepted:** 10 December 2018 **Published:** 13 December 2018**Associate Editor:** Noelle Beckman**Citation:** Nuñez CL, Clark JS, Clark CJ, Poulsen JR. 2018. Low-intensity logging and hunting have long-term effects on seed dispersal but not fecundity in Afrotropical forests. *AoB PLANTS* **10**: ply074; doi: 10.1093/aobpla/ply074

Abstract. Hunting and logging, ubiquitous human disturbances in tropical forests, have the potential to alter the ecological processes that govern population recruitment and community composition. Hunting-induced declines in populations of seed-dispersing animals are expected to reduce dispersal of the tree species that rely on them, resulting in potentially greater distance- and density-dependent mortality. At the same time, selective logging may alter competitive interactions among tree species, releasing remaining trees from light, nutrient or space limitations. Taken together, these disturbances may alter the community composition of tropical forests, with implications for carbon storage, biodiversity conservation and ecosystem function. To evaluate the effects of hunting and logging on tree fecundity and seed dispersal, we use 3 years of seed rain data from a large-scale observational experiment in previously logged, hunted and protected forests in northern Republic of Congo (Brazzaville). We find that low-intensity logging had a meaningful long-term effect on species-specific seed dispersal distances, though the direction and magnitude varied and was not congruent within dispersal vector. Tree fecundity increased with tree diameter, but did not differ appreciably across disturbance regimes. The species-specific dispersal responses to logging in this study point towards the long-lasting toll of disturbance on ecological function and highlight the necessity of conserving intact forest.

Keywords: Anthropocene; dispersal; frugivory; habitat fragmentation; hunting; selective logging; tropical forest.

Introduction

Logging concessions now cover almost 56 million ha of forest in West and Central Africa (FAO 2016). Most concessions are subject to low-intensity, selective logging

intended to reduce the negative ecological impacts of traditional, conventional logging operations. Studies across the tropics have demonstrated that selective logging techniques can substantially reduce the short-term

*Corresponding author's e-mail address: cln12@duke.edu

© The Author(s) 2018. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

effects of logging (Sist 2000; Sist et al. 2003; Medjibe et al. 2013), but few studies have considered the long-term effects of selective logging on critical forest processes (Brown and Gurevitch 2004; Meijaard et al. 2005). Tropical trees respond to environmental disturbance on timescales that usually surpass the duration of ecological studies (Gourlet-Fleury et al. 2013; Edwards et al. 2014; Berdanier and Clark 2015) and changes in tree fecundity and seed dispersal may persist long after disturbance has ended, potentially altering ecosystem function.

Logging directly disturbs tropical forest communities through the extraction of large trees (Laurance et al. 2000), residual damage to remaining trees (Kasenene and Murphy 1991) and disruption of seed-dispersing animal communities (Gutiérrez-Granados 2011; Haurez et al. 2016; Rosin and Poulsen 2016). Road construction fragments the forest and provides hunters access to previously inaccessible areas (Kleinschroth and Healey 2017). Unsustainable hunting is the major cause of defaunation in many parts of the world (Hoffmann et al. 2010), causing over a quarter of the world's vertebrate species to decline in abundance over the last four decades (Dirzo et al. 2014). Reductions in vertebrate dispersers may affect the approximately two-thirds of all woody plants that rely on animals for seed dispersal (Willson and Traveset 2000; Muller-Landau and Hardesty 2005; Beaune et al. 2013). Dispersal failure has consequences for community composition through density-dependent recruitment (Cannon et al. 1994; Bleher and Böhning-Gaese 2001) and competition at later life stages (Nathan and Muller-Landau 2000).

Studies investigating how hunting and logging affect seed dispersal have yielded mixed results (Theimer et al. 2011; Beck et al. 2013; Kurten 2013; Camargo-Sanabria et al. 2014; Comita et al. 2014; Rosin and Poulsen 2016) in part because the interacting effects of hunting and logging have not been quantified beyond their immediate responses to disturbances (Markl et al. 2012). In the short term, intermediate levels of disturbance from selective logging may increase light and nutrients available to survivors (Johns 1988; Kasenene and Murphy 1991; Cannon et al. 1994; Huante et al. 1998; John et al. 2007; Ewel and Mazzarino 2008; Gutiérrez-Granados 2011; Haurez et al. 2016), thereby increasing tree fecundity (Molino and Sabatier 2001; Clark et al. 2010, 2014b). Logging may even increase the dispersal distance of abiotically dispersed species following forest thinning due to greater wind speeds through the canopy (Gardiner 1994; Stacey et al. 1994; Gardiner et al. 1997). However, in the longer term, logging may reduce seed dispersal distance and fecundity through combinations of increased hunting pressure (Kleinschroth and Healey 2017), declines in vertebrate dispersal vectors (Poulsen et al. 2013; Haurez et al. 2016), soil compaction (Pinard et al. 2000) and invasion of fast-growing competitors (Schnitzer and Bongers 2002). Because

declines in dispersal vectors and increases in fecundity can both follow disturbance, investigating the interactions of these processes is essential for understanding the underlying ecological process (Abernethy et al. 2013).

To evaluate the separate and combined effects of hunting and logging on both fecundity and dispersal for animal and abiotically dispersed trees, we collected 3 years of seed rain data from a large-scale observational experiment in previously logged, hunted and protected forests in northern Republic of Congo (Brazzaville). By controlling for logging and hunting in our sampling design, we offer a first opportunity to test their relative effects. We hypothesized that the fecundity and dispersal distances of tropical trees will be sensitive to both hunting and logging. Specifically, we expected that: (i) tree fecundity is greater in logged forests relative to protected forests, regardless of whether trees species are abiotically or animal dispersed; and (ii) hunting reduces dispersal distances of animal-dispersed species, but not the dispersal distances of abiotically (wind or ballistic) dispersed species. Understanding the separate and combined effects of disturbances on seed dispersal is critical to predict changes in forest species composition and diversity.

Materials and Methods

Study area

We conducted the study in the Nouabale Ndoki National Park (NNNP; 400 000 ha) and the Kabo logging concession (267 000 ha) in northern Republic of Congo (Fig. 1). The forests in this area are classified as lowland tropical forest. Dominant tree families include *Meliaceae*, *Euphorbiaceae* and *Annonaceae* (CIB 2006). Rainfall averages ~1700 mm annually and is seasonal with peaks in May and October. The Kabo concession borders the NNNP to the south, and together they include a mosaic of logged and unlogged forest. Twenty years before the study began, the logging concession was selectively logged at low intensity (<2.5 stems per hectare) with four species, *Entandophragma cylindricum*, *E. utile*, *Triplochiton scleroxylon* and *Milicia excelsa*, making up 90 % of the harvest volume (CIB 2006). Although we do not have data on rates of natural disturbance at our study site, a comparison of pantropical data ($n = 65$) report a range of natural stand mortality from 0.86 to 2.02 %, with a best estimate of adjusted stem turnover rate of 1.81 ± 0.16 % (Lewis et al. 2004). Approximately 3000 people inhabited the study site at the time of the study, most residing in the logging town of Kabo. Residents generally hunted with shotguns, and to a lesser extent with wire snares, for consumption and for local trade (Poulsen et al. 2009). A gradient of hunting intensity decreases with distance from Kabo, with some forest types being used more than others (Mockrin 2008).




Figure 1. Location of 30 1-ha study plots in Northern Congo. Protected plots fall within the border of Nouabale-Ndoki National Park (green), whereas plots exposed to hunting and/or logging were located in the Kabo logging concession (grey) in northern Republic of Congo.

Tree census and seed rain data

We established 30 1-ha tree plots comprised of three equal-area groups, including 10 sites that were unlogged and unhunted, 10 sites that were logged and unhunted and 10 sites that were both logged and hunted. Using ArcView 3.2 and a 14-class habitat map (Laporte et al. 2007), we randomly located plots within each disturbance regime in mixed lowland forest, with a buffer of at least 500 m to the nearest primary road and 100 m to the nearest water source. Within each plot, all trees >10 cm diameter at breast height (DBH) were tagged, measured, mapped and identified to species (Wortley and Harris 2014). We additionally recorded canopy status (understory, midstory, canopy and emergent) and presence of lianas in the crown. Canopy openness and light availability were estimated for each plot by averaging values from four hemispherical pictures taken at each quarter of a plot. Seed traps 1 m² in area were centred along three transects at 25, 50 and 75 m from a plot border, with 10 m separating each trap. All traps were at least 20 m from the nearest plot border. Seeds and fruits were collected every 2 weeks and identified to species or genus level. Previous evidence demonstrates that parameter estimates are dominated by the relatively abundant seeds falling from within these distances (Clark et al. 1998).

We used seed rain data from 33 of the most common species to quantify fecundity and seed dispersal dynamics. Although seed rain was collected on many more species, we limited analysis to species that occurred in

at least half of all plots. Tree density, size and species composition were approximately equivalent across plots and disturbance types [see [Supporting Information—Figs S1–S3](#)]. Of the 44 species that contributed seeds to at least half of the plots, 11 were lianas—woody vines that rely on trees for support. We omitted liana species from the present study despite their clear importance for frugivore diets, because they extend laterally tens of metres from their rooting stems, making the attribution of seeds to a censused stem challenging. The number of focal trees per 1-ha plot ranged from 50 to 253 with a median of 155 trees, and the number of seeds per focal species per plot ranged from 16 to 288 with a median of 96.

Plant species trait data

The dispersal mode for each tree species was assigned based on fruit morphology and observations of fruit consumption (Gautier-Hion et al. 1985; Tutin et al. 1997; White and Abernethy 1997; Whitney et al. 1998; Clark et al. 2001; Poulsen et al. 2001, 2002; Hawthorne and Gyakari 2006; Morgan and Sanz 2006) [see [Supporting Information—Table S1](#)]. Because many animal-dispersed species are dispersed by both birds and mammals, we report results by broad classes of animal and abiotic (wind or ballistic) dispersal mode. In addition to dispersal mode, the mean tree DBH (cm) and tree density (stems per hectare) for each species were also calculated by forest type to relate dispersal parameters to species characteristics.

Fecundity estimation and dispersal analysis

We use a state-space model for Mast Inference and Forecasting (available on CRAN as the R package MASTIF, <http://rpubs.com/jimclark/281413>) to determine the relative influence of hunting and logging on the fecundity and dispersal kernel of each tree (Clark, Nuñez and Tomasek, in revision). Mast Inference and Forecasting builds on the rich literature of seed dispersal models that employ a bivariate Student's *t* (2Dt) to relate the size and locations of reproductively active trees to numbers of seeds collected in seed traps in order to probabilistically estimate the seed production of each tree (Fig. 2; Clark et al. 1999, 2010, 2014a). Some authors use a two-parameter version of the 2Dt kernel; we do not fit a shape parameter due to the fact that it is poorly identified in data and it does not respond to the tail of the kernel as was originally hoped (e.g. Clark et al. 1999).

Not all seeds in seed traps must come from trees within the inventory plot. This possibility suggests an intercept proportional to basal area (Clark et al. 2010) or an integral over a large landscape area (Muller-Landau et al. 2008) as a rough accommodation of long-distance dispersal. In our comparisons an intercept can change estimates, without actually being sensitive to seeds outside the plot. This insensitivity to distant trees was demonstrated by Clark et al. (1998) by fitting the model without intercept to increasingly expanded plot areas. An intercept is insensitive to long-distance dispersal because distant trees do not affect the likelihood; the tail of the kernel has no impact on estimates except in cases where seeds are rare (Clark et al. 1999). The converse is also true: standard

errors on estimates of fecundity increase with distance from seed traps. The intercept model further requires a strict assumption about forest composition outside the plot, e.g. extrapolating composition within the plot to infinite distance (Muller-Landau et al. 2008; Clark et al. 2010), which is unrealistic in many forests.

Mast Inference and Forecasting extends the model that has been extensively tested with predictive distributions to allow for uncertainty in seed identification, as well as time-dependence (Clark et al. 2004, 2010) and quasi-periodic variation and synchronicity in seed production (Koenig and Knops 2001; Boutin et al. 2006; Wang et al. 2017). Mast Inference and Forecasting uses Gibbs sampling—a Markov chain Monte Carlo (MCMC) technique—as well as Metropolis and Hamiltonian Markov chain (HMC) for posterior simulations of tree maturation state, fecundity, seed dispersal kernel and parameter estimates. Parameter estimates—the effects of hunting, logging and site-level covariates—are sampled directly from the posterior (Clark, Nuñez and Tomasek, in revision). We used non-informative flat priors for the dispersal parameter and variance in the dispersal parameter with fixed degrees of freedom as detailed in Clark et al. (2004, 2010, 2014a).

The broad dispersion of seed count data is accommodated in at least one of two ways. If accommodated at the data stage with a negative binomial distribution (Clark et al. 1998; Muller-Landau et al. 2008), then the dispersion parameter has no biological interpretation, and it cannot respond to the variables that are known to affect seed variability. Alternatively, a hierarchical specification




Figure 2. A schematic of seed shadow modelling, with spatially distributed trees of varying sizes acting as signal sources of varying strengths, and seed traps acting as stationary detectors through time.

helps to explain that variation, through individual differences in covariates and random effects and year or lag effects (Clark et al. 2004, 2013; Martínez and González-Taboada 2009; Uriarte et al. 2012). In other words, the overdispersion is taken up by the underlying process; the data are conditionally Poisson, but marginally overdispersed (Clark, Nuñez and Tomasek, in revision). Our model incorporates a Poisson likelihood for count data with seed production and dispersal, written as:

$$E(y_s) = A\lambda_s = A \sum_{i=1}^n S_{si} f_i$$

where $E(y_s)$ is the expected number of seeds counted in a trap at location s . λ_s is the expected seed density (seeds per m^2 per year) multiplied by the sampling effort A —the area of a seed trap times the fraction of the fruiting season it was deployed (m^2 per year). S_{si} is the density of seed (m^{-2}) produced by tree i dispersed to seed trap location s ; and f_i is fecundity for an individual tree i at time t , which is the product of maturation status (ρ_{it}) and conditional fecundity (ψ_{it}) of tree i , ($f_{it} = \psi_{it}\rho_{it} \geq 0$). Maturation and conditional fecundity are dynamic processes, modelled with fixed, random and year effects. Coefficients in the vector of fixed effects β^x include tree diameter, exposure to hunting or logging, and interactions (Clark 2010; Clark et al. 2013). Random individual effects accommodate the heterogeneity of responses among individual trees. The effect of year is random across species and within each of the three disturbance types, accommodating seed rain fluctuations that are coherent within, but not among the three groups.

Dispersal is summarized by the mean parameter of the 2Dt dispersal kernel (Clark et al. 1999), here termed the ‘dispersal parameter’. A shape parameter is also sometimes fitted for this model, but we have found it to be unstable and unresponsive to long-distance dispersal (Clark et al. 2004, 2010).

Our modelling did not explicitly incorporate boundary effects because previous analysis demonstrated that trees tens of metres from seed traps have little impact on estimates (Clark et al. 1999). Muller-Landau et al. (2008), however, concluded that failure to account for boundary effects could bias models towards higher fecundity and fat tails (Muller-Landau et al. 2008), leading to overestimated fecundities and dispersal distances. However, this would not change inferences related to the relative effects of vectors or disturbance on seed dispersal patterns.

Gibbs sampling was used for posterior simulation. For each tree species [see Supporting Information—Fig. S5], model estimates were taken from 50 000 iterations, discarding the first 1000 iteration as pre-convergence. We visually inspected trace plots to confirm convergence and adequate mixing [see Supporting

Information—Fig. S6A–C]. Model fit was assessed with root mean squared prediction error (RMSPE) across species [see Supporting Information—Fig. S4]. Variable selection was based on Deviance Information Criterion (DIC). Model estimates reported in the text are posterior means and 95 % credible intervals (CIs) based on the Gibbs sampler realizations.

Results

Hunting and logging influenced the mean distances of dispersal kernels (hereafter average dispersal distance), with the greatest effects on animal-dispersed species, though the direction and magnitude varied. Two-thirds of all species (22/33) in disturbed forests had 95 % CIs for dispersal parameters that did not overlap with estimates from protected plots, indicating a role of disturbance. This trend held true whether a species relied on animals for dispersal entirely (13/18), in part (5/8) or not at all (4/7).

Of the 22 species affected by disturbance, 17 species showed an effect of logging alone: nine species had higher dispersal estimates in logged compared to protected forest (*Celtis mildbraedii*, *Diospyros canaliculata*, *Erythrophleum suaveolens*, *Greenwayodendron suaveolens*, *Lannea welwitschii*, *Pausinystalia macroceras*, *Rinorea oblongifolia*, *Staudtia kamerunensis*, *Strombosia nigropunctata*), and eight species had lower dispersal estimates (*Cleistopholis patens*, *Grossera macrantha*, *Myrianthus arboreus*, *Macaranga barteri*, *Nesogordonia kabingaensis*, *Strombosiaopsis tetrandra*, *Thomandersia hensii*, *Terminalia superba*).

The combined effects of hunting and logging were consistent with logging alone for the majority of species, with the exception of six species that had dispersal estimates greater than (*Pteleopsis hyloidendron*, *S. tetrandra*, *Guarea cedrata*) or less than (*G. macrantha*, *D. canaliculata* and *E. suaveolens*) logging alone. Notably, three species exhibited divergent effects of disturbance regime on dispersal estimates: logging positively affected *D. canaliculata* and *E. suaveolens*, whereas the combination of hunting and logging negatively affected dispersal estimates relative to protected plots. *Strombosiaopsis tetrandra* displayed the opposite pattern (Table 1; Figs 3 and 4).

To reveal potential group-level effects of dispersal vectors, we clustered dispersal parameters from individual species by dispersal vector (i.e. animal, abiotic or mixed dispersal). Predictions were congruent within each dispersal vector, regardless of disturbance type (Fig. 5). Abiotically dispersed species had the greatest dispersal estimates overall, with 51.4 m [2.5th and 97.5th quantiles: 17.9, 75.5]. Species dispersed both by animals and abiotically had dispersal estimates of 41.1

Table 1. Predictive mean and 95 % CI for seed dispersal distances in metres.

	Mean predicted dispersal distance								
	Logged forests			Protected forests			Hunted and logged forests		
	Estimate	2.50 %	97.50 %	Estimate	2.50 %	97.50 %	Estimate	2.50 %	97.50 %
Abiotically dispersed									
<i>Albizia gummifera</i>	51.3	47.7	54.9	52.6	49.4	55.8	53.3	50.3	56.3
<i>Erythrophleum suaveolens</i>	42.6	40.2	45.0	31.8	30.0	33.7	23.4	22.1	24.7
<i>Nesogordonia kabingaensis</i>	12.4	9.5	16.5	41.1	38.1	44.1	37.0	33.5	40.6
<i>Petersianthus macrocarpus</i>	65.9	64.1	67.7	63.3	61.2	65.6	61.1	58.6	63.4
<i>Pteleopsis hyloidendron</i>	43.9	39.0	48.6	36.5	28.4	43.6	57.5	54.0	61.2
<i>Pterocarpus soyauxii</i>	56.9	53.3	60.4	62.9	60.3	65.5	66.9	64.6	69.2
<i>Terminalia superba</i>	68.0	66.6	69.6	75.0	73.4	76.8	75.9	74.3	77.7
Animal dispersed									
<i>Angylocalyx pynaertii</i>	45.0	41.5	48.5	41.0	37.2	45.0	49.9	46.9	52.7
<i>Celtis adolfi-friderici</i>	18.0	16.0	20.3	14.5	13.2	15.9	13.8	12.9	14.8
<i>Celtis mildbraedii</i>	20.1	18.7	21.6	10.3	9.9	10.8	21.0	19.8	22.3
<i>Cleistopholis patens</i>	17.8	13.8	21.7	38.4	30.1	43.6	38.4	34.4	42.3
<i>Diospyros bipindensis</i>	41.9	38.7	45.2	39.8	36.7	43.1	39.5	35.2	43.8
<i>Diospyros canaliculata</i>	45.9	42.8	49.0	36.8	33.2	40.2	13.9	12.8	15.0
<i>Greenwayodendron suaveolens</i>	37.2	35.7	38.7	31.4	30.2	32.7	42.4	40.5	44.2
<i>Guarea cedrata</i>	35.3	31.0	39.7	28.0	19.5	35.1	39.2	35.9	42.6
<i>Guarea thompsonii</i>	40.7	37.5	44.0	40.2	36.4	43.9	46.9	43.7	50.1
<i>Lannea welwitschii</i>	42.5	37.3	47.6	2.2	1.0	8.7	16.9	14.0	20.6
<i>Macaranga barteri</i>	10.2	8.4	12.5	24.4	20.6	28.3	4.5	3.6	6.0
<i>Staudtia kamerunensis</i>	49.9	45.2	55.2	34.2	22.9	42.0	49.0	45.7	52.5
<i>Strombosia nigropunctata</i>	21.1	19.5	22.8	9.8	9.1	10.5	19.6	18.3	21.0
<i>Strombosia pustulata</i>	17.5	15.9	19.2	15.6	14.5	16.9	14.2	12.7	15.9
<i>Strombosiopsis tetrandra</i>	19.2	18.1	20.3	28.7	26.7	30.7	41.4	38.9	43.9
<i>Xylopia chrysophylla</i>	40.4	36.9	44.0	34.4	27.9	40.0	42.9	40.3	45.6
<i>Xylopia hypolampra</i>	98.5	95.3	100.0	98.8	96.1	100.0	98.0	93.6	99.9
<i>Xylopia phloiodora</i>	47.8	44.3	51.2	45.7	42.2	49.2	44.7	41.3	48.1
Abiotic and animal dispersed									
<i>Camptostylus mannii</i>	42.3	38.9	45.6	41.5	38.0	45.1	39.7	36.5	43.0
<i>Grossera macrantha</i>	40.5	35.6	45.0	51.9	49.2	54.5	43.9	40.1	46.9
<i>Lepidobotrys staudtii</i>	35.9	27.9	42.1	45.7	41.9	49.4	51.1	47.9	54.3
<i>Myrianthus arboreus</i>	34.2	28.8	39.0	43.3	39.8	46.6	25.3	22.7	28.1
<i>Pausinystalia macroceras</i>	37.1	34.0	40.3	31.2	28.8	33.7	39.0	35.6	42.4
<i>Radlkofera calodendron</i>	41.3	37.4	45.1	45.7	42.1	49.0	42.4	38.9	46.0
<i>Rinorea oblongifolia</i>	46.7	43.6	49.8	31.7	27.0	36.3	46.7	43.3	50.0
<i>Thomandersia hensii</i>	38.0	31.9	43.2	54.0	50.9	57.2	37.9	31.5	43.1




Figure 3. Comparison of average dispersal distance parameters among species in plots that were hunted and logged, logged, or protected from hunting and logging. Species are ordered by mean dispersal distance parameter in protected plots. Densities on right Y-axis show distribution of the dispersal type for species on left Y-axis.




Figure 4. Comparison of difference in average dispersal distance parameter from protected forests among species in plots that were hunted and logged, or logged. Species are ordered by mean dispersal distance in hunted and logged plots. Densities on right Y-axis show distribution of the dispersal type for species on left Y-axis.




Figure 5. Post hoc comparison of the mean predicted dispersal distances for all species grouped by dispersal vector. Error bars show the 2.5th and 97.5th quantiles of mean dispersal distance in forests that were logged, hunted and logged, or protected from hunting and logging.

m [28.7, 52.8], and animal-dispersed species had the lowest dispersal estimates of 34.4 m [6.2, 98.3].

To evaluate the group-level effects of disturbance, we clustered dispersal estimates of all species by disturbance type, including protected (38.9 m [8.2, 79.8]), hunted and logged (40.5 m [12.0, 80.3]) and logged forests (39.6 m [11.9, 74.1]). The large overlap in dispersal estimates among forest types indicates a lack of consistent effects of disturbance on dispersal distance.

Estimated tree fecundity increased with tree diameter (Fig. 6), but was not affected by disturbance regime (Table 2; Figs 7 and 8). A majority of species (25/33) exhibited a positive effect of tree diameter on fecundity, with the exception of *Radlkofera calodendron*, *Lepidobotrys staudtii*, *S. kamerunensis*, *R. oblongifolia*, *Xylopia chrysophylla*, *Diospyros bipindensis*, *Camptostylus mannii* and *D. canaliculata*. Logging only influenced fecundity estimates of three species (*D. bipindensis*, posterior mean and 95 % CIs: -1.78 [-3.51 , -0.03], *G. macrantha* -1.49 [-2.96 , -0.08] and *M. arboreus* 2.30 [0.67 , 3.84]).

Discussion

We find that low-intensity logging affected seed dispersal two decades after the logging event. Guidelines aimed at reducing the ecological damage stemming from logging can substantially reduce short-term impacts (Sist 2000; Sist et al. 2003), but our study suggests that impacts of low-intensity logging on ecological processes like seed

dispersal are long term and may linger for decades. The difficult-to-detect effects on a key ecological process could have direct consequences for forest species composition through density-dependent recruitment (Janzen 1970; Connell 1971; Cannon et al. 1994; Bleher and Böhning-Gaese 2001) and competition at later life stages (Nathan and Muller-Landau 2000), potentially altering the diversity and function of forest ecosystems.

Contrary to our expectations, the dispersal vector of a seed type, abiotic or animal, was not a reliable indicator of the magnitude or direction of the responses of tree species to disturbance. Our results do not support the argument that dispersal decreases for animal-dispersed species following perturbation of the disperser community (Terborgh et al. 2008; Markl et al. 2012), at least several decades after the fact. It further does not support the notion that dispersal increases for abiotically dispersed species following forest thinning due to increased canopy wind speeds (Gardiner 1994; Stacey et al. 1994; Gardiner et al. 1997). Our results are more consistent with dispersal effects that are species-specific, as might be expected from the fact that each species has a unique relationship to unmeasured abiotic variables that contribute to its response to disturbance.

Despite a design specifically implemented to detect it, our study did not find evidence for an interaction between hunting and logging for most species, suggesting instead that dispersal following disturbance primarily responds to logging, but not hunting. Using the same data set, Poulsen et al. (2013) modelled seed dispersal of nine mammal-dispersed species finding that mean dispersal distance was farther in logged than unlogged forest for five species and farther in unhunted than hunted forest for six species. The disparity between the two studies could be due to the fact that we modelled dispersal for 33 tree species, separating them into animal and abiotic vectors, whereas Poulsen et al. (2013) only modelled nine mammal-dispersed species for which they had adequate seed numbers.

Limited evidence for a hunting effect on dispersal could come from the fact that hunting pressures were too low, even where present in our data set. Although hunting has clearly reduced the abundance of large vertebrates in the area (Poulsen et al. 2011), all species still exist throughout the landscape (Clark et al. 2009)—the vertebrate community is degraded, not defaunated. Alternatively, large frugivorous birds may have replaced the seed dispersal services of large, arboreal mammals. Bird species richness can increase with logging intensity (Burivalova et al. 2014), which can aggravate the negative effects of disturbance on seed dispersal due to the reduction in seed dispersers (Moran et al. 2004; Kirika et al. 2008a, b; Neuschulz et al. 2011) or mitigate the




Figure 6. Comparison of posterior parameter estimates and 95 % CI show a positive effect of tree diameter on tree fecundity for a majority of species. Species names are colour coordinated here as elsewhere in the manuscript to denote dispersal vector: animal dispersed (orange), abiotically dispersed (black) or both animal and abiotically dispersed (grey).

effects of disturbance if generalist bird dispersers replace lost or reduced dispersal services (Putz et al. 2001; Gray et al. 2007; Burivalova et al. 2014; LaManna and Martin 2017; Trolliet et al. 2017). Indeed, in our study area, there was a 77 % increase in the density of large frugivorous birds following logging (Poulsen et al. 2011), a result that is consistent with other sites in the region (Koerner et al. 2017). Birds are not commonly hunted in our study site, and 2/3 of the mammal-dispersed species were also dispersed by birds [see Supporting Information—Fig. S2], meaning that the full effects of hunting could be attenuated by an expanded bird community.

It is also possible that seed trap data inadequately sample long-distance seed dispersal by animals. A majority of seeds fall locally (Clark et al. 1999, 2005; Muller-Landau and Hardesty 2005; Muller-Landau et al. 2008), and studies that have combined seed traps with direct observations of seed counts from the canopy (LaDeau and Clark

2001, 2006) or the ground (Minor and Kobe 2017) find seed traps estimate fecundity well. However, seed dispersers may forage over large areas—over 4000 ha in some hornbills (Holbrook and Smith 2000). Seed trap data do not fully capture the dispersal of seeds that are consumed and dispersed outside of the plot. Although long-distance dispersal events may be rare, fully estimating the effects of disturbance on seed dispersal may require combined methods that can account for both local and long-distance dispersal. Nevertheless, our findings indicate that once a forest is disturbed by logging, seed dispersal may be altered regardless of the effect hunting has on seed disperser communities. This is consistent with other studies that found animal guild densities were negatively affected by logging even in the absence of hunting (Poulsen et al. 2013), but contradicts studies that found hunting and logging amplified the negative effects of either in isolation (Poulsen et al. 2011; Markl et al. 2012).

Table 2. Posterior mean and 95 % CIs of covariate effects on conditional fecundity.

	Covariate effects on conditional fecundity								
	Diameter			Logging			Hunting and logging		
	Posterior mean	2.50 %	97.50 %	Posterior mean	2.50 %	97.50 %	Posterior mean	2.50 %	97.50 %
Abiotically dispersed									
<i>Albizia gummifera</i>	2.4	0.9	3.8	0.2	−3.1	3.7	1.3	−1.1	3.7
<i>Erythrophleum suaveolens</i>	2.3	1.2	3.4	1.4	−0.6	3.4	0.3	−2.4	3.0
<i>Nesogordonia kabingaensis</i>	2.3	1.4	3.2	−0.6	−1.7	0.5	−0.5	−1.5	0.4
<i>Petersianthus macrocarpus</i>	3.3	2.3	4.3	−1.5	−3.1	0.1	−1.0	−2.3	0.3
<i>Pteleopsis hylo dendron</i>	1.5	0.0	3.0	−1.3	−4.0	1.3	−2.0	−5.1	1.2
<i>Pterocarpus soyauxii</i>	2.8	1.5	4.1	−1.3	−3.0	0.5	−1.1	−2.8	0.6
<i>Terminalia superba</i>	3.9	2.8	4.9	0.0	−1.9	1.8	−1.3	−3.0	0.4
Animal dispersed									
<i>Angylocalyx pynaertii</i>	2.4	1.3	3.5	−0.2	−1.8	1.4	−0.2	−1.8	1.4
<i>Celtis adolfi-friderici</i>	2.9	2.0	3.8	0.0	−1.2	1.2	−0.3	−1.4	0.7
<i>Celtis mildbraedii</i>	2.4	2.0	2.9	−0.2	−1.0	0.5	−0.8	−1.4	−0.1
<i>Cleistopholis patens</i>	2.4	0.9	4.0	−1.1	−3.3	1.2	−2.0	−4.1	0.1
<i>Diospyros bipindensis</i>	0.8	−1.4	2.9	−1.8	−3.5	0.0	−1.8	−3.7	0.1
<i>Diospyros canaliculata</i>	0.3	−1.6	2.3	0.1	−1.5	1.8	−1.0	−2.4	0.4
<i>Greenwayodendron suaveolens</i>	4.2	3.4	5.0	−0.1	−0.9	0.7	−0.3	−1.0	0.4
<i>Guarea cedrata</i>	1.8	0.2	3.3	−0.7	−3.0	1.5	−0.1	−3.3	3.3
<i>Guarea thompsonii</i>	2.3	1.3	3.4	−1.1	−2.2	0.1	−1.0	−2.1	0.1
<i>Lannea welwitschii</i>	2.1	0.3	3.8	−0.2	−4.5	4.0	0.6	−4.4	5.2
<i>Macaranga barteri</i>	2.4	1.1	3.7	−1.0	−2.6	0.6	−0.8	−2.4	0.8
<i>Staudtia kamerunensis</i>	1.6	−0.2	3.4	−0.3	−4.1	3.6	0.0	−6.2	6.2
<i>Strombosia nigropunctata</i>	1.6	0.6	2.6	−0.5	−1.5	0.5	−0.6	−1.5	0.4
<i>Strombosia pustulata</i>	2.2	1.4	3.0	−0.6	−1.5	0.2	−0.3	−1.2	0.6
<i>Strombosiaopsis tetrandra</i>	3.2	2.2	4.1	−0.9	−2.1	0.3	−0.6	−1.7	0.6
<i>Xylopia chrysophylla</i>	1.3	−0.4	3.0	0.1	−3.0	3.2	−2.5	−5.1	0.2
<i>Xylopia hypolampra</i>	1.8	0.2	3.4	1.6	−2.0	5.1	0.9	−2.4	4.2
<i>Xylopia phloiodora</i>	1.7	0.0	3.3	0.8	−1.4	2.9	1.6	−0.9	4.0
Abiotic and animal dispersed									
<i>Camptostylus mannii</i>	0.9	−1.1	2.8	0.7	−1.3	2.7	0.3	−1.6	2.1
<i>Grossera macrantha</i>	2.3	0.9	3.7	−1.5	−3.0	−0.1	−0.7	−2.2	0.8
<i>Lepidobotrys staudtii</i>	1.7	−0.1	3.6	−0.2	−3.0	2.6	0.1	−2.2	2.4
<i>Myrianthus arboreus</i>	1.7	0.6	2.7	2.1	−0.3	4.5	2.3	0.7	3.8
<i>Pausinystalia macroceras</i>	1.1	0.1	2.1	−0.1	−1.5	1.3	0.3	−1.0	1.6
<i>Radlkofera calodendron</i>	2.1	0.0	4.2	−1.9	−4.3	0.7	1.5	−2.4	5.4
<i>Rinorea oblongifolia</i>	1.6	−0.5	3.8	1.7	−1.7	5.0	−1.9	−5.2	1.1
<i>Thomandersia hensii</i>	2.9	0.0	5.9	1.1	−4.8	6.9	0.0	−6.2	6.2




Figure 7. Comparison of posterior parameter estimates and 95 % CI show no effect of logging on tree fecundity for a majority of species.

Although dispersal vector was not predictive of how dispersal would respond to hunting or logging, there was a clear distinction in dispersal kernel estimates. Abiotically dispersed seeds moved farthest from the parent tree, animal-dispersed seeds generally fell closest and species dispersed both by animals and abiotically arrived at intermediate distances. Differences in dispersal distance between vectors (Venable and Brown 1988; Greene and Johnson 1989, 1993; Cornelissen et al. 2003; Clark et al. 2005; Thomson et al. 2011) are partly a result of mechanical properties. Abiotically dispersed seeds tend to have small mass that facilitate passive dispersal by wings, plumes, samaras and other adaptations for flight (Greene and Johnson 1989, 1993). Seeds reliant on animal dispersers must develop fleshy fruit mass to entice seed dispersers (Cao et al. 2016) limiting their passive dispersal distance.

Estimated fecundity long after disturbance did not differ across disturbance regimes to the extent found in studies immediately following disturbance (Markl et al.

2012; Uriarte et al. 2012; Berdanier and Clark 2016). Low-intensity logging in resource-limited tropical forest environments may have limited effects on crowding, light and soil moisture levels (Molino and Sabatier 2001; Bongers et al. 2009). However, our results suggest that any fecundity benefits from disturbance are unobservable 20 years post-logging. Lack of a long-term effect on fecundity may also be a result of studying only relatively large trees (≥ 10 cm DBH), which have already made it through the competitive gauntlet of the understory to attain adulthood, and can access resources that facilitate resilience to competitive environments in ways that smaller plants cannot (Clark et al. 2004).

Tree size was an important determinant of fecundity making large trees especially important for forest regeneration (Plumptre 1995; Freitas and Pinard 2008). Fecundity of large trees should encourage their protection during logging campaigns (CIB 2006). In addition to their outsized contribution to longer-distance dispersal events (Norghauer et al. 2011), large trees store a




Figure 8. Comparison of posterior parameter estimates and 95 % CI show no effect of hunting and logging on tree fecundity for a majority of species.

disproportionate amount of above-ground carbon (Clark and Clark 1996; Lutz et al. 2012; Slik et al. 2013; Stephenson et al. 2014) and are crucial for maintenance of forest structure (Lindenmayer et al. 2012; Lutz et al. 2013) and animal habitat (Tews et al. 2004; Lutz et al. 2012, 2013).

Our study demonstrates that disturbances to forests and animal communities contribute to seed dispersal patterns even decades after the initial logging event. In this case, the responses in seed dispersal to disturbance varied across species with weak patterns related to dispersal vector or disturbance type. Our lack of a clear directional effect of hunting and logging on seed dispersal could be partially due to our study design, which was pseudoreplicated: study plots affected by the same disturbance type were geographically grouped together out of necessity. This was a direct result of the study area, particularly the spatial pattern of hunting and logging around the village of Kabo (Poulsen et al. 2011), and means that other, unmeasured environmental gradients could influence our results.

The limitations of our study should serve as a challenge to dispersal ecologists and modelers—what are the best methods or combinations of methods for disentangling the effects of multiple disturbances that can operate over disparate spatial and timescales?

Logging concessions cover much of West and Central Africa (FAO 2016), yet the long-term impacts of low-intensity logging techniques on fundamental ecological processes like seed dispersal have been largely overlooked. This work advances our understanding of how the separate and combined effects of hunting and logging affect seed dispersal in the understudied Afrotropics. Although care needs to be taken before extrapolating our results to other contexts, the species-specific dispersal responses to logging in this study point towards the long-lasting toll of disturbance on ecological function. Whereas the effects of disturbance on forest structure and animal communities are easily measured, the effects on ecological processes may be more cryptic, long-lasting and difficult to decipher.

Data

<https://github.com/chasenunez/2018.AOBP>.

Sources of Funding

The U.S. Fish and Wildlife Service Great Ape Fund generously provided financial support for this research. C.L.N. was supported by National Science Foundation (NSF) (GRF-1106401) and a Neil Williams Presidential fellowship; J.R.P. and C.J.C. were supported by a University of Florida Presidential fellowship (J.R.P.), a School of Natural Resources and Environment alumni fellowship (C.J.C.) and Environmental Protection Agency Science to Achieve Results (STAR) fellowships (91630801-0 to J.R.P. and 91643301-0 to C.J.C.); J.S.C. was supported by NSF-EF-1137364, NSF-EF-1550911 and NASA's AIST programme.

Contributions by the Authors

C.L.N. posed the central questions, wrote the original manuscript, and analyzed the data; J.S.C. wrote the R and C++ code for the MASTIF model with testing and feedback by C.L.N. through development; J.R.P. and C.J.C. collected data with help from those in Acknowledgements section; J.R.P. and J.S.C. edited for content and provided guidance on structure and style.

Conflict of Interest

None declared.

Acknowledgements

We thank Rebecca Maas. We also thank O. Mbani, Y. Nganga, M. Moko and I. Loungouba for their tireless fieldwork, and the Government of Congo and its Ministries of Forestry Economy and Scientific Research for their support. The Wildlife Conservation Society provided logistical support, and we owe thanks to its staff, particularly P. Elkan, S. Elkan, B. Curran, M. Gately, E. Stokes, C. Prevost, J. Beck and J. Mokoko. We also acknowledge the Buffer Zone Project (PROGEPP) and the Congolaise Industrielle des Bois for their collaboration.

Supporting Information

The following additional information is available in the online version of this article—

Figure S1. Boxplots comparing the distribution of tree diameters within each plot type show no systematic difference across plot types.

Figure S2. Boxplots comparing the distributions of total stems per plot show significant overlap across plot type.

Figure S3. Stacked bar plots comparing community composition show a consistent distribution of 33 focal species across plots.

Figure S4. Comparison of standardized root mean squared prediction error (individual RMSPE/average number of seeds per trap) with size of circle indicating relative number of seeds from that species present in the study.

Figure S5. (A–D) Example of individual results (*Nesogordonia kabingaensis*) that were amalgamated across species for in-text summary figures.

Figure S6. (A–C) Examples of model diagnostics for *Nesogordonia kabingaensis*.

Table S1. Table of species information and dispersal vectors.

Literature Cited

- Abernethy KA, Coad L, Taylor G, Lee ME, Maisels F. 2013. Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **368**:20130494.
- Beaune D, Bretagnolle F, Bollache L, Hohmann G, Surbeck M, Fruth B. 2013. Seed dispersal strategies and the threat of defaunation in a Congo forest. *Biodiversity and Conservation* **22**:225–238.
- Beck H, Snodgrass JW, Thebpanya P. 2013. Long-term enclosure of large terrestrial vertebrates: implications of defaunation for seedling demographics in the Amazon rainforest. *Biological Conservation* **163**:115–121.
- Berdanier AB, Clark JS. 2015. Multi-year drought-induced morbidity preceding tree death in Southeastern US forests. *Ecological Applications* **26**:150731093536001.
- Berdanier AB, Clark JS. 2016. Divergent reproductive allocation trade-offs with canopy exposure across tree species in temperate forests. *Ecosphere* **7**:e01313.
- Bleher B, Bohning-Gaese K. 2001. Consequences of frugivore diversity for seed dispersal, seedling establishment and the spatial pattern of seedlings and trees. *Oecologia* **129**:385–394.
- Bongers F, Poorter L, Hawthorne WD, Sheil D. 2009. The intermediate disturbance hypothesis applies to tropical forests, but disturbance contributes little to tree diversity. *Ecology Letters* **12**:798–805.
- Boutin S, Wauters LA, McAdam AG, Humphries MM, Tosi G, Dhondt AA. 2006. Anticipatory reproduction and population growth in seed predators. *Science (New York, N.Y.)* **314**:1928–1930.
- Brown KA, Gurevitch J. 2004. Long-term impacts of logging on forest diversity in Madagascar. *Proceedings of the National Academy of Sciences of the United States of America* **101**:6045–6049.
- Burivalova Z, Sekercioglu CH, Koh LP. 2014. Thresholds of logging intensity to maintain tropical forest biodiversity. *Current Biology* **24**:1893–1898.
- Camargo-Sanabria AA, Mendoza E, Guevara R, Martinez-Ramos M, Dirzo R. 2014. Experimental defaunation of terrestrial mammalian herbivores alters tropical rainforest understorey diversity. *Proceedings of the Royal Society B: Biological Sciences* **282**:20142580.

- Cannon C, Kartawinata K, Leighton M, Peart David R. 1994. The structure of lowland rainforest after selective logging in West Kalimantan, Indonesia. *Forest Ecology and Management* **67**:49–68.
- Cao L, Wang Z, Yan C, Chen J, Guo C, Zhang Z. 2016. Differential foraging preferences on seed size by rodents result in higher dispersal success of medium-sized seeds. *Ecology* **97**:3070–3078.
- Congolaise Industrielle des Bois (CIB). 2006. *Plan d'aménagement de l'unité forestière d'aménagement de Kobo (2005–2034)*. Brazzaville, Republic of the Congo: Ministry of Forest Economy.
- Clark JS. 2010. Individuals and the variation needed for high species diversity in forest trees. *Science (New York, N.Y.)* **327**:1129–1132.
- Clark JS, Bell D, Chu C, Courbaud B, Dietze M, Hersh M, HilleRisLambers J, Ibáñez I, LaDeau S, McMahon S, Metcalf J, Mohan J, Moran E, Pangle L, Pearson S, Salk C, Shen Z, Valle D, Wyckoff P. 2010. High-dimensional coexistence based on individual variation: a synthesis of evidence. *Ecological Monographs* **80**:569–608.
- Clark JS, Bell DM, Kwit M, Powell A, Zhu K. 2013. Dynamic inverse prediction and sensitivity analysis with high-dimensional responses: application to climate-change vulnerability of biodiversity. *Journal of Agricultural, Biological, and Environmental Statistics* **18**:376–404.
- Clark JS, Bell DM, Kwit MC, Zhu K. 2014a. Competition-interaction landscapes for the joint response of forests to climate change. *Global Change Biology* **20**:1979–1991.
- Clark DB, Clark DA. 1996. Abundance, growth and mortality of very large trees in neotropical lowland rain forest. *Forest Ecology and Management* **80**:235–244.
- Clark JS, Gelfand AE, Woodall CW, Zhu K. 2014b. More than the sum of the parts: forest climate response from joint species distribution models. *Ecological Applications* **24**:990–999.
- Clark JS, LaDeau S, Ibanez I. 2004. Fecundity of trees and the colonization-competition hypothesis. *Ecological Monographs* **74**:415–442.
- Clark JS, Macklin E, Wood L. 1998. Stages and spatial scales of recruitment limitation in southern Appalachian forests. *Ecological Monographs* **68**:213–235.
- Clark CJ, Poulsen JR, Bolker BM, Connor EF, Parker VT. 2005. Comparative seed shadows of bird-, monkey-, and wind-dispersed trees. *Ecology* **86**:2684–2694.
- Clark CJ, Poulsen JR, Malonga R, Elkan PW. 2009. Logging concessions can extend the conservation estate for central African tropical forests. *Conservation Biology* **23**:1281–1293.
- Clark CJ, Poulsen JR, Parker VT. 2001. The role of arboreal seed dispersal groups on the seed rain of a lowland tropical forest. *Biotropica* **33**:606–620.
- Clark JS, Silman M, Kern R, Macklin E, Hillerislammers J. 1999. Seed dispersal near and far: patterns across temperate and tropical forests. *Ecology* **80**:1475–1494.
- Comita LS, Queenborough SA, Murphy SJ, Eck JL, Xu K, Krishnadas M, Beckman N, Zhu Y, Gomez-Aparicio L. 2014. Testing predictions of the Janzen-Connell hypothesis: a metaanalysis of experimental evidence for distance- and density dependent seed and seedling survival. *The Journal of Ecology* **102**:845–856.
- Connell JH. 1971. On the role of natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees. In: Den Boer PJ, Gradwell GR, eds. *Dynamics of populations*. Wageningen, The Netherlands: Centre for Agricultural Publishing and Documentation, 298–312.
- Cornelissen JHC, Lavorel S, Garnier E, Díaz S, Buchmann N, Gurvich DE, Reich PB, ter Steege H, Morgan HD, van der Heijden MGA, Pausas JG, Poorter H. 2003. A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany* **51**:335–380.
- Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJ, Collen B. 2014. Defaunation in the Anthropocene. *Science (New York, N.Y.)* **345**:401–406.
- Edwards DP, Tobias JA, Sheil D, Meijaard E, Laurance WF. 2014. Maintaining ecosystem function and services in logged tropical forests. *Trends in Ecology & Evolution* **29**:511–520.
- Ewel JJ, Mazzarino MJ. 2008. Competition from below for light and nutrients shifts productivity among tropical species. *Proceedings of the National Academy of Sciences of the United States of America* **105**:18836–18841.
- FAO. 2016. *The contemporary forest concessions in West and Central Africa: chronicle of a foretold decline?* Forestry Policy and Institutions Working Paper 34. Rome: FAO.
- Freitas JV de, Pinard MA. 2008. Applying ecological knowledge to decisions about seed tree retention in selective logging in tropical forests. *Forest Ecology and Management* **256**:1434–1442.
- Gardiner BA. 1994. Wind and wind forces in a plantation spruce forest. *Boundary-Layer Meteorology* **67**:161–186.
- Gardiner BA, Stagey GR, Belcher RE, Wood CJ. 1997. Field and wind tunnel assessments of the implications of respacing and thinning for tree stability. *Forestry* **70**:233–252.
- Gautier-Hion A, Duplantier J-M, Quris R, Feer F, Sourd C, Decoux JP, Dubost G, Emmons L, Erard C, Heeketsweiler P, Moungazi A, Roussillon C, Thiollay J-M. 1985. Fruit characters as a basis of fruit choice and seed dispersal in a tropical forest vertebrate community. *Oecologia* **65**:324–337.
- Gourlet-Fleury S, Mortier F, Fayolle A, Baya F, Ouedraogo D, Benedet F, Picard N. 2013. Tropical forest recovery from logging: a 24 year silvicultural experiment from Central Africa. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **368**:20120302.
- Gray MA, Baldauf SL, Mayhew PJ, Hill JK. 2007. The response of avian feeding guilds to tropical forest disturbance. *Conservation Biology* **21**:133–141.
- Greene DF, Johnson EA. 1989. A model of wind dispersal of winged or plumed seeds. *Ecology* **70**:339–347.
- Greene D, Johnson E. 1993. Seed mass and dispersal capacity in wind-dispersed diaspores. *Oikos* **67**:69–74.
- Gutierrez-Granados G. 2011. Effect of logging on rodent scatterhoarding dynamics in tropical forests: implications for plant recruitment. *Integrative Zoology* **6**:74–80.
- Haurez B, Tagg N, Petre C-A, Vermeulen C, Doucet J-L. 2016. Short term impact of selective logging on a western lowland gorilla population. *Forest Ecology and Management* **364**:46–51.
- Hawthorne W, Gyakari N. 2006. *Photoguide for the forest trees of Ghana: a tree-spotter's field guide for identifying the largest trees*. Oxford: Oxford Forestry Institute.
- Hoffmann M, Hilton-Taylor C, Angulo A, Böhm M, Brooks TM, Butchart SH, Carpenter KE, Chanson J, Collen B, Cox NA, Darwall WR, Dulvy NK, Harrison LR, Katariya V, Pollock CM, Quader S, Richman NI, Rodrigues AS, Tognelli MF, Vié JC, Aguiar JM, Allen DJ, Allen GR, Amori G, Ananjeva NB, Andreone F, Andrew P, Aquino Ortiz AL, Baillie JE, Baldi R, Bell BD, Biju SD, Bird JP, Black-Decima P, Blanc JJ, Bolaños F, Bolívar-G W,

- Burfield IJ, Burton JA, Capper DR, Castro F, Catullo G, Cavanagh RD, Channing A, Chao NL, Chenery AM, Chiozza F, Clausnitzer V, Collar NJ, Collett LC, Collette BB, Cortez Fernandez CF, Craig MT, Crosby MJ, Cumberlidge N, Cuttelod A, Derocher AE, Diesmos AC, Donaldson JS, Duckworth JW, Dutson G, Dutta SK, Emslie RH, Farjon A, Fowler S, Freyhof J, Garshelis DL, Gerlach J, Gower DJ, Grant TD, Hammerson GA, Harris RB, Heaney LR, Hedges SB, Hero JM, Hughes B, Hussain SA, Icochea MJ, Inger RF, Ishii N, Iskandar DT, Jenkins RK, Kaneko Y, Kottelat M, Kovacs KM, Kuzmin SL, La Marca E, Lamoreux JF, Lau MW, Lavilla EO, Leus K, Lewison RL, Lichtenstein G, Livingstone SR, Lukoschek V, Mallon DP, McGowan PJ, McIvor A, Moehلمان PD, Molur S, Muñoz Alonso A, Musick JA, Nowell K, Nussbaum RA, Olech W, Orlov NL, Papenfuss TJ, Parra-Olea G, Perrin WF, Polidoro BA, Pourkazemi M, Racey PA, Ragle JS, Ram M, Rathbun G, Reynolds RP, Rhodin AG, Richards SJ, Rodríguez LO, Ron SR, Rondinini C, Rylands AB, Sadovy de Mitcheson Y, Sanciangco JC, Sanders KL, Santos-Barrera G, Schipper J, Self-Sullivan C, Shi Y, Shoemaker A, Short FT, Sillero-Zubiri C, Silvano DL, Smith KG, Smith AT, Snoeks J, Stattersfield AJ, Symes AJ, Taber AB, Talukdar BK, Temple HJ, Timmins R, Tobias JA, Tsytulina K, Tweddle D, Ubeda C, Valenti SV, van Dijk PP, Veiga LM, Veloso A, Wege DC, Wilkinson M, Williamson EA, Xie F, Young BE, Akçakaya HR, Bennun L, Blackburn TM, Boitani L, Dublin HT, da Fonseca GA, Gascon C, Lacher TE Jr, Mace GM, Mainka SA, McNeely JA, Mittermeier RA, Reid GM, Rodríguez JP, Rosenberg AA, Samways MJ, Smart J, Stein BA, Stuart SN. 2010. The impact of conservation on the status of the world's vertebrates. *Science* **330**:1503–1509.
- Holbrook KM, Smith TB. 2000. Seed dispersal and movement patterns in two species of *Ceratogymna* hornbills in a West African tropical lowland forest. *Oecologia* **125**:249–257.
- Huante P, Rincon E, Chapin III FS. 1998. Foraging for nutrients, responses to changes in light, and competition in tropical deciduous tree seedlings. *Oecologia* **117**:209–216.
- Janzen DH. 1970. Herbivores and the number of tree species in tropical forests. *The American Naturalist* **104**:501–528.
- John R, Dalling JW, Harms KE, Yavitt JB, Stallard RF, Mirabello M, Hubbell SP, Valencia R, Navarrete H, Vallejo M, Foster RB. 2007. Soil nutrients influence spatial distributions of tropical tree species. *Proceedings of the National Academy of Sciences of the United States of America* **104**:864–869.
- Johns AD. 1988. Effects of “selective” timber extraction on rain forest structure and composition and some consequences for frugivores and folivores. *Source: Biotropica* **20**:31–37.
- Kasenene JM, Murphy PG. 1991. Post-logging tree mortality and major branch losses in Kibale Forest, Uganda. *Forest Ecology and Management* **46**:295–307.
- Kirika JM, Bleher B, Bohning-Gaese K, Chira R, Farwig N. 2008a. Fragmentation and local disturbance of forests reduce frugivore diversity and fruit removal in *Ficus thonningii* trees. *Basic and Applied Ecology* **9**:663–672.
- Kirika JM, Farwig N, Bohning-Gaese K. 2008b. Effects of local disturbance of tropical forests on frugivores and seed removal of a small-seeded Afrotropical tree. *Conservation Biology* **22**:318–328.
- Kleinschroth F, Healey JR. 2017. Impacts of logging roads on tropical forests. *Biotropica* **49**:620–635.
- Koenig WD, Knops JMH. 2001. Seed-crop size and eruptions of North American boreal seed-eating birds. *Journal of Animal Ecology* **70**:609–620.
- Koerner SE, Poulsen JR, Blanchard EJ, Okouyi J, Clark CJ. 2017. Vertebrate community composition and diversity declines along a defaunation gradient radiating from rural villages in Gabon. *Journal of Applied Ecology* **54**:805–814.
- Kurten EL. 2013. Cascading effects of contemporaneous defaunation on tropical forest communities. *Biological Conservation* **163**:22–32.
- LaDeau SL, Clark JS. 2001. Rising CO₂ levels and the fecundity of forest trees. *Science* **292**:95–98.
- LaDeau SL, Clark JS. 2006. Elevated CO₂ and tree fecundity: the role of tree size, interannual variability, and population heterogeneity. *Global Change Biology* **12**:822–833.
- LaManna JA, Martin TE. 2017. Logging impacts on avian species richness and composition differ across latitudes and foraging and breeding habitat preferences. *Biological Reviews of the Cambridge Philosophical Society* **92**:1657–1674.
- Laporte NT, Stabach JA, Grosch R, Lin TS, Goetz SJ. 2007. Expansion of industrial logging in Central Africa. *Science (New York, N.Y.)* **316**:1451.
- Laurance WF, Delamonica P, Laurance SG, Vasconcelos HL, Lovejoy TE. 2000. Rainforest fragmentation kills big trees. *Nature* **404**:836.
- Lewis SL, Phillips OL, Sheil D, Vinceti B, Baker TR, Brown S, Graham AW, Higuchi N, Hilbert DW, Laurance WF, Lejoly J, Malhi Y, Monteagudo A, Vargas PN, Sonké B, Supardi MNN, Terborgh JW, Martínez RV. 2004. Tropical forest tree mortality, recruitment and turnover rates: calculation, interpretation and comparison when census intervals vary. *Journal of Ecology* **92**:929–944.
- Lindenmayer DB, Laurance WF, Franklin JF. 2012. Ecology. Global decline in large old trees. *Science (New York, N.Y.)* **338**:1305–1306.
- Lutz JA, Larson AJ, Freund JA, Swanson ME, Bible KJ. 2013. The importance of large-diameter trees to forest structural heterogeneity. *PLoS One* **8**:e82784.
- Lutz JA, Larson AJ, Swanson ME, Freund JA. 2012. Ecological importance of large-diameter trees in a temperate mixed-conifer forest. *PLoS One* **7**:e36131.
- Markl JS, Schleuning M, Forget PM, Jordano P, Lambert JE, Traveset A, Wright SJ, Bohning-Gaese K. 2012. Meta-analysis of the effects of human disturbance on seed dispersal by animals. *Conservation Biology* **26**:1072–1081.
- Martinez I, Gonzalez-Taboada F. 2009. Seed dispersal patterns in a temperate forest during a mast event: performance of alternative dispersal kernels. *Oecologia* **159**:389–400.
- Medjibe VP, Putz FE, Romero C. 2013. Certified and uncertified logging concessions compared in Gabon: changes in stand structure, tree species, and biomass. *Environmental Management* **51**:524–540.
- Meijaard E, Sheil D, Nasi R, Augeri D, Rosenbaum B, Iskandar D, Setyawati T, Lammertink M, Rachmatika I, Wong A, Soehartono T, Stanley S, O'Brien T. 2005. *Life after logging. Reconciling wild-life conservation and production forestry in Indonesian Borneo*. Bogor, Indonesia: Center for International Forestry Research (CIFOR).

- Minor DM, Kobe RK. 2017. Masting synchrony in northern hardwood forests: super-producers govern population fruit production. *Journal of Ecology* **105**:987–998.
- Mockrin MH. 2008. *The spatial structure and sustainability of subsistence wildlife harvesting in Kaba, Congo*. PhD Thesis, Columbia University of Utrecht, USA.
- Molino JF, Sabatier D. 2001. Tree diversity in tropical rain forests: a validation of the intermediate disturbance hypothesis. *Science (New York, N.Y.)* **294**:1702–1704.
- Moran C, Catterall CP, Green RJ, Olsen MF. 2004. Functional variation among frugivorous birds: implications for rainforest seed dispersal in a fragmented subtropical landscape. *Oecologia* **141**:584–595.
- Morgan D, Sanz C. 2006. Chimpanzee feeding ecology and comparisons with sympatric gorillas in the Goulougo Triangle, Republic of Congo. In: Hohmann G, Robbins M, Boesch C, eds. *Feeding ecology in apes and other primates*. Cambridge: Cambridge University Press, 97–122.
- Muller-Landau HC, Hardesty BD. 2005. Seed dispersal of woody plants in tropical forests : concepts, examples and future directions. In: Burslem DFRP, Pinard MA, Hartley SE, eds. *Biotic interactions in the tropic*. Cambridge, UK: Cambridge University Press, 267–309.
- Muller-Landau HC, Wright SJ, Calderon O, Condit R, Hubbell SP. 2008. Interspecific variation in primary seed dispersal in a tropical forest. *Journal of Ecology* **96**:653–667.
- Nathan R, Muller-Landau HC. 2000. Spatial patterns of seed dispersal, their determinants and consequences for recruitment. *Trends in Ecology & Evolution* **15**:278–285.
- Neuschulz EL, Botzat A, Farwig N. 2011. Effects of forest modification on bird community composition and seed removal in a heterogeneous landscape in South Africa. *Oikos* **120**:1371–1379.
- Norghauer JM, Nock CA, Grogan J. 2011. The importance of tree size and fecundity for wind dispersal of big-leaf mahogany. *PLoS One* **6**:e17488.
- Pinard MA, Barker MG, Tay J. 2000. Soil disturbance and postlogging forest recovery on bulldozer paths in Sabah, Malaysia. *Forest Ecology and Management* **130**:213–225.
- Plumptre AJ. 1995. The importance of “seed trees” for the natural regeneration of selectively logged tropical forest. *Commonwealth Forestry Review* **74**:253–258.
- Poulsen JR, Clark CJ, Bolker BM. 2011. Decoupling the effects of logging and hunting on an Afrotropical animal community. *Ecological Applications* **21**:1819–1836.
- Poulsen JR, Clark CJ, Connor EF, Smith TB. 2002. Differential resource use by primates and hornbills: implications for seed dispersal. *Ecology* **83**:228–240.
- Poulsen JR, Clark CJ, Mavah G, Elkan PW. 2009. Bushmeat supply and consumption in a tropical logging concession in northern Congo. *Conservation Biology* **23**:1597–1608.
- Poulsen JR, Clark CJ, Palmer TM. 2013. Ecological erosion of an Afrotropical forest and potential consequences for tree recruitment and forest biomass. *Biological Conservation* **163**:122–130.
- Poulsen JR, Clark CJ, Smith TB. 2001. Seed dispersal by a diurnal primate community in the Dja Reserve, Cameroon. *Journal of Tropical Ecology* **17**:787–808.
- Putz FE, Blate GM, Redford KH, Fimbel R, Robinson J. 2001. Tropical forest management and conservation of biodiversity: an overview. *Conservation Biology* **15**:7–20.
- Rosin C, Poulsen JR. 2016. Hunting-induced defaunation drives increased seed predation and decreased seedling establishment of commercially important tree species in an Afrotropical forest. *Forest Ecology and Management* **382**:206–213.
- Schnitzer SA, Bongers F. 2002. The ecology of lianas and their role in forests. *Trends in Ecology and Evolution* **17**:223–230.
- Sist P. 2000. Reduced-impact logging in the tropics: objectives, principles and impacts. *International Forestry Review* **2**:3–10.
- Sist P, Fimbel R, Sheil D, Nasi R, Chevallier MH. 2003. Towards sustainable management of mixed dipterocarp forests of Southeast Asia: moving beyond minimum diameter cutting limits. *Environmental Conservation* **30**:364–374.
- Slik JWF, Paoli G, McGuire K, Amaral I, Barroso J, Bastian M, Blanc L, Bongers F, Boundja P, Clark C, Collins M, Dauby G, Ding Y, Doucet JL, Eler E, Ferreira L, Forshed O, Fredriksson G, Gillet JF, Harris D, Leal M, Laumonier Y, Malhi Y, Mansor A, Martin E, Miyamoto K, Araujo-Murakami A, Nagamasu H, Nilus R, Nurtjahya E, Oliveira A, Onrizal O, Parada-Gutierrez A, Permana A, Poorter L, Poulsen J, Ramirez-Angulo H, Reitsma J, Rovero F, Rozak A, Sheil D, Silva-Espejo J, Silveira M, Spironelo W, ter Steege H, Stevart T, Navarro-Aguilar GE, Sunderland T, Suzuki E, Tang J, Theilade I, van der Heijden G, van Valkenburg J, VanDo T, Vilanova E, Vos V, Wich S, Wöll H, Yoneda T, Zang R, Zhang MG, Zweifel N. 2013. Large trees drive forest above-ground biomass variation in moist lowland forests across the tropics. *Global Ecology and Biogeography* **22**:1261–1271.
- Stacey GR, Belcher RE, Wood CJ, Gardiner BA. 1994. Wind flows and forces in a model spruce forest. *Boundary-Layer Meteorology* **69**:311–334.
- Stephenson NL, Das AJ, Condit R, Russo SE, Baker PJ, Beckman NG, Coomes DA, Lines ER, Morris WK, Ruger N, Alvarez E, Blundo C, Bunyavejchewin S, Chuyong G, Davies SJ, Duque A, Ewango CN, Flores O, Franklin JF, Grau HR, Hao Z, Harmon ME, Hubbell SP, Kenfack D, Lin Y, Makana JR, Malizia A, Malizia LR, Pabst RJ, Pongpattananurak N, Su SH, Sun IF, Tan S, Thomas D, van Mantgem PJ, Wang X, Wiser SK, Zavala MA. 2014. Rate of tree carbon accumulation increases continuously with tree size. *Nature* **507**:90–93.
- Terborgh J, Nunez-Iturri G, Pitman NC, Valverde FH, Alvarez P, Swamy V, Pringle EG, Paine CE. 2008. Tree recruitment in an empty forest. *Ecology* **89**:1757–1768.
- Tews J, Brose U, Grimm V, Tielbörger K, Wichmann MC, Schwager M, Jeltsch F. 2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *Journal of Biogeography* **31**:79–92.
- Theimer TC, Gehring CA, Green PT, Connell JH. 2011. Terrestrial vertebrates alter seedling composition and richness but not diversity in an Australian tropical rain forest. *Ecology* **92**:1637–1647.
- Thomson FJ, Moles AT, Auld TD, Kingsford RT. 2011. Seed dispersal distance is more strongly correlated with plant height than with seed mass. *Journal of Ecology* **99**:1299–1307.
- Trollet F, Forget PM, Doucet JL, Gillet JF, Hambuckers A. 2017. Frugivorous birds influence the spatial organization of tropical forests through the generation of seedling recruitment foci under zoochoric trees. *Acta Oecologica* **85**:69–76.
- Tutin CE, Ham RM, White LJ, Harrison MJ. 1997. The primate community of the Lope Reserve, Gabon: diets, responses to fruit scarcity, and effects on biomass. *American Journal of Primatology* **42**:1–24.

Uriarte M, Clark JS, Zimmerman JK, Comita LS, Forero-Montana J, Thompson J. 2012. Multidimensional trade-offs in species responses to disturbance: implications for diversity in a subtropical forest. *Ecology* **93**:191–205.

Venable DL, Brown JS. 1988. The selective interactions of dispersal, dormancy, and seed size as adaptations for reducing risk in variable environments. *The American Naturalist* **131**:360–384.

Wang Y, Zhang J, LaMontagne JM, Lin F, Li B, Ye J, Yuan Z, Wang X, Hao Z. 2017. Variation and synchrony of tree species mast seeding in an old-growth temperate forest. *Journal of Vegetation Science* **28**:413–423.

White LJT, Abernethy K. 1997. *A guide to the vegetation of the Lopé Reserve, Gabon*. 2nd edn. Libreville: Multipress-Gabon.

Whitney KD, Fogiel MK, Lamperti AM, Holbrook KM, Stauffer DJ, Hardesty BD, Thomas Parker V, Smith TB. 1998. Seed dispersal by *Ceratogymna* hornbills in the Dja Reserve, Cameroon. *Journal of Tropical Ecology* **14**:351–371.

Willson M, Traveset A. 2000. The ecology of seed dispersal. In: Fenner M, ed. *Seeds: the ecology of regeneration in plant communities*. Oxfordshire, UK: Centre for Agriculture and Bioscience International, 85–110.

Wortley AH, Harris DJ. 2014. Sangha trees: an identification and training guide to the trees of the northern republic of Congo. In: Watson MF, Lyal CHC, Pendry CA, eds. *Descriptive taxonomy: the foundation of biodiversity research*. Cambridge, UK: Cambridge University Press, 127–145.