

CONFORMAL UNIFORMIZATION OF PLANAR PACKINGS BY DISK PACKINGS

DIMITRIOS NTALAMPEKOS

ABSTRACT. A Sierpiński packing in the 2-sphere is a countable collection of disjoint, non-separating continua with diameters shrinking to zero. We show that any Sierpiński packing by continua whose diameters are square-summable can be uniformized by a disk packing with a packing-conformal map, a notion that generalizes conformality in open sets. Being special cases of Sierpiński packings, Sierpiński carpets and some domains can be uniformized by disk packings as well. As a corollary of the main result, the conformal loop ensemble (CLE) carpets can be uniformized conformally by disk packings, answering a question of Rohde–Werness.

CONTENTS

1. Introduction	1
2. Preliminaries	7
3. Transboundary modulus estimates	15
4. Uniformization of Sierpiński packings	18
5. Topology of planar maps	31
6. Continuous extension	38
7. Homeomorphic extension	45
References	48

1. INTRODUCTION

One of the most intriguing open problems in complex analysis is *Koebe’s conjecture* [Koe08], predicting that every domain in the Riemann sphere is conformally equivalent to a *circle domain*, i.e., a domain whose complementary components are geometric disks or points. This conjecture was established for finitely connected domains by Koebe himself [Koe20] and it took over 70 years until it was established for countably connected domains by He–Schramm [HS93]. This result was proved with a different method by Schramm [Sch95] in a seminal work, where the notion of transboundary modulus was introduced. More recently, Rajala [Raj21] gave another proof of the result, providing a new perspective. Remarkably, in [Sch95], Schramm establishes Koebe’s conjecture for all cofat domains, i.e., domains whose complementary components satisfy a uniform geometric condition that we discuss

Date: July 3, 2023.

2020 Mathematics Subject Classification. Primary 30C20, 30C65; Secondary 30C35, 30C62.

Key words and phrases. Sierpiński carpet, Sierpiński packing, packing-conformal, packing-quasiconformal, conformal loop ensemble, uniformization, Koebe’s conjecture.

The author is partially supported by NSF Grant DMS-2000096.

below in Section 2.2, independently of connectivity. The general case of Koebe's conjecture seems to be far out of reach. Koebe's conjecture and uniformization problems for domains in metric surfaces other than the Riemann sphere has been studied in [MW13, RR21, Reh22].

A topic very closely related to Koebe's conjecture is the uniformization of Sierpiński carpets. A *Sierpiński carpet* is a continuum in the sphere that has empty interior and is obtained by removing from the sphere countably many open Jordan regions, called *peripheral disks*, with disjoint closures and diameters shrinking to zero. The boundaries of the peripheral disks are the *peripheral circles* of the carpet. A fundamental result of Whyburn [Why58] states that all Sierpiński carpets are homeomorphic to each other. Bonk [Bon11] proved that if the peripheral disks of a Sierpiński carpet are uniformly relatively separated, uniform quasidisks, then the carpet can be mapped with a quasimetric map to a *round carpet*, i.e., a carpet whose peripheral disks are geometric disks. Later, in [Nta20b] the author developed a potential theory on Sierpiński carpets of area zero and proved that if the peripheral disks of such a carpet are uniformly fat and uniformly quasiround, then the carpet can be mapped in a natural way to a *square carpet*, defined in the obvious manner, with a map that is *carpet-conformal* in the sense that it preserves a type of modulus. We note that the geometric assumptions in [Nta20b] are weaker than in [Bon11]. However, if one strengthens the assumptions to uniformly relatively separated uniform quasidisks, then the carpet-conformal map of [Nta20b] is upgraded to a quasimetric. Both mentioned works of Bonk and the author depend crucially on the notion of transboundary modulus of Schramm [Sch95].

In this work we push the results of [Bon11, Nta20b] to their limit and we *remove entirely the geometric assumptions* at the cost of weakening the topological properties and the regularity of the uniformizing conformal map. Instead, we only impose the square-summability of the diameters of the peripheral disks. Before stating the results we give the required definitions.

Let $\{p_i\}_{i \in \mathbb{N}}$ be a collection of pairwise disjoint and non-separating continua in the Riemann sphere $\widehat{\mathbb{C}}$ such that $\text{diam}(p_i) \rightarrow 0$ as $i \rightarrow \infty$. The collection $\{p_i\}_{i \in \mathbb{N}}$ is called a *Sierpiński packing* and the set $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ is its *residual set*. When it does not lead to a confusion, we make no distinction between the terms packing and residual set. The continua p_i , $i \in \mathbb{N}$, are called the *peripheral continua* of X . Note that if the peripheral continua of X are closed Jordan regions, then $\overline{X}$ is a Sierpiński carpet, provided that it has empty interior. Thus, Sierpiński packings can be regarded as a generalization of Sierpiński carpets.

The natural spaces that can be used to parametrize a Sierpiński packing are *round Sierpiński packings*, i.e., packings whose peripheral continua are (possibly degenerate) closed disks. We now state our main theorem.

Theorem 1.1. *Let $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be a Sierpiński packing whose peripheral continua are closed Jordan regions or points with diameters in $\ell^2(\mathbb{N})$. Then there exist*

- (A) *a collection of disjoint closed disks $\{p_i\}_{i \in \mathbb{N}}$, where p_i is degenerate if and only if q_i is degenerate, a round Sierpiński packing $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$,*
- (B) *a continuous, surjective, and monotone map $H: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ with the property that $H^{-1}(\text{int}(q_i)) = \text{int}(p_i)$ for each $i \in \mathbb{N}$, and*
- (C) *a non-negative Borel function $\rho_H \in L^2(\widehat{\mathbb{C}})$,*

with the following properties.

- (Transboundary upper gradient inequality) There exists a curve family Γ_0 in $\widehat{\mathbb{C}}$ with $\text{Mod}_2 \Gamma_0 = 0$ such that for all curves $\gamma: [a, b] \rightarrow \widehat{\mathbb{C}}$ outside Γ_0 we have

$$\sigma(H(\gamma(a)), H(\gamma(b))) \leq \int_{\gamma} \rho_H ds + \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

- (Conformality) For each Borel set $E \subset \widehat{\mathbb{C}}$ we have

$$\int_{H^{-1}(E)} \rho_H^2 d\Sigma \leq \Sigma(E \cap Y).$$

Moreover, if Y is cofat, then H may be taken to be a homeomorphism of the sphere.

Here σ denotes the spherical distance and Σ is the spherical measure. The monotonicity of H means that the preimage of every point is a continuum and is equivalent to the statement that H is the uniform limit of homeomorphisms; see Section 2.3. The map H in the conclusion of the theorem is called a *packing-conformal map*. Our definition of a packing-conformal map is motivated by the transboundary modulus of Schramm and by the so-called analytic definition of quasiconformality for maps between metric spaces [Wil12]. Moreover, an analogous definition under the terminology *weakly quasiconformal map* has been used recently by Romney and the author [NR22b, NR22a] in the solution of the problem of quasiconformal uniformization of metric 2-spheres of finite area.

If U is an open subset of $\widehat{\mathbb{C}}$ contained in the packing Y , then the map H of Theorem 1.1 is a conformal map in $H^{-1}(U)$ in the usual sense. However, not every conformal map between domains satisfies the transboundary upper gradient inequality. Nevertheless, one can show that this is always the case for countably connected domains.

Remark 1.2. We remark that although $H^{-1}(\text{int}(q_i)) = \text{int}(p_i)$ in Theorem 1.1 (B), the continuum $H^{-1}(q_i)$ might be larger than the disk p_i when the packing Y is not cofat. It is precisely this phenomenon that prevents us from proving Koebe's conjecture (e.g. for domains with countably many non-degenerate boundary components having diameters in ℓ^2) with this method; the phenomenon is illustrated in Lemma 2.14 and in Example 4.4. However, a non-trivial consequence of the topological and regularity conditions of Theorem 1.1 is that the map H is degenerate on the set $H^{-1}(q_i) \setminus p_i$, in the sense that it maps each continuum $E \subset H^{-1}(q_i) \setminus p_i$ to a point; see Figure 1. We prove this fact in Proposition 6.2.

In fact, Theorem 1.1 is a consequence of a more general uniformization theorem for Sierpiński packings Y without the topological assumption that the peripheral continua are closed Jordan regions or points. To each Sierpiński packing Y we can associate a topological sphere $\mathcal{E}(Y)$ by collapsing all peripheral continua to points, in view of Moore's decomposition theorem [Moo25].

Theorem 1.3. *Let $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be a Sierpiński packing such that the diameters of the peripheral continua lie in $\ell^2(\mathbb{N})$. Then there exists a round Sierpiński packing X and a packing-conformal map from $\mathcal{E}(X)$ onto $\mathcal{E}(Y)$.*

As we see, the uniformizing packing-conformal map exists only at the level of the topological spheres $\mathcal{E}(X), \mathcal{E}(Y)$, and in general does not induce a map between the packings X, Y in the sphere $\widehat{\mathbb{C}}$. For the definition of packing-conformal maps




FIGURE 1. The set $H^{-1}(q_i)$ might be larger than the disk p_i . In this figure it contains the shaded regions and bold “branches”, which are subsets of X . However, the map H is constant in each of them.

between the associated topological spheres see Section 2.6. The above theorem is restated as Corollary 4.2. The statement is proved via an approximation argument. We consider the finitely connected domains $Y_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n q_i$ and we uniformize them conformally by finitely connected circle domains X_n using Koebe’s theorem. Then our task is to show that the conformal maps from X_n to Y_n converge in a uniform sense to the desired limiting map from $\mathcal{E}(X)$ onto $\mathcal{E}(Y)$. This is where the $\ell^2(\mathbb{N})$ -summability of the diameters of the peripheral continua becomes important. In particular, we use this assumption to establish Lemma 3.1, which provides uniform transboundary modulus bounds; these bounds are then used in order to prove convergence. The $\ell^2(\mathbb{N})$ -summability condition is also used later in establishing the regularity properties of the limiting mapping; see Lemma 4.15.

This proof strategy (of using Koebe’s theorem for finitely connected domains and passing to the limit) is also followed by Schramm [Sch95] in showing that cofat domains can be uniformized by circle domains and in [Bon11] in uniformizing Sierpiński carpets by round carpets. The recent developments in the field of analysis on metric spaces and our much more thorough understanding of quasiconformal maps between metric spaces allow us to identify the topological and regularity properties of the limiting map in our more fractal setting, where no uniform geometry is imposed, as in the works of Schramm and Bonk.

We note that in unpublished work, Rohde and Werness [RW15] show that the complementary disks of the circle domain X_n converge in the Hausdorff sense after passing to a subsequence to a collection of pairwise disjoint disks. However, they were not able to identify the limit of the conformal maps from X_n to Y_n .

Theorem 1.1 is proved by showing that the topological assumptions on Y allow one to lift a packing-conformal map between $\mathcal{E}(X)$ and $\mathcal{E}(Y)$ as in Theorem 1.3 to the map H in the sphere $\widehat{\mathbb{C}}$ that has the desired properties. We remark that the regularity properties of packing-conformal maps are used crucially in establishing continuity and injectivity properties of H . The lifting process is achieved through Theorem 6.1, which provides a monotone map H . In the case that Y is cofat, the homeomorphism H as in the last part of Theorem 1.1 is provided by Theorem 7.1.




FIGURE 2. A CLE carpet (simulation by D.B. Wilson [Wil]).

Thus, Theorem 1.1 is a consequence of Corollary 4.2, Theorem 6.1, and Theorem 7.1.

Another generalization is that we do not need to restrict to round Sierpiński packings X in order to parametrize a given packing Y . Instead of using geometric disks as the peripheral continua of X , one can use homothetic images of any countable collection of uniformly fat and non-separating continua, such as squares. See Corollary 4.3 for the precise statement.

As a corollary of the main theorem we give an answer to a question of Rohde–Werness [RW15] regarding the uniformization of the conformal loop ensemble (CLE) carpet. CLE was introduced by Sheffield–Werner [SW12], as a random collection of Jordan curves in the unit disk that combines conformal invariance and a natural restriction property; see Figure 2. Each CLE gives rise to a Sierpiński carpet with non-uniform geometry; hence the current carpet uniformization theory of [Bon11, Nta20b] is not sufficient to treat them. However, Rohde–Werness [RW15] proved in unpublished work that, with probability 1, the diameters of the peripheral disks of a CLE carpet are square-summable. Therefore, we obtain the following corollary of the main theorem.

Corollary 1.4. *If Y is a CLE carpet, almost surely there exists a round Sierpiński packing X and a packing-conformal map that maps $\bar{X}$ onto Y .*

It would be interesting to obtain some stronger statements for the uniformization of CLE carpets. We pose several questions for further study.

Question 1.5. Under what conditions is the uniformizing round Sierpiński packing X and the packing-conformal map H of Theorem 1.1 unique (up to Möbius transformations)?

If one could at least show the uniqueness of X , then this would imply that CLE gives rise to another stochastic process that generates round packings.

Question 1.6. Under what conditions is $\bar{X}$ a carpet whenever $\bar{Y}$ is a carpet?

Theorem 1.1 already shows that a sufficient condition is the cofatness of Y . What about CLE carpets?

Question 1.7. Can one use the present techniques to prove Koebe’s conjecture for domains whose complementary components have diameters in ℓ^2 ?

Another natural question is whether one can obtain alternative proofs of results of [Bon11, Nta20b] upon strengthening the geometric assumptions on the peripheral continua.

Question 1.8. If the peripheral continua of a packing Y are uniformly relatively separated, uniform quasidisks, is the map H of Theorem 1.1 quasimetric?

As discussed above, the *specific* map H arising from the proof of Theorem 1.1 is a limit of conformal maps of finitely connected circle domains onto finitely connected approximations of Y . The same procedure is followed by Bonk [Bon11], who shows that the limiting map is quasimetric under the geometric assumptions of Question 1.8. Thus, the actual question is whether *any* map satisfying the conclusions of Theorem 1.1, and not just the one that arises as a limit, is quasimetric.

Question 1.9. If the peripheral continua of a packing Y are uniformly fat and uniformly quairound, is the map H of Theorem 1.1 carpet-conformal in the sense of [Nta20b]?

Our main theorem relates to the recent work of Hakobyan–Li [HL21], where the quasimetric embedding problem for non-planar carpets is studied. The authors study *dyadic slit carpets*, which are obtained as inverse limits of sequences of vertical slit domains in the unit square, endowed with the inner length metrics. The centers of the slits are located in centers of dyadic squares, whence the name dyadic slit carpets, while the lengths depend on some parameters of the construction and can differ from generation to generation. The slits give rise to the peripheral circles C_i , $i \in \mathbb{N}$, of the dyadic slit carpet Y that is generated by this construction. The main theorem of [HL21], Theorem 1.7, states that the carpet Y is quasimetric to a carpet in $\widehat{\mathbb{C}}$ if and only if $\{\text{diam}(C_i)\}_{i \in \mathbb{N}}$ lies in $\ell^2(\mathbb{N})$. This is precisely the main assumption in Theorem 1.1.

Summarizing, on one hand, in the current work we prove a uniformization result for carpets already embedded in $\widehat{\mathbb{C}}$ with potentially “bad” geometry, and on the other hand, [HL21] establishes a quasimetric embedding result for the specific family of dyadic slit carpets that have “good” geometry; e.g., the peripheral circles are always uniform quasicircles and each such carpet can be embedded in a *linearly locally connected* metric 2-sphere in a natural way. It would be interesting if the current techniques can be used to uniformize non-planar carpets under weaker assumptions; e.g. non-dyadic slit carpets or more generally carpets embedded in a 2-sphere of finite area whose peripheral circles have square-summable diameters. The recent work of Romney and the author [NR22a] on the uniformization of 2-spheres of finite area might be a valuable tool in this direction.

In the subsequent paper [Nta23c], we use the notion of packing-conformal maps and the results of the present paper in order to study the problem of *conformal rigidity of circle domains*, a problem that is closely related to the uniqueness in Koebe’s conjecture. A circle domain is *conformally rigid* if every conformal map from that domain onto another circle domain is the restriction of a Möbius transformation. Specifically, we prove that circle domains whose boundary is *CNED* (*countably negligible for extremal distances*), as introduced and studied in [Nta23b, Nta23a], are conformally rigid. This result unifies and extends all previous works in the subject [HS93, HS94, NY20]. Moreover, it provides strong evidence for a conjecture of

He-Schramm, asserting that a circle domain is rigid if and only if its boundary is *conformally removable*.

Acknowledgments. I would like to thank Dennis Sullivan for many thought-provoking and motivating conversations on the subject of planar uniformization and Koebe's conjecture. I first learned of the problem of the uniformization of CLE from Huy Tran whom I also thank for various discussions. Finally, I am grateful to the anonymous referees for carefully reading the manuscript and providing valuable comments and corrections.

2. PRELIMINARIES

2.1. Notation and terminology. Let (X, d) be a metric space. The open ball of radius $r > 0$, centered at a point $x \in X$ is denoted by $B_d(x, r)$; the corresponding closed ball is $\overline{B}_d(x, r)$. We also denote by $S(x, r)$ the circle $\{y \in X : d(x, y) = r\}$. The diameter of a set E is denoted by $\text{diam}_d(E)$. If the metric is implicitly understood we will often drop the symbol d from the subscript. For the Euclidean metric in the plane we will use the subscript e , when necessary. For example, we write $B_e(x, r)$ and $\text{diam}_e(E)$. Finally, we denote by σ the spherical metric and by Σ the spherical measure on the Riemann sphere $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$.

A continuous function γ from a compact interval $[a, b]$ into X is called a *compact curve*. A continuous function γ from (a, b) into X is called a *non-compact curve*. In this case, if γ extends continuously to a map $\bar{\gamma}: [a, b] \rightarrow X$ then it is called an *open curve*. The *trace* of a curve $\gamma: I \rightarrow X$ is the set $\gamma(I)$ and is denoted by $|\gamma|$. A curve $\gamma: [a, b] \rightarrow X$ is *closed* if $\gamma(a) = \gamma(b)$.

For $s \geq 0$ the *s-dimensional Hausdorff measure* $\mathcal{H}^s(E)$ of a set E in a metric space X is defined by

$$\mathcal{H}^s(E) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^s(E) = \sup_{\delta > 0} \mathcal{H}_\delta^s(E),$$

where

$$\mathcal{H}_\delta^s(E) = \inf \left\{ c(s) \sum_{j=1}^{\infty} \text{diam}(U_j)^s : E \subset \bigcup_j U_j, \text{diam}(U_j) < \delta \right\}$$

for a normalizing constant $c(s) > 0$ so that the n -dimensional Hausdorff measure agrees with Lebesgue measure in $\mathbb{R}^n$. Note that $c(1) = 1$. We will use the notation $\mathcal{H}_d^s$ for the Hausdorff measure $\mathcal{H}^s$ if we wish to emphasize that the metric d is used. We now state the co-area inequality for Lipschitz functions [EHa21, Theorem 2.1].

Proposition 2.1. *Let $L > 0$ and $\psi: \widehat{\mathbb{C}} \rightarrow \mathbb{R}$ be an L -Lipschitz function. Then for each Borel function $\rho: \widehat{\mathbb{C}} \rightarrow [0, \infty]$ we have*

$$\int_{\mathbb{R}} \int_{\psi^{-1}(t)} \rho d\mathcal{H}^1 dt \leq \frac{4L}{\pi} \int_{\widehat{\mathbb{C}}} \rho d\Sigma.$$

The cardinality of a set E is denoted by $\#E$. For quantities A and B we write $A \lesssim B$ if there exists a constant $c > 0$ such that $A \leq cB$. If the constant c depends on another quantity H that we wish to emphasize, then we write instead $A \leq c(H)B$ or $A \lesssim_H B$. Moreover, we use the notation $A \simeq B$ if $A \lesssim B$ and $B \lesssim A$. As previously, we write $A \simeq_H B$ to emphasize the dependence of the implicit constants on the quantity H . All constants in the statements are assumed

to be positive even if this is not stated explicitly and the same letter may be used in different statements to denote a different constant.

Let E be a set in a metric space X . For $r > 0$ we denote by $N_r(E)$ the open r -neighborhood of E . The Hausdorff distance of two sets $E, F \subset X$, denoted by $d_H(E, F)$, is defined as the infimum of all $r > 0$ such that $E \subset N_r(F)$ and $F \subset N_r(E)$. We say that a sequence of sets $E_n \subset X$, $n \in \mathbb{N}$, converges to a set E in the Hausdorff distance if $d_H(E_n, E) \rightarrow 0$ as $n \rightarrow \infty$. If the limiting set E is closed, then it consists precisely of all limit points of sequences $x_n \in E_n$, $n \in \mathbb{N}$.

A *continuum* is a compact and connected set. An elementary property of Hausdorff convergence is that it preserves connectedness; namely, if a sequence of continua E_n , $n \in \mathbb{N}$, converges to a compact set E , then E is also a continuum. See [BBI01, Section 7.3.1] for more background.

Lemma 2.2. *Let X, Y be compact metric spaces and $\pi: X \rightarrow Y$ be a continuous map.*

- (i) *Let $E \subset Y$ (resp. $E \subset X$) be a compact set and $\{E_n\}_{n \in \mathbb{N}}$ be a sequence of compact sets with the property that for each $r > 0$ there exists $N \in \mathbb{N}$ such that $E_n \subset N_r(E)$ for all $n > N$. Then for each $r > 0$ there exists $N \in \mathbb{N}$ such that*

$$\pi^{-1}(E_n) \subset N_r(\pi^{-1}(E)) \quad (\text{resp. } \pi(E_n) \subset N_r(\pi(E)))$$

for all $n > N$.

- (ii) *If $\{E_n\}_{n \in \mathbb{N}}$, $\{F_n\}_{n \in \mathbb{N}}$ are sequences of compact sets in Y converging in the Hausdorff sense to compact sets E, F , respectively, then*

$$\text{dist}(\pi^{-1}(E), \pi^{-1}(F)) \leq \liminf_{n \rightarrow \infty} \text{dist}(\pi^{-1}(E_n), \pi^{-1}(F_n)).$$

Proof. The first part follows from compactness and continuity. For the second part, note that

$$\text{dist}(\pi^{-1}(E), \pi^{-1}(F)) = \lim_{r \rightarrow 0} \text{dist}(N_r(\pi^{-1}(E)), N_r(\pi^{-1}(F))).$$

By the first part, for each $r > 0$, we have

$$\text{dist}(N_r(\pi^{-1}(E)), N_r(\pi^{-1}(F))) \leq \text{dist}(\pi^{-1}(E_n), \pi^{-1}(F_n))$$

for all sufficiently large $n \in \mathbb{N}$. This completes the proof. $\square$

2.2. Fat sets. Let $\tau > 0$. A measurable set $K \subset \widehat{\mathbb{C}}$ is τ -fat if for each $x \in K$ and for each ball $B_\sigma(x, r)$ that does not contain K we have $\Sigma(B_\sigma(x, r) \cap K) \geq \tau r^2$. A set is *fat* if it is τ -fat for some $\tau > 0$. Note that points are automatically τ -fat for every $\tau > 0$. A more modern terminology for fatness is Ahlfors 2-regularity, but we prefer to use the original terminology that was used by Schramm [Sch95] and Bonk [Bon11].

Lemma 2.3. *Let $\tau > 0$. If a connected set $K \subset \widehat{\mathbb{C}}$ is τ -fat and T is a Möbius transformation, then $T(K)$ is $c(\tau)$ -fat.*

Proof. Schramm [Sch95, Theorem 2.1] established the invariance of fatness under Möbius transformations. However, Schramm's definition of fatness uses the Euclidean metric rather than the spherical one, requiring that $\mathcal{H}_e^2(B_e(x, r) \cap K) \geq \tau r^2$ for every $x \in K \cap \mathbb{C}$ and ball $B_e(x, r)$ that does not contain K . Hence, it suffices to show that fatness according to the spherical metric is equivalent to fatness according to the Euclidean metric, quantitatively.

Suppose that K is τ -fat according to the spherical metric and fix $x \in K \cap \mathbb{C}$ and a ball $B_e(x, r)$ that does not contain K . Consider the annuli $A_e(x; (n-1)r/5, nr/5) = B_e(x, nr/5) \setminus \overline{B_e(x, (n-1)r/5)}$, $n \in \{2, \dots, 5\}$. One of these annuli, say A , has the property that $\text{dist}_e(A, 0) \geq r/5$. Since K is connected and not contained in $B_e(x, r)$, which contains A , there exists a point $y \in K$ lying in the circle that is equidistant from the boundary circles of A . The ball $B_e(y, r/10)$ is contained in $A \subset B_e(x, r)$. For all points $z \in B_e(y, r/10)$ we have $|z| \simeq r$. Thus, for all $z, w \in B_e(y, r/10)$ we have $\sigma(z, w) \simeq (1 + r^2)^{-1}|z - w|$. This implies that

$$\frac{\mathcal{H}_e^2(B_e(x, r) \cap K)}{r^2} \gtrsim \frac{\mathcal{H}_e^2(B_e(y, r/10) \cap K)}{\mathcal{H}_e^2(B_e(y, r/10))} \simeq \frac{\Sigma(B_e(y, r/10) \cap K)}{\Sigma(B_e(y, r/10))} \simeq_\tau 1,$$

since a Euclidean ball is also a spherical ball (of possibly different radius). Therefore, we conclude the fatness of K according to the Euclidean metric, quantitatively.

Conversely, suppose that K is fat according to the Euclidean metric and fix $x \in K$ and a ball $B_\sigma(x, r)$ that does not contain K . We apply an isometry P of $\widehat{\mathbb{C}}$ so that $P(x) = 0$. The set $P(K)$ is also fat according to the Euclidean metric by Schramm's result. We have

$$\Sigma(B_\sigma(x, r) \cap K) \geq \Sigma(B_\sigma(x, r/4) \cap K) = \Sigma(B_\sigma(0, r/4) \cap P(K)).$$

Since $B_\sigma(0, r/4)$ is contained in the unit disk in the plane, the identity map from $B_\sigma(0, r/4)$ into $(\mathbb{C}, |\cdot|)$ is uniformly bi-Lipschitz and $B_\sigma(0, r/4)$ corresponds to a Euclidean ball $B_e(0, cr)$ for some constant $c \simeq 1$. Thus,

$$\Sigma(B_\sigma(0, r/4) \cap P(K)) \simeq \mathcal{H}_e^2(B_e(0, cr) \cap P(K)) \gtrsim_\tau r^2.$$

This completes the proof. $\square$

Lemma 2.4. *Let $\tau > 0$ and $K_n \subset \widehat{\mathbb{C}}$, $n \in \mathbb{N}$, be a sequence of τ -fat compact sets. Then every compact limit of $\{K_n\}_{n \in \mathbb{N}}$ in the Hausdorff sense is τ -fat.*

Proof. Let $K \subset \widehat{\mathbb{C}}$ be a compact set that is the Hausdorff limit of a subsequence of K_n , $n \in \mathbb{N}$, which we denote by K_n for the sake of simplicity. If K is a point, then K is trivially τ -fat, so without loss of generality, we assume that $\text{diam}(K) > 0$. Let $x \in K$ and $B(x, r)$ be a ball that does not contain K . Our goal is to show that $\Sigma(B(x, r) \cap K) \geq \tau r^2$. Let $\varepsilon > 0$ and $U \supset K$ be an open set such that $\Sigma(B(x, r) \cap K) \geq \Sigma(B(x, r) \cap U) - \varepsilon$. For all sufficiently large $n \in \mathbb{N}$, we have $K_n \subset U$ by the Hausdorff convergence. Moreover, there exists a sequence $x_n \in K_n$ converging to x such that for each $\delta > 0$ we have $B(x_n, r - \delta) \subset B(x, r)$ for all sufficiently large $n \in \mathbb{N}$. Since $K \not\subset B(x, r)$, we have $K_n \not\subset B(x_n, r - \delta)$ for all sufficiently large $n \in \mathbb{N}$. Altogether, for all sufficiently large $n \in \mathbb{N}$ we have

$$\Sigma(B(x, r) \cap K) \geq \Sigma(B(x, r) \cap U) - \varepsilon \geq \Sigma(B(x_n, r - \delta) \cap K_n) - \varepsilon \geq \tau(r - \delta)^2 - \varepsilon.$$

We let $\delta \rightarrow 0$, and then $\varepsilon \rightarrow 0$ to obtain the desired conclusion. $\square$

The next statement can be found [MN22, Lemma 2.6 (iii)] in a slightly altered form.

Lemma 2.5. *Let $\tau > 0$ and $\{p_i\}_{i \in \mathbb{N}}$ be a collection of disjoint τ -fat continua in $\widehat{\mathbb{C}}$. For each compact set $E \subset \widehat{\mathbb{C}}$ and $a > 0$ the set*

$$\{i : p_i \cap E \neq \emptyset \text{ and } \text{diam}(p_i) \geq a \text{diam}(E)\}$$

has at most $c(\tau, a)$ elements.

We also record an elementary consequence of fatness; see [NY20, Property (F2), p. 154] for a proof.

Lemma 2.6. *Let $\tau > 0$ and $K \subset \widehat{\mathbb{C}}$ be a connected τ -fat set. Then for each $x \in \widehat{\mathbb{C}}$ and $r > 0$ we have*

$$\mathcal{H}^1(\{s \in (0, r) : K \cap S(x, s) \neq \emptyset\})^2 \leq c(\tau) \Sigma(K \cap B(x, r)).$$

A metric measure space (X, d, μ) is *doubling* if every ball in X has positive and finite measure and there exists a constant $L > 0$ such that

$$\mu(B(x, 2r)) \leq L\mu(B(x, r))$$

for each $x \in X$ and $r > 0$. In this case, we say that X is L -doubling.

Lemma 2.7. *Let (X, d, μ) be an L -doubling metric measure space for some $L > 0$. Let $p \geq 1$, $a \geq 1$, and $\{b_i\}_{i \in I}$ be a collection of non-negative numbers. Suppose that $\{D_i\}_{i \in I}$ is a family of measurable sets and $\{B_i = B(x_i, r_i)\}_{i \in I}$ is a family of balls in X with the property that $D_i \subset B_i$ and $\mu(B_i) \leq a\mu(D_i)$ for each $i \in I$. Then*

$$\left\| \sum_{i \in I} b_i \chi_{B_i} \right\|_{L^p(X)} \leq c(L, p, a) \left\| \sum_{i \in I} b_i \chi_{D_i} \right\|_{L^p(X)}$$

Proof. Note that for $p = 1$ the proof of the inequality is straightforward. Suppose that $p > 1$. For a non-negative measurable function ϕ on X consider the centered maximal function $M\phi$. For each $i \in I$ and $x \in D_i \subset B_i$ we have

$$\frac{1}{\mu(B_i)} \int_{B_i} \phi \leq \frac{1}{\mu(B_i)} \int_{B(x, 2r_i)} \phi \leq LM\phi(x).$$

Thus,

$$\int_{D_i} M\phi \geq L^{-1} \mu(D_i) \frac{1}{\mu(B_i)} \int_{B_i} \phi \geq L^{-1} a^{-1} \int_{B_i} \phi.$$

Now, let $f = \sum_{i \in I} b_i \chi_{B_i}$ and ϕ be an arbitrary non-negative function with $\|\phi\|_{L^q(X)} = 1$, where $1/p + 1/q = 1$. Then, by the Hardy–Littlewood maximal inequality for doubling metric measure spaces [HKST15, Theorem 3.5.6, p. 92] we have

$$\begin{aligned} \int f\phi &= \sum_{i \in I} b_i \int_{B_i} \phi \leq c(L, a) \sum_{i \in I} b_i \int_{D_i} M\phi = c(L, a) \int \left(\sum_{i \in I} b_i \chi_{D_i} \right) M\phi \\ &\leq c(L, a) \left\| \sum_{i \in I} b_i \chi_{D_i} \right\|_{L^p(X)} \|M\phi\|_{L^q(X)} \leq c(L, p, a) \left\| \sum_{i \in I} b_i \chi_{D_i} \right\|_{L^p(X)}. \end{aligned}$$

The duality between L^p and L^q shows the desired inequality. $\square$

2.3. Topological preliminaries. Let $\nu: X \rightarrow Y$ be a continuous map between topological spaces. The map ν is *proper* if the preimage of each compact set is compact. The map ν is *monotone* if the preimage of each point is a continuum. Moreover, ν is *cell-like* if the preimage of each point is a continuum that is contractible in all of its open neighborhoods. In 2-manifolds without boundary cell-like continua coincide with sets that have a simply connected neighborhood that they do not separate. In the 2-sphere, this condition is simply equivalent to the condition that the continuum is non-separating.

If ν is a continuous map from the 2-sphere onto itself, then it is monotone if and only if it is cell-like if and only if it is the uniform limit of homeomorphisms; see [You48] and [NR22a, Theorem 6.3]. In analogy, for non-compact manifolds without boundary, we have the Armentrout–Quinn–Siebenmann approximation theorem [Dav86, Corollary 25.1A, p. 189]. We formulate the theorem according to [Sie72].

Theorem 2.8. *A continuous, proper, and cell-like map $\nu: X \rightarrow Y$ between 2-manifolds without boundary can be uniformly approximated by homeomorphisms in the following strong sense. For each continuous function $\epsilon: X \rightarrow (0, \infty)$ and for each metric d on Y compatible with the topology, there exists a homeomorphism $\tilde{\nu}: X \rightarrow Y$ such that*

$$d(\nu(x), \tilde{\nu}(x)) < \epsilon(x)$$

for each $x \in X$.

Lemma 2.9. *Let $\nu: X \rightarrow Y$ be as in Theorem 2.8.*

- (i) *The map ν is surjective.*
- (ii) *For each open set $U \subset Y$ the set $\nu^{-1}(U)$ is homeomorphic to U .*
- (iii) *A compact set $E \subset Y$ is connected if and only if $\nu^{-1}(E)$ is connected.*
- (iv) *A compact set $E \subset Y$ is cell-like if and only if $\nu^{-1}(E)$ is cell-like.*
- (v) *Let $U \subset Y$ be an open set such that ν is injective on $\nu^{-1}(\partial U)$. Then $\nu^{-1}(\partial U) = \partial \nu^{-1}(U)$.*

Proof. For (i), suppose that $y_0 \in Y \setminus \nu(X)$. The properness of ν implies that there exists a ball $B(y_0, r)$ that is disjoint from $\nu(X)$. This contradicts the conclusion of Theorem 2.8. For (ii), let $U \subset Y$ be an open set. The map ν is a continuous, proper, and cell-like map from $\nu^{-1}(U)$ onto U . Theorem 2.8 implies that ν is the uniform limit of homeomorphisms from $\nu^{-1}(U)$ onto U . This proves (ii). The non-trivial direction in (iii) follows from the fact that $\nu|_{\nu^{-1}(E)}$ is a monotone map from the compact set $\nu^{-1}(E)$ onto E ; by [Why42, (2.2), Chap. VIII, p. 138], in this setting, the preimage of a connected set is connected. For part (iv), if E is cell-like, then there exists a simply connected neighborhood $U \supset E$ such that $U \setminus E$ is connected. Then by (ii) $\nu^{-1}(U)$ is a simply connected neighborhood of $\nu^{-1}(E)$ that is not separated by $\nu^{-1}(E)$; hence E is cell-like. Conversely, if $\nu^{-1}(E)$ is cell-like and U is a neighborhood of E , then $\nu^{-1}(E)$ is contractible in $\nu^{-1}(U)$, so E is contractible in U .

For (v), by continuity we have $\partial \nu^{-1}(U) \subset \nu^{-1}(\partial U)$. Moreover, for each $y \in \overline{U}$ the set $\nu^{-1}(y)$ intersects $\overline{\nu^{-1}(U)}$; indeed, if $y_n \in U$ and $y_n \rightarrow y$, then by properness there exists a sequence $x_n \in \nu^{-1}(y_n) \subset \nu^{-1}(U)$ converging to a point $x \in \nu^{-1}(y)$. The injectivity of ν on $\nu^{-1}(\partial U)$ implies that if $x \in \nu^{-1}(\partial U)$, then $x = \nu^{-1}(\nu(x))$. By the previous this point lies in $\overline{\nu^{-1}(U)}$, and thus in $\partial \nu^{-1}(U)$. $\square$

We will also use Moore's theorem [Moo25], which facilitates the study of planar domains, Sierpiński packings, and Sierpiński carpets. Let G be a partition of $\widehat{\mathbb{C}}$ into disjoint continua. We call G a *decomposition* of $\widehat{\mathbb{C}}$. We say that the decomposition G is *upper semicontinuous* if for each $g \in G$ and each open set $U \subset \widehat{\mathbb{C}}$ containing g , there exists an open set $V \subset \widehat{\mathbb{C}}$ containing g such that if $g' \in G$ and $g' \cap V \neq \emptyset$, then $g' \subset U$. Equivalently, if $g_n \in G$, $n \in \mathbb{N}$, is a sequence that converges in the Hausdorff sense to a compact set A , then there exists $g \in G$ such that $A \subset g$. We now state Moore's theorem; see [Dav86, Theorem 25.1] for a modern proof.

Theorem 2.10. *If G is an upper decomposition of $\widehat{\mathbb{C}}$ into non-separating continua, then $\widehat{\mathbb{C}}/G$ is homeomorphic to $\widehat{\mathbb{C}}$.*

2.4. Sierpiński packings. We recall some definitions from the introduction. Let $\{p_i\}_{i \in \mathbb{N}}$ be a collection of pairwise disjoint, non-separating continua in $\widehat{\mathbb{C}}$ such that $\text{diam}(p_i) \rightarrow 0$ as $i \rightarrow \infty$. The collection $\{p_i\}_{i \in \mathbb{N}}$ is called a Sierpiński packing and the set $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ is its residual set. When there is no confusion, we call X a Sierpiński packing and the underlying collection $\{p_i\}_{i \in \mathbb{N}}$ is implicitly understood. The continua p_i , $i \in \mathbb{N}$, are called the peripheral continua of X . A Sierpiński packing (resp. domain) is *cofat* if there exists $\tau > 0$ such that each of its peripheral continua (resp. complementary components) is τ -fat.

Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} p_i$ be a Sierpiński packing or a domain, where in the latter case the collection $\{p_i\}_{i \in I}$ is assumed to comprise the complementary components. We consider the quotient space $\mathcal{E}(X) = \widehat{\mathbb{C}}/\{p_i\}_{i \in I}$, together with the natural projection map $\pi_X: \widehat{\mathbb{C}} \rightarrow \mathcal{E}(X)$. For a set $A \subset \widehat{\mathbb{C}}$ we denote $\pi_X(A)$ by $\widehat{A}$. For a set $E \subset \widehat{\mathbb{C}}$, let $I_E = \{i \in I : p_i \cap E \neq \emptyset\}$. We define $\mathcal{E}(X; E) = \widehat{\mathbb{C}}/\{p_i\}_{i \in I \setminus I_E}$; that is, the sets p_i that intersect E are not collapsed to points.

If X is a Sierpiński packing or a domain, we note that the decomposition of $\widehat{\mathbb{C}}$ into the singleton points of X and the continua p_i , $i \in I$, is always upper semicontinuous. In the case that X is a Sierpiński packing, the fact that $\text{diam}(p_i) \rightarrow 0$ as $i \rightarrow \infty$ implies that for each set $E \subset \widehat{\mathbb{C}}$ the decomposition of $\widehat{\mathbb{C}}$ into the continua p_i , $i \in I \setminus I_E$, and the remaining singleton points is upper semicontinuous. Therefore, a consequence of Moore's theorem (Theorem 2.10) is the following statement.

Theorem 2.11. *Let X be a Sierpiński packing or a domain. Then $\mathcal{E}(X)$ is homeomorphic to $\widehat{\mathbb{C}}$. Moreover, if X is a Sierpiński packing and $E \subset \widehat{\mathbb{C}}$, then $\mathcal{E}(X; E)$ is homeomorphic to $\widehat{\mathbb{C}}$.*

A consequence of Lemma 2.9 that we will often use is that the preimages of continua under the projection maps π_X and $\pi_{X;E}$ are continua.

2.5. Transboundary modulus. First, we give the definition of 2-modulus on the sphere. Let Γ be a family of curves in $\widehat{\mathbb{C}}$. We say that a Borel function $\rho: \widehat{\mathbb{C}} \rightarrow [0, \infty]$ is *admissible* for the curve family Γ if

$$\int_{\gamma} \rho \, ds \geq 1$$

for each locally rectifiable curve $\gamma \in \Gamma$. We then define the 2-modulus, or else *conformal modulus*, of Γ as

$$\text{Mod}_2 \Gamma = \inf_{\rho} \int \rho^2 \, d\Sigma,$$

the infimum taken over all admissible functions ρ . The next lemma is simple consequence of the co-area inequality; see [Nta20b, Lemma 2.4.3] for an argument.

Lemma 2.12. *Let $f: \widehat{\mathbb{C}} \rightarrow \mathbb{R}$ be a Lipschitz function and Γ_0 be a family of curves in $\widehat{\mathbb{C}}$ with $\text{Mod}_2 \Gamma_0 = 0$. Then for a.e. $t \in \mathbb{R}$ every simple curve γ whose trace is contained in $f^{-1}(t)$ lies outside Γ_0 .*

Next, we define transboundary modulus, as introduced by Schramm [Sch95]. Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} p_i$ be a domain. Let $\rho: \mathcal{E}(X) \rightarrow [0, \infty]$ be a Borel function and

$\gamma: [a, b] \rightarrow \mathcal{E}(X)$ be a curve. Then there exist countably many curves γ_j , $j \in J$, such that for each $j \in J$ we have $|\gamma_j| \subset \widehat{X}$ and $\gamma_j = \pi_X \circ \alpha_j$ for some possibly non-compact curve α_j in the domain X . We define

$$\int_{\gamma} \rho ds = \sum_{j \in J} \int_{\alpha_j} \rho \circ \pi_X ds,$$

where this is understood to be infinite if one of the curves α_j is not locally rectifiable. Let Γ be a family of curves in $\mathcal{E}(X)$. We say that a Borel function $\rho: \mathcal{E}(X) \rightarrow [0, \infty]$ is *admissible* for Γ if

$$\int_{\gamma} \rho ds + \sum_{i: |\gamma| \cap \widehat{p}_i \neq \emptyset} \rho(\widehat{p}_i) \geq 1$$

for each $\gamma \in \Gamma$. The *transboundary modulus* of Γ with respect to the domain X is defined to be

$$\text{Mod}_X \Gamma = \inf_{\rho} \left\{ \int_X (\rho \circ \pi_X)^2 d\Sigma + \sum_{i \in \mathbb{N}} \rho(\widehat{p}_i)^2 \right\},$$

where the infimum is taken over all admissible functions ρ .

Let X, Y be domains in $\widehat{\mathbb{C}}$ and $f: X \rightarrow Y$ be a conformal map. Then f induces a homeomorphism $\widehat{f}: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ such that $\widehat{f} = \pi_Y \circ f \circ \pi_X^{-1}$ on $\widehat{X}$; see [NY20, Section 3] for a detailed discussion. It was observed by Schramm [Sch95] that transboundary modulus is invariant under conformal maps.

Lemma 2.13. *Let X, Y be domains in $\widehat{\mathbb{C}}$ and $f: X \rightarrow Y$ be a conformal map. Then for each curve family Γ in $\mathcal{E}(X)$ we have*

$$\text{Mod}_X \Gamma = \text{Mod}_Y \widehat{f}(\Gamma).$$

We also introduce the set function $f^* = \pi_Y^{-1} \circ \widehat{f} \circ \pi_X$ from the powerset of $\widehat{\mathbb{C}}$ into itself. In particular, $f^* = f$ on subsets of X and if A is contained in a boundary component of X , then $f^*(A)$ is the corresponding boundary component of Y . Observe that if $g = f^{-1}$, then

$$(2.1) \quad g_n^*(f_n^*(A)) \supset A$$

for each set $A \subset \widehat{\mathbb{C}}$ with equality if $A \subset X$. The next lemma is an implication of Carathéodory's kernel convergence theorem for multiply connected domains [Gol69, Theorem V.5.1, p. 228].

Lemma 2.14. *Let $\Omega \subset \widehat{\mathbb{C}}$ be a domain and f_n , $n \in \mathbb{N}$, be a sequence of conformal maps in Ω that converges locally uniformly in Ω to a conformal map f . Then for each compact $E \subset \widehat{\mathbb{C}}$ and for each compact limit E^* of $\{f_n^*(E)\}_{n \in \mathbb{N}}$ in the Hausdorff sense we have*

$$f^*(E) \supset E^*.$$

Proof. Suppose first that E is a complementary component of Ω . Since f_n converges to f locally uniformly, $f_n(\Omega)$ converges in the Carathéodory topology to $f(\Omega)$ (with respect to a point of $f(\Omega)$). This implies that each compact Hausdorff limit of $\widehat{\mathbb{C}} \setminus f_n(\Omega)$ is contained in $\widehat{\mathbb{C}} \setminus f(\Omega)$. Thus, if E^* is a compact limit of $f_n^*(E)$, which is a component of $\widehat{\mathbb{C}} \setminus f_n(\Omega)$, then E^* is contained in a component of $\widehat{\mathbb{C}} \setminus f(\Omega)$. One can now see that this component has to be $f^*(E)$.

For the general case, let E be an arbitrary compact set in $\widehat{\mathbb{C}}$. By the definition of f^* , the set $f^*(E)$ is the union of $f(\Omega \cap E)$ with the complementary components $f^*(B)$ of $f(\Omega)$, where B is a complementary component of Ω such that $B \cap E \neq \emptyset$. The set E^* consists of $f(\Omega \cap E)$ and Hausdorff limits of complementary components $f^*(B)$, where $B \cap E \neq \emptyset$. The previous case completes the proof. $\square$

2.6. Packing-quasiconformal maps. For two Sierpiński packings or domains X, Y , we introduce the notion of a packing-quasiconformal map between the associated topological spheres $\mathcal{E}(X), \mathcal{E}(Y)$.

Definition 2.15. Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} p_i$ and $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} q_i$ be Sierpiński packings or domains. Let $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ be a continuous, surjective, and monotone map such that $h(\widehat{p}_i) = \widehat{q}_i$ for each $i \in I$. We say that h is *packing-quasiconformal* if there exists $K \geq 1$ and a non-negative Borel function $\rho_h \in L^2(\widehat{\mathbb{C}})$ with the following properties.

- (*Transboundary upper gradient inequality*) There exists a curve family Γ_0 in $\widehat{\mathbb{C}}$ with $\text{Mod}_2 \Gamma_0 = 0$ such that for all curves $\gamma: [a, b] \rightarrow \widehat{\mathbb{C}}$ outside Γ_0 we have

$$\text{dist}(\pi_Y^{-1} \circ h \circ \pi_X(\gamma(a)), \pi_Y^{-1} \circ h \circ \pi_X(\gamma(b))) \leq \int_{\gamma} \rho_h ds + \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

- (*Quasiconformality*) For each Borel set $E \subset \widehat{\mathbb{C}}$ we have

$$\int_{\pi_X^{-1}(h^{-1}(\pi_Y(E)))} \rho_h^2 d\Sigma \leq K \Sigma(E \cap Y).$$

In this case, we say that h is *packing- K -quasiconformal*. If $K = 1$, then h is called *packing-conformal*.

A Borel function ρ_h satisfying the transboundary upper gradient inequality as above is called a *transboundary weak upper gradient of h* . If the transboundary upper gradient inequality holds for all rectifiable curves in $\widehat{\mathbb{C}}$, without the need to exclude a family of conformal modulus zero, then we say that ρ_h is a *transboundary upper gradient of h* .

Remark 2.16. Note that we are not requiring that $h^{-1}(\widehat{q}_i) = \widehat{p}_i$; in general $h^{-1}(\widehat{q}_i)$ could be much larger than $\widehat{p}_i$. See Example 4.4 for an instance. The quasiconformality condition implies that if $E = q_i$, then $\rho_h = 0$ a.e. on $\pi_X^{-1}(h^{-1}(\widehat{q}_i))$, $i \in \mathbb{N}$. Thus, ρ_h is supported in the set $\pi_X^{-1}(h^{-1}(\widehat{Y}))$. In fact, we can set ρ_h equal to 0 *everywhere* on $\pi_X^{-1}(h^{-1}(\widehat{q}_i))$, $i \in \mathbb{N}$, rather than almost everywhere. Indeed, line integrals are not affected by this change for Mod_2 -a.e. curve [HKST15, Lemma 5.2.16, p. 133]. Hence, by enlarging the exceptional curve family Γ_0 , we may have that the transboundary upper gradient inequality also holds for the modified function ρ_h .

The following lemma is straightforward for finitely connected domains; it is also true for countably connected domains but we will not need that generality.

Lemma 2.17. *Let $X, Y \subset \widehat{\mathbb{C}}$ be finitely connected domains and $f: X \rightarrow Y$ be a conformal map. Then the induced map $\widehat{f}: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ is packing-conformal and the derivative of f in the spherical metric is a transboundary upper gradient of $\widehat{f}$.*

Here, the derivative of f in the spherical metric at a point $z \in \widehat{\mathbb{C}}$ can be given by the following precise formula when $z, f(z) \in \mathbb{C}$.

$$|Df|(z) = \frac{1 + |z|^2}{1 + |f(z)|^2} |f'(z)|.$$

3. TRANSBOUNDARY MODULUS ESTIMATES

Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} p_i$ be a Sierpiński packing or a domain. Let $\widehat{E}, \widehat{F} \subset \mathcal{E}(X)$ be arbitrary sets and $\Omega \subset \mathcal{E}(X)$ be an open set. We denote by $\Gamma(\widehat{E}, \widehat{F}; \Omega)$ the family of open paths in Ω joining $\widehat{E}$ and $\widehat{F}$. That is, $\Gamma(\widehat{E}, \widehat{F}; \Omega)$ contains precisely the open paths $\gamma: (a, b) \rightarrow \Omega$ such that $\bar{\gamma}: [a, b] \rightarrow \bar{\Omega}$ intersects both $\widehat{E}$ and $\widehat{F}$; recall that an open path extends continuously to the endpoints by definition. If $\Omega = \mathcal{E}(X)$, then we simply write $\Gamma(\widehat{E}, \widehat{F})$.

Recall that for $E \subset \widehat{\mathbb{C}}$ we denote $I_E = \{i \in I : p_i \cap E \neq \emptyset\}$ and $\mathcal{E}(X; E) = \widehat{\mathbb{C}} / \{p_i\}_{i \in I \setminus I_E}$. Consider the natural projection $\pi_{X; E}: \widehat{\mathbb{C}} \rightarrow \mathcal{E}(X; E)$, which is injective on $X \cup E$. Recall that if X is a Sierpiński packing, and in particular $\text{diam}(p_i) \rightarrow 0$ as $i \rightarrow \infty$, then the space $\mathcal{E}(X; E)$ is homeomorphic to the sphere $\widehat{\mathbb{C}}$ by Theorem 2.11.

The next lemma is one of the main technical ingredients of the proof of Theorem 1.1.

Lemma 3.1 (Non-degeneracy lemma). *Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ be a Sierpiński packing such that the diameters of the peripheral continua lie in $\ell^2(\mathbb{N})$. Let $\widehat{E} \subsetneq \mathcal{E}(X)$ be a continuum such that $E = \pi_X^{-1}(\widehat{E})$ is non-degenerate and let d be a metric on $\mathcal{E}(X; E)$ that induces the quotient topology. For each $\delta > 0$ and $N \in \mathbb{N} \cup \{0\}$ there exists a constant $C(X, E, d, \delta, N) > 0$ such that the following statement is true.*

Let $F \subset \widehat{\mathbb{C}} \setminus E$ be a continuum such that $\text{diam}_d(\pi_{X; E}(F)) \geq \delta$. For a finite set $J \subset \mathbb{N}$ consider the domain $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in J} p_i$. Then for each set $J_0 \subset J$ with $\#J_0 \leq N$ we have

$$\text{Mod}_Y \Gamma \left(\pi_Y(E), \pi_Y(F \setminus \bigcup_{i \in J_0} p_i); \mathcal{E}(Y) \setminus \bigcup_{i \in J_0} \pi_Y(p_i) \right) \geq C(X, E, d, \delta, N).$$

Surprisingly, this lemma gives uniform lower modulus bounds, although the packing X does not have uniform geometry. The reason is that we freeze a continuum E and we consider curve families connecting relatively large continua F to E . If one varies the continuum E as well, then it is impossible to obtain uniform modulus bounds without some strong uniform geometric assumptions, as in [Bon11, Section 8].

Proof. Note that $E \neq \widehat{\mathbb{C}}$, since $\widehat{E} \neq \mathcal{E}(X)$. The set E is a continuum by Lemma 2.9 (iii). The space $Z = \mathcal{E}(X; E)$ is homeomorphic to $\widehat{\mathbb{C}}$. The projection $\pi = \pi_{X; E}: \widehat{\mathbb{C}} \rightarrow Z$ is injective on the set E , thus, $\pi(E)$ is a non-degenerate continuum in Z . We endow Z with a metric d inducing its topology, as in the statement of the lemma. We also fix $\delta > 0$.

The set $Z \setminus \pi(E \cup \bigcup_{i \in \mathbb{N}} p_i)$ is non-empty by Baire's theorem. Fix a point x_∞ in that set and a ball $B_d(x_\infty, r) \subset Z \setminus \pi(E)$, where $r < \delta/4$. Consider a homeomorphism φ from $Z \setminus \{x_\infty\}$ onto the plane $\mathbb{C}$ with the property that it maps the




FIGURE 3. Left: Case 1a. Middle: Case 1b. Right: Case 2.

complement of the ball $B_d(x_\infty, r)$ into the unit ball $B_e(0, 1)$. Let $E' = \varphi(\pi(E))$, which is a non-degenerate continuum in $B_e(0, 1)$. By uniform continuity, there exists a constant $c_0 > 0$ such that if $\widehat{F} \subset Z$ is a continuum with $\text{diam}_d(\widehat{F}) \geq \delta$, then $\varphi(\widehat{F} \setminus B_d(x_\infty, r))$ contains a continuum $F' \subset B_e(0, 1)$ with $\text{diam}_e(F') \geq c_0$. We fix such a continuum F' .

Now, for $\varepsilon > 0$ consider a grid of squares of side length ε in the plane with sides parallel to the coordinate axes such that the 1-skeleton avoids the countably many points $p'_i = \varphi(\pi(p_i))$, $i \in \mathbb{N} \setminus I_E$. Let $\mathcal{G}$ be the intersection of the 1-skeleton with the square $[-2, 2]^2$. Let $\mathcal{B}'$ be the collection of simple paths that are contained in $\mathcal{G}$ and connect all pairs of junction points (i.e., points where four edges meet) of $\mathcal{G}$. Note that $\mathcal{B}'$ contains a bounded number of paths, depending only on ε . Also, if two paths of $\mathcal{B}'$ do not intersect, then their distance is at least ε . For each path $\beta \in \mathcal{B}'$, consider a Jordan region $U(\beta)$ such that $|\beta| \subset U(\beta) \subset N_{\varepsilon/2}(|\beta|)$ and $\partial U(\beta)$ avoids $\bigcup_{i \in \mathbb{N} \setminus I_E} p'_i$. In particular, observe that if $|\beta_1| \cap |\beta_2| = \emptyset$, then $U(\beta_1) \cap U(\beta_2) = \emptyset$.

Let $N \in \mathbb{N} \cup \{0\}$, as in the statement of the lemma. We claim that if the mesh of the grid is sufficiently small, i.e., ε is sufficiently small, depending only on N , E' , and c_0 , but not on F' , then $\mathcal{B}'$ contains $N + 1$ disjoint paths $\beta_1, \dots, \beta_{N+1}$ that connect E' and F' with the additional feature that $\partial U(\beta_1), \dots, \partial U(\beta_{N+1})$ also connect E' and F' . To see this, we consider separate cases.

Case 1. Suppose that the projection of E' to the x -axis has positive diameter equal to c_1 and the projection of F' to the x -axis has diameter larger than $c_0/2$. We set $M = \min\{c_1, c_0/2\}$.

Case 1a. Suppose that the overlap of the projections is larger than or equal to $M/2$. Then by choosing $\varepsilon < 2^{-1}M(N+3)^{-1}$ we may find $N+1$ disjoint vertical paths in $\mathcal{B}'$ intersecting both E' and F' so that their parallel translates by ε to either side also have the same property. See Figure 3.

Case 1b. Suppose that the overlap of the projections is smaller than $M/2$. Then there exist subcontinua E'' and F'' of E' and F' , respectively, such that the projections of E'' and F'' to the x -axis are disjoint and both have diameter $M/2$. If we choose $\varepsilon < 2^{-1}M(N+3)^{-1}$, then there exist $N+1$ disjoint vertical paths in $\mathcal{B}'$ intersecting F'' but not E'' and $N+1$ disjoint vertical paths intersecting E'' but not F'' so that their parallel translates by ε to either side also have the same property. We truncate these paths appropriately outside $B_e(0, 1)$ and then connect them with disjoint horizontal segments in $\mathcal{B}'$ to obtain Π -shaped paths; see Figure 3. The resulting collection of paths has the desired properties.

Case 2. Suppose that the projection of E' to the x -axis has diameter zero and the projection of F' to the x -axis has diameter larger than $c_0/2$. Thus, E' is a vertical line segment that projects to a segment of diameter c_1 in the y -axis. Our goal is to find appropriate subcontinua E'' and F'' of E' and F' , respectively, whose diameters are bounded below depending on c_0 and c_1 and whose projections to both axes are disjoint. First, we consider a subcontinuum F'' of F' with diameter $c_0/8$ whose projection to the x -axis is disjoint from the projection of E' . If the projection of F'' to the y -axis has diameter larger than or equal to $\min\{c_1/2, c_0/16\}$, then we can argue as in Case 1, replacing the x -axis with the y -axis and finding a value of ε that depends on c_0, c_1 , and N . Otherwise, we consider a subsegment E'' of E' of length $c_1/4$ so that the projections of E'' and F'' to both axes are disjoint. Then by choosing a small enough ε , we may find Γ -shaped curves from the collection $\mathcal{B}'$ that join E'' and F'' and have the desired properties; see Figure 3. The remaining cases are symmetric to the ones we treated.

Note that $(\varphi \circ \pi)^{-1}$ is injective on the square grid $\mathcal{G}$ and on the boundaries $\partial U(\beta)$, $\beta \in \mathcal{B}'$, since these sets avoid the set $\bigcup_{i \in \mathbb{N} \setminus I_E} p'_i$. We pull back the collection $\mathcal{B}'$ and the regions $U(\beta)$, $\beta \in \mathcal{B}'$, to $\widehat{\mathbb{C}}$ under the proper and cell-like map $\varphi \circ \pi: \widehat{\mathbb{C}} \setminus \pi^{-1}(x_\infty) \rightarrow \mathbb{C}$. Using Lemma 2.9 (v) we obtain a collection $\mathcal{B}$ of simple curves and Jordan regions $U(\beta)$, $\beta \in \mathcal{B}$, in $\widehat{\mathbb{C}}$ with the following properties.

- (i) For each $\beta \in \mathcal{B}$ the Jordan region $U(\beta)$ contains $|\beta|$ and if $\beta_1, \beta_2 \in \mathcal{B}$ are disjoint curves, then $U(\beta_1)$ and $U(\beta_2)$ are disjoint and intersect disjoint collections of sets p_i , $i \in \mathbb{N} \setminus I_E$.
- (ii) Whenever $F \subset \widehat{\mathbb{C}} \setminus E$ is a continuum with $\widehat{F} = \pi_{X;E}(F)$ and $\text{diam}_d(\widehat{F}) \geq \delta$, there exist disjoint paths $\beta_1, \dots, \beta_{N+1} \in \mathcal{B}$ such that β_i and $\partial U(\beta_i)$ intersect E and F for each $i \in \{1, \dots, N+1\}$.

We let $\eta = \min\{\text{dist}(|\beta|, \partial U(\beta)) : \beta \in \mathcal{B}\}$, which is positive since $\mathcal{B}$ is a finite collection.

Fix a continuum $F \subset \widehat{\mathbb{C}} \setminus E$ and paths $\beta_1, \dots, \beta_{N+1} \in \mathcal{B}$ as in (ii). If $J_0 \subset J$ is a set with $\#J_0 \leq N$ as in the statement of the lemma, then by (i) and the pigeonhole principle there exists $k \in \{1, \dots, N+1\}$ such that $U(\beta_k)$ does not intersect $\bigcup_{i \in J_0 \setminus I_E} p_i$. Note that $F \cap U(\beta_k)$ does not intersect $\bigcup_{i \in J_0} p_i$, but this is not necessarily true for E , since it intersects p_i , whenever $i \in I_E \cap J_0$. Let $\psi(x) = \text{dist}(x, |\beta_k|)$, which is a 1-Lipschitz function on $\widehat{\mathbb{C}}$. For a.e. $t \in (0, \eta)$ the components of $\psi^{-1}(t)$ are points, Jordan curves, or Jordan arcs; see [Nta20a, Theorem 1.5] for a general statement in metric surfaces or [Bro72] for a planar version. Hence, for a.e. $t \in (0, \eta)$, the set $\psi^{-1}(t)$ contains a Jordan curve separating $|\beta_k|$ from $\partial U(\beta_k)$. We fix such a $t \in (0, \eta)$. Since E and $F \setminus \bigcup_{i \in J_0} p_i$ connect β_k and $\partial U(\beta_k)$, we conclude they intersect this Jordan curve. Thus, $\psi^{-1}(t)$ contains an open curve connecting E and $F \setminus \bigcup_{i \in J_0} p_i$ and avoiding $\bigcup_{i \in J_0} p_i$. It follows that $\pi_Y(\psi^{-1}(t))$ contains a curve in $\Gamma = \Gamma(\pi_Y(E), \pi_Y(F \setminus \bigcup_{i \in J_0} p_i); \mathcal{E}(Y) \setminus \bigcup_{i \in J_0} \pi_Y(p_i))$. Thus, if $\rho: \mathcal{E}(Y) \rightarrow [0, \infty]$ is admissible for Γ , then

$$\int_{\psi^{-1}(t) \cap Y} \rho \circ \pi_Y d\mathcal{H}^1 + \sum_{\substack{i: \psi^{-1}(t) \cap p_i \neq \emptyset \\ i \in J}} \rho(\widehat{p}_i) \geq 1$$

for a.e. $t \in (0, \eta)$. Integrating over $t \in (0, \eta)$, and using the fact that ψ is 1-Lipschitz, by the co-area inequality (Proposition 2.1) we obtain

$$\begin{aligned} C\eta &\leq \int_{N_\eta(|\beta_k|) \cap Y} \rho \circ \pi_Y d\Sigma + \sum_{i \in J} \rho(\widehat{p}_i) \text{diam}(p_i) \\ &\leq \left(\int_Y (\rho \circ \pi_Y)^2 d\Sigma + \sum_{i \in J} \rho(\widehat{p}_i)^2 \right)^{1/2} \left(\Sigma(\widehat{\mathbb{C}}) + \sum_{i \in \mathbb{N}} \text{diam}(p_i)^2 \right)^{1/2}. \end{aligned}$$

Infimizing over ρ gives

$$\text{Mod}_Y \Gamma \geq C' \eta^2 \left(1 + \sum_{i \in \mathbb{N}} \text{diam}(p_i)^2 \right)^{-1}.$$

This completes the proof. $\square$

The next lemma is established in [Bon11]. The *relative distance* of two non-degenerate subsets E, F of a metric space is defined as

$$\Delta(E, F) = \frac{\text{dist}(E, F)}{\min\{\text{diam}(E), \text{diam}(F)\}}.$$

Lemma 3.2 ([Bon11, Proposition 8.7]). *For each $\tau > 0$ there exists a number $N_0 > 0$ and a function $\psi: (0, \infty) \rightarrow (0, \infty)$ with $\lim_{t \rightarrow \infty} \psi(t) = 0$ such that the following is true. Let $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in I} p_i$ be a τ -cofat finitely connected domain. Let $\widehat{E}, \widehat{F} \subset \mathcal{E}(X)$ be disjoint sets and let $E = \pi_X^{-1}(\widehat{E})$, $F = \pi_X^{-1}(\widehat{F})$. If $\Delta(E, F) \geq 12$, then there exists a set $I_0 \subset I$ with $\#I_0 \leq N_0$ such that*

$$\text{Mod}_X \Gamma \left(\widehat{E}, \widehat{F}; \mathcal{E}(X) \setminus \bigcup_{i \in I_0} \widehat{p}_i \right) \leq \psi(\Delta(E, F)).$$

The statement here is slightly different from [Bon11], but the proof remains unchanged. We point out the main differences. First, Bonk uses continua E, F , while we use arbitrary sets. Note that upper modulus bounds are not affected by this generalization; it is instead that lower modulus bounds require continua so that the curve family that connects them is rich enough. Second, in [Bon11] the continua E, F are chosen in $\overline{X}$, but we choose sets $\widehat{E}, \widehat{F}$ in $\mathcal{E}(X)$ and then take their preimages in $\widehat{\mathbb{C}}$. Third, we use path families in $\mathcal{E}(X)$ rather than in $\widehat{\mathbb{C}}$. Each path in $\mathcal{E}(X)$ corresponds to countably many possibly non-compact paths in the domain X and thus the considerations in [Bon11], which use line integrals in $\widehat{\mathbb{C}}$, are applicable here as well.

4. UNIFORMIZATION OF SIERPIŃSKI PACKINGS

Our goal in this section is to prove the following theorem at the heart of the paper.

Theorem 4.1. *Let $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be a Sierpiński packing such that $\{\text{diam}(q_i)\}_{i \in \mathbb{N}}$ lies in $\ell^2(\mathbb{N})$. Let $\tau > 0$ and $\zeta_\infty, \zeta_0, \zeta_1 \in Y$, and for each $n \in \mathbb{N}$, let f_n be a conformal map from the domain $Y_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n q_i$ onto a τ -cofat domain $X_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n p_{i,n}$ such that $f_n(\zeta_\infty) = \infty$, $f_n(\zeta_0) = 0$, $|f_n(\zeta_1)| = 1$, and $f_n^*(q_i) = p_{i,n}$ for each $i \in \{1, \dots, n\}$. Suppose, in addition, that for each $i \in \mathbb{N}$ all Hausdorff limits of*

the sequence $\{p_{i,n}\}_{n \geq i}$ are non-separating. Then there exists a sequence $\{k_n\}_{n \in \mathbb{N}}$ increasing to ∞ with the following properties.

- (i) For each $i \in \mathbb{N}$, p_{i,k_n} converges as $n \rightarrow \infty$ to a τ -fat non-separating continuum p_i in the Hausdorff sense.
- (ii) The set $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ is a τ -cofat Sierpiński packing.
- (iii) For each $i \in \mathbb{N}$, p_i is non-degenerate if and only if q_i is non-degenerate.
- (iv) The sequence $g_{k_n} = f_{k_n}^{-1}: X_{k_n} \rightarrow Y_{k_n}$ gives rise to a sequence of set functions $h_n = \pi_Y \circ g_{k_n}^* \circ \pi_X^{-1}$ from subsets of $\mathcal{E}(X)$ to subsets of $\mathcal{E}(Y)$ that converges uniformly to a packing-conformal map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$.

The proof of the theorem is given in Sections 4.1–4.3. We now record two immediate corollaries. Suppose that the maps f_n are given by Koebe’s uniformization theorem for finitely connected domains [Koe20]. In this case, the sets $p_{i,n}$ are geometric disks and points. Each of their Hausdorff limits is a geometric disk (allowing the case of the entire sphere) or a point, so it is non-separating. Thus, we obtain the following corollary.

Corollary 4.2. *Let $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be a Sierpiński packing such that $\{\text{diam}(q_i)\}_{i \in \mathbb{N}}$ lies in $\ell^2(\mathbb{N})$. Then there exists a round Sierpiński packing $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ and a packing-conformal map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$. Moreover, for each $i \in \mathbb{N}$, the disk p_i is non-degenerate if and only if q_i is non-degenerate.*

More generally, one can consider a collection $\{P_i\}_{i \in \mathbb{N}}$ of non-degenerate, τ -fat, and non-separating continua in the plane $\mathbb{C}$, which is regarded as a subset of $\widehat{\mathbb{C}}$. According to the Brandt–Harrington uniformization theorem [Bra80, Har82] there exists a conformal map f_n from the finitely connected domain $Y_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n q_i$ onto a domain $X_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n p_{i,n}$, where $p_{i,n}$ either is a point or is homothetic to P_i ; that is, $p_{i,n}$ is the image of P_i under a transformation $z \mapsto az + b$, $a > 0$, $b \in \mathbb{C}$. By postcomposing f_n with a homothetic transformation, we may assume that it satisfies the normalizations of Theorem 4.1. By Lemma 2.3, the sets $p_{i,n}$ are $c(\tau)$ -fat. Also, note that if $p_{i,n}$ is homothetic to P_i , then each non-degenerate Hausdorff limit of $p_{i,n}$ as $n \rightarrow \infty$ is also homothetic to P_i , provided that it is not the entire sphere. Therefore, Theorem 4.1 gives the following corollary.

Corollary 4.3. *Let $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be a Sierpiński packing such that $\{\text{diam}(q_i)\}_{i \in \mathbb{N}}$ lies in $\ell^2(\mathbb{N})$. For $\tau > 0$ consider a collection $\{P_i\}_{i \in \mathbb{N}}$ of non-degenerate, τ -fat, and non-separating continua in the plane $\mathbb{C}$. Then there exists a Sierpiński packing $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$, where p_i is a point when q_i is a point and p_i is homothetic to P_i when q_i is non-degenerate, and there exists a packing-conformal map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$.*

Example 4.4. We provide an example showing that in general we only expect to have $h^{-1}(\widehat{q}_i) \supset \widehat{p}_i$, $i \in \mathbb{N}$, for the map h of Theorem 4.1 and we cannot guarantee equality.

The following example is presented in [Nta23c, Section 6]. There exists a countably connected domain $Y = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ whose complementary components $\{q_i\}_{i \in \mathbb{N}}$ have diameters in $\ell^2(\mathbb{N})$ and a conformal map F from Y onto a circle domain Z such that q_1 is a singleton, $F^*(q_1)$ is a non-degenerate disk, and each component q_i , $i \geq 2$, is isolated.

Let $Y_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n q_i$ and consider normalized conformal maps f_n from Y_n onto a finitely connected circle domain $X_n = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^n p_{i,n}$ as in the statement of Theorem

4.1. After passing to a subsequence, f_n converges locally uniformly in the domain Y to a conformal embedding f from Y into $\widehat{\mathbb{C}}$. The fact that each component q_i , $i \geq 2$, is isolated implies that $f^*(q_i)$ is a geometric disk p_i that is a Hausdorff limit of $p_{i,n} = f_n^*(q_i)$; this can be justified by using Carathéodory's kernel convergence [Gol69, Theorem V.5.1]. However, we have no information about $f^*(q_1)$. The map $F \circ f^{-1}$ is a conformal map from $f(Y)$ onto Z . Based on transboundary modulus arguments (see e.g. [Sch95, Theorem 6.1]), and given that $F^*(q_1)$ is a non-degenerate disk, one can show that $f^*(q_1)$ is a non-degenerate continuum. Observe that $f(Y) = \widehat{\mathbb{C}} \setminus (f^*(q_1) \cup \bigcup_{i=2}^{\infty} p_i)$.

If we apply Theorem 4.1, after passing to a subsequence, we obtain a round Sierpiński packing $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$, where p_1 is a Hausdorff limit of $p_{1,n}$. Since q_1 is a singleton, we conclude that p_1 is a singleton. Also, by the last part of the theorem, the sequence $h_n = \pi_Y \circ g_n^* \circ \pi_X^{-1}$ (where $g_n = f_n^{-1}$) converges uniformly to a packing-conformal map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$.

The convergence of f_n implies that the sequence $g_n = f_n^{-1}$ of conformal maps converges locally uniformly in $f(Y)$ to $g = f^{-1}$ and $g^*(f^*(q_1)) = q_1$. One can show that $\pi_Y \circ g^* \circ \pi_X^{-1} = h$. Using that, we have $h(\pi_X(f^*(q_1))) = \widehat{q}_1$. Note that $f^*(q_1) \supsetneq p_1$ by Lemma 2.14, so $\pi_X(f^*(q_1)) \supsetneq \widehat{p}_1$. It follows that $h^{-1}(\widehat{q}_1) \supsetneq \widehat{p}_1$.

4.1. Existence of limiting Sierpiński packing. We now initiate the proof of Theorem 4.1. Recall that the conformal map $f_n: Y_n \rightarrow X_n$ gives rise to a set function $f_n^* = \pi_{X_n}^{-1} \circ \widehat{f}_n \circ \pi_{Y_n}$ from the powerset of $\widehat{\mathbb{C}}$ into itself. Namely, $f_n^* = f_n$ in Y_n and $f_n^*(A) = p_{i,n}$ whenever $A \subset q_i$. By Lemma 2.9 (iii), f_n^* maps continua to continua.

Lemma 4.5 (Non-degeneracy). *Let $\widehat{E} \subset \mathcal{E}(Y)$ be a continuum such that $E = \pi_Y^{-1}(\widehat{E})$ is non-degenerate. Then*

$$\liminf_{n \rightarrow \infty} \text{diam}(f_n^*(E)) > 0.$$

Proof. Without loss of generality $\widehat{E} \neq \mathcal{E}(Y)$, since in that case we have $f_n^*(E) = \widehat{\mathbb{C}}$ for all $n \in \mathbb{N}$. The set E is a continuum and has the property that it contains all peripheral continua of Y that intersect E . We set $E'_n = f_n^*(E)$, which is a non-degenerate continuum by the conformality of f_n , and suppose that $\text{diam}(E'_n) \rightarrow 0$ along a subsequence. We wish to derive a contradiction. After passing to a further subsequence, we assume that E'_n converges to a point $x_0 \in \widehat{\mathbb{C}}$ in the Hausdorff sense. Consider the great circle through the points $0 = f_n(\zeta_0)$, $f_n(\zeta_1)$, and $\infty = f_n(\zeta_\infty)$. The point x_0 does not lie on at least one of the arcs (of the great circle) from 0 to $f_n(\zeta_1)$, $f_n(\zeta_1)$ to ∞ , or ∞ to 0. Without loss of generality, after passing to a subsequence, suppose that E'_n stays away from the arc α_n joining 0 to $f_n(\zeta_1)$. That is, there exists $c > 0$ such that $\text{dist}(E'_n, |\alpha_n|) \geq c$ for all $n \in \mathbb{N}$. Let $F'_n = \pi_{X_n}^{-1}(\pi_{X_n}(|\alpha_n|))$, i.e., the union of the arc $|\alpha_n|$, together with all the sets $p_{j,n}$, $j \in \{1, \dots, n\}$, that intersect $|\alpha_n|$. Note that F'_n might be very close to E'_n , although they are disjoint, because they both contain the complementary components of X_n that they intersect.

The τ -cofatness of X_n and Lemma 2.5 imply that there exists a constant $N_0 \in \mathbb{N}$, depending only on τ and c , such that at most N_0 of the sets $p_{i,n}$, $i \in \{1, \dots, n\}$, with diameter larger than $c/2$ intersect the arc $|\alpha_n|$. Thus, there exists a set

$J_n \subset \{1, \dots, n\}$ with $\#J_n \leq N_0$ such that

$$\text{dist}(E'_n, F'_n \setminus \bigcup_{i \in J_n} p_{i,n}) \geq c/2$$

for each $n \in \mathbb{N}$. Since $\text{diam}(F'_n \setminus \bigcup_{i \in J_n} p_{i,n}) \geq \text{diam}(\{0, f_n(\zeta_1)\})$, we have

$$\Delta(E'_n, F'_n \setminus \bigcup_{i \in J_n} p_{i,n}) \rightarrow \infty$$

as $n \rightarrow \infty$. Finally, we observe that the sets E'_n and $F'_n \setminus \bigcup_{i \in J_n} p_{i,n}$ are invariant under the set function $\pi_{X_n}^{-1} \circ \pi_{X_n}$ because they contain all complementary components of X_n that they intersect. By Lemma 3.2 there exists a constant $N'_0 \in \mathbb{N}$ that depends only on τ and a set $J'_n \subset \{1, \dots, n\}$ with $\#J'_n \leq N'_0$ such that

$$\text{Mod}_{X_n} \Gamma \left(\pi_{X_n}(E'_n), \pi_{X_n}(F'_n \setminus \bigcup_{i \in J_n} p_{i,n}); \mathcal{E}(X_n) \setminus \bigcup_{i \in J'_n} \pi_{X_n}(p_{i,n}) \right) \rightarrow 0$$

as $n \rightarrow \infty$. Note that by the monotonicity of modulus, if we set $J''_n = J_n \cup J'_n$, then we also obtain

$$\text{Mod}_{X_n} \Gamma \left(\pi_{X_n}(E'_n), \pi_{X_n}(F'_n \setminus \bigcup_{i \in J''_n} p_{i,n}); \mathcal{E}(X_n) \setminus \bigcup_{i \in J''_n} \pi_{X_n}(p_{i,n}) \right) \rightarrow 0$$

as $n \rightarrow \infty$.

Consider the set $F_n = (f_n^{-1})^*(F'_n)$; that is, F_n is the union of $f_n^{-1}(|\alpha_n| \cap X_n)$ together with the sets q_j , $j \in \{1, \dots, n\}$, that intersect its closure. Note that in the space $\mathcal{E}(Y; E)$ (endowed with any fixed metric that induces its topology) the projection of F_n has diameter uniformly bounded from below, since it connects the projections of the points ζ_0 and ζ_1 , which do not lie on E . Also, note that $\#J''_n \leq N_0 + N'_0$. Thus, by Lemma 3.1, we have

$$\text{Mod}_{Y_n} \Gamma \left(\pi_{Y_n}(E), \pi_{Y_n}(F_n \setminus \bigcup_{i \in J''_n} q_i); \mathcal{E}(Y_n) \setminus \bigcup_{i \in J''_n} \pi_{Y_n}(q_i) \right) \geq C$$

for all $n \in \mathbb{N}$. The conformal invariance of transboundary modulus (Lemma 2.13) leads to a contradiction. $\square$

Corollary 4.6. *For each $i \in \mathbb{N}$, the following statements are true.*

- (i) *If q_i is non-degenerate, then the sequence of sets $\{p_{i,n}\}_{n \geq i}$ does not degenerate to a point as $n \rightarrow \infty$.*
- (ii) *The set q_i is a point if and only if $p_{i,n}$ is a point for all $n \geq i$.*

The second part of the lemma follows from the fact that f_n is a conformal map. After taking a diagonal sequence, assume that for each $i \in \mathbb{N}$ the sequence $p_{i,n}$ converges in the Hausdorff sense as $n \rightarrow \infty$ to a compact set p_i that is connected, does not separate the sphere by the assumption in Theorem 4.1, and is τ -fat by Lemma 2.4. We have already established parts (i) and (iii) of Theorem 4.1.

Next, we wish to show that the sets p_i , $i \in \mathbb{N}$, are pairwise disjoint. First we establish a preliminary lemma.

Lemma 4.7 (No clustering). *There exists $N(\tau) > 0$ such that for each $x \in \widehat{\mathbb{C}}$ there are at most $N(\tau)$ non-degenerate sets p_i , $i \in \mathbb{N}$, containing x .*

Proof. Let $N \in \mathbb{N}$ and suppose that the sets $p_{i_1}, \dots, p_{i_N}$ are non-degenerate and contain x . Fix $r > 0$ be such that p_{i_l} is not contained in $B(x, r)$ for each $l \in \{1, \dots, N\}$ and consider points $x_{i_l, n} \in p_{i_l, n}$ converging to x as $n \rightarrow \infty$. For fixed $\delta \in (0, r)$ note that the sets $B(x_{i_l, n}, r - \delta) \cap p_{i_l, n}$, $l \in \{1, \dots, N\}$, are pairwise disjoint and are contained in $B(x, r)$ for all sufficiently large $n \in \mathbb{N}$. Moreover, $p_{i_l, n}$ is not contained in $B(x_{i_l, n}, r - \delta)$ for all sufficiently large $n \in \mathbb{N}$. By the τ -fatness of $p_{i_l, n}$, for large $n \in \mathbb{N}$ we have

$$\Sigma(B(x, r)) \geq \sum_{l=1}^N \Sigma(B(x_{i_l, n}, r - \delta) \cap p_{i_l, n}) \geq N\tau(r - \delta)^2.$$

Letting $\delta \rightarrow 0$, gives $N \leq \tau^{-1}\Sigma(B(x, r))r^{-2} \leq C\tau^{-1}$. $\square$

Lemma 4.8 (No collisions). *Let $\widehat{F}_1, \widehat{F}_2 \subset \mathcal{E}(Y)$ be disjoint continua and let $F_1 = \pi_Y^{-1}(\widehat{F}_1)$, $F_2 = \pi_Y^{-1}(\widehat{F}_2)$. Then*

$$\liminf_{n \rightarrow \infty} \text{dist}(f_n^*(F_1), f_n^*(F_2)) > 0.$$

Proof. We may assume that $\widehat{F}_1$ and $\widehat{F}_2$ are, in addition, non-separating continua, after replacing them if necessary with larger continua that are disjoint and non-separating. Suppose that $\text{dist}(f_n^*(F_1), f_n^*(F_2)) \rightarrow 0$ along a subsequence. After passing to a subsequence, we assume that $f_n^*(F_i)$ converges in the Hausdorff sense to a continuum F_i^* , $i = 1, 2$, and $F_1^* \cap F_2^* \neq \emptyset$. Note that the sets F_i , $i = 1, 2$, contain all of the peripheral continua of Y that they intersect. We consider two cases.

Case 1. The set q_i is non-degenerate for only finitely many $i \in \mathbb{N}$. Let $J \subset \mathbb{N}$ be the set of those indices. The conformal map $f_n: Y_n \rightarrow X_n$ extends conformally to the isolated point q_i whenever $i \in \{1, \dots, n\} \setminus J$. Since F_1 and F_2 are disjoint, non-separating, and contain all peripheral continua that they intersect, the open set $\Omega = \widehat{\mathbb{C}} \setminus (F_1 \cup F_2 \cup \bigcup_{i \in J} q_i)$ is connected and the sets F_1, F_2 are complementary components of Ω . The three-point normalization of f_n implies that, after passing to a further subsequence, f_n converges locally uniformly in $\widehat{\mathbb{C}} \setminus (F_1 \cup F_2 \cup \bigcup_{i \in J} q_i)$ to a map f that is either constant or conformal. By Lemma 4.5, f is non-constant, so it is conformal. Lemma 2.14 implies that $f^*(F_i) \supset F_i^*$ for $i = 1, 2$. This leads to a contradiction, since the sets $f^*(F_1), f^*(F_2)$ are disjoint, but the sets F_1^*, F_2^* are not.

Case 2. There are infinitely many non-degenerate sets q_i , $i \in \mathbb{N}$. By Corollary 4.6, p_i is non-degenerate if and only if q_i has this property. Let $x \in F_1^* \cap F_2^*$. By Lemma 4.7, there exists $N(\tau) > 0$ such that at most $N(\tau)$ non-degenerate sets p_i , $i \in \mathbb{N}$, contain the point x . Thus, there exists $i_0 \in \mathbb{N}$ such that $E = q_{i_0}$ is non-degenerate and $x \notin p_{i_0}$. The sets $E'_n = f_n^*(E)$ converge to p_{i_0} so they have diameters uniformly bounded away from 0 and they stay away from the point x . Let α_n be a curve in $\widehat{\mathbb{C}}$ joining $f_n^*(F_1)$ and $f_n^*(F_2)$ such that $|\alpha_n|$ converges to x in the Hausdorff sense. Let G'_n be the union of $|\alpha_n|$ together with the sets $p_{j, n}$, $j \in \{1, \dots, n\}$, that intersect $|\alpha_n|$. Note that the diameter of G'_n might be large.

We now argue as in the proof of Lemma 4.5. The cofatness of X_n and Lemma 2.5 imply that if J_n is the set of indices j such that $p_{j, n} \cap |\alpha_n| \neq \emptyset$ and $\text{diam}(p_{j, n}) \geq \text{diam}(|\alpha_n|)$, then $\#J_n$ is uniformly bounded, depending only on τ . We conclude that $\Delta(E'_n, G'_n \setminus \bigcup_{i \in J_n} p_{i, n}) \rightarrow \infty$ as $n \rightarrow \infty$. Lemma 3.2 implies that there exists

a set of natural numbers $J''_n \supset J_n$ with uniformly bounded cardinality such that

$$\text{Mod}_{X_n} \Gamma \left(\pi_{X_n}(E'_n), \pi_{X_n}(G'_n \setminus \bigcup_{i \in J''_n} p_{i,n}); \mathcal{E}(X_n) \setminus \bigcup_{i \in J''_n} \pi_{X_n}(p_{i,n}) \right) \rightarrow 0$$

as $n \rightarrow \infty$.

On the other hand, we consider the sets E and $G_n = (f_n^{-1})^*(G'_n)$. We note that the continuum G_n joins F_1 and F_2 , so its projection to $\mathcal{E}(Y; E)$ has diameter uniformly bounded below away from 0, in any given metric. By Lemma 3.1 we have

$$\text{Mod}_{Y_n} \Gamma \left(\pi_{Y_n}(E), \pi_{Y_n}(G_n \setminus \bigcup_{i \in J''_n} q_i); \mathcal{E}(Y_n) \setminus \bigcup_{i \in J''_n} \pi_{Y_n}(q_i) \right) \geq C$$

for each $n \in \mathbb{N}$. This contradicts the conformal invariance of transboundary modulus. $\square$

Corollary 4.9. *The sets p_i , $i \in \mathbb{N}$, are pairwise disjoint. Moreover, the set $X = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^{\infty} p_i$ is a Sierpiński packing.*

Proof. The sets p_i , $i \in \mathbb{N}$, are pairwise disjoint by Lemma 4.8. As we have discussed, each of the sets p_i , $i \in \mathbb{N}$, is a non-separating continuum and is τ -fat. A trivial consequence of Lemma 2.5 is that $\text{diam}(p_i) \rightarrow 0$ as $i \rightarrow \infty$. These facts imply that X is a Sierpiński packing. $\square$

Thus, we have established part (ii) of Theorem 4.1. In the following sections we prove the existence of the limiting map of part (iv) and its regularity.

4.2. Existence of limiting map. Fix metrics on the spaces $\mathcal{E}(X), \mathcal{E}(Y)$ that induce their topology. The particular metrics are not of importance. We set $g_n = f_n^{-1}$ and consider the sequence of set functions

$$h_n = \pi_Y \circ g_n^* \circ \pi_X^{-1} = \pi_Y \circ \pi_{Y_n}^{-1} \circ \widehat{g}_n \circ \pi_{X_n} \circ \pi_X^{-1}, \quad n \in \mathbb{N}$$

from subsets of $\mathcal{E}(X)$ to subsets of $\mathcal{E}(Y)$.

Lemma 4.10 (Equicontinuity). *For each $\varepsilon > 0$ there exists $\delta > 0$ and $N \in \mathbb{N}$ such that if $\widehat{E}'$ is a set in $\mathcal{E}(X)$ with $\text{diam}(\widehat{E}') < \delta$, then $\text{diam}(h_n(\widehat{E}')) < \varepsilon$ for all $n > N$.*

Roughly speaking, this lemma says that if a set has small diameter in $\mathcal{E}(X)$, then its image under g_n^* projects to a set that has small diameter in $\mathcal{E}(Y)$. We note that the diameter need not be small if we do not project to $\mathcal{E}(Y)$. As an analogy, consider a conformal map f from a simply connected domain $\mathcal{Y}$ whose boundary is not locally connected onto the unit disk $\mathcal{X}$. Then small sets near $\partial\mathcal{X}$ need not be mapped to small sets in $\mathcal{Y}$ under $g = f^{-1}$.

Proof. By the local connectivity of $\mathcal{E}(X) \simeq \widehat{\mathbb{C}}$, it suffices to show the statement for continua $\widehat{E}'$, rather than arbitrary sets. We argue by contradiction, assuming that there exists $\varepsilon_0 > 0$ and a sequence of positive integers $\{k_n\}_{n \in \mathbb{N}}$ with $k_n \rightarrow \infty$ such that there exists a sequence of continua $\widehat{E}'_n \subset \mathcal{E}(X)$ converging to a point $\widehat{x} \in \mathcal{E}(X)$ as $n \rightarrow \infty$, but $\text{diam}(\pi_Y(E_n)) \geq \varepsilon_0$ for all $n \in \mathbb{N}$, where $E_n = g_{k_n}^*(\pi_X^{-1}(\widehat{E}'_n))$. After passing to a subsequence, we assume that E_n converges to a continuum $E \subset \widehat{\mathbb{C}}$, and by continuity $\pi_Y(E_n)$ converges to a continuum $\pi_Y(E)$ in $\mathcal{E}(Y)$ with $\text{diam}(\pi_Y(E)) \geq \varepsilon_0$.

Consider distinct points $\hat{y}, \hat{z} \in \pi_Y(E) \setminus \bigcup_{i \in \mathbb{N}} \pi_Y(q_i)$ and let $y = \pi_Y^{-1}(\hat{y})$, $z = \pi_Y^{-1}(\hat{z}) \in E \cap Y$. Consider disjoint closed Jordan regions U_y, U_z containing y, z in their interior, whose boundaries lie in Y ; these regions can be obtained as preimages under π_Y of appropriate Jordan regions in $\mathcal{E}(Y)$. By Lemma 4.8, the distance of the images $f_{k_n}^*(U_y), f_{k_n}^*(U_z)$ is uniformly bounded away from zero. By the Hausdorff convergence of E_n to E , for all sufficiently large $n \in \mathbb{N}$ the set E_n intersects both U_y and U_z . Since $y, z \in Y$, there exist points $y_n, z_n \in E_n \cap Y$ converging to y, z , respectively. The points $y'_n = f_{k_n}(y_n), z'_n = f_{k_n}(z_n)$ lie in $\pi_X^{-1}(\hat{E}'_n)$ and stay away from each other as $n \rightarrow \infty$. Since the points $\pi_X(y'_n)$ and $\pi_X(z'_n)$ converge to the point $\hat{x}$, but the points y'_n and z'_n stay away from each other, by Lemma 2.2 we must have $\hat{x} = \pi_X(p_{i_0})$ for some $i_0 \in \mathbb{N}$. We conclude that y'_n, z'_n accumulate at p_{i_0} as $n \rightarrow \infty$.

Fix a closed Jordan region U containing y in its interior such that U is disjoint from q_{i_0} and $\partial U \subset Y$; this is possible because $y \in Y$. We have $y_n \in U$ for all sufficiently large $n \in \mathbb{N}$. On the other hand, the sets $f_{k_n}^*(U)$ and $f_{k_n}^*(q_{i_0}) = p_{i_0, k_n}$ come arbitrarily close to each other. This contradicts Lemma 4.8. $\square$

The following statement is a version of the Arzelà–Ascoli theorem for equicontinuous families of set functions. The proof is a straightforward adaptation of the classical argument and the experienced reader may safely skip it.

Lemma 4.11 (Compactness). *Let $\mathcal{X}, \mathcal{Y}$ be compact metric spaces and $h_n: \mathcal{P}(\mathcal{X}) \rightarrow \mathcal{P}(\mathcal{Y})$, $n \in \mathbb{N}$, be a sequence of set functions with the following properties:*

- (i) *Surjectivity: for each $n \in \mathbb{N}$, $h_n(\mathcal{X}) = \mathcal{Y}$.*
- (ii) *Setwise monotonicity: for each $n \in \mathbb{N}$, if $A, B \subset \mathcal{X}$ and $A \subset B$, then $h_n(A) \subset h_n(B)$.*
- (iii) *Inverse image property: for each $n \in \mathbb{N}$, if $A \subset \mathcal{X}$ and $y \in h_n(A)$, then there exists $x \in A$ such that $y \in h_n(x)$.*
- (iv) *Equicontinuity: for each $\varepsilon > 0$ there exists $\delta > 0$ and $N \in \mathbb{N}$ such that for each set $E \subset \mathcal{X}$ with $\text{diam}(E) < \delta$ we have $\text{diam}(h_n(E)) < \varepsilon$ for all $n > N$.*

Then there exists a subsequence $\{h_{k_n}\}_{n \in \mathbb{N}}$ of $\{h_n\}_{n \in \mathbb{N}}$ that converges uniformly to a continuous and surjective map $h: \mathcal{X} \rightarrow \mathcal{Y}$ in the following sense: for each $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that for each set $E \subset \mathcal{X}$ we have

$$d_H(h_{k_n}(E), h(E)) < \varepsilon$$

for all $n > N$.

Proof. Note that the space of compact subsets of a compact metric space is compact in the Hausdorff metric [BBI01, Theorem 7.3.8, p 253]. Consider a countable dense set $\{x_l\}_{l \in \mathbb{N}}$ in $\mathcal{X}$ and a diagonal subsequence h_{k_n} such that for each $l \in \mathbb{N}$ the sequence of sets $h_{k_n}(x_l)$, $n \in \mathbb{N}$, converges, and thus is a Cauchy sequence in the Hausdorff metric. For simplicity, we denote h_{k_n} by h_n .

We fix $\varepsilon > 0$. For each $l \in \mathbb{N}$ and $\varepsilon > 0$ there exists $N(\varepsilon, l) > 0$ such that for all $n, m > N(\varepsilon, l)$ we have

$$d_H(h_n(y_l), h_m(y_l)) < \varepsilon/3.$$

By equicontinuity, there exists $\delta > 0$ and $N \in \mathbb{N}$ such that if $E \subset \mathcal{X}$ is a set with $\text{diam}(E) < \delta$, then $\text{diam}(h_n(E)) < \varepsilon/3$ for $n > N$. By the compactness of $\mathcal{X}$, we may find $M \in \mathbb{N}$ such that every point $x \in \mathcal{X}$ is within distance δ from the set $\{x_1, \dots, x_M\}$. We define $N_0 = \max\{N, N(\varepsilon, 1), \dots, N(\varepsilon, M)\}$.

Let $x \in \mathcal{X}$ be arbitrary and consider $l \in \{1, \dots, M\}$ such that $d(x, x_l) < \delta$. Let $E = \{x, x_l\}$, so $\text{diam}(\mathfrak{h}_n(E)) < \varepsilon/3$ for all $n > N$. For $n, m > N_0$ we now have

$$\begin{aligned} d_H(\mathfrak{h}_n(x), \mathfrak{h}_m(x)) &\leq d_H(\mathfrak{h}_n(x), \mathfrak{h}_n(x_l)) + d_H(\mathfrak{h}_n(x_l), \mathfrak{h}_m(x_l)) + d_H(\mathfrak{h}_m(x_l), \mathfrak{h}_m(x)) \\ &\leq \text{diam}(\mathfrak{h}_n(E)) + \varepsilon/3 + \text{diam}(\mathfrak{h}_m(E)) < \varepsilon. \end{aligned}$$

Here we used the monotonicity property of $\mathfrak{h}_n$, which implies that $\mathfrak{h}_n(x) \cup \mathfrak{h}_n(x_l) \subset \mathfrak{h}_n(E)$. Since the space of compact subsets of $\mathcal{X}$ is complete with the Hausdorff metric, we conclude that $\mathfrak{h}_n(x)$ (as well as, its closure) converges in the Hausdorff sense to a compact subset of $\mathcal{X}$. By the equicontinuity condition, we have $\text{diam}(\mathfrak{h}_n(x)) \rightarrow 0$, so the limit has to be a point $\mathfrak{h}(x)$.

Summarizing, we have shown that for each $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that for each $x \in \mathcal{X}$ we have $d_H(\mathfrak{h}_n(x), \mathfrak{h}(x)) < \varepsilon$ for all $n > N$. The equicontinuity of $\mathfrak{h}_n$ also implies that $\mathfrak{h}$ is continuous.

For the surjectivity, we use the surjectivity of $\mathfrak{h}_n$ and the inverse image property. Note that for each $y \in \mathcal{Y} = \mathfrak{h}_n(\mathcal{X})$ there exists $\tilde{x}_n \in \mathcal{X}$ such that $y \in \mathfrak{h}_n(\tilde{x}_n)$. By the uniform convergence, we have $d_H(\mathfrak{h}_n(\tilde{x}_n), \mathfrak{h}(\tilde{x}_n)) \rightarrow 0$ as $n \rightarrow \infty$. After passing to a subsequence, we assume that $\tilde{x}_n \rightarrow x \in \mathcal{X}$. Thus, $d_H(\mathfrak{h}_n(\tilde{x}_n), \mathfrak{h}(x)) \rightarrow 0$. Since $y \in \mathfrak{h}_n(\tilde{x}_n)$, we have $\mathfrak{h}(x) = y$.

Finally, we prove the uniform convergence for images of sets as in the end of the statement of the lemma. Let $\varepsilon > 0$ and $N \in \mathbb{N}$ be such that for each $x \in \mathcal{X}$ we have $d_H(\mathfrak{h}_n(x), \mathfrak{h}(x)) < \varepsilon$ for all $n > N$. Let $E \subset \mathcal{X}$ be any set and fix $n > N$. We will show that $d_H(\mathfrak{h}_n(E), \mathfrak{h}(E)) < \varepsilon$. First, we show that $\mathfrak{h}_n(E) \subset N_\varepsilon(\mathfrak{h}(E))$. Let $y \in \mathfrak{h}_n(E)$, so by the inverse image property there exists $\tilde{x}_n \in E$ such that $y \in \mathfrak{h}_n(\tilde{x}_n) \subset B(\mathfrak{h}(\tilde{x}_n), \varepsilon) \subset N_\varepsilon(\mathfrak{h}(E))$. Conversely, we will show that $\mathfrak{h}(E) \subset N_\varepsilon(\mathfrak{h}_n(E))$. Let $y \in \mathfrak{h}(E)$. By the surjectivity of $\mathfrak{h}$, there exists $x \in E$ such that $y = \mathfrak{h}(x) \in N_\varepsilon(\mathfrak{h}_n(x)) \subset N_\varepsilon(\mathfrak{h}_n(E))$; here we also used the setwise monotonicity. $\square$

Lemma 4.12. *The sequence of set functions $h_n = \pi_Y \circ g_n^* \circ \pi_X^{-1}$, $n \in \mathbb{N}$, satisfies the assumptions of Lemma 4.11. In particular, after passing to a subsequence, h_n converges uniformly as $n \rightarrow \infty$ to a continuous and surjective map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$.*

Proof. Recall that $h_n = \pi_Y \circ \pi_{Y_n}^{-1} \circ \hat{g}_n \circ \pi_{X_n} \circ \pi_X^{-1}$. The surjectivity is immediate. The equicontinuity follows from Lemma 4.10. In general, if ϕ is a function between any sets, then the induced set functions ϕ and ϕ^{-1} have the monotonicity and inverse image properties. Also, note that if ϕ and ψ are set functions that have the monotonicity and inverse image properties, then $\phi \circ \psi$ also does so. This shows that h_n has these properties. $\square$

Lemma 4.13 (Topological properties). *The map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ is monotone and $h(\hat{p}_i) = \hat{q}_i$ for each $i \in \mathbb{N}$.*

Proof. For the monotonicity we argue by contradiction. Let $\hat{y} \in \mathcal{E}(Y)$ and suppose that $h^{-1}(\hat{y})$ is disconnected. Let $\hat{x}, \hat{z}$ be points lying on distinct components of $h^{-1}(\hat{y})$ and let $\hat{E}'$ be a continuum in $\mathcal{E}(X) \setminus h^{-1}(\hat{y})$ that separates them. Consider the continuum $E' = \pi_X^{-1}(\hat{E}')$, which separates the sets $x = \pi_X^{-1}(\hat{x})$ and $z = \pi_X^{-1}(\hat{z})$.

In $\mathcal{E}(Y)$ consider the continuum $\hat{E} = h(\hat{E}')$, which is disjoint from $\hat{y}$. By the uniform convergence, the sets $h_n(\hat{E}')$ converge to $\hat{E}$ in the Hausdorff sense, so there exists a closed Jordan region $\hat{U}$, disjoint from $\hat{y}$, containing $h_n(\hat{E}')$ for all

sufficiently large $n \in \mathbb{N}$. Since $h_n(\hat{x})$ and $h_n(\hat{z})$ converge to $\hat{y}$, there exists a closed Jordan region $\hat{V}$, disjoint from $\hat{U}$, containing these sequences for all large $n \in \mathbb{N}$.

Let U, V be the preimages of $\hat{U}, \hat{V}$ under π_Y , respectively. So, U, V are disjoint continua in $\hat{\mathbb{C}}$. Using (2.1), we have

$$f_n^*(U) \supset f_n^*(\pi_Y^{-1}(h_n(\hat{E}')))) = f_n^*(\pi_Y^{-1}(\pi_Y(g_n^*(E')))) \supset f_n^*(g_n^*(E')) \supset E'$$

and similarly $f_n^*(V)$ contains x, z for all large $n \in \mathbb{N}$. Since E' separates x, z and $f_n^*(V)$ joins them, we conclude that $f_n^*(U) \cap f_n^*(V) \neq \emptyset$ for all large $n \in \mathbb{N}$. However, by Lemma 4.8, $f_n^*(U)$ and $f_n^*(V)$ have distance uniformly bounded below away from 0 as $n \rightarrow \infty$. This is a contradiction. Therefore $h^{-1}(\hat{y})$ is connected.

Next we show that $h(\hat{p}_i) = \hat{q}_i$ for each $i \in \mathbb{N}$. Recall that $p_{i,n} \rightarrow p_i$ as $n \rightarrow \infty$. This implies that the sequence of sets $E_n = \pi_X(p_{i,n})$ converges to the point $\pi_X(p_i) = \hat{p}_i$. By uniform convergence, $h(\hat{p}_i)$ is precisely the limit of $h_n(E_n)$. Note that $\pi_X^{-1}(E_n)$ contains $p_{i,n}$. Thus, $g_n^*(\pi_X^{-1}(E_n))$ contains q_i and $h_n(E_n)$ contains $\hat{q}_i$ for all sufficiently large $n \in \mathbb{N}$. It follows that the limit of $h_n(E_n)$, which is the point $h(\hat{p}_i)$, is precisely equal to $\hat{q}_i$. $\square$

This completes the proof of the existence and of the topological properties of the limiting map $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$. In the next section we establish the analytic properties of h as required in the definition of a packing-conformal map, completing the proof of Theorem 4.1.

4.3. Regularity of limiting map. For each conformal map $g_n: X_n \rightarrow Y_n$ consider the derivative (with respect to the spherical metrics) $|Dg_n|$, which satisfies the following relations by Lemma 2.17. First, we have the transboundary upper gradient inequality:

$$(4.1) \quad \text{dist}(g_n^*(\gamma(a)), g_n^*(\gamma(b))) \leq \int_{\gamma} |Dg_n| ds + \sum_{i: p_{i,n} \cap |\gamma| \neq \emptyset} \text{diam}(q_i)$$

for all rectifiable paths $\gamma: [a, b] \rightarrow \hat{\mathbb{C}}$. Second, for all Borel sets $E \subset Y_n$ we have

$$(4.2) \quad \int_{g_n^{-1}(E)} |Dg_n|^2 d\Sigma = \Sigma(E).$$

We extend $|Dg_n|$ to $\hat{\mathbb{C}}$ by setting it to be zero in $\bigcup_{i=1}^n p_{i,n}$. We first establish a preliminary result.

Lemma 4.14. *Let $\{\lambda_i\}_{i \in \mathbb{N}}$ be a non-negative sequence in $\ell^2(\mathbb{N})$. Then*

$$\limsup_{n \rightarrow \infty} \sum_{i: p_{i,n} \cap |\gamma| \neq \emptyset} \lambda_i \leq \sum_{i: p_i \cap |\gamma| \neq \emptyset} \lambda_i.$$

for all compact curves γ in $\hat{\mathbb{C}}$ outside a curve family Γ_0 with $\text{Mod}_2 \Gamma_0 = 0$.

The proof relies on the fact that for each $n \in \mathbb{N}$ the sets $p_{i,n}$, $i \in \{1, \dots, n\}$, are pairwise disjoint and τ -fat, and they converge in the Hausdorff sense as $n \rightarrow \infty$ to the sets p_i , $i \in \mathbb{N}$, which are also pairwise disjoint and τ -fat.

Proof. Suppose that γ is a compact path so $|\gamma|$ is compact. For each $i \in \mathbb{N}$, if $p_i \cap |\gamma| = \emptyset$, then p_i has a positive distance from $|\gamma|$ by compactness. Thus,

$p_{i,n} \cap |\gamma| = \emptyset$ for all sufficiently large $n \in \mathbb{N}$. It follows that for each $M \in \mathbb{N}$ we have

$$(4.3) \quad \limsup_{n \rightarrow \infty} \sum_{\substack{i \in \{1, \dots, M\} \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \lambda_i \leq \sum_{i: p_i \cap |\gamma| \neq \emptyset} \lambda_i.$$

We will show that there exists a curve family Γ_0 with $\text{Mod}_2 \Gamma_0 = 0$ such that if γ is path outside Γ_0 , then

$$(4.4) \quad \lim_{M, n \rightarrow \infty} \sum_{\substack{i > M \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \lambda_i = 0.$$

Combined with (4.3), this gives the desired conclusion.

Let J be the set of indices $i \in \mathbb{N}$ such that $\text{diam}(p_i) > 0$. By Corollary 4.6, if $i \in J$ then $p_{i,n}$ is non-degenerate for all $n \geq i$, and if $i \notin J$ then $p_{i,n}$ is a point for all $n \geq i$. For each $n \in \mathbb{N}$ and $i \in \{1, \dots, n\} \cap J$ consider points $x_{i,n} \in p_{i,n}$ and $x_i \in p_i$ such that $x_{i,n} \rightarrow x_i$ as $n \rightarrow \infty$. We let $B_{i,n} = B(x_{i,n}, \text{diam}(p_{i,n}))$ and $B_i = B(x_i, \text{diam}(p_i))$. Note that $\Sigma(2B_{i,n}) \leq c(\tau)\Sigma(p_{i,n})$ and $\Sigma(2B_i) \leq c(\tau)\Sigma(p_i)$ by the τ -fatness, and observe that for each $i \in J$ the characteristic functions $\chi_{2B_{i,n}}$ converge as $n \rightarrow \infty$ pointwise a.e. to χ_{2B_i} . We define the functions

$$\begin{aligned} \phi(x) &= \sum_{i \in J} \frac{\lambda_i}{\text{diam}(p_i)} \chi_{2B_i}(x), \\ \psi_{i,n}(x) &= \left| \frac{\lambda_i \chi_{\{1, \dots, n\}}(i)}{\text{diam}(p_{i,n})} \chi_{2B_{i,n}}(x) - \frac{\lambda_i}{\text{diam}(p_i)} \chi_{2B_i}(x) \right|, \quad i \in J, \quad n \in \mathbb{N}, \\ \phi_n(x) &= \sum_{i \in J} \psi_{i,n}(x), \quad n \in \mathbb{N}. \end{aligned}$$

Using Lemma 2.7 and the fact that the sets p_i , $i \in J$, are pairwise disjoint and τ -fat, we obtain

$$\int \phi^2 \leq c(\tau) \int \sum_{i \in J} \frac{\lambda_i^2}{\text{diam}(p_i)^2} \chi_{p_i} \leq c'(\tau) \sum_{i=1}^{\infty} \lambda_i^2$$

and similarly, for each $n, M \in \mathbb{N}$ we have

$$(4.5) \quad \int \left(\sum_{i > M, i \in J} \psi_{i,n}(x) \right)^2 \leq c(\tau) \sum_{i > M} \lambda_i^2.$$

Note that for each $i \in J$ the sequence of functions $\{\psi_{i,n}\}_{n \in \mathbb{N}}$ is uniformly bounded and converges to 0 pointwise a.e. The dominated convergence theorem implies that for each fixed $M \in \mathbb{N}$, the sequence $\{\sum_{i \leq M, i \in J} \psi_{i,n}\}_{n \in \mathbb{N}}$ converges to 0 in $L^2(\widehat{\mathbb{C}})$.

This fact, combined with (4.5) gives that $\{\phi_n\}_{n \in \mathbb{N}}$ converges to 0 in $L^2(\widehat{\mathbb{C}})$.

By Fuglede's lemma [HKST15, p. 131], there exists a curve family Γ_1 with $\text{Mod}_2 \Gamma_1 = 0$ such that for all curves γ outside Γ_1 we have

$$\lim_{n \rightarrow \infty} \int_{\gamma} \phi_n ds = 0.$$

Moreover, since $\phi \in L^2(\widehat{\mathbb{C}})$, there exists a curve family Γ_2 with $\text{Mod}_2 \Gamma_2 = 0$ such that $\int_{\gamma} \phi ds < \infty$ for each $\gamma \notin \Gamma_2$. Finally, there exists a family Γ_3 with $\text{Mod}_2 \Gamma_3 = 0$ that contains all non-constant curves intersecting the countable collection of points

$p_i, i \notin J$, and $p_{i,n}, i \notin J, n \geq i$; see [Väi71, §7.9, p. 23]. We define $\Gamma_0 = \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$, which satisfies $\text{Mod}_2 \Gamma_0 = 0$, and fix a curve $\gamma \notin \Gamma_0$. Note that if γ is a constant curve, then the claim (4.4) holds trivially, since the sum contains at most one term and $\lambda_i \rightarrow 0$ as $i \rightarrow \infty$. Thus, we assume that γ is non-constant.

Since $\gamma \notin \Gamma_1$, for each $\varepsilon > 0$ there exists $N_0 \in \mathbb{N}$ such that

$$(4.6) \quad \int_{\gamma} \sum_{i \in J} \left| \frac{\lambda_i \chi_{\{1, \dots, n\}}(i)}{\text{diam}(p_{i,n})} \chi_{2B_{i,n}}(x) - \frac{\lambda_i}{\text{diam}(p_i)} \chi_{2B_i} \right| ds < \varepsilon$$

for all $n > N_0$. By Lemma 2.5, there exists a number $N_1 \in \mathbb{N}$, depending only on τ , such that for each $n \in \mathbb{N}$ there exists a set $I_n \subset \mathbb{N}$ with $\#I_n \leq N_1$ that contains precisely the indices $i \in \mathbb{N}$ with the property that $p_{i,n} \cap |\gamma| \neq \emptyset$ and $\text{diam}(2B_{i,n}) \geq \text{diam}(|\gamma|)$. Hence, $\text{diam}(|\gamma|) > \text{diam}(2B_{i,n})$ for each $i \in \{1, \dots, n\} \cap J \setminus I_n$ with $p_{i,n} \cap |\gamma| \neq \emptyset$. By the properties of the set I_n , if $p_{i,n} \cap |\gamma| \neq \emptyset$ and $i \in \{1, \dots, n\} \cap J \setminus I_n$, then

$$\lambda_i \leq \int_{\gamma} \frac{\lambda_i}{\text{diam}(p_{i,n})} \chi_{2B_{i,n}} ds.$$

Thus, for each $M \in \mathbb{N}$ and $n > N_0$, using (4.6), we have

$$\begin{aligned} \sum_{\substack{i > M, i \in J \setminus I_n \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \lambda_i &\leq \int_{\gamma} \sum_{\substack{i > M, i \in J \setminus I_n \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \frac{\lambda_i \chi_{\{1, \dots, n\}}(i)}{\text{diam}(p_{i,n})} \chi_{2B_{i,n}} ds \\ &< \varepsilon + \int_{\gamma} \sum_{i > M, i \in J} \frac{\lambda_i}{\text{diam}(p_i)} \chi_{2B_i} ds. \end{aligned}$$

Since $\gamma \notin \Gamma_2$, we have $\int_{\gamma} \phi ds < \infty$; thus, by the dominated convergence theorem, for all sufficiently large $M \in \mathbb{N}$ the latter line integral term in the above inequalities is less than ε . Therefore, for all sufficiently large $M \in \mathbb{N}$ and for $n > N_0$ we have

$$\sum_{\substack{i > M, i \in J \setminus I_n \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \lambda_i < 2\varepsilon.$$

Since $\lambda_i \rightarrow 0$ as $i \rightarrow \infty$, we also have

$$\sum_{i > M, i \in I_n} \lambda_i \leq N_1 \cdot \max\{\lambda_i : i > M\} < \varepsilon$$

for all sufficiently large $M \in \mathbb{N}$. Altogether,

$$\sum_{\substack{i > M, i \in J \\ p_{i,n} \cap |\gamma| \neq \emptyset}} \lambda_i < 3\varepsilon$$

for all sufficiently large $M \in \mathbb{N}$ and $n > N_0$. Finally, since $\gamma \notin \Gamma_3$, we have $i \in J$ whenever $p_{i,n} \cap |\gamma| \neq \emptyset$. This implies that we may remove the restriction $i \in J$ in the summation range of the latter sum. This completes the proof of (4.4). $\square$

Lemma 4.15 (Upper gradient). *The sequence $\{|Dg_n|\}_{n \in \mathbb{N}}$ has a subsequence that converges weakly in $L^2(\widehat{\mathbb{C}})$ to a function ρ_h with the property that*

$$\text{dist}(\pi_Y^{-1} \circ h \circ \pi_X(\gamma(a)), \pi_Y^{-1} \circ h \circ \pi_X(\gamma(b))) \leq \int_{\gamma} \rho_h ds + \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i)$$

for all curves $\gamma: [a, b] \rightarrow \widehat{\mathbb{C}}$ outside a curve family Γ_0 with $\text{Mod}_2 \Gamma_0 = 0$.

Proof. Recall that $h_n = \pi_Y \circ g_n^* \circ \pi_X^{-1}$, so for each set $A \subset \widehat{\mathbb{C}}$ we have

$$(4.7) \quad \pi_Y^{-1} \circ h_n \circ \pi_X(A) \supset g_n^*(A).$$

Let $\gamma: [a, b] \rightarrow \widehat{\mathbb{C}}$ be an arbitrary rectifiable path. If we set $\alpha = \pi_X(\gamma(a))$ and $\beta = \pi_X(\gamma(b))$, the transboundary upper gradient inequality of g_n , as stated in (4.1), and (4.7) imply that

$$(4.8) \quad \text{dist}(\pi_Y^{-1} \circ h_n(\alpha), \pi_Y^{-1} \circ h_n(\beta)) \leq \int_{\gamma} |Dg_n| ds + \sum_{i: p_{i,n} \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

Our goal is to show that we can take limits in this expression and derive the claimed upper gradient inequality of h for all curves γ outside an exceptional family Γ_0 with $\text{Mod}_2 \Gamma_0 = 0$.

First, we treat the line integral terms. By (4.2), the sequence $|Dg_n|$ is uniformly bounded in $L^2(\widehat{\mathbb{C}})$. Consider a weak limit $\rho_h \in L^2(\widehat{\mathbb{C}})$ of $|Dg_n|$, given by the Banach–Alaoglu theorem [HKST15, Theorem 2.4.1]. By Mazur’s lemma [HKST15, p. 19], there exist convex combinations of $|Dg_n|$ that converge strongly in $L^2(\widehat{\mathbb{C}})$ to ρ_h . Specifically, these convex combinations have the form

$$\rho_n = \sum_{i=n}^{M_n} \lambda_{i,n} |Dg_i|$$

for some $M_n > n$ and $0 \leq \lambda_{i,n} \leq 1$, $i \in \{n, \dots, M_n\}$, where $\sum_{i=n}^{M_n} \lambda_{i,n} = 1$. By Fuglede’s lemma [HKST15, p. 131], there exists a curve family Γ_1 with $\text{Mod}_2 \Gamma_1 = 0$ such that for all curves γ outside Γ_1 we have

$$(4.9) \quad \lim_{n \rightarrow \infty} \int_{\gamma} \rho_n ds = \int_{\gamma} \rho_h ds.$$

Next, we treat the left-hand side of (4.8). Observe that as $n \rightarrow \infty$, the sets $h_n(\alpha)$ and $h_n(\beta)$ converge to the points $h(\alpha)$ and $h(\beta)$ respectively; recall Lemma 4.12. By Lemma 2.2 (ii), this implies that

$$\text{dist}(\pi_Y^{-1} \circ h(\alpha), \pi_Y^{-1} \circ h(\beta)) \leq \liminf_{n \rightarrow \infty} \text{dist}(\pi_Y^{-1} \circ h_n(\alpha), \pi_Y^{-1} \circ h_n(\beta)).$$

Note that if we set $d_n = \text{dist}(\pi_Y^{-1} \circ h_n(\alpha), \pi_Y^{-1} \circ h_n(\beta))$ and consider the convex combinations

$$\tilde{d}_n = \sum_{i=n}^{M_n} \lambda_{i,n} d_i$$

then we obtain immediately

$$(4.10) \quad \text{dist}(\pi_Y^{-1} \circ h(\alpha), \pi_Y^{-1} \circ h(\beta)) \leq \liminf_{n \rightarrow \infty} \tilde{d}_n.$$

Finally, we treat the summation term in (4.8). We apply Lemma 4.14 to conclude that there exists a curve family Γ_2 with $\text{Mod}_2 \Gamma_2 = 0$ such that for all compact curves γ outside Γ_2 we have

$$\limsup_{n \rightarrow \infty} \sum_{i: p_{i,n} \cap |\gamma| \neq \emptyset} \text{diam}(q_i) \leq \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

We set

$$s_n = \sum_{i: p_{i,n} \cap |\gamma| \neq \emptyset} \text{diam}(q_i) \quad \text{and} \quad \tilde{s}_n = \sum_{i=n}^{M_n} \lambda_{i,n} s_i$$

and observe that

$$(4.11) \quad \limsup_{n \rightarrow \infty} \tilde{s}_n \leq \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

We now define $\Gamma_0 = \Gamma_1 \cup \Gamma_2$ and let $\gamma \notin \Gamma_0$ be a compact curve. Taking convex combinations in (4.8), we obtain

$$\tilde{d}_n \leq \int_{\gamma} \rho_n ds + \tilde{s}_n.$$

By (4.9), (4.10), and (4.11), we can take limits to obtain

$$\text{dist}(\pi_Y^{-1} \circ h(\alpha), \pi_Y^{-1} \circ h(\beta)) \leq \int_{\gamma} \rho_h ds + \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

This completes the proof. $\square$

Lemma 4.16 (Conformality). *For all Borel sets $E \subset Y$ we have*

$$\int_{\pi_X^{-1}(h^{-1}(\pi_Y(E)))} \rho_h^2 d\Sigma \leq \Sigma(E).$$

Proof. Let $E \subset Y$ be a Borel set. Since π_Y is injective on Y , by the Lusin–Souslin theorem [Kec95, Theorem 15.1, p. 89] $\pi_Y(E)$ is also a Borel set. By continuity, $\pi_X^{-1}(h^{-1}(\pi_Y(E)))$ is a Borel set. Let K be a compact subset of $\pi_X^{-1}(h^{-1}(\pi_Y(E)))$. Since π_Y is injective on E , we have $\pi_Y^{-1}(h(\pi_X(K))) \subset E$. We will show that

$$\int_K \rho_h^2 d\Sigma \leq \Sigma(\pi_Y^{-1}(h(\pi_X(K)))) \leq \Sigma(E).$$

Since K is an arbitrary compact subset of $\pi_X^{-1}(h^{-1}(\pi_Y(E)))$, the proof will be completed by the inner regularity of Σ .

Let $\hat{K} = \pi_X(K)$. By Lemma 2.2 (i), for each $\delta > 0$, there exists an open neighborhood $\hat{V}$ of $h(\hat{K})$ such that the open set $\pi_Y^{-1}(\hat{V})$ contains $\pi_Y^{-1}(h(\hat{K}))$ and is contained in the open δ -neighborhood of $\pi_Y^{-1}(h(\hat{K}))$. In particular, by the compactness of $\pi_Y^{-1}(h(\hat{K}))$, for each $\varepsilon > 0$ we may find such an open set $\hat{V}$ with the additional property that

$$(4.12) \quad \Sigma(\pi_Y^{-1}(\hat{V})) \leq \Sigma(\pi_Y^{-1}(h(\hat{K}))) + \varepsilon.$$

Since $h_n(\hat{K})$ converges in the Hausdorff sense to $h(\hat{K})$, we have $h_n(\hat{K}) \subset \hat{V}$ for all sufficiently large $n \in \mathbb{N}$. This implies that $g_n^*(\pi_X^{-1}(\hat{K})) \subset \pi_Y^{-1}(\hat{V})$ for all sufficiently large $n \in \mathbb{N}$. This inclusion, the conformality of g_n , and (4.12), give

$$\begin{aligned} \int_K |Dg_n|^2 d\Sigma &= \int_{K \cap X_n} |Dg_n|^2 d\Sigma \leq \int_{\pi_X^{-1}(\hat{K}) \cap X_n} |Dg_n|^2 d\Sigma \\ &= \Sigma(g_n(\pi_X^{-1}(\hat{K}) \cap X_n)) \leq \Sigma(\pi_Y^{-1}(\hat{V})) \leq \Sigma(\pi_Y^{-1}(h(\hat{K}))) + \varepsilon \end{aligned}$$

for all sufficiently large $n \in \mathbb{N}$. Since $|Dg_n|$ converges weakly to ρ_h in $L^2(\widehat{\mathbb{C}})$, we see that $|Dg_n|\chi_K$ also converges weakly to $\rho_h\chi_K$. Thus,

$$\int_K \rho_h^2 d\Sigma \leq \liminf_{n \rightarrow \infty} \int_K |Dg_n|^2 d\Sigma \leq \Sigma(\pi_Y^{-1}(h(\widehat{K}))) + \varepsilon.$$

Finally, we let $\varepsilon \rightarrow 0$. □

Lemma 4.17. *The function ρ_h is supported in the set $\pi_X^{-1}(h^{-1}(\pi_Y(Y)))$.*

Proof. It suffices to show that $\rho_h = 0$ on $\pi_X^{-1}(h^{-1}(\pi_Y(q_i)))$ for each $i \in \mathbb{N}$. The argument is similar to the one used in the previous lemma. Let K be a non-empty compact subset of $\pi_X^{-1}(h^{-1}(\pi_Y(q_i)))$. It suffices to show that $\rho_h = 0$ a.e. on K . We set $\widehat{K} = \pi_X(K)$ and note that $h(\widehat{K})$ is the singleton $\pi_Y(q_i)$. By Lemma 2.2, for each $\delta > 0$, there exists an open neighborhood $\widehat{V}$ of $h(\widehat{K})$ such that the open set $\pi_Y^{-1}(\widehat{V})$ contains $\pi_Y^{-1}(h(\widehat{K}))$ and is contained the open δ -neighborhood of $\pi_Y^{-1}(h(\widehat{K})) = q_i$. Therefore, for each $\varepsilon > 0$, we may find such an open set $\widehat{V}$ with the additional property that

$$\Sigma(\pi_Y^{-1}(\widehat{V}) \setminus q_i) < \varepsilon.$$

As in the proof of Lemma 4.16, $g_n^*(\pi_X^{-1}(\widehat{K})) \subset \pi_Y^{-1}(\widehat{V})$ for all sufficiently large $n \in \mathbb{N}$, and particularly, $g_n^*(\pi_X^{-1}(\widehat{K}) \setminus p_{i,n}) \subset \pi_Y^{-1}(\widehat{V}) \setminus q_i$. We now obtain

$$\begin{aligned} \int_K |Dg_n|^2 d\Sigma &= \int_{(K \setminus p_{i,n}) \cap X_n} |Dg_n|^2 d\Sigma \leq \int_{(\pi_X^{-1}(\widehat{K}) \setminus p_{i,n}) \cap X_n} |Dg_n|^2 d\Sigma \\ &= \Sigma(g_n((\pi_X^{-1}(\widehat{K}) \setminus p_{i,n}) \cap X_n)) \leq \Sigma(\pi_Y^{-1}(\widehat{V}) \setminus q_i) < \varepsilon. \end{aligned}$$

Taking limits and using the weak convergence of $|Dg_n|\chi_K$ to $\rho_h\chi_K$, we obtain

$$\int_K \rho_h^2 d\Sigma \leq \varepsilon.$$

We let $\varepsilon \rightarrow 0$, so $\rho_h = 0$ a.e. on K . □

A consequence of Lemma 4.16 and Lemma 4.17 is the following statement, which concludes the proof that h is packing-conformal and the proof of Theorem 4.1.

Corollary 4.18. *For all Borel sets $E \subset \widehat{\mathbb{C}}$ we have*

$$\int_{\pi_X^{-1}(h^{-1}(\pi_Y(E)))} \rho_h^2 d\Sigma \leq \Sigma(E \cap Y).$$

5. TOPOLOGY OF PLANAR MAPS

Our goal in this section is to study continuous, proper, and cell-like maps between simply connected domains in the sphere. In view of the approximation theorem, Theorem 2.8, these maps behave like homeomorphisms. Specifically, we wish to understand when these maps have an extension to the boundaries and to the whole sphere satisfying certain properties. This section can be read independently of the other sections. The results are used in the proof of Theorem 6.1 in the next section.

5.1. Conditions for continuous extension. Let $f: \Omega \rightarrow D$ be a map between domains in $\widehat{\mathbb{C}}$. For a point $z_0 \in \partial\Omega$ we define the *cluster set* $\text{Clu}(f, z_0)$ to be the set of accumulation points of $\{f(z_n)\}_{n \in \mathbb{N}}$ over all sequences $\{z_n\}_{n \in \mathbb{N}}$ in Ω converging to z_0 . Recall from Section 2.3 that a continuous map between open subsets of the sphere is cell-like if the preimage of each point is a non-separating continuum.

Lemma 5.1. *Let $\Omega, D \subset \widehat{\mathbb{C}}$ be simply connected regions such that ∂D is a Peano continuum and let $f: \Omega \rightarrow D$ be a continuous, proper, and cell-like map.*

- (i) *For each $z_0 \in \partial\Omega$ and $\varepsilon > 0$ there exists $\delta > 0$ such that if γ is a closed curve in $B(z_0, \delta) \setminus \{z_0\}$ that is not null-homotopic, then*

$$\text{diam}(f(|\gamma| \cap \Omega)) \geq \text{diam}(\text{Clu}(f, z_0)) - \varepsilon.$$

- (ii) *The map f extends to a continuous map on $\overline{\Omega}$ if and only if for each $z_0 \in \partial\Omega$ and $z_\infty \in \widehat{\mathbb{C}} \setminus \{z_0\}$ there exists a sequence of curves γ_n in $\widehat{\mathbb{C}} \setminus \{z_0, z_\infty\}$ that are not null-homotopic and converge to z_0 such that*

$$\lim_{n \rightarrow \infty} \text{diam}(f(|\gamma_n| \cap \Omega)) = 0.$$

In this case $f(\overline{\Omega}) = \overline{D}$ and $f^{-1}(\partial D) = \partial\Omega$.

Proof. This first part of the lemma is established in [NY20], as Lemma 3.10, under the assumption that f is a homeomorphism and the complements of Ω, D are points or closed Jordan regions; see the first two paragraphs in [NY20, p. 152]. The proof applies with few changes to this more general setting so we omit it, but we make instead a few remarks. One can reduce the statement to homeomorphisms via the approximation theorem, Theorem 2.8. In particular, one can replace f by a homeomorphism without altering the cluster sets and so that $\text{diam}(f(|\gamma| \cap \Omega))$ is altered very slightly. The assumption that Ω is a Jordan region or a point is not used in [NY20]. Finally, for D all we need is that it has the property that any two points that are close to each other can be connected with a path in D that is small in diameter; this is true since ∂D is a Peano continuum [Why42, Theorem (4.2), p. 112].

We prove part (ii). If there is a continuous extension then the conclusion holds trivially for any sequence of curves γ_n as in the statement. Conversely, suppose that for each z_0 there exists a sequence of curves γ_n as in the statement. By (i) we see that $\text{Clu}(f, z_0)$ contains only one point. Thus, f extends continuously to $\overline{\Omega}$. The surjectivity of $f: \Omega \rightarrow D$ from Lemma 2.9 (i) implies the surjectivity of the extension onto $\overline{D}$. The properness of f implies that $\text{Clu}(f, z_0) \subset \partial D$ for each $z_0 \in \partial\Omega$. Hence $f^{-1}(\partial D) = \partial\Omega$. $\square$

5.2. Implications of continuous extension. We state the main result of the section.

Theorem 5.2. *Let $\Omega \subset \widehat{\mathbb{C}}$ be a simply connected region and $Y, D \subset \widehat{\mathbb{C}}$ be Jordan regions such that $\Omega \subset Y$. Assume that $f: \Omega \rightarrow D$ is a continuous, proper, and cell-like map that extends to a continuous map from $\overline{\Omega}$ onto $\overline{D}$ with the property that for each component V of $Y \setminus \overline{\Omega}$, $f|_{\partial V \setminus \partial Y}$ is constant. Then there exists a unique extension of f to a continuous and cell-like map $\tilde{f}: \overline{Y} \rightarrow \overline{D}$. The map $\tilde{f}$ is constant in each continuum $E \subset Y \setminus \Omega$ and can be further extended to a continuous and cell-like map $\tilde{f}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ such that $\tilde{f}^{-1}(\widehat{\mathbb{C}} \setminus \overline{D}) = \widehat{\mathbb{C}} \setminus \overline{Y}$.*

The proof will be completed in several steps. We establish some preliminary statements.

Lemma 5.3. *Let $U \subsetneq \widehat{\mathbb{C}}$ be an open set with $\partial\overline{U} = \partial U$ and let $P \subset U$ be a Jordan region. Then for each component V of $U \setminus \overline{P}$ the set $\partial V \setminus \partial P$ is a non-empty dense subset of $\partial V \cap \partial U$.*

The assumption $\partial\overline{U} = \partial U$ holds automatically if U is a connected component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$ for some open set Ω .

Proof. Using the Schoenflies theorem, we may assume that ∂P is a circle. Suppose that $\partial V \setminus \partial P = \emptyset$, so $\partial V \subset \partial P$. The set ∂V is a non-empty closed subset of ∂P . Let $x \in \partial V$ and consider a ball $B(x, r)$ that does not contain P . Then ∂P separates the ball $B(x, r)$ into two connected open sets, one contained in P , and one outside $\overline{P}$. The latter set does not intersect ∂V and contains points of V , since $x \in \partial V$. By the connectedness of V , we have $B(x, r) \setminus \overline{P} \subset V$. This implies that $B(x, r) \cap \partial P \subset \partial V$, so ∂V is open in the relative topology of ∂P . We conclude that $\partial V = \partial P$. The Jordan curve theorem implies that $V = \widehat{\mathbb{C}} \setminus \overline{P}$. Thus, $\widehat{\mathbb{C}} = V \cup \overline{P} \subset U$, which contradicts the assumption that $U \subsetneq \widehat{\mathbb{C}}$. Therefore, $\partial V \setminus \partial P \neq \emptyset$.

We show that $\partial V \setminus \partial P \subset \partial V \cap \partial U$. If not, there exists $x \in \partial V \setminus \partial P \subset \partial V \subset \overline{U}$ with $x \in U$. Since $x \notin \overline{P}$, there exists a ball $B(x, r) \subset U \setminus \overline{P}$. This implies that $V \cup B(x, r)$ is a connected open subset of $U \setminus \overline{P}$, which contradicts the assumption that V is a connected component of $U \setminus \overline{P}$.

It remains to show that each point of $\partial V \cap \partial U \cap \partial P$ can be approximated by points of $\partial V \setminus \partial P$. For the sake of contradiction, suppose that there exists $x \in \partial V \cap \partial U \cap \partial P$ and an open ball $B(x, r)$ that does not intersect $\partial V \setminus \partial P$ and does not contain ∂P (upon choosing a small enough $r > 0$). Thus, $B(x, r) \cap \partial V \subset \partial P$. Consider the arc $\alpha = B(x, r) \cap \partial P$. The arc α separates the ball $B(x, r)$ into precisely two regions: one region contained in P and hence not intersecting V , and one region outside $\overline{P}$ that does not intersect ∂V . The latter region contains points of V near x , since $x \in \partial V$. By the connectedness of V , this region is contained in V . Therefore, $\alpha \subset \partial V$. We conclude that $B(x, r)$ is contained in $P \cup \overline{V} \subset \overline{U}$. This contradicts that $x \in \partial U = \partial\overline{U}$. $\square$

Lemma 5.4. *Let $E \subset \widehat{\mathbb{C}}$ be a compact set, $D \subset \widehat{\mathbb{C}}$ be a Jordan region, and $g: E \rightarrow \overline{D}$ be a continuous, surjective, and cell-like map. Then $E \neq \widehat{\mathbb{C}}$, E is a non-separating continuum, and $g(\partial E) = \partial D$. Moreover, there exists an extension of g to a continuous and cell-like map $\tilde{g}: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ such that $\tilde{g}^{-1}(\widehat{\mathbb{C}} \setminus \overline{D}) = \widehat{\mathbb{C}} \setminus E$.*

Proof. We consider a decomposition of $\widehat{\mathbb{C}}$ into the non-separating continua $g^{-1}(z)$, $z \in \overline{D}$, and the remaining singleton points. This decomposition is upper semicontinuous, as follows from the continuity of g . By Moore's theorem (Theorem 2.10), $\widehat{\mathbb{C}}/\sim$ is a topological 2-sphere. Consider the projection $\pi: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}/\sim$. We define the map $G = g \circ \pi^{-1}$ on $\pi(E)$ by mapping each point $\pi(g^{-1}(z))$ to z , where $z \in \overline{D}$. Trivially, this map is injective. Using Lemma 2.2 and the continuity of g one can show that G is continuous from $\pi(E)$ onto $\overline{D}$. Therefore, G is a homeomorphism, which implies that $\pi(E)$ is a closed Jordan region, whose interior we denote by W ; moreover, $G(\partial W) = \partial D$. The region $\widehat{\mathbb{C}} \setminus E$ projects under π homeomorphically onto the complement of $\overline{W}$. In particular, $\widehat{\mathbb{C}} \setminus E \neq \emptyset$ and $\pi(\partial E) = \pi(\partial(\widehat{\mathbb{C}} \setminus E)) = \partial W$.




FIGURE 4. The region U_0 is the unbounded component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$. The map $\tilde{f}$ is constant on each component $U \neq U_0$ of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$. Here the set A_z is the closure of the shaded regions.

This implies that $g(\partial E) = G(\pi(\partial E)) = \partial D$. Finally, consider an arbitrary extension of G to a homeomorphism of the sphere $\widehat{\mathbb{C}}/\sim$. Then $\tilde{g} = G \circ \pi$ gives the desired extension of g . $\square$

In the next lemmas the standing assumptions are that $\Omega \subset \widehat{\mathbb{C}}$ is a simply connected region, $D \subset \widehat{\mathbb{C}}$ is a Jordan region, and $f: \Omega \rightarrow D$ is a continuous, proper, and cell-like map that has an extension to a continuous map from Ω onto $\overline{D}$, which we also denote by f .

Lemma 5.5. *There exists a unique component U_0 of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$ such that $f(\partial U_0) = \partial D$. Moreover, there exists an extension of f to a continuous and cell-like map $\tilde{f}: \widehat{\mathbb{C}} \setminus U_0 \rightarrow \overline{D}$ such that $\tilde{f}|_{\overline{U}}$ is constant for each component $U \neq U_0$ of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$.*

See Figure 4 for an illustration of the conclusions of the lemma.

Proof. By postcomposing f with a homeomorphism of $\widehat{\mathbb{C}}$ via the Schoenflies theorem, we assume that D is the unit disk. For each $z \in \partial D$ the set $f^{-1}(z)$ is a subset of $\partial \Omega$ by Lemma 5.1 (ii). Moreover, we have

$$f^{-1}(z) = \bigcap_{r>0} \overline{f^{-1}(B(z, r) \cap D)}$$

by the continuity of f . Since f is cell-like and the set $B(z, r) \cap D$ is connected, by Lemma 2.9 (ii) the same is true for $f^{-1}(B(z, r) \cap D)$, and thus for $\overline{f^{-1}(B(z, r) \cap D)}$. We conclude that $f^{-1}(z)$ is a connected subset of $\partial \Omega$.

For each $z \in D$, the set $A_z = f^{-1}(z)$ is a non-separating continuum in Ω , since f is cell-like and $f^{-1}(\partial D) = \partial \Omega$. For $z \in \partial D$ the set $f^{-1}(z)$ is a connected subset of $\partial \Omega$. If $f^{-1}(z)$ does not separate the sphere, then we set $A_z = f^{-1}(z)$. If $f^{-1}(z)$ separates the sphere, then $\widehat{\mathbb{C}} \setminus f^{-1}(z)$ contains at least two components. Precisely one of the components of $\widehat{\mathbb{C}} \setminus f^{-1}(z)$ contains the connected set Ω . We define A_z to be the complement of that component; see Figure 4. Then A_z is non-separating by its definition and has the property that

$$(5.1) \quad \emptyset \neq \partial A_z \subset f^{-1}(z) \subset \partial \Omega.$$

The set $\text{int}(A_z)$ is disjoint from $\overline{\Omega}$, so

$$(5.2) \quad A_z \cap \partial \Omega = \partial A_z \cap \partial \Omega \subset f^{-1}(z).$$

We claim that $A_z \cap A_w = \emptyset$ for each $w \neq z, z, w \in \overline{D}$. If $z \in D$, then $A_z = f^{-1}(z) \subset \Omega$ and the claim is trivial. Suppose that $z, w \in \partial D$. By (5.2), we have

$$A_z \cap f^{-1}(w) = A_z \cap f^{-1}(w) \cap \partial\Omega \subset f^{-1}(z) \cap f^{-1}(w) = \emptyset.$$

Thus, the connected set A_z is contained in a component of $\widehat{\mathbb{C}} \setminus f^{-1}(w)$. The definition of A_w implies that $A_z \subset A_w$ or $A_z \cap A_w = \emptyset$. If $A_z \subset A_w$, then then by reversing the roles of z and w one obtains $A_w \subset A_z$, so $A_z = A_w$. This contradicts (5.1).

Consider the set $\mathcal{A} = \bigcup_{z \in \overline{D}} A_z$, which contains $\overline{\Omega}$. We observe that this set is closed in $\widehat{\mathbb{C}}$. Indeed, if x_n is a sequence in A_{z_n} , where $z_n \in \overline{D}$ are distinct points, then there exists $x'_n \in \partial A_{z_n} \subset \overline{\Omega}$ such that $\sigma(x_n, x'_n) \rightarrow 0$. Thus, x'_n , and hence x_n as well, must accumulate at points of $\overline{\Omega} \subset \mathcal{A}$. Another observation is that

$$(5.3) \quad \partial\mathcal{A} \subset \partial\Omega.$$


Indeed, if $x \in \partial\mathcal{A} \subset \mathcal{A}$, then $x \in \partial A_z$ for some $z \in \overline{D}$. If $z \in D$, then $A_z \subset \Omega \subset \text{int}(\mathcal{A})$, so we must have $z \in \partial D$. In this case, $\partial A_z \subset \partial\Omega$ by (5.1), as desired.

We extend f to $\mathcal{A} = \bigcup_{z \in \overline{D}} A_z$ by defining $\tilde{f}(x) = z$ for $x \in A_z$. Observe that if $x \in A_z$, then for each point $x' \in \partial A_z \subset f^{-1}(z)$ we have $\tilde{f}(x) = z = f(x')$. We claim that the extension is continuous. If not, there exists $\varepsilon > 0$ and a sequence $x_n \rightarrow x$, such that $\sigma(\tilde{f}(x_n), \tilde{f}(x)) \geq \varepsilon$ for each $n \in \mathbb{N}$. Note that the point x cannot lie in the interior of A_z for any $z \in \overline{D}$, since $\tilde{f}$ is continuous there. Thus, we necessarily have $x \in \partial A_z$ for some $z \in \overline{D}$. By (5.1), $x \in \overline{\Omega}$, so $\tilde{f}(x) = f(x)$. Assume that $x_n \in A_{z_n}$, $n \in \mathbb{N}$. For each $n \in \mathbb{N}$ we connect x_n to x with a small path in $\widehat{\mathbb{C}}$ and we see that there exists a point $x'_n \in \partial A_{z_n} \subset f^{-1}(z_n)$ such that $\tilde{f}(x_n) = f(x'_n)$ and $x'_n \rightarrow x$. This contradicts the continuity of f on $\overline{\Omega}$. Therefore, $\tilde{f}$ is continuous. By construction, $\tilde{f}^{-1}(z) = A_z$ is a non-separating continuum for each $z \in \overline{D}$; thus $\tilde{f}$ is also cell-like.

By Lemma 5.4, $\widehat{\mathbb{C}} \setminus \mathcal{A} \neq \emptyset$, $\mathcal{A}$ is a non-separating continuum, and $\tilde{f}(\partial\mathcal{A}) = \partial D$. We define $U_0 = \widehat{\mathbb{C}} \setminus \mathcal{A}$, so $\tilde{f}(\partial U_0) = \partial D$. The connected set U_0 is contained in a component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$. By (5.3) we have $\partial U_0 = \partial\mathcal{A} \subset \partial\Omega$, which implies that U_0 is a component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$. Let U be a component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$ that is different from U_0 . Then $U \subset \mathcal{A}$, so $U \cap A_z \neq \emptyset$ for some $z \in \overline{D}$; in fact, $z \in \partial D$, since $U \cap \Omega = \emptyset$. Note that U is a connected set that is disjoint from $f^{-1}(z)$. By the definition of A_z we have $U \subset A_z$. In particular, $\tilde{f}$ is constant on $\overline{U}$. $\square$

Lemma 5.6. *Let $Y \subset \widehat{\mathbb{C}}$ be a Jordan region such that $\Omega \subset Y$. Assume that for each component V of $Y \setminus \overline{\Omega}$ the restriction $f|_{\partial V \setminus \partial Y}$ is constant. Then there exists a unique extension of f to a continuous and cell-like map $\tilde{f}: \overline{Y} \rightarrow \overline{D}$. The map $\tilde{f}|_{\overline{Y} \setminus \overline{\Omega}}$ attains countably many values and $\tilde{f}^{-1}(D) = \Omega$.*

Proof. Consider the region U_0 and the extension $\tilde{f}: \widehat{\mathbb{C}} \setminus U_0 \rightarrow \overline{D}$ as provided by Lemma 5.5. Let $Y \supset \Omega$ as in the statement. The connected region $\widehat{\mathbb{C}} \setminus \overline{Y}$ is contained in a component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$. If $\widehat{\mathbb{C}} \setminus \overline{Y}$ is contained in a component $U \neq U_0$ of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$, then $\tilde{f}|_{\overline{U}}$ is constant; moreover, $U_0 \subset Y$ and U_0 is a component of $Y \setminus \overline{\Omega}$. By assumption, $f|_{\partial U_0 \setminus \partial Y}$ is constant. Since $\partial U_0 \cap \partial Y \subset \partial U_0 \cap \overline{U}$, f is also constant on $\partial U_0 \cap \partial Y$. This contradicts the surjectivity of $f|_{\partial U_0}$ onto ∂D from Lemma 5.5. Therefore $\widehat{\mathbb{C}} \setminus \overline{Y} \subset U_0$ and $\partial Y \subset \overline{U_0}$.

FIGURE 5. A portion of $\partial\Omega$ and ∂Y .

Next, we extend $\tilde{f}$ to $\overline{Y} \setminus \overline{\Omega}$ as follows. If V is a component of $Y \setminus \overline{\Omega}$, then either V is a component of $\widehat{\mathbb{C}} \setminus \overline{\Omega}$ that is disjoint from U_0 , or V is a component of $Y \cap U_0$. If V is disjoint from U_0 , then $\tilde{f}$ has already been defined in $\overline{V}$ and is constant with value equal to $f(\partial V) \subset f(\partial\Omega) \subset \partial D$. Suppose that V is a component of $Y \cap U_0$. By Lemma 5.3 (applied to U_0 and $P = \widehat{\mathbb{C}} \setminus \overline{Y}$), the set $\partial V \setminus \partial Y$ is a non-empty dense subset of $\partial V \cap \partial U_0$. By assumption, $f|_{\partial V \setminus \partial Y}$ is constant, so by continuity $f|_{\partial V \cap \partial U_0}$ is constant. We define

$$\tilde{f}|_{\overline{V}} = f|_{\partial V \cap \partial U_0}.$$

Since $f(\partial U_0) \subset \partial D$, we see that $\tilde{f}(\overline{V})$ is a point of ∂D .

First, we ensure that this gives a well-defined map. Let V_1, V_2 be distinct components of $Y \cap U_0$ such that $\overline{V_1} \cap \overline{V_2} \neq \emptyset$. If $\overline{V_1} \cap \overline{V_2}$ contains a point $x \in U_0$, then that point would have to lie on ∂Y ; however, ∂Y has exactly two “sides” near x , one contained in the region Y , and one contained in precisely one of V_1, V_2 . This is a contradiction. Therefore, $\overline{V_1} \cap \overline{V_2} \subset \partial U_0$. Since f is constant on $\partial V_i \cap \partial U_0$, $i = 1, 2$, we conclude that $\tilde{f}|_{\overline{V_1}} = \tilde{f}|_{\overline{V_2}}$, as desired.

Next, we note that

$$\overline{Y} = \overline{\Omega} \cup \bigcup_{V \subset Y \setminus \overline{\Omega}} \overline{V},$$

where the union is over all components V of $Y \setminus \overline{\Omega}$. Hence, the above definition provides an extension of $\tilde{f}$ to all of $\overline{Y}$. Indeed, if $x \in Y \setminus \overline{\Omega}$, then x lies in a component V of $Y \setminus \overline{\Omega}$. If $x \in \partial Y \setminus \overline{\Omega}$, then $x \in \partial Y \cap U_0$, so x lies in the boundary of a component V of $Y \setminus \overline{\Omega}$, because ∂Y is “two-sided” near x .

Since there are countably many components V of $Y \setminus \overline{\Omega}$, we observe that $\tilde{f}$ attains countably many values in $\overline{Y} \setminus \overline{\Omega}$, as required in the conclusion of Lemma 5.6. Moreover, by the definition of the extension, we have $\tilde{f}^{-1}(D) = \Omega$.

Now, we show that $\tilde{f}$ is continuous. Since f is already continuous in the complement of U_0 by Lemma 5.5, it suffices to show that if x_n is a sequence in $\overline{Y} \cap U_0$ converging to some point x , then $\tilde{f}(x_n)$ converges to $\tilde{f}(x)$. For the sake of contradiction, suppose that there exists $\varepsilon > 0$ such that $\sigma(\tilde{f}(x_n), \tilde{f}(x)) \geq \varepsilon$ for all $n \in \mathbb{N}$. If $x \in U_0$, then $x, x_n \in \overline{V}$ for all sufficiently large $n \in \mathbb{N}$, for some component V of $Y \cap U_0$; thus, $\tilde{f}(x) = \tilde{f}(x_n)$ for all large $n \in \mathbb{N}$ and we obtain a contradiction. Suppose $x \in \partial U_0$. We have $x_n \in \overline{V_n}$ for a sequence of components V_n of $Y \cap U_0$. If $\overline{V_n}$ is a specific component $\overline{V}$ for infinitely many $n \in \mathbb{N}$, then $x \in \overline{V}$, so $\tilde{f}(x_n) = \tilde{f}(x)$

for infinitely many $n \in \mathbb{N}$, a contradiction. Thus, we may assume that the components V_n , $n \in \mathbb{N}$, are distinct. One may find points $y_n \in \partial V_n$ with $\sigma(y_n, x_n) \rightarrow 0$ and $\tilde{f}(y_n) = \tilde{f}(x_n)$. By [Pom92, Prop. 2.13] for each V_n there exists an open arc $\alpha_n = \partial V_n \cap U_0$ that is a component of $\partial Y \cap U_0$, so that ∂V_n is the union of α_n and $\partial V_n \cap \partial U_0$; see Figure 5. Moreover, for distinct sets V_n the arcs α_n are disjoint, since ∂Y is “two-sided”. The local connectivity of ∂Y implies that $\text{diam}(\alpha_n) \rightarrow 0$ as $n \rightarrow \infty$. If $y_n \in \partial V_n \cap \partial U_0$, we set $z_n = y_n$ and if $y_n \in \alpha_n$, we set z_n to be an endpoint of α_n (Figure 5). Therefore, one may find points $z_n \in \partial V_n \cap \partial U_0$ such that $\tilde{f}(y_n) = \tilde{f}(z_n) = f(z_n)$ and $\sigma(y_n, z_n) \rightarrow 0$. Since $z_n \rightarrow x$, the continuity of f on $\partial U_0 \subset \bar{\Omega}$ implies that $f(z_n) \rightarrow f(x)$. This contradiction completes the proof of the continuity of $\tilde{f}$.

Next, we show that for each $z \in \bar{D}$, the set $\tilde{f}^{-1}(z)$ is a non-separating continuum. If $z \in D$, then $\tilde{f}^{-1}(z) = f^{-1}(z) \subset \Omega$, so $f^{-1}(z)$ is a non-separating continuum by the cell-likeness of f . Next, suppose that $z \in \partial D$. By Lemma 5.5, $\tilde{f}|_{\hat{\mathbb{C}} \setminus U_0}$ is cell-like, so the set $E = \tilde{f}^{-1}(z) \cap (\hat{\mathbb{C}} \setminus U_0)$ is a non-separating continuum. By the definition of $\tilde{f}$ on $\bar{Y}$, $\tilde{f}^{-1}(z)$ is the union of E with the closures of components V of $Y \cap U_0$ such that $\tilde{f}|_{\bar{V}} \equiv z$. By the previous, for each such component we have $\partial V \cap \partial U_0 \neq \emptyset$, so $\bar{V} \cap E \neq \emptyset$. This shows that $\tilde{f}^{-1}(z)$ is connected.

In order to show that $\tilde{f}^{-1}(z)$ is non-separating, we observe that $\hat{\mathbb{C}} \setminus \tilde{f}^{-1}(z)$ is the union of $\bar{\Omega} \setminus \tilde{f}^{-1}(z)$ with $\hat{\mathbb{C}} \setminus \bar{Y}$ and with the sets $\bar{V}$ where V is a component of $Y \setminus \bar{\Omega}$ with $\bar{V} \cap \tilde{f}^{-1}(z) = \emptyset$. Let W be the component of $\hat{\mathbb{C}} \setminus \tilde{f}^{-1}(z)$ that contains the connected set Ω and note that $\bar{\Omega} \setminus \tilde{f}^{-1}(z)$ is connected, so it is contained in W . The sets V as above are either components of $\hat{\mathbb{C}} \setminus \bar{\Omega}$ or components of $Y \cap U_0$. In the first case we have $\partial V \subset \partial \Omega$ so $\bar{V} \cap (\bar{\Omega} \setminus \tilde{f}^{-1}(z)) \neq \emptyset$, while in the second case, $\partial V \cap \partial U_0 \neq \emptyset$, so $\bar{V} \cap (\bar{\Omega} \setminus \tilde{f}^{-1}(z)) \neq \emptyset$. Therefore, in both cases, $\bar{V} \subset W$. Since $\tilde{f}|_{\partial U_0}$ is surjective onto ∂D , there exists a point $x \in \partial U_0 \setminus \tilde{f}^{-1}(z) \subset \bar{\Omega} \setminus \tilde{f}^{-1}(z)$. If $x \in \partial Y$, then this shows that $\hat{\mathbb{C}} \setminus \bar{Y}$ lies in W . If $x \in \partial V$ for some V as above, then $\bar{V} \subset W$ by the previous and ∂V contains an arc of ∂Y . Thus, $\hat{\mathbb{C}} \setminus \bar{Y} \subset W$. This completes the proof that $W = \hat{\mathbb{C}} \setminus \tilde{f}^{-1}(z)$, and thus $\tilde{f}^{-1}(z)$ is non-separating.

Now, we show the uniqueness statement. Suppose that $\tilde{f}_1$ is another extension of f to a continuous and cell-like map from $\bar{Y}$ onto $\bar{D}$. We first observe that $\tilde{f}_1^{-1}(D) = \Omega$; indeed, if $\tilde{f}_1^{-1}(z)$ intersects the complement of Ω for some $z \in D$, then $\tilde{f}_1^{-1}(z)$ is a continuum that intersects both Ω and its complement, so it intersects $\partial \Omega$. However, $\tilde{f}_1(\partial \Omega) = f(\partial \Omega) = \partial D$, a contradiction.

Let V be a component of $Y \setminus \bar{\Omega}$, such that $\tilde{f}|_{\bar{V}} \equiv z$ for some $z \in \partial D$. Our goal is to show that $\tilde{f}_1 \equiv z$ on $\bar{V}$. Suppose that $\tilde{f}_1^{-1}(w) \cap \bar{V} \neq \emptyset$ for some $w \in \bar{D}$; then we necessarily have $w \in \partial D$ by the previous paragraph. The continuum $E = \tilde{f}_1^{-1}(w)$ intersects ∂U_0 , since $\tilde{f}_1|_{\partial U_0} = f|_{\partial U_0} : \partial U_0 \rightarrow \partial D$ is surjective. This implies that $\partial V \cap E \neq \emptyset$.

If $V \cap U_0 = \emptyset$, then V is a component of $\hat{\mathbb{C}} \setminus \bar{\Omega}$, so $\partial V \subset \partial \Omega$. Hence, $w = \tilde{f}_1|_{\partial V \cap E} = f|_{\partial V \cap E} = \tilde{f}|_{\bar{V}} = z$. Now, suppose that $V \subset U_0$, so V is a component of $Y \cap U_0$. We claim that $E \cap \partial V \cap \partial U_0 \neq \emptyset$. Suppose that $E \cap \partial V \cap \partial U_0 = \emptyset$. Note that $E \cap \bar{V}$ is a closed subset of E . We will show that $E \cap \bar{V}$ is open in the relative topology of E . It suffices to show that each $x \in E \cap \partial V$ has an open neighborhood

in the topology of E that is contained in $E \cap \bar{V}$. By assumption, we have

$$x \in E \cap \partial V = E \cap \partial V \cap (\partial Y \cup \partial U_0) = E \cap \partial V \cap \partial Y.$$

Since Y is “two-sided” and $E \subset \bar{Y}$, all points of E near x are contained in $\bar{V}$, as desired. Since $E \cap \bar{V}$ is clopen in E , the connectedness of E implies now that $E \cap \bar{V} = E$. Thus,

$$E \cap \partial V \cap \partial U_0 = E \cap \bar{V} \cap \partial U_0 = E \cap \partial U_0 \neq \emptyset,$$

a contradiction. Therefore, $E \cap \partial V \cap \partial U_0 \neq \emptyset$, which implies that $w = \tilde{f}_1|_{E \cap \partial V \cap \partial U_0} = \tilde{f}|_{\partial V \cap \partial U_0} = \tilde{f}|_{\bar{V}} = z$. Hence, we conclude that $w = z$ in all cases, as desired. $\square$

Proof of Theorem 5.2. Consider the unique extension of f to a continuous and cell-like map $\tilde{f}: \bar{Y} \rightarrow \bar{D}$ as provided by Lemma 5.6. By Lemma 5.4 the map $\tilde{f}$ can be extended to a continuous and cell-like map of the sphere such that $\tilde{f}^{-1}(\hat{\mathbb{C}} \setminus \bar{D}) = \hat{\mathbb{C}} \setminus \bar{Y}$, as required in the theorem. Moreover, $\tilde{f}(\partial Y) = \partial D$. By Lemma 5.6, $\tilde{f}$ takes countably many values in $\bar{Y} \setminus \bar{\Omega}$. If $\partial Y \cap \partial \Omega = \emptyset$, then $\partial Y \subset \bar{Y} \setminus \bar{\Omega}$, so $\tilde{f}$ is constant on the connected set ∂Y . This contradicts the fact that $\tilde{f}(\partial Y) = \partial D$. Therefore, $\partial Y \cap \partial \Omega \neq \emptyset$.

Let E be a continuum in $Y \setminus \Omega$ as in the statement of the theorem. Consider the component A of $\hat{\mathbb{C}} \setminus E$ that contains the connected set $\hat{\mathbb{C}} \setminus Y$. Since $\partial Y \cap \partial \Omega \neq \emptyset$ and A is open, we see that Ω is also contained in A . Let E' be the complement of A . Then $E' \supset E$ and E' is a non-separating continuum disjoint from Ω and $\hat{\mathbb{C}} \setminus Y$. We claim that $\tilde{f}$ is constant on E' . We argue by contradiction, assuming that $\tilde{f}(E')$ is a non-degenerate continuum. By Lemma 5.6, $\tilde{f}^{-1}(D) = \Omega$, so $\tilde{f}(E')$ is an arc of ∂D . Then there exist distinct points $z_1, z_2 \in \partial D$ such that $\tilde{f}^{-1}(z_i) \cap E' \neq \emptyset$, $i \in \{1, 2\}$. Since $\tilde{f}(\partial Y) = \partial D$, we see that $\tilde{f}^{-1}(z_i) \cap \partial Y \neq \emptyset$ for $i \in \{1, 2\}$.

We collapse topologically the non-separating continua $K = \hat{\mathbb{C}} \setminus Y$ and E' to points (e.g. via Moore’s theorem) and consider a continuous projection map $\lambda: \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ such that $\lambda(K) = w_1$, $\lambda(E') = w_2$ for some points $w_1 \neq w_2$, and λ is a homeomorphism from $\hat{\mathbb{C}} \setminus (K \cup E')$ onto $\hat{\mathbb{C}} \setminus \{w_1, w_2\}$. The continua $G_i = \lambda(\tilde{f}^{-1}(z_i))$, $i \in \{1, 2\}$, meet at the points w_1 and w_2 , but otherwise they are disjoint. For $i \in \{1, 2\}$ there exists a continuum $G'_i \subset G_i$ that is *irreducible* between the points w_1 and w_2 ; that is, there is no proper subcontinuum of G'_i that contains both w_1 and w_2 . See [Wil70, Theorem 28.4] for the existence.

A result of Moore [Moo28, Theorems 1 and 2] implies that $\hat{\mathbb{C}} \setminus (G'_1 \cup G'_2)$ is the union of precisely two domains and the boundary of each is $G'_1 \cup G'_2$. Let Z be the component of $\hat{\mathbb{C}} \setminus (G'_1 \cup G'_2)$ that is disjoint from $\lambda(\Omega)$ and thus from $\lambda(\bar{\Omega})$. Since $\tilde{f}$ attains countably many values in $\bar{Y} \setminus \bar{\Omega}$, we conclude that $\tilde{f} \circ \lambda^{-1}$ is constant in Z . However, the boundary values of $\tilde{f} \circ \lambda^{-1}|_Z$ are non-constant, a contradiction. $\square$

6. CONTINUOUS EXTENSION

In this section we provide sufficient conditions so that a packing-quasiconformal map between the topological spheres associated to two Sierpiński packings as in Definition 2.15 can be lifted to a map between the actual packings.

Theorem 6.1. *Let $X = \hat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ and $Y = \hat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} q_i$ be Sierpiński packings such that the peripheral continua of X are uniformly fat closed Jordan regions or points and the peripheral continua of Y are closed Jordan regions or points with*

diameters lying in $\ell^2(\mathbb{N})$. Let $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ be a packing- K -quasiconformal map for some $K \geq 1$. Then there exists a continuous, surjective, and monotone map $H: \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ such that $\pi_Y \circ H = h \circ \pi_X$ and $H^{-1}(\text{int}(q_i)) = \text{int}(p_i)$ for each $i \in \mathbb{N}$. Moreover, there exists a non-negative Borel function $\rho_H \in L^2(\widehat{\mathbb{C}})$ with the following properties.

- There exists a curve family Γ_0 in $\widehat{\mathbb{C}}$ with $\text{Mod}_2 \Gamma_0 = 0$ such that for all curves $\gamma: [a, b] \rightarrow \widehat{\mathbb{C}}$ outside Γ_0 we have

$$\sigma(H(\gamma(a)), H(\gamma(b))) \leq \int_{\gamma} \rho_H ds + \sum_{i: p_i \cap |\gamma| \neq \emptyset} \text{diam}(q_i).$$

- For each Borel set $E \subset \widehat{\mathbb{C}}$ we have

$$\int_{H^{-1}(E)} \rho_H^2 d\Sigma \leq K \Sigma(E \cap Y).$$

Moreover, we show that any map H as in the theorem has a further topological property.

Proposition 6.2. *Let H be a map satisfying the conclusions of Theorem 6.1. Then for each $i \in \mathbb{N}$ and for each continuum $E \subset H^{-1}(q_i) \setminus p_i$ the map $H|_E$ is constant.*

Remark 6.3. The assumptions of Theorem 6.1 can be relaxed, allowing the possibility that finitely many peripheral continua of X are not fat. See Remark 6.10 below for more details.

The proofs of both statements are given in Section 6.3. We first establish several preliminary statements.

6.1. Preliminaries.

Lemma 6.4. *Let $\tau > 0$ and $X = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} p_i$ be a τ -cofat Sierpiński packing. Let $\rho: \widehat{\mathbb{C}} \rightarrow [0, \infty]$ be a Borel function in $L^2(\widehat{\mathbb{C}})$ and $\{\lambda_i\}_{i \in \mathbb{N}}$ be a non-negative sequence in $\ell^2(\mathbb{N})$. For each $x, y \in \widehat{\mathbb{C}}$ and $0 < r_0 < \text{diam}(\widehat{\mathbb{C}})$ we have*

$$\begin{aligned} & \text{ess inf}_{r \in (r_0/2, r_0)} \left(\int_{S(x, r)} \rho ds + \int_{S(y, r)} \rho ds + \sum_{i: p_i \cap S(x, r) \neq \emptyset} \lambda_i + \sum_{i: p_i \cap S(y, r) \neq \emptyset} \lambda_i \right) \\ & \leq c(\tau) \left(\int_{B(x, r_0) \cup B(y, r_0)} \rho^2 d\Sigma + \sum_{i: p_i \cap B(x, r_0) \neq \emptyset} \lambda_i^2 + \sum_{i: p_i \cap B(y, r_0) \neq \emptyset} \lambda_i^2 \right)^{1/2}. \end{aligned}$$

It is straightforward to obtain the conclusion of the lemma if $\lambda_i \equiv 0$ by integrating over $r \in (r_0/2, r_0)$ and using the co-area inequality of Proposition 2.1. For the general statement one uses the cofatness of X as well, in order to treat the summation terms. The proof of the lemma follows from the argument used in [Nta20b, Lemma 2.4.7 (a)] and we omit it.

Lemma 6.5. *Let Γ_0 be a path family in $\widehat{\mathbb{C}}$ with $\text{Mod}_2 \Gamma_0 = 0$ and let $x, y \in \widehat{\mathbb{C}}$. Define Γ_1 to be the family of paths γ such that the set $\{r > 0 : \text{there exists a simple path } \gamma_r \in \Gamma_0 \text{ whose trace is contained in } |\gamma| \cup S(x, r) \cup S(y, r)\}$ has positive Lebesgue outer measure. Then $\text{Mod}_2 \Gamma_1 = 0$. In particular, for Mod_2 -a.e. path γ and for a.e. $r > 0$, every simple path γ_r whose trace is contained in $|\gamma| \cup S(x, r) \cup S(y, r)$ lies outside Γ_0 .*

Proof. Since $\text{Mod}_2 \Gamma_0 = 0$, there exists a non-negative Borel function $\rho \in L^2(\widehat{\mathbb{C}})$ such that $\int_\gamma \rho ds = \infty$ for each $\gamma \in \Gamma_0$ (see [HKST15, Lemma 5.2.8, p. 129]). Let $\gamma \in \Gamma_1$. Then for a set of $r > 0$ of positive Lebesgue outer measure there exists a simple path $\gamma_r \in \Gamma_0$ whose trace is contained in $|\gamma| \cup S(x, r) \cup S(y, r)$. Thus,

$$\infty = \int_{\gamma_r} \rho ds \leq \int_\gamma \rho ds + \int_{S(x, r)} \rho ds + \int_{S(y, r)} \rho ds.$$

By the co-area inequality of Proposition 2.1, the latter two line integrals are finite for a.e. $r > 0$. Thus, we conclude that $\int_\gamma \rho ds = \infty$. This implies that $\text{Mod}_2 \Gamma_1 = 0$. $\square$

Remark 6.6. We will use the following standard fact about modulus. If $\text{Mod}_2 \Gamma_0 = 0$, then the family of paths that have a subpath in Γ_0 also has modulus zero [Väi71, Theorem 6.4, p. 17]. Thus, we may apply the transboundary upper gradient inequality of packing-quasiconformal maps (see Definition 2.15) not only for paths outside a family of modulus zero, but also for all subpaths of such paths, upon enlarging the exceptional curve family Γ_0 .

6.2. Continuous extension to each peripheral continuum. Let X, Y be as in Theorem 6.1 and $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ be a packing-quasiconformal map as in Definition 2.15. The proof of Theorem 6.1 will be completed in several steps. Consider the map $g = h \circ \pi_X: \widehat{\mathbb{C}} \rightarrow \mathcal{E}(Y)$, which is continuous and cell-like; indeed, the preimage of each point under h is a non-separating continuum and the preimage of each non-separating continuum under π_X is a non-separating continuum by Lemma 2.9 (iv). Recall that

$$\mathcal{E}(Y) = \pi_Y(\widehat{\mathbb{C}}) = \pi_Y(Y) \cup \bigcup_{i \in \mathbb{N}} \pi_Y(q_i).$$

Let $\tilde{X} = g^{-1}(\pi_Y(Y))$ and note that $\tilde{X} \subset X$ since $g(p_i) = h(\pi_X(p_i)) = \pi_Y(q_i)$ for each $i \in \mathbb{N}$, by the definition of a packing-quasiconformal map. If we set $\tilde{p}_i = g^{-1}(\pi_Y(q_i))$, we see that $\tilde{X} = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} \tilde{p}_i$. Note that $\tilde{p}_i$ is non-separating, since g is cell-like.

Lemma 6.7. *The map $H = \pi_Y^{-1} \circ g: \tilde{X} \rightarrow Y$ is continuous, surjective, and cell-like.*

Proof. Note that $g(\tilde{X}) = \pi_Y(Y)$ and that each point of $\pi_Y(Y)$ has a unique preimage under π_Y , so $H = \pi_Y^{-1} \circ g$ gives a well-defined map from $\tilde{X}$ onto Y . The continuity of the maps π_Y, g , together with the injectivity of π_Y on Y , imply that H is continuous on $\tilde{X}$. Finally, the fact that g is cell-like implies immediately that H is cell-like. $\square$

We fix $i \in \mathbb{N}$ and consider the space $\mathcal{E}(Y; q_i)$. Recall that for a set $E \subset \widehat{\mathbb{C}}$ the space $\mathcal{E}(Y; E)$ is defined by collapsing to points all peripheral continua of Y that do not intersect E . The space $\mathcal{E}(Y; q_i)$ is a topological 2-sphere by Moore's theorem (Theorem 2.11). The projection $\pi_{Y; q_i}: \widehat{\mathbb{C}} \rightarrow \mathcal{E}(Y; q_i)$ maps each q_j , $j \neq i$, to a point and is injective otherwise.

Lemma 6.8. *For each $i \in \mathbb{N}$ the open set $\Omega_i = \widehat{\mathbb{C}} \setminus \tilde{p}_i$ is simply connected and $D_i = \mathcal{E}(Y; q_i) \setminus \pi_{Y; q_i}(q_i)$ is a Jordan region or the complement of a point. Moreover, the map*

$$g_i = \pi_{Y; q_i} \circ \pi_Y^{-1} \circ g: \Omega_i \rightarrow D_i,$$

defined alternatively by $g_i = \pi_{Y;q_i} \circ H$ on $\tilde{X}$ and $g_i(\tilde{p}_j) = \pi_{Y;q_i}(q_j)$ for $j \neq i$, is continuous, proper, and cell-like.

Proof. For simplicity, we set $p = p_i$, $\tilde{p} = \tilde{p}_i$, $q = q_i$, and $\pi_i = \pi_{Y;q_i}$. Since $\tilde{p}$ is a non-separating continuum, the domain $\Omega = \hat{\mathbb{C}} \setminus \tilde{p}$ is simply connected. Next, since π_i is injective on q , we see that $\pi_i(q)$ is a closed Jordan region or a point in $\mathcal{E}(Y; q)$. Thus, $D = \mathcal{E}(Y; q) \setminus \pi_i(q)$ is a Jordan region or the complement of a point.

Observe that $\pi_{Y;q_i}(\pi_Y^{-1}(g(x)))$ is a point of D for each $x \in \Omega$. We show that the map $g_i = \pi_i \circ \pi_Y^{-1} \circ g$ is continuous on Ω . Let $x \in \Omega$ and let $x_n \in \Omega$ be a sequence converging to x . Then $g(x_n)$ converges to $g(x)$. By Lemma 2.2 (i), $\pi_Y^{-1}(g(x_n))$ is contained in arbitrarily small neighborhoods of $\pi_Y^{-1}(g(x))$ as $n \rightarrow \infty$. By the continuity of π_i , $\pi_i(\pi_Y^{-1}(g(x_n)))$ converges to the point $\pi_i(\pi_Y^{-1}(g(x)))$, as desired.

If a sequence $x_n \in \Omega$ accumulates at $\partial\Omega \subset \tilde{p}$, then $g(x_n)$ accumulates at the point $\pi_Y(q_i)$ by continuity. By Lemma 2.2 (i), $\pi_Y^{-1}(g(x_n))$ accumulates at q_i , so $g_i(x_n)$ accumulates at $\pi_i(q_i) \supset \partial D$. Thus, g_i is proper. Finally, g_i is cell-like, directly from its definition and from the cell-likeness of H from Lemma 6.7. $\square$

In the proofs of the remaining statements of Section 6.2, we adopt the same notational conventions as in the first paragraph of the proof of Lemma 6.8.

Lemma 6.9. *For each $i \in \mathbb{N}$ the map $g_i: \Omega_i \rightarrow D_i$ has an extension to a continuous map from $\overline{\Omega_i}$ onto $\overline{D_i}$.*

Proof. We will show that for each $a \in \partial\Omega = \partial\tilde{p}$ there exists a sequence of closed curves C_n surrounding a and shrinking to a so that $\text{diam}(H(C_n \cap \tilde{X})) \rightarrow 0$. Suppose that this is the case. The continuity of $\pi_i = \pi_{Y;q_i}$ then implies that $\text{diam}(\pi_i(H(C_n \cap \tilde{X}))) \rightarrow 0$ so the relation $g_i = \pi_i \circ H$ from Lemma 6.8 yields

$$\text{diam}(g_i(C_n \cap \tilde{X})) \rightarrow 0.$$

The set $g_i(C_n \cap \tilde{X})$ is dense in $g_i(C_n \cap \Omega)$, so

$$\text{diam}(g_i(C_n \cap \Omega)) \rightarrow 0.$$

Lemma 5.1 (ii) now implies that g_i extends to a continuous map from $\overline{\Omega}$ onto $\overline{D}$, as desired.

Now we show the original claim. Suppose first that $a \in \partial\tilde{p} \setminus p$ or that p is a point and $a = p$. Fix $r_0 > 0$ so that $(B(a, r_0) \setminus \{a\}) \cap p = \emptyset$. Let $C_r = S(a, r)$, $r > 0$. For $r \in (0, r_0)$ the circle C_r does not intersect $p = p_i$. We now apply the transboundary upper gradient inequality of h , which holds along the circle C_r for a.e. $r \in (r_0/2, r_0)$; this is a consequence of Lemma 2.12. By the relation $H = \pi_Y^{-1} \circ h \circ \pi_X$ on $\tilde{X}$, we have

$$\text{diam}(H(C_r \cap \tilde{X})) \leq \int_{C_r} \rho_h ds + \sum_{\substack{j: p_j \cap C_r \neq \emptyset \\ j \neq i}} \text{diam}(q_j).$$

We apply Lemma 6.4 with $x = y = a$, $\lambda_j = \text{diam}(q_j)$ for $j \neq i$ and $\lambda_i = 0$, so we have

$$\text{ess inf}_{r \in (r_0/2, r_0)} \text{diam}(H(C_r \cap \tilde{X})) \lesssim_\tau \left(\int_{B(a, r_0)} \rho_h^2 d\Sigma + \sum_{\substack{j: p_j \cap B(a, r_0) \neq \emptyset \\ j \neq i}} \text{diam}(q_j)^2 \right)^{1/2}.$$




FIGURE 6. Illustration of the proof of Lemma 6.11.

Note that as $r_0 \rightarrow 0$ the right-hand side tends to 0, since $a \notin \bigcup_{j \neq i} p_j$. Thus, we can find a sequence $r_n \rightarrow 0$ such that $\text{diam}(H(C_{r_n} \cap \tilde{X})) \rightarrow 0$ as claimed.

Next, suppose that p is a closed Jordan region and that $a \in \partial \tilde{p} \cap p$, so $a \in \partial p$. Let $r_0 > 0$ such that $B(a, r_0)$ does not contain p . Consider a circle centered at a of radius $r < r_0$. Since ∂p is a Jordan curve, there exists a unique arc α of the circle whose endpoints lie in ∂p that separates (in $\widehat{\mathbb{C}} \setminus \text{int}(p)$) the point a from any point outside $B(a, r_0)$. We let C_r be the Jordan curve formed by the circular arc α , together with a simple arc inside the interior of p that connects the endpoints of α . Then C_r is a Jordan curve that surrounds the point a . If the non-circular part of C_r is chosen appropriately, then we may have that C_r shrinks to the point a as $r \rightarrow 0$. Now, the transboundary upper gradient inequality, applied to the circular part of C_r , gives

$$\text{diam}(H(C_r \cap \tilde{X})) \leq \int_{C_r} \rho_h ds + \sum_{\substack{j: p_j \cap C_r \neq \emptyset \\ j \neq i}} \text{diam}(q_j)$$

for a.e. $r \in (r_0/2, r_0)$. We apply Lemma 6.4 as above with $\lambda_j = \text{diam}(q_j)$ for $j \neq i$ and $\lambda_i = 0$ and the conclusion follows. $\square$

Remark 6.10. The cofatness of X is needed only for the application of Lemma 6.4. In fact, it suffices to apply the lemma to the cofat Sierpiński packing generated by continua p_j , $j \neq i$, lying in a neighborhood of p_i . In particular, the fatness of p_i is not needed.

Lemma 6.11. *For each $i \in \mathbb{N}$ and for each component V of $\text{int}(\tilde{p}_i \setminus p_i)$ the map g_i is constant on the set $\partial V \setminus \partial p_i$.*

See the shaded regions in Figure 1 for a depiction of the components of $\text{int}(\tilde{p}_i \setminus p_i)$.

Proof. Note that if $x \in \partial \tilde{p}$, then the circle $S(x, r)$ intersects $\widehat{\mathbb{C}} \setminus \tilde{p}$, and thus $\tilde{X}$, for all sufficiently small $r > 0$. Indeed, since $\tilde{p}$ is non-separating and $x \in \partial \tilde{p}$, if $S(x, r) \subset \tilde{p}$, then the complementary component of $S(x, r)$ that does not contain x

has to be contained in $\tilde{p}$. However, if this occurs for a sequence of $r \rightarrow 0$, then we see that $\tilde{p} = \widehat{\mathbb{C}}$, a contradiction.

Let V be a component of $\text{int}(\tilde{p} \setminus p)$ and observe that $\partial V \setminus \partial p \subset \partial \tilde{p}$. We fix $x, y \in \partial V \setminus \partial p$ and $\varepsilon > 0$. Let $r_0 > 0$ such that $S(x, r)$ and $S(y, r)$ are disjoint from each other, intersect both V and $\tilde{X}$, and are disjoint from p for all $r < r_0$. In Lemma 6.4 we set $\rho = \rho_h$, $\lambda_j = \text{diam}(q_j)$ for $j \neq i$, and $\lambda_i = 0$. If r_0 is sufficiently small, then there exists a set $S \subset (r_0/2, r_0)$ that has positive Lebesgue measure such that for all $r \in S$ we have

$$(6.1) \quad \int_{S(x,r) \cup S(y,r)} \rho_h ds + \sum_{\substack{j: p_j \cap S(x,r) \neq \emptyset \\ j \neq i}} \text{diam}(q_j) + \sum_{\substack{j: p_j \cap S(y,r) \neq \emptyset \\ j \neq i}} \text{diam}(q_j) < \varepsilon.$$

This uses the fact that $x, y \in \partial \tilde{p}$, so $x, y \notin \bigcup_{j \neq i} p_j$ (see also the proof of Lemma 6.9).

Consider the curve family Γ_0 that has 2-modulus zero as in the transboundary upper gradient inequality of h . Also, consider a curve family Γ_1 with $\text{Mod}_2 \Gamma_1 = 0$ as in Lemma 6.5. Then we may find (e.g. by invoking Lemma 2.12) a curve $\gamma \notin \Gamma_0 \cup \Gamma_1$ contained in the connected open set V and joining the circles $S(x, r_0/2)$ and $S(y, r_0/2)$, so that for a.e. $r \in (r_0/2, r_0)$, every simple path γ_r whose trace is contained in $|\gamma| \cup S(x, r) \cup S(y, r)$ lies outside Γ_0 . In particular, we may find such a path γ_r with $r \in S$ such that γ_r connects a point $x' \in S(x, r) \cap \tilde{X}$ to a point $y' \in S(y, r) \cap \tilde{X}$; see Figure 6.

By the transboundary upper gradient inequality, we have

$$\sigma(H(x'), H(y')) \leq \int_{\gamma_r} \rho_h ds + \sum_{j: p_j \cap |\gamma_r| \neq \emptyset} \text{diam}(q_j).$$

Since $\rho_h = 0$ in V (see Remark 2.16), by (6.1) we have $\sigma(H(x'), H(y')) < \varepsilon$. If we let $r_0 \rightarrow 0$ and then $\varepsilon \rightarrow 0$, we see that there exist sequences $x_n, y_n \in \tilde{X}$ converging to x, y , respectively, such that $\sigma(H(x_n), H(y_n)) \rightarrow 0$. Therefore

$$\begin{aligned} d(g_i(x), g_i(y)) &= \lim_{n \rightarrow \infty} d(g_i(x_n), g_i(y_n)) \\ &= \lim_{n \rightarrow \infty} d(\pi_i(H(x_n)), \pi_i(H(y_n))) = 0. \end{aligned}$$

This shows that $g_i(x) = g_i(y)$ as desired. $\square$

Corollary 6.12. *For each $i \in \mathbb{N}$ the map $g_i: \Omega_i \rightarrow D_i$ has a unique extension to a continuous and cell-like map from $\widehat{\mathbb{C}} \setminus \text{int}(p_i)$ onto $\mathcal{E}(Y; q_i) \setminus \pi_{Y; q_i}(\text{int}(q_i))$. The extension is constant on each continuum $E \subset \tilde{p}_i \setminus p_i$ and can be further extended to a continuous and cell-like map $g_i: \widehat{\mathbb{C}} \rightarrow \mathcal{E}(Y; q_i)$ such that $g_i^{-1}(\pi_{Y; q_i}(\text{int}(q_i))) = \text{int}(p_i)$.*

Proof. If q is a point then the extension is trivial, so assume that q is a closed Jordan region. By Lemma 6.9, g_i extends to a continuous map from $\overline{\Omega}$ onto $\overline{D}$. By Lemma 6.11, for each component V of

$$\text{int}(\tilde{p} \setminus p) = \text{int}(\tilde{p}) \setminus p = (\widehat{\mathbb{C}} \setminus p) \setminus \overline{\Omega}$$

the map $g_i|_{\partial V \setminus \partial p}$ is constant. We now apply Theorem 5.2 (with $Y = \widehat{\mathbb{C}} \setminus p$) to obtain the desired conclusions. $\square$

6.3. Patching continuous extensions.

Corollary 6.13. *The map $H: \tilde{X} \rightarrow Y$ extends to a continuous, surjective, and monotone map $H: \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ such that $\pi_Y \circ H = h \circ \pi_X$ and $H^{-1}(\text{int}(q_i)) = \text{int}(p_i)$ for each $i \in \mathbb{N}$. The extension is constant on each continuum $E \subset \tilde{p}_i \setminus p_i$, $i \in \mathbb{N}$.*

Proof. We apply Corollary 6.12 so we obtain appropriate extensions of g_i for each $i \in \mathbb{N}$. For each $i \in \mathbb{N}$, we extend H to $\tilde{p}_i$ by setting $H = \pi_{Y;q_i}^{-1} \circ g_i$; note that $\pi_{Y;q_i}^{-1}$ is injective on $g_i(\tilde{p}_i) = \pi_{Y;q_i}(q_i)$. We observe that the relation $\pi_{Y;q_i} \circ H = g_i$ (which is already valid on $\tilde{X}$ by the definition of g_i) holds on all of $\hat{\mathbb{C}}$.

This extension is surjective onto $\hat{\mathbb{C}}$. Moreover, the preimage of each point is a continuum; this follows from the facts that the maps $H|_{\tilde{X}}: \tilde{X} \rightarrow Y$ and $g_i|_{\tilde{p}_i}: \tilde{p}_i \rightarrow q_i$ are cell-like. If $E \subset \tilde{p}_i \setminus p_i$ is a continuum, then $g_i|_E$ is constant. Thus, H is constant on E , as claimed. The relation $\pi_Y \circ H = h \circ \pi_X$, which holds on $\tilde{X}$ extends automatically to $\hat{\mathbb{C}}$, since $H(\tilde{p}_i) = q_i$ for each $i \in \mathbb{N}$.

We finally argue that H is continuous. Suppose first that $x \in \tilde{X}$. If $x_n \in \hat{\mathbb{C}}$ is a sequence converging to x , then $\pi_Y(H(x_n)) = h(\pi_X(x_n))$ converges to $h(\pi_X(x))$ by continuity. By Lemma 2.2, the point $H(x_n) \in \pi_Y^{-1}(\pi_Y(H(x_n)))$ is contained in arbitrarily small neighborhoods of $\pi_Y^{-1}(h(\pi_X(x))) = H(x)$ as $n \rightarrow \infty$. Next, suppose that $x \in \tilde{p}_i$ for some $i \in \mathbb{N}$ and consider a sequence $x_n \in \hat{\mathbb{C}}$ converging to x . The sequence $\pi_{Y;q_i}(H(x_n)) = g_i(x_n)$ converges by continuity to $g_i(x)$. Using again Lemma 2.2, we see that $H(x_n)$ is contained in arbitrarily small neighborhoods of $\pi_{Y;q_i}^{-1}(g_i(x)) = H(x)$ as $n \rightarrow \infty$. $\square$

Proof of Theorem 6.1. Consider the extension of H as in Corollary 6.13. It remains to show that H satisfies the transboundary upper gradient inequality, as stated in Theorem 6.1. Suppose that h , ρ_h , and Γ_0 are as in Definition 2.15 and let $\rho_H = \rho_h$. By enlarging the curve family Γ_0 , we may assume that it still has conformal modulus zero and all subcurves of every curve $\gamma \notin \Gamma_0$ also satisfy the transboundary upper gradient inequality (see Remark 6.6). We fix a curve $\gamma: [a, b] \rightarrow \hat{\mathbb{C}}$ outside Γ_0 .

If $\gamma(a)$ and $\gamma(b)$ lie in p_i for some $i \in \mathbb{N}$, then $H(\gamma(a)), H(\gamma(b)) \in q_i$, so $\sigma(H(\gamma(a)), H(\gamma(b))) \leq \text{diam}(q_i)$. In this case there is nothing to show.

Suppose that $\gamma(a)$ and $\gamma(b)$ do not lie on the same peripheral continuum. Then there exists an open subpath $\gamma_1 = \gamma|_{(a_1, b_1)}: (a_1, b_1) \rightarrow \hat{\mathbb{C}}$ of γ that does not intersect the peripheral continua that possibly contain $\gamma(a)$ or $\gamma(b)$, and the points $\gamma(a_1), \gamma(b_1)$ lie on the boundaries of the peripheral continua that possibly contain $\gamma(a), \gamma(b)$, respectively. It suffices to show the desired inequality for the open path γ_1 , in view of the previous case where the endpoints lie in the same peripheral continuum.

There exists a further subpath $\gamma_2 = \gamma|_{(a_2, b_2)}$ of γ_1 such that $\gamma((a_1, a_2])$ (resp. $\gamma([b_2, b_1))$) is contained in some set $\tilde{p}_i$, $i \in \mathbb{N}$, and $\gamma(a_2)$ (resp. $\gamma(b_2)$) can be approximated by points in $|\gamma_2| \cap H^{-1}(Y)$. Note that these conditions could be vacuously true and we could have $\gamma_2 = \gamma_1$. By Corollary 6.13, H is constant on $\gamma((a_1, a_2])$ and on $\gamma([b_2, b_1))$. Therefore, it suffices to show the desired inequality for the open path γ_2 , which has the property that its endpoints can be approximated by points of $|\gamma_2| \cap H^{-1}(Y)$.

We may find parameters $a_3 > a_2$ and $b_3 < b_2$ so that $\gamma(a_3), \gamma(b_3) \in H^{-1}(Y)$ and $\gamma(a_3), \gamma(b_3)$ are arbitrarily close to $\gamma(a_2), \gamma(b_2)$, respectively. Since π_Y is injective on Y , we have $H = \pi_Y^{-1} \circ h \circ \pi_X$ on $H^{-1}(Y)$. The transboundary upper gradient

inequality of h now gives

$$\sigma(H(\gamma(a_3)), H(\gamma(b_3))) \leq \int_{\gamma_2} \rho_H ds + \sum_{i: p_i \cap |\gamma_2| \neq \emptyset} \text{diam}(q_i).$$

By letting $\gamma(a_3) \rightarrow \gamma(a_2)$ and $\gamma(b_3) \rightarrow \gamma(b_2)$, using the continuity of H we obtain the same inequality for $\sigma(H(\gamma(a_2)), H(\gamma(b_2)))$. $\square$

Proof of Proposition 6.2. We fix $i \in \mathbb{N}$ and we use the same notational conventions as in the proof of Lemma 6.8. By Lemma 6.8, there exists a continuous, proper, and cell-like map g_i from Ω onto D such that

$$g_i = \pi_i \circ \pi_Y^{-1} \circ g = \pi_i \circ \pi_Y^{-1} \circ h \circ \pi_X.$$

The relation $\pi_Y \circ H = h \circ \pi_X$ implies that on the set $\tilde{X}$ we have $H = \pi_Y^{-1} \circ h \circ \pi_X$ and that $H(\tilde{p}_j) = q_j$ for each $j \in \mathbb{N}$. Thus, $g_i = \pi_i \circ H$ on all of Ω .

Note that the continuous, surjective, and monotone map $H: \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ is also cell-like (see the comments in Section 2.3). This fact and Lemma 2.9 (iv) imply that the map $\pi_i \circ H$ is continuous and cell-like from $\hat{\mathbb{C}}$ onto $\mathcal{E}(Y; q)$. This shows that g_i extends to a continuous and cell-like map from $\hat{\mathbb{C}}$ onto $\mathcal{E}(Y; q)$. Since $H^{-1}(\text{int}(q)) = \text{int}(p)$, we have $g_i^{-1}(\pi_i(\text{int}(q))) = \text{int}(p)$. Hence g_i is continuous and cell-like from $\hat{\mathbb{C}} \setminus \text{int}(p)$ onto $\mathcal{E}(Y; q) \setminus \pi_i(\text{int}(q))$. By Corollary 6.12, this extension is unique and has the property that it is constant on each continuum $E \subset \tilde{p} \setminus p$. The desired property of H follows. $\square$

7. HOMEOMORPHIC EXTENSION

Theorem 7.1. *Let X, Y be Sierpiński packings and $h: \mathcal{E}(X) \rightarrow \mathcal{E}(Y)$ be a packing- K -quasiconformal map for some $K \geq 1$ as in the statement of Theorem 6.1. If Y is cofat, then the map H of Theorem 6.1 is a homeomorphism from $\bar{X}$ onto $\bar{Y}$. In particular, it can be extended to a homeomorphism of $\hat{\mathbb{C}}$.*

Proof. Assume that H maps $\bar{X}$ homeomorphically onto $\bar{Y}$. From Theorem 6.1 we have $H^{-1}(\text{int}(q_i)) = \text{int}(p_i)$, which implies that $H(\partial p_i) = \partial q_i$. For each $i \in \mathbb{N}$ we replace $H|_{p_i}$ with a homeomorphism onto q_i to obtain a global homeomorphism of $\hat{\mathbb{C}}$; it is important here that $\text{diam}(p_i) \rightarrow 0$ and $\text{diam}(q_i) \rightarrow 0$ as $i \rightarrow \infty$.

We now show that for each point $w_0 \in \bar{Y}$ the set $H^{-1}(w_0) \subset \bar{X}$ is a singleton. Suppose that there exists $w_0 \in \bar{Y}$ such that $E = H^{-1}(w_0)$ is a non-degenerate continuum. Based on the following claim, which we prove afterwards, we complete the proof of Theorem 7.1. See Figure 7 for an illustration.

Claim 7.2. There exists $\delta > 0$, $r_0 > 0$, and a point $z_0 \in E$ such that for all $t \in [\delta/2, \delta]$ and for all $r \in (0, r_0)$ the circle $S(z_0, t)$ has an open arc $\beta_t: (a, b) \rightarrow \hat{\mathbb{C}}$ such that $\beta_t(a) \in E$, $\beta_t(b) \in \partial H^{-1}(B(w_0, r))$, and $\beta_t((a, b)) \subset H^{-1}(B(w_0, r)) \setminus E$. Moreover, if $w_0 \in q_j$ for some $j \in \mathbb{N}$, then $\beta_t((a, b)) \cap p_j = \emptyset$.

For fixed $r > 0$, let $I(r) = \{i \in \mathbb{N} : q_i \cap S(w_0, r) \neq \emptyset\}$. If $i \in I(r)$, we define $\lambda(q_i) = \mathcal{H}^1(\{s \in (0, r) : q_i \cap S(w_0, s) \neq \emptyset\})$ and otherwise we define $\lambda(q_i) = \text{diam}(q_i)$. By the fatness of q_i , $i \in \mathbb{N}$, and Lemma 2.6, we have

$$(7.1) \quad \lambda(q_i)^2 \leq C \Sigma(q_i \cap B(w_0, r))$$

whenever $q_i \cap B(w_0, r) \neq \emptyset$.




FIGURE 7. Left: the circle $S(z_0, t)$ and the arc β_t (red). Right: the path $H \circ \beta_t$ (red) and the subpath $H \circ \gamma_t$ (green) that connects w_0 to a point on $q_{i_0} \cap S(w_0, r_1)$.

Suppose that $H \circ \beta_t$ does not intersect any q_i , $i \in I(r)$. That is, if $q_i \cap |H \circ \beta_t| \neq \emptyset$, then $q_i \subset B(w_0, r)$. By Lemma 2.12, the transboundary upper gradient inequality holds for a.e. $t \in [\delta/2, \delta]$, so we have

$$r \leq \int_{\beta_t} \rho ds + \sum_{i: p_i \cap |\beta_t| \neq \emptyset} \text{diam}(q_i) = \int_{\beta_t} \rho ds + \sum_{i: p_i \cap |\beta_t| \neq \emptyset} \lambda(q_i).$$

If $H \circ \beta_t$ intersects some q_i , $i \in I(r)$, we consider an open subpath γ_t of β_t such that $H \circ \gamma_t$ does not intersect any q_i , $i \in I(r)$, and connects w_0 to a point of q_{i_0} for some $i_0 \in I(r)$ that lies on a circle $S(w_0, r_1)$; see Figure 7. By the transboundary upper gradient inequality, for a.e. $t \in [\delta/2, \delta]$ we have

$$r_1 \leq \int_{\gamma_t} \rho ds + \sum_{i: p_i \cap |\gamma_t| \neq \emptyset} \text{diam}(q_i) = \int_{\gamma_t} \rho ds + \sum_{i: p_i \cap |\gamma_t| \neq \emptyset} \lambda(q_i).$$

Since the continuum q_{i_0} intersects both circles $S(w_0, r_1)$ and $S(w_0, r)$, we have

$$r - r_1 \leq \mathcal{H}^1(\{s \in (0, r) : q_{i_0} \cap S(w_0, s) \neq \emptyset\}) = \lambda(q_{i_0}).$$

Altogether, we have

$$r \leq \int_{\beta_t} \rho ds + \sum_{i: p_i \cap |\beta_t| \neq \emptyset} \lambda(q_i).$$

Since β_t is contained in $S(z_0, t) \cap (H^{-1}(B(w_0, r)) \setminus E)$, for a.e. $t \in [\delta/2, \delta]$ and for all $r < r_0$ we have

$$(7.2) \quad r \leq \int_{S(z_0, t) \cap (H^{-1}(B(w_0, r)) \setminus E)} \rho d\mathcal{H}^1 + \sum_{\substack{i: p_i \cap S(z_0, t) \neq \emptyset \\ q_i \cap B(w_0, r) \neq \emptyset}} \lambda(q_i).$$

We integrate over $t \in [\delta/2, \delta]$ and then apply the co-area (Proposition 2.1) and Cauchy-Schwarz inequalities to obtain

$$\begin{aligned} (\delta/2)r &\leq C \int_{H^{-1}(B(w_0, r)) \setminus E} \rho d\Sigma + \sum_{i: q_i \cap B(w_0, r) \neq \emptyset} \lambda(q_i) \text{diam}(p_i) \\ &\leq C \left(\int_{H^{-1}(B(w_0, r)) \setminus E} \rho^2 d\Sigma + \sum_{i: q_i \cap B(w_0, r) \neq \emptyset} \lambda(q_i)^2 \right)^{1/2} \\ &\quad \cdot \left(\Sigma(H^{-1}(B(w_0, r)) \setminus E) + \sum_{i: q_i \cap B(w_0, r) \neq \emptyset} \text{diam}(p_i)^2 \right)^{1/2} \\ &=: C \cdot A(r) \cdot B(r). \end{aligned}$$

Suppose that $w_0 \notin \bigcup_{i \in \mathbb{N}} q_i$. As $r \rightarrow 0$, the term $B(r)$ converges to 0, since $H^{-1}(B(w_0, r))$ converges to E and $\{i : q_i \cap B(w_0, r) \neq \emptyset\}$ shrinks to the empty set. If $w_0 \in p_j$ for some $j \in \mathbb{N}$, then by the last part of Claim 7.2, β_t does not intersect p_j , so we may exclude the index j from the sum in (7.2). In this case, if we perform the same calculations as above based on the modified version of (7.2), we will obtain a term $B(r)$ such that the summation term does not include the index j . Therefore, $\{i : i \neq j, q_i \cap B(w_0, r) \neq \emptyset\}$ shrinks again to the empty set. This shows that $B(r) \rightarrow 0$ as $r \rightarrow 0$ also in this case.

We now discuss the first term, $A(r)$. By quasiconformality (see the last inequality in Theorem 6.1) we have

$$\int_{H^{-1}(B(w_0, r)) \setminus E} \rho^2 d\Sigma \lesssim \Sigma(B(w_0, r)) \lesssim r^2.$$

Moreover, (7.1) implies that

$$\sum_{i: q_i \cap B(w_0, r) \neq \emptyset} \lambda(q_i)^2 \leq C \sum_{i: q_i \cap B(w_0, r) \neq \emptyset} \Sigma(q_i \cap B(w_0, r)) \lesssim r^2.$$

Therefore, $A(r) \lesssim r$, so

$$\delta r \lesssim r B(r).$$

This is true for all $r < r_0$ so we obtain a contradiction as $r \rightarrow 0$. $\square$

Proof of Claim 7.2. We consider two cases:

- (a) E is not contained in p_i for any $i \in \mathbb{N}$.
- (b) $E \subset \partial p_j$ for some $j \in \mathbb{N}$.

We will treat now the first case. Note that $E = H^{-1}(w_0)$ can intersect at most one set p_i , $i \in \mathbb{N}$, which occurs only if $w_0 \in \partial q_i$. If $\partial E \subset p_i$, then ∂E is an arc of ∂p_i , so $\widehat{\mathbb{C}} \setminus \partial E$ has one or two components. Each of them lies in the interior or exterior of E . Hence, either $E = \partial E \subset \partial p_i$, which is a contradiction to Case (a), or

$E = \widehat{\mathbb{C}} \setminus \text{int}(p_i)$, which is again a contradiction as $E \cap p_j = \emptyset$ for $j \neq i$. Therefore, ∂E is not contained in p_i for any $i \in \mathbb{N}$, so $\partial E \cap X \neq \emptyset$.

Consider a point $z_0 \in \partial E \cap X$ and note that the circle $S(z_0, t)$ intersects E for all sufficiently small $t > 0$. Recall that H is monotone, so it is also cell-like (see Section 2.3). Therefore, $E = H^{-1}(w_0)$ is a non-separating continuum. This implies that the circle $S(z_0, t)$ intersects $\widehat{\mathbb{C}} \setminus E$ for all sufficiently small $t > 0$ (see the argument in the proof of Lemma 6.11). We fix $\delta > 0$ such that for all $t \in [\delta/2, \delta]$ the circle $S(z_0, t)$ intersects both E and $\widehat{\mathbb{C}} \setminus E$. Moreover, if $w_0 \in q_j$, the fact that $z_0 \in X$ allows us to choose an even smaller $\delta > 0$ so that $S(z_0, t)$ does not intersect p_j for all $t \in [\delta/2, \delta]$.

For $r > 0$ consider the ball $B(w_0, r)$ and the preimage $H^{-1}(B(w_0, r))$, which contains the set E in its interior. If $r_0 > 0$ is sufficiently small, then for $r < r_0$ the set $H^{-1}(B(w_0, r))$ does not contain any of the circles $S(z_0, t)$, $t \in [\delta/2, \delta]$; indeed, otherwise, by a limiting argument we see that there would exist $t \in [\delta/2, \delta]$ such that $S(z_0, t) \subset E$, a contradiction. We fix $t \in [\delta/2, \delta]$ and $r < r_0$. Then, the circle $S(z_0, t)$ intersects both E and $\partial H^{-1}(B(w_0, r))$. We consider an open arc $\beta_t: (a, b) \rightarrow \widehat{\mathbb{C}}$ of $S(z_0, t)$ such that $\beta_t(a) \in E$, $\beta_t(b) \in \partial H^{-1}(B(w_0, r))$, and $\beta_t((a, b)) \subset H^{-1}(B(w_0, r)) \setminus E$. See Figure 7. This completes the proof of Claim 7.2 in Case (a).

Suppose that $E \subset \partial p_j$ as in Case (b). Let $z_0 \in \partial p_j$ and fix $\delta_0 > 0$ such that E is not contained in $B(z_0, \delta_0)$. Consider a Jordan arc $\eta: (0, 1) \rightarrow \widehat{\mathbb{C}}$ such that $\eta(0) = z_0$, $\eta(1) \in E \setminus B(z_0, \delta_0)$, and $\eta((0, 1)) \cap p_j = \emptyset$. Let $\delta < \delta_0$. For each $t \in [\delta/2, \delta]$ there exists a unique open arc $\alpha_t \subset S(z_0, t)$ with endpoints on ∂p_j but otherwise disjoint from p_j such that $\eta(s_0) \in \alpha_t$ for some $s_0 \in (0, 1)$ and $\eta((0, s_0)) \cap S(z_0, t) = \emptyset$; in other words, α_t is the “first” arc of $S(z_0, t) \setminus p_j$ that η meets.

We claim that if δ is sufficiently small, then for $t \in [\delta/2, \delta]$ at least one endpoint of α_t lies in E . Consider the arc A_t of ∂p_j containing z_0 and having the same endpoints as α_t . By the local connectivity of ∂p_j , as $t \rightarrow 0$ the diameter of A_t tends to 0. Hence, E cannot be contained in A_t for small $t > 0$. The connectedness of E implies that one of the endpoints of A_t , and thus of α_t , is contained in E for all sufficiently small $t > 0$, as desired.

Next, we show that if $r_0 > 0$ is sufficiently small, then α_t is not contained in $H^{-1}(B(w_0, r))$ for all $r < r_0$ and for all $t \in [\delta/2, \delta]$. To see this, note that there exists an arc $\eta([s_1, s_2])$, disjoint from p_j , intersected by α_t for each $t \in [\delta/2, \delta]$. Since $H^{-1}(B(w_0, r))$ converges to $H^{-1}(w_0) = E$ as $r \rightarrow 0$, there exists $r_0 > 0$ such that $H^{-1}(B(w_0, r_0))$ is disjoint from $\eta([s_1, s_2])$. Therefore, α_t intersects the complement of $H^{-1}(B(w_0, r))$, for all $r < r_0$, as claimed.

Finally, we consider an open subarc $\beta_t: (a, b) \rightarrow \widehat{\mathbb{C}}$ of α_t such that $\beta_t(a) \in E$, $\beta_t(b) \in \partial H^{-1}(B(w_0, r))$, and $\beta_t((a, b)) \subset H^{-1}(B(w_0, r)) \setminus E$. This completes the proof in Case (b). $\square$

REFERENCES

- [BBI01] D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001.
- [Bon11] M. Bonk, *Uniformization of Sierpiński carpets in the plane*, Invent. Math. **186** (2011), no. 3, 559–665.
- [Bra80] M. Brandt, *Ein Abbildungssatz für endlich-vielfach zusammenhängende Gebiete*, Bull. Soc. Sci. Lett. Łódź **30** (1980), no. 3, 12.

- [Bro72] M. Brown, *Sets of constant distance from a planar set*, Michigan Math. J. **19** (1972), 321–323.
- [Dav86] R. J. Daverman, *Decompositions of manifolds*, Pure and Applied Mathematics, vol. 124, Academic Press, Inc., Orlando, FL, 1986.
- [EHa21] B. Esmayli and P. Hajlász, *The coarea inequality*, Ann. Fenn. Math. **46** (2021), no. 2, 965–991.
- [Gol69] G. M. Goluzin, *Geometric theory of functions of a complex variable*, Translations of Mathematical Monographs, vol. 26, American Mathematical Society, Providence, RI, 1969.
- [Har82] A. N. Harrington, *Conformal mappings onto domains with arbitrarily specified boundary shapes*, J. Analyse Math. **41** (1982), 39–53.
- [HKST15] J. Heinonen, P. Koskela, N. Shanmugalingam, and J. T. Tyson, *Sobolev spaces on metric measure spaces. An approach based on upper gradients*, New Mathematical Monographs, vol. 27, Cambridge University Press, Cambridge, 2015.
- [HL21] H. Hakobyan and W.-b. Li, *Quasisymmetric embeddings of slit Sierpiński carpets* (2021), preprint arXiv:1901.05632.
- [HS93] Z.-X. He and O. Schramm, *Fixed points, Koebe uniformization and circle packings*, Ann. of Math. (2) **137** (1993), no. 2, 369–406.
- [HS94] Z.-X. He and O. Schramm, *Rigidity of circle domains whose boundary has σ -finite linear measure*, Invent. Math. **115** (1994), no. 2, 297–310.
- [Kec95] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.
- [Koe08] P. Koebe, *Über die uniformisierung beliebiger analytischer kurven, III*, Nachr. Ges. Wiss. Gött. (1908), 337–358.
- [Koe20] P. Koebe, *Abhandlungen zur Theorie der konformen Abbildung*, Math. Z. **7** (1920), no. 1-4, 235–301.
- [MN22] S. Maio and D. Ntalampekos, *On the Hausdorff dimension of the residual set of a packing by smooth curves*, J. London Math. Soc. **105** (2022), no. 3, 1752–1786.
- [Moo25] R. L. Moore, *Concerning upper semi-continuous collections of continua*, Trans. Amer. Math. Soc. **27** (1925), no. 4, 416–428.
- [Moo28] R. L. Moore, *On the separation of the plane by a continuum*, Bull. Amer. Math. Soc. **34** (1928), no. 3, 303–306.
- [MW13] S. Merenkov and K. Wildrick, *Quasisymmetric Koebe uniformization*, Rev. Mat. Iberoam. **29** (2013), no. 3, 859–909.
- [NR22a] D. Ntalampekos and M. Romney, *Polyhedral approximation and uniformization for non-length surfaces* (2022), preprint arXiv:2206.01128.
- [NR22b] D. Ntalampekos and M. Romney, *Polyhedral approximation of metric surfaces and applications to uniformization*, Duke Math. J. (2022), to appear.
- [Nta20a] D. Ntalampekos, *Monotone Sobolev functions in planar domains: Level sets and smooth approximation*, Arch. Ration. Mech. Anal. **238** (2020), 1199–1230.
- [Nta20b] D. Ntalampekos, *Potential theory on Sierpiński carpets: With applications to uniformization*, Lecture Notes in Mathematics, vol. 2268, Springer, Cham, 2020.
- [Nta23a] D. Ntalampekos, *CNED sets: countably negligible for extremal distances* (2023), preprint arXiv:2303.13187.
- [Nta23b] D. Ntalampekos, *Metric definition of quasiconformality and exceptional sets* (2023), preprint.
- [Nta23c] D. Ntalampekos, *Rigidity and continuous extension for conformal maps of circle domains*, Trans. Amer. Math. Soc. **376** (2023), no. 7, 5221–5239.
- [NY20] D. Ntalampekos and M. Younsi, *Rigidity theorems for circle domains*, Invent. Math. **220** (2020), no. 1, 129–183.
- [Pom92] Ch. Pommerenke, *Boundary behaviour of conformal maps*, Grundlehren der Mathematischen Wissenschaften, vol. 299, Springer-Verlag, Berlin, 1992.
- [Raj21] K. Rajala, *Uniformization of planar domains by exhaustion* (2021), preprint arXiv:2111.00845.
- [Reh22] J. Rehmert, *Quasisymmetric koebe uniformization of metric surfaces with uniformly relatively separated boundary*, Ph.D. Thesis, 2022.
- [RR21] K. Rajala and M. Rasimus, *Quasisymmetric Koebe uniformization with weak metric doubling measures*, Illinois J. Math. **65** (2021), no. 4, 749–767.

- [RW15] S. Rohde and B. M. Werner, *Partial uniformization of CLE* (2015), available at http://www.math.stonybrook.edu/Videos/Colloquium/direct_download.php?file=PDFs/20150212-Werner.pdf. Accessed on August 1, 2022.
- [Sch95] O. Schramm, *Transboundary extremal length*, J. Anal. Math. **66** (1995), 307–329.
- [Sie72] L. C. Siebenmann, *Approximating cellular maps by homeomorphisms*, Topology **11** (1972), 271–294.
- [SW12] S. Sheffield and W. Werner, *Conformal loop ensembles: the Markovian characterization and the loop-soup construction*, Ann. of Math. (2) **176** (2012), no. 3, 1827–1917.
- [Väi71] J. Väisälä, *Lectures on n -dimensional quasiconformal mappings*, Lecture Notes in Mathematics, vol. 229, Springer-Verlag, Berlin-New York, 1971.
- [Why42] G. T. Whyburn, *Analytic Topology*, American Mathematical Society Colloquium Publications, vol. 28, American Mathematical Society, New York, 1942.
- [Why58] G. T. Whyburn, *Topological characterization of the Sierpiński curve*, Fund. Math. **45** (1958), 320–324.
- [Wil12] M. Williams, *Geometric and analytic quasiconformality in metric measure spaces*, Proc. Amer. Math. Soc. **140** (2012), no. 4, 1251–1266.
- [Wil70] S. Willard, *General topology*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970.
- [Wil] D. B. Wilson, *Simulation of CLE_4 gasket*, available at http://dbwilson.com/gallery/gasket-4-512_lowres.png. Accessed on September 1, 2022.
- [You48] J. W. T. Youngs, *Homeomorphic approximations to monotone mappings*, Duke Math. J. **15** (1948), 87–94.

MATHEMATICS DEPARTMENT, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794, USA.
 Email address: dimitrios.ntalampekos@stonybrook.edu