

K-STABILITY AND BIRATIONAL MODELS OF MODULI OF QUARTIC K3 SURFACES

KENNETH ASCHER, KRISTIN DEVLEMING, AND YUCHEN LIU

ABSTRACT. We show that the K-moduli spaces of log Fano pairs (\mathbb{P}^3, cS) where S is a quartic surface interpolate between the GIT moduli space of quartic surfaces and the Baily-Borel compactification of moduli of quartic K3 surfaces as c varies in the interval $(0, 1)$. We completely describe the wall crossings of these K-moduli spaces. As the main application, we verify Laza-O’Grady’s prediction on the Hassett-Keel-Looijenga program for quartic K3 surfaces. We also obtain the K-moduli compactification of quartic double solids, and classify all Gorenstein canonical Fano degenerations of \mathbb{P}^3 .

CONTENTS

1. Introduction	1
2. Preliminaries on K-stability and K-moduli	6
3. Geometry and moduli of quartic K3 surfaces	12
4. Tangent developable surface and unigonal K3 surfaces	16
5. Hyperelliptic K3 surfaces	31
6. Proof of main theorems	42
References	53

1. INTRODUCTION

An important question in algebraic geometry is to construct geometrically meaningful compact moduli spaces for polarized K3 surfaces. The global Torelli theorem indicates that the coarse moduli space \mathfrak{M}_{2d} of primitively polarized K3 surfaces with du Val singularities of degree $2d$ is isomorphic, under the period map, to the arithmetic quotient $\mathcal{F}_{2d} = \mathcal{D}_{2d}/\Gamma_{2d}$ of a Type IV Hermitian symmetric domain \mathcal{D}_{2d} as the period domain. The space \mathcal{F}_{2d} has a natural Baily-Borel compactification \mathcal{F}_{2d}^* , but it is well-known that \mathcal{F}_{2d}^* does not carry a nicely behaved universal family. Thus it is a natural problem to compare \mathcal{F}_{2d}^* with other geometric compactifications, e.g. those coming from geometric invariant theory (GIT), via the period map.

In particular, it is natural to ask if there exists a modular way to resolve the (birational) period map. When the degree $2d = 2$, there is a birational period map between the GIT quotient of sextic plane curves and \mathcal{F}_2^* , since a generic such K3 is the double cover of \mathbb{P}^2 ramified along a sextic. By work of Looijenga and Shah, this map can be resolved by considering either a partial Kirwan desingularization of the GIT quotient, or via a small partial resolution of \mathcal{F}_2^* [Sha80, Loo86]. A realization of Laza-O’Grady (based on work of Looijenga), is that an alternate systematic approach to this problem is via interpolating the Proj of $R(\mathcal{F}_2, \lambda + \beta\Delta)$, where λ is the Hodge line bundle on \mathcal{F}_2 , the divisor Δ is some geometrically meaningful Heegner divisor, and β varies between 0 and 1 (see e.g. [LO18]).

Date: November 14, 2022.

When the degree $2d = 4$ (for simplicity, denoted by $\mathfrak{M} = \mathfrak{M}_4$ and $\mathcal{F} = \mathcal{F}_4$), a distinguished geometric compactification is given by the GIT moduli space $\overline{\mathfrak{M}}^{\text{GIT}}$ of quartic surfaces in \mathbb{P}^3 . There is a birational period map $\mathfrak{p} : \overline{\mathfrak{M}}^{\text{GIT}} \dashrightarrow \mathcal{F}^*$ with much more complicated exceptional loci as compared to the degree two case. In a series of papers [LO19, LO18, LO21], Laza and O’Grady proposed a systematic way to resolve such period maps (when \mathcal{F} is an arithmetic quotient of a Type IV Hermitian symmetric domain) via a sequence of explicit birational transformations governed by the Heegner divisors in \mathcal{F}^* , and predict that they satisfy a natural interpolation. Motivated by the Hassett-Keel program – running the log minimal model program on \overline{M}_g to interpolate between different birational models of the moduli space of curves (see e.g. [HH13]), they named this program the *Hassett-Keel-Looijenga program*. In [LO21], Laza and O’Grady verified their proposal for the moduli of hyperelliptic quartic K3 surfaces, which is an 18-dimensional divisor in \mathcal{F}^* , but their prediction for the 19-dimensional space \mathcal{F} has remained open. One of the main purposes of this paper is to completely verify their prediction for \mathcal{F} using the recently constructed moduli spaces of log Fano pairs from the theory of K-stability. We note that the analogous question in the case of EPW sextics remains open, and it would be interesting to try to use K-moduli to study their compactifications.

For a rational number $c \in (0, 1)$, the pair (\mathbb{P}^3, cS) is a log Fano pair, where $S \subset \mathbb{P}^3$ is a smooth quartic surface. Thus, K-stability provides a natural framework to construct geometrically meaningful compactifications of moduli of quartic K3 surfaces. In recent years, the algebro-geometric theory of constructing projective K-moduli spaces of log Fano pairs has been completed as a combination of the important works [Jia20, LWX21, CP21, BX19, ABHLX20, BLX22, Xu20, XZ20, XZ21, BHLLX21, LXZ22]. Meanwhile, when we vary the coefficient c , the K-moduli spaces of \mathbb{Q} -Gorenstein smoothable log Fano pairs display wall-crossing phenomena as established in [ADL19] (see also [Zho21b]).

In this paper, we show that the K-moduli compactifications of log Fano pairs (\mathbb{P}^3, cS) where S is a smooth quartic surface interpolate naturally between the GIT moduli space $\overline{\mathfrak{M}}^{\text{GIT}}$ and the Baily-Borel compactification \mathcal{F}^* as c varies in the interval $(0, 1)$. As a result, we resolve the period map $\mathfrak{p} : \overline{\mathfrak{M}}^{\text{GIT}} \dashrightarrow \mathcal{F}^*$ where all intermediate birational models have a modular meaning as they parametrize certain K-polystable log Fano pairs. Furthermore, using the positivity of the log CM line bundle [CP21, Pos22, XZ20], we confirm the prediction by Laza and O’Grady on the Hassett-Keel-Looijenga program for moduli space of quartic K3 surfaces [LO18, LO19].

We first fix some notation. Let \mathfrak{M}° and \mathfrak{M}° be the Deligne-Mumford stack and coarse moduli space of quartic surfaces $S \subset \mathbb{P}^3$ with du Val singularities, respectively. Let \mathcal{F} be the locally symmetric variety parametrizing periods of all polarized K3 surfaces of degree 4 with du Val singularities. The global Torelli theorem implies that the period map $\mathfrak{p} : \mathfrak{M}^\circ \hookrightarrow \mathcal{F}$ is an open immersion of quasi-projective varieties. Let \mathcal{F}^* be the Baily-Borel compactification of \mathcal{F} . Let λ be the Hodge line bundle over \mathcal{F} . By [LO19], there are two Heegner divisors H_h and H_u of \mathcal{F} , which parametrize hyperelliptic and unigonal quartic K3 surfaces respectively, such that $\mathfrak{p}(\mathfrak{M}^\circ) = \mathcal{F} \setminus (H_h \cup H_u)$. Let $\overline{\mathfrak{M}}^{\text{GIT}}$ and $\overline{\mathfrak{M}}^{\text{GIT}}$ be the GIT moduli stack and space of quartic surfaces in \mathbb{P}^3 , respectively.

Theorem 1.1. *For $c \in (0, 1) \cap \mathbb{Q}$, let $\overline{\mathfrak{M}}_c^K$ (resp. $\overline{\mathfrak{M}}_c^K$) be the K-moduli stack (resp. K-moduli space) parametrizing K-semistable (resp. K-polystable) log Fano pairs (X, cD) admitting a \mathbb{Q} -Gorenstein smoothing to (\mathbb{P}^3, cS) where S is a quartic surface.*

- (1) *For any $c \in (0, \frac{1}{3}) \cap \mathbb{Q}$, there are isomorphisms $\overline{\mathfrak{M}}_c^K \cong \overline{\mathfrak{M}}^{\text{GIT}}$ and $\overline{\mathfrak{M}}_c^K \cong \overline{\mathfrak{M}}^{\text{GIT}}$.*
- (2) *For any $c \in (0, 1) \cap \mathbb{Q}$, the section ring $R(\mathcal{F}, c\lambda + (1 - c)\Delta^K)$ is finitely generated where $\Delta^K := \frac{1}{4}H_h + \frac{9}{8}H_u$. Moreover, there is an isomorphism $\overline{\mathfrak{M}}_c^K \cong \text{Proj } R(\mathcal{F}, c\lambda + (1 - c)\Delta^K)$*

where the log CM line bundle on $\overline{\mathcal{M}}_c^K$ is proportional to $\mathcal{O}(1)$ on the Proj up to a positive constant.

- (3) For $0 < \epsilon \ll 1$, the K -moduli space $\overline{\mathcal{M}}_{1-\epsilon}^K$ is isomorphic to Looijenga's \mathbb{Q} -Cartierization $\widehat{\mathcal{F}}$ of \mathcal{F}^* associated to H_h and H_u . Moreover, the Hodge line bundle on $\overline{\mathcal{M}}_{1-\epsilon}^K$ is semiample and its ample model is isomorphic to \mathcal{F}^* .
- (4) There are 9 K -moduli walls for $c \in (0, 1)$. Among them, 2 walls are divisorial contractions: contracting a strict transform of H_h to the double quadric surface $[2Q]$ when $c = \frac{1}{3}$, and contracting a strict transform of H_u to the tangent developable surface $[T]$ of a twisted cubic curve when $c = \frac{9}{13}$, respectively. The remaining 7 walls are flips.

For a detailed description of the K -moduli wall crossings, see Theorem 5.16.

That is, by varying the coefficient c , the K -moduli spaces $\overline{\mathcal{M}}_c^K$ provide a natural interpolation between the GIT quotient for quartic surfaces and the Baily-Borel compactification, and explicitly resolve the period map. We note that a special case of Theorem 1.1(1) was proved earlier in [GMGS21, Theorem 1.2] and [ADL19, Theorem 1.4] (see also [Zho21a]). We also note that the two walls which are divisorial contractions are actually weighted blowups of Kirwan type (see Remarks 5.17 and 6.10).

Since these K -moduli spaces $\overline{\mathcal{M}}_c^K$ provide birational models of \mathcal{F} , we are able to confirm Laza-O'Grady's prediction on the Hassett-Keel-Looijenga program for \mathcal{F} [LO19, LO18] by modifying $\overline{\mathcal{M}}_c^K$ and checking ampleness of Laza-O'Grady's line bundle. Indeed, we prove a more general finite generation result and describe a wall-chamber structure for the full-dimensional subcone of $N_{\mathbb{R}}^1(\mathcal{F})$ generated by λ , H_h , and H_u .

Theorem 1.2. *For any $a, b \in \mathbb{Q}_{>0}$, the section ring $R(\mathcal{F}, \lambda + \frac{1}{2}(aH_h + bH_u))$ is finitely generated, which yields a projective birational model $\mathcal{F}(a, b) := \text{Proj } R(\mathcal{F}, \lambda + \frac{1}{2}(aH_h + bH_u))$ of \mathcal{F} . These $\mathcal{F}(a, b)$'s have a wall-chamber structure where the walls are $a = a_i$ or $b = 1$ with*

$$(a_1, a_2, \dots, a_8) = \left(\frac{1}{9}, \frac{1}{7}, \frac{1}{6}, \frac{1}{5}, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1 \right).$$

Moreover, we have the following description of $\mathcal{F}(a, b)$. Here we assume $0 < \epsilon \ll 1$.

- (1) If $a \in (0, \frac{1}{9})$ and $b \in (0, 1)$, then $\mathcal{F}(a, b) \cong \widehat{\mathcal{F}}$.
- (2) If $a, b \in [1, +\infty)$, then $\mathcal{F}(a, b) \cong \overline{\mathcal{M}}^{\text{GIT}}$.
- (3) The birational map $\mathcal{F}(1-\epsilon, b) \rightarrow \mathcal{F}(1, b)$ is a divisorial contraction of the strict transform of H_h to a point, and $\mathcal{F}(1, b) \cong \mathcal{F}(a, b)$ for any $a > 1$.
- (4) The birational map $\mathcal{F}(a, 1-\epsilon) \rightarrow \mathcal{F}(a, 1)$ is a divisorial contraction of the strict transform of H_u to a point, and $\mathcal{F}(a, 1) \cong \mathcal{F}(a, b)$ for any $b > 1$.
- (5) If $1 \leq i \leq 7$, then birational maps $\mathcal{F}(a_i - \epsilon, b) \rightarrow \mathcal{F}(a_i, b) \leftarrow \mathcal{F}(a_i + \epsilon, b)$ form a flip whose flipping locus (resp. flipped locus) is the strict transform of Z^j (resp. of W_{j-1}) where $j = \begin{cases} 9 - i & \text{if } i \geq 4; \\ 10 - i & \text{if } i \leq 3. \end{cases}$ Here $Z^j \subset \mathcal{F}$ is a tower of Shimura subvarieties of codimension j (see (3.2)), and $W_i \subset \overline{\mathcal{M}}^{\text{GIT}}$ is a tower of i -dimensional subvarieties (see (3.1)).

Corollary 1.3. *Laza-O'Grady's prediction for the 19-dimensional locally symmetric variety \mathcal{F} [LO19, Prediction 5.1.1] holds.*

We note that partial results toward Laza-O'Grady's prediction were obtained in [LO19, LO18]. In [LO21] the 18-dimensional case of their prediction was confirmed (see also [ADL20] for a different approach). In [LO21] the authors used an intricate and subtle variation of GIT argument, motivated by their previous arithmetic and hodge theoretic computations in [LO19].

As a consequence of Theorem 1.1, we give an explicit description of the K-moduli space of quartic double solids, i.e. del Pezzo threefolds of degree 2. The smooth quartic double solids are previously known to be K-stable [Der16]. Note that this K-moduli space displays similar behavior to the K-moduli space of del Pezzo surfaces of degree 1 [OSS16] as both are two-step birational modifications (a blow-up followed by a small contraction) of GIT moduli spaces, while K-moduli spaces of del Pezzo threefolds/fourfolds of degree 3 or 4 are identical to GIT moduli spaces [SS17, LX19, Liu22].

Theorem 1.4. *Let $\bar{\mathfrak{Y}}$ be the K-moduli space of quartic double solids. Then the seminormalization of $\bar{\mathfrak{Y}}$ is isomorphic to $\bar{\mathfrak{M}}_2^K$. Moreover, it fits into the following diagram*

$$\bar{\mathfrak{M}}^{\text{GIT}} \xleftarrow{\rho} \widehat{\mathfrak{M}}^{\text{GIT}} \xrightarrow{\psi} \bar{\mathfrak{M}}_2^K \xrightarrow{\iota} \bar{\mathfrak{Y}}$$

where ρ is a divisorial contraction of a birational transform of H_h to the point parametrizing the double quadric surface [2Q], ψ is a small contraction of a rational curve (the strict transform of W_1) to a point p , and ι is the seminormalization obtained by taking fiberwise double covers, where $\iota(p)$ represents the toric \mathbb{Q} -Fano threefold $(x_2^4 = x_3x_4) \subset \mathbb{P}(1, 1, 2, 4, 4)_{[x_0, \dots, x_4]}$.

Another interesting consequence is a classification of all Gorenstein canonical Fano degenerations of \mathbb{P}^3 . Here X_h is the projective anti-canonical cone over $\mathbb{P}^1 \times \mathbb{P}^1$, and X_u is a Gorenstein \mathbb{Q} -Fano threefold constructed in Section 4.2. Their notation is chosen so that X_h (resp. X_u) contains a general hyperelliptic (resp. unigonal) quartic K3 surface as its anti-canonical divisor.

Theorem 1.5. *Let X be a Gorenstein canonical Fano variety that admits a \mathbb{Q} -Gorenstein smoothing to \mathbb{P}^3 . Then X is isomorphic to \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$, or X_u .*

Sketch of proofs. We sketch the proofs of Theorems 1.1 and 1.2. First of all, by [GMGS21, ADL19] we know that $\bar{\mathfrak{M}}_c^K \cong \bar{\mathfrak{M}}^{\text{GIT}}$. If S is a quartic surface in \mathbb{P}^3 with semi-log canonical (slc) singularities (also called insignificant limit singularities), then (\mathbb{P}^3, S) is a K-semistable log Calabi-Yau pair, and \mathbb{P}^3 is K-polystable. Hence by interpolation of K-stability, the log Fano pair (\mathbb{P}^3, cS) is K-semistable for any $0 < c < 1$. Therefore, the birational map $\bar{\mathfrak{M}}_c^K \dashrightarrow \bar{\mathfrak{M}}^{\text{GIT}}$ is isomorphic over the open subset $\mathfrak{M}^{\text{slc}}$ parametrizing quartic surfaces with slc singularities. Thus in order to describe wall-crossings of $\bar{\mathfrak{M}}_c^K$, we only need to understand the K-polystable replacements of $\bar{\mathfrak{M}}^{\text{GIT}} \setminus \mathfrak{M}^{\text{slc}}$ parametrizing quartic surfaces with significant limit singularities. From the GIT of quartic surfaces [Sha81, LO18], we know that $\bar{\mathfrak{M}}^{\text{GIT}} \setminus \mathfrak{M}^{\text{slc}} = W_8 \sqcup \{[T]\}$ where T is the tangent developable surface of a twisted cubic curve, and W_8 is the largest subvariety of $\bar{\mathfrak{M}}^{\text{GIT}}$ in the tower W_i (see (3.1)). Indeed, the K-polystable replacements of $[T]$ (resp. W_i) precisely correspond to unigonal (resp. hyperelliptic) quartic K3 surfaces.

In Section 4, we study the K-stability of (\mathbb{P}^3, cT) and its K-polystable replacements. Using equivariant K-stability from [Zhu21], we show that the K-semistable threshold of (\mathbb{P}^3, T) , i.e. the largest c where (\mathbb{P}^3, cT) is K-semistable, is equal to $\frac{9}{13}$. Then we construct the K-polystable replacement $(X_u, \frac{9}{13}T_0)$ of $(\mathbb{P}^3, \frac{9}{13}T)$ by explicit birational geometry. Here X_u is constructed as a particular Gorenstein \mathbb{Q} -Fano threefold that contains all unigonal quartic K3 surfaces as anti-canonical divisors (see Section 4.2). Then using Paul-Tian criterion type arguments and the deformation theory of Gorenstein toric threefold singularities [Alt97], we show that the K-moduli wall crossing at $c = \frac{9}{13}$ near $[T]$ is a divisorial contraction whose exceptional divisor, birational to H_u , is the GIT moduli space of (X_u, S) where S is a Weierstrass elliptic surface.

In Section 5, we study the K-polystable replacements of the tower W_i in $\bar{\mathfrak{M}}^{\text{GIT}}$. This is the trickiest part of the proof. Our motivation comes from [ADL20] where we show that the K-moduli compactification \bar{K}_c of $(\mathbb{P}^1 \times \mathbb{P}^1, cC)$ where $C \in |\mathcal{O}(4, 4)|$ is identical to the VGIT

moduli space of slope $t = \frac{3c}{2c+2}$. By taking fiberwise double covers, we obtain a family of K-moduli spaces birational to H_h . However, these K-moduli spaces parametrize surface pairs rather than threefold pairs. Nevertheless, we notice that a hyperelliptic quartic K3 surface S as a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ (resp. of $\mathbb{P}(1, 1, 2)$) naturally embeds into the cone X_h (resp. $\mathbb{P}(1, 1, 2, 4)$) as an anti-canonical divisor. Then using a cone construction, a covering trick, and interpolation (see Section 5.1 for more details), we show that $\overline{K}_{\frac{3c-1}{4}}$ admits a closed embedding into $\overline{\mathcal{M}}_c^K$ for $c > \frac{1}{3}$ whose image $H_{h,c}$ is a birational transform of H_h (see Theorem 5.9). Then we construct K-polystable replacements of W_i by first embedding all of \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$ into $\mathbb{P}(1^4, 2)$ as weighted hypersurfaces of degree two, and then finding a particular 1-PS (coming from VGIT in [LO21]) that degenerates (\mathbb{P}^3, cS) to a K-polystable pair in $H_{h,c}$ (see Theorem 5.12). Then we use deformation theory to classify exceptional loci after the walls. In particular, all K-moduli spaces $\overline{\mathcal{M}}_c^K$ are isomorphic outside of the loci $H_{h,c}$ and $H_{u,c}$. We give a complete description of all wall-crossings of $\overline{\mathcal{M}}_c^K$ in Theorem 5.16.

Finally, in Section 6 we prove the main theorems. We observe that $\mathcal{F} \dashrightarrow \overline{\mathcal{M}}_c^K$ is a birational contraction by Theorem 5.16. The upshot to show $\overline{\mathcal{M}}_c^K \cong \text{Proj } R(\mathcal{F}, c\lambda + (1-c)\Delta^K)$ is to use ampleness of log CM line bundles [XZ20], and to compute the variation of log CM line bundles which interpolate between the Hodge line bundle and the absolute CM line bundle (see (6.1)). Then we perform necessary gluing operations and birational modifications on $\overline{\mathcal{M}}_c^K$ to obtain $\mathcal{F}(a, b)$ (see Definition 6.4 and Proposition 6.5), and show that the pushforward of $\lambda + \frac{a}{2}H_h + \frac{b}{2}H_u$ is ample.

Prior and related works. Compactifying the moduli space \mathcal{F}_2 of degree 2 K3 surfaces is a well studied problem. Recall that a general K3 surface of degree two can be realized as a double cover of \mathbb{P}^2 branched along a sextic curve. As such, there is a natural birational period map between the GIT quotient of plane sextics and the Baily-Borel compactification. Shah [Sha80] constructed a partial Kirwan desingularization of the GIT quotient which provides a compactification of \mathcal{F}_2 with a set-theoretic map to \mathcal{F}_2^* . Work of Looijenga [Loo86, Loo03b] shows that Shah's compactification is a \mathbb{Q} -factorialization of \mathcal{F}_2^* and additionally resolves the birational period map. In fact, this case serves as a major motivation for the Hassett-Keel-Looijenga (HKL) program. The case of degree 2 was revisited, in terms of Kollar-Shepherd-Barron (KSB) stable pairs, by Laza [Laz16], and more recently studied from the viewpoint of toroidal compactifications in work of Alexeev-Engel-Thompson [AET19] (see also the more recent [AE21]).

As mentioned above, the Hassett-Keel-Looijenga program was proposed by Laza-O'Grady [LO19] for Type IV locally symmetric varieties associated to the lattice $U^2 \oplus D_{N-2}$. It has been verified in the case of $N = 18$ for hyperelliptic quartic K3 surfaces by Laza-O'Grady using variation of GIT [LO21], and partial results for $N = 19$, i.e. moduli of quartic K3 surfaces were obtained in [LO19, LO18].

The wall-crossing phenomenon for K-moduli spaces of log Fano pairs with varying coefficients was systematically investigated in [ADL19]. One novelty of this strategy is to naturally connect well-studied moduli spaces, such as GIT, moduli of curves, and K3 surfaces, through birational maps between a sequence of K-moduli spaces. In our previous works [ADL19, ADL20], we carried out this strategy for the Hassett-Keel-Looijenga program for \mathcal{F}_2 and the hyperelliptic Heegner divisor H_h of \mathcal{F}_4 . In this paper, we use this novel approach of wall-crossing for K-moduli to solve the problem of Laza-O'Grady (HKL program for \mathcal{F}_4). One of the key benefits of this approach is that it gives a direct solution to a problem which was posed from an entirely different point of view, namely considering the Proj of a ring of automorphic forms and interpolating based on some arithmetic predictions.

Finally, we mention that moduli of pairs (\mathbb{P}^3, cH) have been studied from the point of view of KSB stable pairs by DeVleming [DeV22] where H is a surface in \mathbb{P}^3 of degree $d \geq 5$.

Acknowledgements. We would like to thank Dori Bejleri, Justin Lacini, Zhiyuan Li, Andrea Petracci, David Stapleton, Xiaowei Wang, and Chenyang Xu for helpful discussions, and Yuji Odaka for useful comments. We thank the referee for their helpful comments and suggestions. The authors were supported in part by the American Institute of Mathematics as part of the AIM SQuaREs program. Research of KA was supported in part by the NSF grant DMS-2140781 (formerly DMS-2001408). Research of YL was supported in part by the NSF grant DMS-2148266 (formerly DMS-2001317).

2. PRELIMINARIES ON K-STABILITY AND K-MODULI

Throughout this paper, we work over the field of complex numbers \mathbb{C} . We refer the background of singularities of pairs, such as Kawamata log terminal (klt), purely log terminal (plt), log canonical (lc), and semi-log canonical (slc), to the standard references [KM98, Kol13]. All schemes are assumed to be of finite type over \mathbb{C} .

2.1. K-stability.

2.1.1. Fujita-Li's valuative criteria.

Definition 2.1. Let X be a normal variety. Let D be an effective \mathbb{Q} -divisor on X . We say (X, D) is a *log pair* if $K_X + D$ is \mathbb{Q} -Cartier. A log pair (X, D) is called *log Fano* if X is projective and $-K_X - D$ is ample. A log pair (X, D) is called *log Calabi-Yau* if X is projective and $K_X + D \sim_{\mathbb{Q}} 0$. If $(X, 0)$ is a klt log Fano pair, then we call X a *\mathbb{Q} -Fano variety*.

We refer the definitions of test configurations and K-(poly/semi)stability of log Fano pairs to [ADL19, Section 2.1]. Here we use Fujita-Li's valuative criteria as alternative definitions.

Definition 2.2. Let (X, D) be a log pair. We call E a *prime divisor over X* if there is a proper birational morphism $\mu : Y \rightarrow X$ from a normal variety Y such that E is a prime divisor on Y . We define the *log discrepancy* of E with respect to (X, D) as

$$A_{(X,D)}(E) := 1 + \text{coeff}_E(K_Y - \mu^*(K_X + D)).$$

If, in addition, (X, D) is a log Fano pair, then we define the *pseudo-effective threshold*, the *expected vanishing order* (also known as the *S-functional*), and the β -*invariant* of E with respect to (X, D) as

$$\begin{aligned} T_{(X,D)}(E) &:= \sup\{t \in \mathbb{R}_{\geq 0} \mid \mu^*(-K_X - D) - tE \text{ is big}\}, \\ S_{(X,D)}(E) &:= \frac{1}{\text{vol}_X(-K_X - D)} \int_0^{T_{(X,D)}(E)} \text{vol}_X(-K_X - D - tE) dt, \\ \beta_{(X,D)}(E) &:= A_{(X,D)}(E) - S_{(X,D)}(E). \end{aligned}$$

Here $\text{vol}_X(-K_X - D - tE) := \text{vol}_Y(\mu^*(-K_X - D) - tE)$. The α -*invariant* and the *stability threshold* (also known as the δ -*invariant*) of a klt log Fano pair (X, D) are defined as

$$\alpha(X, D) := \inf_E \frac{A_{(X,D)}(E)}{T_{(X,D)}(E)} \quad \text{and} \quad \delta(X, D) := \inf_E \frac{A_{(X,D)}(E)}{S_{(X,D)}(E)},$$

where the infima run over all prime divisors E over X .

Next, we recall Fujita-Li's valuative criteria for K-(semi)stability and uniform K-stability.

Theorem 2.3 ([Fuj19c, Li17, BX19], see also [FO18, BJ20]). *A klt log Fano pair (X, D) is*

- (1) K -semistable if and only if $\beta_{(X,D)}(E) \geq 0$ for any prime divisor E over X , or equivalently, $\delta(X, D) \geq 1$;
- (2) K -stable if and only if $\beta_{(X,D)}(E) > 0$ for any prime divisor E over X ;
- (3) uniformly K -stable if and only if $\delta(X, D) > 1$.

Note that a K -semistable log Fano pair is always klt by [Oda13]. By a recent result of Liu-Xu-Zhuang [LXZ22], K -stability is equivalent to uniform K -stability for any klt log Fano pair.

If X is a \mathbb{Q} -Fano variety, D is an effective \mathbb{Q} -Cartier \mathbb{Q} -divisor on X , and $c \in \mathbb{Q}_{>0}$, then we say (X, D) is c - K -(poly/semi)stable if (X, cD) is a K -(poly/semi)stable log Fano pair.

2.1.2. Special degenerations and plt blow-ups. Recall that a test configuration $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ of a klt log Fano pair (X, D) is called *special* if $(\mathcal{X}, \mathcal{X}_0 + \mathcal{D})$ is plt and $\mathcal{L} \sim_{\mathbb{Q}} -l(K_{\mathcal{X}/\mathbb{A}^1} + \mathcal{D})$ for some $l \in \mathbb{Z}_{>0}$. In this case, we call $(\mathcal{X}_0, \mathcal{D}_0)$ a *special degeneration* of (X, D) and denote by $(X, D) \rightsquigarrow (\mathcal{X}_0, \mathcal{D}_0)$. By adjunction we know that $(\mathcal{X}_0, \mathcal{D}_0)$ is also a klt log Fano pair.

Next, we recall a result of Li-Wang-Xu which gives a characterization for K -polystability in terms of special degenerations.

Theorem 2.4 ([LWX21]). *A K -semistable log Fano pair (X, D) is K -polystable if and only if any K -semistable special degeneration of (X, D) is isomorphic to itself.*

Definition 2.5. Let (X, D) be a klt log pair. Let E be a prime divisor over X .

- (1) ([Fuj19b]) We say E is of *plt type* over (X, D) if there exists a birational morphism $\mu : Y \rightarrow X$ from a normal projective variety Y such that
 - E is a \mathbb{Q} -Cartier prime divisor on Y , and $-E$ is μ -ample;
 - $\mu|_{Y \setminus E} : Y \setminus E \rightarrow X \setminus \mu(E)$ is an isomorphism;
 - $(Y, E + \mu_*^{-1}D)$ is plt.

Such a morphism μ is called a *plt blow-up*.

- (2) [Pro00, Xu14] A plt type divisor E over (X, D) with center $\mu(E) = x$ being a closed point is called a *Kollar component* over the singularity $x \in (X, D)$.
- (3) We say E is a *special divisor* over (X, D) if there exists a special test configuration $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ of (X, D) and a positive integer d such that $\text{ord}_{\mathcal{X}_0}|_{\mathbb{C}(X)} = d \cdot \text{ord}_E$.

Lemma 2.6 (Zhuang). *Any special divisor over a klt log Fano pair (X, D) is of plt type.*

Proof. By Zhuang's Theorem [Xu21, Theorem 4.12], if E is a special divisor over (X, D) , then there exists a \mathbb{Q} -complement D^+ of (X, D) such that E is the only lc place of (X, D^+) . Thus for $0 < \epsilon \ll 1$, the log pair $(X, (1 - \epsilon)D + \epsilon D^+)$ is klt where E has log discrepancy less than 1. Thus by [BCHM10] there exists a birational morphism $\mu : Y \rightarrow X$ from a normal projective variety Y such that the first two conditions in Definition 2.5(2) hold. Moreover, since E is the only lc place of (X, D^+) , we know that $(Y, E + \mu_*^{-1}D^+)$ is plt. Thus $(Y, E + \mu_*^{-1}D)$ is also plt. \square

2.1.3. Equivariant K -stability. The following theorem is essentially due to Zhuang [Zhu21]. The equivalence between (i) and (iii) was also proved by Fujita [Fuj19b] when G is trivial.

Theorem 2.7 (Zhuang). *Let (X, D) be a klt log Fano pair. Let G be an algebraic group acting on (X, D) . Then the following are equivalent.*

- (i) (X, D) is K -semistable;
- (ii) $\beta_{(X,D)}(E) \geq 0$ for any G -invariant special divisor E over (X, D) ;
- (iii) $\beta_{(X,D)}(E) \geq 0$ for any G -invariant prime divisor E of plt type over (X, D) .

Proof. The (i) \Rightarrow (iii) part is a consequence of Theorem 2.3. The (iii) \Rightarrow (ii) part follows from Lemma 2.6. So we focus on the (ii) \Rightarrow (i) part. Assume to the contrary that (X, D) is K -unstable. By [LXZ22, Theorem 1.2] and [BHLLX21, Theorem 1.2(2)], there exists a non-trivial special test

configuration $(\mathcal{X}, \mathcal{D})$ of (X, D) that minimizes the bi-valued invariant $\left(\frac{\text{Fut}(\mathcal{X}, \mathcal{D})}{\|\mathcal{X}, \mathcal{D}\|_m}, \frac{\text{Fut}(\mathcal{X}, \mathcal{D})}{\|\mathcal{X}, \mathcal{D}\|_2}\right)$ under the lexicographic order among all special test configurations, where $\|\cdot\|_m$ and $\|\cdot\|_2$ represents the minimum norm and the L^2 -norm respectively (see [BHLLX21, Section 2.3] for definitions). Moreover, such a minimizing special test configuration $(\mathcal{X}, \mathcal{D})$ is unique up to rescaling. The minimizing property implies $\text{Fut}(\mathcal{X}, \mathcal{D}) < 0$ as (X, D) is K-unstable. Every $g \in G$ induces a pull-back $(\mathcal{X}_g, \mathcal{D}_g)$ of the test configuration $(\mathcal{X}, \mathcal{D})$ with the same $\text{Fut}(\cdot)$, $\|\cdot\|_m$, and $\|\cdot\|_2$. Since a non-trivial rescaling must change the norms, the uniqueness of $(\mathcal{X}, \mathcal{D})$ implies that $(\mathcal{X}_g, \mathcal{D}_g) \cong (\mathcal{X}, \mathcal{D})$ for every $g \in G$. Thus $(\mathcal{X}, \mathcal{D})$ is G -equivariant. Let E be a special divisor over (X, D) such that $\text{ord}_{\mathcal{X}_0}|_{\mathbb{C}(X)} = d \cdot \text{ord}_E$ for some $d \in \mathbb{Z}_{>0}$. Then $\text{Fut}(\mathcal{X}, \mathcal{D}) = d \cdot \beta_{(X, D)}(E) \geq 0$ by [Fuj19c], a contradiction to $\text{Fut}(\mathcal{X}, \mathcal{D}) < 0$. Thus (X, D) is K-semistable. \square

Proposition 2.8. *Let (X, D) be a K-semistable, but not K-polystable, log Fano pair. Let G be a reductive group acting on (X, D) . Then there exists a G -invariant special divisor E over (X, D) with $\beta_{(X, D)}(E) = 0$ which induces a K-polystable special degeneration of (X, D) .*

Proof. By [LWX21, Theorem 1.3] let (X_0, Δ_0) be the unique K-polystable special degeneration of (X, D) . Then the second paragraph of [Zhu21, Proof of Corollary 4.11] implies that there exists a non-trivial G -equivariant special test configuration $(\mathcal{X}, \mathcal{D})$ of (X, D) such that $(\mathcal{X}_0, \mathcal{D}_0) \cong (X_0, \Delta_0)$. Since (X_0, Δ_0) is K-polystable, we have $\text{Fut}(\mathcal{X}, \mathcal{D}) = 0$. From [Fuj19c] we know that there exists a special divisor E over (X, D) and $d \in \mathbb{Z}_{>0}$ such that $\text{ord}_{\mathcal{X}_0}|_{\mathbb{C}(X)} = d \cdot \text{ord}_E$, and $0 = \text{Fut}(\mathcal{X}, \mathcal{D}) = d \cdot \beta_{(X, D)}(E)$. Since $(\mathcal{X}, \mathcal{D})$ is G -equivariant, we know that E is G -invariant. Thus the proof is finished. \square

2.1.4. *Almost log Calabi-Yau pairs.* The following definition is equivalent to the original definition by [Oda13].

Definition 2.9. A log Calabi-Yau pair (X, D) is called *K-semistable* if (X, D) is log canonical.

The following theorem can be viewed as an algebraic analogue of [JMR16, Corollary 1]. A different proof can be obtained by applying [Zho21b, Lemma 5.3].

Theorem 2.10. *Let X be a \mathbb{Q} -Fano variety of dimension $n \geq 2$. Let $D \sim_{\mathbb{Q}} -K_X$ be an effective \mathbb{Q} -Cartier Weil divisor. Assume that (X, D) is plt. Then there exists $\epsilon_1 \in (0, 1)$ depending only on n , such that for any rational number $c \in (1 - \epsilon_1, 1)$, the log Fano pair (X, cD) is uniformly K-stable. In particular, $\text{Aut}(X, D)$ is a finite group.*

Proof. Consider the pair $(X, \frac{m-1}{m}D)$ where $m \in \mathbb{Z}_{>0}$. By [BLX22, Corollary 3.5] based on Birkar's boundedness of complements [Bir19, Theorem 1.8], there exists $N \in \mathbb{Z}_{>0}$ depending only on n , such that either $(X, \frac{m-1}{m}D)$ is uniformly K-stable, or

$$(2.1) \quad \delta(X, \frac{m-1}{m}D) = \inf_E \frac{A_{(X, \frac{m-1}{m}D)}(E)}{S_{(X, \frac{m-1}{m}D)}(E)},$$

where E runs over lc places of N -complements Δ_m^+ of $(X, \frac{m-1}{m}D)$ satisfying $\Delta_m^+ \geq \frac{m-1}{m}D$. Since $N\Delta_m^+$ is a Weil divisor, we know that $\Delta_m^+ \geq D$ as long as $m > N$. Since $\Delta_m^+ \sim_{\mathbb{Q}} -K_X \sim_{\mathbb{Q}} D$, we have that $\Delta_m^+ = D$ for $m > N$.

We claim that $(X, \frac{N}{N+1}D)$ is uniformly K-stable. If not, from $n \geq 2$ and ampleness of D we know that D is connected. Since $\Delta_{N+1}^+ = D$ and (X, D) is plt, the prime divisor D is the only lc place of N -complements Δ_{N+1}^+ . Hence (2.1) implies that $\delta(X, \frac{N}{N+1}D)$ is computed by $E = D$, and simple computation shows that

$$A_{(X, \frac{N}{N+1}D)}(D) = \frac{1}{N+1} \quad \text{and} \quad S_{(X, \frac{N}{N+1}D)}(D) = \frac{1}{(n+1)(N+1)}.$$

This contradicts the assumption that $\delta(X, \frac{N}{N+1}D) \leq 1$. Thus we prove the claim which implies the uniform K-stability of (X, cD) for any $c \in [\frac{N}{N+1}, 1)$ by [ADL19, Proposition 2.13]. The finiteness of $\text{Aut}(X, D)$ follows from [BX19, Corollary 1.3]. \square

2.2. CM line bundles. The CM line bundle of a flat family of polarized projective varieties is a functorial line bundle over the base which was introduced algebraically by Tian [Tia97]. The following definition of CM line bundles is due to Paul and Tian [PT06, PT09] using the Knudsen-Mumford expansion (see also [FR06]). We use the concept of relative Mumford divisors from [Kol18, Kol19]; see also [ADL20, Definition 2.7].

Definition 2.11 (log CM line bundle). Let $f : \mathcal{X} \rightarrow B$ be a proper flat morphism of connected schemes. Assume that f has S_2 fibers of pure dimension n . Let \mathcal{L} be an f -ample line bundle on \mathcal{X} . Let $\mathcal{D} := \sum_{i=1}^k c_i \mathcal{D}_i$ be a relative Mumford \mathbb{Q} -divisor on \mathcal{X} over B where each \mathcal{D}_i is a relative Mumford divisor and $c_i \in [0, 1] \cap \mathbb{Q}$. We also assume that each \mathcal{D}_i is flat over B (see Remark 2.12).

A result of Knudsen-Mumford [KM76] says that there exist line bundles $\lambda_j = \lambda_j(\mathcal{X}, \mathcal{L})$ on B such that for all k ,

$$\det f_!(\mathcal{L}^k) = \lambda_{n+1}^{\binom{k}{n+1}} \otimes \lambda_n^{\binom{k}{n}} \otimes \cdots \otimes \lambda_0.$$

By flatness, the Hilbert polynomial $\chi(\mathcal{X}_b, \mathcal{L}_b^k) = a_0 k^n + a_1 k^{n-1} + O(k^{n-2})$ for any $b \in B$. Then the *CM line bundle* and the *Chow line bundle* of the data $(f : \mathcal{X} \rightarrow B, \mathcal{L}, \mathcal{D})$ are defined as

$$\lambda_{\text{CM}, f, \mathcal{L}} := \lambda_{n+1}^{\mu+n(n+1)} \otimes \lambda_n^{-2(n+1)}, \quad \lambda_{\text{Chow}, f, \mathcal{L}} := \lambda_{n+1}.$$

where $\mu := \frac{2a_1}{a_0}$. The *log CM \mathbb{Q} -line bundle* of the data $(f : \mathcal{X} \rightarrow B, \mathcal{L}, \mathcal{D})$ is defined as

$$\lambda_{\text{CM}, f, \mathcal{D}, \mathcal{L}} := \lambda_{\text{CM}, f, \mathcal{L}} - \frac{n(\mathcal{L}_b^{n-1} \cdot \mathcal{D}_b)}{(\mathcal{L}_b^n)} \lambda_{\text{Chow}, f, \mathcal{L}} + (n+1) \lambda_{\text{Chow}, f|_{\mathcal{D}}, \mathcal{L}|_{\mathcal{D}}},$$

where $(\mathcal{L}_b^{n-1} \cdot \mathcal{D}_b) := \sum_{i=1}^k c_i (\mathcal{L}_b^{n-1} \cdot \mathcal{D}_{i, b})$ and $\lambda_{\text{Chow}, f|_{\mathcal{D}}, \mathcal{L}|_{\mathcal{D}}} := \bigotimes_{i=1}^k \lambda_{\text{Chow}, f|_{\mathcal{D}_i}, \mathcal{L}|_{\mathcal{D}_i}}^{\otimes c_i}$.

Remark 2.12. In Definition 2.11 we assumed that each \mathcal{D}_i are flat over B . This is guaranteed in our setting – \mathbb{Q} -Gorenstein smoothable log Fano families over reduced base schemes – by [ADL20, Proposition 2.12].

2.3. K-moduli of \mathbb{Q} -Fano varieties. We first recall the moduli stack of \mathbb{Q} -Fano varieties.

Definition 2.13. A \mathbb{Q} -Fano family is a morphism $f : \mathcal{X} \rightarrow B$ between schemes such that

- (1) f is projective and flat of pure relative dimension n for some positive integer n ;
- (2) the geometric fibers of f are \mathbb{Q} -Fano varieties;
- (3) $-K_{\mathcal{X}/B}$ is \mathbb{Q} -Cartier and f -ample;
- (4) f satisfies Kollar's condition.

We recall the following definition from [BHLLX21].

Definition 2.14. [BHLLX21, Section 4.1] Let n be a positive integer and V a positive rational number. We define the moduli pseudo-functor $\mathcal{M}_{n, V}^{\text{Fano}}$ that sends a scheme B to

$$\mathcal{M}_{n, V}^{\text{Fano}}(B) := \{\mathbb{Q}\text{-Fano families } f : \mathcal{X} \rightarrow B \mid \dim(X_b) = n, (-K_{X_b})^n = V \text{ for all } b \in S\},$$

Fix an $0 < \epsilon \leq 1$, and let $\mathcal{M}_{n, V}^{\delta \geq \epsilon} \subseteq \mathcal{M}_{n, V}^{\text{Fano}}$ denote the subfunctor defined by

$$\mathcal{M}_{n, V}^{\delta \geq \epsilon}(B) := \{[\mathcal{X} \rightarrow B] \in \mathcal{M}_{n, V}^{\text{Fano}}(B) \mid \delta(\mathcal{X}_b) \geq \epsilon \text{ for all } b \in B\}.$$

We also define

$$\mathcal{M}_{n,V}^{\text{Kss}}(B) := \{[\mathcal{X} \rightarrow B] \in \mathcal{M}_{n,V}^{\text{Fano}}(B) \mid \mathcal{X}_{\bar{b}} \text{ is K-semistable for all } b \in B\}.$$

Then by Theorem 2.3 we know that $\mathcal{M}_{n,V}^{\text{Kss}} = \mathcal{M}_{n,V}^{\delta \geq 1}$.

By [BHLLX21, Section 4.1], the pseudo-functor $\mathcal{M}_{n,V}^{\delta \geq \epsilon}$ is represented by an Artin stack of finite type with affine diagonal (indeed, a quotient stack $[Z/\text{PGL}_{m+1}]$ where Z is a quasi-projective scheme). The CM \mathbb{Q} -line bundle on $\mathcal{M}_{n,V}^{\delta \geq \epsilon}$ is defined as the CM \mathbb{Q} -line bundle of its universal family.

By [BL22, BLX22], we know that for a \mathbb{Q} -Fano family $\mathcal{X} \rightarrow B$, the function $b \mapsto \min\{1, \delta(\mathcal{X}_{\bar{b}})\}$ is constructible and lower semi-continuous. Thus for any $0 < \epsilon < \epsilon' \leq 1$ there are canonical open immersions $\mathcal{M}_{n,V}^{\delta \geq \epsilon'} \hookrightarrow \mathcal{M}_{n,V}^{\delta \geq \epsilon}$.

The following result, known as the K-moduli theorem, is a combination of many recent important algebraic works [Jia20, LWX21, CP21, BX19, ABHLX20, BLX22, Xu20, XZ20, XZ21, BHLLX21, LXZ22].

Theorem 2.15 (K-moduli theorem). *Let n be a positive integer and V a positive rational number. Then there exists an Artin stack $\mathcal{M}_{n,V}^{\text{Kss}}$ of finite type with affine diagonal parametrizing K-semistable \mathbb{Q} -Fano varieties of dimension n and volume V . Moreover, $\mathcal{M}_{n,V}^{\text{Kss}}$ admits a projective good moduli space $M_{n,V}^{\text{Kps}}$ parametrizing K-polystable \mathbb{Q} -Fano varieties, and the CM \mathbb{Q} -line bundle on $\mathcal{M}_{n,V}^{\text{Kss}}$ descends to an ample \mathbb{Q} -line bundle on $M_{n,V}^{\text{Kps}}$.*

We call $\mathcal{M}_{n,V}^{\text{Kss}}$ and $M_{n,V}^{\text{Kps}}$ a *K-moduli stack* and a *K-moduli space*, respectively.

In this paper, we are mainly interested in the \mathbb{Q} -Gorenstein smoothable case.

Definition 2.16. Let n and V be positive integers, and fix any $0 < \epsilon \leq 1$. Let $\mathcal{M}_{n,V}^{\text{sm}, \delta \geq \epsilon}$ be the open substack of $\mathcal{M}_{n,V}^{\delta \geq \epsilon}$ parametrizing smooth Fano varieties X of dimension n and volume V with $\delta(X) \geq \epsilon$. Let $\overline{\mathcal{M}}_{n,V}^{\text{sm}, \delta \geq \epsilon}$ be the Zariski closure of $\mathcal{M}_{n,V}^{\text{sm}, \delta \geq \epsilon}$ in $\mathcal{M}_{n,V}^{\delta \geq \epsilon}$ with reduced structure. We call $\overline{\mathcal{M}}_{n,V}^{\text{sm}, \delta \geq \epsilon}$ a *moduli stack of \mathbb{Q} -Gorenstein smoothable \mathbb{Q} -Fano varieties*. A \mathbb{Q} -Fano variety X is called \mathbb{Q} -Gorenstein smoothable if $[X] \in \overline{\mathcal{M}}_{n,V}^{\text{sm}, \delta \geq \epsilon}$ for some n, V, ϵ .

Let $\overline{\mathcal{M}}_{n,V}^{\text{sm}, \text{Kss}} := \overline{\mathcal{M}}_{n,V}^{\text{sm}, \delta \geq 1}$ be a reduced closed substack of $\mathcal{M}_{n,V}^{\text{Kss}}$. According to Theorem 2.15, the stack $\overline{\mathcal{M}}_{n,V}^{\text{sm}, \text{Kss}}$ admits a projective good moduli space $\overline{M}_{n,V}^{\text{sm}, \text{Kps}}$ as a reduced closed subscheme of $M_{n,V}^{\text{Kps}}$. Note that prior to the algebraic approach in Theorem 2.15, it was shown using analytic methods that there exists a proper good moduli space $\overline{M}_{n,V}^{\text{sm}, \text{Kps}}$ of $\overline{\mathcal{M}}_{n,V}^{\text{sm}, \text{Kss}}$ by [LWX19] (see also [Oda15]).

2.3.1. *\mathbb{Q} -Gorenstein smoothable log Fano pairs.* We will consider the following class of pairs.

Definition 2.17. Let c, r be positive rational numbers such that $c < \min\{1, r^{-1}\}$. A log Fano pair (X, cD) is \mathbb{Q} -Gorenstein smoothable if there exists a \mathbb{Q} -Fano family $\pi : \mathcal{X} \rightarrow C$ over a pointed smooth curve ($0 \in C$) and a relative Mumford divisor \mathcal{D} on \mathcal{X} over C such that the following holds:

- \mathcal{D} is \mathbb{Q} -Cartier, π -ample, and $\mathcal{D} \sim_{\mathbb{Q}, \pi} -rK_{\mathcal{X}/C}$;
- Both π and $\pi|_{\mathcal{D}}$ are smooth morphisms over $C \setminus \{0\}$;
- $(\mathcal{X}_0, c\mathcal{D}_0) \cong (X, cD)$, in particular X has klt singularities.

A \mathbb{Q} -Gorenstein smoothable log Fano family $f : (\mathcal{X}, c\mathcal{D}) \rightarrow B$ over a reduced scheme B consists of a \mathbb{Q} -Fano family $f : \mathcal{X} \rightarrow B$ and a \mathbb{Q} -Cartier relative Mumford divisor \mathcal{D} on \mathcal{X} over B , such that all fibers $(\mathcal{X}_b, c\mathcal{D}_b)$ are \mathbb{Q} -Gorenstein smoothable log Fano pairs, and $\mathcal{D} \sim_{\mathbb{Q}, f} -rK_{\mathcal{X}}$.

For a \mathbb{Q} -Gorenstein smoothable log Fano family, we define its Hodge line bundle as follows.

Definition 2.18. For $c, r \in \mathbb{Q}_{>0}$ with $cr < 1$, let $f : (\mathcal{X}, c\mathcal{D}) \rightarrow B$ be a \mathbb{Q} -Gorenstein smoothable log Fano family over a reduced scheme B where $\mathcal{D} \sim_{\mathbb{Q}, f} -rK_{\mathcal{X}/B}$. The *Hodge \mathbb{Q} -line bundle* $\lambda_{\text{Hodge}, f, r^{-1}\mathcal{D}}$ is defined as the \mathbb{Q} -linear equivalence class of \mathbb{Q} -Cartier \mathbb{Q} -divisors on T such that

$$K_{\mathcal{X}/B} + r^{-1}\mathcal{D} \sim_{\mathbb{Q}} f^*\lambda_{\text{Hodge}, f, r^{-1}\mathcal{D}}.$$

In [ADL19] we define the Artin stacks $\mathcal{KM}_{\chi_0, r, c}$ and prove that they admit proper good moduli spaces $KM_{\chi_0, r, c}$, where the projectivity of such K-moduli spaces is proven by Xu and Zhuang [XZ20]. Note that the K-moduli theorem also holds for all log Fano pairs without the \mathbb{Q} -Gorenstein smoothable assumption as a generalization of Theorem 2.15 (see e.g. [LXZ22, Theorem 1.3]), though we restrict to the \mathbb{Q} -Gorenstein smoothable case in this article.

Theorem 2.19 ([ADL19, Theorem 3.1 and Remark 3.25] and [XZ20]). *Let χ_0 be the Hilbert polynomial of an anti-canonically polarized Fano manifold. Fix $r \in \mathbb{Q}_{>0}$ and a rational number $c \in (0, \min\{1, r^{-1}\})$. Consider the following moduli pseudo-functor over reduced schemes B :*

$$\mathcal{KM}_{\chi_0, r, c}(B) = \left\{ (\mathcal{X}, \mathcal{D})/B \left| \begin{array}{l} (\mathcal{X}, c\mathcal{D})/B \text{ is a } \mathbb{Q}\text{-Gorenstein smoothable log Fano family,} \\ \mathcal{D} \sim_{B, \mathbb{Q}} -rK_{\mathcal{X}/B}, \text{ each fiber } (\mathcal{X}_b, c\mathcal{D}_b) \text{ is } K\text{-semistable,} \\ \text{and } \chi(\mathcal{X}_b, \mathcal{O}_{\mathcal{X}_b}(-kK_{\mathcal{X}_b})) = \chi_0(k) \text{ for } k \text{ sufficiently divisible.} \end{array} \right. \right\}.$$

Then there exists a reduced Artin stack $\mathcal{KM}_{\chi_0, r, c}$ (called a K-moduli stack) of finite type over \mathbb{C} representing the above moduli pseudo-functor. In particular, the \mathbb{C} -points of $\mathcal{KM}_{\chi_0, r, c}$ parametrize K -semistable \mathbb{Q} -Gorenstein smoothable log Fano pairs (X, cD) with Hilbert polynomial $\chi(X, \mathcal{O}_X(-mK_X)) = \chi_0(m)$ for sufficiently divisible m and $D \sim_{\mathbb{Q}} -rK_X$.

Moreover, the Artin stack $\mathcal{KM}_{\chi_0, r, c}$ admits a good moduli space $KM_{\chi_0, r, c}$ (called a K-moduli space) as a projective reduced scheme of finite type over \mathbb{C} , whose closed points parametrize K -polystable log Fano pairs, and the CM \mathbb{Q} -line bundle on $\mathcal{KM}_{\chi_0, r, c}$ descends to an ample \mathbb{Q} -line bundle on $KM_{\chi_0, r, c}$.

Lemma 2.20. *Let n and V be positive integers. Let r be a positive rational number. Let χ_0 be the Hilbert polynomial of an anti-canonically polarized Fano manifold of dimension n and volume V . Then there exists $\epsilon_0 \in (0, 1]$ depending only on n and r such that the forgetful map $\mathcal{KM}_{\chi_0, r, c} \rightarrow \overline{\mathcal{M}}^{\text{sm}, \delta \geq \epsilon_0}$ that assigns $[(X, D)] \mapsto [X]$ is well-defined for every $c \in (0, \min\{1, r^{-1}\})$. Moreover, if r is an integer, then this forgetful map is a smooth morphism with connected or empty fibers.*

Proof. We fix a positive integer n and a positive rational number r . Since smooth Fano manifolds of dimension n are bounded [KMM92, Cam92], there are finitely many choices of V and χ_0 . By [ADL19, Theorem 1.2] we know that the collection of n -dimensional \mathbb{Q} -Fano varieties X such that $[(X, D)] \in \mathcal{KM}_{\chi_0, r, c}$ for some D , χ_0 , V , and $c \in (0, \min\{1, r^{-1}\})$ is bounded. Thus by [BJ20, Theorem A] and [BL22, Proposition 5.3], there exists $\epsilon_0 \in (0, 1]$ depending only on n and r such that $\delta(X) \geq \epsilon_0$ for every X in this collection. This shows that the forgetful map $\mathcal{KM}_{\chi_0, r, c} \rightarrow \overline{\mathcal{M}}^{\text{sm}, \delta \geq \epsilon_0}$ is well-defined.

Next, we assume $r \in \mathbb{Z}_{>0}$. For the last statement, following the last paragraph of the proof of [ADL20, Theorem 2.21], it suffices to show that for any \mathbb{Q} -Gorenstein smoothable \mathbb{Q} -Fano variety X and any effective Weil divisor D on X satisfying $D \sim_{\mathbb{Q}} -rK_X$, we have $D \sim -rK_X$. Let $\pi : (\mathcal{X}, \mathcal{D}) \rightarrow B$ be a \mathbb{Q} -Gorenstein smoothing over a pointed curve $0 \in B$ with $(\mathcal{X}_0, \mathcal{D}_0) \cong (X, D)$

and $\mathcal{D} \sim_{B, \mathbb{Q}} -rK_{\mathcal{X}/B}$. Since π is smooth over $B \setminus \{0\}$, the class group of \mathcal{X}_b is torsion free for $b \in B \setminus \{0\}$. Thus we have $\mathcal{D}|_{\mathcal{X} \setminus \mathcal{X}_0} \sim_B -rK_{\mathcal{X}/B}|_{\mathcal{X} \setminus \mathcal{X}_0}$ as r is an integer. Since \mathcal{X}_0 is integral and $\mathcal{X}_0 \sim_B 0$, we know that $\mathcal{D} \sim_B -rK_{\mathcal{X}/B}$. This implies that $\mathcal{D}_0 \sim -rK_{\mathcal{X}_0}$. The proof is finished. \square

3. GEOMETRY AND MODULI OF QUARTIC K3 SURFACES

3.1. Geometry of quartic K3 surfaces. Our goal in this section is to review the Hassett-Keel-Looijenga program (Section 3.4). Before doing so, we introduce some terminology from the geometry of quartic K3 surfaces as studied by Mayer in [May72]. A *K3 surface* S is a connected projective surface with du Val singularities such that $\omega_S \cong \mathcal{O}_S$ and $H^1(S, \mathcal{O}_S) = 0$. A *polarized K3 surface* (S, L_S) consists of a K3 surface S and an ample line bundle L_S which is primitive. A *quartic K3 surface* is a polarized K3 surface (S, L_S) of degree 4, i.e. $(L_S^2) = 4$. Consider the map $S \dashrightarrow |L_S|^\vee \cong \mathbb{P}^3$ induced by the linear system $|L_S|$.

Definition 3.1. Generically, the linear system $|L_S|$ defines an isomorphism onto a quartic surface in \mathbb{P}^3 with du Val singularities.

- (1) We say that S is *hyperelliptic* if $|L_S|$ induces a $2 : 1$ map onto a quadric surface in \mathbb{P}^3 . In this case S is isomorphic to a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$ ramified along a $(4, 4)$ curve or a degree 8 curve, respectively.
- (2) We say that S is *unigonal* if $|L_S|$ defines a rational map from S onto a twisted cubic curve in \mathbb{P}^3 with general fiber a smooth elliptic curve.

By [May72], we know that any quartic K3 surface belongs to one of the three classes above.

3.2. K-moduli of quartic surfaces. We define the K-moduli stacks $\overline{\mathcal{M}}_c^K$ and spaces $\overline{\mathfrak{M}}_c^K$.

Definition 3.2. Let χ_0 be the Hilbert polynomial of $(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(4))$. Let $c \in (0, 1) \cap \mathbb{Q}$ be a rational number. We define the K-moduli stacks $\overline{\mathcal{M}}_c^K$ and spaces $\overline{\mathfrak{M}}_c^K$ as

$$\overline{\mathcal{M}}_c^K := \mathcal{KM}_{\chi_0, 1, c} \quad \text{and} \quad \overline{\mathfrak{M}}_c^K := KM_{\chi_0, 1, c}.$$

By Theorem 2.19 we know that $\overline{\mathcal{M}}_c^K$ is a reduced Artin stack of finite type, and $\overline{\mathfrak{M}}_c^K$ is a reduced projective scheme.

Lemma 3.3. Let X be a \mathbb{Q} -Fano variety in $\overline{\mathcal{M}}_{3, 64}^{\text{sm}, \delta \geq \epsilon}$ for some $\epsilon \in (0, 1]$. Then X admits a \mathbb{Q} -Gorenstein smoothing to \mathbb{P}^3 . Moreover, there exists an ample \mathbb{Q} -Cartier Weil divisorial sheaf L on X such that the following conditions hold.

- (1) $L^{[m]}$ is Cohen-Macaulay for any $m \in \mathbb{Z}$;
- (2) $\omega_X \cong L^{[-4]}$ and $(L^3) = 1$;
- (3) $h^i(X, L^{[m]}) = h^i(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(m))$ for any $m \in \mathbb{Z}$ and $i \geq 0$;

Proof. From the Iskovskikh-Mori-Mukai classification of smooth Fano threefolds [IP99], we know that \mathbb{P}^3 is the only smooth Fano threefold with anti-canonical volume 64. Hence X admits a \mathbb{Q} -Gorenstein smoothing $\pi : \mathcal{X} \rightarrow B$ over a smooth pointed curve $0 \in B$ such that $\mathcal{X}_0 \cong X$ and $\mathcal{X}_b \cong \mathbb{P}^3$ for $b \in B \setminus \{0\}$. Denote by $\mathcal{X}^\circ := \mathcal{X} \setminus \mathcal{X}_0$ and $B^\circ := B \setminus \{0\}$. After a quasi-finite base change of π , we may assume that $\mathcal{X}^\circ \cong \mathbb{P}^3 \times B^\circ$. Let \mathcal{L}° be a Weil divisor on \mathcal{X}° in $|\mathcal{O}_{\mathbb{P}^3_{B^\circ}}(1)|$. Let \mathcal{L} be the Zariski closure of \mathcal{L}° in \mathcal{X} . Since $4\mathcal{L}^\circ \sim_B -K_{\mathcal{X}^\circ/B^\circ}$ and \mathcal{X}_0 is a Cartier prime divisor, we know that $4\mathcal{L} \sim_B -K_{\mathcal{X}/B}$, in particular \mathcal{L} is \mathbb{Q} -Cartier. Since \mathcal{X} is klt and \mathcal{L} is \mathbb{Q} -Cartier, the sheaf $\mathcal{O}_{\mathcal{X}}(m\mathcal{L})$ is Cohen-Macaulay for any $m \in \mathbb{Z}$ by [KM98, Corollary 5.25]. Let $L := \mathcal{L}|_{\mathcal{X}_0}$, then L is a \mathbb{Q} -Cartier Weil divisor on X . Moreover, we have $\mathcal{O}_{\mathcal{X}}(m\mathcal{L}) \otimes \mathcal{O}_{\mathcal{X}_0}$ and $\mathcal{O}_X(mL)$ are isomorphic on a big open subset of \mathcal{X}_0 , hence they are isomorphic everywhere since $\mathcal{O}_{\mathcal{X}}(m\mathcal{L}) \otimes \mathcal{O}_{\mathcal{X}_0}$ is Cohen-Macaulay. Thus part (1) is proved.

For part (2), notice that $4\mathcal{L} \sim_B -K_{\mathcal{X}/B}$ implies $4L \sim -K_X$. We have $(L^3) = 1$ since $(-K_X)^3 = 64$. Part (3) follows from Kawamata-Viehweg vanishing similar to [Liu22, Proof of Theorem 3.1]. \square

From now on, we fix a number $\epsilon_0 \in (0, 1]$ from Lemma 2.20 with $n = 3$ and $r = 1$.

Next we recall a result connecting K-stability and GIT stability as a special case of [GMGS21, Theorem 1.2] and [ADL19, Theorem 1.4] (see also [Zho21a]).

Theorem 3.4 ([GMGS21, ADL19]). *There exists $\epsilon_2 \in (0, 1)$ such that for any rational number $c \in (0, \epsilon_2)$, a quartic surface $S \subset \mathbb{P}^3$ is GIT (poly/semi)stable if and only if (\mathbb{P}^3, cS) is K-(poly/semi)stable.*

Lemma 3.5. *Let $S \subset \mathbb{P}^3$ be a quartic surface.*

- (1) *If S has only ADE singularities, then (\mathbb{P}^3, cS) is K-stable for any $c \in [0, 1] \cap \mathbb{Q}$.*
- (2) *If S is semi-log canonical, then (\mathbb{P}^3, cS) is K-semistable for any $c \in [0, 1] \cap \mathbb{Q}$.*

Proof. We first prove part (1). Since S has ADE singularities, it is GIT stable by [Sha81, Theorem 2.4]. Hence Theorem 3.4 implies that $(\mathbb{P}^3, \epsilon S)$ is K-stable for $0 < \epsilon \ll 1$. Moreover, by adjunction we have that (\mathbb{P}^3, S) is plt. Hence part (1) follows from [ADL19, Proposition 2.13].

Next we prove (2). In fact, inversion of adjunction implies that (\mathbb{P}^3, S) is log canonical since S is slc. Then the result follows from [ADL19, Proposition 2.13] since \mathbb{P}^3 is K-polystable. \square

Recall that \mathcal{M}° and \mathfrak{M}° denote the modui stack and coarse moduli space of ADE quartic surfaces in \mathbb{P}^3 , respectively. Denote by \mathcal{M}^{slc} the open substack of the GIT moduli stack $\overline{\mathcal{M}}^{\text{GIT}}$ parametrizing quartic surfaces S that are semi-log canonical (i.e. (\mathbb{P}^3, S) is log canonical).

Proposition 3.6. *For any rational number $c \in (0, 1)$, both $\overline{\mathcal{M}}_c^K$ and $\overline{\mathfrak{M}}_c^K$ are irreducible. Moreover, there are open immersions $\mathcal{M}^\circ \hookrightarrow \mathcal{M}^{\text{slc}} \hookrightarrow \overline{\mathcal{M}}_c^K$ whose images in $\overline{\mathcal{M}}_c^K$ are saturated open substacks. Taking good moduli spaces yields open immersions $\mathfrak{M}^\circ \hookrightarrow \mathfrak{M}^{\text{slc}} \hookrightarrow \overline{\mathfrak{M}}_c^K$ where $\mathfrak{M}^{\text{slc}}$ parametrizes GIT polystable slc quartic surfaces $S \subset \mathbb{P}^3$. In particular, both \mathcal{M}° and \mathcal{M}^{slc} are saturated open substacks of $\overline{\mathcal{M}}^{\text{GIT}}$.*

Proof. By Lemma 3.3 we know that $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ is irreducible. Thus $\overline{\mathcal{M}}_c^K$ is irreducible since the forgetful map $\overline{\mathcal{M}}_c^K \rightarrow \overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ is smooth with connected fibers by Lemma 2.20. By Lemma 3.5 (2), we know that any $[S] \in \mathcal{M}^{\text{slc}}$ satisfies that (\mathbb{P}^3, cS) is K-semistable for any $c \in (0, 1)$. Thus by openness of klt and lc (see [Kol97, Corollary 7.6]), we know that both \mathcal{M}° and \mathcal{M}^{slc} are dense open substacks of $\overline{\mathcal{M}}_c^K$.

Next, we show saturatedness of \mathcal{M}° in $\overline{\mathcal{M}}_c^K$. If S is an ADE quartic surface, then Lemma 3.5 implies that (\mathbb{P}^3, cS) is K-stable for any $c \in (0, 1)$. Thus all \mathbb{C} -points in \mathcal{M}° are closed with finite stabilizers, which implies that \mathcal{M}° is saturated in $\overline{\mathcal{M}}_c^K$.

Finally, we show saturatedness of \mathcal{M}^{slc} in $\overline{\mathcal{M}}_c^K$. Let S be a slc quartic surface. Since (\mathbb{P}^3, cS) is K-semistable for any $c \in (0, 1)$ from the above discussion, by Theorem 3.4 we know that S is GIT semistable. Let S_0 be the unique GIT polystable quartic surface in the orbit closure of S . Then by Theorem 3.4 we know that $(\mathbb{P}^3, \epsilon S_0)$ is K-polystable for $0 < \epsilon \ll 1$. Denote by $(\mathbb{P}^3 \times \mathbb{A}^1, \mathcal{S})$ the test configuration of (\mathbb{P}^3, S) degenerating to (\mathbb{P}^3, S_0) . Hence we have $\text{Fut}(\mathbb{P}^3 \times \mathbb{A}^1, \epsilon \mathcal{S}) = 0$. Since Fut is linear in coefficients, we know that $\text{Fut}(\mathbb{P}^3 \times \mathbb{A}^1, c\mathcal{S}) = 0$ for any $c \in (0, 1)$. Hence (\mathbb{P}^3, cS_0) is K-semistable for any $c \in (0, 1)$ by [LWX21, Lemma 3.1] which implies that S_0 is slc. Thus interpolation of K-stability [ADL19, Proposition 2.13] implies that (\mathbb{P}^3, cS_0) is the unique K-polystable degeneration of (\mathbb{P}^3, cS) for any $c \in (0, 1)$. Since $S_0 \in \mathcal{M}^{\text{slc}}$, we have that \mathcal{M}^{slc} is saturated in $\overline{\mathcal{M}}_c^K$. The last statement follows from $\overline{\mathcal{M}}_c^K \cong \overline{\mathcal{M}}^{\text{GIT}}$ by Theorem 3.4. \square

Definition 3.7. The *K-semistable threshold* of a quartic surface $S \subset \mathbb{P}^3$ is defined as

$$\text{kst}(\mathbb{P}^3, S) := \sup\{c \in [0, 1] \mid (\mathbb{P}^3, cS) \text{ is K-semistable}\}.$$

If S is GIT semistable, then by Theorem 3.4 and [ADL19, Theorem 3.15] we know that $\text{kst}(\mathbb{P}^3, S) \in (0, 1]$ is a rational number, and the supremum is a maximum.

By Lemma 3.5, we know that $\text{kst}(\mathbb{P}^3, S) = 1$ if and only if S is slc. If, in addition, S is GIT polystable, then by interpolation of K-stability [ADL19, Proposition 2.13] and Theorem 3.4, we know that for any $c \in [0, \text{kst}(\mathbb{P}^3, S))$, the log pair (\mathbb{P}^3, cS) is K-polystable. If a GIT polystable quartic surface S is not slc, i.e. $\text{kst}(\mathbb{P}^3, S) < 1$, then $(\mathbb{P}^3, \text{kst}(\mathbb{P}^3, S)S)$ is K-semistable but not K-polystable by [ADL19, Proposition 3.18].

The following theorem follows directly from [ADL19, Theorem 1.2] and Proposition 3.6.

Theorem 3.8. *There exist rational numbers $0 = c_0 < c_1 < c_2 < \dots < c_k = 1$ such that for each $0 \leq i \leq k-1$, both the K-moduli stack $\overline{\mathcal{M}}_c^K$ and the K-moduli space $\overline{\mathfrak{M}}_c^K$ are independent of the choice of the rational number $c \in (c_i, c_{i+1})$. Moreover, for each $1 \leq i \leq k-1$ and $0 < \epsilon \ll 1$ we have open immersions*

$$\overline{\mathcal{M}}_{c_i-\epsilon}^K \hookrightarrow \overline{\mathcal{M}}_{c_i}^K \hookleftarrow \overline{\mathcal{M}}_{c_i+\epsilon}^K.$$

which induce projective birational morphisms

$$\overline{\mathfrak{M}}_{c_i-\epsilon}^K \xrightarrow{\phi_i^-} \overline{\mathfrak{M}}_{c_i}^K \xleftarrow{\phi_i^+} \overline{\mathfrak{M}}_{c_i+\epsilon}^K.$$

In addition, all the above morphisms have local VGIT presentations in terms of [AFS17, (1.2)].

3.3. GIT stratification of quartic surfaces. By Proposition 3.6, the GIT moduli space $\overline{\mathfrak{M}}^{\text{GIT}}$ has an open subset $\mathfrak{M}^{\text{slc}}$ parametrizing GIT polystable quartic surfaces with slc singularities. From [Sha81, LO18], we know that the complement $\overline{\mathfrak{M}}^{\text{GIT}} \setminus \mathfrak{M}^{\text{slc}}$ (denoted by \mathfrak{M}^{IV} therein) has two connected components, where one of them is an isolated point $\{[T]\}$ representing the tangent developable surface T (see Definition 4.1), and the other component has the following stratification

$$(3.1) \quad (\overline{\mathfrak{M}}^{\text{GIT}} \setminus \mathfrak{M}^{\text{slc}}) \setminus \{[T]\} = W_8 \supset W_7 \supset W_6 \supset W_4 \supset W_3 \supset W_2 \supset W_1 \supset W_0.$$

Here W_i is an i -dimensional closed integral subvariety of $\overline{\mathfrak{M}}^{\text{GIT}}$, and $W_0 = \{[2Q]\}$ is the single point representing the double quadric surface. For each $i \in \{0, 1, 2, 3, 4, 6, 7, 8\}$, we denote by $W_i^\circ := W_i \setminus W_{i-1}$ when $i \notin \{0, 6\}$, $W_6^\circ := W_6 \setminus W_4$, and $W_0^\circ = W_0$.

The following result follows from the classification of Shah [Sha81, S-4.3 on Page 282] (see also [LO18, Section 4.3]).

Theorem 3.9 ([Sha81]). *Let $[S] \in W_i^\circ$ for $i \in \{0, 1, 2, 3, 4, 6, 7, 8\}$. Then in suitable projective coordinates $[x_0, x_1, x_2, x_3]$ of \mathbb{P}^3 , the equation of S has the form $q^2 + g = 0$ where q and g are given as follows.*

$$q = x_0x_2 + x_1^2 + ax_3^2,$$

$$g = \begin{cases} x_3^3(x_0 + \beta_1(x_1, x_2, x_3)) + x_2(x_3^2f_1(x_1, x_2) + x_2x_3g_1(x_1, x_2) + x_2^2h_1(x_1, x_2)), & \text{if } i \geq 3; \\ x_3^3l_1(x_1, x_3), & \text{if } i \leq 2. \end{cases}$$

Here β_1 , f_1 , g_1 , h_1 , and l_1 are homogeneous linear polynomials in corresponding variables. More precisely, we have the following classification.

- (1) $[S] \in W_8^\circ$ if and only if $x_2 \nmid h_1$;
- (2) $[S] \in W_7^\circ$ if and only if $x_2 \mid h_1$ and $x_2 \nmid g_1$;
- (3) $[S] \in W_6^\circ$ if and only if $x_2 \mid h_1 \neq 0$ and $x_2 \mid g_1$;

- (4) $[S] \in W_4^\circ$ if and only if $h_1 = 0$, and either $x_2 \mid g_1 \neq 0$ or $x_2 \nmid f_1$;
- (5) $[S] \in W_3^\circ$ if and only if $h_1 = g_1 = 0$ and $x_2 \mid f_1 \neq 0$;
- (6) $[S] \in W_2^\circ$ if and only if $x_3 \nmid l_1$;
- (7) $[S] \in W_1^\circ$ if and only if $x_3 \mid l_1 \neq 0$;
- (8) $[S] \in W_0^\circ$ if and only if $g = 0$ and $a \neq 0$.

3.4. Laza-O'Grady and the Hassett-Keel-Looijenga Program. The moduli space of quartic K3 surfaces can be constructed as a Type IV locally symmetric variety \mathcal{F} , and comes with a natural Baily-Borel compactification \mathcal{F}^* . The global Torelli theorem for K3 surfaces implies that the period map $\mathfrak{p} : \overline{\mathcal{M}}^{\text{GIT}} \dashrightarrow \mathcal{F}^*$ is birational. Building off of previous work of Shah [Sha80] and Looijenga [Loo03a, Loo03b], in a series of papers [LO19, LO18, LO21] Laza and O'Grady propose a conjectural method to resolve the period map \mathfrak{p} whenever \mathcal{F} is a Type IV locally symmetric variety associated to a lattice of the form $U^2 \oplus D_{N-2}$. When $N = 18$ this is the case of hyperelliptic quartic K3 surfaces, and when $N = 19$ this is the case of quartic K3 surfaces.

Recall that Baily-Borel showed that (for any N) one has $\mathcal{F}^* \cong \text{Proj } R(\mathcal{F}, \lambda)$, where λ denotes the hodge line bundle. Based on observations of Looijenga, Laza-O'Grady predict that in many cases there is an isomorphism $\overline{\mathcal{M}}^{\text{GIT}} \cong \text{Proj } R(\mathcal{F}, \lambda + \Delta)$, for some geometrically meaningful boundary divisor Δ depending on \mathcal{F} . Moreover, they predict that more generally the rings $R(\mathcal{F}, \lambda + \beta\Delta)$ are finite generated and so the schemes $\mathcal{F}(\beta) = \text{Proj } R(\mathcal{F}, \lambda + \beta\Delta)$ interpolate between $\overline{\mathcal{M}}^{\text{GIT}}$ and \mathcal{F}^* . Their work also predicts the location of the walls, i.e. the values β where the moduli spaces change.

3.4.1. Hyperelliptic quartic K3 surfaces. When $N = 18$, i.e. the *hyperelliptic case*, Laza and O'Grady confirm their conjecture in [LO19]. By Definition 3.1, this case occurs when the K3 surface is a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ ramified along a $(4, 4)$ curve. If we let $\overline{\mathcal{M}}_{(4,4)}^{\text{GIT}}$ denote the GIT quotient of $(4, 4)$ curves on $\mathbb{P}^1 \times \mathbb{P}^1$, then Laza and O'Grady show that the period map $\mathfrak{p} : \overline{\mathcal{M}}_{(4,4)}^{\text{GIT}} \dashrightarrow \mathcal{F}^*(18)$ can be resolved via a series of explicit wall crossings arising from variation of GIT. Let $H_{h,18} \subset \mathcal{F}(18)$ denote the divisor which parametrizes periods of hyperelliptic K3s which are the double cover of a quadric cone.

If $\text{Reg}(\mathfrak{p}) \subset \overline{\mathcal{M}}_{(4,4)}^{\text{GIT}}$ denotes the regular locus of \mathfrak{p} , then $\mathfrak{p}(\text{Reg}(\mathfrak{p})) \cap \mathcal{F}(18) \cong \mathcal{F}(18) \setminus H_{h,18}$. Laza and O'Grady prove that $R(\beta) := R(\mathcal{F}(18), \lambda + \beta \cdot \frac{H_{h,18}}{2})$ is a finitely generated \mathbb{C} -algebra, and that $\mathcal{F}_{18}(\beta) := \text{Proj } R(\beta)$ is a projective variety which interpolates between $\mathcal{F}_{18}(0) \cong \mathcal{F}(18)^*$, the Baily-Borel compactification, and $\mathcal{F}_{18}(1) \cong \overline{\mathcal{M}}_{(4,4)}^{\text{GIT}}$, the GIT quotient. Moreover, the period map can be explicitly described as a composition of elementary birational maps. The first step $\mathcal{F}_{18}(\epsilon) \rightarrow \mathcal{F}_{18}(0)$ can be realized as the \mathbb{Q} -factorialization of $\mathcal{F}(18)^*$, which fails to be \mathbb{Q} -factorial along $H_{h,18}$. The remainder of the birational transformations are flips, finally followed by a divisorial contraction $\mathcal{F}_{18}(1 - \epsilon) \rightarrow \overline{\mathcal{M}}_{(4,4)}^{\text{GIT}}$.

In [ADL20], we show that the wall crossings resolving the period map \mathfrak{p} can be interpreted as wall-crossings in a suitable K-moduli space of log Fano pairs. Let $\overline{\mathcal{K}}_c$ denote the connected component of the moduli stack parametrizing K-semistable log Fano pairs admitting \mathbb{Q} -Gorenstein smoothings to $(\mathbb{P}^1 \times \mathbb{P}^1, cC)$ where C is a $(4, 4)$ curve. Denote the good moduli space of $\overline{\mathcal{K}}_c$ by \overline{K}_c . Then, varying the weight c , the K-moduli spaces \overline{K}_c interpolate between $\overline{\mathcal{M}}_{(4,4)}^{\text{GIT}}$ and $\mathcal{F}(18)^*$. Moreover, the explicit intermediate spaces constructed in [LO19] using variation of GIT are all isomorphic to K-moduli spaces, and the walls coincide.

3.4.2. General conjecture. In fact, Laza and O'Grady conjecture that a similar behavior that is shown in [LO19] is true for many classes of varieties whose moduli space can be constructed as a Type IV locally symmetric variety. They expect that, as in the hyperelliptic case, if

$\mathfrak{p} : \overline{\mathfrak{M}}^{\text{GIT}} \dashrightarrow \mathcal{F}^*$ represents the relevant birational period map from a GIT compactification to Baily-Borel, then $\mathfrak{p}(\text{Reg}(\mathfrak{p})) \cap \mathcal{F} \cong \mathcal{F} \setminus \Delta$, for some geometrically meaningful divisor Δ . Moreover, $R(\beta) := R(\mathcal{F}, \lambda + \beta \cdot \Delta)$ should be finitely generated and so $\mathcal{F}(\beta) := \text{Proj } R(\beta)$ should interpolate between $\overline{\mathfrak{M}}^{\text{GIT}}$ and \mathcal{F}^* . Moreover, they conjecture that interpolating from \mathcal{F}^* to $\overline{\mathfrak{M}}^{\text{GIT}}$ should consist of birational transformations related to Δ . More precisely, let $\mathcal{H} := \pi^{-1} \text{Supp} \Delta$, where $\pi : \mathcal{D} \rightarrow \mathcal{F}$. Then \mathcal{H} is a union of hyperplane sections of \mathcal{D} , and has a stratification by closed subsets, where the stratification is given by the number of independent sheets of \mathcal{H} containing the general point. Then, the stratification of \mathcal{H} induces a stratification of $\text{Supp} \Delta$.

With this in mind, the prediction for resolving \mathfrak{p} is as follows: \mathbb{Q} -factorialize Δ , followed by a series of explicit flips of strata inside Δ , followed by a divisorial contraction of the strict transform of Δ to obtain $\overline{\mathfrak{M}}^{\text{GIT}}$.

3.4.3. Quartic K3 surfaces. Now let $\overline{\mathfrak{M}}^{\text{GIT}}$ denote the GIT moduli space of quartic surfaces. We recall some notation from [LO19]. For quartic K3 surfaces, we have that the K3 lattice $\Lambda \cong U^2 \oplus D_{17}$. Consider

$$\mathcal{D} = \{[\sigma] \in \mathbb{P}(\Lambda \otimes \mathbb{C}) \mid \sigma^2 = 0, (\sigma + \overline{\sigma})^2 > 0\}^+,$$

where the superscript $+$ indicates that we have taken one of the two connected components. Let $O^+(\Lambda)$ denote the subgroup of isometries $O(\Lambda)$ of Λ which fixes \mathcal{D} . Then $\mathcal{F} \cong \mathcal{D}/O^+(\Lambda)$ is the period space for quartic K3 surfaces (see [LO19, Section 1.2]).

Definition 3.10. The *hyperelliptic divisor* $H_h \subset \mathcal{F}$ is the image of $v^\perp \cap \mathcal{D}$ for $v \in \Lambda$ such that $q(v) = -4$ and $\text{div}(v) = 2$. The *unigonal divisor* $H_u \subset \mathcal{F}$ is the image of $v^\perp \cap \mathcal{D}$ for $v \in \Lambda$ such that $q(v) = -4$ and $\text{div}(v) = 4$.

Remark 3.11. Note that $\mathfrak{p}(S, L_S) \in H_h$ (resp. H_u) if and only if (S, L_S) is hyperelliptic (resp. unigonal) in the sense of Definition 3.1.

For quartic K3 surfaces, Laza-O'Grady predict that the regular locus of \mathfrak{p} is the complement of H_u and H_h . If $\Delta = (H_u + H_h)/2$, they predict that the predicted critical β values are $\beta \in \{1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \frac{1}{7}, \frac{1}{9}, 0\}$ (see [LO19, Prediction 5.1.1]). Recall, in (3.1), we discussed a stratification of $\overline{\mathfrak{M}}^{\text{GIT}}$. They predict that, under \mathfrak{p} , this stratification is related to a stratification of \mathcal{F}^* . This is made more precise as follows. Let $\Delta^{(k)} \subset \text{Supp} \Delta$ be the k -th stratum of the stratification defined above, and consider

$$(3.2) \quad Z^9 \subset Z^8 \subset Z^7 \subset Z^5 \subset Z^4 \subset Z^3 \subset Z^2 \subset Z^1 = H_u \cup H_h \subset \mathcal{F},$$

where $Z^k = \Delta^{(k)}$ for $k \leq 5$, and then

- $Z^7 = \text{Im} \mathcal{F}(\text{II}_{2,10} \oplus A_2) \hookrightarrow \mathcal{F}$,
- $Z^8 = \text{Im} \mathcal{F}(\text{II}_{2,10} \oplus A_1) \hookrightarrow \mathcal{F}$, and
- $Z^9 = \text{Im} \mathcal{F}(\text{II}_{2,10}) \hookrightarrow \mathcal{F}$.

Then, they predict that each birational map occurring at one of the critical values above corresponds to a flip with center Z^k , and that each Z^k is replaced by $W_{k-1} \subset \overline{\mathfrak{M}}^{\text{GIT}}$.

4. TANGENT DEVELOPABLE SURFACE AND UNIGONAL K3 SURFACES

In this section, we show that if T denotes the tangent developable surface (see Definition 4.1), then the pair (\mathbb{P}^3, cT) is K-polystable if and only if $c < \frac{9}{13}$. Moreover, the c -K-polystable replacements of (\mathbb{P}^3, T) for $c > \frac{9}{13}$ are log pairs (X_u, S) where X_u is a Gorenstein canonical Fano threefold (see Proposition 4.5 for the construction), and S is a GIT polystable unigonal K3 surface.

4.1. Destabilizing divisor.

Definition 4.1. [EL20, Pages 30 - 31] The *tangent developable* to the twisted cubic curve C_0 in \mathbb{P}^3 is the surface $T \subset \mathbb{P}^3$ defined to be the union of all the embedded projective tangent lines to C_0 . The surface T is a quartic surface that has cuspidal singularities along C_0 , and whose normalization is $\mathbb{P}^1 \times \mathbb{P}^1$ such that the diagonal $\Delta_{\mathbb{P}^1} \subset \mathbb{P}^1 \times \mathbb{P}^1$ is the preimage of C_0 .

Consider the group $G := \mathrm{PGL}(2, \mathbb{C})$ acting linearly on \mathbb{P}^3 where C_0 is G -invariant, since clearly $\mathrm{Aut}(\mathbb{P}^3, C_0) \cong \mathrm{Aut}(C_0) \cong G$. A simple analysis shows that there are precisely three G -orbits in \mathbb{P}^3 , which are C_0 , $T \setminus C_0$, and $\mathbb{P}^3 \setminus T$.

Proposition 4.2. *Let $\iota : \mathbb{P}^1 \xrightarrow{\cong} C_0$ be a parametrization. Then $\iota^* \mathcal{N}_{C_0/\mathbb{P}^3} \cong \mathcal{O}_{\mathbb{P}^1}(5) \oplus \mathcal{O}_{\mathbb{P}^1}(5)$. Moreover, there exists a G -equivariant sub-line bundle \mathcal{N}_1 of $\mathcal{N}_{C_0/\mathbb{P}^3}$ such that $\iota^* \mathcal{N}_1 \cong \mathcal{O}_{\mathbb{P}^1}(4)$. We denote by $\mathcal{N}_2 := \mathcal{N}_{C_0/\mathbb{P}^3}/\mathcal{N}_1$ the quotient line bundle. Then locally analytically along C_0 we may split $\mathcal{N}_{C_0/\mathbb{P}^3}$ as $\mathcal{N}_1 \oplus \mathcal{N}_2$ such that the surface T has analytic equation $(y^2 = x^3)$ where $\mathcal{N}_1 = \langle \partial/\partial x \rangle$ and $\mathcal{N}_2 = \langle \partial/\partial y \rangle$.*

Proof. By [EVdV81], $\iota^* \mathcal{N}_{C_0/\mathbb{P}^3} \cong \mathcal{O}_{\mathbb{P}^1}(5) \oplus \mathcal{O}_{\mathbb{P}^1}(5)$. Denote by $\mathcal{O}_{C_0}(m) := \mathcal{O}_{\mathbb{P}^3}(m) \otimes \mathcal{O}_{C_0}$, then we have $\iota^* \mathcal{O}_{C_0}(m) \cong \mathcal{O}_{\mathbb{P}^1}(3m)$. Denote by \mathcal{I}_{C_0} the ideal sheaf of C_0 in \mathbb{P}^3 . Since C_0 is projectively normal, we have a short exact sequence

$$0 \rightarrow H^0(\mathbb{P}^3, \mathcal{I}_{C_0}(2)) \rightarrow H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(2)) \rightarrow H^0(C_0, \mathcal{O}_{C_0}(2)) \rightarrow 0.$$

Hence a dimension computation shows that $h^0(\mathbb{P}^3, \mathcal{I}_{C_0}(2)) = 3$. Since $G = \mathrm{PGL}(2, \mathbb{C})$ has fundamental group $\mathbb{Z}/2\mathbb{Z}$, the line bundle $\mathcal{O}_{\mathbb{P}^3}(2)$ has a natural G -linearization. Thus G acts on $H^0(\mathbb{P}^3, \mathcal{I}_{C_0}(2))$. From the classification of G -orbits, we know that there does not exist any G -invariant quadric surface in \mathbb{P}^3 . Thus $H^0(\mathbb{P}^3, \mathcal{I}_{C_0}(2))$ is a 3-dimensional irreducible representation of G . Next we consider the restriction map

$$r : H^0(\mathbb{P}^3, \mathcal{I}_{C_0}(2)) \rightarrow H^0(C_0, (\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2)(2)).$$

It is clear that r is non-zero and G -equivariant. Since $\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2$ is the conormal bundle $\mathcal{N}_{C_0/\mathbb{P}^3}^\vee$, we have that $\iota^* (\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2)(2) \cong \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2}$ which implies that $h^0(C_0, (\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2)(2)) = 4$. Since G is reductive, the cokernel of r provides a non-zero G -invariant section $s \in H^0(C_0, (\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2)(2))$. Since $(\mathcal{I}_{C_0}/\mathcal{I}_{C_0}^2)(2) \cong \mathrm{Hom}(\mathcal{N}_{C_0/\mathbb{P}^3}, \mathcal{O}_{C_0}(2))$, the section s induces a non-zero G -equivariant morphism $\mathcal{N}_{C_0/\mathbb{P}^3} \rightarrow \mathcal{O}_{C_0}(2)$ which has to be surjective since G acts transitively on C_0 . Thus we define $\mathcal{N}_2 := \mathcal{O}_{C_0}(2)$ and \mathcal{N}_1 as the kernel of the surjection $\mathcal{N}_{C_0/\mathbb{P}^3} \rightarrow \mathcal{N}_2$. Computations on degrees show that $\iota^* \mathcal{N}_1 \cong \mathcal{O}_{\mathbb{P}^1}(4)$ and $\iota^* \mathcal{N}_2 \cong \mathcal{O}_{\mathbb{P}^1}(6)$.

It is clear that T has equation $(y^2 = x^3)$ in some analytic coordinate (x, y, z) of \mathbb{P}^3 . Moreover, the tangent vector $\partial/\partial x$ spans a G -invariant sub-line bundle \mathcal{N}'_1 of $\mathcal{N}_{C_0/\mathbb{P}^3}$. Since the G -action on $C_0 \cong \mathbb{P}^1$ is transitive, we know that either $\mathcal{N}'_1 = \mathcal{N}_1$ or $\mathcal{N}'_1 \oplus \mathcal{N}_1 \cong \mathcal{N}_{C_0/\mathbb{P}^3}$. The latter case is not possible since $\mathcal{O}_{\mathbb{P}^1}(4) \cong \mathcal{N}_1 \hookrightarrow \mathcal{N}_{C_0/\mathbb{P}^3} \cong \mathcal{O}_{\mathbb{P}^1}(5)^{\oplus 2}$ does not split. Thus we have $\langle \partial/\partial x \rangle = \mathcal{N}'_1 = \mathcal{N}_1$. \square

Theorem 4.3. *Let T be the tangent developable surface of twisted cubic curve C_0 in \mathbb{P}^3 . Let $c \in [0, 1)$ be a rational number. Then (\mathbb{P}^3, cT) is K-semistable (resp. K-polystable) if and only if $c \leq \frac{9}{13}$ (resp. $c < \frac{9}{13}$).*

Proof. We first show the statement for K-semistability. For the “only if” part, we use Fujita-Li’s valuative criteria Theorem 2.3. Denote by $\mu : Y_0 \rightarrow \mathbb{P}^3$ the $(2, 3)$ -weighted blow up of \mathbb{P}^3 along the twisted cubic curve C_0 in the local coordinates (x, y) defined by \mathcal{N}_1 and \mathcal{N}_2 from Proposition 4.2. Let E_0 be the exceptional divisor of μ . Then we know that $E_0 \cong \mathrm{Proj}_{\mathbb{P}^1} \mathrm{Sym}(\mathcal{E}_2 \oplus \mathcal{E}_3)$ where $\mathcal{E}_2 := \mathcal{O}_{\mathbb{P}^1}(-4) \cong \iota^* \mathcal{N}_1^\vee$, $\mathcal{E}_3 := \mathcal{O}_{\mathbb{P}^1}(-6) \cong \iota^* \mathcal{N}_2^\vee$, $\mathrm{Sym}(\mathcal{E}_2 \oplus \mathcal{E}_3)$ is a $\mathbb{Z}_{\geq 0}$ -graded $\mathcal{O}_{\mathbb{P}^1}$ -algebra, and

each \mathcal{E}_i has degree i . Denote by \tilde{T} the strict transform of T in Y_0 . From the local computation that the $(2, 3)$ blow-up normalizes the cusp, we see that \tilde{T} is the normalization of T , so by [EL20, Page 31], $\tilde{T} \cong \mathbb{P}^1 \times \mathbb{P}^1$. Denote the two pencil of rulings on \tilde{T} by R and C respectively, such that μ_*R is a tangent line of C_0 and μ_*C is a conic curve. Denote by $H := \mu^*\mathcal{O}(1)$. Then we have the following relations:

$$(4.1) \quad (H \cdot R) = 1, \quad (H \cdot C) = 2, \quad (E_0 \cdot R) = 1, \quad (E_0 \cdot C) = 1.$$

Moreover, we have $\tilde{T} = 4H - 6E_0$ and $-K_{Y_0} = 4(H - E_0)$.

Claim. $H - E_0$ (and hence $-K_{Y_0}$) is nef and big on Y_0 .

Proof of claim. We first show that $H - E_0$ is nef. Indeed, since $4H - 4E_0 \sim \tilde{T} + 2E_0$, it suffices to show that $(H - E_0)|_{\tilde{T}}$ and $(H - E_0)|_{E_0}$ are both nef. Since $\tilde{T} \cong \mathbb{P}^1 \times \mathbb{P}^1$ with R and C as two rulings, (4.1) implies that $((H - E_0) \cdot R) = 0$, $((H - E_0) \cdot C) = 1$. Hence $(H - E_0)|_{\tilde{T}}$ is nef. Denote by F the fiber of the \mathbb{P}^1 -bundle $E_0 \rightarrow C_0$. Let $\Delta_{\tilde{T}} := \tilde{T} \cap E_0$. It is clear that $E_0 \cong \mathbb{P}^1 \times \mathbb{P}^1$ with F and $\Delta_{\tilde{T}}$ as the two rulings. Since $\Delta_{\tilde{T}} \sim R + C$ in \tilde{T} , (4.1) implies that $((H - E_0) \cdot \Delta_{\tilde{T}}) = 1$. Moreover, we know that $(H \cdot F) = (\mathcal{O}_{\mathbb{P}^3}(1) \cdot \mu_*F) = 0$ and $-E_0|_{E_0} \sim \mathcal{O}_{E_0}(1)$, hence $(-E_0 \cdot F) = 1$. Thus $((H - E_0) \cdot F) = 1$. This implies that $(H - E_0)|_{E_0}$ is also nef.

Next we show that $H - E_0$ is big. This is done by computing the self intersection number of $H - tE_0$ for $t \in [0, 1]$ and observing that the computation in the previous paragraph also shows that $H - tE_0$ is nef for $t \in [0, 1]$. It is clear that

$$H|_{\tilde{T}} \sim 2R + C, \quad E_0|_{\tilde{T}} \sim R + C, \quad \tilde{T}|_{\tilde{T}} \sim 2R - 2C.$$

Hence

$$(H^3) = 1, \quad (H^2 \cdot \tilde{T}) = 4, \quad (H \cdot \tilde{T}^2) = -2, \quad (\tilde{T}^3) = -8.$$

Using the fact that $E_0 = \frac{1}{6}(4H - \tilde{T})$, we see

$$(H - tE_0)^3 = ((1 - \frac{2}{3}t)H + \frac{t}{6}\tilde{T})^3 = 1 - \frac{3}{2}t^2 + \frac{2}{3}t^3.$$

When $0 \leq t \leq 1$ this is also the volume of $H - tE_0$ since it is nef. Thus $\text{vol}_{Y_0}(H - E_0) = \frac{1}{6} > 0$ which implies that $H - E_0$ is big. The claim is proved.

Next we compute the S -invariant of E_0 . Since $-K_{Y_0}$ is nef and big, and Y_0 has only quotient singularities which are klt, we know that $-K_{Y_0}$ (and hence $H - E_0$) is semi-ample. The ample model of $H - E_0$ gives a divisorial contraction $g : Y_0 \rightarrow Y'_0$ which contracts \tilde{T} to a smooth rational curve by contracting R to a point. By computation (using the intersection theory above), we have

$$g^*g_*(H - tE_0) = H - tE_0 + \left(\frac{1-t}{2}\right)\tilde{T} = H - tE_0 + \left(\frac{1-t}{2}\right)(4H - 6E_0) = (3 - 2t)(H - E_0).$$

Therefore, if $1 < t < \frac{3}{2}$, the divisor $H - tE_0$ is big and by Zariski decomposition we know that

$$\text{vol}_{Y_0}(H - tE_0) = \text{vol}_{Y_0}(g^*g_*(H - tE_0)) = \text{vol}((3 - 2t)(H - E_0)) = \frac{(3 - 2t)^3}{6}.$$

Therefore,

$$\int_0^\infty \text{vol}(H - tE_0)dt = \int_0^1 (1 - \frac{3}{2}t^2 + \frac{2}{3}t^3)dt + \int_1^{3/2} \frac{(3 - 2t)^3}{6}dt = \frac{11}{16}.$$

Thus we know that

$$A_{(\mathbb{P}^3, cT)}(E_0) = 5 - 6c, \quad S_{(\mathbb{P}^3, cT)}(E_0) = \frac{11}{16}(4 - 4c).$$

So if (\mathbb{P}^3, cT) is K-semistable, then Theorem 2.3 implies that $5 - 6c \geq \frac{11}{16}(4 - 4c)$ which is equivalent to $c \leq \frac{9}{13}$. Thus we have shown the “only if” part for K-semistability.

Next, we show the “if” part for K-semistability. By interpolation of K-stability [ADL19, Proposition 2.13], it suffices to show that $(\mathbb{P}^3, \frac{9}{13}T)$ is K-semistable. Consider the $G = \mathrm{PGL}(2, \mathbb{C})$ -action on \mathbb{P}^3 which preserves the twisted cubic curve C_0 . By Theorem 2.7, it suffices to show that $\beta_{(\mathbb{P}^3, \frac{9}{13}T)}(F) \geq 0$ for any G -invariant prime divisor F of plt type over $(\mathbb{P}^3, \frac{9}{13}T)$. Since there are only three G -orbits in \mathbb{P}^3 : C_0 , $T \setminus C_0$, and $\mathbb{P}^3 \setminus T$, we know that either $F = T$ or F is centered at C_0 . The first case is easy. For the second case, notice that localizing $(\mathbb{P}^3, \frac{9}{13}T)$ at the generic point of C_0 produces a singularity analytically isomorphic to $(\mathbb{A}^2, \frac{9}{13}(y^2 = x^3))$. Hence the $(2, 3)$ -weighted blow up in (x, y) produces the only G -invariant divisor E_0 of plt type centered at C_0 by Lemma 4.4. Therefore, the above computations show that $\beta_{(\mathbb{P}^3, \frac{9}{13}T)}(E_0) = 0$. Thus $(\mathbb{P}^3, \frac{9}{13}T)$ is K-semistable. This finishes the proof for K-semistability.

Finally, we show the statement for K-polystability. Since T is GIT polystable by [Sha81, LO18], Theorem 3.4 implies that $(\mathbb{P}^3, \epsilon T)$ is GIT polystable for $0 < \epsilon \ll 1$. Thus [ADL19, Proposition 2.13] implies that (\mathbb{P}^3, cT) is K-polystable when $0 \leq c < \frac{9}{13}$. Moreover, since (\mathbb{P}^3, cT) is K-unstable for any $c > \frac{9}{13}$ from the “only if” part for K-semistability, [ADL19, Proposition 3.18] implies that $(\mathbb{P}^3, \frac{9}{13}T)$ is not K-polystable. Thus the proof is finished. \square

Lemma 4.4. *The klt singularity $0 \in (\mathbb{A}^2, \frac{9}{13}(y^2 = x^3))$ admits a unique plt blow-up given by the $(2, 3)$ -weighted blow-up in (x, y) .*

Proof. Let $E \subset Y \xrightarrow{\pi} \mathbb{A}^2$ be the $(2, 3)$ -weighted blow-up in (x, y) . Let Γ be the different divisor on E , i.e. $K_E + \Gamma = (K_Y + E + \pi_*^{-1}D)|_E$ where $D = \frac{9}{13}(y^2 = x^3)$. Then it is not hard to see that $(E, \Gamma) \cong (\mathbb{P}^1, \frac{1}{2}[0] + \frac{9}{13}[1] + \frac{2}{3}[\infty])$. Hence $\alpha(E, \Gamma) = (1 - \max\{\frac{1}{2}, \frac{9}{13}, \frac{2}{3}\})(2 - \deg(\Gamma)) = \frac{24}{11} > 1$, which means that (E, Γ) is an exceptional log Fano pair, i.e. there is at most exceptional divisor over it with log discrepancy one (see [Pro00, Definition 4.1]). Thus [Pro00, Section 4] implies that E is the unique Kollar component over $0 \in (\mathbb{A}^2, D)$. \square

4.2. Construction of X_u . Let $\mathcal{E} := \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-4) \oplus \mathcal{O}_{\mathbb{P}^1}(-6)$ be a rank 3 vector bundle over \mathbb{P}^1 . Denote by \mathcal{E}_i the i -th direct summand line bundle of \mathcal{E} for $i = 1, 2, 3$. Denote by $\mathbb{P}\mathcal{E} := \mathrm{Proj}_{\mathbb{P}^1} \mathrm{Sym} \mathcal{E}$ where $\mathrm{Sym} \mathcal{E}$ is a $\mathbb{Z}_{\geq 0}$ -graded $\mathcal{O}_{\mathbb{P}^1}$ -algebra such that each line bundle \mathcal{E}_i has degree i . Thus $p : \mathbb{P}\mathcal{E} \rightarrow \mathbb{P}^1$ is a $\mathbb{P}(1, 2, 3)$ -bundle. It is clear that

$$\begin{aligned} p_* \mathcal{O}_{\mathbb{P}\mathcal{E}}(1) &\cong \mathcal{O}_{\mathbb{P}^1}, \\ p_*(\mathcal{O}_{\mathbb{P}\mathcal{E}}(2) \otimes p^* \mathcal{O}_{\mathbb{P}^1}(4)) &\cong \mathcal{O}_{\mathbb{P}^1}(4) \oplus \mathcal{O}_{\mathbb{P}^1}, \\ p_*(\mathcal{O}_{\mathbb{P}\mathcal{E}}(3) \otimes p^* \mathcal{O}_{\mathbb{P}^1}(6)) &\cong \mathcal{O}_{\mathbb{P}^1}(6) \oplus \mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}_{\mathbb{P}^1}. \end{aligned}$$

Let $x \in H^0(\mathbb{P}\mathcal{E}, \mathcal{O}_{\mathbb{P}\mathcal{E}}(1))$, $y \in H^0(\mathbb{P}\mathcal{E}, \mathcal{O}_{\mathbb{P}\mathcal{E}}(2) \otimes p^* \mathcal{O}_{\mathbb{P}^1}(4))$, and $z \in H^0(\mathbb{P}\mathcal{E}, \mathcal{O}_{\mathbb{P}\mathcal{E}}(3) \otimes p^* \mathcal{O}_{\mathbb{P}^1}(6))$ be non-zero sections in the last direct summand of the right-hand-side in each isomorphism above.

By the Euler sequence for weighted projective bundles (see [Har77, Ex. III.8.4] and [Dol82]), we have

$$(4.2) \quad \mathcal{O}(-K_{\mathbb{P}\mathcal{E}}) \sim \pi^*(\mathcal{E}_1^\vee \otimes \mathcal{E}_2^\vee \otimes \mathcal{E}_3^\vee) \otimes \pi^* \mathcal{O}_{\mathbb{P}^1}(-K_{\mathbb{P}^1}) \otimes \mathcal{O}_{\mathbb{P}\mathcal{E}}(1+2+3) \cong p^* \mathcal{O}_{\mathbb{P}^1}(12) \otimes \mathcal{O}_{\mathbb{P}\mathcal{E}}(6).$$

It is clear that

$$p_* \mathcal{O}_{\mathbb{P}\mathcal{E}}(6) \cong \mathcal{E}_3^{\otimes 2} \oplus (\mathcal{E}_1 \otimes \mathcal{E}_2 \otimes \mathcal{E}_3) \oplus (\mathcal{E}_1^{\otimes 3} \otimes \mathcal{E}_3) \oplus \mathcal{E}_2^{\otimes 3} \oplus (\mathcal{E}_1^{\otimes 2} \otimes \mathcal{E}_2^{\otimes 2}) \oplus (\mathcal{E}_1^{\otimes 3} \otimes \mathcal{E}_2) \oplus \mathcal{E}_1^{\otimes 6}.$$

Thus any section $s \in H^0(\mathbb{P}\mathcal{E}, \mathcal{O}_{\mathbb{P}\mathcal{E}}(-K_{\mathbb{P}\mathcal{E}}))$ can be uniquely expressed as

$$(4.3) \quad s = az^2 + f_2xyz + f_6x^3z + by^3 + f_4x^2y^2 + f_8x^4y + f_{12}x^6,$$

where $f_j \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(j))$.

Let $H_{\mathcal{E}} \in |\mathcal{O}_{\mathbb{P}\mathcal{E}}(1)|$ be the Weil divisor defined by $(x = 0)$. Let $T_{\mathcal{E}}$ be the anti-canonical divisor on $\mathbb{P}\mathcal{E}$ defined by $(z^2 - y^3 = 0)$. Then $T_{\mathcal{E}}$ and $H_{\mathcal{E}}$ intersect transversally along a smooth curve $C_{\mathcal{E}} = V(x, z^2 - y^3)$ which is a section of $\mathbb{P}\mathcal{E} \rightarrow \mathbb{P}^1$. Let $h : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow \mathbb{P}\mathcal{E}$ be the $(2, 1)$ -weighted blow up along the divisors $(T_{\mathcal{E}}, H_{\mathcal{E}})$. Let $\widetilde{E}_{\mathcal{E}}$ be the exceptional divisor of h . Denote by $\widetilde{H}_{\mathcal{E}}$ and $\widetilde{T}_{\mathcal{E}}$ the strict transforms of $H_{\mathcal{E}}$ and $T_{\mathcal{E}}$ in $\widetilde{\mathbb{P}\mathcal{E}}$ respectively. It is clear that $\widetilde{H}_{\mathcal{E}} \cong H_{\mathcal{E}} \cong \mathbb{P}^1 \times \mathbb{P}^1$.

Proposition 4.5. *With the above notation, $\widetilde{\mathbb{P}\mathcal{E}}$ is a Gorenstein canonical weak Fano threefold, i.e. $-K_{\widetilde{\mathbb{P}\mathcal{E}}}$ is nef and big. We call its anti-canonical model X_u . Then X_u is a Gorenstein canonical Fano threefold. Moreover, the birational morphism $\psi : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow X_u$ contracts $\widetilde{H}_{\mathcal{E}}$ to an isolated singularity $o \in X_u$, and is isomorphic elsewhere.*

Proof. From the geometry of the weighted blow-up h we know that $\widetilde{\mathbb{P}\mathcal{E}}$ has quotient singularities along three disjoint smooth rational curves in $\widetilde{H}_{\mathcal{E}}$, where two curves are of type $\frac{1}{2}(1, 1, 0)$, and the rest is of type $\frac{1}{3}(1, 2, 0)$. Thus $\widetilde{\mathbb{P}\mathcal{E}}$ is Gorenstein canonical.

Let $S_{\mathcal{E}}$ be an anti-canonical divisor on $\mathbb{P}\mathcal{E}$ defined by $(z^2 = y^3 + f_8x^4y + f_{12}x^6)$ where f_8 and f_{12} are general degree 8 and 12 binary forms respectively. By [Huy16, Chapter 11 §2] we have that $S_{\mathcal{E}} \rightarrow \mathbb{P}^1$ is a smooth elliptic K3 surface with a section $C_{\mathcal{E}}$, i.e. a smooth unigonal K3 surface, as the discriminant does not vanish for general binary forms f_8 and f_{12} . Denote by F a general elliptic fiber on $S_{\mathcal{E}}$. Then since $(C_{\mathcal{E}}^2) = -2$ we know that $mF + C_{\mathcal{E}}$ is ample on $S_{\mathcal{E}}$ whenever $m \geq 3$. It is easy to see that $S_{\mathcal{E}}$ and $T_{\mathcal{E}}$ are tangent along $C_{\mathcal{E}}$ which implies that $\text{ord}_{\widetilde{E}_{\mathcal{E}}}(S_{\mathcal{E}}) = 2$. Denote by $\widetilde{S}_{\mathcal{E}}$ the strict transform of $S_{\mathcal{E}}$ in $\widetilde{\mathbb{P}\mathcal{E}}$. Thus $\widetilde{S}_{\mathcal{E}} = h^*S_{\mathcal{E}} - 2\widetilde{E}_{\mathcal{E}} \sim h^*(-K_{\mathbb{P}\mathcal{E}}) - 2\widetilde{E}_{\mathcal{E}} \sim -K_{\widetilde{\mathbb{P}\mathcal{E}}}$ because $A_{\mathbb{P}\mathcal{E}}(\widetilde{E}_{\mathcal{E}}) = 3$. By (4.2), we know that $-K_{\mathbb{P}\mathcal{E}}|_{S_{\mathcal{E}}} \sim 12F + 6C_{\mathcal{E}}$. Thus

$$-K_{\widetilde{\mathbb{P}\mathcal{E}}}|_{\widetilde{S}_{\mathcal{E}}} \sim h^*(12F + 6C_{\mathcal{E}}) - 2\widetilde{E}_{\mathcal{E}}|_{\widetilde{S}_{\mathcal{E}}} \sim 4h^*(3F + C_{\mathcal{E}}).$$

Since $h : \widetilde{S}_{\mathcal{E}} \xrightarrow{\cong} S_{\mathcal{E}}$, we know that $-K_{\widetilde{\mathbb{P}\mathcal{E}}}|_{\widetilde{S}_{\mathcal{E}}}$ is ample. Because $-K_{\widetilde{\mathbb{P}\mathcal{E}}} \sim \widetilde{S}_{\mathcal{E}}$, it has non-negative intersection with any curve not contained in $\widetilde{S}_{\mathcal{E}}$, so this implies that $-K_{\widetilde{\mathbb{P}\mathcal{E}}}$ is nef. Furthermore, this implies $(-K_{\widetilde{\mathbb{P}\mathcal{E}}})^3 = (-K_{\widetilde{\mathbb{P}\mathcal{E}}})^2 \cdot \widetilde{S}_{\mathcal{E}} = (-K_{\widetilde{\mathbb{P}\mathcal{E}}}|_{\widetilde{S}_{\mathcal{E}}})^2 > 0$, so by [Laz04, Theorem 2.2.16], $-K_{\widetilde{\mathbb{P}\mathcal{E}}}$ is big. By the Kawamata-Shokurov basepoint free theorem [KM98, Theorem 3.3], $-K_{\widetilde{\mathbb{P}\mathcal{E}}}$ is semiample with ample model X_u .

Moreover, $\widetilde{S}_{\mathcal{E}}$ is disjoint from $\widetilde{H}_{\mathcal{E}}$ which implies that $-K_{\widetilde{\mathbb{P}\mathcal{E}}}|_{\widetilde{H}_{\mathcal{E}}} \sim 0$. Thus $\widetilde{H}_{\mathcal{E}}$ is contracted under ψ to a single point. Since $h \circ p$ realizes $\widetilde{H}_{\mathcal{E}}$ as a \mathbb{P}^1 -bundle over \mathbb{P}^1 , we know that the curve classes $\mathcal{O}(1, 0)$ and $\mathcal{O}(0, 1)$ on $\widetilde{H}_{\mathcal{E}}$ are not proportional in $N_1(\widetilde{\mathbb{P}\mathcal{E}})_{\mathbb{R}}$. Since $\widetilde{\mathbb{P}\mathcal{E}}$ has Picard rank 3, if we only contract $\widetilde{H}_{\mathcal{E}}$ from $\widetilde{\mathbb{P}\mathcal{E}}$ then the resulting variety has Picard rank 1 which has to be X_u . This implies that ψ is isomorphic away from $\widetilde{H}_{\mathcal{E}}$.

Denote by $o = \psi(\widetilde{H}_{\mathcal{E}})$ the unique singular point of X_u . Since $\widetilde{S}_{\mathcal{E}} \sim -K_{\widetilde{\mathbb{P}\mathcal{E}}}$ is disjoint from $\widetilde{H}_{\mathcal{E}} = \text{Exc}(\psi)$, we know that $\psi_*\widetilde{S}_{\mathcal{E}} \sim -K_{X_u}$ does not pass through o . Thus $-K_{X_u}$ is Cartier. Moreover, we have $\psi^*(-K_{X_u}) = \psi^*\psi_*(-K_{\widetilde{\mathbb{P}\mathcal{E}}}) = -K_{\widetilde{\mathbb{P}\mathcal{E}}}$ as $-K_{\widetilde{\mathbb{P}\mathcal{E}}}|_{\widetilde{H}_{\mathcal{E}}} \sim 0$. Thus $-K_{X_u}$ is ample on X_u , and ψ is crepant birational. Thus X_u is canonical as $\widetilde{\mathbb{P}\mathcal{E}}$ is canonical. Therefore, X_u is a Gorenstein canonical Fano threefold. \square

Proposition 4.6. *With the above notation, E_0 induces a K-polystable degeneration $(X_u, \frac{9}{13}T_0)$ of $(\mathbb{P}^3, \frac{9}{13}T)$.*

Proof. We perform the following birational transformations,

$$\begin{array}{ccccc}
& & \mathcal{X} & \dashrightarrow & \mathcal{X}^+ \\
& \swarrow \mu & & \searrow g & \swarrow \psi \\
\mathbb{P}^3 \times \mathbb{A}^1 & & \mathcal{Y} & \xleftarrow{h} & \mathcal{Z}
\end{array}$$

where in the central fiber we have

$$\begin{array}{ccccc}
& & Y_0 \cup \mathbb{P}\mathcal{E} & \dashrightarrow & Y'_0 \cup \widetilde{\mathbb{P}\mathcal{E}} \\
& \swarrow \mu & & \searrow g & \swarrow \psi \\
\mathbb{P}^3 & & Y'_0 \cup \mathbb{P}\mathcal{E} & \xleftarrow{h} & X_u
\end{array}$$

Recall that $C_0 \subset T$ denotes the twisted cubic curve. Using notation from Theorem 4.3 and Proposition 4.5, the maps are as follows:

- (1) $\mu : \mathcal{X} \rightarrow \mathbb{P}^3 \times \mathbb{A}^1$ is the $(2, 3, 1)$ -weighted blow up along C_0 in the central fiber, where Y_0 is the strict transform of $\mathbb{P}^3 \times \{0\}$ and $\mathbb{P}\mathcal{E}$ is the exceptional divisor.
- (2) $f : \mathcal{X} \dashrightarrow \mathcal{X}^+$ is the flip of \tilde{T} induced by the contraction g of the rulings R of \tilde{T} in the central fiber. The strict transform of Y_0 is Y'_0 and the strict transform of $\mathbb{P}\mathcal{E}$ is $\widetilde{\mathbb{P}\mathcal{E}}$.
- (3) $\psi : \mathcal{X}^+ \rightarrow \mathcal{Z}$ is the divisorial contraction of Y'_0 .

We elaborate on these maps below.

The first morphism $\mu : \mathcal{X} \rightarrow \mathbb{P}^3 \times \mathbb{A}^1$ is the $(2, 3, 1)$ -weighted blow up of $\mathbb{P}^3 \times \mathbb{A}^1$ along C_0 , in local coordinates (x, y, t) , where x, y are as in Theorem 4.3, whose exceptional divisor $\mathbb{P}\mathcal{E}$ is a $\mathbb{P}(1, 2, 3)$ bundle over C_0 . In the central fiber, this is the $(2, 3)$ -weighted blow up of \mathbb{P}^3 along $C_0 \subset T$.

Let R and C denote the two pencils of rulings ($\mu_* R$ is a tangent line of C_0 , and $\mu_* C$ is a conic curve). With this notation, g is the flipping contraction obtained by contracting the rulings R of \tilde{T} , the strict transform of T in the central fiber, and the flip f flips $\tilde{T} \subset \mathcal{X}$. Indeed, the rulings R are contractible, and the computations in Theorem 4.3 imply that they generate a $K_{\mathcal{X}} + \mathcal{T}$ -negative extremal ray in \mathcal{X} , where \mathcal{T} is the family of surfaces. Using the local structure of \mathcal{X} as a fibration over C_0 , we can obtain an explicit description of the flip. Along C_0 , using the local coordinates (x, y) in the central fiber, T has equation $y^2 = x^3$ and μ normalizes the cusp. Consider \mathcal{X} as a fibration of surfaces in a neighborhood of $p \in C_0$. In each fiber $\mathcal{X}_p = Y_{0p} \cup \mathbb{P}(1, 2, 3)$, the normal bundle of R is $\mathcal{N}_{R/\mathcal{X}_p} = \mathcal{O}_R(-2) \oplus \mathcal{O}_R(-1)$ by computation, so the contraction g creates a $\frac{1}{2}(1, 1)$ singularity at $q \in Y'_{0p}$ and the map h is the $(2, 1)$ weighted blow up of q in $\mathbb{P}(1, 2, 3)$. The structure of the central fiber of \mathcal{X}^+ is therefore exactly as described in Proposition 4.5, so in the central fiber, the strict transform of \mathcal{T} is $\tilde{T}_{\mathcal{E}}$ in $\widetilde{\mathbb{P}\mathcal{E}}$. Recall from Proposition 4.5 that $T_{\mathcal{E}}$ is the anti-canonical divisor on $\mathbb{P}\mathcal{E}$ defined by $(z^2 - y^3 = 0)$, and by construction, the strict transform of \mathcal{T} under the first morphism μ in $\mathbb{P}\mathcal{E}$ is an anti-canonical divisor with this equation.

The last morphism ψ is the divisorial contraction of Y'_0 , and X_u is the image of $\widetilde{\mathbb{P}\mathcal{E}}$ under ψ . Indeed, an easy computation shows that curves in Y'_0 are $K_{\mathcal{X}^+}$ -negative, and Proposition 4.5 shows that $Y'_0 \cap \widetilde{\mathbb{P}\mathcal{E}}$ is contractible in $\widetilde{\mathbb{P}\mathcal{E}}$, but Y'_0 has Picard rank one, hence is contractible in \mathcal{X}^+ to \mathcal{Z} . Let T_0 be the central fiber of the strict transform of \mathcal{T} in \mathcal{Z} , which is the image of $\tilde{T}_{\mathcal{E}} \subset \widetilde{\mathbb{P}\mathcal{E}}$ under ψ .

Finally, we show that $(X_u, \frac{9}{13}T_0)$ is K-polystable. By Proposition 4.3, the log Fano pair $(\mathbb{P}^3, \frac{9}{13}T)$ is K-semistable but not K-polystable. Let $G := \mathrm{PGL}(2, \mathbb{C})$ as in the proof of Theorem 4.3. Then Proposition 2.8 implies that there exists a G -invariant special divisor F over $(\mathbb{P}^3, \frac{9}{13}T)$

which induces a K-polystable degeneration and $\beta_{(\mathbb{P}^3, \frac{9}{13}T)}(F) = 0$. Since F is G -invariant of plt type by Lemma 2.6, from the proof of Theorem 4.3 there are only two such divisors: T or E_0 , and the β -invariant of E_0 (resp. of T) is zero (resp. non-zero). Hence $F = E_0$ and the proof is finished. \square

In particular, a consequence of Proposition 4.6 is that X_u admits a \mathbb{Q} -Gorenstein smoothing to \mathbb{P}^3 . Let L be the \mathbb{Q} -Cartier Weil divisor on X_u obtained as the limit of $\mathcal{O}_{\mathbb{P}^3}(1)$ according to Lemma 3.3. The next result determines the Cartier index of L .

Lemma 4.7. *With the above notation, the Cartier index of L on X_u is 4.*

Proof. By Lemma 3.3 and Proposition 4.5, we know that L is a \mathbb{Q} -Cartier Weil divisor on X_u such that $4L \sim -K_{X_u}$ is Cartier. Thus the Cartier index of L divides 4. Recall that $h : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow \mathbb{P}\mathcal{E}$ is a $(2, 1)$ -weighted blow up along $(T_{\mathcal{E}}, H_{\mathcal{E}})$ where the center $C_{\mathcal{E}}$ is a section of $\mathbb{P}\mathcal{E} \rightarrow \mathbb{P}^1$. Hence on $\widetilde{\mathbb{P}\mathcal{E}}$ the divisor $\widetilde{T}_{\mathcal{E}} := h_*^{-1}T_{\mathcal{E}}$ and the h -exceptional divisor $\widetilde{E}_{\mathcal{E}}$ have transverse intersection along a section of $\widetilde{\mathbb{P}\mathcal{E}} \rightarrow \mathbb{P}^1$. Denote by $R_{\mathcal{E}}$ a fiber of $E_{\mathcal{E}} \rightarrow \mathbb{P}^1$. Then the above argument implies that $(\widetilde{T}_{\mathcal{E}} \cdot R_{\mathcal{E}}) = 1$. By the proof of Proposition 4.5, we know that $\widetilde{T}_{\mathcal{E}} \sim -K_{\widetilde{\mathbb{P}\mathcal{E}}} = \psi^*(-K_{X_u}) \sim 4\psi^*L$. Thus we have $(L \cdot \psi_*R_{\mathcal{E}}) = \frac{1}{4}(\widetilde{T}_{\mathcal{E}} \cdot R_{\mathcal{E}}) = \frac{1}{4}$ which implies that L has Cartier index 4. \square

4.3. GIT for Weierstrass models of unigonal K3 surfaces. By construction of X_u , the surface T_0 has the structure of a Weierstrass elliptic fibration over the twisted cubic curve. Repeating these birational transforms for any family $(\mathbb{P}^3 \times \mathbb{A}^1, \mathcal{T})$ whose central fiber T_0 is the tangent developable surface, the new central fiber also has this structure. These elliptic surfaces naturally live inside (birational models of) $\mathbb{P}\mathcal{E}$, a $\mathbb{P}(1, 2, 3)$ bundle over C_0 . Next, we relate this construction to GIT for Weierstrass elliptic surfaces.

We recall some notation from Section 4.2, namely the construction of Weierstrass elliptic surfaces inside a $\mathbb{P}(1, 2, 3)$ -bundle. As above, let $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-4) \oplus \mathcal{O}_{\mathbb{P}^1}(-6)$ and let \mathcal{E}_i denote the i th direct summand line bundle where \mathcal{E}_i has degree i . Then we obtain the $\mathbb{P}(1, 2, 3)$ bundle $p : \mathbb{P}\mathcal{E} \rightarrow \mathbb{P}^1$, where $\mathbb{P}\mathcal{E} := \text{Proj}_{\mathbb{P}^1} \text{Sym} \mathcal{E}$. Consider the affine space $\mathbf{A} := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(8)) \oplus H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(12))$ of dimension 22 parametrizing pairs (A, B) of degree 8 and 12 binary forms respectively. In this way, for each pair $(A, B) \in \mathbf{A}$ one can associate a *Weierstrass elliptic surface* $S_{(A, B)} \in |-K_{\mathbb{P}\mathcal{E}}|$ whose equation is given by

$$S_{(A, B)} = (z^2 = y^3 + Ax^4y + Bx^6),$$

where x, y, z have degrees one, two, and three respectively. Note that $S_{(0, 0)} = T_{\mathcal{E}}$ by definition.

The discriminant divisor $4A^3 + 27B^2$ vanishes at the singular fibers, and for every point $p \in \mathbb{P}^1$, one has $\text{ord}_p(A) \leq 3$ or $\text{ord}_p(B) \leq 5$. The latter condition is equivalent to the surface having only ADE singularities. Furthermore, one can show that a pair (A, B) satisfying these conditions yields a unigonal K3 surface with ADE singularities, and vice versa.

Next, we introduce the GIT set-up for Weierstrass elliptic surfaces. Clearly the affine space \mathbf{A} admits an $(\text{SL}(2, \mathbb{C}) \times \mathbb{G}_m)$ -action given by

$$(g, t) \cdot (A, B) := (t^4 A \circ g, t^6 B \circ g).$$

Let $\mathcal{P} := [(\mathbf{A} \setminus \{0\})/\mathbb{G}_m]$ be the weighted projective stack where \mathbb{G}_m acts as above. Let \mathbf{P} be the weighted projective space as the coarse moduli space of \mathcal{P} . Then we have $\mathbf{P} \cong \mathbb{P}(2^9, 3^{13})$. The $\text{SL}(2, \mathbb{C})$ -action on \mathbf{A} descends to an $\text{SL}(2, \mathbb{C})$ -action on $(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(1))$. We say that a non-zero pair $(A, B) \in \mathbf{A} \setminus \{0\}$ is GIT (poly/semi)stable if $[A, B] \in \mathbf{P}$ is GIT (poly/semi)stable with respect to the $\text{SL}(2, \mathbb{C})$ -action on $(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(1))$. Denote by $\mathbf{A}^{\text{ss}} \subset \mathbf{A} \setminus \{0\}$ the GIT semistable open locus.

In [Mir81], Miranda constructs a compact moduli space parametrizing Weierstrass fibrations over \mathbb{P}^1 using GIT following the above discussion. In particular, Miranda gives (see [Mir81, Proposition 5.1]) a criterion for when a pair (A, B) is GIT (semi)stable based on the valuations.

Remark 4.8. Before stating the following theorem, we point out that Miranda's setup (and the discussion that follows in [OO21] and [AB19]) considers Weierstrass elliptic surfaces embedded in a \mathbb{P}^2 bundle, instead of a $\mathbb{P}(1, 2, 3)$ bundle, which is more convenient for our work. The same results hold in our setup, as the underlying surfaces are the same.

Theorem 4.9. *If $(A, B) \in \mathbf{A} \setminus \{0\}$ is GIT semistable, then the surface $S_{(A, B)}$ is slc.*

Proof. See [OO21, Proposition 7.4] for an explicit description of the GIT compactification of Weierstrass elliptic surfaces (see also [AB19, Section 10]). \square

Lemma 4.10. *Let $S \in |-K_{X_u}|$ be an anti-canonical divisor. If S does not pass through the singular point $o \in X_u$, then there exists an automorphism $\varphi \in \text{Aut}(X_u)$ such that (X_u, φ^*S) is the birational transform of $(\mathbb{P}\mathcal{E}, S_{(A, B)})$ for some $(A, B) \in \mathbf{A}$. Moreover, S is isomorphic to the Weierstrass elliptic surface $S_{(A, B)}$.*

Proof. Recall from Proposition 4.5 that there is a diagram $\mathbb{P}\mathcal{E} \xleftarrow{h} \widetilde{\mathbb{P}\mathcal{E}} \xrightarrow{\psi} X_u$ where h is a weighted blow-up and ψ is taking the anti-canonical model. Let $S_{\mathcal{E}} := h_*\psi^*S$. Then we know that $\text{ord}_{\widetilde{\mathbb{P}\mathcal{E}}}(S_{\mathcal{E}}) = 2$ which implies that $S_{\mathcal{E}}$ is tangent to $T_{\mathcal{E}}$ along the curve $C_{\mathcal{E}}$. Thus by (4.3) we know that $S_{\mathcal{E}}$ has the equation $(a(z^2 - y^3) + f_6x^3z + f_4x^2y^2 + f_8x^4y + f_{12}x^6 = 0)$. Since $o \notin S$, we know that $S_{\mathcal{E}}$ does not contain $H_{\mathcal{E}} = (x = 0)$ which implies $a \neq 0$. Hence we may assume that $a = 1$. Let $\varphi_{\mathcal{E}} \in \text{Aut}(\mathbb{P}\mathcal{E})$ be $\varphi_{\mathcal{E}}(x, y, z) := (x, y + \frac{f_4}{3}x^2, z - \frac{f_6}{2}x^3)$. Then clearly $\varphi_{\mathcal{E}}^*S_{\mathcal{E}}$ is defined by the equation $(z^2 = y^3 + Ax^4y + Bx^6)$ which is a Weierstrass elliptic surface, i.e. $\varphi_{\mathcal{E}}^*S_{\mathcal{E}} = S_{(A, B)}$. Hence $\varphi_{\mathcal{E}}$ induces the desired $\varphi \in \text{Aut}(X_u)$. Since $S_{\mathcal{E}}$ is tangent to $T_{\mathcal{E}}$ along $C_{\mathcal{E}}$, we know that $h_*^{-1}S_{\mathcal{E}}$ is isomorphic to $S_{\mathcal{E}}$ and disjoint with $\tilde{H}_{\mathcal{E}}$, which implies $S = \psi_*(h_*^{-1}S_{\mathcal{E}}) \cong h_*^{-1}S_{\mathcal{E}} \cong S_{\mathcal{E}}$. \square

By Lemma 4.10, every unigonal K3 surface $S_{(A, B)}$ with ADE singularities is isomorphic to an anti-canonical divisor $S \in |-K_{X_u}|$ not passing through o . Since the ample \mathbb{Q} -Cartier divisor L on X_u is Cartier away from o , and $(L^2 \cdot S) = \frac{1}{16}(-K_{X_u})^3 = 4$, we know that every polarized unigonal K3 surface of degree 4 is isomorphic to some $(S, L|_S)$.

Theorem 4.11. *Let $S, S' \in |-K_{X_u}|$ be two divisors where $o \notin S$ and $o \in S'$. Let $(A, B) \in \mathbf{A}$ be a pair such that φ^*S is the birational transform of $S_{(A, B)}$ for some $\varphi \in \text{Aut}(X_u)$.*

- (1) $(X_u, \frac{9}{13}S)$ is always K-semistable, and it is K-polystable if and only if $\varphi^*S = T_0$;
- (2) if $c \in (\frac{9}{13}, 1) \cap \mathbb{Q}$, then (X_u, cS) is K-(poly/semi)stable if and only if $(A, B) \in \mathbf{A} \setminus \{0\}$ is GIT (poly/semi)stable;
- (3) if $c \in (0, \frac{9}{13}) \cap \mathbb{Q}$, then (X_u, cS) is K-unstable;
- (4) (X_u, cS') is K-unstable for any rational $c \in (0, 1)$.

Proof. For simplicity, we always assume that φ is the identity as guaranteed by Lemma 4.10, i.e. $S = \psi_*(h_*^{-1}S_{(A, B)})$ for some $(A, B) \in \mathbf{A}$.

(1) Since $\lim_{t \rightarrow 0} (t^4 A, t^6 B) = (0, 0)$, the \mathbb{G}_m -action on \mathbf{A} induces a special degeneration of $(X_u, \frac{9}{13}S)$ to $(X_u, \frac{9}{13}T_0)$ which is K-polystable by Proposition 4.6. Hence it follows from openness of K-semistability [BLX22, Xu20] and the fact that $(X_u, S) \not\cong (X_u, T_0)$ whenever $(A, B) \neq (0, 0)$.

(2) To start with, notice that (\mathbb{P}^3, cT) admits a special degeneration to (X_u, cT_0) for $c \leq \frac{9}{13}$. Since (\mathbb{P}^3, cT) is K-polystable for $c < \frac{9}{13}$ by Theorem 4.3, we know that (X_u, cT_0) is K-unstable for $c < \frac{9}{13}$. Since $(X_u, \frac{9}{13}T_0)$ is K-polystable by Proposition 4.6, we know that (X_u, cT_0) is K-unstable for any $c \neq \frac{9}{13}$ by [ADL19, Proposition 3.18]. Thus we may assume $(A, B) \neq 0$.

We first show the “only if” part, that is, K implying GIT. This follows from the Paul-Tian criterion [ADL19, Theorem 2.22]. There is a universal family of Weierstrass elliptic surfaces $(\mathbb{P}\mathcal{E} \times \mathbf{A}, \mathcal{S}_{\mathcal{E}, \mathbf{A}}) \rightarrow \mathbf{A}$ where the fiber over $(A, B) \in \mathbf{A}$ is $(\mathbb{P}\mathcal{E}, (z^2 = y^3 + Ax^4y + Bx^6))$. By

performing the weighted blow-up h and anti-canonical morphism ψ in families, we obtain a universal family $(X_u \times \mathbf{A}, \mathcal{S}_\mathbf{A}) \rightarrow \mathbf{A}$. Clearly the \mathbb{G}_m -action on \mathbf{A} lifts naturally to $\mathbb{P}\mathcal{E} \times \mathbf{A}$ (where $t \cdot ([x, y, z], (A, B)) := ([tx, y, z], (t^4 A, t^6 B))$) and $X_u \times \mathbf{A}$ such that both $\mathcal{S}_{\mathcal{E}, \mathbf{A}}$ and $\mathcal{S}_\mathbf{A}$ are \mathbb{G}_m -invariant. Hence taking the stacky quotient of \mathbb{G}_m over $\mathbf{A} \setminus \{0\}$, we obtain a \mathbb{Q} -Gorenstein log Fano family $\pi_{\mathcal{P}} : (\mathcal{X}_{\mathcal{P}}, \mathcal{S}_{\mathcal{P}}) \rightarrow \mathcal{P}$. We show that the assumptions of [ADL19, Theorem 2.22] are satisfied for $\pi_{\mathcal{P}} : (\mathcal{X}_{\mathcal{P}}, c\mathcal{S}_{\mathcal{P}}) \rightarrow \mathcal{P}$ when $c \in (\frac{9}{13}, 1)$. Assumptions (a) and (b) are straightforward. For assumption (c), it suffices to show that the CM line bundle $\lambda_{\text{CM}, \pi_{\mathcal{P}}, c\mathcal{S}_{\mathcal{P}}}$ descends to an ample \mathbb{Q} -line bundle $\Lambda_{\mathbf{P}, c}$ on \mathbf{P} for $c \in (\frac{9}{13}, 1)$. For a general choice of (A, B) , the surface S is klt which implies that (X_u, S) is plt. Hence Theorem 2.10 implies that $(X, (1 - \epsilon)S)$ uniformly K-stable for $0 < \epsilon \ll 1$. Thus interpolation [ADL19, Proposition 2.13] implies that (X_u, cS) is uniformly K-stable for any $c \in (\frac{9}{13}, 1)$. Thus [CP21, Pos22, XZ20] implies that $\Lambda_{\mathbf{P}, c}$ is big, hence it is ample since \mathbf{P} has Picard rank 1. Thus the proof of the “only if” part is finished by [ADL19, Theorem 2.22].

For the “if” part, notice that by Theorem 4.9 any GIT semistable S is slc. Thus (X_u, cS) is K-semistable for any $c \in (\frac{9}{13}, 1)$ by part (1) and interpolation for K-stability [ADL19, Proposition 2.13]. For a general choice of (A, B) , the above discussion shows that (X_u, cS) is K-stable for any $c \in (\frac{9}{13}, 1)$. If S is GIT polystable, then it suffices to show that $(X_u, (\frac{9}{13} + \epsilon)S)$ is K-polystable, since then (X_u, cS) is K-polystable by interpolation [ADL19, Proposition 2.13]. Let $(X_0, (\frac{9}{13} + \epsilon)S_0)$ be the K-polystable degeneration of $(X_u, (\frac{9}{13} + \epsilon)S)$. Then we know that $(X_0, \frac{9}{13}S_0)$ is K-semistable. Since $(X_u, \frac{9}{13}T_0)$ is the K-polystable degeneration of $(X_u, \frac{9}{13}S)$, by [LWX21] we have a sequence of special degenerations $(X_u, \frac{9}{13}S) \rightsquigarrow (X, \frac{9}{13}S_0) \rightsquigarrow (X_u, \frac{9}{13}T_0)$. This implies that $X \cong X_u$ and S_0 does not pass through the singular point on X . We may assume that $X = X_u$ for simplicity. By Lemma 4.10, we can find a 1-parameter family of automorphisms $(\varphi_t) \in \text{Aut}(X_u)$ over a pointed curve $(B, 0)$ such that $\varphi_t^* S$ ($t \neq 0$) and $\varphi_0^* S_0$ are all in Weierstrass form, and $\lim_{t \rightarrow 0} \varphi_t^* S = \varphi_0^* S_0$. Since $(X_u, (\frac{9}{13} + \epsilon)\varphi_0^* S_0)$ is K-polystable, we know that $\varphi_0^* S_0 \neq T_0$, i.e. $\varphi_0^* S_0$ corresponds to a non-zero pair $(A_0, B_0) \in \mathbf{A}$. By the “only if” part, (A_0, B_0) is GIT polystable. Hence the separatedness of GIT quotient implies that $[A, B]$ and $[A_0, B_0]$ belong to the same $\text{SL}(2, \mathbb{C})$ -orbit in \mathbf{P} . In particular, $(X_u, (\frac{9}{13} + \epsilon)S) \cong (X_u, (\frac{9}{13} + \epsilon)S_0)$ are K-polystable. The proof is finished.

(3) Assume to the contrary that (X_u, cS) is K-semistable for some $c < \frac{9}{13}$. By openness of K-semistability [BLX22, Xu20], we may choose S such that (A, B) is a general pair of binary forms. Thus (A, B) is GIT stable which implies that $(X, (\frac{9}{13} + \epsilon)S)$ is K-stable by part (2). By interpolation [ADL19, Proposition 2.13] we know that $(X, \frac{9}{13}S)$ is K-stable, contradicting part (1). Thus part (3) is proved.

(4) By Proposition 4.5, the divisor S' is Cartier as X_u is Gorenstein. Besides, the birational morphism $\psi : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow X_u$ contracts the prime divisor $\widetilde{H}_{\mathcal{E}}$ to $o \in X_u$, where $\widetilde{H}_{\mathcal{E}} \cong \mathbb{P}^1 \times \mathbb{P}^1$ and $K_{\widetilde{\mathbb{P}\mathcal{E}}} \mid_{\widetilde{H}_{\mathcal{E}}} \sim 0$. Hence by adjunction we have $\widetilde{H}_{\mathcal{E}} \mid_{\widetilde{H}_{\mathcal{E}}} = K_{\widetilde{H}_{\mathcal{E}}} + \Gamma$ where Γ is the different divisor. Denote the two families of fibers of $\mathbb{P}^1 \times \mathbb{P}^1$ by F_a and F'_b for $a, b \in \mathbb{P}^1$. Then by construction we have that $\Gamma = \frac{1}{2}F_0 + \frac{1}{2}F_1 + \frac{2}{3}F_\infty$ under a suitable choice of coordinates. Thus $A_{X_u}(\widetilde{H}_{\mathcal{E}}) = 1$ and $\text{vol}_{X_u, o}(\widetilde{H}_{\mathcal{E}}) = (-K_{\widetilde{H}_{\mathcal{E}}} - \Gamma)^2 = \frac{4}{3}$. Here we refer to the survey article [LLX20] for the background of normalized volumes. Since S is Cartier and passes through $o = \psi(\widetilde{H}_{\mathcal{E}})$, we have $\text{ord}_{\widetilde{H}_{\mathcal{E}}}(S) \geq 1$. Hence we have

$$\widehat{\text{vol}}_{(X_u, cS), o}(\widetilde{H}_{\mathcal{E}}) = (A_{X_u}(\widetilde{H}_{\mathcal{E}}) - \text{cord}_{\widetilde{H}_{\mathcal{E}}}(S))^3 \text{vol}_{X_u, o}(\widetilde{H}_{\mathcal{E}}) \leq \frac{4}{3}(1 - c)^3.$$

On the other hand, we have $\frac{27}{64}(-K_{X_u} - cS)^3 = 27(1 - c)^3 > \widehat{\text{vol}}_{(X_u, cS), o}(\widetilde{H}_{\mathcal{E}})$, which implies that (X_u, cS) is K-unstable by [LL19]. \square

4.4. Deformation theory of X_u . In this subsection, we show that the allowed deformations of X_u are unobstructed, and any such small deformation of X_u is isomorphic to itself or \mathbb{P}^3 .

In the moduli stack $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$, we only consider deformations induced by the index one cover of the \mathbb{Q} -Cartier Weil divisor L that is the limit of $\mathcal{O}_{\mathbb{P}^3}(1)$, see Lemma 3.3. In the case of X_u , the divisor L is 4-Cartier by Lemma 4.7.

Lemma 4.12. *The unique singular point $o \in X_u$ admits a special degeneration to an orbifold cone singularity $x_0 \in X_{u,0}$. Under this special degeneration, the divisor L degenerates to a \mathbb{Q} -Cartier Weil divisor L_0 on $X_{u,0}$, whose Cartier index is 4. Moreover, the index one cover $Y \rightarrow X_{u,0}$ of L_0 is a toric Gorenstein canonical singularity of the form $(xy - zw = 0) \subset \mathbb{A}^4/\mu_3$, where the action of μ_3 is given by $\zeta_3 \cdot (x, y, z, w) = (\zeta_3 x, \zeta_3^2 y, \zeta_3^2 z, \zeta_3 w)$ with $\zeta_3 = e^{\frac{2\pi i}{3}}$.*

Proof. By construction from Section 4.2 and the proof of Theorem 4.11(4), we know that $(\tilde{H}_{\mathcal{E}}, \Gamma) \cong (\mathbb{P}^1 \times \mathbb{P}^1, \frac{1}{2}F_0 + \frac{1}{2}F_1 + \frac{2}{3}F_{\infty})$ and $\tilde{H}_{\mathcal{E}}|_{\tilde{H}_{\mathcal{E}}} = K_{\tilde{H}_{\mathcal{E}}} + \Gamma$ where Γ is the different divisor. Thus $\tilde{H}_{\mathcal{E}}$ induces a special degeneration of $(o \in X_u)$ to the orbifold cone $X_{u,0} := C_a(\mathbb{P}^1 \times \mathbb{P}^1, -K_{\mathbb{P}^1 \times \mathbb{P}^1} - \Gamma)$ (as defined in [LX20, Section 2.4] and [LZ22, Section 2.4]). The notion of special degeneration in the local setting comes from Kollar components over the singular point (c.f. [Xu21, Definition 4.24]). Since $X_{u,0}$ is an anti-canonical cone over a klt log Fano pair with standard coefficients, it is Gorenstein canonical ([Kol13, Lemma 3.1, Proposition 3.14]).

Let L_0 be the degeneration of L to $X_{u,0}$. Then clearly $4L_0 \sim -K_{X_{u,0}}$ is Cartier, and $2L_0$ is not Cartier because $2L$ is not Cartier. Thus we can take an index 1 cover $Y \rightarrow X_{u,0}$ of L_0 which has degree 4. In fact, there is a local universal cover of $o \in X_{u,0}$ as follows:

$$C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 1)) \rightarrow C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2, 2)) \rightarrow C_a(\mathbb{P}^1 \times \mathbb{P}^1, -K_{\mathbb{P}^1 \times \mathbb{P}^1} - \Gamma).$$

Here the first map is raising the polarization to the second tensor power, and the second map comes from the quotient map $\mathbb{P}^1 \rightarrow (\mathbb{P}^1, \frac{1}{2}[0] + \frac{1}{2}[1] + \frac{2}{3}[\infty])$ of degree 6, induced by the \mathfrak{S}_3 -action on \mathbb{P}^1 generated by $[u_0, u_1] \mapsto [u_0, \zeta_3 u_1]$ and $[u_0, u_1] \mapsto [u_1, u_0]$. Thus we have a group of order 12 (isomorphic to a binary dihedral group) acting on the singularity $C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 1))$ whose quasi-étale quotient is precisely $x_0 \in X_{u,0}$.


Since L_0 has Cartier index 4, the index 1 cover of L_0 on $X_{u,0}$ is a μ_3 -quotient of the singularity $C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 1))$. Let $([u_0, u_1], [v_0, v_1])$ be the projective coordinate on $\mathbb{P}^1 \times \mathbb{P}^1$, then we may identify $C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 1))$ with $\tilde{Y} := (xy - zw = 0) \subset \mathbb{A}^3$, where

$$(x, y, z, w) = (u_0 v_0, u_1 v_1, u_1 v_0, u_0 v_1).$$

The group μ_3 acts on \mathbb{P}^1 as generated by $[u_0, u_1] \mapsto [u_0, \zeta_3 u_1]$. After lifting to the anti-canonical cone $C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2, 2))$, the μ_3 -action becomes $(u_0^2, u_0 u_1, u_1^2) \mapsto (\zeta_3^2 u_0^2, u_0 u_1, \zeta_3 u_1^2)$ where v_0^2 , $v_0 v_1$, and v_1^2 are μ_3 -invariant. Thus the only lifting of μ_3 -action on \tilde{Y} is given by $(x, y, z, w) \mapsto (\zeta_3 x, \zeta_3^2 y, \zeta_3^2 z, \zeta_3 w)$. \square

Therefore, we must understand the deformation theory of these singularities. Because they are toric Gorenstein threefold singularities, Altmann's method described in [Altmann97] applies to compute the miniversal deformation space.

Lemma 4.13. *The singularity $0 \in Y = (xy - zw) \subset \mathbb{A}^4/\mu_3$ with above action of μ_3 is isomorphic to the affine toric Gorenstein threefold singularity V defined by the cone in \mathbb{R}^3 with two dimensional polytope Q as the hyperplane section $t = 1$ pictured below.*



Proof. Let $M' = \mathbb{Z}^3$ be the standard lattice. Let $\omega \subset M'_{\mathbb{R}} = \mathbb{R}^3$ be the cone generated by $(0, 0, 1)$, $(1, 1, 1)$, $(1, 0, 1)$, and $(0, 1, 1)$ corresponding to the variables x, y, z , and w . Then the toric variety $\tilde{Y} := \mathbb{C}[x, y, z, w]/(xy - zw)$ is isomorphic to $\text{Spec } \mathbb{C}[\omega \cap M']$. Let $N' := \text{Hom}(M', \mathbb{Z}) = \mathbb{Z}^3$ and $\sigma = \omega^{\vee} \subset N'_{\mathbb{R}} = \mathbb{R}^3$, then computations shows that σ is generated by $(1, 0, 0)$, $(0, 1, 0)$, $(0, -1, 1)$, and $(-1, 0, 1)$. The μ_3 -action on \tilde{Y} induces an index 3 sublattice $M \subset M'$ corresponding to μ_3 -invariant monomials. It is clear that xz , $\frac{w}{x}$, and x^3 are μ_3 -invariant. Thus M is generated by $(1, 0, 2)$, $(0, 1, 0)$, $(0, 0, 3)$ which correspond to the above three μ_3 -invariant monomials. Let $N = \text{Hom}(M, \mathbb{Z})$ be the dual lattice. Then computing dual basis shows that N is generated by $(1, 0, 0)$, $(0, 1, 0)$, and $(-\frac{2}{3}, 0, \frac{1}{3})$. We pick a new basis $\vec{v}_1 = (1, -1, 0)$, $\vec{v}_2 = (-\frac{2}{3}, 0, \frac{1}{3})$, and $\vec{v}_3 = (0, 1, 0)$ of N . Then under the basis \vec{v}_i , the cone σ is generated by

$$\vec{v}_1 + \vec{v}_3 = (1, 0, 0), \quad \vec{v}_3 = (0, 1, 0), \quad 2\vec{v}_1 + 3\vec{v}_2 + \vec{v}_3 = (0, -1, 1), \quad \text{and } \vec{v}_1 + 3\vec{v}_2 + \vec{v}_3 = (-1, 0, 1).$$

This shows that $Y = \tilde{Y}/\mu_3$ corresponds to the polytope Q from the picture as

$$Q = \{(a, b) \in \mathbb{R}^2 \mid a\vec{v}_1 + b\vec{v}_2 + \vec{v}_3 \in \sigma\}.$$

The proof is finished. \square

Proposition 4.14. *The base of the miniversal deformation space for the singularity $0 \in Y$ is a smooth curve. In particular, the singularity $0 \in Y$ has unobstructed deformations.*

Proof. A more general result is proved in [Pet22a, Proposition 4.4 (ii)]. We provide a proof here for readers' convenience. The miniversal base space of a toric Gorenstein threefold singularity is determined by the corresponding two-dimensional polytope. Indeed, in [Alt97, Theorem 5.1], Altmann constructs a flat deformation over a base space \tilde{M} from the polytope, and proves it is the miniversal deformation space in [Alt97, Corollary 7.2]. For more information on the construction of \tilde{M} , see [Alt97, Definition 2.2, Theorem 2.4].

To obtain equations for the space \tilde{M} , label the edges of the polytope Q in a counterclockwise fashion starting at the origin: $d^1 = (1, 0)$, $d^2 = (1, 3)$, $d^3 = (-1, 0)$, $d^4 = (-1, -3)$. Define the vector valued polynomial $g_k(t) = \sum_{i=1}^4 t_i^k d^i$. The inner products $\langle g_k(t), (1, 0) \rangle$ and $\langle g_k(t), (0, 1) \rangle$ define two polynomials $g_{k,x}(t)$ and $g_{k,y}(t)$. We define the ideal

$$\mathcal{J} = \{g_{k,x}(t), g_{k,y}(t) \mid k \geq 1\} \subset \mathbb{C}[t_1, t_2, t_3, t_4].$$

Let $M = \text{Spec } \mathbb{C}[t_1, t_2, t_3, t_4]/\mathcal{J} \subset \mathbb{A}^4$. By [Alt97, Theorem 7.4], M is defined by equations in $\mathbb{C}[t_i - t_j]$, and the miniversal base space \tilde{M} is defined by $\mathcal{J} \cap \mathbb{C}[t_i - t_j]$, or the pre-image of M under the canonical projection $\mathbb{C}^4 \rightarrow \mathbb{C}^4/\mathbb{C} \cdot (1, 1, 1, 1)$.

Plugging in the values of d^i , we see that \mathcal{J} is defined by the equations $t_1^k + t_2^k - t_3^k - t_4^k$ and $3t_2^k - 3t_4^k$, for $k \geq 1$ which reduces to $t_1 - t_3$ and $t_2 - t_4$. Hence, $M = \text{Spec } \mathbb{C}[t_1, t_2]$, and $\tilde{M} = \text{Spec } \mathbb{C}[t_1 - t_2] \cong \mathbb{A}^1$. Therefore, the miniversal base space of the singularity Y is a smooth curve, so Y has unobstructed deformations. \square

The miniversal base space of the singular point of $X_{u,0}$ where L_0 deforms in a \mathbb{Q} -Gorenstein family is given by the $\mathbb{Z}/(4)$ -invariant part of the miniversal base space of Y , which is a proper

subspace of \widetilde{M} . However, by construction $x_0 \in X_{u,0}$ deforms to $o \in X_u$ which admits a smoothing to \mathbb{P}^3 , so the miniversal base space must be at least one-dimensional. Therefore, it must be all of \widetilde{M} and we obtain the following corollary.

Corollary 4.15. *The singularities $x_0 \in X_{u,0}$ and $o \in X_u$ are formally isomorphic. As a result, the singular point $o \in X_u$ has unobstructed deformations where L deforms in a \mathbb{Q} -Gorenstein family.*

To finish the study of the deformation theory of X_u , we must show that there are no local-to-global obstructions in extending the local deformation to a global deformation of X_u , which is done by the following lemma.

Lemma 4.16. *There are no local-to-global obstructions for deformations of X_u .*

Proof. By [Pet22b, Proposition 2.3], it suffices to show that $H^2(X_u, T_{X_u}) = 0$.

For simplicity, write $X = X_u$. By construction of X , there is a small \mathbb{Q} -factorialization $\pi : Z \rightarrow X$ such that Z has D_5 singularities along an irreducible curve, contracted to the singular point $o \in X$. The map $\psi : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow X$ factors through π and the fibration structure of $\widetilde{\mathbb{P}\mathcal{E}} \rightarrow \mathbb{P}^1$ descends to a fibration $f : Z \rightarrow \mathbb{P}^1$. Furthermore, $f : Z \rightarrow \mathbb{P}^1$ is an isotrivial fibration of \mathbb{Q} -Fano surfaces S each with an isolated D_5 singularity. In a neighborhood of any point p in the singular locus of Z , the threefold looks like $S \times T$, where T is a smooth curve.

From the five-term exact sequence from the Leray spectral sequence

$$H^p(X, R^q \pi_* T_Z) \Rightarrow H^{p+q}(Z, T_Z)$$

and the fact that $\pi_* T_Z = T_X$, there is a sequence

$$0 \rightarrow H^1(X, T_X) \rightarrow H^1(Z, T_Z) \rightarrow H^0(X, R^1 \pi_* T_Z) \rightarrow H^2(X, T_X) \rightarrow H^2(Z, T_Z).$$

In fact, this can be extended to

$$0 \rightarrow H^1(X, T_X) \rightarrow H^1(Z, T_Z) \rightarrow H^0(X, R^1 \pi_* T_Z) \rightarrow H^2(X, T_X) \rightarrow H^2(Z, T_Z) \rightarrow H^1(X, R^1 \pi_* T_Z)$$

as in general the second-to-last term is $\ker H^2(Z, T_Z) \rightarrow H^0(X, R^2 \pi_* T_Z)$, but the latter is 0 as the fibers of π have dimension at most 1. In the following Lemma, we will show that $R^1 \pi_* T_Z = 0$, so we find that $H^1(X, T_X) = H^1(Z, T_Z)$ and $H^2(X, T_X) = H^2(Z, T_Z)$.

Finally, we show that $H^2(Z, T_Z) = 0$. First note that T_Z and its dual $\Omega_Z^{[1]}$ are Cohen-Macaulay: their restriction to any (Cartier) fiber S of $f : Z \rightarrow \mathbb{P}^1$ is S_2 and $\dim S = 2$, so the restriction to the Cartier fiber is Cohen Macaulay. (To see that the restriction is S_2 , one may use the sequence $0 \rightarrow \mathcal{O}_S(-S) \rightarrow \Omega_Z^{[1]}|_S \rightarrow \Omega_S^{[1]} \rightarrow 0$; shown to be exact using the description of Z as a fibration over \mathbb{P}^1 .) By Serre Duality, then $H^2(Z, T_Z) = H^1(Z, \Omega_Z^{[1]}(K_Z))^\vee$.

The proof that $H^1(Z, \Omega_Z^{[1]}(K_Z)) = 0$ then follows by the same logic used in [Nam97, Proposition 4]. The input [Nam97, Proposition 2] holds (and the proof holds verbatim), replacing Z by X and Y by X_u in the author's notation, i.e. if D is an anticanonical section of X_u and D' its strict transform on Z , we have $\text{Pic}(Z) \rightarrow \text{Pic}(D')$ is injective. Therefore, from the standard exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O} \rightarrow \mathcal{O}^\times \rightarrow 0$$

and vanishing of $H^i(Z, \mathcal{O}_Z)$ for $i > 0$, we have an injection $H^2(Z, \mathbb{Z}) \rightarrow H^2(D', \mathbb{Z})$. Because Z has only quotient singularities, it is a V -manifold as in [Ste77b, Theorem pg. 4], and the Hodge structure is pure and $H^{1,1}(Z)$ is identified with $H^1(Z, \Omega_Z^{[1]})$. By Hodge theory, the injection on cohomology then induces an injection on the parts of the Hodge decomposition $H^{p,q}(Z) \rightarrow H^{p,q}(D')$ with $p+q = 2$. Therefore, we have an injection $H^1(Z, \Omega_Z^{[1]}) \rightarrow H^1(D', \Omega_{D'}^1)$.

Noting that the exact sequences in the proof of [Nam97, Proposition 4] are exact replacing Ω_Z^1 with $\Omega_Z^{[1]}$ because D' does not intersect Z , the argument shows that $H^2(Z, T_Z) = 0$. \square

Lemma 4.17. *In the notation as in the proof of Lemma 4.16, we have $R^1\pi_*T_Z = 0$.*

Proof. Recall that $\psi : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow X_u$ is taking the anti-canonical model. From the construction of Z , we know that ψ factors as $\widetilde{\mathbb{P}\mathcal{E}} \xrightarrow{\theta} Z \xrightarrow{\pi} X_u$ where both θ and π are crepant birational. Since $\widetilde{\mathbb{P}\mathcal{E}}$ and Z are both Gorenstein canonical with quotient singularities, applying [Ste77a, Lemma 1.11] (see also [GKKP11, Theorem 1.4]) to a common log resolution of them yields that $\theta_*\Omega_{\widetilde{\mathbb{P}\mathcal{E}}}^{[2]} \cong \Omega_Z^{[2]}$. Thus we have

$$\theta_*T_{\widetilde{\mathbb{P}\mathcal{E}}} = \theta_*(\Omega_{\widetilde{\mathbb{P}\mathcal{E}}}^{[2]} \otimes \omega_{\widetilde{\mathbb{P}\mathcal{E}}}^\vee) = \theta_*(\Omega_{\widetilde{\mathbb{P}\mathcal{E}}}^{[2]} \otimes \theta^*\omega_Z^\vee) \cong (\theta_*\Omega_{\widetilde{\mathbb{P}\mathcal{E}}}^{[2]}) \otimes \omega_Z^\vee \cong \Omega_Z^{[2]} \otimes \omega_Z^\vee = T_Z.$$

From the first two terms of Leray spectral sequence, we have that $R^1\pi_*T_Z$ injects into $R^1\psi_*T_{\widetilde{\mathbb{P}\mathcal{E}}}$. Thus it suffices to show $R^1\psi_*T_{\widetilde{\mathbb{P}\mathcal{E}}} = 0$.

Recall that $X_{u,0} = C_a(\mathbb{P}^1 \times \mathbb{P}^1, -K_{\mathbb{P}^1 \times \mathbb{P}^1} - \Gamma)$ is the orbifold cone where $\Gamma = \frac{1}{2}F_0 + \frac{1}{2}F_1 + \frac{2}{3}F_\infty$. Let W be the total space of the orbifold line bundle $K_{\mathbb{P}^1 \times \mathbb{P}^1} + \Gamma$, i.e.

$$W := \text{Spec }_{\mathbb{P}^1 \times \mathbb{P}^1} \bigoplus_{m=0}^{\infty} \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(\lfloor m(-K_{\mathbb{P}^1 \times \mathbb{P}^1} - \Gamma) \rfloor).$$

Let Σ be the zero section of W . Denote by Δ_Σ the different of (W, Σ) on Σ . Thus we know that $\psi_0 : W \rightarrow X_{u,0}$ provides the Kollar component $(\Sigma, \Delta_\Sigma) \cong (\mathbb{P}^1 \times \mathbb{P}^1, \Gamma)$. From the proof of Lemma 4.12 and Corollary 4.15 we know that the formal isomorphism between X_u and $X_{u,0}$ lifts to a formal isomorphism between $\widetilde{\mathbb{P}\mathcal{E}}$ and W along $\widetilde{H}_{\mathcal{E}}$ and Σ . Thus it suffices to prove $R^1\psi_{0,*}T_W = 0$ which is equivalent to $H^1(W, T_W) = 0$ as $X_{u,0}$ is affine.

Recall from the proof of Lemma 4.12 that there is a finite Galois morphism $\tau_{\widetilde{X}} : \widetilde{X} \rightarrow X_{u,0}$ where $\widetilde{X} := C_a(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2, 2))$ and the Galois group is \mathfrak{S}_3 . Let \widetilde{W} be the total space of the line bundle $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-2, -2)$ with zero section $\widetilde{\Sigma}$. Then $\tilde{\psi} : \widetilde{W} \rightarrow \widetilde{X}$ is the blow-up of the cone vertex providing the Kollar component $\widetilde{\Sigma} \cong \mathbb{P}^1 \times \mathbb{P}^1$. Then the quotient map $\tau_{\widetilde{X}}$ lifts to a quotient map $\tau : \widetilde{W} \rightarrow W$ by the action of \mathfrak{S}_3 which gives the following commutative diagram:

$$\begin{array}{ccc} \widetilde{W} & \xrightarrow{\tau} & W \\ \downarrow \tilde{\psi} & & \downarrow \psi_0 \\ \widetilde{X} & \xrightarrow{\tau_{\widetilde{X}}} & X_{u,0} \end{array}$$

From the proof of Lemma 4.12, we know that τ is quasi-étale, and $\tau|_{\widetilde{\Sigma}} : \widetilde{\Sigma} \rightarrow \Sigma$ is the quotient map of the effective \mathfrak{S}_3 -action on $\widetilde{\Sigma} \cong \mathbb{P}^1 \times \mathbb{P}^1$. Thus τ is also quasi-étale. By [Kni73, Theorem 3], we know that $\Omega_{\widetilde{W}}^{[1]} \cong (\tau_*\Omega_{\widetilde{W}}^{[1]})^{\mathfrak{S}_3}$. Since τ is quasi-étale and $\Omega_{\widetilde{W}}^{[1]}$ is reflexive, we know that $\tau_*\Omega_{\widetilde{W}}^{[1]}$ is also reflexive. This implies that $((\tau_*\Omega_{\widetilde{W}}^{[1]})^{\mathfrak{S}_3})^\vee \cong ((\tau_*\Omega_{\widetilde{W}}^{[1]})^\vee)^{\mathfrak{S}_3}$. Since $\psi : \widetilde{\mathbb{P}\mathcal{E}} \rightarrow X_{u,0}$ is crepant between Gorenstein normal varieties, so is $\psi_0 : W \rightarrow X_{u,0}$. In particular, W is normal and Gorenstein. By construction, \widetilde{W} is smooth hence also normal and Gorenstein. Applying Lemma 4.18 to the morphism τ and sheaf $\Omega_{\widetilde{W}}^{[1]}$, we have that

$$T_W = (\Omega_W^{[1]})^\vee \cong ((\tau_*\Omega_{\widetilde{W}}^{[1]})^{\mathfrak{S}_3})^\vee \cong ((\tau_*\Omega_{\widetilde{W}}^{[1]})^\vee)^{\mathfrak{S}_3} \cong (\tau_*((\Omega_{\widetilde{W}}^{[1]})^\vee))^{\mathfrak{S}_3} = (\tau_*T_{\widetilde{W}})^{\mathfrak{S}_3}.$$

Since $T_W = (\tau_*T_{\widetilde{W}})^{\mathfrak{S}_3}$ is a direct summand of $\tau_*T_{\widetilde{W}}$, we have that $H^1(W, T_W)$ is a direct summand of $H^1(W, \tau_*T_{\widetilde{W}}) \cong H^1(\widetilde{W}, T_{\widetilde{W}})$. Thus it suffices to show $H^1(\widetilde{W}, T_{\widetilde{W}}) = 0$.

Denote by $f_{\widetilde{W}} : \widetilde{W} \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ the \mathbb{A}^1 -bundle structure. Then we have a short exact sequence

$$0 \rightarrow T_{\widetilde{W}/\mathbb{P}^1 \times \mathbb{P}^1} \rightarrow T_{\widetilde{W}} \rightarrow f_{\widetilde{W}}^* T_{\mathbb{P}^1 \times \mathbb{P}^1} \rightarrow 0.$$

By the long exact sequence of cohomology, it suffices to show the vanishing of both $H^1(\widetilde{W}, T_{\widetilde{W}/\mathbb{P}^1 \times \mathbb{P}^1})$ and $H^1(\widetilde{W}, f_{\widetilde{W}}^* T_{\mathbb{P}^1 \times \mathbb{P}^1})$.

Denote by $L_{\widetilde{W}} := \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2, 2)$. Since \widetilde{W} is the total space of the line bundle $L_{\widetilde{W}}^\vee$, we know that $f_{\widetilde{W},*} \mathcal{O}_{\widetilde{W}} \cong \bigoplus_{m=0}^{\infty} L_{\widetilde{W}}^{\otimes m}$, and $T_{\widetilde{W}/\mathbb{P}^1 \times \mathbb{P}^1} = f_{\widetilde{W}}^* L_{\widetilde{W}}^\vee$. Thus we have

$$H^1(\widetilde{W}, T_{\widetilde{W}/\mathbb{P}^1 \times \mathbb{P}^1}) = H^1(\widetilde{W}, f_{\widetilde{W}}^* L_{\widetilde{W}}^\vee) \cong H^1(\mathbb{P}^1 \times \mathbb{P}^1, f_{\widetilde{W},*} f_{\widetilde{W}}^* L_{\widetilde{W}}^\vee) \cong H^1(\mathbb{P}^1 \times \mathbb{P}^1, \bigoplus_{m=-1}^{\infty} L_{\widetilde{W}}^{\otimes m}).$$

By Kodaira vanishing we know that $H^1(\mathbb{P}^1 \times \mathbb{P}^1, L_{\widetilde{W}}^{\otimes m}) = H^1(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2m, 2m)) = 0$ for every $m \in \mathbb{Z}$. Thus we get the vanishing of $H^1(\widetilde{W}, T_{\widetilde{W}/\mathbb{P}^1 \times \mathbb{P}^1})$. On the other hand, we have

$$H^1(\widetilde{W}, f_{\widetilde{W}}^* T_{\mathbb{P}^1 \times \mathbb{P}^1}) \cong H^1(\mathbb{P}^1 \times \mathbb{P}^1, f_{\widetilde{W},*} f_{\widetilde{W}}^* T_{\mathbb{P}^1 \times \mathbb{P}^1}) \cong H^1(\mathbb{P}^1 \times \mathbb{P}^1, \bigoplus_{m=0}^{\infty} L_{\widetilde{W}}^{\otimes m} \otimes T_{\mathbb{P}^1 \times \mathbb{P}^1}).$$

Since $T_{\mathbb{P}^1 \times \mathbb{P}^1} \cong T_{\mathbb{P}^1} \boxtimes T_{\mathbb{P}^1} \cong \mathcal{O}(2, 0) \oplus \mathcal{O}(0, 2)$, by Kodaira vanishing we have

$$H^1(\mathbb{P}^1 \times \mathbb{P}^1, L_{\widetilde{W}}^{\otimes m} \otimes T_{\mathbb{P}^1 \times \mathbb{P}^1}) \cong H^1(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2m+2, 2m) \oplus \mathcal{O}(2m, 2m+2)) = 0$$

for every $m \geq 0$. Thus the vanishing of $H^1(\widetilde{W}, f_{\widetilde{W}}^* T_{\mathbb{P}^1 \times \mathbb{P}^1})$ is proved. The proof is finished. \square

Lemma 4.18. *Let $f : X \rightarrow Y$ be a quasi-étale finite morphism between normal Gorenstein varieties. Let \mathcal{G} be a coherent sheaf on X . Then we have $(f_* \mathcal{G})^\vee = f_* (\mathcal{G}^\vee)$.*

Proof. Since f is quasi-étale and both X and Y are Gorenstein, we know that $\omega_X = f^* \omega_Y = f^! \omega_Y$. By Grothendieck duality for finite morphisms (see e.g. [Sta18, Tag 0AU3]), we have

$$(4.4) \quad f_* \mathcal{H}om_{\mathcal{O}_X}(\mathcal{G} \otimes \omega_X, \omega_X) = \mathcal{H}om_{\mathcal{O}_Y}(f_*(\mathcal{G} \otimes \omega_X), \omega_Y).$$

Since ω_X is invertible, the left-hand side of (4.4) is

$$f_* \mathcal{H}om_{\mathcal{O}_X}(\mathcal{G} \otimes \omega_X, \omega_X) = f_* \mathcal{H}om_{\mathcal{O}_X}(\mathcal{G}, \mathcal{O}_X) = f_*(\mathcal{G}^\vee).$$

For the right-hand side of (4.4), we get

$$\begin{aligned} \mathcal{H}om_{\mathcal{O}_Y}(f_*(\mathcal{G} \otimes \omega_X), \omega_Y) &= \mathcal{H}om_{\mathcal{O}_Y}(f_*(\mathcal{G} \otimes f^* \omega_Y), \omega_Y) \\ &= \mathcal{H}om_{\mathcal{O}_Y}((f_* \mathcal{G}) \otimes \omega_Y, \omega_Y) \\ &= \mathcal{H}om_{\mathcal{O}_Y}(f_* \mathcal{G}, \mathcal{O}_Y) \\ &= (f_* \mathcal{G})^\vee. \end{aligned}$$

Here we use projection formula and the fact that ω_Y is invertible. The proof is finished by combining the above equalities. \square

Corollary 4.19. *The \mathbb{Q} -Fano threefold X_u has unobstructed deformations where L deforms in a \mathbb{Q} -Gorenstein family, and the miniversal base space is a smooth curve. Moreover, any small deformation of X_u in $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ is isomorphic to \mathbb{P}^3 or X_u .*

Proof. This follows from Corollary 4.15 and Lemma 4.16. \square

Definition-Proposition 4.20. Let $c \in (\frac{9}{13}, 1)$ be a rational number. Let $\mathcal{H}_{u,c}$ be the locally closed substack of $\overline{\mathcal{M}}_c^K$ with reduced structure parametrizing K-semistable pairs (X, cS) where $X \cong X_u$. Then $\mathcal{H}_{u,c}$ is a closed substack of $\overline{\mathcal{M}}_c^K$.

Denote by $H_{u,c}$ the good moduli space of $\mathcal{H}_{u,c}$. Then $H_{u,c}$ is a closed subscheme of $\overline{\mathcal{M}}_c^K$ that is isomorphic to the GIT quotient $\mathbf{P} // \mathrm{SL}(2, \mathbb{C})$.

Proof. We first show that $\mathcal{H}_{u,c}$ is closed which follows from the existence part of valuative criterion for properness of the map $\mathcal{H}_{u,c} \hookrightarrow \overline{\mathcal{M}}_c^K$. Let $(\mathcal{X}, c\mathcal{S}) \rightarrow C$ be a \mathbb{Q} -Gorenstein family of K-semistable log Fano pairs over a smooth pointed curve $0 \in C$ such that $K_{\mathcal{X}/C} + c\mathcal{S} \sim_C 0$ and $\mathcal{X}_t \cong X_u$ for any $t \in C \setminus \{0\}$. It suffices to show that $\mathcal{X}_0 \cong X_u$ as well.

Denote by $C^\circ := C \setminus \{0\}$, $\mathcal{X}^\circ := \mathcal{X} \times_C C^\circ$, and $\mathcal{S}^\circ := \mathcal{S}|_{\mathcal{X}^\circ}$. After replacing $(0 \in C)$ by a quasi-finite cover if necessary, we may assume that $\mathcal{X}^\circ \cong X_u \times C^\circ$. Recall from the proof of Theorem 4.11(2) that there is a universal family $(X_u \times \mathbf{A}, \mathcal{S}_\mathbf{A}) \rightarrow \mathbf{A}$ parametrizing $(X_u, S_{(A,B)})$ for $(A, B) \in \mathbf{A}$. Hence by a family version of Lemma 4.10, we can find a map $\gamma^\circ : C^\circ \rightarrow \mathbf{A}$ such that $(\mathcal{X}^\circ, \mathcal{S}^\circ) \cong (X_u \times \mathbf{A}, \mathcal{S}_\mathbf{A}) \times_{\gamma^\circ} C^\circ$. Since $(\mathcal{X}_t, c\mathcal{S}_t)$ is K-semistable for $t \in C^\circ$, Theorem 4.11(2) implies that \mathcal{S}_t is the birational transform of $S_{(A,B)}$ where $(A, B) \in \mathbf{A} \setminus \{0\}$ is GIT semistable. Hence γ° factors as $\gamma^\circ : C^\circ \rightarrow \mathbf{A}^{\mathrm{ss}} \hookrightarrow \mathbf{A}$. Since $\mathbf{A}^{\mathrm{ss}} // (\mathrm{SL}(2, \mathbb{C}) \times \mathbb{G}_m) \cong \mathbf{P} // \mathbb{G}_m$ is proper, after replacing $(0 \in C)$ by a further quasi-finite base change we can find $g : C^\circ \rightarrow \mathrm{SL}(2, \mathbb{C}) \times \mathbb{G}_m$ and $\gamma' : C \rightarrow \mathbf{A}^{\mathrm{ss}}$ such that $\gamma'|_{C^\circ} = g \cdot \gamma^\circ$. In particular, we have a K-semistable log Fano family $(X_u \times C, c\mathcal{S}') := (X_u \times \mathbf{A}, c\mathcal{S}_\mathbf{A}) \times_{\gamma'} C$ over C such that $(X_u \times C, c\mathcal{S}') \times_C C^\circ \cong (\mathcal{X}^\circ, c\mathcal{S}^\circ)$. Thus $(X_u, c\mathcal{S}'_0)$ and $(\mathcal{X}_0, c\mathcal{S}_0)$ are S-equivalent K-semistable log Fano pairs, and by [BX19] they admit a common K-polystable degeneration (X'', cS'') . By Theorem 4.11(2), we have that $X'' \cong X_u$ and S'' is the GIT polystable degeneration of \mathcal{S}'_0 . Hence \mathcal{X}_0 is isomorphic to X_u as it not only specially degenerates to X_u but also comes from an isotrivial degeneration of X_u . Thus $\mathcal{H}_{u,c}$ is a closed substack of $\overline{\mathcal{M}}_c^K$.

Finally, we show that $H_{u,c}$ is isomorphic to $\mathbf{P} // \mathrm{SL}(2, \mathbb{C})$. In fact, the universal family $(X_u \times \mathbf{A}, \mathcal{S}_\mathbf{A}) \times_{\mathbf{A}} \mathbf{A}^{\mathrm{ss}}$ parameterizes c -K-semistable log Fano pairs by Theorem 4.11(2). After taking quotient of $\mathrm{SL}(2, \mathbb{C}) \times \mathbb{G}_m$, we get a stack morphism $[\mathbf{A}^{\mathrm{ss}} // (\mathrm{SL}(2, \mathbb{C}) \times \mathbb{G}_m)] \rightarrow \mathcal{H}_{u,c}$. Thus taking good moduli spaces yields a morphism $\mathbf{P} // \mathrm{SL}(2, \mathbb{C}) \rightarrow H_{u,c}$ which is bijective by Theorem 4.11. By Corollary 4.19 we know that $\mathcal{H}_{u,c}$ is smooth, so $H_{u,c}$ is normal. Therefore, $\mathbf{P} // \mathrm{SL}(2, \mathbb{C}) \rightarrow H_{u,c}$ is an isomorphism. \square

Theorem 4.21. The K-moduli spaces $\overline{\mathcal{M}}_c^K$ has a wall at $c = \frac{9}{13}$. Moreover, we have

- (1) The wall crossing morphism $\phi^- : \overline{\mathcal{M}}_{\frac{9}{13}-\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{9}{13}}^K$ replaces $[(\mathbb{P}^3, T)]$ by $[(X_u, T_0)]$, and is isomorphic near $[(X_u, T_0)]$.
- (2) The wall crossing morphism $\phi^+ : \overline{\mathcal{M}}_{\frac{9}{13}+\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{9}{13}}^K$ replaces $[(X_u, T_0)]$ by the divisor $H_{u, \frac{9}{13}+\epsilon}$.
- (3) For any $c \in (\frac{9}{13}, 1)$, the birational map $\overline{\mathcal{M}}_c^K \dashrightarrow \overline{\mathcal{M}}_{\frac{9}{13}+\epsilon}^K$ is an isomorphism over a neighborhood of $H_{u, \frac{9}{13}+\epsilon}$.

Proof. (1) Let $U_T := \overline{\mathcal{M}}^{\mathrm{GIT}} \setminus W_8$ be an open neighborhood of $[T]$. By (3.1), we know that any $[S] \in U_T \setminus \{[T]\}$ is slc, thus (\mathbb{P}^3, cS) is K-stable for any $c \in (0, 1)$. Since $\mathrm{kst}(\mathbb{P}^3, T) = \frac{9}{13}$ by Theorem 4.3, there are open immersions $U_T \hookrightarrow \overline{\mathcal{M}}_c^K$ when $c \in (0, \frac{9}{13})$ and $U_T \setminus \{[T]\} \hookrightarrow \overline{\mathcal{M}}_c^K$ when $c \in [\frac{9}{13}, 1)$. Thus the map $\phi^- : \overline{\mathcal{M}}_{\frac{9}{13}-\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{9}{13}}^K$ is isomorphic on $U_T \setminus \{[T]\}$. On the other hand, we know that $\phi^-([(P^3, T)]) = [(X_u, T_0)]$ by Proposition 4.6. By Corollary 4.19, we know that $\overline{\mathcal{M}}_{3,64}^{\mathrm{sm}, \delta \geq \epsilon_0}$ is smooth in a neighborhood of $[X_u]$. Thus $\overline{\mathcal{M}}_{\frac{9}{13}}^K$ is normal near $[(X_u, T_0)]$ by

Lemma 2.20. By Zariski's main theorem, we know that $(\phi^-)^{-1}([(X_u, T_0)])$ is connected, thus it has to be the singleton $\{[(\mathbb{P}^3, T)]\}$. Hence ϕ^- is an isomorphism near $[(X_u, T_0)]$.

(2) By Theorem 4.11(1) we know that ϕ^+ contracts $H_{u, \frac{9}{13}+\epsilon}$ to the point $[(X_u, T_0)]$. It suffices to show that any $[(X, S)] \in \overline{\mathcal{M}}_{\frac{9}{13}+\epsilon}^K$ whose $\frac{9}{13}$ -K-polystable replacement is $[(X_u, T_0)]$ satisfies $X \cong X_u$. By Corollary 4.19, we know that X is isomorphic to \mathbb{P}^3 or X_u . If $X \cong \mathbb{P}^3$, then by interpolation [ADL19, Proposition 2.13] we know that $(X, \frac{9}{13}S)$ is also K-polystable, a contradiction to the uniqueness of K-polystable degenerations [LWX21]. Hence we have $X \cong X_u$.

(3) Let $U_u := (\phi^+)^{-1}(\phi^-(U_T))$. By parts (1) and (2), we have $U_u = (U_T \setminus \{[T]\}) \sqcup H_{u, \frac{9}{13}+\epsilon}$ as sets. Since every $[S] \in U_T \setminus \{[T]\}$ is slc and GIT polystable, Proposition 3.6 implies that (\mathbb{P}^3, cS) is always K-polystable for $c \in (0, 1)$. Moreover, Theorem 4.11(2) implies that any pair (X_u, S) in $H_{u, \frac{9}{13}+\epsilon}$ is c -K-polystable for any $c \in (\frac{9}{13}, 1)$. Thus there are open immersions $U_u \hookrightarrow \overline{\mathcal{M}}_c^K$ for any $c \in (\frac{9}{13}, 1)$, which implies that the birational map $\overline{\mathcal{M}}_c^K \dashrightarrow \overline{\mathcal{M}}_{\frac{9}{13}+\epsilon}^K$ is an isomorphism over U_u . The proof is finished. \square

5. HYPERELLIPTIC K3 SURFACES

In this section, we will use the results from [ADL20] to study K-polystable replacements of the locus W_i in $\overline{\mathcal{M}}^{\text{GIT}}$ for $i \in \{0, 1, 2, 3, 4, 6, 7, 8\}$ (see Section 3.3 for the definition). We will show that the first K-moduli wall crossing extracts the divisor H_h birationally over the point $W_0 = \{[2Q]\}$, and subsequent wall crossings precisely replace W_i ($i \geq 1$) by $Z^{i+1} \subset \mathcal{F}$ inside the hyperelliptic divisor H_h as introduced in Section 3.4.

5.1. A cone construction for hyperelliptic K3 surfaces. In this subsection, we provide a cone construction to produce K-polystable threefold pairs from K-polystable surface pairs. This is very useful in constructing the K-polystable replacements of the locus W_i in the GIT moduli space based on the replacements from [ADL20].

Definition 5.1. Let V be a Gorenstein log Del Pezzo surface, that is, a \mathbb{Q} -Fano variety of dimension 2 with K_V Cartier.¹ Let $C \in |-2K_V|$ be an effective Cartier divisor defined by a section $s_C \in H^0(V, -2K_V)$ on V . Let $X := C_p(V, -K_V) = \text{Proj}(\bigoplus_{m \geq 0} \bigoplus_{r=0}^m H^0(V, -rK_V)t^{m-r})$ be the projective cone. Let S be the double cover of V branched along C ; i.e. $S = (t^2 = s_C)$. Then S is naturally embedded into X as an anti-canonical divisor. We denote this construction by $\mathcal{C}(V, cC) := (X, \frac{4c+1}{3}S)$ where $c \in [0, \frac{1}{2}]$ is a rational number.

Theorem 5.2. *With the above notation, let $c \in [0, \frac{1}{2}]$ be a rational number. Then (V, cC) is K-semistable (resp. K-polystable) if and only if $(X, \frac{4c+1}{3}S) = \mathcal{C}(V, cC)$ is K-semistable (resp. K-polystable).*

Proof. We first treat the “only if” part. Assume that (V, cC) is K-semistable. Let $V_X \subset X$ be the section at infinity. Let $Y := C_p(V, -2K_V)$ be a new projective cone with V_Y the section at infinity. Then there exists a finite morphism $\pi : X \rightarrow Y$ as a double cover branched along V_Y and $\pi^*V_Y = 2V_X$. Denote by $\tau : X \rightarrow X$ the involution induced by π . Then it is clear that S is τ -invariant. Denote by $D := S/\tau$ as a divisor in Y . Clearly D corresponds to a section of Y such that $D|_{V_Y} = C$. The finite morphism π is crepant between $(X, \frac{4c+1}{3})$ and $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$. Hence by [LZ20, Zhu21] it suffices to show that $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ is K-semistable. The natural \mathbb{G}_m -action on Y degenerates D to D_0 as the cone over C . Let $r := \frac{1}{2} - c$ be a positive rational number, hence $-2K_V \sim_{\mathbb{Q}} r^{-1}(-K_V - cC)$. By [LX20, Proposition 5.3], we know that $(Y, (1 - \frac{r}{3})V_Y + cD_0)$ is K-semistable. Since $1 - \frac{r}{3} = \frac{5}{6} + \frac{c}{3}$, we know that

¹In later discussions, we will very often assume that V is isomorphic to either $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$.

$(Y, (\frac{5}{6} + \frac{c}{3})V_Y + cD)$ is K-semistable since it admits a K-semistable special degeneration. Similarly, since D is also section of Y , the roles of V_Y and D are interchangeable and we could alternatively degenerate V_Y to V_{Y0} , we know that $(Y, cV_Y + (\frac{5}{6} + \frac{c}{3})D)$ is also K-semistable. We know that $(\frac{1}{2}, \frac{4c+1}{3})$ is a convex linear combination of $((\frac{5}{6} + \frac{c}{3}), c)$ and $(c, (\frac{5}{6} + \frac{c}{3}))$ since the sum of two components are the same and $c < \frac{1}{2} < \frac{5}{6} + \frac{c}{3}$. Hence by interpolation ([ADL19, e.g. Proposition 2.13]), we conclude that $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ is K-semistable. Hence $(X, \frac{4c+1}{3}S)$ is K-semistable.

Next we assume that (V, cC) is K-polystable. Since π is a Galois morphism, by [LZ20, Zhu21] it suffices to show that $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ is K-polystable. By [LWX21] we can choose a special test configuration $(\mathcal{Y}, \frac{1}{2}\mathcal{V} + \frac{4c+1}{3}\mathcal{D})$ of $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ whose central fiber $(Y', \frac{1}{2}V' + \frac{4c+1}{3}D')$ is K-polystable. In particular, $\text{Fut}(\mathcal{Y}, \frac{1}{2}\mathcal{V} + \frac{4c+1}{3}\mathcal{D}) = 0$. Denote by $b := \frac{5}{6} + \frac{c}{3}$. By linearity of the generalized Futaki invariant in coefficients and K-semistability of $(Y, bV_Y + cD)$ and $(Y, cV_Y + bD)$, we know that

$$(5.1) \quad \text{Fut}(\mathcal{Y}, b\mathcal{V} + c\mathcal{D}) = \text{Fut}(\mathcal{Y}, c\mathcal{V} + b\mathcal{D}) = 0.$$

By [LWX21, Theorem 1.4], we know that the analogous statement of [LX20, Proposition 5.3] for K-polystability is true (see also [LZ22, Proposition 2.11]). Hence $(Y, bV_Y + cD_0)$ is the K-polystable since (V, cC) is K-polystable. In particular, $(Y, bV_Y + cD_0)$ is the K-polystable special degeneration of $(Y, bV_Y + cD)$. By [LX14] and [LWX21, Lemma 3.1], (5.1) implies that $(Y', bV' + cD')$ is a K-semistable special degeneration of $(Y, bV_Y + cD)$. Thus [LWX21, Theorem 1.3] implies that $(Y, bV_Y + cD_0)$ is isomorphic to the K-polystable special degeneration of $(Y', bV' + cD')$. Thus we have a sequence of special degenerations

$$(5.2) \quad (Y, bV_Y + cD) \rightsquigarrow (Y', bV' + cD') \rightsquigarrow (Y, bV_Y + cD_0).$$

By forgetting D , D' , and D_0 , we obtain $(Y, bV_Y) \rightsquigarrow (Y', bV') \rightsquigarrow (Y, bV_Y)$. This implies that $(Y', V') \cong (Y, V_Y)$. Similarly, since V_Y and D are symmetric, using the second equality in (5.1) we have that $(Y', D') \cong (Y, V_Y)$. Thus $Y' \cong Y = C_p(V, -2KV)$ where both V' and D' are sections in Y' . Moreover, since $D|_{V_Y} = D_0|_{V_Y}$, after restricting (5.2) to V_Y and V' we see that $(V, C) \cong (V_Y, D|_{V_Y}) \cong (V', D'|_{V'})$. Hence $(Y', V' + D') \cong (Y, V_Y + D)$ which implies that $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ is K-polystable.

Next we treat the “if” part for K-semistability. Assume that $(X, \frac{4c+1}{3}S)$ is K-semistable, which implies the K-semistability of $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ from the above discussion. It suffices to show that (V, cC) is K-semistable. Assume to the contrary that (V, cC) is K-unstable. By Theorem 2.3, there exists a prime divisor E over V such that $\beta_{(V, cC)}(E) < 0$. Let v_t be the quasi-monomial valuation on Y obtained by taking the $(1, t)$ -linear combination of ord_{V_Y} and ord_{E_∞} where E_∞ is a prime divisor over Y by taking cone over E . For simplicity, denote by $\Delta := \frac{1}{2}V_Y + \frac{4c+1}{3}D$. Then a simple computation shows that $A_Y(v_t) = 1 + tA_V(E)$, $v_t(V_Y) = 1$, and $v_t(D) = \min\{1, \text{tord}_E(C)\}$. Hence we have

$$(5.3) \quad A_{(Y, \Delta)}(v_t) = \frac{1}{2} + tA_{(V, cC)}(E) + ct\text{ord}_E(C) - \frac{4c+1}{3}\min\{1, \text{tord}_E(C)\}.$$

Next we compute $S_{(Y, \Delta)}(v_t)$. Let $L_Y := \mathcal{O}_Y(V_Y)$. Then for $m \in \mathbb{Z}_{>0}$ it is clear that

$$H^0(Y, mL_Y) \cong \bigoplus_{i=0}^m H^0(V, -2iKV) \cdot s_0^{m-i},$$

where $(s_0 = 0)$ represents the divisor V_Y , and $s \in H^0(V, -2iKV)$ corresponds a section in $H^0(Y, iL_Y)$ by taking the cone. We have that $v_t(s \cdot s_0^{m-i}) = \text{tord}_E(s) + (m - i)$ for each non-zero section $s \in H^0(V, -2iKV)$. Hence we have

$$(5.4) \quad S_{L_Y, m}(v_t) = \frac{1}{mh^0(Y, mL_Y)} \sum_{i=0}^m h^0(V, -2iKV)(2itS_{-KV, m}(\text{ord}_E) + (m - i)).$$

Here we refer to [BJ20] for the definition and properties of S_m -invariants. It is clear that $h^0(Y, mL_Y) \sim \frac{1}{6}\text{vol}_V(-2K_V)m^3$ and $h^0(V, -2iK_V) \sim \frac{1}{2}\text{vol}_V(-2K_V)i^2$ as $m, i \rightarrow \infty$. By taking limit of (5.4) as $m \rightarrow \infty$, we have that

$$(5.5) \quad S_{L_Y}(v_t) = \frac{3}{2}tS_{-K_V}(E) + \frac{1}{4}.$$

Since $-K_Y - \Delta \sim_{\mathbb{Q}} \frac{2-4c}{3}L_Y$ and $-K_V - cC \sim_{\mathbb{Q}} (1-2c)(-K_V)$, (5.5) implies that

$$(5.6) \quad S_{(Y, \Delta)}(v_t) = tS_{(V, cC)}(E) + \frac{1-2c}{6}.$$

Combining (5.3) and (5.6), we have that

$$(5.7) \quad \beta_{(Y, \Delta)}(v_t) = t\beta_{(V, cC)}(\text{ord}_E) + ct\text{ord}_E(C) + \frac{c+1}{3} - \frac{4c+1}{3} \min\{1, t\text{ord}_E(C)\}.$$

Recall that $\beta_{(V, cC)}(\text{ord}_E) < 0$ by our assumption. If $\text{ord}_E(C) = 0$, then we see that $\beta_{(Y, \Delta)}(v_t) < 0$ for $t \gg 0$ which implies that (Y, Δ) is K-unstable by Theorem 2.3, a contradiction. If $\text{ord}_E(C) \neq 0$, we choose $t = \frac{1}{\text{ord}_E(C)}$. Then (5.7) implies that $\beta_{(Y, \Delta)}(v_t) = t\beta_{(V, cC)}(\text{ord}_E) < 0$, again a contradiction. Thus the “if” part for K-semistability is proven.

Finally, we treat the “if” part for K-polystability. Assume that $(X, \frac{4c+3}{3}S)$ is K-polystable, which implies the K-polystability of $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ from the above discussion. Assume to the contrary that (V, cC) is not K-polystable. By the “if” part for K-semistability, we know that (V, cC) is K-semistable. Let (V', cC') be a K-polystable degeneration of (V, cC) . By the “only if” part, we may use the cone construction over (V', cC') to obtain a K-polystable log Fano pair $(Y', \frac{1}{2}V'_Y + \frac{4c+1}{3}D')$. Then we may take the cone of $(V, cC) \rightsquigarrow (V', cC')$ as in [LZ22, Proof of Proposition 2.11] to produce a K-polystable degeneration $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D) \rightsquigarrow (Y', \frac{1}{2}V'_Y + \frac{4c+1}{3}D')$ which has to be a product test configuration since $(Y, \frac{1}{2}V_Y + \frac{4c+1}{3}D)$ is K-polystable. By restricting to $V_Y \rightsquigarrow V'_Y$, we have that $(V, cC) \cong (V', cC')$ is K-polystable. The proof is finished. \square

We will apply the cone construction to hyperelliptic degree 4 K3 surfaces, when V is either $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$. In this case, the cone constructed in Definition 5.1 is either the cone over the anticanonical embedding of the smooth quadric, which we denote by $X_h = C_p(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2, 2))$, or the cone over the anticanonical embedding of the singular quadric, which is the weighted projective space $\mathbb{P}(1, 1, 2, 4) = C_p(\mathbb{P}(1, 1, 2), \mathcal{O}(4))$.

Proposition 5.3. *Let X be a \mathbb{Q} -Fano threefold that is isomorphic to either X_h or $\mathbb{P}(1, 1, 2, 4)$. Let $S \sim -K_X$ be an effective Cartier divisor on X . If S passes through the cone vertex o of X , then (X, cS) is K-unstable for any $c \in [0, 1)$.*

Proof. Let $\tilde{X} \rightarrow X$ be the partial resolution by blowing up the cone vertex. Denote by $E \subset \tilde{X}$ the exceptional divisor. Since $X \cong C_p(V, -K_V)$ where V is $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$, we know that $A_X(E) = 1$ and $\text{vol}_{X, o}(E) = (-K_V)^2 = 8$. Since S is Cartier and passes through $o = c_X(E)$, we know that $\text{ord}_E(S) \geq 1$. Hence we have

$$\widehat{\text{vol}}_{(X, cS), o}(E) = (A_X(E) - \text{cord}_E(S))^3 \text{vol}_{X, o}(E) \leq 8(1-c)^3.$$

On the other hand, we have $\frac{27}{64}(-K_X - cS)^3 = 27(1-c)^3 > \widehat{\text{vol}}_{(X, cS), o}(E)$, which implies that (X, cS) is K-unstable by [LL19]. \square

5.2. Deformation theory of cones. In this subsection, we show that the allowed deformations of X_h or $\mathbb{P}(1, 1, 2, 4)$ are unobstructed, and any such small deformation of X_h (resp. $\mathbb{P}(1, 1, 2, 4)$) is isomorphic to itself or \mathbb{P}^3 (resp. itself, X_h , or \mathbb{P}^3). Recall that X_h is the projective anti-canonical cone over $\mathbb{P}^1 \times \mathbb{P}^1$.

Lemma 5.4. *The \mathbb{Q} -Gorenstein deformations of X_h or $\mathbb{P}(1, 1, 2, 4)$ in the moduli stack $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ of \mathbb{Q} -Gorenstein smoothable \mathbb{Q} -Fano varieties are unobstructed.*

Proof. We consider the deformations only in the smoothable locus, in particular, we only consider deformations induced by the index 1 cover of L , the limit of $\mathcal{O}_{\mathbb{P}^3}(1)$. In this setting, the deformation theory in [Hac01, Section 3] applies and the obstructions are contained in $T_{QG,X}^2$, defined as follows. Let $\pi: Z \rightarrow X$ be local index 1 cover of L near $x \in X$, with group G , and let $p: \mathcal{Z} \rightarrow X$ be the index 1 cover stack. Then, define $\mathcal{T}_Z^i = \mathcal{E}xt^i(\Omega_Z^1, \mathcal{O}_Z)$ and $T_Z^i = \text{Ext}^i(\Omega_Z^1, \mathcal{O}_Z)$. The \mathbb{Q} -Gorenstein smoothable deformations of X are controlled by $\mathcal{T}_{QG,X}^i = \pi_*(\mathcal{T}_Z^i)^G$ (locally) and $T_{QG,X}^i = \text{Ext}^i(\mathbb{L}_{\mathcal{Z}}, \mathcal{O}_X)$, where $\mathbb{L}_{\mathcal{Z}}$ is the cotangent complex of the stack.

By definition, $\mathcal{T}_{QG,X}^0 = T_X$, the sheaf \mathcal{T}_Z^1 is supported on the singular locus of Z and, by [Ser06, Corollary 3.1.13(ii)] the sheaf \mathcal{T}_Z^2 is supported on the non-lci locus of Z . Furthermore, there is a local-to-global spectral sequence $H^p(\mathcal{T}_{QG,X}^q) \Rightarrow T_{QG,X}^{p+q}$, so it suffices to show that $H^p(\mathcal{T}_{QG,X}^q) = 0$ for $p + q = 2$.

First, consider $X = X_h$. The divisor L is 2-Cartier and passes through the vertex of the cone. A computation shows $H^2(\mathcal{T}_{QG,X_h}^0) = H^2(T_{X_h}) = 0$, and \mathcal{T}_{QG,X_h}^1 is supported on the singular locus of X_h , a single point, hence $H^1(\mathcal{T}_{X_h}^1) = 0$. Finally, the index 1 cover of L on X_h has only hypersurface singularities, hence $\mathcal{T}_Z^2 = 0$, so $H^0(\mathcal{T}_{QG,X_h}^2) = 0$. Therefore, $T_{QG,X_h}^2 = 0$ and the deformations are unobstructed.

Now, let $X = \mathbb{P}(1, 1, 2, 4)$. The divisor $L = \mathcal{O}_X(2)$ is 2-Cartier. From the Euler sequence and cohomology of weighted projective space, we also obtain $H^2(\mathcal{T}_{QG,X}^0) = H^2(T_X) = 0$. Let $o \in X$ be the $\frac{1}{4}(1, 1, 2)$ singularity. Away from o , L is Cartier, and near o , we may compute the index one covering of L : if the coordinates on X are $[x : y : z : w]$, near o , the section ($w = 0$) is a non-vanishing section of $L^{[2]}$. So, one can compute the index one cover is given by the map $p: \mathbb{P}(1, 1, 2, 2) \rightarrow X$, where, if the coordinates on $Z = \mathbb{P}(1, 1, 2, 2)$ are $[x : y : u : v]$, the map is $[x : y : u : v] \mapsto [x : y : u : v^2]$. Noting that Z is a (singular) quadric threefold in \mathbb{P}^4 , we have an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^4}(-2)|_Z \rightarrow \Omega_{\mathbb{P}^4}^1|_Z \rightarrow \Omega_Z^1 \rightarrow 0.$$

Dualizing, we obtain

$$0 \rightarrow T_Z \rightarrow T_{\mathbb{P}^4}|_Z \rightarrow \mathcal{O}_{\mathbb{P}^4}(2)|_Z \rightarrow \mathcal{T}_Z^1 \rightarrow 0.$$

Because \mathcal{T}_Z^1 is mapped onto by the line bundle $\mathcal{O}_{\mathbb{P}^4}(2)|_Z = \mathcal{O}_Z(4)$ in the weighted coordinates on Z , and is supported on the singular locus $x = y = 0$ of Z , $\mathbb{P}_{[u:v]}^1$, we obtain that $\mathcal{T}_Z^1 = \mathcal{O}_{\mathbb{P}^1}(2)$. Let us assume the branch locus of p is given by $w = 0$. By definition and because the canonical covering stack is uniquely determined in the étale topology, away from the branch locus of $p: Z \rightarrow X$, we have $\mathcal{T}_{QG,X}^1 = p_*(T_Z^1)^G$, where G is the action $v \rightarrow -v$ on the singular locus $\mathbb{P}_{[u:v]}^1$. As $T_{QG,X}^1$ is supported on the singular locus $x = y = 0$ of X , which is $\mathbb{P}(2, 4)_{[z:w]} \cong \mathbb{P}_{[z^2:w]}^1$, we can explicitly compute $p_*(T_Z^1)^G$, where $p := p|_{\mathbb{P}_{[u:v]}^1}: \mathbb{P}_{[u:v]}^1 \rightarrow \mathbb{P}_{[z^2:w]}^1$. By computation, since $T_Z^1 = \mathcal{O}_{\mathbb{P}^1}(2)$, $p_*(T_Z^1) = \mathcal{O}(1) \oplus \mathcal{O}$. As p is given by the map $[u : v] \mapsto [u : v^2]$, we can compute the local charts and transition functions for $p_*(T_Z^1)$. On the local chart where $u \neq 0$, computation shows that the module $\mathcal{O}_{\mathbb{P}^1}(2) = \mathbb{C}[v/u]$ can be viewed as the $\mathcal{O}_{\mathbb{P}_{[z^2:w]}^1}$ -module $\mathbb{C}[z^2/w] \oplus v/u\mathbb{C}[z^2/w]$, where $z^2/w = v^2/u^2$. Similarly, on the local chart where

$v \neq 0$, computation shows that the module $\mathcal{O}_{\mathbb{P}^1}(2) = \mathbb{C}[u/v]$ can be viewed as the $\mathcal{O}_{\mathbb{P}^1_{[z^2:w]}}$ -module $\mathbb{C}[z/z^2] \oplus u/v\mathbb{C}[w/z^2]$, where $w/z^2 = u^2/v^2$. Furthermore, the transition function on $\mathcal{O}_{\mathbb{P}^1}(2)_{u \neq 0} = \mathbb{C}[v/u]$ to $\mathcal{O}_{\mathbb{P}^1}(2)_{v \neq 0} = \mathbb{C}[u/v]$ is given by multiplication by u^2/v^2 (and v^2/u^2 in the other direction), gluing together to give three global sections, u^2 , uv , and v^2 . With the $\mathcal{O}_{\mathbb{P}^1_{[z^2:w]}}$ -module structure, we see that the transition functions (from the $w \neq 0$ chart to the $z^2 \neq 0$ chart, and vice versa) become multiplication by $z^2/w = v^2/u^2$ and multiplication by $w/z^2 = u^2/v^2$, so we see that first summand of $p_*(T_Z^1)$ is G -invariant and the second is not. Therefore, $p_*(T_Z^1)^G = \mathcal{O}(1)$.

However, this only computes $T_{QG,X}^1$ on the chart $w \neq 0$. In order to compute $T_{QG,X}^1$ on the entire singular locus, we can compute the canonical covering with a different section (branched over a different point), and use the explicitly computed transition functions to glue them together.

Indeed, consider index-one covering using the section $w - z^2$, giving a map $p' : Z \rightarrow X$ such that the branch locus is $w = z^2$. This computes $T_{QG,X}^1$ on the chart $w \neq z^2$. By the same computation as above, $p'_*(T_Z^1)^G = \mathcal{O}(1)$, given on charts $w - z^2 \neq 0$ by $\mathbb{C}[z^2/(w - z^2)]$ and $z^2 \neq 0$ by $\mathbb{C}[(w - z^2)/z^2]$ with transition function multiplication by $(w - z^2)/z^2$ from $w - z^2 \neq 0$ to $z^2 \neq 0$ and $z^2/(w - z^2)$ in the opposite direction.

From the computation of $T_{QG,X}^1$ on both charts, now it is a matter of gluing the charts together. Note that the singular locus is covered completely by the charts $w \neq 0$ (accurately computing $T_{QG,X}^1$ at all points) and $w - z^2 \neq 0$ (accurately computing $T_{QG,X}^1$ at all points). Noting that these coincide on their common intersection, we use the previous descriptions to determine the gluing and transition functions. We then see that $T_{QG,X}^1$ is given by $\mathbb{C}[z^2/w]$ on the $w \neq 0$ chart and $\mathbb{C}[z^2/(w - z^2)]$ on the $w - z^2 \neq 0$ chart, and the transition function is given by multiplication by $w/z^2 \cdot z^2/w - z^2 = w/w - z^2$ from $w \neq 0$ to $w - z^2 \neq 0$, and by $w - z^2/w$ in the other direction. Then, it is easy to see that there are two global sections w and z^2 , so in particular, $T_{QG,X}^1 = \mathcal{O}(1)$ and $H^1(T_{QG,X}^1) = 0$.

Finally, the index 1 cover of L on $\mathbb{P}(1,1,2,4)$ has only hypersurface singularities, so $\mathcal{T}_Z^2 = 0$ and $H^0(\mathcal{T}_{QG,X}^2) = 0$. Hence, $T_{QG,X}^2 = 0$ and the deformations are unobstructed. \square

Lemma 5.5. *Let $\pi : \mathcal{X} \rightarrow B$ be a \mathbb{Q} -Gorenstein smoothable \mathbb{Q} -Fano family over a smooth pointed curve $0 \in B$.*

- (1) *If $\mathcal{X}_0 \cong X_h$, then a general fiber \mathcal{X}_b is isomorphic to \mathbb{P}^3 or X_h .*
- (2) *If $\mathcal{X}_0 \cong \mathbb{P}(1,1,2,4)$, then a general fiber \mathcal{X}_b is isomorphic to \mathbb{P}^3 , X_h , or $\mathbb{P}(1,1,2,4)$.*

Proof. Since both X_h and $\mathbb{P}(1,1,2,4)$ belong to $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$, we know that π is obtained by pulling back the universal family over $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ under some morphism $B \rightarrow \overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$. Let η be the generic point of B . Since the geometric geometric fiber $[\mathcal{X}_{\bar{\eta}}] \in \overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$, by Lemma 3.3 it admits a \mathbb{Q} -Cartier Weil divisor $\mathcal{L}_{\bar{\eta}}$ such that $4\mathcal{L}_{\bar{\eta}} \sim -K_{\mathcal{X}_{\bar{\eta}}}$. After replacing $0 \in B$ by a quasi-finite cover, we may assume that $\mathcal{L}_{\bar{\eta}}$ is the base change of a \mathbb{Q} -Cartier Weil divisor \mathcal{L}_{η} on the generic fiber \mathcal{X}_{η} . Then we can take the Zariski closure \mathcal{L} of \mathcal{L}_{η} as a Weil divisor on \mathcal{X} . By similar arguments to the proof of Lemma 3.3, we have $\mathcal{L}^{[-4]} \cong_B \omega_{\mathcal{X}/B}$, and the sheaves $\pi_* \mathcal{L}$ and $\pi_* \mathcal{L}^{[2]}$ are locally free over B of rank 4 and 10 respectively.

Consider the sheaf $\mathcal{F} := \text{coker}(\text{Sym}^2 \pi_* \mathcal{L} \rightarrow \pi_* \mathcal{L}^{[2]})$. Denote by $\mathcal{F}_b := \mathcal{F} \otimes \mathcal{O}_B/\mathfrak{m}_b$. If $(\mathcal{X}_0, \mathcal{L}_0)$ is isomorphic to $(\mathbb{P}(1,1,2,4), \mathcal{O}_{\mathbb{P}(1,1,2,4)}(2))$, we know that $\dim \mathcal{F}_0 = 1$. Since $b \mapsto \dim \mathcal{F}_b$ is upper semi-continuous, we know that there exists an open neighborhood $0 \in B' \subset B$ such that $\dim \mathcal{F}_b \leq 1$ for any $b \in B'$. In particular, because $\mathbb{P}(1,1,2,4)$ can be partially smoothed to X_h in a \mathbb{Q} -Gorenstein smoothable family, we may assume that $\dim \mathcal{F}_0 \leq 1$ if either $(\mathcal{X}_0, \mathcal{L}_0)$ is

isomorphic to $(\mathbb{P}(1, 1, 2, 4), \mathcal{O}_{\mathbb{P}(1,1,2,4)}(2))$ or isomorphic to $(X_h, \mathcal{O}_{X_h}(2))$. Moreover, since $\mathcal{L}_0^{[2]}$ is base point free and $\pi_* \mathcal{L}^{[2]}$ is locally free, we may assume that $\mathcal{L}_b^{[2]}$ is base point free (in particular Cartier) for any $b \in B'$. If $\dim \mathcal{F}_b = 0$, then \mathcal{L}_b is base point free hence Cartier, which implies that $\mathcal{X}_b \cong \mathbb{P}^3$ because \mathcal{X}_b is a canonical Gorenstein threefold with Fano index 4 [Shi89, Theorem 3.9]. If $\dim \mathcal{F}_b = 1$, then we may pick a basis s_0, \dots, s_3 of $H^0(\mathcal{X}_b, \mathcal{L}_b)$ and $s_4 \in H^0(\mathcal{X}_b, \mathcal{L}_b^{[2]})$ whose image in \mathcal{F}_b is non-zero. Since $\mathcal{L}_b^{[2]}$ is base point free, we know that $[s_0, \dots, s_3, s_4]$ defines a finite morphism $\mathcal{X}_b \rightarrow \mathbb{P}(1^4, 2)$. Since this map is a closed embedding when $b = 0$, it is of degree 1 for general b . Thus its image is a weighted hypersurface of degree 2 that is not isomorphic to projective space. Then a simple analysis shows that $\mathcal{X}_b \cong X_h$ or $\mathbb{P}(1, 1, 2, 4)$: a degree 2 weighted hypersurface ($g = 0$) in $\mathbb{P}(1^4, 2)$ with coordinates $[x_0 : \dots : x_3 : y]$ is not isomorphic to \mathbb{P}^3 if and only if $g(0, 0, 0, 0, y) = 0$, so has equation $g(x_0, \dots, x_3)$ consisting of monomials of degree 2 in the x_i s. As $\mathbb{P}(1^4, 2) = C_p(\mathbb{P}^3, \mathcal{O}(2))$, such an equation defines the anticanonical cone over a (possibly singular) quadric in \mathbb{P}^3 , so \mathcal{X}_b is either X_h or $\mathbb{P}(1, 1, 2, 4)$. The proof is complete by observing that, if $\mathcal{X}_0 \cong X_h$, then \mathcal{X}_b cannot be $\mathbb{P}(1, 1, 2, 4)$ as the singular locus cannot have larger dimension on the generic fiber. \square

The next result shows that there is a smooth open substack of $\overline{\mathcal{M}}_c^K$ parametrizing \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$, or X_u . Later on we will see that this open substack is indeed the entire stack $\overline{\mathcal{M}}_c^K$ (see Proposition 5.15).

Corollary 5.6. *The subset of $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ that parametrizes \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$, and X_u is a smooth open substack. In particular, there exists a smooth open substack of $\overline{\mathcal{M}}_c^K$ parametrizing (X, S) where X is isomorphic to \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$, or X_u .*

Proof. This follows from Corollary 4.19, Lemma 5.4, and Lemma 5.5. \square

Definition 5.7. Consider the reduced locally closed substack \mathcal{T} of $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ parametrizing X_h or $\mathbb{P}(1, 1, 2, 4)$. Let \mathbf{H} be the locally closed subscheme of $\text{Hilb}(\mathbb{P}^9)$ parametrizing X_h and $\mathbb{P}(1, 1, 2, 4)$ embedded by $2L$. Then $\mathcal{T} \cong [\mathbf{H}/\text{PGL}(10, \mathbb{C})]$ (see e.g. [ADL19, Section 3.6] or [ADL20, Proof of Proposition 5.9]). Let $\pi_c : \overline{\mathcal{M}}_c^K \rightarrow \overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$ be the forgetful map where $\pi_c([(X, S)]) = [X]$. We define $\mathcal{H}_{h,c}$ to be the locally closed substack $\mathcal{H}_{h,c} := \pi_c^{-1}(\mathcal{T})$ of $\overline{\mathcal{M}}_c^K$.

Lemma 5.8. *Notation as in Definition 5.7. The stack $\mathcal{H}_{h,c}$ is smooth.*

Proof. The strategy of showing smoothness of \mathcal{T} is similar to [ADL19, Section 3.6] and [ADL20, Proof of Proposition 5.9]. Let $\mathcal{X}_{\mathbf{H}} \rightarrow \mathbf{H}$ be the universal family. Since the embedded X_h and $\mathbb{P}(1, 1, 2, 4)$ are projective cones, there exists a section $\sigma : \mathbf{H} \rightarrow \mathcal{X}_{\mathbf{H}}$ taking fiberwise cone vertices. Therefore we have a morphism $g : \mathbf{H} \rightarrow \mathbb{P}^9$ as the composition $\mathbf{H} \xrightarrow{\sigma} \mathcal{X}_{\mathbf{H}} \hookrightarrow \mathbb{P}^9 \times \mathbf{H} \rightarrow \mathbb{P}^9$. It is clear that g is an isotrivial fibration with fiber isomorphic to the Hilbert scheme \mathbf{H}' of anti-canonically embedded $\mathbb{P}^1 \times \mathbb{P}^1$ and $\mathbb{P}(1, 1, 2)$ in \mathbb{P}^8 . By [ADL20, Proof of Proposition 5.9] we know that \mathbf{H}' is smooth, hence \mathbf{H} is also smooth. Thus we obtain the smoothness of \mathcal{T} , thereby obtaining the smoothness of $\mathcal{H}_{h,c}$. \square

Theorem 5.9. *Let $\overline{\mathcal{K}}_c$ be the K -moduli stack constructed from [ADL20]. Let $\tilde{\mathcal{K}}_c \rightarrow \overline{\mathcal{K}}_c$ be the μ_2 -gerbe obtained by taking fiberwise double covers. Then $\tilde{\mathcal{K}}_{\frac{3c-1}{4}} \cong \mathcal{H}_{h,c}$ for any $c \in (\frac{1}{3}, 1)$.*

Proof. Let $f_c : \mathcal{H}_{h,c} \rightarrow \tilde{\mathcal{K}}_{\frac{3c-1}{4}}$ be the forgetful functor $f_c([X, D]) = [D]$. We will show that f_c is separated, stabilizer preserving on \mathbb{C} -points, and an isomorphism on \mathbb{C} -points. Using this, by Lemma 5.8 and [AI19, Theorem A.5] (a version of Zariski's main theorem for Artin stacks), we will conclude that f_c is an isomorphism of stacks.

By the valuative criterion (see e.g. [LMB00, Chapter 7]), we consider a diagram

$$\begin{array}{ccc}
U & \longrightarrow & \mathcal{H}_{h,c} \\
\downarrow & \nearrow & \downarrow f_c \\
T & \longrightarrow & \widetilde{\mathcal{K}}_{\frac{3c-1}{4}}
\end{array}$$

where $T = \text{Spec } R$ is a DVR and U is the complement of the closed point $0 \in T$. We must show there is at most one dashed arrow completing the diagram. Suppose for contradiction there are two, i.e. there exist two families $(\mathcal{X}, \mathcal{D}) \rightarrow T$ and $(\mathcal{X}', \mathcal{D}') \rightarrow T$. Because the diagram commutes, the maps agree on \mathcal{D} , so we may assume $\mathcal{D} = \mathcal{D}'$, and \mathcal{X} and \mathcal{X}' are isomorphic away from the central fiber. If they are not isomorphic in the central fiber, then by considering the graph of the rational map $\mathcal{X} \dashrightarrow \mathcal{X}'$, the image of \mathcal{X}_0 in \mathcal{X}' is a proper subvariety of \mathcal{X}'_0 containing \mathcal{D}_0 , as the map is an isomorphism on \mathcal{D} . Consider a generic ruling R of the cone \mathcal{X}_0 , and let \mathbb{A}^1 be the ruling R minus the cone point. Because the image of \mathbb{A}^1 in \mathcal{X}'_0 must be 0-dimensional and the map was an isomorphism on \mathcal{D}_0 , the image must be the two points of intersection of \mathbb{A}^1 and \mathcal{D}_0 . However, this means the image of the connected variety \mathbb{A}^1 is disconnected, a contradiction. Therefore, $\mathcal{X} \cong \mathcal{X}'$ and the map is separated.

Next, we show the forgetful functor is stabilizer preserving on \mathbb{C} -points. Suppose $\sigma \in \text{Aut}(X, D)$ is an automorphism that is the identity on D . Then, we claim that σ is the identity. Note that σ must take rulings of the cone X to rulings of X : in the universal family of the threefolds in \mathbb{P}^9 , the rulings are lines and hence curves of lowest degree, and intersection numbers are preserved in the automorphism. Furthermore, any line on X must be a ruling. Fix any ruling $R \cong \mathbb{P}^1$ of X . By definition, σ fixes the cone point and the two points of intersection with D , and σ takes rulings to rulings, so $\sigma|_{\mathbb{P}^1}$ fixes three points on \mathbb{P}^1 , and hence must be the identity. Therefore, σ is the identity.

Now we show that f_c is an isomorphism on \mathbb{C} -points. By [ADL20, Theorem 4.8], we know that for any point $[(V, C)] \in \widetilde{\mathcal{K}}_{\frac{3c-1}{4}}$, the underlying surface V is isomorphic to either $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$. Hence by taking the cone construction $\mathcal{C}(V, \frac{3c-1}{4}C) = (X, cD)$ (c.f. Definition 5.1), we know that X is isomorphic to either X_h or $\mathbb{P}(1, 1, 2, 4)$. On the other hand, if $(X, D) \in \mathcal{H}_{h,c}$, then Proposition 5.3 implies that D does not pass through the cone vertex of X . Hence after an automorphism of X we have $(X, cD) \cong \mathcal{C}(V, \frac{3c-1}{4}C)$ where $V \cong \mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$. Since (X, cD) is K-semistable, Theorem 5.2 implies that $(V, \frac{3c-1}{4}C)$ is also K-semistable hence belongs to $\widetilde{\mathcal{K}}_{\frac{3c-1}{4}}$. As a result, the forgetful map $f_c : \mathcal{H}_{h,c}(\mathbb{C}) \rightarrow \widetilde{\mathcal{K}}_{\frac{3c-1}{4}}(\mathbb{C})$ sending $[(X, D)]$ to D has inverse given by the cone construction $\widetilde{\mathcal{K}}_{\frac{3c-1}{4}}(\mathbb{C}) \rightarrow \mathcal{H}_{h,c}(\mathbb{C})$.

By Lemma 5.8, the stack $\mathcal{H}_{h,c}$ is smooth, and by [ADL20, Theorem 2.21], the stack $\widetilde{\mathcal{K}}_{\frac{3c-1}{4}}$ is smooth. Therefore, we conclude that f_c is actually an isomorphism by [AI19, Theorem A.5], noting that being isomorphic and stabilizer preserving on \mathbb{C} -points implies fully faithful and essentially surjective. \square

Definition 5.10. Let $c \in (\frac{1}{3}, 1)$ be a rational number. By Theorem 5.9, there exists a closed immersion $\widetilde{\mathcal{K}}_{\frac{3c-1}{4}} \hookrightarrow \overline{\mathcal{M}}_c^K$ of Artin stacks. By taking good moduli spaces, we obtain a closed immersion $\overline{\mathcal{K}}_{\frac{3c-1}{4}} \hookrightarrow \overline{\mathcal{M}}_c^K$. Let $H_{h,c}$ (resp. $\mathcal{H}_{h,c}$) be the image of the closed embedding in the K-moduli space (resp. K-moduli stack). Equivalently, by Theorem 5.9 $H_{h,c}$ (resp. $\mathcal{H}_{h,c}$) is the locus parametrizing K-polystable (resp. K-semistable) pairs (X, cD) where $X \cong X_h$ or $\mathbb{P}(1, 1, 2, 4)$.

5.3. K-polystable replacements. In this section, we describe all K-polystable replacements of the locus W_8 inside $\overline{\mathcal{M}}^{\text{GIT}}$. In particular, we show that the replacements of (\mathbb{P}^3, S) where $[S] \in W_8$ is either (X_h, S') or $(\mathbb{P}(1, 1, 2, 4), S')$. Here S' is a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ or $\mathbb{P}(1, 1, 2)$.

By Theorem 3.9, every $[S] \in W_8$ is defined by $S = (q^2 + g = 0)$ where q is a degree 2 polynomial and g is a degree 4 polynomial. There is a natural closed embedding of such a log pair (\mathbb{P}^3, S) with into the weighted projective space $\mathbb{P}(1^4, 2)$ with coordinates $[x_0, x_1, x_2, x_3, z]$ such that the image is given by $(V(z - q), V(z - q, z^2 + g))$.

We start with the K-polystable replacement of $[2Q]$.

Proposition 5.11. *Let $S = 2Q$ where $Q \subset \mathbb{P}^3$ is a smooth quadric surface. Let $c \in (0, 1)$ be a rational number. Then (\mathbb{P}^3, cS) is K-semistable (resp. K-polystable) if and only if $c \leq \frac{1}{3}$ (resp. $< \frac{1}{3}$). Moreover, the K-polystable degeneration of $(\mathbb{P}^3, \frac{1}{3}S)$ is isomorphic to $(X_h, \frac{1}{3}S_0)$ where $S_0 = 2Q_\infty$ and Q_∞ is the section of $X_h = C_p(Q, -K_Q)$ at infinity.*

Proof. We first show that if (\mathbb{P}^3, cS) is K-semistable, then $c \leq \frac{1}{3}$. Computation shows that

$$A_{(\mathbb{P}^3, cS)}(\text{ord}_Q) = 1 - 2c, \quad S_{(\mathbb{P}^3, cS)}(\text{ord}_Q) = (4 - 4c) \int_0^{1/2} (1 - 2t)^3 dt = \frac{1 - c}{2}.$$

By Theorem 2.3, we have $A_{(\mathbb{P}^3, cS)}(\text{ord}_Q) \geq S_{(\mathbb{P}^3, cS)}(\text{ord}_Q)$ which implies that $c \leq \frac{1}{3}$.

Next, we show that $(\mathbb{P}^3, \frac{1}{3}S)$ special degenerates to $(X_h, \frac{1}{3}S_0)$. We may embed (\mathbb{P}^3, S) into $\mathbb{P}(1^4, 2)$ with image $(V(z - q), V(z - q, z^2))$ where $q = x_0x_1 + x_2^2 + x_3^2$. Consider a 1-PS $\tilde{\sigma}$ in $\text{SL}(4, \mathbb{C}) \times \mathbb{G}_m$ of weight $(0, 0, 0, 0, -1)$ acting diagonally on $\mathbb{P}(1^4, 2)$. Then $\tilde{\sigma}$ specially degenerates $(\mathbb{P}^3, \frac{1}{3}S)$ to $(V(q), \frac{1}{3}V(q, z^2))$ which is isomorphic to $(X_h, \frac{1}{3}S_0)$. Since $Q \cong \mathbb{P}^1 \times \mathbb{P}^1$ is K-polystable and $X_h \cong C_p(Q, -K_Q)$, by [LZ22, Proposition 2.11(2)] we know that $(X_h, \frac{1}{3}S_0)$ is K-polystable. This implies that $(\mathbb{P}^3, \frac{1}{3}S)$ is K-semistable by openness of K-semistability [BLX22, Xu20]. Since $S = 2Q$ is GIT polystable, Theorem 3.4 implies that $(\mathbb{P}^3, \epsilon S)$ is K-polystable for $0 < \epsilon \ll 1$. Hence interpolation for K-stability [ADL19, Proposition 2.13] implies that (\mathbb{P}^3, cS) is K-polystable for any $c \in (0, \frac{1}{3})$. The proof is finished. \square

Theorem 5.12. *Let $[S] \in W_8 \setminus \{[2Q]\}$ be a GIT polystable point with $S = (q^2 + g = 0)$ as in Theorem 3.9. Let $V = (q = 0)$ and $C = V(q, g)$ be a quadric surface and a $(2, 4)$ -complete intersection curve in \mathbb{P}^3 , respectively. Denote by $c := \text{kst}(\mathbb{P}^3, S)$. Then we have the following.*

- (1) *The log Fano pair $(V, \frac{3c-1}{4}C)$ is K-semistable but not K-polystable.*
- (2) *There exists a 1-PS σ in $\text{SL}(4, \mathbb{C})$ that induces a K-polystable degeneration $(V_0, \frac{3c-1}{4}C_0)$ of $(V, \frac{3c-1}{4}C)$ in \mathbb{P}^3 .*
- (3) *There exists a 1-PS $\tilde{\sigma}$ in $\text{SL}(4, \mathbb{C}) \times \mathbb{G}_m$ acting diagonally on $\mathbb{P}(1^4, 2)$ such that the $\text{SL}(4, \mathbb{C})$ -component of $\tilde{\sigma}$ is a positive rescaling of σ , and $\tilde{\sigma}$ induces a K-polystable degeneration $\mathcal{C}(V_0, \frac{3c-1}{4}C_0)$ of (\mathbb{P}^3, cS) in $\mathbb{P}(1^4, 2)$.*

Moreover, the K-semistable thresholds and the destabilizing 1-PS in $\text{SL}(4, \mathbb{C}) \times \mathbb{G}_m$ are given in Table 1, where $0 < \alpha \ll 1$ and $\text{kst}(W_i^\circ) := \text{kst}(\mathbb{P}^3, S)$ for any $[S] \in W_i^\circ$.

Proof. Let $S = (q^2 + 2g = 0)$ be a GIT polystable quartic surface in W_i° for $i \in \{1, 2, 3, 4, 6, 7, 8\}$. As mentioned earlier, the pair (\mathbb{P}^3, S) admits an embedding into $\mathbb{P}(1^4, 2)$ given by the equations $(V(z - q), V(z - q, z^2 + g))$. The pair $(V, C) = (V(q), V(q, g)) \subset \mathbb{P}^3$ provides a threefold pair $(X, S') := \mathcal{C}(V, C)$. It is clear that (X, S') admits a embedding into $\mathbb{P}(1^4, 2)$ given by the equations $(V(q), V(q, z^2 + g))$. According to [LO21, Theorem 6.2, Proposition 6.6, and Table 2], for each $i \in \{1, 2, 3, 4, 6, 7, 8\}$ and $[S] \in W_i^\circ$, there exists a 1-PS σ of $\text{SL}(4, \mathbb{C})$ acting diagonally on \mathbb{P}^3 and a rational number t_{i-1} depending on i such that the following properties hold in the context of VGIT from [LO21, Section 6].

TABLE 1. K-polystable replacements of W_i°

i	$\text{kst}(W_i^\circ)$	Sing. of S in W_i°	Destabilizing 1-PS
0	$\frac{1}{3}$	double quadric	$(0, 0, 0, 0, -1)$
1	$\frac{1}{2}$	two quadrics tangent along a conic	$(1, \alpha, 2\alpha - 1, -3\alpha, -6\alpha)$
2	$\frac{3}{5}$	cuspidal along a conic	$(1, \alpha, 2\alpha - 1, -3\alpha, -4\alpha)$
3	$\frac{2}{3}$	$J_{4,\infty}$	$(7, 3, -1, -9, -10)$
4	$\frac{5}{7}$	$J_{3,0}, J_{3,r}$, or $J_{3,\infty}$	$(3, 1, -1, -3, -3)$
6	$\frac{3}{4}$	E_{14}	$(17, 5, -7, -15, -14)$
7	$\frac{7}{9}$	E_{13}	$(11, 3, -5, -9, -8)$
8	$\frac{9}{11}$	E_{12}	$(8, 2, -4, -6, -5)$

- (V, C) is GIT polystable at slope $(t_{i-1} - \epsilon)$ for $0 < \epsilon \ll 1$, and is GIT semistable but not polystable at slope t_{i-1} ;
- σ degenerates (V, C) to a GIT polystable pair (V_0, C_0) at slope t_{i-1} .

Here the correspondence between i and t_{i-1} are given in the following table. Indeed, the σ from [LO21, Table 2] matches the $\text{SL}(4, \mathbb{C})$ -component of $\tilde{\sigma}$ from Table 1 up to positive rescaling.

TABLE 2. VGIT slopes and K-semistable thresholds

i	1	2	3	4	6	7	8
VGIT slope: t_{i-1}	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{3}{10}$	$\frac{1}{3}$	$\frac{5}{14}$	$\frac{3}{8}$	$\frac{2}{5}$
$\text{kst}(W_i^\circ)$: $\frac{1+2t_{i-1}}{3-2t_{i-1}}$	$\frac{1}{2}$	$\frac{3}{5}$	$\frac{2}{3}$	$\frac{5}{7}$	$\frac{3}{4}$	$\frac{7}{9}$	$\frac{9}{11}$

By [ADL20, Theorem 1.1(2)], we know that $(V, \frac{2t_{i-1}}{3-2t_{i-1}}C)$ is K-semistable but not K-polystable, whose K-polystable degeneration is $(V_0, \frac{2t_{i-1}}{3-2t_{i-1}}C_0)$. Let $c := \frac{1+2t_{i-1}}{3-2t_{i-1}}$ which depends on the choice of i . Hence Theorem 5.2 implies that $(X_0, cS_0) := \mathcal{C}(V_0, \frac{2t_{i-1}}{3-2t_{i-1}}C_0)$ is K-polystable. Suppose $(V_0, C_0) = (V(q_0), V(q_0, g_0)) \subset \mathbb{P}^3$. Then clearly $(X_0, S_0) = (V(q_0), V(q_0, z^2 + g_0)) \subset \mathbb{P}(1^4, 2)$. We choose the \mathbb{G}_m -component of $\tilde{\sigma}$ so that the weight of z^2 is the same as the weight of g_0 , while the weight of z is smaller than the weight of q_0 . It is straightforward to check that the column of $\tilde{\sigma}$ from Table 1 satisfies this property. Thus we know that $\tilde{\sigma}$ degenerates both (\mathbb{P}^3, cS) and (X, cS') to the K-polystable pair (X_0, cS_0) . By openness of K-semistability [BLX22, Xu20], this implies that (\mathbb{P}^3, cS) is K-semistable but not K-polystable, as X_0 is either X_h or $\mathbb{P}(1, 1, 2, 4)$ so $\mathbb{P}^3 \not\cong X_0$. Thus $c = \text{kst}(\mathbb{P}^3, S) = \frac{1+2t_{i-1}}{3-2t_{i-1}}$ is listed in the last row of Table 2 for every quartic surface $[S] \in W_i^\circ$. The proof is finished. \square

5.4. K-moduli wall crossings. Recall from Theorem 3.8 that K-moduli spaces $\overline{\mathcal{M}}_c^K$ display wall crossing phenomena as we vary c . The goal of this subsection is to completely describe all K-moduli walls $\{c_i\}_{1 \leq i \leq k-1}$ and all wall crossing birational morphisms $\phi_i^\pm : \overline{\mathcal{M}}_{c_i \pm \epsilon}^K \rightarrow \overline{\mathcal{M}}_{c_i}^K$. We first give a definition of the exceptional loci of ϕ_i^\pm .

Definition 5.13. For each K-moduli wall $c_i \in (0, 1)$, the *exceptional locus* E_i^\pm is defined as

$$E_i^\pm := \{[(X, S)] \in \overline{\mathcal{M}}_{c_i \pm \epsilon}^K \mid (X, S) \text{ is } (c_i \pm \epsilon)\text{-K-polystable but not } c_i\text{-K-polystable}\}.$$

By [ADL19, Theorem 1.2], we know that E_i^\pm is a Zariski closed subset of $\overline{\mathcal{M}}_{c_i \pm \epsilon}^K$.

We start from describing the first wall crossing.

Theorem 5.14 (First wall crossing). *The first wall of K-moduli stacks $\overline{\mathcal{M}}_c^K$ is $c_1 = \frac{1}{3}$. The first wall crossing has the following description.*

- (1) *For any $0 < c < \frac{1}{3}$, we have $\overline{\mathcal{M}}_c^K \cong \overline{\mathcal{M}}^{\text{GIT}}$ and $\overline{\mathcal{M}}_c^K \cong \overline{\mathcal{M}}^{\text{GIT}}$.*
- (2) *The wall crossing morphism $\phi_1^- : \overline{\mathcal{M}}_{\frac{1}{3}-\epsilon}^K \xrightarrow{\cong} \overline{\mathcal{M}}_{\frac{1}{3}}^K$ is an isomorphism which only replaces $(\mathbb{P}^3, 2Q)$ by $(X_h, 2Q_\infty)$.*
- (3) *The wall crossing morphism $\rho : \overline{\mathcal{M}}_{\frac{1}{3}+\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{1}{3}-\epsilon}^K = \overline{\mathcal{M}}^{\text{GIT}}$ is a divisorial contraction with image $[2Q]$ such that $\phi_1^+ = \phi_1^- \circ \rho : \overline{\mathcal{M}}_{\frac{1}{3}+\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{1}{3}}^K$. The exceptional locus E_1^+ of ρ parametrizes $[(X_h, S)]$ where S is a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along a GIT polystable $(4, 4)$ -curve D .*

Proof. We first prove part (1). It follows from Theorem 3.4 that we have $\overline{\mathcal{M}}_\epsilon^K \cong \overline{\mathcal{M}}^{\text{GIT}}$ for $0 < \epsilon \ll 1$. By Proposition 3.6, the K-moduli stack $\overline{\mathcal{M}}_c^K$ is irreducible. Hence it suffices to show that any GIT semistable quartic surface $S \subset \mathbb{P}^3$ satisfies $\text{kst}(\mathbb{P}^3, S) \geq \frac{1}{3}$. Let S_0 be the GIT polystable degeneration of S . Then by openness of K-semistability [BLX22, Xu20], we know that $\text{kst}(\mathbb{P}^3, S_0) \leq \text{kst}(\mathbb{P}^3, S)$. If S_0 is slc, then Lemma 3.5 implies that $\text{kst}(\mathbb{P}^3, S_0) = 1$. If S_0 is not slc, then either $[S_0] = [T]$ or $[S_0] \in W_8$ in $\overline{\mathcal{M}}^{\text{GIT}}$. Thus we have $\text{kst}(\mathbb{P}^3, S_0) \geq \frac{1}{3}$ by Theorems 4.3 and 5.12. This finishes the proof of part (1).

For part (2), Theorem 3.8 implies that ϕ_1^\pm are projective and birational, hence surjective. By Theorems 4.3 and 5.12, we know that the only GIT polystable quartic surface S with $\text{kst}(\mathbb{P}^3, S) = \frac{1}{3}$ is the double quadric surface $[S] = [2Q]$. Hence ϕ_1^- only replaces $(\mathbb{P}^3, 2Q)$ by $(X_h, 2Q_\infty)$ by Proposition 5.11. Since $\overline{\mathcal{M}}_c^K$ is irreducible, by [ADL19, Theorem 1.2] the morphism ϕ_1^- is projective and bijective. Since any closed point $[(X, S)]$ of $\overline{\mathcal{M}}_{\frac{1}{3}}^K$ satisfies that $X \cong \mathbb{P}^3$ or X_h , Corollary 5.6 implies that $\overline{\mathcal{M}}_{\frac{1}{3}}^K$ is smooth. Hence $\overline{\mathcal{M}}_{\frac{1}{3}}^K$ is normal, and ϕ_1^- is an isomorphism by Zariski's main theorem.

For part (3), we use the deformation theory of X_h . By Theorem 5.9 and Definition 5.10, there is a closed embedding $\overline{K}_{\epsilon/4} \hookrightarrow \overline{\mathcal{M}}_{\frac{1}{3}+\epsilon}^K$ whose image we denote by $H_{h, \frac{1}{3}+\epsilon}$. Note that $H_{h, \frac{1}{3}+\epsilon}$ is a divisor since the locus of hyperelliptic K3s forms a divisor (see e.g. [LO21]). By [ADL20, Theorem 1.1(1)], we know that $H_{h, \frac{1}{3}+\epsilon}$ parametrizes (X_h, S) where S is a double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along a GIT polystable $(4, 4)$ -curve D . Since (X_h, S) admits a special degeneration to $(X_h, 2Q_\infty)$, we know that $\phi_1^+(H_{h, \frac{1}{3}+\epsilon}) = [(X_h, 2Q_\infty)]$. Thus $H_{h, \frac{1}{3}+\epsilon}$ is contained in $E_1^+ = (\phi_1^+)^{-1}([(X_h, 2Q_\infty)])$. On the other hand, if $(X, S) \in E_1^+$ i.e. $\phi_1^+([(X, S)]) = [(X_h, 2Q_\infty)]$, then X admits a special degeneration to X_h . Thus Lemma 5.5 implies that $X \cong \mathbb{P}^3$ or X_h . If $X \cong \mathbb{P}^3$, then $(X, \frac{1}{3}S)$ is K-polystable by interpolation, a contradiction. Thus we have $X \cong X_h$ and hence $[(X, S)] \in H_{h, \frac{1}{3}+\epsilon}$ by Theorem 5.9. Thus we have $H_{h, \frac{1}{3}+\epsilon} = E_1^+$. \square

The following result shows that all but one K-moduli wall crossings of $\overline{\mathcal{M}}_c^K$ for $c \in (\frac{1}{3}, 1)$ are directly induced by those of $H_{h,c}$, i.e. the divisor parametrizing hyperelliptic K3 surfaces.

Proposition 5.15. *Let $c \in (\frac{1}{3}, 1)$ be a rational number. Then any point in $\overline{\mathcal{M}}_c^K \setminus H_{h,c}$ is either $[(\mathbb{P}^3, S)]$ where S is a GIT polystable quartic surface, or $[(X_u, S')]$ where S' is an anti-canonical divisor on X_u .*

Proof. We do induction on the K-moduli walls. When $c = \frac{1}{3} + \epsilon$, by Theorem 5.14 we know that $\overline{\mathcal{M}}_c^K = (\overline{\mathcal{M}}^{\text{GIT}} \setminus \{[2Q]\}) \cup H_{h,c}$. Assume that we hit a K-moduli wall c_i such that the statement is true for any $\frac{1}{3} < c < c_i$. Then we analyze the wall crossing morphisms

$$\overline{\mathcal{M}}_{c_i-\epsilon}^K \xrightarrow{\phi_i^-} \overline{\mathcal{M}}_{c_i}^K \xleftarrow{\phi_i^+} \overline{\mathcal{M}}_{c_i+\epsilon}^K.$$

We first show that the statement is true for $c = c_i$. Indeed, all c_i -K-polystable replacements of $H_{h,c_i-\epsilon}$ belongs to H_{h,c_i} by Theorem 5.9. For $[(\mathbb{P}^3, S)] \in \overline{\mathcal{M}}_{c_i-\epsilon}^K \setminus H_{h,c_i-\epsilon}$ and $[S] \in W_8$, its K-polystable replacement belongs to H_{h,c_i} by Theorem 5.12. For (\mathbb{P}^3, T) , its K-polystable replacement at $c_i = \frac{9}{13}$ is (X_u, T_0) . Since any Fano threefold appearing in H_{h,c_i} is either X_h or $\mathbb{P}(1, 1, 2, 4)$, Lemma 5.4 implies that $\overline{\mathcal{M}}_{c_i}^K$ is a smooth stack. Thus the statement holds for $c = c_i$.

Next, we show that the statement is true for $c = c_i + \epsilon$. Since $\overline{\mathcal{M}}_{c_i+\epsilon}^K$ is an open substack of $\overline{\mathcal{M}}_{c_i}^K$, it is also a smooth stack. Assume that $[(X, S)] \in \overline{\mathcal{M}}_{c_i+\epsilon}^K \setminus H_{h,c_i+\epsilon}$. If (X, S) is c_i -K-polystable, then it is also $(c_i - \epsilon)$ -K-polystable by [ADL19, Proposition 3.18]. Thus $X \cong \mathbb{P}^3$ or X_u . If (X, S) is not c_i -K-polystable, then let (X_0, S_0) be its c_i -K-polystable replacement. Then either $(X_0, S_0) \in H_{h,c_i}$ or $(X_0, S_0) \cong (X_u, T_0)$. In the first case, we know that $X_0 \cong X_h$ or $\mathbb{P}(1, 1, 2, 4)$ which implies that $X \cong \mathbb{P}^3$, X_h , or $\mathbb{P}(1, 1, 2, 4)$ by Lemma 5.5. In fact, X cannot be isomorphic to \mathbb{P}^3 since otherwise $(X, c_i S)$ is K-polystable by interpolation, a contradiction. Thus $X \cong X_h$ or $\mathbb{P}(1, 1, 2, 4)$ which implies that $[(X, S)] \in H_{h,c_i+\epsilon}$ by Theorem 5.9, again a contradiction. In the second case, by Corollary 4.19 we have that $X \cong \mathbb{P}^3$ or X_u . Hence the proof is finished. \square

The following theorem summarizes the results we have obtained, which provides a detailed description of wall crossings for K-moduli spaces $\overline{\mathcal{M}}_c^K$.

Theorem 5.16. *The K-moduli space $\overline{\mathcal{M}}_c^K$ (resp. K-moduli stack $\overline{\mathcal{M}}_c^K$) is irreducible and normal (resp. smooth) for any $c \in (0, 1)$. Moreover, the list of K-moduli walls of $\overline{\mathcal{M}}_c^K$ is given as follows.*

$$(5.8) \quad (c_1, c_2, \dots, c_9) = \left(\frac{1}{3}, \frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{9}{13}, \frac{5}{7}, \frac{3}{4}, \frac{7}{9}, \frac{9}{11} \right).$$

In the below, we give precise description of the wall crossing morphisms.

- (1) When $c = c_1 = \frac{1}{3}$, the K-moduli wall crossing map decreasing from $c = \frac{1}{3} + \epsilon$ to $c = \frac{1}{3} - \epsilon$ is a divisorial contraction of the exceptional divisor E_1^+ , which is the birational transform of H_h , to the point $[(\mathbb{P}^3, 2Q)]$.
- (2) When $c = c_5 = \frac{9}{13}$, the K-moduli wall crossing map decreasing from $c = \frac{9}{13} + \epsilon$ to $\frac{9}{13} - \epsilon$ is a divisorial contraction of the exceptional divisor E_5^+ , which is the birational transform of H_u , to the point $[(\mathbb{P}^3, T)]$.
- (3) When $c = c_i \in \{\frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{5}{7}, \frac{3}{4}, \frac{7}{9}, \frac{9}{11}\}$, i.e. $2 \leq i \leq 9$ and $i \neq 5$, K-moduli wall crossings are flips. Moreover, if $i \in \{2, 3, 4, 7, 8, 9\}$ (resp. if $i = 6$) then the exceptional locus E_i^- is the birational transform of W_{i-1} (resp. W_{i-2}), while E_i^+ is the birational transform of Z^i (resp. Z^{i-1}).

Remark 5.17. Using techniques similar to [ADL19, Section 5.2], one should be able to show that the wall crossing morphisms in Theorem 5.16 (1) and (2) are weighted blow-ups of Kirwan type. Since this is not necessary for our main results, we omit the calculation. By Remark 6.10 (a combination of [LO18, Sections 5.1 and 5.2] and the results of Section 6, where we identify our

K-moduli spaces with the spaces defined in Laza-O'Grady), we will see that the wall crossing morphisms in Theorem 5.16 (1) and (2) are indeed weighted blowups of Kirwan type at the point $[2Q]$ and $[T]$ respectively.

Proof. By Theorem 5.14 and Proposition 5.15, we know that the only possible Fano threefolds appearing in $\overline{\mathcal{M}}_c^K$ are \mathbb{P}^3 , X_h , $\mathbb{P}(1, 1, 2, 4)$, or X_u for any $c \in (0, 1)$. Thus the smoothness of $\overline{\mathcal{M}}_c^K$ and normality of $\overline{\mathcal{M}}_c^K$ follow from Corollary 5.6, while irreducibility is proven in Proposition 3.6.

Next, we turn to the list of K-moduli walls. By Proposition 5.15, a K-moduli wall c_i of $\overline{\mathcal{M}}_c^K$ either satisfies $c_i = \text{kst}(\mathbb{P}^3, S)$ for some $[S] \in W_8 \cup \{[T]\}$, or it is a wall of the K-moduli spaces $\overline{K}_{\frac{3c-1}{4}} \cong H_{h,c}$ from [ADL20], as there are no wall crossings on $H_{u,c}$ when $c \in (\frac{9}{13}, 1)$ by Theorem 4.21. In the former case, we precisely obtain the right-hand-side of (5.8). In the latter case, by [ADL20, Remark 5.13] the collection of walls is $\{c \mid \frac{3c-1}{4} \in \{\frac{1}{8}, \frac{1}{5}, \frac{1}{4}, \frac{2}{7}, \frac{5}{16}, \frac{1}{3}, \frac{4}{11}\}\}$, which equals $\{\frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{5}{7}, \frac{3}{4}, \frac{7}{9}, \frac{9}{11}\}$ as a subset of the right-hand-side of (5.8). Thus we have verified the list of all K-moduli walls (5.8).

Next, we characterize the K-moduli wall crossing morphisms. Part (1) follows from Theorem 5.14. Part (2) follows from Theorems 4.11 and 4.21. We focus on part (3). Let $j := i - 1$ (resp. $j := i - 2$) when $i \in \{2, 3, 4, 7, 8, 9\}$ (resp. when $i = 6$). By Theorem 5.12 and Proposition 5.15, we know that $E_i^- = W_j^\circ \sqcup (E_i^- \cap H_{h,c_i-\epsilon})$ as sets, and $E_i^+ \subset H_{h,c_i+\epsilon}$. Since $H_{h,c} \cong \overline{K}_{\frac{3c-1}{4}}$ by Theorem 5.9, we know that $E_i^\pm \cap H_{h,c_i\pm\epsilon}$ is isomorphic (via the operation $\mathcal{C}(\cdot, \cdot)$) to the exceptional locus $E_i'^\pm$ of $\overline{K}_{c_i'\pm\epsilon} \rightarrow \overline{K}_{c_i'}$ where $c_i' := \frac{3c_i-1}{4}$. By [ADL20, Theorem 1.1], the locus $E_i'^\pm$ is the same as the VGIT exceptional locus of slope $t_i := \frac{3c_i'}{2c_i'+2}$ for $(2, 4)$ complete intersections in \mathbb{P}^3 . By [LO21, Theorem 1.1], we know that $E_i'^-$ is the strict transform of $\rho_*^{-1}W_j \cap H_{h,\frac{1}{3}+\epsilon}$, while $E_i'^+$ is the strict transform of Z^{j+1} . Thus we see that E_i^- (resp. E_i^+) is the strict transform of W_j (resp. Z^{j+1}). The fact that these morphisms are flips will follow from the calculations in Theorem 6.2, as the morphisms are shown to be MMP with scaling with respect to the CM line bundle. \square

6. PROOF OF MAIN THEOREMS

In this section we present proofs of main theorems.

6.1. CM line bundles on K-moduli spaces. In this subsection, we compute the log CM \mathbb{Q} -line bundles on the K-moduli spaces $\overline{\mathcal{M}}_c^K$, and prove Theorem 1.1.

The following result describes the locus of K3 surfaces with Du Val singularities inside K-moduli stacks and spaces.

Proposition 6.1. *Let $c \in (0, 1)$ be a rational number. There exists a saturated open substack $\mathcal{M}_c^{K,\circ}$ of $\overline{\mathcal{M}}_c^K$ consisting of c -K-stable log pairs $[(X, S)]$ where S has Du Val singularities. Moreover, $\mathcal{M}_c^{K,\circ}$ is a smooth Deligne-Mumford stack with coarse moduli space $\mathcal{M}_c^{K,\circ}$. The birational period map $\mathfrak{p}_c : \overline{\mathcal{M}}_c^K \dashrightarrow \mathcal{F}$ for boundary divisors induces an open immersion $\mathfrak{p}_c^\circ : \mathcal{M}_c^{K,\circ} \hookrightarrow \mathcal{F}$ satisfying the following properties.*

- (1) $\mathcal{M}_c^{K,\circ}$ is a big open subset of $\overline{\mathcal{M}}_c^K$ for any $c \in (0, 1)$.
- (2) The divisorial components of $\mathcal{F} \setminus \mathfrak{p}_c^\circ(\mathcal{M}_c^{K,\circ})$ are $\begin{cases} \text{none} & \text{if } c \in (\frac{9}{13}, 1); \\ H_u & \text{if } c \in (\frac{1}{3}, \frac{9}{13}]; \\ H_h \cup H_u & \text{if } c \in (0, \frac{1}{3}]. \end{cases}$
- (3) \mathfrak{p}_c° is an isomorphism if and only if $c \in (\frac{9}{11}, 1)$.

Proof. We first show that $\mathcal{M}_c^{K,\circ}$ is a saturated open substack of $\overline{\mathcal{M}}_c^K$. By the openness of K-stability [BLX22] and lower semi-continuity of lct, there exists an open substack $\mathcal{M}_c^{\text{lc}}$ of $\overline{\mathcal{M}}_c^K$ parametrizing c -K-stable log pairs (X, S) that are log canonical. By applying adjunction, we obtain a \mathbb{Q} -Gorenstein universal family $\mathcal{S} \rightarrow \mathcal{M}_c^{\text{lc}}$ with fibers being semi-log-canonical surfaces with trivial canonical divisor. Since klt is an open condition in \mathbb{Q} -Gorenstein families, and Gorenstein klt is the same as having ADE singularities for surfaces, we know that $\mathcal{M}_c^{K,\circ}$ is an open substack of $\mathcal{M}_c^{\text{lc}}$ and hence an open substack of $\overline{\mathcal{M}}_c^K$. Since $\mathcal{M}_c^{K,\circ}$ consists of K-stable log pairs, every point is closed with finite stabilizers. Hence we know that $\mathcal{M}_c^{K,\circ}$ is a saturated open Deligne-Mumford substack of $\overline{\mathcal{M}}_c^K$. By taking period map for boundary divisors, we obtain a morphism $\mathcal{M}_c^{K,\circ} \rightarrow \mathcal{F}$ which descends to a morphism $\mathfrak{p}_c^\circ : \mathcal{M}_c^{K,\circ} \rightarrow \mathcal{F}$.

Since \mathfrak{p}_c is birational, so is \mathfrak{p}_c° . Next we show that \mathfrak{p}_c° is injective on closed points. By global Torelli theorem it suffices to show that if (X, S) and (X', S') both belong to $\mathcal{M}_c^{K,\circ}$ and $S \cong S'$ as polarized K3 surfaces, then $(X, S) \cong (X', S')$. This clearly holds when S and S' are quartic surfaces in \mathbb{P}^3 . If S and S' are hyperelliptic, then their quotients (V, C) and (V', C') by the hyperelliptic involutions are isomorphic. By Theorem 5.9 and Proposition 5.15, we know that $(X, cS) \cong \mathcal{C}(V, \frac{3c-1}{4}C) \cong \mathcal{C}(V', \frac{3c-1}{4}C') \cong (X', cS')$. If S and S' are unigonal, then Proposition 5.15 implies that $X \cong X' \cong X_u$, and hence $(X, S) \cong (X', S')$ by Lemma 4.10. Thus \mathfrak{p}_c° is injective on closed points. By Zariski's main theorem and normality of K-moduli spaces (Theorem 5.16), we conclude that \mathfrak{p}_c° is an open immersion.

Next we turn to parts (1) – (3). For part (1), if $c \in (0, \frac{1}{3}]$, then by Lemma 3.5 and Theorem 5.14 we know that $\mathcal{M}_c^{K,\circ} \cong \mathcal{M}^\circ$ under the isomorphism $\overline{\mathcal{M}}_c^K \cong \overline{\mathcal{M}}^{\text{GIT}}$, which implies that $\mathcal{M}_c^{K,\circ}$ is a big open subset of $\overline{\mathcal{M}}_c^K$. If $c > \frac{1}{3}$, then a general hyperelliptic K3 surface $S \subset X_h$ satisfies that (X_h, cS) is K-stable by Theorem 5.9 and [ADL20, Theorem 1.1]. If $c > \frac{9}{13}$, then a general unigonal K3 surface $S' \subset X_u$ satisfies that (X_u, cS') is K-stable by Theorem 4.21. Thus we know that $\mathcal{M}_c^{K,\circ}$ has non-empty intersection with $H_{h,c}$ (resp. $H_{u,c}$) if $c > \frac{1}{3}$ (resp. if $c > \frac{9}{13}$). Since there is an open immersion $\mathcal{M}^\circ \hookrightarrow \mathcal{M}_c^{K,\circ}$ for any c by Lemma 3.5, and all but two wall crossings are flips by Theorem 5.16, we know that $\mathcal{M}_c^{K,\circ}$ is a big open subset of $\overline{\mathcal{M}}_c^K$ for any $c \in (0, 1)$. Part (2) follows for similar reasons to part (1) as $\mathcal{M}^\circ \cong \mathcal{F} \setminus (H_h \cup H_u)$ under \mathfrak{p} . For part (3), since \mathfrak{p}_c° is an open immersion, it suffices to show that it is surjective if and only if $c \in (\frac{9}{11}, 1)$. We know that $\mathcal{F} \setminus (H_h \cup H_u) \cong \mathcal{M}^\circ \hookrightarrow \mathcal{M}_c^{K,\circ}$ for any c , and $H_u \subset \mathcal{M}_c^{K,\circ}$ if and only if $c > \frac{9}{13}$ by Theorem 4.21. Thus the surjectivity of \mathfrak{p}_c° is equivalent to the containment $H_h \subset H_{h,c}$. Since $H_{h,c} \cong \overline{K}_{\frac{3c-1}{4}}$ by Theorem 5.9, from the explicit wall crossings for K-moduli of hyperelliptic quartic K3 surfaces (see [ADL20, Remarks 5.13 and 5.14] and [LO21, Section 6]) we know that $H_h \subset H_{h,c}$ if and only if $\frac{3c-1}{4} > \frac{4}{11}$, i.e. $c > \frac{9}{11}$. Thus the proof is finished. \square

Next, we recall the definition of log CM line bundles on K-moduli stacks and spaces from [ADL19, Definition 3.34]. By the construction of K-moduli stacks from [ADL19, Section 3], we may write $\overline{\mathcal{M}}_c^K = [Z_c^{\text{red}}/\text{PGL}(N+1)]$ where Z_c^{red} is a reduced locally closed subscheme of certain relative Hilbert scheme in \mathbb{P}^N parametrizing c -K-semistable pairs in $\overline{\mathcal{M}}_c^K$. Let $\pi_c : (\mathcal{X}_c, \mathcal{D}_c) \rightarrow Z_c^{\text{red}}$ be the universal family. Let $c' \in (0, 1)$ be a rational number. Then the log CM \mathbb{Q} -line bundle $\lambda_{\text{CM}, \pi_c, c' \mathcal{D}_c}$ (resp. the Hodge \mathbb{Q} -line bundle $\lambda_{\text{Hodge}, \pi_c, \mathcal{D}_c}$) on Z_c^{red} descends to a \mathbb{Q} -line bundle $\lambda_{c, c'}$ (resp. $\lambda_{c, \text{Hodge}}$) on $\overline{\mathcal{M}}_c^K$. We simply denote $\lambda_c := \lambda_{c, c'}$. By [ADL19, Proposition 3.35], we know that λ_c descends to a \mathbb{Q} -line bundle Λ_c on $\overline{\mathcal{M}}_c^K$ for any $c \in (0, 1)$. Moreover, if $c \in (0, 1)$ is not a K-moduli wall listed in Theorem 5.16, then both $\lambda_{c, c'}$ and $\lambda_{c, \text{Hodge}}$ descend to \mathbb{Q} -line bundles $\Lambda_{c, c'}$ and $\Lambda_{c, \text{Hodge}}$ on $\overline{\mathcal{M}}_c^K$. By [XZ20], we know that the CM \mathbb{Q} -line bundle Λ_c is ample

on $\overline{\mathfrak{M}}_c^K$ for any $c \in (0, 1)$. Moreover, the Hodge \mathbb{Q} -line bundle $\Lambda_{c, \text{Hodge}}$ is nef on $\overline{\mathfrak{M}}_c^K$ for any $c \in (\frac{9}{11}, 1)$ by Theorem 5.16 and [ADL19, Proposition 3.35].

Theorem 6.2. *Let λ be the Hodge line bundle on \mathcal{F} . Let $\Delta^K := \frac{1}{4}H_h + \frac{9}{8}H_u$. Then for any $c \in (0, 1) \cap \mathbb{Q}$ we have*

$$\overline{\mathfrak{M}}_c^K \cong \text{Proj } R(\mathcal{F}, c\lambda + (1 - c)\Delta^K).$$

Moreover, the CM \mathbb{Q} -line bundle Λ_c on $\overline{\mathfrak{M}}_c^K$ is proportional to $(\mathfrak{p}_c^{-1})_*(c\lambda + (1 - c)\Delta^K)$ up to a positive constant.

Proof. From Theorem 5.16, we know that the birational map $\overline{\mathfrak{M}}_{c'}^K \dashrightarrow \overline{\mathfrak{M}}_c^K$ is a birational contraction for any $0 < c < c' < 1$. Moreover, by [XZ20] we know that Λ_c is ample on $\overline{\mathfrak{M}}_c^K$. Thus similar arguments to [ADL19, Theorem 9.4] imply that for any $c \in (0, 1) \cap \mathbb{Q}$ and $0 < \epsilon \ll 1$ we have

$$\overline{\mathfrak{M}}_c^K \cong \text{Proj } R(\overline{\mathfrak{M}}_{1-\epsilon}^K, \Lambda_{1-\epsilon, c}),$$

and Λ_c is the same as the pushforward of $\Lambda_{1-\epsilon, c}$ under $\overline{\mathfrak{M}}_{1-\epsilon}^K \dashrightarrow \overline{\mathfrak{M}}_c^K$. By Proposition 6.1(3), we know that $\mathfrak{p}_{1-\epsilon}^{-1} : \mathcal{F} \hookrightarrow \overline{\mathfrak{M}}_{1-\epsilon}^K$ is a regular open immersion whose image $\overline{\mathfrak{M}}_{1-\epsilon}^{K, \circ}$ is a big open subset of $\overline{\mathfrak{M}}_{1-\epsilon}^K$. Since $\overline{\mathfrak{M}}_{1-\epsilon}^K$ is normal by Theorem 5.16, to prove the theorem it suffices to show that $(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, c}$ is proportional to $c\lambda + (1 - c)\Delta^K$ up to a positive constant.

By [ADL19, Proposition 3.35] we know that

$$(6.1) \quad (1 - c)^{-3}\Lambda_{1-\epsilon, c} = (1 - c)\Lambda_{1-\epsilon, 0} + 4^4 c\Lambda_{1-\epsilon, \text{Hodge}}.$$

By adjunction, we have $(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, \text{Hodge}} = \lambda$. Since $\mathfrak{p}_{1-\epsilon}^{-1}(\mathcal{F} \setminus (H_h \cup H_u))$ parametrizes pairs (\mathbb{P}^3, S) , we know that the pullback of $\Lambda_{1-\epsilon, 0}$ to $\mathcal{F} \setminus (H_h \cup H_u)$ is trivial as the underlying family of Fano threefolds is an isotrivial \mathbb{P}^3 -fibration. Thus the support of $4^{-4}(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, 0}$ is contained in $H_h \cup H_u$, and we may write $4^{-4}(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, 0} = b_h H_h + b_u H_u$ for some $b_h, b_u \in \mathbb{Q}$. Hence we have

$$(6.2) \quad 4^{-4}(1 - c)^{-3}(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, c} = c\lambda + (1 - c)(b_h H_h + b_u H_u).$$

Therefore, the theorem reduces to showing $4^{-4}(\mathfrak{p}_{1-\epsilon}^{-1})^*\Lambda_{1-\epsilon, 0} = \Delta^K$, i.e. $b_h = \frac{1}{4}$ and $b_u = \frac{9}{8}$.

Let Λ^{GIT} be the ample \mathbb{Q} -line bundle on $\overline{\mathfrak{M}}^{\text{GIT}}$ induced by the hyperplane line bundle on $|\mathcal{O}_{\mathbb{P}^3}(4)|$. Then by [LO19, (4.1.2)] we have

$$(6.3) \quad \mathfrak{p}^*\Lambda^{\text{GIT}} = \lambda + \frac{H_h}{2} + \frac{H_u}{2},$$

where $\mathfrak{p} : \overline{\mathfrak{M}}^{\text{GIT}} \dashrightarrow \mathcal{F}$ is the birational period map.

Next we compute b_u . By Theorem 4.21, for every $c \in (\frac{9}{13}, 1)$ we have that $H_u \subset H_{u,c}$, and $\overline{\mathfrak{M}}_c^K \dashrightarrow \overline{\mathfrak{M}}^{\text{GIT}}$ is regular near $H_{u,c}$ which contracts $H_{u,c}$ to the point $[T]$. Thus $(\mathfrak{p}^*\Lambda^{\text{GIT}})|_{H_u}$ is \mathbb{Q} -trivial. Since $H_h \cap H_u = \emptyset$ by [LO19, Lemma 1.7.3], (6.3) implies that

$$(6.4) \quad \left(\lambda + \frac{H_u}{2} \right) \Big|_{H_u} = 0.$$

On the other hand, by [ADL19, Theorem 3.36] we know that $\Lambda_{\frac{9}{13}+\epsilon, \frac{9}{13}} = (\phi_5^+)^*\Lambda_{\frac{9}{13}}$ which is trivial along $H_u \subset H_{u, \frac{9}{13}+\epsilon}$. Since $\overline{\mathfrak{M}}_{1-\epsilon}^K \dashrightarrow \overline{\mathfrak{M}}_{\frac{9}{13}+\epsilon}^K$ is isomorphic near H_u , we have that $0 = \Lambda_{\frac{9}{13}+\epsilon, \frac{9}{13}}|_{H_u} = \Lambda_{1-\epsilon, \frac{9}{13}}|_{H_u}$. By (6.2) we get

$$(6.5) \quad 0 = \left(\frac{9}{13}\lambda + \frac{4}{13}(b_h H_h + b_u H_u) \right) \Big|_{H_u} = \frac{9}{13} \left(\lambda + \frac{4}{9}b_u H_u \right) \Big|_{H_u}$$

Since $\lambda|_{H_u}$ is ample but not \mathbb{Q} -trivial, equations (6.4) and (6.5) imply that $b_u = \frac{9}{8}$.

Finally, we compute b_h . Denote by $H_h^\circ := H_h \setminus Z^2$ parametrizing hyperelliptic K3 surfaces S with ADE singularities that are double covers of $\mathbb{P}^1 \times \mathbb{P}^1$. By Theorem 5.14, we know that $(X_h, (\frac{1}{3} + \epsilon)S)$ is K-stable. Thus interpolation [ADL19, Proposition 2.13] implies that (X_h, cS) is K-stable for any $c \in (\frac{1}{3}, 1)$, i.e. $H_h^\circ \subset H_{h,c}$. Similar to the H_u case, we have $\Lambda_{\frac{1}{3} + \epsilon, \frac{1}{3}} = (\phi_1^+)^* \Lambda_{\frac{1}{3}}$ which is trivial along H_h° , and $\overline{\mathcal{M}}_{1-\epsilon}^K \dashrightarrow \overline{\mathcal{M}}_{\frac{1}{3} + \epsilon}^K$ is isomorphic near H_h° . Thus (6.2) implies

$$(6.6) \quad 0 = \left(\frac{1}{3}\lambda + \frac{2}{3}(b_h H_h + b_u H_u) \right) \Big|_{H_h^\circ} = \frac{1}{3}(\lambda + 2b_h H_h)|_{H_h^\circ}.$$

Meanwhile, (6.3) implies that

$$(6.7) \quad (\lambda + \frac{1}{2}H_h)|_{H_h^\circ} = 0.$$

It is clear that H_h° is a big open subset of $H_{h,\frac{1}{3}+\epsilon}$ which is isomorphic to the GIT moduli space of $(4, 4)$ -curves on $\mathbb{P}^1 \times \mathbb{P}^1$. Thus $\lambda|_{H_h^\circ}$ is ample and not \mathbb{Q} -trivial. This combining with (6.6) and (6.7) implies that $b_h = \frac{1}{4}$. \square

Lemma 6.3. *For any $\epsilon \in (0, \frac{2}{11}) \cap \mathbb{Q}$, $a \in (0, \frac{1}{9}) \cap \mathbb{Q}$, and $b \in (0, \frac{1}{2}) \cap \mathbb{Q}$, we have*

$$\overline{\mathcal{M}}_{1-\epsilon}^K \cong \text{Proj } R(\mathcal{F}, \lambda + \frac{a}{2}H_h + \frac{b}{2}H_u) \cong \widehat{\mathcal{F}}.$$

Proof. We focus on the first isomorphism, as the second isomorphism is an easy consequence of [LO18, Proposition 17] where it is shown that $\widehat{\mathcal{F}} \cong \text{Proj } R(\mathcal{F}, \lambda + \epsilon\Delta)$ for $0 < \epsilon \ll 1$. By Theorem 5.16, we know that $\overline{\mathcal{M}}_{1-\epsilon}^K$ is independent of the choice of $\epsilon \in (0, \frac{2}{11}) \cap \mathbb{Q}$. Since $\overline{\mathcal{M}}_{1-\epsilon}^K$ and \mathcal{F} are isomorphic in codimension 1, it suffices to show that $(\mathfrak{p}_{1-\epsilon}^{-1})_*(\lambda + \frac{1}{2}(aH_h + bH_u))$ is ample on $\overline{\mathcal{M}}_{1-\epsilon}^K$ for $\epsilon \in (0, \frac{2}{11}) \cap \mathbb{Q}$, $a \in (0, \frac{1}{9}) \cap \mathbb{Q}$, and $b \in (0, \frac{1}{2}) \cap \mathbb{Q}$. We split into two cases.

Case 1: $b \leq \frac{9}{2}a$. Since $a < \frac{1}{9}$, we may choose $\epsilon := \frac{2a}{1+2a} < \frac{2}{11}$. As $\Delta^K = \frac{1}{4}H_h + \frac{9}{8}H_u$, we have

$$(6.8) \quad \lambda + \frac{1}{2}(aH_h + bH_u) = \frac{2b}{9a}(\lambda + 2a\Delta^K) + (1 - \frac{2b}{9a})(\lambda + \frac{a}{2}H_h).$$

By Theorems 5.16 and 6.2, $(\mathfrak{p}_{1-\epsilon}^{-1})_*(\lambda + 2a\Delta^K) = (1 + 2a)(\mathfrak{p}_{1-\epsilon}^{-1})_*((1 - \epsilon)\lambda + \epsilon\Delta^K)$ is a positive multiple of the CM \mathbb{Q} -line bundle $\Lambda_{1-\epsilon}$ on $\overline{\mathcal{M}}_{1-\epsilon}^K$ hence is ample by [XZ20]. As we mentioned earlier, $(\mathfrak{p}_{1-\epsilon}^{-1})_*\lambda = \Lambda_{1-\epsilon, \text{Hodge}}$ is nef on $\overline{\mathcal{M}}_{1-\epsilon}^K$. Thus by (6.8) it suffices to show that $(\mathfrak{p}_{1-\epsilon}^{-1})_*(\lambda + \frac{a}{2}H_h) = \Lambda_{1-\epsilon, \text{Hodge}} + \frac{a}{2}H_{h,1-\epsilon}$ is nef on $\overline{\mathcal{M}}_{1-\epsilon}^K$. Since $\Lambda_{1-\epsilon, \text{Hodge}}$ is nef and $H_{h,1-\epsilon}$ is effective \mathbb{Q} -Cartier, it suffices to show that $(\Lambda_{1-\epsilon, \text{Hodge}} + \frac{a}{2}H_{h,1-\epsilon})|_{H_{h,1-\epsilon}}$ is nef. By Theorem 5.16, we know that $H_{h,1-\epsilon}$ and $H_{u,1-\epsilon}$ are disjoint. Thus

$$(\Lambda_{1-\epsilon, \text{Hodge}} + \frac{a}{2}H_{h,1-\epsilon})|_{H_{h,1-\epsilon}} = (\Lambda_{1-\epsilon, \text{Hodge}} + \frac{a}{2}H_{h,1-\epsilon} + \frac{9a}{4}H_{u,1-\epsilon})|_{H_{h,1-\epsilon}}$$

is a positive multiple of $\Lambda_{1-\epsilon}|_{H_{h,1-\epsilon}}$ which is ample. Thus Case 1 is proved.

Case 2: $b > \frac{9}{2}a$. Since $b < \frac{1}{2}$, we may choose $\epsilon := \frac{4b}{9+4b} < \frac{2}{11}$. Then we have

$$\lambda + \frac{1}{2}(aH_h + bH_u) = \frac{9a}{2b}(\lambda + \frac{4b}{9}\Delta^K) + (1 - \frac{9a}{2b})(\lambda + \frac{b}{2}H_u).$$

Similarly to Case 1, $(\mathfrak{p}_{1-\epsilon}^{-1})_*(\lambda + \frac{4b}{9}\Delta^K) = (1 + \frac{4b}{9})(\mathfrak{p}_{1-\epsilon}^{-1})_*((1 - \epsilon)\lambda + \epsilon\Delta^K)$ is ample as a positive multiple of the CM \mathbb{Q} -line bundle $\Lambda_{1-\epsilon}$. Thus it suffices to show the nefness of $(\Lambda_{1-\epsilon, \text{Hodge}} + \frac{b}{2}H_{u,1-\epsilon})|_{H_{u,1-\epsilon}}$ as $H_{u,1-\epsilon}$ is also effective \mathbb{Q} -Cartier. This again follows from the disjointness of $H_{h,1-\epsilon}$ and $H_{u,1-\epsilon}$ and the ampleness of $\Lambda_{1-\epsilon}|_{H_{u,1-\epsilon}}$. Thus Case 2 is proved. \square

Proof of Theorem 1.1. Part (1) is a consequence of Theorem 5.14. Part (2) is exactly Theorem 6.2. Part (4) follows from Theorem 5.16. Hence we only need to prove part (3). By construction, \mathcal{F} , $\widehat{\mathcal{F}}$, and \mathcal{F}^* are all isomorphic in codimension 1. Hence Lemma 6.3 implies $(\widehat{\mathcal{F}}, \widehat{\lambda}) \cong (\overline{\mathcal{M}}_{1-\epsilon}^K, \Lambda_{1-\epsilon, \text{Hodge}})$ where $\widehat{\lambda}$ is the unique extension of λ on $\widehat{\mathcal{F}}$. Clearly, λ uniquely extends to an ample \mathbb{Q} -line bundle λ^* on \mathcal{F}^* whose pullback under the morphism $\widehat{\mathcal{F}} \rightarrow \mathcal{F}^*$ is exactly $\widehat{\lambda}$. Thus (3) is proved. \square

6.2. Proof of Laza-O'Grady's prediction. In this section, we prove Theorem 1.2 which implies Laza-O'Grady's prediction. The key idea is to construct $\mathcal{F}(a, b)$ from modifications of K-moduli spaces $\overline{\mathcal{M}}_c^K$, use positivity of the log CM line bundle [CP21, Pos22, XZ20], and follow the MMP with scaling from [KKL16].

We recall some notation and consequences from the proof of Theorem 4.21. We define $U_T = \overline{\mathcal{M}}^{\text{GIT}} \setminus W_8$ and $U_u = (\phi_5^+)^{-1}(\phi_5^-(U_T)) \subset \overline{\mathcal{M}}_{\frac{9}{13}+\epsilon}^K$ where $\phi_5^\pm : \overline{\mathcal{M}}_{\frac{9}{13}\pm\epsilon}^K \rightarrow \overline{\mathcal{M}}_{\frac{9}{13}}^K$ are the K-moduli wall crossing morphisms. Moreover, there are canonical open immersions $U_T \hookrightarrow \overline{\mathcal{M}}_c^K$ and $U_u \hookrightarrow \overline{\mathcal{M}}_c^K$ for any $0 < c < \frac{9}{13} < c' < 1$. Let $\rho_u : U_u \rightarrow U_T$ be the composition $\rho_u := ((\phi_5^-)^{-1} \circ \phi_5^+)|_{U_u}$. Then ρ_u is a projective birational morphism that contracts the divisor $H_{u, \frac{9}{13}+\epsilon}$ to the point $[T] \in U_T$. In addition, the restriction of ρ_u to $U_u \setminus H_{u, \frac{9}{13}+\epsilon}$, denoted by ρ_u° , is an isomorphism onto $U_T \setminus \{[T]\}$.

Definition 6.4. Let $a, b \in \mathbb{Q}_{>0}$. Denote by $c = c(a) := \frac{1}{1+2a}$. We define the schemes $\mathcal{F}(a, b)$ as follows.

- (1) For $a \in (0, \frac{2}{9})$ and $b \in (0, 1)$, we define $\mathcal{F}(a, b) := \overline{\mathcal{M}}_c^K$;
- (2) For $a \in (\frac{2}{9}, +\infty)$ and $b \in (0, 1)$, we define $\mathcal{F}(a, b)$ as the gluing of $\overline{\mathcal{M}}_c^K \setminus \{[(\mathbb{P}^3, T)]\}$ and U_u through the isomorphism of open subschemes $U_T \setminus \{[T]\} \xrightarrow[\cong]{(\rho_u^\circ)^{-1}} U_u \setminus H_{u, \frac{9}{13}+\epsilon}$, and $\mathcal{F}(\frac{2}{9}, b) := \mathcal{F}(\frac{2}{9} + \epsilon, b)$ for $0 < \epsilon \ll 1$;
- (3) For $a \in (0, \frac{2}{9})$ and $b \in [1, +\infty)$, we define $\mathcal{F}(a, b)$ as the gluing of $\overline{\mathcal{M}}_c^K \setminus H_{u, c}$ and U_T through the isomorphism of open subschemes $U_u \setminus H_{u, \frac{9}{13}+\epsilon} \xrightarrow[\cong]{\rho_u^\circ} U_T \setminus \{[T]\}$;
- (4) For $a \in [\frac{2}{9}, +\infty)$ and $b \in [1, +\infty)$ we define $\mathcal{F}(a, b) := \overline{\mathcal{M}}_c^K$.

Proposition 6.5. For every $a, b \in \mathbb{Q}_{>0}$, the scheme $\mathcal{F}(a, b)$ is an irreducible normal proper scheme. Denote by $c = c(a) := \frac{1}{1+2a}$. Moreover, if $a \in (\frac{2}{9}, +\infty)$ and $b \in (0, 1)$, then there is a birational morphism $\sigma_{a, b} : \mathcal{F}(a, b) \rightarrow \overline{\mathcal{M}}_c^K$ which contracts $H_{u, \frac{9}{13}+\epsilon}$ to the point $[(\mathbb{P}^3, T)]$, and is isomorphic elsewhere; if $a \in (0, \frac{2}{9})$ and $b \in [1, +\infty)$, then there is a birational morphism $\sigma'_{a, b} : \overline{\mathcal{M}}_c^K \rightarrow \mathcal{F}(a, b)$ which contracts $H_{u, c}$ to the point $[T]$, and is isomorphic elsewhere.

Proof. We first construct the birational morphisms.

Suppose $a \in (\frac{2}{9}, +\infty)$ and $b \in (0, 1)$ which implies $c \in (0, \frac{9}{13})$. Hence by Theorem 4.21 we know that $\overline{\mathcal{M}}_c^K$ is the gluing of $\overline{\mathcal{M}}_c^K \setminus \{[(\mathbb{P}^3, T)]\}$ and U_T through the common open subschemes $U_T \setminus \{[T]\}$. Thus by Definition 6.4(2) we know that there is a birational morphism $\sigma_{a, b} : \mathcal{F}(a, b) \rightarrow \overline{\mathcal{M}}_c^K$ obtained by gluing the identity map on $\overline{\mathcal{M}}_c^K \setminus \{[(\mathbb{P}^3, T)]\}$ and $\rho_u : U_u \rightarrow U_T$. Since ρ_u is a projective morphism, so is $\sigma_{a, b}$. Thus $\mathcal{F}(a, b)$ is a projective scheme.

Suppose $a \in (0, \frac{2}{9})$ and $b \in [1, +\infty)$ which implies $c \in (\frac{9}{13}, 1)$. Hence by Theorem 4.21 we know that $\overline{\mathcal{M}}_c^K$ is the gluing of two $\overline{\mathcal{M}}_c^K \setminus H_{u, c}$ and U_u through the common open subschemes $U_u \setminus H_{u, \frac{9}{13}+\epsilon}$. Thus by Definition 6.4(3) we know that there is a birational morphism

$\sigma'_{a,b} : \overline{\mathfrak{M}}_c^K \rightarrow \mathcal{F}(a,b)$ obtained by gluing the identity map on $\overline{\mathfrak{M}}_c^K \setminus H_{u,c}$ and $\rho_u : U_u \rightarrow U_T$. Then it is clear that $\sigma_{a,b}$ is a surjective proper morphism, which implies that $\mathcal{F}(a,b)$ is a proper scheme by [Sta18, Tag 03GN and Tag 09MQ].

Finally, the irreducibility and normality of $\mathcal{F}(a,b)$ come from the corresponding properties of $\overline{\mathfrak{M}}_c^K$ by Theorem 5.16. \square

Denote by $\psi_{a,b} : \mathcal{F} \dashrightarrow \mathcal{F}(a,b)$ the inverse of the birational period map. Let $\lambda(a,b)$, $H_h(a,b)$, and $H_u(a,b)$ be the pushforward of λ , H_h , and H_u under $\psi_{a,b}$, respectively. By Theorem 5.16, Definition 6.4 and Proposition 6.5, we know that such $\mathcal{F}(a,b)$'s undergo wall crossings at $a = a_i$ or $b = 1$ where

$$(a_1, \dots, a_8) = \left(\frac{1}{9}, \frac{1}{7}, \frac{1}{6}, \frac{1}{5}, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1 \right)$$

We denote the wall crossing morphisms for a fixed $b \in (0,1)$ by $\mathcal{F}(a_i - \epsilon, b) \xrightarrow{\varphi_i^-} \mathcal{F}(a_i, b) \xleftarrow{\varphi_i^+} \mathcal{F}(a_i + \epsilon, b)$.

Lemma 6.6. *Recall the birational morphisms $\rho : \overline{\mathfrak{M}}_{\frac{1}{3}+\epsilon}^K \rightarrow \overline{\mathfrak{M}}^{\text{GIT}}$ from Theorem 5.14 and $\rho_u : U_u \rightarrow U_T$ from Definition 6.4. Then $H_{h,\frac{1}{3}+\epsilon}$ is ρ -anti-ample, and $H_{u,\frac{9}{13}+\epsilon}$ is ρ_u -anti-ample.*

Proof. Since ρ and ρ_u come from wall-crossing morphisms of $\overline{\mathfrak{M}}_c^K$ at $c = \frac{1}{3}$ and $c = \frac{9}{13}$ respectively, by Theorem 6.2 we know that $(\mathfrak{p}_{c+\epsilon})_* \Delta^K$ on $\overline{\mathfrak{M}}_{c+\epsilon}^K$ is relatively anti-ample over $\overline{\mathfrak{M}}_c^K$ for $c \in \{\frac{1}{3}, \frac{9}{13}\}$. For $c = \frac{1}{3}$, we know that $(\mathfrak{p}_{\frac{1}{3}+\epsilon})_* \Delta^K = \frac{1}{4} H_{h,\frac{1}{3}+\epsilon}$ which implies the first statement as $\overline{\mathfrak{M}}_{\frac{1}{3}}^K \cong \overline{\mathfrak{M}}^{\text{GIT}}$. For $c = \frac{9}{13}$, we know that $(\mathfrak{p}_{\frac{9}{13}+\epsilon})_* \Delta^K = \frac{1}{4} H_{h,\frac{9}{13}+\epsilon} + \frac{9}{8} H_{u,\frac{9}{13}+\epsilon}$. From the definition we know that $H_{h,\frac{9}{13}+\epsilon}$ is disjoint from U_u , hence $\frac{9}{8} H_{u,\frac{9}{13}+\epsilon}$ is relatively anti-ample over U_T which implies the second statement. \square

Lemma 6.7. *Let $a, b \in (0,1) \cap \mathbb{Q}$. Then the birational map $\psi_{a,b} : \mathcal{F} \dashrightarrow \mathcal{F}(a,b)$ is an isomorphism in codimension 1. Moreover, we have the following.*

- (1) *If $a \in (0, \frac{2}{9})$, then $\lambda(a,b) + \frac{a}{2} H_h(a,b) + \frac{9a}{4} H_u(a,b)$ is ample on $\mathcal{F}(a,b)$;*
- (2) *If $a \in [\frac{2}{9}, 1)$, then $\lambda(a,b) + \frac{a}{2} H_h(a,b) + \frac{1-\epsilon}{2} H_u(a,b)$ is ample on $\mathcal{F}(a,b)$ for $0 < \epsilon \ll 1$.*
- (3) *$\lambda(a,b) + \frac{a}{2} H_h(a,b)$ is nef for any $a, b \in (0, 1) \cap \mathbb{Q}$.*

Proof. By Theorem 6.2 we know that $(\mathfrak{p}_c^{-1})_*(\lambda + \frac{1-c}{c} \Delta^K)$ is ample on $\overline{\mathfrak{M}}_c^K$.

(1) If $a \in (0, \frac{2}{9})$, we know that $\mathfrak{p}_c^{-1} = \psi_{a,b}$, hence $(\psi_{a,b})_*(\lambda + \frac{1-c}{c} \Delta^K)$ is ample on $\mathcal{F}(a,b)$ for $c = \frac{1}{1+2a}$. It is clear that

$$\lambda + \frac{1-c}{c} \Delta^K = \lambda + 2a \left(\frac{1}{4} H_h + \frac{9}{8} H_u \right) = \lambda + \frac{a}{2} H_h + \frac{9a}{4} H_u.$$

Hence $\lambda(a,b) + \frac{a}{2} H_h(a,b) + \frac{9a}{4} H_u(a,b)$ is ample.

(2) If $a \in [\frac{2}{9}, 1)$, by Theorem 5.16 we know that \mathfrak{p}_c^{-1} is a birational contraction which only contracts the divisor H_u since $c = \frac{1}{1+2a} \in (\frac{1}{3}, \frac{9}{13}]$. Hence we know

$$(\mathfrak{p}_c^{-1})_*(\lambda + \frac{a}{2} H_h) = (\mathfrak{p}_c^{-1})_*(\lambda + \frac{1-c}{c} \Delta^K)$$

is ample on $\overline{\mathfrak{M}}_c^K$. By Proposition 6.5 and Lemma 6.6, there is a birational morphism $\sigma_{a,b} : \mathcal{F}(a,b) \rightarrow \overline{\mathfrak{M}}_c^K$ with exceptional divisor $H_u(a,b)$ which is anti-ample over $\overline{\mathfrak{M}}_c^K$. Since $(\lambda + \frac{1}{2} H_u)|_{H_u} \sim_{\mathbb{Q}} 0$ and $H_h \cap H_u = \emptyset$ by (6.4), we know that

$$(\sigma_{a,b})^* (\mathfrak{p}_c^{-1})_*(\lambda + \frac{a}{2} H_h) = (\psi_{a,b})_*(\lambda + \frac{a}{2} H_h + \frac{1}{2} H_u).$$

Hence $(\psi_{a,b})_*(\lambda + \frac{a}{2}H_h + \frac{1-\epsilon}{2}H_u)$ is ample as it is the pull back of an ample \mathbb{Q} -divisor on $\overline{\mathfrak{M}}_c^K$ twisted by a small multiple of a $\sigma_{a,b}$ -ample divisor.

(3) Since the statement is independent of the choice of $b \in (0, 1)$, we may assume $b \in (0, \frac{1}{2})$. We prove nefness by induction on the walls of a . To start with, we assume $a \in (0, a_1)$ where $a_1 = \frac{1}{9}$. Then we have $\mathcal{F}(a, b) \cong \overline{\mathfrak{M}}_c^K \cong \widehat{\mathcal{F}}$ by Lemma 6.3 as $c = \frac{1}{1+2a} \in (\frac{9}{11}, 1)$. By Lemma 6.3, we know that $\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{b}{2}H_u(a, b)$ is ample. Hence we get nefness of $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ for $a \in (0, a_1)$ by letting $b \rightarrow 0$. Next, we divide the induction into two parts. Note that we always assume $0 < \epsilon \ll 1$ in this proof.

Assume that $\lambda(a_i - \epsilon, b) + \frac{a_i - \epsilon}{2}H_h(a_i - \epsilon, b)$ is nef. Since $\mathcal{F}(a_i - \epsilon, b)$ is independent of the choice of ϵ , by letting $\epsilon \rightarrow 0$ we have that $\lambda(a_i - \epsilon, b) + \frac{a_i}{2}H_h(a_i - \epsilon, b)$ is nef. As all $\mathcal{F}(a, b)$'s with $a, b \in (0, 1) \cap \mathbb{Q}$ are isomorphic in codimension 1, we have that

$$(6.9) \quad \lambda(a_i \pm \epsilon, b) + \frac{a_i}{2}H_h(a_i \pm \epsilon, b) = (\varphi_i^\pm)^*(\lambda(a_i, b) + \frac{a_i}{2}H_h(a_i, b)).$$

Hence we obtain that $\lambda(a_i, b) + \frac{a_i}{2}H_h(a_i, b)$ is also nef.

Assume that $\lambda(a_i, b) + \frac{a_i}{2}H_h(a_i, b)$ is nef and $a \in (a_i, a_{i+1})$. By (6.9) and $\mathcal{F}(a, b) \cong \mathcal{F}(a_i + \epsilon, b)$, we know that $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ is the φ_i^+ -pull-back of $\lambda(a_i, b) + \frac{a_i}{2}H_h(a_i, b)$ hence is nef. Since $a > a_i$, in order to show nefness of $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ it suffices to show that $(\lambda(a, b) + \frac{a}{2}H_h(a, b))|_{H_h(a, b)}$ is nef. By parts (1) and (2), there exists $b' = b'(a) \in (0, 1)$ such that $\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{b'}{2}H_u(a, b)$ is ample. Since $H_h(a, b) \cap H_u(a, b) = \emptyset$,

$$(\lambda(a, b) + \frac{a}{2}H_h(a, b))|_{H_h(a, b)} = (\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{b'}{2}H_u(a, b))|_{H_h(a, b)}$$

is ample. Thus $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ is nef for any $a \in (a_i, a_{i+1})$. As a result, the induction steps are validated which yield the nefness of $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ for any $a, b \in (0, 1) \cap \mathbb{Q}$. \square

Definition 6.8. Let $\widetilde{\mathfrak{M}}^{\text{GIT}} := \mathcal{F}(1-\epsilon, 1-\epsilon)$ for $0 < \epsilon \ll 1$. Then the birational map $\mathcal{F} \dashrightarrow \widetilde{\mathfrak{M}}^{\text{GIT}}$ is isomorphic in codimension 1. Denote the pushforwards of λ , H_h , and H_u under this map by $\widetilde{\lambda}$, \widetilde{H}_h , and \widetilde{H}_u , respectively. By Definition 6.4 and Proposition 6.5, there is a birational morphism $\tilde{\rho} : \widetilde{\mathfrak{M}}^{\text{GIT}} \rightarrow \overline{\mathfrak{M}}^{\text{GIT}}$ induced by $\sigma_{1-\epsilon, 1-\epsilon}$ that contracts \widetilde{H}_h and \widetilde{H}_u to $[2Q]$ and $[T]$ respectively and is isomorphic elsewhere.

From Definition 6.4 and Theorem 5.16, we see that $\mathcal{F}(a, b) \cong \overline{\mathfrak{M}}^{\text{GIT}}$ if $a, b \in [1, +\infty)$, and $\mathcal{F}(1-\epsilon, b_1) \cong \widetilde{\mathfrak{M}}^{\text{GIT}}$ if $0 < \epsilon \ll 1$ and $b_1 \in (0, 1)$.

Theorem 6.9. For any $a, b \in \mathbb{Q}_{>0}$, the section ring $R(\mathcal{F}, \lambda + \frac{a}{2}H_h + \frac{b}{2}H_u)$ is finitely generated, and $\mathcal{F}(a, b) \cong \text{Proj } R(\mathcal{F}, \lambda + \frac{a}{2}H_h + \frac{b}{2}H_u)$. In particular, every $\mathcal{F}(a, b)$ is projective.

Proof. We split into four cases based on values of a and b .

Case 1: $a, b \in (0, 1)$. By Lemma 6.7, $\psi_{a,b} : \mathcal{F} \dashrightarrow \mathcal{F}(a, b)$ is an isomorphism in codimension 1. Thus it suffices to show that the \mathbb{Q} -divisor $\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{b}{2}H_u(a, b)$ is ample on $\mathcal{F}(a, b)$ for $a, b \in (0, 1) \cap \mathbb{Q}$.

By Lemma 6.7(1)(2), there exists $b' = b'(a) \in (0, 1)$ such that $\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{b'}{2}H_u(a, b)$ is ample. By Lemma 6.7(3), we know that $\lambda(a, b) + \frac{a}{2}H_h(a, b)$ is nef. Since a strict convex combination of a nef \mathbb{Q} -divisor and an ample \mathbb{Q} -divisor is ample, it suffices to show the nefness of $\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{1}{2}H_u(a, b)$. Moreover, since $\frac{b'}{2} < \frac{1}{2}$ and $H_h(a, b) \cap H_u(a, b) = \emptyset$, it suffices to show that

$$(\lambda(a, b) + \frac{a}{2}H_h(a, b) + \frac{1}{2}H_u(a, b))|_{H_u(a, b)} = (\lambda(a, b) + \frac{1}{2}H_u(a, b))|_{H_u(a, b)}$$

is nef. Indeed, by Definition 6.4, Theorem 4.21(3), and Lemma 6.3, the birational map $\widehat{\mathcal{F}} \dashrightarrow \mathcal{F}(a, b)$ is isomorphic near \widehat{H}_u and $H_u(a, b)$. Since H_u is a big open subset of \widehat{H}_u and $(\lambda + \frac{H_u}{2})|_{H_u} = 0$ by (6.4), we know that $(\widehat{\lambda} + \frac{\widehat{H}_u}{2})|_{\widehat{H}_u} = 0$. This implies $(\lambda(a, b) + \frac{1}{2}H_u(a, b))|_{H_u(a, b)}$ is \mathbb{Q} -linearly trivial and hence nef. Thus Case 1 is proved.

Case 2: $a \in (0, 1)$ and $b \in [1, +\infty)$. We have the diagram $\mathcal{F}(a, 1 - \epsilon) \xrightarrow{\rho_a} \mathcal{F}(a, 1) \xleftarrow{\cong} \mathcal{F}(a, b)$ from Definition 6.4, where ρ_a is the birational morphism that contracts the divisor $H_u(a, 1 - \epsilon)$ to a point. We first show that $\lambda(a, 1) + \frac{a}{2}H_h(a, 1)$ is ample on $\mathcal{F}(a, 1)$. Indeed, from Case 1 we have the ampleness of $\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{1-\epsilon}{2}H_u(a, 1 - \epsilon)$ on $\mathcal{F}(a, 1 - \epsilon)$. By letting $\epsilon \rightarrow 0$, we know that $\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{1}{2}H_u(a, 1 - \epsilon)$ is big and nef, whose restrict to $H_u(a, 1 - \epsilon)$ is \mathbb{Q} -linearly trivial. Thus

$$\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{1}{2}H_u(a, 1 - \epsilon) = \rho_a^*(\lambda(a, 1) + \frac{a}{2}H_h(a, 1)).$$

This shows that $\lambda(a, 1) + \frac{a}{2}H_h(a, 1)$ is big and nef. By the Nakai-Moishezon criterion, to show the ampleness of $\lambda(a, 1) + \frac{a}{2}H_h(a, 1)$, it suffices to show that $(\lambda(a, 1) + \frac{a}{2}H_h(a, 1))|_V$ is big for any positive dimensional closed subvariety $V \subset \mathcal{F}(a, 1)$. Let $\tilde{V} \subset \mathcal{F}(a, 1 - \epsilon)$ be the birational transform of V . Then clearly $\tilde{V} \not\subset H_u(a, 1 - \epsilon)$ as ρ_a contracts $H_u(a, 1 - \epsilon)$ to a point. Since $(\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{1-\epsilon}{2}H_u(a, 1 - \epsilon))|_{\tilde{V}}$ is ample, we know that

$$(\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{1}{2}H_u(a, 1 - \epsilon))|_{\tilde{V}} = \rho_a^*((\lambda(a, 1) + \frac{a}{2}H_h(a, 1))|_V)$$

is big. Thus $(\lambda(a, 1) + \frac{a}{2}H_h(a, 1))|_V$ is big which implies the ampleness of $\lambda(a, 1) + \frac{a}{2}H_h(a, 1)$.

Since $\mathcal{F}(a, 1 - \epsilon)$ and \mathcal{F} are isomorphic in codimension 1, Case 2 reduces to showing

$$\text{Proj } R(\mathcal{F}(a, 1 - \epsilon), \lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{b}{2}H_u(a, 1 - \epsilon)) \cong \mathcal{F}(a, 1).$$

This is true because $\lambda(a, 1) + \frac{a}{2}H_h(a, 1)$ is ample, $H_u(a, 1 - \epsilon)$ is ρ_a -exceptional, and

$$\lambda(a, 1 - \epsilon) + \frac{a}{2}H_h(a, 1 - \epsilon) + \frac{b}{2}H_u(a, 1 - \epsilon) = \rho_a^*(\lambda(a, 1) + \frac{a}{2}H_h(a, 1)) + \frac{b-1}{2}H_u(a, 1 - \epsilon).$$

Thus Case 2 is proved.

Case 3: $a \in [1, +\infty)$ and $b \in (0, 1)$. Since \mathcal{F} and $\widetilde{\mathcal{M}}^{\text{GIT}}$ are isomorphic in codimension 1, it suffices to show that

$$(6.10) \quad \text{Proj } R(\widetilde{\mathcal{M}}^{\text{GIT}}, \tilde{\lambda} + \frac{a}{2}\tilde{H}_h + \frac{b}{2}\tilde{H}_u) \cong \mathcal{F}(a, b).$$

By Definition 6.4, we know that $\widetilde{\mathcal{M}}^{\text{GIT}}$ is the gluing of U_u and $\overline{\mathcal{M}}_{\frac{1}{3}+\epsilon}^K \setminus \{[(\mathbb{P}^3, T)]\}$, while $\mathcal{F}(a, b)$ is the gluing of U_u and $\overline{\mathcal{M}}_c^K \setminus \{[(\mathbb{P}^3, T)]\}$ with $c = \frac{1}{1+2a} \leq \frac{1}{3}$. Thus we have $\overline{\mathcal{M}}_c^K \cong \widetilde{\mathcal{M}}^{\text{GIT}}$ by Theorem 5.14. Thus the birational morphism $\tilde{\rho} : \widetilde{\mathcal{M}}^{\text{GIT}} \rightarrow \overline{\mathcal{M}}^{\text{GIT}}$ can be decomposed into

$$\widetilde{\mathcal{M}}^{\text{GIT}} \xrightarrow{\rho_1} \mathcal{F}(a, b) \xrightarrow{\rho_2} \overline{\mathcal{M}}^{\text{GIT}},$$

where ρ_1 and ρ_2 contracts \tilde{H}_h and $(\rho_1)_*\tilde{H}_u$ respectively. Note that ρ_2 is also induced by $\sigma_{a,b}$.

We first show ampleness of $(\rho_1)_*(\tilde{\lambda} + \frac{b}{2}\tilde{H}_u)$. Indeed, by (6.3) we know that $(\rho_1)_*(\tilde{\lambda} + \frac{1}{2}\tilde{H}_u)$ is the pull-back of the ample \mathbb{Q} -line bundle Λ^{GIT} on $\overline{\mathcal{M}}^{\text{GIT}}$ under ρ_2 . Since $(\rho_1)_*\tilde{H}_u$ is ρ_2 -anti-ample by Lemma 6.6, we have that $(\rho_1)_*(\tilde{\lambda} + \frac{1-\epsilon}{2}\tilde{H}_u)$ is ample for $0 < \epsilon \ll 1$. On the other hand, we know that $(\tilde{\lambda} + \frac{1}{2}\tilde{H}_h)|_{\tilde{H}_h}$ is \mathbb{Q} -linearly trivial by (6.7). Thus $(\rho_1)^*(\rho_1)_*\tilde{\lambda} = \tilde{\lambda} + \frac{1}{2}\tilde{H}_h$. By Definition 6.8 and Lemma 6.7(3), we know that $\widetilde{\mathcal{M}}^{\text{GIT}} = \mathcal{F}(1 - \epsilon, 1 - \epsilon)$ and $\tilde{\lambda} + \frac{1-\epsilon}{2}\tilde{H}_h$ is

nef. By letting $\epsilon \rightarrow 0$ we obtain nefness of $\tilde{\lambda} + \frac{1}{2}\tilde{H}_h$ which implies the nefness of $(\rho_1)_*\tilde{\lambda}$. This together with ampleness of $(\rho_1)_*(\tilde{\lambda} + \frac{1-\epsilon}{2}\tilde{H}_u)$ implies that $(\rho_1)_*(\tilde{\lambda} + \frac{b}{2}\tilde{H}_u)$ is ample.

Now we prove (6.10). From the above arguments, we have that $(\rho_1)_*(\tilde{\lambda} + \frac{b}{2}\tilde{H}_u)$ is ample on $\mathcal{F}(a, b)$, and

$$\tilde{\lambda} + \frac{a}{2}\tilde{H}_h + \frac{b}{2}\tilde{H}_u = (\rho_1)^*(\rho_1)_*(\tilde{\lambda} + \frac{b}{2}\tilde{H}_u) + \frac{a-1}{2}\tilde{H}_h.$$

Since $a \geq 1$ and \tilde{H}_h is ρ_1 -exceptional, (6.10) follows. Thus Case 3 is proved.

Case 4: $a, b \in [1, +\infty)$. By Definition 6.8 we know that $\mathcal{F}(a, b) \cong \overline{\mathfrak{M}}^{\text{GIT}}$. Similarly to Case 3, it suffices to show that

$$(6.11) \quad \text{Proj } R(\overline{\mathfrak{M}}^{\text{GIT}}, \tilde{\lambda} + \frac{a}{2}\tilde{H}_h + \frac{b}{2}\tilde{H}_u) \cong \overline{\mathfrak{M}}^{\text{GIT}}.$$

By (6.3), we know that $\tilde{\rho}^*\Lambda^{\text{GIT}} = \tilde{\lambda} + \frac{1}{2}\tilde{H}_h + \frac{1}{2}\tilde{H}_u$. Since Λ^{GIT} is ample on $\overline{\mathfrak{M}}^{\text{GIT}}$, both \tilde{H}_h and \tilde{H}_u are $\tilde{\rho}$ -exceptional, and $a, b \geq 1$, we conclude that (6.11) holds. Thus Case 4 is proved. \square

Remark 6.10. From the identification of the K-moduli spaces with $\mathcal{F}(a, b)$, and the results of [LO18, Sections 5.1 and 5.2], where Laza-O’Grady prove that $\tilde{\rho} : \widetilde{\mathfrak{M}}^{\text{GIT}} = \mathcal{F}(1-\epsilon, 1-\epsilon) \rightarrow \overline{\mathfrak{M}}^{\text{GIT}}$ (denoted by $\widetilde{\mathfrak{M}} \rightarrow \mathfrak{M}$ in their notation) is a composition of weighted blowups of Kirwan type at $[2Q]$ and $[T]$, we can conclude that the morphisms ρ in Theorem 5.14(3) and ρ_u in Definition 6.4 are weighted blowups at $[2Q]$ and $[T]$ respectively.

Proof of Theorem 1.2. This follows from Definition 6.4, Theorems 5.16, and 6.9. \square

Proof of Corollary 1.3. This is a direct consequence of Theorem 1.2 by letting $a = b$. \square

Remark 6.11. It is reasonable to expect that $R(\mathcal{F}, \lambda + \frac{a}{2}H_h + \frac{b}{2}H_u)$ is finitely generated if one of a or b is zero and the other is positive. Taking Proj of these rings should extend the wall-crossing picture as described in Theorem 1.2 to $a, b \in \mathbb{Q}_{\geq 0}$ where $\mathcal{F}(0, 0) = \mathcal{F}^*$.

6.3. Quartic double solids. Recall that a smooth quartic double solid Y is a double cover of \mathbb{P}^3 branched along a smooth quartic surface S . Denote the double cover map by $\pi : Y \rightarrow \mathbb{P}^3$. A quartic double solid is the same as a del Pezzo threefold of degree 2 (see e.g. [Fuj90]). By [Der16, Example 4.2], we know that Y is Kähler-Einstein and hence K-stable as $\text{Aut}(Y)$ is finite. Since any smooth deformation of a del Pezzo threefold is still del Pezzo of the same degree, there exists an open substack \mathcal{Y} of $\mathcal{M}_{3,16}^{\text{Kss}}$ parametrizing all smooth quartic double solids. Let $\overline{\mathcal{Y}}$ be the Zariski closure of \mathcal{Y} in $\mathcal{M}_{3,16}^{\text{Kss}}$ with reduced structure. By definition, we know that $\overline{\mathcal{Y}}$ is also a closed substack of $\overline{\mathcal{M}}_{3,16}^{\text{sm}, \text{Kss}}$. Let $\overline{\mathfrak{Y}}$ be the good moduli space of $\overline{\mathcal{Y}}$, then $\overline{\mathfrak{Y}}$ is a closed subscheme of $\overline{\mathcal{M}}_{3,16}^{\text{sm}, \text{Kps}}$ and $M_{3,16}^{\text{Kps}}$. Since \mathcal{Y} parametrizes K-stable pairs, it is a saturated Deligne-Mumford open substack of $\overline{\mathfrak{Y}}$. Hence \mathcal{Y} admits a coarse moduli space \mathfrak{Y} as an open subscheme of $\overline{\mathfrak{Y}}$. We call $\overline{\mathfrak{Y}}$ the *K-moduli space of quartic double solids*. We know that $\overline{\mathfrak{Y}}$ is an irreducible component of $M_{3,16}^{\text{Kps}}$ since \mathfrak{Y} is open in $M_{3,16}^{\text{Kps}}$.

Proposition 6.12. *There exists a bijective morphism $\iota : \overline{\mathfrak{M}}_{\frac{1}{2}}^{\text{K}} \rightarrow \overline{\mathfrak{Y}}$.*

Proof. Consider the Hilbert scheme \mathbb{H} of pairs $(X, D) \subset \mathbb{P}^9$ with Hilbert polynomial $\chi(\mathbb{P}^3, \mathcal{O}(4))$ and $D \sim -K_X$. Let \mathcal{H} denote the locally closed subscheme of \mathbb{H} parameterizing K-semistable pairs $(X, \frac{1}{2}D) \subset \mathbb{P}^9$ (see [ADL19, Definition 3.7] or [ADL20, Theorem 2.21]). Because $c = \frac{1}{2}$, the wall crossing results in Section 5.4 prove that, if $[(X, \frac{1}{2}D)] \in \mathcal{H}$, then $X \cong \mathbb{P}^3, X_h$, or $\mathbb{P}(1, 1, 2, 4)$.

By Lemma 5.4, the \mathbb{Q} -Gorenstein deformations of $X = \mathbb{P}^3, X_h$, or $\mathbb{P}(1, 1, 2, 4)$ are unobstructed in $\overline{\mathcal{M}}_{3,64}^{\text{sm}, \delta \geq \epsilon_0}$, and there is no torsion in the class group for $\mathcal{X}_t = \mathbb{P}^3, X_h$, or $\mathbb{P}(1, 1, 2, 4)$ (see e.g.

[Kol13, Proposition 3.14]). Hence $\mathcal{O}_{\mathbb{P}^9}(1)|_X$ is the unique Weil divisor class that is \mathbb{Q} -linearly equivalent to $-\frac{1}{2}K_X$. Therefore, \mathcal{H} is smooth, so the quotient stack $[\mathcal{H}/\mathrm{PGL}(10, \mathbb{C})]$ is smooth.

By construction of \mathcal{H} which parameterizes K-semistable pairs, universality of K-moduli gives a map $[\mathcal{H}/\mathrm{PGL}(10, \mathbb{C})] \rightarrow \overline{\mathcal{M}}_{\frac{1}{2}}^K$. This map is separated, stabilizer preserving, and bijective on \mathbb{C} -points. Therefore, by [AI19, Theorem A.5], we have that $[\mathcal{H}/\mathrm{PGL}(10, \mathbb{C})] \cong \overline{\mathcal{M}}_{\frac{1}{2}}^K$.

We now construct a morphism $\iota : \overline{\mathcal{M}}_{\frac{1}{2}}^K \rightarrow \overline{\mathcal{Y}}$. Consider the universal family $(\mathcal{X}, \mathcal{D}) \rightarrow \overline{\mathcal{M}}_{\frac{1}{2}}^K$. By the isomorphism of $[\mathcal{H}/\mathrm{PGL}(10, \mathbb{C})]$ and $\overline{\mathcal{M}}_{\frac{1}{2}}^K$, there is a line bundle $\mathcal{O}_{\mathcal{X}}(1)$ on \mathcal{X} obtained as the pull-back of the line bundle $\mathcal{O}_{\mathbb{P}^9}(1)$ from the universal family on the Hilbert scheme. Since $\mathcal{O}_{\mathcal{X}_t}(2) \sim -K_{\mathcal{X}_t} \sim \mathcal{D}_t$ on each fiber \mathcal{X}_t for any $t \in |\overline{\mathcal{M}}_{\frac{1}{2}}^K|$, we know that $\mathcal{O}_{\mathcal{X}}(\mathcal{D}) \otimes \mathcal{O}_{\mathcal{X}}(-2)$ is trivial on every fiber \mathcal{X}_t , which implies that it is the pull-back of a line bundle \mathcal{F} on $\overline{\mathcal{M}}_{\frac{1}{2}}^K$. Let $\phi_{\mathcal{Z}} : \mathcal{Z} \rightarrow \overline{\mathcal{M}}_{\frac{1}{2}}^K$ be the μ_2 -gerbe obtained as the second root stack of \mathcal{F} (see e.g. [AGV08, Appendix B.1]), i.e. $\mathcal{Z} := \overline{\mathcal{M}}_{\frac{1}{2}}^K \times_{B\mathbb{G}_m} B\mathbb{G}_m$ where $\overline{\mathcal{M}}_{\frac{1}{2}}^K \rightarrow B\mathbb{G}_m$ is the classifying morphism of \mathcal{F} , and $B\mathbb{G}_m \rightarrow B\mathbb{G}_m$ is the second power map. Hence there is a line bundle \mathcal{G} on \mathcal{Z} such that $\mathcal{G}^{\otimes 2}$ is the pull-back of \mathcal{F} on \mathcal{Z} . Denote by $\pi_{\mathcal{Z}} : (\mathcal{X}_{\mathcal{Z}}, \mathcal{D}_{\mathcal{Z}}) \rightarrow \mathcal{Z}$ the base change of the family $(\mathcal{X}, \mathcal{D})$ to \mathcal{Z} . Then there is a line bundle $\mathcal{N}_{\mathcal{Z}} := \mathcal{O}_{\mathcal{X}_{\mathcal{Z}}}(1) \otimes \pi_{\mathcal{Z}}^* \mathcal{G}$ on \mathcal{Z} satisfying $\mathcal{N}_{\mathcal{Z}}^{\otimes 2} \cong \mathcal{O}_{\mathcal{X}_{\mathcal{Z}}}(\mathcal{D}_{\mathcal{Z}})$.

Consider the double cover $\mathcal{Y}_{\mathcal{Z}}$ of $\mathcal{X}_{\mathcal{Z}}$ branched along $\mathcal{D}_{\mathcal{Z}}$, i.e.

$$\mathcal{Y}_{\mathcal{Z}} := \mathrm{Spec}_{\mathcal{X}_{\mathcal{Z}}} \mathcal{O}_{\mathcal{X}_{\mathcal{Z}}} \oplus \mathcal{N}_{\mathcal{Z}}^{\otimes -1},$$

where the $\mathcal{O}_{\mathcal{X}_{\mathcal{Z}}}$ -algebra structure is induced by the sheaf homomorphism $\mathcal{N}_{\mathcal{Z}}^{\otimes -2} \xrightarrow{s} \mathcal{O}_{\mathcal{X}_{\mathcal{Z}}}$ where s is a section of $\mathcal{N}_{\mathcal{Z}}^{\otimes 2}$ such that $(s = 0) = \mathcal{D}_{\mathcal{Z}}$. Since $\mathcal{N}_{\mathcal{Z}}$ is locally free, the double cover $\mathcal{Y}_{\mathcal{Z}} \rightarrow \mathcal{X}_{\mathcal{Z}}$ is also a fiberwise double cover. In particular, each fiber $(\mathcal{X}_t, \frac{1}{2}\mathcal{D}_t)$ for $t \in |\overline{\mathcal{M}}_{\frac{1}{2}}^K|$ is the μ_2 -quotient of \mathcal{Y}_z where $z \in |\mathcal{Z}|$ is the unique point lying over t . Thus, by [LZ20, Zhu21] we know that $\mathcal{Y}_{\mathcal{Z}} \rightarrow \mathcal{Z}$ is a \mathbb{Q} -Fano family with K-semistable fibers, where a general fiber is a smooth quartic double solid. Therefore, this gives $\mathcal{Z} \rightarrow \overline{\mathcal{Y}}$ by the universality of the K-moduli stack.

Next we prove that the composition $\mathcal{Z} \xrightarrow{\phi_{\mathcal{Z}}} \overline{\mathcal{M}}_{\frac{1}{2}}^K \rightarrow \overline{\mathcal{M}}_{\frac{1}{2}}^K$ provides a good moduli space of \mathcal{Z} . By [Alp13] it suffices to show that $(\phi_{\mathcal{Z}})_* \mathcal{O}_{\mathcal{Z}} = \mathcal{O}_{\overline{\mathcal{M}}_{\frac{1}{2}}^K}$ and $\phi_{\mathcal{Z}}$ is cohomologically affine. The first statement follows from the fact that $\phi_{\mathcal{Z}}$ is a μ_2 -gerbe. For the second statement, applying [Alp13, Proposition 3.10(vii)] to the Cartesian diagram in the fiber product construction of \mathcal{Z} , it suffices to show that the second power map $f : B\mathbb{G}_m \rightarrow B\mathbb{G}_m$ is cohomologically affine. A quasi-coherent sheaf V over $B\mathbb{G}_m$ corresponds via the weight decomposition to a family $(V_i)_{i \in \mathbb{Z}}$ of \mathbb{C} -vector spaces. It is clear that $W := f_* V$ corresponds to $(W_j)_{j \in \mathbb{Z}}$ where $W_j = V_{2j}$. Since $V \mapsto V_i$ is exact for every $i \in \mathbb{Z}$, we know that f_* is exact. Hence $\phi_{\mathcal{Z}}$ is cohomologically affine, which implies that $\mathcal{Z} \rightarrow \overline{\mathcal{M}}_{\frac{1}{2}}^K$ is a good moduli space morphism. Descending the map $\mathcal{Z} \rightarrow \overline{\mathcal{Y}}$ to level of good moduli spaces gives the desired morphism $\iota : \overline{\mathcal{M}}_{\frac{1}{2}}^K \rightarrow \overline{\mathcal{Y}}$.

Next, we will show that $\iota : \overline{\mathcal{M}}_{\frac{1}{2}}^K \rightarrow \overline{\mathcal{Y}}$ is bijective. Clearly, \mathcal{Y} is contained in the image of ι , so properness of $\overline{\mathcal{M}}_{\frac{1}{2}}^K$ implies the surjectivity of ι . It suffices to show injectivity of ι , i.e. for any two points $[(X, D)], [(X', D')] \in \overline{\mathcal{M}}_{\frac{1}{2}}^K$, if their double covers Y and Y' are isomorphic, then $(X, D) \cong (X', D')$. First of all, we know that X (resp. X') is isomorphic to \mathbb{P}^3 , X_h , or $\mathbb{P}(1, 1, 2, 4)$ by Theorem 5.16, where in the latter two cases D (resp. D') does not pass through the cone vertex. By Lemma 3.3, we know that there exist ample \mathbb{Q} -Cartier Weil divisorial sheaves L and L' on X and X' respectively, such that $-K_X = 4L$ and $-K_{X'} = 4L'$. In addition, if X (resp.

X' is a cone, then L (resp. L') is Cartier away from the cone vertex, and it has Cartier index 2 at the cone vertex. Let \tilde{L} and \tilde{L}' be the pull-back of L and L' to Y and Y' respectively. If X (resp. X') is smooth, then clearly Y (resp. Y') has local complete intersection singularities. If X (resp. X') is singular, then Y (resp. Y') has precisely two singularities, the preimage of the cone vertex, that are not local complete intersections, since D (resp. D') is away from the cone vertex by Theorem 5.16. Since $Y \cong Y'$, we know that X is smooth if and only if X' is smooth. We split into two cases. For simplicity we assume $Y = Y'$, and denote the double cover maps by $\pi : Y \rightarrow X$ and $\pi' : Y \rightarrow X'$.

Case 1: $X \cong X' \cong \mathbb{P}^3$. In this case, both \tilde{L} and \tilde{L}' are Cartier on Y , and $-K_Y = 2\tilde{L} = 2\tilde{L}'$ which implies that $\tilde{L} - \tilde{L}'$ is a torsion Cartier divisor on Y . Since Y is \mathbb{Q} -Fano, it is rationally connected by [Zha06] and hence simply connected. Thus any torsion line bundle on Y is trivial which implies $\tilde{L} = \tilde{L}'$. By [KM98, Definition 2.50] we know that $\pi_*\mathcal{O}_Y(\tilde{L}) \cong \mathcal{O}_X(L) \oplus \mathcal{O}_X(-L)$ and $\pi'_*\mathcal{O}_Y(\tilde{L}') \cong \mathcal{O}_{X'}(L') \oplus \mathcal{O}_{X'}(-L')$. Thus

$$(6.12) \quad H^0(X, \mathcal{O}_X(L)) \cong H^0(Y, \tilde{L}) = H^0(Y, \tilde{L}') \cong H^0(X', \mathcal{O}_{X'}(L')).$$

Hence the linear system $|\tilde{L}|$ induces a map $Y \rightarrow \mathbb{P}^3$ isomorphic to both π and π' . By taking ramification divisors, we obtain $(X, D) \cong (X', D')$.

Case 2: both X and X' are cones. In this case, denote the unique non-lci singularity in X and X' by x and x' , respectively. Then $\pi^{-1}(x) = \pi'^{-1}(x') =: \{y_1, y_2\}$. From the geometry of X_h and $\mathbb{P}(1, 1, 2, 4)$, we know that $\text{Pic}(x \in X) \cong \text{Pic}(x' \in X') \cong \mathbb{Z}/2\mathbb{Z}$ where L and L' are generators respectively. Since π (resp. π') is étale over a neighborhood of x (resp. of x'), we know that $\tilde{L} - \tilde{L}'$ is a torsion Cartier divisor on Y , and as above we conclude that $\tilde{L} = \tilde{L}'$. We have (6.12), and also $\pi_*\mathcal{O}_Y(2\tilde{L}) = \mathcal{O}_X(2L) \oplus \mathcal{O}_X$ and $\pi'_*\mathcal{O}_Y(2\tilde{L}') \cong \mathcal{O}_{X'}(2L') \oplus \mathcal{O}_{X'}$ which implies

$$(6.13) \quad H^0(X, \mathcal{O}_X(2L)) \oplus \mathbb{C} \cong H^0(Y, 2\tilde{L}) = H^0(Y, 2\tilde{L}') \cong H^0(X', \mathcal{O}_{X'}(2L')) \oplus \mathbb{C}.$$

By choosing a basis (s_0, s_1, s_2, s_3) of $H^0(X, L)$ and an element $s_4 \in H^0(Y, 2\tilde{L}) \setminus H^0(X, \mathcal{O}_X(2L))$, we obtain a morphism $[s_0, \dots, s_4] : Y \rightarrow \mathbb{P}(1^4, 2)$ which is isomorphic to π after taking the image. Similarly, we have $[s'_0, \dots, s'_4] : Y \rightarrow \mathbb{P}(1^4, 2)$ isomorphic to π' after taking the image. From the construction, (6.12), and (6.13), we know that π and π' only differ by an automorphism of $\mathbb{P}(1^4, 2)$, so they are isomorphic to each other. Thus $(X, D) \cong (X', D')$. \square

Proof of Theorem 1.4. The first statement follows from Proposition 6.12. The diagram follows from Theorem 5.16 where $p = [(\mathbb{P}(1, 1, 2, 4), (x_3^2 = x_2^4))]$. Then $\tau(p)$ represents the weighted hypersurface $(x_4^2 = x_3^2 - x_2^4) \subset \mathbb{P}(1, 1, 2, 4, 4)$ which is isomorphic to $(x_2^4 = x_3 x_4) \subset \mathbb{P}(1, 1, 2, 4, 4)$ after an automorphism of $\mathbb{P}(1, 1, 2, 4, 4)$. \square

Remark 6.13. If we take quadruple cyclic covers instead of double covers, then similar arguments show that there is a finite birational morphism $\overline{\mathfrak{M}}_{3,4}^K \rightarrow M_{3,4}^{\text{Kps}}$ whose image is the Zariski closure of the moduli space of smooth quartic threefolds that are cyclic covers of \mathbb{P}^3 under linear projections. Note that smooth quartic threefolds are known to be K-stable by [Che01, CP02, Fuj19a], while their K-moduli compactification is currently unknown.

6.4. Gorenstein \mathbb{Q} -Fano degenerations of \mathbb{P}^3 .

In this subsection, we prove Theorem 1.5.

Proof of Theorem 1.5. Let X be a Gorenstein \mathbb{Q} -Fano degeneration of \mathbb{P}^3 . By the effective non-vanishing theorem of Ambro [Amb99, Main Theorem] and Kawamata [Kaw00, Theorem 5.1], there exists an effective Cartier divisor $S \in |-K_X|$ such that (X, S) is plt. Hence Theorem 2.10 implies that $(X, (1 - \epsilon)S)$ is uniformly K-stable for $0 < \epsilon \ll 1$. Let $\pi : \mathcal{X} \rightarrow B$ be a \mathbb{Q} -Fano family over a smooth pointed curve $0 \in B$ such that $\mathcal{X}_0 \cong X$ and $\mathcal{X}_b \cong \mathbb{P}^3$ for any $b \in B \setminus \{0\}$. By

Lemma 3.3 we know that $\pi_*\omega_{\mathcal{X}/B}^\vee$ is a vector bundle over B whose fiber over $b \in B$ is precisely $H^0(\mathcal{X}_b, \omega_{\mathcal{X}_b}^\vee)$. In particular, we know that (X, S) admits a \mathbb{Q} -Gorenstein smoothing to (\mathbb{P}^3, S_b) where S_b is a smooth quartic surface. Hence $[(X, S)] \in \overline{\mathcal{M}}_{1-\epsilon}^K$, and the statement follows from Proposition 5.15. \square

REFERENCES

- [AB19] Kenneth Ascher and Dori Bejleri. Compact moduli of elliptic K3 surfaces. *Geom. Topol., to appear*, 2019. [arXiv:1902.10686](https://arxiv.org/abs/1902.10686).
- [ABHLX20] Jarod Alper, Harold Blum, Daniel Halpern-Leistner, and Chenyang Xu. Reductivity of the automorphism group of K-polystable Fano varieties. *Invent. Math.*, 222(3):995–1032, 2020.
- [ADL19] Kenneth Ascher, Kristin DeVleming, and Yuchen Liu. Wall crossing for K-moduli spaces of plane curves. 2019. [arXiv:1909.04576](https://arxiv.org/abs/1909.04576).
- [ADL20] Kenneth Ascher, Kristin DeVleming, and Yuchen Liu. K-moduli of curves on a quadric surface and K3 surfaces. *J. Inst. Math. Jussieu, to appear*, 2020. [arXiv:2006.06816](https://arxiv.org/abs/2006.06816).
- [AE21] Valery Alexeev and Philip Engel. Compact moduli of K3 surfaces, 2021. [arXiv:2101.12186](https://arxiv.org/abs/2101.12186).
- [AET19] Valery Alexeev, Philip Engel, and Alan Thompson. Stable pair compactification of moduli of K3 surfaces of degree 2. 2019. [arXiv:1903.09742](https://arxiv.org/abs/1903.09742).
- [AFS17] Jarod Alper, Maksym Fedorchuk, and David Ishii Smyth. Second flip in the Hassett-Keel program: existence of good moduli spaces. *Compos. Math.*, 153(8):1584–1609, 2017.
- [AGV08] Dan Abramovich, Tom Graber, and Angelo Vistoli. Gromov-Witten theory of Deligne-Mumford stacks. *Amer. J. Math.*, 130(5):1337–1398, 2008.
- [AI19] Shamil Asgarli and Giovanni Inchiostro. The Picard group of the moduli of smooth complete intersections of two quadrics. *Trans. Amer. Math. Soc.*, 372(5):3319–3346, 2019.
- [Alp13] Jarod Alper. Good moduli spaces for Artin stacks. *Ann. Inst. Fourier (Grenoble)*, 63(6):2349–2402, 2013.
- [Alt97] Klaus Altmann. The versal deformation of an isolated toric Gorenstein singularity. *Invent. Math.*, 128(3):443–479, 1997.
- [Amb99] Florin Ambro. Ladders on Fano varieties. *J. Math. Sci. (New York)*, 94(1):1126–1135, 1999. Algebraic geometry, 9.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23(2):405–468, 2010.
- [BHLLX21] Harold Blum, Daniel Halpern-Leistner, Yuchen Liu, and Chenyang Xu. On properness of K-moduli spaces and optimal degenerations of Fano varieties. *Selecta Math. (N.S.)*, 27(4):Paper No. 73, 2021.
- [Bir19] Caucher Birkar. Anti-pluricanonical systems on Fano varieties. *Ann. of Math. (2)*, 190(2):345–463, 2019.
- [BJ20] Harold Blum and Mattias Jonsson. Thresholds, valuations, and K-stability. *Adv. Math.*, 365:107062, 57, 2020.
- [BL22] Harold Blum and Yuchen Liu. Openness of uniform K-stability in families of \mathbb{Q} -Fano varieties. *Ann. Sci. Éc. Norm. Supér. (4)*, 55(1):1–41, 2022.
- [BLX22] Harold Blum, Yuchen Liu, and Chenyang Xu. Openness of K-semistability for Fano varieties. *Duke Math. J.*, 171(13):2753 – 2797, 2022.
- [BX19] Harold Blum and Chenyang Xu. Uniqueness of K-polystable degenerations of Fano varieties. *Ann. of Math. (2)*, 190(2):609–656, 2019.
- [Cam92] F. Campana. Connexité rationnelle des variétés de Fano. *Ann. Sci. École Norm. Sup. (4)*, 25(5):539–545, 1992.
- [Che01] I. A. Cheltsov. Log canonical thresholds on hypersurfaces. *Mat. Sb.*, 192(8):155–172, 2001.
- [CP02] I. A. Cheltsov and Jihun Park. Global log-canonical thresholds and generalized Eckardt points. *Mat. Sb.*, 193(5):149–160, 2002.
- [CP21] Giulio Codogni and Zsolt Patakfalvi. Positivity of the CM line bundle for families of K-stable klt Fano varieties. *Invent. Math.*, 223(3):811–894, 2021.
- [Der16] Ruadhai Dervan. On K-stability of finite covers. *Bull. Lond. Math. Soc.*, 48(4):717–728, 2016.
- [DeV22] Kristin DeVleming. Moduli of surfaces in \mathbb{P}^3 . *Compositio Mathematica*, 158(6):1329–1374, 2022.
- [Dol82] Igor Dolgachev. Weighted projective varieties. In *Group actions and vector fields (Vancouver, B.C., 1981)*, volume 956 of *Lecture Notes in Math.*, pages 34–71. Springer, Berlin, 1982.

[EL20] Lawrence Ein and Robert Lazarsfeld. Tangent developable surfaces and the equations defining algebraic curves. *Bull. Amer. Math. Soc. (N.S.)*, 57(1):23–38, 2020.

[EVdV81] D. Eisenbud and A. Van de Ven. On the normal bundles of smooth rational space curves. *Math. Ann.*, 256(4):453–463, 1981.

[FO18] Kento Fujita and Yuji Odaka. On the K-stability of Fano varieties and anticanonical divisors. *Tohoku Math. J. (2)*, 70(4):511–521, 2018.

[FR06] J. Fine and J. Ross. A note on positivity of the CM line bundle. *Int. Math. Res. Not.*, pages Art. ID 95875, 14, 2006.

[Fuj90] Takao Fujita. On singular del Pezzo varieties. In *Algebraic geometry (L’Aquila, 1988)*, volume 1417 of *Lecture Notes in Math.*, pages 117–128. Springer, Berlin, 1990.

[Fuj19a] Kento Fujita. K-stability of Fano manifolds with not small alpha invariants. *J. Inst. Math. Jussieu*, 18(3):519–530, 2019.

[Fuj19b] Kento Fujita. Uniform K-stability and plt blowups of log Fano pairs. *Kyoto J. Math.*, 59(2):399–418, 2019.

[Fuj19c] Kento Fujita. A valuative criterion for uniform K-stability of \mathbb{Q} -Fano varieties. *J. Reine Angew. Math.*, 751:309–338, 2019.

[GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell. Differential forms on log canonical spaces. *Publ. Math. Inst. Hautes Études Sci.*, (114):87–169, 2011.

[GMGS21] Patricio Gallardo, Jesus Martinez-Garcia, and Cristiano Spotti. Applications of the moduli continuity method to log K-stable pairs. *J. Lond. Math. Soc. (2)*, 103(2):729–759, 2021.

[Hac01] Paul Hacking. *A compactification of the space of plane curves*. 2001. Thesis (Ph.D.)– Cambridge Univserty.

[Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.

[HH13] Brendan Hassett and Donghoon Hyeon. Log minimal model program for the moduli space of stable curves: the first flip. *Annals of Mathematics*, 177:1–58, 2013.

[Huy16] Daniel Huybrechts. *Lectures on K3 surfaces*, volume 158 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016.

[IP99] V. A. Iskovskikh and Yu. G. Prokhorov. Fano varieties. In *Algebraic geometry, V*, volume 47 of *Encyclopaedia Math. Sci.*, pages 1–247. Springer, Berlin, 1999.

[Jia20] Chen Jiang. Boundedness of \mathbb{Q} -Fano varieties with degrees and alpha-invariants bounded from below. *Ann. Sci. Éc. Norm. Supér. (4)*, 53(5):1235–1248, 2020.

[JMR16] Thalia Jeffres, Rafe Mazzeo, and Yanir A. Rubinstein. Kähler-Einstein metrics with edge singularities. *Ann. of Math. (2)*, 183(1):95–176, 2016.

[Kaw00] Yujiro Kawamata. On effective non-vanishing and base-point-freeness. *Asian J. Math.*, 4(1):173–181, 2000. Kodaira’s issue.

[KKL16] Anne-Sophie Kaloghiros, Alex Küronya, and Vladimir Lazić. Finite generation and geography of models. In *Minimal models and extremal rays (Kyoto, 2011)*, volume 70 of *Adv. Stud. Pure Math.*, pages 215–245. Math. Soc. Japan, [Tokyo], 2016.

[KM76] Finn Faye Knudsen and David Mumford. The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”. *Math. Scand.*, 39(1):19–55, 1976.

[KM98] János Kollar and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.

[KMM92] János Kollar, Yoichi Miyaoka, and Shigefumi Mori. Rational connectedness and boundedness of Fano manifolds. *J. Differential Geom.*, 36(3):765–779, 1992.

[Kni73] Carol M. Knighten. Differentials on quotients of algebraic varieties. *Trans. Amer. Math. Soc.*, 177:65–89, 1973.

[Kol97] János Kollar. Singularities of pairs. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 221–287. Amer. Math. Soc., Providence, RI, 1997.

[Kol13] János Kollar. *Singularities of the minimal model program*, volume 200 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013. With a collaboration of Sándor Kovács.

[Kol18] János Kollar. Mumford divisors. 2018. [arXiv:1803.07596](https://arxiv.org/abs/1803.07596).

[Kol19] János Kollar. Families of divisors. 2019. [arXiv:1910.00937](https://arxiv.org/abs/1910.00937).

[Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics*

and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.

[Laz16] Radu Laza. The KSBA compactification for the moduli space of degree two $K3$ pairs. *J. Eur. Math. Soc. (JEMS)*, 18(2):225–279, 2016.

[Li17] Chi Li. K-semistability is equivariant volume minimization. *Duke Math. J.*, 166(16):3147–3218, 2017.

[Liu22] Yuchen Liu. K-stability of cubic fourfolds. *J. Reine Angew. Math.*, 786:55–77, 2022.

[LL19] Chi Li and Yuchen Liu. Kähler-Einstein metrics and volume minimization. *Adv. Math.*, 341:440–492, 2019.

[LLX20] Chi Li, Yuchen Liu, and Chenyang Xu. A guided tour to normalized volume. In *Geometric analysis*, volume 333 of *Progr. Math.*, pages 167–219. Birkhäuser/Springer, Cham, 2020.

[LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.

[LO18] Radu Laza and Kieran G. O’Grady. GIT versus Baily-Borel compactification for quartic $K3$ surfaces. In *Geometry of moduli*, volume 14 of *Abel Symp.*, pages 217–283. Springer, Cham, 2018.

[LO19] Radu Laza and Kieran O’Grady. Birational geometry of the moduli space of quartic $K3$ surfaces. *Compos. Math.*, 155(9):1655–1710, 2019.

[LO21] Radu Laza and Kieran O’Grady. GIT versus Baily-Borel compactification for $K3$ ’s which are double covers of $\mathbb{P}^1 \times \mathbb{P}^1$. *Adv. Math.*, 383:107680, 63, 2021.

[Loo86] Eduard Looijenga. New compactifications of locally symmetric varieties. In *Proceedings of the 1984 Vancouver conference in algebraic geometry*, volume 6 of *CMS Conf. Proc.*, pages 341–364. Amer. Math. Soc., Providence, RI, 1986.

[Loo03a] Eduard Looijenga. Compactifications defined by arrangements. I. The ball quotient case. *Duke Math. J.*, 118(1):151–187, 2003.

[Loo03b] Eduard Looijenga. Compactifications defined by arrangements. II. Locally symmetric varieties of type IV. *Duke Math. J.*, 119(3):527–588, 2003.

[LWX19] Chi Li, Xiaowei Wang, and Chenyang Xu. On the proper moduli spaces of smoothable Kähler-Einstein Fano varieties. *Duke Math. J.*, 168(8):1387–1459, 2019.

[LWX21] Chi Li, Xiaowei Wang, and Chenyang Xu. Algebraicity of the metric tangent cones and equivariant K-stability. *J. Amer. Math. Soc.*, 34(4):1175–1214, 2021.

[LX14] Chi Li and Chenyang Xu. Special test configuration and K-stability of Fano varieties. *Ann. of Math. (2)*, 180(1):197–232, 2014.

[LX19] Yuchen Liu and Chenyang Xu. K-stability of cubic threefolds. *Duke Math. J.*, 168(11):2029–2073, 2019.

[LX20] Chi Li and Chenyang Xu. Stability of valuations and Kollar components. *J. Eur. Math. Soc. (JEMS)*, 22(8):2573–2627, 2020.

[LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to K-stability. *Ann. of Math. (2)*, 196(2):507–566, 2022.

[LZ20] Yuchen Liu and Ziwen Zhu. Equivariant K-stability under finite group action. 2020. [arXiv:2001.10557](https://arxiv.org/abs/2001.10557).

[LZ22] Yuchen Liu and Ziquan Zhuang. On the sharpness of Tian’s criterion for K-stability. *Nagoya Math. J.*, 245:41–73, 2022.

[May72] Alan L. Mayer. Families of $K - 3$ surfaces. *Nagoya Mathematical Journal*, 48:1–17, 1972.

[Mir81] Rick Miranda. The moduli of Weierstrass fibrations over \mathbb{P}^1 . *Math. Ann.*, 255(3):379–394, 1981.

[Nam97] Yoshinori Namikawa. Smoothing Fano 3-folds. *J. Algebraic Geom.*, 6(2):307–324, 1997.

[Oda13] Yuji Odaka. The GIT stability of polarized varieties via discrepancy. *Ann. of Math. (2)*, 177(2):645–661, 2013.

[Oda15] Yuji Odaka. Compact moduli spaces of Kähler-Einstein Fano varieties. *Publ. Res. Inst. Math. Sci.*, 51(3):549–565, 2015.

[OO21] Yuji Odaka and Yoshiki Oshima. *Collapsing $K3$ surfaces, tropical geometry and moduli compactifications of Satake, Morgan-Shalen type*, volume 40 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2021.

[OSS16] Yuji Odaka, Cristiano Spotti, and Song Sun. Compact moduli spaces of del Pezzo surfaces and Kähler-Einstein metrics. *J. Differential Geom.*, 102(1):127–172, 2016.

[Pet22a] Andrea Petracci. On deformation spaces of toric singularities and on singularities of K-moduli of Fano varieties. *Trans. Amer. Math. Soc.*, 375(8):5617–5643, 2022.

[Pet22b] Andrea Petracci. On deformations of toric Fano varieties. In *Interactions with lattice polytopes*, volume 386 of *Springer Proc. Math. Stat.*, pages 287–314. Springer, Cham, [2022] ©2022.

[Pos22] Quentin Posva. Positivity of the CM line bundle for K-stable log Fano varieties. *Trans. Amer. Math. Soc.*, 375(7):4943–4978, 2022.

[Pro00] Yu. G. Prokhorov. Blow-ups of canonical singularities. In *Algebra (Moscow, 1998)*, pages 301–317. de Gruyter, Berlin, 2000.

[PT06] Sean Timothy Paul and Gang Tian. CM stability and the generalized Futaki invariant I. 2006. [arXiv:math/0605278](https://arxiv.org/abs/math/0605278).

[PT09] Sean Timothy Paul and Gang Tian. CM stability and the generalized Futaki invariant II. *Astérisque*, (328):339–354 (2010), 2009.

[Ser06] Edoardo Sernesi. *Deformations of algebraic schemes*, volume 334 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2006.

[Sha80] Jayant Shah. A complete moduli space for $K3$ surfaces of degree 2. *Ann. of Math. (2)*, 112(3):485–510, 1980.

[Sha81] Jayant Shah. Degenerations of $K3$ surfaces of degree 4. *Trans. Amer. Math. Soc.*, 263(2):271–308, 1981.

[Shi89] Kil-Ho Shin. 3-dimensional Fano varieties with canonical singularities. *Tokyo J. Math.*, 12(2):375–385, 1989.

[SS17] Cristiano Spotti and Song Sun. Explicit Gromov-Hausdorff compactifications of moduli spaces of Kähler-Einstein Fano manifolds. *Pure Appl. Math. Q.*, 13(3):477–515, 2017.

[Sta18] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2018.

[Ste77a] J. H. M. Steenbrink. Mixed Hodge structure on the vanishing cohomology. In *Real and complex singularities (Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976)*, pages 525–563. Sijhoff and Noordhoff, Alphen aan den Rijn, 1977.

[Ste77b] Joseph Steenbrink. Intersection form for quasi-homogeneous singularities. *Compositio Math.*, 34(2):211–223, 1977.

[Tia97] Gang Tian. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.*, 130(1):1–37, 1997.

[Xu14] Chenyang Xu. Finiteness of algebraic fundamental groups. *Compos. Math.*, 150(3):409–414, 2014.

[Xu20] Chenyang Xu. A minimizing valuation is quasi-monomial. *Ann. of Math. (2)*, 191(3):1003–1030, 2020.

[Xu21] Chenyang Xu. K-stability of Fano varieties: an algebro-geometric approach. *EMS Surv. Math. Sci.*, 8(1-2):265–354, 2021.

[XZ20] Chenyang Xu and Ziquan Zhuang. On positivity of the CM line bundle on K-moduli spaces. *Ann. of Math. (2)*, 192(3):1005–1068, 2020.

[XZ21] Chenyang Xu and Ziquan Zhuang. Uniqueness of the minimizer of the normalized volume function. *Camb. J. Math.*, 9(1):149–176, 2021.

[Zha06] Qi Zhang. Rational connectedness of log \mathbf{Q} -Fano varieties. *J. Reine Angew. Math.*, 590:131–142, 2006.

[Zho21a] Chuyu Zhou. Log K-stability of GIT-stable divisors on Fano manifolds. 2021. [arXiv:2102.07458](https://arxiv.org/abs/2102.07458).

[Zho21b] Chuyu Zhou. On wall crossing for K-stability. 2021. [arXiv:2103.04925](https://arxiv.org/abs/2103.04925).

[Zhu21] Ziquan Zhuang. Optimal destabilizing centers and equivariant K-stability. *Invent. Math.*, 226(1):195–223, 2021.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA, 92697, USA

Email address: `kascher@uci.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003-9305, USA

Email address: `kdevleming@umass.edu`

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208, USA

Email address: `yuchenl@northwestern.edu`