

Smoothability of relative stable maps to stacky curves

Kenneth Ascher and Dori Bejleri

Abstract. Using log geometry, we study smoothability of genus 0 twisted stable maps to stacky curves relative to a collection of marked points. One application is to smoothing semi-log canonical fibered surfaces with marked singular fibers.

Keywords. Moduli of curves, moduli of stable maps, log geometry, log stable maps

2020 Mathematics Subject Classification. 14A21, 14D23, 14D20, 14H10

Received by the Editors on November 11, 2021, and in final form on July 23, 2022.
 Accepted on September 17, 2022.

Kenneth Ascher
 Department of Mathematics, University of California, Irvine, CA 92697-3875, USA
 e-mail: kascher@uci.edu

Dori Bejleri
 Department of Mathematics, Harvard University, Cambridge, MA 02138, USA
 e-mail: bejleri@math.harvard.edu

K.A. is supported in part by funds from NSF Grant DMS-2140781 (formerly DMS-2001408). D.B. is supported in part by an NSF postdoctoral fellowship.

Contents

1. Introduction	2
2. Genus 0 relative stable maps to (\mathbb{P}^1, ∞)	5
3. Log twisted curves and maps	9
4. Log maps to 1-dimensional targets	12
5. Smoothing twisted stable maps	16
6. Proofs of the main theorems	20
References	21

1. Introduction

Consider the moduli space Maps_n° of degree n maps $f: C \rightarrow \mathbb{P}^1$ from a smooth rational curve C such that f is unramified over infinity. Marking the preimage $f^{-1}(\infty) = \{p_1, \dots, p_n\}$ of such a map induces a locally closed embedding

$$\widetilde{\text{Maps}}_n^\circ \hookrightarrow \overline{\mathcal{M}}_{0,n}(\mathbb{P}^1, n)$$

of the \mathfrak{S}_n -torsor $\widetilde{\text{Maps}}_n^\circ \rightarrow \text{Maps}_n^\circ$ parametrizing such $f: C \rightarrow \mathbb{P}^1$ as well as a labeling of $f^{-1}(\infty)$ into the space of n -pointed degree n genus 0 stable maps to \mathbb{P}^1 . The image of this embedding is the locus of stable maps $(f: C \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ such that C is smooth and $f(p_i) = \infty$ for all i . Let $\overline{\text{Maps}}_n$ denote the closure of this locus.

Question 1. Is there a combinatorial description of the boundary $\overline{\text{Maps}}_n \setminus \widetilde{\text{Maps}}_n^\circ$?

There are several natural combinatorial conditions (see Proposition 2.4 and the discussion preceding it) that are necessary for a stable map $(f: C \xrightarrow{n:1} \mathbb{P}^1, p_1, \dots, p_n)$ to lie inside $\overline{\text{Maps}}_n$:

- (1) The evaluations satisfy $f(p_i) = \infty$ for all i .
- (2) Every point of $f^{-1}(\infty) \subset C$ either is a marked point or is on a contracted component.
- (3) For each maximal connected closed subvariety $T \subset C$ contracted to ∞ by f , we have

$$\# \text{ marked points on } T = \sum \text{ ramification index of } f \text{ at the nodes } T \cap (\overline{C \setminus T}).$$

More generally, for any tuple of non-negative integers (d_1, \dots, d_n) with $d = \sum d_i$, one can consider the locally closed subset $\mathcal{M}_{(d_1, \dots, d_n)} \subset \overline{\mathcal{M}}_{0,n}(\mathbb{P}^1, d)$ of n -pointed degree d genus 0 maps $(f: C \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ such that $f(p_i) = \infty$ and f is ramified of order d_i at p_i if $d_i > 0$. If $d_i = 0$, there is no condition imposed on the f at p_i . Denoting by $\overline{\mathcal{M}}_{(d_1, \dots, d_n)}$ the closure of $\mathcal{M}_{(d_1, \dots, d_n)}$, we can ask the variant of Question 1 for this space. The necessary conditions above naturally generalize to Conditions (*) below.

Definition 1.1. Let $(f: C \rightarrow X, p_1, \dots, p_n)$ be a prestable map to a smooth curve, and fix a tuple of positive integers (d_1, \dots, d_n) and a point $x \in X$. We say that f is a **relative map to (X, x) with tangency (d_1, \dots, d_n)** if it satisfies the following conditions:

Combinatorial conditions (*). ([Gat02, Definition 1.1 & Remark 1.7])

- (1) *The evaluations satisfy $f(p_i) = x$ for all i .*
- (2) *Every point of $f^{-1}(x) \subset C$ either is a marked point or is on a contracted component.*

(3) For each maximal connected closed subvariety $T \subset f^{-1}(x)$, we have

$$\sum_{p_i \in T} d_i = \sum_{q \in T \cap (\overline{C \setminus T})} e_f(q),$$

where the first sum is over marked points contained in T and $e_f(q)$ denotes the ramification of $f|_{\overline{C \setminus T}}$ at the point q (see Figure 1).

This question in much greater generality was studied by Gathmann [Gat02, Proposition 1.14], building off previous work of Vakil [Vak00, Theorem 6.1]. In particular, Gathmann showed that Conditions $(*)$ relative to ∞ are both necessary and sufficient for a stable map to lie in $\overline{\mathcal{M}}_{(d_1, \dots, d_n)}$. As a consequence, the set of points of $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^1, d)$ satisfying Conditions $(*)$ relative to ∞ are the points of an irreducible closed substack. See also the balancing condition of Gross–Siebert [GS13, Definition 1.12 and Lemma 1.15].

In [AV02], Abramovich and Vistoli introduced moduli spaces of *twisted stable maps*, which allows the target to instead be a Deligne–Mumford stack. To form a compact moduli space, the source curves obtain a stacky structure. Let $\mathcal{K}_{0,n}(\mathcal{X}, d)$ denote the moduli space of n -pointed genus 0 degree d twisted stable maps to a Deligne–Mumford stack \mathcal{X} . The goal of this paper is to study the analogue of Question 1 for genus 0 twisted stable maps to a weighted projective line $\mathcal{P}(a, b)$, or more generally a genus 0 Deligne–Mumford curve.

When \mathcal{X} is $\mathcal{P}(a, b)$ and $\infty \in \mathcal{P}(a, b)$ is a fixed point away from $[0 : 1]$ and $[1 : 0]$, we have the following (see Theorem 1.5 for a more general result). For this example, the reader can keep in mind $\overline{\mathcal{M}}_{1,1} = \mathcal{P}(4, 6)$ and the point $j = \infty$ parametrizing a nodal elliptic curve.

Theorem 1.2. *Let $(f: \mathcal{C} \rightarrow \mathcal{P}(a, b), p_1, \dots, p_n)$ be an n -pointed genus 0 twisted stable map such that the coarse map $(g: \mathcal{C} \rightarrow \mathbb{P}^1, q_1, \dots, q_n)$ satisfies Conditions $(*)$ with respect to ∞ . Then f is smoothable in a family with generic fiber satisfying: $f(p_i) = \infty$ and f is ramified to order d_i at p_i for all $i = 1, \dots, n$.*

In what follows, we set up notation needed to state our more general results.

Let $(\mathcal{X}, x_1, \dots, x_r)$ be a smooth and proper 1-dimensional genus 0 Deligne–Mumford curve, and suppose the $x_i \in \mathcal{X}$ are points where the coarse moduli space map is étale (equivalently, the x_i have the same stabilizer as the generic point of \mathcal{X}). For each $j = 1, \dots, r$, let $\Gamma_j = (d_{j1}, \dots, d_{jn_j})$ be a tuple of positive integers, and fix $n_0 \geq 0$. Set $n = \sum_{j=0}^r n_j$ and

$$d = \sum_{j=1}^r \sum_{k=1}^{n_j} d_{jk},$$

and let $\Gamma = (n_0, \{\Gamma_1, x_1\}, \dots, \{\Gamma_r, x_r\})$ be the tuple of combinatorial data.

Definition-Notation 1.3. For any tuple of combinatorial data Γ , let $\mathcal{M}_\Gamma(\mathcal{X})$ be the locally closed substack of $\mathcal{K}_{0,n}(\mathcal{X}, d)$ parametrizing n -pointed genus 0 degree d twisted stable maps $(f: \mathcal{C} \rightarrow \mathcal{X}, \{p_{jk}\}_{k=1}^{n_j})_{j=0}^r$ such that

- (1) the p_{jk} are marked points with stabilizer of order a_{jk} lying over smooth points q_{jk} of the coarse space \mathcal{C} ,
- (2) for each $j > 0$, the image of p_{jk} is $f(p_{jk}) = x_j$,
- (3) for each $j > 0$, the coarse map $h: \mathcal{C} \rightarrow \mathcal{X}$ is ramified to order d_{jk} at q_{jk} , and
- (4) \mathcal{C} is smooth.

Similarly, we let $\mathcal{N}_\Gamma(\mathcal{X})$ be the locally closed substack of $\mathcal{K}_{0,n}(\mathcal{X}, d)$ of maps satisfying conditions (1), (2) and (3) above as well as the condition

- (4') \mathcal{C} is smooth in a neighborhood of $f^{-1}(x_j)$ for all j .

We denote by $\overline{\mathcal{M}}_\Gamma(\mathcal{X})$ (resp. $\overline{\mathcal{N}}_\Gamma(\mathcal{X})$) the closure of $\mathcal{M}_\Gamma(\mathcal{X})$ (resp. $\mathcal{N}_\Gamma(\mathcal{X})$) inside $\mathcal{K}_{0,n}(\mathcal{X}, d)$.

Note that here and throughout, we are fixing a bijection between the set $\{1, \dots, n\}$ and the set $\{(j, k)\}_{k=1}^{n_j} \}_{j=0}^r$ indexing the marked points. Note also that the marked points with $j = 0$ are the ones with no tangency conditions, so in the situation of $\mathcal{M}_{(d_1, \dots, d_n)}$, we may assume for notational convenience that the first n_0 entries (d_1, \dots, d_{n_0}) of (d_1, \dots, d_n) are 0 so that $\mathcal{M}_{(d_1, \dots, d_n)} = \mathcal{M}_\Gamma(\mathbb{P}^1)$, where $\Gamma = (n_0, \{(d_{n_0+1}, \dots, d_n), \infty\})$.

Definition-Notation 1.4. For any tuple of combinatorial data Γ , let $\mathcal{K}_\Gamma(\mathcal{X})$ denote the subset of $\mathcal{K}_{0,n}(\mathcal{X}, d)$ parametrizing those twisted stable maps $(f: \mathcal{C} \rightarrow \mathcal{X}, \{(p_{jk})\}_{k=1}^{n_j}\}_{j=0}^r)$ such that for each $j = 1, \dots, r$, the coarse moduli map $(h: C \rightarrow X, q_{j1}, \dots, q_{jn_j})$ is a relative map to (X, x_j) with tangency Γ_j .

Theorem 1.5. *Let \mathcal{X} and Γ be as above. Then we have an equality $\overline{\mathcal{N}}_\Gamma(\mathcal{X}) = \mathcal{K}_\Gamma(\mathcal{X})$. That is, every twisted stable map whose coarse moduli map satisfies the relative condition for $\{\Gamma_j, x_j\}$ for each $j = 1, \dots, r$ is smoothable to a twisted stable map parametrized by $\mathcal{N}_\Gamma(\mathcal{X})$. In particular, $\mathcal{K}_\Gamma(\mathcal{X})$ is the set of points of a closed substack of $\mathcal{K}_{0,n}(\mathcal{X}, d)$.*

Remark 1.6. Note that Theorem 1.5 is local on the target. Indeed, the definitions of both $\mathcal{K}_\Gamma(\mathcal{X})$ and $\mathcal{N}_\Gamma(\mathcal{X})$ are in terms of local conditions around the points $x_i \in \mathcal{X}$. Therefore, the theorem has a natural generalization to higher-genus maps to a higher-genus target \mathcal{X} provided we only consider those maps for which the 1-dimensional components of the preimages $f^{-1}(x_i)$ are rational curves. In this case, the theorem reads that any such map which is also contained in $\mathcal{K}_\Gamma(\mathcal{X})$ is smoothable in a family with generic fiber contained in $\mathcal{N}_\Gamma(\mathcal{X})$.

When the target is a weighted projective line, we obtain the following stronger statement.

Theorem 1.7. *Let $\mathcal{X} = \mathbb{P}(a, b)$ be a weighted projective line, and let Γ be a tuple of combinatorial data as above. Then we have an equality $\overline{\mathcal{M}}_\Gamma(\mathcal{X}) = \mathcal{K}_\Gamma(\mathcal{X})$. That is, every twisted stable map which satisfies the relative condition for $\{\Gamma_j, x_j\}$ for each $j = 1, \dots, r$ is smoothable to a family of stable maps from a smooth rational curve satisfying $f(p_{jk}) = x_j$ and with ramification d_{jk} at p_{jk} .*

1.1. Applications to moduli of fibered surfaces

Our original motivation for writing this paper came from studying compactifications of the moduli space of fibered surfaces. Twisted stable maps are used in [AV00, AB19b] to construct a compactification $\mathcal{F}_{g,n}^\nu(\gamma, \nu)$ of the moduli space of genus γ fibrations over a genus g curve with ν marked sections and n marked fibers. The objects of the boundary are certain semi-log canonical unions of birationally fibered surfaces called *twisted surfaces*. This compactification is closely related to the compactification via stable log varieties from the minimal model program. In [AB19b, Section 1.4], we proposed the problem of using log geometry to give the main component a moduli-theoretic interpretation and classify the boundary components. The present paper solves this problem for elliptic fibrations with marked singular fibers (we refer the reader to [AB21] and [AB19a, Section 4]). A key observation is that Conditions $(*)$ relative to $\infty \in \overline{\mathcal{M}}_{1,1}$ as well as the choice of stabilizers on the marked points translate to conditions on the configuration of singular fibers on the components of the twisted elliptic surface; see [AB19a, Propositions 4.1 and 4.4].

Theorem 1.8. *Theorem 1.7 gives necessary and sufficient combinatorial conditions for a twisted elliptic surface over a genus 0 curve with marked singular fibers to be smoothable to an elliptic surface over \mathbb{P}^1 with marked singular fibers.*

For convenience, we work over an algebraically closed field of characteristic 0.

Acknowledgements

We thank Dan Abramovich, Qile Chen, Martin Olsson, Dhruv Ranganathan, Ravi Vakil, Jonathan Wise, and David Yang for helpful discussions. We thank the referees for their helpful comments.




Figure 1. In this example, the contracted component is attached to a double branched, unbranched, and triply branched point, respectively, and thus must have $2 + 1 + 3 = 6$ marked points.




Figure 2. Maps of this type are given by choosing the map $f_1 = f|_{C_1}$ (which is parametrized by $\overline{\text{Maps}}_n$), a point of the finite set $f_1^{-1}(\infty)$ along which to attach C_2 , and the configuration of the $n+1$ special points on C_2 (which is parametrized by $\mathcal{M}_{0,n+1}$), yielding the dimension count $\dim \overline{\text{Maps}}_n + n - 2$.

2. Genus 0 relative stable maps to (\mathbb{P}^1, ∞)

In this section, we prove the special case of Theorem 1.2 where the target is \mathbb{P}^1 ; *i.e.*, $a = b = 1$. This special case was originally proved by Gathmann [Gat02]. Our approach differs from that of [Gat02] in that we give a direct construction of smoothings of comb-type maps (see Propositions 2.7 and 2.11) rather than appealing to [Vak00, Theorem 6.1]. This will be a key step in the proof of the general case of Theorem 1.2. For the remainder of this section, fix positive integers $\Gamma = (d_1, \dots, d_n)$ with $d = \sum d_i$.

Example 2.1. We begin with some examples motivating Conditions $(*)$ when $\Gamma = (1, \dots, 1)$. First, it is clear that Condition $(*)$ (1) is required as the evaluation condition is closed. However, consider a degree n map $f_1: (C_1, q_1) \rightarrow \mathbb{P}^1$ from a smooth rational curve C_1 with $f(q_1) = \infty$. Let $C = C_1 \cup_{q_1, q_2} C_2$ be a nodal union of two rational curves, and let $p_1, \dots, p_n \in C_2 \setminus q_2$ be n marked points. Then there is an n -marked degree n stable map $(f: C \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ given by taking the map f_1 on C_1 and the constant map with image ∞ on C_2 . While this map satisfies Condition $(*)$ (1), a simple dimension count shows that the dimension of this locus inside $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^1, n)$ is equal to $\dim \overline{\text{Maps}}_n + n - 2$, and so this condition alone is not enough to cut out the locus $\overline{\text{Maps}}_n$ at least for $n \geq 3$ (see Figure 2).

The above example motivates Condition $(*)$ (2), which in this case requires that the map f_1 be totally ramified at q_1 so that there are no other points of C_1 in $f^{-1}(\infty)$. Requiring f_1 to be totally ramified at q_1 means that we impose the vanishing of n derivatives, which is a codimension n condition on $\overline{\text{Maps}}_n$. Thus the locus of maps of this combinatorial type satisfying both Conditions $(*)$ (1) and $(*)$ (2) has dimension $\dim \overline{\text{Maps}}_n - 2$. Note that such maps automatically satisfy Condition $(*)$ (3) as well. However, the following example illustrates that Conditions $(*)$ (1) and $(*)$ (2) do not imply Condition $(*)$ (3) in general.

Example 2.2. Consider a degree 3 map $f_1: (C_1, q_1, q_2) \rightarrow \mathbb{P}^1$ from a smooth rational curve such that $f(q_i) = \infty$, and suppose that f has ramification index i at q_i . Let (C_2, q'_1, p_1, p_2) be a smooth pointed rational curve, and let $C = C_1 \cup_{q_1, q'_1} C_2$. Then we have a 3-pointed degree 3 stable map $(f: C \rightarrow \mathbb{P}^1, q_2, p_1, p_2)$




Figure 3. The map in Example 2.2 which satisfies Conditions (1) and (2) but not (3) since the map is unramified where the contracted component C_2 is attached and ramified over ∞ elsewhere.

where $f|_{C_1} = f_1$ and $f|_{C_2}$ is constant ∞ . This map satisfies Conditions $(*)$ (1) and $(*)$ (2) but not $(*)$ (3). Note furthermore that this stable map cannot lie in $\overline{\text{Maps}}_n$ as f_1 is ramified to order 2 at the marked point q_2 , but the only way this can happen in the limit of a family in Maps_n° is if two marked points collided at q_2 .

Remark 2.3. Note that a degenerate special case of Condition $(*)$ (3) is when $T = q$ is itself a point. Since T must be a maximal closed subvariety contracted by f , this means that q cannot lie on a contracted component. Therefore, by Condition $(*)$ (2), q must be a marked point. Then Condition $(*)$ (3) reads that f must be unramified at q . Indeed, we saw this was necessary in the above Example 2.2 (see Figure 3).

We now prove that conditions (1), (2), and (3) of Conditions $(*)$ are necessary.

Proposition 2.4. *Conditions $(*)$ (1), (2), and (3) are necessary for a stable map to lie in $\overline{\mathcal{M}}_{(d_1, \dots, d_n)}$.*

Proof. It is clear that (1) is necessary as evaluation is continuous so the condition $f(p_i) = \infty$ is closed, so we proceed to (2) and (3). Consider a 1-parameter family of stable maps

$$\begin{array}{ccc} C & \xrightarrow{f} & \mathbb{P}^1 \\ \sigma_i \swarrow \pi & & \\ S & & \end{array}$$

over the spectrum S of a DVR with generic fiber lying in Maps_n° , and denote by $(f_0: C_0 \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ the central fiber.

First we show (2). Let $U \subset C$ denote the open complement of the locus of components contracted by the map f , and consider $D := f^{-1}(\infty)$. As the total space C is normal and f is non-constant, D is a Cartier divisor. Moreover, note that any π -vertical component of D is contracted by f and therefore is not in U . Thus, restricting to U , we see that $D|_U$ is a Cartier divisor which is horizontal over S , and therefore any point of $D|_U$ lying over $0 \in S$ must be in the closure of the marked points of the generic fiber.

Finally, we show (3). Ramification corresponds to a polynomial having a multiple root at a point, and this multiplicity takes into account precisely how many points collided, *i.e.*, the number of marked points on the relevant contracted component. More formally, we can consider the intersection product $T \cdot D$. By the projection formula, $T \cdot D = 0$. On the other hand, $D = \sum \sigma_i + E$, where E is the π -vertical component. Computing $T \cdot (\sum \sigma_i + E) = 0$ in terms of local multiplicities gives exactly the equality in Condition $(*)$ (3). \square

Our task now is to show that Conditions $(*)$ (1)–(3) are sufficient for a stable map $(f: C \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ to lie in $\overline{\mathcal{M}}_{(d_1, \dots, d_n)}$. We will do this by constructing a smoothing of the marked curve and a linear series on the total space which restricts to f on the central fiber but whose generic fiber lies in $\mathcal{M}_{(d_1, \dots, d_n)}$.

We begin with a preliminary lemma that shows that nodal curves can be smoothed to surfaces admitting A_m singularities for any m at the nodes.

Lemma 2.5. *Let C_0 be a genus 0 nodal curve with k nodes q_1, \dots, q_k , and let $m_1, \dots, m_k \geq 1$. Then there exists a smooth $C \rightarrow S$, where $S = \text{Spec}(R)$ is the spectrum of a DVR, such that the total space C has an A_{m_i-1} singularity at q_i for each $i = 1, \dots, k$.*

Proof. The miniversal deformation space of a pointed prestable curve of genus 0 is $k[[t_1, \dots, t_k]]$, where t_i is the smoothing parameter of the i^{th} node; *i.e.*, formally locally around the i^{th} node, the miniversal family of curves looks like $xy = t_i$ inside $\mathbb{A}_{k[[t_1, \dots, t_k]]}^2$. Let $R = [[z]]$ be the DVR in the statement. To prove the lemma, it suffices to note we can construct a map of rings $k[[t_1, \dots, t_k]] \rightarrow k[[z]]$ such that $t_i \mapsto z^{m_i}$. Then formally locally around the i^{th} node, the family of curves over $\text{Spec}(R)$ will be isomorphic to $xy = z^{m_i}$, as required. \square

We now show that it suffices to consider the case of a stable map where every maximal connected subtree contracted by f (as in Condition $(*)$ (3)) is irreducible.

Lemma 2.6. *Let $(f_0: C_0 \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ be a stable map satisfying Conditions $(*)$ (1), (2), and (3). Then there exists a deformation to a family of stable maps $(f: C \rightarrow \mathbb{P}^1, \sigma_i) \rightarrow \text{Spec}(R)$ over the spectrum of a DVR such that*

- (1) *the generic fiber $(f_\eta, C_\eta \rightarrow \mathbb{P}^1, (\sigma_i)_\eta)$ satisfies Conditions $(*)$ (1), (2), and (3), and*
- (2) *every connected component of $f_\eta^{-1}(\infty) \subset C_\eta$ is irreducible.*

Proof. Let E^1, \dots, E^k be the 1-dimensional connected components of $f_0^{-1}(\infty)$, and write C^1, \dots, C^r for the connected components of the closure of the complement $C_0 \setminus \{E^i\}_{i=1}^k$. Each E^i is pointed by (q_{ij}, p_{il}) , where the p_{il} are the marked points that lie on E^i and the q_{ij} are the points of E^i along which E^i is glued to the non-contracted components of C . Then each E_i is a pointed genus 0 prestable curve, and there exists a smoothing $\widetilde{E}^i \rightarrow \text{Spec}(R)$ with sections τ_{ij} smoothing the q_{ij} and σ_{il} smoothing the p_{il} . Now consider the constant family $\bigsqcup_l C_R^l$ with constant sections corresponding to the marked points and the points q'_{ij} along which the q_{ij} are glued.

We can glue $\bigsqcup_l C_R^l$ with $\bigsqcup_t \widetilde{E}^t$ by identifying the sections $q'_{ij} \times \text{Spec}(R)$ with τ_{ij} for all (i, j) and call the result $C \rightarrow \text{Spec}(R)$, which is now a family of pointed curves. By construction, this is a partial smoothing of the pointed curve (C_0, p_1, \dots, p_n) . Moreover, there is a stable map $f: C \rightarrow \mathbb{P}^1$ constructed by taking $f_R|_{C_R^l}: C_R^l \rightarrow \mathbb{P}^1$ for each l and taking the constant map ∞ on each \widetilde{E}^t . This descends to a map f as desired since $f(q'_{ij}) = \infty$. Conditions $(*)$ (1), (2) are satisfied by f_η by construction, and $(*)$ (3) is satisfied since the maximal irreducible components \widetilde{E}_η^i of $f_\eta^{-1}(\infty)$ have the same number of marked points as E^i and f_η has the same ramification at $(q'_{ij})_\eta$ as f_0 at q'_{ij} . \square

We now restrict ourselves to the case where the connected components of $f^{-1}(\infty)$ are irreducible. Let $(f: C_0 \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ be such a map, suppose that E_1, \dots, E_k are the 1-dimensional connected components of $f^{-1}(\infty)$, and write $E = \bigsqcup_i E_i$. By the above reduction, each E_i is a smooth rational curve.

We write the closure of the complement of E in C_0 as a union of connected components C_1, \dots, C_r . Thus each C_i is a tree of rational curves such that $(f_0)|_{C_i}$ is non-constant and such that the preimage of ∞ is 0-dimensional. For each i , let $I_i \subset \{1, \dots, n\}$ be the subset of indices k such that p_k lies on E_i . If E_i and C_j intersect, we let $q_{ij} \in E_i$ and $t_{ij} \in C_j$ be the points of E_i and C_j , respectively, at which they are glued, and we let e_{ij} be the ramification index of $(f_0)|_{C_j}$ at t_{ij} . Note that Condition $(*)$ (3) then reads that for each i ,

$$n_i := \sum_{k \in I_i} d_k = \sum_j e_{ij},$$

where the left side is a definition and the right sum is over the j such that E_i meets C_j .

Consider a smoothing $(C \rightarrow S, \sigma_1, \dots, \sigma_n)$ of (C_0, p_1, \dots, p_n) over the spectrum of a DVR, and let $D = \sum d_i \sigma_i$ denote the divisor of Γ -weighted marked sections on this surface. We wish to construct a rank 1 linear series on such a smoothing so that the central fiber agrees with f_0 and the generic fiber satisfies that $D|_\eta$ is the preimage of ∞ . Our strategy is to consider the map $\pi: C \rightarrow C'$ which contracts E to a point and instead construct the appropriate linear series on C' .

Proposition 2.7. *In the setting above, there exist a smoothing $(C \rightarrow S, \sigma_1, \dots, \sigma_n)$ and a Cartier divisor D' on C' , where $\pi: C \rightarrow C'$ is the contraction of the E_i such that*

- (1) $\pi^*D' = D + \sum a_i E_i$ for some a_i , where D is the divisor of marked sections, and
- (2) $a_i E_i \cdot C_j = e_{ij} t_{ij}$ for all i and j such that E_i meets C_j .

Proof. For each i , we let $a_i = \prod_{j=1}^r e_{ij}$ and let $m_{ij} = \prod_{k \neq j} e_{ik}$. By Lemma 2.6, there exists a smoothing $(C \rightarrow S, \sigma_1, \dots, \sigma_n)$ of (C_0, p_1, \dots, p_n) such that C has an $A_{m_{ij}-1}$ singularity at the node t_{ij} . Computing intersection products in the surface C , we have $E_i \cdot C_0 = 0$ since C_0 is a fiber containing E_i . Moreover, if E_i and C_j intersect, then

$$E_i \cdot C_j = \frac{1}{m_{ij}} t_{ij}$$

since locally around the node t_i , the curves E_i and C_j are distinct lines through an $A_{m_{ij}-1}$ singularity. Therefore,

$$E_i^2 = \sum_j -\frac{1}{m_{ij}},$$

where the sum is over those j for which C_j and E_i meet. Here, we go back and forth between viewing these intersection products as numbers or as divisors on the curves depending on whether it is convenient to emphasize the particular intersection points.

Now consider the divisor $D + \sum a_i E_i$. We can compute that $D \cdot E_i = \sum_{k \in I_i} d_k = n_i$ since D is the divisor of weighted marked sections and E_i contains p_k for $k \in I_i$. On the other hand,

$$a_i E_i \cdot C_j = \frac{a_i}{m_{ij}} = \frac{\prod_j e_{ij}}{\prod_{k \neq j} e_{ik}} = e_{ij}.$$

Finally,

$$\left(D + \sum a_k E_k \right) \cdot E_i = n_i - \sum_j \frac{a_i}{m_{ij}} = n_i - \sum_j \frac{\prod_i e_{ij}}{\prod_{k \neq j} e_{ik}} = n_i - \sum_j e_{ij} = 0$$

by Condition (3). Here we have used that $E_k \cdot E_i = 0$ if $i \neq k$.

Therefore, we need to show that $D + \sum a_i E_i$ descends to a Cartier divisor D' along the contraction $\pi: C \rightarrow C'$. Note that π exists, C' is a normal quasiprojective surface, and $\pi_* \mathcal{O}_C = \mathcal{O}_{C'}$ by [KM98, Theorem 3.7]. Moreover, we have an exact sequence

$$(2.1) \quad 0 \longrightarrow \text{Pic}(C') \xrightarrow{\pi^*} \text{Pic}(C) \longrightarrow \mathbb{Z},$$

where the map $\text{Pic}(C) \rightarrow \mathbb{Z}$ is the restriction $L \mapsto L|_E$; see [KM98, Corollary 3.17]. In particular, since $\mathcal{O}_C(D + \sum a_i E_i)$ is in the kernel of this map, there exists a line bundle L on C' such that $\pi^* L = \mathcal{O}_C(D + aE)$. On the other hand, $\pi_* \mathcal{O}_C = \mathcal{O}_{C'}$, and so by the projection formula,

$$\pi_* \mathcal{O}_C \left(D + \sum a_i E_i \right) = L.$$

Therefore, there exists a section $s \in H^0(L)$ which pulls back to the section cutting out $D + \sum a_i E_i$ on C , and so $D' = \text{div}(s)$ is the required Cartier divisor. \square

Remark 2.8. Fix a list of positive integers r_{ij} for i and j such that E_i meets C_j . Then in the construction of the smoothing in Proposition 2.7, we can replace e_{ij} with $r_{ij} e_{ij}$ in the definitions of a_i and m_{ij} . In this way, we get a smoothing satisfying the properties of Proposition 2.7, where the total space has $A_{m_{ij}-1}$ singularities with m_{ij} divisible by any fixed choice of integers.

Let $0 \in \mathbb{P}^1$ be a general point of the target, and consider the divisor $B_0 \subset C_0$ given by $f_0^{-1}(0)$. Then B_0 consists of a union of points on C_0 which are disjoint from E and contained in the locus where f_0 is étale.

Since $C \rightarrow S$ is a family of genus 0 curves birational to \mathbb{P}_S^1 , we can extend the collection of points B_0 into a Cartier divisor B on C which is horizontal and satisfies $B|_{C_0} = B_0$.

Proposition 2.9. *With notation as above, $D + \sum a_i E_i \sim B$ are linearly equivalent Cartier divisors on C .*

Proof. Let $\psi: C \rightarrow S$ denote the morphism to the base, and consider $\psi_* N$, where $N = \mathcal{O}_C(D + \sum a_i E_i - B)$. First we claim that $H^1(C_0, N|_{C_0}) = 0$. Indeed, since C_0 is a nodal genus 0 curve, it suffices to show that the degree of N restricted to each component of C_0 is at least 0. Now $N|_E = \mathcal{O}_E$ by construction since $(D + \sum a_i E_i) \cdot E = 0$ and B avoids E . On the other hand, for each i , we have $(D + \sum a_i E_i) \cdot C_j = f_0^{-1}(\infty)|_{C_j}$ and $B \cdot C_j = f_0^{-1}(0)|_{C_j}$, so these divisors are linearly equivalent. Therefore, we in fact have that N is trivial on each component of C_0 . Hence, by cohomology and base change, $\psi_* N$ is a vector bundle whose formation commutes with base change. Moreover, over the generic fiber, $C_\eta \cong \mathbb{P}_\eta^1$, and B and $D + \sum a_i E_i$ are divisors of the same degree n by construction. Therefore, $H^0(C_\eta, N|_{C_\eta}) = 1$. Since any line bundle on the spectrum of a DVR is trivial, we conclude that $\psi_* N \cong \mathcal{O}_S$. Consequently, N has a non-vanishing section which exhibits the required linear equivalence $D + \sum a_i E_i \sim B$. \square

Remark 2.10. Note that the vanishing claimed in the above proof does not hold if we replace C_0 with a genus 0 Deligne–Mumford stack. See Remark 5.7 for an example.

Now let B' be the image of B under π , which is also a Cartier divisor since B is contained in the locus where π is an isomorphism. By the exact sequence (2.1), the linear equivalence in Proposition 2.9 is equivalent to a linear equivalence $D' \sim B'$. In particular, D' and B' form a basepoint-free rank 1 linear subseries of $H^0(C', L)$, where $L = \mathcal{O}_{C'}(D')$.

Proposition 2.11. *With notation as above, let $g: C' \rightarrow \mathbb{P}^1$ be the morphism induced by the basepoint-free linear series $\langle B', D' \rangle$. Then the composition $C \xrightarrow{\pi} C' \xrightarrow{g} \mathbb{P}^1$ satisfies $(g \circ \pi)|_{C_0} = f_0$.*

Proof. The map $(g \circ \pi)$ contracts E to ∞ by construction. Moreover, on each C_j , the restrictions of B' and D' satisfy

$$B'|_{C_j} = f_0^{-1}(0)|_{C_j}$$

and

$$D'|_{C_j} = f_0^{-1}(\infty)|_{C_j}$$

by Proposition 2.7. Therefore, $(g \circ \pi)|_{C_j} = (f_0)|_{C_j}$ for all $j = 1, \dots, k$. We conclude that $(g \circ \pi)|_{C_0} = f_0$ as it agrees with f_0 on each component of C_0 . \square

Putting these together, we conclude the following.

Theorem 2.12. *Let $(f_0: C_0 \rightarrow \mathbb{P}^1, p_1, \dots, p_n)$ be a stable map satisfying Conditions (1), (2), and (3). Then there exist a smoothing $(C \rightarrow S, \sigma_1, \dots, \sigma_n)$ over $S = \text{Spec}(R)$, the spectrum of a DVR, and a stable map $f: C \rightarrow \mathbb{P}^1$ such that the generic fiber $(f_\eta, C_\eta \rightarrow \mathbb{P}_\eta^1, (\sigma_i)_\eta)$ is contained in $\mathcal{M}_{(d_1, \dots, d_n)}$ and $f|_{C_0} = f_0$. In particular, Conditions (1), (2), and (3) are both necessary and sufficient for a stable map to be contained in $\overline{\mathcal{M}}_{(d_1, \dots, d_n)}$.*

Proof. By Propositions 2.7 and 2.11, there exist a smoothing $(C \rightarrow S, \sigma_i)$ and a map $f: C \rightarrow \mathbb{P}^1$ such that $f|_{C_0} = f_0$ and $f^{-1}(\infty) = D + \sum a_i E_i$, where D is the divisor of weighted marked sections and the E_i are contained on C_0 . Therefore, $f|_{C_\eta}$ is in $\mathcal{M}_{(d_1, \dots, d_n)}$. \square

3. Log twisted curves and maps

We refer the reader to [ACG⁺13] for the standard definitions in log geometry and to [CJR⁺18, Section 2] for log twisted stable maps. We always work with fine and saturated (fs) log structures. We will use M_X to denote the log structure of X , which will be implicit if no confusion arises. The characteristic of the log structure will be denoted by \overline{M}_X .

3.1. Log geometry

We will need the following results regarding log smooth morphisms.

Remark 3.1.

- If a morphism of log schemes $f: X \rightarrow \text{pt}$ is log smooth (where the point has the trivial log structure), then $U \subset X$ is a toroidal embedding, where U is the locus with trivial log structure and M_X is the divisorial log structure.
- If C is a curve, then log smooth is equivalent to nodal; see [ACG⁺13, Theorem 5].
- Given a family $f: C \rightarrow S$ of nodal curves, there is a *minimal* log structure M_S on S such that any other log structure that makes f log smooth is pulled back from it; see [ACG⁺13, Section 7.3]. We call the corresponding log structure and structure morphism $f^\flat: f^*M_S \rightarrow \tilde{M}_C$ the *canonical* log structure of $f: C \rightarrow S$. Note here that \tilde{M}_C does not include marked points. If f is equipped with marked points, then we denote the natural log structure by

$$M_C := \tilde{M}_C \bigoplus_{\mathcal{O}_C^*} \left(\oplus_{\mathcal{O}_C^*} N_i \right),$$

where the sum is over marked points and N_i is the divisorial log structure associated to the i^{th} marked point; we call M_C the *canonical log structure* associated to a pointed curve if there is no confusion.

There exists a log cotangent complex $\mathbb{L}_{X/Y}^{\log}$ (see [Ols05] or [ACG⁺13, Section 7]) for morphisms of log schemes $f: X \rightarrow Y$, and deformation theory of log schemes is controlled by the log cotangent complex (see [Ols05, Theorem 5.2]).

Remark 3.2. If f is a log smooth morphism, then the log cotangent complex $\mathbb{L}_{X/Y}^{\log}$ is represented by the sheaf of log differentials (see [Ols05, Section 1.1(iii)]). There does not exist a distinguished triangle in general; however, Olsson constructs a distinguished triangle for log flat or integral morphisms (see [Ols05, 1.3]).

3.2. (log) Twisted curves

Our main reference is [Ols07]. To compactify moduli spaces of maps $f: C \rightarrow \mathcal{M}$, where \mathcal{M} is a Deligne–Mumford stack, one needs to allow C to be a stack as well, known as a *twisted curve* (see e.g. [AV02, AB19b]).

Definition 3.3. A *twisted curve* is a purely 1-dimensional Deligne–Mumford stack \mathcal{C} , with at most nodes as singularities, satisfying the following conditions:

- (1) If $\pi: \mathcal{C} \rightarrow C$ denotes the coarse space morphism, then $\mathcal{C}^{\text{sm}} = \pi^{-1}C^{\text{sm}}$, and π is an isomorphism over a dense open subset of C .
- (2) If $\bar{x} \rightarrow C$ is a node such that the strictly henselian local ring $\mathcal{O}_{C, \bar{x}}$ is the strict henselization of $k[x, y]/(xy)$, then

$$\mathcal{C} \times_C \text{Spec}(\mathcal{O}_{C, \bar{x}}) \cong \left[\text{Spec}(\mathcal{O}_{C, \bar{x}}[z, w]/(zw, z^m - x, w^m - y)) / \mu_m \right],$$

where $\xi \in \mu_m$ acts by $(z, w) \mapsto (\xi z, \xi^{-1}w)$.

Definition 3.4. An *n -pointed twisted curve* \mathcal{C}/S marked by disjoint closed substacks $\{\Sigma_i\}_{i=1}^n$ in \mathcal{C} is assumed to satisfy the following:

- (1) The Σ_i are contained in \mathcal{C}^{sm} .
- (2) Each Σ_i is a tame étale gerbe over S .
- (3) The map $\mathcal{C}^{\text{sm}} \setminus \cup \Sigma_i \rightarrow C$ is an open embedding.

Let \mathcal{X} be a Deligne–Mumford stack. We say that a fine log structure $M_{\mathcal{X}}$ is *locally free* if for every geometric point $\bar{x} \rightarrow \mathcal{X}$, the characteristic sheaf satisfies $\overline{M}_{\mathcal{X}} \cong \mathbb{N}^r$ for some r .

Definition 3.5. In the above situation, we say that a morphism of sheaves of monoids $M \rightarrow M'$ is *simple* if for every geometric point $\bar{x} \rightarrow \mathcal{X}$, we have

$$\begin{array}{ccc} \overline{M}_{\mathcal{X}} & \longrightarrow & \overline{M}'_{\mathcal{X}} \\ \downarrow & & \downarrow \\ \mathbb{N}^r & \xrightarrow{\phi} & \mathbb{N}^r, \end{array}$$

where ϕ is given by (m_1, \dots, m_r) .

Definition 3.6. An *n-pointed twisted log curve over S* is the data

$$(C/S, \{\sigma_i, a_i\}, l: M_S \rightarrow M'_S),$$

where

- $(C, \{\sigma_i\})/S$ is an *n*-pointed nodal curve,
- M_S is the minimal log structure for the family $C \rightarrow S$,
- the $a_i: S \rightarrow \mathbb{Z}_{>0}$ are locally constant, and
- l is a simple morphism.

Theorem 3.7 (cf. [Ols07, Theorem 1.8]). *The fibered category of n-pointed twisted curves is naturally equivalent to the stack of n-pointed log twisted curves.*

There is a natural map from the stack of twisted curves to the stack of (pre)stable curves induced by taking the coarse space $\pi: \mathcal{C} \rightarrow C$. The induced map on local deformation spaces $\text{Def}(\mathcal{C}) \rightarrow \text{Def}(C)$ can be described as a root stack of order m_i along the boundary divisor $\{t_i = 0\}$, where t_i is the deformation parameter of the i^{th} -node of C and m_i is the stabilizer order of \mathcal{C} at the i^{th} node. The stabilizer orders m_i correspond via Theorem 3.7 to the data of the simple extension l as in Definition 3.5. For more details, see the discussion following [Ols07, Theorem 1.9 and Remark 1.10].

Example 3.8. Consider a smoothing of a nodal curve as in Lemma 2.5, and let $(C, M_C) \rightarrow (S, M_S)$ be the minimal log structure. The appearance of A_{m_i-1} singularities on the total space of the smoothing at the nodes of C_0 is equivalent to the existence of a simple extension $M_S \hookrightarrow M'_S$ with $\phi = (m_1, \dots, m_k)$. Thus, by Olsson’s Theorem 3.7, such a smoothing is the coarse space of a smoothing of the twisted curve with stabilizer μ_{m_i} at the i^{th} node of C_0 . Note that we can see this directly by taking the canonical stack of the total space C which introduces a stabilizer μ_{m_i} at the A_{m_i-1} singularity.

Let us briefly explicate the construction of Theorem 3.7 as we will use it further on. If $\pi: \mathcal{C} \rightarrow C$ is the coarse map of a twisted curve over S , then the simple extension $l: M_S \hookrightarrow M'_S$ is the map between the minimal log structures of $h: C \rightarrow S$ and $f: \mathcal{C} \rightarrow S$, respectively. The pushforward log structure $\pi_* \tilde{M}_{\mathcal{C}}$ sits in a pushout square

$$\begin{array}{ccc} h^* M_S & \xrightarrow{h^* l} & h^* M'_S \\ h^b \downarrow & & \downarrow \pi_* f^b \\ \tilde{M}_C & \longrightarrow & \pi_* \tilde{M}_{\mathcal{C}}. \end{array}$$

Each a_i keeps track of the stabilizer of \mathcal{C}/C along σ_i , and we have simple extensions $N_{C,i} \hookrightarrow \pi_* N_{\mathcal{C},i}$ corresponding to multiplication by a_i on characteristic monoids.

Definition 3.9. Let $(\mathcal{X}, M_{\mathcal{X}})$ be a separated Deligne–Mumford stack with fs log structure. A *log twisted prestable map* is a diagram

$$\begin{array}{ccc} (\mathcal{C}, M_{\mathcal{C}}) & \xrightarrow{(f, f^b)} & (\mathcal{X}, M_{\mathcal{X}}) \\ \downarrow & & \\ (S, M_S), & & \end{array}$$

where $\mathcal{C} \rightarrow S$ is a twisted curve and $(\mathcal{C}, M_{\mathcal{C}}) \rightarrow (S, M_S)$ is log smooth. A log twisted prestable map is *stable* if the underlying map f is representable and the coarse map $h: C/S \rightarrow X$ is a stable map.

We will also make use of the *root stack* construction of Borne–Vistoli [BV12, Section 4.2] in the case of a locally free log structure. Suppose (X, M_X) is a locally free log scheme and $\phi: \overline{M}_X \rightarrow \overline{M}'$ is a simple extension. The root stack $\sqrt[\ell]{(X, M_X)}$ is a log algebraic stack over Sch_X which associates to $s: T \rightarrow X$ the groupoid of pairs (l, φ) , where $l: s^*M_X \rightarrow M_T'$ is a simple extension of log structures and φ is an isomorphism of maps of characteristic monoids $\bar{l} \cong s^*\phi$.

We conclude this section by comparing the sheaves of log differentials between \mathcal{C} and C . This simple comparison is one of the advantages of using log geometry to study twisted curves. Let $(f, f^b): (\mathcal{C}, M_{\mathcal{C}}) \rightarrow (S, M'_S)$ and $(h, h^b): (C, M_C) \rightarrow (S, M_S)$ denote the natural log smooth maps, where $M_S \hookrightarrow M'_S$ is a simple extension.

Notation 3.10. We set $\Omega_{\mathcal{C}/S}^{\log} := \Omega_{(f, f^b)}^{\log}$ and $\Omega_{C/S}^{\log} := \Omega_{(h, h^b)}^{\log}$ and similarly for the cotangent complex.

Proposition 3.11. *Let $\pi: \mathcal{C} \rightarrow C$ be the coarse space of a twisted curve. Then*

$$\mathbb{L}_{\mathcal{C}/S}^{\log} \simeq_{\mathrm{qis}} \Omega_{\mathcal{C}/S}^{\log}[0] \cong \pi^* \Omega_{C/S}^{\log}[0] \simeq_{\mathrm{qis}} L\pi^* \mathbb{L}_{C/S}^{\log}.$$

Proof. Since both (f, f^b) and (h, h^b) are log smooth, $\mathbb{L}_{\mathcal{C}/S}^{\log} \simeq_{\mathrm{qis}} \Omega_{\mathcal{C}/S}^{\log}$ and $\mathbb{L}_{C/S}^{\log} \simeq_{\mathrm{qis}} \Omega_{C/S}^{\log}$. Since $\Omega_{C/S}^{\log}$ is locally free, $L\pi^* \Omega_{C/S}^{\log}[0] \simeq_{\mathrm{qis}} \pi^* \Omega_{C/S}^{\log}[0]$. Therefore, it suffices to show that the natural map

$$\rho: \pi^* \Omega_{C/S}^{\log} \rightarrow \Omega_{\mathcal{C}/S}^{\log}$$

is an isomorphism which we can compute étale locally. Toward that end, suppose $S = \mathrm{Spec}(R)$. In an étale neighborhood of a marked gerbe with stabilizer μ_a , the map π can be written as $R[u] \mapsto R[x]$ with $u \mapsto x^a$, so $\rho: d\log u \mapsto ad\log x$. In a neighborhood of a node with stabilizer μ_a , we have $R[u, v]/(uv - t^a) \rightarrow R[x, y]/(xy - t)$ with $(u, v) \mapsto (x^a, y^a)$, so ρ is given by $d\log u \mapsto ad\log x$, $d\log v \mapsto ad\log y$. In either case, ρ is an isomorphism since $a \in R$ is invertible. \square

4. Log maps to 1-dimensional targets

In this section, we collect some results on log twisted maps to stacky curves which we will use to lift the smoothings from Section 2 to the twisted case.

4.1. Introducing log structures

We begin by giving a criterion to lift a prestable map with 1-dimensional target to a log map. More precisely, suppose $g: C \rightarrow X$ is a prestable map to a smooth curve X with log structure M_X such that (X, M_X) is log smooth over the trivial log point. We wish to find a log structure M_C on C such that (C, M_C) is a log smooth curve and g lifts to a log map. For the purpose of constructing smoothings, it suffices by Lemma 2.6 to consider the case where the connected components of C which are contracted to the marked points of (X, M_X) are smooth (see also Remark 4.2).

Proposition 4.1. *Let $g: C \rightarrow X$ be a prestable map with (X, M_X) as above. Suppose that the 1-dimensional connected components of the support of $f^*(\overline{M}_X)$ are smooth genus 0 curves. Then g lifts to a prestable logarithmic map. That is, there exist log structures M_C on C and M on $S = \text{Spec}(k)$ and a map $g^\flat: g^*M_X \rightarrow M_C$ such that $(C, M_C) \rightarrow (S, M)$ is log smooth.*

Proof. The question is étale local on the target, so we may pull back to a chart for the log structure (X, M_X) . Hence without loss of generality, we may assume that the target is $(\mathbb{A}^1, 0) = (X, D)$ with its toric log structure and that $g: C \rightarrow \mathbb{A}^1$ is a proper map satisfying the assumptions of the proposition. Moreover, we write $g^{-1}(0) = \{p_1, \dots, p_k, E_1, \dots, E_r\}$ as a disjoint union of points and smooth rational curves. Let us denote by q_{ij} the nodes of C along E_i and by e_{ij} the ramification of g at these points. First we claim that g factors through an expansion of the target

$$C \xrightarrow{h} \widetilde{X} \xrightarrow{\varphi} X.$$

Indeed, consider $\widetilde{X} = E \cup_0 \mathbb{A}^1$, where $E = \mathbb{P}(N_{D/X} + \mathcal{O}_D)$. To construct such a factorization, it suffices to construct non-constant maps $h_i: E_i \rightarrow E$ with $h_i(q_{ij}) = 0$ and such that the order of tangency of h_i at q_{ij} is equal to e_{ij} . Since the E_i are smooth rational curves, such maps exist.

Therefore, we have a factorization $h: C \rightarrow \widetilde{X}$, and it suffices to show that h can be lifted to a log map where

$$M_{\widetilde{X}} = \widetilde{M}_{\widetilde{X}} \oplus_{\mathcal{O}_{\widetilde{X}}^*} \varphi^* M_X$$

since φ underlies a log map $(\widetilde{X}, M_{\widetilde{X}}) \rightarrow (X, M_X)$ and a composition of log maps is a log map. Now we have reduced to the case where the support of $h^*(\overline{M}_{\widetilde{X}}/\mathbb{N})$ is $\{p_1, \dots, p_k, p_{k+1}, \dots, p_n\}$, a union of isolated points. Here the copy of \mathbb{N} in the quotient is the pullback of the minimal log structure on $\text{Spec}(k)$ along $\widetilde{X} \rightarrow \text{Spec}(k)$. In a neighborhood of the smooth p_i , there is a unique divisorial log structure such that the map lifts, encoding the order of tangency. In a neighborhood of the nodal p_i , the map lifts to a log map by the tangency condition above, which is precisely the predeformability condition for a map to an expansion (see for example [ACG⁺13, Section 12]). \square

Remark 4.2. Proposition 4.1 may be generalized to the case where the 1-dimensional connected components of the support of $f^*\overline{M}_X$ are trees of rational curves at the expense of allowing C to be blown up at the nodes. To do this, one needs to consider the map g^{trop} of tropical curves and subdivide the source and target to make g^{trop} tropically transverse (see [Ran22, Sections 2.5 and 2.6] for more details). Such a subdivision corresponds to choosing a possibly larger expansion $\widetilde{X} \rightarrow X$ and a sequence of log blowups $\widetilde{C} \rightarrow C$ after which we may lift g to a map $h: \widetilde{C} \rightarrow \widetilde{X}$ which underlies a log map as in the proposition. Since this is not strictly necessary for our stated goal of constructing smoothings, we leave the details to the reader.

Next we address the question of lifting the log structure to a twisted map from its coarse space. More precisely, consider a proper morphism $f: \mathcal{C}/S \rightarrow \mathcal{X}$ from a twisted curve to a smooth Deligne–Mumford curve \mathcal{X} . Let $M_{\mathcal{X}}$ be a log structure making $(\mathcal{X}, M_{\mathcal{X}})$ a log twisted curve with coarse space $(X, M_X \hookrightarrow M'_X)$. Here (X, M_X) is log smooth, and $M_X \hookrightarrow M'_X$ is the simple extension encoding the stabilizers of \mathcal{X} . Suppose furthermore that there exist log structures M_C and M_S on the coarse space C of \mathcal{C} and on S , respectively, such that

$$\begin{array}{ccc} (C, M_C) & \xrightarrow{g} & (X, M_X) \\ & \downarrow & \\ & & (S, M_S) \end{array}$$

is a prestable log map.

Proposition 4.3. *Let $f: \mathcal{C}/S \rightarrow \mathcal{X}$ and $f: (C, M_C)/(S, M_S) \rightarrow (X, M_X)$ be as above. Then there exist a root stack $\tilde{\mathcal{C}} \rightarrow \mathcal{C}$ and log structures $M_{\tilde{\mathcal{C}}}$ on $\tilde{\mathcal{C}}$ and M'_S on S such that $(\tilde{\mathcal{C}}, M_{\tilde{\mathcal{C}}}) \rightarrow (S, M'_S)$ is a log smooth curve, and there exists a map $f^\flat: (f \circ \pi)^* M_{\mathcal{X}} \rightarrow M_{\tilde{\mathcal{C}}}$ making f into a prestable log map.*

Proof. Let $g: C \rightarrow X$ be the coarse map. By Theorem 3.7, we have simple extensions of log structures $M_C \hookrightarrow M'_C$, $M_S \hookrightarrow M'_S$, and $M_X \hookrightarrow M'_X$ on C , S , and X , respectively, such that $(C \rightarrow S, \{a_i, \sigma_i\}, M_S \hookrightarrow M'_S)$ and $(X \rightarrow \text{Spec}(k), \{b_i, x_i\}, k^* \rightarrow k^*)$ are log twisted curves. Here the $\sigma_i: S \rightarrow C$ correspond to the marked gerbes of $\mathcal{C} \rightarrow S$ with order of stabilizer given by a_i . Similarly, the $x_i \in X$ correspond to the marked gerbes of \mathcal{X} with stabilizer order b_i .

The question is local on X , so we can restrict to a neighborhood of one of the marked points x_i with stabilizer μ_{b_i} on X and suppose without loss of generality that there is a single marked point $x \in X$ with stabilizer μ_b . This point is exactly the support of $\overline{M}_X = \mathbb{N}$ and $\overline{M}'_X = \mathbb{N}$, and the simple extension is simply the map $1 \mapsto b$.

By assumption, there exists a map $g^\flat: g^*M_X \rightarrow M_C$ making (g, g^\flat) into a log map. By Theorem 3.7, it suffices to show that there exist a further simple extension $M'_C \hookrightarrow M''_C$ which is an isomorphism away from finitely many smooth points and a map of log structures $f^\flat: g^*M'_X \rightarrow M'_C$ such that the square

$$\begin{array}{ccc} g^*M_X & \xrightarrow{g^\flat} & M_C \\ \downarrow & & \downarrow s \\ g^*M'_X & \xrightarrow{f^\flat} & M''_C \end{array}$$

commutes. Note that $M'_X = M_X \oplus \mathbb{N}/(e = be')$, where e is the element corresponding to a local parameter of $x \in X$ and e' is the generator of the copy of \mathbb{N} . By the universal property of this pushout, to construct f^\flat , it suffices to show that $s(g^\flat(e))$ is divisible by b . However, we can guarantee this by taking a large enough simple extension s . \square

4.2. Deformations of log twisted maps and their coarse spaces

Let

$$\begin{array}{ccc} (\mathcal{C}, M_{\mathcal{C}}) & \xrightarrow{(f, f^\flat)} & (\mathcal{X}, M_{\mathcal{X}}) \\ \downarrow & & \\ (S, M_S) & & \end{array}$$

be a stable log map from a log twisted curve, and suppose that $(\mathcal{X}, M_{\mathcal{X}})$ is a log smooth curve. We define the *critical locus* of f to be the support of the cokernel of the natural map

$$f^*\Omega_{\mathcal{X}}^{\log} \rightarrow \Omega_{\mathcal{C}/S}^{\log}.$$

Note that this map, viewed as a two-term complex, is a presentation of the log cotangent complex \mathbb{L}_f^{\log} relative to S , so we may equivalently define the critical locus to be the support of $\mathcal{H}^0(\mathbb{L}_f^{\log})$.

The critical locus is a union of nodes of \mathcal{C} , branch points of f , and contracted components of f . By contrast, the kernel of the above map, or equivalently $\mathcal{H}^{-1}(\mathbb{L}_f^{\log})$, is supported solely on the contracted components of f .

Let A_1, \dots, A_k be the connected components of the critical locus and U_1, \dots, U_k be affine étale neighborhoods of the A_i with log structure pulled back from \mathcal{C} . Suppose each U_i avoids A_j for $j \neq i$. Let $\text{Def}_{(f, f^\flat)}$ be the versal deformation space of the log map (f, f^\flat) . Since the moduli space is a Deligne–Mumford stack, $\text{Def}_{(f, f^\flat)}$ is pro-representable. Let Def_i be the miniversal deformation space of the restriction $(f_i := f|_{U_i}: U_i \rightarrow \mathcal{X}, f^\flat|_{U_i})$. We have the following proposition (see also [Vak00, Proposition 4.3]).

Proposition 4.4. *In the setting above, the natural map*

$$\text{Def}_{(f, f^\flat)} \rightarrow \text{Def}_1 \times \dots \times \text{Def}_k$$

is an isomorphism.

Proof. The deformation space $\text{Def}_{(f, f^\flat)}$ (resp. Def_i) is constructed via the complex

$$R\text{Hom}(\mathbb{L}_f^{\log}, \mathcal{O}_{\mathcal{C}}) \text{ (resp. } R\text{Hom}(\mathbb{L}_{f_i}^{\log}, \mathcal{O}_{U_i})).$$

Since the $g_i: U_i \rightarrow \mathcal{C}$ are strict étale, $Lg_i^* \mathbb{L}_f^{\log} = \mathbb{L}_{f_i}^{\log}$. Let K_i be the component of $K = \mathcal{H}^{-1}(\mathbb{L}_f^{\log})$ supported on $A_i \subset U_i$ and Q_i the component of $Q = \mathcal{H}^0(\mathbb{L}_f^{\log})$ supported on $A_i \subset U_i$. Since the A_i are disjoint, we have $K = \bigoplus K_i$ and $Q = \bigoplus Q_i$ and $K_i = \mathcal{H}^{-1}(\mathbb{L}_{f_i}^{\log})$ and $Q_i = \mathcal{H}^0(\mathbb{L}_{f_i}^{\log})$.

Since both \mathbb{L}_f^{\log} and $\mathbb{L}_{f_i}^{\log}$ are two-term complexes, we have distinguished triangles

$$K[1] \rightarrow \mathbb{L}_f^{\log} \rightarrow Q \rightarrow \quad \text{and} \quad K_i[1] \rightarrow \mathbb{L}_{f_i}^{\log} \rightarrow Q_i \rightarrow.$$

There is a natural map $R\text{Hom}(\mathbb{L}_f^{\log}, \mathcal{O}_{\mathcal{C}}) \rightarrow R\text{Hom}(\mathbb{L}_{f_i}^{\log}, \mathcal{O}_{U_i})$ given by restricting f to U_i . This map corresponds to the map on deformation spaces $\text{Def}_{(f, f^\flat)} \rightarrow \text{Def}_i$. Taking direct sums yields the diagram of distinguished triangles

$$(4.1) \quad \begin{array}{ccccccc} R\text{Hom}(Q, \mathcal{O}_{\mathcal{C}}) & \longrightarrow & R\text{Hom}(\mathbb{L}_f^{\log}, \mathcal{O}_{\mathcal{C}}) & \longrightarrow & R\text{Hom}(K[1], \mathcal{O}_{\mathcal{C}}) & \longrightarrow & \\ \parallel & & \downarrow & & \parallel & & \\ \bigoplus R\text{Hom}(Q_i, \mathcal{O}_{U_i}) & \longrightarrow & \bigoplus R\text{Hom}(\mathbb{L}_{f_i}^{\log}, \mathcal{O}_{U_i}) & \longrightarrow & \bigoplus R\text{Hom}(K_i[1], \mathcal{O}_{U_i}) & \longrightarrow & \end{array}$$

where the vertical maps are equalities via the identifications $K = \bigoplus K_i$ and $Q = \bigoplus Q_i$. The middle map corresponds to the product of the restriction map on deformation spaces

$$\text{Def}_{(f, f^\flat)} \rightarrow \text{Def}_1 \times \cdots \times \text{Def}_k.$$

Since the vertical maps on the ends are equalities, the middle map is an isomorphism by the properties of distinguished triangles, or equivalently the five lemma, from which we conclude. \square

Remark 4.5. Via Diagram (4.1), we see that in fact the deformation spaces Def_i of an étale neighborhood U_i of the component A_i of the critical locus is independent of the choice of U_i since the Q_i - and K_i -terms are independent of U_i .

Now we address the question of lifting log smooth deformations along the coarse space map of a log twisted curve. Let

$$\begin{array}{ccc} (C_0, M_{C_0}) & \xrightarrow{h_0} & (X, M_X) \\ \downarrow & & \\ (S_0, M_{S_0}) & & \end{array}$$

be a prestable log map, and suppose that there exists a simple extension of log structures $M_S \rightarrow M'_S$ such that $(C_0/S_0, (\sigma_i, a_i), l_0: M_{S_0} \rightarrow M'_{S_0})$ is a log twisted curve. Let $(\mathcal{C}_0, \Sigma_i) \rightarrow S_0$ be the corresponding twisted curve with log map $(\mathcal{C}_0, M_{\mathcal{C}_0}) \rightarrow (S_0, M'_{S_0})$, and let $g_0: (\mathcal{C}_0, M_{\mathcal{C}_0}) \rightarrow (X, M_X)$ be the composition of h_0 with the coarse space map $\pi_0: \mathcal{C}_0 \rightarrow C$. Let $(S_0, l_0: M_{S_0} \rightarrow M'_{S_0}) \hookrightarrow (S, l: M_S \rightarrow M'_S)$ be a strict closed immersion of log schemes with simple extensions defined by a square 0 ideal $I \subset \mathcal{O}_S$.

Proposition 4.6. *With notation as above, suppose we have a log smooth deformation*

$$(4.2) \quad \begin{array}{ccc} (C, M_C) & \xrightarrow{h} & (X, M_X) \\ \downarrow & & \\ (S, M_S) & & \end{array}$$

of h_0 over (S, M_S) . Then h lifts uniquely to a log smooth deformation

$$\begin{array}{ccc} (\mathcal{C}, M_{\mathcal{C}}) & \xrightarrow{g} & (X, M_X) \\ \downarrow & & \\ (S, M'_S) & & \end{array}$$

of g_0 factoring through h .

Proof. We can summarize the situation with the commutative diagram below:

$$\begin{array}{ccccc} (\mathcal{C}_0, M_{\mathcal{C}_0}) & & & & (X, M_X) \\ \pi_0 \searrow & & \nearrow g_0 & & \\ & (C_0, (\pi_0)_* M_{\mathcal{C}_0}) & \xrightarrow{(\text{id}, s_0)} & (C_0, M_{C_0}) & \xrightarrow{h_0} (X, M_X) \\ \downarrow & & & & \downarrow \\ (S_0, M'_{S_0}) & \xrightarrow{(\text{id}, l_0)} & (S_0, M_{S_0}) & & \end{array}$$

where the middle square is a pullback in the category of fs log schemes by [Ols07, Lemma 4.6, Diagram 4.6.2, and Lemma 4.7]. Given a strict square 0 thickening $(S, l: M_S \rightarrow M'_S)$ of $(S_0, l_0: M_{S_0} \rightarrow M'_{S_0})$ with ideal I and a log smooth deformation h as in Diagram (4.2), we can form the pullback

$$\begin{array}{ccc} (C, M'_C) & \longrightarrow & (C, M_C) \xrightarrow{h} (X, M_X) \\ \downarrow & & \downarrow \\ (S, M'_S) & \longrightarrow & (S, M_S) \end{array}$$

in the category of fs log schemes to obtain a log smooth deformation of $h_0 \circ (\text{id}, s_0)$. Then $M_C \hookrightarrow M'_C$ is a simple extension, and by Theorem 3.7, this is equivalent to a log smooth deformation g . \square

5. Smoothing twisted stable maps

Our goal now is to lift the smoothings constructed in Section 2 to the case of twisted maps. More precisely, suppose we have a genus 0 twisted stable map $(f_0: \mathcal{C}_0 \rightarrow \mathcal{X}, p_1, \dots, p_n)$ such that

- \mathcal{X} is a smooth stacky curve with coarse map $\tau: \mathcal{X} \rightarrow X = \mathbb{P}^1$,
- the coarse map $(h_0: C_0 \rightarrow X, q_1, \dots, q_n)$ satisfies Conditions (*) relative to $\infty \in X$, and
- τ is étale over ∞ .

The situation can be summarized in the following diagram, where $S_0 = \text{Spec}(k)$:

$$\begin{array}{ccc} (\mathcal{C}_0, p_1, \dots, p_n) & \xrightarrow{f_0} & \mathcal{X} \\ \pi_0 \downarrow & & \downarrow \tau \\ (C_0, q_1, \dots, q_n) & \xrightarrow{h_0} & X \\ \downarrow & & \\ S_0. & & \end{array}$$

By Theorem 2.12, there is a smoothing $h: (C, \sigma_i)/S \rightarrow X$ of h_0 over $S = \text{Spec}(R)$, the spectrum of a DVR, with generic fiber contained in $\mathcal{M}_{(d_1, \dots, d_n)}$, and we wish to lift this to a smoothing $f: (\mathcal{C}, \Sigma_i)/S \rightarrow \mathcal{X}$ of f_0

with generic fiber contained in $\mathcal{M}_\Gamma(\mathcal{X})$. That is, we wish to fill in the dotted arrows in the diagram

$$\begin{array}{ccccc}
 (\mathcal{C}_0, p_i) & \dashrightarrow & (\mathcal{C}, \Sigma_i) & \xrightarrow{f} & \mathcal{X} \\
 \pi_0 \downarrow & & \pi \downarrow & & \tau \downarrow \\
 (C_0, q_i) & \longrightarrow & (C, \sigma_i) & \xrightarrow{h} & X \\
 \downarrow & & \downarrow & & \\
 S_0 & \longrightarrow & S. & &
 \end{array}$$

The difficulty is that there are global obstructions to lifting from a map to X to a map to \mathcal{X} . For example, the j -invariant of a map to $\overline{\mathcal{M}}_{1,1}$ must satisfy that the discriminant is a sum of a square and a cube which cuts out a high-codimension locus inside the space of maps to \mathbb{P}^1 . The key observation is that τ is étale. Thus we can lift such a smoothing locally around ∞ (see Theorem 5.1) to obtain a partial smoothing of f_0 . Then we can construct a further global smoothing when \mathcal{X} has some positivity (see Theorem 5.6).

Theorem 5.1. *In the situation above, there exists a partial smoothing $f: (\mathcal{C}, \Sigma_i)/S \rightarrow \mathcal{X}$ of f_0 such that*

- $f(\Sigma_i) = \infty$,
- \mathcal{C}_η is smooth in a neighborhood of $f_\eta^{-1}(\infty)$, and
- the coarse map on the generic fiber h_η is ramified to order d_i at $(\sigma_i)_\eta$.

Proof. We wish to use the log deformation theory results of Section 4.2. To do this, we begin by showing that the smoothing of the coarse map $h: (C, \sigma_i)/S \rightarrow X$ can be endowed with the structure of a stable log map.

Lemma 5.2. *Let $(h_0: C_0 \rightarrow \mathbb{P}^1, q_1, \dots, q_n)$ be as above, and let*

$$\begin{array}{ccc}
 C & \xrightarrow{h} & \mathbb{P}^1 \\
 \sigma_i \swarrow \downarrow & & \\
 S & &
 \end{array}$$

*be a smoothing constructed in Theorem 2.12. Denote by M the divisorial log structure on \mathbb{P}^1 corresponding to the point ∞ . Then there exist log structures M_C and M_S on C and S and a map $h^\flat: h^*M \rightarrow M_C$ such that h is a stable log map over S .*

Proof. First note that $C \rightarrow S$ is log smooth, where we equip $S = \text{Spec}(R)$ with the standard log structure and C with the divisorial log structure induced by the union of the central fiber and marked sections. Indeed, this follows since $C \rightarrow S$ is a toroidal morphism of the corresponding toroidal embeddings. Endow \mathbb{P}^1 with the divisorial log structure for $\infty \subset \mathbb{P}^1$; the pullback of the monomial x_∞ cutting out ∞ is locally a sum of monomials for the toroidal structure on C since $h^*[\infty]$ is supported on the toroidal boundary. Thus, there is a natural map h^\flat of divisorial log structures recording the exponents of the monomial h^*x_∞ . \square

Suppose (R, \mathfrak{m}) is complete, and let $h_n: (C_n, \sigma_{i,n})/S_n \rightarrow X$ be the compatible system of deformations over $S_n = \text{Spec}(R/\mathfrak{m}^n)$ obtained by truncating the smoothing over S . We will lift these to a compatible system of deformations of h_0 in two steps.

First let $(g_0, g_0^\flat): (\mathcal{C}_0, M_{\mathcal{C}_0}) \rightarrow (X, M)$ be the composition $h_0 \circ \pi_0$, where we note that h_0 is a log map by Lemma 5.2 and π_0 is a log map by the definition of log twisted curves. By Lemma 4.6 and induction on n , the map h_n extends to a log smooth deformation $g_n: (\mathcal{C}_n, M_{\mathcal{C}_n}) \rightarrow (X, M_X)$ over (S_n, M'_{S_n}) , where $M_S \hookrightarrow M'_S$ a simple extension and $(S_n, M'_{S_n}) \hookrightarrow (S, M'_S)$ is a strict closed embedding. Moreover, the g_i are compatible by construction. In this way, we get an R -point of the formal deformation space $\text{Spf } R \rightarrow \text{Def}_{(g_0, g_0^\flat)}$.

Next we will use Proposition 4.4 to lift this R -point of $\text{Def}_{(g_0, g_0^\flat)}$, at least in a neighborhood of ∞ , to a smoothing of f_0 . Note that f_0 lifts to a log map (f_0, f_0^\flat) by Proposition 4.3, where we endow \mathcal{X} with the

pullback log structure τ^*M . Equivalently, this is the divisorial log structure for ∞ but is different from the canonical log structure on a log twisted curve. Let $U \subset \mathbb{P}^1$ be an étale neighborhood of ∞ such that $\tau|_V: V \rightarrow U$ is étale, where $V = \tau^{-1}(U)$, and such that U avoids all the components of the critical loci of both f_0 and g_0 away from ∞ . Let $\text{Def}_{f_0, \infty}$ be the miniversal deformation space of the restriction of (f_0, f_0^\flat) to an étale neighborhood of $f_0^{-1}(\infty)$ and similarly for $\text{Def}_{g_0, \infty}$. Note that by the independence of the chosen étale neighborhood (see Remark 4.5), we have that these are the miniversal deformation spaces of $(f_0)|_U: f_0^{-1}(U) \rightarrow U \subset \mathbb{P}^1$ and $(g_0)|_V: g_0^{-1}(V) \rightarrow V \subset \mathcal{X}$. We set $U' = f_0^{-1}(U) = g_0^{-1}(V)$ and denote by U'' a neighborhood of the critical loci of both f_0 and g_0 away from ∞ .

By Proposition 4.4, we have

$$\text{Def}_{(f_0, f_0^\flat)} \cong \text{Def}_{f_0, \infty} \times \text{Def}_{f_0, \neq \infty}$$

and

$$\text{Def}_{(g_0, g_0^\flat)} \cong \text{Def}_{g_0, \infty} \times \text{Def}_{g_0, \neq \infty},$$

where $\text{Def}_{f_0, \neq \infty}$ (resp. $\text{Def}_{g_0, \neq \infty}$) is the miniversal deformation space of a neighborhood of the critical loci of (f_0, f_0^\flat) (resp. (g_0, g_0^\flat)) which are away from ∞ .

Lemma 5.3. *The natural map $\text{Def}_{f_0, \infty} \rightarrow \text{Def}_{g_0, \infty}$ induced by composition with τ is an isomorphism.*

Proof. We proceed as in Proposition 4.4 and see that there is a natural isomorphism

$$R\text{Hom}(\mathbb{L}_{f_0}^{\log}, \mathcal{O}_{\mathcal{C}}) = R\text{Hom}(\mathbb{L}_{f_0'}^{\log}, \mathcal{O}_{U'}) \oplus R\text{Hom}(\mathbb{L}_{f_0''}^{\log}, \mathcal{O}_{U''})$$

and similarly

$$R\text{Hom}(\mathbb{L}_{g_0}^{\log}, \mathcal{O}_{\mathcal{C}}) = R\text{Hom}(\mathbb{L}_{g_0'}^{\log}, \mathcal{O}_{U'}) \oplus R\text{Hom}(\mathbb{L}_{g_0''}^{\log}, \mathcal{O}_{U''}).$$

We now compute

$$\mathbb{L}_{f_0'}^{\log} = [f_0^* \Omega_U \rightarrow \Omega_{U'}^{\log}],$$

$$\mathbb{L}_{g_0'}^{\log} = [g_0^* \Omega_V \rightarrow \Omega_{U'}^{\log}].$$

Since $\tau|_V$ is étale, we have that $g_0^* \Omega_V \cong g_0^* \tau^* \Omega_U = f_0^* \Omega_U$ by the functoriality of pullbacks. Therefore, the natural map

$$\mathbb{L}_{g_0'}^{\log} \rightarrow \mathbb{L}_{f_0'}^{\log}$$

is an isomorphism, so we conclude that the natural map

$$R\text{Hom}(\mathbb{L}_{f_0'}^{\log}, \mathcal{O}_{U'}) \rightarrow R\text{Hom}(\mathbb{L}_{g_0'}^{\log}, \mathcal{O}_{U'})$$

is an isomorphism. □

Now let $\text{Spf } R \rightarrow \text{Def}_{f_0, \infty}$ be the composition of the formal R -point of $\text{Def}_{(g_0, g_0^\flat)}$ constructed above with composition

$$\text{Def}_{(g_0, g_0^\flat)} \rightarrow \text{Def}_{g_0, \infty} \rightarrow \text{Def}_{f_0, \infty},$$

where the first map is the projection from Proposition 4.4 and the second map is the inverse of the isomorphism in Lemma 5.3. There is a canonical splitting of the projection $\text{Def}_{(f_0, f_0^\flat)} \rightarrow \text{Def}_{f_0, \infty}$ via the isomorphism of Proposition 4.4 given by picking the constant deformation in $\text{Def}_{f_0, \neq \infty}$. Further composing with this splitting gives us a formal deformation $\text{Spf } R \rightarrow \text{Def}_{(f_0, f_0^\flat)}$. Since $\text{Def}_{(f_0, f_0^\flat)}$ is the formal neighborhood of a point of a locally of finite type algebraic stack (see [Che14, GS13]), this formal deformation algebraizes to a family $(f, f^\flat): (\mathcal{C}, M_{\mathcal{C}})/(S, M'_S) \rightarrow (\mathcal{X}, \tau^* M)$ of log twisted stable maps (see for example [Bha16, Theorem 1.1] and [BHL17, Corollary 1.5]). Let us denote by $f: (\mathcal{C}, \Sigma_i)/S \rightarrow \mathcal{X}$ the underlying twisted stable map. By construction, the coarse map of f agrees with h over an étale neighborhood of ∞ , so we conclude that f satisfies the required conditions. □

Remark 5.4. By construction, the smoothing of Theorem 5.1 induces the constant infinitesimal deformation of the restriction of f_0 away from ∞ . More precisely, suppose A is a component of the critical locus of f_0 which is disjoint from $f_0^{-1}(\infty)$, and let $U \rightarrow C$ be an étale neighborhood of A which is disjoint from all other critical loci. Then $f|_U \bmod \mathfrak{m}^n$ is the constant infinitesimal deformation of $f_0|_{U \times_C C_0}$.

Remark 5.5. In Theorem 5.1 and the situation above, we can drop the assumption that τ is étale over ∞ at the cost of replacing \mathcal{C}_0 with a root stack. Indeed, if τ is not étale, then τ factors through a non-trivial root stack along ∞ . In this case, one obtains a corresponding simple extension of log structures $M \rightarrow M'$ on \mathbb{P}^1 . Then by Proposition 4.3, the map f_0 can be extended to a log map after taking a further root stack $\mathcal{C}_1 \rightarrow \mathcal{C}_0$. The rest of the argument goes through as written to produce a partial smoothing of the composition $\mathcal{C}_1 \rightarrow \mathcal{C}_0 \rightarrow \mathcal{X}$ with the required properties as in the theorem. We leave the details to the reader since our main interest is in the case of $(\overline{\mathcal{M}}_{1,1}, \infty)$, where τ is étale over ∞ .

5.1. Global smoothings for target weighted projective lines

Next we consider the problem of smoothing twisted stable maps to a weighted projective line $\mathcal{P} = \mathcal{P}(a, b)$ without any tangency conditions.

Theorem 5.6. *Let $(f_0: \mathcal{C}_0 \rightarrow \mathcal{P}, \Sigma_{i,0})$ be a genus 0 twisted stable map to a weighted projective line $\mathcal{P} = \mathcal{P}(a, b)$. Then there exists a smoothing*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{f} & \mathcal{P} \\ \Sigma_i \downarrow & & \\ S & & \end{array}$$

over $S = \text{Spec}(R)$, the spectrum of a DVR.

Proof. Suppose (R, \mathfrak{m}) is complete. We will construct such a smoothing by building a compatible system of deformations over R/\mathfrak{m}^n . Since the stack of not necessarily representable twisted stable maps to \mathcal{P} is a locally of finite type algebraic stack, such a compatible system automatically algebraizes as in the proof of Theorem 5.1. First note that if $a = ka'$ and $b = kb'$ with a', b' coprime, then the canonical map $\rho: \mathcal{P}(a, b) \rightarrow \mathcal{P}(a', b')$ is étale, and so infinitesimal deformations lift uniquely along ρ . Thus, without loss of generality, we may assume that a and b are coprime and \mathcal{P} is a twisted curve.

First we reduce to the case where every connected component in the fibers of $f_0: \mathcal{C}_0 \rightarrow \mathcal{P}$ is irreducible. We can argue as in Lemma 2.6. If E is a connected component of the preimage $f_0^{-1}(p)$ for some p , then we can view $f_0|_E$ as a map to the residual gerbe \mathcal{G}_p where E is pointed by the nodes. Then since \mathcal{G}_p is étale over $\text{Spec}(k)$, it suffices to smooth the composition $E \rightarrow \text{Spec}(k)$ as a marked twisted curve and then glue together the smoothing of E with the constant family of maps $f|_{\overline{\mathcal{C}_0 \setminus E}}$ along the markings corresponding to nodes exactly as in the proof of Lemma 2.6 to obtain a partial smoothing where each connected component of the contracted locus is irreducible. Therefore, without loss of generality, we may suppose that every connected component of the contracted locus of f_0 is irreducible.

By Propositions 4.1 and 4.3, there exists a twisted curve $\tilde{\mathcal{C}}_0$ with a partial coarse space map $\pi_0: \tilde{\mathcal{C}}_0 \rightarrow \mathcal{C}_0$ such that $g_0 := f_0 \circ \pi_0: \tilde{\mathcal{C}}_0 \rightarrow \mathcal{P}$ can be endowed with a log structure where \mathcal{P} has the toric log structure and $(\tilde{\mathcal{C}}_0, M_{\tilde{\mathcal{C}}_0}) \rightarrow (\text{Spec}(k), M)$ is log smooth. Since $(\mathcal{P}, M_{\mathcal{P}})$ is toric, the log cotangent complex $\mathbb{L}_{\mathcal{P}}^{\log} \cong \mathcal{O}_{\mathcal{P}}$ is trivial. Moreover, since $\tilde{\mathcal{C}}_0$ is genus 0,

$$(5.1) \quad \text{Ext}^1(g_0^* \mathbb{L}_{\mathcal{P}}^{\log}, \mathcal{O}_{\tilde{\mathcal{C}}_0}) = H^1(\tilde{\mathcal{C}}_0, \mathcal{O}_{\tilde{\mathcal{C}}_0}) = 0.$$

Therefore, the log deformation space of (g_0, g_0^b) is log smooth over the log deformation space of the curve $(\tilde{\mathcal{C}}_0, M_{\tilde{\mathcal{C}}_0})$. By Theorem 3.7, $\text{Def}(\tilde{\mathcal{C}}_0, M_{\tilde{\mathcal{C}}_0})$ is a root stack over $\text{Def}(\mathcal{C}_0, p_1, \dots, p_n)$, where the p_i are the smooth marked points lying under $\Sigma_{i,0}$. In particular, we can build a smoothing $(C/S, \sigma_i)$ of (\mathcal{C}_0, p_i) whose

truncations $(C_n/\text{Spec}(R/\mathfrak{m}^n), \sigma_{i,n})$ lift to a formal smoothing of $(\tilde{\mathcal{C}}_0, M_{\tilde{\mathcal{C}}_0})$, after possibly extending the log structure on S . By the vanishing obstruction group (5.1), the map (g_0, g_0^b) can then be lifted to a compatible family $(g_n: \tilde{\mathcal{C}}_n \rightarrow \mathcal{P}, g_n^b)$ of log smooth deformations over R/\mathfrak{m}^n .

Thus, after algebraizing, we have a smoothing $(g: \tilde{\mathcal{C}}/S \rightarrow \mathcal{P}, \Sigma_i)$ of g_0 . Taking the relative coarse space of g gives a flat family of maps $f: \mathcal{C}/\text{Spec}(R) \rightarrow \mathcal{P}$ whose formation commutes with base change (see [AOV11, Proposition 3.4] and [AOV08, Corollary 3.3]). Therefore, we conclude that the central fiber of f agrees with f_0 and that the generic fiber is smooth since it is a partial coarse space of the smooth generic fiber of $\tilde{\mathcal{C}}/S$. \square

Remark 5.7. For the proof of Theorem 5.6, it is necessary to use log deformation theory even though the statement does not make any reference to log geometry. The reason is that the stack \mathcal{P} is *not* convex in the classical sense. The tangent bundle is not necessarily effective even though it has positive degree, and there are examples of genus 0 twisted stable maps where the vanishing (5.1) does not hold for the usual tangent bundle. For example, consider $\mathcal{C} = [E/\tau]$, where E is an elliptic curve with hyperelliptic involution $\tau: E \rightarrow E$, $\pi: \mathcal{C} \rightarrow \mathbb{P}^1$ is the coarse space, and $p_i \in \mathbb{P}^1$ are the points lying under the fixed points of τ . Then the line bundle $L = \mathcal{O}_{\mathcal{C}}\left(\frac{1}{2}p_1 - \frac{1}{2}p_2 + \frac{1}{2}p_3 - \frac{1}{2}p_4\right)$ with sections $0 \in L$ and $1 \in L^{\otimes 2}$ induces a map $f: \mathcal{C} \rightarrow \mathcal{P}(1, 2)$ with $f^*T_{\mathcal{P}} = L$ and $h^1(\mathcal{C}, L) = h^1(\mathbb{P}^1, \pi_*L) = h^1(\mathbb{P}^1, \mathcal{O}(-2)) \neq 0$.

6. Proofs of the main theorems

We use the notation from the introduction. Let $(\mathcal{X}, x_1, \dots, x_r)$ be a smooth and proper pointed stacky curve, and suppose that the coarse map $\tau: \mathcal{X} \rightarrow X$ is étale at the points x_i (see also Remark 5.5). Fix a tuple of discrete data $\Gamma = (n_0, \{\Gamma_1, x_1\}, \dots, \{\Gamma_r, x_r\})$.

Proposition 6.1. *The loci $\mathcal{M}_{\Gamma}(\mathcal{X})$ and $\mathcal{N}_{\Gamma}(\mathcal{X})$ (see Definition 1.3) are locally closed substacks of $\mathcal{K}_{0,n}(\mathcal{X}, d)$.*

Proof. We run through the conditions of Definition 1.3. Condition (1) is both open and closed by the definition of a twisted curve, and condition (2) is closed. Conditions (4) and (4') are open by the openness of the smooth locus. For condition (3), we consider the cohomology sheaves of the cotangent complex \mathbb{L}_f^{\log} . The locus where f is constant is given by the support of $\mathcal{H}^{-1}(\mathbb{L}_f^{\log})$. Since the universal family of curves over the moduli space is proper, the locus where the marked points avoid $\text{Supp}\mathcal{H}^{-1}(\mathbb{L}_f^{\log})$ is open, and therefore so is the condition of f being non-constant along the marked points. Finally, under this condition, the order of tangency of f at x_i is measured by the length of $\text{coker}(f^*\Omega_{\mathcal{X}}^1 \rightarrow \Omega_{\mathcal{C}}^1)_{x_i}$, and by semicontinuity, there is a locally closed subset where this length is constant d_i . \square

Proof of Theorem 1.5. The marked points p_{0k} for $k = 1, \dots, n_0$ with no tangency conditions do not change the outcome of the problem. Indeed, if we can construct such a smoothing with $n_0 > 0$, then by forgetting and stabilizing, we obtain a smoothing with $n_0 = 0$, and, conversely, if we have a partial 1-parameter smoothing over a base S with $n_0 = 0$, then up to a finite base change $S' \rightarrow S$, we can pick generic sections passing through p_{0k} to obtain a smoothing with $n_0 > 0$. So without loss of generality, suppose $n_0 = 0$.

We proceed by induction on r . The base case is precisely Theorem 5.1. In general, let $(f: \mathcal{C} \rightarrow \mathcal{X}, \{p_{jk}\}_{k=1}^{n_j} \}_{j=1}^r)$ be a map in $\mathcal{K}_{\Gamma}(\mathcal{X})$. Applying Theorem 5.1 to $(f: \mathcal{C} \rightarrow \mathcal{X}, p_{r1}, \dots, p_{rn_r})$ produces a 1-parameter deformation whose generic fiber is a relative map to (\mathcal{X}, x_r) with tangency Γ_r which also satisfies that the curve is smooth along the preimage of x_r . Moreover, by Remark 5.4, the generic fiber is contained in $\mathcal{K}_{\Gamma}(\mathcal{X})$. Therefore, without loss of generality, we may assume that \mathcal{C} is smooth in a neighborhood of $f^{-1}(x_r)$. By the inductive hypothesis, there exists a 1-parameter deformation of f with generic fiber contained in $\mathcal{N}_{\Gamma'}(\mathcal{X})$, where $\Gamma' = (\{\Gamma_1, x_1\}, \dots, \{\Gamma_{r-1}, x_{r-1}\})$. Moreover, by Remark 5.4, this partial smoothing induces the constant deformation in an étale neighborhood of $f^{-1}(x_r)$, and so the generic fiber is contained in $\mathcal{N}_{\Gamma}(\mathcal{X})$. \square

Proof of Theorem 1.7. Let $(f: \mathcal{C} \rightarrow \mathcal{X}, p_{jk})$ be a map contained in $\mathcal{K}_\Gamma(\mathcal{X})$. By Theorem 1.5, there exists a partial smoothing of $(f: \mathcal{C} \rightarrow \mathcal{X}, p_{jk})$ with generic fiber contained in $\mathcal{N}_\Gamma(\mathcal{X})$, so without loss of generality, suppose that $(f: \mathcal{C} \rightarrow \mathcal{X}, p_{jk})$ is contained in $\mathcal{N}_\Gamma(\mathcal{X})$. By Theorem 5.6, there exists a smoothing of the stable map into the interior of $\mathcal{K}_{0,n}(\mathcal{X}, d)$ since $\mathcal{X} = \mathcal{P}$ is a weighted projective line. Let $\text{Spf}R \rightarrow \text{Def}_f$ be the corresponding formal deformation. By Proposition 4.4, we can write

$$\text{Def}_f = \text{Def}_{f,x_1} \times \cdots \times \text{Def}_{f,x_r} \times \text{Def}_{f,\neq},$$

where Def_{f,x_i} is a miniversal deformation space of a small étale neighborhood of the fiber $f^{-1}(x_i)$ and $\text{Def}_{f,\neq}$ is a miniversal deformation space of the critical loci of f which are disjoint from the fibers $f^{-1}(x_i)$ for each i . Then projecting onto $\text{Def}_{f,\neq}$ and then composing with the section $\text{Def}_{f,\neq} \rightarrow \text{Def}_f$ which picks the constant deformation on each of the Def_{f,x_i} factors produces a new formal deformation $\text{Spf}R \rightarrow \text{Def}_f$ which agrees with the smoothing from Theorem 5.6 away from the fibers $f^{-1}(x_i)$ but is the constant deformation in a neighborhood of $f^{-1}(x_i)$ for all i . After algebraizing this formal deformation as in the proof of Theorem 5.1, we obtain a 1-parameter family over $\text{Spec}(R)$ such that the generic fiber is contained in $\mathcal{N}_\Gamma(\mathcal{X})$ but is also smooth away from the union of the fibers $f^{-1}(x_i)$. Therefore, the generic fiber is smooth everywhere and thus contained in $\mathcal{M}_\Gamma(\mathcal{X})$. \square

References

- [ACG⁺13] D. Abramovich, Q. Chen, D. Gillam, Y. Huang, M. Olsson, M. Satriano, and S. Sun, *Logarithmic geometry and moduli*, in: *Handbook of moduli. Vol. I*, pp. 1–61, Adv. Lect. Math. (ALM), vol. 24, Int. Press, Somerville, MA, 2013.
- [AOV08] D. Abramovich, M. Olsson, and A. Vistoli, *Tame stacks in positive characteristic*, Ann. Inst. Fourier (Grenoble) **58** (2008), no. 4, 1057–1091.
- [AOV11] ———, *Twisted stable maps to tame Artin stacks*, J. Algebraic Geom. **20** (2011), no. 3, 399–477.
- [AV00] D. Abramovich and A. Vistoli, *Complete moduli for fibered surfaces*, in: *Recent progress in intersection theory* (Bologna, 1997), pp. 1–31, Trends Math., Birkhäuser Boston, Boston, MA, 2000.
- [AV02] ———, *Compactifying the space of stable maps*, J. Amer. Math. Soc. **15** (2002), no. 1, 27–75.
- [AB19a] K. Ascher and D. Bejleri, *Compact moduli of elliptic K3 surfaces*, preprint [arXiv:1902.10686](https://arxiv.org/abs/1902.10686) (2019). To appear in Geometry & Topology.
- [AB19b] ———, *Moduli of fibered surface pairs from twisted stable maps*, Math. Ann. **374** (2019), no. 1–2, 1007–1032.
- [AB21] ———, *Moduli of weighted stable elliptic surfaces and invariance of log plurigenera*, Proc. Lond. Math. Soc **122** (2021), no. 5, 617–677.
- [Bhal16] B. Bhatt, *Algebraization and Tannaka duality*, Camb. J. Math. **4** (2016), no. 4, 403–461.
- [BHL17] B. Bhatt and D. Halpern-Leistner, *Tannaka duality revisited*, Adv. Math. **316** (2017), 576–612.
- [BV12] N. Borne and A. Vistoli, *Parabolic sheaves on logarithmic schemes*, Adv. Math. **231** (2012), no. 3–4, 1327–1363.
- [Che14] Q. Chen, *Stable logarithmic maps to Deligne-Faltings pairs I*, Ann. of Math. (2) **180** (2014), no. 2, 455–521.
- [CJR⁺18] Q. Chen, F. Janda, Y. Ruan, and A. Sauvaget, *Towards a Theory of Logarithmic GLSM Moduli Spaces*, preprint [arXiv:1805.02304](https://arxiv.org/abs/1805.02304) (2018). To appear in Geometry & Topology.

- [Gat02] A. Gathmann, *Absolute and relative Gromov-Witten invariants of very ample hypersurfaces*, Duke Math. J. **115** (2002), no. 2, 171–203.
- [GS13] M. Gross and B. Siebert, *Logarithmic Gromov-Witten invariants*, J. Amer. Math. Soc. **26** (2013), no. 2, 451–510.
- [KM98] J. Kollar and S. Mori, *Birational geometry of algebraic varieties*, Translated from the 1998 Japanese original, With the collaboration of C. H. Clemens and A. Corti, Cambridge Tracts in Math., vol. 134, Cambridge Univ. Press, Cambridge, 1998.
- [Ols05] M. C. Olsson, *The logarithmic cotangent complex*, Math. Ann. **333** (2005), no. 4, 859–931.
- [Ols07] ———, *(Log) twisted curves*, Compos. Math. **143** (2007), no. 2, 476–494.
- [Ran22] D. Ranganathan, *Logarithmic Gromov-Witten theory with expansions*, Algebr. Geom. **9** (2022), no. 6, 714–761.
- [Vak00] R. Vakil, *The enumerative geometry of rational and elliptic curves in projective space*, J. reine angew. Math. **529** (2000), 101–153.