

Three approaches for the classification of protoneutron star oscillation modes

M. C. Rodriguez,^{1,2} Ignacio F. Ranea-Sandoval,^{1,2} C. Chirenti^{3,4,5,6} and D. Radice^{7,8,9†}

¹Grupo de Gravitación, Astrofísica y Cosmología, Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque S/N, 1900 La Plata, Argentina

²CONICET, Godoy Cruz 2290, 1425 CABA, Argentina

³Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

⁴Astroparticle Physics Laboratory NASA/GSFC, Greenbelt, MD 20771, USA

⁵Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771, USA

⁶Center for Mathematics, Computation, and Cognition, UFABC, Santo André, SP 09210-170, Brazil

⁷Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

⁸Department of Physics, The Pennsylvania State University, University Park, PA 16802, USA

⁹Department of Astronomy & Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

Accepted 2023 May 10. Received 2023 May 2; in original form 2023 March 31

ABSTRACT

The future detection of gravitational waves (GWs) from a Galactic core-collapse supernova will provide information on the physics inside protoneutron stars (PNS). In this work, we apply three different classification methods for the PNS non-radial oscillation modes: Cowling classification, Generalized Cowling Nomenclature (GCN), and a classification based on modal properties (CBMP). Using PNS models from 3D simulations of core-collapse supernovae, we find that in the early stages of the PNS evolution, typically 0.4 s after the bounce, the Cowling classification is inconsistent, but the GCN and the CBMP provide complementary information that helps to understand the evolution of the modes. In the GCN, we note several avoided crossings as the mode frequencies evolve at early times, while the CBMP tracks the modes across the avoided crossings. We verify that the strongest emission of GWs by the PNS corresponds to the f mode in the GCN, indicating that the mode trapping region alternates between the core and the envelope at each avoided crossing. At later times, approximately 0.4 s after the bounce, the three classification methods present a similar description of the mode spectrum. We use our results to test universal relations for the PNS modes according to their classification and find that the behaviour of the universal relations for f and p modes is remarkably simple in the CBMP.

Key words: asteroseismology – gravitational waves – stars: neutron – stars: oscillations.

1 INTRODUCTION

In recent years, the first detection of gravitational waves (GWs) emitted by the merger of binary systems of black holes (BHs; Abbott et al. 2016) and neutron stars (NSs; Abbott et al. 2017b) has given rise to a new field of research that could offer information about the physics of these compact objects. The double NS coalescence GW170817 has also been detected through the entire electromagnetic spectrum and thus added GWs to the landscape of multimessenger astronomy. These events, together with the astronomical properties of the ~ 100 mergers of compact binaries detected so far (Abbott et al. 2021), help to constrain the properties of BHs and NSs. The GWTC-2 and GWTC-3 catalogues contain the properties inferred from the detection of GWs (Abbott et al. 2021a, b).

A complementary view of the physics of compact objects will emerge upon the detection of GWs from a Galactic core-collapse supernova. One of the possible remnants of a core-collapse supernova is a protoneutron star (PNS; see e.g. Burrows & Vartanyan 2021, and references therein). The resulting GW emission is associated with PNS oscillations and dynamics (Marek, Janka & Müller 2009; Murphy, Ott & Burrows 2009; Müller, Janka & Marek 2013). Unlike NS binary coalescences, for which the stars are effectively cold, inside a PNS temperatures are initially above 5×10^{11} K. Over a time-scale of a few seconds, PNSs become more compact and cool down until they reach an equilibrium in which they are considered NSs (Pons et al. 1999). During this time, the microphysics inside the star also changes due to processes such as electron capture and, later, neutrino emission. As a PNS cools, its central densities rapidly become higher than nuclear saturation density. At such densities, the equation of state (EoS), which gives us information about the relation between the pressure and the energy density inside the star, is still largely unknown.

The study of the GW signal from simulations of core-collapse supernovae has been performed in two dimensions with axial

* E-mail: mcrodriguez@fcaglp.unlp.edu.ar (MCR); iranea@fcaglp.unlp.edu.ar (IFRS); cecilia.chirenti@ufabc.edu.br (CC)

† Alfred P. Sloan fellow.

symmetry, e.g. Marek et al. (2009), Murphy et al. (2009), Cerdá-Durán et al. (2013), Müller et al. (2013), Abdikamalov et al. (2014), Yakunin et al. (2015), Morozova et al. (2018), and Pan et al. (2018), among others, and, in the last years, in three dimensions (see e.g. Hayama et al. 2016; Kuroda, Kotake & Takiwaki 2016; Andresen et al. 2017; Kuroda et al. 2017; Hayama et al. 2018; O'Connor & Couch 2018; Powell & Müller 2019, 2022; Mezzacappa et al. 2020b, 2023; Nakamura, Takiwaki & Kotake 2022; Vartanyan, Coleman & Burrows 2022; Vartanyan et al. 2023), in which the treatment is much more precise and thus computationally more expensive than in two dimensions.

The complexity of the processes during core collapse renders it impossible to predict with high precision the resulting gravitational waveforms. None the less, the community has reached some agreement on the key aspects of the signal morphology after several years of numerical simulations (O'Connor et al. 2018; Mezzacappa et al. 2020a). Therefore, a future detection of GWs from Galactic core-collapse supernovae will add more information about the physics inside the PNS and will facilitate tests of the simulated models (Abdikamalov, Pagliaroli & Radice 2022; Wolfe et al. 2023).

However, detecting the emitted GWs produced in core-collapse supernovae will be challenging. Knowledge of the morphology of the emitted waveform is of paramount importance for developing effective algorithms capable of extracting GW data from the detected signal. Different methods have been proposed: examples include principal component analysis (Heng 2009; Röver et al. 2009; Powell, Szczepanczyk & Heng 2017; Roma et al. 2019), inference using Bayesian analysis (Summerscales et al. 2008; Röver et al. 2009; Powell et al. 2017; Gill et al. 2018; Roma et al. 2019), denoising techniques (Mukherjee et al. 2017), and machine learning (Astone et al. 2018; Cavaglià et al. 2020; Chan, Heng & Messenger 2020; Iess et al. 2020; López et al. 2021). One of the most promising ways to detect GWs from nearby core-collapse supernovae uses the excess-power coherent waveburst search, which has already detected GWs produced in mergers of compact objects (Abbott et al. 2019; Abbott et al. 2021a, b), even in low-latency [e.g. for GW150914 (Abbott et al. 2017a)].

Here, however, we focus on a different approach, using a theoretical framework which has been developed to study stellar perturbations. We use this framework to describe the GW signal of a supernova as the superposition of the oscillation modes of the PNS (Marek et al. 2009; Murphy et al. 2009; Müller et al. 2013). Oscillation modes of stellar configurations can be classified by taking into account their main restoring force. Generally speaking, the dominant restoring force is pressure for p modes (and f modes) and buoyancy for g modes (Cowling 1941; Kokkotas & Schmidt 1999). In the standard picture, two local quantities determine the character of a mode with given frequency $\sigma/(2\pi)$: the Lamb frequency, \mathcal{L} , and the Brünt–Väisälä frequency, \mathcal{N} . Sound waves can propagate in regions of the star in which σ^2 is greater than both \mathcal{L}^2 and \mathcal{N}^2 , while gravity waves are possible in regions of the star in which σ^2 is smaller than both \mathcal{L}^2 and \mathcal{N}^2 (see e.g. Unno et al. 1989, for a review).

Usually, the classification of non-radial oscillation modes is based on the classification scheme proposed by Cowling (1941), in which the modes are defined by their restoring force and the number of nodes in the radial eigenfunction. But, for stars in which the stellar dynamics is different from main-sequence stars (or cold NSs), such as evolved stars and newly born PNS, the classification is not so trivial and the Cowling classification is not enough to correctly identify the non-radial oscillation modes. For this reason, in Section 3 we

explore alternative classification methods of PNS modes, which were developed in the context of the study of non-radial pulsations of evolved stars and give complementary information about the modes, enabling a proper classification in this setting.

Following Unno et al. (1989), we use the Generalized Cowling Nomenclature (GCN; Eckart 1960; Scuflaire 1974; Osaki 1975) and the classification based on modal properties (CBMP) of Shibahashi & Osaki (1976). The GCN was also applied to PNS simulations by Torres-Forné et al. (2019b) and other subsequent works. The CBMP provides qualitatively similar results to the matching classification based on the shape of the radial eigenfunction proposed by Torres-Forné et al. (2019b) and to the ramp-up mode description discussed by Mori, Suwa & Takiwaki (2023), but the CBMP is based on a more rigorous evaluation of the kinetic energy of the mode in its trapped region. These complementary classification schemes help us to understand and identify different features of the modes, especially in the early stages of the PNS after the bounce.

The aim of our work is to be able to identify correctly the non-radial oscillations modes of PNSs. By using complementary classification schemes, we are able to capture information from GW signals that can help to infer properties of the pulsating PNS. To obtain the modes, we apply linear perturbation theory to the PNS and use a Lagrangian description of the fluid perturbation, following Morozova et al. (2018).

In Section 2, we present the set-up for calculating the non-radial oscillation modes from PNS described by data from core-collapse supernova simulations (Radice et al. 2019). In Section 3, we describe the three different mode classification schemes. In Section 4, we present the results obtained using the three different classification methods and a brief analysis of the universal relations of PNSs. Finally, in Section 5 we summarize our work and present our conclusions.

2 PROTONEUTRON STAR NON-RADIAL OSCILLATIONS

We use hydrostatic PNS configurations obtained from 3D core-collapse supernovae simulations of Radice et al. (2019) following the approach of Sotani & Takiwaki (2016). Namely, the results of the 3D simulations are averaged to construct an effective barotropic EoS, which we use to solve the Tolman–Oppenheimer–Volkoff equations to reconstruct the structure of the PNS and the space–time geometry. See also Westernacher-Schneider (2020) for an alternative approach that is fully consistent with the pseudo-Newtonian formalism employed in the dynamical simulations. The line element that describes a static spherically symmetric conformally flat space–time in isotropic coordinates (t, x_i) is given by

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = -\alpha^2 dt^2 + \psi^4 f_{ij}dx^i dx^j, \quad (1)$$

where α is the lapse function, ψ is the conformal factor whose value is set to 1, and f_{ij} is the flat spatial 3-metric. The conformal flatness approximation incorporates the space–time dynamics evolution, but it ignores gravitational radiation (Wilson & Mathews 1989; Isenberg 2008).

To describe the fluid perturbations we used the Lagrangian displacement vector

$$\xi^i = (\xi^r, \xi^\theta, \xi^\phi), \quad (2)$$

with

$$\begin{aligned}\xi^r &= \eta_r Y_{lm} e^{-i\sigma t}, \\ \xi^\theta &= \eta_\perp \frac{1}{r^2} \partial_\theta Y_{lm} e^{-i\sigma t}, \\ \xi^\phi &= \eta_\perp \frac{1}{r^2 \sin^2 \theta} \partial_\phi Y_{lm} e^{-i\sigma t},\end{aligned}\quad (3)$$

where Y_{lm} are the spherical harmonics and the functions η_r and η_\perp depend only on the radial coordinate r . For each of the perturbing functions, a time dependence $e^{i\sigma t}$ is assumed.

In order to find the quasi-normal modes, we need to solve a set of four first-order coupled differential equations for η_r , η_\perp and two additional functions f_α and $\delta\hat{\alpha}$, with appropriate values at the centre and a boundary condition, as done in Morozova et al. (2018). The scalar function $\delta\hat{\alpha}$ is the amplitude of the lapse function perturbation that depends only on the radial coordinate, i.e. $\delta\alpha = \delta\hat{\alpha}(r)Y_{lm} e^{-i\sigma t}$ and f_α is defined as $f_\alpha = \partial_r(\delta\hat{\alpha}/\alpha)$. The equations for the four unknown functions η_r , η_\perp , f_α , and $\delta\hat{\alpha}$ are

$$0 = \partial_r \eta_r + \left[\frac{2}{r} + \frac{1}{\Gamma_1} \frac{\partial_r P}{P} + 6 \frac{\partial_r \psi}{\psi} \right] \eta_r + \frac{\psi^4}{\alpha^2 c_s^2} (\sigma^2 - \mathcal{L}^2) \eta_\perp - \frac{1}{\alpha c_s^2} \delta\hat{\alpha}, \quad (4)$$

$$0 = \partial_r \eta_\perp - \left(1 - \frac{\mathcal{N}^2}{\sigma^2} \right) \eta_r + \left[\partial_r \ln q - \tilde{G} \left(1 + \frac{1}{c_s^2} \right) \right] \eta_\perp - \frac{1}{\alpha \tilde{G}} \frac{\mathcal{N}^2}{\sigma^2} \delta\hat{\alpha}, \quad (5)$$

$$0 = \partial_r f_\alpha + \frac{2}{r} f_\alpha + 4\pi \left[\partial_r \rho - \frac{\rho}{P \Gamma_1} \partial_r P \right] \eta_r - \frac{4\pi \rho}{P \Gamma_1} q \sigma^2 \eta_\perp + \left[\frac{4\pi \rho^2 h}{P \Gamma_1 \alpha} - \frac{1}{\alpha} \frac{l(l+1)}{r^2} \right] \delta\hat{\alpha}, \quad (6)$$

$$0 = \partial_r \delta\hat{\alpha} - f_\alpha \alpha + \tilde{G} \delta\hat{\alpha}, \quad (7)$$

where Γ_1 is the adiabatic index, P is the pressure, ρ is the mass density, h is the specific enthalpy, c_s is the speed of sound and $q = \rho h \alpha^{-2} \psi^4$; all of these quantities refer to the interior of the PNS. The Brunt–Väisälä and Lamb frequencies are

$$\mathcal{N}^2 = \frac{\alpha \partial_r \alpha}{\psi^4} \left(\frac{1}{\Gamma_1} \frac{\partial_r P}{P} - \frac{\partial_r e}{\rho h} \right), \quad (8)$$

$$\mathcal{L}^2 = \frac{\alpha^2}{\psi^4} c_s^2 \frac{l(l+1)}{r^2}. \quad (9)$$

where e is the energy density. The radial component of the gravitational acceleration \tilde{G} is defined as $\tilde{G} = \frac{1}{\rho h} \partial_r P$.¹

To solve equations (4)–(7), the boundary condition $\Delta P = 0$ on the Lagrangian perturbation of the pressure has to be satisfied at the stellar radius, R . Following Morozova et al. (2018), we define the stellar radius where $\rho = 10^{10} \text{ g cm}^{-3}$. Different values of the mass density for the stellar radius do not have a large impact on the mode frequencies; g modes are essentially unchanged. The boundary condition is given by

$$q \sigma^2 \eta_\perp - \frac{\rho h}{\alpha} \delta\hat{\alpha} + \partial_r P \eta_r = 0. \quad (10)$$

From equations (4)–(7), regularity of the solutions at $r = 0$ requires

$$\eta_r|_{r=0} = f_\alpha|_{r=0} \sim r^{l-1}, \quad (11)$$

¹An alternative definition is given by $\tilde{G} = -\partial_r \ln \alpha$. Both definitions of \tilde{G} were tested in our code giving very similar results as expected.

Table 1. 3D simulations of core-collapse supernovae from Radice et al. (2019) corresponding to the PNS evolution models used in this work. Models 3 and 7 ignore the virial correction to the neutrino–nucleon scattering from Horowitz et al. (2017), which is used in all other models.

Model	ZAMS mass (M_\odot)	Explode	Symbol
1	9	Yes	▲
2	10	Yes	●
3	11	Yes	○
4	11	Yes	■
5	12	Yes	□
6	13	No	*
7	19	Yes	×
8	19	Yes	+
9	25	Yes	▽
10	60	Yes	△

$$\delta\hat{\alpha}|_{r=0} = \alpha(0) \eta_\perp|_{r=0} \sim \frac{\alpha(0)}{l} r^l. \quad (12)$$

We note that Morozova et al. (2018) set the boundary conditions for $\delta\hat{\alpha}$ and f_α to zero. However, the difference in the resulting mode frequencies is negligible.

We computed the oscillation frequencies and their respective eigenfunctions in FORTRAN90. Our code uses a shooting method with Ridders' root-finding algorithm to look for eigenfrequencies in the range from 100 to 2000 Hz until the boundary condition is satisfied, solving equations (4)–(7) (or equations 13 and 14 below, in the Cowling approximation) with a Runge–Kutta–Fehlberg integrator. This is done at every time-step of the evolution of the PNS, as described in Section 2.1.

2.1 Data

The profiles of evolving PNS were calculated from the 3D simulations of core-collapse supernovae performed by Radice et al. (2019). The simulations were performed using the FORNAX code (Skinner et al. 2019), which solves the equations of neutrino radiation hydrodynamics using a sophisticated, multidimensional transport scheme. Gravity was handled using an effective general relativistic potential following Marek et al. (2009). The neutrino treatment accounted for gravitational redshift, Doppler effects, and inelastic scattering (Burrows et al. 2018). NS matter was treated using the SFHo EoS of Steiner, Hempel & Fischer (2013). In this work, we use models with $9 M_\odot$, $10 M_\odot$, $11 M_\odot$, $12 M_\odot$, $13 M_\odot$, $19 M_\odot$, $25 M_\odot$, and $60 M_\odot$ zero-age main sequence (ZAMS) mass. In Table 1, we list the details of each model. For more details, see Radice et al. (2019) and Burrows et al. (2020).

2.2 The relativistic Cowling approximation

As the classification methods described in Section 3 below are strictly valid only when the metric perturbations are neglected, we need to use the relativistic Cowling approximation (McDermott, van Horn & Scholl 1983), in which only fluid perturbations are considered. Under this approximation, the set of differential equations (4)–(7) becomes

$$0 = \partial_r \eta_r + \left[\frac{2}{r} + \frac{1}{\Gamma_1} \frac{\partial_r P}{P} \right] \eta_r + \frac{1}{\alpha^2 c_s^2} (\sigma^2 - \mathcal{L}^2) \eta_\perp, \quad (13)$$




Figure 1. Propagation diagram for a PNS at 0.4 s after the bounce, produced by the model with $9 M_{\odot}$ progenitor mass (model 1 in Table 1), for $l = 2$. The region above $\log_{10}(N^2)$ and $\log_{10}(L^2)$ is the p-propagation zone and the region below $\log_{10}(N^2)$ and $\log_{10}(L^2)$ is the g-propagation zone. Both N and L are normalized by $\sqrt{GM/R^3}$. The horizontal lines show the frequencies of three different modes; diamonds indicate the position of the nodes in the radial eigenfunction. Filled (empty) diamonds correspond to p-(g-)nodes, see Section 3.2.

$$0 = \partial_r \eta_{\perp} - \left(1 - \frac{N^2}{\sigma^2}\right) \eta_r + \left[\partial_r \ln q - \tilde{G} \left(1 + \frac{1}{c_s^2}\right) \right] \eta_{\perp}. \quad (14)$$

In the relativistic Cowling approximation, these are the only equations needed to find the mode frequency, σ , whose associated eigenfunctions are η_r and η_{\perp} .

Boundary conditions need to be imposed to solve equations (13) and (14). Under this approximation, the values of σ^2 we are looking for will be the ones that satisfy

$$q\sigma^2 \eta_{\perp} + \partial_r P \eta_r = 0. \quad (15)$$

3 CLASSIFICATION OF MODES

The analysis of the stellar oscillations gives information about the properties of the modes, which can be inferred in a propagation diagram like the one shown in Fig. 1. This propagation diagram shows, for a PNS model, the normalized Brunt–Väisälä and the Lamb frequencies as functions of the normalized radius of the star, and corresponds to perturbation modes with $l = 2$. In this diagram, the p-propagation zone is defined as the region above both $\log_{10}(N^2)$ and $\log_{10}(L^2)$ (represented with khaki colour pattern) and the g-propagation zone is defined as the region below both $\log_{10}(N^2)$ and $\log_{10}(L^2)$ (represented with orange colour pattern). Frequencies of different modes are shown with horizontal lines and the position of the nodes of the corresponding eigenfunctions are represented with diamonds.

In the hot and dynamically evolving environment of a PNS, mode classification becomes more challenging than in the case of cold NSs. To classify the non-radial modes of oscillation, we apply different classification schemes that provide complementary physical information about the mode characteristics and insights into their evolution. The three different approaches we use are described in the following subsections.

3.1 Cowling classification

This classification scheme introduced by Cowling (1941) is based on the restoring force (pressure or buoyancy), acting on a fluid element when it is displaced from its equilibrium position inside the star. The number of nodes, n , in the radial eigenfunction η_r , i.e. the number of times η_r changes sign inside the star, is used to classify the modes. In this classification, the fundamental f mode has 0 nodes, and each p_n mode and g_n mode have n nodes, with $n \geq 1$. The way to distinguish a p mode from a g mode with the same number of nodes is by comparing their frequencies: p modes have higher frequencies than g modes, and f modes are in the middle.

In the top panel of Fig. 2, we show as an example the radial eigenfunction profiles of the modes depicted in Fig. 1. According to this classification, the green curve corresponds to a p_1 mode, with 1 node in the p-propagation zone represented with a filled diamond. The yellow curve corresponds to a g_2 mode with 2 nodes, one in the p-propagation zone and the other in the evanescent zone represented with filled and empty diamonds, respectively. Finally, a g_1 mode is represented with the purple curve, having 1 node in the g-propagation zone, represented with an empty diamond. In this example, there is no f mode between p modes and g modes. This classification is also used in Ferrari, Miniutti & Pons (2003), Sotani & Takiwaki (2016), Morozova et al. (2018), Torres-Forné et al. (2018), Sotani et al. (2019), and Sotani & Takiwaki (2020a, b, c) for the modes of PNS, among others. Other works adopting this classification scheme for cold NS are, for example, Lindblom & Detweiler (1983), McDermott, van Horn & Hansen (1988), Benhar, Ferrari & Gualtieri (2004), Lasky (2015), Vásquez Flores, Hall & Jaikumar (2017), Ranea-Sandoval et al. (2018), and Kumar, Mishra & Malik (2023).

3.2 Generalized Cowling nomenclature

The GCN classification (Eckart 1960; Scuflaire 1974; Osaki 1975), takes into account the fact that, for evolved stars, the relation between the number of nodes in the eigenfunction and the mode number is not direct, in other words, modes do not necessarily have the same number of nodes along the stellar evolution, instead they can lose or gain nodes. The central panel of Fig. 2 illustrates the phase diagram (η_r, η_{\perp}) for the modes depicted in the top panel with the same colours. The solid black circle indicates the stellar centre, and the nodes of the eigenfunctions are classified into g nodes and p nodes, depending on whether the curve in the phase diagram is travelling clockwise (g nodes) or counterclockwise (p nodes) at the crossing of the axis $\eta_r = 0$, as the radial coordinate, r , increases (Unno et al. 1989). Accordingly, the empty diamonds represent g nodes, and the filled diamonds represent p nodes. The number of p nodes and g nodes are N_p and N_g , respectively. Their difference

$$\tilde{n} \equiv N_p - N_g, \quad (16)$$

increases monotonically with the mode frequency, and it is conserved for each mode as the star evolves. The modes are classified depending on the sign of \tilde{n} , which defines whether the mode is a g mode ($\tilde{n} < 0$) or a p mode ($\tilde{n} > 0$). For the fundamental mode, $\tilde{n} = 0$. Hence, using this classification in the middle panel of Fig. 2, the green curve shows a p_1 mode as it has $N_p = 1$ and $N_g = 0$ ($\tilde{n} = 1$). The yellow curve belongs to the fundamental mode as it has $N_p = 1$ and $N_g = 1$ ($\tilde{n} = 0$). Finally, the purple curve shows a g_1 mode as it has $N_p = 0$ and $N_g = 1$ ($\tilde{n} = -1$). Therefore, unlike in the Cowling Classification, there is a fundamental mode with two nodes between the g modes and p modes.




Figure 2. Representative example of the classification of PNS oscillation modes (see also the propagation diagram in Fig. 1). From top to bottom, the mode identifications are obtained with the Cowling classification, the GCN, and the CBMP, as described in the main text. The same three modes are represented in every panel with the same colours, but they are classified differently according to each method. On the top and middle panels, filled (empty) diamonds correspond to p -(g)-nodes; on the bottom panel, the shaded range corresponds to the PNS core.

3.3 Classification based on modal property

Even though the GCN is more appropriate to classify modes of evolved stars than the Cowling classification, it does not necessarily represent the character of the mode. Thereby, a second complementary classification, the CBMP, was introduced by Shibahashi & Osaki (1976). This classification is based on the characteristics of the main trapping zone of the oscillation mode. In the propagation diagram, if the two propagation zones are widely separated by an intermediate evanescent zone, oscillation modes can be classified into two types by identifying their main trapping zones. In general, evolved main-sequence stars can consist of four layers starting at the centre: a homogeneous convective core, a μ -gradient zone with




Figure 3. Time evolution of mode frequencies after the bounce for the PNS models shown in Table 1. The same modes are shown in every panel, but from top to bottom the classifications used are the Cowling Classification, GCN, and CBMP, respectively. For times greater than approximately 0.4 s after the bounce, all classifications agree on the identification of f and p modes. For earlier times, the Cowling classification is not well defined to the disappearance of the f mode; the GCN shows several avoided crossings and the CBMP identifies an additional mode, G_0 . See the main text for more details.

varying chemical potential μ , a thin convective shell that acts like an evanescent zone and the radiative envelope, see e.g. fig. 1 of Shibahashi & Osaki (1976). In analogy to the case of an evolved main-sequence star, we use the CBMP to classify modes according to whether they are trapped in the core or the envelope of the PNS (see Fig. 1, where the corresponding boundary between core and envelope is $r \sim 0.3R$). In order to describe this classification, some terminology needs to be defined

- (i) G_n : indicates a mode trapped mainly in the core zone that has n g nodes.
- (ii) p_n : denotes a mode trapped mainly in the p -propagation zone of the envelope and having n p nodes.
- (iii) g_n : a mode trapped mainly in the g -propagation zone of the envelope with n g nodes.

A key quantity for this classification scheme is the ratio of the kinetic energy of the oscillation mode within the core to that of the whole star, defined as:

$$\Delta = \frac{\int_0^{r_s} \rho |\xi|^2 4\pi r^2 dr}{\int_0^R \rho |\xi|^2 4\pi r^2 dr}, \quad (17)$$




Figure 4. Top: time evolution of mode frequencies classified with the GCN for PNS model 1 in Table 1. Middle: time evolution of the ratio between the kinetic energy of the oscillations within the core and their total kinetic energy inside the star, for the p_1 , f , and g_1 modes in the top panel. The horizontal line shows the threshold in Δ used for the CBMP, and the shaded region represents possible thresholds of Δ that give qualitatively the same result (see also Appendix A). Bottom: time evolution of the mode frequencies for the same model shown in the top panel, now classified using the CBMP.

where r_s is the outer radius of the core and the square of the displacement vector, ξ^2 , is given by

$$\xi^2 = \eta_r^2 + l(l+1)\eta_{\perp}^2/r^2. \quad (18)$$

According to this classification, if Δ is close to 0 then the mode is trapped in the envelope. Otherwise, the mode is trapped in the core. The threshold, Δ_{critic} , used to define whether a mode is trapped in the core or in the envelope is an arbitrary parameter that we will discuss in Section 4. The number of nodes of each mode is determined by the number of nodes in the region where the mode is trapped.

In the bottom panel of Fig. 2, we show the integrand in equation (17) as a function of the radius of the PNS for the modes shown in the top and middle panels. In this example, the p_1 mode in the GCN (green curve) has $\Delta = 0.66$, which means the mode is oscillating mostly in the core, without nodes in the core, so it is classified as a G_0 mode in this classification. On the other hand, the f mode in the GCN (orange curve) has $\Delta = 0.04$ and 1 node in the p -propagation zone, so it is classified as a p_1 mode in this classification. Something similar occurs with the g_1 mode in the GCN (purple curve), which is classified as an f mode because $\Delta = 0.15$, and it has no nodes in the envelope. Unlike the other methods, the advantage of this

classification is that it gives us information about the region of the star which contributes the most to the GW emission.

4 RESULTS

4.1 PNS modes

We consider only the quadrupolar ($l = 2$) modes as they should be the most energetic in the gravitational-wave radiation from PNSs.

To classify non-radial modes in PNSs, we consider 10 models of PNS evolution, shown in Table 1. We calculate the frequencies of the modes in the Cowling approximation at every 10 ms in each of the models and we classify the modes using the three different classification methods described in Section 3, focusing on the frequency range 100–2000 Hz.

In the Cowling classification, the modes are classified by counting the number of nodes of the amplitude of the radial Lagrangian displacement, η_r , and by comparing their frequency with the f mode (with zero nodes) as discussed in Section 3.1. The results are presented in the top panel of Fig. 3. The different symbols represent the PNS models introduced in Table 1 and the colours represent each mode classification, also indicated with labels in the figure (There are modes with 4 and 5 nodes in other regions of the figure but we only represent those necessary to complete the mode evolution). Strictly speaking, the Cowling approximation is only valid after $t \gtrsim 0.35$ s, when the f mode is present.² Going backwards in time from $t \sim 0.35$ s, we can see that modes gain nodes along the PNS evolution towards the bounce, when their frequencies get very close to each other, changing their Cowling classification. This feature is also seen in Morozova et al. (2018), Sotani et al. (2019), and Torres-Forné et al. (2019b). Additionally, Sotani & Sumiyoshi (2019) and Sotani & Takiwaki (2020c) found that the amplitude of the modes involved in each ‘close approach’ (see the avoided crossing discussion below) changes while the amplitude of the rest of the modes remains almost unchanged.

The mode classification obtained with the GCN is presented in the middle panel of Fig. 3. The different colours now represent the classification through fixed values of $\tilde{n} = N_p - N_g$, which increases monotonically with the frequency at every time-step. Progressing upward from the lowest frequency, the modes are uniquely classified as g_3 , g_2 , g_1 , f , p_1 , p_2 and p_3 (only modes with $|\tilde{n}| \leq 3$ are represented). Compared with the Cowling classification, in the GCN case we find that the description of the mode evolution is much simpler. The ‘close approaches’ noted in the top panel now appear as avoided crossings. For instance, as the f mode evolves, it shows avoided crossings as it bounces between the p_1 and g_1 modes: at $t \simeq 0.05$ s with p_1 , at $t \simeq 0.15$ with g_1 , at $t \simeq 0.3$ s with p_1 and finally at $t \simeq 0.5$ s with g_1 .

The results obtained in the CBMP are presented in the bottom panel of Fig. 3. The colours represent different mode classifications. Modes with upper case (G_i modes) and lower case (p_i , f , and g_i modes) letters correspond to modes trapped in the core and in the envelope, respectively. The sub-index in each label represents the number of nodes in the region where the mode is trapped. For instance, the G_2 mode is trapped in the core and has 2 nodes in the core, the f mode is trapped in the envelope and has no nodes in the envelope (but it could

²We choose to follow Morozova et al. (2018) and classify the mode with 2 nodes at $0.1 \text{ s} \lesssim t \lesssim 0.4 \text{ s}$ as a g_2 mode, but the classification is not unique (it could also be classified as a p_2 mode) due to the absence of the f mode at early times.




Figure 5. Evolution of the parameters used in each mode classification method, corresponding to the f mode (as classified by the GCN). Times are specified on the left-hand side and mode classifications are shown as labels in each panel. Left: evolution of the radial eigenfunction η_r , used in the Cowling classification method. Centre: mode trajectories in the (η_r, η_\perp) phase space, used in the GCN. Right: evolution of Δ as a function of the radius of the PNS.

have nodes in the core), and the p_2 mode is trapped in the envelope and has 2 nodes in the p -propagation zone in the envelope. We note that the G_i modes decrease in frequency over time, and there are no more avoided crossings. In particular, the G_0 mode appears, crossing all p modes and the f mode.

It is remarkable that all of the PNS models we analyse show very similar results. As a representative example, we choose model 1 (with a $9 M_\odot$ progenitor star) to explore the complementary descriptions provided by the GCN and CBMP. In Fig. 4, we present the mode evolution only for model 1, using the GCN and the CBMP in the top and bottom panels, respectively. In the central panel, we show the evolution of Δ for the g_1 , f , and p_1 modes, as classified by the GCN. The horizontal black line corresponds to the Δ threshold we use in this work. The grey band represents a Δ range for which the classification would be qualitatively the same (see more details in Appendix A). Values of Δ above the horizontal black line indicate the mode is trapped in the core. Otherwise, the mode is trapped in the envelope of the PNS. Near each avoided crossing in the GCN, Δ changes abruptly and the character of the mode changes (trapped in the core or trapped in the envelope).

We take a closer look at the evolution of the f mode (as classified by the GCN) in Fig. 5. For 8 time slices, we present the radial eigenfunction in the left-hand columns, the (η_r, η_\perp) phase space in the centre and the integrand used in the definition of Δ in the right-hand column (note that the PNS becomes more compact with time, as it cools and approaches the cold NS stage). The f mode gains and loses pairs of p and g nodes, changing its Cowling classification but keeping \tilde{n} constant in the GCN. The CBMP shows the changes in the mode character, alternating between oscillations trapped in the core (G_i classifications) and in the envelope (p_i and f classifications).

4.2 Emission of gravitational waves

It is relevant to ask how most of the GW emission by the PNS is generated. A comparison between the time-evolving spectrum of modes and the spectrogram of the GWs for model 9 is presented in Fig. 6.




Figure 6. Spectrogram of the GW emission for model 9 (see Table 1), shown together with the f-mode sequences identified with the GCN and the CBMP. For earlier times, the GCN shows better agreement with the simulated GWs. For later times, both methods agree on the f-mode classification.




Figure 7. Relation between macroscopic quantities (mass, M and radius, R) and the frequencies of modes classified using the GCN (top panel) and the CBMP (bottom panel). The CBMP sequences show simpler trends than the GCN sequences, due to the absence of the avoided crossings. The black lines represent the fits to each mode in the CBMP (see Table 2) and the shaded areas represent the $\pm 1\sigma$ bands of each fit. Note that the frequencies corresponding to the G_0 modes below the f modes in the bottom panel are not included in the top panel because they are classified as g_1 modes in the GCN.

We show with lines the evolution of the f mode obtained following the GCN classification and the CBMP. In this analysis, we use the modes calculated with the full equations (4)–(7) in order to obtain a better agreement with the simulated GWs. As the GCN and CBMP are strictly valid only in the Cowling approximation of equations (13)

Table 2. Best-fitting parameters for quasi-universal relations of the form $f(x) = a + bx + cx^2$, describing the CBMP mode frequency f in Hz as a function of $x \equiv \sqrt{M/R^3}$ in units of $M_\odot^{1/2}/\text{km}^{3/2}$, shown in Fig. 7.

Mode	a (Hz)	b (Hz $\text{km}^{3/2}/M_\odot^{1/2}$)	c (Hz km^3/M_\odot)
f	-0.050 ± 0.004	174.3 ± 1.9	-2590 ± 195
p_1	0.067 ± 0.003	184.6 ± 1.6	2194 ± 176
p_2	0.062 ± 0.004	313.2 ± 2.5	-6213 ± 326
p_3	0.057 ± 0.006	452.6 ± 4.9	-16780 ± 807
G_0	0.94 ± 0.01	-4.8 ± 5.7	-4715 ± 596

and (14), the classifications shown here were obtained in comparison with the results for the Cowling approximation in the same model. The GCN f mode shows very good agreement with the strongest GW emission and not very prominent avoided crossings at least at $t \simeq 0.15$ and 0.4 s. This result, together with the detailed analysis of the GCN f mode presented in Fig. 5, indicates that the PNS spectrum evolves with avoided crossings. Consequently, it emits GWs mainly through an f mode that rapidly alternates between being trapped in the core and in the envelope of the PNS. We note that, despite the discrepancy between the CBMP f mode and the GW emission at early times, the CBMP classification is still applicable and can provide interesting information, for example helping in the characterization of the modal properties of the GCN f mode and in the identification of avoided crossings.

4.3 Quasi-universal relations

Universal relations between mode frequencies and macroscopic parameters, such as mass and radius, have been proposed as a possible tool to infer properties of NSs using the detection of GWs, see e.g. Andersson & Kokkotas (1998), Benhar et al. (2004), Tsui & Leung (2005), Lau, Leung & Lin (2010), Chirenti, de Souza & Kastaun (2015), Sotani & Kumar (2021), and Ranea-Sandoval et al. (2023).

Here, we analyse some quasi-universal relations proposed in the literature for cold NSs in the context of the dynamic PNS. As all of our PNS models have been produced with the SFHo EoS for dense matter derived by Steiner et al. (2013), we are only able to probe the influence of the progenitor mass.

It has been shown that the frequency of the fundamental mode of cold NSs scales with $\sqrt{M/R^3}$ with minor dependence on the EoS (see e.g. Andersson & Kokkotas 1998; Benhar et al. 2004; Chirenti et al. 2015). Torres-Forné et al. (2019a) proposed a similar dependence for the frequencies of f and p modes of a PNS, see also Sotani & Sumiyoshi (2021) and Mori et al. (2023).

However, note that there is no unique definition for the radius of PNS. Torres-Forné et al. (2019a) consider that the PNS extends up to the radial position of the shock wave, while we follow Morozova et al. (2018) and use $\rho = 10^{10} \text{ g cm}^{-3}$ as a condition to define the radius, which gives somewhat larger values (but both choices produce similar values for $\sqrt{M/R^3}$).

In the top panel of Fig. 7, we show sequences of modes classified with the GCN method. As previously, the symbols represent the different models, and each colour corresponds to a mode. The avoided crossings spread the curves, which are mostly independent of the progenitor mass (time increases to the right). In the bottom panel of Fig. 7, we present the sequences of modes classified with the CBMP. Now f and p modes present a very simple monotonic behaviour which is independent of the progenitor mass. The G_0 mode also shows a simple trend but with a larger spread. The fits to this novel

quasi-universal relation for the modes classified with the CBMP is presented in Table 2.

5 CONCLUSIONS AND DISCUSSION

The asteroseismology of PNSs is complicated by their dynamic and temperature-dependent nature. In this paper, we consider three different methods to classify their non-radial modes of oscillation: the Cowling classification, the GCN, and the CBMP.

The standard Cowling classification is not well defined for a PNS at early times after the bounce. It relies on the identification of a fundamental f mode without nodes to distinguish between p modes (with higher frequencies) and g modes (with lower frequencies). This mode is not present at early times, causing the mode identifications to be underdetermined.

The GCN relies on the identification of *p* and *g* nodes and allows us to follow the mode evolution. It shows marked avoided crossing at early times, at which two modes exchange character. The f mode identified with the GCN has very good agreement with the simulated GW emission of a PNS. Interestingly, this implies that the region responsible for most of the GWs (where the mode is trapped), quickly alternates between the core and envelope of the star at early times.

The CBMP follows the time evolution of sequences of modes with the same properties (trapped in the core or trapped in the envelope) and it is applied here for the first time to the study of PNS oscillation modes. The behaviour of these sequences is remarkably simple and monotonic, as the avoided crossings are absent, and we present fits for quasi-universal relations (independent of progenitor mass) describing the most relevant mode frequencies as functions of the PNS average density. We note that the matching classification scheme of Torres-Forné et al. (2019b), in which modes are classified according to the similarities between their eigenfunctions, produces results qualitatively similar to the CBMP, showing no avoided crossings at early times and a mode compatible with the G_0 mode, classified there as a g_2 mode. Additionally, the ramp-up mode discussed by Mori et al. (2023) is similar to the f mode described by the CBMP. One of the main advantages of the CBMP is that it gives physical information relative to the part of the PNS where GWs are most likely being produced.

There are a few different promising directions for extensions of our work. First, formal work should be done to extend the GCN and CBMP criteria beyond the Cowling approximation ($\delta\alpha \neq 0$)³ and to calculate the modes using time-dependent perturbation theory. Secondly, the quasi-universal behaviour of the mode frequencies should be checked against simulations with different progenitor masses and additional EoSs.

Finally, an assessment of the detectability (and reliable identification) of these modes in a galactic core-collapse supernova would be of great interest. Recently, Bruel et al. (2023) have estimated that current GW detectors could infer properties of such events up to the Large Magellanic Cloud. The mode evolution provides the rate of contraction of the PNS, which depends not only on the EOS but also on transport properties in dense matter, related to the rate of deleptonization of the star. Abundant neutrinos detected from a galactic core-collapse supernova will provide complementary information (Nagakura & Vartanyan 2022), potentially allowing for the independent determination of the time-evolving mass and radius of the PNS.

³See also an alternative approach by Takata (2012) for taking the perturbation of the gravitational potential into account by using a continuous integral index.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge Viktoriya Morozova for guidance related to the PNS models and QNM calculations during the early stages of this project, and Adam Burrows and Shin Yoshida for useful comments and discussions. MCR and IFR acknowledge CONICET: Consejo Nacional de Investigaciones Científicas y Técnicas (National Scientific and Technical Research Council) and UNLP: Universidad Nacional de La Plata (National University of La Plata) for financial support under grants G157 and G007. IFR was also partially supported by PICT 2019–0366 from ANPCyT (Argentina) and by the National Science Foundation (USA) under grant no. PHY-2012152. CC acknowledges support by NASA: National Aeronautics and Space Administration (USA) under award numbers 80GSFC17M0002 and TCAN80NSSC18K1488. DR acknowledges funding from the U.S. Department of Energy, Office of Science, Division of Nuclear Physics under award number(s) DE-SC0021177 and from the National Science Foundation under grant nos PHY-2011725, PHY-2020275, PHY-2116686, and AST-2108467.

DATA AVAILABILITY

The computed data presented and discussed in this paper will be shared upon reasonable request.

REFERENCES

Abbott B. P. et al., 2016, *ApJ*, 818, L22
 Abbott B. P. et al., 2017a, *Phys. Rev. D*, 96, 122004
 Abbott B. P. et al., 2017b, *ApJ*, 848, L12
 Abbott B. P. et al., 2019, *Phys. Rev. X*, 9, 031040
 Abbott R. et al., 2021, *ApJ*, 913, L7
 Abbott R. et al., 2021a, preprint ([arXiv:2111.03606](https://arxiv.org/abs/2111.03606))
 Abbott R. et al., 2021b, *Phys. Rev. X*, 11, 021053
 Abdikamalov E., Gossan S., DeMaio A. M., Ott C. D., 2014, *Phys. Rev. D*, 90, 044001
 Abdikamalov E., Pagliaroli G., Radice D., 2022, in Bambi C., Katsanevas S., Kokkotas K. D., eds, *Handbook of Gravitational Wave Astronomy*. Springer, Singapore, p. 21
 Andersson N., Kokkotas K. D., 1998, *MNRAS*, 299, 1059
 Andresen H., Müller B., Müller E., Janka H. T., 2017, *MNRAS*, 468, 2032
 Astone P., Cerdá-Durán P., Di Palma I., Drago M., Muciaccia F., Palomba C., Ricci F., 2018, *Phys. Rev. D*, 98, 122002
 Benhar O., Ferrari V., Gualtieri L., 2004, *Phys. Rev. D*, 70, 124015
 Bruel T. et al., 2023, *Phys. Rev. D*, 107, 083029
 Burrows A., Vartanyan D., 2021, *Nature*, 589, 29
 Burrows A., Vartanyan D., Dolence J. C., Skinner M. A., Radice D., 2018, *Space Sci. Rev.*, 214, 33
 Burrows A., Radice D., Vartanyan D., Nagakura H., Skinner M. A., Dolence J. C., 2020, *MNRAS*, 491, 2715
 Cavaglià M., Gaudio S., Hansen T., Staats K., Szczepańczyk M., Zanolin M., 2020, *Mach. Learn.: Sci. Technol.*, 1, 015005
 Cerdá-Durán P., DeBrye N., Aloy M. A., Font J. A., Obergaulinger M., 2013, *ApJ*, 779, L18
 Chan M. L., Heng I. S., Messenger C., 2020, *Phys. Rev. D*, 102, 043022
 Chirenti C., de Souza G. H., Kastaun W., 2015, *Phys. Rev. D*, 91, 044034
 Cowling T. G., 1941, *MNRAS*, 101, 367
 Eckart C., 1960, *Hydrodynamics of Oceans and Atmospheres*. Pergamon Press, New York
 Ferrari V., Miniutti G., Pons J. A., 2003, *Class. Quantam Gravity*, 20, S841
 Gill K., Wang W., Valdez O., Szczepańczyk M., Zanolin M., Mukherjee S., 2018, preprint ([arXiv:1802.07255](https://arxiv.org/abs/1802.07255))
 Hayama K., Kuroda T., Nakamura K., Yamada S., 2016, *Phys. Rev. Lett.*, 116, 151102
 Hayama K., Kuroda T., Kotake K., Takiwaki T., 2018, *MNRAS*, 477, L96
 Heng I. S., 2009, *Class. Quantam Gravity*, 26, 105005

Horowitz C. J., Caballero O. L., Lin Z., O'Connor E., Schwenk A., 2017, *Phys. Rev. C*, 95, 025801

Iess A., Cuoco E., Morawski F., Powell J., 2020, *Mach. Learn.: Sci. Technol.*, 1, 025014

Isenberg J. A., 2008, *Int. J. Mod. Phys. D*, 17, 265

Kokkotas K. D., Schmidt B. G., 1999, *Living Rev. Relativ.*, 2, 2

Kumar D., Mishra H., Malik T., 2023, *J. Cosmol. Astropart. Phys.*, 2023, 015

Kuroda T., Kotake K., Takiwaki T., 2016, *ApJ*, 829, L14

Kuroda T., Kotake K., Hayama K., Takiwaki T., 2017, *ApJ*, 851, 62

Lasky P. D., 2015, *Publ. Astron. Soc. Aust.*, 32, e034

Lau H. K., Leung P. T., Lin L. M., 2010, *ApJ*, 714, 1234

Lindblom L., Detweiler S. L., 1983, *ApJS*, 53, 73

López M., Di Palma I., Drago M., Cerdá-Durán P., Ricci F., 2021, *Phys. Rev. D*, 103, 063011

McDermott P. N., van Horn H. M., Scholl J. F., 1983, *ApJ*, 268, 837

McDermott P. N., van Horn H. M., Hansen C. J., 1988, *ApJ*, 325, 725

Marek A., Janka H. T., Müller E., 2009, *A&A*, 496, 475

Mezzacappa A., Endeve E., Messer O. E. B., Bruenn S. W., 2020a, *Liv. Rev. Comput. Astrophys.*, 6, 4

Mezzacappa A. et al., 2020b, *Phys. Rev. D*, 102, 023027

Mezzacappa A. et al., 2023, *Phys. Rev. D*, 107, 043008

Mori M., Suwa Y., Takiwaki T., 2023, *Phys. Rev. D*, 107, 083015

Morozova V., Radice D., Burrows A., Vartanyan D., 2018, *ApJ*, 861, 10

Mukherjee S., Salazar L., Mittelstaedt J., Valdez O., 2017, *Phys. Rev. D*, 96, 104033

Müller B., Janka H.-T., Marek A., 2013, *ApJ*, 766, 43

Murphy J. W., Ott C. D., Burrows A., 2009, *ApJ*, 707, 1173

Nagakura H., Vartanyan D., 2022, *MNRAS*, 512, 2806

Nakamura K., Takiwaki T., Kotake K., 2022, *MNRAS*, 514, 3941

O'Connor E. P., Couch S. M., 2018, *ApJ*, 865, 81

O'Connor E. et al., 2018, *J. Phys. G: Nucl. Part. Phys.*, 45, 104001

Osaki Y., 1975, *PASJ*, 27, 237

Pan K.-C., Liebendörfer M., Couch S. M., Thielemann F.-K., 2018, *ApJ*, 857, 13

Pons J. A., Reddy S., Prakash M., Lattimer J. M., Miralles J. A., 1999, *ApJ*, 513, 780

Powell J., Müller B., 2019, *MNRAS*, 487, 1178

Powell J., Müller B., 2022, *Phys. Rev. D*, 105, 063018

Powell J., Szczepański M., Heng I. S., 2017, *Phys. Rev. D*, 96, 123013

Radice D., Morozova V., Burrows A., Vartanyan D., Nagakura H., 2019, *ApJ*, 876, L9

Ranea-Sandoval I. F., Guilera O. M., Mariani M., Orsaria M. G., 2018, *J. Cosmol. Astropart. Phys.*, 2018, 031

Ranea-Sandoval I. F., Mariani M., Lugones G., Guilera O. M., 2023, *MNRAS*, 519, 3194

Roma V., Powell J., Heng I. S., Frey R., 2019, *Phys. Rev. D*, 99, 063018

Röver C., Bizouard M.-A., Christensen N., Dimmelmeier H., Heng I. S., Meyer R., 2009, *Phys. Rev. D*, 80, 102004

Scuflaire R., 1974, *A&A*, 36, 107

Shibahashi H., Osaki Y., 1976, *PASJ*, 28, 199

Skinner M. A., Dolence J. C., Burrows A., Radice D., Vartanyan D., 2019, *ApJS*, 241, 7

Sotani H., Kumar B., 2021, *Phys. Rev. D*, 104, 123002

Sotani H., Sumiyoshi K., 2019, *Phys. Rev. D*, 100, 083008

Sotani H., Sumiyoshi K., 2021, *MNRAS*, 507, 2766

Sotani H., Takiwaki T., 2016, *Phys. Rev. D*, 94, 044043

Sotani H., Takiwaki T., 2020a, *Phys. Rev. D*, 102, 023028

Sotani H., Takiwaki T., 2020b, *Phys. Rev. D*, 102, 063025

Sotani H., Takiwaki T., 2020c, *MNRAS*, 498, 3503

Sotani H., Kuroda T., Takiwaki T., Kotake K., 2019, *Phys. Rev. D*, 99, 123024

Steiner A. W., Hempel M., Fischer T., 2013, *ApJ*, 774, 17

Summerscales T. Z., Burrows A., Finn L. S., Ott C. D., 2008, *ApJ*, 678, 1142

Takata M., 2012, *PASJ*, 64, 66

Torres-Forné A., Cerdá-Durán P., Passamonti A., Font J. A., 2018, *MNRAS*, 474, 5272

Torres-Forné A., Cerdá-Durán P., Obergaulinger M., Müller B., Font J. A., 2019a, *Phys. Rev. Lett.*, 123, 051102

Torres-Forné A., Cerdá-Durán P., Passamonti A., Obergaulinger M., Font J. A., 2019b, *MNRAS*, 482, 3967

Tsui L. K., Leung P. T., 2005, *MNRAS*, 357, 1029

Unno W., Osaki Y., Ando H., Saio H., Shibahashi H., 1989, Nonradial Oscillations of Stars, 2nd edn. Univ. of Tokyo Press, Tokyo

Vartanyan D., Coleman M. S. B., Burrows A., 2022, *MNRAS*, 510, 4689

Vartanyan D., Burrows A., Wang T., Coleman M. S. B., White C. J., 2023, *Phys. Rev. D*, 107, 103015

Vásquez Flores C., Hall Z. B., Jaikumar P., 2017, *Phys. Rev. C*, 96, 065803

Westernacher-Schneider J. R., 2020, *Phys. Rev. D*, 101, 083021

Wilson J. R., Mathews G. J., 1989, in Evans C. R., Finn L. S., Hobill D. W., eds, Frontiers in Numerical Relativity. Cambridge University Press, p. 306

Wolfe N. E., Frohlich C., Miller J. M., Torres-Forné A., Cerdá-Durán P., 2023, preprint (arXiv:2303.16962)

Yakunin K. N. et al., 2015, *Phys. Rev. D*, 92, 084040

APPENDIX A: RESULTS FOR DIFFERENT Δ THRESHOLDS

In the CBMP, modes are classified by the region in the star where they are trapped. With this method, the modes change character when the ratio of the kinetic energy of the oscillations within the core to that of the whole star, Δ , significantly changes from ~ 0 to ~ 1 , and vice versa. In this appendix, we test different values of Δ_{critic} from which we determine if the mode is trapped in the core or in the envelope of the PNS.

In Fig. A1, we show the time evolution of the mode frequencies classified with the CBMP. In each panel, we used a different value of Δ_{critic} from 0.2 to 0.5. The modes change character at different times for each case: for larger Δ_{critic} , modes trapped in the core (envelope) become trapped in the envelope (core) earlier (later) compared with smaller Δ_{critic} . Therefore, for $0.2 \leq \Delta_{\text{critic}} \leq 0.5$, the results are qualitatively the same.




Figure A1. Time evolution of mode frequencies after the bounce using the CMBP for values of Δ_{critic} between 0.2 and 0.5 (see middle panel of Fig. 4). The results are quantitatively the same in all cases; the core-trapped G_0 mode becomes slightly more prominent with decreasing Δ_{critic} .

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.