

ALESQP: AN AUGMENTED LAGRANGIAN EQUALITY-CONSTRAINED SQP METHOD FOR OPTIMIZATION WITH GENERAL CONSTRAINTS*

HARBIR ANTIL[†], DREW P. KOURI[‡], AND DENIS RIDZAL[‡]

Abstract. We present a new algorithm for infinite-dimensional optimization with general constraints, called ALESQP. In short, ALESQP is an augmented Lagrangian method that penalizes inequality constraints and solves equality-constrained nonlinear optimization subproblems at every iteration. The subproblems are solved using a matrix-free trust-region sequential quadratic programming (SQP) method that takes advantage of iterative, i.e., inexact linear solvers, and is suitable for large-scale applications. A key feature of ALESQP is a constraint decomposition strategy that allows it to exploit problem-specific variable scalings and inner products. We analyze convergence of ALESQP under different assumptions. We show that strong accumulation points are stationary. Consequently, in finite dimensions ALESQP converges to a stationary point. In infinite dimensions we establish that weak accumulation points are feasible in many practical situations. Under additional assumptions we show that weak accumulation points are stationary. We present several infinite-dimensional examples where ALESQP shows remarkable discretization-independent performance in all of its iterative components, requiring a modest number of iterations to meet constraint tolerances at the level of machine precision. Also, we demonstrate a fully matrix-free solution of an infinite-dimensional problem with nonlinear inequality constraints.

Key words. ALESQP, augmented Lagrangian, composite step trust-region method, SQP, convergence analysis, constraint decomposition, nonlinear constraints

MSC codes. 49M37, 90C30, 90C39, 93C20, 49K20, 49J20

DOI. 10.1137/20M1378399

1. Introduction. In this paper, we develop a provably convergent algorithm for solving optimization problems of the form

$$(1.1) \quad \min_{x \in X} f(x) \quad \text{subject to} \quad g(x) = 0, \quad Tx \in \bigcap_{i=1}^m C_i,$$

* Received by the editors November 4, 2020; accepted for publication (in revised form) September 2, 2022; published electronically February 3, 2023.

<https://doi.org/10.1137/20M1378399>

Funding: The first author was partially supported by NSF grants DMS-2110263 and DMS-1913004 and by the Air Force Office of Scientific Research under awards FA9550-19-1-0036 and FA9550-22-1-0248. The second and third authors were supported by the U.S. Air Force Office of Scientific Research, Optimization Program under award F4FGA09135G001. The third author was also supported by Sandia National Laboratories' LDRD program. This article has been authored by an employee of National Technology & Engineering Solutions of Sandia, LLC under Contract No. DE-NA0003525 with the U.S. Department of Energy (DOE). The employee owns all right, title and interest in and to the article and is solely responsible for its contents. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this article or allow others to do so, for United States Government purposes. The DOE will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan.

[†] Department of Mathematical Sciences and Center for Mathematics and Artificial Intelligence CMAI, George Mason University, Fairfax, VA 22030 USA (hantil@gmu.edu).

[‡] Optimization and Uncertainty Quantification, Sandia National Laboratories, Albuquerque, NM 87185 USA (dpkouri@sandia.gov, dridzal@sandia.gov).

where X and Y are real Banach spaces, Z is a real Hilbert space, $f : X \rightarrow \mathbb{R}$, $g : X \rightarrow Y$, and $T : X \rightarrow Z$ is a linear operator. Moreover, the sets C_1, \dots, C_m are nonempty, closed, and convex, where $C_i \subseteq Z$ for $i = 1, \dots, m$. The optimization problem (1.1) encompasses many finite-dimensional and infinite-dimensional nonlinear optimization problems. Our proposed algorithm employs m individual augmented Lagrangian penalties to handle the constraints $Tx \in C_i$ for $i = 1, \dots, m$. The resulting subproblems are *equality constrained* and can be solved efficiently using modern sequential quadratic programming (SQP) methods. By separately penalizing the constraints $Tx \in C_i$, for $i = 1, \dots, m$, our algorithm increases the augmented Lagrangian penalty parameters according to the associated infeasibility of the current iterate, adapting the i th penalty parameter to the specific scaling of the constraint $Tx \in C_i$. We call the proposed algorithm the *augmented Lagrangian equality-constrained SQP (ALESQP)* method.

To motivate (1.1) and ALESQP, we note that optimization problems constrained by partial differential equations (PDEs) can be written in the form (1.1), where $x = (u, z)$ is split into state u and control z variables and the equality constraint $g(x) = 0$ represents the governing PDE. For such problems, it is often difficult to prove regularity of multipliers, especially when constraints of the type $Tx \in C_i$ are enforced on the PDE solution [10, 22, 31]. As a result, nonlinear programming methods developed in finite dimensions often exhibit *mesh dependence* when applied to discretizations of such problems. More specifically, their algorithmic performance degrades with refinement of the PDE discretization—in other words, with increasing problem size. We tackle this particular challenge by deriving and analyzing ALESQP in an *infinite-dimensional* setting. A common approach to solving PDE-constrained optimization problems is to reformulate (1.1), with the aforementioned splitting $x = (u, z)$, by eliminating the PDE solution variable u . When the PDE is nonlinear, this approach requires a nonlinear solver, e.g., a Newton-type iteration, to solve the PDE and evaluate the objective function at every optimization iteration. In contrast, optimization formulation (1.1) allows us to maintain the PDE as an explicit constraint. In doing so, ALESQP does not require an accurate solution of the equality constraint $g(x) = 0$ until convergence, and, in fact, it balances the PDE solution accuracy with other feasibility and optimality metrics as the algorithm iterates. This further allows us to approximately solve equality-constrained subproblems using inexact matrix-free SQP methods that take advantage of iterative linear system solves [19, 20] and mesh adaptivity [47].

The augmented Lagrangian method, or the method of multipliers, was originally introduced in [21, 34] for finite-dimensional, equality-constrained optimization and subsequently extended and analyzed by numerous authors; see [6, 36, 37, 38]. Augmented Lagrangian also serves as the backbone of numerous successful numerical optimization software packages. For example, the MINOS solver uses an augmented Lagrangian penalty for linearized equality constraints and solves linearly constrained subproblems, the LANCELOT solver employs an augmented Lagrangian penalty for equality constraints and solves bound-constrained subproblems [14], and the ALGENCAN solver has the ability to use augmented Lagrangian penalties to handle both equality and inequality constraints [8]. The ALESQP method is closely related to two existing augmented Lagrangian approaches: LANCELOT and sequential equality-constrained optimization (SECO) [7]. Our approach generalizes the problem formulation of SECO and solves a sequence of penalized equality-constrained subproblems. An important addition to the SECO algorithm is in the use of *multiple augmented Lagrangian penalties* to handle disparate constraint scalings.

The general mechanics of the ALESQP algorithm are borrowed from the LANCELOT solver described in [12], including the penalty parameter and multiplier update procedures, with extensions to support multiple penalties. A principal difference between ALESQP and both SECO and LANCELOT is that we prove convergence for infinite-dimensional problems. This advance enables discretization-independent performance of ALESQP on such problems, including mesh-based discretizations. Notably, we observe mesh-independent performance in all iterative components of ALESQP, including the augmented Lagrangian iteration, its SQP subproblem solver, and SQP's quadratic optimization solver.

In contrast to the extensive body of work on augmented Lagrangian methods and software for the solution of finite-dimensional optimization problems, there has been little work on solving general infinite-dimensional optimization problems using the augmented Lagrangian. For instance, the references [4, 5, 23, 24, 26] are limited to specific convex optimization problems, treat only finite-dimensional constraints, or require strong assumptions and therefore do not support the solution of the general problem (1.1). In addition, the augmented Lagrangian SQP methods developed in [25, 43, 44] add an additional SQP step to the augmented Lagrangian algorithm to accelerate the convergence of the Lagrange multiplier estimates. However, these methods apply only to equality-constrained optimization problems. Only recently Börgens, Kanzow, and Steck [9] introduced and analyzed a generally applicable infinite-dimensional augmented Lagrangian method. There are four major differences between ALESQP and the method presented in [9]. First, we consider a different problem formulation, with an emphasis on maintaining the explicit constraint $g(x) = 0$. In the context of PDE-constrained optimization, where $g(x) = 0$ encompasses the PDE constraint, this choice crucially enables an *inexact and therefore efficient* solution of the governing PDE, through rigorous use of iterative linear solvers [19] and mesh adaptivity [47]. Second, we treat all constraints of the type $Tx \in C_i$ in a unified fashion, through multiple penalties and the corresponding multiplier and penalty updates, and we solve equality-constrained subproblems. In contrast, due to strong regularity assumptions on the constraint function in [9] (complete continuity of the mapping G [9, Assump. 5.1]), certain inequality constraints must be treated implicitly, as part of the subproblem, while others are penalized using the augmented Lagrangian. Third, we provide a complete algorithmic framework, with a discussion of methods that are chosen specifically for their suitability as ALESQP subproblem solvers. We demonstrate excellent performance on a variety of infinite-dimensional problems, with nearly constant iteration counts *in all algorithmic components* of the ALESQP framework, independent of problem size. Fourth, we do not employ a multiplier safeguard (also used in, e.g., [2, 8]). Rather, we use the multiplier update from LANCELOT; see [12].

The remainder of the paper is organized as follows. In sections 2 and 3 we introduce the notation and describe the assumptions on (1.1), recalling the associated optimality conditions. In sections 4 and 5 we introduce the augmented Lagrangian algorithm and prove asymptotic stationarity and asymptotic feasibility of the generated sequence of iterates. We build on these results and show that, under additional assumptions, weak accumulation points of the sequence of iterates are stationary points for (1.1). In section 6 we extend the augmented Lagrangian formulation to handle nonlinear constraint operators T . In section 7 we briefly discuss the remaining components of the ALESQP framework, including the SQP algorithm and its subroutines. We conclude with a variety of numerical results including statistical estimation and PDE-constrained optimization in section 8.

2. Notation. Given a Banach space $(X, \|\cdot\|_X)$, we denote the topological dual space of X by X^* and the associated dual pairing by $\langle \cdot, \cdot \rangle_{X^*, X}$. If X is a Hilbert space, we denote by $(\cdot, \cdot)_X$ the inner product on X and we assume that $\|\cdot\|_X$ is the usual norm on X . We denote by B_ρ^X for $\rho > 0$ the closed norm ball on X with radius ρ . For two Banach spaces X and Y , we denote the space of bounded linear operators that map X into Y by $\mathcal{L}(X, Y)$. For a closed, convex subset C of the Banach space X , we denote the projection of a point $x \in X$ onto C by $\mathbf{P}_C(x)$ and the distance from x to C by $d_C(x)$. That is, $\mathbf{P}_C(x)$ and $d_C(x)$ satisfy

$$d_C(x) := \min_{y \in C} \|x - y\|_X = \|x - \mathbf{P}_C(x)\|_X.$$

In addition, we denote the normal cone to C at the point $x \in C$ by

$$N_C(x) := \{\lambda \in X^* \mid \langle \lambda, y - x \rangle_{X^*, X} \leq 0 \quad \forall y \in C\}$$

with $N_C(x) = \emptyset$ if $x \notin C$. Finally, we denote convergence with respect to the weak topology by \rightharpoonup , convergence with respect to the weak* topology by \rightharpoonup^* , and convergence with respect to the norm topology by \rightarrow .

3. Problem formulation and assumptions. Let X and Y be real Banach spaces and let Z be a real Hilbert space. To simplify the presentation, we will associate Z^* with Z . Given the problem data $f : X \rightarrow \mathbb{R}$, $g : X \rightarrow Y$, $T \in \mathcal{L}(X, Z)$ and a nonempty, closed, and convex set $C \subseteq Z$, we consider the optimization problem

$$(3.1) \quad \min_{x \in X} f(x) \quad \text{subject to} \quad g(x) = 0, \quad Tx \in C.$$

When f and g are Fréchet differentiable, we say that $\bar{x} \in X$ is a first-order stationary point of (3.1) if there exists $\bar{\zeta} \in Y$ such that

$$(3.2) \quad -(f'(\bar{x}) + g'(\bar{x})^* \bar{\zeta}) \in T^* N_C(T\bar{x}) \quad \text{and} \quad g(\bar{x}) = 0.$$

Note that this presumes $T\bar{x} \in C$ since the normal cone is empty otherwise.

Remark 3.1 (Banach space valued constraints). As in [9], we could consider the case where Z is a real Banach space that is densely embedded in a real Hilbert space. However, this would complicate the presentation with little added benefit.

To prove convergence of our algorithm, we will require the following assumptions on the objective function f , the constraint operators g and T , and the constraint set C . In our subsequent analysis, we will explicitly state when each assumption is required. Assumptions (A0) (feasibility) and (A1) (differentiability) will be required throughout, whereas (A2), (A3), and (A4) will only be required to prove convergence.

Assumption 3.2 (Regularity of problem data).

- (A0) There exists $\bar{x} \in X$ such that $g(\bar{x}) = 0$ and $T\bar{x} \in C$.
- (A1) The functions f and g are continuously Fréchet differentiable.
- (A2) The adjoint operator T^* is injective.
- (A3) The functions f and $\|g(\cdot)\|_Y$ are weakly lower semicontinuous.
- (A4) There exist $C_i \subset Z$ for $i = 1, \dots, m$ that are nonempty, closed, and convex for which $C = C_1 \cap \dots \cap C_m \neq \emptyset$ and $\{C_1, \dots, C_m\}$ is boundedly regular in the sense that

$$\max_{i=1, \dots, m} d_{C_i}(Tx_k) \rightarrow 0 \quad \implies \quad d_C(Tx_k) \rightarrow 0$$

as $k \rightarrow \infty$ for every bounded sequence $\{x_k\} \subset X$.

Remark 3.3 (assumption (A2)). Recall that the operator T is surjective if and only if T^* is injective and the range of T^* is norm-closed [39, Thm. 4.15]. As a consequence, assumption (A2) is satisfied if T is surjective. In addition, recall that T^* is injective if and only if the kernel of T^* is trivial, i.e., $\ker T^* := \{z \in Z \mid T^*z = 0\} = \{0\}$.

4. Augmented Lagrangian with explicit equality constraints. To develop the augmented Lagrangian portion of our algorithm, we recall that (3.1) is equivalent to the equality-constrained problem

$$(4.1) \quad \min_{x \in X} \{f(x) + I_C(Tx)\} \quad \text{subject to} \quad g(x) = 0,$$

where $I_C(y) = 0$ if $y \in C$ and $I_C(y) = \infty$ if $y \notin C$. Here, $I_C(Tx)$ enforces the constraint $Tx \in C$. Given the constraint decomposition in assumption (A4), we can rewrite I_C as

$$I_C = \sum_{i=1}^m I_{C_i}.$$

As is typically done in augmented Lagrangian methods [38], we replace the indicator functions $I_{C_i}(Tx)$ with the relaxations $\Psi_i(x, \lambda, r)$, where $\Psi_i : X \times Z \times (0, \infty) \rightarrow \mathbb{R}$ is defined as

$$(4.2) \quad \Psi_i(x, \lambda, r) := \sup_{\mu \in Z} \{(\mu, Tx)_Z - I_{C_i}^*(\mu) - \frac{1}{2r} \|\mu - \lambda\|_Z^2\}$$

and $I_{C_i}^*$ is the Fenchel conjugate of I_{C_i} , i.e.,

$$I_{C_i}^*(\mu) := \sup_{z \in C_i} (\mu, z)_Z.$$

The augmented Lagrangian functional is given by

$$(4.3) \quad L(x, \lambda_1, \dots, \lambda_m, r_1, \dots, r_m) := f(x) + \sum_{i=1}^m \Psi_i(x, \lambda_i, r_i),$$

where $\lambda_i \in Z$ and $r_i > 0$ for $i = 1, \dots, m$. The function being maximized in (4.2) is strongly concave and has the unique maximizer

$$(4.4) \quad \Lambda_i(x, \lambda, r) := r((r^{-1}\lambda + Tx) - \mathbf{P}_{C_i}(r^{-1}\lambda + Tx)).$$

Substituting $\Lambda_i(x, \lambda, r)$ into (4.2) and rearranging terms yields the usual augmented Lagrangian penalty function

$$\Psi_i(x, \lambda, r) = \frac{1}{2r} \|\Lambda_i(x, \lambda, r)\|_Z^2 - \frac{1}{2r} \|\lambda\|_Z^2.$$

Motivated by [7, 12], we formulate Algorithm 4.1 using the augmented Lagrangian L .

Remark 4.1 (penalty parameter and multiplier update). The penalty parameter update in Algorithm 4.1 is completely decoupled for the first K_0 iterations. Here, K_0 can be taken arbitrarily large, but finite, e.g., $K_0 = 1000$. This allows each $r_i^{(k)}$ to be calibrated to the scaling associated with the i th constraint. The penalty parameter and multiplier update schemes are adapted from [12, Alg. 2]. In particular, ν_i and

Algorithm 4.1. Multipenalty equality-constrained augmented Lagrangian.

Input: Initial multiplier estimates $\{\lambda_1^{(1)}, \dots, \lambda_m^{(1)}\} \in Z$, positive penalty parameters $\{r_1^{(1)}, \dots, r_m^{(1)}\}$, nonnegative null sequences $\{\delta^{(k)}\}$ and $\{\varepsilon^{(k)}\}$, $K_0 \in \mathbb{N}$, and positive constants $\{\nu_1, \dots, \nu_m\}$, $\{\gamma_1, \dots, \gamma_m\}$ with $\gamma_i < 1/2$, $\{\tau_1^{(0)}, \dots, \tau_m^{(0)}\}$, $\{\theta_1, \dots, \theta_m\}$ with $\theta_i < 1$, $\tau_*, \delta_*, \varepsilon_*$, $\{\eta_1, \dots, \eta_m\}$ with $\eta_i > 1$, $\bar{\eta} > 1$, $\{\alpha_1, \dots, \alpha_m\}$ and $\{\beta_1, \dots, \beta_m\}$. Set $\theta_i^{(1)} = \min\{1/r_i^{(1)}, \theta_i\}$ and $\tau_i^{(1)} = \tau_i^{(0)}(\theta_i^{(1)})^{\alpha_i}$.

- 1: **for** $k = 1, 2, 3, \dots$ **do**
- 2: Compute $(x^{(k)}, \zeta^{(k)}) \in X \times Y^*$ that satisfies $\|g(x^{(k)})\|_Y \leq \delta^{(k)}$ and $\|f'(x^{(k)}) + \sum_i T^* \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}\|_{X^*} \leq \varepsilon^{(k)}$
- 3: **if** $\|g(x^{(k)})\|_Y \leq \delta_*$, $\|f'(x^{(k)}) + \sum_i T^* \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}\|_{X^*} \leq \varepsilon_*$ and $\max_i d_{C_i}(Tx^{(k)}) \leq \tau_*$ **then**
- 4: **return** $x^{(k)}$ as the approximate solution
- 5: **end if**
- 6: **if** $k = K_0 + 1$ **then**
- 7: $\eta_i = \bar{\eta}$ for $i = 1, \dots, m$
- 8: **end if**
- 9: **update** = false
- 10: **if** $k > K_0$ and $\exists i$ such that $\|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z > r_i^{(k)} \tau_i^{(k)}$ **then**
- 11: **update** = true
- 12: **end if**
- 13: **for** $i = 1, \dots, m$ **do**
- 14: **if** $\|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z > r_i^{(k)} \tau_i^{(k)}$ or **update** = true **then**
- 15: $r_i^{(k+1)} = \eta_i r_i^{(k)}$
- 16: $\theta_i^{(k+1)} = \min\{1/r_i^{(k+1)}, \theta_i\}$
- 17: $\tau_i^{(k+1)} = \tau_i^{(0)}(\theta_i^{(k+1)})^{\alpha_i}$
- 18: **else**
- 19: $r_i^{(k+1)} = r_i^{(k)}$
- 20: $\theta_i^{(k+1)} = \min\{1/r_i^{(k+1)}, \theta_i\}$
- 21: $\tau_i^{(k+1)} = \tau_i^{(k)}(\theta_i^{(k+1)})^{\beta_i}$
- 22: **end if**
- 23: **if** $\|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)})\|_Z \leq \nu_i(r_i^{(k+1)})^{\gamma_i}$ **then**
- 24: $\lambda_i^{(k+1)} = \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)})$
- 25: **else**
- 26: $\lambda_i^{(k+1)} = \lambda_i^{(k)}$
- 27: **end if**
- 28: **end for**
- 29: **end for**

γ_i are used to ensure that the i th multiplier update does not grow too rapidly while η_i is used to increase the penalty parameter r_i . After K_0 iterations, Algorithm 4.1 switches schemes and updates all penalty parameters in unison. This penalty update scheme is a safeguard for the case in which the algorithm produces an infinite sequence of iterations, forcing the sequence to accumulate at a feasible point (under certain assumptions), and is typically never active in practice. By decoupling the penalty parameter updates for the first K_0 iterations, Algorithm 4.1 adaptively learns the appropriate constraint scalings, which are generally not known a priori.

Remark 4.2 (subproblem tolerance sequences). In Algorithm 4.1, we update the feasibility tolerances $\tau_i^{(k)}$ for $i = 1, \dots, m$ in an analogous fashion as [12, Alg. 2] using the parameters α_i , β_i , and θ_i . In contrast, the sequences of optimality tolerances $\{\varepsilon^{(k)}\}$ and equality constraint feasibility tolerances $\{\delta^{(k)}\}$ are only required to be nonnegative and converge to zero. A basic choice is

$$\delta^{(k+1)} = \begin{cases} \delta_0 \delta^{(k)} & \text{if } \exists i \text{ such that } r_i^{(k+1)} \neq r_i^{(k)}, \\ \delta_1 \delta^{(k)} & \text{otherwise,} \end{cases} \quad \text{and}$$

$$\varepsilon^{(k+1)} = \begin{cases} \varepsilon_0 \varepsilon^{(k)} & \text{if } \exists i \text{ such that } r_i^{(k+1)} \neq r_i^{(k)}, \\ \varepsilon_1 \varepsilon^{(k)} & \text{otherwise,} \end{cases}$$

for $0 < \delta_1 < \delta_0 < 1$ and $0 < \varepsilon_1 < \varepsilon_0 < 1$. More complicated updating strategies are possible. For example, adapting the tolerance sequence from [12] yields

$$\delta^{(k+1)} = \begin{cases} \delta^{(0)}(\theta^{(k+1)})^{\alpha_\delta} & \text{if } \exists i \text{ such that } r_i^{(k+1)} \neq r_i^{(k)}, \\ \delta^{(k)}(\theta^{(k+1)})^{\beta_\delta} & \text{otherwise,} \end{cases} \quad \text{and}$$

$$\varepsilon^{(k+1)} = \begin{cases} \varepsilon^{(0)}(\theta^{(k+1)})^{\alpha_\varepsilon} & \text{if } \exists i \text{ such that } r_i^{(k+1)} \neq r_i^{(k)}, \\ \varepsilon^{(k)}(\theta^{(k+1)})^{\beta_\varepsilon} & \text{otherwise,} \end{cases}$$

for positive constants $\delta^{(0)}, \varepsilon^{(0)}, \alpha_\delta, \alpha_\varepsilon, \beta_\delta$, and β_ε , where, e.g., $\theta^{(k+1)} = (\max_i r_i^{(k+1)})^{-1}$, whereas adapting the sequences from [7] yields

$$\delta^{(k+1)} = \min\{\delta_0 \delta^{(k)}, \delta_1 \|g(x^{(k)}\|_Y\} \quad \text{and}$$

$$\varepsilon^{(k+1)} = \min\{\varepsilon_0 \varepsilon^{(k)}, \varepsilon_1 \|f'(x^{(k)}) + c'(x^{(k)})^* \zeta^{(k)}\|_{X^*}\}$$

for constants $\delta_0, \delta_1, \varepsilon_0, \varepsilon_1 \in (0, 1)$.

4.1. Properties of the augmented Lagrangian. We note that the first step in Algorithm 4.1 seeks an approximate stationary point of the equality-constrained subproblem

$$(4.5) \quad \min_{x \in X} L_k(x) \quad \text{subject to} \quad g(x) = 0,$$

where $L_k(x) := L(x, \lambda_1^{(k)}, \dots, \lambda_m^{(k)}, r_1^{(k)}, \dots, r_m^{(k)})$. To facilitate the solution of (4.5), we first show that the penalty function $\Psi_i(\cdot, \lambda, r)$ is convex and continuously Fréchet differentiable with Lipschitz continuous gradient.

PROPOSITION 4.3. *For all $\lambda \in Z$ and $r > 0$, the penalty function $\Psi_i(\cdot, \lambda, r)$ is convex and continuous for $i = 1, \dots, m$. Additionally, $\Psi_i(\cdot, \lambda, r)$ is continuously Fréchet differentiable with gradient*

$$(4.6) \quad \nabla_x \Psi_i(x, \lambda, r) = T^* \Lambda_i(x, \lambda, r),$$

which is Lipschitz continuous with modulus $r \|T\|_{\mathcal{L}(X, Z)}^2$.

Proof. Notice that $\Psi_i(\cdot, \lambda, r)$ is the Fenchel conjugate of $I_{C_i}^*(\cdot) + \frac{1}{2r} \|\cdot - \lambda\|_X^2$ composed with T . Therefore, $\Psi_i(\cdot, \lambda, r)$ is equal to the infimal convolution of the conjugates of $I_{C_i}^*$ and $\frac{1}{2r} \|\cdot - \lambda\|_X^2$ composed with T [3, Prop. 13.21(i)], i.e.,

$$\begin{aligned} \Psi_i(x, \lambda, r) &= \inf_{y \in Z} \{I_{C_i}(y) + (\lambda, Tx - y)_Z + \frac{r}{2} \|Tx - y\|_Z^2\} \\ &= \inf_{y \in C_i} \{(\lambda, Tx - y)_Z + \frac{r}{2} \|Tx - y\|_Z^2\}. \end{aligned}$$

Consequently, $\Psi_i(\cdot, \lambda, r)$ is continuous convex and Fréchet differentiable with Lipschitz continuous gradient (cf. Propositions 9.5 and 12.29 and Corollary 9.15 in [3]). \square

COROLLARY 4.4. *For any fixed $\lambda_i \in Z$ and $r_i > 0$ for $i = 1, \dots, m$, the augmented Lagrangian $L(\cdot, \lambda_1, \dots, \lambda_m, r_1, \dots, r_m)$ is (i) weakly lower semicontinuous if f is, (ii) convex if f is, and (iii) continuously Fréchet differentiable if f is. Moreover, in case (iii), the derivative of $L(\cdot, \lambda_1, \dots, \lambda_m, r_1, \dots, r_m)$ is given by*

$$(4.7) \quad L_x(x, \lambda_1, \dots, \lambda_m, r_1, \dots, r_m) = f'(x) + \sum_{i=1}^m T^* \Lambda_i(x, \lambda_i, r_i),$$

and if f' is Lipschitz continuous, then so is $L_x(\cdot, \lambda_1, \dots, \lambda_m, r_1, \dots, r_m)$.

Proof. By Proposition 4.3, $\Psi_i(\cdot, \lambda_i, r_i)$ is convex and continuous and therefore weakly lower semicontinuous. In addition, $\Psi_i(\cdot, \lambda_i, r_i)$ is continuously Fréchet differentiable with Lipschitz gradients. The desired results then follow from the assumed properties of f . \square

5. Convergence theory. In the subsequent results, we consider infinite sequences of iterates generated by Algorithm 4.1 ignoring the stopping conditions, i.e.,

$$(5.1a) \quad \|g(x^{(k)})\|_Y \leq \delta_*,$$

$$(5.1b) \quad \|L_k'(x^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}\|_{X^*} \leq \varepsilon_*,$$

$$(5.1c) \quad \max_{i=1, \dots, m} d_{C_i}(Tx^{(k)}) \leq \tau_*$$

with $\delta_* = \varepsilon_* = \tau_* = 0$. We denote by $\mathbb{P}_i \subseteq \mathbb{N}$ the set of indices k that satisfy

$$(5.2) \quad \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z > r_i^{(k)} \tau_i^{(k)}.$$

We further denote by $\mathbb{M}_i \subseteq \mathbb{N}$ the subsets of indices for which

$$(5.3) \quad \lambda_i^{(k+1)} = \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)})$$

holds. For any set $\mathbb{S} \subset \mathbb{N}$, we denote the complement of \mathbb{S} by $\mathbb{S}^c := \mathbb{N} \setminus \mathbb{S}$ and the cardinality of \mathbb{S} by $|\mathbb{S}|$. The set \mathbb{P}_i encapsulates the iterations for which the penalty parameter for the i th constraint is increased, while the set \mathbb{M}_i encapsulates the iterations for which the multipliers for the i th constraint are changed. We note that $r_i^{(k)} \rightarrow \infty$ if and only if there exists at least one $j = 1, \dots, m$ such that $|\mathbb{P}_j| = \infty$; in this case, all penalty parameters are increased for any iteration $k \in \mathbb{P}_j$ with $k \geq K_0$ (see lines 9 through 22 in Algorithm 4.1). Our first result is technical and relates the satisfaction of the constraints in C_i to the multiplier updates in Algorithm 4.1.

LEMMA 5.1. *Let $x \in X$, $\lambda \in Z$, and $r > 0$ be arbitrary. Then,*

$$(5.4) \quad d_{C_i}(Tx) \leq \frac{1}{r} \|\Lambda_i(x, \lambda, r) - \lambda\|_Z \leq d_{C_i}(Tx) + \frac{1}{r} \|\lambda\|_Z.$$

Proof. We first prove the lower bound. By definition of Λ_i , we have that

$$d_{C_i}(Tx) = \min_{y \in C_i} \|y - Tx\|_Z \leq \|Tx - \mathbf{P}_{C_i}(r^{-1}\lambda + Tx)\|_Z = \frac{1}{r} \|\Lambda_i(x, \lambda, r) - \lambda\|_Z.$$

Similarly, the Lipschitz continuity (with unit modulus) of the projection operator [3, Prop. 4.8] ensures that

$$\begin{aligned} \frac{1}{r} \|\Lambda_i(x, \lambda, r) - \lambda\|_Z &\leq \|Tx - \mathbf{P}_{C_i}(Tx)\|_Z + \|\mathbf{P}_{C_i}(Tx) - \mathbf{P}_{C_i}(r^{-1}\lambda + Tx)\|_Z \\ &\leq d_{C_i}(Tx) + \frac{1}{r} \|\lambda\|_Z \end{aligned}$$

as desired. \square

5.1. Dual convergence. In this subsection, we analyze the sequence of dual variables $\{\lambda_i^{(k)}\}$ generated by Algorithm 4.1. Our first result is motivated by Lemma 4.2 in [12] and shows that the sequence $\{\|\lambda_i^{(k)}\|_Z\}$ does not grow too fast if $r_i^{(k)} \rightarrow \infty$. The second result relates the first to the asymptotic feasibility of the iterates $\{x^{(k)}\}$. Finally, we show that $\{\lambda_i^{(k)}\}$ converges strongly if $|\mathbb{P}_i| < \infty$.

LEMMA 5.2. *Let $\{\lambda_i^{(k)}\}$ be an infinite sequence of multipliers for the i th constraint generated by Algorithm 4.1, ignoring the stopping conditions (5.1). If $r_i^{(k)} \rightarrow \infty$, then*

$$\lim_{k \rightarrow \infty} \frac{1}{(r_i^{(k)})^\alpha} \|\lambda_i^{(k)}\|_Z = 0 \quad \forall \alpha > \gamma_i,$$

where $\gamma_i < 1/2$ is defined in Algorithm 4.1.

Proof. We note that the proof of this fact is similar to the proof of [12, Lem. 4.2] for equality-constrained augmented Lagrangian methods. If $\mathbb{M}_i = \{k_1, k_2, \dots\}$ is finite, then the result clearly holds since $\lambda_i^{(k)}$ is fixed after finitely many iterations. Now suppose that \mathbb{M}_i is infinite. For any $k_j < k \leq k_{j+1}$, we have that $\lambda_i^{(k)} = \lambda_i^{(k_{j+1})}$ and $r_i^{(k)} \geq r_i^{(k_{j+1})}$. Therefore,

$$\frac{1}{(r_i^{(k)})^\alpha} \|\lambda_i^{(k)}\|_Z \leq \frac{1}{(r_i^{(k_{j+1})})^\alpha} \|\lambda_i^{(k_{j+1})}\|_Z \leq \nu_i(r_i^{(k_{j+1})})^{\gamma_i - \alpha}.$$

The upper bound follows from line 23 in Algorithm 4.1. Since $\alpha > \gamma_i$, the right-hand side converges to zero and the desired result follows. \square

Our next lemma builds on Lemma 5.2 and provides equivalent conditions that imply that the sequence $\{x^{(k)}\}$ is asymptotically feasible for the i th constraint.

LEMMA 5.3. *Let $\{x^{(k)}\}$ be an infinite sequence of iterates generated by Algorithm 4.1, ignoring the stopping conditions (5.1), with the associated sequence of multipliers $\{\lambda_i^{(k)}\}$ for the i th constraint. If $r_i^{(k)} \rightarrow \infty$, then the following three conditions are equivalent:*

- (a) $\liminf_{k \rightarrow \infty} d_{C_i}(Tx^{(k)}) = 0$.
- (b) $\liminf_{k \rightarrow \infty} \frac{1}{r_i^{(k)}} \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z = 0$.
- (c) $\liminf_{k \rightarrow \infty} \frac{1}{r_i^{(k)}} \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)})\|_Z = 0$.

These equivalences remain true if the limit inferiors are replaced by limits (or equivalently limit superiors).

Proof. By Lemma 5.1 with $y = x^{(k)}$, $\lambda = \lambda_i^{(k)}$, and $r = r_i^{(k)}$, we have that

$$(5.5) \quad d_{C_i}(Tx^{(k)}) \leq \frac{1}{r_i^{(k)}} \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z \leq d_{C_i}(Tx^{(k)}) + \frac{1}{r_i^{(k)}} \|\lambda_i^{(k)}\|_Z.$$

This and Lemma 5.2 yield (a) \iff (b). The implication (b) \iff (c) follows from Lemma 5.2 and the triangle and reverse triangle inequalities. \square

Before proving our main finite-termination result, we provide situations for which the sequence of multiplier estimates converges strongly. Strong convergence will be useful for later results as it will allow us to demonstrate that weak accumulation points are stationary under certain assumptions.

THEOREM 5.4. *Let $\{x^{(k)}\}$ be an infinite sequence of iterates generated by Algorithm 4.1, ignoring the stopping conditions (5.1), with the associated sequence of multipliers $\{\lambda_i^{(k)}\}$ for the i th constraint. If $|\mathbb{M}_i| < \infty$ or $|\mathbb{P}_j| < \infty$ for all $j = 1, \dots, m$, then the sequence of multipliers $\{\lambda_i^{(k)}\}$ converges strongly to some $\bar{\lambda}_i \in Z$.*

Proof. First note that if $|\mathbb{M}_i| < \infty$, then $\lambda_i^{(k)} = \bar{\lambda}_i$ is constant for all $k > \max \mathbb{M}_i$ and the result follows. Now, consider the case when $|\mathbb{P}_j| < \infty$ for $j = 1, \dots, m$. Let $k' = \max_j \max \mathbb{P}_j + 1$. We will first show that $\{\lambda_i^{(k)}\}$ is Cauchy (and hence converges). Let $\epsilon > 0$ be arbitrary and choose $k_\epsilon \geq k'$ such that

$$\tau_i^{(k_\epsilon)} < \frac{1 - (\theta_i^{(k')})^{\beta_i}}{r_i^{(k')}} \epsilon.$$

Such a k_ϵ exists since $r_i^{(k)} = r_i^{(k')}$ and $\theta_i^{(k)} = \theta_i^{(k')}$ for all $k \geq k'$, and $\{\tau_i^{(k)}\}_{k \geq k'}$ is strictly decreasing to zero. For any $k \geq k_\epsilon$ and any $h \in \mathbb{N}$, we have that

$$\lambda_i^{(k+h)} - \lambda_i^{(k)} = \sum_{j=k}^{k+h-1} \lambda_i^{(j+1)} - \lambda_i^{(j)}.$$

Since $k \geq k_\epsilon$, we have that $\|\lambda_i^{(j+1)} - \lambda_i^{(j)}\|_Z$ either is equal to zero or is bounded above by $r_i^{(j)} \tau_i^{(j)} > 0$. Therefore, the triangle inequality ensures that

$$\|\lambda_i^{(k+h)} - \lambda_i^{(k)}\|_Z \leq \sum_{j=k}^{k+h-1} \|\lambda_i^{(j+1)} - \lambda_i^{(j)}\|_Z \leq \sum_{j=k}^{k+h-1} r_i^{(j)} \tau_i^{(j)} = r_i^{(k')} \tau_i^{(k)} \sum_{j=0}^h (\theta_i^{(k')})^{j \beta_i}.$$

Since $\theta_i^{(k')} < 1$, we have that the sum on the right-hand side of the above inequality converges geometrically and thus

$$\|\lambda_i^{(k+h)} - \lambda_i^{(k)}\|_Z \leq \frac{\tau_i^{(k)} r_i^{(k')}}{1 - (\theta_i^{(k')})^{\beta_i}} < \frac{\tau_i^{(k_\epsilon)} r_i^{(k')}}{1 - (\theta_i^{(k')})^{\beta_i}} < \epsilon.$$

Consequently, $\{\lambda_i^{(k)}\}$ is Cauchy and hence converges strongly to some $\bar{\lambda}_i \in Z$. \square

COROLLARY 5.5. *Consider the setting of Theorem 5.4. If $|\mathbb{P}_j| < \infty$ for $j = 1, \dots, m$, then*

$$\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) \rightarrow \bar{\lambda}_i, \quad i = 1, \dots, m.$$

Proof. The triangle inequality ensures that

$$\begin{aligned} \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \bar{\lambda}_i\|_Z &\leq \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) - \lambda_i^{(k)}\|_Z + \|\bar{\lambda}_i - \lambda_i^{(k)}\|_Z \\ &\leq r_i^{(k)} \tau_i^{(k)} + \|\bar{\lambda}_i - \lambda_i^{(k)}\|_Z \quad \forall k \geq \max_j \max \mathbb{P}_j + 1. \end{aligned}$$

Since $r_i^{(k)}$ is constant for all k sufficiently large and $\tau_i^{(k)} \rightarrow 0$, the result then follows from Theorem 5.4. \square

5.2. Finite termination. In this subsection, we investigate the scenarios for which Algorithm 4.1 terminates in a finite number of iterations. As the following result suggests, it is uncommon for the algorithm to produce infinitely many iterations without satisfying the stopping conditions (5.1). In particular, this result states that Algorithm 4.1 will satisfy the stopping conditions (5.1) after a finite number of iterations if there are infinitely many successful iterations (i.e., $|\mathbb{P}_i^c| = \infty$) or if the multiplier are updated infinitely often (i.e., $|\mathbb{M}_i| = \infty$). In fact, this result shows that the only case for which Algorithm 4.1 may not terminate in finitely many iterations is if $|\mathbb{P}_i^c| < \infty$ and the associated multiplier estimates are only updated a finite number of times, i.e., $|\mathbb{M}_i| < \infty$.

THEOREM 5.6. *Let $\{(x^{(k)}, \zeta^{(k)}, \lambda_i^{(k)}, r_i^{(k)})\}$ be an infinite sequence of iterates generated by Algorithm 4.1, ignoring the stopping conditions (5.1). Then, the sequence satisfies*

$$(5.6) \quad \lim_{k \rightarrow \infty} \|g(x^{(k)})\|_Y = 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} \|L_k'(x^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}\|_{X^*} = 0.$$

Consider arbitrary $i \in \{1, \dots, m\}$. If $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$, then

$$(5.7) \quad \lim_{j \rightarrow \infty} d_{C_i}(Tx^{(k_j)}) = 0, \quad \text{where} \quad \mathbb{P}_i^c \cup \mathbb{M}_i = \{k_j\}_{j=1}^{\infty}.$$

In particular, if $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$, then

$$(5.8) \quad \liminf_{k \rightarrow \infty} d_{C_i}(Tx^{(k)}) = 0,$$

and if either $|\mathbb{P}_i| < \infty$ or $|\mathbb{M}_i^c| < \infty$, then $d_{C_i}(Tx^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$.

Proof. We first note that the tolerance update rules in Algorithm 4.1 ensure that

$$(5.9) \quad \lim_{k \rightarrow \infty} \tau_i^{(k)} = 0, \quad \lim_{k \rightarrow \infty} \delta^{(k)} = 0, \quad \text{and} \quad \lim_{k \rightarrow \infty} \varepsilon^{(k)} = 0.$$

As a result (5.6) holds. Now, let $i \in \{1, \dots, m\}$ be arbitrary. By Lemma 5.1, we have that $d_{C_i}(Tx^{(k)}) \leq \tau_i^{(k)}$ for all $k \in \mathbb{P}_i^c$ and therefore $\{d_{C_i}(Tx^{(k)})\}_{k \in \mathbb{P}_i^c}$ converges to zero if $|\mathbb{P}_i^c| = \infty$. In particular, if $|\mathbb{P}_i| < \infty$, then we have that $d_{C_i}(Tx^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$ since $d_{C_i}(Tx^{(k)}) \leq \tau_i^{(k)}$ for all k sufficiently large. Now, suppose that $|\mathbb{P}_i| = \infty$. The multiplier update rule in Algorithm 4.1 ensures that

$$(5.10) \quad \frac{1}{r_i^{(k)}} \|\Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)})\|_Z \leq \frac{\nu_i}{r_i^{(k)}} (r_i^{(k+1)})^{\gamma_i} \leq \nu_i \eta_i^{\gamma_i} (r_i^{(k)})^{\gamma_i - 1} \quad \forall k \in \mathbb{M}_i.$$

Note that if $k > K_0$, then η_i is replaced by $\bar{\eta}$ in (5.10). Therefore, $\{d_{C_i}(Tx^{(k)})\}_{k \in \mathbb{M}_i}$ converges to zero by Lemma 5.3 if $|\mathbb{M}_i| = \infty$ since $\gamma_i < 1/2$. Consequently, if $|\mathbb{M}_i^c| < \infty$, then $d_{C_i}(Tx^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$ since (5.10) holds for all k sufficiently large. Combining these results, we see that (5.7) holds if $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$. Finally, (5.8) is a direct consequence of (5.7). \square

Theorem 5.6 provides conditions under which Algorithm 4.1 terminates in a finite number of iterations. However, it does not address the asymptotic satisfaction of the first-order optimality conditions (3.2). Our next result demonstrates that the sequence

of iterates generated by Algorithm 4.1 asymptotically satisfies (3.2) as long as (5.8) holds.

PROPOSITION 5.7. *Consider the setting of Theorem 5.6. Then the iterates satisfy*

$$(5.11) \quad \limsup_{k \rightarrow \infty} \langle T^* \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}), y - x^{(k)} \rangle_{X^*, X} \leq 0 \quad \forall y \in T^{-1}(C_i)$$

for $i = 1, \dots, m$.

Proof. The following proof was motivated by the proof of [28, Lem. 5.2]. For the subsequent arguments, it will be convenient to define

$$s^{(k)} = \mathbf{P}_{C_i}((r^{(k)})^{-1} \lambda_i^{(k)} + T x^{(k)}) \quad \text{and} \quad t^{(k)} = \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}).$$

We have that $s^{(k)} \in C_i$ and $t^{(k)} \in N_{C_i}(s^{(k)})$ by [3, Prop. 6.46]. In addition, by [3, Thm. 6.29], we can write

$$T x^{(k)} = \frac{t^{(k)} - \lambda_i^{(k)}}{r_i^{(k)}} + s^{(k)}.$$

Consequently, for any $y \in T^{-1}(C_i)$, we have that

$$(5.12) \quad \begin{aligned} \langle T^* t^{(k)}, y - x^{(k)} \rangle_{X^*, X} &= (t^{(k)}, T y - (r_i^{(k)})^{-1} (t^{(k)} - \lambda_i^{(k)}) - s^{(k)})_Z \\ &\leq \frac{1}{r_i^{(k)}} ((t^{(k)}, \lambda_i^{(k)})_Z - \|t^{(k)}\|_Z^2), \end{aligned}$$

where the upper bound follows from the fact that $t^{(k)} \in N_{C_i}(s^{(k)})$. If $|\mathbb{P}_j| < \infty$ for $j = 1, \dots, m$, then Theorem 5.4 and Corollary 5.5 ensure that the upper bound in (5.12) converges to zero since $\lambda_i^{(k)} \rightarrow \bar{\lambda}_i$ and $t^{(k)} \rightarrow \bar{\lambda}_i$. Now, consider the case when $|\mathbb{P}_i| = \infty$. By maximizing the quadratic form on the right-hand side of the above equation with respect to $t^{(k)}$, we see that

$$\langle T^* t^{(k)}, y - x^{(k)} \rangle_{X^*, X} \leq \frac{1}{4r_i^{(k)}} \|\lambda_i^{(k)}\|_Z^2.$$

After passing to the limit superior, the desired result follows as a consequence of Lemma 5.2 with $\alpha = 1/2 > \gamma_i$. \square

Remark 5.8 (relation to first-order optimality conditions). Let

$$(5.13) \quad e^{(k)} := \sum_{i=1}^m T^* \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}) + (f'(x^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}).$$

By Theorem 5.6, we have that $g(x^{(k)}) \rightarrow 0$ and

$$(5.14) \quad -(f'(x^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}) + e^{(k)} = \sum_{i=1}^m T^* \Lambda_i(x^{(k)}, \lambda_i^{(k)}, r_i^{(k)}),$$

where $e^{(k)} \rightarrow 0$. If $\langle e^{(k)}, x^{(k)} \rangle_{X^*, X} \rightarrow 0$, then Proposition 5.7 ensures that

$$(5.15) \quad \limsup_{k \rightarrow \infty} \langle -(f'(x^{(k)}) + g'(x^{(k)})^* \zeta^{(k)}), y - x^{(k)} \rangle_{X^*, X} \leq 0 \quad \forall y \in T^{-1}(C)$$

and therefore the sequence of iterates $\{(x^{(k)}, \zeta^{(k)})\}$ asymptotically satisfies the first-order optimality conditions (3.2), as long as (5.8) holds.

5.3. Convergence to feasible points and asymptotic stationarity. In this subsection, we show that weak accumulation points of the iterates $\{x^{(k)}\}$ generated by Algorithm 4.1 are nearly feasible. The assumptions required for this result follow from standard assumptions in the convergence theory for SQP and augmented Lagrangian methods [12, 15, 19, 20].

THEOREM 5.9. *Consider the setting of Theorem 5.6 and let assumptions (A1)–(A4) hold. Suppose there exists a weakly converging subsequence $\{x^{(k_j)}\}$ with limit $\bar{x} \in X$ such that*

$$(5.16) \quad \theta^{(k_j)} := \left(\sum_{i=1}^m r_i^{(k_j)} \right)^{-1} \rightarrow 0 \quad \Rightarrow \quad \theta^{(k_j)} (f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}) \rightharpoonup^* 0.$$

If $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$ for all $i = 1, \dots, m$, then $T\bar{x} \in C$. On the other hand, if there is at least one $i \in \{1, \dots, m\}$ for which $|\mathbb{P}_i^c \cup \mathbb{M}_i| < \infty$, then there exists $\bar{t}_i \in (0, 1)$ with $\bar{t}_1 + \dots + \bar{t}_m = 1$ and $\bar{y}_i \in C_i$ such that

$$(5.17) \quad T\bar{x} = \sum_{i=1}^m \bar{t}_i \bar{y}_i.$$

In addition, the subsequence $\{(x^{(k_j)}, \zeta^{(k_j)})\}$ satisfies

$$(5.18) \quad \limsup_{k \rightarrow \infty} \langle -(f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}), y - x^{(k_j)} \rangle_{X^*, X} \leq 0 \quad \forall y \in T^{-1}(C).$$

Proof. Assumption (A3) and Theorem 5.6 ensure that $g(\bar{x}) = 0$. If $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$, then Theorem 5.6 and the weak lower semicontinuity of $d_{C_i} \circ T$ [46, Lem. 1.5] imply that $T\bar{x} \in C_i$. As a result, if $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$ for all $i = 1, \dots, m$, then $T\bar{x} \in C$. Now, assume that there exists at least one i for which $|\mathbb{P}_i^c \cup \mathbb{M}_i| < \infty$. For such i , $|\mathbb{P}_i| = \infty$, which implies that $r_\ell^{(k_j)} \rightarrow \infty$ for all $\ell = 1, \dots, m$. Lemma 5.2 then ensures that $\theta^{(k_j)} \lambda_\ell^{(k_j)} \rightarrow 0$ for $\ell = 1, \dots, m$. Now, by (5.14) and (5.16), we have

$$(5.19) \quad \begin{aligned} & \theta^{(k_j)} \sum_{i=1}^m T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) \\ &= \theta^{(k_j)} (e^{(k_j)} - (f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)})) \rightharpoonup^* 0. \end{aligned}$$

Expanding the left-hand side yields

$$(5.20) \quad \theta^{(k_j)} \sum_{i=1}^m \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) = T x^{(k_j)} + \sum_{i=1}^m \{ \theta^{(k_j)} \lambda_i^{(k_j)} - t_i^{(k_j)} \mathbf{P}_{C_i}(z_i^{(k_j)}) \},$$

where $t_i^{(k_j)} := \theta^{(k_j)} r_i^{(k_j)} > 0$, $t_1^{(k_j)} + \dots + t_m^{(k_j)} = 1$, and $z_i^{(k_j)} := (r_i^{(k_j)})^{-1} \lambda_i^{(k_j)} + T x^{(k_j)}$. Note that for all $k_j > K_0$, we have $t_i^{(k_j)} = t_i^{(K_0)} = \bar{t}_i$. By assumption (A2), we have that $\ker T^* = \{0\}$. Since $\{x^{(k_j)}\}$ converges weakly, it is bounded and hence $\{\mathbf{P}_{C_i}(z_i^{(k_j)})\}$ is also bounded for $i = 1, \dots, m$. Therefore, $\{\mathbf{P}_C(z_i^{(k_j)})\}$ has a weakly converging subsequence (that we do not relabel) with limit $\bar{y}_i \in C_i$ since Z is a Hilbert space and C_i is closed and convex (hence, weakly closed) for $i = 1, \dots, m$. Consequently, the sequence on the left-hand side of (5.19) converges weakly* to $T^*(T\bar{x} - \sum_i \bar{t}_i \bar{y}_i)$. Owing to the uniqueness of weak* limits, we have that $T^*(T\bar{x} - \sum_i \bar{t}_i \bar{y}_i) = 0$ and hence (5.17) holds since $\ker T^* = \{0\}$. Moreover, (5.18) follows from (5.14) and Proposition 5.7 since $\{x^{(k_j)}\}$ is bounded. \square

The next result is a simple consequence of Theorem 5.9 that employs common assumptions from the convergence theory for SQP (cf. [30] for more details) that ensure the results of Theorem 5.9 hold.

COROLLARY 5.10. *Suppose there exists a set $\Omega \subseteq X$ such that $x^{(k)} \in \Omega$ for all k and for which f' and g' are uniformly bounded on Ω . Moreover, assume that $\{\zeta^{(k)}\}$ is bounded. Then, any weak accumulation point of $\{x^{(k)}\}$ satisfies (5.17), and (5.18) holds for any bounded subsequence of $\{x^{(k)}\}$. In particular, Algorithm 4.1 either terminates in a finite number of iterations with an approximate stationary point or produces an infinite sequence $\{x^{(k)}\}$ for which all bounded subsequences satisfy (5.18) and all weak accumulation points of $\{x^{(k)}\}$ satisfy (5.17).*

Remark 5.11 (weak limits and feasibility). A consequence of Corollary 5.10 is that if $\{x^{(k)}\}$ has a weak accumulation point, then the feasible set for optimization problem (3.1) is nonempty. Notably, if X is reflexive and Ω is bounded, then $\{x^{(k)}\}$ has a weakly converging subsequence. The assumption that Ω is bounded is used to prove convergence of the augmented Lagrangian algorithm in [12, Assump. AS2].

Theorem 5.9 does not ensure that weak accumulation points \bar{x} of $\{x^{(k)}\}$ are feasible. It only shows that $T\bar{x}$ is an element of the convex hull of $C_1 \cup \dots \cup C_m$. We conclude this section with some common situations for which $T\bar{x}$ is guaranteed to be feasible. In these cases, Algorithm 4.1 either terminates in a finite number of iterations or generates a sequence $\{x^{(k)}\}$ that satisfies the asymptotic stationarity condition (5.15) and whose weak accumulation points are feasible.

COROLLARY 5.12. *Let the assumptions of Theorem 5.9 hold and suppose one of the following conditions holds:*

- (a) $m = 1$;
- (b) \mathbf{P}_{C_i} is weakly continuous for $i = 1, \dots, m$;
- (c) T is completely continuous;
- (d) $x^{(k_j)}$ converges strongly to \bar{x} ;
- (e) there exists a Banach space X_0 that is compactly embedded in X such that $\{x^{(k_j)}\} \subset X_0$ and $x^{(k_j)} \rightharpoonup \bar{x}$ in X_0 ;
- (f) X is finite dimensional.

Then, $\{x^{(k_j)}\}$ satisfies the asymptotic stationarity condition (5.18) and $T\bar{x} \in C$.

Proof. Case (a) is obvious. For cases (b)–(d), we have that $\bar{y}_i = \mathbf{P}_{C_i}(T\bar{x})$. Therefore, (5.17) shows that $T\bar{x}$ is a fixed point of the map $\sum_i \bar{t}_i \mathbf{P}_{C_i}(\cdot)$ and it follows from [3, Prop. 4.34] that the fixed points of this map are exactly the set C . Moreover, if (e) holds, then the compact embedding of X_0 in X ensures that $x^{(k_j)} \rightarrow \bar{x}$ in X and the result follows from (d). Finally, if (f) holds, then (b), (c), (d), and (e) also hold. \square

Remark 5.13 (second-order optimality conditions). It may be possible to prove strong convergence of $\{x^{(k_j)}\}$ under additional conditions such as second-order optimality conditions. See [9], where this is done for a related augmented Lagrangian algorithm in Banach space.

5.4. Convergence to stationary points. Theorem 5.6 gives sufficient conditions for Algorithm 4.1 to terminate in a finite number of iterations. However, it does not ensure that the sequence of iterates $\{x^{(k)}\}$ satisfies the first-order stationary conditions (3.2). In this subsection, we address this question. The next

results demonstrate the limiting behavior of the sequence of iterates $\{x^{(k)}\}$ generated by Algorithm 4.1. In particular, if a weak accumulation point of $\{x^{(k)}\}$ exists, then that point must be a first-order stationary point. This result requires additional regularity assumptions on the problem data f and g . In particular, we assume the following.

Assumption 5.14 (regularity of derivatives).

(A5) The derivative of the equality-constrained Lagrangian satisfies, If $x_k \rightharpoonup x$ in X , $\zeta_k \rightharpoonup^* \zeta$ in Y^* , and

$$\limsup_{k \rightarrow \infty} \langle f'(x_k) + g'(x_k)^* \zeta_k, x_k - x \rangle_{X^*, X} \leq 0,$$

then for all $y \in T^{-1}(C)$, the following holds:

$$\langle f'(x) + g'(x)^* \zeta, x - y \rangle_{X^*, X} \leq \limsup_{k \rightarrow \infty} \langle f'(x_k) + g'(x_k)^* \zeta_k, x_k - y \rangle_{X^*, X}.$$

A brief discussion of assumption (A5) is in order. If X is finite dimensional, then weak and strong convergence coincide, and therefore the continuity of $f' : X \rightarrow X^*$ and $g' : X \rightarrow \mathcal{L}(X, Y)$ (assumption (A1)) ensures that

$$f'(x_k) + g'(x_k)^* \zeta_k \rightarrow f'(x) + g'(x)^* \zeta \quad \text{and} \quad x_k \rightarrow x.$$

Consequently, (A5) is satisfied. In infinite dimensions, it may require quite strong assumptions to satisfy (A5). Following our next result, we provide assumptions that ensure that assumption (A5) holds.

THEOREM 5.15. *Consider the setting of Theorem 5.6 and let assumptions (A1)–(A5) hold. Let $(\bar{x}, \bar{\zeta}) \in X \times Y^*$ be a weak/weak* accumulation point of $\{(x^{(k)}, \zeta^{(k)})\}$. If $T\bar{x} \in C$, then \bar{x} is a first-order stationary point of (3.1). That is, \bar{x} satisfies (3.2).*

Proof. Let $\{(x^{(k_j)}, \zeta^{(k_j)})\}$ denote a subsequence such that $x^{(k_j)} \rightharpoonup \bar{x}$ and $\zeta^{(k_j)} \rightharpoonup^* \bar{\zeta}$ and suppose $T\bar{x} \in C$. Assumption (A3) ensures that $g(\bar{x}) = 0$. We now prove the first condition in (3.2). Proposition 5.7, the fact that $T\bar{x} \in C$, and the strong convergence of $\{e^{(k_j)}\}$ (see (5.13) for the definition of $e^{(k_j)}$) ensure that

$$\begin{aligned} & \limsup_{k_j \rightarrow \infty} \langle -(f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}), \bar{x} - x^{(k_j)} \rangle_{X^*, X} \\ &= \limsup_{k_j \rightarrow \infty} \langle -(f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}) + e^{(k_j)}, \bar{x} - x^{(k_j)} \rangle_{X^*, X} \\ &= \limsup_{k_j \rightarrow \infty} \sum_{i=1}^m (T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}), \bar{x} - x^{(k_j)})_Z \leq 0. \end{aligned}$$

The result then follows from assumption (A5) and (5.15). \square

The next result provides strong assumptions on the derivatives f' and g' that ensure assumption (A5) holds. In addition, the result employs a constraint qualification to ensure that $\{\zeta^{(k)}\}$ is bounded and hence has a weakly* converging subsequence. These are then used to show that the results of Theorem 5.9 hold. The assumptions used in this result were motivated by [9, sect. 5].

COROLLARY 5.16. *Consider the setting of Theorem 5.6, and let assumptions (A1)–(A4) hold. Moreover, assume that f' and g' satisfy the following assumptions:*

(A6) The derivative $f' : X \rightarrow X^*$ is pseudomonotone, i.e.,

$$\begin{aligned} x_k \rightharpoonup x \quad \text{and} \quad \limsup_{k \rightarrow \infty} \langle f'(x_k), x_k - x \rangle_{X^*, X} \leq 0 \\ \implies \langle f'(x), x - y \rangle_{X^*, X} \leq \liminf_{k \rightarrow \infty} \langle f'(x_k), x_k - y \rangle_{X^*, X} \quad \forall y \in X. \end{aligned}$$

(A7) The Jacobian $g' : X \rightarrow \mathcal{L}(X, Y)$ is sequentially weak-to-strong continuous and $g'(x)^* \in \mathcal{L}(Y^*, X^*)$ is sequentially weak*-to-strong continuous for all $x \in X$. Then assumption (A5) holds. If, in addition,

(A8) for any bounded set $D \subset X$, the set $\{f'(x) \mid x \in D\} \subseteq X^*$ is bounded, and \bar{x} is a weak accumulation point of $\{x^{(k)}\}$ that satisfies the extended Robinson constraint qualification,

$$(5.21) \quad 0 \in \text{int } g'(\bar{x})(T^{-1}(C) - \bar{x}),$$

then, \bar{x} satisfies (3.2) as long as $T\bar{x} \in C$.

Proof. The proof of this result is similar to the proof of [9, Thm. 5.4]. We first show that assumption (A5) holds. Let $x_k \rightharpoonup x$ in X and $\zeta_k \rightharpoonup^* \zeta$ in Y^* . By assumption (A7), we have that $g'(x_k) \rightarrow g'(x)$ and $g'(x)^*$ is sequentially weak*-to-strong continuous. As a result, we have that $g'(x)^*\zeta_k \rightarrow g'(x)^*\zeta$, which yields $g'(x_k)^*\zeta_k \rightarrow g'(x)^*\zeta$. Assumption (A5) then follows from assumption (A6).

Now, suppose (5.21) holds at \bar{x} and let $\{x^{(k_j)}\}$ denote a subsequence of $\{x^{(k)}\}$ that weakly converges to \bar{x} with associated multiplier subsequence $\{\zeta^{(k_j)}\}$. The generalized open mapping theorem [48, Thm. 2.1] ensures that there exists $\rho > 0$ such that

$$B_\rho^Y \subset g'(\bar{x})[(T^{-1}(C) - \bar{x}) \cap B_1^X].$$

Let $y^{(k_j)} \in B_1^Y$ be a sequence of unit vectors satisfying $\langle \zeta^{(k_j)}, y^{(k_j)} \rangle_{Y^*, Y} \geq \frac{1}{2} \|\zeta^{(k_j)}\|_{Y^*}$. As a result of the above inclusion, we have that $-\rho y^{(k_j)} \in B_\rho^Y$ and there exists a bounded sequence $\{v^{(k_j)}\}$ in $T^{-1}(C)$ such that

$$-\rho y^{(k_j)} = g'(\bar{x})(v^{(k_j)} - \bar{x}).$$

Assumption (A7) ensures that $-\rho y^{(k_j)} = g'(x^{(k_j)})(v^{(k_j)} - x^{(k_j)}) + \eta^{(k_j)}$, where $\eta^{(k_j)} \rightarrow 0$. Now, for k_j sufficiently large so that $\|\eta^{(k_j)}\|_Y \leq \frac{\rho}{4}$, we have that

$$\begin{aligned} (5.22) \quad \frac{\rho}{2} \|\zeta^{(k_j)}\|_{Y^*} &\leq \langle \zeta^{(k_j)}, \rho y^{(k_j)} \rangle_{Y^*, Y} \\ &\leq \langle \zeta^{(k_j)}, -g'(x^{(k_j)})(v^{(k_j)} - x^{(k_j)}) \rangle_{Y^*, Y} + \frac{\rho}{4} \|\zeta^{(k_j)}\|_{Y^*}. \end{aligned}$$

With $e^{(k)}$ as defined in (5.14), we can rewrite

$$-g'(x^{(k_j)})^* \zeta^{(k_j)} = f'(x^{(k_j)}) + \sum_{i=1}^m T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) - e^{(k_j)}.$$

Substituting this expression into (5.22) and rearranging terms gives

$$\begin{aligned} (5.23) \quad \frac{\rho}{4} \|\zeta^{(k_j)}\|_{Y^*} &\leq \langle f'(x^{(k_j)}), v^{(k_j)} - x^{(k_j)} \rangle_{X^*, X} \\ &\quad + \sum_{i=1}^m \langle T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) - e^{(k_j)}, v^{(k_j)} - x^{(k_j)} \rangle_{X^*, X}. \end{aligned}$$

Notice that Proposition 5.7 ensures that there exists a sequence $\{n^{(k_j)}\} \in (0, \infty)$ such that $n^{(k_j)} \searrow 0$ and

$$\sum_{i=1}^m \langle T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}), v^{(k_j)} - x^{(k_j)} \rangle_{X^*, X} \leq n^{(k_j)}$$

since $v^{k_j} \in T^{-1}(C)$. Therefore, the right-hand side of (5.23), and hence $\{\zeta^{(k_j)}\}$, is bounded by (A8) and the fact that $e^{(k)} \rightarrow 0$. Since $\{\zeta^{(k_j)}\}$ is bounded, it has a weakly* convergent subsequence by the Banach–Alaoglu theorem [16, Thm. 5.18]. The desired result then follows from Theorem 5.15. \square

6. Algorithmic extensions. We now present extensions to Algorithm 4.1. Our first extension allows us to separately penalize multiple individual constraints. The second extension allows us to handle nonlinear constraint operators T in (3.1).

6.1. Finitely many linear constraints. In this subsection, we consider the common setting in which there are finitely many constraints of the form $T_i x \in \widehat{C}_i$, where $T_i \in \mathcal{L}(X, Z_i)$ and $\widehat{C}_i \in Z_i$ is nonempty, closed, and convex for $i = 1, \dots, m$. Here, Z_i denotes a real Hilbert space for $i = 1, \dots, m$. We first note that this setting can be stated in the more general setting of (3.1) by defining

$$(6.1) \quad Z := Z_1 \oplus \dots \oplus Z_m, \quad Tx := (T_1 x, \dots, T_m x), \quad \text{and} \quad C := \widehat{C}_1 \times \dots \times \widehat{C}_m.$$

Consequently, it is straightforward to apply Algorithm 4.1 with T and C defined above, either by treating $Tx \in C$ as a single constraint or by handling $T_i x \in \widehat{C}_i$ individually. In particular, define

$$C_i := \left(\prod_{j=1}^{i-1} Z_j \right) \times \widehat{C}_i \times \left(\prod_{j=i+1}^m Z_j \right), \quad i = 1, \dots, m,$$

where \prod denotes the Cartesian product and the first and last products are void if $i = 1$ and $i = m$, respectively. It is clear from the definition of C and C_i that assumption (A4) holds. Moreover, using these definitions, we see that

$$I_C(Tx) = \sum_{i=1}^m I_{C_i}(Tx) = \sum_{i=1}^m I_{\widehat{C}_i}(T_i x)$$

and hence, Ψ_i and Λ_i only depend on $T_i x$ and \widehat{C}_i for $i = 1, \dots, m$. Unfortunately, T as defined above need not satisfy assumption (A2). In particular, the adjoint operator T^* , which is given by $T^* z = T_1^* z_1 + \dots + T_m^* z_m$ for $z = (z_1, \dots, z_m)$ and $z_i \in Z_i$ for $i = 1, \dots, m$, need not be injective even if T_i^* is for $i = 1, \dots, m$. Fortunately, there are practical situations where T^* is in fact injective. One such situation is when the optimization space X is a direct sum of Banach spaces. This is often the case for optimal control problems, in which case X is typically composed of the state and control spaces.

Suppose there exist real Banach spaces X_i and operators $\widehat{T}_i \in \mathcal{L}(X_i, Z_i)$ satisfying (A2), $i = 1, \dots, m$, for which

$$(6.2) \quad X = X_1 \oplus \dots \oplus X_m \quad \text{and} \quad T_i x = \widehat{T}_i x_i \quad \text{for} \quad i = 1, \dots, m.$$

In this case, $T = \widehat{T}_1 \oplus \cdots \oplus \widehat{T}_m$ satisfies assumption (A2). Therefore, Theorem 5.9 applies directly to problems of this type. In fact, we can show that $T\bar{x} \in C$.

THEOREM 6.1. *Consider the setting of Theorem 5.6 and let assumptions (A1)–(A4) hold. Let X , Z , T , and C be defined as in (6.1) and (6.2), and suppose there exists a weakly converging subsequence $\{x^{(k_j)}\}$ with limit $\bar{x} \in X$ such that*

$$(6.3) \quad r_i^{(k_j)} \rightarrow \infty \quad \Rightarrow \quad \frac{1}{r_i^{(k_j)}} (f_{x_i}(x^{(k_j)}) + g_{x_i}(x^{(k_j)})^* \zeta^{(k_j)}) \rightharpoonup^* 0$$

for $i = 1, \dots, m$. Here, f_{x_i} and g_{x_i} denote the partial derivatives of f and g with respect to x_i , $i = 1, \dots, m$, respectively. Then, \bar{x} satisfies $T\bar{x} \in C$.

Proof. Clearly if $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$, then $\widehat{T}_i \bar{x}_i \in \widehat{C}_i$ for $i = 1, \dots, m$. We assume that at least one $i = 1, \dots, m$ satisfies $|\mathbb{P}_i| = \infty$. Using the product structure of the problem, (6.3) ensures that

$$\frac{1}{r_i^{(k_j)}} T_i^* \Lambda_i(x_i^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) = \frac{1}{r_i^{(k_j)}} \left(e_i^{(k_j)} - (f_{x_i}(x^{(k_j)}) + g_{x_i}(x^{(k_j)})^* \zeta^{(k_j)}) \right) \rightharpoonup^* 0,$$

for which the left-hand side can be expanded as

$$\frac{1}{r_i^{(k_j)}} T_i^* \Lambda_i(x_i^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) = T_i^* (z_i^{(k_j)} - \mathbf{P}_{\widehat{C}_i}(z_i^{(k_j)})),$$

where $z_i^{(k_j)} = (r_i^{(k_j)})^{-1} \lambda_i^{(k_j)} + \widehat{T}_i x_i^{(k_j)}$. Since $(r_i^{(k_j)})^{-1} \lambda_i^{(k_j)} \rightarrow 0$ by Lemma 5.2 and $\widehat{T}_i x_i^{(k_j)} \rightharpoonup \widehat{T}_i \bar{x}_i$, the sequence $\{z_i^{(k_j)}\}$ is bounded and hence so is $\{\mathbf{P}_{\widehat{C}_i}(z_i^{(k_j)})\}$. Furthermore, since Z_i is a Hilbert space, $\{\mathbf{P}_{\widehat{C}_i}(z_i^{(k_j)})\}$ has a weakly converging subsequence (that we do not relabel) with limit $\bar{y}_i \in \widehat{C}_i$. The injectivity of \widehat{T}_i^* and the uniqueness of weak* limits then ensure that $\widehat{T}_i \bar{x}_i = \bar{y}_i \in \widehat{C}_i$ for $i = 1, \dots, m$. \square

6.2. Nonlinear constraints. We now consider the addition of the nonlinear constraint $T_0(x) \in C_0$ to (3.1). Here, $T_0 : X \rightarrow Z_0$, where Z_0 is a real Hilbert space and $C_0 \subseteq Z_0$ is nonempty, closed, and convex. We define the penalty function Ψ_0 and the multiplier update functions Λ_0 analogously to Ψ_i and Λ_i for $i = 1, \dots, m$. Additionally, we define $r_0^{(k)}$, $\lambda_0^{(k)}$, \mathbb{P}_0 , and \mathbb{M}_0 analogously. With these definitions, the results in section 5.1 and Theorem 5.6 hold with no modifications. In the next theorem, we demonstrate how the result of Theorem 5.9 changes in this setting.

THEOREM 6.2. *Consider the setting of Theorem 5.9. Let $\bar{x} \in X$ be a weak accumulation point of $\{x^{(k)}\}$ with associated subsequence $\{x^{(k_j)}\}$ and suppose that there exists $\alpha \in (0, 1)$ such that*

$$(6.4) \quad \theta^{(k_j)} := \left(\sum_{i=0}^m r_i^{(k_j)} \right)^{-1} \rightarrow 0 \quad \Rightarrow \quad (\theta^{(k_j)})^\alpha (f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}) \rightharpoonup^* 0.$$

Moreover, assume that

(A9) T_0 is completely continuous and continuously Fréchet differentiable, and the derivative T_0' satisfies

$$x_k \rightharpoonup x, \quad y_k \rightharpoonup y \quad \text{in } X \quad \Rightarrow \quad T_0'(x_k)y_k \rightharpoonup T_0'(x)y.$$

If $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$ for $i = 0, \dots, m$, then $T\bar{x} \in C$ and $T_0(\bar{x}) \in C_0$. On the other hand, if there exists at least one $i = 0, \dots, m$ for which $|\mathbb{P}_i^c \cup \mathbb{M}_i| < \infty$, then \bar{x} satisfies

$$(6.5) \quad \langle T_0'(\bar{x})^*(T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x}))), y - \bar{x} \rangle_{X^*, X} \geq 0 \quad \forall y \in T^{-1}(C).$$

Finally, if \bar{x} satisfies the extended Robinson constraint qualification

$$(6.6) \quad 0 \in \text{int} \{T_0(\bar{x}) + T_0'(\bar{x})(T^{-1}(C) - \bar{x}) - C_2\},$$

then $T_0(\bar{x}) \in C_0$ and (5.17) holds.

Proof. If $|\mathbb{P}_i^c \cup \mathbb{M}_i| = \infty$ for $i = 0, \dots, m$, then the feasibility follows from the arguments in the proof of Theorem 5.9. To prove the remaining results, suppose that $|\mathbb{P}_i^c \cup \mathbb{M}_i| < \infty$ for at least one $i = 0, \dots, m$. In this case, (6.4) ensures that

$$(6.7) \quad \begin{aligned} & \theta^{(k_j)} \sum_{i=1}^m T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}) + \theta^{(k_j)} T_0'(x^{(k_j)})^* \Lambda_0(x^{(k_j)}, \lambda_0^{(k_j)}, r_0^{(k_j)}) \\ &= \theta^{(k_j)} \left(e^{(k_j)} - (f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}) \right) \rightharpoonup^* 0 \end{aligned}$$

and assumption (A9) and Lemma 5.2 ensure that the left-hand side of (6.7) weakly* converges to

$$(6.8) \quad \sum_{i=1}^m \bar{t}_i T^* (T\bar{x} - \bar{y}_i) + \bar{t}_0 T_0'(\bar{x})^* (T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x}))) = 0$$

for $\bar{y}_i \in C_i$, $i = 1, \dots, m$, as in Theorem 5.9. Assumption (A9) further ensures that

$$\begin{aligned} & \lim_{k_j \rightarrow \infty} \theta^{(k_j)} \langle T_0'(x^{(k_j)})^* \Lambda_0(x^{(k_j)}, \lambda_0^{(k_j)}, r_0^{(k_j)}), y - x^{(k_j)} \rangle_{X^*, X} \\ &= \bar{t}_0 \langle T_0'(\bar{x})^* (T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x}))), y - \bar{x} \rangle_{X^*, X} \quad \forall y \in X \end{aligned}$$

and therefore Proposition 5.7 applied to the $i = 1, \dots, m$ constraints combined with (6.4) and (6.7) implies (6.5). In particular, $(\theta^{(k_j)})^{1-\alpha}(y - x^{(k_j)}) \rightarrow 0$ since $\{x^{(k_j)}\}$ converges weakly (and hence is bounded) and

$$\begin{aligned} & -\bar{t}_0 \langle T_0'(\bar{x})^* (T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x}))), y - \bar{x} \rangle_{X^*, X} \\ &= \lim_{k_j \rightarrow \infty} \theta^{(k_j)} \left\{ -\langle T_0'(x^{(k_j)})^* \Lambda_0(x^{(k_j)}, \lambda_0^{(k_j)}, r_0^{(k_j)}), y - x^{(k_j)} \rangle_{X^*, X} \right. \\ & \quad \left. + \langle e^{(k_j)} - (f'(x^{(k_j)}) + g'(x^{(k_j)})^* \zeta^{(k_j)}), y - x^{(k_j)} \rangle_{X^*, X} \right\} \\ & \leq \limsup_{k_j \rightarrow \infty} \theta^{(k_j)} \sum_{i=1}^m \langle T^* \Lambda_i(x^{(k_j)}, \lambda_i^{(k_j)}, r_i^{(k_j)}), y - x^{(k_j)} \rangle_{X^*, X} \leq 0 \quad \forall y \in T^{-1}(C). \end{aligned}$$

To conclude, suppose that \bar{x} satisfies (6.6); then there exists $\rho > 0$ such that $B_\rho^{Z_0} \subseteq T_0(\bar{x}) + T_0'(\bar{x})(T^{-1}(C) - \bar{x}) - C_0$. In particular, for any $z \in B_\rho^{Z_0}$, there exists $y \in T^{-1}(C)$ and $c \in C_0$ such that $z = T_0(\bar{x}) + T_0'(\bar{x})(y - \bar{x}) - c$. Therefore,

$$\begin{aligned} (T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x})), z)_{Z_0} &= \langle T_0'(\bar{x})^*(T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x}))), y - \bar{x} \rangle_{X^*, X} \\ & \quad + (T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x})), T_0(\bar{x}) - c)_{Z_0}. \end{aligned}$$

The first term on the right-hand side is nonnegative by the above arguments and the second term is nonnegative by [3, Thm. 3.14]. Since this holds for all $z \in B_\rho^{Z_0}$, we have that $T_0(\bar{x}) - \mathbf{P}_{C_0}(T_0(\bar{x})) = 0$ and consequently $T_0(\bar{x}) \in C_0$. Combining this with (6.8) and the injectivity of T^* shows that (5.17) holds. \square

Remark 6.3 (optimality of \bar{x}). Note that if $T\bar{x} \in C$, then the variational inequality (6.5) is the first-order optimality condition for the optimization problem

$$\min_{Tx \in C} d_{C_0}(T_0(x))^2.$$

7. Solution of the subproblem. An important motivation for the ALESQP method is to enable iterative, and therefore inexact, solution of linear systems involving the discretizations of $g'(x)$. Therefore, a good choice for the solution of the augmented Lagrangian subproblem (4.5) is the inexact, matrix-free trust-region SQP method [19, 30]. To provide context for some modifications related to its use with the augmented Lagrangian, we give a short summary of the method. For this, we assume that X and Y are Hilbert spaces and that f and g have Lipschitz continuous derivatives. We define the SQP Lagrangian $\mathcal{L} : X \times Y^* \rightarrow \mathbb{R}$ for (4.5), which includes the augmented Lagrangian,

$$\mathcal{L}(x, \zeta) := L(x) + \langle \zeta, g(x) \rangle_{Y^*, Y},$$

where $L(x) := L_k(x)$, and k denotes the k th augmented Lagrangian iteration. The SQP method [19] extends the composite-step approach of [32] to rigorously handle inexact linear system solves. In the context of (4.5), the method comprises the following steps at its j th iteration. We start with an iterate x_j , the corresponding Lagrange multiplier ζ_j , a trust-region radius Δ_j , and a self-adjoint approximation of $\nabla_{xx}\mathcal{L}(x_j, \zeta_j)$, denoted by $H_j = H(x_j, \zeta_j)$, with $H \in \mathcal{L}(X, X)$. First, to reduce the linear infeasibility, $\|g'(x_j)n + g(x_j)\|_Y$, we approximately solve the *quasi-normal subproblem*,

$$(7.1) \quad \min_{n \in X} \|g'(x_j)n + g(x_j)\|_Y^2 \quad \text{subject to} \quad \|n\|_X \leq 0.8\Delta_j,$$

using Powell's dogleg method [35], where we compute the second-order (Newton) step by iteratively solving an *augmented system*, subject to the stopping condition provided in [30, eq. 34]. Second, given a solution n_j of (7.1), to improve optimality we solve the *tangential subproblem*,

$$(7.2) \quad \begin{aligned} & \min_{\tilde{t} \in X} \frac{1}{2}(H_j \tilde{t}, \tilde{t})_X + (\widetilde{W}_j(\nabla_x \mathcal{L}(x_j, \zeta_j) + H_j n_j), \tilde{t})_X \\ & \text{subject to } \tilde{t} \in \text{Range}(\widetilde{W}_j), \quad \|\tilde{t} + n_j\|_X \leq \Delta_j, \end{aligned}$$

using the projected conjugate gradient (CG) method with Steihaug–Toint termination criteria [40]. In (7.2) the linear operator $\widetilde{W}_j \in \mathcal{L}(X, X)$ represents an approximate projection onto the null-space of $g'(x_j)$, $\ker(g'(x_j))$. Its action on a vector is given by the solution of another augmented system, with the stopping conditions [30, eq. 37 and 39]. Third, to ensure that the trial step remains sufficiently close to $\ker(g'(x_j))$, i.e., maintains linearized feasibility, we additionally project the solution \tilde{t}_j of (7.2) onto $\ker(g'(x_j))$ to compute the *tangential step* t_j . To accomplish this, we solve an augmented system with the stopping condition [30, eq. 41]. This yields the trial step $s_j = n_j + t_j$. Fourth, we compute the Lagrange multipliers ζ_{j+1} by solving

another augmented system, with the stopping conditions [30, eq. 43]. Finally, we apply trust-region acceptance and update criteria; see [30, Alg. 4, Steps 3 and 4].

The aforementioned augmented systems are optimality systems of the form

$$(7.3) \quad \begin{pmatrix} R_{X,X^*} & g'(x_j)^* \\ g'(x_j) & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} + \begin{pmatrix} e_1 \\ e_2 \end{pmatrix},$$

where $R_{X,X^*} \in \mathcal{L}(X, X^*)$ is the inverse Riesz map [18]. If the system (7.3) is solved directly, the residual $(e_1 \ e_2)^\top$ is ignored. If it is solved iteratively, the size of $(e_1 \ e_2)^\top$ can be controlled using stopping conditions of the form

$$\|e_1\|_{X^*} + \|e_2\|_Y \leq \mathcal{T}(\|y_1\|_X, \|b_1\|_{X^*}, \|b_2\|_Y, \Delta_j, \tau_{\text{nom}}),$$

where $\mathcal{T} : \mathbb{R}_+^5 \rightarrow \mathbb{R}_+$ is a tolerance function specific to each step, as referenced above, and $\tau_{\text{nom}} > 0$ is a chosen nominal tolerance. The SQP method adjusts linear system tolerances based on its progress, in order to ensure global convergence under standard assumptions. A discussion of the theoretical assumptions, the nominal tolerance choice, the Riesz map, and some modifications to the augmented system (7.3) follows.

Function-space setting. The SQP algorithm [19] requires that X and Y be Hilbert spaces. In section 8 all numerical examples satisfy these assumptions, justifying the application of the algorithm. However, our augmented Lagrangian algorithm and the corresponding convergence theory are developed in the more general setting of Banach spaces. In order to apply ALESQP in Banach space, extensions to the SQP algorithm are necessary. For instance, different notions of Cauchy points and Cauchy decrease conditions are needed (see [13, sect. 8.3.2]), projections onto $\ker(g'(x))$ as discussed previously do not apply, the objective function in (7.2) must be modified, etc. The required extensions, while plausible, are beyond the scope of this paper.

Lipschitz continuous derivatives. In [19] it is assumed that the functions L and g are twice continuously differentiable. This is an appropriate assumption for all numerical examples in section 8 *in the absence* of the constraints $Tx \in C$, i.e., when $L = f$. Once the constraints are included, the constraint penalty terms in L render L' Lipschitz continuous; see Corollary 4.4. The proof of Theorem 3.5 in [19] is easily extended to handle \mathcal{L} with Lipschitz continuous derivatives. Specifically, the second-order Taylor expansion used on page 1537 of [19] can be replaced with the first-order expansion, followed by the use of Lipschitz continuity of \mathcal{L}' ; see the Assumption AW.1c for the composite-step algorithm analyzed in [13], and Theorem 3.1.4 in [13].

Nominal linear solver stopping tolerance. The theory in [19] permits an arbitrary choice of the nominal tolerance $\tau_{\text{nom}} > 0$. For good numerical performance, we choose $\tau_{\text{nom}} = \min\{\sqrt{\delta^{(k)}}, \sqrt{\epsilon^{(k)}}\}$. The same value is used for all augmented system solves, i.e., the nominal tolerances $\tau^{qn}, \tau^{pg}, \tau^{proj}, \tau^{tang}$, and τ^{lmh} from [19, 30].

Implementation of the Riesz maps. The SQP algorithm is posed in Hilbert space and therefore naturally supports the use of Riesz maps, such as $R_{X^*,X}$. However, in large-scale applications the Riesz map may require an iterative solution of additional linear systems—nested within the iterative augmented system solve, or in other components of the SQP algorithm. Inexact or variable Riesz maps are not supported by [19]. To circumvent this challenge, in section 8 we use diagonally weighted inner products, which are exact to within machine precision.

Preconditioning of the projected CG method. In certain applications we can accelerate the projected CG iteration for the solution of (7.2) by replacing the augmented system solve that yields the constraint null-space projection with the solution of a

related linear system. Motivated by a comment about the “perfect preconditioner” for projected CG [17, p. 1381], we solve a system of the form

$$(7.4) \quad \begin{pmatrix} B(x_j) + T^* \left(\sum_{i=1}^m r_i^{(k)} (R_{Z,Z^*} - D_{ij}) \right) T & g'(x_j)^* \\ g'(x_j) & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ 0 \end{pmatrix} + \begin{pmatrix} e_1 \\ e_2 \end{pmatrix},$$

where $B(x_j) \in \mathcal{L}(X, X^*)$ approximates $f''(x_j)$ and D_{ij} denotes the Newton derivative [11, Def. 2.1] of $\mathbf{P}_{C_i}((r_i^{(k)})^{-1} \lambda_i^{(k)} + T x_j)$. Our assumptions for the use of this system are as follows: (i) f is twice continuously differentiable; (ii) X is a Hilbert space and B is nonnegative; and (iii) $(e_1 \ e_2)^\top = (0 \ 0)^\top$, i.e., only a direct solve is permitted. It is possible to relax the third assumption and allow iterative solves of (7.4). However, this leads to two challenges, namely the derivation of stopping conditions for the iterative solve to replace conditions [30, eq. 37 and 39] and, more importantly, the question of efficient preconditioning of (7.4). Both are beyond the scope of this paper, which is why we enforce the third assumption whenever (7.4) is used in section 8.

8. Applications. In this section, we demonstrate Algorithm 4.1 on three infinite-dimensional optimization problems. Our first problem computes a probability density function (pdf) by maximizing the Rényi entropy. The second and third problems are optimization problems constrained by PDEs. Throughout, $\Omega \subset \mathbb{R}^d$ is Lipschitz with $d = 1, 2$, and $L^p(\Omega)$, $p \in [1, \infty]$, denotes the usual p -order Lebesgue space. Moreover, $L_+^p(\Omega)$ will denote the subset of nonnegative $L^p(\Omega)$ functions. We denote by $\partial\Omega$ the boundary of Ω and by $W^{1,s}(\Omega)$ and $H^1(\Omega) := W^{1,2}(\Omega)$ the Sobolev spaces of weakly differentiable functions [1]. Furthermore, we denote by $W_0^{1,s}(\Omega)$ the subspace of $W^{1,s}(\Omega)$ -functions that are zero on the boundary in the trace sense, and $H_0^1(\Omega) := W_0^{1,2}(\Omega)$. All examples are discretized using continuous piecewise linear finite elements on regular simplicial meshes. We use diagonal Riesz map discretizations in all components of ALESQP, associated with the function spaces X , Y , and Z . In particular, we use the lumped mass matrix for both $L^2(\Omega)$ and $H_0^1(\Omega)$.

We choose the following parameters for Algorithm 4.1:

- (a) zero initial guesses throughout, i.e., $x^{(0)} = 0$, $\zeta^{(0)} = 0$, and $\lambda_i^{(1)} = 0$, for $i = 0, \dots, m$, with the exception of the initial guess $x^{(0)} = 1$ for the Rényi entropy example (due to the presence of the log function);
- (b) initial SQP subproblem stopping tolerances

$$\delta^{(0)} = \epsilon^{(0)} = \max\{10^{-3} \|L_k'(x^{(0)}) + g'(x^{(0)})^* \zeta^{(0)}\|_{X^*}, 10^{-6}\};$$

- (c) the basic tolerance update (see Remark 4.2), with reduction factors $\delta_0 = 0.25$, $\delta_1 = 0.9$, $\epsilon_0 = 0.25$, and $\epsilon_1 = 0.9$;
- (d) the g -feasibility, optimality, and T - feasibility stopping tolerances $\delta_* = \epsilon_* = \tau_* = 10^{-6}$, respectively;
- (e) update factors $\eta_i = 5$, for $i = 0, 1, \dots, m$, for the augmented Lagrangian penalties; and
- (f) $\bar{\eta} = 5$, $K_0 = 10^3$, $\theta_i = 0.1$, $\alpha_i = 0.1$, $\beta_i = 0.9$, $\tau_i^{(0)} = 1$, $\nu_i = 10^6$, $\gamma_i = 0.49$ for $i = 0, 1, \dots, m$.

Initial augmented Lagrangian penalty parameters. As in all augmented Lagrangian methods, the choice of the initial penalty parameters is important for good performance, and ALESQP is no exception. We use two general guidelines when choosing the initial parameters. First, they should be chosen as large as possible, without detriment to the convergence of the SQP subproblem solver. A conservative choice is $r_i^{(1)} = 10$ for $i = 0, 1, \dots, m$. This is the default choice in ALESQP. Second, they

should be chosen so that all terms comprising the augmented Lagrangian functional are well balanced. In our first example, the inequality constraint scaling is such that the augmented Lagrangian terms are well balanced, and we can use the default penalty parameter choice. In the second and third examples, the problem structure—specifically the splitting of the variables into states and controls, combined with the PDE nature of the equality constraint linking the states and controls—dictates a more subtle choice, described in more detail in section 8.2.

In the presented results, AL denotes the total number of augmented Lagrangian iterations, SQP the total number of SQP iterations, CG the total number of CG iterations, normg the equality constraint violation $\|g(\bar{x})\|_Y$, grad-lag the norm of the gradient of the subproblem Lagrangian $\|L_k'(\bar{x}) + g'(\bar{x})^* \bar{\zeta}\|_{X^*}$, and feas the constraint violation $\max_i d_{C_i}(T\bar{x})$. We implemented the entire ALESQP framework in MATLAB (R2019a) and studied its performance using a single core of a 2.9 GHz Intel Core i9 processor and 32 GB of RAM. The problem instances studied here range in size from 4,225 to 524,801 optimization variables.

8.1. Maximum entropy. The purpose of this example is to demonstrate mesh independent performance using direct and iterative linear system solves. Our maximum entropy problem seeks a pdf, x , that satisfies certain moment constraints. Let $\Omega = [0, 1]^2$, $X = L^p(\Omega)$, $Y = \mathbb{R}^3$, $Z = L^2(\Omega)$, $Z_0 = \mathbb{R}$, $C = L_+^2(\Omega)$, and $C_0 = [0, 1]$. We solve

$$(8.1a) \quad \min_{X \in X} \left\{ f(x) := \frac{1}{p-1} \log \left(\int_{\Omega} x(\omega)^p d\omega \right) \right\}$$

$$(8.1b) \quad \text{subject to} \quad Tx := x \geq 0 \text{ a.e.},$$

$$(8.1c) \quad g_1(x) := \int_{\Omega} x(\omega) d\omega - 1 = 0,$$

$$(8.1d) \quad g_2(x) := \int_{\Omega} x(\omega) \omega d\omega - \mu = 0,$$

$$(8.1e) \quad T_0(x) := \sigma^{-1} \det \left(\int_{\Omega} x(\omega) (\omega - \mu) (\omega - \mu)^{\top} d\omega \right) \leq 1,$$

where $\sigma > 0$ and $\mu \in \mathbb{R}^2$ are given, $g(x) = (g_1(x), g_2(x))$, and the objective function is the negative p -order Rényi entropy [42] with $p = 2.5$. Constraints (8.1b) and (8.1c) ensure that x is a pdf, (8.1d) ensures that the expected value associated with x is μ , and (8.1e) ensures that the generalized variance [45] associated with x is smaller than σ . A straightforward computation shows that T_0 satisfies assumption (A9) and therefore Theorem 6.2 applies. We use the problem data

$$\mu = (0.45, 0.45) \quad \text{and} \quad \sigma = \frac{1}{2} \det \left(\int_{\Omega} (\omega - \mu) (\omega - \mu)^{\top} d\omega \right) \approx 0.00368,$$

where the latter is chosen so the generalized variance associated with the optimal pdf is less than half of the generalized variance associated with the uniform density.

For our numerical results, we use the default initial penalty parameters, $r_0^{(1)} = 10$ and $r_1^{(1)} = 10$, because the constraint values at the initial guess are well balanced. In particular, $\|Tx^{(0)}\|_Z = 1$ and $|T_0(x^{(0)})| = 2$. Table 1 documents ALESQP performance as the problem size grows, using direct solutions of the augmented systems

TABLE 1

Maximum entropy, direct solution of (7.3). ALESQP performance for varying spatial discretization (Mesh). The AL, SQP, and CG iteration numbers are nearly mesh independent.

Mesh	AL	SQP	CG	normg	grad-lag	feas
64×64	8	44	249	1.36e-16	8.32e-08	2.06e-07
128×128	7	42	239	7.85e-17	1.61e-07	4.60e-07
256×256	7	47	246	6.23e-16	3.47e-07	8.98e-07
512×512	6	43	261	5.27e-16	9.35e-07	4.37e-07

TABLE 2

Maximum entropy, iterative solution of (7.3). ALESQP performance for varying spatial discretization (Mesh). The AL, SQP, and CG iteration numbers are nearly mesh independent. The tot.aug column gives the total number of MINRES iterations in augmented system solves, for the entire run of ALESQP. The avg.aug column gives the average number of MINRES iterations per augmented system solve. The tot.aug and avg.aug iterations vary little as the mesh is refined.

Mesh	AL	SQP	CG	normg	grad-lag	feas	tot.aug	avg.aug
64×64	8	42	222	7.78e-14	2.00e-07	5.05e-07	2969	7.9
128×128	9	47	207	4.12e-15	3.01e-07	4.21e-08	2859	7.6
256×256	10	54	234	3.24e-16	8.72e-07	3.66e-08	3217	7.5
512×512	8	57	273	5.27e-16	9.57e-07	1.63e-07	3516	7.4

(7.3). We observe nearly mesh-independent iteration numbers for the augmented Lagrangian loop and all its iterative components. We note that for this example the penalty parameters do not increase; e.g., for the 128×128 mesh the final values are $r_0^{(7)} = 10$ and $r_1^{(7)} = 10$. Table 2 documents ALESQP performance with iterative augmented system solves, where we have used unpreconditioned MINRES [33] to solve (7.3). Again, we observe nearly mesh-independent iteration numbers for the augmented Lagrangian loop and all its iterative components. Most notably, the total number of MINRES iterations is around 3,000, and it does not change significantly as the mesh is refined. In other words, we have demonstrated discretization-independent algorithmic performance of a fully matrix-free framework on an infinite-dimensional optimization problem with nonlinear inequality constraints. Finally, we note that the solution time for the matrix-free approach increases linearly with problem size, with the wallclock time of 5 seconds on the smallest mesh and 358 seconds on the largest mesh.

8.2. Semilinear elliptic PDE with control and state constraints. The purpose of this example is to demonstrate nearly mesh-independent performance of ALESQP on a PDE-constrained optimization problem with control and state constraints. From now on, we solve (7.4) to accelerate the projected CG method. As mentioned earlier, with (7.4) we only use direct linear system solves. Additionally, we demonstrate that ALESQP meets constraint tolerances at the level of machine precision with only marginally increased iteration counts. Let $\Omega = (0, 1)^2$, $X = X_1 \oplus X_2$ with $X_1 = H_0^1(\Omega) \cap C_0(\Omega)$ and $X_2 = L^2(\Omega)$, $Y = H^{-1}(\Omega)$, and $Z_i = L^2(\Omega)$ for $i = 1, 2, 3$. Here, $C_0(\Omega)$ is the space of continuous functions on $\bar{\Omega}$ that vanish on the boundary $\partial\Omega$. We consider the problem

$$(8.2a) \quad \min_{u \in X_1, z \in X_2} \left\{ f(u, z) := \frac{1}{2} \|u - u_d\|_{L^2(\Omega)}^2 + \frac{\alpha}{2} \|z\|_{L^2(\Omega)}^2 \right\},$$

$$(8.2b) \quad \text{subject to } u_a \leq u \quad \text{a.e. in } \Omega,$$

$$(8.2c) \quad z_a \leq z \leq z_b \quad \text{a.e. in } \Omega,$$

$$(8.2d) \quad g(u, z) := \begin{cases} -\Delta u + u^p - z = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $u_d \in L^2(\Omega)$, $z_a, z_b \in L^2(\Omega)$ with $z_a \leq z_b$ a.e. in Ω , $u_a \in C(\bar{\Omega})$ with $u_a \leq 0$ on $\partial\Omega$, $p \in \{3, 5\}$, and $\alpha > 0$ is the penalty parameter, are given. Moreover, $g(u, z) = 0$ is the weak form of (8.2d), \widehat{T}_1 is the compact embedding of $H_0^1(\Omega)$ into $L^2(\Omega)$, \widehat{T}_2 and \widehat{T}_3 are the identity operator on $L^2(\Omega)$, $\widehat{C}_1 := \{u \in Z_1 \mid u_a \leq u\}$, $\widehat{C}_2 := \{z \in Z_2 \mid z_a \leq z\}$, and $\widehat{C}_3 := \{z \in Z_3 \mid z \leq z_b\}$. One can show that the solution to (8.2d), $u \in H_0^1(\Omega)$, in fact satisfies $u \in C_0(\Omega)$ [10]. In addition, from [31, Thm. 2.14], we recall that the Lagrange multiplier ζ associated with the constraint $g(u, z) = 0$ satisfies $\zeta \in W_0^{1,s}(\Omega)$ with $s \in [1, 2)$. Consequently, if $z_a, z_b \in W^{1,s}(\Omega)$, then we can conclude that the optimal control to (8.2) satisfies $z \in W^{1,s}(\Omega)$, where $W^{1,s}(\Omega)$ is compactly embedded in $L^2(\Omega)$. As a consequence of Theorem 6.1 and Corollary 5.12(e), any weak accumulation point in $(H_0^1(\Omega) \cap C_0(\Omega)) \oplus W^{1,s}(\Omega)$ of the sequence of iterates generated by Algorithm 4.1 is feasible, so long as (6.3) holds. When only state constraints are present (i.e., $z_a = -\infty$ and $z_b = +\infty$), Corollary 5.12(c) ensures that the sequence of iterates generated by Algorithm 4.1 is asymptotically stationary and accumulates at a feasible point. Moreover, if (5.21) holds (which is possible since $T^{-1}(C) = \{u \in H_0^1(\Omega) \cap C_0(\Omega) \mid u_a \leq u\}$ has a nonempty interior), then any accumulation point is stationary by Corollary 5.16.

We investigate ALESQP performance on three scenarios: (i) only control constraints (i.e., $u_a = -\infty$), (ii) only state constraints (i.e., $z_a = -\infty$ and $z_b = \infty$), and (iii) both control and state constraints. For our numerical studies, we set $r_1^{(1)} = 10^3$, $r_2^{(1)} = \alpha$, and $r_3^{(1)} = \alpha$. The choice of initial penalty parameters is important to account for the differences in regularity and scaling of the associated multipliers. More precisely, we use α for the control-constraint parameters to balance the control penalty term $\frac{\alpha}{2}\|z\|_{L^2(\Omega)}^2$ in the objective function. Additionally, we note that the control-constraint multipliers are in $L^2(\Omega)$ and that we expect the corresponding penalty parameters to remain bounded. In contrast, the state-constraint multiplier is a measure, which suggests that the sequence $\{\|\lambda_1^{(k)}\|_Z\}_{k=1}^\infty$ is unbounded. Consequently, Lemma 5.2 suggests that sequence of penalty parameters $\{r_1^{(k)}\}_{k=1}^\infty$ is also unbounded. Therefore, it is appropriate to choose a very large initial parameter, here $r_1^{(1)} = 10^3$. In our studies, considerably larger values of $r_1^{(1)}$ had little impact on overall performance, including the solution of the SQP subproblems. Smaller values delayed the convergence of the outer augmented Lagrangian loop somewhat.

The problem data in (8.2) is motivated by [9]. In particular, we set $p = 3$, $u_d \equiv -1$, $\alpha = 10^{-3}$, $z_a \equiv -10$, $z_b \equiv 10$, and

$$u_a(x) = -\frac{2}{3} + \frac{1}{2} \min\{x_1 + x_2, \min\{1 + x_1 - x_2, \min\{1 - x_1 + x_2, 2 - x_1 - x_2\}\}\}.$$

For scenario (i), we replace $z_b = 10$ by $z_b = -1$, to ensure that the constraints are active. In Table 3 we observe nearly mesh-independent performance of ALESQP for all three scenarios. Additionally, we note that the state-constraint penalty parameter increases significantly and that the growth of the control-constraint penalty parameters is more moderate; e.g., for the 128×128 mesh in scenario (iii) the final values are $r_1^{(14)} = 3.12 \cdot 10^6$, $r_2^{(14)} = 1.25 \cdot 10^{-1}$, and $r_3^{(14)} = 6.25 \cdot 10^{-1}$. Moreover, if we tighten the outer stopping tolerances to 10^{-12} , as we do later in Table 4,

TABLE 3

Semilinear. Control: only control constraints; State: only state constraints; Control + State: control and state constraints. ALESQP performance for varying spatial discretization (Mesh). In all cases, we observe that the AL, SQP, and CG iterations are nearly mesh independent.

Mesh	AL	Control		State		Control + State		
		SQP	CG	AL	SQP	CG	AL	SQP
64×64	9	20	41	11	28	57	12	35
128×128	9	20	40	12	33	66	14	44
256×256	9	22	45	14	40	80	16	50
512×512	9	22	47	16	47	95	18	60
								123

TABLE 4

Semilinear (Control + State). ALESQP performance for varying stopping tolerances, $\varepsilon_* = \delta_* = \tau_* = \text{tol}$. We observe that it is possible to achieve machine precision for all convergence measures with a very mild increase in the iteration counts.

tol	AL	SQP	CG	normg	grad-lag	feas
1e-6	14	44	90	9.51e-13	7.99e-12	1.35e-08
1e-8	17	46	94	9.56e-13	1.31e-13	7.43e-10
1e-10	21	48	98	9.58e-13	8.66e-14	2.77e-12
1e-12	25	50	102	9.67e-13	8.20e-14	2.49e-14

the final penalty parameter values are $r_1^{(25)} = 3.91 \cdot 10^8$, $r_2^{(25)} = 6.25 \cdot 10^{-1}$, and $r_3^{(25)} = 6.25 \cdot 10^{-1}$. In other words, the state-constraint penalty continues to grow, while the control-constraint penalties stagnate. The discrepancy between $r_1^{(25)}$ and $r_2^{(25)}$ or $r_3^{(25)}$ strongly underlines the need for multiple penalties.

In Table 4, we illustrate a remarkable feature of our algorithm. We consider a fixed mesh of size 128×128 and vary the outer stopping tolerances, including the T -feasibility tolerance τ_* . We observe that it is possible to achieve machine precision for all convergence measures with almost no increase in the total number of projected CG iterations.

8.2.1. Comparisons with related methods. We provide comparisons with related augmented Lagrangian methods, developed in [27] and [9]. We note that in [29], these methods are shown to be competitive with Moreau–Yosida regularization. Our first comparison is based on the semilinear optimal control example with state constraints and a quintic nonlinear term from [27, sect. 6.1]. Specifically, we set $p = 5$ in (8.2d), and we set the auxiliary function data u_d , f_d , and u_b as in [27]. Here, $u_b \in C(\bar{\Omega})$ is an upper bound on u that fulfills $u_b \geq 0$ on $\partial\Omega$. Additionally, we have $\alpha = 1$, $u_a = -\infty$, $z_a = -\infty$, and $z_b = +\infty$. The setup for our numerical simulations, based on finite element discretizations, is similar to the finite element framework used in [27]. We compare the numbers of outer (augmented Lagrangian) and inner (semismooth Newton) iterations reported in [27] with the numbers of outer (augmented Lagrangian) and inner (smooth equality-constrained SQP) ALESQP iterations. Table 5 demonstrates that ALESQP is a competitive method for PDE-constrained optimization problems with state constraints.

Our second comparison is based on the semilinear optimal control example with state and control constraints and a cubic nonlinear term, studied earlier in this section. This problem was originally studied in [9, sect. 8.1]. In [9] the authors use a five-point finite difference discretization, while our numerical studies use finite elements. To enable a more direct solution comparison, we match the computational grids used in

TABLE 5

Comparison (state constraints). ALESQP performance in comparison with the augmented Lagrangian method from [27]. The numbers of outer and inner ALESQP iterations are modest.

Mesh	Aug Lagr [27]		ALESQP		
	Outer	Inner	Outer	Inner	CG
16×16	10	21	6	7	9
32×32	9	23	2	6	10
64×64	10	27	4	8	14
128×128	11	33	7	11	17
256×256	12	38	11	16	26
512×512	—	—	8	14	21

TABLE 6

Comparison (control and state constraints). ALESQP performance in comparison with the augmented Lagrangian method from [9]. ALESQP is a competitive alternative.

Mesh	Aug Lagr [9]		ALESQP		
	Outer	Inner	Outer	Inner	CG
16×16	27	211	12	23	50
32×32	30	238	11	28	61
64×64	38	289	11	35	73
128×128	46	332	12	47	97
256×256	52	370	14	49	101
512×512	—	—	16	54	109

[9], i.e., change our triangular meshes to quadrilateral meshes, and change linear simplicial elements to bilinear quadrilateral elements. This results in a slight change of ALESQP iteration numbers compared to those reported in Table 3. More importantly, there are significant algorithmic differences between the method of [9] and ALESQP. The method of [9] solves subproblems with control inequality constraints. Specifically, the authors use `fmincon` in MATLAB. In contrast, ALESQP handles all inequality constraints through the augmented Lagrangian, and solves smooth subproblems constrained only by equalities, using SQP. Therefore, we expect the inner iterations in ALESQP to be computationally cheaper compared to the inner iterations in [9]. Table 6 demonstrates that ALESQP is a competitive alternative.

8.3. Burgers' PDE with control and state constraints. This example showcases ALESQP in the context of *dynamic optimization*. Let $\Omega = (0, 1)$, $Q := \Omega \times (0, T)$, $\Sigma := \partial\Omega \times (0, T)$, $X = X_1 \oplus X_2$ with $X_1 = L^2(0, T; H_0^1(\Omega))$ and $X_2 = L^2(Q)$, $Y = L^2(0, T; H^{-1}(\Omega))$, and $Z_1 = Z_2 = Z_3 = L^2(Q)$. We consider the problem

$$(8.3a) \quad \min_{u \in X_1, z \in X_2} \left\{ f(u, z) := \frac{1}{2} \|u - u_d\|_{L^2(Q)}^2 + \frac{\alpha}{2} \|z\|_{L^2(Q)}^2 \right\},$$

$$(8.3b) \quad \text{subject to } u_a \leq u \quad \text{a.e. in } Q,$$

$$(8.3c) \quad z_a \leq z \leq z_b \quad \text{a.e. in } Q,$$

$$(8.3d) \quad g(u, z) := \begin{cases} \partial_t u + u \partial_x u - \nu \partial_{xx}^2 u - z = 0 & \text{in } Q, \\ u = 0 & \text{on } \Sigma, \\ u(\cdot, 0) - u_0 = 0 & \text{in } \Omega \end{cases}$$

TABLE 7

Burger's equation. Control: only control constraints; State: only state constraints; Control + State: control and state constraints. ALESQP performance for varying spatial and temporal discretization (Mesh). In all cases, the AL, SQP, and CG iterations are nearly mesh independent.

Mesh	Control			State			Control + State		
	AL	SQP	CG	AL	SQP	CG	AL	SQP	CG
64×64	10	22	86	6	28	78	11	43	135
128×128	10	25	91	8	31	89	11	54	166
256×256	10	29	104	8	33	92	14	72	210
512×512	10	29	105	9	34	95	14	68	201

with datum $u_d \in L^2(Q)$, $u_a \in C(\bar{Q})$ with $u_a(x, t) \leq 0$ for all $(x, t) \in \partial\Omega \times [0, T]$, $z_a, z_b \in L^2(Q)$ with $z_a \leq z_b$ a.e. in Q . Moreover, $g(u, z) = 0$ is the weak form of (8.3d), \hat{T}_1 is the embedding of $L^2(0, T; H_0^1(\Omega))$ into $L^2(Q)$, \hat{T}_2 and \hat{T}_3 are the identity operators on $L^2(Q)$, and the constraint sets \hat{C}_i , $i = 1, 2, 3$, are defined similarly to the ones in section 8.2. It is unclear if the above problem fully satisfies our theory. In principle, one may be able to use regularity arguments similar to the ones from section 8.2. However, such regularity results are not known for ζ associated with (8.3d), and the required study is beyond the scope of the paper. Nevertheless, we observe that our algorithm solves this problem efficiently. We refer to [41] for regularity results involving the control-constrained case, case (i) below.

Similar to the previous example, we test ALESQP in three different scenarios: (i) control constraints, (ii) state constraints, and (iii) mixed constraints. For our numerical results, we set $r_1^{(1)} = 10^3$, $r_2^{(1)} = \alpha$, and $r_3^{(1)} = \alpha$. The choice of the initial penalty parameters is justified by the problem structure and closely follows the considerations given in section 8.2. For $t \in (0, 1)$, we set $u_d = 1$ for $x \in (0, 1/2)$ and $u_d = 0$ otherwise, $\alpha = 5 \times 10^{-2}$, $z_a = -1$, $z_b = 2$, and $u_a = 0$. Table 7 shows ALESQP iteration counts. As in all previous examples, we observe nearly mesh-independent performance. Similar to section 8.2 we note that the state-constraint penalty parameter increases significantly, while the control-constraint penalty parameters increase moderately; e.g., for the 128×128 case in scenario (iii) the final values are $r_1^{(15)} = 1.25 \cdot 10^5$, $r_2^{(15)} = 1.25 \cdot 10^0$, and $r_3^{(15)} = 2.5 \cdot 10^{-1}$.

REFERENCES

- [1] R. ADAMS AND J. F. FOURNIER, *Sobolev Spaces*, Pure Appl. Math. (Amsterdam) 140, 2nd ed., Elsevier/Academic Press, Amsterdam, 2003.
- [2] R. ANDREANI, E. G. BIRGIN, J. M. MARTÍNEZ, AND M. L. SCHUVERDT, *On augmented Lagrangian methods with general lower-level constraints*, SIAM J. Optim., 18 (2007), pp. 1286–1309, <https://doi.org/10.1137/060654797>.
- [3] H. H. BAUSCHKE AND P. L. COMBETTES, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, CMS Books Math., Springer, New York, 2011.
- [4] M. BERGOUNIOUX, *Augmented Lagrangian method for distributed optimal control problems with state constraints*, J. Optim. Theory Appl., 78 (1993), pp. 493–521, <https://doi.org/10.1007/BF00939879>.
- [5] M. BERGOUNIOUX AND K. KUNISCH, *Augmented Lagrangian techniques for elliptic state constrained optimal control problems*, SIAM J. Control Optim., 35 (1997), pp. 1524–1543, <https://doi.org/10.1137/S036301299529330X>.
- [6] D. P. BERTSEKAS, *Constrained Optimization and Lagrange Multiplier Methods*, Compu. Sci. Appl. Math., Academic Press, New York, 1982.
- [7] E. G. BIRGIN, L. F. BUENO, AND J. M. MARTÍNEZ, *Sequential equality-constrained optimization for nonlinear programming*, Comput. Optim. Appl., 65 (2016), pp. 699–721, <https://doi.org/10.1007/s10589-016-9849-6>.

- [8] E. G. BIRGIN AND J. M. MARTÍNEZ, *Practical Augmented Lagrangian Methods for Constrained Optimization*, Fundam. Algorithms 10, SIAM, Philadelphia, 2014, <https://doi.org/10.1137/1.9781611973365>.
- [9] E. BÖRGENS, C. KANZOW, AND D. STECK, *Local and global analysis of multiplier methods for constrained optimization in Banach spaces*, SIAM J. Control Optim., 57 (2019), pp. 3694–3722, <https://doi.org/10.1137/19M1240186>.
- [10] E. CASAS, *Control of an elliptic problem with pointwise state constraints*, SIAM J. Control Optim., 24 (1986), pp. 1309–1318, <https://doi.org/10.1137/0324078>.
- [11] X. CHEN, Z. NASCHED, AND L. QI, *Smoothing methods and semismooth methods for non-differentiable operator equations*, SIAM J. Numer. Anal., 38 (2000), pp. 1200–1216, <https://doi.org/10.1137/S0036142999356719>.
- [12] A. R. CONN, N. I. M. GOULD, AND P. L. TOINT, *A globally convergent augmented lagrangian algorithm for optimization with general constraints and simple bounds*, SIAM J. Numer. Anal., 28 (1991), pp. 545–572, <https://doi.org/10.1137/0728030>.
- [13] A. R. CONN, N. I. M. GOULD, AND P. L. TOINT, *Trust-Region Methods*, SIAM, Philadelphia, 2000.
- [14] A. R. CONN, N. I. M. GOULD, AND P. L. TOINT, *Lancelot: A Fortran Package for Large-Scale Nonlinear Optimization (Release A)*, Springer Ser. Comput. Math. 17, Springer, Berlin, 2013.
- [15] J. E. DENNIS, M. EL-ALEM, AND M. C. MACIEL, *A global convergence theory for general trust-region-based algorithms for equality constrained optimization*, SIAM J. Optim., 7 (1997), pp. 177–207, <https://doi.org/10.1137/S1052623492238881>.
- [16] G. B. FOLLAND, *Real Analysis: Modern Techniques and Their Applications*, Pure Appl. Math., Wiley, New York, 1999.
- [17] N. I. M. GOULD, M. E. HRIBAR, AND J. NOCEDAL, *On the solution of equality constrained quadratic programming problems arising in optimization*, SIAM J. Sci. Comput., 23 (2002), pp. 1376–1395, <https://doi.org/10.1137/S1064827598345667>.
- [18] A. GÜNNEL, R. HERZOG, AND E. SACHS, *A note on preconditioners and scalar products in Krylov subspace methods for self-adjoint problems in Hilbert space*, Electron. Trans. Numer. Anal., 41 (2014), pp. 13–20.
- [19] M. HEINKENSCHLOSS AND D. RIDZAL, *A matrix-free trust-region SQP method for equality constrained optimization*, SIAM J. Optim., 24 (2014), pp. 1507–1541, <https://doi.org/10.1137/130921738>.
- [20] M. HEINKENSCHLOSS AND L. N. VICENTE, *Analysis of inexact trust-region SQP algorithms*, SIAM J. Optim., 2 (2001), pp. 283–302, <https://doi.org/10.1137/S1052623499361543>.
- [21] M. R. HESTENES, *Multiplier and gradient methods*, J. Optim. Theory Appl., 4 (1969), pp. 303–320, <https://doi.org/10.1007/BF00927673>.
- [22] M. HINZE, R. PINNAU, M. ULRICH, AND S. ULRICH, *Optimization with PDE Constraints*, Math. Model. Theory Appl. 23, Springer, New York, 2009.
- [23] K. ITO AND K. KUNISCH, *The augmented Lagrangian method for equality and inequality constraints in Hilbert spaces*, Math. Program., 46 (1990), pp. 341–360, <https://doi.org/10.1007/BF01585750>.
- [24] K. ITO AND K. KUNISCH, *An augmented Lagrangian technique for variational inequalities*, Appl. Math. Optim., 21 (1990), pp. 223–241, <https://doi.org/10.1007/BF01445164>.
- [25] K. ITO AND K. KUNISCH, *Augmented Lagrangian-SQP-methods in Hilbert spaces and application to control in the coefficients problems*, SIAM J. Optim., 6 (1996), pp. 96–125, <https://doi.org/10.1137/0806007>.
- [26] K. ITO AND K. KUNISCH, *Lagrange Multiplier Approach to Variational Problems and Applications*, Adv. Des. Control 15, SIAM, Philadelphia, 2008, <https://doi.org/10.1137/1.9780898718614>.
- [27] C. KANZOW, V. KARL, D. STECK, AND D. WACHSMUTH, *Safeguarded augmented Lagrangian methods in Banach spaces*, in Non-Smooth and Complementarity-Based Distributed Parameter Systems, Internat. Ser. Numer. Math. 172, Springer, Cham, 2022, pp. 241–282, <https://doi.org/10.1007/978-3-030-79393-7-10>.
- [28] C. KANZOW AND D. STECK, *Quasi-variational inequalities in banach spaces: Theory and augmented Lagrangian methods*, SIAM J. Optim., 29 (2019), pp. 3174–3200.
- [29] C. KANZOW, D. STECK, AND D. WACHSMUTH, *An augmented Lagrangian method for optimization problems in Banach spaces*, SIAM J. Control Optim., 56 (2018), pp. 272–291, <https://doi.org/10.1137/16M1107103>.
- [30] D. P. KOURI AND D. RIDZAL, *Inexact trust-region methods for PDE-constrained optimization*, in *Frontiers in PDE-Constrained Optimization*, Springer, New York, 2018, pp. 83–121.
- [31] I. NEITZEL, J. PFEFFERER, AND A. RÖSCH, *Finite element discretization of state-constrained elliptic optimal control problems with semilinear state equation*, SIAM J. Control Optim., 53 (2015), pp. 874–904, <https://doi.org/10.1137/140960645>.

- [32] E. O. OMOJOKUN, *Trust-Region Algorithms for Optimization with Nonlinear Equality and Inequality Constraints*, Ph.D. thesis, University of Colorado, Boulder, 1989.
- [33] C. C. PAIGE AND M. A. SAUNDERS, *Solution of sparse indefinite systems of linear equations*, SIAM J. Numer. Anal., 12 (1975), pp. 617–629.
- [34] M. J. D. POWELL, *A method for nonlinear constraints in minimization problems*, in Optimization, Academic Press, London, 1969, pp. 283–298.
- [35] M. J. D. POWELL, *A new algorithm for unconstrained optimization*, in Nonlinear Programming, Academic Press, New York, 1970, pp. 31–65.
- [36] R. T. ROCKAFELLAR, *The multiplier method of Hestenes and Powell applied to convex programming*, J. Optim. Theory Appl., 12 (1973), pp. 555–562, <https://doi.org/10.1007/BF00934777>.
- [37] R. T. ROCKAFELLAR, *Augmented Lagrange multiplier functions and duality in nonconvex programming*, SIAM J. Control, 12 (1974), pp. 268–285.
- [38] R. T. ROCKAFELLAR, *Augmented Lagrangians and applications of the proximal point algorithm in convex programming*, Math. Oper. Res., 1 (1976), pp. 97–116, <https://doi.org/10.1287/moor.1.2.97>.
- [39] W. RUDIN, *Functional Analysis*, Int. Ser. Pure Appl. Math., McGraw-Hill, New York, 1991.
- [40] T. STEIHAUG, *The conjugate gradient method and trust regions in large scale optimization*, SIAM J. Numer. Anal., 20 (1983), pp. 626–637, <https://doi.org/10.1137/0720042>.
- [41] F. TRÖLTZSCH AND S. VOLKWEIN, *The SQP method for control constrained optimal control of the Burgers equation*, ESAIM Control Optim. Calc. Var., 6 (2001), pp. 649–674, <https://doi.org/10.1051/cocv:2001127>.
- [42] T. VAN ERVEN AND P. HARREMOS, *Rényi divergence and Kullback-Leibler divergence*, IEEE Trans. Inform. Theory, 60 (2014), pp. 3797–3820.
- [43] S. VOLKWEIN, *Mesh-independence for an augmented lagrangian-SQP method in Hilbert spaces*, SIAM J. Control Optim., 38 (2000), pp. 767–785.
- [44] S. VOLKWEIN AND M. WEISER, *Affine invariant convergence analysis for inexact augmented Lagrangian-SQP methods*, SIAM J. Control Optim., 41 (2002), pp. 875–899.
- [45] S. S. WILKS, *Certain generalizations in the analysis of variance*, Biometrika, 24 (1932), pp. 471–494, <https://doi.org/10.1093/biomet/24.3-4.471>.
- [46] E. H. ZARANTONELLO, *Projections on convex sets in Hilbert space and spectral theory*, in Contributions to Nonlinear Functional Analysis, Academic Press, New York, 1971, pp. 237–424.
- [47] J. ZIEMS AND S. ULRICH, *Adaptive multilevel inexact SQP methods for PDE-constrained optimization*, SIAM J. Optim., 21 (2011), pp. 1–40, <https://doi.org/10.1137/080743160>.
- [48] J. ZOWE AND S. KURCYUSZ, *Regularity and stability for the mathematical programming problem in Banach spaces*, Appl. Math. Optim., 5 (1979), pp. 49–62.