

Dicke Superradiance Requires Interactions beyond Nearest Neighbors

Wai-Keong Mok^{1,2}, Ana Asenjo-Garcia,³ Tze Chien Sum⁴, and Leong-Chuan Kwek^{1,5,6,7}

¹Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543, Singapore

²California Institute of Technology, Pasadena, California 91125, USA

³Department of Physics, Columbia University, New York, New York 10027, USA

⁴Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

⁵MajuLab, CNRS-UNS-NUS-NTU International Joint Research Unit, Singapore UMI 3654, Singapore

⁶National Institute of Education, Nanyang Technological University, Singapore 637616, Singapore

⁷Quantum Science and Engineering Centre (QSec), Nanyang Technological University, 50 Nanyang Ave, Singapore 639798, Singapore



(Received 3 November 2022; revised 18 April 2023; accepted 1 May 2023; published 25 May 2023)

Photon-mediated interactions within an excited ensemble of emitters can result in Dicke superradiance, where the emission rate is greatly enhanced, manifesting as a high-intensity burst at short times. The superradiant burst is most commonly observed in systems with long-range interactions between the emitters, although the minimal interaction range remains unknown. Here, we put forward a new theoretical method to bound the maximum emission rate by upper bounding the spectral radius of an auxiliary Hamiltonian. We harness this tool to prove that for an arbitrary ordered array with only nearest-neighbor interactions in all dimensions, a superradiant burst is not physically observable. We show that Dicke superradiance requires minimally the inclusion of next-nearest-neighbor interactions. For exponentially decaying interactions, the critical coupling is found to be asymptotically independent of the number of emitters in all dimensions, thereby defining the threshold interaction range where the collective enhancement balances out the decoherence effects. Our findings provide key physical insights to the understanding of collective decay in many-body quantum systems, and the designing of superradiant emission in physical systems for applications such as energy harvesting and quantum sensing.

DOI: 10.1103/PhysRevLett.130.213605

Introduction.—Collective spontaneous emission of N initially inverted atoms with identical all-to-all interactions mediated by the electromagnetic vacuum results in a burst of light with intensity scaling as N^2 [1–3]. This phenomenon is commonly referred to as “Dicke superradiance” or “superradiant burst.” Over the past decades, this many-body phenomenon has attracted a lot of interest in both theoretical [4–24] and experimental studies [25,26] using a multitude of physical platforms such as trapped ions [27], molecular aggregates [28–31], solid-state emitters [32–36], cold atoms and molecules [37–40], and superconducting qubits [41–43], with wide-ranging applications including the generation of multiphoton states with improved metrological properties [18,44–47], energy harvesting [48–50], ultrabright LEDs [51] and quantum sensing [52,53].

The atoms in Dicke’s original model were assumed to be confined within a spatial extent smaller than the emission wavelength λ . Consequently, the atoms become indistinguishable with respect to the absorption or emission of photons, such that their quantum state $|j = N/2, m\rangle$ (with $-N/2 \leq m \leq N/2$) is permutation invariant. This permutation symmetry greatly reduces the complexity of the problem, as it constrains the dynamics to $N + 1$ states, instead of exploring the full Hilbert space (which scales as

2^N). Recently, there has been substantial research progress with extended systems where atoms are distributed over a region larger than λ , thus breaking this symmetry. Of particular interest are ordered atomic arrays [54,55], in which the superradiant properties can be greatly affected by the geometry and dimensionality of the lattice [16,19,20,22,24,56]. The interactions between the emitters are typically modeled by long-range dipole-dipole interactions mediated via the electromagnetic vacuum [57,58].

A long-standing fundamental question is the *minimal* interaction range required for the occurrence of a superradiant burst. Intuitively, superradiance can be thought of as a competition between (transient) phase synchronization, which leads to the buildup of atomic correlations, and decoherence [59]. Both effects stem from the same dissipative interactions [8,22]. Since synchronization of non-linear classical phase oscillators has been demonstrated with nearest-neighbor (NN) coupling [60], one may expect the atomic phases to synchronize for sufficiently strong NN interactions resulting in a superradiant burst [59]. Moreover, for a fixed interaction range, higher dimensionality was reported to result in stronger superradiance due to long-range order [19,24]. On the flip side, it could also be argued that for short-range interactions, the buildup of

correlations is not strong enough to overcome decoherence, thereby preventing superradiance.

In this Letter, we prove that superradiant burst is impossible in an arbitrary D -dimensional array with only nearest-neighbor interactions, for arbitrary times and initial states. That is, we show that, in all cases, the emission rate is upper bounded by that of independent emitters, resulting in no enhancement from collective dynamics. Including next-nearest-neighbor interactions, we show that a superradiant burst can be physically observed for certain values of the interaction strengths, thereby defining a minimal interaction range for superradiance. Another question is the *threshold* interaction range, which we define to be such that the critical coupling required for a burst becomes independent of the number of emitters, for any D . We show that exponentially decaying interactions lie on the threshold interaction range for which the synchronization of the dipoles arising from the emission balances the decoherence effects.

Model.—The dynamics of an undriven ensemble of N emitters can be described by the Lindblad master equation (setting $\hbar = 1$)

$$\dot{\rho} = -i \sum_{i,j=1}^N [J_{ij}\sigma_i^+\sigma_j^-, \rho] + \sum_{i,j=1}^N \gamma_{ij}\mathcal{D}[\sigma_i^-, \sigma_j^-]\rho \equiv \mathcal{L}[\rho], \quad (1)$$

with $J_{ij} = J_{ji}^*$ and $\gamma_{ij} = \gamma_{ji}^*$ to ensure Hermiticity. The raising and lowering operators for the j th emitter are denoted as $\sigma_j^+ \equiv |e_i\rangle\langle g_i|$ and $\sigma_j^- \equiv |g_i\rangle\langle e_i|$ which describe transitions between the ground state $|g_i\rangle$ and excited state $|e_i\rangle$. The first term contains the coherent Hamiltonian interactions between the emitters, while the second term captures processes such as collective and local dissipation of the emitters via the superoperator $\mathcal{D}[\sigma_i^-, \sigma_j^-]\rho = \sigma_i^-\rho\sigma_j^+ - \{\sigma_j^+\sigma_i^-, \rho\}/2$. We assume J_{ij} and γ_{ij} to be time independent, such that the superoperator \mathcal{L} generates a dynamical semigroup describing the dynamics of a Markovian open quantum system.

For a physically valid evolution (i.e., a completely positive and trace-preserving map), the matrix Γ containing the elements γ_{ij} (which we will refer to as the *decoherence matrix*) must be positive semidefinite [61–63]. The decoherence matrix can be diagonalized to yield N decay rates $\Gamma_\nu \geq 0$, with $\nu \in \{1, \dots, N\}$ and the corresponding collective jump operators \hat{c}_ν . The total photon emission rate of the emitters, integrated over all emission directions, is defined for any state ρ as

$$R_\rho \equiv \sum_{\nu=1}^N \Gamma_\nu \langle \hat{c}_\nu^\dagger \hat{c}_\nu \rangle = \sum_{\nu=1}^N \Gamma_\nu \text{Tr}(\hat{c}_\nu^\dagger \hat{c}_\nu \rho). \quad (2)$$

For independent emitters with $\gamma_{ij} = \gamma_0 \delta_{ij}$, the total emission rate has a maximum of $N\gamma_0$ (saturated by the fully




FIG. 1. Dynamics of the photon emission rate $R(t)$ for emitter arrays with only nearest-neighbor interactions of strength γ , normalized by the individual emitter decay rate γ_0 . For $\gamma/\gamma_0 < \gamma_s$, $\dot{R}(0) < 0$ and the photon emission rate decays monotonically without a superradiant burst (blue). Superradiance occurs for $\gamma/\gamma_0 > \gamma_s$ (red). The physically valid regime is defined by $0 < \gamma/\gamma_0 \leq \gamma_p$. For nearest-neighbor interactions, $\gamma_p < \gamma_s$ (with a finite gap between γ_p and γ_s) for any arbitrary emitter array in all dimensions, rendering Dicke superradiance physically impossible.

excited state), and $R(t) \equiv R_{\rho(t)}$ decays exponentially. However, interactions between the emitters can cause $R(t)$ to increase beyond its initial value. This speedup in emission is commonly referred to as the superradiant burst, first discovered by Dicke [1] (see Fig. 1). Throughout this work, we refer to superradiant burst as the increase in the total emission rate beyond $N\gamma_0$, but the peak intensity need not scale as N^2 . In general, characterizing the burst at arbitrary times can be difficult, hence one typically uses

$$\dot{R}_\rho = i \sum_\nu \langle [H, \hat{c}_\nu^\dagger \hat{c}_\nu] \rangle - \sum_{\mu, \nu} \Gamma_\mu \Gamma_\nu \langle \hat{c}_\mu^\dagger [\hat{c}_\mu, \hat{c}_\nu^\dagger] \hat{c}_\nu \rangle \quad (3)$$

evaluated at the fully excited initial state $\rho(0)$, with $\dot{R}(0) \equiv \dot{R}_{\rho(0)} > 0$ a sufficient condition for a superradiant burst. While we consider the burst at $t = 0$, we will provide physical justification on why this is sufficient.

Here, we put forward a new (and complementary) criterion to preclude any possibility of a burst: by a simple change of basis, one can write Eq. (2) as the expectation value of an auxiliary spin Hamiltonian

$$H_\Gamma = \sum_{j,k=1}^N \gamma_{kj} \sigma_j^+ \sigma_k^-, \quad (4)$$

with $R_\rho = \text{tr}(H_\Gamma \rho)$. The maximum photon emission rate can thus be calculated by bounding the spectral radius of the auxiliary spin Hamiltonian. If the upper bound is equal to or smaller than $N\gamma_0$, no burst can occur for all times and

arbitrary initial states. While finding the largest eigenvalue of H_Γ may be nontrivial, this criterion allows one to definitively prove the absence of a burst for arbitrary times, thus going beyond the condition $\dot{R}(0) \equiv \dot{R}_{\rho(0)} > 0$. Furthermore, this approach opens up the possibility of finding theoretical limits for the emission rate arising from superradiant dynamics, as we show below and in Supplemental Material [64].

No superradiance for nearest-neighbor coupling.—Let us consider a hypercube array of N emitters with arbitrary dimension D ($N = n^D$). For the case of NN interactions, $\gamma_{ii} \equiv \gamma_0 = 1$ and $\gamma_{ij} = \gamma$ if emitters i and j are nearest neighbor ($\gamma_{ij} = 0$ otherwise). The coupling $\gamma \in [0, 1]$ is required for the matrix Γ to be positive semidefinite. Without loss of generality, we have assumed γ_{ij} to be real and positive. We prove that for this model, superradiant burst cannot occur for any $t > 0$, for any arbitrary initial state and for any Hamiltonian coupling J_{ij} . To determine the physically valid regime, we impose the condition that Γ is positive semidefinite. Notice that the decoherence matrix can be expressed as $\Gamma = \mathbf{I}_N + \gamma \mathbf{A}$, where \mathbf{I}_N is the $N \times N$ identity matrix, and \mathbf{A} is the adjacency matrix of a $n \times n$ grid graph. Using the fact that the grid graph is the Cartesian product of D path graphs $P_n \square \cdots \square P_n$, it can be shown that the smallest eigenvalue of Γ is [64]

$$\Gamma_{\min} = 1 - 2D\gamma \cos\left(\frac{\pi}{N^{1/D} + 1}\right), \quad (5)$$

which gives the physically valid regime as $\gamma \leq \gamma_p$,

$$\gamma_p = \left[2D \cos\left(\frac{\pi}{N^{1/D} + 1}\right)\right]^{-1}. \quad (6)$$

This rate reduces to $\gamma_p = 1/(2D)$ in the $N \rightarrow \infty$ limit, or when imposing periodic boundary conditions for a finite N . This can be regarded as coming from the coordination number for each emitter, which approaches $2D$ in the infinite-array limit. We now state our main result.

Theorem 1.—Let Γ be the decoherence matrix for a nearest-neighbor interaction model, with $\gamma_{ij} = \delta_{ij} + \gamma\delta_{\langle ij \rangle}$, where $\gamma \in [0, 1]$, and $\delta_{\langle ij \rangle} = 1$ if the emitters indexed by i and j are nearest neighbor on the D -dimensional regular lattice, and 0 otherwise. For $\gamma \leq (2D)^{-1}$, the emission rate R_ρ is maximized by the fully excited state $|e\rangle^{\otimes N}$ with $R_\rho = N$.

We provide a sketch of the proof here, while the details can be found in Supplemental Material [64]. By expressing H_Γ in the product-state basis and using the Gershgorin circle theorem [75], we can upper bound $\max_t R(t) \leq N$ in the physically valid regime $\gamma < 1/(2D)$. This is saturated by N independent emitters in the fully excited state, with eigenvalue N . Hence, Theorem 1 implies that superradiant burst is impossible at all times. To gain a deeper physical

understanding, we evaluate the superradiant regime $\gamma > \gamma_s$ for the fully excited initial state, characterized by the transition at $\dot{R}(0) = 0$, for which [64]

$$\gamma_s = [2D(1 - N^{-1/D})]^{-1/2}. \quad (7)$$

For all $2 < N^{1/D} < \infty$, it can be shown that $\gamma_p < \gamma_s^2$ and therefore $\gamma_p < \gamma_s$. Hence, the superradiant regime does not overlap with the physically valid regime. Generalization to the hyperrectangle configuration where the number of sites along each dimension can be different is straightforward, and the same conclusion is obtained [64]. While our analysis of the NN model is valid for any initial state, we consider a fully inverted initial state for the next two sections: the analysis of next-nearest neighbor and exponentially decaying interactions.

Next-nearest neighbor coupling.—Including the NNN interactions, we now show that a superradiant burst is indeed possible. For simplicity, let us consider a 1D ring of N emitters with periodic boundary conditions. In this configuration, Γ turns out to be a circulant matrix with the first column given by $(1, \gamma_1, \gamma_2, 0, \dots, 0, \gamma_2, \gamma_1)^T$ with $0 \leq \{\gamma_1, \gamma_2\} \leq 1$. The subsequent columns are simply cyclic permutations of the first column. Diagonalizing Γ exactly yields the eigenvalues

$$\Gamma_\nu = 1 + 2\gamma_1 \cos\left(\frac{2\pi\nu}{N}\right) + 2\gamma_2 \cos\left(\frac{4\pi\nu}{N}\right) \quad (8)$$

for $\nu = 0, \dots, N - 1$. In the infinite-array limit $N \rightarrow \infty$, the eigenvalues form a continuous band in momentum space $\Gamma(k) = 1 + 2\gamma_1 \cos(k) + 2\gamma_2 \cos(2k)$, with the dimensionless wave vector $0 \leq k < 2\pi$. At the turning points where $\partial_k \Gamma = 0$, we have: $\Gamma(0) = 1 + 2(\gamma_1 + \gamma_2)$ which is always positive, $\Gamma(\pi) = 1 - 2(\gamma_1 - \gamma_2)$ and $\Gamma(k_*) = 1 - (\gamma_1^2 + 8\gamma_2^2)/4\gamma_2$ where $\cos k_* = -\gamma_1/4\gamma_2$. Demanding that $\Gamma(k) > 0$ thus produces the physically valid regimes: (I) $\gamma_1 - \gamma_2 \leq \frac{1}{2}$, $\gamma_1 > 4\gamma_2$ and (II) $\gamma_1^2 + 8\gamma_2^2 \leq 4\gamma_2$, $\gamma_1 \leq 4\gamma_2$, together with the bounds $\gamma_1, \gamma_2 \in [0, 1]$ (blue regions in Fig. 2). The superradiant condition can be obtained from $\dot{R}(0) = 0$ as (III) $\gamma_1^2 + \gamma_2^2 > 1/2$.

There is an overlap region with the physically valid regime, as shown by the red shaded region in Fig. 2. For certain values of γ_1, γ_2 , superradiant burst can occur. Moreover, the fact that this overlap region requires $\gamma_2 > (4 - \sqrt{2})/14 \approx 0.185$ is consistent with our previous conclusion of no superradiance using only NN coupling (i.e., $\gamma_2 = 0$). Superradiance is also forbidden by having only NNN coupling (i.e., $\gamma_1 = 0$). Results from numerical simulations of $N = 9$ emitters are presented in Fig. 3, which show that the NNN model has a small superradiant burst compared to the Dicke model, and no superradiance for NN models. We remark that this superradiance arises from destructive interference leading to dark decay




FIG. 2. Region of superradiant burst in the γ_2 - γ_1 plane. The physically valid (superradiant) regime is contained within the blue (red) boundary lines, with the conditions stated in the main text. Blue shaded region: physically valid, but not superradiant. Regions I, II, and III are defined in the main text. Red shaded region: physically valid with superradiant burst. Gray shaded region: unphysical regime. The red shaded region requires a minimum of $\gamma_2 \approx 0.185$. All shaded regions here are obtained from numerical calculations for $N = 100$, which agree very well with the analytical results obtained in the infinite-array limit.

channels with suppressed decay rates $\Gamma_\nu \approx 0$ while the dominant decay channel has a rate that does not scale with N . This mechanism is generally true for all models with a sharp interaction cutoff beyond a certain range.

Threshold interaction range for a superradiant burst.—In many previous works [16,19,20,22,24], Γ is obtained from a realistic modeling of the atomic interactions mediated by electromagnetic vacuum using the appropriate Green's function. Our goal here, however, is to shed light

on the essential physics of superradiance by considering analytically tractable models that still exhibit interesting behaviors. Consider an interaction which decays exponentially with the separation r_{ij} between the emitters: $\gamma_{ij} \propto e^{-\kappa r_{ij}}$, where κ controls the decay of the interaction strength with emitter separation. We set the diagonal elements of Γ as 1, and define $\gamma \equiv e^{-\kappa d}$ with d the emitter NN separation such that $\gamma_{ij} = \gamma^{|\vec{x}_i - \vec{x}_j|}$, where $\vec{x}_i \in \mathbb{Z}^D$ is the position vector of the i th lattice site. Physically, this model describes exponentially decaying interactions between the atoms. For a sufficiently large N in D dimensions such that $\gamma^N \ll 1$, $\dot{R}(0)$ is approximately given by the asymptotic form

$$\dot{R}(0) \sim N \left(\frac{2D\gamma^2}{1-\gamma^2} - 1 + \frac{C}{(-\ln \gamma)^D} \right) \quad (9)$$

for some constant C [64]. Interestingly, this suggests that the critical coupling parameter γ_s for superradiance is independent of N as $N \rightarrow \infty$ for all dimension, agreeing with the numerical results shown in Fig. 4. This is in stark contrast with previous results (primarily using long-range power-law interactions such as $\gamma_{ij} \propto 1/r_{ij}$), which predict that the critical emitter separation increases with N in 2D and 3D arrays [19,24]. Figure 4 also shows that for large N , $\gamma_s \sim D^{-0.793}$ exhibits a power-law scaling with the spatial dimension. This is intuitive as the average coupling per emitter increases with D which in turn lowers the critical coupling required for superradiance [24]. The N independence of γ_s for our short-range exponential model can be physically interpreted as the threshold interaction range where the synchronization effects due to collective interactions scales similarly with N as the local decoherence, such that adding more emitters does not affect the onset of




FIG. 3. Differential emission rate $\Delta R = R(t)/R(0) - 1$ against time (in units of emitter lifetime), for $N = 9$ emitters. $\Delta R > 0$ indicates superradiance. (a) Dynamical behavior of ΔR for the Dicke model (red), Next-nearest neighbor 1D ring (NNN, orange), nearest-neighbor 1D ring (NN, blue), and nearest-neighbor 2D square (NN, green) [see labels in (b)]. The coupling parameters are chosen to maximize $g^{(2)}(0)$. (b) Short-time behavior obtained by zooming into the gray region of (a). Only the Dicke and the next-nearest neighbor models exhibit superradiance. The curve for the Dicke model is scaled down by a factor of 10 for visualization purposes.




FIG. 4. Critical coupling γ_s for exponentially decaying interactions in a 1D chain, a 2D square array, and a 3D cubic array with N emitters. Superradiance occurs for $\gamma > \gamma_s$. For all dimension D , γ_s becomes independent of N for large N . Inset: log-log plot of γ_s against D for $N \approx 10^6$. γ_s decreases as D increases with a power-law scaling $\gamma_s \sim D^{-0.793}$.

the superradiant regime. For even shorter-range interactions such as the NN model, the local decoherence dominates, which prevents superradiance. Longer-range models such as power-law interactions favor synchronization and thus enhance superradiance as N increases.

Scaling of the peak emission rate with number of emitters.—Equation (4) shows that the problem of calculating the emission rate is equivalent to finding the average energy of a state under the Hamiltonian H_Γ . This enables us to find upper bounds on the scaling of the peak emission rate with N , for arbitrary geometries and types of interactions. As we have shown before in Theorem 1, the maximum emission rate for arbitrary NN models is $N\gamma_0$. For 1D arrays with an exponentially decaying interaction, the upper bound on the emission rate is found to scale as $O(N)$ for $\gamma < 1$ [64]. This bound increases to $O(N \log N)$ for 1D arrays with a power-law interaction of the form $1/r$ [64]. This latter scaling is consistent with the numerical results obtained in the literature which, in contrast to our bound, have only been obtained for relatively small systems and under certain approximations [20,22,24]. While finding exact bounds may be exponentially hard, one could in principle upper-bound other models, as well as tighten the currently obtained bounds.

Discussion.—In this Letter, we addressed the fundamental problem of the minimal interaction range required for superradiance. Crucially, we proved that nearest-neighbor interactions cannot induce emitter correlations faster than the decoherence, resulting in the impossibility of superradiance. As shown, the minimal interaction range is therefore next-nearest neighbor, and longer-range interactions generally lead to stronger superradiance. We also found that the short-range exponential interaction marks the threshold interaction range in all dimensions where the emitter correlations and local decoherence scale similarly

with the number of emitters such that the critical coupling required for superradiant burst becomes independent of the number of emitters, in stark contrast with previous conclusions using longer-range power-law interactions. We stress that, apart from the nearest-neighbor model, our classification of a superradiant burst is, strictly speaking, only valid at short times up to $\mathcal{O}[(\gamma_0 t)^2]$ [if $\ddot{R}(0) < 0$ which is true for the models considered here [64]], where the dynamics of the fully excited emitters do not depend on the Hamiltonian. This can be physically justified for later times using second-order mean field theory [64].

The techniques used in this work have broader applications in determining the theoretical bounds for the emission rate of different models, thereby exposing the ultimate limitations of superradiance beyond the NN model. Beyond providing fundamental insights to the physics of superradiance, our results can also motivate the design of atomic lattices in engineered baths such as nanophotonic crystals with engineered interactions or superconducting resonator arrays for qubits. Moreover, hypercube geometries should be within reach of state-of-the-art quantum simulators, given the recent advances in generating arbitrary networks in cavity [76] and circuit [77] quantum electrodynamics platforms.

The authors are grateful to Ben Grossmann, McCoy Lim, Chris Chen, Jasen Zion, Kishor Bharti, Davit Aghamalyan, Lewis Ruks, Thi Ha Kyaw, Tobias Haug, Steven Touzard, Klaus Mølmer and Stuart Masson for insightful discussions. This research is supported by the National Research Foundation (NRF), Singapore, under its Competitive Research Program (CRP) (NRF-CRP25-2020-0004). A. A.-G. gratefully acknowledges support from the Air Force Office of Scientific Research through their Young Investigator Prize (Grant No. 21RT0751), the National Science Foundation through their CAREER Award (No. 2047380), the A. P. Sloan Foundation, and the David and Lucile Packard Foundation.

- [1] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [2] M. Gross and S. Haroche, Superradiance: An essay on the theory of collective spontaneous emission, *Phys. Rep.* **93**, 301 (1982).
- [3] N. E. Rehler and J. H. Eberly, Superradiance, *Phys. Rev. A* **3**, 1735 (1971).
- [4] C. R. Stroud, J. H. Eberly, W. L. Lamo, and L. Mandel, Superradiant effects in systems of two-level atoms, *Phys. Rev. A* **5**, 1094 (1972).
- [5] R. Friedberg, S. Hartmann, and J. Manassah, Limited superradiant damping of small samples, *Phys. Lett.* **40A**, 365 (1972).
- [6] B. Coffey and R. Friedberg, Effect of short-range Coulomb interaction on cooperative spontaneous emission, *Phys. Rev. A* **17**, 1033 (1978).

[7] G. Banfi and R. Bonifacio, Superfluorescence and cooperative frequency shift, *Phys. Rev. A* **12**, 2068 (1975).

[8] J. P. Clemens, L. Horvath, B. C. Sanders, and H. J. Carmichael, Collective spontaneous emission from a line of atoms, *Phys. Rev. A* **68**, 023809 (2003).

[9] G.-D. Lin and S. F. Yelin, Superradiance in spin- j particles: Effects of multiple levels, *Phys. Rev. A* **85**, 033831 (2012).

[10] D. Bhatti, J. von Zanthier, and G. S. Agarwal, Superbunching and nonclassicality as new hallmarks of superradiance, *Sci. Rep.* **5**, 17335 (2015).

[11] P. Longo, C. H. Keitel, and J. Evers, Tailoring superradiance to design artificial quantum systems, *Sci. Rep.* **6**, 23628 (2016).

[12] N. E. Nefedkin, E. S. Andrianov, A. A. Zyablovsky, A. A. Pukhov, A. P. Vinogradov, and A. A. Lisyansky, Superradiance of non-Dicke states, *Opt. Express* **25**, 2790 (2017).

[13] M.-O. Pleinert, J. von Zanthier, and G. S. Agarwal, Hyper-radiance from collective behavior of coherently driven atoms, *Optica* **4**, 779 (2017).

[14] N. Shammah, N. Lambert, F. Nori, and S. De Liberato, Superradiance with local phase-breaking effects, *Phys. Rev. A* **96**, 023863 (2017).

[15] M. Houde, A. Mathews, and F. Rajabi, Explaining fast radio bursts through Dicke's superradiance, *Mon. Not. R. Astron. Soc.* **475**, 514 (2017).

[16] S. J. Masson, I. Ferrier-Barbut, L. A. Orozco, A. Browaeys, and A. Asenjo-Garcia, Many-Body Signatures of Collective Decay in Atomic Chains, *Phys. Rev. Lett.* **125**, 263601 (2020).

[17] S. Fuchs, A. Vukics, and S. Y. Buhmann, Superradiance from nonideal initial states: A quantum trajectory approach, *Phys. Rev. A* **103**, 043712 (2021).

[18] B. Lemberger and K. Mølmer, Radiation eigenmodes of Dicke superradiance, *Phys. Rev. A* **103**, 033713 (2021).

[19] F. Robicheaux, Theoretical study of early-time superradiance for atom clouds and arrays, *Phys. Rev. A* **104**, 063706 (2021).

[20] O. Rubies-Bigorda and S. F. Yelin, Superradiance and subradiance in inverted atomic arrays, *Phys. Rev. A* **106**, 053717 (2022).

[21] D. Malz, R. Trivedi, and J. I. Cirac, Large- n limit of Dicke superradiance, *Phys. Rev. A* **106**, 013716 (2022).

[22] S. J. Masson and A. Asenjo-Garcia, Universality of Dicke superradiance in arrays of quantum emitters, *Nat. Commun.* **13**, 2285 (2022).

[23] A. Piñeiro Orioli, J. K. Thompson, and A. M. Rey, Emergent Dark States from Superradiant Dynamics in Multilevel Atoms in a Cavity, *Phys. Rev. X* **12**, 011054 (2022).

[24] E. Sierra, S. J. Masson, and A. Asenjo-Garcia, Dicke superradiance in ordered lattices: Dimensionality matters, *Phys. Rev. Res.* **4**, 023207 (2022).

[25] N. Skribanowitz, I. P. Herman, J. C. MacGillivray, and M. S. Feld, Observation of Dicke Superradiance in Optically Pumped HF Gas, *Phys. Rev. Lett.* **30**, 309 (1973).

[26] J. M. Raimond, P. Goy, M. Gross, C. Fabre, and S. Haroche, Collective Absorption of Blackbody Radiation by Rydberg Atoms in a Cavity: An Experiment on Bose Statistics and Brownian Motion, *Phys. Rev. Lett.* **49**, 117 (1982).

[27] R. G. DeVoe and R. G. Brewer, Observation of Superradiant and Subradiant Spontaneous Emission of Two Trapped Ions, *Phys. Rev. Lett.* **76**, 2049 (1996).

[28] F. C. Spano and S. Mukamel, Superradiance in molecular aggregates, *J. Chem. Phys.* **91**, 683 (1989).

[29] F. C. Spano, J. R. Kuklinski, and S. Mukamel, Temperature-Dependent Superradiant Decay of Excitons in Small Aggregates, *Phys. Rev. Lett.* **65**, 211 (1990).

[30] F. C. Spano, J. R. Kuklinski, and S. Mukamel, Cooperative radiative dynamics in molecular aggregates, *J. Chem. Phys.* **94**, 7534 (1991).

[31] T. Meier, Y. Zhao, V. Chernyak, and S. Mukamel, Polarons, localization, and excitonic coherence in superradiance of biological antenna complexes, *J. Chem. Phys.* **107**, 3876 (1997).

[32] M. Scheibner, T. Schmidt, L. Worschech, A. Forchel, G. Bacher, T. Passow, and D. Hommel, Superradiance of quantum dots, *Nat. Phys.* **3**, 106 (2007).

[33] V. I. Yukalov and E. P. Yukalova, Dynamics of quantum dot superradiance, *Phys. Rev. B* **81**, 075308 (2010).

[34] C. Bradac, M. T. Johnsson, M. v. Breugel, B. Q. Baragiola, R. Martin, M. L. Juan, G. K. Brennen, and T. Volz, Room-temperature spontaneous superradiance from single diamond nanocrystals, *Nat. Commun.* **8**, 1205 (2017).

[35] G. Haider, K. Sampathkumar, T. Verhagen, L. Nádvorník, F. J. Sonia, V. Valeš, J. Sýkora, P. Kapusta, P. Němec, M. Hof, O. Frank, Y.-F. Chen, J. Vejpravová, and M. Kalbáč, Superradiant emission from coherent excitons in van der Waals heterostructures, *Adv. Funct. Mater.* **31**, 2102196 (2021).

[36] O. Mello, Y. Li, S. A. Camayd-Muñoz, C. DeVault, M. Lobet, H. Tang, M. Lonçar, and E. Mazur, Extended many-body superradiance in diamond epsilon near-zero metamaterials, *Appl. Phys. Lett.* **120**, 061105 (2022).

[37] T. Wang, S. F. Yelin, R. Côté, E. E. Eyler, S. M. Farooqi, P. L. Gould, M. Koštrun, D. Tong, and D. Vrinceanu, Superradiance in ultracold Rydberg gases, *Phys. Rev. A* **75**, 033802 (2007).

[38] G. Ferioli, A. Glicenstein, F. Robicheaux, R. T. Sutherland, A. Browaeys, and I. Ferrier-Barbut, Laser-Driven Superradiant Ensembles of Two-Level Atoms near Dicke Regime, *Phys. Rev. Lett.* **127**, 243602 (2021).

[39] C. Liedl, S. Pucher, F. Tebbenjohanns, P. Schneeweiss, and A. Rauschenbeutel, Collective Excitation and Decay of Waveguide-Coupled Atoms: From Timed Dicke States to Inverted Ensembles, *Phys. Rev. Lett.* **130**, 163602 (2023).

[40] J.-B. Trebbia, Q. Deplano, P. Tamarat, and B. Lounis, Tailoring the superradiant and subradiant nature of two coherently coupled quantum emitters, *Nat. Commun.* **13**, 2962 (2022).

[41] N. Lambert, Y. Matsuzaki, K. Kakuyanagi, N. Ishida, S. Saito, and F. Nori, Superradiance with an ensemble of superconducting flux qubits, *Phys. Rev. B* **94**, 224510 (2016).

[42] Z. Wang, H. Li, W. Feng, X. Song, C. Song, W. Liu, Q. Guo, X. Zhang, H. Dong, D. Zheng, H. Wang, and D.-W. Wang, Controllable Switching between Superradiant and Subradiant States in a 10-Qubit Superconducting Circuit, *Phys. Rev. Lett.* **124**, 013601 (2020).

[43] T. Orell, M. Zanner, M. L. Juan, A. Sharafiev, R. Albert, S. Oleschko, G. Kirchmair, and M. Silveri, Collective bosonic effects in an array of transmon devices, *Phys. Rev. A* **105**, 063701 (2022).

[44] A. González-Tudela, V. Paulisch, D. E. Chang, H. J. Kimble, and J. I. Cirac, Deterministic Generation of Arbitrary Photonic States Assisted by Dissipation, *Phys. Rev. Lett.* **115**, 163603 (2015).

[45] V. Paulisch, M. Perarnau-Llobet, A. González-Tudela, and J. I. Cirac, Quantum metrology with one-dimensional superradiant photonic states, *Phys. Rev. A* **99**, 043807 (2019).

[46] C. Groiseau, A. E. J. Elliott, S. J. Masson, and S. Parkins, Proposal for a Deterministic Single-Atom Source of Quasisuperradiant n -Photon Pulses, *Phys. Rev. Lett.* **127**, 033602 (2021).

[47] M. Perarnau-Llobet, A. González-Tudela, and J. I. Cirac, Multimode Fock states with large photon number: Effective descriptions and applications in quantum metrology, *Quantum Sci. Technol.* **5**, 025003 (2020).

[48] R. Monshouwer, M. Abrahamsson, F. van Mourik, and R. van Grondelle, Superradiance and exciton delocalization in bacterial photosynthetic light-harvesting systems, *J. Phys. Chem. B* **101**, 7241 (1997).

[49] G. D. Scholes, Designing light-harvesting antenna systems based on superradiant molecular aggregates, *Chem. Phys.* **275**, 373 (2002).

[50] G. L. Celardo, P. Poli, L. Lussardi, and F. Borgonovi, Cooperative robustness to dephasing: Single-exciton superradiance in a nanoscale ring to model natural light-harvesting systems, *Phys. Rev. B* **90**, 085142 (2014).

[51] G. Rainò, H. Utzat, M. Bawendi, and M. Kovalenko, Superradiant emission from self-assembled light emitters: From molecules to quantum dots, *MRS Bull.* **45**, 841 (2020).

[52] D. Yang, S.-h. Oh, J. Han, G. Son, J. Kim, J. Kim, M. Lee, and K. An, Realization of superabsorption by time reversal of superradiance, *Nat. Photonics* **15**, 272 (2021).

[53] K. D. B. Higgins, S. C. Benjamin, T. M. Stace, G. J. Milburn, B. W. Lovett, and E. M. Gauger, Superabsorption of light via quantum engineering, *Nat. Commun.* **5**, 4705 (2014).

[54] S. J. Masson and A. Asenjo-Garcia, Atomic-waveguide quantum electrodynamics, *Phys. Rev. Res.* **2**, 043213 (2020).

[55] L. Ruks and T. Busch, Green's functions of and emission into discrete anisotropic and hyperbolic baths, *Phys. Rev. Res.* **4**, 023044 (2022).

[56] S. Cardenas-Lopez, S. J. Masson, Z. Zager, and A. Asenjo-Garcia, Many-body superradiance and dynamical symmetry breaking in waveguide QED, [arXiv:2209.12970](https://arxiv.org/abs/2209.12970).

[57] H. Carmichael and K. Kim, A quantum trajectory unraveling of the superradiance master equation, *Opt. Commun.* **179**, 417 (2000).

[58] F. Mattiotti, M. Kuno, F. Borgonovi, B. Jankó, and G. L. Celardo, Thermal decoherence of superradiance in lead halide perovskite nanocrystal superlattices, *Nano Lett.* **20**, 7382 (2020).

[59] B. Bellomo, G. L. Giorgi, G. M. Palma, and R. Zambrini, Quantum synchronization as a local signature of super- and subradiance, *Phys. Rev. A* **95**, 043807 (2017).

[60] H. F. El-Nashar, Y. Zhang, H. A. Cerdeira, and A. F. Ibiyinka, Synchronization in a chain of nearest neighbors coupled oscillators with fixed ends, *Chaos* **13**, 1216 (2003).

[61] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **48**, 119 (1976).

[62] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2007), p. 656.

[63] D. A. Lidar, Lecture notes on the theory of open quantum systems, [arXiv:1902.00967](https://arxiv.org/abs/1902.00967).

[64] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.130.213605> for additional calculation details, which includes Refs. [65–74].

[65] J. S. Douglas, H. Habibian, C.-L. Hung, A. V. Gorshkov, H. J. Kimble, and D. E. Chang, Quantum many-body models with cold atoms coupled to photonic crystals, *Nat. Photonics* **9**, 326 (2015).

[66] E. Munro, A. Asenjo-Garcia, Y. Lin, L. C. Kwek, C. A. Regal, and D. E. Chang, Population mixing due to dipole-dipole interactions in a one-dimensional array of multilevel atoms, *Phys. Rev. A* **98**, 033815 (2018).

[67] P. Solano, P. Barberis-Blostein, F. K. Fatemi, L. A. Orozco, and S. L. Rolston, Super-radiance reveals infinite-range dipole interactions through a nanofiber, *Nat. Commun.* **8**, 1857 (2017).

[68] P. Lodahl, S. Mahmoodian, S. Stobbe, A. Rauschenbeutel, P. Schneeweiss, J. Volz, H. Pichler, and P. Zoller, Chiral quantum optics, *Nature (London)* **541**, 473 (2017).

[69] H. Pichler, T. Ramos, A. J. Daley, and P. Zoller, Quantum optics of chiral spin networks, *Phys. Rev. A* **91**, 042116 (2015).

[70] W.-K. Mok, D. Aghamalyan, J.-B. You, T. Haug, W. Zhang, C. E. Png, and L.-C. Kwek, Long-distance dissipation-assisted transport of entangled states via a chiral waveguide, *Phys. Rev. Res.* **2**, 013369 (2020).

[71] W.-K. Mok, J.-B. You, L.-C. Kwek, and D. Aghamalyan, Microresonators enhancing long-distance dynamical entanglement generation in chiral quantum networks, *Phys. Rev. A* **101**, 053861 (2020).

[72] S. Mahmoodian, G. Calajó, D. E. Chang, K. Hammerer, and A. S. Sørensen, Dynamics of Many-Body Photon Bound States in Chiral Waveguide QED, *Phys. Rev. X* **10**, 031011 (2020).

[73] F. Robicheaux and D. A. Suresh, Beyond lowest order mean-field theory for light interacting with atom arrays, *Phys. Rev. A* **104**, 023702 (2021).

[74] O. Rubies-Bigorda, S. Ostermann, and S. F. Yelin, Characterizing superradiant dynamics in atomic arrays via a cumulant expansion approach, *Phys. Rev. Res.* **5**, 013091 (2023).

[75] G. H. Golub and C. F. van Loan, *Matrix Computations*, 4th ed. (JHU Press, Baltimore, 2013).

[76] A. Periwal, E. S. Cooper, P. Kunkel, J. F. Wienand, E. J. Davis, and M. Schleier-Smith, Programmable interactions and emergent geometry in an array of atom clouds, *Nature (London)* **600**, 630 (2021).

[77] A. J. Kollár, M. Fitzpatrick, and A. A. Houck, Hyperbolic lattices in circuit quantum electrodynamics, *Nature (London)* **571**, 45 (2019).