ANNUAL REVIEWS

Annual Review of Earth and Planetary Sciences
Reconstructing the
Environmental Context of
Human Origins in Eastern
Africa Through Scientific
Drilling

Andrew S. Cohen, 1,* Christopher J. Campisano, 2,*
J. Ramón Arrowsmith, 3 Asfawossen Asrat, 4,5
Catherine C. Beck, 6 Anna K. Behrensmeyer, 7
Alan L. Deino, 8 Craig S. Feibel, 9 Verena Foerster, 10
John D. Kingston, 11 Henry F. Lamb, 12,13
Tim K. Lowenstein, 14 Rachel L. Lupien, 15
Veronica Muiruri, 16 Daniel O. Olago, 17
R. Bernhart Owen, 18 Richard Potts, 19
James M. Russell, 20 Frank Schaebitz, 10
Jeffery R. Stone, 21 Martin H. Trauth, 22
and Chad L. Yost 21

¹Department of Geosciences, University of Arizona, Tucson, Arizona, USA; email: cohen@email.arizona.edu

²Institute of Human Origins, School of Human Evolution and Social Change, Arizona State University, Tempe, Arizona, USA

³School of Earth and Space Exploration, Arizona State University, Tempe, Arizona, USA

⁴Department of Mining and Geological Engineering, Botswana International University of Science and Technology, Palapye, Botswana

⁵School of Earth Science, Addis Ababa University, Addis Ababa, Ethiopia

⁶Geosciences Department, Hamilton College, Clinton, New York, USA

⁷National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

⁸Berkeley Geochronology Center, Berkeley, California, USA

⁹Department of Earth and Planetary Sciences, Rutgers University, Piscataway, New Jersey, USA

¹⁰Institute of Geography Education, University of Cologne, Cologne, Germany

¹¹Department of Anthropology, University of Michigan, Ann Arbor, Michigan, USA

 $^{12}\mbox{Department}$ of Geography and Earth Science, Aberystwyth University, Aberystwyth, United Kingdom

ANNUAL CONNECT

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Earth Planet. Sci. 2022. 50:451-76

First published as a Review in Advance on February 25, 2022

The Annual Review of Earth and Planetary Sciences is online at earth.annualreviews.org

https://doi.org/10.1146/annurev-earth-031920-081947

Copyright © 2022 by Annual Reviews. All rights reserved

*These authors contributed equally to this article

Keywords

continental scientific drilling, human evolution, East African Rift, paleoclimate

Abstract

Paleoanthropologists have long speculated about the role of environmental change in shaping human evolution in Africa. In recent years, drill cores of late Neogene lacustrine sedimentary rocks have yielded valuable high-resolution records of climatic and ecosystem change. Eastern African Rift sediments (primarily lake beds) provide an extraordinary range of data in close proximity to important fossil hominin and archaeological sites, allowing critical study of hypotheses that connect environmental history and hominin evolution. We review recent drill-core studies spanning the Plio–Pleistocene boundary (an interval of hominin diversification, including the earliest members of our genus *Homo* and the oldest stone tools), and the Mid–Upper Pleistocene (spanning the origin of *Homo sapiens* in Africa and our early technological and dispersal history). Proposed drilling of Africa's oldest lakes promises to extend such records back to the late Miocene.

- High-resolution paleoenvironmental records are critical for understanding external drivers
 of human evolution.
- African lake basin drill cores play a critical role in enhancing hominin paleoenvironmental records given their continuity and proximity to key paleoanthropological sites.
- The oldest African lakes have the potential to reveal a comprehensive paleoenvironmental context for the entire late Neogene history of hominin evolution.

1. INTRODUCTION

For nearly 150 years, since the time of Charles Lyell (1863) and Charles Darwin (1871), scientists have hypothesized about the role of environmental factors, specifically climate, in shaping the evolution of the hominins (humans and our close relatives). What kinds of climate, vegetation, land-scape, and resources did early hominins encounter? Did environmental conditions remain stable over long periods, change directionally, or vary in time and space? How might these changes have impacted selection for characteristics that define humans and our close relatives? From the evolutionary divergence between hominins and our closest relatives, the African great apes, some 8–6 million years ago until the early Pleistocene, this history was exclusively an African chronicle. It is unsurprising, then, that since the early twentieth-century discovery of the first early hominins in South Africa (Dart 1925), extraordinary effort has gone into extracting records of African environmental history relevant to understanding evolutionary innovations, phylogeny, behavioral

¹³Botany Department, School of Natural Sciences, Trinity College Dublin, Ireland

¹⁴Department of Geological Sciences and Environmental Studies, Binghamton University, Binghamton, New York, USA

¹⁵Lamont-Doherty Earth Observatory, Columbia University, Pallisades, New York, USA

¹⁶ National Museums of Kenya, Nairobi, Kenya

¹⁷Department of Geology, University of Nairobi, Nairobi, Kenya


¹⁸Department of Geography, Hong Kong Baptist University, Hong Kong

¹⁹Human Origins Program, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

²⁰Department of Earth, Environmental, and Planetary Sciences, Brown University, Providence, Rhode Island, USA

²¹Department of Earth and Environmental Systems, Indiana State University, Terre Haute, Indiana, USA

²²Department of Geosciences, University of Potsdam, Potsdam, Germany



Conceptual model showing temporal linkages between environmental drivers and diversification/extinction events in human evolution (time advances upward in each case). (a) Null hypothesis: tempo of evolutionary history largely unrelated to environmental history. (b) Speciation/extinction/dispersal linked to strong temporal environmental gradients. (c) Speciation/extinction/dispersal linked to pulses of increased environmental variability. Note that whereas this diagram explicitly references correlations of cladogenetic

events (or lack thereof) to environmental processes, these conceptual linkages could be extended to

noncladogenetic processes such as acquisition of novel behaviors or dispersal events.

change, and dispersal within the hominin lineage. Understanding the relationship between Earth system history and human evolution remains an enduring challenge of broad scientific and public interest.


Possible linkages between hominin evolution and environmental history can be broadly placed in three categories (**Figure 1**). A null hypothesis is that environmental history has had little to do with hominin evolution and that the major drivers of selection, adaptation, diversification, and extinction in hominins were interspecific biotic interactions and intraspecific cultural interactions, rather than changes in climate and environment. If hominin evolution had been generally independent of large-scale external environmental forcing [variants of the Red Queen hypothesis (Van Valen 1973, Foley 1994)], we might expect little or no temporal correlation

between major hominin evolutionary events and environmental histories, beyond those occurring randomly (Figure 1a). However, if environmental change did play a major role in shaping our evolution, we would expect significant correlations between external forcing mechanisms and evolutionary events. These might entail directional shifts in climate, for example toward more arid conditions, driving evolutionary events (Washburn 1960), or as coordinated pulses of diversification and extinction (Vrba 1988) (Figure 1b). Alternatively, adaptive change might have responded to increased environmental variability (Potts 1996) (Figure 1c). Correlation in time alone is insufficient to infer a causal linkage between records of environmental and phylogenetic history; a theoretical understanding of how landscapes and resources would have regulated evolutionary change must underpin the connection (Faith et al. 2021). Whereas current alternative hypotheses make predictions about the temporal linkages between environmental history and hominin evolution, data to test or support different hypotheses remain limited. These temporal correlations, however, are a necessary starting point for supporting or refuting particular hypotheses about environmental history's role in human evolution.

Over the past 25 years science has made significant advances in improving the chronology of both the environmental context of human evolution (deMenocal 1995, 2004; Behrensmeyer 2006; Kingston et al. 2007; Trauth et al. 2007; Donges et al. 2011; Blome et al. 2012; Levin 2015; Potts & Faith 2015) and African hominin phylogeny (Collard & Wood 2000, Kimbel 2015, Haile-Selassie et al. 2016, Strait & Grine 2016, Foley 2017). Our understanding of the former comes from three sources of information: (a) outcrop records at key fossil and archaeological sites (e.g., Bonnefille et al. 2004, Kingston et al. 2007, Cerling et al. 2011, Potts et al. 2018), (b) marine drill cores from offshore Africa (e.g., deMenocal 1995, Bonnefille 2010, Feakins et al. 2013, Uno et al. 2016a), and (c) terrestrial (primarily lacustrine) sediment drill cores from Africa and surrounding regions (e.g., Cohen et al. 2007, 2016; Campisano et al. 2017) (Figure 2). Each of these sources of information has strengths and weaknesses for addressing the hypotheses discussed previously. In this review, we consider the major insights in understanding the environmental context of human evolution that have been recently achieved using drill-core records derived from lacustrine sedimentary successions. These records come from a combination of extant, ancient lakes, such as those of the East African Rift System (EARS), and sediments deposited in lake basins that have long since dried up, filled in, or been destroyed through tectonic processes. Ultimately, understanding the environment/evolution nexus in hominin evolution must rely on a synthesis from all three sources, as they provide complementary information at different temporal and spatial scales, appropriate for addressing different aspects of these hypotheses.

2. INTEGRATING LAKE SEDIMENT CORES WITH OTHER RECORDS OF AFRICAN ENVIRONMENTAL HISTORY

Outcrop-based records documenting conditions at paleoanthropological sites are critical for understanding hominin evolution but present researchers with some fundamental problems in contextualizing this evolutionary history. Most eastern African fossil hominin and archaeological occurrences are in terrestrial deposits, particularly in paleosols and fluvial deposits, recording conditions near where hominins actually lived (e.g., Cerling et al. 2011, Uno et al. 2016b) (Figure 3). However, these deposits are highly episodic and/or time averaged over ecologically long intervals, and they typically do not provide high-resolution records of seasonality or even decadal- or centennial-scale climate events relevant to local hominin populations and their ecosystems. In regions of limited erosion, discontinuous outcrop records also limit our ability to reconstruct long climate records and precisely correlate records regionally. Nevertheless, a paleoenvironmental



Key African and nearby Eurasian terrestrial drilling site locations and prior offshore records overlain on African regional topography and bathymetry. Dates indicate when drilling took place. Abbreviations: DSDP, Deep Sea Drilling Project; HSPDP, Hominin Sites and Paleolakes Drilling Project; ODP, Ocean Drilling Program; OLO, Olorgesailie Drilling Project.




Figure 3

Relative strengths (+ blue text) and weaknesses (- red text) of lake bed drill core, marine drill core, and outcrop paleoenvironmental records and how they can complement each other for contextualizing the role of environmental history in human evolution. Photo of R/V JOIDES Resolution courtesy of Arito Sakaguchi (photographer) and IODP/TAMU; other photos courtesy of Hominin Sites and Paleolakes Drilling Project.

record from the precise deposit in which a fossil was recovered provides highly relevant localized information about a specific paleolandscape in which early hominins lived.

Other sources of paleoenvironmental information likewise have a combination of advantages and disadvantages in terms of contextualizing human origins. Marine drill-core records from the Indian Ocean, the Mediterranean Sea, and other areas surrounding the African continent provide relatively continuous records of climate change over the entire time span of human evolution (e.g., deMenocal 2004, Liddy et al. 2017, Caley et al. 2018, Pollisar et al. 2019). However, their utility is limited by the slow sedimentation rates of the deep ocean and distance from hominin fossil and archaeological sites in the EARS, creating a filter of temporal and spatial averaging in these archives.

It was this combination of archival limitations of outcrops and marine cores that led researchers more than 40 years ago to suggest that drill cores from the large lakes of the EARS might offer an exceptional opportunity to obtain records of humanity's deep-time environmental history, particularly because much of the Plio–Pleistocene paleoanthropological record in Africa has been recovered from these deposystems (Lewin 1981). Drill cores from lacustrine sites that are close to key fossil and archaeological sites provide a wide array of paleoenvironmental records of nearby hominin habitats, typically at annual to decadal resolution (Cohen et al. 2016, Campisano et al. 2017).


Lake deposits offer important advantages for assembling information about hominin habitats. First, because of the calm and often anoxic conditions present in tropical lake floor environments, sediments typically accumulate in laminated beds, undisturbed by intensive bioturbation or current reworking, and thus register annual- or even subannual-scale environmental conditions. Second, an extraordinary range of records can be extracted from lake deposits, reflecting watershed-scale vegetation, geomorphology, internal lake hydrology and ecology, and regional-scale climate. These include paleoecological, geochemical/mineralogical, and sedimentological/facies analysis records, as well as petrophysical data obtained using multi-sensor core loggers. Associated nonlacustrine fluvial, wetland, aeolian, or paleosol deposits, commonly encountered in lake basin drill cores, can also provide information on environmental heterogeneity, although these typically lack the temporal resolution of lake beds. Unweathered drill cores frequently preserve a level of stratigraphic detail and unaltered geochemical information that is difficult or impossible to obtain from nearby outcrops (e.g., Maxbauer et al. 2016) (Figure 4).

This recognition of the value of lake drill-core records for paleoanthropology provided impetus for early drilling campaigns in African lakes. At the meteorite impact crater lake at Tswaing, South Africa (the Pretoria Saltpan), Partridge et al. (1993) demonstrated the role of orbital cyclicity on precipitation cycles in the region of importance to early modern humans. And at Lake Malawi, the significance of documented megadroughts for early modern human demography was an important outcome of drill-core studies (Cohen et al. 2007, Scholz et al. 2007). Efforts to obtain lacustrine drill cores specifically to address paleoanthropological questions began in the mid-2000s, involving a series of conceptual planning workshops (e.g., Cohen et al. 2006). These workshops spurred the development of the multisite Hominin Sites and Paleolakes Drilling Project (HSPDP) (sites in green in **Figure 2**) starting in the late 2000s, along with other, similar drilling projects near the paleoanthropological sites at Olorgesailie, Kenya (Potts et al. 2020), and Olduvai Gorge, Tanzania (Colcord et al. 2019, Deino et al. 2020, Rodríguez-Cintas et al. 2020, Njau et al. 2021).

Following nearly a decade of data collection, our collective experience has shown that our best hope for reconstructing the environments our earliest ancestors inhabited and how those conditions shaped the course of human evolution lies in creative synthesis of outcrop, marine, and lacustrine drill-core records.

3. IMPORTANCE OF GEOCHRONOLOGY FOR TESTING HUMAN EVOLUTION AND ENVIRONMENTAL HISTORY LINKAGES USING LAKE BED CORES

A highly resolved chronostratigraphic framework is essential for developing causal linkages between global climate change, regional environmental responses, and faunal evolution. However, these attempts have often relied on broad temporal correlations with an incomplete fossil or archaeological record. As in outcrop studies, a variety of geochronological techniques are available to study drill-core records, depending on the time span covered by those cores. But dating material derived from drill cores can be complicated compared to outcrops due to uncertainties about drill core—outcrop correlations, limited sample size, and sample artefacts and uncertainties created by the drilling process itself (e.g., core deformation, recoring, and cave-ins). These challenges have led researchers to expand and refine dating techniques to increase the geochronological toolkit available for scientific drilling, particularly addressing the geologic constraints of equatorial (e.g., for paleomagnetic studies), rift-lake settings. For example, Owen et al. (2018) successfully applied U-series dating to authigenic cherts in the HSPDP Lake Magadi drill core. U-series dating was previously applied to Lake Magadi cores, but these results provided only minimum ages for lacustrine sediments, as there was no way at the time to determine whether the dated minerals



Outcrop (a) versus drill core sample (b) (portion of WTK-1A-35Q2; core scale bar in centimeters) from approximate correlative Nachukui Formation lacustrine deposits at West Turkana, Kenya (outcrop is less than 1 km from WTK-1 drill site and \sim 2.5 m high), demonstrating the much greater detail in information extractable from the unweathered core segments, even when compared with shallowly excavated (i.e., archaeologically trenched) surfaces. The core segment shows clear evidence of desiccation surfaces and incipient paleosol formation (intraclastic horizons, marked P) overlain by evidence of abrupt transgressive events marked by finer, laminated muds (T). Only vague lithologic alternations in color and degree of oxidative weathering (responsible for orange and brown outcrop coloration), lacking the millimeter-scale facies changes when observed at the same scale, were visible in the equivalent outcrop section. Photo courtesy of Craig Feibel.

were syndepositional or diagenetic (Goetz & Hillaire-Marcel 1992). Recently, the HSPDP team developed methods for determining which chert units formed syndepositionally and thereby refine the geochronology (Leet et al. 2021). Ultimately, these U-series dates were integrated with ¹⁴C, ⁴⁰Ar/³⁹Ar, and paleomagnetic age control points to develop a comprehensive age model for the core (Owen et al. 2018). Similarly, at Chew Bahir (CHB), multiple geochronometers (¹⁴C, ⁴⁰Ar/³⁹Ar, optically stimulated luminescence, and geochemical fingerprinting of known tephra) were used to develop that core's age model (Roberts et al. 2021).

Another methodological breakthrough for the HSPDP team involved the challenge of determining polarity orientation of samples from near-equatorial drill cores, where inclination is close to zero. This problem is compounded by the rotation of core segments during drilling, which can randomly redistribute declination. Sier et al. (2017, 2021) developed two independent drill-core

reorientation methods for paleomagnetic analysis of samples from the HSPDP West Turkana (WTK) and Baringo-Tugen Hills-Barsemoi (BTB) drill sites.

Tephra from cores that can be geochemically correlated to nearby outcrops and/or well-dated eruptions are extremely valuable, even if those tephra cannot be directly dated (e.g., Lowe 2011; Lane et al. 2017, and references therein). There is a long history of tephrostratigraphic research at fossil sites in eastern Africa (WoldeGabriel et al. 2005, 2013; Brown et al. 2006), and increasingly tephra correlations are being incorporated into drill-core age modeling in eastern African basins (e.g., Lane et al. 2013, 2018; McNamara et al. 2018; Deino et al. 2019, 2020; Roberts et al. 2021). Eastern African tephra have also been correlated to marine drill cores in the Gulf of Aden and the Arabian Sea, providing radiometric or orbitally tuned ages for the horizons (Brown et al. 1992, deMenocal et al. 1999). Several of the HSPDP site age models have incorporated tephra correlations to ⁴⁰Ar/³⁹Ar-dated outcrop horizons. This can be advantageous as, unlike laterally extensive outcrops, drill-core sample sizes are limited by core diameter. Ages derived from thin lake-core tephra that may be poorly preserved in weathered subaerial exposures can also provide age control to outcrops. Four 40Ar/39Ar ages based on outcrop tephra correlations were incorporated into the BTB age model, compared to twelve 40Ar/39Ar ages directly from core material (Deino et al. 2019), whereas all but one of the six WTK 40Ar/39Ar ages were based on tephra correlations (Lupien et al. 2018). Core-to-outcrop tephra correlations have also been established for the Northern Awash HSPDP site (Garello 2019) as well as the Olduvai Gorge Coring Project (Deino et al. 2020, McHenry et al. 2020).

Additional geochronological tools can be deployed to determine exhumation and erosion rates in the watersheds of the lake basins. These provide a means of assessing the roles of topographic relief, tectonics, and climate drivers on sediment sources, and erosion rates over thousand- to million-year timescales during the sampled time periods. Zawacki et al. (2022) used detrital zircon U-Pb double dating of high-temperature crystallization ages and low-temperature exhumation ages on the same samples to characterize sedimentary provenance and transport. Cosmogenic ¹⁰Be derived from quartz-bearing sands at the Northern Awash and WTK sites indicates millennial-scale variability in erosion rates.

Although sedimentation in large, modern lakes is often assumed to be relatively continuous, stratigraphic discontinuities are evident in most lake cores, including those of the HSPDP. Discontinuities can be associated with fluvial incisions, lava flows, erosive volcaniclastics, and subaerial exposure, often with associated pedogenesis of lacustrine sediments (Campisano et al. 2017; Lupien et al. 2018; Deino et al. 2019, 2021). These challenges led Deino et al. (2019) to develop a Bayesian age model for the Olorgesailie drill core that corrected for variable rates of deposition and accommodated different scales of discontinuities. For example, identified paleosols were classified based on a maturity index ensuring that paleosol-based indicator records (e.g., pedogenic carbonates) were not treated as coincident with the lacustrine records (e.g., diatoms) upon which the soil developed. Additionally, to account for the rapid deposition of volcanic material, sedimentary units that included a high proportion of tephra were variably temporally compressed based on sedimentary characteristics.

4. WHAT ARE DRILL-CORE RECORDS TELLING US ABOUT PAST EASTERN AFRICAN ENVIRONMENTS?

4.1. The Plio-Pleistocene Transition in Tropical Africa and Its Significance for Hominin Diversification

The intensification of Northern Hemispheric glaciation associated with the Plio-Pleistocene transition has long been posited as a driving force behind environmental change and mammalian

evolution in Africa, tethered to a shift toward more arid conditions or greater climate variability (e.g., Vrba 1988, 1995; deMenocal 1995, 2004; Potts & Faith 2015). This interval coincides with the first appearance of two hominin genera (*Homo* and *Paranthropus*) and Oldowan stone tools (Harrison 2011, Villmoare et al. 2015, Braun et al. 2019). Outcrop studies from the Baringo Basin demonstrated that equatorial Africa responded to precessional forcing between ~2.7 and 2.55 Ma (Deino et al. 2006, Kingston et al. 2007). The HSPDP BTB and Northern Awash Osi Isi (NAO) and Northern Awash Weranso (NAW) drill-core sites, selected to span the Plio–Pleistocene transition, thus provide unique opportunities to explore how tropical terrestrial communities responded across a major global climate transition and the relative roles of high-latitude boundary conditions and low-latitude insolation forcing of climate.

Rather than showing the most profound change at the Plio-Pleistocene boundary, evidence from the high-resolution BTB core (**Figure 5**) indicates significant changes in multiple records ~3.04 Ma, near the end of the mid-Piacenzian Warm Period (Dowsett et al. 2011), when a major shift in Baringo Basin hydrology occurred (Deino et al. 2021, Westover et al. 2021). Before ~3.04 Ma, environments were dominated by alluvial, fluvial, deltaic, and wetland environments with only brief intervals of deep lakes and probably overfilled depositional systems, as recorded in the high-resolution lithofacies, mineralogy, gamma density, magnetic susceptibility, elemental [X-ray fluorescence (XRF)] geochemistry, and diatom data from the BTB core (Billingsley et al. 2021, Yost et al. 2021). After ~3.04 Ma, a major environmental shift occurred, featuring dramatic swings between deep, freshwater lakes and shallow/marginal lacustrine or subaerial exposure environments, synchronous with precessional periodicity cycles during a high-eccentricity phase (Kingston et al. 2007, Billingsley et al. 2021, Westover et al. 2021). Various records indicate a dramatic change in character at ~3 Ma, from high-frequency, low-amplitude oscillations in the earlier sequence to higher-amplitude, lower-frequency oscillations that intensified over time (Westover et al. 2021).

Vegetation records from the BTB core also show a significant shift at ~3.04 Ma, demonstrating that these changes were not primarily a consequence of rift basin reconfiguration but rather reflect broad climatic changes. Change-point analysis of leaf wax carbon isotopes shows a shift from an average of over 50% C₃ vegetation (trees/shrubs) relative to C₄ grasses prior to ~3.04 Ma to ~40% afterward, suggesting increasing aridity/seasonality. The younger sequence also includes high-amplitude variability that oscillated between ~20% and 100% C₄ plants (Lupien et al. 2019). Phytoliths, which can distinguish C₃ and C₄ grasses and can further differentiate between mesophytic and xerophytic C4 grasses, indicate a shift toward increasing C4 grassland vegetation at ~3.04 Ma in the BTB record (Yost et al. 2021). This record is probably the most detailed, highresolution, and long-duration vegetation reconstruction ever developed from phytoliths. In the pre-3.04 Ma sequence, phytoliths show that woody cover fluctuated at precessional periodicity between open savanna/wooded grassland and woodland/forest habitats (Figure 5). In contrast, the younger sequence is dominated by grasslands and wooded grasslands that include precession-scale oscillations between mesic tall-grass and xeric short-grass habitats (Figure 5). These oscillations were important as less nutritious tall grasses support a different herbivore community than short grasses, hide predators, and promote higher fire frequency and intensity (Dobson 2009; Pays et al. 2012; Hempson et al. 2015a,b).

The Plio–Pleistocene BTB records suggest the onset of major environmental change in eastern Africa was more likely linked to the end of the mid-Piacenzian Warm Period and subsequent cooling than the intensification of Northern Hemisphere glaciation at \sim 2.7 Ma. Furthermore, not only is precession-scale variability observed in these equatorial records, but also such fluctuations appear to have been much larger than the long-term secular changes and responses to high-latitude climate, an observation consistent with a recent synthesis that argues for the




Figure 5

Comparison of the paleoanthropological record across the Plio–Pleistocene transition with stratigraphic, paleoecologic, and geochemical records from the BTB (central Kenya) drill core and Northern Awash (Ethiopia) outcrop records. BTB phytolith reconstructions are shown for C_4 grass type [phytolith index (Iph)] and percent tree cover (D/P° index; grassland category includes grasslands with 0–10% tree cover and wooded grasslands with 10–40% tree cover) (Yost et al. 2021). The BTB leaf wax biomarker ($\delta^{13}C_{\text{wax}}$, dashed green line) is the linearly resampled record of C_3 versus C_4 vegetation (showing a major change point at 3.04 Ma) and independently derived bulk organic matter $\delta^{13}C$ record (brown line) (Lupien et al. 2021). The BTB gamma density is -20 pt. running median with higher values indicative of increased terrigenous inputs and lower values indicative of deeper water and lacustrine accumulation of nonmagnetic and light components (e.g., organic matter or biogenic silica) (Billingsley et al. 2021). The dark red vertical bar is the duration of the mid-Piacenzian Warm Period. Dietary classification of Hadar and Ledi-Geraru fauna from tooth enamel $\delta^{13}C$ shows increasing C_4 consumption, reflecting more open habitats over time (Wynn et al. 2016, Robinson et al. 2017). Comparisons are made with the benthic foraminiferan ^{18}O stack (Lisiecki & Raymo 2005), calculated insolation at June 30°N, and the ODP 721 (Arabian Sea) terrigenous (dust) flux (deMenocal 1995). Abbreviations: BTB, Baringo-Tugen Hills-Barsemoi; FAD, first appearance datum; ODP, Ocean Drilling Program.

importance of low-latitude forcing of Plio–Pleistocene African climate (Trauth et al. 2021a). These extreme, short-term oscillations would have impacted terrestrial resources that mammals, including hominins, relied upon. Forthcoming results of the HSPDP cores from the Northern Awash sites (\sim 3.3–2.9 Ma) will provide a useful comparison to test whether changes noted in the BTB core at \sim 3.04 Ma are local or regional (Campisano et al. 2017). The fossil record from near the Northern Awash site suggests that significant environmental changes also occurred in that area around the same time, as a distinct faunal turnover has been documented at \sim 3.1 Ma with an increase in more arid-adapted mammalian taxa (Reed 2008) and a further shift toward more open, drier, and seasonally variable environments by 2.8 Ma that may have played a role in the emergence of early *Homo* from *Australopithecus* (Robinson et al. 2017).




Figure 6

Comparison of Middle–Late Pleistocene eastern African drill-core and outcrop records, organized from south to north, indicative of changing paleohydrology and paleoclimate over the past 700 ka with key components of the hominin fossil and archaeological record. The Lake Malawi, Malawi (MAL05–1B/1C), Poaceae (grass) pollen record largely reflects aridity. The Olorgesailie Basin, Kenya, land-lake cycles are based on outcrop records discussed in Potts et al. (2020) and the diatom Correspondence Analysis (CA) Axis 1 record of lake depth from the hydrologically connected downstream Koora Basin ODP-OLO12-1A drill-core record (Potts et al. 2020). The Lake Magadi, Kenya (HSPDP-MAG-1A/C), pollen Principal Component Analysis (PCA) Axis 1 aridity record is shown (Owen et al. 2018). The Chew Bahir, southern Ethiopia (HSPDP-CHB14), elemental K/Zr aridity record is driven by the inverse correlation of K flux with precipitation, as illitization is controlled by evaporation (Foerster et al. 2018); data from V. Foerster et al. (unpublished paper). Comparisons are made with calculated insolation at 1°S, the European Project for Ice Coring in Antarctica (EPICA) ice core CO₂ record (Bereiter et al. 2015), and the benthic foraminiferan ¹⁸O stack, including marked timing of the Mid-Brunhes event (*black asterisk*) at ~430 ka (Lisiecki & Raymo 2005). Abbreviations: FAD, first appearance datum; LSA, Late Stone Age; MSA, Middle Stone Age.

4.2. Middle-Late Pleistocene Lake Records and Their Implications for the Evolution of Modern Humans

Numerous critical transitions in African hominin evolution have been documented in the past half-million years, notably the Acheulian to Middle Stone Age (MSA) change in stone tool technology and the origin of our own species, *Homo sapiens* (Stringer 2016, Hublin et al. 2017, Richter et al. 2017, Brooks et al. 2018) (**Figure 6**). Lacustrine cores spanning this interval provide an excellent opportunity to understand how Middle to Late Pleistocene environmental change may have influenced the physical and cultural evolution of *H. sapiens* in eastern Africa. Results from these studies suggest that over the past several hundred thousand years, hominins and other fauna experienced temporal and spatial heterogeneity in ecological resources (e.g., freshwater supply,

dietary breadth, vegetation structure) coincident with critical transitions in their evolutionary history.

Lake Malawi, in the southern subtropics, was the first African rift lake to be drilled (Scholz et al. 2007). The core record from this lake provides evidence of extraordinary lake-level and climatic fluctuations (Lyons et al. 2015), some of which were in response to eccentricity-modulated precession cycles over the past 800 kyr (Ivory et al. 2016). Malawi pollen data over the past 600 kyr mirror lake-level records, with forest expansion coincident with increases in water level, demonstrating these fluctuations were climatically driven. Rapid fluctuations occurred between forested and semiarid phases, particularly during high precession variability phases (Ivory et al. 2018). In conjunction with nearby (25–150 km) MSA archaeological and geomorphic data, the Malawi pollen and charcoal records also document how early anthropogenic fire impacts, coupled with climate changes, drove landscape and ecological changes (Thompson et al. 2021). Early human use of fire appears to have altered the preagricultural anthropogenic landscape through changes in the seasonality and intensity of ignitions, which in turn influenced both vegetation composition and erosion rates.

Southern Kenyan Rift drill sites, ~900 km north of Lake Malawi, highlight the importance of spatial variation for regional ecosystem change. Olorgesailie, in southern Kenya, has a rich archaeological and paleontological record spanning the past 1.2 million years (Potts et al. 2018). Outcrops preserve evidence of the replacement of the Acheulean by MSA innovations between 500 and 320 ka, including smaller, more diverse, and complex tool types, a change in raw material type and transport distance, and the use of pigments (Brooks et al. 2018, Deino et al. 2018, Potts et al. 2018). A contemporaneous faunal turnover is recorded by the loss of large and mega-herbivores, particularly water-dependent mega-grazers, and an increase in smaller, water-independent mixed grazing or browsing herbivores (Potts & Deino 1995, Potts et al. 2018). An erosional disconformity in the Olorgesailie Basin outcrops spans 500-320 ka, coincident with these transitions, but this interval is covered by Olorgesailie Drilling Project cores in the adjacent Koora Basin, so the precise timing of these cultural and biotic transitions remains uncertain (Deino et al. 2019, Potts et al. 2020). This record shows relatively stable environmental conditions until ~450-400 ka, followed by significantly more variable and arid conditions. This includes changes in basin-scale water balance that produced substantial fluctuations in lake depth and water quality, from large, fresh, and deep to fluctuating saline and shallow. The frequency of short, dry episodes also increased, represented by evidence of subaerial exposure and soil formation. Vegetation structure also shifted, from predominantly C₄ short-grass (xeric) woodland to stronger fluctuations between woody cover and C₄ grass dominance, and then a shift to more mesic C₄ tall-grass dominance (Potts et al. 2020, Lupien et al. 2021).

The nearby Lake Magadi drill-core site (~20 km from the Koora core) was chosen because it likely experienced the same regional climatic conditions, but with a different tectonic and hydrologic history. Comparing the two core sites allows us to untangle tectonic from climatic signals. The Magadi core records a shift from early wetter conditions and a freshwater lake stage (particularly ~750–500 ka) to a stratified, saline lake reflecting a long-term trend toward a more arid climate (Owen et al. 2018, Muiruri et al. 2021a). The geochemical, diatom, and pollen data indicate a directional trend over the past 500 kyr toward more arid conditions, but with many wet/dry cycles and of increasing variability superimposed upon the longer-term drying trend. Intense aridification took place between ~525 and 400 ka, coincident with the Mid-Brunhes increase in amplitude of glacial/interglacial cycles (430 ka), with relatively persistent arid conditions from ~350 ka to present (Owen et al. 2018, 2019; Muiruri et al. 2021a,b). Drier episodes, reflected in the diatom and pollen data, mostly occurred during periods of low-amplitude insolation variability (Muiruri et al. 2021a,b). Notably, the Magadi wet/dry cycles are generally antiphased with the

Malawi record (Muiruri et al. 2021b). Such interhemispheric antiphased relationships have also been recorded and modeled for the Late Pleistocene (Blome et al. 2012, Kutzbach et al. 2020).

The timing of environmental shifts at Magadi is consistent with shifts in the Olorgesailie core record, but the Olorgesailie record does not exhibit a directional trend in aridity after 400 ka (Potts et al. 2020). The difference between these records and some of the changes in water availability and vegetation at Olorgesailie may have been the result of spatial heterogeneity related to local tectonics and increased topographic relief (Behrensmeyer et al. 2018, Potts et al. 2020). Moreover, as a regional drainage sump and terminal saline lake, Magadi may have been more sensitive to shifts in water balance. Nonetheless, the overall similarity in timing of environmental change recorded in the two basins suggests they were responding to changes in regional and global climate (**Figure 6**).

Records from the Chew Bahir basin (southern Ethiopia, ~5°N) core provide another key data point for understanding regional and latitudinal variation of environmental change with implications for hominin evolution and dispersal in eastern Africa since ~620 ka (Foerster et al. 2012; Viehberg et al. 2018; Duesing et al. 2021; Roberts et al. 2021; Schaebitz et al. 2021; Trauth et al. 2021b; V. Foerster et al., unpublished paper). A change-point analysis of the CHB K/Zr record demonstrates a three-phased history of significant changes in hydroclimate driven by orbitally forced variations in low-latitude insolation with general trends similar to those observed in the southern Kenya HSPDP site records (Schaebitz et al. 2021; V. Foerster et al., unpublished paper). The interval between ~620 and 410 ka exhibits a long-term drying trend culminating in fluctuations between hydrological extremes. A wetter phase between ~410 and 210 ka, increasingly pulsed by rapid oscillations on precessional timescales, coincides with the emergence of modern H. sapiens and the MSA in Africa (V. Foerster et al., unpublished paper). The most recent phase, \sim 210 ka to present, shows a long-term drying trend with increasing intensity since \sim 125 ka, superimposed on the highest variability between wet and dry pulses observed in the whole record. The humid pulses of this phase (until ~ 60 ka) may have opened migration corridors for H. sapiens out of Africa. The driest phase (~60-14 ka) during lowest eccentricity shows millennial- to centennial-scale wet/dry fluctuations, associated with Dansgaard-Oeschger and Heinrich events, which could have spurred the development of new adaptive strategies by H. sapiens, such as the innovation of tool technologies (Late Stone Age) or migration to the adjoining highlands (Schaebitz et al. 2021).

A pan-African synthesis of multiple high-resolution core records from both lake and marine settings spanning the past 620 kyr demonstrates an east-west alternation in precipitation regimes across Africa (Kaboth-Bahr et al. 2021). Generally wet conditions prevailed in western Africa and more arid conditions in eastern Africa between ~525 and 279 ka and ~128 and 0 ka, with a reversal of this mode during the intervening period. Kaboth-Bahr et al. argue that these alternations were driven by warming and cooling in the tropical Pacific, driving east-west positional changes in the Walker circulation on eccentricity-linked timescales. This continental-scale alternation in climate and vegetation patterns coincides with molecular phylogenetic evidence for diversification events in pan-African mammalian clades, and would have had profound effects on the resources upon which mammals including early *H. sapiens* depended, as well as their biogeography as those resources shifted.

4.3. Testing the Hypothesized Impact of the Toba Supereruption on Terrestrial Ecosystems and Anatomically Modern Human Populations in Africa Using the Lake Malawi Drill-Core Record

Our ability to resolve environmental events down to annual or even subannual timescales in laminated lake beds makes these records particularly well suited for answering questions about the impact of extreme events on human populations, how seasonality may have influenced early hominin habitats, and the nature of the landscapes particular hominins occupied. One of the most intriguing ideas linking early modern human demography and dispersal to environmental catastrophes is the volcanic winter/weak Garden of Eden hypothesis, linking the Mt. Toba (Indonesia) supercruption to a global climate catastrophe, and subsequently to the collapse of human populations in Africa (Ambrose 1998). The youngest (~74 ka) eruption of Mt. Toba is the largest Quaternary eruption known. Its fingerprinted tephra [the Youngest Toba Tuff (YTT)] can be found across a wide swath of Africa and Eurasia in marine and continental sediments (Smith et al. 2011, 2018), and possibly in Greenland and Antarctic ice core aerosols as well (Svensson et al. 2013). The approximate concurrence of this eruption with molecular genetic evidence for a possible early modern human population bottleneck, as well as the presumed timing of the dispersal of modern humans out of Africa, prompted a lively debate on the possible connection between these events (e.g., Rampino & Ambrose 2000, Oppenheimer 2002, Robock et al. 2009, Williams 2012, Smith et al. 2018). This is difficult to test in most terrestrial occurrences, however, because the linkage between the tephra and coincident paleoenvironmental records is limited by episodic deposition.

Lane et al. (2013) recovered YTT cryptotephra from two of the Lake Malawi drill cores and argued, based on both TEX₈₆ paleotemperature reconstructions and scanning XRF data, that there is no evidence supporting a major environmental perturbation or volcanic winter associated with the tephra horizon. Subsequently, Jackson et al. (2015) and Yost et al. (2018) took advantage of the wide range of paleoecological data preserved in the laminated Malawi cores at subdecadal resolution for several hundred years before and after the eruption to evaluate the volcanic winter hypothesis. They confirmed that neither lacustrine nor terrestrial ecosystems suffered the types of impacts predicted by a hypothesized volcanic winter severe enough to have catastrophic effects on human populations and generate a genetic bottleneck (Rampino & Ambrose 2000, Robock et al. 2009) (Figure 7). Given the degree of hypothesized cooling proposed by the volcanic winter hypothesis, Lake Malawi, a thermally stratified tropical lake, would have undergone a complete limnological overturn. Complete water column mixing would have had an immediate and longlasting impact on a lake of this size by nutrients released into the epilimnion and oxygenation of lake floor. Jackson et al. (2015) and Yost et al. (2018) found no such major perturbations in the fossil record of dominant phytoplankton or zooplankton following the YTT, nor evidence of colonization of the normally anoxic (and thus unbioturbated) lake floor by benthic invertebrates at this time. The terrestrial records of vegetation (from phytolith fossils) and fire (from charcoal abundance) across the YTT horizon are similarly instructive. There is no evidence for the types of wholesale environmental degradation required by the volcanic winter hypothesis to cause a collapse of the ecosystems upon which contemporaneous modern humans depended, as documented by archaeological sites in this region (Thompson et al. 2018, 2021).

The two Malawi drill cores in which the YTT was encountered also allowed Yost et al. (2018) to differentiate low from high elevation impacts of the ashfall. In the central Lake Malawi core (MAL-1C), adjacent to low-elevation miombo woodland, no changes were observed immediately following the YTT in tree cover, or in either cool climate C₃ or warmth-indicating C₄ grasses that exceeded background (i.e., pre-Toba) variability. In the north basin (MAL-2A), adjacent to the Rungwe and Livingstone Mountains, there is evidence of increased charcoal and C₄ grasses immediately after the YTT, signaling some burning and aridification predicted by the volcanic winter hypothesis. However, the levels of these indicators remained well within the background levels of variability prior to the YTT, making it extremely unlikely these climatic effects were sufficiently severe to cause major human population declines across Africa.




Figure 7

Indicators of lake ecology and mixing, terrestrial vegetation, and fire from the Lake Malawi drill core spanning an interval of \sim 100 years before and \sim 200 years after the Mt. Toba supereruption [Youngest Toba Tuff (YTT) horizon] recovered from the GLAD7-MAL05-1C-8H-1 core section with cryptotephra chemically matched to the YTT (Lane et al. 2013). Data are from Jackson et al. (2015) and Yost et al. (2018) and are plotted on a depth scale. Plotted data include counts per dry gram of *Pediastrum* (green algae–photic zone mixing); *Chaoborus* (phantom midge larvae, deep water mixing plus overall secondary productivity); a diatom mixing index [(*Aulacoseira* + *Stephanodiscus*)/*Cyclotella*]; and microcharcoal (<45 μ m) and macrocharcoal (>45 μ m) (regional and local fire activity, respectively). Phytolith-derived climate and vegetation indices are based on grass phytoliths [climate index (Ic), phytolith index (Iph), and the ratio of woody plant to grass phytoliths (D/P)]. Greenland Ice Sheet Project 2 (GISP2) δ 18O values are from Grootes & Stuiver (1997). All records show an absence of significant perturbations in lake or land ecology associated with the YTT beyond background variability. The red dashed line marks the position of first occurrence of YTT cryptotephra in the Lake Malawi core and links it to the position of the GISP2 H₂SO₄ spike attributed to the Toba supereruption (Zielinski et al. 1996), which shows a minor cooling event associated with this eruption against a background decline in temperatures underway long before the eruption.

5. EVALUATING THE ROLE OF ENVIRONMENTAL VARIABILITY IN HUMAN EVOLUTION WITH CORE RECORDS

Since the late 1990s there has been considerable interest among paleoanthropologists in the potential role that environmental variability may have played as a driver of evolution, technological change, and dispersal of hominins. In a series of papers, Potts (1996, 1998) argued that increased variability (as opposed to long-term directional shifts, for example toward drier conditions) resulted in adaptations that conferred greater flexibility during times of increased environmental uncertainty, potentially leading to significant species turnover. Potts termed this variability selection. As examples in the hominin lineage, Potts pointed to the expansion of arguably more adaptively flexible *Homo erectus* at the expense of dietarily specialized *Paranthropus*

(during the climatically variable early Pleistocene) or the replacement of cold climate–adapted Neanderthals by rapidly expanding and behaviorally diverse modern *H. sapiens* in Eurasia through the most extreme climate swings of the late Pleistocene. Potts (1998) recognized that testing this hypothesis required evidence that the evolution of adaptations for increased environmental flexibility be both temporally and logically (in terms of specific adaptations) associated with evidence of enhanced environmental variability. Highly resolved drill-core records are well suited for providing evidence of changes in environmental variability.

Potts & Faith (2015) evaluated the variability selection hypothesis's predictions using orbital insolation cycles to predict periods of high and low climate variability. The insolation record was then compared quantitively against Arabian Sea and Mediterranean dust records, which demonstrated that high- and low-variability orbital insolation intervals corresponded to actual increases and decreases in the two indicator records. They then showed that qualitative changes in paleoen-vironmental records from a series of hominin fossil-bearing basins in eastern Africa corresponded to the timing of variability stages predicted by the orbital-control model of variability. Finally, they compared the times of high variability, as predicted by the variability selection hypothesis, with first and last appearance datums (FADs and LADs) for a variety of hominin evolutionary transitions to see if these corresponded in time. Their null model, comparable to Figure 1a, predicts no statistical relationship between these events, but they found a significant relationship between the two, as would be predicted if evolutionary novelty were driven by increased climate variability (Figure 1c).

More recently, Potts and colleagues (2018, 2020) have emphasized the possible role of nonclimatic aspects of environmental variability in affecting resource fluctuations and their temporal association with hominin technological change. They note that the Acheulian to MSA transition at Olorgesailie, linked to the greater technological capacity of early modern humans to procure resources over greater areas and with greater cognitive flexibility, is temporally associated in the nearby Koora drill-core record with increasing hydroclimate variability. However, the Koora record of water resource variability is itself not strongly associated with orbital cyclicity, but rather with possible tectonically and volcanically induced changes in local hydrology and spatial fragmentation of resources.

Several terrestrial core records show correspondences between important human evolutionary transitions and evidence of high hydroclimate, vegetation, or landscape variability. Lupien et al. (2018) identified a dramatic episode of high variability in δD and $\delta^{13}C$ of leaf waxes in the WTK drill-core record at 1.73 Ma, a time of high eccentricity. This timing corresponds with the last appearance of *Homo rudolfensis*, the first appearance of *H. erectus*, the dispersal of *Homo* spp. out of Africa, and the earliest evidence for Acheulian (e.g., hand axe) technology (Spoor et al. 2007, 2015; Lepre et al. 2011; Antón et al. 2014). Abrupt, high-amplitude climate variability, as registered in the Magadi (Owen et al. 2018), Koora (Potts et al. 2020), and CHB (Duesing et al. 2021; V. Foerster et al., unpublished paper) drill-core records, accompanied the technological transition from the Acheulian to the MSA. This is marked by evidence of increasing human cognition through features such as production of smaller and more complex tools, evidence for greater transport distances of raw materials, and the appearance of pigments, possibly for adornment. The significance of these temporal correspondences needs to be tempered by the fact that hominin fossil sample sizes remain small and geographically restricted; thus, uncertainties of the timing of specific hominin evolutionary transitions remain large (Faith et al. 2021).

Environmental variability is not a singular concept, and different tempos and modes of variability have been theorized to have different effects on hominin resources and populations. Trauth et al. (2021b) quantified different types of variability using recurrence quantification analysis, which allows the recognition of tipping points in hydroclimate (and thus ecosystems and hominin

resources) that may result from relatively small changes in the climate state. They analyzed the CHB potassium (aridity) and Ocean Drilling Project ODP Site 967 wet/dry index records and recognized two types of variability. One of these modes shows maximum variability at precessional frequencies and is accompanied by shorter (centennial- and millennial-scale) changes between wet and dry states. The second type of variability shows a weaker association with orbital parameters and occurs during insolation variability minima by rapid (years to decades) changes between wet and dry conditions, which increase in frequency through the entire phase of this type of variability. Trauth et al. (2021b) argued that the long-term variability associated with the first type would have corresponded to longer timescales, potentially associated with human evolutionary events and large-scale dispersal, whereas the short-term fluctuations of the second type would have more directly influenced differential mortality and fertility and population occupation of more favorable nearby habitats, as discussed by Schaebitz et al. (2021) and V. Foerster et al. (unpublished paper).

Modeling experiments are useful for understanding these potential effects of environmental variability on population growth and dispersal. Grove (2014) suggested that phenotypic plasticity (i.e., variability in phenotypes within a population allowing a broadening of habitat or dietary tolerances) could be expected to increase during periods of environmental instability. This accumulated plasticity could provide a pump for successful dispersal when a period of low climatic instability immediately follows a period of high instability. Grove et al. (2015) modeled favorable periods of dispersal for early H. sapiens out of Africa based on episodes of stability following extreme instability registered in the Lake Tana, Ethiopia, drill core Ca/Ti record (an indicator of relative moisture). They argued that the most favorable intervals for dispersal would have been between ~103 and 97 ka, consistent with evidence for early H. sapiens occupation in the Levant at the time. Lamb et al. (2018), evaluating the same Lake Tana paleoclimate record, noted three intervals favorable to dispersal (150-144, 125-93, and 82-73 ka) that also predate the latest age (~65–55 ka) from genetic evidence for the common ancestry of non-African H. sapiens (Nielsen et al. 2017). Some of these intervals were not marked by records of stability following extreme instability elsewhere in Africa, suggesting that if Grove's (2014) accumulated plasticity model is correct, the dispersal pump must have operated erratically around Africa.

Two important concerns can be raised about these comparisons between individual paleoclimate records of variability and the fossil/archaeological record of hominin evolution. First, it is unknown whether the individual records of variability reflect environmental conditions in the area(s) where the evolutionary transitions occurred. Second, it is well-known that FADs and LADs provide only minimal temporal bounds for the duration of any species or technology (e.g., Faith et al. 2021). This is problematic for extinct hominin species, for which the confidence intervals surrounding the FADs and LADs are very large. To address these uncertainties and further test the long-term relationships between environmental variability and hominin evolution, future studies will investigate how regionally stacked variability records from around Africa compare with the tempo of mammalian evolution and extinction during the Plio–Pleistocene, recognizing the inherent uncertainties of the fossil record.

6. FUTURE DIRECTIONS APPLYING LACUSTRINE DRILL-CORE RECORDS TO UNDERSTANDING HUMAN EVOLUTION

Research integrating paleoenvironmental reconstructions from drill-core records into our understanding of human evolution in Africa is still a work in progress. The cores described here are still under study, and as new approaches to dating these cores or reconstructing past climates and ecosystems are developed, we expect all of them to be mined more intensively for clues about the habitats of early humans. Thus, careful preservation of these valuable materials in facilities such as the Continental Scientific Drilling Facility (https://cse.umn.edu/csd) is essential for maintaining their future value.

Beyond retrieving new information from existing cores, projects such as HSPDP provide a proof of concept for the transformative value of scientific drilling in ancient lake deposits for paleoanthropology. Three future directions for this science hold particular promise. First, ancient African lakes provide an extraordinary geographic range across which records can be collected and compared. It is evident even from the limited data gathered from lacustrine drill cores to date that large spatiotemporal differences exist between these records. To what extent can these be understood using existing paleoclimate models (e.g., Kutzbach et al. 2020) in terms of the relative impacts of low-latitude insolation versus high-latitude, glacial/interglacial forcing? While the evidence from lake core records makes the importance of orbital forcing for long-term climate variability in tropical Africa clear (e.g., Deino et al. 2021), strong evidence from African lake cores and climate models also exists for high-latitude forcing, particularly for abrupt/high-impact events (e.g., Tierney et al. 2008, Otto-Bliesner et al. 2014). Broad-scale syntheses are needed to evaluate the relative importance of these drivers across and adjacent to Africa (i.e., integration with core records from western Asia). From the paleoanthropological perspective, understanding the dynamics of these changes will clarify how and when various parts of the African continent, as well as corridors of dispersal out of Africa, became more or less favorable to hominins.

Second, the value of scientific drilling for addressing paleoanthropological questions is by no means restricted to studies in Africa. Many sites already drilled in Eurasia (**Figure 2**) have been incorporated into our understanding of hominin expansion out of Africa (e.g., Miebach et al. 2019). In the future we can expect similar integration of lacustrine drill-core records in deeper time, drawing on the many ancient lakes of Eurasia, to explore the habitats of Pleistocene hominins other than anatomically modern humans (AMHs), or the environmental context of the expansion of AMHs out of Africa.

Third, hopefully in the near future we can expect much longer records of African environmental history to be obtained from lake beds, spanning the entire 8–6 Ma time frame of hominin evolution. Projects in advanced stages of development from the oldest African lakes, Lakes Tanganyika (Cohen & Salzburger 2016, Russell et al. 2020) and Chad (Sylvestre et al. 2018), promise to provide records of African paleoclimate and ecosystem history since the Late Miocene from two very different regions of the continent (**Figure 2**). Along with other possible future drilling sites in other African Great Lakes, we anticipate these long records will yield tremendous advances in the emerging science of using scientific drilling to further our understanding of the deep-time environmental history of our distant ancestors. Understanding this history of environment and human evolution holds important implications for understanding our own adaptability to climate change into the future.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

AUTHOR CONTRIBUTIONS

A.S.C. and C.J.C. conceived the framework and coordinated the writing of this review and drafted the initial version of the text. Figures were produced by A.S.C., C.J.C., C.C.B., C.S.F., R.B.O., M.H.T., and C.L.Y. All authors made critical comments and substantive changes throughout the review.

ACKNOWLEDGMENTS

We thank our many colleagues of the Hominin Sites and Paleolakes Drilling Project (HSPDP) and Lake Malawi Drilling Project, whose research has contributed extensively to this review. We also thank the International Continental Drilling Program (ICDP) Operations Support Group, the National Lacustrine Core Facility (LacCore, now the Continental Scientific Drilling Facility), DOSECC Inc. and DOSECC Exploration Services for technical support with these projects, and the governments of Malawi, Kenya, and Ethiopia for authorizing this research. Financial support for this research was provided by ICDP, the National Science Foundation (EAR-0602350, EAR-1123942, BCS-1241859, EAR-1338553, BCS-1241790, and EAR-1322017), the Natural Environment Research Council (NE/K014560/1), Deutsche Forschungsgemeinschaft (DFG) priority program SPP 1006 (F.S.: SCHA 472/13 and /18; M.H.T.: TR 419/8, /10 and /16) and the Collaborative Research Center (CRC) 806, DFG Project Number 57444011 at the University of Cologne, the Hong Kong Research Grants Council (HKBU12300815; 12304018), the Peter Buck Fund for Human Origins Research (Smithsonian Institution), the William H. Donner Foundation, the Ruth and Vernon Taylor Foundation, Whitney and Betty MacMillan, the Institute of Human Origins (IHO, Arizona State University), and the Smithsonian's Human Origins Program. This is publication #45 of the HSPDP. This paper is dedicated to our deceased colleagues Andrew Hill, John Kutzbach, Charlie Lockwood, Jean-Jacques Tiercelin, and Mohammed Umer, all of whose contributions have been critical to the success of HSPDP at various stages of this project.

LITERATURE CITED

- Ambrose S. 1998. Late Pleistocene human population bottlenecks, volcanic winter, and differentiation of modern humans. 7. Hum. Evol. 34:623–51
- Antón SC, Potts R, Aiello LC. 2014. Evolution of early Homo: an integrated biological perspective. Science 345(6192):1236828
- Behrensmeyer AK. 2006. Climate change and human evolution. Science 311:476-78
- Behrensmeyer AK, Potts R, Deino A. 2018. The Oltulelei Formation of the southern Kenyan Rift Valley: a chronicle of rapid landscape transformation over the last 500 k.y. Geol. Soc. Am. Bull. 130(9/10):1474–92
- Bereiter B, Eggleston S, Schmitt J, Nehrbass-Ahles C, Stocker TF, et al. 2015. Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophys. Res. Lett.* 42:542–49
- Billingsley A, Reinthal P, Dettman DL, Kingston JD, Deino AL, et al. 2021. 8¹³C records from fish fossils as paleo-indicators of ecosystem response to lake levels in the Plio-Pleistocene lakes of Tugen Hills, Kenya. Palaeogeogr: Palaeoclimatol. Palaeoccol. 570:109320
- Blome MW, Cohen AS, Tryon CA, Brooks AS, Russell J. 2012. The environmental context for the origins of modern human diversity: a synthesis of regional variability in African climate 150,000–30,000 years ago. *7. Hum. Evol.* 62:563–92
- Bonnefille R. 2010. Cenozoic vegetation, climate changes and hominid evolution in tropical Africa. *Glob. Planet. Change* 72:390–411
- Bonnefille R, Potts R, Chalié F, Jolly D, Peyron O. 2004. High-resolution vegetation and climate change associated with Pliocene *Australopithecus afarensis*. PNAS 101:12125–29
- Braun DR, Aldeias V, Archer W, Arrowsmith JR, Barakif N, et al. 2019. Earliest known Oldowan artifacts at >2.58 Ma from Ledi-Geraru, Ethiopia, highlight early technological diversity. *PNAS* 116:11712–17
- Brooks AS, Yellen JE, Potts R, Behrensmeyer AK, Deino AL, et al. 2018. Long-distance stone transport and pigment use in the earliest Middle Stone Age. *Science* 360:90–94
- Brown FH, Haileab B, McDougall I. 2006. Sequence of tuffs between the KBS Tuff and the Chari Tuff in the Turkana Basin, Kenya and Ethiopia. 7. Geol. Soc. 163:185–204
- Brown FH, Sarna-Wojcicki AM, Meyer CE, Haileab B. 1992. Correlation of Pliocene and Pleistocene tephra layers between the Turkana Basin of East Africa and the Gulf of Aden. *Quat. Int.* 13–14:55–67
- Caley T, Extier T, Collins JA, Schefuß E, Dupont L, et al. 2018. A two-million-year-long hydroclimatic context for hominin evolution in southeastern Africa. Nature 560:76–79

- Campisano CJ, Cohen AS, Arrowsmith JR, Asrat A, Behrensmeyer AK, et al. 2017. The Hominin Sites and Paleolakes Drilling Project: acquiring high-resolution paleoclimate records from the East African Rift System and their implications for understanding the environmental context of hominin evolution. Paleoanthropology 2017:1–43
- Cerling TE, Wynn JG, Andanje SA, Bird MI, Korir DK, et al. 2011. Woody cover and hominin environments in the past 6 million years. *Nature* 476:51–56
- Cohen A, Campisano C, Arrowsmith R, Asrat A, Behrensmeyer AK, et al. 2016. The Hominin Sites and Paleolakes Drilling Project: inferring the environmental context of human evolution from eastern African rift lake deposits. Sci. Drill. 21:1–16
- Cohen AS, Ashley GM, Potts R, Behrensmeyer AK, Feibel C, Quade J. 2006. Paleoclimate and human evolution workshop. Eos 87:161–62
- Cohen AS, Salzburger W. 2016. Scientific drilling at Lake Tanganyika, Africa: a transformative record for understanding evolution in isolation and the biological history of the African continent, University of Basel, 6–8 June 2016. Sci. Drill. 22:43–48
- Cohen AS, Stone JR, Beuning KR, Park LE, Reinthal PN, et al. 2007. Ecological consequences of early Late Pleistocene megadroughts in tropical Africa. PNAS 104:16422–27
- Colcord DE, Shilling AM, Freeman KH, Njau JK, Stanistreet IG, et al. 2019. Aquatic biomarkers record Pleistocene environmental changes at Paleolake Olduvai, Tanzania. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 524:250–61
- Collard M, Wood BA. 2000. How reliable are human phylogenetic hypotheses? PNAS 97:5003-6
- Dart R. 1925. Australopithecus africanus: the man-ape of South Africa. Nature 115:195-99
- Darwin C. 1871. The Descent of Man and Selection in Relation to Sex. London: John Murray
- Deino AL, Behrensmeyer AK, Brooks AS, Yellen JE, Sharp WD, Potts R. 2018. Chronology of the Acheulean to Middle Stone Age transition in eastern Africa. *Science* 360(6384):95–98
- Deino AL, Dommain R, Keller CB, Potts R, Behrensmeyer AK, et al. 2019. Chronostratigraphic model of a high-resolution drill core record of the past million years from the Koora Plain, south Kenya Rift: overcoming the difficulties of variable sedimentation rate and hiatuses. *Quat. Sci. Rev.* 215:213–31
- Deino AL, Heil CJ, King J, McHenry LJ, Stanistreet IG, et al. 2020. Chronostratigraphy and age modeling of Pleistocene drill cores from the Olduvai Basin, Tanzania (Olduvai Gorge Coring Project). Palaeogeogr. Palaeoclimatol. Palaeoecol. 571:109990
- Deino AL, Kingston JD, Glen JM, Edgar RK, Hill A. 2006. Precessional forcing of lacustrine sedimentation in the late Cenozoic Chemeron Basin, Central Kenya Rift. Earth Planet. Sci. Lett. 247:41–60
- Deino AL, Sier MJ, Garello D, Keller B, Kingston J, et al. 2021. Chronostratigraphy of the Baringo-Tugen-Barsemoi (HSPDP-BTB13-1A) core—40 Ar/39 Ar dating, magnetostratigraphy, tephrostratigraphy, sequence stratigraphy and Bayesian age modeling. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 570:109519
- deMenocal PB. 1995. Plio-Pleistocene African climate. Science 270:53-59
- deMenocal PB. 2004. African climate change and faunal evolution during the Pliocene–Pleistocene. *Earth Planet. Sci. Lett.* 220:3–24
- deMenocal PB, Brown FH, Agusti J, Rook L, Andrews P. 1999. Pliocene tephra correlations between East African hominid localities, the Gulf of Aden, and the Arabian Sea. In *Hominid Evolution and Climatic Change in Europe*, ed. J Agusti, L Rook, P Andrews, pp. 23–54. Cambridge, UK: Cambridge Univ. Press
- Dobson A. 2009. Food-web structure and ecosystem services: insights from the Serengeti. *Philos. Trans. R. Soc.* B 364:1665–82
- Donges JF, Donner RV, Trauth M, Marwan N, Schnellnhuber HJ, Kurths J. 2011. Nonlinear detection of paleoclimate-variability transitions possibly related to human evolution. PNAS 108:20422–27
- Dowsett HJ, Haywood AM, Valdees PJ, Robinson MM, Lunt DJ, et al. 2011. Sea surface temperatures of the mid-Piacenzian Warm Period: a comparison of PRISM3 and HadCM3. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 309:83–91
- Duesing W, Kaboth-Bahr S, Asrat A, Cohen AS, Foerster V, et al. 2021. Changes in the cyclicity and variability of the eastern African paleoclimate over the last 620 kyrs. *Quat. Sci. Rev.* 273:107219
- Faith JT, Du A, Behrensmeyer AK, Davies B, Patterson DB, et al. 2021. Rethinking the ecological drivers of hominin evolution. Trends Ecol. Evol. 36:797–807

- Feakins SJ, Levin NE, Liddy HM, Sieracki A, Eglinton TI, Bonnefille R. 2013. Northeast African vegetation change over 12 m.y. Geology 41:295–98
- Foerster V, Deocampo DM, Asrat A, Günter C, Junginger A, et al. 2018. Towards an understanding of climate proxy formation in the Chew Bahir basin, southern Ethiopian Rift. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 501:111–23
- Foerster V, Junginger A, Langkamp O, Gebru T, Asrat A, et al. 2012. Climatic change recorded in the sediments of the Chew Bahir basin, southern Ethiopia, during the last 45,000 years. *Quat. Int.* 274:25–37
- Foley RA. 1994. Speciation, extinction and climatic change in hominid evolution. J. Hum. Evol. 26:275-89
- Foley RA. 2017. Evolutionary ecology of fossil hominids. In *Evolutionary Ecology and Human Behavior*, ed. EA Smith, B Winterhalder, pp. 131–64. Chicago: Aldine de Gruyter
- Garello D. 2019. Tephrostratigraphy of Pliocene drill cores from Kenya and Ethiopia, and Pleistocene exposures in the Ledi-Geraru Research Project area, Ethiopia: geological context for the evolution of Australopithecus and Homo. PhD Diss., Ariz. State Univ., Tempe
- Goetz C, Hillaire-Marcel C. 1992. U-series disequilibria in early diagenetic minerals from Lake Magadi sediments: dating potential. Geochim. Cosmochim. Acta 56:1331–41
- Grootes PM, Stuiver M. 1997. Oxygen 18/16 variability in Greenland snow and ice with 10⁻³- to 10⁵-year time resolution. *J. Geophys. Res.* 102(C12):26455–70
- Grove M. 2014. Evolution and dispersal under climatic instability: a simple evolutionary algorithm. Adapt. Behav. 22:235–54
- Grove M, Lamb HF, Roberts H, Davies S, Marshall M, Bates R, Huws D. 2015. Climatic variability, plasticity, and dispersal: a case study from Lake Tana, Ethiopia. 7. Hum. Evol. 87:32–47
- Haile-Selassie Y, Melillo SM, Su DF. 2016. The Pliocene hominin diversity conundrum: Do more fossils mean less clarity? PNAS 113(23):6364–71
- Harrison T. 2011. Hominins. In *Paleontology and Geology of Laetoli: Human Evolution in Context*, Vol. 2, ed. T Harrison, pp. 141–88. Dordrecht, Neth.: Springer
- Hempson GP, Archibald S, Bond WJ. 2015a. A continent-wide assessment of the form and intensity of large mammal herbivory in Africa. Science 350:1056–61
- Hempson GP, Archibald S, Bond WJ, Ellis RP, Grant CC, et al. 2015b. Ecology of grazing lawns in Africa. Biol. Rev. 90:979–94
- Hublin JJ, Abdelouahed BN, Bailey SE, Freidline SE, Neubauer S, et al. 2017. New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens. Nature 546:289–92
- Ivory SJ, Blome MW, King JW, McGlue MM, Cole JE, Cohen AS. 2016. Environmental change explains cichlid adaptive radiation at Lake Malawi over the past 1.2 million years. *PNAS* 113:11895–900
- Ivory SJ, Lézine A, Vincens A, Cohen AS. 2018. Waxing and waning of forests: late Quaternary biogeography of southeast Africa. Glob. Change Biol. 24:2939–51
- Jackson LJ, Stone JR, Cohen AS, Yost CL. 2015. High resolution paleoecological records from Lake Malawi show no significant cooling associated with the Mount Toba supereruption at ca. 75 ka. Geology 43(9):823– 26
- Kaboth-Bahr S, Gosling WD, Vogelsang R, Bahr A, Scerri EM, et al. 2021. Paleo-ENSO influence on African environments and early modern humans. PNAS 118(23):e2018277118
- Kimbel W. 2015. The species and diversity of australopiths. In *Handbook of Paleoantbropology*, Vol. 3, ed. W Henke, I Tattersall, pp. 2071–105. New York: Springer. 2nd ed.
- Kingston JD, Deino AL, Edgar RK, Hill A. 2007. Astronomically forced climate change in the Kenyan Rift Valley 2.7–2.55 Ma: implications for the evolution of early hominin ecosystems. *7. Hum. Evol.* 53:478–503
- Kutzbach JE, Guan J, He F, Cohen AS, Orland IJ, Chen G. 2020. African climate response to insolation and glaciation: 140,000-year climate simulation with implications for early human environments. PNAS 117:2255–64
- Lamb HF, Bates R, Bryant CL, Davies S, Huws DG, et al. 2018. 150,000-year palaeoclimate record from northern Ethiopia supports early, multiple dispersals of modern humans from Africa. Sci. Rep. 8:1077
- Lane CS, Chorn BT, Johnson TC. 2013. Ash from the Toba supercruption in Lake Malawi shows no volcanic winter in East Africa at 75 ka. PNAS 110:8025–29
- Lane CS, Lowe DJ, Blockley SPE, Suzuki T, Smith VC. 2017. Advancing tephrochronology as a global dating tool: applications in volcanology, archaeology, and palaeoclimatic research. *Quat. Geochron.* 40:1–7

- Lane CS, Martin-Jones C, Johnson TC. 2018. A cryptotephra record from the Lake Victoria sediment core record of Holocene palaeoenvironmental change. Holocene 28:1909–17
- Leet K, Lowenstein TK, Renaut RW, Owen RB, Cohen AS. 2021. Labyrinth patterns in Magadi cherts: evidence for early formation from siliceous gels. Geology 49:1137–42
- Lepre CJ, Roche H, Kent DV, Harmand S, Quinn RL, et al. 2011. An earlier origin for the Acheulian. *Nature* 477:82–85
- Levin N. 2015. Environment and climate of early human evolution. *Annu. Rev. Earth Planet. Sci.* 43:405–29 Lewin R. 1981. Lake bottoms linked with human origins. *Science* 211:564–66
- Liddy HM, Feakins SJ, Tierney JE. 2017. Cooling and drying in northeast Africa across the Pliocene. Earth Planet. Sci. Lett. 449:430–38
- Lisiecki LE, Raymo ME. 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic δ¹⁸O records. Paleoceanography 20:PA1003
- Lowe DJ. 2011. Tephrochronology and its application: a review. Quat. Geochron. 6:107-53
- Lupien RL, Russell JM, Feibel C, Beck C, Castañeda I, et al. 2018. A leaf wax biomarker record of early Pleistocene hydroclimate from West Turkana, Kenya. Quat. Sci. Rev. 186:225–35
- Lupien RL, Russell JM, Subramanian A, Kinyanjui R, Beverly EJ, et al. 2021. Eastern African environmental variation and its role in the evolution and cultural change in *Homo* over the last 1 million years. *J. Hum.* Evol. 157:103028
- Lupien RL, Russell JM, Yost CL, Kingston JD, Deino AL, et al. 2019. Vegetation change in the Baringo Basin, East Africa across the onset of Northern Hemisphere glaciation 3.3–2.6 Ma. Palaeogeogr. Palaeoclimatol. Palaeoecol. 570:109426
- Lyell C. 1863. The Geological Evidences of the Antiquity of Man. London: John Murray
- Lyons RP, Scholz CA, Cohen AS, King JW, Brown ET, et al. 2015. A continuous 1.3 million year record of East African hydroclimate, and implications for patterns of evolution and biodiversity. PNAS 112:15568–73
- Maxbauer DP, Feinberg JM, Fox DL, Clyde WC. 2016. Magnetic minerals as recorders of weathering, diagenesis, and paleoclimate: a core-outcrop comparison of Paleocene–Eocene paleosols in the Bighorn Basin, WY, USA. Earth Planet. Sci. Lett. 452:15–26
- McHenry LJ, Stainstreet IG, Stollhofen H, Njau JK, Toth N, Schick K. 2020. Tuff fingerprinting and correlations between OGCP cores and outcrops for Pre-Bed I and Beds I/II at Olduvai Gorge, Tanzania. Palaeogeogr: Palaeoclimatol. Palaeoecol. 548:109630
- McNamara K, Cashman KV, Rust AC, Fontijn K, Chalié F, et al. 2018. Using lake sediment cores to improve records of volcanism at Aluto volcano in the Main Ethiopian Rift. Geochem. Geophys. Geosyst. 19:3164–88
- Miebach A, Stolzenberger S, Wacker L, Hense A, Litt T. 2019. A new Dead Sea pollen record reveals the last glacial paleoenvironment of the southern Levant. *Quat. Sci. Rev.* 214:98–116
- Muiruri VM, Owen RB, de Cort G, Renaut RW, Rabideaux NM, et al. 2021a. Middle Pleistocene to recent diatoms and stratigraphy of the Magadi Basin, south Kenya Rift. 7. Paleolimnol. 65:315–33
- Muiruri VM, Owen RB, Lowenstein TK, Renaut RW, Marchant R, et al. 2021b. A million year vegetation history and palaeoenvironmental record from the Lake Magadi Basin, Kenya Rift Valley. Palaeogeogr: Palaeoclimatol. Palaeoecol. 567:110247
- Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E. 2017. Tracing the peopling of the world through genomics. Nature 541:302–10
- Njau J, Toth N, Schick K, Stainstreet IG, McHenry LJ, Stollhofen H. 2021. The Olduvai Gorge Coring Project: drilling high resolution palaeoclimatic and palaeoenvironmental archives to constrain hominin evolution. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 561:110059
- Oppenheimer C. 2002. Limited global change due to the largest known Quaternary eruption, Toba ≈74 kyr BP? *Quat. Sci. Rev.* 21:1593–609
- Otto-Bliesner BL, Russell JM, Clark PU, Liu Z, Overpeck JT, et al. 2014. Coherent changes of southeastern equatorial and northern African rainfall during the last deglaciation. *Science* 346:1223–27
- Owen RB, McNulty E, Muiruri V, Billingsley A, Luo S, et al. 2019. Quaternary history of the Lake Magadi basin, southern Kenya Rift: evidence from outcrop and cores. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 518:97–118
- Owen RB, Muiruri VM, Lowenstein TK, Renaut RW, Rabideaux N, et al. 2018. Progressive aridification in East Africa over the last half million years and implications for human evolution. *PNAS* 115:11174–79

- Partridge T, Kerr SJ, Metcalfe SE, Scott L, Talma AS, Vogel JC. 1993. The Pretorial Saltpan: a 200,000 year southern African lacustrine sequence. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 101:317–37
- Pays O, Blanchard P, Valeix M, Chamaille-Jammes S, Duncan P, Periquet S. 2012. Detecting predators and locating competitors while foraging: an experimental study of a medium-sized herbivore in an African savanna. *Oecologia* 169:419–30
- Pollisar PJ, Rose C, Uno K, Phelps SR, deMenocal P. 2019. Synchronous rise of African C₄ ecosystems 10 million years ago in the absence of aridification. *Nat. Geosci.* 12:657–60
- Potts R. 1996. Evolution and climate variability. Science 273:922-23
- Potts R. 1998. Variability selection in hominid evolution. Evol. Anthropol. 7:81-96
- Potts R, Behrensmeyer AK, Faith JT, Tryon CA, Brooks AS, et al. 2018. Environmental dynamics during the onset of the Middle Stone Age in eastern Africa. Science 360:86–90
- Potts R, Deino AL. 1995. Mid-Pleistocene change in large mammal faunas of East Africa. *Quat. Res.* 43:106–13 Potts R, Dommain R, Moerman JW, Behrensmeyer AK, Deino AL, et al. 2020. Increased ecological resource variability during a critical transition in hominin evolution. *Sci. Adv.* 6:eabc8975
- Potts R, Faith JT. 2015. Alternating high and low climate variability: the context of natural selection and speciation in Plio-Pleistocene hominin evolution. 7. Hum. Evol. 87:5–20
- Rampino MR, Ambrose SH. 2000. Volcanic winter in the Garden of Eden: the Toba supereruption and the late Pleistocene human population crash. In *Volcanic Hazards and Disasters in Human Antiquity*, ed. FW McCoy, G Heiken, pp. 71–82. Boulder, CO: Geol. Soc. Am.
- Reed KE. 2008. Paleoecological patterns at the Hadar hominin site, Afar Regional State, Ethiopia. J. Hum. Evol. 54:743–68
- Richter D, Grün R, Joannes-Boyau R, Steele TE, Amani F, et al. 2017. The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. *Nature* 546:293–96
- Roberts HM, Ramsey CB, Chapot MS, Deino AL, Lane CS, et al. 2021. Using multiple chronometers to establish a long, directly-dated lacustrine record: constraining >600,000 years of environmental change at Chew Bahir, Ethiopia. *Quat. Sci. Rev.* 266:107025
- Robinson JR, Rowan J, Campisano CJ, Wynn JG, Reed KE. 2017. Late Pliocene environmental change during the transition from *Australopithecus* to *Homo. Nat. Ecol. Evol.* 1:0159
- Robock A, Ammann CM, Oman L, Shindell D, Levis S, Stenchikov G. 2009. Did the Toba volcanic eruption of ~74 ka B.P. produce widespread glaciation? *J. Geophys. Res.* 114(D10):D10107
- Rodríguez-Cintas A, Albert RM, Bamford MK, Stanistreet IG, Stollhofen H, et al. 2020. Palaeovegetation changes recorded in Palaeolake Olduvai OGCP Core 2A (2.09–2.12 Ma) Naibor Soit Formation Olduvai Gorge, Tanzania. Palaeogeogr. Palaeoclimatol. Palaeoecol. 557:109928
- Russell JM, Barker P, Cohen AS, Ivory S, Kimirei I, et al. 2020. ICDP Workshop on the Lake Tanganyika Scientific Drilling Project: a late Miocene–present record of climate, rifting, and ecosystem evolution from the world's oldest tropical lake. *Sci. Drill.* 27:53–60
- Schaebitz F, Asrat A, Lamb HF, Cohen AS, Foerster V, et al. 2021. Hydroclimate changes in eastern Africa over the past 200,000 years may have influenced early human dispersal. Comm. Earth Environ. 2:123
- Scholz CA, Johnson TC, Cohen AS, King JW, Peck J, et al. 2007. East African megadroughts between 135–75 kyr ago and implications for early human history *PNAS* 104:16416–21
- Sier MJ, Dupont-Nivet G, Langereis C, Deino AL, Kingston JD, Cohen AS. 2021. Magnetostratigraphy of the Hominin Sites and Paleolakes Drilling Project (HSPDP) Baringo-Tugen Hills Barsemoi core (Kenya). Palaeogeogr. Palaeoclimatol. Palaeoecol. 570:110190
- Sier MJ, Langereis CG, Dupont-Nivet G, Feibel CS, Joordens JCA, et al. 2017. The top of the Olduvai Subchron in a high resolution magnetostratigraphy from the West Turkana core WTK13, Hominin Sites and Paleolakes Drilling Project (HSPDP). *Quat. Geochron.* 42:117–29
- Smith EI, Jacobs Z, Johnsen R, Ren M, Fisher E, et al. 2018. Humans thrived in South Africa through the Toba eruption about 74,000 years ago. Nature 555:511–15
- Smith VC, Pearce NJ, Matthews N, Westgate JA, Petraglia MD, et al. 2011. Geochemical fingerprinting of the widespread Toba tephra using biotite compositions. *Quat. Int.* 246:97–104
- Spoor F, Gunz P, Neubauer S, Stelzer S, Scott N, et al. 2015. Reconstructed Homo habilis type OH 7 suggests deep-rooted species diversity in early Homo. Nature 519:83–86

- Spoor F, Leakey MG, Gathogo PN, Brown FH, Antón SC, et al. 2007. Implications of new early Homo fossils from Ileret, east of Lake Turkana, Kenya. Nature 448:688–91
- Strait DS, Grine FE. 2016. Inferring hominoid and early hominid phylogeny using craniodental characters: the role of fossil taxa. In *Human Evolution Source Book*, ed. R Ciochon, J Fleagle, pp. 399–452. Abingdon, UK: Routledge. 2nd ed.
- Stringer C. 2016. The origin and evolution of Homo sapiens. Philos. Trans. R. Soc. B 371:20150237
- Svensson A, Bigler M, Blunier T, Clausen HB, Dahl-Jensen D, et al. 2013. Direct linking of Greenland and Antarctic ice cores at the Toba eruption (74 ka BP). Clim. Past 9:749–66
- Sylvestre F, Schuster M, Vogel H, Abdheramane M, Ariztegui D, et al. 2018. The Lake CHAd Deep DRILLing project (CHADRILL)—targeting ~10 million years of environmental and climate change in Africa. Sci. Drill. 24:71–78
- Thompson JC, Mackay A, Nightingale S, Wright D, Choi J-H, et al. 2018. Ecological risk, demography and technological complexity in the Late Pleistocene of northern Malawi: implications for geographical patterning in the Middle Stone Age. *J. Quat. Sci.* 33:261–84
- Thompson JC, Wright DK, Ivory SJ, Choi JH, Nightingale S, et al. 2021. Early human impacts and ecosystem reorganization in southern-central Africa. Sci. Adv. 7:eabf9776
- Tierney JE, Russell JM, Huang Y, Sinninghe Damsté JS, Hopmans EC, Cohen AS. 2008. Northern Hemisphere controls on tropical southeast African climate during the past 60,000 years. *Science* 322:252–55
- Trauth MH, Asrat A, Berner N, Bibi F, Foerster V, et al. 2021a. Northern Hemisphere Glaciation, African climate and human evolution. *Quat. Sci. Rev.* 268:107095
- Trauth MH, Asrat A, Cohen AS, Duesing W, Foerster V, et al. 2021b. Recurring types of variability and transitions in the ~620 kyr record of climate change from the Chew Bahir basin, southern Ethiopia. *Quat. Sci. Rev.* 266:106777
- Trauth MH, Maslin MA, Deino AL, Strecker MR, Bergner AG, Duhnforth M. 2007. High- and low-latitude forcing of Plio-Pleistocene East African climate and human evolution. *J. Hum. Evol.* 53:475–86
- Uno KT, Polissar PJ, Jackson KE, deMenocal PB. 2016a. Neogene biomarker record of vegetation change in eastern Africa. PNAS 113:6355–63
- Uno KT, Polissar PJ, Kahle E, Feibel C, Harmand S, et al. 2016b. A Pleistocene palaeovegetation record from plant wax biomarkers from the Nachukui Formation, West Turkana, Kenya. PNAS 371:20150235
- Van Valen L. 1973. A new evolutionary law. Evol. Theory 1:1-30
- Viehberg FA, Just J, Dean JR, Wagner B, Franz SO, et al. 2018. Environmental change during MIS4 and MIS 3 opened corridors in the Horn of Africa for *Homo sapiens* expansion. *Quat. Sci. Rev.* 202:139–53
- Villmoare B, Kimbel WH, Seyoum C, Campisano CJ, DiMaggio EN, et al. 2015. Early Homo at 2.8 Ma from Ledi-Geraru, Afar, Ethiopia. Science 347:1352–55
- Vrba ES. 1988. Late Pliocene climatic events and hominid evolution. In The Evolutionary History of the Robust Australopithecines, ed. F Grine, pp. 405–26. New York: Aldine de Gruyter
- Vrba ES. 1995. The fossil record of African antelopes (Mammalia, Bovidae) in relation to human evolution and paleoclimate. In *Paleoclimate and Evolution with Emphasis on Human Origins*, ed. ES Vrba, GH Denton, TC Partridge, LH Burckle, pp. 285–24. New Haven, CT: Yale Univ. Press
- Washburn S. 1960. Tools and human evolution. Sci. Am. 3:62–75
- Westover KS, Stone JR, Yost CL, Scott JJ, Cohen AS, et al. 2021. Diatom paleolimnology of late Pliocene Baringo Basin (Kenya) paleolakes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 570:109382
- Williams M. 2012. The ~73 ka Toba super-eruption and its impact: history of a debate. Quat. Int. 258:19–29
- WoldeGabriel G, Endale T, White TD, Thouveny N, Hart WK, et al. 2013. The role of tephra studies in African paleoanthropology as exemplified by the Sidi Hakoma Tuff. J. Afr. Earth Sci. 77:41–58
- WoldeGabriel G, Hart WK, Katoh S, Beyene Y, Suwa G. 2005. Correlation of Plio–Pleistocene Tephra in Ethiopian and Kenyan rift basins: temporal calibration of geological features and hominid fossil records. 7. Volcanol. Geotherm. Res. 147:81–108
- Wynn JG, Reed KE, Sponheimer M, Kimbel WH, Alemseged Z, et al. 2016. Dietary flexibility of Australopithecus afurensis in the face of paleoecological change during the middle Pliocene: faunal evidence from Hadar, Ethiopia. J. Hum. Evol. 99:93–106

- Yost CL, Cohen AS, Jackson L, Stone J. 2018. Subdecadal phytolith and charcoal records from Lake Malawi, East Africa imply minimal effects on human evolution from the ∼74ka Toba supereruption. *J. Hum. Evol.* 116:75–94
- Yost CL, Ivory SJ, Deino AL, Rabideaux N, Kingston JD, Cohen AS. 2021. Phytoliths, pollen, and microcharcoal from the Baringo Basin, Kenya reveal savanna dynamics during the Plio-Pleistocene transition. Palaeogeogr. Palaeoclimatol. Palaeoecol. 570:109779
- Zawacki EE, van Soest MC, Hodges KV, Scott JJ, Barboni M, et al. 2022. Sediment provenance and silicic volcano-tectonic evolution of the northern East African Rift System from U/Pb and (U-Th)/He laser ablation double dating of detrital zircons. *Earth Planet. Sci. Lett.* 580:117375
- Zielinski GA, Mayewski PA, Meeker LD, Whitlow S, Twickler MS, Taylor K. 1996. Potential atmospheric impact of the Toba mega-eruption ~71,000 years ago. *Geophys. Res. Lett.* 23:837–40



Annual Review of Earth and Planetary Sciences

Volume 50, 2022

Contents

Civilization-Saving Science for the Twenty-First Century Marcia K. McNutt	1
Application of Light Hydrocarbons in Natural Gas Geochemistry of Gas Fields in China Shipeng Huang, Jianzhong Li, Tongshan Wang, Qingchun Jiang, Hua Jiang, Xiaowan Tao, Bin Bai, and Ziqi Feng	13
Where Has All the Carbon Gone? A. Scott Denning	55
Volcanic Outgassing of Volatile Trace Metals Marie Edmonds, Emily Mason, and Olivia Hogg	79
Dynamos in the Inner Solar System Sonia M. Tikoo and Alexander J. Evans	99
Deciphering Temperature Seasonality in Earth's Ancient Oceans Linda C. Ivany and Emily J. Judd	123
Shear Properties of Earth's Inner Core Hrvoje Tkalčić, Sheng Wang, and Thanh-Son Phạm	153
Seismic Advances in Process Geomorphology Kristen L. Cook and Michael Dietze	183
Molar-Tooth Structure as a Window into the Deposition and Diagenesis of Precambrian Carbonate Agustin Kriscautzky, Linda C. Kah, and Julie K. Bartley	205
Determining the State of Activity of Transcrustal Magmatic Systems and Their Volcanoes G. Giordano and L. Caricchi	231
Carbonatites: Classification, Sources, Evolution, and Emplacement Gregory M. Yaxley, Michael Anenburg, Sebastian Tappe, Sophie Decree, and Tibor Guzmics	261

Tectonics of the Colorado Plateau and Its Margins Karl E. Karlstrom, Justin Wilgus, Jacob O. Thacker, Brandon Schmandt, David Coblentz, and Micael Albonico
Fracture, Friction, and Permeability of Ice Erland M. Schulson and Carl E. Renshaw
Geodetic and Geological Deformation of the Island Arc in Northeast Japan Revealed by the 2011 Tohoku Earthquake Takeshi Sagiya and Angela Meneses-Gutierrez
Biomarker Approaches for Reconstructing Terrestrial Environmental Change Gordon N. Inglis, Tripti Bhattacharya, Jordon D. Hemingway, Emily H. Hollingsworth, Sarah J. Feakins, and Jessica E. Tierney
The Isotopic Ecology of the Mammoth Steppe Dorothée G. Drucker
Macrostratigraphy: Insights into Cyclic and Secular Evolution of the Earth-Life System Shanan E. Peters, Daven P. Quinn, Jon M. Husson, and Robert R. Gaines
Reconstructing the Environmental Context of Human Origins in Eastern Africa Through Scientific Drilling Andrew S. Cohen, Christopher J. Campisano, J. Ramón Arrowsmith, Asfawossen Asrat, Catherine C. Beck, Anna K. Behrensmeyer, Alan L. Deino, Craig S. Feibel, Verena Foerster, John D. Kingston, Henry F. Lamb, Tim K. Lowenstein, Rachel L. Lupien, Veronica Muiruri, Daniel O. Olago, R. Bernhart Owen, Richard Potts, James M. Russell, Frank Schaebitz, Jeffery R. Stone, Martin H. Trauth, and Chad L. Yost
Toward Understanding Deccan Volcanism Stephen Self, Tushar Mittal, Gauri Dole, and Loÿc Vanderkluysen
Physics of Melt Extraction from the Mantle: Speed and Style Richard F. Katz, David W. Rees Jones, John F. Rudge, and Tobias Keller
Pleistocene Periglacial Processes and Landforms, Mid-Atlantic Region, Eastern United States Dorothy J. Merritts and Michael A. Rahnis
Carbon Fluxes in the Coastal Ocean: Synthesis, Boundary Processes, and Future Trends Minhan Dai, Jianzhong Su, Yangyang Zhao, Eileen E. Hofmann, Zhimian Cao, Wei-Jun Cai, Jianping Gan, Fabrice Lacroix, Goulven G. Laruelle, Feifei Meng, Jens Daniel Müller, Pierre A.G. Regnier, Guizhi Wang, and Zhixuan Wang

Reckoning with the Rocky Relationship Between Eruption Size and	
Climate Response: Toward a Volcano-Climate Index	
Anja Schmidt and Benjamin A. Black	627

Errata

An online log of corrections to *Annual Review of Earth and Planetary Sciences* articles may be found at http://www.annualreviews.org/errata/earth

Related Articles

From the Annual Review of Astronomy and Astrophysics, Volume 59 (2021)

Transneptunian Space

Brett Gladman and Kathryn Volk

Exoplanet Statistics and Theoretical Implications

Wei Zhu and Subo Dong

Carrington Events

Hugh S. Hudson

From the Annual Review of Chemical and Biomolecular Engineering, Volume 12 (2021)

Storage of Carbon Dioxide in Saline Aquifers: Physicochemical Processes, Key

Constraints, and Scale-Up Potential

Philip S. Ringrose, Anne-Kari Furre, Stuart M.V. Gilfillan, Samuel Krevor, Martin Landrø, Rory Leslie, Tip Meckel, Bamshad Nazarian, and Adeel Zahid

Reactive Flows in Porous Media: Challenges in Theoretical and Numerical Methods

Anthony J.C. Ladd and Piotr Szymczak

From the Annual Review of Ecology, Evolution, and Systematics, Volume 52 (2021)

Ancient Invaders: How Palaeogenetic Tools Help to Identify and Understand Biological Invasions of the Past

Michael Knapp, Catherine 7. Collins, and Elizabeth Matisoo-Smith

A Genetic Perspective on Cetacean Evolution

Andrea A. Cabrera, Martine Bérubé, Xênia M. Lopes, Marie Louis, Tom Oosting, Alba Rey-Iglesia, Vania E. Rivera-León, Dóra Székely, Eline D. Lorenzen, and Per 7. Palsbøll

Biotic and Abiotic Controls on the Phanerozoic History of Marine Animal Biodiversity

Andrew M. Bush and Jonathan L. Payne

The Emerging Phylogenetic Perspective on the Evolution of Actinopterygian Fishes

Alex Dornburg and Thomas J. Near

From the Annual Review of Environment and Resources, Volume 46 (2021)

Land Use and Ecological Change: A 12,000-Year History Erle C. Ellis

Insights from Time Series of Atmospheric Carbon Dioxide and Related Tracers Ralph F. Keeling and Heather D. Graven

The Cold Region Critical Zone in Transition: Responses to Climate Warming and Land Use Change

Kunfu Pi, Magdalena Bieroza, Anatoli Brouchkov, Weitao Chen, Louis J.P. Dufour, Konstantin B. Gongalsky, Anke M. Herrmann, Eveline J. Krab, Catherine Landesman, Anniet M. Laverman, Natalia Mazei, Yuri Mazei, Mats G. Öquist, Matthias Peichl, Sergey Pozdniakov, Fereidoun Rezanezhad,

Céline Roose-Amsaleg, Anastasia Shatilovich, Andong Shi, Christina M. Smeaton, Lei Tong, Andrey N. Tsyganov, and Philippe Van Cappellen

Freshwater Scarcity

Peter H. Gleick and Heather Cooley

From the Annual Review of Fluid Mechanics, Volume 54 (2022)

Drop Impact Dynamics: Impact Force and Stress Distributions

Xiang Cheng, Ting-Pi Sun, and Leonardo Gordillo

Rotating Horizontal Convection

Bishakdatta Gayen and Ross W. Griffiths

Spontaneous Aggregation of Convective Storms

Caroline Muller, Da Yang, George Craig, Timothy Cronin, Benjamin Fildier, Jan O. Haerter, Cathy Hohenegger, Brian Mapes, David Randall, Sara Shamekh, and Steven C. Sherwood

Mass Transfer at the Ocean–Atmosphere Interface: The Role of Wave Breaking, Droplets, and Bubbles

Luc Deike

Dynamic Mode Decomposition and Its Variants

Peter J. Schmid

Flood Inundation Prediction

Paul D. Bates

From the Annual Review of Marine Science, Volume 14 (2022)

The Balance of Nature: A Global Marine Perspective Constantin W. Arnscheidt and Daniel H. Rothman Earth, Wind, Fire, and Pollution: Aerosol Nutrient Sources and Impacts on Ocean Biogeochemistry

Douglas S. Hamilton, Morgane M.G. Perron, Tami C. Bond, Andrew R. Bowie, Rebecca R. Buchholz, Cecile Guieu, Akinori Ito, Willy Maenhaut, Stelios Myriokefalitakis, Nazlı Olgun, Sagar D. Rathod, Kerstin Schepanski, Alessandro Tagliabue, Robert Wagner, and Natalie M. Mahowald

The History of Ocean Oxygenation

Christopher T. Reinhard and Noah 7. Planavsky

Argo—Two Decades: Global Oceanography, Revolutionized

Gregory C. Johnson, Shigeki Hosoda, Steven R. Jayne, Peter R. Oke, Stephen C. Riser, Dean Roemmich, Tohsio Suga, Virginie Thierry, Susan E. Wijffels, and Jianping Xu

Modeling the Morphodynamics of Coastal Responses to Extreme Events: What Shape Are We In?

Christopher R. Sherwood, Ap van Dongeren, James Doyle, Christie A. Hegermiller, Tian-Jian Hsu, Tarandeep S. Kalra, Maitane Olabarrieta, Allison M. Penko, Yashar Rafati, Dano Roelvink, Marlies van der Lugt, Jay Veeramony, and John C. Warner

From the *Annual Review of Materials Research*, Volume 51 (2021)

Chemistry Under Shock Conditions
Brenden W. Hamilton, Michael N. Sakano, Chunyu Li, and Alejandro Strachan

From the Annual Review of Statistics and Its Application, Volume 9 (2022)

Basis-Function Models in Spatial Statistics
Noel Cressie, Matthew Sainsbury-Dale, and Andrew Zammit-Mangion