

Azolation of Benzylic C–H Bonds via Photoredox-Catalyzed Carbocation Generation

Mrinmoy Das, Leila Zamani, Christopher Bratcher, and Patricia Z. Musacchio*

Cite This: *J. Am. Chem. Soc.* 2023, 145, 3861–3868

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: A visible-light photoredox-catalyzed method is reported that enables the coupling between benzylic C–H substrates and N–H azoles. Classically, medicinally relevant *N*-benzyl azoles are produced via harsh substitution conditions between the azole and a benzyl electrophile in the presence of strong bases at high temperatures. Use of C–H bonds as the alkylating partner streamlines the preparation of these important motifs. In this work, we report the use of *N*-alkoxyppyridinium salts as a critically enabling reagent for the development of a general C(sp³)–H azolation. The platform enables the alkylation of electron-deficient, -neutral, and -rich azoles with a range of C–H bonds, most notably secondary and tertiary partners. Moreover, the protocol is mild enough to tolerate benzyl electrophiles, thus offering an orthogonal approach to existing S_N2 and cross-coupling methods.

N-Benzyl azoles are an abundant motif in drug discovery,¹ with key examples including letrozole,² bifenazole,³ and carboetomidate⁴ (Figure 1). Such motifs exhibit widespread utility as active compounds across a number of disease areas and medical uses. Generally, azoles, such as pyrazoles, possess tempered nucleophilicity relative to halides or carboxylates.⁵ Thus, their alkylation via substitution reactions typically requires harsh conditions. Indeed, the preparation of the C–N bond of letrozole relies on an S_N2-type protocol with triazole and a benzyl electrophile (halide/tosylate) at 100 °C with strong base.⁶ Additionally, benzyl electrophiles are prone to hydrolysis and often require extra preparatory steps. Thus, a need exists for new mild and streamlined protocols to be developed. In addition, a strategy that differs from classical transition metal cross-coupling conditions would also allow for an expanded substrate scope to include electron-rich azoles, organoboranes, and alkyl, aryl, and benzyl/allyl halides to maximize synthetic route opportunities.⁷


Recently, alternative alkylating partners have been investigated, with carboxylic acids being a particularly attractive option.⁸ Although they are abundant and provide a handle for accessing radicals and carbocations, their activation requires acyl group manipulations or strong oxidants to facilitate the decarboxylation. By contrast, C–H bonds represent the most prevalent functionality in organic compounds; accordingly, rendering them reactive for C–X bond formation would be greatly advantageous toward the goal of a streamlined reaction platform.⁹ The primary advantage of C–H functionalization methods is a decrease in preparatory steps of reagents and eventual use for the rapid diversification of late-stage targets.

A handful of methods have demonstrated that single-electron oxidation of arenes can activate the benzylic C–H positions toward subsequent functionalization, albeit requiring high oxidation potentials.¹⁰ In the past few years, a mechanistic strategy for expanding the C–H scope employs a hydrogen-atom transfer (HAT) event prior to a (radical) oxidation (i.e., radical–polar crossover, RPC).¹¹ Notably, the combination of


HAT and RPC has been engineered into one catalytic cycle utilizing different HAT species: Lei and co-workers used the phenoxy radical of 2,3-dichloro-5,6-dicyano-*p*-benzoquinone (DDQ);¹² the Noël group employed a UV-activated decatungstate catalyst;¹³ and the Stahl group leveraged a sulfonamide radical derived from *N*-fluorobenzenesulfonamide (NFSI) in conjugation with copper catalysis.¹⁴ While all three methods proceed via a carbocation intermediate, noticeably lacking in the scope of each protocol is the formation of fully substituted centers arising from *in situ*-generated tertiary carbocations. Recently, our group,¹⁵ concurrently with the Doyle group,¹⁶ published a visible-light photoredox-catalyzed [HAT+RPC] mechanism that can engage classically weak nucleophiles, including fluoride, and readily forge fully substituted centers from tertiary C–H precursors. We hypothesized that the established platform could be extended to the formation of an array of *N*-benzyl azoles, including those bearing fully substituted centers. In addition, these works demonstrated the power of photoredox catalysis to mediate a formal hydride abstraction with two different types of HAT reagents: *tert*-butyl peroxybenzoate (TBPB) and *N*-acyloxypthalimide. Thus far, only photoredox platforms have exhibited such modularity in the examination of stereoelectronically diverse HAT reagents for the [HAT+RPC] process. Herein we report the implementation of *N*-alkoxyppyridinium salts in this process. Simple *N*-alkoxyppyridinium salts can be readily prepared in one step and offer an electronically tunable HAT scaffold.^{17,18} Finally, mechanistic evidence suggests that an electron-donor–acceptor (EDA) complex may be operable for

Received: December 2, 2022


Published: February 9, 2023



A. Importance of N-benzylic azole motif in drug discovery



B. Mechanistic strategy for C–H azolation



C. Current C–H azolation methods via [HAT+RPC] design






Figure 1. Background on C–H azolation methods. (A) *N*-benzylic azoles are attractive pharmaceutical motifs. (B) Recent mechanism for C–H functionalization. (C) Current C–H azolation methods utilizing [HAT+RPC].

activating the pyridinium reagents for certain electron-rich benzylic partners.¹⁹

Our efforts toward the development of a C–H azolation protocol initiated with the use of TBPB to facilitate the HAT event. Although successful in our prior work for C–H fluorination with nucleophilic fluoride ($N = 10.8$ – 13.2),²⁰ switching to less nucleophilic coupling partners such as pyrazoles ($N = 8.9$ – 9.6) resulted in competitive trapping of the carbocation by both benzoate ($N = 16.8$) and *tert*-butanol ($N = 5.4$)⁵ byproducts (Figure 2A). Work by Hong,²¹ Li,²² and Lakhdar²³ has demonstrated that *N*-alkoxy pyridinium reagents can facilitate intermolecular HAT processes at P–H, C(sp²)–H, Si–H, and α -oxy C(sp³)–H bonds.²⁴ We hypothesized that this reagent could also be used for the intermolecular abstraction of H• at benzylic C(sp³)–H bonds in the desired transformation (Figure 2B). While *N*-

A. Inherent issue with prior HAT reagents

Prior HAT reagent:



B. Proposed mechanism and Stern–Volmer studies



C. Survey of *N*-alkoxy pyridinium salts as HAT reagents

Figure 2. Optimization with *N*-alkoxy pyridinium reagents. (A) Use of TBPB can lead to competitive nucleophiles. (B) Proposed mechanism and Stern–Volmer experiment. (C) Optimization of the *N*-alkoxy pyridinium scaffold.

alkoxy pyridiniums bearing long-chain alkyl groups have been used for intramolecular 1,5-HAT processes,²⁵ they have not been used widely for intermolecular efforts at C(sp³)–H bonds.²² The pyridinium reagents would be particularly attractive in the desired transformation, as reductive fragmentation would reduce the generation of competitive nucleophilic byproducts (Figure 2B). Accordingly, we rapidly synthesized a suite of *N*-methoxy- and *N*-ethoxy pyridinium reagents.²⁶ With indane and 4-bromopyrazole as our model substrates, we delightfully observed the desired C–H azolation product (Figure 2C).

Electron-withdrawing *p*-cyanopyridinium **pyr-1** was discovered to result in higher product formation, possibly due to a lower reduction potential (see the Supporting Information). Stern–Volmer experiments corroborated an interaction between the excited state of the photocatalyst and **pyr-1**. Next, we explored the generality of the substrate scope.

Starting with an exploration of azoles (Figure 3), an array of electron-withdrawing groups at the 4-position of pyrazole were well-tolerated, including other halides (2–4), esters (5), and trifluoromethyl (6), cyano (7), and nitro groups (8). Substitution at the 3-position of pyrazole also resulted in




Figure 3. Azole scope for secondary and tertiary benzylic C–H substrates. Reactions were run on a 0.5 mmol scale. DCE = 1,2-dichloroethane. HFIP = hexafluoroisopropanol. ^a48 h. ^b72 h. ^c60 °C. ^d96 h.

good to excellent yields (9–13). Difunctionalized pyrazoles afforded high yields of the products (14–17), and an extended heterocycle was also successful in the protocol (18). Excitingly, more electron-rich pyrazoles were successfully alkylated with our system, representing a class of substrates that are not compatible with base-metal-catalyzed strategies due to potential catalyst poisoning (19–22). Furthermore, 1,2,3- and 1,2,4-triazoles ($N \approx 7.7$) were also viable nucleophiles, as was benzotriazole (23–25).²⁶ Substituted tetrazoles, imidazoles, and benzimidazoles were also viable substrates (26–33).

Given the minimal effect of steric hindrance observed, we questioned whether the photoredox-catalyzed [HAT+RPC] platform could enable the functionalization of tertiary benzylic C–H bonds. To the best of our knowledge, the synthesis of fully substituted carbon centers has not been reported in prior C–H azolation methods with the [HAT+RPC] formula, despite the enhanced carbocation stability. Classically, tertiary benzyl halides/tosylates are unstable and/or prepared with harsh reagents (strong acids) and expensive oxidants.²⁷ Moreover, the carboxylic acid equivalent of 2-isopropyl-naphthalene is not widely available from commercial vendors. The use of C–H alternatives thus represents an advantage in terms

of synthetic ease and available resources. Gratifyingly, 2-isopropyl-naphthalene could be readily functionalized with a wide array of azoles using TBPB as the HAT reagent. Currently, we postulate that a methyl radical could be the most effective H-atom abstractor for tertiary benzylic C–H sites, which can be derived from $\cdot\text{OtBu}$ via a facile β -scission.^{16,28} Various 4- and 3-substituted pyrazoles were successfully alkylated (34–41), with the latter being functionalized at the less sterically hindered nitrogen. Electron-rich pyrazoles also afforded the products in modest yields (42–44). Notably, 44 contains a nucleophilic arylboron functionality that would not be tolerated by transition metal approaches. Moreover, the strength of the carbocation strategy was highlighted with the alkylation of 3,5-dimethylpyrazole to give sterically congested adduct 45, albeit in low yield. Finally, the protocol was also successful at producing fully substituted tetrazole, imidazole, and (benzo)triazole C–N adducts (46–49).²⁹

Next, we examined the generality of the benzylic C–H scope. Methylene sites on both cyclic and acyclic precursors afforded appreciable yields of C–N products (50–57). Diphenylmethane, a common motif in drug targets (Figure 1a), worked in good yield (51). Electron-deficient functional




Figure 4. Scope of the C–H reaction partner. ^a60 °C. ^b48 h. ^cTetrahydrofuran (5 equiv). ^dConditions: hydrocarbon (3 equiv), azole (1 equiv), and TBPB (3 equiv) as indicated in Figure 3. ^e1 equiv of C–H precursor, azole (3 equiv), and pyr-1 (1.5 equiv). ^fEosin Y (5 mol %).

groups at the para position did not significantly hinder the reaction efficiency (52–56); nevertheless, higher yields were observed with electron-donating groups (57). Notably, our carbocation-generating protocol is permissible of aryl bromide and chloride motifs, allowing for the retention of functional handles for further derivatization via classical cross-coupling catalysis. Meta substitution was also well-tolerated (58). The primary benzylic substrate leading to 59 represents another class of substrates not demonstrated in other [HAT+RPC] strategies for azolation. C–H functionalization of allylic positions was achieved, giving 60 and 61 in 18% and 49% yield, respectively. Lastly, α -oxy C–H sites were also viable substrates for the visible-light-mediated azolation (62).

Next, we sought to apply the method to late-stage functionalization of pharmaceutical scaffolds. The seven-

membered cyclic core of ivabradine, used in the treatment of heart failure, was successfully elaborated at the α -benzylic site in 48% yield (63). Celestolide (64, 70% yield) and the core of donepezil (65, 45% yield) also underwent C–H azolation with the [HAT+RPC] protocol in appreciable yields. Lastly, we demonstrated that azole derivatives of the antifungal agent bifonazole (66) can be readily prepared in one step from commercially available 4-benzylbiphenyl. These examples demonstrate the utility of direct functionalization of benzylic C–H bonds as opposed to the established multistep processes involving conversion of a benzyl alcohol to a benzyl chloride followed by harsh conditions.³⁰

Next, an array of other tertiary benzylic C–H partners were translated to fully substituted products (67–71). γ -Phenyl-lactone gave sterically congested 68 in good yield, as did

phenylcyclohexane (69). Monofunctionalization of compounds containing two tertiary benzylic sites was successful (70), and last, installation of a congested C–N center on 9-methylfluorene was realized (71). Notably, for substrates 68–71, the corresponding benzyl chlorides or carboxylic acids are either commercially unavailable or prohibitively expensive. Excitingly, compound 72 was successfully prepared, demonstrating potential for the azolation of heteroarene C–H substrates.


To probe regioselectivity, a competition experiment was conducted. Using **pyr-1**, a preference for azolation at indane over 2-isopropynaphthalene was observed. High selectivity for the secondary position of 73 (16.3:1) was congruent with this finding. The protocol appears to be selective for secondary benzylic C–H sites over primary benzylic (74) and aliphatic tertiary (75) sites and is completely selective for α -oxy benzylic positions over secondary benzylic ones (76; see the [Supporting Information](#)).

Finally, we sought to evaluate the specificity and orthogonality of our photocatalytic C–H azolation. Classically, letrazole and bifonazole are prepared via S_N2 -type reactions on benzyl chlorides with heat and strong bases.^{6,30} As depicted in [Figure 4](#), this platform is sufficiently mild to tolerate the preparation of 77 in 40% yield with no detection of the S_N2 product (>20:1 regioselectivity). Subsequently, high yields were achieved for S_N2 azidation (78), thiolation (79), and esterification (80) substitution reactions. Furthermore, the benzyl chloride functionality could also serve as an electrophile in a Pd-catalyzed cross-coupling (81).³¹

Lastly, pyridinium salts have been reported to participate in EDA complexes.^{18,23b,c,32} Control experiments suggested that when HFIP is added as a cosolvent, an EDA complex could be operable, on the basis of the observation of product without photocatalyst and an observed bathochromic shift in the UV–vis spectra ([Figure 5b](#)). Presumably, the EDA complex facilitates oxidation of the arene, which triggers fragmentation of **pyr-1** to release a methoxy radical. Subsequent HAT at the benzylic position of the resultant arene radical cation can generate the benzylic carbocation.³³ Further UV–vis studies indicated that only electron-rich substrates form an EDA complex (see the [Supporting Information](#)), suggesting that other substrates may follow a photocatalyst-mediated mechanism.


In conclusion, we have successfully developed a benzylic C–H azolation reaction via a photoredox-catalyzed formal hydride abstraction mechanism. An *N*-methoxy-pyridinium salt was used as an effective intermolecular HAT reagent at benzylic C(sp³)–H bonds. We have demonstrated the generality of the method, as it includes the alkylation of a plethora of azoles. Additionally, a broad C–H partner scope was established, including the alkylation of secondary and tertiary benzylic C–H sites, the latter of which afford a direct and simplified route to *N*-tert-alkyl azole motifs. Importantly, we showcase the complementary nature of our reaction conditions to classical S_N2 and transition metal cross-coupling conditions, which are two current state-of-the-art technologies for forging alkylated azole products. Lastly, mechanistic studies suggest that a plausible EDA mechanism is operable for certain C–H substrates.

Mechanistic insights into the *N*-methoxy-pyridinium Reagent



a. Control Experiments

Rxn	Change to Conditions	Yield
1	No change	80%
2	no light	0%
3	no photocatalyst	40%
4	no pyr-1 reagent	0%
5	1 equiv. TEMPO	6%
6	DCE	45%
7	DCE, no photocatalyst	0%



b. UV-vis experiments and evidence of EDA complex




Figure 5. Mechanistic studies. (a) Control experiments for optimized conditions. (b) UV–vis experiments to probe for EDA complex formation and yields of substrates without photocatalyst (PC). ^a48 h.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c12850>.

Additional experimental details, characterization data, optimization, and relevant spectra ([PDF](#))

■ AUTHOR INFORMATION

Corresponding Author

Patricia Z. Musacchio – Department of Chemistry, Worcester Polytechnic Institute, Worcester, Massachusetts 01609, USA; orcid.org/0000-0002-0211-1893; Email: pzmusacchio@wpi.edu

Authors

Mrimoy Das – Department of Chemistry, Worcester Polytechnic Institute, Worcester, Massachusetts 01609, USA

Leila Zamani – Department of Chemistry, Worcester Polytechnic Institute, Worcester, Massachusetts 01609, USA
Christopher Bratcher – Department of Chemistry, Worcester Polytechnic Institute, Worcester, Massachusetts 01609, USA

Complete contact information is available at:

<https://pubs.acs.org/10.1021/jacs.2c12850>

Funding

Research reported in this publication was supported by a startup grant from Worcester Polytechnic Institute and the National Institute of Health (R35GM147021). C.B. was supported by the National Science Foundation for summer research through the REU Program (Grant CHE-1950512).

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

C.B. thanks the NSF REU Program for summer support. We thank C. C. Le, J. M. Lipshultz, and D. A. Nagib for helpful conversations about the manuscript, Y. Zhang (WPI) and A. Ali (UMass Medical) for NMR assistance, and M. Xatse and C. Olsen (WPI) for HRMS assistance.

REFERENCES

- (1) (a) Bennani, F. E.; Doudach, L.; Cherrah, Y.; Ramli, Y.; Karrouchi, K.; Ansar, M.; Faouzi, M. E. A. Overview of Recent Developments of Pyrazole Derivatives as an Anticancer Agent in Different Cell Line. *Bioorg. Chem.* **2020**, *97*, 103470. (b) Baumann, M.; Baxendale, I. R.; Ley, S. V.; Nikbin, N. An Overview of the Key Routes to the Best Selling 5-Membered Ring Heterocyclic Pharmaceuticals. *Beilstein J. Org. Chem.* **2011**, *7*, 442–495. (c) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among U.S. FDA approved pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274. (d) Küçükgüzel, Ş. G.; Şenkardeş, S. Recent Advances in Bioactive Pyrazoles. *Eur. J. Med. Chem.* **2015**, *97*, 786–815. (e) Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. Organic Synthesis Provides Opportunities to Transform Drug Discovery. *Nat. Chem.* **2018**, *10*, 383–394. (2) Bhatnagar, A. S. The discovery and mechanism of action of letrozole. *Breast Cancer Res. Treat.* **2007**, *105*, 7–17. (3) El Hage, S.; Lajoie, B.; Feuillolay, C.; Roques, C.; Baziard, G. Synthesis, Antibacterial and Antifungal Activities of Bifonazole Derivatives. *Arch. Pharm. Pharm. Med. Chem.* **2011**, *344*, 402–410. (4) Ode, K. Intravenous Anaesthetic Agents. *Anaesth. Intensive Care Med.* **2019**, *20*, 118–125. (5) Mayr, H. Mayr's Database of Reactivity Parameters. <https://www.cup.lmu.de/oc/mayr/reaktionsdatenbank/fe/> (accessed 2022-12-01). (6) (a) Palle, V. R. A.; Kalaria, A. J.; Shelke, S. A. Process for Preparing Letrozole. US 2007/0100149 A1, May 3, 2007. (b) Friedman, O.; Freger, B.; Etlin, O.; Ditzkovitch, J.; Danon, E.; Seryi, Y.; Davidi, G.; Arad, O.; Kaspi, J. Letrozole production process. US 7,538,230 B2, May 26, 2009. (c) Doiron, J.; Soultan, A. H.; Richard, R.; Touré, M. M.; Picot, N.; Richard, R.; Čuperlović-Culfi, M.; Robichaud, G. A.; Touaibia, M. Synthesis and structure–activity relationship of 1- and 2-substituted-1,2,3-triazole letrozole-based analogues as aromatase inhibitors. *Eur. J. Med. Chem.* **2011**, *46*, 4010–4024. (d) Long, Y.; Zheng, Y.; Xia, Y.; Qu, L.; Yang, Y.; Xiang, H.; Zhou, X. Nickel-Catalyzed Synthesis of an Aryl Nitrile via Aryl Exchange between an Aromatic Amide and a Simple Nitrile. *ACS Catal.* **2022**, *12*, 4688–4695. (7) For transition-metal-catalyzed approaches toward *N*-benzyl/allyl azole synthesis, see: (a) Kong, D.; Moon, P. J.; Bsharat, O.; Lundgren, R. J. Direct catalytic decarboxylative amination of aryl acetic acids. *Angew. Chem., Int. Ed.* **2020**, *59*, 1313–1319. (b) Jiu, A. Y.; Slocumb, H. S.; Yeung, C. S.; Yang, X.-H.; Dong, V. M. Enantioselective Addition of Pyrazoles to Dienes. *Angew. Chem., Int. Ed.* **2021**, *60*, 19660–19664. (c) Ye, Y.; Kim, S.-T.; Jeong, J.; Baik, M.-H.; Buchwald, S. L. CuH-Catalyzed Enantioselective Alkylation of Indole Derivatives with Ligand-Controlled Regiodivergence. *J. Am. Chem. Soc.* **2019**, *141*, 3901–3909. (d) Chen, H.; Yang, Y.; Li, N.; Xue, X.; He, Z.; Zeng, Q. Palladium Catalyzed C–N Cross-Coupling of NH-Heteroarenes and Quaternary Ammonium Salts via C–N Bond Cleavage. *Org. Process Res. Dev.* **2019**, *23*, 1679–1685. (e) Lacker, C. R.; Delano, T. J.; Chen, E. P.; Kong, J.; Belyk, K. M.; Piou, T.; Reisman, S. E. Enantioselective Synthesis of *N*-Benzylic Heterocycles by Ni/Photoredox Dual Catalysis. *J. Am. Chem. Soc.* **2022**, *144*, 20190–20195. (8) For decarboxylation strategies for azole alkylation, see: (a) Sheng, T.; Zhang, H.-J.; Shang, M.; He, C.; Vantourout, J. C.; Baran, P. S. Electrochemical Decarboxylative *N*-Alkylation of Heterocycles. *Org. Lett.* **2020**, *22*, 7594–7598. (b) Liang, Y.; Zhang, X.; MacMillan, D. W. C. Decarboxylative sp^3 C–N coupling via dual copper and photoredox catalysis. *Nature* **2018**, *559*, 83–88. (c) Kobayashi, R.; Shibutani, S.; Nagao, K.; Ikeda, Z.; Wang, J.; Ibáñez, I.; Reynolds, M.; Sasaki, Y.; Ohmiya, H. Decarboxylative *N*-alkylation of azoles through visible-light-mediated organophotoredox catalysis. *Org. Lett.* **2021**, *23*, 5415–5419. (d) Li, P.; Zbieg, J. R.; Terrett, J. A. The Direct Decarboxylative *N*-Alkylation of Azoles, Sulfonamides, Ureas, and Carbamates with Carboxylic Acids via Photoredox Catalysis. *Org. Lett.* **2021**, *23*, 9563–9568. (e) Zhao, W.; Wurz, R. P.; Peters, J. C.; Fu, G. C. Photoinduced, Copper-Catalyzed Decarboxylative C–N Coupling to Generate Protected Amines: An Alternative to the Curtius Rearrangement. *J. Am. Chem. Soc.* **2017**, *139*, 12153–12156. (f) Shao, X.; Zheng, Y.; Tian, L.; Martín-Torres, I.; Echavarren, A. M.; Wang, Y. Decarboxylative C_{sp^3} –N Bond Formation by Electrochemical Oxidation of Amino Acids. *Org. Lett.* **2019**, *21*, 9262–9267. (g) Tang, Z.-L.; Ouyang, X.-H.; Song, R.-J.; Li, J.-H. Decarboxylative $C(sp^3)$ –N Cross-Coupling of Diacyl Peroxides with Nitrogen Nucleophiles. *Org. Lett.* **2021**, *23*, 1000–1004. (h) Nguyen, V. T.; Nguyen, V. D.; Haug, G. C.; Vuong, N. T. H.; Dang, H. T.; Arman, H. D.; Larionov, O. V. Visible-light-enabled direct decarboxylative *N*-alkylation. *Angew. Chem., Int. Ed.* **2020**, *59*, 7921–7927. (i) Merchant, R. R.; Lang, S. B.; Yu, T.; Zhao, S.; Qi, Z.; Suzuki, T.; Bao, J. A General One-Pot Protocol for Hindered *N*-Alkyl Azaheterocycles from Tertiary Carboxylic Acids. *Org. Lett.* **2020**, *22*, 4180–4184. (9) For selected reviews on C–H functionalization, see: (a) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The Medicinal Chemist's Toolbox for Late Stage Functionalization of Drug-Like Molecules. *Chem. Soc. Rev.* **2016**, *45*, 546–576. (b) Abrams, D. J.; Provencher, P. A.; Sorensen, E. J. Recent applications of C–H functionalization in complex natural product synthesis. *Chem. Soc. Rev.* **2018**, *47*, 8925–8967. (c) Govaerts, S.; Nyuchev, A.; Noel, T. Pushing the boundaries of C–H bond functionalization chemistry using flow technology. *J. Flow Chem.* **2020**, *10*, 13–71. (d) Chu, J. C. K.; Rovis, T. Complementary Strategies for Directed $C(sp^3)$ –H Functionalization: A Comparison of Transition-Metal-Catalyzed Activation, Hydrogen Atom Transfer, and Carbene/Nitrene Transfer. *Angew. Chem., Int. Ed.* **2018**, *57*, 62–101. (10) For examples relying on arene oxidation, see: (a) Pandey, G.; Laha, R.; Singh, D. Benzylic $C(sp^3)$ –H Functionalization for C–N and C–O Bond Formation via Visible Light Photoredox Catalysis. *J. Org. Chem.* **2016**, *81*, 7161–7171. (b) Ruan, Z.; Huang, Z.; Xu, Z.; Zeng, S.; Feng, P.; Sun, P.-H. Late-Stage Azolation of Benzylic C–H Bonds Enabled by Electrooxidation. *Sci. China. Chem.* **2021**, *64*, 800–807. (c) Hou, Z.-W.; Liu, D.-J.; Xiong, P.; Lai, X.-L.; Song, J.; Xu, H.-C. Site Selective Electrochemical Benzylic C–H Amination. *Angew. Chem., Int. Ed.* **2021**, *60*, 2943–2947. (d) Shao, X.; Tian, L.; Wang, Y. C–N Coupling of Azoles or Imides with Carbocations Generated by Electrochemical Oxidation. *Eur. J. Org. Chem.* **2019**, *2019*, 4089–4094. (e) Yang, Y.-Z.; Song, R.-J.; Li, J.-H. Intermolecular Anodic Oxidative Cross-Dehydrogenative $C(sp^3)$ –N Bond-Coupling Reactions of Xanthenes with Azoles. *Org. Lett.* **2019**, *21*, 3228–3231.

- (11) For early examples of the [HAT+RPC] mechanism applied to C–H functionalization, see: (a) Dai, C.; Meschini, F.; Narayanam, J. M. R.; Stephenson, C. R. J. Friedel-Crafts Amidokylation via Thermolysis and Oxidative Photocatalysis. *J. Org. Chem.* **2012**, *77*, 4425–4431. (b) Pandey, G.; Laha, R. Visible-Light-Catalyzed Direct Benzylic C(sp³)–H Amination Reaction by Cross-Dehydrogenative Coupling. *Angew. Chem., Int. Ed.* **2015**, *54*, 14875–14879. (c) Li, G.-X.; Morales-Rivera, C. A.; Gao, F.; Wang, Y.; He, G.; Liu, P.; Chen, G. A unified photoredox-catalysis strategy for C(sp³)–H hydroxylation and amidation using hypervalent iodine. *Chem. Sci.* **2017**, *8*, 7180–7185. (d) Hu, H.; Chen, S.-J.; Mandal, M.; Pratik, S. M.; Buss, J. A.; Krska, S. W.; Cramer, C. J.; Stahl, S. S. Copper-catalysed benzylic C–H coupling with alcohols via radical relay enabled by redox buffering. *Nat. Catal.* **2020**, *3*, 358–367.
- (12) Song, C.; Dong, X.; Yi, H.; Chiang, C.-W.; Lei, A. DDQ-Catalyzed Direct C(sp³)–H Amination of Alkylheteroarenes: Synthesis of Biheteroarenes Under Aerobic and Metal-Free Conditions. *ACS Catal.* **2018**, *8*, 2195–2199. For an electrochemical version, see: Hou, Z.-W.; Li, L.; Wang, L. Organocatalytic electrochemical amination of benzylic C–H bonds. *Org. Chem. Front.* **2021**, *8*, 4700–4705.
- (13) Wan, T.; Capaldo, L.; Laudadio, G.; Nychev, A. V.; Rincón, J. A.; García-Losada, P.; Mateos, C.; Frederick, M. O.; Nuño, M.; Noël, T. Decatungstate-Mediated C(sp³)–H Heteroarylation via Radical-Polar Crossover in Batch and Flow. *Angew. Chem., Int. Ed.* **2021**, *60*, 17893–17897.
- (14) Chen, S.-J.; Golden, D. L.; Krska, S. W.; Stahl, S. S. Copper-Catalyzed Cross-Coupling of Benzylic C–H Bonds and Azoles with Controlled N-Site Selectivity. *J. Am. Chem. Soc.* **2021**, *143*, 14438–14444.
- (15) (a) Zhang, Y.; Fitzpatrick, N. A.; Das, M.; Bedre, I.; Yayla, H. G.; Lall, M. S.; Musacchio, P. Z. A photoredox-catalyzed approach for formal hydride abstraction to enable C_{sp}³–H functionalization with nucleophilic partners (F, C, O, N, and Br/Cl). *Chem. Catal.* **2022**, *2*, 292–308. (b) Fitzpatrick, N. A.; Zamani, L.; Das, M.; Yayla, H. G.; Lall, M. S.; Musacchio, P. Z. A S_N1 mechanistic approach to the Williamson ether reaction via photoredox catalysis applied to benzylic C(sp³)–H bonds. *Tetrahedron* **2022**, *125*, 132986.
- (16) Leibler, I. N.-M.; Tekle-Smith, M.; Doyle, A. G. A general strategy for C(sp³)–H functionalization with nucleophiles using methyl radical as a hydrogen atom abstractor. *Nat. Commun.* **2021**, *12*, 6950.
- (17) He, F.-S.; Ye, S.; Wu, J. Recent Advances in Pyridinium Salts as Radical Reservoirs in Organic Synthesis. *ACS Catal.* **2019**, *9*, 8943–8960.
- (18) For another example of an electronically tunable pyridinium system, see: McClain, E. J.; Wortman, A. K.; Stephenson, C. R. J. Radical generation enabled by photoinduced N–O bond fragmentation. *Chem. Sci.* **2022**, *13*, 12158.
- (19) Crisenza, G. E. M.; Mazzarella, D.; Melchiorre, P. Synthetic Methods Driven by the Photoactivity of Electron Donor–Acceptor Complexes. *J. Am. Chem. Soc.* **2020**, *142*, 5461–5476.
- (20) The nucleophilicity parameter as established by Mayr is represented as *N*.
- (21) (a) Kim, I.; Min, M.; Kang, D.; Kim, K.; Hong, S. Direct Phosphonation of Quinolinones and Coumarins Driven by the Photochemical Activity of Substrates and Products. *Org. Lett.* **2017**, *19*, 1394–1397. (b) Kim, I.; Kang, G.; Lee, K.; Park, B.; Kang, D.; Jung, H.; He, Y.-T.; Baik, M.-H.; Hong, S. Site-Selective Functionalization of Pyridinium Derivatives via Visible-Light-Driven Photocatalysis with Quinolinone. *J. Am. Chem. Soc.* **2019**, *141*, 9239–9248.
- (22) Zheng, M.; Hou, J.; Zhan, L.-W.; Huang, Y.; Chen, L.; Hua, L.-L.; Li, Y.; Tang, W.-Y.; Li, B.-D. Visible-Light-Driven, Metal-Free Divergent Difunctionalization of Alkenes Using Alkyl Formates. *ACS Catal.* **2021**, *11*, 542–553.
- (23) (a) Rammal, F.; Gao, D.; Boujnah, S.; Hussein, A. A.; Lalevée, J.; Gaumont, A.-C.; Morlet-Savary, F.; Lakhdar, S. Photochemical C–H Silylation and Hydroxymethylation of Pyridines and Related Structures: Synthetic Scope and Mechanisms. *ACS Catal.* **2020**, *10*, 13710–13717. (b) Quint, V.; Morlet-Savary, F.; Lohier, J.-F.; Lalevée, J.; Gaumont, A.-C.; Lakhdar, S. Metal-Free, Visible Light-Photocatalyzed Synthesis of Benzo[b]phosphole Oxides: Synthetic and Mechanistic Investigations. *J. Am. Chem. Soc.* **2016**, *138*, 7436–7441. (c) Quint, V.; Chouchene, N.; Askri, M.; Lalevée, J.; Gaumont, A.-C.; Lakhdar, S. Visible-light-mediated α -phosphorylation of *N*-aryl tertiary amines through the formation of electron-donor–acceptor complexes: synthetic and mechanistic studies. *Org. Chem. Front.* **2019**, *6*, 41–44.
- (24) Chang, L.; An, Q.; Duan, L.; Feng, K.; Zuo, Z. Alkoxy Radicals See the Light: New Paradigms of Photochemical Synthesis. *Chem. Rev.* **2022**, *122*, 2429–2486.
- (25) (a) Bao, X.; Wang, Q.; Zhu, J. Remote C(sp³)–H Arylation and Vinylation of *N*-Alkoxyppyridinium Salts to *d*-Aryl and *d*-Vinyl Alcohols. *Chem. - Eur. J.* **2019**, *25*, 11630–11634. (b) Kim, I.; Park, B.; Kang, G.; Kim, J.; Jung, H.; Lee, H.; Baik, M.-H.; Hong, S. Visible-Light-Induced Pyridylation of Remote C(sp³)–H Bonds by Radical Translocation of *N*-Alkoxyppyridinium Salts. *Angew. Chem., Int. Ed.* **2018**, *57*, 15517–15522. (c) Bao, X.; Wang, Q.; Zhu, J. Dual Photoredox/CopperCatalysis for the Remote C(sp³)–H Functionalization of Alcohols and Alkyl Halides by *N*-Alkoxyppyridinium Salts. *Angew. Chem., Int. Ed.* **2019**, *58*, 2139–2143.
- (26) Barthelemy, A. L.; Tuccio, B.; Magnier, E.; Dagousset, G. Alkoxy Radicals Generated under Photoredox Catalysis: A Strategy for anti-Markovnikov Alkoxylation Reactions. *Angew. Chem., Int. Ed.* **2018**, *57*, 13790–13794.
- (27) (a) Olah, G. A.; Comisarow, M. B.; Kim, C. J. Stable carbonium ions. LXXIII. 3-Aryl-2,3-dimethyl-2-butyl cations and factors controlling phenonium ion-phenethyl cation-benzyl cation equilibria. *J. Am. Chem. Soc.* **1969**, *91*, 1458–1469. (b) Canestrari, D.; Lancianesi, S.; Badiola, E.; Strinna, C.; Ibrahim, H.; Adamo, M. F. A. Desulfurative Chlorination of Alkyl Phenyl Sulfides. *Org. Lett.* **2017**, *19*, 918–921. (c) Cantillo, D.; de Frutos, O.; Rincon, J. A.; Mateos, C.; Kappe, C. O. A Scalable Procedure for Light-Induced Benzylic Brominations in Continuous Flow. *J. Org. Chem.* **2014**, *79*, 223–229.
- (28) It is plausible that a methyl radical operates as the HAT reagent. When TBPB is used with a polar solvent, β -scission of the tBuO $\cdot$ radical is likely (see ref 16).
- (29) This example is particularly remarkable given the propensity of benzotriazolyl moieties to undergo reductive cleavage. See: (a) Teders, M.; Gómez-Suárez, A.; Pitzer, L.; Hopkinson, M. N.; Glorius, F. Diverse Visible-Light-Promoted Functionalizations of Benzotriazoles Inspired by Mechanism-Based Luminescence Screening. *Angew. Chem., Int. Ed.* **2017**, *56*, 902–906. (b) Jian, Y.; Chen, M.; Huang, B.; Jia, W.; Yang, C.; Xia, W. Visible-Light-Induced C(sp²)–P Bond Formation by Denitrogenative Coupling of Benzotriazoles with Phosphites. *Org. Lett.* **2018**, *20*, 5370–5374. (c) Xie, T.; Zhou, L.; Shen, M.; Li, J.; Lv, X.; Wang, X. Diastereoselective synthesis of *cis*-1,2-disubstituted cyclopropanols and cyclopent-3-enols via SmI₂ mediated C–N(Bt) bond cleavage. *Tetrahedron Lett.* **2015**, *56*, 3982–3987.
- (30) Franckowiak, G.; Hess, C.; Weidemann, K. Preparation of bifonazole, useful as antimycotic agent, comprises reacting biphenyl-4-yl(phenyl) methanol with chlorinating agent in cyclohexane and then with imidazole. DE 10332684 B3, April 21, 2005.
- (31) Molander, G. A.; Elia, M. D. Suzuki-Miyaura Cross-Coupling Reactions of Benzyl Halides with Potassium Aryltrifluoroborates. *J. Org. Chem.* **2006**, *71*, 9198–9202.
- (32) For selected examples of pyridinium salts in EDA complexes, see: (a) Wu, J.; Grant, P. S.; Li, X.; Noble, A.; Aggarwal, V. K. Catalyst Free Deaminative Functionalizations of Primary Amines by Photo-induced Single-Electron Transfer. *Angew. Chem., Int. Ed.* **2019**, *58*, 5697–5701. (b) Katritzky, A. R.; de Ville, G. Z.; Patel, R. C. The S_{RN}2 Mechanism of Nucleophilic Substitution. *Tetrahedron Lett.* **1980**, *21*, 1723–1726. (c) Katritzky, A. R.; de Ville, G. Z.; Patel, R. C. Carbon Alkylation of Simple Nitronate Anions by *N*-Substituted Pyridiniums. *Tetrahedron* **1981**, *37*, 25–30. (d) Katritzky, A. R.; Chen, J.-L.; Marson, C. M.; Maia, A.; Kashmiri, M. A. The Non-Chain Radicaloid

C-Alkylation of Nitronate Anions: Further Evidence for the Mechanism. *Tetrahedron* **1986**, 42, 101–108. (e) Yang, M.; Cao, T.; Xu, T.; Liao, S. Visible-Light-Induced Deaminative Thioesterification of Amino Acid Derived Katritzky Salts via Electron Donor-Acceptor Complex Formation. *Org. Lett.* **2019**, 21, 8673–8678. (f) James, M. J.; Strieth-Kalthoff, F.; Sandfort, F.; Klauck, F. J. R.; Wagener, F.; Glorius, F. Visible-Light-Mediated Charge Transfer Enables C-C Bond Formation with Traceless Acceptor Groups. *Chem. - Eur. J.* **2019**, 25, 8240–8244. (g) Wu, J.; He, L.; Noble, A.; Aggarwal, V. K. Photoinduced Deaminative Borylation of Alkylamines. *J. Am. Chem. Soc.* **2018**, 140, 10700–10704. (h) Sandfort, F.; Strieth-Kalthoff, F.; Klauck, F. J. R.; James, M. J.; Glorius, F. Deaminative Borylation of Aliphatic Amines Enabled by Visible Light Excitation of an Electron Donor-Acceptor Complex. *Chem. - Eur. J.* **2018**, 24, 17210–17214. (i) Fu, M.-C.; Shang, R.; Zhao, B.; Wang, B.; Fu, Y. Photocatalytic decarboxylative alkylations mediated by triphenylphosphine and sodium iodide. *Science* **2019**, 363, 1429–1434.

(33) See the [Supporting Information](#) for the mechanism scheme with EDA complex involvement.

Recommended by ACS

Radical Hydrocarboxylation of Unactivated Alkenes via Photocatalytic Formate Activation

Sara N. Alektiar, Zachary K. Wickens, *et al.*

MAY 15, 2023
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ !\[\]\(f95dab70c751fda7d824b8b03650f7aa_img.jpg\)](#)

Photo- and Metal-Mediated Deconstructive Approaches to Cyclic Aliphatic Amine Diversification

David M. Soro, Richmond Sarpong, *et al.*

MAY 12, 2023
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ !\[\]\(e9474ce1d70442456f8fe9c393ea149c_img.jpg\)](#)

N-Allylation of Azoles with Hydrogen Evolution Enabled by Visible-Light Photocatalysis

Chang-Ming You, Hu Cai, *et al.*

MARCH 04, 2023
ORGANIC LETTERS

[READ !\[\]\(9db214d549b9aeebe72aa11d3a5c4b1a_img.jpg\)](#)

Photoredox-Nickel Dual-Catalyzed C-Alkylation of Secondary Nitroalkanes: Access to Sterically Hindered α -Tertiary Amines

Sina Rezazadeh, Donald A. Watson, *et al.*

FEBRUARY 16, 2023
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ !\[\]\(a05a1b59a958625e01d770867ed2a42e_img.jpg\)](#)

[Get More Suggestions >](#)