



Note

On the existence of balanced generalized de Bruijn sequences [☆]Matthew Baker^{a,*}, Bhumika Mittal^b, Haran Mouli^c, Eric Tang^d^a School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, United States of America^b Ashoka University, Sonapat, Haryana 131029, India^c University of Wisconsin-Madison, Madison, WI 53706, United States of America^d Harvard University, Cambridge, MA 02138, United States of America

ARTICLE INFO

Article history:

Received 29 June 2022

Received in revised form 10 April 2023

Accepted 20 April 2023

Available online xxxx

Keywords:

De Bruijn sequence

Eulerian digraph

ABSTRACT

A balanced generalized de Bruijn sequence with parameters (n, l, k) is a cyclic sequence of n bits such that (a) the number of 0's equals the number of 1's, and (b) each substring of length l occurs at most k times. We determine necessary and sufficient conditions on n, l , and k for the existence of such a sequence.

© 2023 Elsevier B.V. All rights reserved.

1. Statement of the main theorem

De Bruijn sequences, named after Nicolaas Govert de Bruijn (who first wrote about them in 1946) but first explored systematically by Camille Flye Sainte-Marie in 1894, are well-studied in the mathematical literature. A (binary) **de Bruijn sequence of order m** is a cyclic sequence where every possible m -bit substring occurs exactly once. It is well known ([2], see also [4]) that there are 2^{2^m-1-m} distinct de Bruijn sequences of order m . In this paper, we generalize de Bruijn sequences in the following manner:

Definition 1. A **generalized de Bruijn sequence** with parameters (n, l, k) is a cyclic sequence of n bits such that each substring of length l occurs at most k times. Such a sequence is called *balanced* if the number of 0s and 1s are equal.

In our terminology, a (classical) de Bruijn sequence of order m is a generalized de Bruijn sequence with parameters $(2^m, m, 1)$. Such a sequence is automatically balanced.

The main result of this paper characterizes those parameters (n, l, k) for which a balanced generalized de Bruijn sequence with parameters (n, l, k) exists:

[☆] The second, third, and fourth authors completed this research under the direction of the first author as part of the summer 2021 PROMYS program. We thank the PROMYS program for their support and for creating such a stimulating work environment. We also thank Persi Diaconis for helpful feedback on an earlier draft. The first author was supported by Simons Foundation Collaboration Grant 708029.

* Corresponding author.

E-mail addresses: mbaker@math.gatech.edu (M. Baker), bhumika.mittal_ug24@ashoka.edu.in (B. Mittal), hmouli@wisc.edu (H. Mouli), etang@college.harvard.edu (E. Tang).

Theorem 2. Given positive integers n, l , and k , a balanced generalized de Bruijn sequence with parameters (n, l, k) exists if and only if n is even and $k \geq \frac{n}{2}$.

In Section 3, we discuss our original motivation for introducing balanced generalized de Bruijn sequences, which came from the world of mathematical card tricks.

2. Proof of the main theorem

Our proof of Theorem 2 used ideas from graph theory. Specifically, we will be concerned with certain subgraphs of the following family of directed graphs:

Definition 3. For $l \geq 2$, the **de Bruijn graph** of rank l , denoted by G_l , is a directed graph on 2^{l-1} vertices, labeled by the elements of $\{0, 1\}^{l-1}$, and 2^l edges, labeled by the elements of $\{0, 1\}^l$. For any l bits $b_0, b_1, \dots, b_{l-1}$, the edge $e = b_0b_1 \dots b_{l-1}$ connects $v_1 = b_0b_1 \dots b_{l-2}$ to $v_2 = b_1b_2 \dots b_{l-1}$.

We begin with some simple lemmas.

Lemma 4. Every vertex in the de Bruijn graph G_l has both in-degree and out-degree 2.

Proof. Consider the vertex $v = b_0b_1 \dots b_{l-2} \in V(G_l)$. By inspection, v has in-edges $bb_0b_1 \dots b_{l-2}$ and out-edges $b_0b_1 \dots b_{l-2}b$ where $b \in \{0, 1\}$. Thus, v has both in-degree and out-degree 2. $\square$

Lemma 5. The de Bruijn graph G_l is Eulerian.

Proof. We know from Lemma 4 that every vertex in G_l has equal in-degree and out-degree. Thus, by a well-known criterion for the existence of Eulerian circuits in digraphs (see for example [1, Theorem 13.1.2]) it suffices to show that G_l is connected. This follows from the fact that given vertices $v = b_0b_1 \dots b_{l-2}$ and $v' = b'_0b'_1 \dots b'_{l-2}$, the following is a walk from v to v' :

$$b_0b_1 \dots b_{l-2} \rightarrow b_1 \dots b_{l-2}b'_0 \rightarrow \dots \rightarrow b'_0b'_1 \dots b'_{l-2}. \quad \square$$

It will be convenient to color each edge of G_l either red or blue according to the following rule:

Definition 6. We define an edge in G_l to be **red** if its last bit is 0 and **blue** if its last bit is 1. A subgraph of G_l is **balanced** if it contains an equal number of red and blue edges.

Lemma 7. Each vertex $v \in V(G_l)$ has one red and one blue out-edge, and the two in-edges of v have the same color.

Proof. Let $v = b_0b_1 \dots b_{l-2}$ be a vertex in G_l . Then by definition, the out-edges $b_0b_1 \dots b_{l-2}0$ and $b_0b_1 \dots b_{l-2}1$ are red and blue, respectively. The in-edges $0b_0b_1 \dots b_{l-2}$ and $1b_0b_1 \dots b_{l-2}$ must end with the same bit, and are thus of the same color. $\square$

Recall that a **circuit** in a graph is a path which begins and ends at the same vertex, and a **cycle** is a circuit in which no vertex (other than the initial and final one) is repeated.

By symmetry, the deBruijn graph G_l itself is balanced for all $l \geq 2$. We are interested in certain balanced subgraphs of G_l because of the following simple observation:

Lemma 8. There is a one-to-one correspondence between balanced circuits of length n in G_l and balanced generalized de Bruijn sequences with parameters $(n, l, 1)$.

Proof. Let the edges of the circuit of length n be denoted as:

$$b_0b_1 \dots b_{l-1} \rightarrow b_1b_2 \dots b_l \rightarrow \dots \rightarrow b_{n-1}b_0 \dots b_{l-2}.$$

The l -bit substrings of the n -bit sequence $b_0b_1 \dots b_{n-1}$ precisely correspond to the edges of the circuit of length n . Thus, as the edges on the circuit are distinct, so are the l -bit substrings. Moreover, since the number of red and blue edges in the circuit are equal, the number of 0s and 1s in the string are equal as well. Hence, the n -bit string is a balanced generalized de Bruijn sequence with parameters $(n, l, 1)$. $\square$

Lemma 9. If the graph G_l has a balanced circuit of length n , then the graph G_{l+1} has a balanced cycle of length n .

Proof. The edges of G_l and vertices of G_{l+1} have the same labelling. This gives us a natural bijection between the edges of G_l and the vertices of G_{l+1} . Let $e_1, e_2 \in G_l$ correspond to $v_1, v_2 \in G_{l+1}$ under this bijection. It follows from the definition of a de Bruijn graph that the head of e_1 is the tip of e_2 in G_l if and only if there is an edge connecting v_1 to v_2 in G_{l+1} . Now, consider the edges of the circuit of length n in G_l :

$$b_0 b_1 \cdots b_{l-1} \rightarrow b_1 b_2 \cdots b_l \rightarrow \cdots \rightarrow b_{n-1} b_0 \cdots b_{l-2}.$$

We consider these edges as the vertices of a walk in G_{l+1} . As all the edges of the circuit are distinct, the walk in G_{l+1} contains distinct vertices. Thus, the bijection maps a circuit of length n in G_l to an n -cycle in G_{l+1} . Moreover, since the circuit is balanced, there are equal number of 0s and 1s among $b_0, b_1, \dots, b_{n-1}$. These will be the last bits in the edges of the cycle in G_{l+1} as well, and hence, the cycle is balanced. $\square$

Lemma 10. Let $l \geq 2$, and let n be an even positive integer at most 2^l . Then the graph G_l contains a balanced circuit of length n .

Proof. We proceed by induction on l . First, for $l = 2$, we can either have $n = 2$ or $n = 4$. We may take the circuits $0 \rightarrow 1$ and $0 \rightarrow 1 \rightarrow 1 \rightarrow 0$, respectively.

Now, assume that the claim holds true for l . We prove it holds true for $l + 1$. For even $n \leq 2^l$, we may use the induction hypothesis to find a balanced circuit of length n in G_l . From Lemma 9, there exists a balanced cycle (also a circuit) of length n in G_{l+1} .

We now show the claim for $2^l < n < 2^{l+1}$ where n is even. Since $2 \leq 2^{l+1} - n < 2^l$, there exists a balanced $(2^{l+1} - n)$ -cycle in G_{l+1} , say H . Then the graph $G_{l+1} - H$ contains n edges. Assume that $G_{l+1} - H = H_1 \cup H_2 \cup \cdots \cup H_t$, where each H_i for $1 \leq i \leq t$ is a component of $G_{l+1} - H$. Since both G_{l+1} and H are balanced and Eulerian, $G_{l+1} - H$ must be balanced and each H_i must be Eulerian.

If $G_{l+1} - H$ is connected, we are done, since it is a circuit in G_{l+1} of length n . Otherwise, since G_{l+1} is connected, there exists an edge e in H which connects vertices in two different components of $G_{l+1} - H$, say H_1 and H_2 . Let the edge e go from $v_1 \in V(H_1)$ to $v_2 \in V(H_2)$. Without loss of generality, assume e is red. Now, let e_1 be an edge from v_1 to u_1 in H_1 , and let e_2 be an edge from u_2 to v_2 in H_2 . Then the last $l - 1$ bits of v_1 and u_2 are the first $l - 1$ bits of v_2 and u_1 . Hence, there must be an edge e' from u_2 to u_1 in H . We deduce from Lemma 7 that since e is red, e_1 must be blue, e_2 must be red, and e' must be blue.

Consider the new subgraph obtained by replacing e_1 and e_2 with e and e' . One checks easily that the degrees of all vertices, as well as the number of red and blue edges, are preserved. Moreover, the number of connected components has been reduced. Thus, we end up with a balanced subgraph with n edges and fewer components, all of which are Eulerian. By repeating this process, we will eventually end up with a balanced subgraph of n edges which is a circuit. Finally, for $n = 2^{l+1}$, all of the edges of the graph form a circuit since G_{l+1} is Eulerian. The result follows. $\square$

With these preliminaries out of the way, we now turn to the proof of our main result:

Proof of Theorem 2. ($\Rightarrow$) Since a balanced generalized de Bruijn sequence has an equal number of 0s and 1s, n must be even. Moreover, by the Pigeonhole Principle, we must have $k \geq \frac{n}{2^l}$, since there are only 2^l possible l -bit substrings and n total substrings.

($\Leftarrow$) Now, we show that these constraints are sufficient for a balanced generalized de Bruijn sequence to exist. We may assume that $k = \lceil \frac{n}{2^l} \rceil$ since a balanced generalized de Bruijn sequence with parameters (n, l, k) is also one with parameters (n, l, k') where $k' > k$.

For $l = 1$, the claim is trivial since the substrings are simply bits, and we may use any balanced string of length n . Now, let $l \geq 2$. We will now prove the claim by induction on k . The base case $k = 1$ follows from Lemmas 8 and 10. It suffices to show that the existence of a balanced generalized de Bruijn sequence with parameters (n, l, k) implies the existence of one with parameters $(n + 2^l, l, k + 1)$. Let $b_0 b_1 \cdots b_{n-1}$ be a balanced generalized de Bruijn sequence with parameters (n, l, k) , and let $b'_0 b'_1 \cdots b'_{2^l-1}$ be one with parameters $(2^l, l, 1)$. The latter contains every l -bit string as a substring exactly once. We may therefore assume that $b'_0 b'_1 \cdots b'_{l-1} = b_0 b_1 \cdots b_{l-1}$. Thus

$$b_0 b_1 \cdots b_{n-1} b'_0 b'_1 \cdots b'_{2^l-1}$$

is a balanced sequence in which each substring of length l appears at most k times in the first part of the sequence and exactly once in the second part. It is therefore a balanced generalized de Bruijn sequence with parameters $(n + 2^l, l, k + 1)$. The theorem follows by induction. $\square$

Remark 11. Theorem 2 can be generalized as follows. Say that a (finite) binary sequence is **almost-balanced** if the number of 0's and the number of 1's differ by at most one. Since the complement of an almost-balanced sequence is also almost-balanced, the same argument used to prove Theorem 2 yields the following:

Theorem 12. Given positive integers n, l , and k , an almost-balanced generalized de Bruijn sequence with parameters (n, l, k) exists if and only if $k \geq \frac{n}{2^l}$.

3. Motivation

One of our motivations for introducing balanced generalized de Bruijn sequences comes from the following card trick, a 32-card version of which is described on p.18 of [3]. The 52-card variant described here is alluded to on p.29 of [3].

Effect and presentation.

A regular deck of 52 cards, secured with rubber bands in its case, is tossed out into the audience. The deck is tossed around a few times until finally one audience member is instructed to remove the rubber bands, open the case, and give the deck a cut at some random position. This audience member takes the top card and then passes it to another spectator who removes the current top card. This is repeated until five members of the audience have taken cards. The performer instructs the five participants who have taken cards to look at their card, make a mental picture of it, and try to transmit an image of their card to the performer. The performer says, "You're all doing great, but some of your signals are crossing and I can't quite make everything out. Would everyone who has a red card please stand up and try extra hard to project the mental image of their card?" Turning to the first participant that took a card, the performer says, "I'm having trouble making out the suit of your card...it's not a heart, is it?" The participant shakes his head no. "No, I didn't think so...no, it's a diamond. It's the 9 of Diamonds!" The participant nods in agreement. Without asking any more questions, the performer goes on to identify the exact card that each of the other four participants is thinking of.

Method and explanation.

The trick is based on a balanced generalized de Bruijn sequence with parameters (52, 5, 2), for example the following one:

000001110101001000101100110111110000010110111101001

Each 5-bit string corresponds to either one or two cards in a standard deck of 52 cards; the color of the corresponding card is red if the first bit is a 0 and black if the first bit is a 1.

When the five spectators signal the color of their respective cards, this produces a 5-bit string (0 = red, 1 = black, going in order from spectator 1 to spectator 5). The magician looks up this 5-bit string in the following table:

00000: A♥, A♦	01000: 10♦, K♥	10000: 4♣, 4♠	11000: 7♠
00001: 7♥, 7♦	01001: K♦, 5♥	10001: 6♣	11001: J♠
00010: 6♥, 6♦	01010: 2♥	10010: 3♣, A♠	11010: 8♠, J♣
00011: 3♦	01011: 9♥, 9♦	10011: 5♣	11011: A♣, 2♠
00100: 5♦, 10♥	01100: J♥	10100: 7♣, 3♠	11100: 10♣
00101: 8♦, 8♥	01101: Q♥, 4♦	10101: 10♠	11101: K♠, 8♣
00110: 3♥	01110: 2♦	10110: Q♣, Q♠	11110: 9♠, K♣
00111: Q♦	01111: 4♥, J♦	10111: 6♠, 2♣	11111: 5♠, 9♣

If there are two possibilities listed, for example 9H or 9D, the magician feints hesitation, saying (in this example) to the first spectator, "I'm having trouble seeing the suit...it's not a heart, is it?" If the spectator says yes, the magician responds, "Yes, I thought so! It's the 9 of Hearts!" If the spectator says no, the magician responds, "No, I didn't think so...it's a diamond. It's the 9 of Diamonds!"

The magician now knows the first spectator's card, and is able to reveal all of the other selections by finding the first spectator's card in the following list (the next four cards – wrapping around cyclically, if necessary – are, in that order, the cards which the next four spectators are holding):

A♥ 7♥ 3♦ Q♦ 2♦ K♠ 8♠ 10♠ 2♥ 7♣ K♦ 3♣ 5♦ 10♦ 6♣ 6♥ 8♦ 9♥ Q♣ J♥ J♠ 5♣ 3♥ Q♥ A♣ 2♣
 4♦ 5♠ 9♠ 10♣ 7♠ 4♣ A♦ 7♦ 6♦ 8♥ 9♦ Q♠ 4♥ 2♠ 6♠ J♦ 9♣ K♣ 8♣ J♣ 3♠ 5♥ A♠ 10♥ K♥ 4♠

To disguise the glimpsing of the necessary information, once the spectators have signaled the color of their cards, grab a clipboard and a pen and say that you're going to try to write down the exact cards held by each of the five participants. Unbeknownst to the participants, on the clipboard you have a crib sheet containing the above information.

4. Some open problems

1. In lieu of binary bits, consider an alphabet A of size m where $m > 2$. Does a suitable analogue of Theorem 2 still hold? That is, given positive integers n, l, k , and m , does an A -balanced generalized de Bruijn sequence of length n exist if and only if m divides n and $k \geq \frac{n}{m^l}$? As considering the complement of such a sequence no longer suffices, the induction step used to prove Theorem 2 does not apply in this more general context.
2. Suppose n is even and $k \geq \frac{n}{2^l}$. How many balanced generalized de Bruijn sequences with parameters (n, l, k) are there? We know there is at least 1, and when $(n, l, k) = (2^m, m, 1)$ there are precisely $2^{2^{m-1}-m}$ such sequences. Is there an easily described general formula – at least in some special cases?

3. Are there algebraic ‘shift register’ type constructions that would permit the performer to do the $(52, 5, 2)$ trick described in Section 3 without mnemonics or a list? The trick described on p.18 of [3] (with parameters $(32, 5, 1)$) has this property. For analogous constructions related to a 6-card version of our $(52, 5, 2)$ trick, see [5] or [6].

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

- [1] Richard A. Brualdi, *Introductory Combinatorics*, fifth edition, Pearson Prentice Hall, Upper Saddle River, NJ, 2010.
- [2] N.G. de Bruijn, A combinatorial problem, *Nederl. Akad. Wetensch., Proc.* 49 (1946) 758–764 = *Indag. Math.* 8 (1946) 461–467.
- [3] Diaconis Persi, Ron Graham, *Magical mathematics*, in: *The Mathematical Ideas That Animate Great Magic Tricks*, Princeton University Press, Princeton, NJ, 2012. With a foreword by Martin Gardner.
- [4] Marshall Hall Jr., *Combinatorial Theory*, second edition, Wiley-Interscience Series in Discrete Mathematics, John Wiley & Sons, Inc., New York, 1986. A Wiley-Interscience Publication.
- [5] Christian Lawson-Perfect, David’s de Bruijn sequence card trick, Available at <https://aperiodical.com/2012/08/davids-de-bruijn-sequence-card-trick/>, 2012.
- [6] Siang Wun, Song Magic, Discover 6 in 52 cards knowing their colors based on the de Bruijn sequence, Available at <https://www.ime.usp.br/~song/magic/de-bruijn-magic52cards.pdf>.