



Gaussian unitary ensemble in random lozenge tilings

Amol Aggarwal¹ · Vadim Gorin²

Received: 25 June 2021 / Revised: 18 August 2022 / Accepted: 17 September 2022 /

Published online: 6 October 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

This paper establishes a universality result for scaling limits of uniformly random lozenge tilings of large domains. We prove that whenever the boundary of the domain has three adjacent straight segments inclined under 120 degrees (measured in the direction internal to the domain) to each other, the asymptotics of tilings near the middle segment is described by the GUE–corners process of random matrix theory. An important step in our argument is to show that fluctuations of the height function of random tilings on essentially arbitrary simply-connected domains of diameter N have magnitude smaller than $N^{1/2}$.

Keywords Lozenge tilings · GUE corners process · Universality

Mathematics Subject Classification Primary 82B20; Secondary 60K35 · 60B20

Contents

1 Introduction	1140
2 Setup and main results	1142
3 Concentration estimate	1149
4 Convergence to the GUE–corners process: proof	1159
A uniformity in convergence of local statistics	1163
References	1164

✉ Amol Aggarwal
amolaggarwal@math.columbia.edu

Vadim Gorin
vadicgor@gmail.com

¹ Columbia University, New York, NY, USA

² University of California, Berkeley, CA, USA

1 Introduction

Random lozenge tilings attracted substantial interest in the recent years following a 25 years old mathematical discovery that they exhibit highly non-trivial limit shapes with flat “frozen” facets; see [4, 7, 24, 29] for the pioneer results, [12] for an extensive review of the area, and Figs. 1 and 4 for simulations.

One intriguing feature of random tilings is their link to random matrix theory. The definition of the random tilings model does not have any obvious matrices involved, yet, the answers to the asymptotic questions about tilings turn out to involve distributions which are also encountered in the studies of eigenvalues of random matrices. In this paper we concentrate on the most direct of such connections and discuss how local limits near straight boundaries of tilings of general domains are described by the GUE–corners process—a multilevel version of the celebrated Gaussian Unitary Ensemble.

Definition 1 Let $X = [X_{ij}]_{i,j=1}^{\infty}$ be an infinite matrix of i.i.d. complex Gaussian random variables of the form $N(0, 1) + iN(0, 1)$ and set $M = \frac{X+X^*}{2}$ to be its Hermitian part. Define $\xi_1^k \leq \xi_2^k \leq \dots \leq \xi_k^k$ to be the eigenvalues of the principal $k \times k$ top-left corner of M as in (1.1). The array $\{\xi_i^k\}_{1 \leq i \leq k}$ is called the *GUE–corners process*.

$$\begin{pmatrix} M_{11} & M_{12} & M_{13} & M_{14} \\ M_{21} & M_{22} & M_{23} & M_{24} \\ M_{31} & M_{32} & M_{33} & M_{34} \\ M_{41} & M_{42} & M_{43} & M_{44} \end{pmatrix} \quad (1.1)$$

Let us explain the connections between tilings and the GUE–corners process on the example of uniformly random tilings of a hexagon, where it was first proved in [14, 22]. Consider a regular hexagon of side length N drawn on the triangular grid. We tile




Fig. 1 Left panel: A lozenge tiling of a regular hexagon and coordinates of horizontal lozenges along the left boundary: $y_1^1 = 2$ on the first vertical, $(y_1^2, y_2^2) = (0, 4)$ on the second vertical, $(y_1^3, y_2^3, y_3^3) = (0, 2, 6)$ on the third vertical. Right panel: A uniformly random tiling of the hexagon for $N = 50$ and a point where we see the GUE–corners process. Three types of lozenges are shown in three different colors

the hexagon with three types of lozenges (rhombuses) obtained by gluing two adjacent triangles on the grid; one possible tiling for $N = 5$ is shown in the left panel of Fig. 1. For fixed N there are finitely many such tilings and we are interested in a uniformly random lozenge tiling of the hexagon. A fascinating feature of random tilings is how their boundary conditions lead to inhomogeneous density profiles for three types of lozenges. The right panel of Fig. 1 shows a simulation for a large hexagon, where this aspect is clearly visible. In particular, one sees that inside the inscribed circle all three types of lozenges are present, while outside it there are six regions with only one type of lozenges in each. The points where the circle is tangent to the sides of the hexagon serve as the most drastic exemplification of this phenomenon since, even locally, the density of tiles there is maximally inhomogeneous. Simultaneously, these are the only points through which the tilings non-trivially interact with boundaries. Therefore, a question of interest is to mathematically understand these special points and to figure out the conditions of their appearance.

Let us distinguish one type of lozenge, say, we focus on the horizontal lozenges shown in gray in Fig. 1 and trace their positions near the left vertical boundary of the hexagon. Due to combinatorics of the model, there will be one horizontal lozenge with coordinate y_1^1 on the first vertical line (in our coordinate system shown in left panel of Fig. 1, a lozenge at the lower boundary of the hexagon has coordinate 0). Further, there will be two horizontal lozenges with coordinates $y_1^2 < y_2^2$ on the second vertical line, three horizontal lozenges with coordinates $y_1^3 < y_2^3 < y_3^3$ on the third line, etc. Since the tiling is random, y_i^k are random variables. We are interested in their asymptotic behavior as $N \rightarrow \infty$. [14, 22] showed that the limit is governed by the object of Definition 1:

$$\lim_{N \rightarrow \infty} \left(\frac{y_i^k - m_N}{\sqrt{N}} \right)_{1 \leq i \leq k} = (\sigma \xi_i^k)_{1 \leq i \leq k}, \quad \text{where } m_N = \frac{N}{2} \quad \text{and} \quad \sigma = \sqrt{\frac{3}{8}}, \quad (1.2)$$

in the sense of convergence of finite-dimensional distributions (jointly over all i and k such that $1 \leq i \leq k$). In particular, (1.2) implies that y_1^1 is asymptotically Gaussian as $N \rightarrow \infty$ and its variance grows proportionally to $\sqrt{N}$. In words, (1.2) says that discontinuity in densities of lozenges near a boundary leads to the appearance of a random matrix object.

Beyond tilings of the hexagon, convergence to the GUE–corners process near the boundaries, generalizing (1.2), was proven for several classes of domains; see [13, 21, 23, 25, 27]. The only necessary change is the expression for the constants m_N and σ . Simultaneously, Okounkov and Reshetikhin [25] explained that if the scaling limit of tilings near a boundary exists and satisfies certain very natural assumptions, then by invoking a classification theorem for Gibbs measures on interlacing arrays [26], one can conclude that the only possible candidate is the GUE–corners process. Hence, a universality prediction arose in [14, 25]: whenever a domain’s boundary has a straight segment, the GUE–corners process should asymptotically arise near this segment.

Our main result, Theorem 4, proves this prediction. While a possible path to a proof might have been through checking that the assumptions of [25] always hold, we did not know how to make such a check and found another approach. In our proof, we rely

on the results of [13, 23] which show that the GUE–corners process always appears in tilings of a special class of domains called trapezoids. We further observe that locally, near a straight segment of the boundary, any planar domain looks like a trapezoid. However (in contrast to the setting of [13, 23]) one boundary of the trapezoid is random and its fluctuations could a priori spoil the GUE–corners asymptotic. To resolve this issue, it must be shown that these fluctuations are $o(N^{1/2})$, that is, they have smaller scale than the GUE–corners limit—while widely expected,¹ such a bound for tilings of general domains has not been rigorously proven up to now. Known concentration estimates on arbitrary domains only apply on scale $O(N^{1/2})$; proving the improved $o(N^{1/2})$ estimate occupies a substantial portion of the manuscript. This is done by first using a (known) interpretation for the height variance through the double-dimer model, and then by patching together multiple instances of bulk local limit results of [1] for random lozenge tilings of arbitrary domains.

We expect that universal appearance of the GUE–corners process extends beyond lozenge tilings with the first candidates for probing the extensions given by the domino tilings and the six-vertex model, see [9–11, 14] for partial results in this direction. An interesting and important project for a future work is to figure out whether our approach can be helpful beyond lozenges.

Notations

Throughout the text, we let $|u - v|$ denote the Euclidean distance between any two points $u, v \in \mathbb{R}^2$; we also let $\text{dist}(\mathcal{S}, \mathcal{S}') = \inf_{s \in \mathcal{S}} \inf_{s' \in \mathcal{S}'} |s - s'|$ denote the distance between any sets $\mathcal{S}, \mathcal{S}' \subset \mathbb{R}^2$. We adopt a convention to write $C > 1$ for constants which are large and $c > 0$ for constants which are small. In general, these constants change from statement to statement.

2 Setup and main results

2.1 Tilings and height functions

Let us introduce the coordinate system on the triangular grid. We use x and y coordinate axes, which are inclined to each other by 120 degrees: y –coordinate grows up and x –coordinate grows in down-right direction, as in the left panel of Fig. 1. The *triangular lattice* $\mathbb{T}$ in this coordinate system becomes a graph whose vertex set is $\mathbb{Z}^2$ and whose edge set consists of edges connecting $(x, y), (x', y') \in \mathbb{Z}^2$ if and only if $(x' - x, y' - y) \in \{(1, 0), (-1, 0), (0, 1), (0, -1), (1, 1), (-1, -1)\}$. The faces of $\mathbb{T}$ are therefore triangles with vertices of the form $\{(x, y), (x+1, y), (x+1, y+1)\}$ or $\{(x, y), (x, y+1), (x+1, y+1)\}$. A *domain* $R \subseteq \mathbb{T}$ is a simply-connected² induced subgraph of $\mathbb{T}$. The *boundary* $\partial R \subseteq R$ is the set of all vertices $v \in R$ adjacent to a vertex in $\mathbb{T} \setminus R$.

¹ See Lectures 11–12 in [12] for the heuristics explaining that the fluctuations should grow logarithmically in the size of the domain.

² Tilings of multiply-connected regions in $\mathbb{T}$ can give rise to height functions that are multi-valued; hence, making sense of a limit shape (see Lemma 15 below) in this context requires some additional effort. This has been done recently in [19], so our methods should also extend to the multiply-connected setting as well, but we do not address this here.

A pair of adjacent triangular faces of $\mathbb{T}$ forms a parallelogram, which we refer to as a *lozenge* or *tile*. Lozenges can be oriented in one of three ways; see the left panel of Fig. 2 for all three orientations. We refer to the topmost lozenge there (that is, one with vertices of the form $\{(x, y), (x, y + 1), (x + 1, y + 2), (x + 1, y + 1)\}$) as a *type 1* lozenge. Similarly, we refer to the middle (with vertices of the form $\{(x, y), (x + 1, y), (x + 2, y + 1), (x + 1, y + 1)\}$) and bottom (vertices of the form $\{(x, y), (x, y + 1), (x + 1, y + 1), (x + 1, y)\}$) ones there as *type 2* and *type 3* lozenges, respectively. The horizontal lozenges discussed in the introduction are of type 2. We deal with tilings of R by lozenges of types 1, 2, and 3. Let $\Omega(R)$ denote the set of all tilings of R ; if $\Omega(R)$ is nonempty, we say that R is *tileable*.

A *height function*³ $H : R \rightarrow \mathbb{Z}$ is a function on the vertices of R that satisfies

$$\begin{aligned} H(v) - H(u) &\in \{0, 1\}, \quad \text{whenever } u = (x, y) \text{ and} \\ v &\in \{(x + 1, y), (x, y + 1), (x + 1, y + 1)\}, \end{aligned} \quad (2.1)$$

for some $(x, y) \in \mathbb{Z}^2$. We refer to the restriction $h = H|_{\partial R}$ as a *boundary height function*. For a fixed vertex $v \in R$ and integer $m \in \mathbb{Z}$, one can associate with any tiling of R a height function $H : R \rightarrow \mathbb{Z}$ as follows. First, set $H(v) = m$, and then define H at the remaining vertices of R in such a way that the height functions along the four vertices of any lozenge in the tiling are of the form depicted on the left panel of Fig. 2. In particular, we require that $H(x + 1, y) - H(x, y) = 1$ if and only if (x, y) and $(x + 1, y)$ are vertices of the same type 1 lozenge, and that $H(x, y + 1) - H(x, y) = 1$ if and only if (x, y) and $(x, y + 1)$ are vertices of the same type 2 lozenge. Since R is simply-connected, a height function on R is uniquely determined by these conditions and the value of $H(v) = m$.

We refer to the right panel of Fig. 2 for an example of a height function; there we chose v to be the bottom-left corner of the hexagon and $m = 0$. We can also view a lozenge tiling as a projection of a stepped surface in three-dimensional space: the three types of lozenges then become three types of faces of a unit cube, as shown in the middle panel of Fig. 2. The height function then counts the (signed) distance between a given point in three-dimensional space and coordinate plane corresponding to type 3 lozenges. In particular, the group of vertices with height 0 around the left corner of the right panel of Fig. 2 lies in this plane.

One important observation is that, if there exists a tiling $\mathcal{M}$ of R associated with some height function H , then the boundary height function $h = H|_{\partial R}$ is independent of $\mathcal{M}$ and is uniquely determined by R (except for a global shift, which was above fixed by the value of $H(v) = m$).

It will be useful to introduce continuum analogs of the above notions. So, set $\mathcal{T} = \{(s, t) : 0 < s + t < 1\} \subset \mathbb{R}^2$ and its closure $\overline{\mathcal{T}} = \{(s, t) : 0 \leq s + t \leq 1\}$. We interpret $\overline{\mathcal{T}}$ as the set of possible gradients, also called *slopes*, for a continuum height function; $\mathcal{T}$ is then the set of *liquid* slopes, whose associated tilings still “appear random.” For any simply-connected domain $\mathfrak{R} \subset \mathbb{R}^2$, we say that a function $\mathfrak{H} :$

³ There are other definitions of the height function; see, e.g., [12, Section 1.4] for a more symmetric one. All definitions contain the same information, because they differ from each other by explicit affine transformations.




Fig. 2 Left panel: three types of lozenges and corresponding changes of the height function. Middle panel: a tiled unit hexagon can be viewed as a projection of three faces of a unit cube in 3D. Right panel: a lozenge tiling and corresponding values of the height function

$\mathfrak{R} \rightarrow \mathbb{R}$ is *admissible* if $\mathfrak{h}$ is 1-Lipschitz and $\nabla \mathfrak{h}(z) \in \overline{\mathcal{T}}$ for almost all (with respect to Lebesgue measure) $z \in \mathfrak{R}$. We further say a function $\mathfrak{h} : \partial \mathfrak{R} \rightarrow \mathbb{R}$ *admits an admissible extension to $\mathfrak{R}$* if $\text{Adm}(\mathfrak{R}; \mathfrak{h})$, the set of admissible functions $\mathfrak{h} : \mathfrak{R} \rightarrow \mathbb{R}$ with $\mathfrak{h}|_{\partial \mathfrak{R}} = \mathfrak{h}$, is not empty.

2.2 Approximation of domains

Let us now explain the sense in which a sequence of discrete domains “converges” to a continuum one. To that end, for any connected subset $\mathcal{S} \subseteq \mathbb{R}^2$ and points $x_1, x_2 \in \mathcal{S}$, let $d_{\mathcal{S}}(x_1, x_2) = \inf_{\gamma} |\gamma|$, where $\gamma \subseteq \mathcal{S}$ is taken over all paths in $\mathcal{S}$ connecting x_1 and x_2 , and $|\gamma|$ denotes the length of γ . Next, we say that a sequence of connected subsets $\mathfrak{R}_1, \mathfrak{R}_2, \dots \subset \mathbb{R}^2$ *converges to $\mathfrak{R} \subset \mathbb{R}^2$* , and write $\lim_{N \rightarrow \infty} \mathfrak{R}_N = \mathfrak{R}$, if for any $\delta > 0$ there exists an integer $N_0 = N_0(\delta) > 1$ such that the following two properties hold whenever $N > N_0$. First, for any $x \in \mathfrak{R}$, there exist $x_N \in \mathfrak{R}_N$ and $x' \in \mathfrak{R} \cap \mathfrak{R}_N$ such that

$$\max \{d_{\mathfrak{R}}(x, x'), d_{\mathfrak{R}_N}(x_N, x')\} < \delta. \quad (2.2)$$

Second, for any $x_N \in \mathfrak{R}_N$, there exist $x \in \mathfrak{R}$ and $x' \in \mathfrak{R} \cap \mathfrak{R}_N$ such that (2.2) again holds.

In this case, we moreover say that a sequence of Lipschitz functions $\mathfrak{h}_N : \partial \mathfrak{R}_N \rightarrow \mathbb{R}$ *converges to $\mathfrak{h} : \partial \mathfrak{R} \rightarrow \mathbb{R}$* , and write $\lim_{N \rightarrow \infty} \mathfrak{h}_N = \mathfrak{h}$, if for every real number $\delta > 0$ there exists an integer $N_0 = N_0(\delta) > 1$ such that the following two properties hold whenever $N > N_0$. First, for each $x \in \partial \mathfrak{R}$, there exist $x_N \in \partial \mathfrak{R}_N$ and $x' \in \mathfrak{R} \cap \mathfrak{R}_N$ such that (2.2) and $|\mathfrak{h}_N(x_N) - \mathfrak{h}(x)| < \delta$ both hold. Second, for each $x_N \in \partial \mathfrak{R}_N$, there exist $x \in \partial \mathfrak{R}$ and $x' \in \mathfrak{R} \cap \mathfrak{R}_N$ such that the same inequalities are satisfied.

Definition 2 Suppose we are given a sequence $R_1, R_2, \dots \subset \mathbb{T}$ of domains and boundary height functions $h_N : \partial R_N \rightarrow \mathbb{Z}$, $N = 1, 2, \dots$. For each integer $N \geq 1$, define the domain $\mathfrak{R}_N = \frac{1}{N} R_N \subset \mathbb{R}^2$. We also define the function $\mathfrak{h}_N : \partial \mathfrak{R}_N \rightarrow \mathbb{R}$ by

setting $\mathfrak{h}_N(x) = \frac{1}{N}h_N(\frac{1}{N}x)$ for any point x on the lattice $\frac{1}{N}\partial R_N$, and by linear interpolation to all of $\partial\mathfrak{R}_N$. We say that $(R_N; h_N)$ converges to $(\mathfrak{R}; \mathfrak{h})$, for some $\mathfrak{R} \subset \mathbb{R}^2$ and $\mathfrak{h} : \partial\mathfrak{R} \rightarrow \mathbb{R}$, if $\lim_{N \rightarrow \infty} \mathfrak{R}_N = \mathfrak{R}$ and $\lim_{N \rightarrow \infty} \mathfrak{h}_N = \mathfrak{h}$.

Example 3 Let $\mathfrak{R}$ be a tileable polygon drawn on $\mathbb{T}$. Taking any tiling of $\mathfrak{R}$ and using definitions of Section 2.1, we get the corresponding boundary height function $\mathfrak{h} : \partial\mathfrak{R} \rightarrow \mathbb{Z}$. For $N = 1, 2, \dots$, let $R_N = N \cdot \mathfrak{R}$. It is straightforward to check that R_N is tileable; let h_N be the corresponding boundary height function. In this situation $(R_N; h_N)$ converges to $(\mathfrak{R}; \mathfrak{h})$, because we can choose $x = x' = x_N$ in (2.2).

Another point of view on this example is that a polygon $\mathfrak{R}$ is fixed and we are tiling it with lozenges of smaller and smaller side lengths $1/N$.

2.3 Convergence to the GUE–corners: statements

We are now ready to state our main theorem about convergence of random tilings to the GUE–corners process. We first explain a simpler version of the theorem, and then proceed to the most general setting.

Throughout this section we fix a sequence of tileable domains $R_N \subset \mathbb{T}$ with corresponding boundary height functions h_N , $N = 1, 2, \dots$ and assume that they converge to $(\mathfrak{R}; \mathfrak{h})$. We are going to choose a specific straight segment in the boundary of each R_N and assume that there are two adjacent straight segments inclined to it under an angle of 120 degrees in the internal direction of the domain: There are six options depending on the orientation of the straight segment, as shown in Fig. 3. In each of these six situations we trace a particular type of lozenges (whose shorter diagonal is parallel to the straight segment of interest) and get an interlacing configuration of lozenges, in the same way as in Fig. 1 we were tracing horizontal lozenges near a vertical segment of the boundary. We investigate the coordinates of these lozenges counted in the direction parallel to the straight segment of interest; in this way in Fig. 1 the segment was vertical and, hence, we were studying the vertical coordinates y_i^k . For notational simplicity, we state the next theorem for the vertical boundaries, as in the top-left panel of Fig. 3; applying rotations, exactly the same statement holds for other five orientations.

Theorem 4 Take a sequence of tileable simply-connected domains $R_N \subset \mathbb{R}$ and corresponding boundary height functions h_N . Suppose that (R_N, h_N) converges to some $(\mathfrak{R}, \mathfrak{h})$, where $\mathfrak{R} \subset \mathbb{R}^2$ is a simply-connected, bounded domain with piecewise smooth boundary. Further, suppose that the boundary of each R_N has a distinguished vertical segment I_N with two adjacent segments $I_N^{(l)}, I_N^{(r)}$ inclined by 120 degrees to I_N in the internal direction of the domain, as in the top-left panel of Fig. 3. Assume that as $N \rightarrow \infty$, rescaled by N , the positions and lengths of I_N , $I_N^{(l)}$, and $I_N^{(r)}$ converge to positions and lengths (which must remain positive) of straight segments of the boundary of $\mathfrak{R}$. Let $\{y_i^k(N)\}_{1 \leq i \leq k}$ denote the interlacing array of the vertical coordinates of horizontal lozenges adjacent to I_N in uniformly random tiling of R_N . Then there exist




Fig. 3 Six possible orientations of a straight segment and corresponding types of lozenges that we trace

constants $m_N \in \mathbb{R}$ and $\sigma \geq 0$, such that

$$\lim_{N \rightarrow \infty} \left(\frac{y_i^k(N) - m_N}{\sqrt{N}} \right)_{1 \leq i \leq k} = (\sigma \xi_i^k)_{1 \leq i \leq k}, \quad (2.3)$$

in the sense of convergence of finite-dimensional distributions (jointly over all i and k such that $1 \leq i \leq k$), and where $\{\xi_i^k\}$ is the GUE-corners process of Definition 1.

Remark 5 There are two ways to think about the constants m_N and σ . First, we can set $m_N = \mathbb{E}y_1^1(N)$ and $\sigma = \lim_{N \rightarrow \infty} N^{-1/2} \sqrt{\text{Var}(y_1^1(N))}$, i.e. they are the mean and asymptotic standard deviation of y_1^1 . Second, σ and the leading order⁴ behavior of m_N can be reconstructed through the limit shape (Law of Large Numbers) for tilings: informally, m_N is the average vertical coordinate of horizontal lozenges along a vertical slice of domain on small, but macroscopic distance to the right from the boundary segment of interest, while σ is a properly defined “variance” of these coordinates,

⁴ Subleading terms, i.e. $m_N - N \cdot \left[\lim_{N \rightarrow \infty} \frac{m_N}{N} \right]$ might depend on the exact way (R_N, h_N) approximates $(\mathfrak{R}, h)$, and there is no way to reconstruct them only from $(\mathfrak{R}, h)$ and the corresponding tiling limit shape.




Fig. 4 Left panel: Sample of a uniformly random lozenge tiling of a large S-shaped domain. Three types of tiles are shown in three colors. Right panel: Theorem 4 covers convergence to eight instances of the GUE–corners process shown in green circles. It does not cover four other instances shown by pink crosses. One embedded trapezoid is shown in red. The simulations are taken from [15] and we are grateful to the authors of that paper for allowing us to use their software (color figure online)

see Lemma 24 below for the precise formulas. We also mention that there might be degenerate situations, such as frozen domains, in which $\sigma = 0$.

Figure 4 illustrates Theorem 4 by showing the situations in which it applies. The same figure contains several situations where Theorem 4 does not apply because there are no three adjacent straight segments of the boundary inclined by 120 degrees to each other; yet, we believe that the GUE–corners process still appears asymptotically in those situations as well. The reason is that the system creates the straight boundaries on its own inside the frozen regions containing with high probability only one type of lozenges; once we know that an auxiliary straight boundary is there, we can apply a version of Theorem 4 again. Let us state this version.

Definition 6 Take a lozenge tiling $\mathcal{L} \in \Omega(R)$ of a domain $R \subset \mathbb{T}$. We say that a trapezoid with straight sides $I^{(l)}, I, I^{(r)}$ is embedded in $\mathcal{L}$, if:

- $I^{(l)}, I, I^{(r)}$ are three segments drawn on the triangular grid and inclined to each other by 120 degrees in one of the six configurations of Fig. 3; and
- $I^{(l)}, I$, and $I^{(r)}$ are inside (or on the boundary) of R ; and
- No lozenges of $\mathcal{L}$ intersect $I^{(l)}, I$, and $I^{(r)}$. In other words, these segments are formed by unions of various sides of lozenges of the tiling.

We refer to the right panel of Fig. 4 for an example of the embedded trapezoid.

Theorem 7 Take a sequence of tileable simply-connected domains $R_N \subset \mathbb{R}$ and corresponding boundary height functions h_N . Suppose that (R_N, h_N) converges to some $(\mathfrak{R}, \mathfrak{h})$, where $\mathfrak{R} \subset \mathbb{R}^2$ is a simply-connected, bounded domain with piecewise smooth boundary. Further, suppose that with probability tending to 1 as $N \rightarrow \infty$, each tiling of R_N has an embedded trapezoid (which depends on N , but not on the tiling) with straight sides $I_N^{(l)}, I_N, I_N^{(r)}$, oriented as in top-left panel of Fig. 3; in particular, I_N

is vertical. Assume that as $N \rightarrow \infty$, rescaled by N , the lengths of I_N , $I_N^{(l)}$, and $I_N^{(r)}$ converge to positive constants. Let $\{y_i^k(N)\}_{1 \leq i \leq k}$ denote the interlacing array of the vertical coordinates of horizontal lozenges adjacent to I_N in uniformly random tiling of R_N . Then there exist constants $m_N \in \mathbb{R}$ and $\sigma \geq 0$, such that

$$\lim_{N \rightarrow \infty} \left(\frac{y_i^k(N) - m_N}{\sqrt{N}} \right)_{1 \leq i \leq k} = (\sigma \xi_i^k)_{1 \leq i \leq k}, \quad (2.4)$$

in the sense of convergence of finite-dimensional distributions (jointly over all i and k such that $1 \leq i \leq k$), and where $\{\xi_i^k\}$ is the GUE–corners process of Definition 1.

Remark 8 We stated Theorem 7 for the orientation of the top-left panel of Fig. 3. Applying rotations, we conclude that it also applies for five other orientations.

Theorem 4 is a particular case of Theorem 7: indeed, in this situation the embedded trapezoid is formed by three straight segments of the boundary of the domain. This trapezoid is embedded in each tiling of R_N with probability 1, because lozenges are never allowed to cross the boundary of the domain.

The conditions of Theorem 7 are not only sufficient, but, in a sense, they are almost necessary: the only part which can be potentially weakened is the linear (in N) growth of the lengths of $I_N^{(l)}$, I_N , and $I_N^{(r)}$; there might be situations in which these lengths grow sublinearly and convergence to the GUE–corners process still holds.⁵ Let us explain necessity of the existence with high probability of the embedded trapezoid.

Imagine that we expect convergence of lozenge tilings to the GUE–corners process in a certain situation (still near a straight segment of a boundary of a domain). The eigenvalues ξ_i^k from Definition 1 almost surely interlace, see e.g., [12, Lecture 20], which means that

$$\xi_i^k < \xi_i^{k+1} < \xi_{i+1}^{k+1}, \quad 1 \leq i \leq k.$$

Hence, convergence to the GUE–corners process implies the existence of a similarly interlacing configuration of lozenges. Combinatorially, once we see an interlacing configuration of horizontal lozenges near a vertical segment of the boundary, far up and down along this boundary we should observe only lozenges of types 1 and 3, respectively, cf. the left panel of Fig. 5. In turn, the presence of lozenges of only one type far up and far down, readily leads to the appearance of an embedded trapezoid.

The conditions under which families of domains have the property that they admit embedded trapezoids are quite subtle. In general, such conditions cannot be read off of the macroscopic shape $\mathfrak{R}$; indeed, as indicated by the right panel of Fig. 5, a single microscopic defect along the boundary has the capacity to destroy this embedded

⁵ However, in some situations it fails. For instance, consider $N \times \lfloor N^\alpha \rfloor \times \lfloor N^\alpha \rfloor$ hexagon with $0 < \alpha < 1$; it obviously has an embedded trapezoid with $I_N^{(l)} = I_N^{(r)} = \lfloor N^\alpha \rfloor$. For this domain, as N tends to ∞ , we need to rescale the positions of lozenges near the longer side by factor $N^{1-\alpha/2}$ in order to see the GUE–corners process in the limit, as can be shown by analyzing explicit formulas of [7] and [22, Section 4]. Hence, if we rescale by $\sqrt{N}$, as in Theorems 4 and 7, the convergence fails. One can also design more complicated examples, in which there is no way to adjust rescaling factors to restore the GUE convergence.




Fig. 5 Left panel: Interlacing configuration of horizontal lozenges near a vertical segment of the boundary automatically leads to the appearance of an embedded trapezoid. Right panel: A microscopic defect in the boundary can prevent the GUE–corners asymptotics, because we would no longer have an interlacing array of horizontal lozenges

trapezoid property entirely. Therefore, we make no attempt in this paper (beyond Theorem 4) to classify under what circumstances this property holds. However, let us mention that the situation for polygonal domains, as in Example 3 and Fig. 4, seems promising: one could hope to find embedded trapezoids near most of the straight segments of the boundaries. On the technical level, proving this fact requires a careful analysis of the frozen regions of tilings and showing that with high probability the region occupied by lozenges of one type has not just a small amount, but actually no lozenges of other two types. We will not address this point here; for domains satisfying a technical assumption (conjecturally, this assumption is generic in side lengths of the polygonal domain; in particular it applies to the domain of Fig. 4) it follows as a consequence of [2].

3 Concentration estimate

3.1 Statement of the concentration bound

An important step in our proof of Theorems 4 and 7 is the following concentration estimate (possibly of independent interest) for height functions of random tilings on general simply-connected domains.

Proposition 9 Fix a simply-connected, bounded domain $\mathfrak{R} \subset \mathbb{R}^2$ with piecewise smooth boundary; a function $\mathfrak{h} : \partial\mathfrak{R} \rightarrow \mathbb{R}$ possessing an admissible extension to $\mathfrak{R}$; and any real number $\varepsilon > 0$. Let $R_1, R_2, \dots$ denote a family of simply-connected, tileable domains of $\mathbb{T}$, and for each integer $N \geq 1$ let $h_N : \partial R_N \rightarrow \mathbb{Z}$ denote an associated boundary height function; assume that $(R_N; h_N)$ converges to $(\mathfrak{R}, \mathfrak{h})$ in the sense of Definition 2. For each integer $N \geq 1$, let $H_N : R_N \rightarrow \mathbb{Z}$ denote the height function of a uniformly random tiling from $\Omega(R_N)$. Then there exists a constant $C > 1$ (depending on all the above data) such that for any integer $N \geq C$ and any vertex $v \in R_N$, we have $\text{Var } H_N(v) < \varepsilon N$.

Remark 10 [5, Theorem 21] implies that $\text{Var } H_N(v) \leq cN$ for a certain explicit constant $c > 0$. Our results shows that this constant can, in fact, be chosen to be arbitrarily small as long as N is large enough.

We deduce Proposition 9 from the following statement, which is proven in the rest of this section.

Proposition 11 In the setting of Proposition 9, let H'_N denote the height function of another uniformly random tiling from $\Omega(R_N)$, independent of H_N . For $N > C$, we have $\text{Var } [H_N(v) - H'_N(v)] < \varepsilon N$ for any $v \in R_N$.

Proof of Proposition 9 assuming Proposition 11 Since H_N and H'_N are independent, we have that $2 \text{Var } H_N(v) = \text{Var } [H_N(v) - H'_N(v)] < \varepsilon N$, where Proposition 11 was applied to deduce the inequality. $\square$

Throughout the remainder of this section, we adopt the setting of Proposition 11; abbreviate $H := H_N$ and $H' := H'_N$; define $F : R_N \rightarrow \mathbb{Z}$ by $F(v) = H(v) - H'(v)$; and also set $F(v) = 0$ for any $v \in \mathbb{T} \setminus R_N$. Observe from (2.1) that

$$F(u) - F(v) \in \{-1, 0, 1\}, \quad \text{whenever } u, v \in \mathbb{T} \text{ are adjacent.} \quad (3.1)$$

We refer to the left side of Fig. 6 for an example.

3.2 Distance graphs


A family $\mathcal{S} = (S_1, S_2, \dots, S_m)$ of mutually disjoint, connected subsets with $\bigcup_{i=1}^m S_i = R_N$ is called a *level set decomposition of R_N with respect to F* if the following hold.

1. If $v, v' \in S_i$ for some $1 \leq i \leq m$, then $F(v) = F(v')$.
2. If $v \in S_i$ and $v' \in S_j$ for some $1 \leq i < j \leq m$, and if v and v' are adjacent, then $F(v) \neq F(v')$.

We then set $F(S_i) = F(v)$ for any $v \in S_i$. We further say that $S_i, S_j \in \mathcal{S}$ are *adjacent* if $i \neq j$ and there exist $v \in S_i$ and $v' \in S_j$ such that v and v' are adjacent in $\mathbb{T}$. In particular, observe from (3.1) that

$$|F(S_i) - F(S_j)| = 1, \quad \text{whenever } S_i \text{ and } S_j \text{ are adjacent.} \quad (3.2)$$

We additionally say that S_j is *exterior adjacent* to S_i if S_j is contained in the infinite connected component of $\mathbb{T} \setminus S_i$; we otherwise call S_j *interior adjacent* to S_i .



Lemma 12 *For any $i \in [1, m]$, there is at most one element of $\mathbb{S}$ that is exterior adjacent to S_i .*

Observe by Lemma 12 that $\text{dist}(S_i, \partial R_N) > \text{dist}(S_j, \partial R_N)$ whenever S_j is exterior adjacent to S_i (since any path connecting ∂R_N to S_i must intersect S_j). In particular, if S_j is exterior adjacent to S_i , then S_i is interior adjacent to S_j . Given the above notation and Lemma 12, we define the distance graph.

Observe that $\mathcal{G}_S$ is a (deterministic) function of S . An example is depicted on the right side of Fig. 6. In what follows, we let $\text{dist}_{\mathcal{G}}(S_i, S_j)$ denote the distance between S_i and S_j under $\mathcal{G}$, and let $\text{dist}_{\mathcal{G}}(u, v) = \text{dist}_{\mathcal{G}}(S_i, S_j)$ whenever $v \in S_i$ and $u \in S_j$.

Lemma 14 *For any $u, v \in R_N$, we have $\text{Var}[F(u) - F(v)|\mathcal{S}] = \text{dist}_{\mathcal{G}}(u, v)$.*

1. The value of F on the root of $\mathcal{G}_S$ is 0. (Because $F|_{\partial R_N} = H|_{\partial R_N} - H'|_{\partial R_N} = 0$.)

2. $|F(S_i) - F(S_j)| = 1$ holds whenever S_i is adjacent to S_j (by (3.2)).

This description follows from the uniformity of the measure on lozenge tilings that we started from. Because $\mathcal{G}_S$ is a tree, we can sample F as follows: Associate with each edge e of $\mathcal{G}$ a Bernoulli random variable B_e such that $\mathbb{P}[B_e = 1] = \frac{1}{2} = \mathbb{P}[B_e = -1]$. Then, letting $S_i \in \mathcal{S}$ denote the level set containing v , set $F(v) = F(S_i)$ equal to $\sum_{e \in \mathcal{P}} B_e$, where $\mathcal{P}$ is the unique path of edges from the root of $\mathcal{G}$ to S_i . Taking variances in this equality, we deduce the lemma. $\square$

In view of Lemma 14, to show Proposition 11 it suffices to bound $\mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] = o(N)$; let us mention that the bound $\mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] = O(N)$ follows from the (deterministic) fact that $\text{dist}_{\mathcal{G}}(u, v) \leq |u - v|$.

3.3 Preliminary results

In this section we recall several known results from [1, 3, 6, 8] that will assist in the proof of the improved $o(N)$ estimate on $\mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)]$. We start from the results concerning the limiting height function and local statistics of random tilings. To that end, for any $x \in \mathbb{R}_{\geq 0}$ and $(s, t) \in \overline{\mathcal{T}}$, set

$$L(x) = - \int_0^x \log |2 \sin z| dz; \quad \sigma(s, t) = \frac{1}{\pi} \left(L(\pi s) + L(\pi t) + L(\pi(1 - s - t)) \right).$$

For any $\mathfrak{F} \in \text{Adm}(\mathfrak{R})$, define

$$\mathcal{E}(\mathfrak{F}) = \int_{\mathfrak{R}} \sigma(\nabla \mathfrak{F}(z)) dz.$$

Letting $\mathfrak{H} \in \text{Adm}(\mathfrak{R}; \mathfrak{h})$ denote the maximizer of $\mathcal{E}$ on $\text{Adm}(\mathfrak{R})$ with boundary data $\mathfrak{h}$, the following result of [6] claims that H converges to $\mathfrak{H}$ with high probability.

Lemma 15 ([6, Theorem 1.1]). *In the setting of Proposition 9, for any $\varepsilon > 0$, there exists a constant $C > 1$ such that if $N > C$ then*

$$\mathbb{P} \left[\max_{v \in R_N \cap N\mathfrak{R}} \left| \frac{1}{N} H_N(v) - \mathfrak{H}(\frac{1}{N}v) \right| > \varepsilon \right] < \varepsilon.$$

We will later use the following lemma from [8] that provides a deterministic property for the limit shape $\mathfrak{H}$ concerning the discontinuities of its gradient. It essentially states that at any discontinuity of $\nabla \mathfrak{H}$ there is a path to $\partial \mathfrak{R}$, parallel to an axis of $\mathbb{T}$, along which $\nabla \mathfrak{H}$ is minimal or maximal.

Lemma 16 ([8, Theorem 1.3]). *For any $z \in \mathfrak{R}$, we either have that $\nabla \mathfrak{H}$ is continuous at z or there exists some $z' \in \partial \mathfrak{R}$, with $z' - z = tw$ for some $t \in \mathbb{R}_{>0}$ and $w \in \{(1, 0), (0, 1), (1, 1)\}$, such that the following holds. If $w \in \{(1, 0), (0, 1)\}$, then $\mathfrak{H}(z + sw) = \mathfrak{H}(z)$ for each $s \in [0, t]$. If $w = (1, 1)$, then $\mathfrak{H}(z + sw) = \mathfrak{H}(z) + s$ for each $s \in [0, t]$.*

Next, although in this paper we are primarily interested in uniformly random lozenge tilings of finite domains, it will be useful to state several properties about certain infinite-volume measures on tilings of $\mathbb{T}$ that will arise as limit points of local statistics for our model. Therefore, let us briefly recall a two-parameter family of such measures from [16, 24, 28] (see also [17, Section 5] and [12, Lecture 13]). To that end, let $\mathfrak{P}(R)$ denote the space of probability measures on the set $\Omega(R)$ of all lozenge tilings of some domain $R \subseteq \mathbb{T}$. We say that $\mu \in \mathfrak{P}(\mathbb{T})$ is a *Gibbs measure* if it satisfies the following property for any finite domain $R \subset \mathbb{T}$. The probability under μ of selecting any $\mathcal{M} \in \Omega(\mathbb{T})$, conditional on the restriction of $\mathcal{M}$ to $\mathbb{T} \setminus R$, is uniform.

Moreover, for any $w \in \mathbb{Z}^2$, define the translation map $\mathfrak{S}_w : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ by setting $\mathfrak{S}_w(v) = v - w$ for any $v \in \mathbb{Z}^2$. Then $\mathfrak{S}_w$ induces an operator on $\mathfrak{P}(\mathbb{T})$ that translates a tiling by $-w$; we also refer to it by $\mathfrak{S}_w$. A measure $\mu \in \mathfrak{P}(\mathbb{T})$ is called *translation-invariant* if $\mathfrak{S}_w\mu = \mu$, for any $w \in \mathbb{Z}^2$. We further call a Gibbs translation-invariant measure $\mu \in \mathfrak{P}(R)$ *ergodic* if, for any $p \in (0, 1)$ and Gibbs translation-invariant measures $\mu_1, \mu_2 \in \mathfrak{P}(R)$ such that $\mu = p\mu_1 + (1 - p)\mu_2$, we have $\mu_1 = \mu = \mu_2$.

Let $\mu \in \mathfrak{P}(\mathbb{T})$ be a translation-invariant measure, and let H_μ denote the height function associated with a randomly sampled lozenge tiling under μ , such that $H_\mu(0, 0) = 0$. Setting $s = \mathbb{E}[H_\mu(1, 0) - H_\mu(0, 0)]$ and $t = \mathbb{E}[H_\mu(0, 1) - H_\mu(0, 0)]$, the *slope* of μ is defined to be the pair (s, t) . We must then have $(s, t) \in \overline{\mathcal{T}}$, since s , t , and $1 - s - t$ denote the probabilities of a given lozenge in a random tiling (under μ) being of types 1, 2, and 3, respectively.

It was shown in [28, Theorem 9.1.1] that, for any $(s, t) \in \overline{\mathcal{T}}$, there exists a unique ergodic Gibbs translation-invariant measure $\mu = \mu_{s,t} \in \mathfrak{P}(\mathbb{T})$ of slope (s, t) . We then have the following two results concerning this measure; the first one bounds the variance of the height function under any such $\mu_{s,t}$, and the second one shows convergence of local statistics of H to $\mu_{s,t}$.

In the below, we let $\mathcal{M}$ denote the law of the tiling of R_N associated with H and, for any $v \in \mathbb{T}$ and $D \in \mathbb{Z}_{\geq 0}$, we let $\mathcal{B}_D(v) = \{u \in \mathbb{T} : |u - v| \leq D\}$ denote the disk centered at v with radius D . By $\mathcal{M}|_{\mathcal{B}_D(v)}$ we mean the law of the tiling inside the disk $\mathcal{B}_D(v)$. Moreover, for any $\varepsilon > 0$, we let $\mathcal{T}_\varepsilon = \{(s, t) \in \mathbb{R}_{>0}^2 : \varepsilon < s + t < 1 - \varepsilon\}$, which describes the set of slopes that are “uniformly liquid.” Additionally, for any measures μ and ν on a probability space Ω with σ -algebra $\mathcal{A}$, we let

$$d_{\text{TV}}(\mu, \nu) = \max_{A \in \mathcal{A}} |\mu(A) - \nu(A)|$$

denote the total variation distance between μ and ν .

Lemma 17 ([3, Proposition 4.1]). *There exists a constant $C > 1$ such that the following holds. For any $(s, t) \in \mathcal{T}$ and $u, v \in \mathbb{T}$ we have under $\mu = \mu_{s,t}$ that $\text{Var}(H_\mu(u) - H_\mu(v)) \leq C \log |u - v|$.*

Let us mention that a logarithmic variance bound as in Lemma 17 was also established as Theorem 4.5 of [18], but there the uniformity in the slope (s, t) was not mentioned. [3] did not mention lozenge tilings explicitly, but the discrete sine process studied there is another face of the same object, see [24, Section 3.1.1] or [12, Section 13.4].

Lemma 18 ([1]). *In the setting of Proposition 9, for any real numbers $\varepsilon, \omega > 0$ and $D > 1$, there exists a constant $C > 1$ such that the following holds for any $N > C$. Let $v \in R_N$ be a vertex such that $\mathfrak{v} = \frac{1}{N}v \in \mathfrak{R}$ satisfies $\text{dist}(\mathfrak{v}, \partial\mathfrak{R}) \geq \varepsilon$ and $(s, t) = \nabla\mathfrak{H}(\mathfrak{v}) \in \mathcal{T}_\varepsilon$. Then, $d_{\text{TV}}(\mathcal{M}|_{\mathcal{B}_D(v)}, \mu_{s,t}|_{\mathcal{B}_D(v)}) < \omega$.*

Remark 19 The uniformity in v (and $\mathfrak{v}$) of the constant C stated in Lemma 18 was not directly claimed in (Theorem 1.5 of) [1] but follows from its proof. We outline this in Appendix 4.2 below.

3.4 Proof of Proposition 11

In this section we establish Proposition 11. We begin with the following lemma that bounds $\text{dist}_{\mathcal{G}}(u, v) = o(|u - v|)$ when $u - v$ are nearby.

Lemma 20 *There exists a constant $C_1 > 1$ such that for any real numbers $\varepsilon \in (0, 1)$ and $D > 2$, there is another constant $C_2 > 1$ (depending on all the data) and the following holds for any $N \geq C_2$: Fix $u, v \in R_N$ with $|u - v| \leq D$ satisfying $\text{dist}(\frac{1}{N}v, \partial\mathfrak{R}) \geq \varepsilon$ and $\nabla\mathfrak{H}(\frac{1}{N}u) \in \mathcal{T}_\varepsilon$. Then*

$$\mathbb{P}[\text{dist}_{\mathcal{G}}(u, v) > C_1(\log D)^2] \leq (\log D)^{-1}.$$

Proof Let $\omega = D^{-1}$ and set $(s, t) = \nabla\mathcal{H}(\frac{1}{N}u)$. Letting $\mathcal{M}$ and $\mathcal{M}'$ denote the tilings of R_N associated with H and H' , respectively, Lemma 18 implies (see [20, Proposition 4.7]) that $\mathcal{M}$ and $\mathcal{M}'$ can be coupled with $\mu_{s,t}$ such that they all coincide on $\mathcal{B}_D(v)$ with probability at least $1 - 2\omega$ for sufficiently large N . Hence, using the fact that H is 1-Lipschitz, Lemma 17 implies the existence of a constant $C > 1$ such that, for any fixed $D > 2$ and sufficiently large N , we have

$$\text{Var}[H(u) - H(v)] < C \log D + 2\omega D, \quad \text{and} \quad \text{Var}[H'(u) - H'(v)] < C \log D + 2\omega D,$$

whenever $|u - v| \leq D$. Adding these bounds, using the independence of H and H' , and $\omega = D^{-1}$, we obtain

$$\mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] = \text{Var}[F(u) - F(v)] < 2C \log D + 4,$$

where to deduce the first equality we used Lemma 14. It remains to use the Markov's inequality. $\square$

The next step is to deduce a lemma bounding $\text{dist}_{\mathcal{G}}(u, v) = o(N)$ on global scales in the liquid region; it will follow from applying Lemma 20 on all bounded intervals on a path between u and v .

Lemma 21 *For any $\varepsilon \in (0, 1)$, there exist constants $C = C(\varepsilon) > 1$ and $C' > 1$ (depending on all the data) such that the following holds for any $N \geq C'$: Fix $u, v \in R_N$ with $|u - v| > C$, such that $\text{dist}(\frac{1}{N}u, \partial\mathfrak{R}) \geq \varepsilon$ and $\text{dist}(\frac{1}{N}v, \partial\mathfrak{R}) \geq \varepsilon$. Further*

suppose that there exists an open disk $\mathfrak{U} \subset \mathfrak{R}$ containing $\frac{1}{N}u$ and $\frac{1}{N}v$, with $\nabla \mathfrak{H}(z) \in \mathcal{T}_\varepsilon$ for each $z \in \mathfrak{U}$. Then, $\mathbb{P}[\text{dist}_{\mathcal{G}}(u, v) > \varepsilon|u - v|] \leq \varepsilon$.

Proof Let $D > 2$ denote some integer, which will only depend on ε and $\mathfrak{R}$, and will be fixed later. Since $\text{dist}(\frac{1}{N}u, \partial\mathfrak{R}) \geq \varepsilon$; $\text{dist}(\frac{1}{N}v, \partial\mathfrak{R}) \geq \varepsilon$; and the disk $\mathfrak{U} \subset \mathfrak{R}$ contains $\frac{1}{N}u$ and $\frac{1}{N}v$, it follows that the line segment ℓ_0 connecting $\frac{1}{N}u$ and $\frac{1}{N}v$ is of distance at least ε from ∂R . Hence, for sufficiently large N , the line segment $N\ell_0$ connecting u and v is contained inside R_N and of distance at least $\frac{\varepsilon N}{2}$ from ∂R_N . Thus, there exists a sequence $u = v_0, v_1, \dots, v_k = v \in N\mathfrak{U} \cap R_N$ such that $|v_i - v_{i-1}| \leq D$ for each $i \in [1, k]$ and $k \leq \lceil 2D^{-1}|u - v| \rceil$ (for example, one may take the v_j to be k equally spaced points on $N\ell_0$, rounded to the nearest lattice points in R_N). Recall the constant $C_1 > 1$ from Lemma 20, and define for each $i \in [1, k]$ the event

$$\mathcal{E}_i = \{ \text{dist}_{\mathcal{G}}(v_{i-1}, v_i) > C_1(\log D)^2 \}.$$

Then, Lemma 20 implies $\mathbb{P}[\mathcal{E}_i] \leq (\log D)^{-1}$. So, since denoting by E^c the complement of any event E we have

$$\text{dist}_{\mathcal{G}}(u, v) \leq \sum_{i=1}^k (|v_i - v_{i+1}| \mathbf{1}_{\mathcal{E}_i} + C_1(\log D)^2 \mathbf{1}_{\mathcal{E}_i^c}),$$

it follows that

$$\begin{aligned} \mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] &\leq \sum_{i=1}^k D\mathbb{P}[\mathcal{E}_i] + C_1 k(\log D)^2 \\ &\leq kD(\log D)^{-1} + C_1 k(\log D)^2 \\ &\leq D(\log D)^{-1} \lceil 2D^{-1}|u - v| \rceil + C_1(\log D)^2 \lceil 2D^{-1}|u - v| \rceil \\ &\leq 4|u - v|((\log D)^{-1} + C_1 D^{-1}(\log D)^2), \end{aligned}$$

whenever $|u - v| \geq D$. Letting $C = C(\varepsilon) > 1$ so that $4(\log C)^{-1} + 4C_1 C^{-1}(\log C)^2 < \varepsilon^2$, we deduce by taking $D = C$ that $\mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] < \varepsilon^2|u - v|$, and so the lemma follows from the Markov's inequality. $\square$

The next lemma will be used in settings complimentary to Lemma 21. The claim is that if $u, v \in R_N$ are such that $v - u$ is parallel to an axis of $\mathbb{T}$ and H has approximately maximal or minimal slope along $v - u$, then $\text{dist}_{\mathcal{G}}(u, v)$ is small.

Lemma 22 Let $\delta \in (0, 1)$ be a real number, and let $u, v \in R_N$ denote vertices such that $v - u = tw$ for some $t \in \mathbb{Z}_{\geq 0}$ and $w \in \{(1, 0), (0, 1), (1, 1)\}$. Define the events

$$\begin{aligned} \mathcal{F}_1 &= \mathcal{F}_1(u, v; \delta) = \{H(v) - H(u) \leq \delta t\} \cap \{H'(v) - H'(u) \leq \delta t\}; \\ \mathcal{F}_2 &= \mathcal{F}_2(u, v; \delta) = \{H(v) - H(u) \geq (1 - \delta)t\} \cap \{H'(v) - H'(u) \geq (1 - \delta)t\}. \end{aligned} \quad (3.3)$$

On $\mathcal{F}_1 \cup \mathcal{F}_2$, we have $\text{dist}_{\mathcal{G}}(u, v) \leq 2\delta t$.

Proof In this proof, let us only assume that $\mathcal{F}_1$ holds, for the verification in the case when $\mathcal{F}_2$ holds is entirely analogous. Set $u_j = u + jw \in R_N$ for each $j \in [0, t]$, so that $u_t = v$. Note that both $H(u_j)$ and $H'(u_j)$ depend on j in a monotone way by our definition of the height function and the choice of w . Hence, using (3.3), we conclude that there exist at most δt indices $i \in [1, t]$ such that $H(u_i) - H(u_{i-1}) \neq 0$ and at most δt indices $i \in [1, t]$ such that $H'(u_i) - H'(u_{i-1}) \neq 0$. Thus, there exist at most $2\delta t$ indices $i \in [1, t]$ for which $F(u_i) = H(u_i) - H'(u_i) \neq H(u_{i-1}) - H'(u_{i-1}) = F(u_{i-1})$. Hence, for all but $(1 - 2\delta)t$ indices $i \in [1, t]$, we must have that u_i and u_{i-1} are in the same connected component, which implies that at most $2\delta t$ connected components separate u from v . $\square$

Using the above results, the following proposition bounds the (graph) distance between any vertex in R_N and the root of $\mathcal{G}$.

Proposition 23 *For any real number $\varepsilon > 0$, there exists a constant $C > 1$ such that for any $N > C$, we have that $\mathbb{P}[\text{dist}_{\mathcal{G}}(v, \partial R) > \varepsilon N] < \varepsilon$ for each $v \in R_N$.*

Proof Fix a real number $\delta > 0$ with $\delta < \frac{\varepsilon}{25}(1 + \text{diam } \mathfrak{R})^{-1}$, and let $\mathfrak{R}_\delta = \{z \in \mathfrak{R} : d(z, \partial \mathfrak{R}) > \delta\}$. Let $\mathfrak{v} = \frac{1}{N}v$. We may assume that $\mathfrak{v} \in \mathfrak{R}_\delta$, for otherwise for large N we have a deterministic bound

$$\text{dist}_{\mathcal{G}}(v, \partial R_N) \leq d(v, \partial R_N) \leq 2\delta N < \varepsilon N.$$

Our plan is to define a path $\{\zeta_0, \zeta_1, \dots, \zeta_k\} \subset \mathfrak{R} \cap \frac{1}{N}R_N$ from $\mathfrak{v} = \zeta_0$ to $\zeta_k \in \mathfrak{R} \setminus \mathfrak{R}_\delta$, so that⁶ $\text{dist}_{\mathcal{G}}(N\zeta_{i-1}, N\zeta_i) \leq 2\delta N|\zeta_i - \zeta_{i-1}|$ with high probability, for each $i \in [1, k]$. To that end, for each $z \in \mathfrak{R}$, we define an open subset $\mathcal{U}(z) \subset \mathfrak{R}$ as follows. If $\nabla \mathfrak{h}$ is continuous at z then let $\mathcal{U}(z)$ denote a disk, of radius at most $\frac{\delta}{2}$, centered at z such that $|\nabla \mathfrak{h}(z') - \nabla \mathfrak{h}(z)| < \delta$ for each $z' \in \mathcal{U}(z)$. If $\nabla \mathfrak{h}$ is discontinuous at z , then Lemma 16 implies that there exists $z' \in \partial \mathfrak{R}$ such that the conditions there hold; then let $\mathcal{U}(z)$ denote the δ -neighborhood of the segment connecting z and z' .

Because $\bigcup_{z \in \mathfrak{R}_\delta} \mathcal{U}(z)$ is an open cover of the compact domain $\overline{\mathfrak{R}_\delta}$, it admits a finite subcover $\bigcup_{i=1}^K \mathcal{U}(z_i)$ (where K depends on $\mathfrak{R}$ but is bounded independently of N). We call $z = z_i$ of type 1 if $\nabla \mathfrak{h}$ is discontinuous at z_i ; of type 2 if $\nabla \mathfrak{h}(z) \in \mathcal{T}_{2\delta}$ and $\nabla \mathfrak{h}$ is continuous at z ; and of type 3 if $\nabla \mathfrak{h}(z) \notin \mathcal{T}_{2\delta}$ and $\nabla \mathfrak{h}$ is continuous at z . If z is of type 3, then $\nabla \mathfrak{h}(z)$ is of distance at most 2δ from either the segment $[(0, 0), (1, 0)]$, $[(0, 0), (0, 1)]$, or $[(1, 0), (0, 1)]$ on $\partial \mathcal{T}$; we call z of types 3(a), 3(b), and 3(c) in these cases, respectively.

Now we define the sequence $\mathfrak{v} = \zeta_0, \zeta_1, \zeta_2, \dots, \zeta_k \in \mathfrak{R} \cap \frac{1}{N}R_N$ inductively as follows. First set $\zeta_0 = \mathfrak{v}$, and suppose ζ_i is defined for some $i \geq 0$. If $\zeta_i \in \mathfrak{R} \setminus \mathfrak{R}_\delta$, then set $k = i$ and stop. If otherwise $\zeta_i \in \mathfrak{R}_\delta$, then there exists some $z_j = z_{j(i)}$ such that $\zeta_i \in \mathcal{U}(z_j)$. We will define ζ_{i+1} depending on the type of z_j .

⁶ The points $N\zeta_i$ in $\text{dist}_{\mathcal{G}}(\cdot, \cdot)$ need to be vertices in $\mathbb{T}$, i.e., they should be confined to the integer lattice. Hence, we formally should write $\text{dist}_{\mathcal{G}}(\lfloor N\zeta_{i-1} \rfloor, \lfloor N\zeta_i \rfloor)$. For notational convenience we are going to omit the integer parts throughout this proof; this will not affect the validity of the argument.

1. If z_j is of type 1, then let $z' \in \partial\mathfrak{R}$ and $w \in \{(1, 0), (0, 1), (1, 1)\}$ be the point and vector corresponding to $z = z_j$ under Lemma 16, respectively. Set $k = i + 1$ and $\zeta = \zeta_k = z' - \frac{\delta}{2}w$.
2. If z_j is of type 2, then let $\zeta_{i+1} \in \partial\mathfrak{U}(z_j)$ denote the point such that $\zeta_{i+1} - \zeta_i$ is a positive multiple of $(1, 1)$.
3. Suppose z_j is of type 3.
 - (a) If z_j is of type 3(a), then let $\zeta_{i+1} \in \partial\mathfrak{U}(z_j)$ denote the point such that $\zeta_{i+1} - \zeta_i$ is a positive multiple of $(0, 1)$.
 - (b) If z_j is of type 3(b), then let $\zeta_{i+1} \in \partial\mathfrak{U}(z_j)$ denote the point such that $\zeta_{i+1} - \zeta_i$ is a positive multiple of $(1, 0)$.
 - (c) If z_j is of type 3(c), then let $\zeta_{i+1} \in \partial\mathfrak{U}(z_j)$ denote the point such that $\zeta_{i+1} - \zeta_i$ is a positive multiple of $(1, 1)$.

Let us make three comments about this definition. First, observe when z_j is of type 3 that ζ_{i+1} is chosen so that $\zeta_{i+1} - \zeta_i$ is approximately orthogonal to $\nabla\mathfrak{H}(z_i)$, namely, $\mathfrak{H}(\zeta_i) \approx \mathfrak{H}(\zeta_{i+1})$. Second, we have $\zeta_{i+1} \notin \mathfrak{U}(z_{j(i)})$ (since $\mathfrak{U}(z_{j(i)})$ is open), so $\zeta_{i+1} \neq \zeta_i$. In particular, since $\zeta_{i+1} - \zeta_i \in \mathbb{R}_{\geq 0}^2$, this implies that the process described above to define the sequence $\{\zeta_0, \zeta_1, \dots, \zeta_k\}$ eventually ends. Third, as can be verified by induction on j , we have $\text{dist}(\zeta_j, \partial\mathfrak{R}_\delta) \geq \frac{\delta}{2}$ for each $j \in [0, k]$, so that $N\zeta_k \in R_N = R$ for sufficiently large N .

Next, fix some $i \in [0, k - 1]$, and set $\tilde{u} = N\zeta_i$ and $\tilde{v} = N\zeta_{i+1}$; further denote $z_j = z_{j(i)}$. We claim that

$$\begin{aligned} \mathbb{P}[\text{dist}_{\mathcal{G}}(\tilde{u}, \tilde{v}) > 4\delta N(\text{diam } \mathfrak{R} + 1)] &< \delta k^{-1}, & \text{if } z_j \text{ is of type 1;} \\ \mathbb{P}[\text{dist}_{\mathcal{G}}(\tilde{u}, \tilde{v}) > 8\delta|\tilde{u} - \tilde{v}|] &< \delta k^{-1}, & \text{if } z_j \text{ is of type 2 or 3.} \end{aligned} \quad (3.4)$$

Before proving (3.4), let us show how it implies the statement of Proposition 23. To that end, since $k = i + 1$ whenever $z_{j(i)}$ is of type 1, there is at most one index i such that $z_j = z_{j(i)}$ is of type 1. Hence, applying (3.4) and a union bound, we deduce

$$\begin{aligned} \mathbb{P}[\text{dist}_{\mathcal{G}}(v, \partial R_N) > \varepsilon N] &\leq \mathbb{P}[\text{dist}_{\mathcal{G}}(v, N\zeta_k) > (\varepsilon - 2\delta)N] \\ &\leq \sum_{i=1}^{k-1} \mathbb{P}[\text{dist}_{\mathcal{G}}(N\zeta_{i-1}, N\zeta_i) > 8\delta|\zeta_i - \zeta_{i-1}|N] \\ &\quad + \mathbb{P}[\text{dist}_{\mathcal{G}}(N\zeta_{k-1}, N\zeta_k) > 4\delta N(\text{diam } \mathfrak{R} + 1)] < \delta < \varepsilon. \end{aligned} \quad (3.5)$$

Here, for the first inequality, we used the fact that

$$\text{dist}_{\mathcal{G}}(v, \partial R_N) \leq \text{dist}_{\mathcal{G}}(v, N\zeta_k) + \text{dist}_{\mathcal{G}}(N\zeta_k, \partial R_N) \leq \text{dist}_{\mathcal{G}}(v, N\zeta_k) + 2\delta N,$$

which holds for large N since $\zeta_k \in \mathfrak{R} \setminus \mathfrak{R}_\delta$. For the second estimate, we used the union bound and the fact that

$$\begin{aligned} \sum_{i=1}^{k-1} 8\delta |\zeta_i - \zeta_{i-1}| + 4\delta(\text{diam } \mathfrak{R} + 1) &\leq 16|\zeta_{k-1} - \zeta_0| \\ + 4\delta(\text{diam } \mathfrak{R} + 1) &\leq 20\delta(\text{diam } \mathfrak{R} + 1) \leq \varepsilon - 2\delta, \end{aligned}$$

where the first bound holds since $\zeta_i - \zeta_{i-1}$ is for each $i \in [1, k-1]$ a positive multiple of some vector among $\{(1, 0), (0, 1), (1, 1)\}$; the second holds since $|\zeta_{k-1} - \zeta_0| \leq \text{diam } \mathfrak{R}$; and the third holds since $25\delta(\text{diam } \mathfrak{R} + 1) < \varepsilon$. Since (3.5) verifies the proposition, it remains to establish (3.4), which we do by analyzing each case for z_j individually.

Case 1: z_j is of type 1, meaning that $\mathfrak{U}(z_j)$ is the δ -neighborhood of the line $\ell = \ell(z_j, \zeta_{i+1})$ connecting z_j to the point $z' \in \partial\mathfrak{R}$, and that $z' - \frac{\delta}{2}w = \zeta_{i+1} = \frac{1}{N}\tilde{v}$. Here the analysis is based on Lemma 22.

Denoting $z' - z_j = tw$ for some $t \in \mathbb{R}_{>0}$ and $w \in \{(1, 0), (0, 1), (1, 1)\}$, Lemma 16 implies that $\mathfrak{H}(\zeta_{i+1}) - \mathfrak{H}(z_j) = 0$ if $w \in \{(1, 0), (0, 1)\}$ and $\mathfrak{H}(\zeta_{i+1}) - \mathfrak{H}(z_j) = t - \frac{\delta}{2}$ if $w = (1, 1)$. Let $\tilde{z} = z_j + sw \in \ell$, for some $s \in [0, t]$, denote the point on ℓ closest to $\frac{1}{N}\tilde{u}$. Then, Lemma 15 implies for sufficiently large N that

$$\begin{aligned} \mathbb{P}\left[|H(\tilde{v}) - N\mathfrak{H}(\zeta_{i+1})| \leq \delta N\right] &> 1 - \frac{\delta}{4k}; & \mathbb{P}\left[|H(N\tilde{z}) - N\mathfrak{H}(\tilde{z})| \leq \delta N\right] &> 1 - \frac{\delta}{4k}; \\ \mathbb{P}\left[|H'(\tilde{v}) - N\mathfrak{H}(\zeta_{i+1})| \leq \delta N\right] &> 1 - \frac{\delta}{4k}; & \mathbb{P}\left[|H'(N\tilde{z}) - N\mathfrak{H}(\tilde{z})| \leq \delta N\right] &> 1 - \frac{\delta}{4k}. \end{aligned}$$

It follows from a union bound that

$$\begin{aligned} \mathbb{P}\left[\left|H(\tilde{v}) - H(N\tilde{z}) - N(\mathfrak{H}(\zeta_{i+1}) - \mathfrak{H}(\tilde{z}))\right| \leq 2\delta N\right] &> 1 - \frac{\delta}{2k}; \\ \mathbb{P}\left[\left|H'(\tilde{v}) - H'(N\tilde{z}) - N(\mathfrak{H}(\zeta_{i+1}) - \mathfrak{H}(\tilde{z}))\right| \leq 2\delta N\right] &> 1 - \frac{\delta}{2k}. \end{aligned}$$

So, since z' is given through Lemma 16 and $\zeta_{i+1} = z' - \frac{\delta}{2}$, we deduce from a union bound that

$$\mathbb{P}[\mathcal{F}_1(N\tilde{z}, \tilde{v}; 2\delta) \cup \mathcal{F}_2(N\tilde{z}, \tilde{v}; 2\delta)] > 1 - \delta k^{-1},$$

where we recall the events $\mathcal{F}_1$ and $\mathcal{F}_2$ from (3.3). Hence, Lemma 22 gives

$$\mathbb{P}[\text{dist}_{\mathcal{G}}(N\tilde{z}, \tilde{v}) \leq 4\delta N|\tilde{z} - \zeta_{i+1}|] \geq \mathbb{P}[\text{dist}_{\mathcal{G}}(N\tilde{z}, \tilde{v}) \leq 4\delta sN] > 1 - \delta k^{-1}. \quad (3.6)$$

Since $\frac{1}{N}\tilde{u} = \zeta_i \in \mathfrak{U}(z_j)$, we have $|\tilde{z} - \frac{1}{N}\tilde{u}| < \delta$. So,

$$\text{dist}_{\mathcal{G}}(\tilde{u}, \tilde{v}) \leq \text{dist}_{\mathcal{G}}(\tilde{u}, N\tilde{z}) + \text{dist}_{\mathcal{G}}(N\tilde{z}, \tilde{v}) \leq |\tilde{u} - N\tilde{z}| + \text{dist}_{\mathcal{G}}(N\tilde{z}, \tilde{v}) \leq \text{dist}_{\mathcal{G}}(N\tilde{z}, \tilde{v}) + N\delta,$$

which together with (3.6) (and the bound $|\tilde{z} - \zeta_{i+1}| \leq \text{diam } \mathfrak{R}$) implies the first statement of (3.4).

Case 2: z_j is of type 2, in which case $\nabla \mathfrak{H}(z) \in \mathcal{T}_\delta$ for each $z \in \overline{\mathfrak{U}(z_j)}$. Here Lemma 21 applies directly. Indeed, because $\zeta_i, \zeta_{i+1} \in \overline{\mathfrak{U}(z_j)}$ and $|\tilde{u} - \tilde{v}| = N|\zeta_i - \zeta_{i+1}| \geq c_1 N$ for some constant $c_1 = c_1(\mathfrak{R}, \mathfrak{h}) > 0$, we deduce the second statement of (3.4) from Lemma 21 applied with the ε there equal to δk^{-1} .

Case 3: z_j is of type 3. In this situation we again rely on Lemma 22. Without loss of generality we assume that z_j is of type 3(a), as the cases when it is of type 3(b) or type 3(c) are very similar. Then, $\nabla \mathfrak{H}(z)$ is of distance at most 3δ from the segment $[(0, 0), (1, 0)] \subset \partial \mathcal{T}$ for any $z \in \overline{\mathfrak{U}(z_j)}$. Hence, since $\zeta_{i+1} - \zeta_i$ is a multiple of $(0, 1)$, we have $|(\zeta_{i+1} - \zeta_i) \cdot \nabla \mathfrak{H}(z)| \leq 3\delta$ for each $z \in \mathfrak{U}(z_j)$. Integrating z from ζ_i to ζ_{i+1} thus gives $|\mathfrak{H}(\zeta_i) - \mathfrak{H}(\zeta_{i+1})| < 3\delta|\zeta_i - \zeta_{i+1}|$. Hence, Lemma 15 implies for sufficiently large N that

$$\mathbb{P}[H(\tilde{v}) - H(\tilde{u}) < 4\delta|\tilde{u} - \tilde{v}|] \geq 1 - \frac{\delta}{2k}; \quad \mathbb{P}[H'(\tilde{v}) - H'(\tilde{u}) \leq 4\delta|\tilde{u} - \tilde{v}|] \geq 1 - \frac{\delta}{2k}.$$

Recalling the event $\mathcal{F}_1$ from (3.3), it follows from a union bound that $\mathbb{P}[\mathcal{F}_1(\tilde{u}, \tilde{v}; 4\delta)] \geq 1 - \delta k^{-1}$, and so the second statement of (3.4) follows from Lemma 22. This verifies (3.4) in all cases; as mentioned above, this implies the proposition. $\square$

We have now collected all the ingredients for the proof of Proposition 11.

Proof of Proposition 11 Fix $\delta > 0$ such that $\delta(1 + \text{diam } \mathfrak{R}) = \frac{\varepsilon}{2}$, and fix a vertex $u \in \partial R_N$ closest to v . Then, for sufficiently large N we have

$$\begin{aligned} \text{Var}[F(v)] &= \mathbb{E}[\text{dist}_{\mathcal{G}}(u, v)] \leq \delta N + \mathbb{P}[\text{dist}_{\mathcal{G}}(u, v) > \delta N] \text{diam } R_N \\ &\leq \delta N + \delta \text{diam } R_N \leq \delta N(1 + \text{diam } \mathfrak{R}) + \frac{\delta N}{4} < \varepsilon N. \end{aligned}$$


Here, the first statement holds by Lemma 14 and the deterministic fact that $F(u) = 0$ (as $u \in \partial R_N$); the second by the deterministic bound $\text{dist}_{\mathcal{G}}(u, v) \leq |u - v| \leq \text{diam } R_N$; the third by Proposition 23; the fourth by the fact that $\frac{1}{N} \text{diam } R_N \leq \text{diam } \mathfrak{R} + \frac{\delta}{4}$ holds for sufficiently large N (since $\lim_{N \rightarrow \infty} \frac{1}{N} R_N = \mathfrak{R}$); and the fifth by the definition of δ . This establishes the proposition. $\square$

4 Convergence to the GUE–corners process: proof

4.1 Tilings of fixed trapezoids

While the first ingredient of our proof of Theorem 7 is Proposition 9, the second equally important component is the asymptotic analysis of lozenge tilings of trapezoids from [13, 23]. Let us recall the result from these texts which we need. By a *fixed trapezoid* we mean a domain of Fig. 7: it is parameterized by the width L , the length A of the

Fig. 7 A tiling of a fixed trapezoid and corresponding triangular array of coordinates of horizontal lozenges



vertical boundary, and by L -tuple of integers $0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_L < A + L$, which encodes the positions of the horizontal lozenges along the right boundary of the domain. We refer to these L lozenges as *dents*. The lozenge tilings of a fixed trapezoid are in bijection with triangular arrays $(y_i^k)_{1 \leq i \leq k \leq L}$ of $L(L+1)/2$ integers, which satisfy:

$$y_i^{k+1} \leq y_i^k < y_{i+1}^k, \quad 1 \leq i \leq k < L, \quad \text{and} \quad (y_1^L, \dots, y_L^L) = (\lambda_1, \dots, \lambda_L). \quad (4.1)$$

The integers of the array encode positions of horizontal lozenges, as shown in Fig. 7. The role of the parameter A for the array is very limited: it only appears in the inequality $\lambda_L < A + L$. However, this parameter becomes more important when we want to reconstruct a tiling: the same array can lead to slightly different tilings depending on the value of A .

We are interested in the behavior of uniformly random lozenge tilings of a fixed trapezoid with dents $\lambda_1, \dots, \lambda_L$ as L and A tend to infinity. In this situation $(y_i^k)_{1 \leq i \leq k \leq L}$ become random variables and we would like to understand their asymptotics. For finite i and k the limiting behavior is summarized in the following lemma, which is a slight reformulation of results from [13] and [23].

Lemma 24 ([13, Theorem 1.7], [23, Theorem 1]). *Suppose that $\lambda = (\lambda_1, \dots, \lambda_L)$ and A depend on L , with $0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_L < A + L$. Then, in the sense of convergence in finite-dimensional distributions, we have*

$$\lim_{L \rightarrow \infty} \frac{y_i^k - m(\lambda)}{\sigma(\lambda)\sqrt{L}} = \xi_i^k, \quad \text{jointly over } 1 \leq i \leq k, \quad (4.2)$$

where ξ_i^k is the GUE-corners process of Definition 1,

$$m(\lambda) = \left\lfloor \sum_{i=1}^L \frac{\lambda_i}{L} \right\rfloor - \frac{L}{2}, \quad \sigma(\lambda)^2 = \frac{1}{L} \sum_{i=1}^L \left(\frac{\lambda_i}{L} \right)^2 - \left(\frac{1}{L} \sum_{i=1}^L \frac{\lambda_i}{L} \right)^2 - \frac{1}{12}, \quad (4.3)$$

and the convergence is uniform over A such that A/L stays bounded as $L \rightarrow \infty$ and λ such that $\sigma(\lambda)$ stays bounded away from 0 as $L \rightarrow \infty$. If instead $\sigma(\lambda)$ tends to 0

as $L \rightarrow \infty$, then (4.2) should be interpreted as convergence of $\frac{y_i^k - m(\lambda)}{\sqrt{L}}$ to the zero vector.

Remark 25 The minimal value that $\sigma(\lambda)^2$ can take is achieved when $(\lambda_1, \dots, \lambda_L) = (B, B+1, \dots, B+L-1)$. In this situation the lozenge tiling is completely frozen (i.e. there exists only one array satisfying (4.1)) and

$$\sigma(\lambda)^2 = \frac{L(L-1)(2L-1)}{6L^3} - \left(\frac{L(L-1)}{2L^2} \right)^2 - \frac{1}{12} = \frac{L-1}{L} \left(\frac{2L-1}{6L} - \frac{L-1}{4L} \right) - \frac{1}{12} \xrightarrow{L \rightarrow \infty} 0.$$

On the other hand, if we assume that the empirical measures $\frac{1}{L} \sum_{i=1}^L \delta_{\lambda_i/L}$ converge as $L \rightarrow \infty$ to a probability measure⁷ other than a uniform measure on an interval of length one, then σ^2 converges to a positive constant.

4.2 Proof of Theorem 7

In Lemma 24 we dealt with fixed trapezoids, which means that the L -tuples of dents $(\lambda_1, \dots, \lambda_L)$ were deterministic. Let us now introduce a notion of *fluctuating trapezoid*: this is a trapezoid in which the dents are allowed to be random. When we discuss tilings of a fluctuating trapezoid, we assume that the law of the dents can be arbitrary, but conditionally on the positions of the dents, the probability distribution on lozenge tilings inside the trapezoid is uniform.

For each $N = 1, 2, \dots$, let us look at the embedded trapezoid of Theorem 7. With probability tending to 1 as $N \rightarrow \infty$, each tiling of R_N gives a rise to a lozenge tilings of the (embedded) fluctuating trapezoid. Let us denote the width of this trapezoid through $L = L(N)$ and its dents through $\lambda_i = \lambda_i(N)$, $1 \leq i \leq L$. We would like to set $m_N = \mathbb{E}m(\lambda)$ and $\sigma^2 = \lim_{N \rightarrow \infty} \sigma(\lambda)^2$, as given by (4.3), and apply Lemma 24 to get (2.4) (note that A/L in Lemma 24 is uniformly bounded, since all the domains involved in Theorem 7 and the ratio N/L are bounded). The caveat is that (4.2) involves *random* $m(\lambda)$ and $\sigma(\lambda)$, rather than deterministic m_N and σ of (2.4). Hence, we need to prove the following two claims:

Claim I: $\lim_{N \rightarrow \infty} \frac{m(\lambda) - \mathbb{E}m(\lambda)}{\sqrt{N}} = 0$, in probability.

Claim II: There exists a deterministic limit in probability $\lim_{N \rightarrow \infty} \sigma(\lambda)^2$.

For Claim I we need to rewrite $m(\lambda)$ in terms of the height function. Let $H_L(y)$ denote the height function of the tiling along the vertical line passing through the dents of the embedded trapezoid. In more detail, following the notation of Sect. 2.1, we choose the origin of the coordinate system to be in the bottom-right corner of the embedded trapezoid and set $H_L(y) := H(0, y)$. As the local rules of Fig. 2 show, this function starts as a constant C at the bottom of the trapezoid and as we move up it grows by 1 whenever we cross a horizontal lozenge and stays constant otherwise.

⁷ $\lambda_{i+1} - \lambda_i \geq 1$ implies that the Lebesgue density of the limiting measure is at most one.

Hence, we can write

$$\sum_{i=1}^L \lambda_i = \sum_{y=0}^{A+L-1} y(H_L(y+1) - H_L(y)) = (A+L-1)H_L(A+L) - \sum_{y=1}^{A+L-1} H_L(y). \quad (4.4)$$

Because $H_L(A+L)$ is deterministic, we conclude that

$$m(\lambda) - \mathbb{E}m(\lambda) = \frac{1}{L} \left(\sum_{i=1}^L \lambda_i - \mathbb{E} \sum_{i=1}^L \lambda_i \right) = -\frac{1}{L} \sum_{y=1}^{A+L-1} (H_L(y) - \mathbb{E}H_L(y)) \quad (4.5)$$

Using a bound on the variance of a sum of arbitrary random variables

$$\text{Var}(\alpha_1 + \alpha_2 + \cdots + \alpha_K) \leq K(\text{Var} \alpha_1 + \text{Var} \alpha_2 + \cdots + \text{Var} \alpha_K),$$

the result of Proposition 9, and (4.5), we get as $N \rightarrow \infty$

$$\mathbb{E}(m(\lambda) - \mathbb{E}m(\lambda))^2 \leq \frac{A+L-1}{L^2} \sum_{y=1}^{A+L-1} \text{Var} H_L(y) \leq \frac{(A+L-1)^2}{L^2} o(N).$$

Because both L and A linearly depend on N , the last bound and the Markov's inequality imply Claim I.

For Claim II we notice that (4.4) together with Lemma 15 implies that $\sum_{i=1}^L \lambda_i$ can be approximated using the limit shape given by the variational principle. Recalling that we have (without loss of generality) assumed that the origin of the coordinate system is at the bottom-right corner of the embedded trapezoid, we have⁸

$$\frac{1}{L} \sum_{i=1}^L \frac{\lambda_i}{L} = \frac{N(A+L-1)}{L^2} \mathfrak{H}\left(0, \frac{A+L}{N}\right) - \frac{N}{L^2} \sum_{y=1}^{A+L-1} \mathfrak{H}\left(0, \frac{y}{N}\right) + o(1), \quad (4.6)$$

where $\mathfrak{H}$ is the limit shape and $o(1)$ is random asymptotically vanishing term as $N \rightarrow \infty$. The first term in the right-hand side of (4.6) has a straightforward $N \rightarrow \infty$ limit, while the second one is a Riemann sum, approximating the integral of $\mathfrak{H}$ along the right boundary of the embedded trapezoid as $N \rightarrow \infty$. We conclude that (4.6) converges to a deterministic constant as $N \rightarrow \infty$. A very similar computation shows that

$$\frac{1}{L} \sum_{i=1}^L \left(\frac{\lambda_i}{L} \right)^2$$

⁸ In (4.6) we implicitly assumed that the entire segment $(0, 0) - (0, \frac{A+L}{N})$ lies in $\mathfrak{R}$. This segment has to be inside R_N by definition, but $\mathfrak{R}$ might be slightly different from R_N ; this can introduce at most $o(N) = o(L)$ additional terms λ_i to the sum on the left side of (4.6). Since each $\lambda_i < A+L$, the total contribution of these extra terms is at most $o(L) \cdot \frac{1}{L} \cdot \frac{A+L}{L} = o(1)$, meaning that this difference only introduces another $o(1)$ error, which we can ignore.

also has a deterministic limit as $N \rightarrow \infty$, which can be directly expressed through the limit shape $\mathfrak{H}$. We conclude that all terms in the definition of $\sigma(\lambda)^2$ converge as $N \rightarrow \infty$, which is the desired statement of Claim II.

The proof of Theorem 7 is finished.

A uniformity in convergence of local statistics

In this section we explain how Lemma 18 is a consequence of results from [1]. In what follows, we let $\mathcal{B}_R(z) = \{z' \in \mathbb{R}^2 : |z' - z| = R\}$ denote the disk of radius R centered at $z \in \mathbb{R}^2$.

Proof of Lemma 18 As mentioned in Remark 19, Lemma 18 essentially coincides with Theorem 1.5 of [1], except that the uniformity of the constant $C > 1$ in the vertex $v \in R_N$ was not explicitly claimed there. On the other hand, it was stated in [1] that such uniformity holds if R_N is a disk containing v . More specifically, suppose that there exists some vertex $u \in \mathbb{T}$ such that the following holds.

1. R_N approximates a disk $\mathcal{B}_N(u)$ centered at u . That is, we have $R_N = \mathcal{B}_N(u) \cap \mathbb{T}$.
2. We have $\mathcal{B}_{\varepsilon N}(v) \subset R_N$ (that is, R_N contains a disk centered at v).
3. For each $z \in \mathcal{B}_{\varepsilon}(\frac{1}{N}v)$, we have $\nabla \mathcal{H}(z) \in \mathcal{T}_{\varepsilon}$ (that is, $\nabla \mathcal{H}$ is uniformly liquid on that disk).

Then, Theorem 3.15 (see also Assumption 3.5) of [1] states that Lemma 18 holds, with C uniform in v . Let us mention that Theorem 3.15 of [1] does not require R_N to be tileable. If it is not, then H_N is a uniformly random height function on R_N with (some) boundary height function h_N defined on ∂R_N , and $\mathcal{M}$ is the unique *free tiling* associated with H_N , in which tiles are permitted to extend beyond the boundary of R_N .

Although in Lemma 18 the domain R_N does not have to approximate a disk, we may apply the above result on a subdisk of it containing v in the following way. Since $\nabla \mathcal{H}(v) \in \mathcal{T}_{\varepsilon}$, [8, Proposition 4.1] implies that $\nabla \mathcal{H}$ is continuous in a neighborhood of $v = \frac{1}{N}v \in \mathfrak{R}$. Letting $\kappa = \kappa(\varepsilon) > 0$ denote a sufficiently small real number to be fixed later, there then exists $\rho = \rho(\varepsilon, \kappa, \mathfrak{R}) \in (0, \varepsilon)$ such that $B_{\rho}(v) \subset \mathfrak{R} \cap \frac{1}{N}R_N$ and

$$|\nabla \mathcal{H}(z) - \nabla \mathcal{H}(v)| < \kappa, \quad \text{for each } z \in \mathcal{B}_{\rho}(v). \quad (\text{A.1})$$

Let $\tilde{R}_N = \mathcal{B}_{\rho N}(v) \cap \mathbb{T}$ and $\tilde{\mathfrak{R}}_N = \frac{1}{N}\tilde{R}_N$, and condition on the restriction $\mathcal{M}_N|_{R_N \setminus \tilde{R}_N}$ of the random tiling $\mathcal{M}_N$ to $R_N \setminus \tilde{R}_N$. This induces (random) boundary data $\tilde{h}_N : \partial \tilde{\mathfrak{R}}_N \rightarrow \mathbb{R}$ defined by setting $\tilde{h}_N(z) = \frac{1}{N}H_N(Nz)$ for each $z \in \partial \tilde{\mathfrak{R}}_N$. Let $\tilde{\mathcal{H}} \in \text{Adm}(\tilde{\mathfrak{R}}_N; \tilde{h}_N)$ denote the maximizer of $\mathcal{E}$ on $\tilde{\mathfrak{R}}_N$ with boundary data $\tilde{h}_N$. In order to apply Theorem 3.15 of [1] on the domain $\tilde{\mathfrak{R}}_N$ with boundary data $\tilde{h}_N$, we must verify that $\nabla \tilde{\mathcal{H}}$ is likely uniformly liquid around v (for example, $\nabla \tilde{\mathcal{H}}(z) \in \mathcal{T}_{\varepsilon/2}$ for $z \in \mathcal{B}_{\rho/4}(v)$ with high probability).

To that end, observe for any fixed $\delta = \delta(\varepsilon, \omega, D) > 0$ that the variational principle, Lemma 15, yields a constant $C_0 = C_0(\delta, \mathfrak{R}, \mathfrak{h}) > 1$ such that for $N > C_0$ we have

$$\mathbb{P}[\mathcal{E}_N] > 1 - \delta, \quad \text{where} \quad \mathcal{E}_N = \mathcal{E}_N(\delta) = \left\{ \sup_{z \in \partial \tilde{\mathfrak{R}}_N} |\tilde{\mathfrak{h}}_N(z) - \mathcal{H}(z)| < \delta \right\}. \quad (\text{A.2})$$

Next, for sufficiently small $\kappa = \kappa(\varepsilon) > 0$, (A.1) and Proposition 2.13 of [1] together yield a constant $C_1 = C_1(\varepsilon) > 1$ such that

$$\mathbf{1}_{\mathcal{E}} \sup_{z \in \mathcal{B}_{\rho/4}(\mathfrak{v})} |\nabla \tilde{\mathcal{H}}(z) - \nabla \mathcal{H}(z)| < C_1 \mathbf{1}_{\mathcal{E}} \sup_{z \in \partial \tilde{\mathfrak{R}}_N} |\tilde{\mathfrak{h}}_N(z) - \mathcal{H}(z)| < C_1 \delta. \quad (\text{A.3})$$

Hence, if $\delta = \delta(\varepsilon, \omega, D) > 0$ and $\kappa = \kappa(\varepsilon)$ are chosen sufficiently small so that $C_1 \delta < \frac{\varepsilon}{4}$ and $\kappa < \frac{\varepsilon}{4}$, then it follows from (A.1) and (A.3) that $\nabla \tilde{\mathcal{H}}(z) \in \mathcal{T}_{\varepsilon/2}$ for each $z \in \mathcal{B}_{\rho/4}(\mathfrak{v})$ (since $\nabla \mathcal{H}(\mathfrak{v}) \in \mathcal{T}_{\varepsilon}$).

In particular, Theorem 3.15 of [1] applies. Denoting $(\tilde{s}, \tilde{t}) = \nabla \tilde{\mathcal{H}}(\mathfrak{v})$, it yields a constant $C_2 = C_2(\varepsilon, \omega, D, \mathfrak{R}, \mathfrak{h}) > 1$ such that for $N > C_2$ we have

$$\mathbf{1}_{\mathcal{E}} d_{\text{TV}}(\mathcal{M}|_{\mathcal{B}_D(v)}, \mu_{\tilde{s}, \tilde{t}}|_{\mathcal{B}_D(v)}) < \frac{\omega}{2}. \quad (\text{A.4})$$

By (A.3), we have for sufficiently small $\delta = \delta(\varepsilon, \omega, D) > 0$ that

$$\mathbf{1}_{\mathcal{E}} d_{\text{TV}}(\mu_{\tilde{s}, \tilde{t}}|_{\mathcal{B}_D(v)}, \mu_{s, t}|_{\mathcal{B}_D(v)}) < \frac{\omega}{4}. \quad (\text{A.5})$$

Thus, by (A.2), (A.4), (A.5), and further imposing that $\delta < \frac{\omega}{4}$, we have for $N > \max\{C_0, C_2\}$ that

$$\begin{aligned} d_{\text{TV}}(\mathcal{M}|_{\mathcal{B}_D(v)}, \mu_{s, t}|_{\mathcal{B}_D(v)}) &\leq \mathbf{1}_{\mathcal{E}} d_{\text{TV}}(\mathcal{M}|_{\mathcal{B}_D(v)}, \mu_{s, t}|_{\mathcal{B}_D(v)}) + \mathbb{P}[\mathcal{E}^c] \\ &< \mathbf{1}_{\mathcal{E}} d_{\text{TV}}(\mathcal{M}|_{\mathcal{B}_D(v)}, \mu_{\tilde{s}, \tilde{t}}|_{\mathcal{B}_D(v)}) + d_{\text{TV}}(\mu_{\tilde{s}, \tilde{t}}|_{\mathcal{B}_D(v)}, \mu_{s, t}|_{\mathcal{B}_D(v)}) \\ &\quad + \frac{\omega}{4} < \omega, \end{aligned}$$

which implies the lemma. $\square$

Acknowledgements The authors are grateful to David Keating and Ananth Sridhar for the simulation for Fig. 4. We also thank Alexander I. Bufetov for bringing the work [3] to our attention. We thank anonymous referees for helpful comments and suggestions. The work of A.A. was partially supported by a Clay Research Fellowship. The work of V.G. was partially supported by NSF Grants DMS-1664619, DMS-1949820, by BSF grant 2018248, and by the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin–Madison with funding from the Wisconsin Alumni Research Foundation.

References

1. Aggarwal, A.: Universality for lozenge tilings local statistics. Preprint. [arXiv:1907.09991](https://arxiv.org/abs/1907.09991)

2. Aggarwal, A., Huang, J.: Edge statistics for lozenge tilings of polygons, II: Airy line ensemble. Preprint, [arXiv:2108.12874](https://arxiv.org/abs/2108.12874)
3. Bufetov, A.I.: On the Vershik–Kerov conjecture concerning the Shannon–McMillan–Breiman theorem for the Plancherel family of measures on the space of young diagrams. *Geom. Funct. Anal.* **22**(4), 938–975 (2012)
4. Cerf, R., Kenyon, R.: The low-temperature expansion of the Wulff crystal in the 3D Ising model. *Commun. Math. Phys.* **222**(1), 147–179 (2001)
5. Cohn, H., Elkies, N., Propp, J.: Local statistics for random domino tilings of the Aztec diamond. *Duke Math. J.* **85**(1), 117–166 (1996)
6. Cohn, H., Kenyon, R., Propp, J.: A variational principle for domino tilings. *J. Am. Math. Soc.* **14**(2), 297–346 (2001)
7. Cohn, H., Larsen, M., Propp, J.: The shape of a typical boxed plane partition. *New York J. Math.* **4**, 137–165 (1998). [arXiv:math/9801059](https://arxiv.org/abs/math/9801059)
8. De Silva, D., Savin, O.: Minimizers of convex functionals arising in random surfaces. *Duke Math. J.* **151**(3), 487–532 (2010)
9. Dimitrov, E.: Six-vertex models and the GUE-corners process. *Int. Math. Res. Notices* 2020(6), 1794–1881. (2020). <https://arxiv.org/abs/1610.06893>
10. Dimitrov, E., Rychkovsky, M.: GUE corners process in boundary-weighted six-vertex models. [arXiv:2005.06836](https://arxiv.org/abs/2005.06836) (2020)
11. Gorin, V.: From alternating sign matrices to the Gaussian unitary ensemble. *Commun. Math. Phys.* **332**(1), 437–447 (2014). [arXiv:1306.6347](https://arxiv.org/abs/1306.6347)
12. Gorin, V.: *Lectures on Random Lozenge Tilings*. Cambridge University Press, Cambridge (2021)
13. Gorin, V., Panova, G.: Asymptotics of symmetric polynomials with applications to statistical mechanics and representation theory. *Ann. Probab.* **43**(6), 3052–3132 (2015). [arXiv:1301.0634](https://arxiv.org/abs/1301.0634)
14. Johansson, K., Nordenstam, E.: Eigenvalues of GUE minors. *Electron. J. Probab.* **11**(50), 1342–1371 (2006)
15. Keating, D., Sridhar, A.: Random Tilings with the GPU. *J. Math. Phys.* **59**, 091420 (2018). [arXiv:1804.07250](https://arxiv.org/abs/1804.07250)
16. Kenyon, R.: Local statistics of lattice dimers. *Annales Institut Henri Poincaré Probability and Statistics* **33**(5), 591–618 (1997)
17. Kenyon, R.: Lectures on dimers. In *Statistical mechanics*, volume 16 of IAS/Park City Math. Ser., pp. 191–230. Am. Math. Soc., Providence, RI (2009)
18. Kenyon, R., Okounkov, A., Sheffield, S.: Dimers and amoebae. *Ann Math (2)* **163**(3), 1019–1056 (2006)
19. Kuchmov, N.: A variational principle for domino tilings of multiply-connected domains. Preprint, [arXiv:2110.06896](https://arxiv.org/abs/2110.06896)
20. Levin, D.A., Peres, Y., Wilmer, E.L.: *Markov chains and mixing times*. American Mathematical Society, Providence, RI: With a chapter by James G. Propp and David B. Wilson (2009)
21. Mkrtchyan, S., Petrov, L.: GUE corners limit of q -distributed lozenge tilings. *Electron. J. Probab.* **22**, 1–24 (2017)
22. Nordenstam, E.: *Interlaced particles in tilings and random matrices*, PhD thesis, KTH (2009)
23. Novak, J.: Lozenge Tilings and Hurwitz Numbers. *J. Stat. Phys.* **161**, 509–517 (2015)
24. Okounkov, A., Reshetikhin, N.: Correlation function of Schur process with application to local geometry of a random 3-dimensional Young diagram. *J. Am. Math. Soc.* **16**(3), 581–603 (2003). [arXiv:math/0107056](https://arxiv.org/abs/math/0107056)
25. Okounkov, A.Y., Reshetikhin, N.Y.: The birth of a random matrix. *Moscow Math. J.* **6**(3), 553–566 (2006)
26. Olshanski, G., Vershik, A.: Ergodic unitarily invariant measures on the space of infinite Hermitian matrices. In: *Contemporary mathematical physics*, volume 175 of American Mathematical Society Translations: Series 2, pp. 137–175. American Mathematical Society, Providence, RI (1996)
27. Panova, G.: Lozenge tilings with free boundaries. *Lett. Math. Phys.* **105**(11), 1551–1586 (2015)
28. Sheffield, S.: Random surfaces. *Astérisque*, (304):vi+175, (2005)
29. Vershik, A.: Talk at the 1997 conference on formal power series and algebraic combinatorics, Vienna

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.