

Decomposition, trivially-acting symmetries, and topological operators

Daniel G. Robbins,^{1,*} Eric Sharpe^{2,†} and Thomas Vandermeulen^{3,‡}

¹*Department of Physics, University at Albany, Albany, New York 12222, USA*

²*Department of Physics, MC 0435, 850 West Campus Drive, Virginia Tech, Blacksburg, Virginia 24061, USA*

³*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Texas A & M University, College Station, Texas 77843, USA*



(Received 7 December 2022; accepted 5 April 2023; published 25 April 2023)

Trivially-acting symmetries in two-dimensional conformal field theory include twist fields of dimension zero which are local topological operators. We investigate the consequences of regarding these operators as part of the global symmetry of the theory. That is, we regard such a symmetry as a mix of topological defect lines (TDLs) and topological point operators (TPOs). TDLs related by a trivially-acting symmetry can join at a TPO to form nontrivial two-way junctions. Upon gauging, the local operators at those junctions can become vacua in a disjoint union of theories. Examining the behavior of the TPOs under gauging therefore allows us to refine decomposition by tracking the trivially-acting symmetries of each universe. Mixed anomalies between the TDLs and TPOs provide discrete torsionlike phases for the partition functions of these orbifolds, modifying the resulting decomposition. This framework also readily allows for the consideration of trivially-acting noninvertible symmetries.

DOI: [10.1103/PhysRevD.107.085017](https://doi.org/10.1103/PhysRevD.107.085017)

I. INTRODUCTION

A common assumption made in the study of global symmetries is that they act effectively on the states of a field theory. The purpose of this work is to systematically explore the consequences of relaxing this assumption in a modern framework along the lines of [1–4] where symmetries are associated to extended objects, specifically nonlocal topological operators. Fortunately, there already exist a number of results in this direction to build upon. It has been long known that gauging trivially-acting symmetries in conformal field theory (CFT) leads to a violation of cluster decomposition, but in a mild manner; the resulting theories are equivalent to direct sums of local theories, a phenomenon known as decomposition.¹

More formally, decomposition is the observation that in $d > 1$ dimensions a local quantum field theory with a $(d - 1)$ -form symmetry is equivalent to (decomposes into) a disjoint union of other theories, known as universes. Decomposition was first described in 2006 in [5] as part of efforts to understand string compactifications on generalizations of spaces known as stacks, where it resolved some of the apparent physical inconsistencies of those theories. Since that time, it has been applied in a wide variety of areas, including Gromov-Witten theory (see e.g., [6–11]), to give nonperturbative constructions of geometries in gauged linear sigma models (see e.g., [12–23]), to understand IR limits of supersymmetric pure gauge theories (see e.g., [24]), in two-dimensional adjoint QCD (see e.g., [25]), and in [26–29] to understand the Wang-Wen-Witten anomaly resolution of [30]. See also e.g., [31–48] for other applications and tests in theories in dimensions two, three, and four (see [49–53] for some reviews of the subject).

Prototypical examples of decomposition include gauge theories and orbifolds in which a noneffectively-acting higher-form symmetry group is gauged, see e.g., [46,47,54–56]. However, these studies have so far largely focused on the decomposition of local operators. In this paper, we will begin a study of how extended objects behave under decomposition. For concreteness we will focus on two-dimensional theories, as that is the context in which decomposition has already been studied most heavily, so we can most easily relate the story told here to existing results. That said, we expect the picture

*dgrobbins@albany.edu

†ersharpe@vt.edu

‡tvand@tamu.edu

¹For more details on how such theories violate cluster decomposition, see Sec. 2 in Ref. [5]. The fact that violating cluster decomposition in this manner is itself called decomposition may be confusing, but the term is by now standard in the literature.

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.




FIG. 1. The fusion of two grouplike line operators.

proposed here to generalize straightforwardly to the case of gauging a trivially-acting $(d-2)$ -form symmetry in d spacetime dimensions.

Throughout this paper we begin with a two-dimensional theory with an effectively-acting symmetry group G (which we will assume finite and nonanomalous), generated by TDLs² which are labeled by elements of G and fuse according to the group relations in G , as shown in Fig. 1. We then extend the TDLs to a distinguished set describing a group extension Γ ,

$$1 \rightarrow K \rightarrow \Gamma \rightarrow G \rightarrow 1, \quad (1.1)$$

with K acting trivially. By trivially-acting we mean that an element $k \in K$ acts on any local operator $\mathcal{O}$ as

$$k \cdot \mathcal{O} = \mathcal{O} \quad (1.2)$$

for all $k \in K$ and all local operators $\mathcal{O}$. As we are in two dimensions, (1.1) and (1.2) fully characterize the K lines—they act trivially on local operators and fuse with other topological line operators according to the Γ group law. Because trivially-acting symmetries are in this sense “invisible” to the spectrum of the theory, it is common to restrict one’s attention exclusively to effective symmetries. In this work, however, we will attempt to elucidate some of the structure carried by these trivially-acting symmetries and their interplay with a theory’s effective symmetries.³

Note that the group extension (1.1) describes the total zero-form symmetry of the system. In Sec. III we will add local topological operators to the story, which can be associated with one-form symmetries. One might then be tempted to assign a higher group-type algebraic structure to the resulting operators—Appendix B discusses this interpretation. As many readers would be familiar with the concept of a 2-group (the extension of a zero-form symmetry by a one-form symmetry) it may be worth mentioning that the operators we consider here do not form such a structure.

²For a physics-oriented review of symmetries as topological operators in two dimensions, see [1].

³We will also use the term “noneffectively-acting” symmetry to mean a symmetry such as Γ for which only a subgroup acts trivially.

Each TDL in the resulting theory⁴ is labeled by an element of Γ , whose underlying set is that of $K \times G$. When this extension is trivial, Γ is in fact equal to $K \times G$, and each symmetry element is some (noninteracting) mix of the effective and trivial symmetries. Possible nontrivial extensions describe how these symmetries can mix together, in that we could fuse two G lines and end up with a K line. Note that non-trivial extensions will in general prevent G from being a subgroup of Γ , which means that the theory under consideration may no longer have a standalone G symmetry.

In ungauged theories, if a TDL describes a noneffectively-acting group, then the lines for the trivially-acting subgroup can end on TPOs (which are bound to the lines and do not define standalone one-form symmetries). After gauging, those TPOs become unbound point operators generating one-form symmetries. Tracking such point operators determines TDL structures, which enables one to follow TDLs and TPOs through gauging. In this fashion, we refine decomposition by explicitly tracking such extended objects.

After a review of the requisite basic notions in Sec. II, Sec. III introduces the notion that trivially-acting symmetries are controlled by a mix of topological line and point operators. This perspective allows us to consider the effect of a mixed anomaly when gauging the line operators. Alternatively, in the absence of such anomalies, we could simultaneously gauge the lines and points. Equipped with this technology, Sec. IV runs through a handful of examples, the final of which incorporates noninvertible symmetries in the form of a trivially-acting categorical symmetry. We will see that this framework allows us to track the fate of trivially-acting symmetries under decomposition, with the upshot being that they divide themselves among the various universes.

II. BACKGROUND

This section reviews the various concepts that will enter into our full treatment of trivially-acting symmetries. The material presented here is not new, so readers who are familiar with these subjects may wish to skim the section for notation or simply skip ahead to Sec. III.

A. Higher-form symmetries

We have presented a notion of symmetries as actions on local quantities generated by topological operators of codimension one. In field theory, however, we often consider extended objects in addition to local ones. It would be natural to consider an analog of symmetries for these objects—transformations between extended objects. In fact, analogously to ordinary symmetries, transformations of

⁴The underlying philosophy here is that a “theory” is defined not only by local data, but also by its nonlocal spectrum, as in [2].

n -dimensional objects are naturally described by operators of codimension $n + 1$. Such transformations are termed n -form symmetries.

One realization of higher-form symmetries involves “groups” in which properties such as associativity are weakened. Higher-form symmetries have a long history in physics, e.g., [57–61], in discussions of the String group (see e.g., [62–64]), in the Yetter model (see e.g., [65,66]), and in lattice gauge theories (see e.g., [67,68]), to name a few examples. Additional categorical generalizations of (orbifold) groups, via an application of defects to generalize the ordinary orbifold construction, are discussed in [3,4,69–72]. See also [73] for related ideas and applications of discrete gauge theories and group cohomology in condensed matter physics.

In this paper, we will focus on two dimensions. There, ordinary symmetries are controlled by TDLs. There exists one type of higher-form symmetry; one-form symmetries, controlled by topological point operators (TPOs). These describe how line operators transform into each other. Already in two dimensions we have a fairly rich structure—the generators of the two types of symmetries naturally act on each other. This interplay, as we will see, will be key to fully describing trivially-acting symmetries.

B. Noninvertible symmetries

Broadly speaking, topological defect lines do not obey group laws, but rather more general fusion relations. For instance, there is a noninvertible operator that implements Kramers-Wannier duality in the critical Ising model [74]. Additionally, and more relevant to this paper, in order to fully capture the quantum dual to a non-Abelian symmetry, one is forced to examine nongrouplike symmetries [75].

The fusion of general TDLs is described mathematically by unitary fusion categories; see [75,76]. In such a category, we assume that there exists a set of “simple” objects, and all other objects in the category are expressible as non-negative integral linear combinations of these simple objects. There are a set of fusion rules which govern how the simple objects (and hence the composite ones) combine, which along with addition gives the structure of a ring. There is a map $X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$ known as the associator, and a number of consistency conditions which will not be important for our purposes.

A group G defines a special case of this structure; there is a simple line for each element $g \in G$, and the fusion rules are simply the group relations. To describe anomalies, one can add a nontrivial associator, given by a representative of an element of $H^3(G, U(1))$. Less trivial examples include quantum symmetries of orbifolds. When we gauge a symmetry given by a group G , the gauged theory has a quantum $\text{Rep}(G)$ symmetry where simple lines can be labeled by irreducible representations of G . Since the

$\otimes$	1	X	Y
1	1	X	Y
X	X	1	Y
Y	Y	Y	$1 + X + Y$

FIG. 2. The fusion products for $\text{Rep}(S_3)$.

irreducible representations of a non-Abelian group can have dimension greater than one, irreducible representations in general do not form groups under tensor products, but there does exist a fusion law for products of irreducible representations.

An example which we will use repeatedly is the symmetric group S_3 . It has three irreducible representations; the trivial representation, a nontrivial representation of dimension one, and a single irreducible representation of dimension two. Labeling these irreducible representations respectively as 1, X and Y , their fusion products are as shown in Fig. 2. In particular, the simple line Y corresponding to the two-dimensional irrep has the fusion relation $Y \otimes Y = 1 + X + Y$ —in the grouplike case, fusion of simple lines never produces a sum. Correspondingly, one can see that there is no object Y^{-1} satisfying $Y \otimes Y^{-1} = 1$, hence the term “noninvertible”.

C. Gauging symmetries

We will use a notion of gauging which is sufficiently general to include the categorical symmetries described above. In order to gauge a general symmetry, we select an object A , known as the algebra object, along with suitable multiplication and identity morphisms to define a symmetric Frobenius algebra [74,75].⁵

Intuitively A serves as the identity operator for the gauged theory. We calculate quantities in the gauged theory from ungauged objects by inserting a “sufficiently fine mesh” of A lines, along with appropriate normalization. As an example, when gauging a grouplike symmetry, we can take A to be the sum of lines labeled by elements of the group, i.e.,

$$A = \sum_{g \in G} L_g, \quad (2.1)$$

where L_g denotes a TDL labeled by g . In order to calculate e.g., the torus partition function in the gauged theory, we wrap both cycles with an A line and normalize by the dimension of A , which will simply be the order of the group. Breaking the two A lines into sums over group elements, this reduces to the familiar prescription for the orbifold torus partition function,

⁵In the condensed matter literature, this generalized form of gauging is known as anyon condensation [77].

$$\frac{1}{|A|} Z_{A,A} = \frac{1}{|G|} \sum_{\substack{g_1, g_2 \in G \\ [g_1, g_2] = 1}} Z_{g_1, g_2}. \quad (2.2)$$

The advantage to such a general formalism is that we can also gauge nongrouplike symmetries. For example, returning to $\text{Rep}(S_3)$, there exist suitable choices of multiplication and unit morphisms such that $1 + X$, $1 + Y$ and $1 + X + 2Y$ constitute algebra objects. Taking $A = 1 + X$ gives us a grouplike $\mathbb{Z}_2$ gauging, while $1 + Y$ and $1 + X + 2Y$ lead to nongrouplike gaugings (at least superficially—these two gaugings are “dual” to gauging $\mathbb{Z}_3 \subset S_3$ and all of S_3 , respectively).

The algebra object also has a role to play in determining the local operators that appear in the gauged theory—in particular we will be interested in operators of weight zero. Acting with A on TPOs will have the effect of “projecting out” linear combinations that do not survive as bulk local operators, such that the result is a linear combination of TPOs which are freestanding in the gauged theory.

D. Decomposition

In this section we will review decomposition in two-dimensional orbifold CFTs. Decomposition for two-dimensional orbifolds without discrete torsion can be summarized as follows [5,26]. Let T be a theory with an action of Γ , where $K \subset \Gamma$ acts trivially and $G = \Gamma/K$. Assume that there is no discrete torsion in the Γ orbifold. (Decomposition for two-dimensional orbifolds including discrete torsion is described in [26], with mixed anomalies in [28], and other miscellaneous two-dimensional orbifold cases are in [27]). Then,

$$\text{QFT}([T/\Gamma]) = \text{QFT}\left(\left[\frac{T \times \hat{K}}{G}\right]_{\hat{\omega}}\right), \quad (2.3)$$

where in general the effectively-acting coset G acts on both T and $\hat{K}$, and $\hat{\omega}$ denotes discrete torsion appearing in the universes, as explained in [5,26]. The right-hand side is a disjoint union of theories, as many as orbits of G on $\hat{K}$. The elements of that disjoint union are known as the universes of the decomposition.

For later use, it will be helpful to give explicitly the projectors⁶ onto the universes of decomposition, referring specifically to decomposition in two-dimensional orbifolds without discrete torsion, as described in (2.3). These are essentially a consequence of Wedderburn’s theorem, and are given explicitly in Sec. 2.2.2 in Ref. [44] in terms of TPOs. For Γ orbifolds without discrete torsion, we give

⁶The state-operator correspondence maps the vacuum state in each universe to an operator which projects onto that universe. Due to this mapping, we will be somewhat abusive with notation and use Π to refer to both the vacuum state and the projection operator.

them as follows. Let R be a representation of K associated to a universe, meaning that

$$R = \bigoplus_i R_i \quad (2.4)$$

for $R_i \in \hat{K}$ the irreducible representations forming an orbit of G . Then,

$$\Pi_R = \sum_i \Pi_{R_i}, \quad (2.5)$$

where

$$\Pi_{R_i} = \frac{\dim R_i}{|K|} \sum_{k \in K} \chi_{R_i}(k^{-1}) \sigma_k, \quad (2.6)$$

and where χ_{R_i} denotes a character of R_i and the σ_k are ungauged local operators which fuse according to the group law in K .

To make this paper self-contained, it may be helpful to briefly illustrate decomposition in some simple examples. Let T denote a theory, and K an Abelian group with a trivial action on T . In terms of the prediction (2.3), $G = 1$ so $\Gamma = K$ and decomposition predicts in this case that

$$\text{QFT}([T/K]) = \text{QFT}(T \times \hat{K}), \quad (2.7)$$

as many copies of T as irreducible representations of K , namely $|K|$ since K is Abelian. We can check this by computing partition functions. In calculating the partition function for the orbifold $[T/K]$, each sector Z_{k_1, k_2} is simply going to be equivalent to the parent theory partition function. Therefore, we have

$$Z_{[T/K]} = \frac{1}{|K|} \sum_{k_1, k_2 \in K} Z_{k_1, k_2} = \frac{1}{|K|} \sum_{k_1, k_2 \in K} Z_T = |K| Z_T, \quad (2.8)$$

which confirms the statement of decomposition in this case.

As with orbifolds by effective symmetries, there is a quantum symmetry in this orbifold which we can gauge to return the original theory, given by $\hat{K}$. In this context we can interpret $\hat{K}$ as a subgroup⁷ of the exchange symmetry $S_{|K|}$ acting on the $|K|$ copies appearing in the decomposition.

If K is non-Abelian, a similar story applies, and the orbifold $[T/K]$ is equivalent to as many copies of T as irreducible representations of K . As before, we can check using partition functions. Partition functions on T^2 are computed by summing over commuting pairs, which gives

$$Z_{[T/K]} = \frac{1}{|K|} \sum_{\substack{k_1, k_2 \in K \\ [k_1, k_2] = 1}} Z_{k_1, k_2} = N_K Z_T, \quad (2.9)$$

⁷By virtue of Cayley’s theorem.

where N_K is the number of conjugacy classes in K . There is once again a quantum symmetry we can gauge to return to the original theory, except it is now given by the fusion category $\text{Rep}(K)$ described by the fusion structure of the irreducible representations of K , which will fail to be grouplike due to the presence of irreducible representations with dimension greater than one.

As an example, consider orbifolding a theory by the symmetric group S_3 , acting trivially (so that $G = 1$ and $\Gamma = K = S_3$). This group has three conjugacy classes, hence three irreducible representations, labeled 1, X , Y earlier, so from (2.9) and (2.3) (and the fact that the effectively-acting coset is $G = 1$) we expect three universes (i.e., a decomposition into a direct sum of three theories).

We expect there to be a quantum $\text{Rep}(S_3)$ symmetry acting on the vacua, the fusion products for which were presented in Fig. 2. We can see this as follows: First, label the projectors into these universes as Π_1 , Π_X , Π_Y . The quantum symmetry defines a product of the form

$$R \cdot \sigma_k = \chi_R(k) \sigma_k, \quad (2.10)$$

for a representation R and $k \in S_3$ (where σ_k corresponds to an ungauged local operator), which can be applied to compute

$$1\Pi_a = \Pi_a \quad \text{for all } a, \quad (2.11)$$

$$X\Pi_1 = \Pi_X, \quad X\Pi_X = \Pi_1, \quad (2.12)$$

$$X\Pi_Y = \Pi_Y, \quad (2.13)$$

$$Y\Pi_1 = \frac{1}{2}\Pi_Y = Y\Pi_X, \quad (2.14)$$

$$Y\Pi_Y = 2\Pi_1 + 2\Pi_X + \Pi_Y. \quad (2.15)$$

We will give explicit expressions for the projectors and other details later in Sec. IV D.

Here we see a qualitative difference between invertible and noninvertible symmetries. The $\mathbb{Z}_2$ subsymmetry generated by X exchanges two of the three copies, acting on the sum of the universes as $X(\Pi_1 + \Pi_X + \Pi_Y) = (\Pi_1 + \Pi_X + \Pi_Y)$. The weight two line Y , however, has nonintegral action on the individual universes, and acts on their sum as $Y(\Pi_1 + \Pi_X + \Pi_Y) = 2(\Pi_1 + \Pi_X + \Pi_Y)$.

We have illustrated decomposition in a handful of simple examples to make this paper self-contained. Many more examples in two-dimensional orbifolds are discussed in [5,26–29,48].

III. TRIVIALY-ACTING SYMMETRIES

We begin the analysis of extended operators and trivially-acting symmetries by looking at junctions between two TDLs, focusing on the topological point operators which




FIG. 3. Two effective symmetry lines joined at a topological junction.




FIG. 4. Two line operators joined by a point operator.

live there. In the effective, grouplike case, this relatively uninteresting configuration is pictured in Fig. 3. Because junctions between lines are constrained by the group action, the only weight zero operator that can sit at a junction between a g_1 and g_2 line is the identity, and only when $g_1 = g_2$. Otherwise such a junction can only exist if it contains a higher-weight operator.⁸

In the presence of trivially-acting symmetries, such junctions can support nontrivial-weight zero operators. More precisely, let the total zero-form symmetry of our theory be $\Gamma = K.G$,⁹ where the normal subgroup K acts trivially. Elements of Γ that differ by an element of K can be joined at topological junctions, as shown in Fig. 4. Similarly, a line labeled by an element of the trivially-acting subgroup K can end at a corresponding point operator (which is to say it can have a two-way junction with the identity TDL). While the information is redundant with the labelings of the lines, we will find it convenient to label such point operators by elements of K .

In the same way that the TDLs of our theory satisfy a fusion algebra (the group laws in the grouplike case or a unitary fusion category in the more general case), the TPOs associated with these trivial symmetries can be given a fusion algebra structure. Figure 5 illustrates this—the TPOs there labeled by σ_{k_1} and σ_{k_2} can be brought together, and we can use the operator product expansion (OPE) to fuse them into a single-point operator joining the γ and $k_2 k_1 \gamma$ lines. As we might have expected, the σ_{k_1} and σ_{k_2} operators fuse to the $\sigma_{k_2 k_1}$ operator, and in general the fusion rules for these TPOs are simply the group law in K .

⁸To expand on this point, one often considers twist fields on which TDLs can end. In the effective case, these are local operators, but are not topological. Our main concern in this work is to point out that when the associated symmetry acts trivially, there will be twist fields of weight zero, which are therefore local topological operators, which one can regard as being part of the global symmetry of the theory.

⁹Here we are using the notation of [78] where a dot indicates a general group extension extension. That is, K is a normal subgroup of Γ and $\Gamma/K \simeq G$, but K need not be central and the extension need not be split.





FIG. 5. The fusion of two point operators.

FIG. 6. The action of K on local operators is equivalent to a trivial action.

We can see how such a configuration of point and line operators describes a trivially-acting symmetry by looking at the action on local operators, as in Fig. 6. Figure 6(a) depicts a line labeled by k acting on a local operator. In Fig. 6(b) we insert an identity operator, which in Fig. 6(c) we split into a k and k^{-1} pair,¹⁰ between which runs an identity line. In Fig. 6(d) we bring these operators together from the other side, and finally in Fig. 6(e) we re-fuse them into an identity operator. The remaining identity line and point operators can be removed, and the net result of this process is that $k \cdot \mathcal{O} = \mathcal{O}$, i.e., K acts trivially.

In passing, the computations of this section are formally similar to computations in Sec. IV (Ref. [79]), although we believe our use differs.

¹⁰One might object that in moving from Fig. 6(a) to Fig. 6(e) there is the potential to pick up a k -dependent phase. Such a phase should only arise from a gauge anomaly in K [1], and we are assuming here that there are no such anomalies in our trivially-acting symmetries.


A. Mixed anomalies

As we have discussed, line operators naturally act on local operators. In this section we focus on the action of our TDLs (labeled by Γ) on our TPOs (labeled by K). We will write these TPOs as σ_k . There is a natural action of Γ on these objects; conjugation. However, there exists the possibility of adding nontrivial phases into this action. We can make an ansatz that the general form of Γ acting on σ_k should be

$$\gamma \cdot \sigma_k = \sigma_{\gamma k \gamma^{-1}} B(\gamma, \gamma k \gamma^{-1}) \quad (3.1)$$

where $B(\gamma, k) \in U(1)$ is known as a mixed anomaly between the point and line operators.

Compatibility between fusion of TDLs and TPOs will constrain these phases. More specifically, Fig. 7 illustrates these consistency conditions for the K argument. In Fig. 7(a) we begin with two TPOs labeled by k_1 and k_2 , circled by a TDL labeled by γ . We can shrink the TDL and pinch it off into a circle surrounding each TPO, as in Fig. 7(b). Each of these circles can be further shrunk to give the action of γ on each point, leading to Fig. 7(c). From (3.1), Figs. 7(b) and 7(c) should be related to each other by a factor of

FIG. 7. The fusion of two TPOs acted on by a TDL constrains the dependence of B on its K argument.




FIG. 8. The fusion of two TDLs acting on a TPO constrains the dependence of B on its Γ argument.

$B(\gamma, \gamma k_1 \gamma^{-1}) B(\gamma, \gamma k_2 \gamma^{-1})$. Finally these two TPOs can fuse into one, shown in Fig. 7(e). Alternatively, we could have proceeded from Fig. 7(a) to Fig. 7(d) by fusing the TPOs before shrinking the TDL. This route takes us from Fig. 7(a) to Fig. 7(d) and finally to Fig. 7(e), while picking up the phase $B(\gamma, \gamma k_1 k_2 \gamma^{-1})$ along the way. These two paths to the same result should produce the same phase, so we find the condition

$$B(\gamma, \gamma k_1 k_2 \gamma^{-1}) = B(\gamma, \gamma k_1 \gamma^{-1}) B(\gamma, \gamma k_2 \gamma^{-1}), \quad (3.2)$$

which tells us that B is a homomorphism in its K argument.

We can constrain B 's behavior in Γ similarly, by examining two TDLs γ_1 and γ_2 acting on a single TPO σ_k , shown in Fig. 8(a). Shrinking the γ_2 line to act on σ_k brings us to Fig. 8(b) and introduces a phase $B(\gamma_2, \gamma_2 k \gamma_2^{-1})$. Then we could act on the resulting TPO with the remaining γ_1 line to arrive at Fig. 8(d), and gain the phase $B(\gamma_1, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1})$. Alternatively, in Fig. 8(a) we could have fused the TDLs to arrive at Fig. 8(c). Then, acting on σ_k with the $\gamma_1 \gamma_2$ TDL would bring us to Fig. 8(d) while introducing a phase $B(\gamma_1 \gamma_2, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1})$. Again, the requirement that these two ways of reaching Fig. 8(d) produce the same phase leads to the condition

$$B(\gamma_1 \gamma_2, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1}) = B(\gamma_1, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1}) B(\gamma_2, \gamma_2 k \gamma_2^{-1}). \quad (3.3)$$

Noting that an action of Γ on K taking $k \rightarrow \gamma k \gamma^{-1}$ induces an action $(\gamma^{-1} \cdot \varphi)(k) = \varphi(\gamma k \gamma^{-1})$ on $\varphi: K \rightarrow U(1)$, we can

act on both sides with $\gamma_1 \gamma_2$ to write the above as

$$B(\gamma_1 \gamma_2, k) = B(\gamma_1, k) (\gamma_1 \cdot B)(\gamma_2, k), \quad (3.4)$$

which is the condition for B to be a crossed homomorphism in Γ and $H^1(K, U(1))$. Considering the properties (3.2) and (3.4) of B , we can regard mixed symmetries as cocycles valued in $Z^1(\Gamma, H^1(K, U(1)))$. The following section will confirm that physical effects of these anomalies are only sensitive to the class of this cocycle in $H^1(\Gamma, H^1(K, U(1)))$ (in general with nontrivial action on the coefficients), agreeing with the results of [38,80].

Here we have examined mixed anomaly phases for freestanding TPOs. Since many of the TPOs we will work with are bound to lines and cannot stand freely, for completeness appendix A repeats these consistency arguments for bound TPOs. The results match those of the freestanding case.

B. Application to orbifolds

We would like to take the orbifold by the zero-form symmetry $\Gamma = K.G$, where as before K acts trivially (for simplicity we do not turn on any discrete torsion in Γ). We know to expect this orbifold to exhibit decomposition, which is to say that the result should be a direct sum of orbifolds by G or its subgroups. Each nonvanishing contribution Z_{γ_1, γ_2} (where $[\gamma_1, \gamma_2] = 1$) to the torus partition function should contribute a term Z_{g_1, g_2} to the decomposed partition function, where $g_i = \pi(\gamma_i)$ and $\pi: \Gamma \rightarrow G$ is the projection. In the absence of any phases, we could determine the coefficient of Z_{g_1, g_2} by simply counting how many commuting pairs γ_1, γ_2 exist in Γ .

However, it may be that two sectors Z_{γ_1, γ_2} and $Z_{\gamma'_1, \gamma'_2}$, for

$$\pi(\gamma_1) = g_1 = \pi(\gamma'_1), \quad \pi(\gamma_2) = g_2 = \pi(\gamma'_2), \quad (3.5)$$

which both “project” to Z_{g_1, g_2} in the G orbifold, are related by phases. The framework developed so far in this section will allow us to determine the contribution of such phases to the decomposed partition function. In order to determine the coefficient of Z_{g_1, g_2} , the strategy will be to fix a commuting pair γ_1, γ_2 of elements in Γ , and use the action of the TPOs labeled by K to map each $Z_{\gamma'_1, \gamma'_2}$ to Z_{γ_1, γ_2} (possibly accruing phases along the way). Note that since elements of Γ which project to the same element in G differ by an element of K , we can set $\gamma'_1 = k_1 \gamma_1$ and $\gamma'_2 = k_2 \gamma_2$. Once all of the contributions take this form, the sum of their coefficients should be the coefficient of Z_{g_1, g_2} in the decomposed theory. Naturally one might worry that the results of this procedure would depend on the choice of γ_1 and γ_2 —we will return to this concern shortly.

Figure 9 illustrates this procedure. We begin in Fig. 9(a) by representing $Z_{k_1 \gamma_1, k_2 \gamma_2}$ as two TDLs labeled by $k_1 \gamma_1$ and $k_2 \gamma_2$ wrapping the cycles of a torus, pictured as its




FIG. 9. We map the $k_1\gamma_1, k_2\gamma_2$ sector of the orbifold partition function into the $k_1\gamma_1, \gamma_2$ sector.

fundamental domain in $\mathbb{C}$. In Fig. 9(b) we insert an identity TPO on the $k_2\gamma_2$ line. Moving to Fig. 9(c), we break the identity operator into an inverse pair σ_{k_2} and $\sigma_{k_2^{-1}}$. This is chosen such that the line connecting them is labeled by γ_2 alone. Moving to Fig. 9(d), we drag σ_{k_2} across the $k_1\gamma_1$ line. This will effectively act on the TPO to produce $\sigma_{k_1\gamma_1 k_2\gamma_2^{-1} k_1^{-1}}$, with the incoming line labeled by $k_1\gamma_1 k_2^{-1} \gamma_1^{-1} k_1^{-1} k_2\gamma_2$. Moving from Fig. 9(c) to Fig. 9(d) will introduce the phase $B(k_1\gamma_1, k_1\gamma_1 k_2\gamma_2^{-1} k_1^{-1}) = (g_1^{-1} \cdot B)(k_1\gamma_1, k_2)$ due to a potential mixed anomaly.

In anticipation of the upcoming steps, we split $\sigma_{k_1\gamma_1 k_2\gamma_2^{-1} k_1^{-1}}$ into $\sigma_{[k_2^{-1}, k_1\gamma_1]}$ and σ_{k_2} . The result of this is shown in Fig. 9(e), where the TPO $\sigma_{[k_2^{-1}, k_1\gamma_1]}$ has been moved to the intersection. This allows the horizontal lines on both sides of the intersection to be labeled by γ_2 , and we have effectively moved σ_{k_2} across the junction. In Fig. 9(f) we approach $\sigma_{k_2^{-1}}$ with its inverse from the other side, joining them once again into an identity operator. This leaves us with $Z_{k_1\gamma_1, \gamma_2}$ with the point operator $\sigma_{[k_2^{-1}, k_1\gamma_1]}$ sitting at the junction. The reason for this extra insertion is that, while by assumption $[\gamma_1, \gamma_2] = [k_1\gamma_1, k_2\gamma_2] = 1$, it is not necessarily true that $[k_1\gamma_1, \gamma_2] = 1$.

This is not an issue, as the next move would be to repeat the process and map $Z_{k_1\gamma_1, \gamma_2}$ to Z_{γ_1, γ_2} . The steps are essentially the same, so we will not illustrate this part separately. The important fact is that in doing so, we would generate the phase $B^{-1}(\gamma_2, \gamma_2 k_1 \gamma_2^{-1}) = [(g_2^{-1} \cdot B)(\gamma_2, k_1)]^{-1}$. The inverse appears because we must either move σ_{k_1} across γ_2 with orientation opposite to $k_1\gamma_1$ in the previous steps, or equivalently drag $\sigma_{k_1^{-1}}$ across with the same orientation. In moving σ_{k_1} across γ_2 , we generate a point operator which is inverse to $\sigma_{[k_2^{-1}, k_1\gamma_1]}$, such that the resulting diagram for Z_{γ_1, γ_2} contains no more nonidentity TPOs (as is consistent with the group law, since γ_1 and γ_2 commute).

Of course, the order in which we perform this reduction— $Z_{k_1\gamma_1, k_2\gamma_2} \rightarrow Z_{k_1\gamma_1, \gamma_2} \rightarrow Z_{\gamma_1, \gamma_2}$ versus $Z_{k_1\gamma_1, k_2\gamma_2} \rightarrow Z_{\gamma_1, k_2\gamma_2} \rightarrow Z_{\gamma_1, \gamma_2}$ —should not matter.¹¹ The order would seem to affect the Γ argument of B . In order to guarantee that these two paths to Z_{γ_1, γ_2} are equivalent, we take B to be a pullback from an element of $H^1(G, H^1(K, U(1)))$. Heuristically, this means that we are free to modify how the effective part of the symmetry acts on the TPOs, but not the noneffective part.

In summary, we have used manipulations with TPOs to derive a relationship between different contributions to a Γ

¹¹More specifically, because all of these steps are invertible, we could take one path and then the inverse of the other to map $Z_{k_1\gamma_1, k_2\gamma_2}$ to itself. Because the phase we assign to a given sector should be unambiguous, the total phase accrued this way should be unity. Therefore, the two ways of reaching Z_{γ_1, γ_2} should produce the same phase.

orbifold partition function with trivially-acting normal subgroup K ,¹²

$$Z_{k_1\gamma_1, k_2\gamma_2} = \frac{(g_1^{-1} \cdot B)(g_1, k_2)}{(g_2^{-1} \cdot B)(g_2, k_1)} Z_{\gamma_1, \gamma_2}, \quad (3.6)$$

where $g_i = \pi(\gamma_i)$, with π the projection map $\Gamma \rightarrow G$ appearing in the extension (1.1). In [27–29], where these mixed anomaly phases were studied under the name ‘quantum symmetry phases’, a special version of this relation was taken as an ansatz. Here we see how it follows naturally from the topological operator formulation of trivial symmetries. In the following section we check that these phases produce reasonable results for the decomposed orbifold theory.

C. Consistency checks

Once each sector with fixed $g_1 = \pi(\gamma_1), g_2 = \pi(\gamma_2)$ is mapped to a phase times Z_{γ_1, γ_2} , we expect that the coefficient of Z_{g_1, g_2} appearing in the decomposed partition function should be given by the sum of those phases. Explicitly, this is

$$F(\gamma_1, \gamma_2) = \sum_{\substack{k_1, k_2 \in K \\ [k_1\gamma_1, k_2\gamma_2] = 1}} \frac{(g_1^{-1} \cdot B)(g_1, k_2)}{(g_2^{-1} \cdot B)(g_2, k_1)} \quad (3.7)$$

for a commuting pair γ_1, γ_2 .

1. Reference (in)dependence

Now we return to the question of dependence on γ_1, γ_2 . In some cases, there will be a canonical choice of γ_1, γ_2 . Consider a split extension, i.e., the extension class is trivial. Then there is always a choice of section such that $[(1, g_1), (1, g_2)] = 1$. In such a case, the relative phases that arise from B when mapping each sector in the Γ orbifold to its corresponding $Z_{(1, g_1), (1, g_2)}$ are in fact redundant with phases arising from a choice of discrete torsion in the Γ orbifold.¹³ There will be other cases, however, where there is no obvious choice of reference, and we must be more careful in treating the decomposition process.

In order to see how coefficients in the decomposed partition function depend on γ_1, γ_2 , suppose we had picked a different commuting pair, $\tilde{\gamma}_1, \tilde{\gamma}_2$ (with the same G

¹²It may be helpful to note that, for $\Gamma = K.G$, the action of G on $\varphi \in H^1(K, U(1))$ can be written as $(g \cdot \varphi)(k) = \varphi(\gamma^{-1} k \gamma)$ for any γ satisfying $\pi(\gamma) = g$. In particular, this makes it clear that such an action does not depend on any choice of section for Γ .

¹³The precise condition for mixed anomaly phases to be redundant with discrete torsion is that the cup product of $B \in H^1(G, H^1(K, U(1)))$ and the extension class $c \in H^2(G, K)$ must be trivial in $H^3(G, U(1))$ (see Sec. 2.3 in Ref. [28]). This is manifestly satisfied for split extensions.

projection). Then we would have found the coefficient of Z_{g_1, g_2} to be

$$F(\tilde{\gamma}_1, \tilde{\gamma}_2) = \sum_{\substack{\tilde{k}_1, \tilde{k}_2 \in K \\ [\tilde{k}_1 \tilde{\gamma}_1, \tilde{k}_2 \tilde{\gamma}_2] = 1}} \frac{(g_1^{-1} \cdot B)(g_1, \tilde{k}_2)}{(g_2^{-1} \cdot B)(g_2, \tilde{k}_1)}. \quad (3.8)$$

Since $\pi(\gamma_1) = \pi(\tilde{\gamma}_1) = g_1$ and $\pi(\gamma_2) = \pi(\tilde{\gamma}_2) = g_2$, we can find κ_1 and κ_2 such that $\tilde{\gamma}_1 = \kappa_1 \gamma_1$ and $\tilde{\gamma}_2 = \kappa_2 \gamma_2$. By redefining our summation variable to $k_i = \tilde{k}_i \kappa_i$ in (3.8), we see that these two expressions relate to each other as

$$F(\gamma_1, \gamma_2) = F(\tilde{\gamma}_1, \tilde{\gamma}_2) \frac{(g_2^{-1} \cdot B)(g_2, \kappa_1)}{(g_1^{-1} \cdot B)(g_1, \kappa_2)}, \quad (3.9)$$

which is to say that choosing a different “reference” sector in the Γ orbifold would change the computed coefficient for Z_{g_1, g_2} by a phase—for consistency—such a choice should not affect the result.

There are two ways in which $F(\gamma_1, \gamma_2)$ could depend only on g_1 and g_2 . It could be that the summand of (3.7) is identically trivial, in which case the value of F counts the number of commuting pairs in Γ with fixed G projection; this number is clearly independent of any choice of reference. Alternatively, if B provides any nontrivial phases to the sum, it should be that those phases sum to zero. From (3.9) we see that because changing our reference sector multiplies the entire sum by a phase, $F(\gamma_1, \gamma_2) = 0$ would also be a reference-independent statement.

Consider the case where K is central in Γ . Then the condition $[k_1 \gamma_1, k_2 \gamma_2] = 1$ is equivalent to $[\gamma_1, \gamma_2] = 1$, which γ_1 and γ_2 were specifically chosen to satisfy. This means that the sum in (3.7) is unconstrained and runs over two full copies of K . If we fix an element k_1 , the sum over k_2 is a sum over a homomorphism from K into $U(1)$, so either the homomorphism is trivial or the sum vanishes. The same is true for any value of k_1 , which means that for central K we would calculate the same coefficient no matter which sector was chosen as the reference point.

Unfortunately, such reference independence may not be present for noncentral K . Consider, for example, $\Gamma = S_3 \times \mathbb{Z}_2$ with $K = S_3$. We see from

$$H^1(\mathbb{Z}_2, H^1(S_3, U(1))) = H^1(\mathbb{Z}_2, \mathbb{Z}_2) = \mathbb{Z}_2 \quad (3.10)$$

that there is the possibility of a nontrivial mixed anomaly in this example. With this mixed anomaly present, one can apply decomposition to express the Γ orbifold sectors in terms of $\mathbb{Z}_2$ orbifold sectors as

$$\frac{1}{12} [18Z_{0,0} + 6(a_1 Z_{0,1} + a_2 Z_{1,0} + a_3 Z_{1,1})], \quad (3.11)$$

where choice of reference sector allows for any choice of the coefficients a_i satisfying $a_1^2 = a_2^2 = a_3^2 = 1$. As discussed

above, because this is a split extension, we have the canonical choice of using $Z_{(1, g_1), (1, g_2)}$ as the reference point in each sector. Doing so, we get $a_1 = a_2 = a_3 = 1$, which indeed is the only choice that corresponds to a sensible CFT partition function—the decomposed orbifold is then equivalent to one copy of the parent theory plus one copy of the orbifold by the effectively-acting $\mathbb{Z}_2$. This choice of reference is also the one which for which the mixed anomaly phases are redundant with discrete torsion in Γ .

In more general situations where the extension is neither central nor split, we do not have a general proof of consistency of this procedure, so we would need to verify case by case that choice of reference sector does not lead to any issues. In the examples presented in the following section, only the one examined in Sec. IV C is neither split nor central. However, in calculating F for that example, k_1 and k_2 will always fill out a subgroup of K^2 ; this means that any nontrivial contributions to the sum will cause it to vanish, as in the central case.

2. Coboundary invariance

Next, we can check that B that are exact in $H^1(G, H^1(K, U(1)))$ do not contribute. Such a cocycle takes the form

$$B(g, k) = \frac{(g \cdot \varphi)(k)}{\varphi(k)} \quad (3.12)$$

for $\varphi \in H^1(K, U(1))$. With such a choice of B , the relative phase (3.6) between $Z_{k_1 \gamma_1, k_2 \gamma_2}$ and Z_{γ_1, γ_2} becomes

$$\frac{\varphi(k_2) \varphi(\gamma_2 k_1 \gamma_2^{-1})}{\varphi(k_1) \varphi(\gamma_1 k_2 \gamma_1^{-1})}. \quad (3.13)$$

Recall that the values of k are such that $\gamma'_1 = k_1 \gamma_1$ and $\gamma'_2 = k_2 \gamma_2$ is a commuting pair. Expressing (3.13) in terms of these variables and collecting terms produces

$$\varphi(\gamma'_2 \gamma'_1 \gamma_1^{-1} \gamma_2^{-1} \gamma_1 \gamma_2 \gamma_2'^{-1} \gamma_1'^{-1}), \quad (3.14)$$

which is manifestly trivial since $[\gamma_1, \gamma_2] = [\gamma'_1, \gamma'_2] = 1$, so the phases appearing between sectors from B are invariant under coboundary shifts. B then only matters up to its class in $H^1(G, H^1(K, U(1)))$, as claimed earlier.

3. Modular invariance

Finally, we can check that coefficients calculated in this way do not disrupt modular invariance in the decomposed orbifold. When the mixed anomaly phases are redundant with discrete torsion in Γ , we simply have decomposition with discrete torsion as studied in [26], and modular invariance should hold automatically. It remains to verify modular invariance when the mixed anomaly phases cannot be described by discrete torsion. Following the discussion

of Sec. III C 1 above, let us assume that we have already verified reference independence in the example at hand.

As a recap, we have the coefficient of Z_{g_1, g_2} in the postdecomposition partition function calculated in (3.7) as

$$F(\gamma_1, \gamma_2) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \gamma_1, k_2 \gamma_2] = 1}} \frac{(g_1^{-1} \cdot B)(g_1, k_2)}{(g_2^{-1} \cdot B)(g_2, k_1)}. \quad (3.15)$$

The input here is a pair γ_1, γ_2 of elements of Γ satisfying

$$\pi(\gamma_1) = g_1, \quad \pi(\gamma_2) = g_2, \quad [\gamma_1, \gamma_2] = 1, \quad (3.16)$$

though as argued above we are free to choose any such γ_1, γ_2 satisfying these conditions without changing the answer. Now imagine that we apply a modular transformation to g_1 and g_2 —call the result $\bar{g}_1$ and $\bar{g}_2$. We would calculate the coefficient of $Z_{\bar{g}_1, \bar{g}_2}$ as

$$F(\bar{\gamma}_1, \bar{\gamma}_2) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \bar{\gamma}_1, k_2 \bar{\gamma}_2] = 1}} \frac{(\bar{g}_1^{-1} \cdot B)(\bar{g}_1, \bar{k}_2)}{(\bar{g}_2^{-1} \cdot B)(\bar{g}_2, \bar{k}_1)} \quad (3.17)$$

for any $\bar{\gamma}_1, \bar{\gamma}_2$ satisfying

$$\pi(\bar{\gamma}_1) = \bar{g}_1, \quad \pi(\bar{\gamma}_2) = \bar{g}_2, \quad [\bar{\gamma}_1, \bar{\gamma}_2] = 1. \quad (3.18)$$

Take

$$\bar{g}_1 = g_2, \quad \bar{g}_2 = g_1^{-1}. \quad (3.19)$$

This is a pair related by the modular S transformation. Our goal will be to relate $F(\bar{\gamma}_1, \bar{\gamma}_2)$ to $F(\gamma_1, \gamma_2)$. It is then quite natural to make the choice

$$\bar{\gamma}_1 = \gamma_2, \quad \bar{\gamma}_2 = \gamma_1^{-1}. \quad (3.20)$$

Thanks to the stipulated properties (3.16) of γ_1, γ_2 , this choice of $\bar{\gamma}_1, \bar{\gamma}_2$ manifestly satisfies (3.18). With these choices, $F(\bar{\gamma}_1, \bar{\gamma}_2)$ takes the form

$$F(\bar{\gamma}_1, \bar{\gamma}_2) = F(\gamma_2, \gamma_1^{-1}) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \gamma_2, k_2 \gamma_1^{-1}] = 1}} \frac{(g_2^{-1} \cdot B)(g_2, \bar{k}_2)}{(g_1 \cdot B)(g_1^{-1}, \bar{k}_1)}. \quad (3.21)$$

Of course, we are free to redefine our summation variables, at the cost of changing the commutation constraint to compensate. We will shift to a sum over k_1 and k_2 given by

$$\bar{k}_1 = \gamma_1 k_2 \gamma_1^{-1}, \quad \bar{k}_2 = k_1^{-1}. \quad (3.22)$$

With this redefinition, the commutation constraint appearing in the sum takes the form $[\gamma_1 k_2 \gamma_1^{-1} \gamma_2, k_1^{-1} \gamma_1^{-1}] = 1$. Writing out this constraint and keeping in mind that γ_1 commutes with γ_2 , it is straightforward to rearrange it to

produce $[k_1 \gamma_1, k_2 \gamma_2] = 1$, as appearing in the original $F(\gamma_1, \gamma_2)$ sum. Now that we are summing over the same set of k_1, k_2 in each version, it remains to compare the summands. With all of the choices made so far, we have

$$F(\gamma_2, \gamma_1^{-1}) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \gamma_1, k_2 \gamma_2] = 1}} \frac{(g_2^{-1} \cdot B)(g_2, k_1^{-1})}{(g_1 \cdot B)(g_1^{-1}, \gamma_1 k_2 \gamma_1^{-1})}. \quad (3.23)$$

Keeping in mind the properties (3.2) and (3.4) of B , one readily transforms the above summand to match the one in (3.7), giving $F(\gamma_1, \gamma_2) = F(\gamma_2, \gamma_1^{-1})$.

Now we would like to repeat this process for pairs in G related by the modular T transformation. We can achieve this by choosing

$$\bar{g}_1 = g_1, \quad \bar{g}_2 = g_1 g_2. \quad (3.24)$$

As before, a convenient choice of $\bar{\gamma}_1$ and $\bar{\gamma}_2$ mimics the choice of $\bar{g}$ and is given by

$$\bar{\gamma}_1 = \gamma_1, \quad \bar{\gamma}_2 = \gamma_1 \gamma_2. \quad (3.25)$$

With these choices,

$$F(\bar{\gamma}_1, \bar{\gamma}_2) = F(\gamma_1, \gamma_1 \gamma_2) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \gamma_1, k_2 \gamma_1 \gamma_2] = 1}} \frac{(g_1^{-1} \cdot B)(g_1, \bar{k}_2)}{[(g_1 g_2)^{-1} \cdot B](g_1 g_2, \bar{k}_1)}. \quad (3.26)$$

Shifting our $\bar{k}$ to

$$\bar{k}_1 = k_1, \quad \bar{k}_2 = k_2 \gamma_2 k_1 \gamma_2^{-1} \quad (3.27)$$

gives us the commutation constraint $[k_1 \gamma_1, k_1 \gamma_1 k_2 \gamma_2] = 1$, again equivalent to $[k_1 \gamma_1, k_2 \gamma_2] = 1$. At this point we have

$$F(\gamma_1, \gamma_1 \gamma_2) = \sum_{\substack{k_1, k_2 \in K \\ [k_1 \gamma_1, k_2 \gamma_2] = 1}} \frac{(g_1^{-1} \cdot B)(g_1, k_2 \gamma_2 k_1 \gamma_2^{-1})}{[(g_1 g_2)^{-1} \cdot B](g_1 g_2, k_1)}. \quad (3.28)$$

Manipulations similar to those used in the previous case restore the initial form of the summand, leading to $F(\gamma_1, \gamma_1 \gamma_2) = F(\gamma_1, \gamma_2)$.

4. Summary

To recap the meaning of these many calculations, when we apply decomposition to express the genus one partition function of an orbifold by Γ with trivially-acting subgroup K in terms of a direct sum of orbifolds by subgroups of $G = \Gamma/K$, a mixed anomaly between the K TPOs and the Γ TDLs can contribute relative phases between the sectors of the Γ orbifold. These phases are given explicitly in (3.6), and depend only on the cohomology class of B in $H^1(G, H^1(K, U(1)))$ pulled back to $H^1(\Gamma, H^1(K, U(1)))$.

These phases are sometimes, but not always, redundant with discrete torsion in Γ . The coefficient of Z_{g_1, g_2} in the decomposed orbifold is given by the sum of these phases as (3.7)—this function F gives the coefficient of $Z_{\pi(\gamma_1), \pi(\gamma_2)} = Z_{g_1, g_2}$ for a fixed commuting pair γ_1, γ_2 . Sometimes there will be a canonical choice of γ_1, γ_2 ; when there is not, one may need to verify that the results will not depend on such a choice. The coefficients obtained this way are invariant under the generators of the modular group, and therefore modular invariance is not compromised by the phases arising from B . This formulation not only extends the results of studying such phases in [27–29], it provides physical insight to their origin as mixed anomalies.

D. Mixed gauging

So far we have discussed gauging a zero-form symmetry Γ with trivially-acting subgroup K . In general, the resulting theory has multiple vacua, and a one-form symmetry. Gauging that one-form symmetry projects the theory onto the component in the direct sum associated with that vacuum, as discussed in [34]. In general, we expect subsequent gaugings of a theory to be composable, so this operation should be expressible as a gauging of the original theory. In fact, we can give a relatively simple prescription for the result of such a gauging, motivated by a condensation defect construction in Sec. 3 [48].

For clarity, let us refer schematically to the original theory as T . Gauging the zero-form symmetry $\Gamma_{[0]}$ yields a theory denoted $[T/\Gamma_{[0]}]$, and gauging the one-form symmetry¹⁴ $K_{[1]}$ of the resulting theory brings us to $[[T/\Gamma_{[0]}]/K_{[1]}]$. A correlation function in $[[T/\Gamma_{[0]}]/K_{[1]}]$ can be thought of as a correlation function in $[T/\Gamma_{[0]}]$ with a projector Π_R inserted, given by

$$\Pi_R = \sum_i \frac{\dim R_i}{|K|} \sum_{k \in K} \chi_{R_i}(k^{-1}) \sigma_k, \quad (3.29)$$

following (2.5) and (2.6). Here each R_i is an irreducible representation, and we sum over orbits of irreducible representations related by the action of G on K .

A correlation function of $[T/\Gamma_{[0]}]$ with Π_R inserted is in turn calculated as a sum [with coefficients from (3.29)] of correlation functions of T , each with an insertion of the TPO σ_k . Applying this logic to the genus one partition function, we expect the $[[T/\Gamma_{[0]}]/K_{[1]}]$ partition function to be expressible as a sum of the form¹⁵

¹⁴In the mathematics literature, one-form symmetries have been denoted BK for several decades; here, we use a more recent convention to denote them by $K_{[1]}$.

¹⁵Schematically, we might regard this as a gauging of T by an extension of a one-form symmetry by a zero-form symmetry. Appendix B provides some additional insight to the interpretation of the symmetry of the ungauged theory as such a mixed extension.




FIG. 10. The visual representation of $Z_{\gamma_1, \gamma_2, \sigma_k}$.

$$\sum_i \frac{\dim R_i}{|K||\Gamma|} \sum_{\substack{\gamma_1, \gamma_2 \in \Gamma \\ k \in K}} \chi_{R_i}(k^{-1}) Z_{\gamma_1, \gamma_2, \sigma_k}, \quad (3.30)$$

where $Z_{\gamma_1, \gamma_2, \sigma_k}$ represents a correlation function taken on a torus with γ_1 and γ_2 TDLs wrapping the cycles and the TPO σ_k inserted at their intersection, as shown in Fig. 10. Analyzing these partition functions will be made easier due to the fact that many of the $Z_{\gamma_1, \gamma_2, \sigma_k}$ will vanish. Consistency with the group law gives us the condition

$$\gamma_1 \gamma_2 = k \gamma_2 \gamma_1. \quad (3.31)$$

Let us examine what this construction yields when $\Gamma = K$, i.e., the entire zero-form symmetry is trivially acting. To begin, assume K is Abelian. Then the condition (3.31) is only satisfied for $k = 1$. Since G is trivial, there is no action of G on K and hence each irreducible representation is its own orbit, getting rid of the i sum. Each character appearing in (3.30) is evaluated on the identity, giving 1. Since all of Γ acts trivially, the remaining $Z_{\gamma_1, \gamma_2, \sigma_1}$ are all equivalent to the parent theory partition function $Z_{1,1,\sigma_1}$ and we have

$$\frac{1}{|\Gamma|^2} \sum_{\gamma_1, \gamma_2 \in \Gamma} Z_{1,1,\sigma_1} = Z_{1,1,\sigma_1}. \quad (3.32)$$

Now assume $\Gamma = K$ is non-Abelian—for ease, we will begin by examining the universe corresponding to the identity representation. There will now be pairs γ_1, γ_2 that do not commute (hence $Z_{\gamma_1, \gamma_2, \sigma_1}$ will vanish), but for each such pair there is a nonvanishing contribution to the sum from $Z_{\gamma_1, \gamma_2, \sigma_{[\gamma_1, \gamma_2]}}$, so we once again end up with a single copy of the parent theory partition function.

Moving to a generic irreducible representation R , we have

$$\frac{\dim R}{|\Gamma|^2} \sum_{\gamma_1, \gamma_2 \in \Gamma} \chi_R([\gamma_1, \gamma_2]^{-1}) Z_{\gamma_1, \gamma_2, \sigma_{[\gamma_1, \gamma_2]}}. \quad (3.33)$$

Once again each factor of Z will reduce to the parent theory partition function (since the group is entirely trivially acting), so we simply need to tally up the coefficients. This requires us to evaluate

$$\sum_{\gamma_1, \gamma_2 \in \Gamma} \chi_R([\gamma_1, \gamma_2]^{-1}). \quad (3.34)$$

To do so, we use a classical result of Frobenius that the number $N(\gamma)$ of pairs γ_1, γ_2 satisfying $[\gamma_1, \gamma_2] = \gamma$ is given by

$$N(\gamma) = |\Gamma| \sum_R \frac{\chi_R(\gamma)}{\chi_R(1)} = |\Gamma| \sum_R \frac{\chi_R(\gamma)}{\dim R}. \quad (3.35)$$

This allows us to rewrite (3.34) as

$$\sum_{\gamma \in \Gamma} N(\gamma) \chi_R(\gamma^{-1}) = |\Gamma| \sum_{\substack{\gamma \in \Gamma \\ R' \in \text{Irr}^*}} \frac{\chi_R(\gamma^{-1}) \chi_{R'}(\gamma)}{\dim R'} = \frac{|\Gamma|^2}{\dim R}, \quad (3.36)$$

where we have used orthonormality of characters to arrive at the final result. This calculation shows that (3.33) reduces to a single copy of the parent theory partition function, independent of R . Of course, this is exactly the result we would have expected; when we gauge a totally trivially-acting symmetry, each universe in the decomposition has the same local spectrum as the parent theory, so the result of projecting onto a single copy would be (a single copy of) the parent theory partition function.

In cases where $\Gamma \neq K$ (so there is an effective symmetry in play), we cannot produce a general answer as easily, so instead we will examine a particular example: the symmetric group S_3 with trivially-acting $\mathbb{Z}_3$ subgroup. Again we begin by examining the projector associated to the identity representation. Since the $\mathbb{Z}_3$ subgroup of S_3 is also its commutator subgroup, we can once again find exactly one $Z_{\gamma_1, \gamma_2, \sigma[\gamma_1, \gamma_2]}$ for each pair γ_1, γ_2 . The sum in (3.30) will then have the maximum number of contributions (thirty six). If we let g be the generator of the effective $S_3/\mathbb{Z}_3 \simeq \mathbb{Z}_2$ symmetry, (3.30) in this case becomes

$$\frac{1}{2} [Z_{1,1} + Z_{1,g} + Z_{g,1} + Z_{g,g}], \quad (3.37)$$

which is the partition function of a single copy of the $\mathbb{Z}_2$ orbifold of the corresponding theory with no trivially-acting symmetry. Again this is in line with expectation, since the S_3 orbifold with trivially-acting $\mathbb{Z}_3$ decomposes into a copy of the parent theory plus a copy of its $\mathbb{Z}_2$ orbifold.

Of course, we should also be able to obtain the other universe present in the decomposition from mixed gauging. The remaining two nonidentity irreducible representations are exchanged by the action of $S_3/\mathbb{Z}_3 \simeq \mathbb{Z}_2$ on $\mathbb{Z}_3$, so the i sum in (3.30) has two terms. The relevant coefficients come from the projector which is built as

$$\begin{aligned} & \frac{1}{3} (\sigma_1 + e^{2\pi i/3} \sigma_b + e^{4\pi i/3} \sigma_{b^2}) + \frac{1}{3} (\sigma_1 + e^{4\pi i/3} \sigma_b + e^{2\pi i/3} \sigma_{b^2}) \\ &= \frac{1}{3} (2\sigma_1 - \sigma_b - \sigma_{b^2}), \end{aligned} \quad (3.38)$$

where b generates the $\mathbb{Z}_3$ orbifold group. Evaluating (3.30) with these coefficients causes each contribution to the partition function with a g to cancel out, and we are left with a single copy of the (unorbifolded) parent theory partition function $Z_{1,1}$, which is precisely the other universe in the decomposition.

In this context, the mixed anomalies of Sec. III A are exactly what their name suggests; an obstruction to the mixed gauging of TDLs and TPOs. As the examples of the following section will make clear, the phases (3.6) introduced by a mixed anomaly into the $[T/\Gamma_{[0]}]$ partition function obstruct its decomposition, which would prevent us from consistently being able to form projectors.

IV. EXAMPLES

Using the framework developed in the previous section, we now examine TDLs and TPOs in examples of orbifolds by noneffectively-acting symmetries. We will pay particular attention to the fate of TPOs under gauging to help us track the symmetries possessed by each theory. We introduce the notation $T(\Gamma, K)$ to denote a theory T with distinguished TDLs corresponding to total zero-form symmetry Γ of which the normal subgroup $K \subset \Gamma$ acts trivially.

The key idea in each example will be to track TPOs which are not associated to vacua. For one simple example, consider decomposition in an orbifold $[X/G]$ where the Abelian group Γ is a central extension

$$1 \rightarrow K \rightarrow \Gamma \rightarrow G \rightarrow 1 \quad (4.1)$$

by trivially-acting (and central) K . In the absence of mixed anomalies or discrete torsion, the statement of decomposition (2.3) simplifies to the form

$$\text{QFT}([T/\Gamma]) = \text{QFT}\left(\prod_{\rho \in \hat{K}} [X/G]_{\hat{\omega}(\rho)}\right). \quad (4.2)$$

Here, there are as many universes ($|K|$) as topological operators, so there are no “excess” TPOs, and so no additional trivially-acting symmetries are expected amongst the universes of the decomposition.

The story will become more interesting when the number of universes is smaller than the order of K . This can happen, for instance, if there are phases in the partition function which cause sectors to cancel with each other upon decomposition. These could be present in the Γ orbifold from the beginning (discrete torsion in Γ), or could appear as part of the decomposition process (due to a mixed anomaly). The other way to reduce the number of universes

is for Γ to be non-Abelian. Then a number of sectors Z_{γ_1, γ_2} will simply be absent from the start, since there will be pairs γ_1, γ_2 that do not commute. In each of these cases, there will be (linear combinations of) TPOs which do not form vacua, and instead remain bound to TDLs, providing some universes with trivially-acting symmetries. Exploring how the TPOs distribute themselves among universes will be the focus of this section.

A. $\Gamma = \mathbb{Z}_2 \times \mathbb{Z}_2$ or $\mathbb{Z}_4, K = \mathbb{Z}_2$

We begin by considering a theory $T(\mathbb{Z}_2, 1)$, with an effectively-acting $\mathbb{Z}_2$ symmetry. There exist two ways to supplement $T(\mathbb{Z}_2, 1)$ with a trivially-acting $\mathbb{Z}_2$ symmetry, denoted $T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2)$ and $T(\mathbb{Z}_4, \mathbb{Z}_2)$, both of which have the same local spectrum as T . In $T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2)$ the two symmetries combine trivially to give $\Gamma = \mathbb{Z}_2 \times \mathbb{Z}_2$, while we take $T(\mathbb{Z}_4, \mathbb{Z}_2)$ to have nontrivial extension class, giving $\Gamma = \mathbb{Z}_4$.

As local theories, decomposition predicts that in both of these cases the Γ orbifolds are identical,

$$\text{QFT}([T/\mathbb{Z}_2 \times \mathbb{Z}_2]) = \text{QFT}\left(\coprod_2 [T/\mathbb{Z}_2]\right) = \text{QFT}([T/\mathbb{Z}_4]). \quad (4.3)$$

as is easily checked by e.g., partition functions. However, one might also ask about extended objects in these theories. Schematically, because there are no further TPOs beyond the vacua of the two universes, we do not expect either universe to have trivially-acting symmetries, hence

$$\begin{aligned} [T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2)/(\mathbb{Z}_2 \times \mathbb{Z}_2)] \\ = [T(\mathbb{Z}_2, 1)/\mathbb{Z}_2] \coprod [T(\mathbb{Z}_2, 1)/\mathbb{Z}_2], \end{aligned} \quad (4.4)$$

$$= [T(\mathbb{Z}_4, \mathbb{Z}_2)/\mathbb{Z}_4]. \quad (4.5)$$

While the resulting theories are given by the same direct sum, there is additional information not captured in the notation above that distinguishes them. We know to expect a Γ orbifold to have a $\hat{\Gamma}$ quantum symmetry. This means that (4.4) should possess a $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry. This is given by the product of the symmetry that acts as the effective $\mathbb{Z}_2$ on both copies with the $\mathbb{Z}_2$ exchange symmetry possessed by the two copies. The story is similar for $[T(\mathbb{Z}_4, \mathbb{Z}_2)/\mathbb{Z}_4]$, except we expect a quantum $\mathbb{Z}_4$ symmetry. In this case the two symmetries are combined nontrivially; schematically, we have $(\text{exchange} \mathbb{Z}_2, \text{effective} \mathbb{Z}_2) \simeq \mathbb{Z}_4$.

In both cases above, the TPO that implements the trivially-acting $\mathbb{Z}_2$ symmetry in the ungauged theory becomes the vacuum of a second universe in the gauged theory. Let us see how this changes if we take these theories to have nontrivial mixed anomaly. From the analysis of Secs. III A and III B, we expect mixed anomalies for these theories to be classified by elements of $H^1(\Gamma, \hat{K})$ which are

pulled back from $H^1(G, \hat{K})$, which in this case is simply $\text{Hom}(\mathbb{Z}_2, \mathbb{Z}_2) = \mathbb{Z}_2$. Therefore, there is one nontrivial choice of mixed anomaly.

Such mixed anomaly phases were denoted ‘quantum symmetry phases’ in [28], and from the decomposition statement in Sec. III [28], we expect that for a $\Gamma = \mathbb{Z}_2 \times \mathbb{Z}_2$ orbifold, with $G = K = \mathbb{Z}_2$ and nontrivial $B \in H^1(G, \hat{K}) = \mathbb{Z}_2$ corresponding to an isomorphism $G \rightarrow K$, we have

$$\text{QFT}([T/\mathbb{Z}_2 \times \mathbb{Z}_2]) = \text{QFT}\left(\left[\frac{T \times \text{Coker} B}{\text{Ker} B}\right]\right) = \text{QFT}(T), \quad (4.6)$$

as $\text{Ker} B = 1 = \text{Coker} B$.

Now, let us refine this statement to include TDLs and TPOs. With this mixed anomaly, our prescription (3.6) for computing orbifold partition functions suggests that we reobtain the parent theory,

$$[T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2)/(\mathbb{Z}_2 \times \mathbb{Z}_2)|_{\text{M.A.}}] = T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2) \quad (4.7)$$

[refining (4.6)] and

$$[T(\mathbb{Z}_4, \mathbb{Z}_2)/\mathbb{Z}_4|_{\text{M.A.}}] = T(\mathbb{Z}_4, \mathbb{Z}_2). \quad (4.8)$$

Note that in the $\mathbb{Z}_2 \times \mathbb{Z}_2$ orbifold we could turn on discrete torsion, as well, and decomposition in orbifolds with discrete torsion was discussed in [26]. The phases that would result from this addition are in fact equivalent to those of the mixed anomaly, as discussed in [28]. Using the results there, decomposition predicts

$$\text{QFT}([T/\mathbb{Z}_2 \times \mathbb{Z}_2]|_{\text{D.T.}}) = \text{QFT}(T), \quad (4.9)$$

see e.g., see Sec. 5.1 in Ref. [26].

If we refine the analysis to track trivially-acting symmetries, then from the same reasoning as above, we have

$$[T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2)/(\mathbb{Z}_2 \times \mathbb{Z}_2)|_{\text{D.T.}}] = T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2). \quad (4.10)$$

In this case the mixed anomaly and discrete torsion have redundant effects. In the $\mathbb{Z}_4$ orbifold, of course, there is no choice of discrete torsion, as $H^2(\mathbb{Z}_4, U(1))$ is trivial. The phases from the mixed anomaly are something wholly dependent on the presence of a trivially-acting subgroup.

B. $\Gamma = S_3, K = \mathbb{Z}_3$

Next we take on the simplest non-Abelian example, a theory with TDLs that fuse as the symmetric group S_3 with a trivially-acting, noncentral $\mathbb{Z}_3$ subgroup. We will present S_3 as $\langle a, b | a^2 = b^3 = 1, abab = 1 \rangle$. The ungauged theory has three TPOs, which we can label $\sigma_1, \sigma_b, \sigma_{b^2}$, and using the fact that $G = \mathbb{Z}_2$ exchanges two irreducible representations

of $K = \mathbb{Z}_3$, leaving the third invariant, from (2.3) we see that decomposition predicts

$$\text{QFT}([T/S_3]) = \text{QFT}\left([T/\mathbb{Z}_2] \coprod T\right). \quad (4.11)$$

Now, let us consider extended objects in this decomposition. The algebra object for gauging is simply given by the direct sum of the group elements $A = 1 + a + b + ab + b^2 + ab^2$. We can quickly determine how many bulk TPOs exist in the gauged theory by acting on each ungauged TPO with A ,

$$\frac{1}{6}A \cdot \sigma_1 = \sigma_1, \quad (4.12)$$

$$\frac{1}{6}A \cdot \sigma_b = \frac{1}{2}(\sigma_b + \sigma_{b^2}), \quad (4.13)$$

$$\frac{1}{6}A \cdot \sigma_{b^2} = \frac{1}{2}(\sigma_b + \sigma_{b^2}). \quad (4.14)$$

Due to the nontrivial action of $\mathbb{Z}_2$ on $\mathbb{Z}_3$, we find that there are only two distinct bulk TPOs, consistent with the decomposition into two universes described above.

Next, we compute projectors. From (2.5), corresponding to the three irreducible representations of $K = \mathbb{Z}_3$ we have the three constituents,

$$\Pi_0 = \frac{1}{3}(\sigma_1 + \sigma_b + \sigma_{b^2}), \quad (4.15)$$

$$\Pi_1 = \frac{1}{3}(\sigma_1 + \xi\sigma_b + \xi^2\sigma_{b^2}), \quad (4.16)$$

$$\Pi_2 = \frac{1}{3}(\sigma_1 + \xi^2\sigma_b + \xi\sigma_{b^2}). \quad (4.17)$$

The universe $[T/\mathbb{Z}_2]$ corresponds to the trivial representation of $\mathbb{Z}_3$, and the universe T corresponds to each of the two nontrivial representations. If we denote T by L and $[T/\mathbb{Z}_2]$ by R , then we have the projectors

$$\Pi_L = \frac{1}{3}(2\sigma_1 - \sigma_b - \sigma_{b^2}) = \Pi_1 + \Pi_2, \quad (4.18)$$

$$\Pi_R = \frac{1}{3}(\sigma_1 + \sigma_b + \sigma_{b^2}) = \Pi_0. \quad (4.19)$$

Next, we consider the fate of the third TPO under gauging. We can see the answer by defining a quantity

$$g_L = \frac{i}{\sqrt{3}}(\sigma_b - \sigma_{b^2}) = \Pi_1 - \Pi_2. \quad (4.20)$$

Note that we have $g_L \cdot \Pi_R = 0$, meaning that g_L is an operator localized to the L universe. Also, it satisfies the relations $g_L \cdot \Pi_L = \Pi_L \cdot g_L = g_L$ and $g_L \cdot g_L = \Pi_L$, i.e., it

obeys a $\mathbb{Z}_2$ fusion law with that universe's vacuum. Thus, the third TPO is bound to TDLs in the L universe, and implements a trivially-acting $\mathbb{Z}_2$ symmetry there. In our schematic notation, the full decomposition including TDLs and TPOs is

$$[T(S_3, \mathbb{Z}_3)/S_3] = T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2) \coprod [T(\mathbb{Z}_2, 1)/\mathbb{Z}_2], \quad (4.21)$$

which refines (4.11).

C. $\Gamma = Q_8, K = \mathbb{Z}_4$

In this example we once again have a noncentral subgroup, $\mathbb{Z}_4$ in the group Q_8 of unit quaternions. We present Q_8 in the usual manner as

$$\{i, j, k | i^2 = j^2 = k^2 = ijk = -1\}. \quad (4.22)$$

Taking the trivially-acting $\mathbb{Z}_4$ to be generated by k , there are four TPOs in the ungauged theory: $\sigma_1, \sigma_k, \sigma_{-1}$, and σ_{-k} . The decomposition of the Q_8 orbifold was discussed in Sec [5], where it was argued that

$$\text{QFT}([T/Q_8]) = \text{QFT}\left([T/\mathbb{Z}_2] \coprod [T/\mathbb{Z}_2] \coprod T\right). \quad (4.23)$$

This can also be seen from the formula (2.3), using the fact that of the four irreducible representations of the trivially-acting $\mathbb{Z}_4$, two are invariant under $G = \mathbb{Z}_2$ (corresponding to the two copies of $[T/\mathbb{Z}_2]$) and the other two are exchanged (corresponding to the one copy of T). In this section we will analyze the decomposition of the corresponding TDLs and TPOs, refining the decomposition statement above.

First, consider the algebra object A for the gauging above, which is the sum of the eight group elements. The action on the ungauged TPOs is readily verified to be

$$\frac{1}{8}A \cdot \sigma_1 = \sigma_1, \quad (4.24)$$

$$\frac{1}{8}A \cdot \sigma_{-1} = \sigma_{-1}, \quad (4.25)$$

$$\frac{1}{8}A \cdot \sigma_k = \frac{1}{2}(\sigma_k + \sigma_{-k}), \quad (4.26)$$

$$\frac{1}{8}A \cdot \sigma_{-k} = \frac{1}{2}(\sigma_k + \sigma_{-k}), \quad (4.27)$$

which signals that we should expect three universes in the decomposition. Their vacua correspond to the projection operators from (2.5) (see also Sec. 5.4.2 in Ref. [5] and Sec. 4.2 in Ref. [44])

$$\Pi_a = \frac{1}{2}(\sigma_1 - \sigma_{-1}), \quad (4.28)$$

$$\Pi_b = \frac{1}{4}(\sigma_1 + \sigma_{-1} + \sigma_k + \sigma_{-k}), \quad (4.29)$$

$$\Pi_c = \frac{1}{4}(\sigma_1 + \sigma_{-1} - \sigma_k - \sigma_{-k}). \quad (4.30)$$

Based on the previous examples we would expect the fourth TPO to have become localized to one of the three universes. Indeed, the quantity

$$g_a = \frac{i}{2}(\sigma_k - \sigma_{-k}) \quad (4.31)$$

lives in the a universe and generates a trivially-acting $\mathbb{Z}_2$ symmetry there. The full decomposition at the level of TDLs and TPOs then takes the form

$$\begin{aligned} [T(Q_8, \mathbb{Z}_4)/Q_8] &= T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2) \\ &\times \coprod [T(\mathbb{Z}_2, 1)/\mathbb{Z}_2] \coprod [T(\mathbb{Z}_2, 1)/\mathbb{Z}_2], \end{aligned} \quad (4.32)$$

refining the decomposition described in Sec. 5.4 in Ref. [5] and above in (4.23) by telling us that the universe corresponding to the unorbifolded theory carries an extra trivially-acting symmetry.

We can also take into account a mixed anomaly in this example. The zero-form symmetry fits into the short exact sequence

$$1 \rightarrow \mathbb{Z}_4 \rightarrow Q_8 \rightarrow \mathbb{Z}_2 \rightarrow 1, \quad (4.33)$$

and $H^1(\mathbb{Z}_2, \hat{\mathbb{Z}}_4) = \mathbb{Z}_2$, with nontrivial action on the coefficients. Decomposition in this example with non-trivial mixed anomaly was discussed in Sec. 5.1.3 [27], where it was argued that

$$\text{QFT}([T/Q_8]) = \text{QFT}(T \coprod T). \quad (4.34)$$

We shall next track TDLs and TPOs through this decomposition.

Pulling back the nontrivial element of $H^1(\mathbb{Z}_2, \hat{\mathbb{Z}}_4)$ to an element of $H^1(Q_8, \hat{\mathbb{Z}}_4)$ and using that to modify the action of the TDLs on the TPOs, we find the bulk gauged TPOs to be

$$\frac{1}{8}A \cdot \sigma_1 = \sigma_1, \quad (4.35)$$

$$\frac{1}{8}A \cdot \sigma_{-1} = 0, \quad (4.36)$$

$$\frac{1}{8}A \cdot \sigma_k = \frac{1}{2}(\sigma_k + i\sigma_{-k}), \quad (4.37)$$

$$\frac{1}{8}A \cdot \sigma_{-k} = \frac{1}{2i}(\sigma_k + i\sigma_{-k}). \quad (4.38)$$

Since we now only have two independent combinations of TPOs, we expect a decomposition into two universes. The two vacua are given by

$$\Pi_{\pm} = \frac{1}{2} \left[\sigma_1 \pm \left(\frac{1-i}{2}\sigma_k + \frac{1+i}{2}\sigma_{-k} \right) \right]. \quad (4.39)$$

Each universe carries a single TPO beyond the vacuum, hence a distinguished TDL describing a trivially-acting $\mathbb{Z}_2$ symmetry controlled by

$$g_{\pm} = \frac{1}{2\sqrt{2}i}[(1+i)\sigma_{-1} \pm i\sigma_k \pm \sigma_{-k}]. \quad (4.40)$$

The full decomposition, then, takes the form

$$[T(Q_8, \mathbb{Z}_4)/Q_8]_{\text{M.A.}} = T(\Delta, \mathbb{Z}_2) \coprod T(\Delta, \mathbb{Z}_2), \quad (4.41)$$

where Δ is either $\mathbb{Z}_2 \times \mathbb{Z}_2$ or $\mathbb{Z}_4$. This refines the decomposition result from Sec. 5.1.3 in Ref. [27] and above in (4.34).

D. $\Gamma = K = S_3$

In this example we take a theory with TDLs describing a completely trivially-acting S_3 symmetry. Once again there are TPOs σ_i in the ungauged theory for each trivially-acting group element. The qualitative difference here is that since K is non-Abelian, the trivial symmetries in the postdecomposition universes can be non-Abelian as well. As also discussed in Sec. II D, after gauging the trivially-acting S_3 , one gets three copies of the original theory:

$$\text{QFT}([T/S_3]) = \text{QFT}(T \coprod T \coprod T), \quad (4.42)$$

corresponding to the fact that there are three irreducible representations of S_3 .

Labeling the projectors into these universes as Π_1, Π_X, Π_Y , from the general formula (2.5) and the character Table I we have the projectors

$$\Pi_1 = \frac{1}{6}(\sigma_1 + \sigma_b + \sigma_{b^2} + \sigma_a + \sigma_{ab} + \sigma_{ab^2}), \quad (4.43)$$

$$\Pi_X = \frac{1}{6}(\sigma_1 + \sigma_b + \sigma_{b^2} - \sigma_a - \sigma_{ab} - \sigma_{ab^2}), \quad (4.44)$$

$$\Pi_Y = \frac{2}{6}(2\sigma_1 - \sigma_b - \sigma_{b^2}). \quad (4.45)$$

TABLE I. Character table for S_3 .

Representation	$\{1\}$	$\{b, b^2\}$	$\{a, ab, ab^2\}$
1	1	1	1
X	1	1	-1
Y	2	-1	0

In passing, we can use the explicit expressions above and a product of the form

$$R \cdot \sigma = \chi_R(\sigma)\sigma, \quad (4.46)$$

for any representation R and corresponding character χ_R , corresponding to the quantum symmetry, to justify the products given earlier in equations (2.12)–(2.15).

The three remaining linearly independent combinations form TPOs local to the Y universe, which take the form

$$i_Y = \frac{1}{\sqrt{3}}(\sigma_b - \sigma_{b^2}), \quad (4.47)$$

$$j_Y = \frac{i}{\sqrt{3}}(\sigma_{ab} - \sigma_{ab^2}), \quad (4.48)$$

$$k_Y = \frac{i}{3}[2\sigma_a - (\sigma_{ab} + \sigma_{ab^2})]. \quad (4.49)$$

These can be checked to satisfy

$$i_Y^2 = j_Y^2 = k_Y^2 = i_Y j_Y k_Y = -\Pi_Y, \quad (4.50)$$

group relations which closely resemble those of Q_8 . However, as $-\Pi_Y$ is not a distinct operator from Π_Y , this should be understood as the nontrivial projective representation of $\mathbb{Z}_2 \times \mathbb{Z}_2$. The decomposition including explicitly the distinguished TDLs and TPOs is then

$$[T(S_3, S_3)/S_3] = T(1, 1) \coprod T(1, 1) \times \coprod T(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2 \times \mathbb{Z}_2)_{\text{proj}}, \quad (4.51)$$

which refines the decomposition example in Sec. II D and above in (4.42).

This example is an ideal illustration of the additional information gained by examining decomposition with TPOs in mind; the traditional partition-function analysis of Sec. II D told us that gauging a trivially-acting S_3 symmetry gives us three copies of the parent theory. While it is true that all three theories in the direct sum (4.51) have identical local spectra, we see that they do in fact differ in their extended spectra. This also helps make sense of the asymmetric manner (2.12)–(2.15) in which the fusion category $\text{Rep}(S_3)$ acts on the gauged theory.

This has also been our first example of TPOs which fuse projectively, a phenomenon which is quite readily

understandable in the context of discrete torsion. Recall that for an effective grouplike symmetry K , networks of TDLs will decompose into three-way junctions, where e.g., a $k_1 k_2$ line splits into k_1 and k_2 . The Hilbert space of topological operators at this junction is isomorphic to $\mathbb{C}$. Whatever phase sits at this junction, however, is not unique—we are free to redefine the contribution of a line labeled by k by a cochain $\lambda: K \rightarrow U(1)$, which we will take to obey $\lambda^{-1}(k) = \lambda(k^{-1})$. Doing so changes the junction phase by $\lambda(k_1 k_2) \lambda^{-1}(k_1) \lambda^{-1}(k_2)$, which we recognize as a coboundary shift. Thus, the phases at such junctions are meaningful up to their class in $H^2(K, U(1))$, and this choice provides the discrete torsion when we gauge K .

Now consider the case where K is trivially acting. We could now build the three-way junction described above out of lines that end on TPOs, as illustrated in Fig. 11. The initial setup is shown in Fig. 11(a). In Fig. 11(b) we fuse the σ_{k_1} and σ_{k_2} TPOs, allowing the possibility that they are in a projective representation of K , such that the result $\sigma_{k_1 k_2}$ is multiplied by some $\omega(k_1, k_2) \in H^2(K, U(1))$. In Fig. 11(c) we fuse $\sigma_{k_1 k_2}$ and $\sigma_{(k_1 k_2)^{-1}}$ to the identity TPO. In principle this could produce another phase $\omega(k_1 k_2, (k_1 k_2)^{-1})$, but our earlier insistence to work with cochains satisfying $\lambda^{-1}(k) = \lambda(k^{-1})$ is equivalent to the condition that $\omega(k, k^{-1}) = 1$ (and there is always such a choice of representative). Therefore, we end up with the situation described above, where a $k_1 k_2$ line splits into k_1 and k_2 and the junction is multiplied by a phase given by a class in $H^2(K, U(1))$. This demonstrates that projective fusion of TPOs corresponding to trivially-acting symmetries is equivalent to turning on discrete torsion in that symmetry. Note the similarity here to the correspondence between symmetry-protected topological (SPT) phases and anomalies of theories on the boundary [81].

E. $\Gamma = K = \text{Rep}(S_3)$

In our final example, we look at a theory with a trivially-acting fusion categorical (as opposed to grouplike) symmetry. We can construct such a theory by beginning with the direct sum of n copies of a theory. These copies should have a zero-form S_n exchange symmetry. Gauging this should bring us to a single copy of the theory, which we expect on general grounds to have a quantum symmetry given by the fusion category $\text{Rep}(S_n)$ [75]. As we have seen in previous examples, as the quantum dual to an exchange symmetry, this $\text{Rep}(S_n)$ should act “trivially” on the theory (and we will shortly see how one adapts the notion of a trivial action to a nongrouplike scenario).

The simplest nontrivial example of such a construction occurs for $n = 3$, where our theory has a trivially-acting $\text{Rep}(S_3)$ symmetry. This fusion category was used as a running example throughout Sec. II; its fusion products were given in Fig. 2. Such a theory has three simple TDLs labeled by 1, X, and Y. 1 and X, which form a $\mathbb{Z}_2$




FIG. 11. We build a three-way junction out of TPOs that fuse projectively.

subsymmetry of $\text{Rep}(S_3)$, have the familiar trivial action on local operators $\mathcal{O}$:

$$1 \cdot \mathcal{O} = \mathcal{O}, \quad (4.52)$$

$$X \cdot \mathcal{O} = \mathcal{O}. \quad (4.53)$$

The remaining simple line Y , which is of dimension 2, acts on operators as

$$Y \cdot \mathcal{O} = 2\mathcal{O}. \quad (4.54)$$

The easiest way to see that this must be the case is to note that this “trivial” action should still respect the fusion rules, which in particular means that $(Y \otimes Y) \cdot \mathcal{O} = (1 + X + Y) \cdot \mathcal{O}$. In general, taking a line of dimension d to act as multiplication by d will respect the fusion rules (and this agrees with the notion that in the grouplike case we can take all trivially-acting TDLs to act as multiplication by 1). Additionally, taking $\mathcal{O}$ trivial tells us that an empty loop labeled by a line A contributes a factor of its dimension d_A to a correlation function, which is also a standard result [76].

As in grouplike examples, the triviality of this symmetry should mean that there exist TPOs which live on the TDLs in our theory. There should be three distinct TPOs; σ_1 , σ_X , and σ_Y , which fuse as the simple objects of $\text{Rep}(S_3)$. Their action on lines is quite similar to the grouplike case—if we have a line A that is incoming to a TPO σ_B , the outgoing line is labeled by $B \otimes A$. One consequence of including nongrouplike behavior is that we don’t quite have access to the same tricks. For instance, inserting the identity TPO and splitting it into a pair of inverse operators only works when the TDLs in question are invertible. This is not the case for e.g., Y in $\text{Rep}(S_3)$ —there is no TDL Y^{-1} satisfying $Y \otimes Y^{-1} = Y^{-1} \otimes Y = 1$.

Gauging the $\text{Rep}(S_3)$ symmetry will entail selecting the algebra object $A = 1 + X + 2Y$ (which corresponds to the regular representation of S_3). As before, the vacua of the various universes in the gauged theory should be expressible as linear combinations of the TPOs from the ungauged theory. This example will prove no exception. Projectors onto our three universes are

$$\Pi_1 = \frac{1}{6}(\sigma_1 + \sigma_X + 2\sigma_Y), \quad (4.55)$$

$$\Pi_{[a]} = \frac{1}{2}(\sigma_1 - \sigma_X), \quad (4.56)$$

$$\Pi_{[b]} = \frac{1}{3}(\sigma_1 + \sigma_X - \sigma_Y), \quad (4.57)$$

where we have identified projectors with conjugacy classes [formally, representations of $\text{Rep}(S_3)$] using the character Table I. One can check using the $\text{Rep}(S_3)$ fusion products that these are orthonormal combinations. In fact, we can write down a version of the formula (2.6) for the projectors in $\text{Rep}(K)$ orbifolds. Letting ξ_i denote the set of conjugacy classes of K ,

$$\Pi_{\xi_i} = \frac{|\xi_i|}{|K|} \sum_{R \in \text{Rep}(K)} \chi_R(\xi_i) \sigma_R, \quad (4.58)$$

where, in contrast to (2.6), we hold a conjugacy class fixed and sum over irreducible representations. Of course these are the same formula, and the rewriting is merely a reflection of the fact that swapping K for $\text{Rep}(K)$ exchanges the role of irreducible representations and conjugacy classes. As we should have expected, gauging $\text{Rep}(S_3)$ simply “undoes” the procedure described above in which the $\text{Rep}(S_3)$ symmetry was constructed as a quantum

symmetry—we have recovered the direct sum of three universes with which this construction began.

V. CONCLUSION

While two-dimensional theories proved convenient for the calculations in this paper, many of the statements regarding topological operators should apply to general dimension. For instance, one can engineer decomposition in three-dimensional theories by gauging trivially-acting one-form symmetries [46], and in general dimension by gauging a trivially-acting $(d-2)$ -form symmetry. Of course, the explanation in terms of topological operators is much the same in each of these cases: $(d-2)$ -form symmetries are controlled by topological operators of codimension $d-1$, i.e., TDLs. The trivial action of these TDLs is controlled by a set of operators of codimension d (TPOs) which live bound to those lines. Gauging the $(d-2)$ -form symmetry (potentially) “frees” the TPOs, which act as vacua of disjoint universes. In dimensions above two, we have access to a greater variety of ingredients—there will be various topological operators of dimension greater than one, and it would be interesting to extend the analysis of this paper to such cases, where there is potential for more varied mixed gaugings, mixed anomalies, etc.

Even in two dimensions, there remains work to be done. The analysis of the examples of Sec. IV provides a refinement of the usual decomposition story, in which we track how the TPOs from the ungauged theory divide themselves among the universes. This tells us not only how many universes we have, but provides information about the symmetries that are localized to each universe. Does this provide a full characterization of the symmetries of each universe, or are there further refinements available? Also, are we able to create heuristics to predict how these TPOs should split without doing the full computation, as we are for e.g., which symmetries are gauged in each universe [5,26]? Finally, we could consider how the story changes when our zero-form symmetries have gauge anomalies. These are points we plan to pursue in future work.

ACKNOWLEDGMENTS

We would like to thank D. Ben-Zvi, T. Pantev, and Y. Tachikawa for useful discussions. D.R. was partially supported by NSF Grant No. PHY-1820867. E.S. was partially supported by NSF Grant No. PHY-2014086.

APPENDIX A: CONSISTENCY CONDITIONS OF B FOR BOUND TPOs

We repeat the analysis of Sec. III A in the case that the TPOs are bound to lines. In this case, the mixed anomaly phase $B(\gamma, \gamma k \gamma^{-1})$ would arise when moving from the configuration shown in Fig. 12(a) to that of Fig. 12(b).




FIG. 12. The action of γ on σ_k for a bound TPO.

For readability of the figures, it will be convenient in this section to use a convention where a TPO labeled by k multiplies the label of an incoming TDL from the right rather than the left.

In order to constrain B in its K argument, we begin with the configuration shown in Fig. 13(a). We can then fuse the σ_{k_1} and σ_{k_2} TPOs to arrive at Fig. 13(b). Shrinking the circle then brings us to Fig. 13(e), while introducing a phase $B(\gamma, \gamma k_1 k_2 \gamma^{-1})$. Alternatively, we could have deformed Fig. 13(a) to Fig. 13(c) by pinching off a circle containing σ_{k_1} . Shrinking both circles in Fig. 13c leads to Fig. 13(d), while producing the phase $B(\gamma, \gamma k_1 \gamma^{-1}) B(\gamma, \gamma k_2 \gamma^{-1})$. Finally, fusing the TPOs in Fig. 13(d) leads to Fig. 13(e). Consistency then requires

$$B(\gamma, \gamma k_1 k_2 \gamma^{-1}) = B(\gamma, \gamma k_1 \gamma^{-1}) B(\gamma, \gamma k_2 \gamma^{-1}), \quad (\text{A1})$$

matching (3.2).

For the Γ argument, we begin with Fig. 14(a). Shrinking the rightmost circle would produce the TPO $\sigma_{\gamma_2 k \gamma_2^{-1}}$ and the phase $B(\gamma_2, \gamma_2 k \gamma_2^{-1})$, leading to Fig. 14(b). We can then shrink the remaining circle to produce Fig. 14(c) and pick up $B(\gamma_1, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1})$. Alternatively, we could have directly shrunk everything to the right of the $\gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1}$ TDL in Fig. 14(a) to arrive directly at Fig. 14(c), giving the phase $B(\gamma_1 \gamma_2, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1})$. Demanding equality of these paths leads to


$$B(\gamma_1 \gamma_2, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1}) = B(\gamma_1, \gamma_1 \gamma_2 k \gamma_2^{-1} \gamma_1^{-1}) B(\gamma_2, \gamma_2 k \gamma_2^{-1}), \quad (\text{A2})$$

reproducing (3.4) from the main text.

APPENDIX B: TRIVIAL SYMMETRIES AS A MIXED EXTENSION

In [82], Y. Tachikawa explores the idea that higher-form symmetries of varying degrees can mix in generalizations of group extensions, i.e., a theory could have a theory with symmetry Ω fitting into a short exact sequence

$$1 \rightarrow M_{[p]} \rightarrow \Omega \rightarrow N_{[q]} \rightarrow 1, \quad (\text{B1})$$


FIG. 13. K argument consistency.

where a subscript $[n]$ denotes an n -form symmetry. The conjecture of [82] is that such an extension should be associated to a fibration of Eilenberg-MacLane spaces

$$K(M, p+1) \rightarrow B\Omega \rightarrow K(N, q+1), \quad (\text{B2})$$

and that $B\Omega$ should be the classifying space for the mixed symmetry Ω .

As this paper focuses on a mix of zero- and one-form symmetries, we might wonder if our framework can be cast in this language. Qualitatively, we have seen that a theory with trivially-acting symmetries includes zero-form

FIG. 14. Γ argument consistency.




FIG. 15. The general case of a point operator acting by a hom $\delta: K \rightarrow \Gamma$.

symmetries which can stand on their own, i.e., they are proper subsymmetries of the system which we can gauge. It also includes one-form symmetries (in the form of a nontrivial set of TPOs) which can not be separately gauged, as they are bound to the zero-form symmetries. Examining (B1), this would be consistent with $p = 0$ and $q = 1$, which is to say our one-form symmetry is extended nontrivially by our zero-form symmetry. Matching the notation used elsewhere in the paper, then, we would formally regard a theory with zero-form symmetry Γ and trivially-acting Abelian subgroup K as having a mixed symmetry described by Ω in¹⁶

$$1 \rightarrow \Gamma_{[0]} \rightarrow \Omega \rightarrow K_{[1]} \rightarrow 1. \quad (\text{B3})$$

Note that such an extension is opposite to the case which is most often considered in the literature, in which a higher-form symmetry extends a zero-form symmetry [38,46,63,80,83–85].

1. Interpretation of the extension class

In the following we focus on the case where Γ is Abelian, as that allows for the most concrete statements. Extensions of the form (B3), interpreted in the category of complexes of Abelian groups, induce¹⁷ a homomorphism $K \rightarrow \Gamma$ [though not every such homomorphism may necessarily arise from an extension (B3)].

As with group extensions, a nontrivial extension class should in general obstruct $K_{[1]}$ from being a subsymmetry of the system. We can see this physically as follows: Letting the extension class be $\delta: K \rightarrow \Gamma$, a point labeled by k affects an incoming line labeled by some γ as in Fig. 15.

When δ is nontrivial, the K points are “bound” to the Γ lines. If it happens that $\ker(\delta)$ is all of K , however, we could alter Fig. 15 as shown in Fig. 16, so a trivial extension class “frees” the points from the lines, allowing each symmetry to stand on its own. In the cases examined in this paper, we always have K as a normal subgroup of Γ , and so there is a natural candidate for δ ; the inclusion map.

¹⁶ $K_{[1]}$ is also often written as BK , particularly in the mathematical literature.

¹⁷We would like to thank T. Pantev for results on such extensions. Note in passing that the assignment of such homomorphisms to extensions (B3) is natural, as one might suspect that they should correspond to elements of

$$H_{\text{sing}}^2(B^2K, \Gamma) = \text{Hom}(K(K, 2), K(\Gamma, 2)) = \text{Hom}(K, \Gamma), \quad (\text{B4})$$

a computation described in e.g., Sec. 3.2.1 [48] and Ref. [86].

2. Orbifolds and mixed extensions

Suppose we gauge $\Gamma_{[0]}$. Given our assumption that the total symmetry of the system is an extension of $K_{[1]}$ by $\Gamma_{[0]}$, then from the formal considerations discussed so far, gauging $\Gamma_{[0]}$ should, on its face, bring us to a theory whose symmetry is an extension of $K_{[1]}$ by $\text{Rep}(\Gamma)_{[0]}$.

We will describe the theory both before and after gauging as completely as we can, still under the assumption that Γ is Abelian. Our setup, as it has been before, is a 2d CFT with zero-form symmetry $\Gamma = K.G$, where K acts trivially. The total symmetry should be an extension of $K_{[1]}$ by $\Gamma_{[0]}$, with the extension class given by the inclusion map $\delta: K \rightarrow \Gamma$. This map is injective, so δ has trivial kernel, and therefore all of the nontrivial TPOs are bound to lines. The theory therefore has a unique vacuum and does not exhibit decomposition.

When we orbifold by $\Gamma_{[0]}$, if there is no mixed anomaly between the points and the lines, each TPO from the ungauged theory becomes a freestanding operator in the gauged theory. The extension class $\tilde{\delta}$ of the gauged theory then has all of K as its kernel. The zero-form symmetry of the gauged theory is given by $\hat{\Gamma}$, the Pontryagin dual to Γ . In order to figure out what mixed anomaly the gauged theory has, we can note that $\hat{\Gamma}$ acts on local operators through phases given by characters,

$$\hat{\gamma} \cdot \sigma_k = \sigma_k \times \chi_{\hat{\gamma}}(\delta(k)), \quad (\text{B5})$$

from which we see that the mixed anomaly is given by $\chi_{\hat{\gamma}}(\delta(k))$.

In fact, we can repeat this argument to learn more about the mixed anomaly in the ungauged theory. Note that gauging $\hat{\Gamma}$ brings us to a theory with symmetry $\hat{\Gamma}$, which by Pontryagin duality is canonically isomorphic to Γ . This is the sense in which gauging the quantum symmetry $\hat{\Gamma}$ undoes the first gauging to return us to the original theory. But we can repeat the argument from earlier—the mixed anomaly in the doubly gauged theory should be given by a character of $\hat{\Gamma}$ as $\chi_{\hat{\gamma}}(\tilde{\delta}(k))$. Of course we can map this by isomorphism to a character of Γ , so we expect that the mixed anomaly in the ungauged theory can be written as $\chi_{\gamma}(\delta(k))$. For ease of comparison, let us summarize these conclusions in tabular form:

	Ungauged theory	Gauged theory
Total symmetry	$1 \rightarrow \Gamma_{[0]} \rightarrow \Omega \rightarrow K_{[1]} \rightarrow 1$	$1 \rightarrow \hat{\Gamma}_{[0]} \rightarrow \tilde{\Omega} \rightarrow K_{[1]} \rightarrow 1$
Extension class	δ	$\tilde{\delta}$
Mixed anomaly	$\chi_{\gamma}(\delta(k))$	$\chi_{\hat{\gamma}}(\tilde{\delta}(k))$

Note how the extension class in the ungauged theory determines the mixed anomaly in the gauged theory and vice versa. This is entirely in line with the analysis of [82],



FIG. 16. When $\delta(k) = 1$, the point operator labeled by k can be “dragged off” of its line. The resulting identity point and line operators can then be erased, leaving the point operator free floating.

which predicts that this phenomenon should arise any time we gauge subsymmetries fitting into extensions.

These results are also in line with the predictions of decomposition. For instance, it is well known that if we gauge a trivially-acting Abelian group K , we end up with a direct sum of $|K|$ copies of the original theory. From the topological operator point of view, the vacuum in each theory is a local topological operator which originated as one of the TPOs that implemented the trivial symmetry in the ungauged theory (or possibly a linear combination of those TPOs). When we supplement a theory with a trivially-acting symmetry that has nontrivial mixed anomaly, the table above suggests that we should find a nontrivial extension class in the gauged theory. This means that not all of the TPOs in the gauged theory will be freestanding—decomposition will be obstructed (relative to the same theory without mixed anomaly). It also means that the resulting theory should have its own

trivially-acting symmetries implemented by the TPOs that are still bound to lines. The decomposition structure of orbifolds by noneffective symmetries with nontrivial mixed anomaly was investigated (under the name of quantum-symmetry phases) in [27–29], and indeed turning on a mixed anomaly obstructs decomposition in the gauged theory (as can also be seen from the examples in Sec. IV).

If we were to attempt to extend this analysis to non-Abelian Γ , we would quickly run into trouble. The calculation (B4) of the extension class only makes sense when Γ is Abelian, so while it is conceivable that we still have a classifying space given as a fibration, that fact alone would not get us far. In this case the approach taken in the main text where we focus on the physical behavior of the topological operators seems preferable, as that tactic allows us to work out the behavior of non-Abelian (and even nongrouplike) examples.

-
- [1] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, Topological defect lines and renormalization group flows in two dimensions, *J. High Energy Phys.* **01** (2019) 026.
 - [2] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Generalized global symmetries, *J. High Energy Phys.* **02** (2015) 172.
 - [3] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, Defect lines, dualities, and generalised orbifolds, in 16th International Congress on Mathematical Physics (2009), [arXiv:0909.5013](#).
 - [4] I. Brunner, N. Carqueville, and D. Plencner, Orbifolds and topological defects, *Commun. Math. Phys.* **332**, 669 (2014).
 - [5] S. Hellerman, A. Henriques, T. Pantev, E. Sharpe, and M. Ando, Cluster decomposition, T-duality, and gerby CFT’s, *Adv. Theor. Math. Phys.* **11**, 751 (2007).
 - [6] E. Andreini, Y. Jiang, and H.-H. Tseng, On Gromov-Witten theory of root gerbes, [arXiv:0812.4477](#).
 - [7] E. Andreini, Y. Jiang, and H.-H. Tseng, Gromov-Witten theory of product stacks, *Commun. Anal. Geom.* **24**, 223 (2016).
 - [8] E. Andreini, Y. Jiang, and H.-H. Tseng, Gromov-Witten theory of root gerbes I: Structure of genus 0 moduli spaces, *J. Diff. Geom.* **99**, 1 (2015).
 - [9] H.-H. Tseng, On degree-0 elliptic orbifold Gromov-Witten invariants, *Int. Math. Res. Not.* **2011**, 2444 (2011).
 - [10] A. Gholampour and H.-H. Tseng, On Donaldson-Thomas invariants of threefold stacks and gerbes, *Proc. Am. Math. Soc.* **141**, 191 (2013).
 - [11] X. Tang and H.-H. Tseng, Duality theorems for étale gerbes on orbifolds, *Adv. Math.* **250**, 496 (2014).
 - [12] A. Caldararu, J. Distler, S. Hellerman, T. Pantev, and E. Sharpe, Non-birational twisted derived equivalences in abelian GLSMs, *Commun. Math. Phys.* **294**, 605 (2010).
 - [13] K. Hori, Duality in two-dimensional (2,2) supersymmetric non-abelian gauge theories, *J. High Energy Phys.* **10** (2013) 121.
 - [14] E. Sharpe, Predictions for Gromov-Witten invariants of noncommutative resolutions, *J. Geom. Phys.* **74**, 256 (2013).
 - [15] N. M. Addington, E. P. Segal, and E. Sharpe, D-brane probes, branched double covers, and noncommutative resolutions, *Adv. Theor. Math. Phys.* **18**, 1369 (2014).
 - [16] J. Halverson, V. Kumar, and D. R. Morrison, New methods for characterizing phases of 2d supersymmetric gauge theories, *J. High Energy Phys.* **09** (2013) 143.

- [17] K. Hori and J. Knapp, Linear sigma models with strongly coupled phases—one parameter models, *J. High Energy Phys.* **11** (2013) 070.
- [18] K. Hori and J. Knapp, A pair of Calabi-Yau manifolds from a two parameter non-Abelian gauged linear sigma model, [arXiv:1612.06214](#).
- [19] K. Wong, Two-dimensional gauge dynamics and the topology of singular determinantal varieties, *J. High Energy Phys.* **03** (2017) 132.
- [20] M. Kapustka and M. Rampazzo, Torelli problem for Calabi-Yau threefolds with GLSM description, *Commun. Num. Theor. Phys.* **13**, 725 (2019).
- [21] Z. Chen, T. Pantev, and E. Sharpe, Landau-Ginzburg models for certain fiber products with curves, *J. Geom. Phys.* **137**, 204 (2019).
- [22] Z. Chen, J. Guo, and M. Romo, A GLSM view on homological projective duality, *Commun. Math. Phys.* **394**, 355 (2022).
- [23] J. Guo and M. Romo, Hybrid models for homological projective duals and noncommutative resolutions, *Lett. Math. Phys.* **112**, 117 (2022).
- [24] R. Eager and E. Sharpe, Elliptic genera of pure gauge theories in two dimensions with semisimple non-simply-connected gauge groups, *Commun. Math. Phys.* **387**, 267 (2021).
- [25] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, Symmetries and strings of adjoint QCD₂, *J. High Energy Phys.* **03** (2021) 103.
- [26] D. Robbins, E. Sharpe, and T. Vandermeulen, A generalization of decomposition in orbifolds, *J. High Energy Phys.* **21** (2020) 134.
- [27] D. G. Robbins, E. Sharpe, and T. Vandermeulen, Anomalies, extensions, and orbifolds, *Phys. Rev. D* **104**, 085009 (2021).
- [28] D. G. Robbins, E. Sharpe, and T. Vandermeulen, Quantum symmetries in orbifolds and decomposition, *J. High Energy Phys.* **02** (2022) 108.
- [29] D. G. Robbins, E. Sharpe, and T. Vandermeulen, Anomaly resolution via decomposition, *Int. J. Mod. Phys. A* **36**, 2150220 (2021).
- [30] J. Wang, X.-G. Wen, and E. Witten, Symmetric Gapped Interfaces of SPT and SET States: Systematic Constructions, *Phys. Rev. X* **8**, 031048 (2018).
- [31] S. Hellerman and E. Sharpe, Sums over topological sectors and quantization of Fayet-Iliopoulos parameters, *Adv. Theor. Math. Phys.* **15**, 1141 (2011).
- [32] L. B. Anderson, B. Jia, R. Manion, B. Ovrut, and E. Sharpe, General aspects of heterotic string compactifications on stacks and gerbes, *Adv. Theor. Math. Phys.* **19**, 531 (2015).
- [33] E. Sharpe, Decomposition in diverse dimensions, *Phys. Rev. D* **90**, 025030 (2014).
- [34] E. Sharpe, Undoing decomposition, *Int. J. Mod. Phys. A* **34**, 1950233 (2020).
- [35] Y. Tanizaki and M. Ünsal, Modified instanton sum in QCD and higher-groups, *J. High Energy Phys.* **03** (2020) 123.
- [36] M. Nguyen, Y. Tanizaki, and M. Ünsal, Semi-Abelian gauge theories, non-invertible symmetries, and string tensions beyond N -ality, *J. High Energy Phys.* **03** (2021) 238.
- [37] M. Nguyen, Y. Tanizaki, and M. Ünsal, Noninvertible 1-form symmetry and Casimir scaling in 2D Yang-Mills theory, *Phys. Rev. D* **104**, 065003 (2021).
- [38] M. Yu, Symmetries and anomalies of $(1+1)$ d theories: 2-groups and symmetry fractionalization, *J. High Energy Phys.* **08** (2021) 061.
- [39] A. Cherman and T. Jacobson, Lifetimes of near eternal false vacua, *Phys. Rev. D* **103**, 105012 (2021).
- [40] A. Cherman, T. Jacobson, and M. Neuzil, Universal deformations, *SciPost Phys.* **12**, 116 (2022).
- [41] M. Honda, E. Itou, Y. Kikuchi, and Y. Tanizaki, Negative string tension of higher-charge Schwinger model via digital quantum simulation, *Prog. Theor. Exp. Phys.* **2022**, 033B01 (2022).
- [42] W. Gu, D. Pei, and M. Zhang, On phases of 3d $N = 2$ Chern-Simons-matter theories, *Nucl. Phys. B* **973**, 115604 (2021).
- [43] W. Gu, Vacuum structures revisited, [arXiv:2110.13156](#).
- [44] E. Sharpe, Topological operators, noninvertible symmetries and decomposition, [arXiv:2108.13423](#).
- [45] S. Meynet and R. Moscrop, McKay quivers and decomposition, [arXiv:2208.07884](#).
- [46] T. Pantev, D. Robbins, E. Sharpe, and T. Vandermeulen, Orbifolds by 2-groups and decomposition, *J. High Energy Phys.* **09** (2022) 036.
- [47] T. Pantev and E. Sharpe, Decomposition in Chern-Simons theories in three dimensions, *Int. J. Mod. Phys. A* **37**, 2250227 (2022).
- [48] L. Lin, D. G. Robbins, and E. Sharpe, Decomposition, condensation defects, and fusion, *Fortschr. Phys.* **70**, 2200130 (2022).
- [49] E. Sharpe, Derived categories and stacks in physics, *Lect. Notes Phys.* **757**, 249 (2009).
- [50] E. Sharpe, Landau-Ginzburg models, gerbes, and Kuznetsov's homological projective duality, *Proc. Symp. Pure Math.* **81**, 237 (2010).
- [51] E. Sharpe, GLSM's, gerbes, and Kuznetsov's homological projective duality, *J. Phys. Conf. Ser.* **462**, 012047 (2013).
- [52] E. Sharpe, Categorical equivalence and the renormalization group, *Fortschr. Phys.* **67–9**, 1910019 (2019).
- [53] E. Sharpe, An introduction to decomposition, [arXiv:2204.09117](#).
- [54] T. Pantev and E. Sharpe, Notes on gauging noneffective group actions, [arXiv:hep-th/0502027](#).
- [55] T. Pantev and E. Sharpe, GLSM's for gerbes (and other toric stacks), *Adv. Theor. Math. Phys.* **10**, 77 (2006).
- [56] T. Pantev and E. Sharpe, String compactifications on Calabi-Yau stacks, *Nucl. Phys. B* **733**, 233 (2006).
- [57] D. S. Freed, Higher algebraic structures and quantization, *Commun. Math. Phys.* **159**, 343 (1994).
- [58] J. C. Baez and J. Huerta, An invitation to higher gauge theory, *Gen. Relativ. Gravit.* **43**, 2335 (2011).
- [59] H. Sati, Geometric and topological structures related to M-branes, *Proc. Symp. Pure Math.* **81**, 181 (2010).
- [60] H. Sati, Geometric and topological structures related to M-branes II: Twisted string and $String^c$ structures, *J. Aust. Math. Soc.* **90**, 93 (2011).
- [61] H. Sati, U. Schreiber, and J. Stasheff, L_∞ algebra connections and applications to String- and Chern-Simons n -transport, [arXiv:0801.3480](#).

- [62] C. J. Schommer-Pries, Central extensions of smooth 2-groups and a finite-dimensional string 2-group, *Geom. Topol.* **15**, 609 (2011).
- [63] J. C. Baez, D. Stevenson, A. S. Crans, and U. Schreiber, From loop groups to 2-groups, *Homology Homotopy Appl.* **9**, 101 (2007).
- [64] T. Nikolaus, C. Sachse, and C. Wockel, A smooth model for the String group, *Int. Math. Res. Not.* **16**, 3678 (2013).
- [65] D. N. Yetter, TQFTs from homotopy 2-types, *J. Knot Theory Ramifications* **2**, 113 (1993).
- [66] J. a. F. Martins and T. Porter, On Yetter's invariant and an extension of the Dijkgraaf-Witten invariant to categorical groups, *Theory Appl. Categ.* **18**, 118 (2007).
- [67] H. Grosse and K. G. Schlesinger, Duals for nonAbelian lattice gauge theories by categorical methods, *Int. J. Theor. Phys.* **40**, 459 (2001).
- [68] H. Pfeiffer, Higher gauge theory and a nonAbelian generalization of 2-form electrodynamics, *Ann. Phys. (Amsterdam)* **308**, 447 (2003).
- [69] N. Carqueville and I. Runkel, Orbifold completion of defect bicategories, *Quantum Topol.* **7**, 203 (2016).
- [70] N. Carqueville, A. R. Camacho, and I. Runkel, Orbifold equivalent potentials, *J. Pure Appl. Algebra* **220**, 759 (2016).
- [71] I. Brunner, N. Carqueville, and D. Plencner, Discrete torsion defects, *Commun. Math. Phys.* **337**, 429 (2015).
- [72] N. Carqueville and A. Q. Velez, Calabi-Yau completions and orbifold equivalences, [arXiv:1509.00880](https://arxiv.org/abs/1509.00880).
- [73] J. C. Wang, Z.-C. Gu, and X.-G. Wen, Field Theory Representation of Gauge-Gravity Symmetry-Protected Topological Invariants, Group Cohomology and Beyond, *Phys. Rev. Lett.* **114**, 031601 (2015).
- [74] J. Frohlich, J. Fuchs, I. Runkel, and C. Schweigert, Kramers-Wannier Duality from Conformal Defects, *Phys. Rev. Lett.* **93**, 070601 (2004).
- [75] L. Bhardwaj and Y. Tachikawa, On finite symmetries and their gauging in two dimensions, *J. High Energy Phys.* **03** (2018) 189.
- [76] J. Fuchs, I. Runkel, and C. Schweigert, TFT construction of RCFT correlators 1. Partition functions, *Nucl. Phys.* **B646**, 353 (2002).
- [77] L.-Y. Hung and Y. Wan, Generalized ADE classification of topological boundaries and anyon condensation, *J. High Energy Phys.* **07** (2015) 120.
- [78] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups: Maximal Subgroups and Ordinary Characters for Simple Groups*, (Clarendon Press, 1985).
- [79] L. Bhardwaj, S. Schafer-Nameki, and J. Wu, Universal non-invertible symmetries, *Fortschr. Phys.* **70**, 2200143 (2022).
- [80] F. Benini, C. Córdova, and P.-S. Hsin, On 2-group global symmetries and their anomalies, *J. High Energy Phys.* **03** (2019) 118.
- [81] X.-G. Wen, Classifying gauge anomalies through symmetry-protected trivial orders and classifying gravitational anomalies through topological orders, *Phys. Rev. D* **88**, 045013 (2013).
- [82] Y. Tachikawa, On gauging finite subgroups, *SciPost Phys.* **8**, 015 (2020).
- [83] Z. Wan and J. Wang, Adjoint QCD₄, deconfined critical phenomena, symmetry-enriched topological quantum field theory, and higher symmetry extension, *Phys. Rev. D* **99**, 065013 (2019).
- [84] R. Kobayashi, K. Ohmori, and Y. Tachikawa, On gapped boundaries for SPT phases beyond group cohomology, *J. High Energy Phys.* **11** (2019) 131.
- [85] A. Kapustin and R. Thorngren, Higher symmetry and gapped phases of gauge theories, [arXiv:1309.4721](https://arxiv.org/abs/1309.4721).
- [86] <https://math.stackexchange.com/questions/43142/maps-between-eilenberg-macLane-spaces>.