



OPEN ACCESS

EDITED BY

Andrei S. Steindorff,
Berkeley Lab (DOE), United States

REVIEWED BY

Paul Daly,
Jiangsu Academy of Agricultural Sciences
(JAAS), China
Lucia Ramirez,
Public University of Navarre, Spain
Irina Druzhinina,
Royal Botanic Gardens, Kew,
United Kingdom

*CORRESPONDENCE

Zheng Wang

✉ wang.zheng@yale.edu

Oded Yarden

✉ oded.yarden@mail.huji.ac.il

RECEIVED 30 April 2023

ACCEPTED 06 June 2023

PUBLISHED 30 June 2023

CITATION

Wang Z, Kim W, Wang Y-W, Yakubovich E,
Dong C, Trail F, Townsend JP and
Yarden O (2023) The Sordariomycetes:
an expanding resource with Big
Data for mining in evolutionary
genomics and transcriptomics.
Front. Fungal Biol. 4:1214537.
doi: 10.3389/ffunb.2023.1214537

COPYRIGHT

© 2023 Wang, Kim, Wang, Yakubovich,
Dong, Trail, Townsend and Yarden. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

The Sordariomycetes: an expanding resource with Big Data for mining in evolutionary genomics and transcriptomics

Zheng Wang^{1*}, Wonyong Kim², Yen-Wen Wang¹,
Elizabeta Yakubovich³, Caihong Dong⁴, Frances Trail^{5,6},
Jeffrey P. Townsend^{1,7} and Oded Yarden^{3*}

¹Department of Biostatistics, Yale School of Public Health, New Haven, CT, United States, ²Korean Lichen Research Institute, Sunchon National University, Suncheon, Republic of Korea, ³Department of Plant Pathology and Microbiology, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, Israel, ⁴Institute of Microbiology, Chinese Academy of Sciences, Beijing, China, ⁵Department of Plant Biology, Michigan State University, East Lansing, MI, United States, ⁶Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing, MI, United States, ⁷Department of Ecology and Evolutionary Biology, Program in Microbiology, and Program in Computational Biology and Bioinformatics, Yale University, New Haven, CT, United States

Advances in genomics and transcriptomics accompanying the rapid accumulation of omics data have provided new tools that have transformed and expanded the traditional concepts of model fungi. Evolutionary genomics and transcriptomics have flourished with the use of classical and newer fungal models that facilitate the study of diverse topics encompassing fungal biology and development. Technological advances have also created the opportunity to obtain and mine large datasets. One such continuously growing dataset is that of the Sordariomycetes, which exhibit a richness of species, ecological diversity, economic importance, and a profound research history on amenable models. Currently, 3,574 species of this class have been sequenced, comprising nearly one-third of the available ascomycete genomes. Among these genomes, multiple representatives of the model genera *Fusarium*, *Neurospora*, and *Trichoderma* are present. In this review, we examine recently published studies and data on the Sordariomycetes that have contributed novel insights to the field of fungal evolution *via* integrative analyses of the genetic, pathogenic, and other biological characteristics of the fungi. Some of these studies applied ancestral state analysis of gene expression among divergent lineages to infer regulatory network models, identify key genetic elements in fungal sexual development, and investigate the regulation of conidial germination and secondary metabolism. Such multispecies investigations address challenges in the study of fungal evolutionary genomics derived from studies that are often based on limited model genomes and that primarily focus on the aspects of biology driven by knowledge drawn from a few model species. Rapidly accumulating information and expanding capabilities for systems biological analysis of Big Data are setting the stage for the expansion of the concept of model systems from unitary taxonomic species/genera to inclusive clusters of well-studied models that can facilitate both the in-depth study of specific lineages and

also investigation of trait diversity across lineages. The Sordariomycetes class, in particular, offers abundant omics data and a large and active global research community. As such, the Sordariomycetes can form a core omics clade, providing a blueprint for the expansion of our knowledge of evolution at the genomic scale in the exciting era of Big Data and artificial intelligence, and serving as a reference for the future analysis of different taxonomic levels within the fungal kingdom.

KEYWORDS

Sordariomycetes, evolution, genomics, transcriptomics, Big Data, *Neurospora*, *Fusarium*, *Trichoderma*

Traditional models have served as stepping stones in evolutionary genomics

A model system—sometimes also referred to as a model species—is an optimized living non-human platform that can be easily accessed and manipulated (Davis, 2003b; Yarden et al., 2003; Ankeny and Leonelli, 2020). The progress made in experimental and computational biology, which has contributed to strengthening the links between descriptive research and mechanistic understanding—especially by utilizing molecular biology and genetics—has been largely based on the employment of model systems. Traditional fungal models, including unicellular fungi such as *Saccharomyces cerevisiae* (Saccharomycetaceae, Saccharomycetales) and *Schizosaccharomyces pombe* (Schizosaccharomycetaceae, Schizosaccharomycetales), as well as filamentous fungi such as *Neurospora* (Sordariaceae, Sordariales) and *Aspergillus* spp. (Trichocomaceae, Eurotiales), have well-characterized morphology, have been intensively studied genetically, and, in some cases, are also of economic importance (Yarden, 2007; Yarden, 2016). One of the primary bases for the selection of some of these organisms has been their amenability to developmental, physiological, and genetic manipulations. Currently, technological capabilities have progressed to the point where additional species can be studied and manipulated with increasing ease, providing the option to address specific questions and problems in a growing number of organisms (Goldstein and King, 2016).

The classic model systems, including yeasts, *Neurospora*, and *Aspergillus*, were among the first set of targets when genome sequencing became technically and economically feasible two decades ago, and omics data from these early models profoundly reshaped modern fungus research (Feldmann, 1999; Galagan et al., 2005; Sunnerhagen and Piskur, 2006; Dunlap et al., 2007; Jones, 2007; Nowrousian, 2014; Ellena and Steiger, 2022). Although the roles of classic fungal models in general biology have become increasingly confined (Davis, 2003a), new insights from the genomics of fungal models continue to attract interest in basic fungal biology, especially as they relate to evolutionary biology and ecology, including issues of diversity, invasive species, the impact of climate change, speciation and species concept, circadian clocks, and pathogenicity (Dyer and O’Gorman, 2012; Stukenbrock, 2013; Plissonneau et al., 2017; Stajich, 2017; Boekhout et al., 2021; North et al., 2021; Peris et al., 2022; Feurtey et al., 2023; Kelliher et al., 2023). In addition, these well-studied models have served as references for the annotation of less-studied and rarer systems, including many non-model species (Russell et al., 2017; Tworzydło and Bilinski, 2019), which range from taxonomically related species to ecologically and economically relevant species. However, the rapid development of comparative genomics techniques (Box 1) has facilitated the analysis of new fungal genomes as they have been sequenced (Sivashankari and Shanmughavel, 2007) and has also greatly impacted the analysis of genomes from under-represented fungal classes, especially the basal lineages. These advances in knowledge have improved our understanding of the origin and diversity of the fungal kingdom (Stukenbrock and Croll, 2014;

Box 1: Glossary of technical terms

Model organism: a species whose biology has been widely studied at many different levels and from different perspectives, serving as reference or basis of comparison for many other species. A model species often has particular experimental advantages, is usually easy to maintain, has a short generation time, and can breed and be manipulated in a laboratory setting.

Evolutionary genomics: the study of how features or components of a genome change both within and between species over evolutionary timescales, especially along lineages showing the divergence or convergence of interesting phenomes.

Comparative genomics: the direct comparison of the complete genetic material of one organism with that of another, or those of many others, to gain a better understanding of how genomes and species have evolved and to determine the functions of genes, transposable elements, and non-coding regions of genomes.

Omics: a set of methodologies targeting the collective qualification and quantification of pools of biological molecules, including genomic DNA (genomics), RNA (transcriptomics), proteins (proteomics), and metabolites (metabolomics), which translate into the structure, function, and regulation of an organism at different levels of dynamics.

Wang et al., 2016d; Dornburg et al., 2017; Stajich, 2017; Chang et al., 2021). Here, we review recent advances in comparative omics in the class Sordariomycetes, focusing on how model species in this class have contributed to evolutionary biology and ecology, and how this class can be developed as a template for obtaining novel and extended insights concerning the fungal kingdom.

Clusters of well-studied models with Big Data to bridge large gaps in evolutionary genomics

One of the main drivers of progress in large-scale comparative genomics research on fungi has been the fungal research core at the Joint Genome Institute, where the 1000 Fungal Genomes Project has been pursued (Grigoriev et al., 2011; Grigoriev et al., 2014). Investigation of genomics and available transcriptomic data for a highly inclusive sample of fungal genomes has revealed that evolutionary convergence may occur across large phylogenetic distances (Merényi et al., 2020). Comparative genomics in mushroom-forming fungi has produced insights into the evolution of mycorrhizal symbiosis, wood-decay mechanisms, and morphological development (Riley et al., 2014; Nagy et al., 2016; Miyauchi et al., 2020; Sánchez-García et al., 2020; Virágh et al., 2022). For example/instance, using comparative genomics to examine a more focused group within the Ustilaginaceae, researchers have reported that the gain and loss of effector genes, including orphan and lineage-specific selected genes, are probably the most important determinants of the host specificity of smut fungi (Benevenuto et al., 2018).

Widespread advances in omics technologies, such as genomics, transcriptomics (whole-genome RNA expression profiling), proteomics (genome-wide study of proteomes), and even metabolomics, have further enabled the study of evolutionary genomics at an extraordinarily detailed molecular level. A key advantage of these technologies is their ability to provide a more granular understanding of evolution within gene families and functional groups, and to elucidate the roles that these families and groups play in large molecular processes. An example of an omics-based integrative approach is the demonstration of the presence of highly conserved class-dependent sugar metabolism pathways (Li et al., 2022). Such analyses, in addition to substantiating the taxonomic placements of the species involved, can also assist in providing guidelines concerning the challenges and limitations of transferring metabolic pathways between species for industrial applications. In the course of analysis of Bayesian networks derived from transcriptomics data on *Neurospora* and *Fusarium* (Nectriaceae, Hypocreales), dispensable chromosomes related to pathogen specificity, very large effector classes, and the dynamic regulation of meiotic silencing and transcription factor networks during sexual reproduction were identified in these models (Wang et al., 2014; Trail et al., 2017; Wang et al., 2018a). A proteogenomics study on the model *Sordaria macrospora* (Sordariaceae, Sordariales) identified new genes and previously unknown posttranscriptional modifications (Blank-Landeshammer et al., 2019). As omics data from different species is

increasingly accumulated, the evolutionary integration of multiple models will be essential in analyzing genomic divergences between distinct fungal clades. Computational approaches to the processing of large-scale omics data and data integration methods that focus mainly on a single model are continuously being developed (Winkler, 2020; Duruflé and Déjean, 2023; Shave et al., 2023; Vasaikar et al., 2023). In addition, systems biological approaches—especially those that refine networks *via* statistical modeling for large datasets with a small cohort size—have proved to be useful in the interpretation of omics data (Culibrk et al., 2016; Karahalil, 2016; Maghuly et al., 2022). These techniques can improve and ease the analysis of “unmatched” multimodal data and provide enhanced overall performance compared with traditional methods that typically do not enable the visual and quantitative or semiquantitative interpretation of gene–gene interactions.

To the apt phrase “Nothing in biology makes sense except in the light of evolution” (Dobzhansky, 1973) can be added “Nothing in evolution makes sense except in the light of genomics” (Shapiro, 2016). Evolutionary biologists have benefited from advances in diverse aspects of the sciences, and omics methods have become the latest powerful toolbox that can be used to clarify the workings of novel features that have hitherto been inexplicable until the last few decades (Shapiro, 2016). Proceeding from genic phylogenies to phylogenomics, the evolutionary histories of major known lineages in fungi have been successfully and robustly resolved, enabling the retrospective tracking of molecular events along the history of fungal evolution (Nowrousian, 2014; Heitman et al., 2017; Pöggeler and James, 2023). In another example, the *Neurospora* community has produced a systematic gene disruption strain collection (“the *Neurospora* knockout collection”), targeting nearly the entire genome, and made it available to the public. There are now over 11,000 gene-disrupted strains available at the Fungal Genetics Stock Center (Colot et al., 2006; Dunlap et al., 2007; Collopy et al., 2010). Meanwhile, advances in gene-manipulation techniques, especially the recently developed CRISPR-Cas9 gene editing system, have dramatically improved our ability to identify the causal mechanisms that link genes and phenotypes in diverse species. Many functional gene groups and metabolic pathways have also been subjected to systematic gene manipulations in various fungal species, and conserved metabolic pathways and key elements in development regulatory circuits have been identified in individual fungal species, many of which are model species. Thus, the progress made in the areas of fungal phylogeny, genomics, and functional genetics has provided a strong foundation for the use of a broader, multispecies model to address the mechanistic bases of complex traits in a diverse range of organisms within particular environmental, developmental, social, and/or genomic contexts.

Sordariomycetes is a large class exhibiting broad diversity, accumulated over time

Advances in genome sequencing and omics data analysis have facilitated further comparative studies, which have offered new

insights into fungal evolution (Nagy et al., 2016; Merényi et al., 2020; Miyauchi et al., 2020; Sagita et al., 2021; Sierra-Patev et al., 2023). Examples covering a wide range of significant discoveries made through comparative genomics include to resolve evolutionary histories among early fungal lineages (James et al., 2020), to propose an important molecular mechanism associated with flagellated zoospores in the chytrid fungus *Blastocladiella emersonii* (Blastocladiaceae, Blastocladales; Galindo et al., 2022), to identify horizontal transfer of a large and toxic secondary metabolic gene cluster between *Podospora* and *Aspergillus* (Slot and Rokas, 2011), to reveal mobile pathogenicity chromosomes in *Fusarium* (Ma et al., 2010), and to discover a small genome of a species within the mycoparasitic genus *Escovopsis* (Hypocreaceae, Hypocreales) that rely on their hosts for some key cellular functions (de Man et al., 2016). This progress has prompted efforts to identify missing links, e.g., non-model species, which can be sequenced and manipulated to address specifically defined biological and evolutionary questions (Russell et al., 2017). In addition to providing annotation references for non-model species, a set of model species forms the phylogenetic backbone for a better understanding of the ways in which species have evolved at the genomic level. Comparative genomic analysis, aided by the inclusion of well-studied models, identifies divergence in genome content and structure, gene copy number and sequence, gene synteny, and evolutionary history, and also identifies mobile elements and non-coding regions in non-model genomes (Sunnérhagen and Piskur, 2006; Sivashankari and Shanmughavel, 2007).

Fungi in the class Sordariomycetes exhibit high levels of ecological diversity. They include saprotrophs on decaying substrates as well as endophytes and parasites on plants, insects, vertebrates, and even fungi. They are also capable of producing a diverse array of secondary metabolites under different conditions (Charria-Girón et al., 2022). The profiles of the secondary metabolites produced can be closely correlated with their phylogeny. However, genomic and phylogenomic analyses strongly suggest that even the genomes of well-studied species harbor secondary metabolite biosynthesis gene clusters, the structures and functions of which have yet to be determined (Atanasov et al., 2021). It has been suggested that the ancestral status of the ecology for Sordariomycetes is saprotrophic or parasitic (Zhang et al., 2006; Zhang and Wang, 2015). In fact, Sordariomycetes is one of the largest classes within the Ascomycetes, and, along with progress in molecular phylogeny, the systematics of this class has been revised multiple times, reflecting the complexity of evolutionary histories within the class. In the most recent comprehensive review of the class (Figure 1A), a total of seven subclasses, 45 orders, 167 families, and 1,499 genera (with 308 genera *incertae sedis*) are recognized within Sordariomycetes, representing dramatic diversity in ecology and developmental biology (Maharachchikumbura et al., 2015; Maharachchikumbura et al., 2016; Hyde et al., 2020). Among the Sordariomycetes, there are several species that serve as classic models of filamentous fungi in the genera, such as *Neurospora* and *Podospora* (Podosporaceae, Sordariales); *Sordaria* species for developmental genetics; *Fusarium*, *Magnaporthe* (Magnaporthaceae, Magnaporthales), and the *Cordyceps* (Cordycipitaceae, Hypocreales) species for pathogenesis; and *Chaetomium* (Chaetomiaceae, Sordariales) and *Trichoderma*

(Hypocreaceae, Hypocreales) species for metabolics and fermentation.

The need for the evolutionary integration of multiple models

Classic genetic models

Among the several Sordariomycetes fungi that have been utilized as models, *N. crassa* is perhaps the most widely examined, and was the first multicellular fungus to have its genome sequenced and annotated (Borkovich et al., 2004; Dunlap et al., 2007). It is a non-pathogenic saprotroph that grows vegetatively and can be cultured in most defined and natural media, and, as such, remains a widely used model that facilitates research across a worldwide community, which has addressed a broad range of questions in general and fungal biology (Perkins, 1992). The species has a distinguished history, including its role in the formulation of the “one gene—one enzyme” hypothesis proposed by Beadle and Tatum (1941).

Closely related to *N. crassa*, *N. tetrasperma* is pseudohomothallic and employs a mating strategy that combines selfing with occasional outbreeding (Merino et al., 1996; Sun et al., 2017). *N. tetrasperma* provides a model for study of both the genomic and the evolutionary consequences of asexuality and inbreeding (Sun et al., 2012; Samils et al., 2013; Idnurm et al., 2015). These consequences cannot be studied in obligate outbreeders alone, such as *N. crassa* and other popular genetic models. A pangenomic analysis of 92 genomes from eight phylogenetically and reproductively isolated lineages of *N. tetrasperma* has provided experimental evidence for the role of introgression as a mechanism for the maintenance of mating-type-determining chromosomal regions (Corcoran et al., 2016).

Interestingly, *Neurospora* species are also known as postfire fungi, as they are often spotted in the forest after a fire, a unique environment shared with certain other pyrophilous fungi (Hansen and Pfister, 2006; Wicklow, 2018), such as *Geopyxis carbonaria* (Pyrenomataceae, Pezizales) and some species of *Pyronema* and *Peziza* (Pezizales). Another easily maintained and genetically manipulated species, *Podospora anserina*, has long been used as a model to study aging, meiosis, sexual reproduction, and heterokaryon formation in fungi (Philipp et al., 2013; Silar et al., 2019; Hartmann et al., 2021a; Lelandais et al., 2022). *P. anserina* and *N. crassa* genomes share over 60%–70% similarity in orthologous proteins, making them a powerful pair of models that can be used as references for comparative genomics (Espagne et al., 2008; Paoletti and Saupe, 2008).

Yet another intensively studied developmental genetic model in the Sordariomycetes is *S. macrospora*, with over 100 available developmental mutants (Nowrousian et al., 2010; Nowrousian et al., 2012; Blank-Landeshammer et al., 2019; Teichert et al., 2020). Similar to species of *Neurospora* and *Podospora*, *S. macrospora* grows quickly under laboratory conditions. The fact that *S. macrospora* is homothallic can be advantageous for sexual and developmental synchronization and for maintenance of a homogeneous genetic background.




FIGURE 1

Systematics, genome availability, and some model species for the Sordariomycetes. (A) Updated systematics of the class Sordariomycetes, composed of seven subclasses (color-coded, alongside the current numbers of family, genus, and genome sequences available in the JGI fungal genome database for each order). The ordinal phylogeny was adapted from a phyloT tree of NCBI taxonomy in the Sordariomycetes and modified in accordance with the latest arrangement by Hyde et al. (2020). (B–G) Comparative morphology of fruiting bodies (perithecia) of model species in six genera: (B) *Magnaporthe*, (C) *Nectria*, (D) *Fusarium*, (E) *Neurospora*, (F) *Podospira*, and (G) *Chaetomium*, including a longitudinal section through the outer fruiting body, a single representative mature ascus with ascospores, and representative paraphyses (sterile hyphae associated with asci). Each perithecia harbors tens or hundreds of asci. All genera, except *Magnaporthe* and *Chaetomium*, forcibly discharge their spores. *P. anserina* produces only four spores per ascus; all others generally produce eight (Trail, 2013; after Trail and Seminara, 2014).

Models for fungal pathogenesis

A small number of Sordariomycetes fungi, including *Fusarium* and *Magnaporthe* species, are models for fungal pathogenesis; over the past decades, *Fusarium* research has advanced our understanding of these fungi. *Fusarium* is a large genus, including some anamorphic fungi, for which sexual development and reproduction has never been observed (O'Donnell et al., 2015). This genus harbors important plant pathogens, producers of a wide range of fungal chemicals, and causal agents of opportunistic mycoses in humans (Suga and Hyakumachi, 2004). *F. graminearum sensu lato* is a species complex whose members cause devastating diseases in small grains and mycotoxin contamination worldwide (Wang et al., 2010; Wang et al., 2011;

Garmendia et al., 2018; Chai et al., 2022; Del Ponte et al., 2022). *F. graminearum sensu stricto* refers to the major disease-causing species in the USA (and a few other places); as this was the subject of one of the first genomics projects (Cuomo et al., 2007), it has since been frequently used in many comparative studies (Khan et al., 2020; Rampersad, 2020; Teli et al., 2020). This fungus has been intensively studied as a cosmopolitan model for fungal pathogenesis of crop plants. The largest source of inoculum for disease outbreaks originates from the primary inoculum, airborne sexual spores (forcibly discharged from fruiting bodies), with aboveground asexual spores (mainly splash-dispersed) providing a local secondary inoculum (Ingold and Dring, 1957). For this reason, the species *F. graminearum* has become a model for the study of perithecia development on host plants (Kim et al., 2022)

and the mechanism of forcible ascospore discharge (Trail et al., 2005; Trail, 2007; Cavinder et al., 2012; Trail and Seminara, 2014; David et al., 2016). Recently, *F. graminearum* has become the first plant-pathogenic fungus documented to produce biofilms in association with plant colonization (Shay et al., 2022). Although fungicide applications are the main source of control over the disease, there are no strongly resistant varieties of wheat or barley available. In addition to being infamous crop pathogens, *Fusarium* spp. are also considered to be severe threats to human health and are listed in the WHO fungal priority pathogens list to guide research, development, and public health action (World Health Organization, 2022; Rodrigues and Nosanchuk, 2023). As in the case of *Neurospora* spp., comparative genomic analysis of *Fusarium* spp. has also been instrumental to the field's understanding of the evolution of sexual development (Trail et al., 2017). One example is the identification and analysis of horizontally transferred lineage-specific genomic regions related to pathogenicity (Ma et al., 2010).

Magnaporthe species cause devastating crop diseases. *M. grisea*, the causal agent of rice, was sequenced in 2005 (Dean et al., 2005). Rice blast causes significant economic loss: worldwide, enough rice to feed tens of millions of people is destroyed by rice blast every year (Kumar et al., 2020). Closely related to *M. grisea*, *M. oryzae* attacks wheat, causing wheat blast, an emergent fungal disease that probably evolved through a series of “host jumps” and that possesses core chromosomes and mini-chromosomes with distinct evolutionary histories (Hossain, 2022). A genomic surveillance study of over 500 strains from diverse geographic regions and host types has revealed the adaptation and development of fungicide resistance in a pandemic clonal lineage of *M. oryzae* (Latorre et al., 2023; Rhodes, 2023). Infections occur when fungal spores land on and invade wheat leaves using an appressorium—a specialized infection cell formed during spore germination. The process of the development of appressoria has recently been investigated intensively using genome-wide analysis (Lv et al., 2022; Wang et al., 2022a; Fan et al., 2023; Miguel-Rojas et al., 2023; Qian et al., 2023; Rogers and Egan, 2023). Diverse ecologies have been reported for species of the Magnaporthales, ranging from saprotrophs to cereal pathogens to likely root endophytes (Luo et al., 2015). The Sordariomycetes also include diverse entomopathogenic fungi, including the “zombie” fungus genus *Cordyceps* and the species of *Beauveria* (Cordycipitaceae, Hypocreales) and *Metarhizium* (Cordycipitaceae, Hypocreales), commonly used as biopesticides of pathogenic insects and arthropods. Omics-based investigation of these fungi is expanding rapidly (Zheng et al., 2011; Xiao et al., 2012; Pattemore et al., 2014; Valero-Jiménez et al., 2016; Wang et al., 2016a; Chen et al., 2019; Kato et al., 2022; Thananusak et al., 2022; Liu and Dong, 2023). Interestingly, some fungi, such as *Metarhizium* species, are capable of attacking hosts in both the plant and the animal kingdoms.

In addition to plant, arthropod, and human hosts, the Sordariomycetes also include a unique pathogenic group, mycoparasites, capable of parasitizing other fungi, including both pathogenic and economically beneficial species. These include species such as *Clonostachys rosea* (Hypocreales; Bionectriaceae), *Calcarisporium cordycipiticola* (Hypocreales; Calcarisporiaceae), *Lecanicillium fungicola* (Hypocreales; Cordycipitaceae),

Syspastospora parasitica (Hypocreales; Hypocreaceae), and *Escovopsis weberi* (Hypocreales; Hypocreaceae) (Posada et al., 2004; Berendsen et al., 2010; de Man et al., 2016; Sun et al., 2020; Liu et al., 2021). Over the years, significant attention has been given to the study of mycoparasites belonging to the genus *Trichoderma*, which contains many species of significant value to agriculture and industry, and the genomics of which have been comprehensively examined (Schalamun and Schmoll, 2022). *Trichoderma reesei* is a unique species among other industrial fungi (Seidl et al., 2009; Druzhinina et al., 2011). *T. reesei* (teleomorphic; *Hypocrea jecorina*) is a known industrial cellulolytic enzyme producer, making it a model system for the study of the regulatory mechanisms of plant cell wall-degrading enzymes; this work contributes to the biofuel and agricultural waste industries. *T. harzianum*, *T. asperellum*, and *T. asperelloides* are among the most thoroughly analyzed *Trichoderma* spp. and have been incorporated in biocontrol strategies (Druzhinina et al., 2011). These biocontrol species and other *Trichoderma* strains have been demonstrated to successfully control plant diseases by stimulating plant growth and development, increasing plant resistance to biotic and abiotic stresses, and directly parasitizing phytopathogenic fungi (Zeilinger and Omann, 2007; Atanasova, 2014; Guzmán-Guzmán et al., 2019; Poveda, 2021; Di Lelio et al., 2023; Woo et al., 2023). Moreover, *T. reesei* and *T. asperellum* are considered to be models for the industrial production of various secondary metabolites (Mukherjee et al., 2013; Schalamun and Schmoll, 2022) and biofungicides due to their mycoparasitic properties, which benefit plants challenged by other fungal pathogens (Kubicek et al., 2011).

The first full *Trichoderma* genome analyzed was that of *T. reesei* (Martinez et al., 2008). *T. asperellum* and *T. asperelloides* (along with others) were analyzed later (Berka, 2017; Druzhinina et al., 2018; Gortikov et al., 2022a), well after the genomes of *Neurospora* and *Fusarium* species (Druzhinina et al., 2018). The recently sequenced genome of *T. asperelloides* contributed to the determination of its correct taxonomic identity, in contrast to its previous identification as the closely related *T. asperellum*, a model for biological control (Podder and Ghosh, 2019; Gortikov et al., 2022a). Currently, there are more than 90 *Trichoderma* genomes that have been sequenced and released, making it among the most sequenced genera in the class; this is an achievement that can be attributed to a combination of industrial and academic efforts. Comparative genomic insight into *Trichoderma* spp. is fairly recent. However, insights into the core genome of this genus, potential heterothallic reproduction, and the abundance of species-specific orphan genes (Kubicek et al., 2019) have provided a strong foundation for further class-based comparative genomics with other Sordariomycetes. One species of Sordariomycetes, *Calcarisporium cordycipiticola* (Calcarisporiaceae, Hypocreales), is a mycoparasite of *Cordyceps militaris*; mycoparasitism causes devastating diseases of fruiting cultivation. Phylogenomic analysis has highlighted the fact that *C. cordycipiticola* was evolutionarily close to its host *C. militaris*, and that they diverged after a split with the *Trichoderma* genus. Comparative genomic and transcriptome analyses have provided insights into the origin of the pathogen and the mycoparasitic interactions of two species from sister families (Liu et al., 2021; Liu and Dong, 2023).

Promising new models for fungal biology

Dramatic divergences in the development and biology of Sordariomycetes can be observed even among closely related species; thus, the class offers excellent systems for assessment of the molecular genetic basis of the evolution of these species. Among these promising systems, *Chaetomium* spp. fungi exhibit a frequently homothallic lifestyle, a general lack of asexual reproduction of conidia, enriched secondary metabolism synthesis, and an ability to thrive in highly humid environments (Wang et al., 2016b; Zámocký et al., 2016; Wang et al., 2019a). All of these traits are highly divergent in the genus, rendering these species especially informative regarding rapid evolution in a comparative evolution context. The *Chaetomium* spp. are cosmopolitan saprotrophs and endophytes that are capable of dominating diverse ecological niches, including some extreme environments, evoking interest in their genomic adaptation for survival and dispersal. Species of *Chaetomium* are common contaminants in indoor environments and are considered to be health hazards (Andersen et al., 2011; Miller and McMullin, 2014), causing symptoms of rhinitis and asthma when they infect humans (Mackenzie, 1979; Vesper et al., 2007; Hassett et al., 2009; Green et al., 2014). *C. globosum* is a model for the industrial production of secondary metabolites, and is also generally considered to be a species complex (Asgari and Zare, 2011; Wang et al., 2016b). *Chaetomium* species probably shared thermophilic ancestors, consistent with the thermophilic basal nature of its most diverged lineage (Zámocký et al., 2016). The genome sequence of *C. globosum* was made publicly available in 2015 (Cuomo et al., 2015). Since then, the taxonomy and phylogenetics of the genus have been undergoing reassessment and revision (Wang et al., 2022c).

These models represent some of the diverse genera within the Sordariomycetes that have been studied intensively. The availability of many genome sequences that are closely related to established and emerging model species represents a substantial contribution to analyses of the ecological traits of these organisms and their close relatives, which can in turn provide substantial insight. Currently, there are 570 Sordariomycetes genomes (covering 136 genera or high ranks, with more than 330 identified species and 75 unidentified species) in the JGI fungal genome database (mycocosm.jgi.doe.gov/), in addition to the comparative tools available via that portal (Figure 1A). Among these genomes, 189 cover more than 60 species in four genera, including 81 *Fusarium* and 94 *Trichoderma* species. Many additional Sordariomycetes genomes are available via the NCBI (National Center for Biotechnology Information), with 3,574 genomes (nearly one-third of the total of 11,083) being published and drafted ascomycetes genomes; these include 1,359 *Fusarium*, 117 *Neurospora*, 110 *Trichoderma*, and 14 *Chaetomium* genomes. Many of these genomes are for species that are culturable, produce simple morphologies, and are often either pathogenic or economically important. The ecological and industrial relevance of pathogenic members of Sordariomycetes and those producing valuable fungal products has been a major driver of the extensive sequencing of many non-model genomes in this fungal class. Some of these non-model species are emerging as “new” models and,

unlike the classic models that have undergone intensive investigation in biology and genetics, established during the pre-genomics era, these species are often studied under a reverse approach, with investigation originating in genomics and then extending to genetics and the analysis of biological traits. For example, the genomes of a large number of species of Xylariales have recently been sequenced, and comparative genomics has revealed genetic changes that are probably associated with bioactivities, diverse lifestyles, environmental adaptations, and selective pressure in these “mystical”-looking fungi (Robinson et al., 2020; Wibberg et al., 2020; Franco et al., 2022; Fricke et al., 2023). At the same time, the annotation of classic model genomes, such as those of *Neurospora*, *Podospira*, and *Sordaria* genomes, has undergone continuous improvement, in part on the basis of information obtained from newly sequenced genomes of the same or closely related species (Teichert et al., 2014; Gladieux et al., 2015; Blank-Landeshammer et al., 2019; Teichert et al., 2020; Lelandais et al., 2022; Rodriguez et al., 2022; Vittorelli et al., 2023). These accumulating data continue to contribute to a new era of evolutionary genomics and transcriptomics at further taxonomic levels within the Sordariomycetes class.

New details and new omics data on the Sordariomycetes continue to accumulate in an accelerating manner

Representatives of the major fungal lineages on the tree of life have now had their genomes sequenced, and the better-annotated genomes of model species have been used as references. Thus, model species can function as genomic information hubs to bridge the gaps between these well-studied organisms and the relatively novel characteristics of non-model species (Wang et al., 2018a; Teichert et al., 2020). Associations between the various different ecological and developmental models are of special interest, and within Sordariomycetes, it has been suggested that there is a strong correlation between the evolution of senescence and ephemeral substrate usage (Geydan et al., 2012). Sexual reproduction in the Sordariomycetes involves the formation of perithecia, a round-to-flask-shaped structure with a pore through which meiotic spores are discharged. The morphological details of the perithecia can differ dramatically among species within the class (Figures 1B–G). Sexual development has been examined in *Neurospora*, *Podospira*, and *Sordaria* via genomics- and transcriptomics-based analyses for over a decade (Bidard et al., 2011; Ellison, 2011; Dirschnabel et al., 2014; Teichert et al., 2014; Idnurm et al., 2015; Corcoran et al., 2016; Xie et al., 2017; Grognet et al., 2019; Shen et al., 2019; Hartmann et al., 2021b), with emphasis on the functions and evolution of mating loci, secondary metabolism, and stress responses associated with sexual reproduction.

Evolutionary developmental biology

Divergence in gene expression has long been considered to play a critical role in developmental adaptations during organismal

evolution (King and Wilson, 1975; Carroll, 2005; Hodgins-Davis et al., 2015; Diaz et al., 2023). The incorporation of divergence in sexual morphology within comparatively closely related taxa could improve the precision and accuracy of reconstructions of ancestral gene expression, yielding expansions to the number of associations between shared morphologies and shared transcriptional profiles. Comparative genomics and evolutionary transcriptomics focusing on Sordariomycetes models has revealed the genes and regulatory networks that are critical for sexual development in these fungi, including non-coding sequences and genes that were previously uncharacterized (Trail et al., 2017; Kim et al., 2018; Kim et al., 2019; Lütkenhaus et al., 2019). In addition, novel genes that affect perithecial development have been identified in *M. oryzae* and *N. crassa*, which have diverged functionally and transcriptionally from their orthologous counterparts in *F. graminearum* (Kim et al., 2022). These novel genes are predicted to be “young” (i.e., present only in a recently diverged clade of species) and tend to be involved in lineage- or species-specific functions. Phenotypic study guided by comparative genomics among three distantly related filamentous fungi has identified two genes (predicted to be chromatin modifiers) that play roles in the sexual development of *S. macrospora* (Lütkenhaus et al., 2019).

Evolutionary ecology

Comparative genomics approaches using these model species have also been pursued in order to understand the evolution of diverse ecologies among the Sordariomycetes, especially those that are related to pathogenesis (Figure 2). Among the pathogenic Sordariomycetes, species of *Fusarium* have been intensively studied for their diverse genome structures (Brown and Proctor, 2013; Ma et al., 2013; Zhang, 2019; Mir et al., 2023). Research on entomopathogenic Sordariomycetes, represented by *Cordyceps*, *Beauveria*, and *Metarhizium* species, has been greatly enhanced by the availability of genomes of these species (Gao et al., 2011;

Pattemore et al., 2014; Valero-Jiménez et al., 2016; Kato et al., 2022; Liu and Dong, 2023; Sant Anna Iwanicki et al., 2023). *Neurospora* species have been examined as metabolic models of the biodegradation of cellulose and other plant organics (Feldman et al., 2019; Liu et al., 2020; Wu et al., 2020; Huberman et al., 2021), as well as providing some of the most tractable models for the circadian clock and fungal responses to environmental factors, such as light and temperature (Wang et al., 2016c; Dekhang et al., 2017; Wang et al., 2018b; Kelliher et al., 2020; Kelliher et al., 2023). Based on transcriptomics data from *N. crassa* grown on five different crop residues, researchers have discovered roles for a sporulation regulator *rca-1* in lignocellulose production (Wang et al., 2015). Additional roles in allorecognition have also been identified for *cwr-1*, a putative chitin polysaccharide monooxygenase, in *N. crassa* (Detomasi et al., 2022). Three studies have investigated the associations between secondary metabolite clusters (SMCs) and reproduction in two fungal models, *N. crassa* and *C. globosum*, and reported on the activities of 20 and 24 SMCs in these models, respectively—a manageable set for separate investigation of the roles of SMCs across the life cycle of these two models (Wang et al., 2019a; Wang et al., 2019b; Wang et al., 2022d). Bayesian regulatory networks have been reconstructed using transcriptomic data for *N. crassa* and *C. globosum*. These networks exhibit divergences in associations among genes associated with conidiation and heterokaryon incompatibility between *N. crassa* and *C. globosum*, supporting the theory that there is an evolutionary history of loss of conidiation in the latter, putatively due to unfavorable combinations of heterokaryon incompatibility in homothallic species. A recent study focused specifically on conidial germination in two model pathogens, *F. graminearum* and *M. oryzae*, on artificial medium and on hosts, as conidial germination represents the key stage in the initiation of fungal attack (Miguel-Rojas et al., 2023). The authors’ analyses revealed new and important aspects of early fungal ingress in *F. graminearum* that can form the basis for improvement of antifungal strategies, including a comparative gene expression




FIGURE 2

Phylogenetic relationships among seven closely related taxa in the Hypocreales, inferred with concatenated RPB1 and RPB2 protein sequences using a Bayesian approach (Ronquist et al., 2012) to investigate the repeated evolution of parasitism on specific host types. These species include fungi that attack insects (entomopathogenic species), plants (phytopathogenic species), other fungi (mycopathogenic species), and non-pathogenic species, including endophytic and experimental model species. The classification follows Zhang et al. (2006) and Geiser et al. (2006), and the calibration points used are those of Taylor and Berbee (2006). Species in which genetic manipulations have been well developed are marked with a delta symbol, “Δ”.

finding that toxisomes are not fully functional until after penetration of the host cells. Based on a study using near-synchronous germinating cultures of *T. asperelloides* (a strain used in biocontrol that was formerly identified as *T. asperellum*), it has been reported that the transcript abundance of half of the annotated genome is reduced during early conidial germination (Grotikov et al., 2022b). In this study, it was also discovered that the expression of the chitin synthase and glucan elongase families is significantly increased during germination in the presence of a basidiomycete host, *Rhizoctonia solani* (Ceratobasidiaceae, Cantharellales), indicating that host recognition can occur during the early stages of mycoparasite development (Grotikov et al., 2022b).

Genome evolution

Advances in fungal genomics have produced new insights regarding genome evolution in the kingdom. Mycovirus research has greatly benefited from the availability of a large number of fungal genomes and the development of comparative genomics (Kotta-Loizou, 2019; Myers et al., 2020). Mycoviruses have been the focus of biocontrol against fungal pathogens (Yu and Kim, 2020; Schalamun and Schmoll, 2022). More than 10 different families of viruses have been isolated from fungal hosts, including double-stranded RNA (dsRNA), positive-sense and negative-sense single-stranded RNA, and circular single-stranded DNA (ssDNA) viruses (Applen Clancey et al., 2020). The presence of viruses in *Fusarium* and *Trichoderma* spp. has been particularly well documented (Yun et al., 2016; Li et al., 2019a; You et al., 2019; Yu and Kim, 2020; Wang et al., 2022b). A recent study investigated the presence of viruses in multiple *Neurospora* species and identified various RNA viruses from *N. crassa* and other *Neurospora* spp. using transcriptomics data. The study further established *N. crassa* as a virus host model for use in the study of virus–host interactions and virology in fungi (Honda et al., 2020). The authors also demonstrated that *N. crassa* relies on transcriptional and posttranscriptional regulation to restrict virus replication for at least some of its genes.

Posttranscriptional regulation is another intriguing genome-wide editing mechanism, but how has such a widely distributed mechanism evolved and been maintained across the diverse fungal genome? One example of posttranscriptional regulation is A-to-I editing, which has recently been discovered to be associated with sexual development in filamentous fungi (Bian et al., 2019). Some of these A-to-I-editing filamentous fungi are Sordariomycetes, including *N. crassa* (Liu et al., 2017), *S. macrospora* (Teichert et al., 2017), and *F. graminearum* (Liu et al., 2016). In *M. oryzae*, N⁶-Methyladenosine (m⁶A) RNA methylation, the most common modification of RNA at the post-transcriptional level in eukaryotes, has been found to be important for various aspects of fungal biology, such as vegetative growth, conidiation, and pathogenicity (Shi et al., 2019). Although disruption of m⁶A factors does not affect sexual development in *M. oryzae* and *F. graminearum*, overexpression of the m⁶A writer (the ortholog of the gene coding for *IME4* in yeasts) causes delayed sexual development in *F.*

graminearum, suggesting that m⁶A modification may alter developmental processes in filamentous fungi (W. Kim, unpublished). RNA modification, such as m⁶A, can be detected using direct RNA sequencing, whereby the modification sites can be profiled at nucleotide resolution (Leger et al., 2021). All these new discoveries using the Sordariomycetes models can be expected to evoke interest in exploring model fungal clades that can comprehensively illuminate these fundamental features.

Climate change

The application of evolutionary approaches to genomic data from fungi may also help us to understand some impacts of climate change. Fungi are widely distributed and abundant in almost all ecosystems—even in extreme environments—and have been referred to as “climate warriors” in recognition of the important ecological roles that they play in the recycling of carbon, nitrogen, and other nutrients (Stajich, 2017; Averill et al., 2018; Tiquia-Arashiro and Grube, 2019; Pöggeler and James, 2023). Their presence and activity are considered to be useful indicators in relation to long-term climate change due to their sensitivity to surrounding conditions, such as changes in light, temperature, humidity, ROS elements, host type, and organic and inorganic pollution (Gadd et al., 2007; Frankland et al., 2009; Rodriguez-Romero et al., 2010). Several studies have reported on the impacts of climate change on fungus–host interactions, fungal distribution, fungal disease prevalence, and the production of fungal products (Frankland et al., 2009; Merrild, 2013; Botana and Sainz, 2015; Tiquia-Arashiro and Grube, 2019; Charters, 2020; Meyer et al., 2020; Frias-De-León et al., 2022; Miranda-Apodaca et al., 2023). The wide distribution and ecological diversity of Sordariomycetes fungi contributes to their frequent appearances in reports on the impact of climate change on fungal diversity, which occur especially frequently in climate-sensitive regions or populations of organisms that are closely associated with fungi.

Fungal pathogens that infect economically important crops, such as species of *Fusarium* (Vaughan et al., 2016) and *Magnaporthe* (Qiu et al., 2022), are of special concern. Epidemics of fungal pathogens can be enhanced by particularly wet and warm environments occurring during the flowering seasons of some crops. Remote sensing/geographic information system (GIS) technology has the potential to monitor snowline elevation, average temperature, precipitation, and sunshine hours in terms of their relationship with the product of *Cordyceps sinensis* (syn. *Ophiocordyceps sinensis*), which is also known as “yartsa gunbu” or “Dong Chong Xia Cao” (“winter worm summer grass”) in traditional Tibetan and Chinese medicine (Zhu et al., 2017). A comprehensive collection dataset and an ensemble species distribution modeling method have recently been used to investigate whether and how climate change affects the distribution of *C. sinensis*, and to predict potential shifts in the range of the fungus in the medium term of approximately 50–70 years in response to climate change (Yan et al., 2017; Li et al., 2019b). It has also been proposed that *Trichoderma* spp. could be utilized as “plant savers” in the face of climate change, on the basis

of their several traits and genes that confer beneficial effects on crop plants (Kashyap et al., 2017). A limitation of these studies—with regard to their potential as a basis for omics-based analyses—is that they have thus far mainly addressed the ecological and diversity aspects of the global ecosystem. Hence, substantial fungal genomic resources that may illuminate potential responses to climate change have yet to be identified.

This would be a timely moment to revisit the possibility of developing *N. discreta*, along with other Sordariomycetes, as a model species to test the idea that adaptation to warm temperatures could lead to more efficient carbon metabolism (Romero-Olivares et al., 2015). To best adapt to their environments (often microenvironments), fungi have evolved elegant mechanisms for sensing and quickly responding to changes, and some of their genetic elements may be capable of high rates of evolution (e.g., Wang et al., 2016c; Chandra Nayaka et al., 2021; Kusch et al., 2023). Given the ecological and phenotypic diversity among closely related lineages within Sordariomycetes, further study and comparison of their potential for rapid genomic evolution would clarify their potential roles as models to enhance our understanding of long- and short-term climate and environmental impacts on fungi, and associated plant and animal health. As a motivating example, the relative abundance of *Fusarium* spp. has risen almost fivefold alongside the course of global warming (Delgado-Baquerizo et al., 2020).

We have only reviewed a very small portion of the recent comparative genomics and transcriptomics literature focusing on members of the class of Sordariomycetes, but these clearly illustrate the potential scientific rewards of comparing multiple genomes in different experimental settings to understand the evolution of the relevant organisms or traits and the genetics of select biological or ecological phenomena. However, the insights obtained on the basis of genomics/transcriptomics into trait evolution and into the associations with the genetics of various traits are still far from complete. Two possible factors that may significantly contribute to this gap are our underdeveloped understanding of the evolution of gene expression, and our lack of a system for integrating data from trait evolution and genetics that have already been (and can now be) intensively studied. Therefore, a large system encompassing well-studied model species, such as the Sordariomycetes, is a promising platform for the development of tools and the implementation of strategies to develop a systematic genetic understanding of trait evolution.

Challenges and prospects for Sordariomycetes model systems in the omics era

Members of Sordariomycetes are model species and serve as key references for the annotation of newly sequenced non-model genomes. Nevertheless, investigation into the genomics and transcriptomics of the Sordariomycetes continues to reveal deeper layers of mysteries, including yet-to-be-resolved complexity, variation, and surprises. These efforts often encounter difficulties

owing to the inherent limitations of currently available genomics/transcriptomics and bioinformatics technologies. Such limitations can be overcome, in part, by community efforts to invest in improving the annotation of the model genomes. Phylostratigraphy is a methodology that provides statistical descriptions of the origins of genes in a genome through homology-based searching across the species phylogeny (Domazet-Loso et al., 2007) with much more inclusive phylogenetic representation of genomes; the use of this approach has significantly reduced the number of *N. crassa* lineage-specific genes identified, and has demonstrated that they are clustered and aggregated toward the telomeres of each chromosome (Wang et al., 2022e; Wang et al., 2023). A recent examination of transcription factors in fungal genomes using homology searches identified a large number of mis-annotations as a result of sequencing errors and the mis-assembly of reference genomes (Mayer et al., 2023). However, historical genomic mis-annotations have been resolved only in the case of a few genomes, mainly because of the transient nature of the financial support for scientific advancements and accompanying technical difficulties. Usually, comparative genomic analyses focus on annotated gene families, functional groups, and specific questions about those genomes; thus, large amounts of the available data have been overlooked and are not likely to be revisited if this type of approach to the research persists. Although we anticipate the continued provision of resources for the analysis of larger numbers of new genomes, there is a greater need for support for the generation and comparative analysis of genomes of closely related organisms, so as to empower the development of model clades in terms of both a higher density of sampling and the generation of higher-quality comparisons and annotations. Clusters of classic models with a set of high-quality “core genomes” will be critical for a broader and more profound understanding of evolution of fungal and organismal diversity.

In addition to high-quality genomics data, there is a need for a solid basis for the design and development of methods for the analysis of high-quality transcriptomics, proteomics, and metabolomics data from an evolutionary perspective (Jaffé et al., 2004; Nesvizhskii, 2014; Naranjo-Ortiz and Gabaldón, 2020; Sarsaiya et al., 2021). For example, proteogenomics—which combines proteomics, genomics, and transcriptomics—has considerably improved the annotation of *S. macrospora* through the identification of over 104 “hidden” proteins and the correction of annotations for over 500 genes (Blank-Landeshammer et al., 2019). A proteogenomic approach was also used to identify the function of transporter protein in *N. crassa* (Rupa et al., 2018). In a recent study focusing on eukaryotic chromatin evolution, *N. crassa* was sampled, along with two yeast models and other eukaryotes, for phylogenetic and proteomic reconstruction (Grau-Bové et al., 2022). In addition to being a model of clock- and/or light-responsive metabolic regulation, *N. crassa* has also served as a platform for the development of genome-scale metabolic models (Chen and Loros, 2009; Dreyfuss et al., 2013; Hurley et al., 2014; Bayram et al., 2019). The Sordariomycetes genomes represent excellent resources for investigation of the dramatic diversity of ecology and biology within the class. Recently, many studies have generated pangenomics and pantranscriptomics data, including on

multiple strains within species, and developed core elements of fungal genomes and transcriptomes (Plissonneau et al., 2018; Badet et al., 2020; Bao et al., 2022; Hyun et al., 2022; Vaughn et al., 2022). However, the power of omics approaches is limited by a lack of knowledge of baseline gene activities and their functional dynamics in natural settings—even in the case of intensively studied biological processes of well-annotated model species. Therefore, one challenge that requires future effort is the integration of data and knowledge among “closely related” species—noting that fungi referred to as closely related are often divergent from a most recent common ancestor dating back to nearly the same period as the most recent common ancestor of mammals (Figure 2).

Even under standard laboratory conditions, genome-level comparisons across closely related species face the need to deal with many unexpected results—even setting aside data heterogeneity resulting from the use of different experimental platforms by different research groups. In studies revisiting the transcriptomics data collected from three *Neurospora* and two *Fusarium* species over the morphologically conserved sexual reproduction process, large portions of the genomes have been found to exhibit divergent dynamics in terms of gene expression profiles (Figure 3). Such divergent activities have also been observed among different age groups of genes, suggesting that much more evolution of gene roles occurs than the conservation of functional groups implied by the current use of tools for functional annotation based on gene families and orthologous groups. One may argue that mRNA abundance may not directly reflect the functional status of the coding gene, meaning that additional well-sampled proteomics data are required to piece the puzzle together. This additional research is surely warranted: there is a lack of models with well-sampled transcriptomics and proteomics data covering key growth and developmental stages under standard laboratory conditions as well as more realistic conditions that better match natural environments. The establishment of a diverse range of comparative omics studies of the Sordariomycetes could enable intriguing and novel cross-class comparisons to be made. In addition, insights obtained from such analyses could well serve as starting points for novel hypothesis-driven experiments that can be carried out with relative ease in historically and newly amenable model species.

A central problem in genetics research is that of deciphering how genomic variation affects the function of genes and results in altered phenotypes. Results from genome-wide association studies (GWASs) with fungal genome data offer insights into the genetic basis of phenotypes or traits of interest. However, increasingly innovative methods will need to be developed to effectively integrate diverse data types and/or sources of information in order to identify functional genes and variants and understand how they shape the relevant phenotypes. These methods will probably include approaches that enable a move from detection of a genetic association signal in a chromosomal region to the identification of trait-associated genes and causal variants, as a step toward understanding the underlying morphological, ecological, or metabolic processes in fungi. Fine mapping, sequencing, functional studies, and other approaches have also been used to find the causal variants involved in complex traits, facilitated by statistical

approaches to the integration of diverse data sources, including transcriptomics data from different cohorts, to interrogate causal relationships between genetic variation and the associated mechanism in fungal biology. The use of systems and synthetic biology approaches would also be a useful and creative way to investigate large-scale genetic and environmental interactions (G×E) within the genomes, especially among different genera that flourish in different environmental conditions. Additional obstacles to modeling in the class include: (1) the imbalanced sampling of genomes within this large and diverse fungal class, given that most of the available genomes are drawn from only four out of 45 orders, namely, Xylariales, Hypocreales, Sordariales, and Glomerellales; (2) the fact that many Sordariomycetes species are symbiotic and have not yet been successfully cultured under laboratory conditions; and (3) a lack of understanding of the ecology and biology of the model species in their natural settings. These challenges may partly be addressed by promoting collaboration among the relevant experts within and beyond our immediate scientific community and by incorporating developments in artificial intelligence (AI) that promise to accelerate genomics research.

On the latter point, the first critical step is the systematic feeding of diverse and high-quality data into AI tools, enabling computational facilitation based on reference data and data mining for applications ranging from genome annotation to systems biology analysis. Researchers have developed many widely used high-throughput computational methods of data analysis, including single-cell sequencing, to provide start-to-finish analytical ecosystems for large-scale omics datasets. These data are generated from experiments involving a wide range of systems that are considered to be of great interest for society (Hotelling et al., 2023). Models using methods such as manifold learning and deep learning have been developed, employing supervised and unsupervised learning approaches to the processing and visualization of data, the development of the field's understanding of biological processes, and the characterization of phenotypic diversity and its underlying causal mechanisms. With their high-quality model genomes accompanied by high-quality omics data, the Sordariomycetes can be established at the core of omics databases, serving as a powerful set of model species for AI training for the analysis of evolutionary genomics, as has been pioneered in certain other disciplines (Guhlin et al., 2017; Misra et al., 2019; Biswas and Chakrabarti, 2020; Ko et al., 2020; Reel et al., 2021). Nevertheless, it is important to note that just as some erroneous gene annotations have been made and perpetuated over time, the quality of AI output, especially the outputs of new AI tools, which will be trained for new data types, will have to be critically monitored to minimize the embedding of historical misinterpretations and misconceptions that could substantially impede downstream science.

Big Data provides us with the opportunity to address existing and new challenges in evolutionary genomics research. As omics data accumulate, the systematic investigation of genome–phenome relationships has expanded from model-driven gene-by-gene studies to data-driven studies of multiple species and multiple functional groups. This review has provided only a smattering of examples that touch on the substantial successes to be achieved by revealing the rules that underlie the relationships between genomes




FIGURE 3

Comparative gene expression, illustrated in the form of (A) a heatmap and (B) a Transcriptome Age Index (TAI) plot (which combines gene phylogenetic age with expression changes to describe possible evo-developmental transcriptomics within a single genotype) for single-copy orthologous genes among five species during 144 h of sexual development. Gene expression data from our previous publications (Sikhakolli et al., 2012; Lehr et al., 2014; Trail et al., 2017) were revisited. TAI (the weighted mean of phylostrata using gene expression intensities of a given gene) was computed for each developmental (ontogenetic) stage following the methods described in previous studies on plant and animal models (Domazet-Lošo and Tautz, 2010; Drost et al., 2015).

and phenomes. We advocate for the adoption of the class Sordariomycetes as a cluster of high-quality and diverse models that will enable extensive scientific advances capitalizing on Big Data in evolutionary genomics and that can be used as a high-level reference and blueprint for the analysis of other classes in the fungal kingdom and beyond. The richness and diversity of its members; their ecological capabilities, lifestyles, and economic importance; the accumulated research history with amenable model species; the abundance of sequenced genomes; the rapidly accumulating diversity of other omics data; and, last but not least, the immense global research community in this domain promise a synergy that will yield extraordinary and exciting scientific outcomes.

Author contributions

ZW, WK, Y-WW, EY, and OY drafted the manuscript. ZW, CD, FT, JT, and OY edited the manuscript. All authors contributed to the article and approved the submitted version.

Funding

The work of ZW, Y-WW, and JT was supported by funding awarded to JT by the National Institutes of Health R01, grant AI146584, and by the National Science Foundation, grant IOS

1457044; the work of EY and OY was supported by funding award BSF-2018712 to OY. The work of CD was supported by the National Natural Science Foundation of China (32272786 and 31872163). FT was supported by the National Institutes of Health, R01 grant AI146584, and the Michigan State University AgBioResearch. The work of WK and FT was supported by the National Science Foundation, IOS 1456482, awarded to FT. WK was partly supported by the Basic Science Research Program through the National Research Foundation of Korea, funded by the Ministry of Education (2019R11A1A01057502).

Acknowledgments

We are grateful to the reviewers for their comments and suggestions, which significantly contributed to the quality of this review. We also thank the Broad Institute, FungiDB, and JGI for making numerous fungi genomic datasets available.

References

- Andersen, B., Frisvad, J. C., Søndergaard, I., Rasmussen, I. S., and Larsen, L. S. (2011). Associations between fungal species and water-damaged building materials. *Appl. Environ. Microbiol.* 77, 4180–4188. doi: 10.1128/AEM.02513-10
- Ankeny, R., and Leonelli, S. (2020). *Model organisms* (Cambridge, UK: Cambridge University Press).
- Applen Clancey, S., Ruchti, F., LeibundGut-Landmann, S., Heitman, J., and Janiri, G. (2020). A novel mycovirus evokes transcriptional rewiring in the fungus and stimulates beta interferon production in macrophages. *MBio* 11 (5), e01534–20. doi: 10.1128/mBio.01534-20
- Asgari, B., and Zare, R. (2011). The genus chaetomium in Iran, a phylogenetic study including six new species. *Mycologia* 103, 863–882. doi: 10.3852/10-349
- Atanasov, A. G., Zotchev, S. B., Dirsch, V. M., and Supuran, C. T. (2021). Natural products in drug discovery: advances and opportunities. *Nat. Rev. Drug Discov.* 20, 200–216. doi: 10.1038/s41573-020-00114-z
- Atanasova, L. (2014). “Chapter 2 - ecophysiology of trichoderma in genomic perspective,” in *Biotechnology and biology of trichoderma*, vol. 25–40. Eds. V. K. Gupta, M. Schmoll, A. Herrera-Estrella, R. S. Upadhyay, I. Druzhinina and M. G. Tuohy (Amsterdam: Elsevier).
- Averil, C., Dietze, M. C., and Bhatnagar, J. M. (2018). Continental-scale nitrogen pollution is shifting forest mycorrhizal associations and soil carbon stocks. *Glob. Change Biol.* 24, 4544–4553. doi: 10.1111/gcb.14368
- Badet, T., Oggenfuss, U., Abraham, L., McDonald, B. A., and Croll, D. (2020). A 19-isolate reference-quality global pangenome for the fungal wheat pathogen *Zymoseptoria tritici*. *BMC Biol.* 18, 12. doi: 10.1186/s12915-020-0744-3
- Bao, J., Wang, Z., Chen, M., Chen, S., Chen, X., Xie, J., et al. (2022). Pan-genomics reveals a new variation pattern of secreted proteins in *J. Fungi (Basel)* 8 (12), 1238. doi: 10.3390/jof8121238
- Bayram, Ö. S., Dettmann, A., Karahoda, B., Moloney, N. M., Ormsby, T., McGowan, J., et al. (2019). Control of development, secondary metabolism and light-dependent carotenoid biosynthesis by the velvet complex of *Genetics* 212, 691–710. doi: 10.1534/genetics.119.302277
- Beadle, G. W., and Tatum, E. L. (1941). Genetic control of biochemical reactions in *Neurospora*. *Proc. Natl. Acad. Sci. U. S. A.* 27, 499–506. doi: 10.1073/pnas.27.11.499
- Benevenuto, J., Teixeira-Silva, N. S., Kuramae, E. E., Croll, D., and Monteiro-Vitorello, C. B. (2018). Comparative genomics of smut pathogens: insights from orphans and positively selected genes into host specialization. *Front. Microbiol.* 9, 660. doi: 10.3389/fmicb.2018.00660
- Berendsen, R. L., Baars, J. J. P., Kalkhove, S. I. C., Lugones, L. G., Wösten, H. A. B., and Bakker, P. A. H. M. (2010). *Lecanicillium fungicola*: causal agent of dry bubble disease in white-button mushroom. *Mol. Plant Pathol.* 11, 585–595. doi: 10.1111/j.1364-3703.2010.00627.x
- Berka, R. M. (2017). Genus-wide genomics of the biomass-degrading and plant-beneficial *Trichoderma*. doi: 10.25585/1487483
- Bian, Z., Ni, Y., Xu, J.-R., and Liu, H. (2019). A-to-I mRNA editing in fungi: occurrence, function, and evolution. *Cell. Mol. Life Sci.* 76, 329–340. doi: 10.1007/s00018-018-2936-3
- Bidard, F., Ait Benkhali, J., Coppin, E., Imbeaud, S., Grognet, P., Delacroix, H., et al. (2011). Genome-wide gene expression profiling of fertilization competent mycelium in opposite mating types in the heterothallic fungus *Podospora anserina*. *PLoS One* 6, e21476. doi: 10.1371/journal.pone.0021476
- Biswas, N., and Chakrabarti, S. (2020). Artificial intelligence (AI)-based systems biology approaches in multi-omics data analysis of cancer. *Front. Oncol.* 10, 588221. doi: 10.3389/fonc.2020.588221
- Blank-Landeshammer, B., Teichert, L., Märker, R., Nowrousian, M., Kück, U., and Sickmann, A. (2019). Combination of proteogenomics with peptide sequencing identifies new genes and hidden posttranscriptional modifications. *MBio* 10 (5), e02367–19. doi: 10.1128/mBio.02367-19
- Boekhout, T., Aime, M. C., Begerow, D., Gabaldón, T., Heitman, J., Kemler, M., et al. (2021). The evolving species concepts used for yeasts: from phenotypes and genomes to speciation networks. *Fungal Divers.* 109, 27–55. doi: 10.1007/s13225-021-00475-9
- Borkovich, K. A., Alex, L. A., Yarden, O., Freitag, M., Turner, G. E., Read, N. D., et al. (2004). Lessons from the genome sequence of *Neurospora crassa*: tracing the path from genomic blueprint to multicellular organism. *Microbiol. Mol. Biol. Rev.* 68, 1–108. doi: 10.1128/MMBR.68.1.1-108.2004
- Botana, L. M., and Sainz, M. J. (2015). *Climate change and mycotoxins* (Berlin, Germany/Boston, MA, USA: Walter de Gruyter GmbH & Co KG).
- Brown, D. W., and Proctor, R. H. (2013). *Fusarium: genomics, molecular and cellular biology* (Norfolk, UK: Caister Academic Press Limited).
- Carroll, S. B. (2005). Evolution at two levels: on genes and form. *PLoS Biol.* 3, e245. doi: 10.1371/journal.pbio.0030245
- Cavinder, B., Sikhakolli, U., Fellows, K. M., and Trail, F. (2012). Sexual development and ascospore discharge in *Fusarium graminearum*. *J. Vis. Exp.* (61), 3895. doi: 10.3791/3895
- Chai, Y., Senay, S., Horvath, D., and Pardey, P. (2022). Multi-peril pathogen risks to global wheat production: a probabilistic loss and investment assessment. *Front. Plant Sci.* 13, 1034600. doi: 10.3389/fpls.2022.1034600
- Chandra Nayaka, S., Hosahatti, R., Prakash, G., Tara Satyavathi, C., and Sharma, R. (2021). *Blast disease of cereal crops: evolution and adaptation in context of climate change* (Switzerland: Springer Nature).
- Chang, Y., Rochon, D., Sekimoto, S., Wang, Y., Chovatia, M., Sandor, L., et al. (2021). Genome-scale phylogenetic analyses confirm *olpidium* as the closest living zoospore fungus to the non-flagellated, terrestrial fungi. *Sci. Rep.* 11, 3217. doi: 10.1038/s41598-021-82607-4
- Charria-Girón, E., Surup, F., and Marin-Felix, Y. (2022). Diversity of biologically active secondary metabolites in the ascomycete order sordariales. *Mycol. Prog.* 21, 1–33. doi: 10.1007/s11557-022-01775-3
- Charters, M. D. (2020). *Climate change impacts on interactions between wheat, aphids, and arbuscular mycorrhizal fungi*. PhD thesis. University of Leeds.
- Chen, C.-H., and Loros, J. J. (2009). *Neurospora* sees the light: light signaling components in a model system. *Commun. Integr. Biol.* 2, 448–451. doi: 10.4161/cib.2.5.8835
- Chen, Y., Wu, Y., Liu, L., Feng, J., Zhang, T., Qin, S., et al. (2019). Study of the whole genome, methylome and transcriptome of *Cordyceps militaris*. *Sci. Rep.* 9, 898. doi: 10.1038/s41598-018-38021-4
- Collopy, P. D., Colot, H. V., Park, G., Ringelberg, C., Crew, C. M., Borkovich, K. A., et al. (2010). High-throughput construction of gene deletion cassettes for generation of

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- neurospora crassa knockout strains. *Methods Mol. Biol.* 638, 33–40. doi: 10.1007/978-1-60761-611-5_3
- Colot, H. V., Park, G., Turner, G. E., Ringelberg, C., Crew, C. M., Litvinkova, L., et al. (2006). A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc. Natl. Acad. Sci. U. S. A.* 103, 10352–10357. doi: 10.1073/pnas.0601456103
- Corcoran, P., Anderson, J. L., Jacobson, D. J., Sun, Y., Ni, P., Lascoux, M., et al. (2016). Introgression maintains the genetic integrity of the mating-type determining chromosome of the fungus *Neurospora tetrasperma*. *Genome Res.* 26, 486–498. doi: 10.1101/gr.197244.115
- Culibrk, L., Croft, C. A., and Tebbutt, S. J. (2016). Systems biology approaches for host-fungal interactions: an expanding multi-omics frontier. *OMICS* 20, 127–138. doi: 10.1089/omi.2015.0185
- Cuomo, C. A., Güldenier, U., Xu, J.-R., Trail, F., Turgeon, B. G., Di Pietro, A., et al. (2007). The fusarium graminearum genome reveals a link between localized polymorphism and pathogen specialization. *Science* 317, 1400–1402. doi: 10.1126/science.1143708
- Cuomo, C. A., Untereiner, W. A., Ma, L.-J., Grabherr, M., and Birren, B. W. (2015). Draft genome sequence of the cellulolytic fungus *Chaetomium globosum*. *Genome Announc.* 3 (1), e00021–15. doi: 10.1128/genomeA.00021-15
- David, R. F., Marr, L. C., and Schmale, D. G. (2016). Ascospore release and discharge distances of *Fusarium graminearum* under controlled temperature and relative humidity. *Eur. J. Plant Pathol.* 146, 59–69. doi: 10.1007/s10658-016-0891-0
- Davis, R. H. (2003a). *Aspergillus* (New York, NY: Oxford University Press).
- Davis, R. H. (2003b). *Model systems, model organisms* (New York, NY: Oxford University Press).
- Dean, R. A., Talbot, N. J., Ebbole, D. J., Farman, M. L., Mitchell, T. K., Orbach, M. J., et al. (2005). The genome sequence of the rice blast fungus *Magnaporthe oryzae*. *Nature* 434, 980–986. doi: 10.1038/nature03449
- Dekhang, R., Wu, C., Smith, K. M., Lamb, T. M., Peterson, M., Bredeweg, E. L., et al. (2017). The *Neurospora* transcription factor ADV-1 transduces light signals and temporal information to control rhythmic expression of genes involved in cell fusion. *G3* 7, 129–142. doi: 10.1534/g3.116.034298
- Delgado-Baquerizo, M., Guerra, C. A., Cano-Díaz, C., Egidi, E., Wang, J.-T., Eisenhauer, N., et al. (2020). The proportion of soil-borne pathogens increases with warming at the global scale. *Nat. Clim. Change* 10, 550–554. doi: 10.1038/s41558-020-0759-3
- Del Ponte, E. M., Moreira, G. M., Ward, T. J., O'Donnell, K., Nicolli, C. P., Machado, F. J., et al. (2022). Species complex: a bibliographic analysis and web-accessible database for global mapping of species and trichothecene toxin chemotypes. *Phytopathology* 112, 741–751. doi: 10.1094/PHYTO-06-21-0277-RVW
- de Man, T. J. B., Stajich, J. E., Kubicek, C. P., Teiling, C., Chenthamara, K., Atanasova, L., et al. (2016). Small genome of the fungus *Escovopsis weberi*, a specialized disease agent of ant agriculture. *Proc. Natl. Acad. Sci. U. S. A.* 113, 3567–3572. doi: 10.1073/pnas.1518501113
- Detomasi, T. C., Rico-Ramírez, A. M., Saylor, R. I., Gonçalves, A. P., Marletta, M. A., and Glass, N. L. (2022). A moonlighting function of a chitin polysaccharide monooxygenase, CWR-1, in allorecognition. *Elife* 11, e80459. doi: 10.7554/eLife.80459
- Díaz, R., Wang, Z., and Townsend, J. P. (2023). “Chapter 5 - measurement and meaning in gene expression evolution,” in *Transcriptome profiling*. Eds. M. Ajmal Ali and J. Lee (Boston, MA, USA: Academic Press), 111–129.
- Di Lelio, I., Forni, G., Magoga, G., Brunetti, M., Bruno, D., Becchimanzi, A., et al. (2023). A soil fungus confers plant resistance against a phytophagous insect by disrupting the symbiotic role of its gut microbiota. *Proc. Natl. Acad. Sci. U. S. A.* 120, e2216922120. doi: 10.1073/pnas.2216922120
- Dirschnabel, D. E., Nowrousian, M., Cano-Domínguez, N., Aguirre, J., Teichert, I., and Kück, U. (2014). New insights into the roles of NADPH oxidases in sexual development and ascospore germination in *Sordaria macrospora*. *Genetics* 196, 729–744. doi: 10.1534/genetics.113.159368
- Dobzhansky, T. (1973). Nothing in biology makes sense except in the light of evolution. *Am. Biol. Teach.* 35, 125–129. doi: 10.2307/4444260
- Domazet-Lošo, T., Brajković, J., and Tautz, D. (2007). A phylostratigraphy approach to uncover the genomic history of major adaptations in metazoan lineages. *Trends Genet.* 23, 533–539. doi: 10.1016/j.tig.2007.08.014
- Domazet-Lošo, T., and Tautz, D. (2010). A phylogenetically based transcriptome age index mirrors ontogenetic divergence patterns. *Nature* 468, 815–818. doi: 10.1038/nature09632
- Dornburg, A., Townsend, J. P., and Wang, Z. (2017). Maximizing power in phylogenetics and phylogenomics: a perspective illuminated by fungal big data. *Adv. Genet.* 100, 1–47. doi: 10.1016/bs.adgen.2017.09.007
- Dreyfuss, J. M., Zucker, J. D., Hood, H. M., Ocasio, L. R., Sachs, M. S., and Galagan, J. E. (2013). Reconstruction and validation of a genome-scale metabolic model for the filamentous fungus *Neurospora crassa* using FARM. *PLoS Comput. Biol.* 9, e1003126. doi: 10.1371/journal.pcbi.1003126
- Drost, H.-G., Gabel, A., Grosse, I., and Quint, M. (2015). Evidence for active maintenance of phylotranscriptomic hourglass patterns in animal and plant embryogenesis. *Mol. Biol. Evol.* 32, 1221–1231. doi: 10.1093/molbev/msv012
- Druzhinina, I. S., Chenthamara, K., Zhang, J., Atanasova, L., Yang, D., Miao, Y., et al. (2018). Massive lateral transfer of genes encoding plant cell wall-degrading enzymes to the mycoparasitic fungus *Trichoderma* from its plant-associated hosts. *PLoS Genet.* 14, e1007322. doi: 10.1371/journal.pgen.1007322
- Druzhinina, I. S., Seidl-Seiboth, V., Herrera-Estrella, A., Horwitz, B. A., Kenerley, C. M., Monte, E., et al. (2011). *Trichoderma*: the genomics of opportunistic success. *Nat. Rev. Microbiol.* 9, 749–759. doi: 10.1038/nrmicro2637
- Dunlap, J. C., Borkovich, K. A., Henn, M. R., Turner, G. E., Sachs, M. S., Glass, N. L., et al. (2007). Enabling a community to dissect an organism: overview of the *Neurospora* functional genomics project. *Adv. Genet.* 57, 49–96. doi: 10.1016/S0065-2660(06)57002-6
- Durufle, H., and Déjean, S. (2023). Multi-omics data integration in the context of plant abiotic stress signaling. *Methods Mol. Biol.* 2642, 295–318. doi: 10.1007/978-1-0716-3044-0_16
- Dyer, P. S., and O’Gorman, C. M. (2012). Sexual development and cryptic sexuality in fungi: insights from *Aspergillus* species. *FEMS Microbiol. Rev.* 36, 165–192. doi: 10.1111/j.1574-6976.2011.00308.x
- Ellena, V., and Steiger, M. G. (2022). The importance of complete and high-quality genome sequences in *Aspergillus niger* research. *Front. Fungal Biol.* 3. doi: 10.3389/ffunb.2022.935993
- Ellison, C. (2011). *Evolutionary genomics of divergence and adaptation within the model fungi Neurospora crassa and Neurospora tetrasperma*. (UC Berkeley). Available at: <https://escholarship.org/uc/item/5519j6mt>.
- Espagne, E., Lespinet, O., Malagnac, F., Da Silva, C., Jaillon, O., Porcel, B. M., et al. (2008). The genome sequence of the model ascomycete fungus *Podospora anserina*. *Genome Biol.* 9, 1–22. doi: 10.1186/gb-2008-9-5-r77
- Fan, X., Zhang, P., Batool, W., Liu, C., Hu, Y., Wei, Y., et al. (2023). Contribution of the tyrosinase (MoTyr) to melanin synthesis, conidiogenesis, appressorium development, and pathogenicity in *J. Fungi (Basel)* 9. doi: 10.3390/jof9030311
- Feldman, D., Kowbel, D. J., Cohen, A., Glass, N. L., Hadar, Y., and Yarden, O. (2019). Identification and manipulation of genes involved in sensitivity to furfural. *Biotechnol. Biofuels* 12, 210. doi: 10.1186/s13068-019-1550-4
- Feldmann, H. (1999). “The yeast *Saccharomyces cerevisiae*: insights from the first complete eukaryotic genome sequence” in *Molecular Fungal Biology*. Eds. R. Oliver and M. Schweizer. (Cambridge: Cambridge University Press), 78–134. doi: 10.1017/cbo9781139163972.004
- Feurtrey, A., Lorrain, C., McDonald, M. C., Milgate, A., Solomon, P. S., Warren, R., et al. (2023). A thousand-genome panel retraces the global spread and adaptation of a major fungal crop pathogen. *Nat. Commun.* 14, 1059. doi: 10.1038/s41467-023-36674-y
- Franco, M. E. E., Wisecaver, J. H., Arnold, A. E., Ju, Y.-M., Slot, J. C., Ahrendt, S., et al. (2022). Ecological generalism drives hyperdiversity of secondary metabolite gene clusters in xylarialean endophytes. *New Phytol.* 233, 1317–1330. doi: 10.1111/nph.17873
- Frankland, J. C., Magan, N., and Gadd, G. M. (2009). *Fungi and environmental change* (Cambridge, UK: Cambridge University Press).
- Frias-De-León, M. G., Brunner-Mendoza, C., del Rocio Reyes-Montes, M., and Duarte-Escalante, E. (2022). *The impact of climate change on fungal diseases* (Berlin, Germany/Boston, MA, USA: Springer Nature).
- Fricke, J., Schalk, F., Kreuzenbeck, N. B., Seibel, E., Hoffmann, J., Dittmann, G., et al. (2023). Adaptations of pseudoxylaria towards a comb-associated lifestyle in fungus-farming termite colonies. *ISME J.* 17, 733–747. doi: 10.1038/s41396-023-01374-4
- Gadd, G., Watkinson, S. C., and Dyer, P. S. (2007). *Fungi in the environment* (Cambridge, UK: Cambridge University Press).
- Galagan, J. E., Henn, M. R., Ma, L.-J., Cuomo, C. A., and Birren, B. (2005). Genomics of the fungal kingdom: insights into eukaryotic biology. *Genome Res.* 15, 1620–1631. doi: 10.1101/gr.3767105
- Galindo, L. J., Milner, D. S., Gomes, S. L., and Richards, T. A. (2022). A light-sensing system in the common ancestor of the fungi. *Curr. Biol.* 32, 3146–3153.e3. doi: 10.1016/j.cub.2022.05.034
- Gao, Q., Jin, K., Ying, S.-H., Zhang, Y., Xiao, G., Shang, Y., et al. (2011). Genome sequencing and comparative transcriptomics of the model entomopathogenic fungi *Metarhizium anisopliae* and *M. acridum*. *PLoS Genet.* 7, e1001264. doi: 10.1371/journal.pgen.1001264
- Garmendia, G., Umpierrez-Failache, M., Ward, T. J., and Vero, S. (2018). Development of a PCR-RFLP method based on the transcription elongation factor 1- α gene to differentiate *Fusarium graminearum* from other species within the *Fusarium graminearum* species complex. *Food Microbiol.* 70, 28–32. doi: 10.1016/j.fm.2017.08.020
- Geiser, D. M., Gueidan, C., Miadlikowska, J., Lutzoni, F., Kauff, F., Hofstetter, V., et al. (2006). Eurotiomycetes: eurotiomycetidae and chaetothyriomycetidae. *Mycologia* 98, 1053–1064. doi: 10.1080/15572536.2006.11832633
- Geydan, T. D., Debets, A. J. M., Verkley, G. J. M., and van Diepeningen, A. D. (2012). Correlated evolution of senescence and ephemeral substrate use in the sordariomycetes. *Mol. Ecol.* 21, 2816–2828. doi: 10.1111/j.1365-294X.2012.05569.x
- Gladieux, P., Wilson, B. A., Perraudeau, F., Montoya, L. A., Kowbel, D., Hann-Soden, C., et al. (2015). Genomic sequencing reveals historical, demographic and selective factors associated with the diversification of the fire-associated fungus *Neurospora discreta*. *Mol. Ecol.* 24, 5657–5675. doi: 10.1111/mec.13417

- Goldstein, B., and King, N. (2016). The future of cell biology: emerging model organisms. *Trends Cell Biol.* 26, 818–824. doi: 10.1016/j.tcb.2016.08.005
- Gortikov, M., Wang, Z., Steindorff, A. S., Grigoriev, I. V., Druzhinina, I. S., Townsend, J. P., et al. (2022a). Sequencing and analysis of the entire genome of the mycoparasitic bioeffector fungus *trichoderma asperelloides* strain T 203 (Hypocreales). *Microbiol. Resource Announce.* 11, e00995–e00921. doi: 10.1128/mra.00995-21
- Gortikov, M., Yakubovich, E., Wang, Z., López-Giráldez, F., Tu, Y., Townsend, J. P., et al. (2022b). Differential expression of cell wall remodeling genes is part of the dynamic phase-specific transcriptional program of conidial germination of *J. Fungi* (Basel) 8 (8), 854. doi: 10.3390/jof8080854
- Grau-Bové, X., Navarrete, C., Chiva, C., Pribasni, T., Antó, M., Torruella, G., et al. (2022). A phylogenetic and proteomic reconstruction of eukaryotic chromatin evolution. *Nat. Ecol. Evol.* 6, 1007–1023. doi: 10.1038/s41559-022-01771-6
- Green, B. J., Nayak, A. P., Lemons, A. R., Rittenour, W. R., Hettick, J. M., and Beezhold, D. H. (2014). Production of a chaetomium globosum enolase monoclonal antibody. *Monoclon. Antib. Immunodiagn. Immunother.* 33, 428–437. doi: 10.1089/mab.2014.0042
- Grigoriev, I. V., Cullen, D., Goodwin, S. B., Hibbett, D., Jeffries, T. W., Kubicek, C. P., et al. (2011). Fueling the future with fungal genomics. *Mycology* 2, 192–209. doi: 10.1080/21501203.2011.584577
- Grigoriev, I. V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otillar, R., et al. (2014). MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids Res.* 42, D699–D704. doi: 10.1093/nar/gkt1183
- Grognet, P., Timpano, H., Carlier, F., Ait-Benkhalil, J., Berteaux-Lecellier, V., Debuchy, R., et al. (2019). A RID-like putative cytosine methyltransferase homologue controls sexual development in the fungus *podospora anserina*. *PLoS Genet.* 15, e1008086. doi: 10.1371/journal.pgen.1008086
- Guhlin, J., Silverstein, K. A. T., Zhou, P., Tiffin, P., and Young, N. D. (2017). ODG: omics database generator - a tool for generating, querying, and analyzing multi-omics comparative databases to facilitate biological understanding. *BMC Bioinf.* 18, 367. doi: 10.1186/s12859-017-1777-7
- Guzmán-Guzmán, P., Porras-Troncoso, M. D., Olmedo-Monfil, V., and Herrera-Estrella, A. (2019). Trichoderma species: versatile plant symbionts. *Phytopathology* 109, 6–16. doi: 10.1094/PHYTO-07-18-0218-RVW
- Hansen, K., and Pfister, D. H. (2006). Systematics of the pezizomycetes—the operculate discomycetes. *Mycologia* 98, 1029–1040. doi: 10.1080/15572536.2006.11832631
- Hartmann, F. E., Ament-Velásquez, S. L., Vogan, A. A., Gautier, V., Le Prieur, S., Berramane, M., et al. (2021a). Size variation of the nonrecombining region on the mating-type chromosomes in the fungal *podospora anserina* species complex. *Mol. Biol. Evol.* 38, 2475–2492. doi: 10.1093/molbev/msab040
- Hartmann, F. E., Duhamel, M., Carpentier, F., Hood, M. E., Foulongne-Oriol, M., Silar, P., et al. (2021b). Recombination suppression and evolutionary strata around mating-type loci in fungi: documenting patterns and understanding evolutionary and mechanistic causes. *New Phytol.* 229, 2470–2491. doi: 10.1111/nph.17039
- Hassett, C. C., Elliott Horner, W., Levetin, E., Wild, L. G., Edward Davis, W., Lehrer, S. B., et al. (2009). “Fungi as allergens” in *Allergy and allergic diseases*. Eds. A. B. Kay, J. Bousquet, P. G. Holt and A. P. Kaplan (Oxford, United Kingdom: 2nd ed Wiley), 963–983.
- Heitman, J., Howlett, B. J., Crous, P. W., Stukenbrock, E. H., James, T. Y., and Gow, N. A. R. (2017). *The fungal kingdom* (New Jersey, USA: John Wiley & Sons).
- Hodgins-Davis, A., Rice, D. P., and Townsend, J. P. (2015). Gene expression evolves under a house-of-cards model of stabilizing selection. *Mol. Biol. Evol.* 32, 2130–2140. doi: 10.1093/molbev/msv094
- Honda, S., Eusebio-Cope, A., Miyashita, S., Yokoyama, A., Aulia, A., Shahi, S., et al. (2020). Establishment of *neurospora crassa* as a model organism for fungal virology. *Nat. Commun.* 11, 1–13. doi: 10.1038/s41467-020-19355-y
- Hossain, M. M. (2022). Wheat blast: a review from a genetic and genomic perspective. *Front. Microbiol.* 13. doi: 10.3389/fmicb.2022.983243
- Hotaling, S., Wilcox, E. R., Heckenhauser, J., Stewart, R. J., and Frandsen, P. B. (2023). Highly accurate long reads are crucial for realizing the potential of biodiversity genomics. *BMC Genomics* 24, 1–9. doi: 10.1186/s12864-023-09193-9
- Huberman, L. B., Wu, V. W., Lee, J., Daum, C., O'Malley, R. C., and Glass, N. L. (2021). Aspects of the *neurospora crassa* sulfur starvation response are revealed by transcriptomic profiling and DNA affinity purification sequencing. *mSphere* 6, e0056421. doi: 10.1128/mSphere.00564-21
- Hurley, J. M., Dasgupta, A., Emerson, J. M., Zhou, X., Ringelberg, C. S., Knabe, N., et al. (2014). Analysis of clock-regulated genes in *neurospora* reveals widespread posttranscriptional control of metabolic potential. *Proc. Natl. Acad. Sci. U. S. A.* 111, 16995–17002. doi: 10.1073/pnas.1418963111
- Hyde, K. D., Norphanphoun, C., Maharachchikumbura, S. S. N., Bhat, D. J., Jones, E. B. G., Bundhun, D., et al. (2020). Refined families of sordariomycetes. *Mycosphere* 11, 305–1059. doi: 10.5943/mycosphere/11/1/7
- Hyun, J. C., Monk, J. M., and Palsom, B. O. (2022). Comparative pangenomics: analysis of 12 microbial pathogen pangenomes reveals conserved global structures of genetic and functional diversity. *BMC Genomics* 23, 7. doi: 10.1186/s12864-021-08223-8
- Idnurm, A., Hood, M. E., Johannesson, H., and Giraud, T. (2015). Contrasted patterns in mating-type chromosomes in fungi: hotspots versus coldspots of recombination. *Fungal Biol. Rev.* 29, 220–229. doi: 10.1016/j.fbr.2015.06.001
- Ingold, C. T., and Dring, V. J. (1957). An analysis of spore discharge in *sordaria*. *Ann. Bot.* 21, 465–477. doi: 10.1093/oxfordjournals.aob.a083578
- Jaffe, J. D., Berg, H. C., and Church, G. M. (2004). Proteogenomic mapping as a complementary method to perform genome annotation. *Proteomics* 4, 59–77. doi: 10.1002/pmic.200300511
- James, T. Y., Stajich, J. E., Hittinger, C. T., and Rokas, A. (2020). Toward a fully resolved fungal tree of life. *Annu. Rev. Microbiol.* 74, 291–313. doi: 10.1146/annurev-micro-022020-051835
- Jones, M. G. (2007). The first filamentous fungal genome sequences: *aspergillus* leads the way for essential everyday resources or dusty museum specimens? *Microbiology* 153, 1–6. doi: 10.1099/mic.0.2006/001479-0
- Karahalil, B. (2016). Overview of systems biology and omics technologies. *Curr. Med. Chem.* 23, 4221–4230. doi: 10.2174/0929867323666160926150617
- Kashyap, P. L., Rai, P., Srivastava, A. K., and Kumar, S. (2017). Trichoderma for climate resilient agriculture. *World J. Microbiol. Biotechnol.* 33, 155. doi: 10.1007/s12274-017-2319-1
- Kato, T., Nishimura, K., Misu, S., Ikeo, K., and Park, E. Y. (2022). Changes of the gene expression in silkworm larvae and *corydops militaris* at late stages of the pathogenesis. *Arch. Insect Biochem. Physiol.* 111, e21968. doi: 10.1002/arch.21968
- Kelliher, C. M., Lambregts, R., Xiang, Q., Baker, C. L., Loros, J. J., and Dunlap, J. C. (2020). PRD-2 directly regulates and counteracts nonsense-mediated decay in the *neurospora* circadian clock. *Elife* 9, e64007. doi: 10.7554/eLife.64007
- Kelliher, C. M., Stevenson, E.-L., Loros, J. J., and Dunlap, J. C. (2023). Nutritional compensation of the circadian clock is a conserved process influenced by gene expression regulation and mRNA stability. *PLoS Biol.* 21, e3001961. doi: 10.1371/journal.pbio.3001961
- Khan, M. K., Pandey, A., Athar, T., Choudhary, S., Deval, R., Gezgün, S., et al. (2020). Fusarium head blight in wheat: contemporary status and molecular approaches. *3 Biotech.* 10, 172. doi: 10.1007/s13205-020-2158-x
- Kim, W., Cavinder, B., Proctor, R. H., O'Donnell, K., Townsend, J. P., and Trail, F. (2019). Comparative genomics and transcriptomics during sexual development gives insight into the life history of the cosmopolitan fungus *fusarium neocosmosporiellum*. *Front. Microbiol.* 10. doi: 10.3389/fmicb.2019.01247
- Kim, W., Miguel-Rojas, C., Wang, J., Townsend, J. P., and Trail, F. (2018). Developmental dynamics of long noncoding RNA expression during sexual fruiting body formation in *fusarium graminearum*. *mBio* 9, e01292–18. doi: 10.1128/mbio.01292-18
- Kim, W., Wang, Z., Kim, H., Pham, K., Tu, Y., Townsend, J. P., et al. (2022). Transcriptional divergence underpinning sexual development in the fungal class sordariomycetes. *mBio* 13 (3), e0110022. doi: 10.1128/mbio.01100-22
- King, M. C., and Wilson, A. C. (1975). Evolution at two levels in humans and chimpanzees. *Science* 188, 107–116. doi: 10.1126/science.1090005
- Ko, G., Jang, I., Koo, N., Park, S.-J., Oh, S.-H., Kim, M.-S., et al. (2020). Prometheus, An omics portal for interkingdom comparative genomic analyses. *PLoS One* 15, e0240191. doi: 10.1371/journal.pone.0240191
- Kotta-Loizou, I. (2019). *Mycoviruses* (Basel, Switzerland: MDPI).
- Kubicek, C. P., Herrera-Estrella, A., Seidl-Seiboth, V., Martinez, D. A., Druzhinina, I. S., Thon, M., et al. (2011). Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of trichoderma. *Genome Biol.* 12, R40. doi: 10.1186/gb-2011-12-4-r40
- Kubicek, C. P., Steindorff, A. S., Chenthamara, K., Manganiello, G., Henrissat, B., Zhang, J., et al. (2019). Evolution and comparative genomics of the most common trichoderma species. *BMC Genomics* 20, 1–24. doi: 10.1186/s12864-019-5680-7
- Kumar, S., Kashyap, P. L., and Singh, G. P. (2020). *Wheat blast* (Florida, USA: CRC Press).
- Kusch, S., Qian, J., Loos, A., Kümmel, F., Spanu, P. D., and Panstruga, R. (2023). Long-term and rapid evolution in powdery mildew fungi. *Mol. Ecol.* doi: 10.1111/mec.16909
- Latorre, S. M., Were, V. M., Foster, A. J., Langner, T., Malmgren, A., Harant, A., et al. (2023). Genomic surveillance uncovers a pandemic clonal lineage of the wheat blast fungus. *PLoS Biol.* 21, e3002052. doi: 10.1371/journal.pbio.3002052
- Leger, A., Amaral, P. P., Pandolfini, L., Capitanchik, C., Capraro, F., Miano, V., et al. (2021). RNA Modifications detection by comparative nanopore direct RNA sequencing. *Nat. Commun.* 12, 1–17. doi: 10.1038/s41467-021-27393-3
- Lehr, N. A., Wang, Z., Li, N., Hewitt, D. A., López-Giráldez, F., Trail, F., et al. (2014). Gene expression differences among three *neurospora* species reveal genes required for sexual reproduction in *neurospora crassa*. *PLoS One* 9, e110398. doi: 10.1371/journal.pone.0110398
- Lelandais, G., Remy, D., Malagnac, F., and Grognet, P. (2022). New insights into genome annotation in *podospora anserina* through re-exploiting multiple RNA-seq data. *BMC Genomics* 23, 859. doi: 10.1186/s12864-022-09085-4
- Li, P., Bhattacharjee, P., Wang, S., Zhang, L., Ahmed, I., and Guo, L. (2019a). Mycoviruses in *fusarium* species: an update. *Front. Cell. Infect. Microbiol.* 9. doi: 10.3389/fcimb.2019.00257
- Li, J., Chroumpi, T., Garrigues, S., Kun, R. S., Meng, J., Salazar-Cerezo, S., et al. (2022). The sugar metabolic model of *aspergillus niger* can only be reliably transferred to fungi of its phylum. *J. Fungi* 8, 1315. doi: 10.3390/jof8121315

- Li, X., Wang, F., Liu, Q., Li, Q., Qian, Z., Zhang, X., et al. (2019b). Developmental transcriptomics of Chinese cordyceps reveals gene regulatory network and expression profiles of sexual development-related genes. *BMC Genomics* 20, 337. doi: 10.1186/s12864-019-5708-z
- Liu, Q., and Dong, C. (2023). Dual transcriptomics reveals interspecific interactions between the mycoparasite and its host cordyceps militaris. *Microbiol. Spectr.*, e0480022. doi: 10.1128/spectrum.04800-22
- Liu, H., Li, Y., Chen, D., Qi, Z., Wang, Q., Wang, J., et al. (2017). A-to-I RNA editing is developmentally regulated and generally adaptive for sexual reproduction in. *Proc. Natl. Acad. Sci. U. S. A.* 114, E7756–E7765. doi: 10.1073/pnas.1702591114
- Liu, D., Liu, Y., Zhang, D., Chen, X., Liu, Q., Xiong, B., et al. (2020). Quantitative proteome profiling reveals cellobiose-dependent protein processing and export pathways for the lignocellulolytic response in *Neurospora crassa*. *Appl. Environ. Microbiol.* 86 (15), e00653–20. doi: 10.1128/AEM.00653-20
- Liu, H., Wang, Q., He, Y., Chen, L., Hao, C., Jiang, C., et al. (2016). Genome-wide a-to-I RNA editing in fungi independent of ADAR enzymes. *Genome Res.* 26, 499–509. doi: 10.1101/gr.199877.115
- Liu, Q., Xu, Y., Zhang, X., Li, K., Li, X., Wang, F., et al. (2021). Infection process and genome assembly provide insights into the pathogenic mechanism of destructive mycoparasite *Calcarisporium cordycipitcola* with host specificity. *J. Fungi* 7 (11), 918. doi: 10.3390/jof7110918
- Luo, J., Qiu, H., Cai, G., Wagner, N. E., Bhattacharya, D., and Zhang, N. (2015). Phylogenomic analysis uncovers the evolutionary history of nutrition and infection mode in rice blast fungus and other magnaportheales. *Sci. Rep.* 5, 9448. doi: 10.1038/srep09448
- Lützenhaus, R., Traeger, S., Breuer, J., Carreté, L., Kuo, A., Lipzen, A., et al. (2019). Comparative genomics and transcriptomics to analyze fruiting body development in filamentous ascomycetes. *Genetics* 213, 1545–1563. doi: 10.1534/genetics.119.302749
- Lv, W., Xiao, Y., Xu, Z., Jiang, H., Tong, Q., and Wang, Z. (2022). The paxillin MoPax1 activates mitogen-activated protein (MAP) kinase signaling pathways and autophagy through MAP kinase activator MoMka1 during appressorium-mediated plant infection by the rice blast fungus *magnaporthe oryzae*. *MBio* 13, e0221822. doi: 10.1128/mbio.02218-22
- Ma, L.-J., Geiser, D. M., Proctor, R. H., Rooney, A. P., O'Donnell, K., Trail, F., et al. (2013). *Fusarium* pathogenomics. *Annu. Rev. Microbiol.* 67, 399–416. doi: 10.1146/annurev-micro-092412-155650
- Ma, L.-J., van der Does, H. C., Borkovich, K. A., Coleman, J. J., Daboussi, M.-J., Di Pietro, A., et al. (2010). Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464, 367–373. doi: 10.1038/nature08850
- Mackenzie, D. W. R. (1979). Immune responses to fungal infections. *Immunol. Aspects Infect. Dis.*, 21–75. doi: 10.1007/978-94-011-6191-6_2
- Maghuly, F., Marzban, G., and Jankowicz-Cieslak, J. (2022). *Omics technologies toward systems biology* (Lausanne, Switzerland: Frontiers Media SA).
- Maharachchikumbura, S. S. N., Hyde, K. D., Jones, E. B. G., McKenzie, E. H. C., Bhat, J. D., Dayaratne, M. C., et al. (2016). Families of sordariomycetes. *Fungal Divers.* 79, 1–317. doi: 10.1007/s13225-016-0369-6
- Maharachchikumbura, S. S. N., Hyde, K. D., Jones, E. B. G., McKenzie, E. H. C., Huang, S.-K., Abdel-Wahab, M. A., et al. (2015). Towards a natural classification and backbone tree for sordariomycetes. *Fungal Divers.* 72, 199–301. doi: 10.1007/s13225-015-0331-z
- Martinez, D., Berka, R. M., Henrissat, B., Saloheimo, M., Arvas, M., Baker, S. E., et al. (2008). Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *hypocrea jecorina*). *Nat. Biotechnol.* 26, 553–560. doi: 10.1038/nbt1403
- Mayer, C., Vogt, A., Uslu, T., Scalzitti, N., Chennan, K., Poch, O., et al. (2023). CeGAL: redefining a widespread fungal-specific transcription factor family using an in silico error-tracking approach. *J. Fungi* 9 (4), 424. doi: 10.3390/jof9040424
- Merényi, Z., Prasanna, A. N., Wang, Z., Kovács, K., Hegedüs, B., Bálint, B., et al. (2020). Unmatched level of molecular convergence among deeply divergent complex multicellular fungi. *Mol. Biol. Evol.* 37, 2228–2240. doi: 10.1093/molbev/msaa077
- Merino, S. T., Nelson, M. A., Jacobson, D. J., and Natvig, D. O. (1996). Pseudohomothallism and evolution of the mating-type chromosome in *Neurospora tetrasperma*. *Genetics* 143, 789–799. doi: 10.1093/genetics/143.2.789
- Merrill, M. (2013). Responses of mycorrhizal fungi and other root-associated fungi to climate change. PhD thesis. Denmark: Department of Biology, Faculty of Science, University of Copenhagen. p. 150.
- Meyer, V., Basenko, E. Y., Benz, J. P., Braus, G. H., Caddick, M. X., Csukai, M., et al. (2020). Growing a circular economy with fungal biotechnology: a white paper. *Fungal Biol. Biotechnol.* 7, 1–23. doi: 10.1186/s40694-020-00095-z
- Miguel-Rojas, C., Cavinder, B., Townsend, J. P., and Trail, F. (2023). Comparative transcriptomics of *Fusarium graminearum* and *magnaporthe oryzae* spore germination leading up to infection. *mBio* 14 (1), e0244222. doi: 10.1128/mbio.02442-22
- Miller, J. D., and McMullin, D. R. (2014). Fungal secondary metabolites as harmful indoor air contaminants: 10 years on. *Appl. Microbiol. Biotechnol.* 98, 9953–9966. doi: 10.1007/s00253-014-6178-5
- Mir, Z. A., Chandra, T., Saharan, A., Budhlakoti, N., Mishra, D. C., Saharan, M. S., et al. (2023). Recent advances on genome-wide association studies (GWAS) and genomic selection (GS); prospects for *Fusarium* head blight research in durum wheat. *Mol. Biol. Rep.* 50, 3885–3901. doi: 10.1007/s11033-023-08309-4
- Miranda-Apodaca, J., Artetxe, U., Aguado, I., Martín-Souto, L., Ramírez-García, A., Lacuesta, M., et al. (2023). Stress response to climate change and postharvest handling in two differently pigmented lettuce genotypes: impact on invasion and mycotoxin production. *Plants* 12 (6), 1304. doi: 10.3390/plants12061304
- Misra, B. B., Langefeld, C., Olivier, M., and Cox, L. A. (2019). Integrated omics: tools, advances and future approaches. *J. Mol. Endocrinol.* 62, R21–R45. doi: 10.1530/JME-18-0055
- Miyauchi, S., Kiss, E., Kuo, A., Drula, E., Kohler, A., Sánchez-García, M., et al. (2020). Large-scale genome sequencing of mycorrhizal fungi provides insights into the early evolution of symbiotic traits. *Nat. Commun.* 11, 5125. doi: 10.1038/s41467-020-18795-w
- Mukherjee, P. K., Horwitz, B. A., Herrera-Estrella, A., Schmoll, M., and Kenerley, C. M. (2013). *Trichoderma* research in the genome era. *Annu. Rev. Phytopathol.* 51, 105–129. doi: 10.1146/annurev-phyto-082712-102353
- Myers, J. M., Bonds, A. E., Clemons, R. A., Thapa, N. A., Simmons, D. R., Carter-House, D., et al. (2020). Survey of early-diverging lineages of fungi reveals abundant and diverse mycoviruses. *MBio* 11 (5), e02027–20. doi: 10.1128/mbio.02027-20
- Nagy, L. G., Riley, R., Tritt, A., Adam, C., Daum, C., Floudas, D., et al. (2016). Comparative genomics of early-diverging mushroom-forming fungi provides insights into the origins of lignocellulose decay capabilities. *Mol. Biol. Evol.* 33, 959–970. doi: 10.1093/molbev/msv337
- Naranjo-Ortiz, M. A., and Gabaldón, T. (2020). Fungal evolution: cellular, genomic and metabolic complexity. *Biol. Rev. Camb. Philos. Soc.* 95, 1198–1232. doi: 10.1111/brv.12605
- Nesvizhskii, A. I. (2014). Proteogenomics: concepts, applications and computational strategies. *Nat. Methods* 11, 1114–1125. doi: 10.1038/nmeth.3144
- North, H. L., McGaughan, A., and Jiggins, C. D. (2021). Insights into invasive species from whole-genome resequencing. *Mol. Ecol.* 30, 6289–6308. doi: 10.1111/mec.15999
- Nowrousian, M. (2014). *Fungal genomics* (Heidelberg, Germany: Springer Science & Business Media).
- Nowrousian, M., Stajich, J. E., Chu, M., Engh, I., Espagne, E., Halliday, K., et al. (2010). *De novo* assembly of a 40 Mb eukaryotic genome from short sequence reads: sordaria macrospora, a model organism for fungal morphogenesis. *PLoS Genet.* 6, e1000891. doi: 10.1371/journal.pgen.1000891
- Nowrousian, M., Teichert, I., Masloff, S., and Kück, U. (2012). Whole-genome sequencing of sordaria macrospora mutants identifies developmental genes. *G3* 2, 261–270. doi: 10.1534/g3.111.001479
- O'Donnell, K., Ward, T. J., Robert, V. A. R. G., Crous, P. W., Geiser, D., and Kang, M. G. (2015). DNA sequence-based identification of *Fusarium*: Current status and future directions. *Phytoparasitica* 43, 583–595.
- Paoletti, M., and Saupé, S. J. (2008). The genome sequence of *Podospora anserina*, a classic model fungus. *Genome Biol.* 9, 223. doi: 10.1186/gb-2008-9-5-223
- Pattemore, J. A., Hane, J. K., Williams, A. H., Wilson, B. A. L., Stodart, B. J., and Ash, G. J. (2014). The genome sequence of the biocontrol fungus *metarhizium anisopliae* and comparative genomics of *metarhizium* species. *BMC Genomics* 15, 660. doi: 10.1186/1471-2164-15-660
- Peris, D., Lu, D. S., Kinneberg, V. B., Methlie, I.-S., Dahl, M. S., James, T. Y., et al. (2022). Large-scale fungal strain sequencing unravels the molecular diversity in mating loci maintained by long-term balancing selection. *PLoS Genet.* 18, e1010097. doi: 10.1371/journal.pgen.1010097
- Perkins, D. D. (1992). *Neurospora*: the organism behind the molecular revolution. *Genetics* 130, 687–701. doi: 10.1093/genetics/130.4.687
- Philipp, O., Hamann, A., Servos, J., Werner, A., Koch, I., and Osiewacz, H. D. (2013). A genome-wide longitudinal transcriptome analysis of the aging model *Podospora anserina*. *PLoS One* 8, e83109. doi: 10.1371/journal.pone.0083109
- Plissonneau, C., Benevenuto, J., Mohd-Assaad, N., Fouché, S., Hartmann, F. E., and Croll, D. (2017). Using population and comparative genomics to understand the genetic basis of effector-driven fungal pathogen evolution. *Front. Plant Sci.* 8, 119. doi: 10.3389/fpls.2017.00119
- Plissonneau, C., Hartmann, F. E., and Croll, D. (2018). Pangenome analyses of the wheat pathogen *Zymoseptoria tritici* reveal the structural basis of a highly plastic eukaryotic genome. *BMC Biol.* 16, 5. doi: 10.1186/s12915-017-0457-4
- Podder, D., and Ghosh, S. K. (2019). A new application of *Trichoderma asperellum* as an anopheline larvicide for eco friendly management in medical science. *Sci. Rep.* 9, 1108. doi: 10.1038/s41598-018-37108-2
- Pöggeler, S., and James, T. (2023). *Evolution of fungi and fungal-like organisms* (Heidelberg, Germany: Springer).
- Posada, F., Vega, F. E., Rehner, S. A., Blackwell, M., Weber, D., Suh, S.-O., et al. (2004). *Syspastospora parasitica*, a mycoparasite of the fungus *beauveria bassiana* attacking the Colorado potato beetle *leptinotarsa decemlineata*: a tritrophic association. *J. Insect Sci.* 4, 24. doi: 10.1093/jis/4.1.24
- Poveda, J. (2021). *Trichoderma* as biocontrol agent against pests: new uses for a mycoparasite. *Biol. Control* 159, 104634. doi: 10.1016/j.biocontrol.2021.104634
- Qian, B., Su, X., Ye, Z., Liu, X., Liu, M., Zhang, H., et al. (2023). MoErv14 mediates the intracellular transport of cell membrane receptors to govern the appressorial

formation and pathogenicity of magnaporthe oryzae. *PLoS Pathog.* 19, e1011251. doi: 10.1371/journal.ppat.1011251

Qiu, J., Xie, J., Chen, Y., Shen, Z., Shi, H., Naqvi, N. I., et al. (2022). Warm temperature compromises JA-regulated basal resistance to enhance magnaporthe oryzae infection in rice. *Mol. Plant* 15, 723–739. doi: 10.1016/j.molp.2022.02.014

Rampersad, S. N. (2020). Pathogenomics and management of diseases in plants. *Pathogens* 9. doi: 10.3390/pathogens9050340

Reel, P. S., Reel, S., Pearson, E., Trucco, E., and Jefferson, E. (2021). Using machine learning approaches for multi-omics data analysis: a review. *Biotechnol. Adv.* 49, 107739. doi: 10.1016/j.biotechadv.2021.107739

Rhodes, J. (2023). Genomic surveillance urgently needed to control wheat blast pandemic spreading across continents. *PLoS Biol.* 21, e3002090. doi: 10.1371/journal.pbio.3002090

Riley, R., Salamov, A. A., Brown, D. W., Nagy, L. G., Floudas, D., Held, B. W., et al. (2014). Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. *Proc. Natl. Acad. Sci. U. S. A.* 111, 9923–9928. doi: 10.1073/pnas.1400592111

Robinson, A. J., Natvig, D. O., and Chain, P. S. G. (2020). Genomic analysis of diverse members of the fungal genus monosporascus reveals novel lineages, unique genome content and a potential bacterial associate. *G3 Genes/Genomes/Genetics* 10, 2573–2583. doi: 10.1534/g3.120.401489

Rodrigues, M. L., and Nosanchuk, J. D. (2023). Recognition of fungal priority pathogens: what next? *PLoS Negl. Trop. Dis.* 17, e0011136. doi: 10.1371/journal.pntd.0011136

Rodriguez, S., Ward, A., Reckard, A. T., Shtanko, Y., Hull-Crew, C., and Klocko, A. D. (2022). The genome organization of neurospora crassa at high resolution uncovers principles of fungal chromosome topology. *G3* 12 (5), jkac053. doi: 10.1093/g3journal/jkac053

Rodriguez-Romero, J., Hedtke, M., Kastner, C., Müller, S., and Fischer, R. (2010). Fungi, hidden in soil or up in the air: light makes a difference. *Annu. Rev. Microbiol.* 64, 585–610. doi: 10.1146/annurev.micro.112408.134000

Rogers, A. M., and Egan, M. J. (2023). Septum-associated microtubule organizing centers within conidia support infectious development by the blast fungus magnaporthe oryzae. *Fungal Genet. Biol.* 165, 103768. doi: 10.1016/j.fgb.2022.103768

Romero-Olivares, A. L., Taylor, J. W., and Treseder, K. K. (2015). Neurospora discreta as a model to assess adaptation of soil fungi to warming. *BMC Evol. Biol.* 15, 1–8. doi: 10.1186/s12862-015-0482-2

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D. L., Darling, A., Höhna, S., et al. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542. doi: 10.1093/sysbio/sys029

Rupa, P. V. D., Jageswar, G., and Kiranmayi, P. (2018). Zinc transporter protein (ztz-1) may also play a role in conidiation pathway of neurospora crassa: an insight using proteogenomic approach. *Protein Pept. Lett.* 24, 1120–1129. doi: 10.2174/0929866524666170920114503

Russell, J. J., Theriot, J. A., Sood, P., Marshall, W. F., Landweber, L. F., Fritz-Laylin, L., et al. (2017). Non-model model organisms. *BMC Biol.* 15, 1–31. doi: 10.1186/s12915-017-0391-5

Sagita, R., Quax, W. J., and Haslinger, K. (2021). Current state and future directions of genetics and genomics of endophytic fungi for bioprospecting efforts. *Front. Bioeng. Biotechnol.* 9, 649906. doi: 10.3389/fbioe.2021.649906

Samils, N., Gioti, A., Karlsson, M., Sun, Y., Kasuga, T., Bastiaans, E., et al. (2013). Sex-linked transcriptional divergence in the hermaphrodite fungus neurospora tetrasperma. *Proc. Biol. Sci.* 280, 20130862. doi: 10.1098/rspb.2013.0862

Sánchez-García, M., Ryberg, M., Khan, F. K., Varga, T., Nagy, L. G., and Hibbett, D. S. (2020). Fruiting body form, not nutritional mode, is the major driver of diversification in mushroom-forming fungi. *Proc. Natl. Acad. Sci. U. S. A.* 117, 32528–32534. doi: 10.1073/pnas.1922539117

Sant Anna Iwanicki, N., Delalibera Júnior, I., de Carvalho, L. L. B., Eilenberg, J., and De Fine Licht, H. H. (2023). Comparative transcriptomics of growth metabolism and virulence reveal distinct morphogenic profiles of yeast-like cells and hyphae of the fungus metarhizium rileyi. *Fungal Genet. Biol.* 164, 103766. doi: 10.1016/j.fgb.2022.103766

Sarsaiya, S., Jain, A., Shi, J., and Chen, J. (2021). “1 - fungi endophytes for biofactory of secondary metabolites: genomics and metabolism,” in *Biocontrol agents and secondary metabolites*. Ed. S. Jogaiah (Sawston, Cambridge: Woodhead Publishing), 1–21.

Schalamun, M., and Schmoll, M. (2022). Trichoderma – genomes and genomics as treasure troves for research towards biology, biotechnology and agriculture. *Front. Fungal Biol.* 3. doi: 10.3389/ffunb.2022.1002161

Seidl, V., Seibel, C., Kubicek, C. P., and Schmoll, M. (2009). Sexual development in the industrial workhorse trichoderma reesei. *Proc. Natl. Acad. Sci. U. S. A.* 106, 13909–13914. doi: 10.1073/pnas.0904936106

Shapiro, J. A. (2016). Nothing in evolution makes sense except in the light of genomics: read-write genome evolution as an active biological process. *Biology* 5 (2), 27. doi: 10.3390/biology502027

Shave, S., Dawson, J. C., Athar, A. M., Nguyen, C. Q., Kasprovicz, R., and Carragher, N. O. (2023). Phenonaut: multiomics data integration for phenotypic space exploration. *Bioinformatics* 39 (4), btad143. doi: 10.1093/bioinformatics/btad143

Shay, R., Wiegand, A. A., and Trail, F. (2022). Biofilm Formation and Structure in the Filamentous Fungus Fusarium graminearum, a Plant Pathogen. *Mycology* 10 (4), e00171–22. doi: 10.1128/spectrum.00171-22

Shen, L., Porée, F.-H., Gaslonde, T., Lalucque, H., Chapeland-Leclerc, F., and Ruprich-Robert, G. (2019). Functional characterization of the sterigmatocystin secondary metabolite gene cluster in the filamentous fungus podospira anserina: involvement in oxidative stress response, sexual development, pigmentation and interspecific competitions. *Environ. Microbiol.* 21, 3011–3026. doi: 10.1111/1462-2920.14698

Shi, Y., Wang, H., Wang, J., Liu, X., Lin, F., and Lu, J. (2019). N6-methyladenosine RNA methylation is involved in virulence of the rice blast fungus pyricularia oryzae (syn. magnaporthe oryzae). *FEMS Microbiol. Lett.* 366 (1). doi: 10.1093/femsle/fny286

Sierra-Patev, S., Min, B., Naranjo-Ortiz, M., Looney, B., Konkell, Z., Slot, J. C., et al. (2023). A global phylogenomic analysis of the shitake genus. *Proc. Natl. Acad. Sci. U. S. A.* 120, e2214076120. doi: 10.1073/pnas.2214076120

Sikhakolli, U. R., López-Giraldez, F., Li, N., Common, R., Townsend, J. P., and Trail, F. (2012). Transcriptome analyses during fruiting body formation in fusarium graminearum and fusarium verticillioides reflect species life history and ecology. *Fungal Genet. Biol.* 49, 663–673. doi: 10.1016/j.fgb.2012.05.009

Silar, P., Dauget, J.-M., Gautier, V., Grognet, P., Chablat, M., Hermann-Le Denmat, S., et al. (2019). A gene graveyard in the genome of the fungus podospira comata. *Mol. Genet. Genomics* 294, 177–190. doi: 10.1007/s00438-018-1497-3

Sivashankari, S., and Shanmughavel, P. (2007). Comparative genomics - a perspective. *Bioinformation* 1, 376–378. doi: 10.6026/97320630001376

Slot, J. C., and Rokas, A. (2011). Horizontal transfer of a large and highly toxic secondary metabolic gene cluster between fungi. *Curr. Biol.* 21, 134–139. doi: 10.1016/j.cub.2010.12.020

Stajich, J. E. (2017). Fungal genomes and insights into the evolution of the kingdom. *Fungal Kingdom*, 619–633. doi: 10.1128/9781555819583.ch29

Stukenbrock, E. H. (2013). Evolution, selection and isolation: a genomic view of speciation in fungal plant pathogens. *New Phytol.* 199, 895–907. doi: 10.1111/nph.12374

Stukenbrock, E. H., and Croll, D. (2014). The evolving fungal genome. *Fungal Biol. Rev.* 28, 1–12. doi: 10.1016/j.fbr.2014.02.001

Suga, H., and Hyakumachi, M. (2004). “6 - genomics of phytopathogenic fusarium,” in *Applied mycology and biotechnology*. Eds. D. K. Arora and G. G. Khachatourians (Amsterdam, Netherlands: Elsevier), 161–189.

Sun, Y., Corcoran, P., Menkis, A., Whittle, C. A., Andersson, S. G. E., and Johannesson, H. (2012). Large-Scale introgression shapes the evolution of the mating-type chromosomes of the filamentous ascomycete neurospora tetrasperma. *PLoS Genet.* 8, e1002820. doi: 10.1371/journal.pgen.1002820

Sun, Z.-B., Li, S.-D., Ren, Q., Xu, J.-L., Lu, X., and Sun, M.-H. (2020). Biology and applications of clonostachys rosea. *J. Appl. Microbiol.* 129, 486–495. doi: 10.1111/jam.14625

Sun, Y., Svedberg, J., Hiltunen, M., Corcoran, P., and Johannesson, H. (2017). Large-Scale suppression of recombination predates genomic rearrangements in neurospora tetrasperma. *Nat. Commun.* 8, 1140. doi: 10.1038/s41467-017-01317-6

Sunnerhagen, P., and Piskur, J. (2006). *Comparative genomics: using fungi as models* (Heidelberg, Germany: Springer).

Taylor, J. W., and Berbee, M. L. (2006). Dating divergences in the fungal tree of life: review and new analyses. *Mycologia* 98, 838–849. doi: 10.1080/15572536.2006.11832614

Teichert, I., Dahlmann, T. A., Kück, U., and Nowrousian, M. (2017). RNA Editing during development occurs in distantly related filamentous ascomycetes. *Genome Biol. Evol.* 9, 855–868. doi: 10.1093/gbe/evx052

Teichert, I., Nowrousian, M., Pöggeler, S., and Kück, U. (2014). The filamentous fungus sordaria macrospora as a genetic model to study fruiting body development. *Adv. Genet.* 87, 199–244. doi: 10.1016/B978-0-12-800149-3.00004-4

Teichert, I., Pöggeler, S., and Nowrousian, M. (2020). Sordaria macrospora: 25 years as a model organism for studying the molecular mechanisms of fruiting body development. *Appl. Microbiol. Biotechnol.* 104, 3691–3704. doi: 10.1007/s00253-020-10504-3

Teli, B., Purohit, J., Rashid, M. M., Jailani, A. A. K., and Chattopadhyay, A. (2020). Omics insight on fusarium head blight of wheat for translational research perspective. *Curr. Genomics* 21, 411–428. doi: 10.2174/138920292199920062022631

Thananusak, R., Laoteng, K., Raethong, N., Koffas, M., and Vongsangnak, W. (2022). Dissecting metabolic regulation in mycelial growth and fruiting body developmental stages of cordyceps militaris through integrative transcriptome analysis. *Biotechnol. Bioprocess Eng.*, 1–13. doi: 10.1007/s12257-022-0207-5

Tiquia-Arashiro, S. M., and Grube, M. (2019). *Fungi in extreme environments: ecological role and biotechnological significance* (Heidelberg, Germany: Springer).

Trail, F. (2007). Fungal cannons: explosive spore discharge in the ascomycota. *FEMS Microbiol. Lett.* 276, 12–18. doi: 10.1111/j.1574-6968.2007.00900.x

Trail, F. (2013). “Sex and fruiting in fusarium,” in *Fusarium: genomics, molecular and cellular biology*. Eds. D. Brown and R. Proctor (Norwich, UK: Horizon Scientific Press and Caister Academic Press).

- Trail, F., Gaffoor, I., and Vogel, S. (2005). Ejection mechanics and trajectory of the ascospores of *Gibberella zeae* (anamorph *Fusarium graminearum*). *Fungal Genet. Biol.* 42, 528–533. doi: 10.1016/j.fgb.2005.03.008
- Trail, F., and Seminara, A. (2014). The mechanism of ascus firing – merging biophysical and mycological viewpoints. *Fungal Biol. Rev.* 28, 70–76. doi: 10.1016/j.fbr.2014.07.002
- Trail, F., Wang, Z., Stefanko, K., Cubba, C., and Townsend, J. P. (2017). The ancestral levels of transcription and the evolution of sexual phenotypes in filamentous fungi. *PLoS Genet.* 13, e1006867. doi: 10.1371/journal.pgen.1006867
- Tworzydło, W., and Bilinski, S. M. (2019). *Evo-devo: non-model species in cell and developmental biology* (Cham, Switzerland: Springer Nature).
- Valero-Jiménez, C. A., Faino, L., Spring In't Veld, D., Smit, S., Zwaan, B. J., and van Kan, J. A. L. (2016). Comparative genomics of *Beauveria bassiana*: uncovering signatures of virulence against mosquitoes. *BMC Genomics* 17, 986. doi: 10.1186/s12864-016-3339-1
- Vasaikar, S. V., Savage, A. K., Gong, Q., Swanson, E., Talla, A., Lord, C., et al. (2023). A comprehensive platform for analyzing longitudinal multi-omics data. *Nat. Commun.* 14, 1684. doi: 10.1038/s41467-023-37432-w
- Vaughan, M., Backhouse, D., and Ponte, E. M. D. (2016). Climate change impacts on the ecology of *Fusarium graminearum* species complex and susceptibility of wheat to fusarium head blight: a review. *World Mycotoxin J.* 9, 685–700. doi: 10.3920/WMJ2016.2053
- Vaughn, J. N., Branham, S. E., Abernathy, B., Hulse-Kemp, A. M., Rivers, A. R., Levi, A., et al. (2022). Graph-based pangenomics maximizes genotyping density and reveals structural impacts on fungal resistance in melon. *Nat. Commun.* 13, 7897. doi: 10.1038/s41467-022-35621-7
- Vesper, S., McKinstry, C., Ashley, P., Haugland, R., Yeatts, K., Bradham, K., et al. (2007). Quantitative PCR analysis of molds in the dust from homes of asthmatic children in north Carolina. *J. Environ. Monit.* 9, 826–830. doi: 10.1039/b704359g
- Virágh, M., Merényi, Z., Cséretics, Á., Földi, C., Sahu, N., Liu, X.-B., et al. (2022). Evolutionary morphogenesis of sexual fruiting bodies in basidiomycota: toward a new evo-devo synthesis. *Microbiol. Mol. Biol. Rev.* 86, e0001921. doi: 10.1128/MMBR.00019-21
- Vittorelli, N., Rodríguez de la Vega, R. C., Snirc, A., Levert, E., Gautier, V., Lalanne, C., et al. (2023). Stepwise recombination suppression around the mating-type locus in an ascomycete fungus with self-fertile spores. *PLoS Genet.* 19, e1010347. doi: 10.1371/journal.pgen.1010347
- Wang, B., Cai, P., Sun, W., Li, J., Tian, C., and Ma, Y. (2015). A transcriptomic analysis of *Neurospora crassa* using five major crop residues and the novel role of the sporulation regulator *rca-1* in lignocellulase production. *Biotechnol. Biofuels* 8, 21. doi: 10.1186/s13068-015-0208-0
- Wang, X., Cui, Y., Fan, F., Song, Y., Ren, J., Meng, Q., et al. (2010). Phylogenetic, carbendazim sensitivity and mycotoxin genotype analyses of *Fusarium graminearum* complex species isolated from wheat fusarium head blight in China. *J. Phytopathol.* 158, 576–578. doi: 10.1111/j.1439-0434.2009.01662.x
- Wang, Z., Gudibanda, A., Ugwuowo, U., Trail, F., and Townsend, J. P. (2018a). Using evolutionary genomics, transcriptomics, and systems biology to reveal gene networks underlying fungal development. *Fungal Biol. Rev.* 32, 249–264. doi: 10.1016/j.fbr.2018.02.001
- Wang, X. W., Han, P. J., Bai, F. Y., Luo, A., Bensch, K., Meijer, M., et al. (2022c). Taxonomy, phylogeny and identification of with emphasis on thermophilic species. *Stud. Mycol.* 101, 121–243. doi: 10.3114/sim.2022.101.03
- Wang, Z., Li, N., Li, J., Dunlap, J. C., Trail, F., and Townsend, J. P. (2016c). The fast-evolving *phy-2* gene modulates sexual development in response to light in the model fungus *Neurospora crassa*. *MBio* 7, e02148. doi: 10.1128/mBio.02148-15
- Wang, R., Liu, C., Jiang, X., Tan, Z., Li, H., Xu, S., et al. (2022b). The newly identified trichoderma *harzianum* *partitivir* (ThPV2) does not diminish spore production and biocontrol activity of its host. *Viruses* 14, 1532. doi: 10.3390/v14071532
- Wang, X. W., Lombard, L., Groenewald, J. Z., Li, J., Videira, S. I. R., Samson, R. A., et al. (2016b). Phylogenetic reassessment of the chaetomium *globosum* species complex. *Persoonia - Mol. Phyl. Evol. Fungi* 36, 83–133. doi: 10.3767/003158516X689657
- Wang, Z., Lopez-Giraldez, F., Lehr, N., Farré, M., Common, R., Trail, F., et al. (2014). Global gene expression and focused knockout analysis reveals genes associated with fungal fruiting body development in *Neurospora crassa*. *Eukaryot. Cell* 13, 154–169. doi: 10.1128/EC.00248-13
- Wang, Z., Lopez-Giraldez, F., Slot, J., Yarden, O., Trail, F., and Townsend, J. P. (2022d). Secondary metabolism gene clusters exhibit increasingly dynamic and differential expression during asexual growth, conidiation, and sexual development in *Neurospora crassa*. *mSystems* 7, e002322. doi: 10.1128/mSystems.00232-22
- Wang, Z., López-Giráldez, F., Wang, J., Trail, F., and Townsend, J. P. (2019a). Integrative activity of mating loci, environmentally responsive genes, and secondary metabolism pathways during sexual development of *Chaetomium globosum*. *MBio* 10 (6), e02119–19. doi: 10.1128/mBio.02119-19
- Wang, Z., Miguel-Rojas, C., Lopez-Giraldez, F., Yarden, O., Trail, F., and Townsend, J. P. (2019b). Metabolism and development during conidial germination in response to a carbon-Nitrogen-Rich synthetic or a natural source of nutrition in *Neurospora crassa*. *MBio* 10 (2), e00192–19. doi: 10.1128/mBio.00192-19
- Wang, J.-H., Ndoye, M., Zhang, J.-B., Li, H.-P., and Liao, Y.-C. (2011). Population structure and genetic diversity of the *Fusarium graminearum* species complex. *Toxins* 3, 1020–1037. doi: 10.3390/toxins3081020
- Wang, Z., Nilsson, R. H., James, T. Y., Dai, Y., and Townsend, J. P. (2016d). “Future perspectives and challenges of fungal systematics in the age of big data,” in *Biology of microfungi*. Ed. D.-W. Li (Cham: Springer International Publishing), 25–46.
- Wang, J. B., St Leger, R. J., and Wang, C. (2016a). Advances in genomics of entomopathogenic fungi. *Adv. Genet.* 94, 67–105. doi: 10.1016/bs.adgen.2016.01.002
- Wang, J., Wang, Q., Huang, P., Qu, Y., Huang, Z., Wang, H., et al. (2022a). An appressorium membrane protein, Pams1, controls infection structure maturation and virulence via maintaining endosomal stability in the rice blast fungus. *Front. Plant Sci.* 13, 955254. doi: 10.3389/fpls.2022.955254
- Wang, Z., Wang, Y.-W., Kasuga, T., Hassler, H., Lopez-Giraldez, F., Dong, C., et al. (2023). The “evol” is in the details: a rummage-region model for the origins of lineage-specific elements via gene duplication, relocation, and regional rearrangement in *Neurospora crassa*. doi: 10.22541/au.168259153.37423684/v1
- Wang, Z., Wang, Y., Kasuga, T., Lopez-Giraldez, F., Zhang, Y., Zhang, Z., et al. (2022e). Orphan genes are clustered with all-recognition loci and may be involved in incompatibility and speciation in *Neurospora*. *bioRxiv* 2022, 495464. doi: 10.1101/2022.06.10.495464
- Wang, Z., Wang, J., Li, N., Li, J., Trail, F., Dunlap, J. C., et al. (2018b). Light sensing by opsins and fungal ecology: NOP-1 modulates entry into sexual reproduction in response to environmental cues. *Mol. Ecol.* 27, 216–232. doi: 10.1111/mec.14425
- Wibberg, D., Stadler, M., Lambert, C., Bunk, B., Spröer, C., Rückert, C., et al. (2020). High quality genome sequences of thirteen hypoxylaceae (Ascomycota) strengthen the phylogenetic family backbone and enable the discovery of new taxa. *Fungal Divers.* 106, 7–28. doi: 10.1007/s13225-020-00447-5
- Wicklow, D. T. (2018). Fire as an environmental cue initiating ascomycete development in a tallgrass prairie. *Mycologia* 67 (4), 852–862. doi: 10.1080/00275514.1975.12019813
- Winkler, R. (2020). Processing metabolomics and proteomics data with open software: a practical guide. *R. Soc. Chem.* doi: 10.1039/9781788019880
- Woo, S. L., Hermosa, R., Lorito, M., and Monte, E. (2023). *Trichoderma*: a multipurpose, plant-beneficial microorganism for eco-sustainable agriculture. *Nat. Rev. Microbiol.* 21, 312–326. doi: 10.1038/s41579-022-00819-5
- World Health Organization (2022). *WHO fungal priority pathogens list to guide research, development and public health action*. (WHO).
- Wu, V. W., Thieme, N., Huberman, L. B., Dietschmann, A., Kowbel, D. J., Lee, J., et al. (2020). The regulatory and transcriptional landscape associated with carbon utilization in a filamentous fungus. *Proc. Natl. Acad. Sci. U. S. A.* 117, 6003–6013. doi: 10.1073/pnas.1915611117
- Xiao, G., Ying, S.-H., Zheng, P., Wang, Z.-L., Zhang, S., Xie, X.-Q., et al. (2012). Genomic perspectives on the evolution of fungal entomopathogenicity in *Beauveria bassiana*. *Sci. Rep.* 2, 483. doi: 10.1038/srep00483
- Xie, N., Ruprich-Robert, G., Chapeland-Leclerc, F., Coppin, E., Lalucque, H., Brun, S., et al. (2017). Inositol-phosphate signaling as mediator for growth and sexual reproduction in *Podospora anserina*. *Dev. Biol.* 429, 285–305. doi: 10.1016/j.ydbio.2017.06.017
- Yan, Y., Li, Y., Wang, W.-J., He, J.-S., Yang, R.-H., Wu, H.-J., et al. (2017). Range shifts in response to climate change of *Ophiocordyceps sinensis*, a fungus endemic to the Tibetan plateau. *Biol. Conserv.* 206, 143–150. doi: 10.1016/j.biocon.2016.12.023
- Yarden, O. (2007). Model systems for studying the biology of filamentous fungi: rumors of their death should be postponed. *Phytoparasitica* 35, 111–115. doi: 10.1007/bf02981102
- Yarden, O. (2016). Model fungi: engines of scientific insight. *Fungal Biol. Rev.* 30, 33–35. doi: 10.1016/j.fbr.2016.05.002
- Yarden, O., Ebbola, D. J., Freeman, S., Rodriguez, R. J., and Dickman, M. B. (2003). Fungal biology and agriculture: revisiting the field. *Mol. Plant Microbe Interact.* 16, 859–866. doi: 10.1094/MPMI.2003.16.10.859
- You, J., Zhou, K., Liu, X., Wu, M., Yang, L., Zhang, J., et al. (2019). Defective RNA of a novel mycovirus with high transmissibility detrimental to biocontrol properties of *spp. Microorganisms* 7 (11), 507. doi: 10.3390/microorganisms7110507
- Yu, J., and Kim, K.-H. (2020). Exploration of the interactions between mycoviruses and *Fusarium graminearum*. *Adv. Virus Res.* 106, 123–144. doi: 10.1016/bs.avir.2020.01.004
- Yun, S.-H., Lee, S. H., So, K.-K., Kim, J.-M., and Kim, D.-H. (2016). Incidence of diverse dsRNA mycoviruses in *Trichoderma* spp. causing green mold disease of shiitake *Lentinula edodes*. *FEMS Microbiol. Lett.* 363 (19), fnw220. doi: 10.1093/femsle/fnw220
- Zámocký, M., Tafer, H., Chovanová, K., Lopandic, K., Kamlárová, A., and Obinger, C. (2016). Genome sequence of the filamentous soil fungus *Chaetomium cochliodes* reveals abundance of genes for heme enzymes from all peroxidase and catalase superfamilies. *BMC Genomics* 17, 763. doi: 10.1186/s12864-016-3111-6
- Zeilinger, S., and Omann, M. (2007). *Trichoderma* biocontrol: signal transduction pathways involved in host sensing and mycoparasitism. *Gene Regul. Syst. Bio.* 1, 227–234. doi: 10.4137/GRSB.S397
- Zhang, Y. (2019). *Evolution of the pathogenic fusarium oxysporum through the lens of comparative genomics*. Doctoral Dissertations. 1790. doi: 10.7275/15240939

- Zhang, N., Castlebury, L. A., Miller, A. N., Huhndorf, S. M., Schoch, C. L., Seifert, K. A., et al. (2006). An overview of the systematics of the sordariomycetes based on a four-gene phylogeny. *Mycologia* 98, 1076–1087. doi: 10.1080/15572536.2006.11832635
- Zhang, N., and Wang, Z. (2015). “3 pezizomycotina: sordariomycetes and leotiomycetes” in *Systematics and evolution. The Mycota (A comprehensive treatise on fungi as experimental systems for basic and applied research)*. Eds. D. McLaughlin and J. Spatafora (Berlin, Heidelberg: Springer) 7B, 57–88. doi: 10.1007/978-3-662-46011-5_3
- Zheng, P., Xia, Y., Xiao, G., Xiong, C., Hu, X., Zhang, S., et al. (2011). Genome sequence of the insect pathogenic fungus *Cordyceps militaris*, a valued traditional Chinese medicine. *Genome Biol.* 12, R116. doi: 10.1186/gb-2011-12-11-r116
- Zhu, S.-D., Huang, L.-Q., Guo, L.-P., Ma, X.-T., Hao, Q.-X., Le, Z.-Y., et al. (2017). [Climate change impacts on yield of *Cordyceps sinensis* and research on yield prediction model of *C. sinensis*]. *Zhongguo Zhong Yao Za Zhi* 42, 1281–1286. doi: 10.19540/j.cnki.cjcmm.2017.0055