Tunable spin and valley excitations of correlated insulators in Γ -valley moiré bands

Benjamin A. Foutty, 1, 2, * Jiachen Yu, 1, 3, * Trithep Devakul, 4, * Carlos R. Kometter, 1, 2 Yang Zhang, ^{4,5} Kenji Watanabe, ⁶ Takashi Taniguchi, ⁷ Liang Fu, ⁴ and Benjamin E. Feldman^{1,2,8,†} ¹Geballe Laboratory for Advanced Materials, Stanford, CA 94305, USA ²Department of Physics, Stanford University, Stanford, CA 94305, USA ³Department of Applied Physics, Stanford University, Stanford, CA 94305, USA ⁴Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA ⁵Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA ⁶ Research Center for Functional Materials, National Institute for Material Science, 1-1 Namiki, Tsukuba 305-0044, Japan ⁷International Center for Materials Nanoarchitectonics, National Institute for Material Science, 1-1 Namiki, Tsukuba 305-0044, Japan ⁸Stanford Institute for Materials and Energy Sciences, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA

Moiré superlattices formed from transition metal dichalcogenides (TMDs) support a variety of quantum electronic phases that are highly tunable using applied electromagnetic fields. While the valley degree of freedom affects optoelectronic properties in the constituent TMDs, it has yet to be fully explored in moiré systems. Here, we establish twisted double bilayer WSe2 as an experimental platform to study electronic correlations within Γ-valley moiré bands. Through local and global electronic compressibility measurements, we identify charge-ordered phases at multiple integer and fractional moiré fillings. By measuring the magnetic field dependence of their energy gaps and the chemical potential upon doping, we reveal spin-polarized ground states with spin polaron quasiparticle excitations. In addition, an applied displacement field induces a metal-insulator transition driven by tuning between Γ - and K-valley moiré bands. Our results demonstrate control over the spin and valley character of the correlated ground and excited states in this system.

INTRODUCTION

5

6

9

10

11

12

13

14

15

16

17

18

19

20

21 22

23

24

25

Control over the electronic band structure and in-26 ternal quantum degrees of freedom in solid state materials is highly desirable to engineer novel functionality. Moiré van der Waals heterostructures, in which an interlayer twist and/or mismatch between lattice constants produces a long-wavelength spatial modulation, 32 have attracted great interest due to their ability to host flat bands in which electronic interactions dominate over kinetic energy [3, 5, 14]. Moiré superlattices engineered from semiconducting transition metal dichalcogenides (TMDs) are a particularly flexible platform for investigating correlated electronic states begle [4, 40, 41, 47, 48]. Multiple homo- and heterobilayer systems have been explored to date, revealing correlation-driven charge transfer and/or Mott insulating states at integer moiré fillings [18, 19, 28, 32, 37, 45] and a series of generalized Wigner crystals at fractional fillings [6, 13, 16, 19, 28, 33, 44]. These correlated elec-54 correlated materials [14, 26, 32, 40]. This has motivated 85 in the context of moiré bands also remains an open ques-

55 careful measurement of the spin ordering of the ground 56 states [12, 34, 38, 45]. However, experimental studies 57 have primarily focused on low magnetic fields, and little 58 is known about the nature of the lowest energy excitations or their field dependence.

The valley degree of freedom, which captures the mo-61 mentum of the low-energy states, also plays an important 62 role in TMDs, and the valley character of the moiré bands 63 derives from that of the constituent TMDs [4, 21, 36, 42]. $_{64}$ If the moiré bands are localized around the Γ -valley, then 65 these bands will be spin degenerate but have no valley 66 degeneracy. This presents a qualitatively distinct phys-67 ical system relative to K-valley TMDs, which have two 68 degenerate spin-valley-locked bands [4, 37]. Due to the cause the resulting physics can be tuned by adjusting 69 structure of TMDs, there are also two distinct stack-TMD composition and stacking orientation, while flat 70 ings of their moiré superlattices [29]. Twisted Γ -valley bands emerge without stringent constraints on twist an- 71 moiré TMD systems with AA (i.e. near 3R) stacking are 72 predicted to realize effective Hubbard models in honev-73 comb and Kagome geometries [4, 42, 48]. AB (i.e. near 74 2H) stacked Γ-valley systems realize a triangular lattice, 75 which can have flatter bands due to the larger effective ₇₆ mass at Γ , enhancing the importance of interactions. In 77 untwisted TMDs, increasing the number of layers shifts tronic states are often strongly tuned by applied gate 78 the valence band edge from K to Γ [22, 43], and tunvoltages and magnetic fields, which can favor different 79 ing the relative populations of these two valleys with apmany-body ground states and/or modify the underlying so plied electric field was demonstrated in magnetotransport single-particle bands [10, 17, 18, 33?]. As a result, the sum easurements in trilayer WSe₂ [24]. However, few exper-TMD platform is an ideal venue for quantum simula- 12 imental realizations of Γ-valley moiré systems have been tion and study of the Hubbard model, which is thought 83 reported, and their electronic properties are poorly charto capture the essential physics of conventional strongly 84 acterized. The interplay of $\Gamma-K$ -valley-tunable physics tion in experiments.

111

110 states and their excitations.

Γ-VALLEY MOIRÉ BANDS IN tdWSe₂

113 SET, schematically illustrated in Fig. 1a. The SET si- 169 be driven by interactions. Measurements as a function of multaneously and independently probes two related ther- 170 displacement field, discussed in detail below, suggest that 116 ibility $d\mu/dn$ at low a.c. frequency and the chemical 172 a moiré band at the K-valley. Combining this experimenpotential $\mu(n)$ on d.c. timescales (Methods). This tech- 173 tal finding with first-principles calculations that take into 118 nique mitigates the problem of large contact/intrinsic re- 174 account Coulomb and charge transfer energies, we find 119 sistance that often complicates electrical measurements 175 that this K-valley moiré band will be well-localized at 120 in TMD moiré materials [19, 28]. In Sample S1 (Fig. 176 XX' in real space (Supplementary Sec. 3). 1b), we observe a pronounced incompressible state at $n\,\approx\,3.9\times\,10^{12}~{\rm cm}^{-2},$ which we identify with $\nu\,=\,-1.$ A weaker incompressible peak is also evident at $\nu = -\frac{1}{3}$, 177 and it becomes prominent in a small applied out of plane 178 magnetic field B. Both of these features are generically present across a wide range of spatial locations (Supple-179) ₁₃₁ measurement of chemical potential $\mu(n)$ (Fig. 1b, inset), ₁₈₄ early with B throughout the experimentally accessible

139 states, we consider the low energy valence band structure Here we study magnetic and electric displacement 140 at the single-particle level in AB-stacked tdWSe₂. Due field dependence of correlated insulators in Γ-valley hole 141 to strong spin-orbit coupling in WSe₂, interlayer coupling moiré bands in AB-stacked twisted double bilayer WSe₂ 142 between adjacent layers is suppressed by spin mismatch (tdWSe₂; Fig. 1a). We measure the chemical potential 143 at the the K and K' valleys [31, 43, 49]. At the Γ point, $\mu(n)$ and inverse electronic compressibility $d\mu/dn$ both 144 there is no such barrier to interlayer tunneling and hylocally using a scanning single-electron transistor (SET) $_{145}$ bridization can increase the valence band energy at Γ as and globally in dual gated devices. Our measurements re- 146 the layer number increases (Fig. 1c). This has been both veal charge-ordered states at integer and fractional moiré 147 predicted in theoretical calculations and observed in exfilling factors $\nu = -1, -3$ and $-\frac{1}{3}$, where $\nu = -1$ corre- 148 periments in few-layer WSe₂ [20, 24, 39]. Additionally, sponds to one hole per moiré unit cell. The measured 149 a previous ARPES measurement of tdWSe2 found the gaps grow linearly with perpendicular magnetic field in 150 valence band maximum to occur at Γ rather than K [2]. the low-field regime, indicating spin-polarized ground 151 We confirm this picture by performing density funcstates. The $\nu = -\frac{1}{3}$ gap saturates at high fields, and the 152 tional theory (DFT) calculations at zero displacement behavior of the chemical potential in its vicinity provides 153 field on the high symmetry stackings of tdWSe₂ (Fig. evidence that itinerant spin polarons, bound states of 154 1d, Methods). The respective band structures (Fig. 1e) added electrons and spin-flipped holes, are the lowest en- 155 demonstrate that the valence band maximum occurs at ergy charged excitations [7]. The thermodynamic energy $_{156}$ the Γ valley in the MM' stacking and is separated by gaps at integer fillings decrease monotonically with elec- 157 roughly 100 meV from the energies at other stackings. tric displacement field, driven by a single-particle band 158 Using the energies obtained from DFT, we develop an crossing between the lowest energy Γ and K moiré bands. 159 effective continuum model describing moiré bands from Our experiments establish a new approach to engineer $_{160}$ both Γ and K valley bands involving all four layers (Sup-Γ-valley moiré bands and fully characterize the real- and 161 plementary Sec. 3). This continuum model also allows us momentum-space localization of the correlated ground 162 to later incorporate the effect of a perpendicular displace-163 ment field as an effective layer potential difference. The 164 combination of large effective mass at Γ and strong moiré 165 potential results in a very flat first moiré band which is well localized at the MM' sites (Fig. 1f, Supplementary 167 Sec. 3). Due to the two-fold spin degeneracy of the moiré We focus first on local measurements using a scanning 168 bands, the experimentally observed gap at $\nu=-1$ must modynamic quantities, the inverse electronic compress- 171 the lowest-energy charge excitations at $\nu=-1$ populate

MAGNETIC FIELD DEPENDENCE OF CHARGE-ORDERED STATES

The evolution of the charge gaps in a perpendicular mentary Sec. 1), and from the densities at which they oc- 180 magnetic field clarifies the spin physics of the correlated cur, we extract a twist angle of 3.4° (Methods). The cor- 181 insulating ground states and their charged excitations. responding thermodynamic gap Δ_{ν} , defined as the step 182 Both the $\nu = -\frac{1}{3}$ and $\nu = -1$ gaps grow in a magnetic size in $\mu(n)$ at filling ν , can be determined from the d.c. 183 field (Fig. 2a-b, Supplementary Sec. 2). Δ_{-1} grows linyielding $\Delta_{-1} \approx 6 \text{ meV}$ at B = 0 (Supplementary Sec. 2). 185 field range, whereas $\Delta_{-\frac{1}{8}}$ increases up to B = 7 T before We also measured a similar gap at $\nu = -1$ in a second 186 saturating or decreasing slightly at high fields (Fig. 2c). device (Sample S2) with a smaller twist angle of 2.6°. 187 The regime of linear growth reflects Zeeman energy shifts, Both samples show very little ($< 0.1^{\circ}$) twist angle vari- 188 indicating that the insulating state is spin-polarized and ation (Supplementary Sec. 1), so we focus on a single 189 a spin-flip is involved in the lowest energy particle-hole spatial location in each device throughout the main text. 190 excitation. This is consistent with prior reports of small As a first step toward understanding the correlated $_{191}$ exchange coupling J in moiré systems, such that a small




FIG. 1. Γ-valley moiré bands in twisted double bilayer WSe₂ (tdWSe₂). a, Schematic of device and measurement geometry. b. Local measurement of inverse electronic compressibility $d\mu/dn$ of Sample S1 as a function of moiré filling factor ν at magnetic fields B=0 T and B=2 T. The incompressible peak in $d\mu/dn$ is artificially enhanced in the a.c. measurement. Inset, simultaneous d.c. measurement of chemical potential $\mu(n)$ through the $\nu=-1$ gap. c. Schematic of valence band edges on each layer of AB-stacked tdWSe₂ before consideration of moiré bands. Red (blue) denotes spin up (down) at the K and K' points, while purple bands at Γ are spin degenerate. Arrows show interlayer hybridization, which can occur at Γ but is limited by spin mismatch at K and K'. d, High symmetry stackings of tdWSe₂: MM', in which the W atoms are aligned vertically on the two internal layers of the 4-layer heterostructure, MX', in which all four layers realize 2H stacking, and XX', in which the Se atoms on the two internal layers are aligned vertically. e, Density functional theory calculated bands at the high symmetry points shown in \mathbf{d} , with reference energy E=0 chosen to be the absolute vacuum level. The valence band edge occurs at the MM' site at Γ. f, Wannier orbitals of the lowest energy moiré band for 3.4° tdWSe₂ as calculated from a continuum model on the bands shown in e. States are well-localized at the MM' sites (purple dot). There is no charge density on the other high-symmetry sites MX' (blue) or XX' (teal).

192 magnetic field is sufficient to polarize the spins in the 207 est energy excitation just below the gap is the addition 193 gapped ground state [32]. However, we note that the ob- 208 of a hole of opposite spin at low fields, before Zeeman 194 servation of spin-polarized correlated insulators in a mag- 209 coupling overcomes exchange to favor adding an aligned 195 netic field differs from reports in twsited bilayer WSe₂ 210 hole (Supplementary Sec. 5). In this scenario, the ex-196 [10]. We assume that the insulating state at $\nu = -\frac{1}{3}$ is 211 citation above the gap would be the removal of a single a $\sqrt{3} \times \sqrt{3}$ generalized Wigner crystal, so that both of 212 hole, independent of magnetic field. these states realize triangular lattices in real space (Fig. 213 2e-f). 199

204 and therefore requires us to take into account the mag- 219 2i, black inset). Above the gap, the lowest energy exci-205 netic field dependence of the excitations both above and 220 tation below a critical field B^* involves removing a hole

However, recent theory suggests that more exotic spin 214 polaron quasiparticles may be relevant in a half-filled tri-The saturating gap at $\nu = -\frac{1}{3}$ indicates a change in 215 angular lattice with a spin-polarized ground state [7]. In the nature of the excitations at high fields. The ther- 216 this second scenario, the excitation below the gap remodynamic gap that we measure is equivalent to the 217 mains the same at all magnetic fields: addition of an particle-hole excitation energy (Supplementary Sec. 4), 218 aligned hole at an interstitial (charge-transfer) site (Fig. 206 below the gap. One possible explanation is that the low- 221 and flipping the spin of an adjacent remaining hole on




FIG. 2. Magnetic field dependence of correlated insulating ground states and their excitations. a, $d\mu/dn$ as a function ν (bottom axis), density n (top axis), and of out-of-plane magnetic field B. **b**, Linecuts of d.c. measured $\mu(\nu)$ around $\nu = -1$ at various magnetic fields. c, Extracted charge gaps at $\nu = -1$, $\nu = -\frac{1}{3}$ in Sample S1 and $\nu = -1$ in Sample S2. Dashed lines show linear fits to the data. All increase at low field, while $\Delta_{-1/3}$ saturates and decreases slightly beginning at B=7 T. d, Chemical potential μ_+ and μ_- measured at $\nu=-\frac{1}{3}\pm\epsilon$ for $\epsilon>0$ on either side of the measured gap. On the lower hole doping side of the gap, (μ_+) , the slope is discontinuous in B, whereas at higher hole doping the behavior is smooth. e-f, Schematic of real space spin ordering at $\nu = -1$ (e) and $\nu = -\frac{1}{3}$ (f). Ground state hole spins are shown in black, while the lowest energy hole excitations are shown in orange. g,h, Schematic of the expected gaps as a function of B. At low fields, the gap should increase due to the creation of spin polarons on the lower hole doping side of the gap. Above a critical field B^* , spin polarons are no longer favored and the gap is constant in B. This crossover point should be higher at $\nu = -1$ due to higher effective hopping t. i, Schematic of chemical potential of states at half filling of a charge transfer triangular lattice as a function of magnetic field B. Black line and inset denote the lowest energy excitation at higher hole density, adding an aligned hole at an interstitial. Red line and inset denote the spin polaron excitation at lower hole density, which is favored at intermediate fields $B < B^*$, while the blue line and inset denote the excitation of removing a single hole. Solid circles with spins denote holes, while dashed circles denote the absence of a hole.

other spins (Fig. 2i, blue inset).

To distinguish between these two pictures, both of which predict a saturation of the gap size, we examine the 230 behavior of the chemical potential as we dope away from 231 the charge ordered state. The first scenario would involve

222 the lattice, creating an itinerant spin polaron quasipar- 232 a sharp change in the behavior of the chemical potential 223 ticle (Fig. 2i, red inset) with effective spin $\frac{3}{2}$. At mag- 233 μ_- below the gap (Supplementary Sec. 5). In contrast, 224 netic fields exceeding B*, formation of spin polarons is 234 the spin polaron scenario would lead to a change in the no longer favored due to Zeeman coupling, and the exci- 235 chemical potential above the gap, μ_{+} : the theoretical tation involves removing a bare hole without flipping any 236 chemical potentials for each excitation are shown in Fig. 237 2i. In Fig. 2d, we show the magnetic field dependence of 238 the experimentally measured chemical potentials μ_+ and ₂₃₉ μ_{-} on either side of the $\nu = -\frac{1}{3}$ gap (Supplementary ²⁴⁰ Sec. 6). While μ_- is completely smooth, μ_+ changes slope sharply around B = 7 T, consistent only with the evidence for these composite quasiparticles.

there as well (Supplementary Sec. 6).

Finally, we fit the linear growth of the $\nu = -1$ and $\nu = 309$ to noise for |D| < 0.1 V/nm (Fig. 3e) $-\frac{1}{3}$ gaps in a magnetic field with an effective g-factor g^* , defined by $\Delta_{\nu}(B) = \Delta_{\nu}(0) + g^* \mu_B B$. In Sample S1, both 259 gaps at $\nu = -1$ and $\nu = -\frac{1}{3}$ exhibit a similar effective g- 310 factor $g^* = 4.3 \pm 0.8$. In Sample S2, we measure a larger 311 that displacement fields can drive a continuous metal- $_{261}$ $g^*=5.8\pm1.5$ for $\nu=-1$. These values exceed the bare $_{312}$ insulator transition by changing the bandwidth of the ₂₆₂ spin contribution $g_s = 2$ that we would expect from the ₃₁₃ underlying moiré bands [10, 17]. In those experiments, the observation of $g^* > 2$ at $\nu = -\frac{1}{3}$ likely stems from 321 placement field. that effect [11, 23, 25, 35].

Away from the gapped states, our measurements also reveal regions with negative $d\mu/dn$. This is visible on ei-283 side of the correlated insulator gaps.

DISPLACEMENT FIELD TUNING OF MOIRÉ BANDS

284

285

²⁹⁴ Methods). We studied two samples (C1 and C2) using ³⁴⁶ are to K-valley moiré bands.

²⁴² spin polaron picture. We therefore conclude that spin ²⁹⁵ this technique, with twist angles of 2.2° and 3.1°, respecpolarons are favored at lower hole doping of the $\nu = -\frac{1}{3}$ 296 tively. As hole density is increased beyond the mobility insulator below B = 7 T, providing the first experimental 297 edge at $|n| \approx 0.8 \times 10^{12} \text{cm}^{-2}$, a broad region of nega-298 tive $d\mu/dn$ is present before an insulating gap appears at Theoretically, the critical field B^* scales with the hop- $^{299}\nu = -1$ (Fig. 3b, Supplementary Sec. 9). In both sam-₂₄₇ ping t on the triangular lattice [7]. Because the sites ₃₀₀ ples, Δ_{-1} reaches a maximum around 30 meV at D=0are closer together for the insulating state at $\nu = -1$ as 301 and closes monotonically with |D|, disappearing around compared to the generalized Wigner crystal at $\nu = -\frac{1}{3}$, 302 $|D| \approx 0.2 - 0.4$ V/nm (Fig. 3d-e). Despite the large the effective t will be higher at $\nu = -1$, such that the 303 apparent difference in gap sizes, this is largely consiscrossover field B^* at $\nu = -1$ is inaccessible in our mea- 304 tent with the SET measurements, where the single gate surements (Fig. 2e-h). Nonetheless, the observed chemi- 305 geometry leads to a nonzero displacement field (Supplecal potential evolution on either side of the $\nu=-1$ gap 306 mentary Sec. 10). Qualitatively similar behavior occurs suggests that the spin polaron excitations are relevant 307 at $\nu = -3$ (Fig. 3c), whose gap also vanishes at high 308 displacement field but shows a plateau within the signal

Previous works in TMD moiré systems have shown predicted excitation (Supplementary Sec. 4). At $\nu = -1$, 314 the transport gap closes over a small range of displacethis likely reflects additional contributions from orbital 315 ment field (D < 0.1 V/nm) and does not vary strongly effects [1, 11, 24]. At $\nu = -\frac{1}{3}$, we do not expect an orbital 316 with D outside of that range. At $\nu = -1$ in our device, component as the gap does not involve any change in 317 the gap closes smoothly over a much larger range of D, moiré band. However, exchange enhancement of q^* has 318 and changes continuously all the way to D=0. Simultabeen observed previously in TMDs and two-dimensional 319 neously, our continuum modeling suggests the bandwidth electron gases (2DEGs), particularly at low densities, and $_{320}$ of the Γ moiré bands should be largely unaffected by dis-

Instead, our measurements and modeling are consisther side of the gap at $\nu = -1$ in Fig. 2b, and occurs over 323 tent with the lowest energy moiré band switching from Γ wide range of fillings, a signature of strong Coulomb in- 324 to K at sufficient layer potential imbalance (Fig. 3f, Supteractions [8, 9]. This negative compressibility can be un- $_{325}$ plementary Sec. 5). Because the $\nu = -1$ gap is set by the lerstood through consideration of exchange interactions, $_{326}$ charge transfer excitation from Γ to K moiré bands, tunwhich are sizable relative to the density of states contri- 327 ing the relative moiré band energies leads to a continual bution to $\mu(n)$ in this system (see Supplementary Secs. 328 change in the measured gap. Past the critical displace-7-8 for further discussion). The overall size of negative 329 ment field where the K-valley moiré band crosses, we do compressibility increases in a magnetic field, suggesting 330 not measure an incompressible state at $\nu = -1$, likely paramagnetism through much of the density range out- 331 due to band overlap and/or the greater dispersion of the 332 K-valley moiré band. The measured gap sizes and critical displacement field at $\nu = -1$ show weak dependence 334 on twist angle: Samples C1 and C2 have twist angles that differ by about 1° but the extracted gap at D=0336 is similar across devices. This is because the gap size 337 and displacement field dependence is largely set by the We next discuss the displacement field dependence 338 relative energy of the valence band edge of Γ and K at of the correlated insulators, which has previously been 339 the band edge (Fig. 1e) and the effect that displaceshown to induce particularly strong effects in homobi- 340 ment field has on those energies, rather than details of layers where states are extended across layers [37, 45]. 341 the moiré band structure itself. The continuous closing To study this thermodynamically in tdWSe₂, we fabri- 342 of the gap from its maximum at D=0 provides expercate samples with top and bottom gates and measure 343 imental evidence that the lowest energy moiré bands at the sample-gate capacitance as a function of moiré fill- 344 low displacement fields are from the Γ-valley and that ₂₉₃ ing factor and electrical displacement field D (Fig. 3a, ₃₄₅ even at D=0, the lowest energy excitations at $\nu=-1$




FIG. 3. Displacement field dependence of correlated insulators. a, Schematic of a dual-gated capacitance device. b $d\mu/dn$ in Sample C1 as a function of ν and displacement field D as extracted from top gate capacitance. c, $d\mu/dn$ around $\nu=-3$ in Sample C1. d, Linecuts of $d\mu/dn$ at various displacement fields from the data in panel b. e, Measured gaps $\Delta\mu$ at $\nu=-1$ from Samples C1 and C2, with twist angles $\theta=2.2^\circ$ and $\theta=3.1^\circ$, and gap at $\nu=-3$ in Sample C1. The gap size doesn't depend strongly on the twist angle. f, Continuum model predictions of dependence on moiré bands from the Γ-and K-valleys on layer potential difference Δ , defined such that the energy of the topmost layer is constant as Δ is increased. Widths of the shaded regions show the bandwidth of respective moiré bands. At low layer potential difference, the lowest energy states are from the Γ valley, whereas at large layer potential difference, the lowest energy hole state is localized on the topmost layer at the K valley.

OUTLOOK

347

In conclusion, we show prominent effects of electronic $_{349}$ interactions in Γ -valley moiré bands over a wide range of parameter space in tdWSe₂. The correlated insulators we observe exhibit distinct magnetic and displacement field dependence compared to reports of other TMD moiré superlattices. Measurements of the chemical potential upon doping these charge-ordered states reveal evidence for spin polaron quasiparticle excitations. Together, this demonstrates that AB-stacked Γ-valley moiré TMDs realize novel parameter regimes of a mesoscopic triangular lattice model. Our work also suggests a clear path to- 374 ward engineering low-disorder Γ -valley bands to look at 375 new lattice geometries including honeycomb, Kagome, and anisotropic multi-orbital generalizations of the Hubbard model [4], as well as spontaneous ferroelectric states [48]. A relatively modest displacement field is suffi-364 cient to tune the valley character of the lowest TMD 381

moiré band from Γ to K, providing a way to electrically switch between dramatically different moiré band structures within a single device. Further, our d.c. thermodynamic sensing modality enables quantitative measurement of TMD moiré systems down to low densities. This paves the way to study TMD moiré superlattices with longer wavelengths where the relevant experimental carrier density ranges become smaller, but electronic correlations are even stronger relative to bandwidth.

These authors contributed equally bef@stanford.edu

^[1] G. Aivazian, Z. Gong, A. M. Jones, R.-L. Chu, J. Yan, D. G. Mandrus, C. Zhang, D. Cobden, W. Yao, and X. Xu. Magnetic control of valley pseudospin in monolayer WSe2. *Nature Physics*, 11(2):148–152, Feb. 2015. ISSN 1745-2481. doi:10.1038/nphys3201. URL https://www.nature.com/articles/nphys3201. Number: 2

Publisher: Nature Publishing Group.

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

L. An, X. Cai, D. Pei, M. Huang, Z. Wu, Z. Zhou, 447 J. Lin, Z. Ying, Z. Ye, X. Feng, R. Gao, C. Ca- 448 cho, M. Watson, Y. Chen, and N. Wang. teraction effects and superconductivity signatures in $_{450}$ twisted double-bilayer WSe2. Nanoscale Horizons, 5 451 (9):1309-1316, Aug. 2020. ISSN 2055-6764. doi: 10.1039/D0NH00248H URL http://pubs.rsc.org/en/ content/articlelanding/2020/nh/d0nh00248h. Publisher: The Royal Society of Chemistry.

453

456

462

482

483

504

505

506

507

508

509

- E. Y. Andrei and A. H. MacDonald. Graphene bilayers with a twist. Nature Materials, 19(12):1265-1275, Dec. 2020. ISSN 1476-4660. doi:10.1038/s41563-020-00840-0 URL https://www.nature.com/articles/ 459 [13] s41563-020-00840-0. Number: 12 Publisher: Nature 460 Publishing Group.
- M. Angeli and A. H. MacDonald. valley transition metal dichalcogenide moiré bands. Proceedings of the National Academy of Sciences, 118(10): 464 e2021826118, Mar. 2021. ISSN 0027-8424, 1091-6490. 465 doi:10.1073/pnas.2021826118. URL http://www.pnas. 466 org/lookup/doi/10.1073/pnas.2021826118.
- L. Balents, C. R. Dean, D. K. Efetov, and A. F. Young. Superconductivity and strong correlations in 469 moiré flat bands. Nature Physics, 16(7):725–733, 470 July 2020. ISSN 1745-2481. doi:10.1038/s41567- 471 URL https://www.nature.com/articles/ 020-0906-9 s41567-020-0906-9. Number: 7 Publisher: Nature Pub- 473 lishing Group.
- Z. Chu, E. C. Regan, X. Ma, D. Wang, Z. Xu, 475 [6] M. I. B. Utama, K. Yumigeta, M. Blei, K. Watanabe, 476 T. Taniguchi, S. Tongay, F. Wang, and K. Lai. Nanoscale 477 Conductivity Imaging of Correlated Electronic States in 478 ${\rm WSe}_{2}/{\rm WS}}_{2}$ Moir\'e 479 Superlattices. Physical Review Letters, 125(18):186803, 480 [16] Oct. 2020. doi:10.1103/PhysRevLett.125.186803. URL 481 https://link.aps.org/doi/10.1103/PhysRevLett. 125.186803. Publisher: American Physical Society.
- M. Davydova, Y. Zhang, and L. Fu. Itinerant spin 484 polaron and metallic ferromagnetism in semiconductor 485 moir\'e superlattices. arXiv:2206.01221, June 2022. URL 486 http://arxiv.org/abs/2206.01221.
- J. P. Eisenstein, L. N. Pfeiffer, and K. W. West. Nega- 488 tive compressibility of interacting two-dimensional elec- 489 tron and quasiparticle gases. Physical Review Let- 490 [17] ters, 68(5):674–677, Feb. 1992. ISSN 0031-9007. doi: 491 10.1103/PhysRevLett.68.674. URL https://link.aps. org/doi/10.1103/PhysRevLett.68.674.
- J. P. Eisenstein, L. N. Pfeiffer, and K. W. West. Com- 494 pressibility of the two-dimensional electron gas: Mea- 495 surements of the zero-field exchange energy and frac- 496 tional quantum Hall gap. Physical Review B, 50(3): 497 [18] 1760-1778, July 1994. ISSN 0163-1829, 1095-3795. doi: 498 10.1103/PhysRevB.50.1760 URL https://link.aps. org/doi/10.1103/PhysRevB.50.1760.
- A. Ghiotto, E.-M. Shih, G. S. S. G. Pereira, D. A. 501 Rhodes, B. Kim, J. Zang, A. J. Millis, K. Watanabe, 502 T. Taniguchi, J. C. Hone, L. Wang, C. R. Dean, and 503 A. N. Pasupathy. Quantum criticality in twisted transition metal dichalcogenides. Nature, 597(7876):345-349, Sept. 2021. ISSN 1476-4687. doi:10.1038/s41586-021-03815-6 URL https://www.nature.com/articles/ s41586-021-03815-6. Number: 7876 Publisher: Nature Publishing Group.

- 446 [11] M. V. Gustafsson, M. Yankowitz, C. Forsythe, D. Rhodes, K. Watanabe, T. Taniguchi, J. Hone, X. Zhu, and C. R. Dean. Ambipolar Landau levels and strong band-selective carrier interactions in monolayer WSe2. Nature Materials, 17(5):411-415, May 2018. ISSN 1476-4660. doi:10.1038/s41563-018-0036-2. URL https:// www.nature.com/articles/s41563-018-0036-2.
 - [12] N. C. Hu and A. H. MacDonald. Competing magnetic states in transition metal dichalcogenide moir\'e materials. Physical Review B, 104(21):214403, ISSN 2469-9950, 2469-9969. Dec. 2021. 10.1103/PhysRevB.104.214403. URL http://arxiv. org/abs/2108.02159.
 - X. Huang, T. Wang, S. Miao, C. Wang, Z. Li, Z. Lian, Taniguchi, K. Watanabe, S. Okamoto, D. Xiao, S.-F. Shi, and Y.-T. Cui. Correlated insulating states at fractional fillings of the WS2/WSe2 moiré lattice. Nature Physics, 17(6):715-719, June 2021. ISSN 1745-2481. doi:10.1038/s41567-021-01171-w. URL https: //www.nature.com/articles/s41567-021-01171-w.
 - D. M. Kennes, M. Claassen, L. Xian, A. Georges, A. J. Millis, J. Hone, C. R. Dean, D. N. Basov, A. N. Pasupathy, and A. Rubio. Moiré heterostructures as a condensed-matter quantum simulator. Nature Physics, 17(2):155-163, Feb. 2021. ISSN 1745-2481. doi:10.1038/s41567-020-01154-3. URL https://www. nature.com/articles/s41567-020-01154-3. Number: 2 Publisher: Nature Publishing Group.
- G. Kresse and J. Furthmüller. Efficient iterative schemes 474 [15] for ab initio total-energy calculations using a plane-wave basis set. Physical Review B, 54(16):11169-11186, Oct. 1996. doi:10.1103/PhysRevB.54.11169. URL https:// link.aps.org/doi/10.1103/PhysRevB.54.11169. Publisher: American Physical Society.
 - H. Li, S. Li, M. H. Naik, J. Xie, X. Li, J. Wang, E. Regan, D. Wang, W. Zhao, S. Zhao, S. Kahn, K. Yumigeta, M. Blei, T. Taniguchi, K. Watanabe, S. Tongay, A. Zettl, S. G. Louie, F. Wang, and M. F. Crommie. Imaging moiré flat bands in three-dimensional reconstructed WSe2/WS2 superlattices. Nature Materials, 20(7):945-950, July 2021. ISSN 1476-4660. doi:10.1038/s41563-021-00923-6. URL https://www.nature.com/articles/ **s**41563-021-00923-6. Number: 7 Publisher: Nature Publishing Group.
 - T. Li, S. Jiang, L. Li, Y. Zhang, K. Kang, J. Zhu, K. Watanabe, T. Taniguchi, D. Chowdhury, L. Fu, J. Shan, and K. F. Mak. Continuous Mott transition in semiconductor moiré superlattices. ture, 597(7876):350-354, Sept. 2021. ISSN 1476-4687. doi:10.1038/s41586-021-03853-0. URL https://www. nature.com/articles/s41586-021-03853-0.
 - T. Li, S. Jiang, B. Shen, Y. Zhang, L. Li, Z. Tao, T. Devakul, K. Watanabe, T. Taniguchi, L. Fu, J. Shan, and K. F. Mak. Quantum anomalous Hall effect from intertwined moiré bands. Nature, 600(7890):641-646, Dec. 2021. ISSN 0028-0836, 1476-4687. doi:10.1038/s41586-021-04171-1 URL https://www.nature.com/articles/ s41586-021-04171-1.
 - T. Li, J. Zhu, Y. Tang, K. Watanabe, T. Taniguchi, V. Elser, J. Shan, and K. F. Mak. Charge-order-enhanced capacitance in semiconductor moiré superlattices. Nature Nanotechnology, 16(10):1068–1072, Oct. 2021. ISSN 1748-3395. doi:10.1038/s41565-021-00955-8. URL http: //www.nature.com/articles/s41565-021-00955-8.

[20] G.-B. Liu, D. Xiao, Y. Yao, X. Xu, and W. Yao. 574 Electronic structures and theoretical modelling of two- 575 dimensional group-VIB transition metal dichalcogenides. 576 512 Chemical Society Reviews, 44(9):2643-2663, 2015. ISSN 577 513 0306-0012, 1460-4744. doi:10.1039/C4CS00301B. URL 578 514 http://xlink.rsc.org/?DOI=C4CS00301B. 515

579

584

586

587

606

630

511

517

532

533

536

537

549

550

551

552

553

554

555

556

558

559

560

561

562

563

564

565

566

567

568

- S. J. Magorrian, V. V. Enaldiev, V. Zólyomi, 580 [21] 516 F. Ferreira, V. I. Fal'ko, and D. A. Ruiz-Tijerina. 581 Multifaceted moir∖'e superlattice physics in 582 518 twisted ${\mathrm{WSe}}_{-2}$ bilayers. Phys-583 519 ical Review B, 104(12):125440, Sept. 2021. doi: 520 URL 10.1103/PhysRevB.104.125440 https: 585 521 //link.aps.org/doi/10.1103/PhysRevB.104.125440. 522 Publisher: American Physical Society. 523
- [22] K. F. Mak, C. Lee, J. Hone, J. Shan, and T. F. 524 Heinz. Atomically Thin ${\mathbf{MoS}}_{2}$: 525 A New Direct-Gap Semiconductor. Physical Re- 590 526 view Letters, 105(13):136805, Sept. 2010. doi: 527 10.1103/PhysRevLett.105.136805. URL https://link. 528 aps.org/doi/10.1103/PhysRevLett.105.136805. Pub-529 530 lisher: American Physical Society.
- 531 entis, T. Taniguchi, K. Watanabe, S. K. Banerjee, and E. Tutuc. Density-Dependent Quan- 597 tum Hall States and Zeeman Splitting in Mono- 598 layer and Bilayer ${\mathrm{WSe}}_{-2}$. Physi- 599 cal Review Letters, 118(24):247701, June 2017. doi: 10.1103/PhysRevLett.118.247701 URL https://link. aps.org/doi/10.1103/PhysRevLett.118.247701.
- [24] H. C. Movva, T. Lovorn, B. Fallahazad, S. Lar- 603 539 entis, K. Kim, T. Taniguchi, K. Watanabe, S. K. 604 [32] 540 Banerjee, A. H. MacDonald, and E. Tutuc. Tunable 605 541 $\mathrm{\Gamma}}\$ 542 }K\$ Vallev Populations in Hole-Doped Tri- 607 543 laver ${\rm WSe}_{-2}$ PhusicalRe- 608 544 view Letters, 120(10):107703, Mar. 2018. doi: 609 545 10.1103/PhysRevLett.120.107703. URL https:// 610 546 link.aps.org/doi/10.1103/PhysRevLett.120.107703. 611 547 Publisher: American Physical Society. 548
 - Nicholas, Kl- 613 [25]R. R. J. K. v. and G. Weimann. Exchange itzing, en- 614 of the spin splitting in a GaAs- 615 $}\mathrm{x}}$ \$As heterojunction. Physical 617 Review B, 37(3):1294-1302, Jan. 1988. doi: 618 https://link. 10.1103/PhysRevB.37.1294. URL aps.org/doi/10.1103/PhysRevB.37.1294.
- H. Pan, F. Wu, and S. Das Sarma. Quantum phase 621 557 diagram of a Moiré-Hubbard model. Physical Review 622 B, 102(20):201104, Nov. 2020. ISSN 2469-9950, 2469-623 9969. doi:10.1103/PhysRevB.102.201104. URL https: 624 //link.aps.org/doi/10.1103/PhysRevB.102.201104.
 - H. Peng, Z.-H. Yang, J. P. Perdew, and J. Sun. 626 Versatile van der Waals Density Functional Based on 627 a Meta-Generalized Gradient Approximation. Phys- 628 [35] ical Review X, 6(4):041005, Oct. 2016. doi: 629 10.1103/PhysRevX.6.041005. URL https://link. aps.org/doi/10.1103/PhysRevX.6.041005. Publisher: American Physical Society.
- E. C. Regan, D. Wang, C. Jin, M. I. Bakti Utama, 633 569 B. Gao, X. Wei, S. Zhao, W. Zhao, Z. Zhang, K. Yu- 634 570 migeta, M. Blei, J. D. Carlström, K. Watanabe, 635 571 T. Taniguchi, S. Tongay, M. Crommie, A. Zettl, and 636 572 F. Wang. Mott and generalized Wigner crystal states in 637 573

- WSe2/WS2 moiré superlattices. Nature, 579(7799):359-363, Mar. 2020. ISSN 1476-4687. doi:10.1038/s41586-020-2092-4 URL https://www.nature.com/articles/ s41586-020-2092-4. Number: 7799 Publisher: Nature Publishing Group.
- S. Shabani, D. Halbertal, W. Wu, M. Chen, S. Liu, J. Hone, W. Yao, D. N. Basov, X. Zhu, and A. N. Pasupathy. Deep moiré potentials in twisted transition metal dichalcogenide bilayers. Nature Physics, 17(6):720-725, June 2021. ISSN 1745-2481. doi:10.1038/s41567-021-01174-7. URL https://www.nature.com/articles/ **s41567-021-01174-7**. Number: 6 Publisher: Nature Publishing Group.
- Q. Shi, E.-M. Shih, M. V. Gustafsson, D. A. Rhodes, B. Kim, K. Watanabe, T. Taniguchi, Z. Papić, J. Hone, and C. R. Dean. Odd- and even-denominator fractional quantum Hall states in monolayer WSe2. Nature Nanotechnology, 15(7):569-573, July 2020. ISSN 1748-3395. doi:10.1038/s41565-020-0685-6 URL https:// www.nature.com/articles/s41565-020-0685-6. Number: 7 Publisher: Nature Publishing Group.
- H. C. Movva, B. Fallahazad, K. Kim, S. Lar- 595 [31] Q. Shi, E.-M. Shih, D. Rhodes, B. Kim, K. Barmak, K. Watanabe, T. Taniguchi, Z. Papić, D. A. Abanin, J. Hone, and C. R. Dean. Bilayer WSe2 as a natural platform for interlayer exciton condensates in the strong coupling limit. Nature Nanotechnology, pages 1-6, Apr. 2022. ISSN 1748-3395. doi:10.1038/s41565-022-01104-5. URL https://www.nature.com/articles/ s41565-022-01104-5. Publisher: Nature Publishing Group.
 - Y. Tang, L. Li, T. Li, Y. Xu, S. Liu, K. Barmak, K. Watanabe, T. Taniguchi, A. H. MacDonald, J. Shan, and K. F. Mak. Simulation of Hubbard model physics in WSe2/WS2 moiré superlattices. Nature, 579(7799):353-358, Mar. 2020. ISSN 1476-4687. doi:10.1038/s41586-020-2085-3 URL https://www.nature.com/articles/ s41586-020-2085-3. Number: 7799 Publisher: Nature Publishing Group.
- 612 [33] Y. Tang, J. Gu, S. Liu, K. Watanabe, T. Taniguchi, J. C. Hone, K. F. Mak, and J. Shan. tric catastrophe at the Wigner-Mott transition in a moiré superlattice. Nature Communications, 13(1): ${\rm Ga}_{\rm Ga}\$ -{\mathrm{x}}\$\${\mathrm{Al}}_{1\mathrm{42\lambda}ur\lambda}. ISSN 2041-1723. doi:10.1038/s41467-022-32037-1 URL https://www.nature.com/articles/ **s41467-022-32037-1**. Number: 1 Publisher: Nature Publishing Group.
 - Y. Tang, K. Su, L. Li, Y. Xu, S. Liu, K. Watanabe, T. Taniguchi, J. Hone, C.-M. Jian, C. Xu, K. F. Mak, and J. Shan. Evidence of frustrated magnetic interactions in a Wigner-Mott insulator. Nature Nanotechnology, pages 1-5, Jan. 2023. ISSN 1748-3395. doi:10.1038/s41565-022-01309-8. URL https://www.nature.com/articles/ s41565-022-01309-8. Publisher: Nature Publishing Group.
 - E. Tutuc, S. Melinte, and M. Shayegan. Spin Polarization and g Factor of a Dilute GaAs Two-Dimensional Electron System. Physical Review Letters, 88(3): 036805, Jan. 2002. ISSN 0031-9007, 1079-7114. doi: 10.1103/PhysRevLett.88.036805 URL https://link. aps.org/doi/10.1103/PhysRevLett.88.036805.
 - V. Vitale, K. Atalar, A. A. Mostofi, and J. Lischner. Flat band properties of twisted transition metal dichalcogenide homo- and heterobilavers of $MoS \leq sss sub \greater 2 \less \sub \greater \,$

 $MoSe \leq sss \leq sub \leq sss / sub < sss / sub < sss / sub < sub / sub / sub < sub / sub < sub / sub < sub / sub / sub / sub / sub$ $WS\$\less\$sub\$\greater\$2\$\less\$/sub\$\greater\$$ and $WSe \leq sss \leq sub \leq srs \leq sss \leq sss$ 8(4):045010, Aug. 2021. ISSN 2D Materials. 2053-1583. doi:10.1088/2053-1583/ac15d9 URL 706 https://doi.org/10.1088/2053-1583/ac15d9. Publisher: IOP Publishing.

702

703

704

705

707

708

716

638

639

640

641

642

643

644

656

657

658

659

661

674

676

678

680

681 682

694

697

- L. Wang, E.-M. Shih, A. Ghiotto, L. Xian, D. A. 645 Rhodes, C. Tan, M. Claassen, D. M. Kennes, Y. Bai, 710 646 B. Kim, K. Watanabe, T. Taniguchi, X. Zhu, J. Hone, 711 647 A. Rubio, A. N. Pasupathy, and C. R. Dean. Cor-648 related electronic phases in twisted bilayer transition 713 649 metal dichalcogenides. Nature Materials, 19(8):861-650 866, Aug. 2020. ISSN 1476-4660. doi:10.1038/s41563-651 020-0708-6 URL https://www.nature.com/articles/ 652 **s41563-020-0708-6**. Number: 8 Publisher: Nature Pub-653 654
- X. Wang, C. Xiao, H. Park, J. Zhu, C. Wang, 719 655 T. Taniguchi, K. Watanabe, J. Yan, D. Xiao, D. R. 720 [47] Gamelin, W. Yao, and X. Xu. Light-induced ferromagnetism in moiré superlattices. Nature, 604(7906): 722 468-473, Apr. 2022. ISSN 0028-0836, 1476-4687. 723 doi:10.1038/s41586-022-04472-z. URL https://www. 660 nature.com/articles/s41586-022-04472-z.
- [39] N. R. Wilson, P. V. Nguyen, K. Seyler, P. Rivera, 726 662 A. J. Marsden, Z. P. L. Laker, G. C. Constantinescu, 727 663 V. Kandyba, A. Barinov, N. D. M. Hine, X. Xu, and 728 D. H. Cobden. Determination of band offsets, hybridization, and exciton binding in 2D semiconductor het- 730 erostructures. Science Advances, 3(2):e1601832. 667 10.1126/sciadv.1601832 URL https://www.science. 668 org/doi/10.1126/sciadv.1601832. Publisher: Ameri- 733 669 can Association for the Advancement of Science. 670
- F. Wu, T. Lovorn, E. Tutuc, and A. MacDonald. 735 [40] 671 Hubbard Model Physics in Transition Metal Dichalco- 736 672 genide Moiré Bands. Physical Review Letters, 121(2): 737 673 026402, July 2018. ISSN 0031-9007, 1079-7114. doi: 738 10.1103/PhysRevLett.121.026402 URL https://link. 675 aps.org/doi/10.1103/PhysRevLett.121.026402.
- F. Wu, T. Lovorn, E. Tutuc, I. Martin, and A. Mac-677 Topological Insulators in Twisted Transition Metal Dichalcogenide Homobilayers. Physi-679 cal Review Letters, 122(8):086402, Feb. 2019. doi: 10.1103/PhysRevLett.122.086402 URL https://link. aps.org/doi/10.1103/PhysRevLett.122.086402. Publisher: American Physical Society. 683
- L. Xian, M. Claassen, D. Kiese, M. M. Scherer, 684 S. Trebst, D. M. Kennes, and A. Rubio. Realiza-685 tion of nearly dispersionless bands with strong or-686 bital anisotropy from destructive interference in twisted 687 bilayer MoS2. Nature Communications, 12(1):5644, 688 ISSN 2041-1723. doi:10.1038/s41467-Sept. 2021. 689 021-25922-8. URL http://www.nature.com/articles/ 690 **s41467-021-25922-8**. Number: 1 Publisher: Nature 691 Publishing Group. 692
- 693 pseudospins in layered transition metal dichalcogenides. Nature Physics, 10(5):343-350, May 2014. ISSN 1745-695 2473, 1745-2481. doi:10.1038/nphys2942. URL http: 696 //www.nature.com/articles/nphys2942.
- Y. Xu, S. Liu, D. A. Rhodes, K. Watanabe, T. Taniguchi, 698 J. Hone, V. Elser, K. F. Mak, and J. Shan. Corre-690 700 perlattices. Nature, 587(7833):214–218, Nov. 2020. ISSN 757 ples S1 and S2, local Cr/Au "contact" gates (3nm/50 701

- 1476-4687. doi:10.1038/s41586-020-2868-6. URL https: //www.nature.com/articles/s41586-020-2868-6.
- Y. Xu, K. Kang, K. Watanabe, T. Taniguchi, K. F. Mak, and J. Shan. A tunable bilayer Hubbard model in twisted WSe2. Nature Nanotechnology, 17(9):934-939, Sept. 2022. ISSN 1748-3395. doi:10.1038/s41565-022-01180-7. URL https://www.nature.com/articles/ s41565-022-01180-7. Number: 9 Publisher: Nature Publishing Group.
- J. Yu, B. A. Foutty, Z. Han, M. E. Barber, Y. Schattner, K. Watanabe, T. Taniguchi, P. Phillips, Z.-X. Shen, S. A. Kivelson, and B. E. Feldman. Correlated Hofstadter spectrum and flavour phase diagram in magic-angle twisted bilayer graphene. Nature Physics, 18(7):825–831, July 2022. ISSN 1745-2481. doi:10.1038/s41567-022-01589-wURL https://www.nature.com/articles/ s41567-022-01589-w. Number: 7 Publisher: Nature Publishing Group.
- Y. Zhang, N. F. Q. Yuan, and L. Fu. Moir\'e quantum chemistry: Charge transfer in transition metal dichalcogenide superlattices. Physical Review B, 102(20): 201115, Nov. 2020. doi:10.1103/PhysRevB.102.201115. URL https://link.aps.org/doi/10.1103/PhysRevB. 102.201115. Publisher: American Physical Society.
- Y. Zhang, T. Liu, and L. Fu. Electronic structures, charge transfer, and charge order in twisted transition metal dichalcogenide bilayers. Physical Review B, 103 (15):155142, Apr. 2021. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.103.155142 URL https://link. aps.org/doi/10.1103/PhysRevB.103.155142.
- 732 [49] Z. Y. Zhu, Y. C. Cheng, and U. Schwingenschlögl. Giant spin-orbit-induced spin splitting in two-dimensional transition-metal dichalcogenide semiconductors. Physical Review B, 84(15):153402, Oct. 2011. ISSN 1098-0121. 1550-235X. doi:10.1103/PhysRevB.84.153402 URL https://link.aps.org/doi/10.1103/PhysRevB. 84.153402.

METHODS

Sample fabrication

The tdWSe₂ devices were fabricated using standard 742 dry transfer techniques. An exfoliated Bernal bilayer 743 WSe₂ flake (sources: 2DSemiconductors, HQGraphene) vas pre-cut by a conductive AFM probe in contact mode, vith an a.c. excitation of 10 V at 50 kHz in order to fa-746 cilitate the stacking process and alleviate strain. Using a 747 poly(bisphenol A carbonate) (PC)/polydimethylsiloxane 748 (PDMS) stamp, we pick up a thin (15-30 nm) hexagonal 749 boron nitride (hBN) flake, followed by the first half of the X. Xu, W. Yao, D. Xiao, and T. F. Heinz. Spin and 750 bilayer WSe₂ flake, and then the second half rotated to $_{751}$ a controlled angle of $2^{\circ} - 3.5^{\circ}$. Separately, we prepare a 752 stack with a bottom hBN (25-40 nm thick) and a graphite 753 (5-10 nm) back gate, on which we deposit pre-patterned ₇₅₄ Cr/Pt contacts (6-12 nm). This is annealed at ≈ 300 °C 755 overnight to clean polymer and resist residues before delated insulating states at fractional fillings of moiré su- 756 positing the tdWSe2 stack on the Pt contacts. For samSET measurement 10

₇₅₈ nm) were patterned above the Cr/Pt contacts after as-₈₀₇ applied to both sample and back gate at distinct frequen-759 sembly in order to locally dope the contact regions so 808 cies between 200 and 900 Hz. We then measure inverse ₇₆₀ that they achieve Ohmic contact. For Samples C1 and ₈₀₉ compressibility $d\mu/dn \propto I_{\rm BG}/I_{\rm 2D}$ where $I_{\rm BG}$ and $I_{\rm 2D}$ ₇₆₃ secondary hBN is set down on top of the device, followed $_{812}$ citations [46]. A d.c. offset voltage $V_{\rm 2D}$ is applied to 765 e-beam lithography techniques to fabricate contacts and 814 its maximum sensitivity point within a Coulomb block-766 top gates.

Density and twist angle determination

767

799

For Samples S1 and S2, both the band edge and hBN dielectric capacitance are determined by fitting Landau level oscillations (Supplementary Sec. 11). The band edge is taken to be the point to which these oscillations extrapolate at B = 0, and the fitted slopes give the sample-gate capacitance, which specifies the conversion between gate voltage and hole density. In both samples, the location in gate voltage where we see effective charging of the TMD sample are a few 10^{11} cm⁻² more doped than the band edge taken from the Landau fan. We attribute this difference to a mobility edge at 779 low TMD densities [37]. Based on the density of the $_{780}$ $\nu = -1$ gapped state, we convert from the density of one 781 hole per moiré unit cell n_s to an angle θ by the relation $_{782}$ $\frac{1}{n_s} = \frac{\sqrt{3}a}{4 - 4\cos(\theta)}$ where a = .328 nm is the lattice constant 783 of WSe₂. Twist angles measured using this method agree ₇₈₄ to within $(\pm 0.3^{\circ})$ of the targeted rotation during the fabrication process, and are also confirmed to within $(\pm 0.5^{\circ})$ based on edge orientation as measured from optical images and using atomic force microscopy. Further details and support for the twist angle assignment for each sample are provided in Supplementary Sec. 11.

For Samples C1-C2, we do not have magnetic field dependence to fix the band edge. The capacitances are estimated from hBN thickness and average measured dielectric constants for our batch of hBN samples, $\epsilon \approx 3.1$. The location of the band edge in gate voltage is determined by the spacing of the $\nu = -1$ and $\nu = -3$ features (in Sample C1) and lower frequency/higher contact gate capacitance linetraces where the sample charges better closer to the band edge (Supplementary Sec. 11).

SET measurement

804 ning SET measurements were performed in a Unisoku 856 substrate to avoid parasitic capacitances, and the devices 806 microscope head. a.c. excitations of order 5-10 mV were 858 gate to minimize singly gated regions of sample.

C2, an additional Cr/Au top gate is first patterned over 810 are measurements of the SET current demodulated at the device region (not the Pt contacts), after which a 811 respective frequencies of the back gate and sample exby Cr/Au gates over the contacts. We used standard 813 the sample to maintain the working point of the SET at 815 ade oscillation fringe chosen to be near the "flat-band" 816 condition where the tip does not gate the sample. This 817 minimizes tip-induced doping and provides a direct measurement of $\mu(n)$. The contact gates are held at a large, 819 negative voltage throughout the measurement. All SET measurements are taken at T = 330 mK.

Capacitance measurement in dual-gated devices

We use standard high electron mobility transistor 823 (HEMT) techniques to measure the device capacitance 824 [19, 30]. An FHX35X transistor is glued adjacent to the 825 sample with the HEMT gate connected to the sample 826 contacts. The sample voltage V_{2D} is held fixed, setting 827 the HEMT gain, while a small current is sourced between 828 the source and drain leads of the transistor. Throughout 829 the measurement, the contact gates are held fixed at a 830 large, negative voltage $V_{\rm CG}$ to maintain ohmic contact 1831 resistance independent of the applied gate voltages. The $_{832}$ d.c. voltages $V_{\rm BG}$ and $V_{\rm TG}$ applied to top and back gate, $_{833}$ respectively, independently tune the density n and dis- 834 placement field D. We define the displacement field as 835 $D=rac{1}{2\epsilon_0}(c_t(V_{\mathrm{TG}}-\phi_0)-c_b(V_{\mathrm{BG}}-\phi_1))$ where $c_{t(b)}$ is 836 the capacitance of the top (bottom) hBN dielectric and 837 $\phi_{0(1)}$ is the estimated work function difference between 838 our TMD sample and gates. We take $\phi_1 = -0.8 \text{ V}$ and 839 $\phi_0 = -0.1$ V, as checked by SET measurements in sep-840 arate samples, which are sensitive to the work function. 841 a.c. excitation voltages are applied to the top and back 842 gates with amplitude 5-10 mV and frequency ranging 843 from 300 Hz - 1 kHz, and the amplified capacitance (and 844 out-of-phase dissipation) signal is measured by the a.c. 845 voltage drop across the HEMT leads. All capacitance 846 measurements presented in the main text were taken at ₈₄₇ T = 4.2 K (see Supplementary Sec. 12 for temperature dependence up to $T \approx 15$ K). To convert from measured 849 signals to quantitative units, we normalize the signal us-850 ing the known device capacitance and the difference in 851 the measured signal between the fully gapped and highly The SET sensor was fabricated by evaporating alu- 852 doped regions of the sample, before applying a lumped minum onto a pulled quartz rod, with an estimated di- 853 circuit model to convert from the capacitive and dissiameter at the apex of 50 - 100 nm. The SET "tip" is 854 pative components to $d\mu/dn$ (Supplementary Sec. 13). brought to about 50 nm above the sample surface. Scan- 855 Dual gated devices are fabricated on an undoped Si/SiO₂ USM 1300 scanning probe microscope with a customized 857 were etched (CHF₃/O₂ plasma etching) around the top 859

891

Density functional calculations are performed using 896 generalized gradient approximation with SCAN+rVV10 $\,$ Van der Waals density functional [27], as implemented in the Vienna Ab initio Simulation Package [15]. Pseudopotentials are used to describe the electron-ion interactions. We construct the moiré band structure of tdWSe₂ by calculating the band structures of untwisted double bilayers with MM', MX' and XX' stackings and "stitching" the results together with the continuum model approach. The vacuum spacing is larger than 20 Å to avoid artificial interaction between the periodic images along the z direction. The structure relaxation is performed with force on each atom less than 0.01 eV/A. We use $12 \times 12 \times 1$ for structure relaxation and self-consistent calculation. The more accurate SCAN+rVV10 van der Waals density functional gives the relaxed layer distances as 6.64 Å, 6.62 Å and 7.10 Å for MM', MX' and XX' stacking 877 structures, respectively. By calculating the work func-878 tion from electrostatic energy of converged charge den-879 sity, we plot in Fig. 1e the band structure of MM', MX' and XX'-stacked double bilayers, with reference energy E=0 chosen to be the absolute vacuum level.

Our DFT calculation finds that the valence band maximum of tdWSe₂ is at Γ and lies roughly 100 meV above the band edge at K, consistent with a recent angleresolved photoemission spectroscopy measurement [2]. Unlike monolayer and bilayer, the reversed ordering of T and K band edges in four-layer WSe₂ results from the strong interlayer tunneling around Γ , which leads to large energy splittings on the order of 0.1 eV between layerhybridized Γ -valley bands.

DATA AVAILABILITY

The data that supports the findings of this study are 925 available from the corresponding authors upon reasonable request. 926

 896 The codes that support the findings of this study are 897 available from the corresponding authors upon reason- 898 able request.

ACKNOWLEDGEMENTS

We acknowledge helpful conversations with Allan H. 901 MacDonald. We thank Tony Heinz, Aidan O'Beirne, 902 and Henrique Bucker Ribeiro for their assistance with SHG measurements. Experimental work was primarily supported by NSF-DMR-2103910. B.E.F. acknowledges 905 an Alfred P. Sloan Foundation Fellowship and a Cottrell Scholar Award. The work at Massachusetts Institute of Technology is supported by a Simons Investigator Award from the Simons Foundation. L.F. is partly supported by the David and Lucile Packard Foundation. K.W. and 910 T.T. acknowledge support from JSPS KAKENHI (Grant Numbers 19H05790, 20H00354 and 21H05233). B.A.F. 912 acknowledges a Stanford Graduate Fellowship. Part of 913 this work was performed at the Stanford Nano Shared Fa-914 cilities (SNSF), supported by the National Science Foun-915 dation under award ECCS-2026822.

AUTHOR CONTRIBUTION

B.A.F., J.Y., and B.E.F. designed and conducted the scanning SET experiments. B.A.F. and B.E.F. designed and conducted the dual gate capacitance experiments. T.D., Y.Z., and L.F. conducted theoretical calculations. B.A.F. fabricated the samples, with help from C.R.K. B.Z. K.W. and T.T. provided hBN crystals. All authors participated in analysis of the data and writing of the manuscript.

COMPETING INTERESTS

The authors declare no competing interest.

Supplementary Material for:

Tunable spin and valley excitations of correlated insulators in Γ -valley moiré bands

Benjamin A. Foutty, ^{1, 2, *} Jiachen Yu, ^{1, 3, *} Trithep Devakul, ^{4, *} Carlos R. Kometter, ^{1, 2} Yang Zhang, ⁴ Kenji Watanabe, ⁵ Takashi Taniguchi, ⁶ Liang Fu, ⁴ and Benjamin E. Feldman^{1, 2, 7, †}

¹ Geballe Laboratory for Advanced Materials, Stanford, CA 94305, USA

² Department of Physics, Stanford University, Stanford, CA 94305, USA

³ Department of Applied Physics, Stanford University, Stanford, CA 94305, USA

⁴ Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

⁵ Research Center for Functional Materials, National Institute for Material Science, 1-1 Namiki, Tsukuba 305-0044, Japan

⁶ International Center for Materials Nanoarchitectonics,

National Institute for Materials Science, 1-1 Namiki, Tsukuba 305-0044, Japan

⁷ Stanford Institute for Materials and Energy Sciences,

SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA

1. SPATIAL VARIATION MEASURED WITH SCANNING SET

In the main text, we focus on a single spatial location in each sample. In Fig. S1, we present spatial dependence of inverse electronic compressibility $d\mu/dn$ taken in each sample at magnetic field B=11 T, where the incompressible states are strongest. Here, μ is the chemical potential and n is the carrier density, with the convention that increasing hole density is negative. Both devices are highly homogeneous over micron length scales, with little twist angle disorder in the regions examined. We estimate the total twist angle disorder by fitting the positions of the measured incompressible states and comparing their spatial dependence. In Sample S1, we compare the difference in gate voltage between the peak at filling factor $\nu = -1$, which occurs around back gate voltage $V_{\rm BG} = -7$ V and the $\nu = -\frac{1}{3}$ peak, which occurs around $V_{\rm BG} = -3.2$ V. This yields a twist angle that varies between $\theta = 3.38^{\circ}$ and $\theta = 3.45^{\circ}$ in Area 1, where the measurements in the main text were conducted (Fig. S1a-b, Fig. S2a). In a second region (Area 2) in Sample S1 that is about 4 μ m away, the twist angle varies between $\theta = 3.42^{\circ}$ and $\theta = 3.61^{\circ}$ (Fig. S1c-d, Fig. S2a). In Sample S2, we fit the difference between the incompressible $\nu = -1$ gap at $V_{\rm BG} = -3.95$ V and the incompressible peak at $V_{\rm BG} = -3$ V, and estimate a twist angle that varies between $\theta = 2.55^{\circ}$ and $\theta = 2.65^{\circ}$ (Fig. S1e). We note that this peak may not be a charge-ordered state, but appears to occur at a roughly constant filling factor position and therefore can be used as a benchmark to estimate the twist angle disorder that behaves better than the band edge, which can vary due to underlying potential disorder in the sample. If we assume a constant band edge and just treat the position of the $\nu = -1$ gap in gate voltage as the only parameter, this leads to an even smaller estimate of twist angle disorder. Across both devices, we measure significantly less twist angle inhomogeneity than is typically observed across magic-angle twisted bilayer graphene devices [1-3]. This difference likely results from the larger overall twist angles in these devices which limit excessive relaxation/reconstruction and/or increased stiffness due to the presence of more layers.

In Sample S1, the gaps at $\nu = -1$ are prominent at a variety of locations both at B = 0 and B = 11 T. However, the $\nu = -\frac{1}{3}$ gap is much closer to the noise floor of the measurement at B = 0. In Fig. S3, we present spatial dependence of this gap at B = 0 across a 400 nm \times 500 nm region of the sample. We measure a small but nonzero gap at $\nu = -\frac{1}{3}$ between 0.3 meV and 1.1 meV over this area. The variability in gap size likely arises due to local variation in the underlying spatial disorder (from static charge disorder, small amounts of twist angle variability, and/or strain), which can appreciably affect the magnitude of the small gap at B = 0.

2. GAP EXTRACTION

The gaps Δ_{ν} presented in the main text correspond to the step in the chemical potential at filling factor ν , found by taking the difference between the local maximum and local minimum in μ on either side of the gap. In the scanning single-electron transistor (SET) measurements, we use the DC measurement of $\mu(n)$ to extract charge gap sizes, rather than integrating the AC measurement of $d\mu/dn$ because the high resistance of the contacts and/or sample artificially enhances the a.c. signal of the incompressible peaks at high fields.

In Sample S1, the $\nu = -1/3$ gap appears as $\mu(n)$ begins to turn up near the mobility edge even in the d.c. measurement scheme due to high sample resistance at low densities. To accurately extract the gap, we fit a quadratic polynomial curve to the surrounding data and subtract this background before evaluating the gap size, avoiding an overestimation due to the background shape and finite width of the incompressible gap. Variation of the bounds of this fitted background do not affect the measured gap size nor its quantitative trend with B (i.e., the effective g-factor). An example showing data at 5 T is plotted in Fig. S4. After subtracting the quadratic background, the size of the d.c. measured gap agrees within the noise with the simultaneously a.c. measured gap at low fields (Fig. S5), where a nearly density-independent background is straightforward to identify before integrating $d\mu/dn$. This is the only gap for which we perform any background subtraction; other gaps are well separated from the mobility edge.

In the global capacitance measurements, we only have direct access to $d\mu/dn$, rather than the chemical potential itself. To determine the charge gaps, we convert the capacitance signal to $d\mu/dn$ (see Supplementary Sec. 13 below) and then integrate over density to find $\mu(n)$. The charge gaps are then taken as the resulting step on this $\mu(n)$ curve.

3. CONTINUUM MODELING

We derive effective continuum models for twisted double bilayer WSe₂ (tdWSe₂) from first principles calculations at high symmetry stacking configurations.




FIG. S1. Spatial dependence at B=11 T. a, Measured locations in Area 1 of Sample S1. x and y label relative spatial coordinates. b, Inverse electronic compressibility $d\mu/dn$ as a function of back voltage $V_{\rm BG}$ at the positions shown in a, labelled by index. c, Locations of measurements in a far-separated region of Sample S1. X and Y label spatial coordinates within this second region. d, $d\mu/dn$ at positions shown in c. e, Spatial variation of $d\mu/dn$ along a continuous linecut in Sample S2.

The low energy physics of hole-doped monolayer WSe₂ consists of (local) quadratic band maxima at the Γ and $\pm K$ points of the Brillouin zone. The Γ valley band maximum is spin-degenerate, while the $\pm K$ band maxima are spin polarized with spin $s_z = \pm \frac{1}{2}$ due to spin-orbit coupling. The structure of four-layer WSe₂ is such that each layer is rotated 180° relative to its adjacent layers, so that interlayer tunneling is strongly suppressed in the $\pm K$ valley bands.

Three different high-symmetry stacking configurations, MM', MX', and XX', as illustrated in the main text, are considered. First principles density functional theory (DFT) calculations for four-layer WSe₂, shown in Fig. 1e of the main text, reveal that the valence band maximum is at Γ for all three stacking configurations. The energies of the top four valence bands at Γ , and the top two at K, are listed in Table S1.

We first consider an effective single-particle continuum model for the twisted system of the form

$$H_{\Gamma} = -\frac{|\mathbf{k}|^2}{2m_{\Gamma}} + V_{\Gamma}(\mathbf{r}) \tag{1}$$

where m_{Γ} is the Γ valley effective mass, and $V(\mathbf{r})$ describes a periodic potential with the moiré periodicity. The spin degeneracy is implicit. The potential $V(\mathbf{r})$ can be obtained by taking the first harmonic approximation, $V_{\Gamma}(\mathbf{r}) = 2v \sum_{i=1,3,5} \cos(\mathbf{g}_i \cdot \mathbf{r} + \phi) + c$, from which the three parameters, v, c, ϕ , are chosen to match the energy of the first Γ band in Table S1 at the three positions $\mathbf{R}_{\mathrm{MM'}}, \mathbf{R}_{\mathrm{MX'}}$, and $\mathbf{R}_{\mathrm{XX'}}$. Here, \mathbf{g}_i are moiré reciprocal lattice vectors defined $\mathbf{g}_i = \frac{4\pi}{\sqrt{3}a_M}(-\sin\frac{2\pi(i-1)}{6},\cos\frac{2\pi(i-1)}{6})$, and $\mathbf{R}_{\mathrm{MM'}} = 0$, $\mathbf{R}_{\mathrm{MX'}} = \frac{a_M}{\sqrt{3}}(0,-1)$, and $\mathbf{R}_{\mathrm{XX'}} = \frac{a_M}{\sqrt{3}}(0,1)$. With




FIG. S2. Optical images of the samples. a, Optical image of Sample S1. The measurements in the main text are all taken in the vicinity of the purple dot, while similar behavior was observed in the vicinity of the red dot (Fig. S1c-d) and measurements of the adjacent Bernal bilayer region were taken at the orange dot (Fig. S16). b, Optical image of Sample C2. Scale bars in \bf{a} and \bf{b} are both 5 μ m.




FIG. S3. Spatial dependence of the $\Delta_{-1/3}$ at B=0. a, Spatial location of points plotted in panel c, relative to (0,0) as set in Fig. S1. b, Left, $d\mu/dn$ at B=0 around $\nu=-1/3$ and right, integrated chemical potential $\mu(n)$ subtracting out a flat background at 7×10^{-11} meV cm². The resulting gap $\Delta_{-1/3}$ is taken to be the difference in the local maximum and minimum in $\mu(n)$. c, Gap size $\Delta_{-1/3}$ at B=0 as a function of spatial position as enumerated in a.

this definition, $V(\mathbf{R}_{\mathrm{MM'}}) = 6v\cos(\phi) + c$, $V(\mathbf{R}_{\mathrm{MX'}}) = 6v\cos(\phi + 2\pi/3) + c$, and $V(\mathbf{R}_{\mathrm{XX'}}) = 6v\cos(\phi - 2\pi/3) + c$. We find the parameters v = 17.9 meV, $\phi = 0.44$ (radians), and c = -4613.6 meV. We use the moiré period $a_M = 3.32 \text{ Å}/(2\sin(3.4^{\circ}/2)) \approx 56 \text{ Å}$, and effective mass $m_{\Gamma} = 1.2m_e$, where m_e is the electron mass [4]. The hole density obtained from filling the first moiré band of this continuum model is shown in Fig. 1f of the main text.

To incorporate the effect of a vertical displacement field, we now construct a model which includes all four layer degrees of freedom. We assume the form for the model Hamiltonian at zero displacement field $H_{\Gamma} = -\frac{|\mathbf{k}|^2}{2m_{\Gamma}}\mathbb{I} + W_{\Gamma}(\mathbf{r})$




FIG. S4. Background fitting near $\nu = -1/3$. Background subtraction around the $\nu = -1/3$ gap in Sample S1, measured at B = 5 T. Black dashed lines show local maximum and minimum values of μ_+ and μ_- around the gap, the difference of which is taken to be $\Delta_{-1/3}$.




FIG. S5. Comparison of a.c. and d.c. measured gaps at $\nu = -1/3$ at low magnetic fields. The a.c. measurement, which has overall lower noise, falls well within the scatter of the d.c. measured data. The two measurements are performed in the same location simultaneously, and the d.c. gaps are identical to those presented in Fig. 2c. Above B = 1.5 T, the a.c. measurement suffers from artificial enhancement, hence the focus on the d.c. measured gap in the main text.

where I is the identity matrix and

$$W_{\Gamma}(\mathbf{r}) = \begin{pmatrix} V_{1\Gamma}(\mathbf{r}) & T_{1\Gamma}(\mathbf{r}) & 0 & 0 \\ T_{1\Gamma}(\mathbf{r}) & V_{2\Gamma}(\mathbf{r}) & T_{2\Gamma}(\mathbf{r}) & 0 \\ 0 & T_{2\Gamma}(\mathbf{r}) & V_{2\Gamma}(\mathbf{r}) & T_{1\Gamma}(\mathbf{r}) \\ 0 & 0 & T_{1\Gamma}(\mathbf{r}) & V_{1\Gamma}(\mathbf{r}) \end{pmatrix}, \tag{2}$$

in which all T, V, are periodic functions with the moiré periodicity, and are chosen such that the eigenvalues of $W_{\Gamma}(\mathbf{R}_{\alpha})$ with $\alpha = MM', MX', XX'$, match with $E_{\Gamma}(\alpha)$ in Table S1. To uniquely specify T, we have taken it to

Band	E_{Γ} (MM')	E_{Γ} (MX')	E_{Γ} (XX')	$E_K \text{ (MM')}$	$E_K (MX')$	E_K (XX')
1	-4.5164	-4.7022	-4.6223	-4.6523	-4.8350	-4.7064
2	-4.6412	-4.8144	-4.7011	-4.6602	-4.8569	-4.7168
3	-4.9506	-5.1330	-5.0938	-	-	-
4	-5.3199	-5.5013	-5.3622	-	-	-

TABLE S1. Energy (in eV) obtained from DFT at the Γ and K points, for each of the three high symmetry stacking regions MM', MX', and XX'. Note that each band is doubly degenerate due to inversion and time reversal symmetry.

be real and positive. At each \mathbf{R}_{α} , there are four parameters $\{V_{1\Gamma}(\mathbf{R}_{\alpha}), V_{2\Gamma}(\mathbf{R}_{\alpha}), T_{1\Gamma}(\mathbf{R}_{\alpha}), T_{2\Gamma}(\mathbf{R}_{\alpha})\}$, which can be chosen to fit the four eigenvalues E_{Γ} . From these, we can use the first harmonic approximation to obtain

$$V_{1\Gamma}(\mathbf{r})/\text{meV} = -4799.5 + 16.4 \sum_{i=1,3,5} 2\cos(\mathbf{g}_{i} \cdot \mathbf{r} + 0.62)$$

$$V_{2\Gamma}(\mathbf{r})/\text{meV} = -5093.6 + 18.5 \sum_{i=1,3,5} 2\cos(\mathbf{g}_{i} \cdot \mathbf{r} + 0.47)$$

$$T_{1\Gamma}(\mathbf{r})/\text{meV} = 218.6 + 1.2 \sum_{i=1,3,5} 2\cos(\mathbf{g}_{i} \cdot \mathbf{r} + 1.84)$$

$$T_{2\Gamma}(\mathbf{r})/\text{meV} = 220.3 - 7.8 \sum_{i=1,3,5} 2\cos(\mathbf{g}_{i} \cdot \mathbf{r} + 2.18)$$
(3)

which specifies the continuum model.

A similar construction can be made for the K or K' valley bands. We focus on the K valley, which is related to the K' valley by time reversal. The main difference from the Γ bands is that interlayer tunneling is suppressed, and hence $H_K = -\frac{|\mathbf{k}|^2}{2m_K} + \mathrm{diag}(V_{1K}(\mathbf{r}), V_{2K}(\mathbf{r}), V_{1K}(\mathbf{r}), V_{1K}(\mathbf{r}))$, and \mathbf{k} is now measured relative to the K point. The first two energies E_K at each stacking region specify the potentials V_{1K} and V_{2K} . However, there is now an additional ambiguity as the eigenvalues are invariant under interchanging $V_{1K} \leftrightarrow V_{2K}$. Additional information is needed to determine whether the first pair of bands should be associated with V_{1K} (the outer two layers) or V_{2K} (the inner two layers). From direct inspection of the DFT wave function, we find that the first band has larger component on the outer two layers at all stacking regions. Hence, $V_{1K} > V_{2K}$ and the potentials are fully specified in the first harmonic approximation:

$$V_{1K}(\mathbf{r})/\text{meV} = -4731.2 + 18.1 \sum_{i=1,3,5} 2\cos(\mathbf{g}_i \cdot \mathbf{r} + 0.75)$$

$$V_{2K}(\mathbf{r})/\text{meV} = -4744.6 + 19.5 \sum_{i=1,3,5} 2\cos(\mathbf{g}_i \cdot \mathbf{r} + 0.76)$$
(4)

We use the effective mass $m_K = 0.5 m_e$ [4]. The moiré band structure for the Γ and K valley bands is shown in Fig S6. To address the competition between Γ and K valley bands in a displacement field, we model the displacement field by a layer-dependent potential difference Δ via the term $H_{\Delta} = -\Delta \operatorname{diag}(0, 1, 2, 3)$. At $\nu = 0$, the layer potential is related to the displacement field D by $\Delta = dD/\epsilon$, where ϵ is the relative out-of-plane dielectric constant and d is the interlayer spacing.

We remark that the competition between Γ and K band edges from first principles is often delicate, and can depend sensitively on relaxed lattice structure and van der Waals density functional (see, for instance, Refs. [5] and [6], in which different functionals result in different Γ, K band orderings in both twisted homobilayer WSe₂ and MoTe₂). In addition, interaction effects may also result in a renormalization of the band gap measured experimentally. To compare quantitatively with experiments, we thus take as an additional tuning parameter a constant energy offset Δ_K between Γ and K band edges, modifying the Hamiltonian as $H_{\Gamma} \to H_{\Gamma} - \Delta_{\Gamma K}/2$ and $H_K \to H_K + \Delta_{\Gamma K}/2$. The experimentally measured gap is maximum at D=0, and decreases continuously and smoothly with non-zero D: this is consistent with the scenario in which the state at $\nu=-1$ corresponds to filling a spin-polarized Γ moiré band and the gap being to the first K or K' moiré band. We choose $\Delta_{\Gamma K}=0.1$ eV such that the gap between the first Γ and K bands roughly match the experimentally measured value at D=0. Once $\Delta_{\Gamma K}$ is fixed, the dependence of the gap




FIG. S6. Moiré bands from the continuum model. The Γ (left) and K (right) valley moiré bands obtained from the four layer continuum model.





FIG. S7. Schematic of single-particle moiré bands at low and high D. At D=0, the lowest energy single-particle moiré bands are two-fold (spin) degenerate at the Gamma point. The subsequent moiré band, localized at the K/K' points, is eight-fold degenerate (4x layer, 2x spin/valley). At $D>D_c$, the critical displacement field at which the K/K' moiré band is the lowest energy band, the lowest energy moiré band is completely localized on a single layer as favored by the displacement field with a two-fold degeneracy from spin/valley (up at K / down at K', or vice versa depending on the direction of D). The next moiré band will be the same two-fold degenerate Γ moiré band.

on Δ follows from the continuum model, as shown in Fig. 3f of the main text. The closing, rather than opening of the gap as a function of D is only consistent with a Γ moiré band being the lowest energy hole band at D=0. The band degeneracies at each valley and their relative energies are schematically illustrated in Fig. S7.

We note that this theoretical picture is also consistent with the lack of a gap observed at $\nu = -2$ in the experiment. Specifically, at low displacement fields, our calculations suggest the states filled beyond $\nu = -1$ are in a second, more dispersive band localized at the K-valley. At $\nu = -2$, this second band will be partially filled, and its larger bandwidth makes it less likely to form a correlated insulator.






FIG. S8. K-valley band density distribution. Density distribution in the first K valley band without (left) and with (right) the electrostatic potential from the filled Γ band at $\nu = -1$, calculated from the continuum model. Without interactions, the charges are localized at MM', whereas including the electrostatic potential shifts them to XX'.

To conclude this section, we comment on the nature of the lowest energy K valley excitations at $\nu=-1$. In the absence of interactions, it is clear from the moiré potential that holes prefer to localize on the MM' sites for both Γ and K valley. However, holes filling into the K valley beyond $\nu=-1$, will feel an effective potential resulting from electrostatic repulsion with the filled Γ band. This effect can be captured by a Hartree potential in the continuum model, $V_H(\mathbf{r}) = \int d\mathbf{r} V_C(\mathbf{r} - \mathbf{r}') \langle n^{(\Gamma)}(\mathbf{r}') \rangle$ where $\langle n^{(\Gamma)}(\mathbf{r}') \rangle$ is the density distribution of the filled first Γ band (obtained from the non-interacting continuum model), and $V_C(\mathbf{r})$ is the Coulomb potential. The continuum model description of the first K band now includes $V_H(\mathbf{r})$ in addition to the moiré potential. We further simplify by taking \mathbf{r} to be purely in the 2D plane and use a vertically displaced Coulomb potential $V_C(\mathbf{r}) = e^2/(4\pi\epsilon\sqrt{\mathbf{r}^2+d^2})$, where $\epsilon=3.5\epsilon_0$ and d=7 Å. Figure S8 shows the hole density of the first K valley band with and without the Hartree term. As can be seen, holes in the K valley prefer to localize in the XX' regions once electrostatic repulsion from the filled Γ band is taken into account, resulting in an effective honeycomb lattice description near $\nu=-1$.

4. SPIN q-FACTORS

To quantitatively compare the experimental gaps and their increase in a magnetic field to our theoretical model, we must consider the Zeeman contribution to the effective g-factors of the various excitations. Our measurement of a gap at filling ν , $\Delta_{\nu} \equiv \Delta \mu$ is equivalent to the particle-hole excitation energy, given by $E[N_{\nu}+1]+E[N_{\nu}-1]-2E[N_{\nu}]$, where E[N] is the ground state energy at particle number N, and N_{ν} denotes the particle number at filling ν . First, we consider the case at high magnetic fields (i.e., the right side of Fig. 2i in the main text), where the ground state is a spin-polarized insulator with charge transfer energy gap $\Delta_{\rm CT}$, the high-density excitation is an additional interstitial hole with aligned spin, and the low-density excitation is the absence of a hole. Then, we can write down the energies as follows, which results in a gap that is independent of magnetic field:

$$E[N_{\nu}] = -\frac{1}{2}g_sN_{\nu}B, \quad E[N_{\nu}+1] = \Delta_{\text{CT}} - \frac{1}{2}g_s(N_{\nu}+1)B, \quad E[N_{\nu}-1] = -\frac{1}{2}g_s(N_{\nu}-1)B$$

$$\Delta_{\nu} = E[N_{\nu} + 1] + E[N_{\nu} - 1] - 2E[N_{\nu}] = \Delta_{\text{CT}} - \frac{1}{2}g_sB + \frac{1}{2}g_sB = \Delta_{\text{CT}}$$

At intermediate fields (i.e., the left side of Fig. 2i in the main text), where the only change is that the hole excitation is the formation of a spin polaron, we instead have

$$E[N_{\nu}] = -\frac{1}{2}g_s N_{\nu} B, \quad E[N_{\nu} + 1] = \Delta_{\text{CT}} - E_b - \frac{1}{2}g_s (N_{\nu} + 1)B, \quad E[N_{\nu} - 1] = -\frac{1}{2}g_s (N_{\nu} - 2)B + \frac{1}{2}g_s B$$

$$\Delta_{\nu} = E[N_{\nu} + 1] + E[N_{\nu} - 1] - 2E[N_{\nu}] = \Delta_{\text{CT}} - E_b - \frac{1}{2}g_sB + \frac{3}{2}g_sB = \Delta_{\text{CT}} - E_b + g_sB$$

where $E_b > 0$ is the spin polaron binding energy. Given that the bare spin g-factor $g_s = 2$, the measured effective g-factor of our gapped states would be 2 in the absence of orbital/Berry curvature and exchange effects.

As mentioned in the main text, orbital and Berry curvature differences between different moiré bands (at K and Γ) can lead to additional contributions to g^* at $\nu=-1$. The momentum-resolved orbital magnetization m(k) can vary widely across the the moiré Brillouin zone (mBZ) [?]. Theoretical calculation of m(k) requires calculation of the Berry curvature on the single-particle moiré bands and can depend sensitively on shape of the moiré bands, and therefore the twist angle [6?, 7]. Because our probe is only sensitive to the lowest energy excitation, the measured gap will depend sensitively on the extremal value of m(k) across the mBZ, which can lead to different effective g-factors between samples with different twist angles, and may contribute to the slightly different g-factors measured at $\nu=-1$ in Samples S1 and S2.

5. ALTERNATIVE EXCITATIONS AT $\nu = -\frac{1}{3}$

In the main text, we mention an alternative explanation of the saturating gap at $\nu = -\frac{1}{3}$, namely that below B = 7T, exchange coupling causes the lowest energy additional hole to be anti-aligned with the spin-polarized ground state, until Zeeman coupling overcomes this barrier and the additional hole is aligned with the rest of the spins. In this section, we expand on that picture and discuss why it is not supported by our data and calculations. In this picture, the change in behavior around B=7 T occurs on the excitations below the gap (i.e., μ_{-} in the main text). In Fig. S9a, we show the expected chemical potentials for each excitation (above and below the gap) in this scenario, in an analagous way to Fig. 2i in the main text (which is reproduced as S9b for ease of comparison). If the changing spin character of the additional hole excitation were driving the saturating gap, then we would expect the discontinuity in the slope of the chemical potential on the band edge to occur at μ_- , rather than as we observe at μ_+ . Additionally, the energy scale of J that our data would predict in this case does not appear to be consistent with other microscopic parameters of the problem. To lowest order, we could estimate exchange coupling J as $\frac{1}{2}g^*\mu_B B'$ where g^* is the effective g-factor that we measure for the gap, μ_B is the Bohr magneton, B'=7 T is where the behavior of the gap changes, and the factor of $\frac{1}{3}$ is due to the fact that an added hole couples to three nearest neighbors. This yields $J \approx 0.6$ meV, which is quite large compared with other parameters in the problem. For example, if we take $J \sim 4t^2/U$ where t is the hopping and U is the onsite potential, then we would estimate $J \approx 0.01$ meV, as $t \approx 0.2$ meV from the bandwidth of the lowest moiré band from the continuum model while $U \approx 100$ meV due to the strength of Coulomb interactions [8].

The two scenarios of excitations considered in the main text operate off of the understanding of the insulators as spin-polarized throughout most of the measured range of magnetic fields. One possibility to consider as a potential explanation for the behavior of the gap at $\nu = -\frac{1}{3}$ is that rather than a change to the excitations at B=6 T, the spin character of the gapped state itself changes, e.g. by becoming spin polarized above B=6 T but being unpolarized below that field. We rule out this possibility as unlikely both due to theoretical considerations — the next-neighbor spin-spin coupling should be very weak — and experimental ones — if the gapped state were not polarized by the magnetic field, we would not expect the gap to linearly increase in a magnetic field as measured in Fig. 2c.

Further, the scenarios considered in the main text focus broadly on how the magnetic field couples to the spinful gapped states and excitations through Zeeman coupling. However, there are additional non-Zeeman orbital mechanisms by which the magnetic field can couple into the system. We have considered two such possibilities: i) orbital effects within a moiré site can become important when the magnetic length is comparable to the size of the Wannier orbitals, leading to a change in the effective hopping/interaction strengths, ii) orbital effects can lead to a variation in the energy of excitations via formation of Hofstadter/LL bands. However, these effects should be almost negligible for realistic parameters: for our maximal magnetic field B=11 T, the magnetic length $l_B\approx 8$ nm (which is already larger than the moiré period), and the slope of the gap, were it to come from a purely orbital mechanism, suggests an unrealistically small effective mass $m\approx 0.2m_e$ for the hole excitations about the $\nu=-\frac{1}{3}$ state, and further cannot explain the discontinuity in the slope. Additionally, we do not observe any prominent Hofstadter or LL gaps, which would be expected if such orbital effects were dominant. While we cannot rule out the existence of more exotic mechanisms that we are not aware of, these considerations suggest that the role of orbital effects is minor in this system. Nevertheless, further measurements in a tilted magnetic field could help clarify the relative strengths of orbital and Zeeman coupling.

Our measurements of the correlated insulating gaps provide evidence for spin polaron excitations, and more recent theory predicts a finite range of parameter space in which a metallic state involving such quasiparticles will be stabilized [9]. This compressible phase should have some novel physical properties which the measurements in the present manuscript are not sensitive to. For example, tunneling measurements should find that this state has a single particle gap (for tunneling in electrons) despite the fact that it is compressible. Additionally, there should be a simultaneous plateau in the magnetization throughout this phase. These measurements would provide complementary confirmation of novel itinerant quasiparticle excitations, and would be excellent directions for future experiments.




FIG. S9. Schematic of magnetic field dependence of possible excitations in the considered scenarios. a, Analogue of Fig. 2i for the scenario detailed in Supplementary Sec. 5, in which the excitation below the gap changes from an anti-aligned to aligned hole at B=B', while the excitation above the gap is the removal of a hole at all fields. We show the chemical potential μ as a function of B for various excitations of the (spin-polarized) correlated insulating ground state. The blue line and inset show an excitation above the gap in which a single hole is removed. The green line and inset, the favored excitation below the gap at low magnetic fields, show the addition of a spin anti-aligned hole at an interstitial site. The black line and inset show the addition of an aligned spin at an interstitial site, which is the favored excitation below the gap at high magnetic fields. b, Fig. 2i in the main text, reproduced here for ease of comparison.

6. EXTRACTION OF CHEMICAL POTENTIAL μ_+ AND μ_-

The SET tip directly measures changes in chemical potential of the sample by tracking a particular Coulomb blockade oscillation of the SET while the sample density is tuned by a gate. In order to evaluate whether the discontinuity in the slope of the gap at $\nu = -\frac{1}{3}$ comes from a change on the particle or hole side of the gap, we need to evaluate the behavior of the chemical potential on either side of the gap as a function of magnetic field, which is the "slow" axis in our measurement. One approach is to pick a particular point (say the band edge, or some location far from the gap) and choose this as a reference point at which μ is set to zero at every magnetic field. However, this can give different behavior depending on the choice of reference point, especially when the overall behavior of $\mu(n)$ changes smoothly with magnetic field. For example, in Fig. S11, it is apparent that the chemical potential at the band edge is controlled by the magnitude of negative compressibility; if the minimum point is chosen to be the "reference" point where $\mu = 0$ for each value of B, then the trends in both μ_+ and μ_- will mostly be determined by the change in the negative compressibility, rather than identifying the behavior in the vicinity of the gap itself.

To avoid such artifacts, we take advantage of the fact that the gap at $\nu=-\frac{1}{3}$ is small compared to the background chemical potential. We can fit a smooth polynomial background to the chemical potential on either side of the gap at $\nu=-\frac{1}{3}$ and subtract this background, as in Fig. S4. Then the value of the chemical potential on either side of the gap is measured relative to the background chemical potential of filling the lowest energy moiré band between 0 and 1. The precise values of μ_+ and μ_- and can vary slightly with details of the fit, but the fact that the $\mathrm{d}\mu_+/\mathrm{d}B$ changes at B=7 T while $\mathrm{d}\mu_-/\mathrm{d}B$ does not is consistent. We note that this essentially is a statement of the evolution of the chemical potential relative to the background filling of the lowest moiré band.

At $\nu = -1$, the behavior of $\mu(n)$ in both samples is dramatically different on either side of the gap, preventing an equivalent type of subtraction. Focusing on Sample S1, in Fig. S10 we show extracted chemical potential on either side of the $\nu = -1$ gap with two different choices of reference chemical potential. In Fig. S10a, we set the reference chemical potential to $\nu = -1.25$. Practically, this means that we take the average measured chemical potential in a certain region (red shaded box in Fig. S10c) and subtract it from the measurement at each field such that it becomes the "zero" point of the chemical potential, before then reading off μ_+ and μ_- on either side of the gap. In Fig. S10b, we perform the same analysis but with the reference point taken near the middle of the first moiré band (blue

shaded box in Fig. S10c). In both of these cases, μ_{-} does not change very much; most of the gap dependence thus relates to the change of μ_{+} . Given that we choose partially filled moiré bands as references, this suggests that the dependence of μ_{-} on magnetic field is quite close to the overall change in chemical potential of the bands themselves, while μ_{+} varies strongly relative to this. This qualitatively agrees with the theoretical picture described in the main text, as we would expect that at all experimentally accessible fields, the lowest-energy excitation at μ_{-} would be adding another spin-aligned hole, whose dependence on B would be identical to filling the spin-polarized bands. The excitation at μ_{+} on the other hand, would be due to the formation of spin-polarons, which should have a different (stronger) dependence on magnetic field than the surrounding moiré bands (Fig. 2i in the main text). Therefore, our data also suggest spin polaron excitation are relevant at $\nu = -1$, but the smoking gun signature of a change in gap size and chemical potential evolution with magnetic field are not accessible in our measurement setup.




FIG. S10. Behavior of chemical potential around $\nu=-1$. a, Chemical potential on the left (μ_-) and right (μ_+) sides of $\nu=-1$ taking the reference μ at high densities around $\nu=-1.25$. b, Same as a, but with the reference μ set to intermediate densities around $\nu=-0.5$. c, $\mu(n)$ measured at B=0. The average μ within the red shaded box is set to zero for the data in a, while the average μ within the blue shaded box is set to zero for the data in b.

7. NEGATIVE COMPRESSIBILITY FROM EXCHANGE INTERACTIONS

Here, we present evidence of strong interaction effects even away from the charge-ordered states. In Sample S2, we observe negative compressibility over a wide range of hole density, with $d\mu/dn$ becoming increasingly negative as hole density decreases (Fig. S11a). Negative compressibility is also ubiquitous in Sample S1 (Fig. 2b in the main text), but does not extend as close to the band edge due to a smooth, positive background signal for $|n| < 1.5 \times 10^{12}$ cm⁻² (see below). We attribute this difference at low densities to varying TMD quality, possibly due to different commercial sources for the two samples, and focus on Sample S2 first.

Capacitance enhancements (equivalent to negative $d\mu/dn$) have been reported in TMD moiré heterostructures when the sample-gate distance d is comparable to or less than the moiré length scale $L_M \approx a/\theta$ where a is the lattice constant of WSe₂ [10, 11]. In our devices, sample-gate coupling is small as $d/L_M \approx 4-5$, and much of the parameter space where we measure negative $d\mu/dn$ is in the limit of $nd^2 \gg 1$. In this regime, negative $d\mu/dn$ is generally attributed to strong Coulomb interactions [2, 12–15]. We can understand the magnitude of negative compressibility




FIG. S11. Negative compressibility from exchange interactions. a, $\mu(n)$ in Sample S2 at B=0,5,10 T. Dashed line fit is to $\mu(n)=\frac{\pi\hbar^2}{m^*}n-A\sqrt{n}$ where m^* and A are free parameters. We set $\mu=0$ at high hole density for each trace. b, Magnitude of the coefficient A from the fit as a function of B. The red shaded box marks the B=0 prediction from a Hartree-Fock model of a 2D electron gas where the height shows uncertainty in dielectric environment. The blue shaded box marks the predicted growth in exchange due to full spin polarization, namely $\sqrt{2} \times A(B=0 \text{ T})$.

by comparing to the Hartree-Fock (HF) expression for the chemical potential of a 2DEG:

$$\mu(n) = \frac{\pi \hbar^2 n}{m^*} - \left(\frac{8}{\pi}\right)^{\frac{1}{2}} \left(\frac{e^2}{4\pi\epsilon}\right) \sqrt{n}$$

These two oppositely signed terms describe the density of states and exchange contributions to the chemical potential and will always lead to negative $d\mu/dn$ at sufficiently low density [12, 13]. However, the low kinetic energy due to flat electronic bands in tdWSe₂ causes the negative contribution from exchange interactions to dominate $\mu(n)$ to much higher hole densities than in more dispersive electronic systems. This provides experimental evidence that the underlying moiré bands that we are filling in this density range are quite flat, consistent with our DFT calculations. At zero field, the extracted coefficient from a fit to this form (black dashed curve, Fig. S11a, see Supplementary Sec. 10 for fitting details) matches the order of magnitude of predictions from HF exchange based on the hBN dielectric environment (red shaded box, Fig. S11b). This coefficient, which correlates with the magnitude of the negative compressibility, grows monotonically before saturating around 7 T (Fig. S11b). While the exact value of this coefficient is sensitive to details of the fit, the order of magnitude and relative growth in a magnetic field are consistent (Supplementary Sec. 10, below). This increase is due to polarization of the compressible states being filled, as spin polarization will lead to an increase in the exchange interaction. Again in analogy to the 2DEG HF model, this increase saturates when the metallic phase is fully polarized, with a total increase by a factor $\sqrt{2}$, consistent

with our observations (blue shaded box, Fig. S11b) [16]. This suggests that the magnetic ordering is paramagnetic through much of the density range, outside of the correlated insulator gaps.

Though we focus on Sample S2 in the discussion above, we also observe negative compressibility over large regions of hole doping in Sample S1, before an upturn at low hole density (Fig. S12). The magnitude of negative compressibility, as estimated from the average $|d\mu/dn|$ between $\nu=-.6$ and $\nu=-0.95$ also grows in a magnetic field, though no clear saturation is observed.




FIG. S12. Negative $d\mu/dn$ in Sample S1. a, $\mu(n)$ as measured in S1 across the entire density range. b, Average value of $|d\mu/dn|$ between $n = -3.8 \times 10^{12}$ cm⁻² and $n = -2.5 \times 10^{12}$ cm⁻².

8. FITTING NEGATIVE COMPRESSIBILITY $\mu(n)$

Here we expand on the details of the fit to $\mu(n)$ in Sample S2 shown in Fig. S11. While the order of magnitude of the coefficient A is robust to variation of fitting parameters/schemes, as is its increase by a factor of $\sqrt{2}$ from B=0 to high fields, the precise value of A is somewhat sensitive to details of the fitting procedure. In Fig. S13, we present the extracted A from 3 different fits, changing the range of density included in the fit. The blue data is that which is presented in the main text, because fitting over the widest possible density range seems to provide the best fit to the data. However, given the complicated details of the moiré bands, it is not clear which of these is most physically justifiable. Fitting across such a large density range means that we include filling over multiple moiré bands, each of which will have a varying effective density of states as they are filled. Additionally, it involves fitting across the charge-ordered gap at $\nu=-1$, around which the effective density n may be better approximated by $|n-n_{\nu}|$, where n_{ν} is the charge density at moiré band filling ν . It is apparent in these fits that between $n=-2\times 10^{12}$ cm⁻² and $n=-2.5\times 10^{12}$ cm⁻², just beyond $\nu=-1$, the behavior of $\mu(n)$ varies sharply compared to the background. For the red line, which fits to just the lowest moiré band between $\nu=0$ and $\nu=-1$, the fit has no linear component. In the blue and green fits in Fig. S13b, the size of the linear term is $\frac{\pi\hbar^2}{m^*}\approx 1.25\times 10^{-11}$ meV/cm⁻² and $\frac{\pi\hbar^2}{m^*}\approx 1.75\times 10^{-11}$ meV/cm⁻², respectively. Both of these terms would suggest effective masses much smaller than expected for moiré bands $(m^*=0.1-0.15m_e)$, which is inconsistent with the size of the Landau level gaps we observe simultaneously (Fig. S16). Because these fits are over the entirety of multiple moiré bands, the effective mass is not necessarily expected to remain constant across the fit.

The analogy to the Hartree-Fock (HF) model in tdWSe₂ is a simplification, but it nonetheless remains qualitatively accurate, as detailed below. Specifically, the free electron gas calculation does not incorporate electron-electron correlations nor does it account for the moiré potential, which affects the kinetic energy. To determine how these details alter predictions for the chemical potential, we apply DFT to directly study the continuum model of interacting electrons in the periodic moiré potential, as developed in [17]. We use the local spin density functional (LSD) approximation, obtained from fixed node diffusion quantum Monte Carlo simulations of a two dimensional electron

gas. By construction, the LSD approximation is exact for homogeneous systems. In practice, DFT with the LSD approximation provides a better description of moiré systems with smooth potential compared to the inhomogeneous systems: atoms, molecules and natural solids. In moiré materials with a smooth potential landscape and small bandwidth ($\sim 10 \text{ meV}$), the strongly correlated physics happens within the lowest few bands, and we can safely choose the energy cutoff as 10 eV. This leads to using 31×31 plane wave components, independent of the moiré wavelength or number of atoms in one unit cell.

Because the experimental data spans a wide range of moiré fillings, it interpolates between filling Γ and K moiré bands at low and high fillings factors, respectively. In Fig. S14, we show the resulting moiré DFT calculations of $\mu(\nu)$ for both the Γ and K moiré bands. We note that this calculation is only performed for higher filling factors because it becomes unreliable at low filling, due to the possibility of magnetic and/or charge-ordered states. Due to this constraint, the K-valley calculations are more relevant to the corresponding portion of the experimental data, but we present both for full transparency. For each valley, we also present the HF prediction (dashed line) as a comparison as well as the homogeneous electron gas (HEG) prediction from the moiré DFT, which is the limit where the moiré potential is set to zero, but the correlations are calculated via DFT. All three curves are qualitatively similar, overlapping over broad ranges of density despite small quantitative differences at some fillings. The agreement between these theoretical calculations justifies applying the simpler HF expression to fit the negative compressibility over the full density range. The most prominent discrepancies arise near moiré filling factors (see for example, around $\nu = -2$), where gaps appear in the moiré DFT calculations but not in the HF model. This is similar to the experimental data, where there is sharp behavior (both the gap and subsequent sharper negative compressibility) between $\nu = -1.4$ and $\nu = -1$ in Fig. S11 which is not captured by the smooth HF fit.




FIG. S13. Dependence of coefficient A on fit parameters. a, Extracted coefficient A from a fit to $\mu(n) = \frac{\pi \hbar^2}{m^*} n - A \sqrt{n}$ where m^* and A are free parameters. Blue, green, and red data correspond to fitting the entire curve out to $n = -5 \times 10^{12}$ cm⁻², $n = -4 \times 10^{12}$ cm⁻², and $n = -2 \times 10^{12}$ cm⁻² ($\nu = -1$), respectively. b, Measured $\mu(n)$ at B = 5T (black) overlaid with fitted lines corresponding to those described in panel **a**.

9. CAPACITANCE DATA FROM SAMPLE C2

In Fig. S15, we present $d\mu/dn$ of Sample C2 as a function of D and ν . The data are extracted from "top-gate capacitance" in which we apply a 10 mV a.c. excitation at 337 Hz to the top gate and measure the response on the HEMT gate, which is electrically connected to the sample contacts. The data were taken at T=4.2 K. The range that we can access on the lower (upper) left side are limited by top (bottom) gate dielectrics.




FIG. S14. Moiré DFT calculations of chemical potential. $\mu(\nu)$ calculated using moiré DFT for moiré bands at both K and Γ valleys. The red and blue curves, labeled HEG for homogeneous electron gas, show the limit where the moiré potential is set to zero, while the orange and purple curves take realistic values of the moiré potential, in line with DFT calculations for tdWSe₂. The vertical offsets of these curves are arbitrary.

10. ESTIMATING D IN SET MEASUREMENTS

In general, the effective layer potential difference in a singly-gated two-dimensional (2D) heterostructure device is a difficult problem that depends in part on details of the material being gated [18, 19]. Even the usual definition of an applied electric displacement field, which takes the average of the displacement field on either side of the gated 2D material, is an approximation and not necessarily directly convertible to microscopic parameters such as layer potential differences. Comparing our scanning SET measurements, in which the samples have no top gate, to measurements with a controlled displacement field is therefore nontrivial. However, a lowest order estimate can be made by approximating the set-up as a dual gated device where the SET tip is treated as an effective "top gate" held at the same voltage as the sample. This voltage constraint is accurate because the relative sample-tip voltage is held by d.c. feedback in our experiment to minimize tip-induced doping. Additionally, the tip is held $\sim 50-100$ nm away in vacuum, such that the effective "top-gate" capacitance will be quite small compared to that of the back gate, and any small offsets in voltage will not provide very much doping. We note that this is distinct from what has been reported in recent STM measurements, in which the close tip-sample distance requires the displacement field to be approximated by the voltage difference between tip and back gate [20]. Because the SET tip is far-separated and senses a spatial area underneath the tip that convolves both tip size and distance, any voltage separation between tip and sample leads to (small) non-uniform doping of the sample, resulting in broadening features rather than reliably doping the sample, which limits our ability to change the effective displacement field.

Using this approximation, the effective displacement field can be written as $D_{\text{eff}} = \frac{1}{2\epsilon_0}(0 - C_b(V_b - V_0)) = -\frac{1}{2\epsilon_0}en$. Here, C_b is the back gate capacitance, V_b is the back gate voltage, V_0 is offset of the density due to intrinsic sample doping, and we've used that $C_b(V_b - V_0)/e = n$. Applying this expression to relevant densities in our samples, in Sample S1 at $\nu = -1$, $D_{\text{eff}} = 0.35 \text{ V/nm}$, and in sample S2, $D_{\text{eff}} = 0.18 \text{ V/nm}$; this nonzero displacement field explains




FIG. S15. Displacement field dependence in a second device. Measured $d\mu/dn$ in Sample C2 as a function of displacement field D and moiré filling factor ν .

the reduction in gap size relative to the D=0 value we measure in dual gated devices. We note that for Sample S1, this estimated D is near and slightly above the critical displacement field at which we stop seeing insulating behavior at $\nu=-1$ in dual gated capacitance devices, though the precise value varies slightly across the two devices measured. However, the fact that we measure a gap at all at $\nu=-1$ in the singly-gated devices is strongly suggestive that the lowest energy moiré bands being filled are still at Γ , and suggest that the estimated $D_{\rm eff}$ should be taken as a rough approximation. Additionally, global measurements in our dual gated devices inevitably average over disorder, which may lead to lower overall gap sizes relative to local sensing with the scanning SET. This will artificially decrease the critical displacement field at which the $\nu=-1$ gap vanishes in the dual gated devices.

11. TWIST ANGLE AND BAND EDGE DETERMINATION

In order to quote accurate densities, we need to determine the location of the band edge in gate voltage, which is set by both local (global) intrinsic doping in the sample for SET (dual gated) measurements and by work function differences between gates and sample [21]. For both samples measured with the SET, the gate voltages at which we observe Landau level gaps allows for unambiguous determination of the band edge, which also matches well to the mobility edge (Fig. S16). In Sample S1, we do not observe clear Landau levels in the tdWSe₂ region of the device (Fig. 2a), which may reflect the large effective mass of the moiré bands and/or the proximity to charge ordered states at the low fillings we measure (see below). In comparison, Landau levels are present in an adjacent Bernal bilayer WSe₂ region of Sample S1, from one of the "halves" of the flake that makes up the device. Assuming the entire flake has similar intrinsic doping, this sets the gate voltage offset of the band edge in the twisted double bilayer region of the device as well. In Sample S2, we observe two-fold degenerate landau levels between $\nu = -1$ and $\nu = -3$ near the noise floor of our measurement, with gap sizes of order $200 - 300 \ \mu\text{V}$ at $B = 11 \ \text{T}$. This corresponds to a filling factor range which we were unable to explore in Sample S1.

Here, we expand our discussion on the absence of Landau levels at low densities in both samples. The overall static disorder in these samples is consistent with previous reports. From Fig. S16a, we measure a full-width at half-maximum of 4×10^{10} cm⁻² for the incompressible peaks of quantum Hall states in the adjacent Bernal bilayer region in Sample S1. This estimate of disorder is similar to quoted values in previous high-quality measurements of monolayer WSe₂ [21]. Additionally, we measure very little spatial variation in the twisted double bilayer regions

of both devices (Supplementary Sec. 1). Thus, we rule out spatially inhomogeneous twist angle as a reason for the absence of Landau levels. We instead ascribe the lack of quantum Hall states below $\nu=-1$ to the flatness of the underlying bands, such that quantum Hall gaps are suppressed. From our continuum model theory (Supplementary Sec. 3), the bandwidth of the lowest energy moiré band is quite small, with a correspondingly small tight binding model hopping parameter t=0.2 meV. Expanding the dispersion about Gamma gives an effective mass $m^*=\frac{\hbar^2}{2ta^2}$, where a is the lattice constant of the moiré lattice, 5.6 nm for our 3.4° twisted sample. This yields an effective mass $m^*=6.1m_e$ where m_e is the electron mass, and a corresponding Landau level gap size of 0.2 meV at B=11 T. This is essentially at our experimental noise floor. While the additional interface in the twisted region could trap disorder beyond that present in the pristine bilayers, which could suppress Landau levels in the twisted region independent of the effective mass of the flat moiré bands, the widths of the incompressible peaks in each respective region of the sample are of the same order of magnitude. This suggests that the extrinsic effect of interfacial disorder on Landau levels is minimal.

In Samples C1/C2, we did not conduct measurements at nonzero magnetic field and thus must rely on the capacitance measurements themselves to estimate the band edge. In Fig. S17, we show a linetrace of raw capacitance signal taken at a much higher contact gate voltage. This has the effect of improving the sample contact resistance without affecting the device region measured via capacitance to the top gate, as the top gate screens out any stray field from the contact gates which are above it. In most of the measured parameter space, the signal was indeed unaffected. However, it did lead to a nonzero signal extending to lower and lower densities. We take the point where this signal becomes zero within the signal to noise as zero density, and note that this also agrees with the spacing between the $\nu = -1$ and $\nu = -3$ features in Fig. S11 (i.e., the gate voltage range between $\nu = -1$ and $\nu = -3$ is twice that between $\nu = -1$ and the band edge, $\nu = 0$).




FIG. S16. Landau levels and band edge determination. a, Landau levels observed in a Bernal bilayer region in Sample S1. Dotted lines are filling factor $\nu = 3, 4, \dots, 24, 25$. No clear Landau levels were observed in the tdWSe₂ region of the device. b, Landau levels in tdWSe₂ beyond $\nu = -1$ in Sample S2. These show a two-fold degeneracy; dotted lines are filling factor $\nu = 14, 16, \dots, 24$.




FIG. S17. Capacitance signal measured at higher contact gate voltages. A 9 mV a.c. signal is applied to the top gate at a frequency of 911 Hz, and top and back gate voltages are swept to maintain D = 0 V/nm.

We also provide more details confirming the twist angle of all samples studied. A common technique used to measure the twist angle of TMD devices is second harmonic generation (SHG) [22, 23]. However, due to the inversion symmetry of bilayer TMDs, they do not have a SHG response, so that technique for measuring the twist angle is not generally possible in double bilayer samples. Serendipitously, however, the original bilayer WSe₂ flake stacked to obtain Sample S1 also had an adjacent monolayer region (Fig. S18a). Portions of this contiguous monolayer area were also twisted and stacked simultaneously with the bilayer parts of the flake, yielding identical twist angle as in the twisted double bilayer regions. We conducted polarization dependent SHG measurements on a monolayer region from each half of the twisted device (yellow and purple dots, Fig. S18b), and obtained a twist angle of 3.61 \pm 0.27 degrees by fitting to the oscillations of each signal (Fig. S18c). This represents an independent measure of twist angle, and we obtained similar values throughout the monolayer areas. The quoted value in the text (3.4°), which takes the observed incompressible state at $n = -3.9 \times 10^{12}$ cm⁻² to be $\nu = -1$ indeed falls within this range.




FIG. S18. Second harmonic generation (SHG) reflectance on Sample S1.a, Optical image of the WSe₂ flake used to stack Sample S1. The two halves respectively are outlined in dark blue and red. The monolayer pieces of each half are outlined in cyan and lime green, with the rest of the darker portion of the flake being bilayer. At the bottom of the flake, a small area of thicker WSe₂ is present. b, Optical image of Sample S1 with the locations of the SHG signal measured in panel c indicated by purple and yellow dots, with the same pieces of the WSe₂ flakes outlined as in panel a. The scale bar is 10 μ m. c, SHG signal as a function of polarization angle θ on two monolayer regions of Sample S1.

Other measured samples do not have monolayer regions, so SHG is not informative. However, optical and atomic force microscopy (AFM) images support the twist angle assignments we infer from electronic compressibility measurements for all devices. In Table S2, we provide additional images of each sample before and after twisting/stacking, and we enumerate three angles. The first angle is the target twist angle during the stacking process, i.e. the angle to which the stage was rotated between picking up the two halves of the twisted device. We also list the angle estimated from optical and AFM images. The images in Table S2 are annotated with blue and red lines that highlight edges of the parent flake which are also clear in the final stacking images. We present optical images because they provide better contrast for the entire TMD flakes, and therefore give more accurate measurements of twist angle. However, analysis of the edges measured with AFM yield consistent results. We estimate the angle from optical/AFM images based on the twist angle required to make both of these edges align with the original flake, with uncertainty estimated based on how sharp the edges are by eye. Finally, we quote the angle determined from the electronic measurements. These are extracted based on the densities at which we observe incompressible states, which we assume to occur at integer moiré filling.

For each sample, the angles all mutually agree within experimental uncertainty. In all cases, the target angle during stacking matches the angle determined from the density at which we observed incompressible states to within 0.3°. This is comparable to values typically quoted in the literature. In addition, both angles match the range extracted from optical/AFM images within experimental uncertainty. In contrast, alternate filling factor assignments for the observed incompressible states are not consistent with experimental observations. For example, the most compelling and experimentally plausible alternate filling factor assignment for the most prominent incompressible state would be $\nu = -2$, full filling of the lowest energy moiré band. However, the twist angle that would be required for this falls outside the experimental range of twist angle uncertainty for all four devices. For example, the twist angle which would be required for the gap at $n = -3.9 \times 10^{12}$ cm⁻² in Sample S1 to correspond to $\nu = -2$ would be roughly 2.5°. That angle is well below the range of either the experimentally measured SHG or the estimates of twist angle from optical/AFM images. Finally, we note that the high degree of twist angle homogeneity observed over micron length scales in local scanning SET measurements makes local twist angle variations distinct from the device edges in optical/AFM images highly unlikely.

12. TEMPERATURE DEPENDENCE OF CAPACITANCE DATA

The temperature dependence of Sample C1 is shown up to T=15 K in Fig. S19. The magnitude of the incompressible peak at $\nu=-1$ weakens with increasing temperature. In Fig. S19a-b, X-channel (capacitive) data is presented in units of the geometrical top-gate capacitance c_t . As the temperature is increased, the charging of the tdWSe₂ sample at low densities improves. This is particularly notable at the maroon dashed line ($\nu=-\frac{2}{3}$) where at T=4.2 K the capacitance is at $\approx 0.7c_t$ but it quickly rises as the mobility edge extends rightwards. A more modest version of this effect is also seen at $\nu=-1$ (black), whereas in the negative compressibility region (pink), the temperature dependence is nonmonotonic, first increasing (presumably) due to better charging before decreasing at the highest temperatures. In Fig. S19c, we show line cuts at constant D=-0.2 V/nm at a few temperatures. The incompressible peak at $\nu=-1$ is weaker at 12 K relative to 4.2 K, but still clearly evident. In Fig. S20 we show the behavior of the $\nu=-1$ gap as a function of D at a few distinct temperatures. As in previous reports, the gaps show a linear decrease in gap size as the temperature increases, likely due to thermal broadening [10]. The slope of gap size Δ_{-1} with D is similar across various temperatures.

13. ESTIMATION OF QUANTUM CAPACITANCE FROM GLOBAL CAPACITANCE

As discussed in the Methods, we convert the capacitance signal into units of the geometric capacitance c_t by assuming $C_T = 0$ in the gap between the WSe₂ valence and conduction bands and $C_T = c_t$ at high hole dopings $> 5 \times 10^{12} \text{cm}^{-2}$. Using a lumped circuit capacitance model [10, 24], the measured top gate capacitance C_T is given by

$$C_T = \frac{c_t c_q}{c_t + c_b + c_q} \tag{5}$$

where c_t and c_b are the geometric top and bottom gate capacitances and $c_q \equiv e^2(d\mu/dn)^{-1}$ is the quantum capacitance. As discussed in the main text, this approximation will overestimate $d\mu/dn$ in the case of high sample resistance or high frequency. Some of this enhancement can be accounted for by including the simultaneous measurement of the

Sample	Image of original flake	Image after twisting and stacking	Target angle (stage rotation during stacking)	Angle (from optical / AFM image)	Angle (from compressibility)
S1			3.5°	$3.5 \pm 0.3^{\circ}$ (SHG: $3.6 \pm 0.3^{\circ}$)	3.4°
S2			2.5°	$2.6 \pm 0.4^{\circ}$	2.6°
C1	*		2.5°	$2.5 \pm 0.7^{\circ}$	2.2°
C2			3.0°	$3.0 \pm 0.5^{\circ}$	3.1°

TABLE S2. For all samples studied in the manuscript, we provide images of the "parent" bilayer WSe₂ flakes and images of the devices once fully stacked/twisted. Using red and blue lines, we highlight clear edges that appear from each half of the flake in the final stacking; these edges are used to estimate the angle from the optical image. Thin red lines on the image of original flakes (on S1 and S2) are due to annotations and should be ignored. For each device, we provide the angle to which we twisted the stage while stacking, the angle estimated from the optical/AFM images, and the angle extracted from compressibility experiments based on the density of observed features.




FIG. S19. Temperature dependence in Sample C1. a, Temperature dependence of measured capacitance C_T in units of geometric capacitance c_t . An 8 mV excitation at a frequency of 739.1 Hz and a constant d.c. voltage -1.6 V is applied to the top gate. b, Line cuts as a function of temperature from panel a at $\nu = -1$, $\nu = -0.875$, $\nu = -\frac{2}{3}$. c Measured $d\mu/dn$ at D = -0.2 V/nm at various temperatures.




FIG. S20. Temperature dependence of the $\nu = -1$ gap.

out-of-phase quadrature of the measured capacitance, which is equivalent to the dissipation signal [25, 26]. In this

case, the total measured signal is given by

$$V^{\text{complex}} \propto C_T/c_t = \frac{c_q}{c_t + c_b + c_q} \frac{\tanh(\alpha)}{\alpha}$$
 (6)

$$\alpha = \sqrt{i\omega \frac{c_q c_t}{c_t + c_b + c_q} R_s} \tag{7}$$

where ω is the measurement frequency and R_s is the sample resistance. Note that in the small α limit, the imaginary (capacitive) part of this signal agrees with Eq. 5 and the real (dissipative) part is zero. This system can be solved for c_q and R_s using both the x and y quadratures, and the resulting c_q yields $d\mu/dn$ as before. An example of x and y quadrature capacitance signal and the extracted $d\mu/dn$ by both of these methods are shown in Fig. S21. All data presented from capacitance measurements in the main text show $d\mu/dn$ calculated using this correction. The main effect of including the dissipation on the extracted $d\mu/dn$ occurs in the regions of parameter space which exhibit negative compressibility; there is also a consistent small decrease of the broad peak at $\nu = -1$ (Fig. S21).

In Fig. S22 we show a comparison of the measured charge gap at $\nu=-1$ as a function of applied displacement field D at various frequencies. At 4.7 kHz, the measured gap is clearly enhanced above the lower frequency curves. Below 1.7 kHz and 739 Hz, there is little measured change. This lack of change suggests that the extraction of $d\mu/dn$ from Eqs. 6 and 7 performs adequately at these measurement frequencies.




FIG. S21. Example capacitance measurements on Sample C1. a, Line cuts showing the capacitance signal in the X (capacitance) and Y (dissipation) channels. b, Extracted $d\mu/dn$ signal using each respective method discussed in the text. Data were measured at a temperature of 4.2 K with an 8 mV excitation applied to the top gate at frequency 739.1 Hz and a constant d.c. voltage -1.6 V on the top gate.




FIG. S22. Frequency dependence. Measured gap Δ_{-1} in Sample C1 at various frequencies.

* These authors contributed equally

 † bef@stanford.edu

[1] A. Uri, S. Grover, Y. Cao, J. A. Crosse, K. Bagani, D. Rodan-Legrain, Y. Myasoedov, K. Watanabe, T. Taniguchi, P. Moon, M. Koshino, P. Jarillo-Herrero, and E. Zeldov, Nature 581, 47 (2020), number: 7806 Publisher: Nature Publishing Group.

- [2] U. Zondiner, A. Rozen, D. Rodan-Legrain, Y. Cao, R. Queiroz, T. Taniguchi, K. Watanabe, Y. Oreg, F. von Oppen, A. Stern, E. Berg, P. Jarillo-Herrero, and S. Ilani, Nature 582, 203 (2020), number: 7811 Publisher: Nature Publishing Group.
- [3] J. Yu, B. A. Foutty, Z. Han, M. E. Barber, Y. Schattner, K. Watanabe, T. Taniguchi, P. Phillips, Z.-X. Shen, S. A. Kivelson, and B. E. Feldman, arXiv:2108.00009 [cond-mat] (2021), arXiv: 2108.00009.
- [4] H. C. Movva, T. Lovorn, B. Fallahazad, S. Larentis, K. Kim, T. Taniguchi, K. Watanabe, S. K. Banerjee, A. H. MacDonald, and E. Tutuc, Physical Review Letters 120, 107703 (2018), publisher: American Physical Society.
- [5] F. Wu, T. Lovorn, E. Tutuc, I. Martin, and A. MacDonald, Physical Review Letters 122, 086402 (2019), publisher: American Physical Society.
- [6] T. Devakul, V. Crépel, Y. Zhang, and L. Fu, Nature Communications 12, 6730 (2021), number: 1 Publisher: Nature Publishing Group.
- [7] W.-Y. He, D. Goldhaber-Gordon, and K. T. Law, Nature Communications 11, 1650 (2020), number: 1 Publisher: Nature Publishing Group.
- [8] Y. Zhang, N. F. Q. Yuan, and L. Fu, Physical Review B 102, 201115 (2020), publisher: American Physical Society.
- [9] Y. Zhang and L. Fu, (2022), 10.48550/arXiv.2209.05430.
- [10] T. Li, J. Zhu, Y. Tang, K. Watanabe, T. Taniguchi, V. Elser, J. Shan, and K. F. Mak, Nature Nanotechnology 16, 1068 (2021).
- [11] B. Skinner and B. I. Shklovskii, Physical Review B 82, 155111 (2010).
- [12] J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, Physical Review Letters 68, 674 (1992).
- [13] J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, Physical Review B 50, 1760 (1994).

- [14] L. Li, C. Richter, S. Paetel, T. Kopp, J. Mannhart, and R. C. Ashoori, Science **332**, 825 (2011), publisher: American Association for the Advancement of Science.
- [15] F. Yang, A. A. Zibrov, R. Bai, T. Taniguchi, K. Watanabe, M. P. Zaletel, and A. F. Young, Physical Review Letters 126, 156802 (2021), publisher: American Physical Society.
- [16] B. Tanatar and D. M. Ceperley, Physical Review B 39, 5005 (1989).
- [17] Y. Zhang, H. Isobe, and L. Fu, "Density functional approach to correlated moire states: itinerant magnetism," (2020), arXiv:2005.04238 [cond-mat].
- [18] A. Kou, B. E. Feldman, A. J. Levin, B. I. Halperin, K. Watanabe, T. Taniguchi, and A. Yacoby, Science **345**, 55 (2014), publisher: American Association for the Advancement of Science.
- [19] E. McCann, Physical Review B 74, 161403 (2006), publisher: American Physical Society.
- [20] A. Kerelsky, C. Rubio-Verdú, L. Xian, D. M. Kennes, D. Halbertal, N. Finney, L. Song, S. Turkel, L. Wang, K. Watanabe, T. Taniguchi, J. Hone, C. Dean, D. N. Basov, A. Rubio, and A. N. Pasupathy, Proceedings of the National Academy of Sciences 118, e2017366118 (2021), publisher: Proceedings of the National Academy of Sciences.
- [21] M. V. Gustafsson, M. Yankowitz, C. Forsythe, D. Rhodes, K. Watanabe, T. Taniguchi, J. Hone, X. Zhu, and C. R. Dean, Nature Materials 17, 411 (2018).
- [22] L. Wang, E.-M. Shih, A. Ghiotto, L. Xian, D. A. Rhodes, C. Tan, M. Claassen, D. M. Kennes, Y. Bai, B. Kim, K. Watanabe, T. Taniguchi, X. Zhu, J. Hone, A. Rubio, A. N. Pasupathy, and C. R. Dean, Nature Materials 19, 861 (2020), number: 8 Publisher: Nature Publishing Group.
- [23] Y. Tang, L. Li, T. Li, Y. Xu, S. Liu, K. Barmak, K. Watanabe, T. Taniguchi, A. H. MacDonald, J. Shan, and K. F. Mak, Nature 579, 353 (2020), number: 7799 Publisher: Nature Publishing Group.
- [24] Q. Shi, E.-M. Shih, M. V. Gustafsson, D. A. Rhodes, B. Kim, K. Watanabe, T. Taniguchi, Z. Papić, J. Hone, and C. R. Dean, Nature Nanotechnology 15, 569 (2020), number: 7 Publisher: Nature Publishing Group.
- [25] R. K. Goodall, R. J. Higgins, and J. P. Harrang, Physical Review B 31, 6597 (1985), publisher: American Physical Society.
- [26] Q. Shi, E.-M. Shih, D. Rhodes, B. Kim, K. Barmak, K. Watanabe, T. Taniguchi, Z. Papić, D. A. Abanin, J. Hone, and C. R. Dean, Nature Nanotechnology, 1 (2022), publisher: Nature Publishing Group.