

Sustainable irrigation and climate feedbacks

Received: 8 January 2023

Accepted: 6 July 2023

Published online: 17 August 2023

 Check for updates

Yi Yang  ¹, Zhenong Jin  ²✉, Nathaniel D. Mueller  ^{3,4}✉, Avery W. Driscoll  ⁴, Rebecca R. Hernandez ^{5,6}, Steven M. Grodsky ^{7,13}, Lindsey L. Sloat ^{3,4,14}, Mikhail V. Chester ⁸, Yong-Guan Zhu ^{9,10} & David B. Lobell  ^{11,12}

Agricultural irrigation induces greenhouse gas emissions directly from soils or indirectly through the use of energy or construction of dams and irrigation infrastructure, while climate change affects irrigation demand, water availability and the greenhouse gas intensity of irrigation energy. Here, we present a scoping review to elaborate on these irrigation–climate linkages by synthesizing knowledge across different fields, emphasizing the growing role climate change may have in driving future irrigation expansion and reinforcing some of the positive feedbacks. This Review underscores the urgent need to promote and adopt sustainable irrigation, especially in regions dominated by strong, positive feedbacks.

Irrigation expanded substantially across the globe in the twentieth century, contributing to increased crop productivity¹. Without irrigation, global cereal production on irrigated lands would decrease by nearly 50% and total cereal production would decrease by 20% (ref. ²). Irrigation is expected to continue expanding, partly to meet increasing food demand, but notably to improve the adaptability of crop systems to climate change and variability^{3,4}.

The expansion of irrigation might have important consequences for the climate system on global and local scales through greenhouse gas (GHG) emissions and biophysical pathways. Irrigation causes GHG emissions from energy use and facility construction^{5–7}. It can also directly affect nitrous oxide (N₂O), methane (CH₄) and soil carbon emissions from cropland, and indirectly induce these emissions from canals and reservoirs constructed for farm irrigation^{8,9}. In addition, irrigation has a local cooling effect that is well documented in the hydro-climatic literature¹⁰. Another potentially beneficial effect of irrigation on climate change is that by improving crop yields, irrigation can spare natural environments from being cleared for crop production^{11,12}.

Climate change, on the other hand, also affects irrigation. Shifting precipitation patterns, for example, can drive irrigation expansion, but also impact the water and energy systems in which irrigation

is embedded. As climate change continues to intensify^{13,14}, it is crucial to understand how it impacts irrigation and consequently how irrigation-related activities may feed back to the climate system. These impacts can augment the total GHG emissions of the irrigation system and result in potentially meaningful positive climate feedbacks. Overall, these bidirectional feedback loops have not yet been articulated in the large and growing literature on the food–energy–water nexus¹⁵.

Here, by reviewing studies published over the past decade, we synthesize the various irrigation–climate linkages (Fig. 1); evaluate the impacts of climate change on irrigation systems, including irrigation infrastructure and the food–energy–water systems in which it is embedded; and identify areas in which climate change may intensify irrigation-related GHG emissions. Further, we present emerging and innovative solutions that can facilitate the development of sustainable irrigation under climate change. We close by discussing knowledge gaps and future research needs and priorities.

Climate impacts of irrigation

Energy use and associated GHG emissions

Irrigation activities can produce GHG emissions directly when pumps run on diesel or natural gas, or indirectly when pumps use electricity.

¹Key Laboratory of the Three Gorges Reservoir Region's Eco-Environment, Ministry of Education, Chongqing University, Chongqing, China. ²Department of Bioproducts and Biosystems Engineering, University of Minnesota, St. Paul, MN, USA. ³Department of Ecosystem Science and Sustainability, Colorado State University, Fort Collins, CO, USA. ⁴Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA. ⁵Wild Energy Center, Institute of the Environment, Davis, CA, USA. ⁶Department of Land, Air & Water Resources, University of California, Davis, CA, USA. ⁷Institute of the Environment, University of California, Davis, CA, USA. ⁸School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ, USA. ⁹Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China. ¹⁰Research Center for Eco-environmental Sciences, Chinese Academy of Sciences, Beijing, China. ¹¹Center on Food Security and the Environment, Stanford University, Stanford, CA, USA. ¹²Department of Earth System Science, Stanford University, Stanford, CA, USA. ¹³Present address: New York Cooperative Fish and Wildlife Research Unit, US Geological Survey, Ithaca, NY, USA. ¹⁴Present address: Land and Carbon Lab, World Resources Institute, Washington, DC, USA.  e-mail: jinz@umn.edu; nathan.mueller@colostate.edu




Fig. 1 | Illustration of the climate impacts of irrigation. Direct and indirect GHG emissions (or savings), as well as local cooling effects, associated with a conventional irrigation system that uses a mix of groundwater and surface water (partly transferred from other basins), and runs on internal combustion or electric engines with electricity sourced from hydropower and thermopower.

The climate impacts of irrigation can be local (by affecting local temperatures and cropland biogenic GHG emissions), regional (by affecting electricity generation or interbasin water transfer) and global (by affecting land use elsewhere through crop yield changes).

If powered by electricity, the carbon intensity of irrigation depends on the fuel mix of the regional grid; a higher share of fossil fuels in the grid would yield a greater carbon intensity. Additionally, water source is a critical factor of irrigation-associated energy use and emissions. Pumping groundwater is generally much more energy intensive than pumping surface water because of the additional lift needed (for example, 2,100–4,000 kJ m^{-3} versus 3–4 kJ m^{-3} in the Lower Indus Basin of Pakistan⁶). Owing to its ubiquity and consistency, global groundwater use for irrigation has increased substantially in the twentieth century and now supplies ~40% of all irrigated area¹⁶. Irrigation water can also be transferred from other basins. Depending on the distance and elevation change, the energy intensity of interbasin water transfers can be very high (for example, twice that of groundwater in a case from China¹⁷). Interbasin water transfers have also seen substantial growth, and currently dozens of large-scale water transfer projects are planned or under construction globally, with the majority intended for irrigation use¹⁸.

Biogenic emissions

In addition to energy-related emissions, irrigation affects fluxes of CH_4 , N_2O and CO_2 from croplands. Irrigation, particularly in flooded rice production, is an important driver of global CH_4 emissions due to the anaerobic conditions it creates that favour methanogenic bacteria. Research shows that continuous flooding leads to twice the CH_4 emissions as intermittent flooding¹⁹. Irrigation also increases N_2O emissions by increasing soil moisture and, consequently, stimulating the nitrification and denitrification processes that produce soil N_2O emissions⁸. Field-scale comparisons show that N_2O emissions can increase by 50–140% in irrigated versus non-irrigated fields, although the magnitude of change depends on many factors such as N application rates, soil properties and irrigation intensity²⁰. On the other hand, irrigation

may increase soil carbon storage when it increases plant productivity and hence litter into the soil⁸. However, higher moisture due to irrigation also stimulates plant decomposition, resulting in CO_2 emissions⁸. On average, irrigation enhances soil organic carbon storage in arid and semi-arid areas, not in humid environments, with larger increases tied to lower initial carbon stocks and less precipitation⁸. However, site-level effects of irrigation on soil organic carbon are mixed^{8,20,21} and depth-dependent²², and effects on inorganic carbon stocks are relatively understudied despite being an important component of total carbon stocks in many agronomic systems.

Beyond on-farm emissions, additional GHG emissions are incurred from human-constructed bodies of water that transport and store water for irrigation (for example, canals and reservoirs). In the past century, the number of reservoirs for irrigation grew considerably, resulting in an approximately 25-fold increase in irrigation water supply (from 18 to 460 $\text{km}^3 \text{yr}^{-1}$)²³. Irrigation reservoirs vary in size, and building large dams requires a particularly substantial amount of carbon-intensive materials such as concrete. The emissions embedded in materials can be partially offset when the dams generate hydropower. However, water competition in cases of reduced precipitation or irrigation expansion could reduce hydropower output²⁴, which might result in greater thermal power generation. Regardless, artificial reservoirs can release significant amounts of GHGs by converting organic matter in the flooded areas into CH_4 , CO_2 and N_2O , and also by increasing CH_4 bubbling from sediments²⁵. A recent study identified CH_4 from reservoirs as a main contributor to the carbon footprint of irrigation in Spain²⁶. The impacts of irrigation on GHG emissions and other environmental issues may be reconciled and mitigated by optimizing the siting of reservoirs, as shown in a study of Amazonian hydropower dams²⁷.

Biophysical feedback

Irrigation can modify local or regional temperature and humidity through multiple biophysical mechanisms²⁸. By increasing the availability of water to vegetation, irrigation raises evapotranspiration and the associated latent heat flux. This process lowers air temperature because more energy is used for water vaporization rather than heating the air. However, higher evapotranspiration and humid atmosphere resulting from irrigation tend to foster increased cloud cover, which reflects more shortwave radiation and leads to further cooling, but also amplifies the local greenhouse effect and may contribute to heat stress^{29,30}. During the daytime, the dominance of increased latent heat flux among these contrasting effects often leads to a net cooling effect of irrigation^{10,31,32}. For example, a recent modelling analysis suggests that crop canopy temperatures can be as much as 10 °C lower than ambient air temperature under well-irrigated conditions³³. In the Indo-Gangetic Plain, air temperatures in irrigated croplands are significantly cooler than in non-irrigated areas by up to 1–2 °C during the crop-growing season, as inferred from satellite observations³¹. By contrast, the effects of irrigation on nighttime temperatures are not well studied, but some evidence suggests irrigation could warm nighttime temperatures by increasing soil heat storage³⁴ or enhancing the local greenhouse effect associated with increased atmospheric humidity³⁰, and possibly more than offset the daytime cooling effect³⁴. A frontier research area is the investigation of climate teleconnections associated with irrigation^{35,36}.

As climate change progresses, there is growing concern regarding the escalating risk of humid heat extremes caused by intensified irrigation. Recent studies based on regional or global climate model simulations indicate that irrigation increases wet-bulb temperatures and the frequency of dangerous heat extremes in various regions, including the North China Plain, the central USA and the Middle East^{37,38}. Similarly, satellite and in situ observations found that reduced planetary boundary layer height, as a result of irrigation-induced reduction in sensible heat flux, raises humid heat stress in India, Pakistan and Afghanistan³⁹. While humid heat extremes may have had minimal impacts on or even enhanced yields in some regions⁴⁰, they pose a growing health hazard for agricultural workers worldwide²⁸.

Reduced incentives for land clearing

Increased crop yields from irrigation can potentially reduce GHG emissions by decreasing incentives for land clearing. Irrigation is critical to plants in arid or semi-arid regions with limited rainfall, but even in humid regions, irrigation can increase crop yields by compensating for seasonal rainfall variability and deficits⁴¹. The irrigation-induced cooling effect also contributes to yield gains by mitigating canopy heat stress and atmospheric water demand; for example, a recent study on maize in Nebraska shows that 16% of yield increase from irrigation can be attributed to the cooling effect, with the remaining 84% due to other physiological benefits of increased water supply⁴¹. Without irrigation, global cereal production would drop by around 20% (ref. 2), thus requiring more land to meet agricultural demands.

Despite the importance of irrigation to global crop production, the ‘land sparing’ benefits of irrigation-driven yield increases to global GHG emissions are complex and largely unquantified. Nevertheless, studies across a range of modelling complexities support the notion that agricultural intensification, in general, contributes to decreases in agricultural land use at the global scale, as lower prices reduce pressure for land conversion^{11,42,43}. Quantifying the contributions of irrigation to global land sparing would also require accounting for interactions between supply and demand, prices, trade and input substitution using complex economic models subject to considerable uncertainty⁴⁴.

The spatial configuration of spared land associated with irrigation is also important to consider, because aboveground and belowground carbon stocks, as well as crop productivity, vary substantially across the globe⁴⁵. Additional mechanisms, including land-use zoning, economic

instruments, spatially targeted agricultural investments and voluntary standards or certifications are often needed to proactively link yield increases with the protection of natural ecosystems⁴⁶. Further, the biophysical local climate impacts of land clearing also vary in sign depending on latitude, with substantial local warming from tropical deforestation⁴⁷.

Finally, irrigation may be required for the expansion of bioenergy with carbon capture and storage, a pivotal negative emission technology for meeting climate targets⁴⁸, with implications for total agricultural land use and land sparing⁴⁹. While irrigation will boost yields of bioenergy crops and decrease land requirements, it may drive water consumption and increase global water stress⁵⁰. In addition, bioenergy has been criticized for diverting crops and land away from the food supply, thus raising prices and stimulating land-use change⁵¹.

Growing impacts of climate change

Greater irrigation demand

Greater irrigation demand, resulting from climate-driven changes in regional precipitation and evapotranspiration, would trigger most of the irrigation-induced climate effects. Even when the total precipitation remains constant or increases, future shifts in subseasonal precipitation variability may spur more droughts and irrigation use⁵², although moderately intensified heavy rainfall may offset some drought damage⁵³. Rising temperature also increases evaporation of surface water and plant transpiration, and can reduce photosynthetic rates, notably in C3 plants (for example, wheat and soybean). To achieve comparable yields, farmers may respond by increasing irrigation intensity⁸.

On a global scale, the net impact of climate change on irrigation demand remains uncertain. Significant uncertainties remain around (1) how arid lands (a quarter of Earth’s surface) will respond to increases in irrigation; (2) how humans will use irrigation as an adaptation to climate change; and (3) to what extent elevated CO₂ concentrations can mitigate irrigation needs. The effect of CO₂ has been an area of intense research. For example, one study noted an 8–15% global irrigation reduction by the end of the century with elevated CO₂, compared with a 0–5% rise without factoring in CO₂ (ref. 54). Similarly, another study identified net decreases in irrigation demand using the LPJmL model with CO₂, despite regional increases due to local climate change patterns⁵⁵. However, more recent field experiments have found that elevated CO₂ can increase the photosynthesis as well as canopy size for C3 crops, which counteract the water savings from lower stomatal conductance⁵⁶. Thus, additional irrigation may be needed to fully realize the productivity benefits of elevated CO₂ for many major staple crops^{57,58}, although the net climate outcome remains to be investigated. Indeed, satellite observations have shown a global decline of the CO₂ fertilization effect on vegetation productivity since the 1980s, probably as a result of changes in terrestrial water storage⁵⁹.

Despite uncertainties around changes in irrigation water demand, it is clear that many agricultural regions will face climate challenges relevant to irrigation, including decreases in soil moisture⁶⁰, rising vapour pressure deficit⁶¹, and changes in the magnitude and timing of surface water availability for irrigation, particularly in snow-dependent basins⁶². Even if climate change elicits a net-zero impact on future global irrigation use, it might ultimately increase total irrigation-induced energy use and carbon emissions due to a shift towards water sources that are more energy intensive or carbon intensive, such as groundwater or reservoirs, as discussed below. This could more than offset the energy saved in wetter places projected to require less irrigation in the future.

Greater reliance on groundwater

Increases in overall irrigation demand or decreases in surface water availability from changes in hydrological cycles can increase reliance on more energy- and carbon-intensive water sources (for example, from groundwater and interbasin transfers). In particular, climate

change is likely to exacerbate the need for groundwater use⁶³ by reducing precipitation in some regions and decreasing summer flows in snowmelt-dominated basins⁶⁴. In California, groundwater, critical to agricultural and economic resilience, constitutes 40% of total water use in wet years and 60% in droughts⁶⁵. Similar substitutions of groundwater for surface water have been observed in other regions due to hydroclimate variability⁶⁶.

Irrigation using groundwater requires more energy than with surface water – a climate-driven human adaptation that could result in a positive climate feedback (Fig. 2a). This feedback could be further intensified when persistent deficits in annual recharge combine with continuous over-drafting, leading to lower groundwater levels and higher energy costs of pumping⁶⁷. In Punjab, India, groundwater use increased by 23% and the water table dropped by 5.47 m during 1998–2012, resulting in a doubling of annual carbon emissions⁶⁸. In addition, groundwater contains CO₂ and N₂O because of its interactions with subterranean environments such as soil, minerals and bacteria. When exposed to the atmosphere, these GHGs are released or degassed. The magnitude of the degassed GHGs depends on the properties of groundwater, but is probably small compared with other sources of agricultural GHG emissions^{69,70}.

Severe groundwater depletion in regions reliant on heavily over-exploited aquifers can lead to eventual abandonment of irrigation and/or stricter regulation on inefficient pumping⁷¹. These responses may cause their own climate impacts. For instance, irrigation abandonment may reduce local cooling effects and subsequent yield decreases may increase pressure for land conversion elsewhere. Globally, however, the majority of aquifers remain underexploited⁷², indicating that substantial opportunity for increased groundwater reliance remains.

Greater GHG intensities of irrigation energy

When irrigation is powered by grid electricity, climate change can also affect the grid system in ways that increase grid GHG intensities (Fig. 2a), thus increasing the life-cycle GHG emissions of irrigation. Of particular concern are changes in the availability of hydropower because of its vulnerability to climate variability. Shifts in the patterns of rainfall, snowmelt and glacier melt can lead to lower annual runoff and consequently lower hydropower output, which may increase the dependency on coal, oil or natural gas thermopower to make up for supply shortages⁷³. This climate-driven substitution increases the GHG intensity of grid electricity per kilowatt-hour generated. During the recent drought in Western Europe (2016–2017), Spain's hydropower generation dropped by ~50%, resulting in more thermopower from mostly combined-cycle and coal-fired power plants and 18% additional CO₂ emissions compared with the previous year⁷⁴. Across the western USA, repeated droughts over the past two decades have led to increased power generation from coal and natural gas, and substantially increased CO₂ and other air emissions⁷⁵. Increased precipitation and runoff, on the other hand, could create a negative climate feedback by increasing hydropower output, but too much water could result in equipment damage, outage and dam repairment⁷⁶, potentially offsetting the climate benefit from the negative feedback.

In much the same way, expansion of irrigation can increase grid GHG emissions, owing to the competition for water. Globally, about half of hydropower capacity competes with irrigation²⁴. In these regions, irrigation expansion can reduce the amount of water available for hydropower use and lead to more fossil-fuel-based power generation. Furthermore, climate change can significantly exacerbate water competition among irrigation, hydropower and thermopower (as in prolonged and intense droughts), resulting in greater use of groundwater by both irrigation and energy, higher thermoelectric output and, consequently, substantially higher system-wide energy and carbon intensities than without climate change.

Increased biogenic emissions

Climate change can also increase the biogenic emissions associated with irrigation directly and indirectly. First, temperature and water interact to positively affect soil N₂O emissions. Thus, the N₂O emissions intensity of irrigated cropland might increase as temperature increases⁷⁷, all else being equal. Second, climate change is projected to intensify CH₄ emissions from rice paddies owing to both warming and elevated CO₂ levels. Warming increases the rates of plant root decay and soil organic matter decomposition, which stimulates the growth of methanogenic bacteria⁷⁸. A 1°C of rise in temperature has been estimated to increase rice CH₄ emissions by ~10% (ref. 79). Elevated CO₂ promotes rice root growth and root exudates, resulting in more carbon sources for methanogenic bacteria⁷⁹. Research shows that elevated CO₂ levels (550–743 ppm) may increase rice CH₄ emissions intensity by 30–40%, although the effect can be moderated by incorporation of straw into rice fields⁷⁹.

Third, reservoirs, like groundwater, are an important source to help agriculture adapt to hydroclimatic change and variability. Climate change is projected to increase the demand for reservoirs, especially in regions projected to experience reduced rainfall and snowpack⁸⁰. However, not only may the number of reservoirs grow, but higher temperatures will also increase the intensity of biogenic emissions per reservoir (Fig. 2b). Warming increases the rates of aquatic plant decay and soil organic matter decomposition, which, in turn, stimulates the growth of methanogenic bacteria⁷⁸. For irrigation reservoirs that are eutrophic, which are quite common worldwide⁸¹, warming may also aggravate the emission of GHGs, particularly CH₄. In eutrophic reservoirs, excess nutrients already fuel algae growth and decomposition, which creates an oxygen-poor condition that favours methanogenic bacteria⁸², and warming will intensify this process by further stimulating algae growth⁸³. Studies suggest warming could increase CH₄ emissions intensity from lakes globally by 13–40% by the end of this century⁸⁴. Moreover, climate change may increase the extent of eutrophication among reservoirs, owing partly to increased runoff resulting from shifts in precipitation and partly to increased temperatures, further intensifying the process of CH₄ production.

Sustainable irrigation solutions and innovations

That climate change may intensify the climate impacts of irrigation underscores the urgent need to accelerate the development of sustainable irrigation. Various strategies have long been promoted, including enhancing efficiency with drip systems, improved scheduling, leakage reduction and adopting conservation practices. The wide-scale adoption of these strategies will moderate the projected increase in overall irrigation water use and the number of irrigation-oriented reservoirs needed. Here, we emphasize challenges and tradeoffs involved in some of the innovations that have recently emerged. Promoting these strategies is especially important in regions vulnerable to positive climate feedbacks, aridity, increased groundwater reliance, heightened water resource competition between irrigation and energy, and extensive rice cultivation.

Reduce biogenic CH₄ and N₂O emissions

The potency of CH₄ and the significant contribution of flooded rice paddies to global CH₄ emissions, together with the potential intensifying impact of climate change, highlight the urgency to reduce CH₄ emissions from rice production. But existing GHG mitigation methods often involve tradeoffs. For example, intermittent flooding (for example, midseason drainage) can effectively depress CH₄ emissions from rice fields – as well as water use⁸⁵ – and hence is a potentially important climate adaptation strategy. But it might also increase soil N₂O emissions⁸⁶. Straw incorporation can largely moderate the impact of elevated CO₂ levels on rice CH₄ emissions, but the straw itself is also a source of GHG emissions⁷⁹. Reducing rice CH₄ requires a systems




Fig. 2 | Conceptual models of climate-irrigation feedbacks. **a–c**, Climate change can increase GHG emissions from irrigation energy systems (a) and irrigation reservoir systems (b), and reduce GHG emissions through increasing crop yields (c). Panels on the left indicate mechanisms by which irrigation or

agriculture induce GHG emissions under a relatively stable climate. Panels on the right indicate how climate change affects these mechanisms and leads to greater or lower GHG intensity of irrigation (represented by the thicker or thinner arrows, as opposed to those on the left). SOM, soil organic matter.

approach that manages multiple factors simultaneously to minimize these tradeoffs⁸⁶. Emerging technologies such as biochar application may also be helpful⁸⁷. In other crop systems, switching from furrow

or sprinkler irrigation to drip irrigation – which reduces the extent of denitrification via partial wetting of soils⁸⁸ – can decrease soil N₂O emissions by 32–46% (ref. 88). Irrigation coupled with conservation

tillage can also increase soil organic carbon sequestration compared with conventional tillage⁸.

The increasing demand for irrigation-oriented reservoirs and water transfers in response to climate variability and change presents challenges as well as opportunities. Opportunities arise with new reservoirs as they can be designed to minimize potential GHG emissions. Measures to mitigate emissions include limiting the input of nutrients and organic matter, avoiding a rapid drawdown (which promotes CH₄ emissions), and increasing oxygen concentrations in the water⁹. Covering reservoirs or canals with solar panels can deliver carbon, water and land benefits⁸⁹ (see some examples from California in Fig. 3). Irrigation reservoirs covered by floating solar energy with some power clipped to run an aerator have been shown to help reduce GHG emissions via reduced water temperature and increased dissolved oxygen⁹⁰. Large reservoirs covered by solar panels can produce substantial amounts of energy, but there are potential tradeoffs – for example, effects on aquatic biota and terrestrial wildlife, and on the ecological and recreational values of reservoirs – that must be considered and minimized⁸⁹. Large reservoirs with high GHG emissions can be monitored and involved in carbon credit programmes, which provide financial incentives for mitigation. Freshwater systems such as reservoirs, lakes and ponds are now receiving increasing interest and becoming targets of national GHG mitigation commitments.

Power irrigation with renewables

The reciprocal feedback between climate change and irrigation necessitates the expansion of low-carbon irrigation. Diesel- and gasoline-powered irrigation engines, although less efficient than electric ones, are still widely used globally⁹¹. Irrigation electrification, alongside grid decarbonization, can reduce energy consumption and GHG emissions. However, large-scale clean electricity implementation poses challenges and may not benefit off-grid smallholders in remote areas. For them, the strategy is to install renewable energy generators such as solar-, wind- and water-powered pumps. The selection of renewable sources should adapt to site-specific hydrological and socioeconomic conditions and align with the temporal needs of agricultural production.

Widespread adoption of renewable pumps will depend largely on cost reduction, power purchase agreement decisions, innovative business models and long-term community promoter presence⁹². De-risking investments for development partners and unbanked smallholders is a key area to prioritize, as financing these systems will support climate change mitigation and safeguard vulnerable farmers' livelihoods. When there is social cohesion, group-based pump sharing can facilitate access to finance for the initial capital investment, especially for poor farmers. In many sub-Saharan regions, solar pumps may not be cost-effective within a 25-year period without monetizing environmental benefits⁹³, thus requiring new microfinancing mechanisms. Cases with preliminary success were achieved by using Internet of Things and mobile payments technologies to offer flexible payment plans as a way to align payments with farmers' income patterns⁹⁴. GHG reductions from switching to renewable pumps in poor countries can be monetized and, if compensated by rich countries through financial transfers, could facilitate the adoption of renewable pumps in those countries.

To avoid over-abstraction of groundwater that can emerge from reduced irrigation operational costs, renewable pumps should be integrated into strong regulatory frameworks on sustainable water resource use. Feasibility studies for renewable-powered irrigation systems often focus on technical and economic aspects but lack an assessment of water resource availability and impact. However, a drop in groundwater levels, caused by either climate change or overexploitation, can negatively affect agricultural productivity and economic feasibility of those renewable irrigation systems. Opportunities do exist when on-farm generated renewable energy is used for other purposes,

encouraging farmers to make rational decisions about pumping. In western India, a grid-connected scheme has been implemented to buy back surplus solar power from farmers to prevent excessive withdrawals⁹⁵. Besides financial incentives, educating on integrated soil, water and energy management is vital for system viability⁹⁶. Addressing climate risks in climate-vulnerable regions⁹⁷ is also crucial for minimizing downtime and asset loss of renewable pumps.

Create techno–ecological synergies

Beyond providing electric power, on-farm solar energy can be designed to facilitate techno–ecological synergies that deliver broad benefits to humans and nature. Such systems can maintain crop yields while generating benefits, including reduced irrigation water consumption and reduced GHG emissions associated with water pumping. For example, agrivoltaics are a techno–ecological synergy that co-locates solar energy and crop production⁹⁸. In northwestern India, modelling demonstrated that water inputs for cleaning solar panels are the same as those required for annual aloe production, such that the co-location of solar panels and aloe may yield higher returns per cubic metre of water than either system alone⁹⁹. Agrivoltaics may reduce evapotranspiration, retain more soil moisture and hence reduce irrigation demand due to altered microclimatic conditions by solar arrays¹⁰⁰. The partial shade of solar panels may also provide a cooling effect for crops underneath agrivoltaics systems and bolster yield¹⁰⁰. Adoption remains low for agrivoltaics; however, governments including China, France, Germany, Japan and the USA have supported agrivoltaics development via research investments as well as regulatory permitting pathways and/or incentives.

Solar energy production on marginalized and abandoned farmland, as well as on reservoirs⁸⁹, may spare prime agricultural land with comparatively moister and less saline soils¹⁰¹, and facilitate carbon sequestration, especially when coupled with sustainable development practices such as revegetation and soil amendments (for example, biochar)¹⁰², leading to a climate feedback loop with potentially lower irrigation demand and GHG emissions. Additive solar energy in agricultural landscapes may be developed in lands adjacent to farmland and in the negative space (that is, uncultivated areas) of agricultural fields. The groundcover, interspace and borders of ground-mounted solar energy facilities adjacent to agricultural land may be restored with plants comprising pollinator habitat¹⁰³, which can increase pollination services in nearby agricultural fields (for example, within 1.5 km)¹⁰⁴ that may act in conjunction with abiotic factors, including water stress, to affect crop yield. Additionally, farmers can develop solar energy, underlaid by native pollinator habitat¹⁰⁴, in the corners of agricultural fields irrigated with centre-pivot technology to make use of unirrigated, negative space¹⁰⁵ that may bolster food system resilience, biodiversity conservation and land sparing – outcomes that address climate change and biodiversity goals without additional land resources¹⁰⁶.

Implications and outlook

In this Review, we elaborate on the various climate–irrigation feedback loops and identify areas where climate change may tilt the scale by amplifying some positive feedbacks of irrigation via producing more GHG emissions directly or indirectly. It is especially important to understand these feedback effects in regions constrained by freshwater resources, as different adaptation strategies have very different climate implications. In cases where irrigated croplands revert to rain-fed croplands or grazing lands, crop yields will decline, which might result in indirect land-use change and associated carbon loss. A top priority for future research is to quantify both the contribution of agricultural irrigation to global GHG emissions and the feedback effects due to the changing climate at local and global scales. Global estimates that take into consideration the multiple mechanisms reviewed here are currently lacking but could be potentially large.



Fig. 3 | Examples of floating photovoltaics. **a**, A floating photovoltaic system around the University of California at Davis designed to reduce green algae by improving aeration with solar energy. **b**, A floating solar farm producing clean renewable electricity energy and reducing reservoir evaporation in Flevoland, the Netherlands.

Our Review underscores the need to develop an integrated framework around irrigation in future irrigation research and management. An integrated framework can help researchers and planners (1) identify the relative strength and Earth system relevance of various feedbacks; (2) identify climate hotspots, that is, where changes in local or regional climate may necessitate additional irrigation infrastructure and intensify some of the positive climate feedbacks; and (3) prioritize strategies to better harvest the climate benefits of irrigation while minimizing its negative consequences. Such an integrated framework can, for example, help decision-makers invest in irrigation means that are more sustainable, considering the potential feedback loops.

More broadly, greater attention should be paid to climate change in the rapidly growing food–energy–water literature. As climate change intensifies, there is an urgent need to understand not only the effectiveness of different adaptation and mitigation strategies but also how they would feed back to climate change. The integrated nexus thinking and modelling in the food–energy–water literature can be expanded to climate–food–energy–water. This climate-integrated thinking can help us build more climate-resilient food–energy–water systems and better identify opportunities for adaptation and mitigation synergies.

References

1. Wang, X. et al. Global irrigation contribution to wheat and maize yield. *Nat. Commun.* **12**, 1235 (2021).
2. Siebert, S. & Döll, P. Quantifying blue and green virtual water contents in global crop production as well as potential production losses without irrigation. *J. Hydrol.* **384**, 198–217 (2010).
3. Berrang-Ford, L. et al. A systematic global stocktake of evidence on human adaptation to climate change. *Nat. Clim. Change* **11**, 989–1000 (2021).
4. Mayanja, M. N., Rubaire-Akiiki, C., Morton, J. & Kabasa, J. D. Pastoral community coping and adaptation strategies to manage household food insecurity consequent to climatic hazards in the cattle corridor of Uganda. *Clim. Dev.* **12**, 110–119 (2020).
5. Rajan, A., Ghosh, K. & Shah, A. Carbon footprint of India's groundwater irrigation. *Carbon Manag.* **11**, 265–280 (2020).
6. Siyal, A. W., Gerbens-Leenes, P. W. & Nonhebel, S. Energy and carbon footprints for irrigation water in the Lower Indus Basin in Pakistan, comparing water supply by gravity fed canal networks and groundwater pumping. *J. Clean. Prod.* **286**, 125489 (2021).
7. Zou, X. et al. Greenhouse gas emissions from agricultural irrigation in China. *Mitig. Adapt. Strateg. Glob. Change* **20**, 295–315 (2015).
8. Trost, B. et al. Irrigation, soil organic carbon and N_2O emissions. A review. *Agron. Sustain. Dev.* **33**, 733–749 (2013).
9. *Greenhouse Gases from Reservoirs Caused by Biogeochemical Processes* (World Bank, 2017).
10. Mueller, N. D. et al. Cooling of US Midwest summer temperature extremes from cropland intensification. *Nat. Clim. Change* **6**, 317–322 (2016).
11. Burney, J. A., Davis, S. J. & Lobell, D. B. Greenhouse gas mitigation by agricultural intensification. *Proc. Natl Acad. Sci. USA* **107**, 12052–12057 (2010).
12. Hong, C. et al. Global and regional drivers of land-use emissions in 1961–2017. *Nature* **589**, 554–561 (2021).
13. Schewe, J. et al. State-of-the-art global models underestimate impacts from climate extremes. *Nat. Commun.* **10**, 1005 (2019).
14. Diffenbaugh, N. S. Verification of extreme event attribution: using out-of-sample observations to assess changes in probabilities of unprecedented events. *Sci. Adv.* **6**, eaay2368 (2020).
15. D'Odorico, P. et al. The global food–energy–water nexus. *Rev. Geophys.* **56**, 456–531 (2018).
16. Siebert, S. et al. Groundwater use for irrigation – a global inventory. *Hydrol. Earth Syst. Sci.* **14**, 1863–1880 (2010).
17. Li, X., Liu, J., Zheng, C., Han, G. & Hoff, H. Energy for water utilization in China and policy implications for integrated planning. *Int. J. Water Resour. Dev.* **32**, 477–494 (2016).
18. Shumilova, O., Tockner, K., Thieme, M., Koska, A. & Zarfl, C. Global water transfer megaprojects: a potential solution for the water–food–energy nexus? *Front. Environ. Sci.* **6**, 150 (2018).
19. Zhang, B. et al. Methane emissions from global rice fields: magnitude, spatiotemporal patterns, and environmental controls. *Glob. Biogeochem. Cycles* **30**, 1246–1263 (2016).
20. McGill, B. M., Hamilton, S. K., Millar, N. & Robertson, G. P. The greenhouse gas cost of agricultural intensification with groundwater irrigation in a Midwest U.S. row cropping system. *Glob. Change Biol.* **24**, 5948–5960 (2018).
21. Li, Y., Wang, Y.-G., Houghton, R. A. & Tang, L.-S. Hidden carbon sink beneath desert. *Geophys. Res. Lett.* **42**, 5880–5887 (2015).
22. Denev, K., Stewart, C. E., Brenner, J. & Paustian, K. Does long-term center-pivot irrigation increase soil carbon stocks in semi-arid agro-ecosystems? *Geoderma* **145**, 121–129 (2008).
23. Biemans, H. et al. Impact of reservoirs on river discharge and irrigation water supply during the 20th century. *Water Resour. Res.* **47**, W03509 (2011).
24. Zeng, R., Cai, X., Ringler, C. & Zhu, T. Hydropower versus irrigation—an analysis of global patterns. *Environ. Res. Lett.* **12**, 034006 (2017).
25. Linkhorst, A. et al. Comparing methane ebullition variability across space and time in a Brazilian reservoir. *Limnol. Oceanogr.* **65**, 1623–1634 (2020).
26. Aguilera, E. et al. Methane emissions from artificial waterbodies dominate the carbon footprint of irrigation: a study of transitions in the food–energy–water–climate nexus (Spain, 1900–2014). *Environ. Sci. Technol.* **53**, 5091–5101 (2019).
27. Flecker, A. S. et al. Reducing adverse impacts of Amazon hydropower expansion. *Science* **375**, 753–760 (2022).
28. Lesk, C. et al. Compound heat and moisture extreme impacts on global crop yields under climate change. *Nat. Rev. Earth Environ.* **3**, 872–889 (2022).
29. Cook, B. I., Shukla, S. P., Puma, M. J. & Nazarenko, L. S. Irrigation as an historical climate forcing. *Clim. Dyn.* **44**, 1715–1730 (2015).
30. Li, H., Lo, M.-H., Ryu, D., Peel, M. & Zhang, Y. Possible increase of air temperature by irrigation. *Geophys. Res. Lett.* **49**, e2022GL100427 (2022).
31. Ambika, A. K. & Mishra, V. Observational evidence of irrigation influence on vegetation health and land surface temperature in India. *Geophys. Res. Lett.* **46**, 13441–13451 (2019).
32. Bonfils, C. & Lobell, D. Empirical evidence for a recent slowdown in irrigation-induced cooling. *Proc. Natl Acad. Sci. USA* **104**, 13582–13587 (2007).
33. Luan, X. & Vico, G. Canopy temperature and heat stress are increased by compound high air temperature and water stress and reduced by irrigation – a modeling analysis. *Hydrol. Earth Syst. Sci.* **25**, 1411–1423 (2021).
34. Chen, X. & Jeong, S.-J. Irrigation enhances local warming with greater nocturnal warming effects than daytime cooling effects. *Environ. Res. Lett.* **13**, 024005 (2018).
35. te Wierik, S. A., Cammeraat, E. L. H., Gupta, J. & Artzy-Randrup, Y. A. Reviewing the impact of land use and land-use change on moisture recycling and precipitation patterns. *Water Resour. Res.* **57**, e2020WR029234 (2021).
36. Wang-Erlandsson, L. et al. Remote land use impacts on river flows through atmospheric teleconnections. *Hydrol. Earth Syst. Sci.* **22**, 4311–4328 (2018).
37. Kang, S. & Eltahir, E. A. B. North China Plain threatened by deadly heatwaves due to climate change and irrigation. *Nat. Commun.* **9**, 2894 (2018).

38. Krakauer, N. Y., Cook, B. I. & Puma, M. J. Effect of irrigation on humid heat extremes. *Environ. Res. Lett.* **15**, 094010 (2020).

39. Mishra, V. et al. Moist heat stress extremes in India enhanced by irrigation. *Nat. Geosci.* **13**, 722–728 (2020).

40. Ting, M. et al. Contrasting impacts of dry versus humid heat on US corn and soybean yields. *Sci. Rep.* **13**, 710 (2023).

41. Li, Y. et al. Quantifying irrigation cooling benefits to maize yield in the US Midwest. *Glob. Change Biol.* **26**, 3065–3078 (2020).

42. Byerlee, D., Stevenson, J. & Villoria, N. Does intensification slow crop land expansion or encourage deforestation? *Glob. Food Secur.* **3**, 92–98 (2014).

43. Villoria, N. B. Technology spillovers and land use change: empirical evidence from global agriculture. *Am. J. Agric. Econ.* **101**, 870–893 (2019).

44. Hertel, T. W., Ramankutty, N. & Baldos, U. L. C. Global market integration increases likelihood that a future African Green Revolution could increase crop land use and CO₂ emissions. *Proc. Natl Acad. Sci. USA* **111**, 13799–13804 (2014).

45. West, P. C. et al. Trading carbon for food: global comparison of carbon stocks vs. crop yields on agricultural land. *Proc. Natl Acad. Sci. USA* **107**, 19645–19648 (2010).

46. Rosegrant, M. W. & Msangi, S. Consensus and contention in the food-versus-fuel debate. *Annu. Rev. Environ. Resour.* **39**, 271–294 (2014).

47. Lawrence, D., Coe, M., Walker, W., Verchot, L. & Vandecar, K. The unseen effects of deforestation: biophysical effects on climate. *Front. For. Glob. Change* **5**, 756115 (2022).

48. Xu, S. et al. Delayed use of bioenergy crops might threaten climate and food security. *Nature* **609**, 299–306 (2022).

49. Turner, P. A., Field, C. B., Lobell, D. B., Sanchez, D. L. & Mach, K. J. Unprecedented rates of land-use transformation in modelled climate change mitigation pathways. *Nat. Sustain.* **1**, 240–245 (2018).

50. Stenzel, F., Gerten, D. & Hanasaki, N. Global scenarios of irrigation water abstractions for bioenergy production: a systematic review. *Hydrol. Earth Syst. Sci.* **25**, 1711–1726 (2021).

51. Searchinger, T. et al. Use of US croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science* **319**, 1238–1240 (2008).

52. Lesk, C., Coffel, E. & Horton, R. Net benefits to US soy and maize yields from intensifying hourly rainfall. *Nat. Clim. Change* **10**, 819–822 (2020).

53. Christian, J. I. et al. Global distribution, trends, and drivers of flash drought occurrence. *Nat. Commun.* **12**, 6330 (2021).

54. Elliott, J. et al. Constraints and potentials of future irrigation water availability on agricultural production under climate change. *Proc. Natl Acad. Sci. USA* **111**, 3239–3244 (2014).

55. Konzmann, M., Gerten, D. & Heinke, J. Climate impacts on global irrigation requirements under 19 GCMs, simulated with a vegetation and hydrology model. *Hydrol. Sci. J.* **58**, 88–105 (2013).

56. Gray, S. B. et al. Intensifying drought eliminates the expected benefits of elevated carbon dioxide for soybean. *Nat. Plants* **2**, 16132 (2016).

57. Jin, Z., Ainsworth, E. A., Leakey, A. D. B. & Lobell, D. B. Increasing drought and diminishing benefits of elevated carbon dioxide for soybean yields across the US Midwest. *Glob. Change Biol.* **24**, e522–e533 (2018).

58. Ainsworth, E. A. & Long, S. P. 30 years of free-air carbon dioxide enrichment (FACE): what have we learned about future crop productivity and its potential for adaptation? *Glob. Change Biol.* **27**, 27–49 (2021).

59. Wang, S. et al. Recent global decline of CO₂ fertilization effects on vegetation photosynthesis. *Science* **370**, 1295–1300 (2020).

60. Berg, A., Sheffield, J. & Milly, P. C. D. Divergent surface and total soil moisture projections under global warming. *Geophys. Res. Lett.* **44**, 236–244 (2017).

61. Ficklin, D. L. & Novick, K. A. Historic and projected changes in vapor pressure deficit suggest a continental-scale drying of the United States atmosphere. *J. Geophys. Res. Atmos.* **122**, 2061–2079 (2017).

62. Qin, Y. et al. Agricultural risks from changing snowmelt. *Nat. Clim. Change* **10**, 459–465 (2020).

63. Taylor, R. G. et al. Ground water and climate change. *Nat. Clim. Change* **3**, 322–329 (2013).

64. Kundzewicz, Z. W. et al. The implications of projected climate change for freshwater resources and their management. *Hydrol. Sci. J.* **53**, 3–10 (2008).

65. Stokstad, E. Deep deficit. *Science* **368**, 230–233 (2020).

66. Scott, C. A. Electricity for groundwater use: constraints and opportunities for adaptive response to climate change. *Environ. Res. Lett.* **8**, 035005 (2013).

67. Qiu, G. Y., Zhang, X., Yu, X. & Zou, Z. The increasing effects in energy and GHG emission caused by groundwater level declines in North China's main food production plain. *Agric. Water Manag.* **203**, 138–150 (2018).

68. Kaur, S., Aggarwal, R. & Lal, R. Assessment and mitigation of greenhouse gas emissions from groundwater irrigation. *Irrig. Drain.* **65**, 762–770 (2016).

69. Wood, W. W. & Hyndman, D. W. Groundwater depletion: a significant unreported source of atmospheric carbon dioxide. *Earth's Future* **5**, 1133–1135 (2017).

70. Minamikawa, K. et al. Groundwater-induced emissions of nitrous oxide through the soil surface and from subsurface drainage in an Andosol upland field: a monolith lysimeter study. *Soil Sci. Plant Nutr.* **59**, 87–95 (2013).

71. Schipanski, M. E. et al. Moving from measurement to governance of shared groundwater resources. *Nat. Water* **1**, 30–36 (2023).

72. Gleeson, T., Wada, Y., Bierkens, M. F. P. & van Beek, L. P. H. Water balance of global aquifers revealed by groundwater footprint. *Nature* **488**, 197–200 (2012).

73. Gleick, P. *Impacts of California's Five-Year (2012–2016) Drought on Hydroelectricity Generation* (Pacific Institute, 2017).

74. *Renewable Energy in the Spanish Electricity System 2017* (REE, 2017).

75. Herrera-Estrada, J. E., Diffenbaugh, N. S., Wagner, F., Craft, A. & Sheffield, J. Response of electricity sector air pollution emissions to drought conditions in the western United States. *Environ. Res. Lett.* **13**, 124032 (2018).

76. *Adaptation Challenges and Opportunities for the European Energy System: Building a Climate Resilient Low Carbon Energy System* (European Environmental Agency, 2019).

77. Griffis, T. J. et al. Nitrous oxide emissions are enhanced in a warmer and wetter world. *Proc. Natl Acad. Sci. USA* **114**, 12081–12085 (2017).

78. van Groenigen, K. J., van Kessel, C. & Hungate, B. A. Increased greenhouse-gas intensity of rice production under future atmospheric conditions. *Nat. Clim. Change* **3**, 288–291 (2013).

79. Qian, H. et al. Lower-than-expected CH₄ emissions from rice paddies with rising CO₂ concentrations. *Glob. Change Biol.* **26**, 2368–2376 (2020).

80. Rosa, L. et al. Potential for sustainable irrigation expansion in a 3 °C warmer climate. *Proc. Natl Acad. Sci. USA* **117**, 29526–29534 (2020).

81. Padedda, B. M. et al. Consequences of eutrophication in the management of water resources in Mediterranean reservoirs: a case study of Lake Cedrino (Sardinia, Italy). *Glob. Ecol. Conserv.* **12**, 21–35 (2017).

82. Yan, X. et al. Climate warming and cyanobacteria blooms: looks at their relationships from a new perspective. *Wat. Res.* **125**, 449–457 (2017).

83. Paerl, H. W. & Huisman, J. Blooms like it hot. *Science* **320**, 57–58 (2008).

84. Jansen, J. et al. Global increase in methane production under future warming of lake bottom waters. *Glob. Change Biol.* **28**, 5427–5440 (2022).

85. Carrijo, D. R., Lundy, M. E. & Linquist, B. A. Rice yields and water use under alternate wetting and drying irrigation: a meta-analysis. *Field Crops Res.* **203**, 173–180 (2017).

86. Kritee, K. et al. High nitrous oxide fluxes from rice indicate the need to manage water for both long- and short-term climate impacts. *Proc. Natl Acad. Sci. USA* **115**, 9720–9725 (2018).

87. Lehmann, J. et al. Biochar in climate change mitigation. *Nat. Geosci.* **14**, 883–892 (2021).

88. Kuang, W., Gao, X., Tenuta, M. & Zeng, F. A global meta-analysis of nitrous oxide emission from drip-irrigated cropping system. *Glob. Change Biol.* **27**, 3244–3256 (2021).

89. Almeida, R. M. et al. Floating solar power could help fight climate change — let's get it right. *Nature* **606**, 246–249 (2022).

90. Exley, G., Armstrong, A., Page, T. & Jones, I. D. Floating photovoltaics could mitigate climate change impacts on water body temperature and stratification. *Solar Energy* **219**, 24–33 (2021).

91. Rathore, P. K. S., Das, S. S. & Chauhan, D. S. Perspectives of solar photovoltaic water pumping for irrigation in India. *Energy Strategy Rev.* **22**, 385–395 (2018).

92. Hoffacker, M. K. & Hernandez, R. R. Local energy: spatial proximity of energy providers to their power resources. *Front. Sustain.* **1**, 585110 (2020).

93. Xie, H., Ringler, C. & Mondal, M. D. A. H. Solar or diesel: a comparison of costs for groundwater-fed irrigation in sub-Saharan Africa under two energy solutions. *Earth's Future* **9**, e2020EF001611 (2021).

94. Lefore, N., Closas, A. & Schmitter, P. Solar for all: a framework to deliver inclusive and environmentally sustainable solar irrigation for smallholder agriculture. *Energy Policy* **154**, 112313 (2021).

95. Shim, H.-S. *Case Study: Solar-Powered Irrigation Pumps in India — Capital Subsidy Policies and the Water–Energy Efficiency Nexus* (Global Green Growth Institute, 2017).

96. Hartung, H. & Pluschke, L. *The Benefits and Risks of Solar Powered Irrigation - A Global Overview* (Food and Agriculture Organization of the United Nations, 2018).

97. Lesk, C. & Kornhuber, K. An effective clean energy transition must anticipate growing climate disruptions. *Environ. Res. Clim.* **1**, 013002 (2022).

98. Hernandez, R. R. et al. Techno–ecological synergies of solar energy for global sustainability. *Nat. Sustain.* **2**, 560–568 (2019).

99. Ravi, S. et al. Colocation opportunities for large solar infrastructures and agriculture in drylands. *Appl. Energy* **165**, 383–392 (2016).

100. Barron-Gafford, G. A. et al. Agrivoltaics provide mutual benefits across the food–energy–water nexus in drylands. *Nat. Sustain.* **2**, 848–855 (2019).

101. Hoffacker, M. K., Allen, M. F. & Hernandez, R. R. Land-sparing opportunities for solar energy development in agricultural landscapes: a case study of the Great Central Valley, CA, United States. *Environ. Sci. Technol.* **51**, 14472–14482 (2017).

102. Yang, Y. et al. Restoring abandoned farmland to mitigate climate change on a full Earth. *One Earth* **3**, 176–186 (2020).

103. Grodsky, S. M. Matching renewable energy and conservation targets for a sustainable future. *One Earth* **4**, 924–926 (2021).

104. Walston, L. J. et al. Examining the potential for agricultural benefits from pollinator habitat at solar facilities in the United States. *Environ. Sci. Technol.* **52**, 7566–7576 (2018).

105. Roberts, B. *Potential for Photovoltaic Solar Installation in Non-Irrigated Corners of Center Pivot Irrigation Fields in the State of Colorado* (National Renewable Energy Laboratory, 2011).

106. Suraci, J. P. et al. Achieving conservation targets by jointly addressing climate change and biodiversity loss. *Ecosphere* **14**, e4490 (2023).

Author contributions

Y.Y., Z.J. and N.D.M. conceived and designed the experiments, analysed the data and wrote the paper. A.W.D., R.R.H., S.M.G., L.L.S., M.V.C., Y.-G.Z. and D.B.L. contributed materials and wrote the paper.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Zhenong Jin or Nathaniel D. Mueller.

Peer review information *Nature Food* thanks Corey Lesk and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2023