Firefox about:blank

Nuclear Inst. and Methods in Physics Research, A 1048 (2023) 167949



Contents lists available at ScienceDirect


Nuclear Inst. and Methods in Physics Research, A

journal homepage: www.elsevier.com/locate/nima



Performance of BDX-MINI veto systems

M. Spreafico a,b,*, M. Battaglieri a,c, M. Bondí d, A. Celentano a, P.L. Cole e, M. De Napoli d, R. De Vita a, L. Marsicano a, N. Randazzo d, E.S. Smith c, D. Snowden-Ifft f, M. Wood g



b Universitá degli Studi di Genova, 16126 Genova, Italy

ARTICLE INFO

Keywords: Light dark matter Beam dump experiments

ABSTRACT

This paper describes the veto system of the BDX-MINI detector installed at Jefferson Lab (US). The MINI experiment is the first electron beam-dump experiment specifically designed to search for Light Matter (LDM) particles in the MeV-GeV mass range. The core of the BDX-MINI detector is a lead-tung electromagnetic calorimeter, for a total volume of 4 dm³. The calorimeter is surrounded by a multi-layer aimed at rejecting cosmic background: the innermost layer of the veto is made by a passive tungsten shie for low energy radiation, while plastic scintillators make the middle and outer layers for charged cc particles rejection. Being located about 20 m downstream, the dirt between the beam dump and the det provides sufficient shielding from the beam-related background. In 2019–2020, BDX-MINI was expose about six months to weakly interacting particles (neutrinos and, if existing, DM) produced by a 2.176 electron beam incident on the beam dump of experimental Hall-A at Jefferson Lab.

1. Introduction

Light Dark Matter (LDM) is a new compelling hypothesis that identifies Dark Matter with new sub-GeV "Hidden Sector" states. In the most popular model, LDM particles χ interaction with SM is mediated by a massive vector boson called A' ("Dark Photon"). The Dark Photon can kinematically mix with the Standard Model hypercharge field, resulting in SM-DM interactions.

Many experimental techniques can be used to probe Light Dark Matter [1,2]. Among those, Beam Dump Experiments offers unique possibilities in probing different LDM models.

In Beam Dump Experiments DM is produced by an intense beam impinging on a thick target (beam dump). DM can be produced in the beam dump beside the great number of Standard Model (SM) particles. A sizable shielding located downstream the dump absorbs all SM particles (except neutrinos), filtering a secondary DM beam. LDM particles are detected in a downstream electromagnetic calorimeter through χ scattering with detector electrons.

BDX-MINI is a Beam Dump eXperiment searching for DM particles in the MeV-GeV mass range produced by the interaction of Jefferson Lab (JLab) high intensity electron beam impinging on experimental Hall-A beam dump. It used a $2.176 \text{ GeV } e^-$ -beam, with current up to $150 \mu\text{A}$.

The detector was located in a well $26\ m$ downstream the dun beamline height. The entire setup was located inside a sturdy field

BDX-MINI ran for 6 months in 2019–2020, accumulating a charge of about 2.56×10^{21} EOT.

2. BDX-MINI detector

BDX-Mini detector is composed of an electromagnetic calorin (ECal) surrounded by a multi-layer veto system.

BDX-Mini ECal is composed of 44 parallelepiped PbWO $_4$ crysta a total active volume of $\sim 4 \times 10^{-3}$ m 3 (Fig. 1.A). The crystals sc lation light is read by 6×6 mm 2 Hamamatsu MPPCs (S13360-602 yielding 1 p.e./MeV deposited.

The hermetic veto is composed of three layers. The innermost is a passive tungsten shielding 0.8 cm thick (Fig. 1.B and C). middle (Fig. 1.D) and outer (Fig. 1.E) layer of the veto are made pl scintillator, with an octagonal and cylinder shape, respectively, 0.8 cm thick. The Octagonal Inner Veto is composed by 8 scintil paddles coupled with optical glue; the top and bottom are covere two octagonal scintillator caps. The Cylindrical Outer Veto is comp of a single cylindrical scintillator tube; two round scintillator caps (

https://doi.org/10.1016/j.nima.2022.167949

Received 14 July 2022; Accepted 29 November 2022

1 of 4 9/2/23, 3:02 PM

^c Thomas Jefferson National Accelerator Facility, 23606, Newport News, VA, USA

d Istituto Nazionale di Fisica Nucleare. Sezione di Catania. 95125 Catania. Italy

e Lamar University, 4400 MLK Boulevard, P.O. Box 10046, Beaumont, TX 77710, USA

f Occidental College, Los Angeles, CA 90041, USA

g Canisius College, Buffalo, NY, 14208, USA

^{*} Corresponding author at: Universitá degli Studi di Genova, 16126 Genova, Italy. E-mail address: marco.spreafico@ge.infn.it (M. Spreafico).

Firefox about:blank

Available online 19 December 2022 0168-9002/© 2022 Elsevier B.V. All rights reserved.

M. Spreafico, M. Battaglieri, M. Bondí et al.

Nuclear Inst. and Methods in Physics Research, A 1048 (2023) 1




Fig. 1. The different components of BDX-Mini detector are shown. A. The two modules that constitute the ECal. B and C. Tungsten Shielding D. Inner Veto. E. Outer Veto.




Fig. 2. Response of the OV-Cylinder (upper panel) and IV lower cap (bottom panel) as a function of time. The red line represents the average efficiency. The response OV-Cylinder was measured selecting muons leaving a signal in both IV-O, OV-C and two adjacent caps. The response of IV lower cap was evaluated selecting muons leaving in all caps.

the top and bottom. The scintillation light is collected by Wavelength Shifter Fibers inserted in grooves on the scintillator and read by 3×3 mm² Hamamatsu S13360-3075CS SiPMs. For each veto there are 10 readout channels, 8 for the lateral surface and 2 for the bases.

The front-end electronics and data acquisition system was installed outside the well at ground level. Each SiPM was connected via an 8-m long coaxial cable to a custom front-end circuit that provided the bias voltage and amplified the signal from photosensors. For each event, all channel were digitalized by a 250 MHz fADC and the corresponding waveform was saved on disk.

correct any possible response variation over time. The performan each veto system was studied using cosmic-ray data collected die the accelerator down period. In Ref. [3], the analysis of bear data accumulated in over 30 days showed that requiring the coincidence with the veto yields a very high suppression of the cobackground.

To check the long-term stability of each component, the responent of the veto to well defined trajectories produced by cosmic m was studied as a function of time. For each cap, vertical muons selected by requiring a significant deposited energy (corresponding)

2 of 4 9/2/23, 3:02 PM

Firefox about:blank

> For a more detailed review of BDX-MINI experimental setup see Ref. [3]

3. Detector performance

Since BDX-MINI ran inside a field tent for several months, it was of paramount importance to monitor its performance over all measure-

M. Spreafico, M. Battaglieri, M. Bondí et al.

The authors would like to thank Dipartimento di Fisica (Unive

>5 p.e.) in other caps than the one under study and no activi

the other veto channels. The event selection to study the Inner Octagon (Outer Veto Cylinder) performance required hits (at least

SiPM with a signal >5 p.e.) in the cylinder veto (octagon veto) either the two upper or lower caps. Fig. 2 shows the respons

function of time (each point represents a data-taking run) of the

Cylinder (unner nanel) and Rottom IV can (hottom nanel). The re

Nuclear Inst. and Methods in Physics Research, A 1048 (2023) 1

Acknowledgments

degli Studi di Genova) for funding this talk. Special thanks to CL Collaboration and PANDA Collaboration for providing the Pb

References

- [1] M.e.a. Battaglieri, US cosmic visions: New ideas in dark matter 2017: Comn report, 2017, http://dx.doi.org/10.48550/ARXIV.1707.04591, URL https:// org/abs/1707.04591.
- [2] A. Filippi, M. De Napoli, Searching in the dark: the hunt for the photon, Rev. Phys. (ISSN: 2405-4283) 5 (2020) 100042, http://dx.doi.or 1016/j.revip.2020.100042, URL https://www.sciencedirect.com/science/a pii/S2405428320300058.
- [3] M.B. et al., The BDX-MINI detector for light dark matter search at JLab Phys. J. C 81 (2) (2021) http://dx.doi.org/10.1140/epjc/s10052-021-0895

4. Conclusions

BDX-Mini was the first modern electron beam dump experiment aimed specifically at Light Dark Matter search. Its detector was optimized for DM searches at accelerators, having the ECal surrounded by a highly hermetic multi layer veto. The veto demonstrated to have a very high rejection capability. In this paper, we report the performance of each veto system component investigated using cosmic ray data. This study demonstrated that the response of the veto system was extremely stable during the whole data taking period, although the experiment was performed inside a sturdy field tent.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

9/2/23, 3:02 PM

Firefox about:blank

3

4 of 4