

How does building healthy soils impact sustainable use of water resources in irrigated agriculture?

Sara E. Acevedo¹, Hannah Waterhouse², Felipe Barrios-Masias³, Janina Dierks², Leah L.R. Renwick⁴, Timothy M. Bowles^{2*}

¹ Departamento de Ingeniería Hidráulica y Ambiental, Facultad de Ingeniería, Pontificia Universidad Católica de Chile, Macul, Santiago, Chile

² Department of Environmental Science, Policy, and Management, UC Berkeley, Berkeley, CA 94720, USA

³ Department of Agriculture, Veterinary and Rangeland Sciences, University of Nevada, Reno, NV 89557, USA

⁴ Departamento de Ingeniería y Suelos, Facultad de Ciencias Agronómicas, Universidad de Chile, La Pintana, Santiago, Chile

*timothy.bowles@berkeley.edu

Abstract

As blue water resources become increasingly scarce with more frequent droughts and overuse, irrigated agriculture faces significant challenges to reduce its water footprint while maintaining high levels of crop production. Building soil health has been touted as an important means of enhancing the resilience of agroecosystems to drought, mainly with a focus in rainfed systems reliant on green water through increases in infiltration and soil water storage. Yet, green water often contributes only a small fraction of the total crop water budget in irrigated agricultural regions. To scope the potential for how soil health management could impact water resources in irrigated systems, we review how soil health affects soil water flows, plant-soil-microbe interactions, and plant water capture and productive use. We assess how these effects could interact with irrigation management to help make green and blue water use more sustainable. We show how soil health management could: 1) optimize green water availability (e.g., by increasing infiltration and soil water storage), 2) maximize productive water flows (e.g., by reducing evaporation and supporting crop growth), and 3) reduce blue water withdrawals (e.g., by minimizing impacts of water stress on crop productivity). Quantifying the potential of soil health to improve water resource management will require research that focuses on outcomes for green and blue water provisioning and crop production under

different irrigation and crop management strategies. Such information could be used to improve and parameterize finer scale crop, soil, and hydraulic models, which in turn must be linked with larger scale hydrologic models to address critical water-resources management questions at watershed or regional scales. While integrated soil health-water management strategies have considerable potential to conserve water — especially compared to irrigation technologies that enhance field-level water use efficiency but often increase regional water use — transitions to these strategies will depend on more than technical understanding and must include addressing interrelated structural and institutional barriers. By scoping a range of ways enhancing soil health could improve resilience to water limitations and identifying key research directions, we inform research and policy priorities aimed at adapting irrigated agriculture to an increasingly challenging future.

Introduction

Irrigated agriculture faces the dual challenge of adapting to a hotter, more drought-prone climate with higher evaporative demand while also reducing its water footprint. Rising temperatures and increases in heat waves and the intensity and frequency of droughts due to climate change (Hatfield et al., 2011; Trenberth et al., 2014) will likely worsen water limitations on crop productivity (Quinteiro et al., 2018). Multi-year droughts reduce water available for irrigation, cause agricultural sector job losses, reduce agricultural output (Howitt et al., 2015), and increase groundwater consumption, accelerating groundwater overdraft and land subsidence (Faunt et al., 2016). Climate change-driven reductions in the snowpack that often provides irrigation water and recharges groundwater exacerbate these overdrafts worldwide, particularly important in the Western U.S. (López-Moreno et al., 2017; Pathak et al., 2018)

Current withdrawals of blue water—water in streams, rivers, lakes, reservoirs, and groundwater aquifers—for agriculture already exceed planetary environmental boundaries (Sposito, 2013; Steffen et al., 2015). World-wide, irrigated agriculture consumes about 70 - 90% of blue water used by humans (Ward and Pulido-Velazquez, 2008), which impacts non-agricultural ecosystems (Rost et al., 2008) and competes with growing demand from other human uses like drinking water and industry. Groundwater provides 43% of total consumptive irrigation water used globally, and reliance on groundwater for irrigation is increasing (Siebert et al., 2010), despite 6-20% of global groundwater wells already at risk of running dry (Jasechko and Perrone, 2021). Irrigated areas constitute 20% of the total area and 40% of total agricultural production; their continuing expansion in area dominates global crop blue water consumption, which is expected to increase 70% by 2071–2099 compared to 1971–2000 (Huang et al., 2019). Higher blue water consumption

coupled with lower blue water availability due to climate change will lead to ~60% of the world population likely experiencing blue water shortages by 2050 (Rockström et al., 2009). Increasing blue water withdrawals to intensify or expand irrigated agriculture thus poses major risks for humans and the nonhuman biosphere (Sposito, 2013).

Alternative sources of irrigation water, such as recycled water (i.e. greywater) or desalinated water, present modest opportunities to reduce blue water use (Assouline et al., 2015). While recycled water from both municipal and industrial origins irrigates approximately 20M ha globally (~5% of global irrigated land; Jimenez and Asano 2008; Meier et al. 2018), strategies to deal with tradeoffs, such as salinization in arid and semi-arid regions, are needed to make these alternative irrigation water sources more widely viable (Assouline et al 2015; Hamilton et al 2007).

Green water may provide underappreciated opportunities to alleviate pressure on blue water withdrawals in irrigated systems. Green water is rainfall that infiltrates in the upper unsaturated soil layers and flows back to the atmosphere through evapotranspiration, i.e. it is the water retained in soil after deep percolation and runoff (Falkenmark and Rockström, 2010). Green water accounts for 90% of cropland water consumption globally, including both rainfed and irrigated agriculture (Rost et al., 2008; Sposito, 2013). Even in areas where irrigated agriculture relies largely on blue water, such as Mediterranean regions like California, USA, green water still accounts for 6-35% of total crop evapotranspiration (ET) (Fig. 1; Chiarelli et al., 2020; Devine and O'Geen, 2019). Green water has not received much attention from water resources management and planning agencies in irrigated regions in part due to the challenge of quantifying its consumption on a landscape scale, and the difficulty in differentiating between green and blue water flows (Jewitt, 2006). Meeting the global challenge of adapting irrigated agriculture to water limitations will involve: 1) reducing the consumption of blue water for irrigation to achieve sustainable levels of withdrawal; 2) optimizing green water availability to crops; and 3) maximizing productive consumption of water from all sources, that is, water that flows as transpiration through plants rather than soil evaporation, all while maintaining crop production.

Soil characteristics are a major mediator of both green and blue water flows, alongside factors like topography and land use (Quinteiro et al., 2018). Inherent soil characteristics like texture strongly govern physical properties important for soil hydrologic functioning, such as saturated hydraulic conductivity, water storage capacity, and soil structural stability (Bünemann et al., 2018), thus impacting not only the capture and storage of green water, but also green and blue water flows such as runoff and percolation.

Dynamic soil properties like soil structure, that is, those influenced by management on relatively short seasonal to decadal time scales, also affect water flows (Schwen et al., 2011).

With growing multi-sector interest, soil health is a concept focused on managing dynamic soil properties like soil organic matter (SOM), soil structure, and soil organisms to support soil ecosystem functioning and provide multiple ecosystem services beyond just the provisioning of food, fiber and fuel, including water provisioning and effective use of water resources (de Groot et al., 2010). Managing soil health is based on several well-defined *principles* — reducing soil disturbance, keeping the soil covered, increasing plant diversity, and keeping living roots in the ground — that manifest as *practices* such as reduced/no till, cover cropping, organic matter amendments, crop rotation diversification, and strategic integration of perennial plants into key landscape positions. These practices change the dynamic biological, chemical, and physical properties that underlie healthy soils, which affect water provisioning and effective use in multiple ways.

There is a widespread perception among stakeholders such as farmers, government and academic scientists, and policy advocates that building healthier soils could play a large role in adapting to reduced and volatile water availability, due to its potential for increasing water provisioning (Cano et al., 2018; Wade et al., 2021). Based on studies predominantly in rainfed agriculture, proponents most often argue that soil health-promoting management that increases SOM and improves soil structure will increase soil water holding capacity and infiltration (Gaudin et al., 2015; Lotter et al., 2003; Williams et al., 2016), leading to higher capture and storage of green water available for plants, while also reducing harmful runoff. For instance, an informational campaign from the U.S. Natural Resources Conservation Service asserted in an infographic that “For each 1% increase in organic matter, U.S. cropland could store the amount of water that flows over Niagara Falls in 150 days” (Nichols, 2015). In irrigated agriculture, California’s Fourth Climate Assessment notes benefits of increasing SOM on hydrological flows in croplands, including capturing and storing more water and thus mitigating impacts of water deficits (Flint et al., 2018).

While infiltration and water storage capacity are crucial, understanding the full potential for healthy soils to promote effective use of water resources requires going beyond just water storage and capture, especially in irrigated agriculture. Since green water accounts for a relatively small proportion of crop water use in irrigated Mediterranean systems (Devine and O’Geen, 2019), it is unclear how important increasing green water availability is relative to other mechanisms by which soil health could impact water provisioning and sustainable water use. How soil health practices impact the soil water balance in such systems must also be considered, for instance, whether evapotranspiration from cover cropping depletes green water for subsequent crops (DeVincentis et al. 2022). With climate change expected to bring even starker contrasts

between wet and dry seasons in Mediterranean regions (Nunes et al., 2019), and urbanization and population growth increasing competition for blue water supplies (Nouri et al., 2019) scoping the role of soil health management for water conservation is essential for policy making.

Here we assess how healthy soils could affect both green and blue water availability and its productive use for crop production, focusing in particular on the complex case of irrigated cropping systems. We first review how healthy soils affect hydrologic processes and soil hydraulic properties, plant-soil-microbe interactions, and plant water demand and uptake in annual and perennial croplands. We include studies conducted in both irrigated and rainfed systems to draw from a wide range of research. We then focus on irrigated systems specifically by proposing strategies for integrated soil health and water management to conserve water resources, as well as tradeoffs and social-ecological challenges for making this transition. We highlight examples and lessons from California, where we have experience, with applicability to other irrigated regions. Finally, to assess our current potential for quantitative scenario analysis and planning, we also consider key knowledge gaps and priority areas for future research. In particular, we focus on the extent to which predominant agro-hydrological models adequately capture both impacts of management on water-related soil properties and processes, and if these changes in turn impact water resources across scales. By doing so, we aim to inform discussions, improve current knowledge, and suggest ways forward for how to meet the global agricultural water challenge, particularly in ways that could lead to the well-characterized co-benefits of soil health management.

Healthy soils and water provisioning

Soil water flows and storage

Ideally, a healthy soil increases the capture, retention, and provisioning of both green and blue water and thereby enhances crop growth (Fig. 2). The partitioning of green and blue water is governed by the balance between runoff versus infiltration, percolation, evaporation, and soil water storage (Hoekstra, 2019). Infiltration rates dictate how fast water penetrates the soil surface and thus plays a key role in determining the amount of green water stored in the soil after rains and made available to plants and how much blue water can be applied in an irrigation before runoff occurs. Deep infiltration or percolation contributes to blue water reservoirs by recharging underlying groundwater (O’Geen et al., 2015). Infiltration is controlled by soil pore size distribution and geometry (Pahlavan-Rad et al., 2020), and soil structure (the way in which soil particles are grouped together into aggregates bound together by physical, chemical and biological processes which dictate the pore space of the soil system) (Fischer et al., 2015).

Several soil health practices—including cover cropping, reducing tillage, organic amendments, crop residue retention, and cropping system diversification—affect one or more soil physical properties related to infiltration (Bagnall et al., 2022). For example, continuous living cover through perennial herbaceous and woody crops and cover crops consistently and strongly promotes higher infiltration rates compared to bare fallow (Alvarez et al., 2017; Basche and DeLonge, 2019). Plant roots strongly influence the interrelated biological, chemical and physical processes that increase porosity and aggregation (Meurer et al., 2020). For instance, based on a global meta-analysis, perennials increased infiltration rates by 59.2% on average over annuals, with cover crops having the second largest effect size at 34.8% over bare fallow (Basche and DeLonge, 2019). Reducing soil disturbance through no-till may increase or decrease infiltration; a global quantitative synthesis reported no net effect with a wide range of positive and negative outcomes (Basche and DeLonge, 2019). While more consistently positive effects would be expected with the improvement and preservation of soil structure with reduced/no-till, negative outcomes may be due to comparisons being made soon after tillage. In the short term, tillage can temporarily increase infiltration by promoting macropores and reducing soil bulk density (Alegre et al., 1991; Martínez et al., 2008), especially compared to no-till systems that have not been in place for long enough to re-establish highly interconnected pores (Martínez et al., 2008; Strudley et al., 2008). The interaction of soil health practices differentially affects infiltration rates. For example, a meta-analysis found that while no-till in combination with residue retention improves infiltration, tillage in combination with cover cropping does not affect infiltration (Basche and DeLonge, 2019). Organic matter amendments, such as crop residues, straw, biochar, and manure (from integrated crop-livestock systems), often improve soil structure and infiltration (Dong et al., 2022) but effects vary with the sources and degree of decomposition of its components as well as factors like the rate and duration of application and amended soil types (Kranz et al., 2020, Dong et al., 2022). Integrated crop-livestock systems can affect soil structure both positively and negatively (Franzluebbers 2007, Peterson et al., 2019). High stocking rates and/or poorly timed grazing compacts soil and decreases infiltration but careful grazing management, especially in perennial pastures, can increase SOM and promote continuous biopores (Franzluebbers 2007, Peterson et al., 2019). Diversified crop rotations can improve soil structure and infiltration by including crops with deep and shallow roots and by increasing soil cover and root presence in the topsoil (Alhameid et al., 2020).

At the field scale, leveraging soil health management practices to increase infiltration could increase green water storage in the soil during rain events that could be used to delay irrigation during the spring and reduce the amount of runoff from flood irrigation events during the growing season. At the landscape scale, such practices could improve deep percolation and increase blue water in Mediterranean regions via

groundwater recharge during winter rains. Percolation, the movement of water through soil by gravity and capillary forces, is an important soil property for conveying water beyond the influence of evapotranspiration and eventually recharging groundwater. Percolation occurs mainly through interconnected macropores; thus, the texture and structure of a soil, which dictates a soil's pore size distribution and pore connectivity, will strongly control the rate of percolation. Percolation is also influenced by water storage since water potential (the sum of gravitational, osmotic, and matric potentials), and thus water flow, is a function of soil water content (Richards, 1931). Percolation proceeds after soil has reached field capacity and moves toward saturated conditions. Soil health practices that influence SOM, soil structure, and porosity will affect percolation (Basche and DeLonge, 2019; Dahlke et al., 2018). For instance, with higher SOM, clay-organic matter interactions reduce dispersibility of clays and help improve and preserve soil structure, increasing the resilience of aggregates to disturbance, such as runoff, and improving infiltration (and thus, eventually, percolation). Improved soil structure also creates a “sponge-like” pore system, where pore-size is more evenly distributed across micro to meso to macro pore sizes (de Jonge et al., 2009). This type of pore system can improve the soil's capacity to retain water, as well as percolate water once field capacity is reached. The “sponge-like” hydrological behavior depends on SOM and clay content, and changes in soil management could cause changes in the hydrological services provided (Mosquera et al., 2020). Degraded soils—with poor structure, high macroporosity and “downpipe-like” pore systems—may contribute to groundwater recharge, and thus blue water reservoirs, via preferential flow. But such soils can also have a large proportion of “dead-end” pore spaces within the broader soil matrix that reduce the capacity of a soil to store green water for plant uptake and can accelerate the movement of contaminants toward aquifers (de Jonge et al., 2009). In certain contexts, soil health practices could also reduce percolation. A meta-analysis found that cover-cropping reduced percolation by 27 mm on average in temperate systems compared to bare-fallow plots (Meyer et al., 2019). Overall, however, few studies specifically measure soil health management impacts on percolation, highlighting a knowledge gap concerning the link between soil health and blue water resources.

Since soil evaporation is part of non-productive vertical fluxes (sometimes referred to as “white water”; Jewitt, 2006) reducing soil evaporation relative to crop transpiration promotes productive water flow, and increases the percentage of water recharged via percolation when transpiration is low. Plant biomass residues left on the soil surface reduce soil evaporation and increase soil water content and plant available water in drought-prone areas (Turmel et al., 2015). Also, residues can be an effective weed control method (Nichols et al., 2015). However, when precipitation is light and infrequent, surface residues can intercept precipitation and increase subsequent evaporation (Turmel et al., 2015). No-till systems in semi-arid areas

sometimes reduce evaporation (Jokela and Nair, 2016; Schwartz et al., 2010; Soane et al., 2012; Strudley et al., 2008), but the magnitude and direction of the effect depends on tillage type, when tillage occurred, the timing of precipitation following tillage, and the length of the monitoring period of the study (Schwartz et al., 2010; Strudley et al., 2008; Sun et al., 2018). Decreased evaporation under no-till has been attributed to surface residue retention (Sun et al., 2018); however, other physical changes, such as surface roughness and soil porosity, may be partly responsible (O'Brien and Daigh, 2019). Cover crops similarly reduce evaporation (i.e., via increased surface residues, changes to surface roughness, and soil porosity). But most studies on cover crops and their effects on the soil water balance have been conducted in humid regions (Daryanto et al., 2018). In semi-arid regions, where evapotranspiration can be a large part of the water balance, more research is needed on how cover cropping and other soil health practices affect productive versus nonproductive water flows.

Increases in SOM and subsequent improvements to soil aggregation, pore space distribution, and soil water retention have been a focal point of how soil management can influence soil water storage (Rawls et al., 2003). Yet, recently the axiom “more SOM equals more plant available water” has been questioned (Minasny and McBratney, 2018). In general, positive relationships between SOM and plant available water are strongest in soils with lower SOM and coarse textures (Gupta et al., 2020; Hudson, 1984; Rawls et al., 2003; Villamil et al., 2006). Using a large dataset, Minasny and McBratney (2018) reported that the relationship between SOM and plant available water is overall weak, mainly observed near field capacity in sandy soils and negligible in finer soil textures. Similar results have been found in soils from Mediterranean Europe, where higher SOM did not raise plant available water (Bonfante et al., 2020). Higher SOM not translating into higher plant available water could be due to simultaneous increases in water retained at field capacity and permanent wilting point (Renwick et al., 2021).

Soil texture plays a mediating role in the magnitude and direction of soil health practices, namely no-till, continuous living cover, and cover cropping, and their effect on plant available water. Effects of no-till on plant available water were negative in clayey soils, negligible in fine-textured soils with high SOM, and marginally positive in silt loam soils (Kinoshita et al., 2017). Continuous living cover (cover crops, perennial grasses, agroforestry, and managed forestry) showed similarly mixed outcomes on plant available water as with no-till, where no significant effect on plant available water was found for soils containing >25% silt or >30% clay (Basche and DeLonge, 2017). However, for lighter textured soils, a 9.3% increase in water retained at field capacity was found when continuous cover was implemented (Basche and DeLonge, 2017).

Soil health management variably affects soil hydrologic functioning, and more local research is necessary to create regional and site-specific soil health recommendations to improve water provisioning (Devine and O’Geen, 2019). New methods are needed to quantify the partitioning between productive (transpiration) and non-productive (evaporation) flow, such as assessing the components of the surface energy balance and how varying soil health management practices partition water fluxes into evaporation vs. transpiration (O’Brien and Daigh, 2019). Studies are also needed on the potential trade-offs of cover cropping and its effect on the water balance in semi-arid regions (Mitchell et al., 2015). For example, some studies have found that both percolation and water storage were reduced under cover cropping (Gabriel et al., 2012; Meyer et al., 2019; Mitchell et al., 2015; Ward et al., 2012) while others show no or little effect (DeVincentis et al., 2022). Several factors will influence outcomes for the water balance, such as the termination timing of the cover crop prior to the cash crop growing season (Alonso-Ayuso et al., 2018), the type of cover crop (or mixture of cover crops) (Nielsen and Vigil, 2005), the balance between cover crop transpiration vs. improvements to soil physical properties that increase water capture and then provisioning during the cash crop season, and weather (Jones et al., 2020; Mitchell et al., 2015). For instance, cover crops could increase soil moisture after their termination if there is sufficient post-termination rainfall, despite transpiring water during cover crop growth (Burke et al., 2021; Carlson and Stockwell, 2013; Unger and Vigil, 1998). Multiyear studies are needed to identify the local management and weather conditions that favor positive outcomes for water under cover cropping in semi-arid regions.

Plant-soil-microbe interactions

Soil health management strongly affects plant-soil-microbe interactions (Bender et al., 2016; Dias et al., 2015; Mariotte et al., 2018) and as such, impacts the effective use of water. Specifically, practices like conservation tillage, cover cropping, organic matter amendments, and diversification of crops typically enhance microbial biomass relative to conventional practices (Kallenbach et al., 2011; McDaniel et al., 2016; Paustian et al., 1997; Poepflau and Don, 2015) and sometimes increase microbial richness and diversity (Bowles et al., 2017; Guzman et al., 2021; Oehl et al., 2004; Van Geel et al., 2017). Further, soil health management practices impact diversity and density of soil macrofauna such as earthworms (Ernst & Emmerling 2009) and positively affect the biological suppression of soil pathogens (Larkin 2015; Abawi & Widmer 2000). In turn, these changes can promote productive water use through general impacts of soil microbes on soil structure and biogeochemical processes, as well as through specific plant-microbe interactions (Zheng et al., 2018), particularly those mediated by arbuscular mycorrhizal (AM) fungi and plant growth promoting rhizobacteria (PGPR).

The soil microbial community as a whole has mainly indirect effects on productive green water flow and blue water conservation. By driving nutrient cycling and increasing plant nutrient availability (Williams and Vries, 2020), soil microbes - with the exception of soilborne pathogens - help remove limitations to plant growth and allow plants to take full advantage of available water. Greater plant productivity, specifically increased canopy cover, mediated by plant-soil-microbe interactions further reduces evaporation and increases productive green water flow (Nziguheba et al., 2010; Rockström et al., 2009; Sánchez, 2010). Moreover, soil microbes are critical for soil structure formation: soil microbial decomposition processes drive aggregate formation and aggregate turn-over which dictates soil structure (Rillig and Mummey, 2006; Six et al., 1999). Since soil structure affects soil microbial activity by altering soil water retention (Moreno-Espíndola et al., 2018; Schjønning et al., 2011) and soil management affects both structure and microbial activity, complex feedbacks exist among soil microbial activity, soil structure, soil management and green and blue water.

AM fungi are ubiquitous root symbionts that require plant carbohydrates and in return benefit their hosts via various mechanisms such as enhanced access to nutrients and water (Smith and Read, 2010). Intensive agricultural management strategies, such as frequent chemical and physical disturbance, alter soil nutrients and crop diversity, effecting AM fungal abundance, diversity, and community composition, specifically the relative abundance of AM fungal functional groups (Chagnon et al., 2013; Guzman et al., 2021; Oehl et al., 2004; Van Geel et al., 2017). Reduced tillage and winter cover cropping increase AM fungi root colonization of annual cash crops by about 30% on average, while reduced tillage can increase richness of AM fungal taxa by 11% (Bowles et al., 2017).

There are both indirect and direct mechanisms through which AM fungi affect productive green water flow and blue water conservation. AM fungi play an important role in enhancing soil structure (Rillig and Mummey, 2006) through extraradical hyphae that physically enmesh aggregates (Mardhiah et al., 2016). Arbuscular mycorrhizal fungi can enhance plants' drought tolerance by improving access to nutrients and maintaining physiological performance. This in turn, increases plant growth (Smith and Read, 2010), strengthens plants' ability to recover and/or prevents severe root and photosynthetic apparatus damage (Augé, 2001). Further, AM fungi affect plant water relations, such as stomatal regulation (Augé et al., 2015; Duan et al., 1996; Lazcano et al., 2014) in ways that may optimize responsiveness to low or variable soil moisture. AM fungi stimulate antioxidant enzyme activity which alleviates damage caused by reactive oxygen species under drought stress and improves water use efficiency and greater recovery of plants post drought (Chang et al., 2018; Duc et al., 2018; Li et al., 2019; Pedranzani et al., 2016; Thirkell et al., 2017). Lastly, recent findings show that AM fungal hyphae can transport water but relative to the plants'

transpiration demand the AM fungal-mediated water transfer appears to be low (Püschel et al., 2020). Yet, despite this evidence on how AM fungi influence soils and plant performance during water stress, we still lack a robust understanding of how soil health management can achieve optimized AM functioning with respect to enhancing nutrient and water access to crops in real agricultural settings.

Plant growth promoting rhizobacteria (PGPR) are root-colonizing bacteria that enhance plants' ability to withstand biotic and abiotic stresses via the production of a wide range of enzymes and metabolites (Chauhan et al., 2015; Glick et al., 2007; Kim et al., 2009; Mayak et al., 2004). Knowledge on the effect of soil health management on PGPR is scarce but studies suggest PGPR inoculation success may be affected by soil type, temperature, nitrogen content, salt concentration, and moisture content (Adesemoye and Egamberdieva, 2013). PGPR have been shown to enhance plants' ability to tolerate drought stress via various mechanisms (Ngumbi and Kloepper, 2016; Rubin et al., 2017; Vurukonda et al., 2016). PGPR enhances growth promoting and suppresses growth inhibiting phytohormones (Belimov et al., 2009; Bresson et al., 2014; Cohen et al., 2009; Dodd et al., 2010; Figueiredo et al., 2008; Glick et al., 1998) and hence, increase root growth (Somers et al., 2004; Timmusk et al., 2014) and shoot growth (Timmusk et al., 2014; Vardharajula et al., 2011). Further, relative water content in plants is increased with PGPR which is important for metabolic activity in plant tissue (Bano et al., 2013; Grover et al., 2014; Naseem and Bano, 2014; Naveed et al., 2014; Vardharajula et al., 2011). Finally, PGPR induces over-production of antioxidants and thus enhances plants' drought tolerance (Ngumbi and Kloepper, 2016). However, as for AM fungi, work remains to translate the possible effects of PGPR on plant drought tolerance to actionable management information for agricultural systems.

Macrofauna e.g., earthworms and ants also impact soil water dynamics through the galleries and chambers they create and that facilitate enhanced infiltration (Sofo et al., 2020). Further, by redistributing and stabilizing organic matter, macrofauna also contributes to soil aggregation and as such further enhances soil nutrient retention (Sofo et al., 2020; Fonte & Six 2010). Consequently, macrofauna may also improve productive water flow through supporting enhanced plant growth.

Plant water capture and productive use

Soil health management that impacts below- and above-ground plant traits can shift both the plant's ability to access and take up soil water and its transpiration demand. Water movement through the soil-plant-atmosphere continuum is a process driven by water potential gradients and affected by the conductivity or resistance of the pathway, such as soil and root hydraulic conductivity and stomatal conductance (Steudle, 2001). Transpiration generates high water potential gradients through the soil-plant-atmosphere continuum

and may increase with greater canopy cover and environmental evaporative demand, a function of air temperature, humidity, radiation, and wind speed. To what extent plant canopies meet this atmospheric evaporative demand varies with root system architecture, depth, root length density, and other root physiological traits such as hydraulic conductivity (Barrios-Masias et al., 2019). It is likely that for many crops, modern breeding, and selection mainly for aboveground traits in nutrient and water-rich environments have resulted in smaller root systems unable to explore a larger soil volume, diminishing the capacity of crops to respond to water deficit and increasing the dependency on irrigation (Bishopp and Lynch, 2015; Jackson and Koch, 1997). Irrigation management itself influences roots, such as concentration of roots around subsurface drip irrigation lines, and may interact with crop genotype to promote or hinder root proliferation (Li et al., 2010). Root systems have been understudied, and root system architecture and growth habit are important determinants for water uptake. For instance, a three-year old plum tree (*Prunus persica* L.) can occupy a 50 m³ of soil with a root system composed of woody and fine roots, which actively respond to the soil environment (Vercambre et al., 2003), and annual crops tend to increase their resource uptake at deeper soil profiles when reaching maturity (Weaver and Bruner, 1928). Thus, understanding root proliferation and its plasticity in a heterogeneous soil profile, including abiotic and biotic interactions as influenced by soil health management, could result in more effective use of water.

Soil health management can create conditions favorable to root growth such as increasing soil macroporosity, reducing penetration resistance, and maintaining soil moisture, which facilitate root elongation, oxygen diffusivity, and heat transport (Nunes et al., 2019). A generalized framework across management practices is that higher SOM due to soil health management tends to increase aggregation and promote growth and maintenance of roots by alleviating negative impacts on roots from both compaction in drying soils and poor aeration in waterlogged soils (King et al., 2020). Soil health management practices that include deep-rooted crops and continuous living soil cover for most of the year, such as cover crops, perennial forage and grain crops, can further alleviate negative impacts of compaction by creating root channels, or biopores, that subsequent crop roots use at lower metabolic cost to the plant (E. Han et al., 2015; McCallum et al., 2004; Perkons et al., 2014; Williams and Weil, 2004). Soil organic matter derived from living roots in soils are more efficiently stabilized by microbes and more persistent in soil than inputs of shoot residues (Kong and Six, 2010). This may create positive feedbacks over years to decades between greater root growth, belowground carbon input, SOM accrual, microbial activity, and improved soil aggregation and hydraulic properties (see previous sections) (Austin et al., 2017). Tradeoffs for soil health management practices must be considered. For instance, no-till practices can reduce root penetration in the topsoil even as it alleviates subsoil compaction (Munkholm et al., 2013) and negatively impact root

architecture and growth of the main root axes (taproot), limiting the capacity for root exploration and water uptake (Martínez et al., 2008).

Root interactions with rhizosphere microbial communities, particularly AM fungi, may impact not only root water and nutrient access and uptake but also potentially root-to-shoot signaling that influences canopy plant-water relations and helps maintain canopy transpiration in drying soils (see previous section on plant-soil-microbe interactions). Root traits that favor desirable root-microbe interactions may have been lost during selection for increased yields under high input systems, which hinders the positive effect of increased soil health (Barrios-Masias et al., 2019; Schmidt and Gaudin, 2017). Intensive agricultural systems tend to select for high-yielding crops, without regard for robust root systems and soil-plant communities (Pretty and Bharucha, 2014). Selecting adaptive root responses to water availability by breeding have rarely been applied to irrigated systems (Schmidt and Gaudin, 2017). Greater nutrient uptake is usually associated with increased root length density.

Soil health practices such as crop rotation diversification and organic management favor disease-suppressive and pest-suppressive soils, and facilitate rhizosphere soil-microbe-plant interactions, which could help limit yield reductions due to pests and disease (Blundell et al., 2020; Peralta et al., 2018). Soil health management also boosts nutrient cycling by soil microbes and nutrient availability to plants, including nitrogen, which has been shown to improve the water use efficiency of crops such as wheat under low water availability (Cousins et al., 2020) (see previous section on plant-soil-microbe interactions). Higher soil biological health has been shown to increase maize yield even in intensively managed systems (Wade et al., 2020). If soil health management alleviates yield-limiting factors and thus increases crop growth and canopy cover (e.g., percent soil canopy cover), it may increase productive water flow.

Leveraging soil health benefits in irrigated systems

Soil health and water conservation in irrigated systems

Shifting irrigation strategies is essential to realizing potential benefits of building healthy soils for blue water conservation in irrigated cropping systems, especially when blue water is a substantial portion of the crop water budget. If soil health management results in capturing and storing more green water, but managers do not reduce blue water inputs, then greater blue water conservation will not be achieved. Similarly, storage of green water may not be utilized by a crop if the root system cannot access it (e.g., shallow rooted crops). Since irrigation managers must recognize and react to changes in soil health that

affect water dynamics in order to conserve blue water, realizing the full benefits of soil health management is more complex in irrigated systems compared to rainfed systems that only rely on green water. We consider how several irrigation strategies could interact with healthy soils to support water conservation and reduce vulnerability to droughts or policy interventions that impact blue water supplies, which we term “integrated soil health-water management”, revealing many research gaps (Table 1). We focus on regions with Mediterranean-type climates that have distinct wet and dry seasons (e.g. California, Chile, Mediterranean Europe / North Africa, the Middle East, and parts of Western and South Australia).

First, increasing the proportion of precipitation captured and stored as green water during the rainy season could allow for a delay in the onset of irrigation, in turn saving blue water (Devine and O’Geen, 2019). Irrigation can be delayed until just before the onset of plant water stress, which broadly speaking occurs when about half of plant available water in the soil has been consumed, a proportion known as allowable depletion (Devine and O’Geen, 2019; Hanson et al., 2000). For instance, in California, processing tomato fields may not be irrigated for the initial two to three weeks after transplanting in early spring as seedlings establish with stored green water. Increasing the quantity of plant available water and delaying irrigation initiation would allow for stored green water to contribute relatively more to the crop water budget. Roots would also grow more deeply if seedlings are not established with abundant surface water.

Reducing non-productive evaporative losses of water (green or blue) and/or increasing green water capture could also result in less irrigation water applied. When irrigation is applied on the surface or overhead, substantial reductions in evaporation result from reduced tillage and residue retention (Klocke et al., 2009; Peng et al., 2020). For instance, wheat production under semi-arid conditions under no-till and mulching increased yields by over 40% and precipitation use efficiency by at least 39% (Peng et al., 2020). In tomatoes, no-till and residue retention resulted in an estimated 5–10 cm of reduced evaporation in California annual cropping systems (Mitchell et al., 2012), which accounts for ~10 % of crop ET. Since residues also reduce surface runoff (Smith, 2016), they can enable less frequent, deeper irrigations, which maximize benefits of reduced evaporation (Mitchell et al., 2012). Cover crops are highly effective at increasing infiltration of winter rainfall and could act in concert with soil management that increases soil water storage to increase overall green water availability for crops. Yet, perceptions that cover crops can result in a net reduction in water availability for cash crops is a key barrier to cover crop adoption in arid environments (Carlisle, 2016). Net impacts of cover crops in irrigated systems must be carefully considered, i.e. the balance between increasing the capture of precipitation via increased infiltration and reductions in evaporation vs. increases in transpiration and reduction in percolation (Meyer et al., 2019). Careful management for the timing of establishment, termination, and species selection is key to realizing net

benefits. In areas with very low winter rainfall, crop residues are an alternative to cover crops for increasing infiltration and reducing evaporation, since there is no risk of competition with cash crops for stored green water (Downer and Hodel, 2001; Iqbal et al., 2019). However, unlike cover crops, crop residues would not increase infiltration at depth or leave beneficial biopores (see below).

In Mediterranean-type systems with a long dry season, increasing green water capture and storage through soil health management may only be able to play a relatively minor role in reducing irrigation needs. This is because the soil profile can only store so much water and it can be rapidly depleted after rains stop and evapotranspiration increases dramatically, especially in shallow-rooted crops. In such areas, we suggest that other potential benefits of healthy soils for enabling irrigation management that conserves water may be similarly or relatively more important. First, if soil health management promotes plant-microbe interactions and greater root proliferation/penetration, then it may be possible to use various irrigation strategies such as deficit and partial root zone drying (Box 1; Barrios-Masias and Jackson, 2016) that reduce blue water consumption while minimizing impacts on crop productivity. Soil health management that enables deficit irrigation may be particularly important in perennial crops (e.g. nut trees) that require significant capital investment to establish and cannot be fallowed during times of water scarcity. Composted manure application in an almond orchard in California increased soil volumetric water content in the top 1.5 m by 22% during the driest part of the study year (Lepsch et al., 2019), suggesting organic matter-building strategies could be used to reduce impacts of deficit irrigation on tree water status.

Soil health management could also help different types of root systems adapt to irrigation management and improve the effective use of water. With irrigation, crops often invest preferentially in shallow roots, especially with drip or microsprinkler systems in which applied water is more localized and does not penetrate deeply into the soil profile (Li et al., 2020). Well-structured soils with low resistance to root proliferation could help crops rapidly develop shallow axial roots with extensive lateral branching that would enable exploitation of water and nutrient resources concentrated near the soil surface (Schmidt and Gaudin 2017). Yet this could also limit root access to water (and nutrients) available deeper in the full soil profile (Barrios-Masias et al., 2019). Biopores, such as those formed from roots or earthworms and preserved when soil is not disturbed, enable steeper-angled axial roots to penetrate deeper (Stirzaker et al. 1996; Xiong et al. 2022). Selection of cover crops or preceding cash crops that promote biopores help subsequent cash crops develop deeper roots (Huang et al. 2020; Zhang et al. 2022). In turn, this would support reducing applied water, such as through partial root drying (Barrios-Masias and Jackson, 2016) or with less frequent irrigations at drier soil moisture thresholds (Devine and O'Geen, 2019). With careful crop and cultivar selection, promoting deep rooted crops can allow for more consistent access to soil

moisture without a rapid onset of drought stress and even allow for dry farming (Leap et al 2017; Garrett 2019), a strategy involving almost complete reliance on green water during seasons without rainfall and long practiced by desert farmers and Indigenous peoples around the world (Nabhan 2013).

Over larger areas and longer time scales, managing groundwater recharge through enhanced percolation will also be essential to “bank” water available during wet years for drier years. As finite surface water allocations decline in the future, demands on groundwater will continue to increase (Massoud et al., 2018). Could building soil health improve regional water balances by enhancing deep percolation? In California, groundwater overdraft has occurred at a rate of 2.5 billion cubic meter per year since 1960 due to pumping for irrigation and increased reliance on groundwater during times of drought (PPIC, 2018). Recognizing the potential for depleted aquifers to store 44-80 cubic kilometers of water, or three times that of surface water reservoirs, the state’s water managers have accelerated the pursuit of managed aquifer recharge on working landscapes, including croplands (Dahlke et al., 2018). In years with high magnitude stream-flows, above the required environmental flows and surface water allocations, excess water can be diverted onto agricultural fields to recharge underlying aquifers (Kocis and Dahlke, 2017). A field site’s ability to quickly infiltrate and accommodate deep percolation of water below the root zone to groundwater is essential for increasing the recharge capacity of a site and to avoid negative effects on perennial crop physiology (O’Geen et al., 2015). While areas suitable for groundwater recharge are mainly influenced by subsurface soil properties, topography, and crop type (O’Geen et al., 2015), soil health management could help increase percolation below the root zone and reduce potential negative consequences of aquifer recharge. For instance, nitrate pollution of groundwater could occur if significant residual nitrate exists in the soil profile, exacerbating pollution that has already contaminated drinking water wells in many areas of California (Balazs et al., 2011; Bastani and Harter, 2019; Waterhouse et al., 2020). Cover crops could help reduce nitrate levels while also increasing infiltration during managed aquifer recharge.

Tradeoffs and contradictions with irrigation efficiency technologies

Irrigation technologies that increase crop yield per unit of water applied can also reduce pressure on blue water resources. Subsurface drip irrigation, soil water monitoring and other efficiency technologies can lead to significant reductions in water applied with similar or higher levels of productivity.

When benefits outweigh the capital investments required, irrigation efficiency technologies spread rapidly, such as the widespread shift from furrow irrigation to drip and microsprinkler irrigation in California (Taylor and Zilberman 2017). Yet, tradeoffs also must be considered. For instance, subsurface drip irrigation reduces percolation during the summer when irrigation occurs, limiting the ability to store water

during wet years when surface water is more abundant (Niswonger et al., 2017; O'Geen et al., 2015; Perry, 2007; Scanlon et al., 2015). Another major issue occurs when increases in irrigation efficiency at the field scale lead to expansion of irrigated acreage or higher-value but more water-intensive crops, thus actually increasing total blue water use at the regional scale (Alcott, 2005; Sears et al., 2018). This is one example of a widely recognized environmental challenge, called Jevon's paradox, that limits the utility of interventions aimed at increasing resource use efficiency at a small scale without effective governance and policies at broader scales.

There are also possible tradeoffs with soil health. In a comparison of furrow irrigation, which wets a greater volume of soil less frequently, and subsurface drip irrigation, which wets a smaller volume of soil more frequently, Schmidt et. al (2018) found that subsurface drip reduced the soil's C sequestration potential. Specifically, furrow irrigation had larger soil aggregates in the upper 10 cm of the soil, a greater proportion of C in macroaggregates, and a lower proportion of C in the unprotected silt and clay fraction compared to subsurface drip irrigation after just two years. While the drip system had higher blue water productivity (i.e. more crop per drop) than furrow, declining aggregation in the drip irrigated system could affect infiltration and percolation during the rainy season and thus potentially green water storage and blue water recharge (Schmidt et al. 2018). Different soil wetting patterns with subsurface drip irrigation also affects soil bacterial and fungal community composition, though functional implications are not clear (Quach et al 2022). While limited research exists currently, these studies suggest that different wetting patterns of subsurface drip or microsprinklers may impact soil health.

Transitions to integrated soil health-water management

Developing strategies for integrated soil health and water management and quantifying outcomes are important steps to leverage more green water and reduce applied irrigation water, toward a larger goal of sustainable water resource management in irrigated regions. But such knowledge will not be sufficient to actually make the shift (Chartzoulakis and Bertaki 2015; Iwanaga et al 2020). With changes in decision making required across multiple, interacting levels, from individual farms to irrigation water management districts to regional policymaking, driving change in both soil health systems and water resources management is complex and social-ecological in nature. For example, structural factors like the degree of land tenure, access to values-based markets, and public and private policy all influence farmers' adoption of soil health practices (Esquivel and Carlisle et al 2022; Carlisle 2016), with motivation and other personal characteristics playing an important mediating role (Prokopy et al 2019). Shifting to various deficit irrigation management strategies or less water-intensive alternative crops will likely be by necessity only,

such as when water for irrigation is less available and/or becomes more expensive (Trout and Manning, 2019). Deficit irrigation involves more risks for crop productivity, and alternative crops may generate lower returns, which is an issue when the highest value cropping systems in a region drive land values. While some exceptions exist, such as when deficit irrigation leads to profitable increases in crop quality despite lower yields (e.g. with dry farmed tomatoes or winegrapes; Acevedo-Opazo et al 2010), motivation to reduce water inputs will need hard climate or regulatory realities. Yet such realities may actually be a window of opportunity for farmers to try soil health management as a strategy for reducing the negative impacts of less water available. Times of crisis can spur farmers to reevaluate their systems, consider new possibilities, and experiment in novel ways (Folke et al. 2010; Darnhofer 2014). Innovations could spread when farmers that pioneer integration of soil health and irrigation management and find new ways of thriving with reduced blue water availability share their experiences and knowledge through farmer-to-farmer networks.

At broader levels, policies must address common pool resource problems (Ostrom, 1990), or problems in which action at the individual level does not produce optimal outcomes for all users. Participatory policy processes are essential, not only to consider the wide array of values, knowledge and perspectives that farmers, water managers, and other stakeholders bring, but also to increase the scope of possible solutions and generate momentum for collective action (von Korff et al. 2012). As an example, groundwater use in California has resulted in a quintessential common pool resource problem and extreme levels of groundwater overdraft (Faunt et al. 2016). To address this, in 2014 the California legislature enacted the Sustainable Groundwater Management Act (SGMA) which established new basin-level management institutions, Groundwater Sustainability Agencies (GSA), tasked with developing locally-relevant Groundwater Sustainability Plans (GSPs). As a form of collaborative governance, SGMA mandates that all interested water users, and specifically disadvantaged communities, be involved in the process (Lubell et al 2020). GSPs must chart how the basin will achieve groundwater sustainability via several approaches, including voluntary, incentive-based water saving approaches, implementing on farm managed aquifer recharge, and control policies such as implementing pumping quotas (Niles and Hammond Wagner, 2019). Pumping quotas could result in fallowed fields and negative consequences for the agricultural industry. If incentive programs for soil health management were coupled with regulations for water resources like SGMA, then some negative impacts could be avoided. In other words, providing support for soil health practices could also help farmers adapt to regulatory and climate realities of water limitations. Following the California example, the state-level Healthy Soils Program (HSP) currently provides grants for soil health practices and recognizes the many co-benefits of building soil health (CDFA, 2022), but synergies between

HSP and SGMA remain undeveloped. GSAs could engage with technical assistance providers to support farmers' transitions to soil health management in concert with water-saving irrigation practices (Table 1), building on lessons learned from pioneering farmers, much like a regional model for developing conservative agriculture (Mitchell et al 2016). In turn, modelers could quantify how integrated soil health and water management at the field scale affects the regional water balance. For example, could adoption of soil health practices and changes in irrigation strategies, such as deficit irrigation or changes in the timing and duration of irrigation events (Table 1), if broadly incentivized and adopted, be impactful components of GSA Groundwater Sustainability Plans? How could building soil health on low permeability soils improve infiltration and a farmer's ability to participate in managed aquifer recharge programs, expanding beyond only sites with soils already capable of high infiltration rates (O'Geen et al. 2015)? In sum, regulatory intervention and governance is necessary to move water resource use toward sustainability, while incentives and technical assistance for soil health management could help farmers adapt to new realities with fewer negative consequences. Doing so will require the active commitment and collaboration of farmers, policymakers, and other stakeholders.

Quantifying outcomes: opportunities and knowledge gaps

Water resource managers, farmers and other stakeholders require quantitative information on the potentials and limits of integrated soil health-water management to increase productive green water flows, resilience to blue water shortages, and conservation or even regeneration of blue water resources. Most reports on soil health response to management quantify soil physical, chemical, and more rarely, biological indicators of soil health, as we have reviewed here. While such approaches are crucial for establishing linkages between management and these dynamic properties, understanding the impact of such changes on plant-water relations, crop productivity during drought or deficit irrigation, or on green and blue water flows at scales beyond the plot or field is much more challenging. To move beyond optimistic guesses to actionable information, research and development in several areas are needed.

Rather than focus mainly on soil health indicators, field-scale trials must be able to demonstrate outcomes of soil health management for hydrological processes, plant water status, crop production during water limitation, and blue water savings. Doing so may require new experimental approaches (Gilbert and Medina, 2016). For instance, recent field trials have linked organic matter additions in the form of compost and chipped woody biomass to higher soil moisture and measurable mitigation of tree water stress including under deficit irrigation (Jahanzad et al., 2020; Lepsch et al., 2019). When possible, we recommend

experimental designs with a gradient of irrigation water inputs, such as 25%, 50%, 75%, and 100% of crop evapotranspiration, to determine if thresholds in terms of crop stress or yield response to deficit irrigation differ between soil health management treatments and controls. However, standardizing irrigation volume in soil health treatments and controls limits inference about mechanisms other than higher soil moisture. Rather than standardizing by irrigation volume, standardizing soil moisture between soil health treatments and controls would allow testing for soil health management-induced shifts in dynamic soil biological and physical properties, plant-soil microbe interactions, root water acquisition, and plant water status, rather than only soil moisture *per se*. For example, a deficit irrigation experiment with no difference in soil moisture between treatments with and without symbiosis with AM fungi showed that AM fungi alleviate plant water stress under deficit irrigation when soil moisture is similar between treatments with and without AM symbiosis (Bowles et al., 2016). Soil sampling or preferably soil sensors that are used for irrigation management could monitor soil moisture in a treatment or experimental unit and then variable irrigation volumes could be applied to standardize soil moisture between soil health treatments and controls. Another option that could be more feasible than standardizing by soil moisture, and helps avoid challenges of soil heterogeneity and sensor measurement area, is standardizing by plant water status through direct measurement or proximal or remote sensing at the plant, plot, or field scale, which would allow the blue water savings that are hypothetically possible through soil health management to be quantified, regardless of whether higher soil moisture, plant water status, or other mechanisms are responsible. Future manipulative and observational studies could generate insight into the relative importance of key pathways (abiotic and biotic paths) through which soil health management protects plants in water-limited conditions, such as impacts on the soil water balance, soil nutrient cycling, and plant water and nutrient uptake and status with or without the contribution of soil microbes, and target management accordingly.

In regions with irrigated landscapes and large water conveyance systems, scaling up the impacts of soil health management beyond the plot or field scale is essential to quantify potential blue water conservation through increasing green water availability, irrigation savings, and/or aquifer recharge. Quantification is essential for assessing the relative costs and benefits of different interventions (Hoekstra, 2019). For example, in California, ~3.5M ha of irrigated croplands consume 36-46 cubic kilometers of blue water annually (Cooley, 2020). Yet, availability of surface water resources is increasingly volatile as climate change progresses, leading to groundwater overuse and subsequent policies to implement state-mandated Groundwater Sustainability Plans (GSP) (Harter, 2015). Given the uncertainty in surface water availability, reduced pumping likely will have to be included in GSPs to bring groundwater basins back into sustainable use (Escriva-Bou et al., 2020). In an analysis by Devine and O'Geen (2019) based on inherent soil

properties, deficit irrigation (down to 1 meter deep and 50% allowable depletion) in perennial cropland across California could produce blue water savings equal to 30 cubic kilometers and increase green water use by 7 cubic kilometers over 13 years. This is equivalent to filling the largest surface reservoir in the state, Shasta Lake, 6.6 times over.

To what extent could adoption of soil health management increase green water availability and reduce demand for blue water for irrigation? Would incentivizing adoption of such practices at scale be more cost effective than alternatives like increasing surface reservoir capacity, and with more co-benefits? To what extent could soil health management also increase aquifer recharge through enhanced percolation or reduced groundwater withdrawals, thus forming part of GSPs in California or analogous policies elsewhere? Such questions will require interdisciplinary research that supports agro-hydrological modeling across scales. We identify several challenge areas where modeling development is needed.

Capturing agroecosystem changes due to soil health by modeling

The answer to the key question of how building healthy soils impacts the use of water resources in irrigated agriculture depends on interactions among multiple spatial and temporal scales, and ultimately on water resources management itself. For instance, at the plant canopy to field scale, models must be able to capture how agricultural management, including practices to increase soil health and irrigation, affect soil hydraulic properties and hydrologic flows. Beyond the field, at a basin scale, hydrologic and water resource models must be able to capture how these changes in turn affect green and blue water flows and reservoirs. For instance, changes in infiltration and soil water storage at the field scale could affect runoff and percolation, and thereby blue water flows to surface and groundwater (Flint et al., 2013, Lei et al., 2021). Such coupled models could in turn be used to assess different climate change scenarios, including drought, and reduced blue water supply.

Recent work has focused on how to capture the effect of soil health practices in hydrologic models (Evenson et al., 2022, Maharjan et al., 2018). With this work as a starting point, we selected four representative and widely-used models that simulate the movement of water at different scales. We highlight strengths, weaknesses and opportunities for capturing impacts of soil health management on green and blue water flows (Table 2). We consider both approaches for estimation of water savings: less drop per crop (water management strategies) or more crop per drop (closing yield gap at constant water use) (Bao, 2020; Blum, 2009)

At the global scale, the Lund-Potsdam-Jena managed Land (LPJmL [Rost et al., 2008]) model evaluates global blue-green water patterns. LPJmL is a dynamic global vegetation and hydrologic model, which models coupled carbon, nitrogen, and water fluxes, crop growth and yields. Improvements to LPJmL, including irrigation schemes (Rost et al., 2008) and reservoir management, assess the changing role of reservoirs in sustaining water supply for irrigation in agroecosystems (Biemans et al., 2011). Additionally, LPJmL 5.0 takes into account impacts of tillage by focusing on changes in SOM pools and bulk density (Lutz et al., 2019), making LPJmL potentially capable of assessing how tillage affects irrigation demand and crop yields.

At a watershed to basin scale, the Soil & Water Assessment Tool (SWAT [Arnold et al., 1998]) model predicts the impacts of land management practices on water and sediment flow, including crop yield and crop water productivity, and can incorporate irrigation scheduling (Sun and Ren, 2014). The SWAT model has been used to estimate green and blue consumption at basin scale, allowing efficient water and soil management (Msigwa et al., 2022). However, SWAT cannot estimate water losses to evaporation from water reservoirs, channels and after irrigation water application (Luan et al., 2018), limiting blue water flows estimation. SWAT becomes inaccurate at modeling streamflow during drought periods (Veettil and Mishra, 2016), a disadvantage in running climate change scenarios. However, a calibrated SWAT version was developed validating input parameters for wet and dry seasons (Zhang et al., 2015).

At the field scale, the Food and Agriculture Organization (FAO) model AquaCrop estimates green-blue water balance and plant growth, and can incorporate management impacts on blue water, such as using drip irrigation and soil management techniques like mulching (Chukalla et al., 2015; H. Nouri et al., 2019). Currently, AquaCrop does not simulate lateral water flows, fertilization, or nutrient dynamics (Chukalla et al., 2015). AquaCrop describes soil water stress in more detail than SWAT, but SWAT is more detailed for soil fertilization (Van Gaalen, 2016). AquaCrop may perform better for soil water balance and crop growth processes than SWAT (Hunink et al., 2011), and is better able to parameterize irrigation management. In the comparison of these two models, if ‘less drop per crop’ (decreasing water use at stable yield) is more important than ‘more crop per drop’ (closing yield gap at constant water use), AquaCrop would be more suitable.

In contrast to larger scale models, physical models focused on finer scales like HYDRUS 1D (Šimůnek and Genuchten, 2008) better capture soil hydraulic properties. Capturing how soil health management affects soil hydraulic parameters in process-based models is one limitation in accurately estimating water resources across scales (Dang et al., 2020). For instance, while LPJmL, AquaCrop and SWAT use total porosity as

an input of a unimodal pore size distribution, HYDRUS 1D accounts for dual-porosity, i.e., inter-aggregate pores and intra-aggregate pores with contrasting sizes, and hysteresis in the water flow calculation (Šimůnek et al., 2016). Not considering bimodality in topsoils due to soil management derived from tillage may compromise the accuracy of soil-water modeling (Acevedo et al., 2020; Jensen et al., 2019). A HYDRUS 1D module can incorporate the effect of tillage, homogenizing the soil water content and liquid phase concentration in the topsoil layers, although it does not incorporate changes in soil hydraulic properties with tillage (Mallmann et al., 2014).

To realistically capture outcomes for water provisioning, agro-hydrological models must be able to—directly or indirectly via connection to another model—simulate how soil health management affects the dynamic soil properties that influence hydrologic processes and hydraulic properties, shifts in plant growth that influence water access and demand, and eventually, plant-microbe interactions that influence plant responses to water stress. Overall, the agro-hydrological models reviewed here incorporating soil health management have largely focused on reduced or no tillage. Using field data to validate model parameters, other soil health management such as cover crops or organic matter amendments could be directly incorporated or, as a starting point, represented through their impacts on SOM. Realistic representation of soil health management and soil-water processes in hydrological models requires targeted data collection and validation for dynamic soil property inputs (Gómez et al., 2020). SOM dynamics are some of the most important soil health management-induced changes in agroecosystems (Bolinder et al., 2020) and accurate modeling of hydrological outcomes of soil health management thus require comprehensive integration of SOM processes over time (Kemanian et al., 2011). In LPJmL5, all organic matter pools (vegetation, litter, and soil) are subject to decomposition. The decay of organic residues cannot be simulated by standard versions of AquaCrop, HYDRUS-1D or SWAT, but all of these have been coupled to submodels to address this need. In models where SOM pools are subject to decomposition, hydraulic parameters can be dynamically estimated at different depths based on pedotransfer functions using SOM as an input (Lutz et al., 2019), allowing modeling of soil health practices that affect SOM.

Accurate data on soil physical properties can lead to a better understanding of the state of soil health at local and landscape scales (Keesstra et al., 2021), but some assumptions about soil hydraulic properties can oversimplify their effect on models. First, most hydrological models assume that soil hydraulic properties are constant over time, hampering efforts to capture how management practices temporally affect key properties like soil structure and pore size distribution (Chandrasekhar et al., 2019). For example, soil health management can boost pore connectivity (Galdos et al., 2019); however, temporal changes in soil structure are still in early stages of development (Chandrasekhar et al., 2019; Vereecken et al., 2016). Second, another

assumption is that at different starting soil conditions, properties are arbitrarily changed after management, commonly as correction factors. An example is the bulk density after tillage. The intensity and type of tillage can be represented in simulation models as a mixing efficiency parameter (mE, from 0 to 1) corresponding to the fraction of change in bulk density after tillage (White et al., 2010). Third, most process-based models use pedotransfer functions (PTFs), which establish an empirical relationship between easily measurable properties and soil hydraulic parameters, may be a source of uncertainty in the estimation of agroecosystem changes due to soil health. AquaCrop, SWAT and LPJmL 5.0 use the Saxton and Rawls, 2006 PTF for water characteristics for water potential and hydraulic conductivity by default and HYDRUS-1D uses Rosetta, both developed with US soil databases. This could bias models towards soil types and climate conditions found in this geographic area. Management-induced changes in soil hydraulic properties are also not necessarily consistent across different locations, soils, and experiment designs (Schwen et al., 2011), hampering the use of default parameters in accurate modeling runs. PTF can aid in quantifying benefits of soil management and can be used to model the effect of changes in management on drought resilience (Bagnall et al., 2022). As Cueff et al. (2021) found, available PTFs applied equally to soils under conservation or conventional agriculture have shown poor accuracy results.

Green water is stored in the vadose zone, which is the subsurface unsaturated zone between the topsoil surface and the saturated zone below the water table. Modeling unsaturated soil flow properly is thus mandatory for estimating green water use and storage. Favorable soil properties related to water flow in unsaturated, drying soil, including high porosity and a higher fraction of large capillary pores and a lower fraction of gravitational pores (Kodešová et al., 2009), should be incorporated ideally using real measurements. For example, Maharjan et al. (2018) reviewed 16 hydrological models able to integrate tillage effects and concluded that simulating this management requires site-specific empirical measurement of changes in bulk density after tillage. Notably, hydraulic conductivity at varying water contents is not commonly reported in soil health management studies, which tend to focus on soil physical properties that are relatively easily measurable, such as infiltration or saturated hydraulic conductivity.

Agro-hydrological models could better incorporate and enhance flexibility regarding the biotic components of the soil environment. Across agro-hydrological models generally, changes in the root zone by crops (and/or cover crops) must be properly represented to calculate ET accurately. HYDRUS-1D can simulate crop growth status and water uptake by plants (Karandish and Šimůnek, 2019; Šimůnek et al., 2016) but the root system is considered as a static component (without feedback between the root system and the soil conditions), limiting ability to simulate agroecosystems under different soil health management practices (Parihar et al., 2019). Largely due to a lack of data and uncertainty about representation in models,

hydrologic and crop models do not currently explicitly account for soil biology impacts on plant water uptake, status, and transpiration. Quantifying the effect of soil-plant-microbe interactions on model-relevant inputs such as actual evapotranspiration would enable their incorporation into models and, ultimately, more accurate estimation of the outcomes of building soil health not only for soil physical properties but also crop resilience under various climate change and irrigation scenarios.

However, models must be simple to ensure their operability. Thus, the challenge to integrate soil health into agroecosystem and landscape hydrology increases the need for calibration data and coupled models. For example, AquaCrop has been coupled to HYDRUS-1D and 2D to describe water flow and retention within the soil profile to validate optimum management practices for increased water storage capacity (Gómez et al., 2020). In a similar approach HYDRUS-1D has been coupled with SWAT for quantifying the capillary rise from groundwater contribution to crop transpiration (Han et al., 2015). To sum up, none of the reviewed models can alone incorporate interactions between soil health and irrigation management, and then impacts on green water and blue water reservoirs. Additional development will be needed to incorporate soil health into agroecosystem and landscape hydrology at different scales.

Our analysis indicates that current knowledge of soil health and water is robust on soil moisture, infiltration, and retention (Morris and Bucini, 2016), but to the best of our knowledge it does not yet capture soil health's effect on whole-system hydrologic functioning via modeling. Modeling is a promising avenue for quantifying the magnitude of soil health-induced gains in water savings and drought resilience in irrigated agricultural landscapes. However, current models neither capture the full effect of soil health practices on soil hydraulic parameters nor the influence of soil, plant, and microbial processes and interactions on water cycling

Conclusions

We show several pathways through which soil health management could synergize with irrigation and crop management to increase effective use of green and blue water resources in irrigated cropping systems. Widely cited benefits of soil health management in rainfed agriculture, namely higher infiltration rates and soil water storage, are an important way to increase green water availability. Yet, greater beneficial plant-microbe interactions, root water capture, and reducing plant water stress are at least as important for maintaining productivity in irrigated contexts that depend more on blue water than green water, especially in Mediterranean climate regions dominated by perennial crops that cannot be fallowed during periods of water scarcity. We provide a roadmap for how soil health management can enhance the effectiveness and

suitability of irrigation and water resource management strategies that conserve water resources, such as deficit irrigation or agricultural managed aquifer recharge, and related uncertainties and challenges yet to be resolved. Future research priorities include: 1) promising agro-hydrological modeling strategies and gaps for quantitative scenario analysis and 2) planning to discern possible gains in water conservation and drought resilience through integrated soil health-water management. Impacts on water resources, especially increases in aquifer recharge through enhanced percolation or reduced groundwater withdrawals, are important to evaluate. For quantitative information on how soil health management affects water resources at scales relevant to water resource managers, soil, crop, and hydrological models must be better integrated and expanded, including potential physical and biological changes in the soil. Challenges, trade-offs, and unintended consequences of soil health management for water conservation are also possible, for instance if blue water conservation through soil health management at the field scale enables expansion of irrigated acreage. Changes in irrigation and water resource management must accompany and complement soil health management for blue water savings to be realized. A suite of agroecological and technical research approaches may resolve how building healthy soils impacts the effective use of water resources, specifically in irrigated systems, and help meet the global agricultural water challenge.

790 References

- 791 Abatzoglou, 2021. Climatology Lab. URL <https://www.climatologylab.org/gridmet.html>
- 792 Abatzoglou, J.T., 2013. Development of gridded surface meteorological data for ecological applications
793 and modelling. *Int. J. Climatol.* 33, 121–131. <https://doi.org/10.1002/joc.3413>
- 794 Acevedo, S., Contreras, C., Martinez, S., Bonilla, C., 2020. Estimating the unimodal and bimodal van
795 Genuchten model parameters using the simplified evaporation and dewpoint methods (other). oral.
796 <https://doi.org/10.5194/egusphere-egu2020-15082>
- 797 Acevedo-Opazo, C., Ortega-Farias, S., Fuentes, S., 2010. Effects of grapevine (*Vitis vinifera* L.) water
798 status on water consumption, vegetative growth and grape quality: An irrigation scheduling application to
799 achieve regulated deficit irrigation. *Agricultural Water Management* 97, 956–964.
800 <https://doi.org/10.1016/j.agwat.2010.01.025>
- 801 Adesemoye, A.O., Egamberdieva, D., 2013. Beneficial Effects of Plant Growth-Promoting Rhizobacteria
802 on Improved Crop Production: Prospects for Developing Economies, in: Maheshwari, D.K., Saraf, M.,
803 Aeron, A. (Eds.), *Bacteria in Agrobiolgy: Crop Productivity*. Springer Berlin Heidelberg, Berlin,
804 Heidelberg, pp. 45–63. https://doi.org/10.1007/978-3-642-37241-4_2
- 805 Aggarwal, P., Bhattacharyya, R., Mishra, A.K., Das, T.K., Šimůnek, J., Pramanik, P., Sudhishri, S.,
806 Vashisth, A., Krishnan, P., Chakraborty, D., Kamble, K.H., 2017. Modelling soil water balance and root
807 water uptake in cotton grown under different soil conservation practices in the Indo-Gangetic Plain.
808 *Agriculture, Ecosystems & Environment* 240, 287–299. <https://doi.org/10.1016/j.agee.2017.02.028>
- 809 Alcott, B., 2005. Jevons' paradox. *Ecological Economics* 54, 9–21.
810 <https://doi.org/10.1016/j.ecolecon.2005.03.020>

811 Alegre, J.C., Cassel, D.K., Amezquita, E., 1991. Tillage systems and soil properties in Latin America. *Soil*
812 *and Tillage Research* 20, 147–163. [https://doi.org/10.1016/0167-1987\(91\)90037-X](https://doi.org/10.1016/0167-1987(91)90037-X)

813 Alhameid, A., Singh, J., Sekaran, U., Ozlu, E., Kumar, S., Singh, S., 2020. Crop rotational diversity impacts
814 soil physical and hydrological properties under long-term no- and conventional-till soils. *Soil Res.* 58, 84.
815 <https://doi.org/10.1071/SR18192>

816 Alonso-Ayuso, M., Quemada, M., Vanclooster, M., Ruiz-Ramos, M., Rodriguez, A., Gabriel, J.L., 2018.
817 Assessing cover crop management under actual and climate change conditions. *Science of The Total*
818 *Environment* 621, 1330–1341. <https://doi.org/10.1016/j.scitotenv.2017.10.095>

819 Alvarez, R., Steinbach, H.S., De Paepe, J.L., 2017. Cover crop effects on soils and subsequent crops in the
820 pampas: A meta-analysis. *Soil and Tillage Research* 170, 53–65. <https://doi.org/10.1016/j.still.2017.03.005>

821 Arnold, J.G., Srinivasan, R., Muttiah, R.S., Williams, J.R., 1998. LARGE AREA HYDROLOGIC
822 MODELING AND ASSESSMENT PART I: MODEL DEVELOPMENT. *J Am Water Resources Assoc*
823 34, 73–89. <https://doi.org/10.1111/j.1752-1688.1998.tb05961.x>

824 Assouline, S., Russo, D., Silber, A., Or, D., 2015. Balancing water scarcity and quality for sustainable
825 irrigated agriculture: Balancing water scarcity and quality for irrigation. *Water Resour. Res.* 51, 3419–3436.
826 <https://doi.org/10.1002/2015WR017071>

827 Augé, R.M., Toler, H.D., Saxton, A.M., 2015. Arbuscular mycorrhizal symbiosis alters stomatal
828 conductance of host plants more under drought than under amply watered conditions: a meta-analysis.
829 *Mycorrhiza* 25, 13–24. <https://doi.org/10.1007/s00572-014-0585-4>

830 Austin, E.E., Wickings, K., McDaniel, M.D., Robertson, G.P., Grandy, A.S., 2017. Cover crop root
831 contributions to soil carbon in a no-till corn bioenergy cropping system. *Gcb Bioenergy* 9, 1252–1263.

832 Bagnall, D.K., Morgan, C.L.S., Cope, M., Bean, G.M., Cappellazzi, S., Greub, K., Liptzin, D., Norris, C.L.,
833 Rieke, E., Tracy, P., Aberle, E., Ashworth, A., Bañuelos Tavarez, O., Bary, A., Baumhardt, R.L., Borbón
28

834 Gracia, A., Brainard, D., Brennan, J., Briones Reyes, D., Bruhjell, D., Carlyle, C., Crawford, J., Creech, C.,
835 Culman, S., Deen, W., Dell, C., Derner, J., Ducey, T., Duiker, S.W., Dyck, M., Ellert, B., Entz, M., Espinosa
836 Solorio, A., Fonte, S.J., Fonteyne, S., Fortuna, A., Foster, J., Fultz, L., Gamble, A.V., Geddes, C., Griffin-
837 LaHue, D., Grove, J., Hamilton, S.K., Hao, X., Hayden, Z.D., Howe, J., Ippolito, J., Johnson, G., Kautz,
838 M., Kitchen, N., Kumar, S., Kurtz, K., Larney, F., Lewis, K., Liebman, M., Lopez Ramirez, A., Machado,
839 S., Maharjan, B., Martinez Gamiño, M.A., May, W., McClaran, M., McDaniel, M., Millar, N., Mitchell,
840 J.P., Moore, P.A., Moore, A., Mora Gutiérrez, M., Nelson, K.A., Omondi, E., Osborne, S., Alcalá, L.O.,
841 Owens, P., Pena-Yewtukhiw, E.M., Poffenbarger, H., Ponce Lira, B., Reeve, J., Reinbott, T., Reiter, M.,
842 Ritchey, E., Roozeboom, K.L., Rui, I., Sadeghpour, A., Sainju, U.M., Sanford, G., Schillinger, W.,
843 Schindelbeck, R.R., Schipanski, M., Schlegel, A., Scow, K., Sherrod, L., Sidhu, S., Solís Moya, E., St.
844 Luce, M., Strock, J., Suyker, A., Sykes, V., Tao, H., Trujillo Campos, A., Van Eerd, L.L., Verhulst, N.,
845 Vyn, T.J., Wang, Y., Watts, D., Wright, D., Zhang, T., Honeycutt, C.W., 2022. Carbon-sensitive
846 pedotransfer functions for plant available water. *Soil Science Soc of Amer J* 86, 612–629.
847 <https://doi.org/10.1002/saj2.20395>

848 Balazs, C., Morello-Frosch, R., Hubbard, A., Ray, I., 2011. Social Disparities in Nitrate-Contaminated
849 Drinking Water in California’s San Joaquin Valley. *Environmental Health Perspectives* 119, 1272–1278.
850 <https://doi.org/10.1289/ehp.1002878>

851 Bano, Q., Ilyas, N., Bano, A., Zafar, N., Akram, A., Hassan, F., 2013. Effect of Azospirillum inoculation
852 on maize (*Zea mays* L.) under drought stress. *Pak J Bot* 45, 13–20.

853 Bao, W., 2020. Crop-water management to reduce blue water scarcity : a case study for the Yellow River
854 basin. Essay Master Thesis. <https://purl.utwente.nl/essays/85460>

855

856

857 Barrios-Masias, F.H., Lazcano, C., Hernandez-Espinosa, L., 2019. Advances in understanding vegetable
858 physiology: Root systems as the next frontier to improve sustainable vegetable production, in: Achieving
859 Sustainable Cultivation of Vegetables. Burleigh Dodds Science Publishing.

860 Basche, A., DeLonge, M., 2017. The Impact of Continuous Living Cover on Soil Hydrologic Properties: A
861 Meta-Analysis. Soil Science Society of America Journal 81, 1179–1190.
862 <https://doi.org/10.2136/sssaj2017.03.0077>

863 Basche, A.D., DeLonge, M.S., 2019. Comparing infiltration rates in soils managed with conventional and
864 alternative farming methods: A meta-analysis. PLoS ONE 14, e0215702.
865 <https://doi.org/10.1371/journal.pone.0215702>

866 Bastani, M., Harter, T., 2019. Source area management practices as remediation tool to address groundwater
867 nitrate pollution in drinking supply wells. Journal of Contaminant Hydrology 226, 103521.
868 <https://doi.org/10.1016/j.jconhyd.2019.103521>

869 Belimov, A.A., Dodd, I.C., Hontzeas, N., Theobald, J.C., Safronova, V.I., Davies, W.J., 2009. Rhizosphere
870 bacteria containing 1-aminocyclopropane-1-carboxylate deaminase increase yield of plants grown in drying
871 soil via both local and systemic hormone signalling. New Phytologist 181, 413–423.
872 <https://doi.org/10.1111/j.1469-8137.2008.02657.x>

873 Bender, S.F., Wagg, C., van der Heijden, M.G.A., 2016. An Underground Revolution: Biodiversity and
874 Soil Ecological Engineering for Agricultural Sustainability. Trends in Ecology & Evolution 31, 440–452.
875 <https://doi.org/10.1016/j.tree.2016.02.016>

876 Biemans, H., Haddeland, I., Kabat, P., Ludwig, F., Hutjes, R.W.A., Heinke, J., von Bloh, W., Gerten, D.,
877 2011. Impact of reservoirs on river discharge and irrigation water supply during the 20th century: IMPACT
878 OF RESERVOIRS ON DISCHARGE AND IRRIGATION. Water Resour. Res. 47.
879 <https://doi.org/10.1029/2009WR008929>

880 Bishopp, A., Lynch, J.P., 2015. The hidden half of crop yields. *Nature Plants* 1, 15117.
881 <https://doi.org/10.1038/nplants.2015.117>

882 Blum, A., 2009. Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop
883 yield improvement under drought stress. *Field Crops Research* 112, 119–123.
884 <https://doi.org/10.1016/j.fcr.2009.03.009>
885

886 Bolinder, M.A., Crotty, F., Elsen, A., Frac, M., Kismányoky, T., Lipiec, J., Tits, M., Tóth, Z., Kätterer, T.,
887 2020. The effect of crop residues, cover crops, manures and nitrogen fertilization on soil organic carbon
888 changes in agroecosystems: a synthesis of reviews. *Mitig Adapt Strateg Glob Change* 25, 929–952.
889 <https://doi.org/10.1007/s11027-020-09916-3>

890 Bonfante, A., Basile, A., Bouma, J., 2020. Targeting the soil quality and soil health concepts when aiming
891 for the United Nations Sustainable Development Goals and the EU Green Deal. *SOIL* 6, 453–466.
892 <https://doi.org/10.5194/soil-6-453-2020>

893 Bowles, T.M., Jackson, L.E., Loehrer, M., Cavagnaro, T.R., 2017. Ecological intensification and arbuscular
894 mycorrhizas: a meta-analysis of tillage and cover crop effects. *J Appl Ecol* 54, 1785–1793.
895 <https://doi.org/10.1111/1365-2664.12815>

896 Bresson, J., Vasseur, F., Dauzat, M., Labadie, M., Varoquaux, F., Touraine, B., Vile, D., 2014. Interact to
897 Survive: *Phyllobacterium brassicacearum* Improves *Arabidopsis* Tolerance to Severe Water Deficit and
898 Growth Recovery. *PLoS ONE* 9, e107607. <https://doi.org/10.1371/journal.pone.0107607>

899 Bünemann, E.K., Bongiorno, G., Bai, Z., Creamer, R.E., De Deyn, G., de Goede, R., Flesskens, L., Geissen,
900 V., Kuyper, T.W., Mäder, P., Pulleman, M., Sukkel, W., van Groenigen, J.W., Brussaard, L., 2018. Soil
901 quality – A critical review. *Soil Biology and Biochemistry* 120, 105–125.
902 <https://doi.org/10.1016/j.soilbio.2018.01.030>

903 Burke, J.A., Lewis, K.L., Ritchie, G.L., DeLaune, P.B., Keeling, J.W., Acosta-Martinez, V., Moore, J.M.,
 904 McLendon, T., 2021. Net positive soil water content following cover crops with no tillage in irrigated semi-
 905 arid cotton production. *Soil and Tillage Research* 208, 104869. <https://doi.org/10.1016/j.still.2020.104869>

906 Cano, A., Núñez, A., Acosta-Martinez, V., Schipanski, M., Ghimire, R., Rice, C., West, C., 2018. Current
 907 knowledge and future research directions to link soil health and water conservation in the Ogallala Aquifer
 908 region. *Geoderma* 328, 109–118. <https://doi.org/10.1016/j.geoderma.2018.04.027>

909 Carlisle, L., 2016. Factors influencing farmer adoption of soil health practices in the United States: a
 910 narrative review. *Agroecology and Sustainable Food Systems* 40, 583–613.
 911 <https://doi.org/10.1080/21683565.2016.1156596>

912 Carlisle, L., Esquivel, K., Baur, P., Ichikawa, N.F., Olimpi, E.M., Ory, J., Waterhouse, H., Iles, A., Karp,
 913 D.S., Kremen, C., Bowles, T.M., 2022. Organic farmers face persistent barriers to adopting diversification
 914 practices in California’s Central Coast. *Agroecology and Sustainable Food Systems* 46, 1145–1172.
 915 <https://doi.org/10.1080/21683565.2022.2104420>

916 Carlson, S., Stockwell, R., 2013. Research priorities for advancing adoption of cover crops in agriculture-
 917 intensive regions. *Journal of Agriculture, Food Systems, and Community Development* 3, 125-129-125–
 918 129.

919 CDFA, 2022. Healthy Soils Program. URL <https://www.cdfa.ca.gov/oefi/healthysouls/>

920 Chagnon, P.-L., Bradley, R.L., Maherali, H., Klironomos, J.N., 2013. A trait-based framework to
 921 understand life history of mycorrhizal fungi. *Trends in Plant Science* 18, 484–491.
 922 <https://doi.org/10.1016/j.tplants.2013.05.001>

923 Chandrasekhar, P., Kreiselmeier, J., Schwen, A., Weninger, T., Julich, S., Feger, K.-H., Schwärzel, K.,
 924 2019. SPorDyn: A Python code for modeling the evolution of soil pore size distribution after tillage.
 925 *MethodsX* 6, 2118–2126. <https://doi.org/10.1016/j.mex.2019.09.014>

- 926 Chang, W., Sui, X., Fan, X.-X., Jia, T.-T., Song, F.-Q., 2018. Arbuscular Mycorrhizal Symbiosis Modulates
927 Antioxidant Response and Ion Distribution in Salt-Stressed *Elaeagnus angustifolia* Seedlings. *Front.*
928 *Microbiol.* 9, 652. <https://doi.org/10.3389/fmicb.2018.00652>
- 929 Chartzoulakis, K., Bertaki, M., 2015. Sustainable Water Management in Agriculture under Climate Change.
930 *Agriculture and Agricultural Science Procedia* 4, 88–98. <https://doi.org/10.1016/j.aaspro.2015.03.011>
- 931 Chauhan, H., Bagyaraj, D.J., Selvakumar, G., Sundaram, S.P., 2015. Novel plant growth promoting
932 rhizobacteria—Prospects and potential. *Applied Soil Ecology* 95, 38–53.
933 <https://doi.org/10.1016/j.apsoil.2015.05.011>
- 934 Chen, Y., Zhou, Y., Fang, S., Li, M., Wang, Y., Cao, K., 2022. Crop pattern optimization for the
935 coordination between economy and environment considering hydrological uncertainty. *Science of The*
936 *Total Environment* 809, 151152. <https://doi.org/10.1016/j.scitotenv.2021.151152>
- 937 Chiarelli, D.D., Passera, C., Rosa, L., Davis, K.F., D’Odorico, P., Rulli, M.C., 2020. The green and blue
938 crop water requirement WATNEEDS model and its global gridded outputs. *Sci Data* 7, 273.
939 <https://doi.org/10.1038/s41597-020-00612-0>
- 940 Chukalla, A.D., Krol, M.S., Hoekstra, A.Y., 2015. Green and blue water footprint reduction in irrigated
941 agriculture: effect of irrigation techniques, irrigation strategies and mulching. *Hydrol. Earth Syst. Sci.* 19,
942 4877–4891. <https://doi.org/10.5194/hess-19-4877-2015>
- 943 Cohen, A.C., Travaglia, C.N., Bottini, R., Piccoli, P.N., 2009. Participation of abscisic acid and gibberellins
944 produced by endophytic *Azospirillum* in the alleviation of drought effects in maize. *Botany* 87, 455–462.
945 <https://doi.org/10.1139/B09-023>
- 946 Cousins, O.H., Garnett, T.P., Rasmussen, A., Mooney, S.J., Smernik, R.J., Brien, C.J., Cavagnaro, T.R.,
947 2020. Variable water cycles have a greater impact on wheat growth and soil nitrogen response than constant
948 watering. *Plant Science* 290, 110146. <https://doi.org/10.1016/j.plantsci.2019.05.009>

949 Dahlke, H.E., Brown, A.G., Orloff, S., Putnam, D., O'Geen, T., 2018. Managed winter flooding of alfalfa
 950 recharges groundwater with minimal crop damage. *Calif Agr* 72, 1–11.
 951 <https://doi.org/10.3733/ca.2018a0001>

952 Dang, T.D., Chowdhury, A.F.M.K., Galelli, S., 2020. On the representation of water reservoir storage and
 953 operations in large-scale hydrological models: implications on model parameterization and climate change
 954 impact assessments. *Hydrol. Earth Syst. Sci.* 24, 397–416. <https://doi.org/10.5194/hess-24-397-2020>

955 Darnhofer, I., 2014. Resilience and why it matters for farm management. *European Review of Agricultural*
 956 *Economics* 41, 461–484. <https://doi.org/10.1093/erae/jbu012>

957 Daryanto, S., Fu, B., Wang, L., Jacinthe, P.-A., Zhao, W., 2018. Quantitative synthesis on the ecosystem
 958 services of cover crops. *Earth-Science Reviews* 185, 357–373.
 959 <https://doi.org/10.1016/j.earscirev.2018.06.013>

960 de Groot, R.S., Alkemade, R., Braat, L., Hein, L., Willemen, L., 2010. Challenges in integrating the concept
 961 of ecosystem services and values in landscape planning, management and decision making. *Ecological*
 962 *Complexity* 7, 260–272. <https://doi.org/10.1016/j.ecocom.2009.10.006>

963 de Jonge, L.W., Moldrup, P., Schjønning, P., 2009. Soil Infrastructure, Interfaces & Translocation
 964 Processes in Inner Space ("Soil-it-is"): towards a road map for the constraints and crossroads
 965 of soil architecture and biophysical processes. *Hydrol. Earth Syst. Sci.* 13, 1485–1502.
 966 <https://doi.org/10.5194/hess-13-1485-2009>

967 de Vries, F.T., Griffiths, R.I., Knight, C.G., Nicolitch, O., Williams, A., 2020. Harnessing rhizosphere
 968 microbiomes for drought-resilient crop production. *Science* 368, 270–274.
 969 <https://doi.org/10.1126/science.aaz5192>

970 DeVincentis, A., Solis, S.S., Rice, S., Zaccaria, D., Snyder, R., Maskey, M., Gomes, A., Gaudin, A.,
 971 Mitchell, J., 2022. Impacts of winter cover cropping on soil moisture and evapotranspiration in California's

972 specialty crop fields may be minimal during winter months. Calif Agr 1–9.
 973 <https://doi.org/10.3733/ca.2022a0001>

974 Devine, S., O’Geen, A. (Toby), 2019. Climate-smart management of soil water storage: statewide analysis
 975 of California perennial crops. Environ. Res. Lett. 14, 044021. <https://doi.org/10.1088/1748-9326/ab058c>

976 Dias, T., Dukes, A., Antunes, P.M., 2015. Accounting for soil biotic effects on soil health and crop
 977 productivity in the design of crop rotations. J. Sci. Food Agric. 95, 447–454.
 978 <https://doi.org/10.1002/jsfa.6565>

979 Dodd, I.C., Zinovkina, N.Y., Safronova, V.I., Belimov, A.A., 2010. Rhizobacterial mediation of plant
 980 hormone status. Annals of Applied Biology 157, 361–379. [https://doi.org/10.1111/j.1744-](https://doi.org/10.1111/j.1744-7348.2010.00439.x)
 981 [7348.2010.00439.x](https://doi.org/10.1111/j.1744-7348.2010.00439.x)

982 Dong, L., Zhang, W., Xiong, Y., Zou, J., Huang, Q., Xu, X., Ren, P., Huang, G., 2022. Impact of short-
 983 term organic amendments incorporation on soil structure and hydrology in semiarid agricultural lands.
 984 International Soil and Water Conservation Research 10, 457–469.
 985 <https://doi.org/10.1016/j.iswcr.2021.10.003>

986 Downer, J., Hodel, D., 2001. The effects of mulching on establishment of *Syagrus romanzoffiana* (Cham.)
 987 Becc., *Washingtonia robusta* H. Wendl. and *Archontophoenix cunninghamiana* (H. Wendl.) H. Wendl. &
 988 Drude in the landscape. Scientia Horticulturae 87, 85–92. [https://doi.org/10.1016/S0304-4238\(00\)00171-0](https://doi.org/10.1016/S0304-4238(00)00171-0)

989 Duan, X., Neuman, D.S., Reiber, J.M., Green, C.D., Saxton, A.M., Augé, R.M., 1996. Mycorrhizal
 990 influence on hydraulic and hormonal factors implicated in the control of stomatal conductance during
 991 drought. J Exp Bot 47, 1541–1550. <https://doi.org/10.1093/jxb/47.10.1541>

992 Duc, N.H., Csintalan, Z., Posta, K., 2018. Arbuscular mycorrhizal fungi mitigate negative effects of
 993 combined drought and heat stress on tomato plants. Plant Physiology and Biochemistry 132, 297–307.
 994 <https://doi.org/10.1016/j.plaphy.2018.09.011>

- 995 Dumont, A., Mayor, B., López-Gunn, E., 2013. Is the Rebound Effect or Jevons Paradox a Useful Concept
996 for better Management of Water Resources? Insights from the Irrigation Modernisation Process in Spain.
997 *Aquatic Procedia* 1, 64–76. <https://doi.org/10.1016/j.aqpro.2013.07.006>
- 998 Escriva-Bou, A., Hui, R., Maples, S., Medellín-Azuara, J., Harter, T., Lund, J.R., 2020. Planning for
999 groundwater sustainability accounting for uncertainty and costs: An application to California’s Central
1000 Valley. *Journal of Environmental Management* 264, 110426.
1001 <https://doi.org/10.1016/j.jenvman.2020.110426>
- 1002 Evenson, G., Osterholz, W.R., Shedekar, V.S., King, K., Mehan, S., Kalcic, M., 2022. Representing soil
1003 health practice effects on soil properties and nutrient loss in a watershed-scale hydrologic model. *J of Env*
1004 *Quality* jeq2.20338. <https://doi.org/10.1002/jeq2.20338>
- 1005 Falkenmark, M., Rockström, J., 2010. Building Water Resilience in the Face of Global Change: From a
1006 Blue-Only to a Green-Blue Water Approach to Land-Water Management. *J. Water Resour. Plann. Manage.*
1007 136, 606–610. [https://doi.org/10.1061/\(ASCE\)WR.1943-5452.0000118](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000118)
- 1008 Faunt, C.C., Sneed, M., Traum, J., Brandt, J.T., 2016. Water availability and land subsidence in the Central
1009 Valley, California, USA. *Hydrogeol J* 24, 675–684. <https://doi.org/10.1007/s10040-015-1339-x>
- 1010 Figueiredo, M.V.B., Martinez, C.R., Burity, H.A., Chanway, C.P., 2008. Plant growth-promoting
1011 rhizobacteria for improving nodulation and nitrogen fixation in the common bean (*Phaseolus vulgaris* L.).
1012 *World J Microbiol Biotechnol* 24, 1187–1193. <https://doi.org/10.1007/s11274-007-9591-4>
- 1013 Fischer, C., Tischer, J., Roscher, C., Eisenhauer, N., Ravenek, J., Gleixner, G., Attinger, S., Jensen, B., de
1014 Kroon, H., Mommer, L., Scheu, S., Hildebrandt, A., 2015. Plant species diversity affects infiltration
1015 capacity in an experimental grassland through changes in soil properties. *Plant Soil* 397, 1–16.
1016 <https://doi.org/10.1007/s11104-014-2373-5>

- 1017 Flint, L.E., Flint, A.L., Mendoza, J., Kalansky, J., Ralph, F.M., 2018. Characterizing drought in California:
1018 new drought indices and scenario-testing in support of resource management. *Ecol Process* 7, 1.
1019 <https://doi.org/10.1186/s13717-017-0112-6>
- 1020 Flint, L.E., Flint, A.L., Thorne, J.H., Boynton, R., 2013. Fine-scale hydrologic modeling for regional
1021 landscape applications: the California Basin Characterization Model development and performance. *Ecol*
1022 *Process* 2, 25. <https://doi.org/10.1186/2192-1709-2-25>
- 1023 Folke, C., Carpenter, S.R., Walker, B., Scheffer, M., Chapin, T., Rockström, J., 2010. Resilience Thinking:
1024 Integrating Resilience, Adaptability and Transformability. *E&S* 15, art20. [https://doi.org/10.5751/ES-](https://doi.org/10.5751/ES-03610-150420)
1025 [03610-150420](https://doi.org/10.5751/ES-03610-150420)
- 1026 Fonte, S.J., Barrios, E., Six, J., 2010. Earthworms, soil fertility and aggregate-associated soil organic matter
1027 dynamics in the Quesungual agroforestry system. *Geoderma* 155, 320–328.
1028 <https://doi.org/10.1016/j.geoderma.2009.12.016>
- 1029 Franzluebbers, A., 2007. Soil Physical Aspects of Integrated Crop-Livestock Systems.
- 1030 Gabriel, J.L., Muñoz-Carpena, R., Quemada, M., 2012. The role of cover crops in irrigated systems: Water
1031 balance, nitrate leaching and soil mineral nitrogen accumulation. *Agriculture, Ecosystems & Environment*
1032 155, 50–61. <https://doi.org/10.1016/j.agee.2012.03.021>
- 1033 Galdos, M.V., Pires, L.F., Cooper, H.V., Calonego, J.C., Rosolem, C.A., Mooney, S.J., 2019. Assessing
1034 the long-term effects of zero-tillage on the macroporosity of Brazilian soils using X-ray Computed
1035 Tomography. *Geoderma* 337, 1126–1135. <https://doi.org/10.1016/j.geoderma.2018.11.031>
- 1036 García-Tejera, O., López-Bernal, Á., Orgaz, F., Testi, L., Villalobos, F.J., 2018. Are olive root systems
1037 optimal for deficit irrigation? *European Journal of Agronomy* 99, 72–79.
1038 <https://doi.org/10.1016/j.eja.2018.06.012>

- 1039 Garret, Amy, 2019. Dry Farming in the Maritime Pacific Northwest: Intro to Dry Farming Organic
1040 Vegetables. <https://catalog.extension.oregonstate.edu/sites/catalog/files/project/pdf/em9229.pdf>
- 1041 Gaudin, A.C.M., Tolhurst, T.N., Ker, A.P., Janovicek, K., Tortora, C., Martin, R.C., Deen, W., 2015.
1042 Increasing Crop Diversity Mitigates Weather Variations and Improves Yield Stability. PLoS ONE 10,
1043 e0113261. <https://doi.org/10.1371/journal.pone.0113261>
- 1044 Gilbert, M.E., Medina, V., 2016. Drought Adaptation Mechanisms Should Guide Experimental Design.
1045 Trends in Plant Science 21, 639–647. <https://doi.org/10.1016/j.tplants.2016.03.003>
- 1046 Glick, B.R., Cheng, Z., Czarny, J., Duan, J., 2007. Promotion of plant growth by ACC deaminase-producing
1047 soil bacteria. Eur J Plant Pathol 119, 329–339. <https://doi.org/10.1007/s10658-007-9162-4>
- 1048 Glick, B.R., Penrose, D.M., Li, J., 1998. A Model For the Lowering of Plant Ethylene Concentrations by
1049 Plant Growth-promoting Bacteria. Journal of Theoretical Biology 190, 63–68.
1050 <https://doi.org/10.1006/jtbi.1997.0532>
- 1051 Gómez, J.A., Ben-Gal, A., Alarcón, J.J., De Lannoy, G., de Roos, S., Dostál, T., Fereres, E., Intrigliolo,
1052 D.S., Krása, J., Klik, A., Liebhard, G., Nolz, R., Peeters, A., Plaas, E., Quinton, J.N., Miao, R., Strauss, P.,
1053 Xu, W., Zhang, Z., Zhong, F., Zumr, D., Dodd, I.C., 2020. SHui, an EU-Chinese cooperative project to
1054 optimize soil and water management in agricultural areas in the XXI century. International Soil and Water
1055 Conservation Research 8, 1–14. <https://doi.org/10.1016/j.iswcr.2020.01.001>
- 1056 Grover, M., Madhubala, R., Ali, Sk.Z., Yadav, S.K., Venkateswarlu, B., 2014. Influence of *Bacillus* spp.
1057 strains on seedling growth and physiological parameters of sorghum under moisture stress conditions:
1058 *Bacillus* spp. strains influence sorghum under moisture stress. J. Basic Microbiol. 54, 951–961.
1059 <https://doi.org/10.1002/jobm.201300250>
- 1060 Gupta, R., Tyagi, N.K., Abrol, I., 2020. Rainwater Management and Indian Agriculture: A Call for a Shift
1061 in Focus from Blue to Green Water. Agric Res 9, 429–443. <https://doi.org/10.1007/s40003-020-00467-2>

- 1062 Guzman, A., Montes, M., Hutchins, L., DeLaCerde, G., Yang, P., Kakouridis, A., Dahlquist-Willard, R.M.,
1063 Firestone, M.K., Bowles, T., Kremen, C., 2021. Crop diversity enriches arbuscular mycorrhizal fungal
1064 communities in an intensive agricultural landscape. *New Phytol* 231, 447–459.
1065 <https://doi.org/10.1111/nph.17306>
- 1066 Hamilton, A.J., Stagnitti, F., Xiong, X., Kreidl, S.L., Benke, K.K., Maher, P., 2007. Wastewater Irrigation:
1067 The State of Play. *Vadose Zone Journal* 6, 823–840. <https://doi.org/10.2136/vzj2007.0026>
- 1068 Han, E., Kautz, T., Perkons, U., Uteau, D., Peth, S., Huang, N., Horn, R., Köpke, U., 2015. Root growth
1069 dynamics inside and outside of soil biopores as affected by crop sequence determined with the profile wall
1070 method. *Biol Fertil Soils* 51, 847–856. <https://doi.org/10.1007/s00374-015-1032-1>
- 1071 Han, M., Zhao, C., Šimůnek, J., Feng, G., 2015. Evaluating the impact of groundwater on cotton growth
1072 and root zone water balance using Hydrus-1D coupled with a crop growth model. *Agricultural Water*
1073 *Management* 160, 64–75. <https://doi.org/10.1016/j.agwat.2015.06.028>
- 1074 Hanson, B., Orloff, S., Peters, D., 2000. Monitoring soil moisture helps refine irrigation management.
1075 *California Agriculture* 54, 38–42.
- 1076 Harter, T., 2015. California's agricultural regions gear up to actively manage groundwater use and
1077 protection. *California Agriculture* 69, 193–201.
- 1078 Hatfield, J.L., Boote, K.J., Kimball, B.A., Ziska, L.H., Izaurralde, R.C., Ort, D., Thomson, A.M., Wolfe,
1079 D., 2011. Climate Impacts on Agriculture: Implications for Crop Production. *Agron. J.* 103, 351–370.
1080 <https://doi.org/10.2134/agronj2010.0303>
- 1081 Hoekstra, A.Y., 2019. Green-blue water accounting in a soil water balance. *Advances in Water Resources*
1082 129, 112–117. <https://doi.org/10.1016/j.advwatres.2019.05.012>
- 1083 Howitt, R., MacEwan, D., Medellín-Azuara, J., Lund, J., Sumner, D., 2015. Economic analysis of the 2015
1084 drought for California agriculture. UC-Davis Center for Watershed Sciences.

- 1085 Huang, N., Athmann, M., Han, E., 2020. Biopore-Induced Deep Root Traits of Two Winter Crops.
1086 Agriculture 10, 634. <https://doi.org/10.3390/agriculture10120634>
- 1087 Huang, Z., Hejazi, M., Tang, Q., Vernon, C.R., Liu, Y., Chen, M., Calvin, K., 2019. Global agricultural
1088 green and blue water consumption under future climate and land use changes. Journal of Hydrology 574,
1089 242–256. <https://doi.org/10.1016/j.jhydrol.2019.04.046>
- 1090 Hudson, B.D., 1984. Soil organic matter and available water capacity. Journal of Soil and Water
1091 Conservation 49, 189.
- 1092 Hunink, J., Vila, M., Baille, A., 2011. REDSIM: Approach to soil water modelling.
- 1093 Iizumi, T., Wagai, R., 2019. Leveraging drought risk reduction for sustainable food, soil and climate via
1094 soil organic carbon sequestration. Sci Rep 9, 19744. <https://doi.org/10.1038/s41598-019-55835-y>
- 1095 Iqbal, R., Raza, M.A.S., Saleem, M.F., Khan, I.H., Ahmad, S., Zaheer, M.S., Aslam, M.U., Haider, I., 2019.
1096 Physiological and biochemical appraisal for mulching and partial rhizosphere drying of cotton. J. Arid Land
1097 11, 785–794. <https://doi.org/10.1007/s40333-019-0014-9>
- 1098 Iwanaga, T., Partington, D., Ticehurst, J., Croke, B.F.W., Jakeman, A.J., 2020. A socio-environmental
1099 model for exploring sustainable water management futures: Participatory and collaborative modelling in
1100 the Lower Campaspe catchment. Journal of Hydrology: Regional Studies 28, 100669.
1101 <https://doi.org/10.1016/j.ejrh.2020.100669>
- 1102 Jackson, L.E., Koch, G.W., 1997. The Ecophysiology of Crops and Their Wild Relatives, in: Ecology in
1103 Agriculture. Elsevier, pp. 3–37. <https://doi.org/10.1016/B978-012378260-1/50002-6>
- 1104 Jägermeyr, J., Gerten, D., Schaphoff, S., Heinke, J., Lucht, W., Rockström, J., 2016. Integrated crop water
1105 management might sustainably halve the global food gap. Environ. Res. Lett. 11, 025002.
1106 <https://doi.org/10.1088/1748-9326/11/2/025002>

- 1107 Jahanzad, E., Holtz, B.A., Zuber, C.A., Doll, D., Brewer, K.M., Hogan, S., Gaudin, A.C.M., 2020. Orchard
1108 recycling improves climate change adaptation and mitigation potential of almond production systems. PLoS
1109 ONE 15, e0229588. <https://doi.org/10.1371/journal.pone.0229588>
- 1110 Jasechko, S., Perrone, D., 2021. Global groundwater wells at risk of running dry. Science 372, 418–421.
1111 <https://doi.org/10.1126/science.abc2755>
- 1112 Jensen, J.L., Schjønning, P., Watts, C.W., Christensen, B.T., Munkholm, L.J., 2019. Soil Water Retention:
1113 Uni-Modal Models of Pore-Size Distribution Neglect Impacts of Soil Management. Soil Science Society
1114 of America Journal 83, 18–26. <https://doi.org/10.2136/sssaj2018.06.0238>
- 1115 Jewitt, G., 2006. Integrating blue and green water flows for water resources management and planning.
1116 Physics and Chemistry of the Earth, Parts A/B/C 31, 753–762. <https://doi.org/10.1016/j.pce.2006.08.033>
- 1117 Jimenez, B., Asano, T., 2015. Water Reuse: An International Survey of current practice, issues and needs.
1118 Water Intelligence Online 7, 9781780401881–9781780401881. <https://doi.org/10.2166/9781780401881>
- 1119 Jokela, D., Nair, A., 2016. Effects of reduced tillage and fertilizer application method on plant growth,
1120 yield, and soil health in organic bell pepper production. Soil and Tillage Research 163, 243–254.
1121 <https://doi.org/10.1016/j.still.2016.06.010>
- 1122 Jones, C., Olson-Rutz, K., Miller, P., Zabinski, C., 2020. Cover Crop Management in Semi-Arid Regions:
1123 Effect on Soil and Cash Crop. Crops & Soils Mag. 53, 42–51. <https://doi.org/10.1002/crso.20065>
- 1124 Jouma, N., Dadaser-Celik, F., 2021. Assessing hydrologic alterations due to reservoirs and intensified
1125 irrigation in a semi-arid agricultural river basin using SWAT *. Irrig. and Drain. ird.2661.
1126 <https://doi.org/10.1002/ird.2661>
- 1127 Kane, D.A., Bradford, M.A., Fuller, E., Oldfield, E.E., Wood, S.A., 2021. Soil organic matter protects US
1128 maize yields and lowers crop insurance payouts under drought. Environ. Res. Lett. 16, 044018.
1129 <https://doi.org/10.1088/1748-9326/abe492>

- 1130 Karandish, F., Šimůnek, J., 2019. A comparison of the HYDRUS (2D/3D) and SALTMed models to
1131 investigate the influence of various water-saving irrigation strategies on the maize water footprint.
1132 *Agricultural Water Management* 213, 809–820. <https://doi.org/10.1016/j.agwat.2018.11.023>
- 1133 Keesstra, S., Sannigrahi, S., López-Vicente, M., Pulido, M., Novara, A., Visser, S., Kalantari, Z., 2021. The
1134 role of soils in regulation and provision of blue and green water. *Phil. Trans. R. Soc. B* 376, 20200175.
1135 <https://doi.org/10.1098/rstb.2020.0175>
- 1136 Kemanian, A.R., Julich, S., Manoranjan, V.S., Arnold, J.R., 2011. Integrating soil carbon cycling with that
1137 of nitrogen and phosphorus in the watershed model SWAT: Theory and model testing. *Ecological*
1138 *Modelling* 222, 1913–1921. <https://doi.org/10.1016/j.ecolmodel.2011.03.017>
- 1139 Khan, T., Nouri, H., Booi, M., Hoekstra, A., Khan, H., Ullah, I., 2021. Water Footprint, Blue Water
1140 Scarcity, and Economic Water Productivity of Irrigated Crops in Peshawar Basin, Pakistan. *Water* 13, 1249.
1141 <https://doi.org/10.3390/w13091249>
- 1142 Kim, S.G., Khan, Z., Jeon, Y.H., Kim, Y.H., 2009. Inhibitory Effect of *Paenibacillus polymyxa* GBR-462
1143 on *Phytophthora capsici* Causing Phytophthora Blight in Chili Pepper. *Journal of Phytopathology* 157,
1144 329–337. <https://doi.org/10.1111/j.1439-0434.2008.01490.x>
- 1145 King, A.E., Ali, G.A., Gillespie, A.W., Wagner-Riddle, C., 2020. Soil Organic Matter as Catalyst of Crop
1146 Resource Capture. *Front. Environ. Sci.* 8, 50. <https://doi.org/10.3389/fenvs.2020.00050>
- 1147 Kinoshita, R., Schindelbeck, R.R., van Es, H.M., 2017. Quantitative soil profile-scale assessment of the
1148 sustainability of long-term maize residue and tillage management. *Soil and Tillage Research* 174, 34–44.
1149 <https://doi.org/10.1016/j.still.2017.05.010>
- 1150 Kocis, T.N., Dahlke, H.E., 2017. Availability of high-magnitude streamflow for groundwater banking in
1151 the Central Valley, California. *Environ. Res. Lett.* 12, 084009. <https://doi.org/10.1088/1748-9326/aa7b1b>

- 1152 Kodešová, R., Vignozzi, N., Rohošková, M., Hájková, T., Kočárek, M., Pagliai, M., Kozák, J., Šimůnek,
1153 J., 2009. Impact of varying soil structure on transport processes in different diagnostic horizons of three
1154 soil types. *Journal of Contaminant Hydrology* 104, 107–125. <https://doi.org/10.1016/j.jconhyd.2008.10.008>
- 1155 Kong, A.Y.Y., Six, J., 2010. Tracing Root vs. Residue Carbon into Soils from Conventional and Alternative
1156 Cropping Systems. *Soil Sci. Soc. Am. J.* 74, 1201–1210. <https://doi.org/10.2136/sssaj2009.0346>
- 1157 Kroes, J., Supit, I., Van Dam, J., Van Walsum, P., Mulder, M., 2017. Impact of capillary rise and
1158 recirculation on crop yields (preprint). *Vadose Zone Hydrology/Modelling approaches*.
1159 <https://doi.org/10.5194/hess-2017-223>
- 1160 Lazcano, C., Barrios-Masias, F.H., Jackson, L.E., 2014. Arbuscular mycorrhizal effects on plant water
1161 relations and soil greenhouse gas emissions under changing moisture regimes. *Soil Biology and*
1162 *Biochemistry* 74, 184–192. <https://doi.org/10.1016/j.soilbio.2014.03.010>
1163
- 1164 Kallenbach, C., Grandy, A.S., 2011. Controls over soil microbial biomass responses to carbon amendments
1165 in agricultural systems: A meta-analysis. *Agriculture, Ecosystems & Environment* 144, 241–252.
1166 <https://doi.org/10.1016/j.agee.2011.08.020>
- 1167 Leap, J., Yogg-Comerchero, Kirstin, Wong, D., 2017. Organic mixed production blocks for direct markets
1168 on California's Central Coast: A Guide for Beginning Specialty Crop Growers.
1169 <https://agroecology.ucsc.edu/about/publications/grower-guides/pdf-downloads/csa-production-blocks.pdf>
- 1170 Lei, C., Wagner, P.D., Fohrer, N., 2021. Influences of land use changes on the dynamics of water quantity
1171 and quality in the German lowland catchment of the Stör (preprint). *Catchment hydrology/Modelling*
1172 *approaches*. <https://doi.org/10.5194/hess-2021-476>
- 1173 Lepsch, H.C., Brown, P.H., Peterson, C.A., Gaudin, A.C.M., Khalsa, S.D.S., 2019. Impact of organic matter
1174 amendments on soil and tree water status in a California orchard. *Agricultural Water Management* 222,
1175 204–212. <https://doi.org/10.1016/j.agwat.2019.06.002>

- 1176 Li, J., Meng, B., Chai, H., Yang, X., Song, W., Li, S., Lu, A., Zhang, T., Sun, W., 2019. Arbuscular
1177 Mycorrhizal Fungi Alleviate Drought Stress in C3 (*Leymus chinensis*) and C4 (*Hemarthria altissima*)
1178 Grasses via Altering Antioxidant Enzyme Activities and Photosynthesis. *Front. Plant Sci.* 10, 499.
1179 <https://doi.org/10.3389/fpls.2019.00499>
- 1180 Li, M., Schmidt, J.E., LaHue, D.G., Lazicki, P., Kent, A., Machmuller, M.B., Scow, K.M., Gaudin, A.C.M.,
1181 2020. Impact of Irrigation Strategies on Tomato Root Distribution and Rhizosphere Processes in an Organic
1182 System. *Front. Plant Sci.* 11, 360. <https://doi.org/10.3389/fpls.2020.00360>
- 1183 Li, Q., Dong, B., Qiao, Y., Liu, M., Zhang, J., 2010. Root growth, available soil water, and water-use
1184 efficiency of winter wheat under different irrigation regimes applied at different growth stages in North
1185 China. *Agricultural Water Management* 97, 1676–1682. <https://doi.org/10.1016/j.agwat.2010.05.025>
- 1186 Lipan, L., Martín-Palomo, M.J., Sánchez-Rodríguez, L., Cano-Lamadrid, M., Sendra, E., Hernández, F.,
1187 Burló, F., Vázquez-Araújo, L., Andreu, L., Carbonell-Barrachina, Á.A., 2019. Almond fruit quality can be
1188 improved by means of deficit irrigation strategies. *Agricultural Water Management* 217, 236–242.
1189 <https://doi.org/10.1016/j.agwat.2019.02.041>
- 1190 López-Moreno, J.I., Gascoin, S., Herrero, J., Sproles, E.A., Pons, M., Alonso-González, E., Hanich, L.,
1191 Boudhar, A., Musselman, K.N., Molotch, N.P., Sickman, J., Pomeroy, J., 2017. Different sensitivities of
1192 snowpacks to warming in Mediterranean climate mountain areas. *Environ. Res. Lett.* 12, 074006.
1193 <https://doi.org/10.1088/1748-9326/aa70cb>
- 1194 Lotter, D.W., Seidel, R., Liebhardt, W., 2003. The performance of organic and conventional cropping
1195 systems in an extreme climate year. *Am J Alt Ag* 18, 146–154. <https://doi.org/10.1079/AJAA200345>
- 1196 Luan, X.-B., Yin, Y.-L., Wu, P.-T., Sun, S.-K., Wang, Y.-B., Gao, X.-R., Liu, J., 2018. An improved
1197 method for calculating the regional crop water footprint based on a hydrological process analysis. *Hydrol.*
1198 *Earth Syst. Sci.* 22, 5111–5123. <https://doi.org/10.5194/hess-22-5111-2018>

- 1199 Lubell, M., Blomquist, W., Beutler, L., 2020. Sustainable Groundwater Management in California: A Grand
 1200 Experiment in Environmental Governance. *Society & Natural Resources* 33, 1447–1467.
 1201 <https://doi.org/10.1080/08941920.2020.1833617>
- 1202 Lutz, F., Herzfeld, T., Heinke, J., Rolinski, S., Schaphoff, S., von Bloh, W., Stoorvogel, J.J., Müller, C.,
 1203 2019. Simulating the effect of tillage practices with the global ecosystem model LPJmL (version 5.0-
 1204 tillage). *Geosci. Model Dev.* 12, 2419–2440. <https://doi.org/10.5194/gmd-12-2419-2019>
- 1205 Maharjan, G.R., Prescher, A.-K., Nendel, C., Ewert, F., Mboh, C.M., Gaiser, T., Seidel, S.J., 2018.
 1206 Approaches to model the impact of tillage implements on soil physical and nutrient properties in different
 1207 agro-ecosystem models. *Soil and Tillage Research* 180, 210–221.
 1208 <https://doi.org/10.1016/j.still.2018.03.009>
- 1209 Mallmann, F.J.K., Rheinheimer, D. dos S., Ceretta, C.A., Cella, C., Minella, J.P.G., Guma, R.L., Filipović,
 1210 V., van Oort, F., Šimůnek, J., 2014. Soil tillage to reduce surface metal contamination – model development
 1211 and simulations of zinc and copper concentration profiles in a pig slurry-amended soil. *Agriculture,*
 1212 *Ecosystems & Environment* 196, 59–68. <https://doi.org/10.1016/j.agee.2014.06.024>
- 1213 Mardhiah, U., Caruso, T., Gurnell, A., Rillig, M.C., 2016. Arbuscular mycorrhizal fungal hyphae reduce
 1214 soil erosion by surface water flow in a greenhouse experiment. *Applied Soil Ecology* 99, 137–140.
 1215 <https://doi.org/10.1016/j.apsoil.2015.11.027>
- 1216 Mariotte, P., Mehrabi, Z., Bezemer, T.M., De Deyn, G.B., Kulmatiski, A., Drigo, B., Veen, G.F. (Ciska),
 1217 van der Heijden, M.G.A., Kardol, P., 2018. Plant–Soil Feedback: Bridging Natural and Agricultural
 1218 Sciences. *Trends in Ecology & Evolution* 33, 129–142. <https://doi.org/10.1016/j.tree.2017.11.005>
- 1219 Martínez, E., Fuentes, J.-P., Silva, P., Valle, S., Acevedo, E., 2008. Soil physical properties and wheat root
 1220 growth as affected by no-tillage and conventional tillage systems in a Mediterranean environment of Chile.
 1221 *Soil and Tillage Research* 99, 232–244. <https://doi.org/10.1016/j.still.2008.02.001>

- 1222 Massoud, E.C., Purdy, A.J., Miro, M.E., Famiglietti, J.S., 2018. Projecting groundwater storage changes in
1223 California's Central Valley. *Sci Rep* 8, 12917. <https://doi.org/10.1038/s41598-018-31210-1>
- 1224 Mayak, S., Tirosh, T., Glick, B.R., 2004. Plant growth-promoting bacteria that confer resistance to water
1225 stress in tomatoes and peppers. *Plant Science* 166, 525–530. <https://doi.org/10.1016/j.plantsci.2003.10.025>
- 1226 McCallum, M.H., Kirkegaard, J.A., Green, T.W., Cresswell, H.P., Davies, S.L., Angus, J.F., Peoples, M.B.,
1227 2004. Improved subsoil macroporosity following perennial pastures. *Aust. J. Exp. Agric.* 44, 299.
1228 <https://doi.org/10.1071/EA03076>
- 1229 McDaniel, M.D., Grandy, A.S., Tiemann, L.K., Weintraub, M.N., 2016. Eleven years of crop
1230 diversification alters decomposition dynamics of litter mixtures incubated with soil. *Ecosphere* 7.
1231 <https://doi.org/10.1002/ecs2.1426>
- 1232 Meier, J., Zabel, F., Mauser, W., 2018. A global approach to estimate irrigated areas – a comparison
1233 between different data and statistics. *Hydrol. Earth Syst. Sci.* 22, 1119–1133. [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-22-1119-2018)
1234 [22-1119-2018](https://doi.org/10.5194/hess-22-1119-2018)
- 1235 Meurer, K., Barron, J., Chenu, C., Coucheney, E., Fielding, M., Hallett, P., Herrmann, A.M., Keller, T.,
1236 Koestel, J., Larsbo, M., Lewan, E., Or, D., Parsons, D., Parvin, N., Taylor, A., Vereecken, H., Jarvis, N.,
1237 2020. A framework for modelling soil structure dynamics induced by biological activity. *Glob. Change*
1238 *Biol.* 26, 5382–5403. <https://doi.org/10.1111/gcb.15289>
- 1239 Meyer, N., Bergez, J.-E., Constantin, J., Justes, E., 2019a. Cover crops reduce water drainage in temperate
1240 climates: A meta-analysis. *Agron. Sustain. Dev.* 39, 3. <https://doi.org/10.1007/s13593-018-0546-y>
- 1241 Meyer, N., Bergez, J.-E., Constantin, J., Justes, E., 2019b. Cover crops reduce water drainage in temperate
1242 climates: A meta-analysis. *Agron. Sustain. Dev.* 39, 3. <https://doi.org/10.1007/s13593-018-0546-y>
- 1243 Minasny, B., McBratney, A.B., 2018. Limited effect of organic matter on soil available water capacity. *Eur*
1244 *J Soil Sci* 69, 39–47. <https://doi.org/10.1111/ejss.12475>

- 1245 Mitchell, J.P., Shrestha, A., Irmak, S., 2015. Trade-offs between winter cover crop production and soil
1246 water depletion in the San Joaquin Valley, California. *Journal of Soil and Water Conservation* 70, 430–
1247 440. <https://doi.org/10.2489/jswc.70.6.430>
- 1248 Mitchell, J.P., Singh, P.N., Wallender, W.W., Munk, D.S., Wroble, J.F., Horwath, W.R., Hogan, P., Roy,
1249 R., Hanson, B.R., 2012. No-tillage and high-residue practices reduce soil water evaporation. *Cal Ag* 66,
1250 55–61. <https://doi.org/10.3733/ca.v066n02p55>
- 1251 Moreno-Espíndola, I.P., Ferrara-Guerrero, M.J., Luna-Guido, M.L., Ramírez-Villanueva, D.A., De León-
1252 Lorenzana, A.S., Gómez-Acata, S., González-Terreros, E., Ramírez-Barajas, B., Navarro-Noya, Y.E.,
1253 Sánchez-Rodríguez, L.M., Fuentes-Ponce, M., Macedas-Jimenez, J.U., Dendooven, L., 2018. The Bacterial
1254 Community Structure and Microbial Activity in a Traditional Organic Milpa Farming System Under
1255 Different Soil Moisture Conditions. *Front. Microbiol.* 9, 2737. <https://doi.org/10.3389/fmicb.2018.02737>
- 1256 Morris, K.S., Bucini, G., 2016. California’s drought as opportunity: Redesigning U.S. agriculture for a
1257 changing climate. *Elem Sci Anth* 4, 000142. <https://doi.org/10.12952/journal.elementa.000142>
- 1258 Mosquera, G.M., Crespo, P., Breuer, L., Feyen, J., Windhorst, D., 2020. Water transport and tracer mixing
1259 in volcanic ash soils at a tropical hillslope: A wet layered sloping sponge. *Hydrological Processes* 34, 2032–
1260 2047. <https://doi.org/10.1002/hyp.13733>
- 1261 Msigwa, A., Chawanda, C.J., Komakech, H.C., Nkwasa, A., van Griensven, A., 2022. Representation of
1262 seasonal land use dynamics in SWAT+ for improved assessment of blue and green water consumption.
1263 *Hydrol. Earth Syst. Sci.* 26, 4447–4468. <https://doi.org/10.5194/hess-26-4447-2022>
- 1264 Munkholm, L.J., Heck, R.J., Deen, B., 2013. Long-term rotation and tillage effects on soil structure and
1265 crop yield. *Soil and Tillage Research* 127, 85–91. <https://doi.org/10.1016/j.still.2012.02.007>
- 1266 N. L. Klocke, R. S. Currie, R. M. Aiken, 2009. Soil Water Evaporation and Crop Residues. *Transactions*
1267 *of the ASABE* 52, 103–110. <https://doi.org/10.13031/2013.25951>

- 1268 Nabhan, G.P., McKibben, B., 2013. Growing Food in a Hotter, Drier Land: Lessons from Desert Farmers
1269 on Adapting to Climate Uncertainty. Chelsea Green Pub.
- 1270 Naseem, H., Bano, A., 2014. Role of plant growth-promoting rhizobacteria and their exopolysaccharide in
1271 drought tolerance of maize. *Journal of Plant Interactions* 9, 689–701.
1272 <https://doi.org/10.1080/17429145.2014.902125>
- 1273 Naveed, M., Mitter, B., Reichenauer, T.G., Wiczorek, K., Sessitsch, A., 2014. Increased drought stress
1274 resilience of maize through endophytic colonization by Burkholderia phytofirmans PsJN and Enterobacter
1275 sp. FD17. *Environmental and Experimental Botany* 97, 30–39.
1276 <https://doi.org/10.1016/j.envexpbot.2013.09.014>
- 1277 Ngumbi, E., Kloepper, J., 2016. Bacterial-mediated drought tolerance: Current and future prospects.
1278 *Applied Soil Ecology* 105, 109–125. <https://doi.org/10.1016/j.apsoil.2016.04.009>
- 1279 Nichols, V., Verhulst, N., Cox, R., Govaerts, B., 2015. Weed dynamics and conservation agriculture
1280 principles: A review. *Field Crops Research* 183, 56–68. <https://doi.org/10.1016/j.fcr.2015.07.012>
- 1281 Nielsen, D.C., Vigil, M.F., 2005. Legume Green Fallow Effect on Soil Water Content at Wheat Planting
1282 and Wheat Yield. *Agron.j.* 97, 684–689. <https://doi.org/10.2134/agronj2004.0071>
- 1283 Niles, M.T., Hammond Wagner, C.R., 2019. The carrot or the stick? Drivers of California farmer support
1284 for varying groundwater management policies. *Environ. Res. Commun.* 1, 045001.
1285 <https://doi.org/10.1088/2515-7620/ab1778>
- 1286 Niswonger, R.G., Morway, E.D., Triana, E., Huntington, J.L., 2017. Managed aquifer recharge through off-
1287 season irrigation in agricultural regions. *Water Resour. Res.* 53, 6970–6992.
1288 <https://doi.org/10.1002/2017WR020458>

- 1289 Nouri, Hamideh, Chavoshi Borujeni, S., Hoekstra, A.Y., 2019. The blue water footprint of urban green
1290 spaces: An example for Adelaide, Australia. *Landscape and Urban Planning* 190, 103613.
1291 <https://doi.org/10.1016/j.landurbplan.2019.103613>
- 1292 Nouri, H., Stokvis, B., Galindo, A., Blatchford, M., Hoekstra, A.Y., 2019. Water scarcity alleviation
1293 through water footprint reduction in agriculture: The effect of soil mulching and drip irrigation. *Science of*
1294 *The Total Environment* 653, 241–252. <https://doi.org/10.1016/j.scitotenv.2018.10.311>
- 1295 Nunes, M.R., Karlen, D.L., Denardin, J.E., Cambardella, C.A., 2019. Corn root and soil health indicator
1296 response to no-till production practices. *Agriculture, Ecosystems & Environment* 285, 106607.
1297 <https://doi.org/10.1016/j.agee.2019.106607>
- 1298 Nziguheba, G., Palm, C.A., Berhe, T., Denning, G., Dicko, A., Diouf, O., Diru, W., Flor, R., Frimpong, F.,
1299 Harawa, R., Kaya, B., Manumbu, E., McArthur, J., Mutuo, P., Ndiaye, M., Niang, A., Nkhoma, P., Nyadzi,
1300 G., Sachs, J., Sullivan, C., Teklu, G., Tobe, L., Sanchez, P.A., 2010. The African Green Revolution, in:
1301 *Advances in Agronomy*. Elsevier, pp. 75–115. <https://doi.org/10.1016/B978-0-12-385040-9.00003-7>
- 1302 O'Brien, P.L., Daigh, A.L.M., 2019. Tillage practices alter the surface energy balance – A review. *Soil and*
1303 *Tillage Research* 195, 104354. <https://doi.org/10.1016/j.still.2019.104354>
- 1304 Oehl, F., Sieverding, E., Mader, P., Dubois, D., Ineichen, K., Boller, T., Wiemken, A., 2004. Impact of
1305 long-term conventional and organic farming on the diversity of arbuscular mycorrhizal fungi. *Oecologia*
1306 138, 574–583. <https://doi.org/10.1007/s00442-003-1458-2>
- 1307 O'Geen, A.T., Saal, M., Dahlke, H., Doll, D., Elkins, R., Fulton, A., Fogg, G., Harter, T., Hopmans, J.W.,
1308 Ingels, C., Niederholzer, F., Solis, S.S., Verdegaal, P., Walkinshaw, M., 2015. Soil suitability index
1309 identifies potential areas for groundwater banking on agricultural lands. *Cal Ag* 69, 75–84.
1310 <https://doi.org/10.3733/ca.v069n02p75>
- 1311 Ostrom, E., 1990. *Governing the commons: The evolution of institutions for collective action*. Cambridge
1312 university press.

- 1313 Pahlavan-Rad, M.R., Dahmardeh, K., Hadizadeh, M., Keykha, G., Mohammadnia, N., Gangali, M., Keikha,
1314 M., Davatgar, N., Brungard, C., 2020. Prediction of soil water infiltration using multiple linear regression
1315 and random forest in a dry flood plain, eastern Iran. CATENA 194, 104715.
1316 <https://doi.org/10.1016/j.catena.2020.104715>
- 1317 Parihar, C.M., Nayak, H.S., Rai, V.K., Jat, S.L., Parihar, N., Aggarwal, P., Mishra, A.K., 2019. Soil water
1318 dynamics, water productivity and radiation use efficiency of maize under multi-year conservation
1319 agriculture during contrasting rainfall events. Field Crops Research 241, 107570.
1320 <https://doi.org/10.1016/j.fcr.2019.107570>
- 1321 Pathak, T., Maskey, M., Dahlberg, J., Kearns, F., Bali, K., Zaccaria, D., 2018. Climate Change Trends and
1322 Impacts on California Agriculture: A Detailed Review. Agronomy 8, 25.
1323 <https://doi.org/10.3390/agronomy8030025>
- 1324 Paustian, K., Andrén, O., Janzen, H.H., Lal, R., Smith, P., Tian, G., Tiessen, H., Noordwijk, M., Woomer,
1325 P.L., 1997. Agricultural soils as a sink to mitigate CO₂ emissions. Soil Use & Management 13, 230–244.
1326 <https://doi.org/10.1111/j.1475-2743.1997.tb00594.x>
- 1327 Pedranzani, H., Rodríguez-Rivera, M., Gutiérrez, M., Porcel, R., Hause, B., Ruiz-Lozano, J.M., 2016.
1328 Arbuscular mycorrhizal symbiosis regulates physiology and performance of *Digitaria eriantha* plants
1329 subjected to abiotic stresses by modulating antioxidant and jasmonate levels. Mycorrhiza 26, 141–152.
1330 <https://doi.org/10.1007/s00572-015-0653-4>
- 1331 Peng, Z., Wang, L., Xie, J., Li, L., Coulter, J.A., Zhang, R., Luo, Z., Cai, L., Carberry, P., Whitbread, A.,
1332 2020. Conservation tillage increases yield and precipitation use efficiency of wheat on the semi-arid Loess
1333 Plateau of China. Agricultural Water Management 231, 106024.
1334 <https://doi.org/10.1016/j.agwat.2020.106024>
- 1335 Peralta, A.L., Sun, Y., McDaniel, M.D., Lennon, J.T., 2018. Crop rotational diversity increases disease
1336 suppressive capacity of soil microbiomes. Ecosphere 9. <https://doi.org/10.1002/ecs2.2235>

- 1337 Perkons, U., Kautz, T., Uteau, D., Peth, S., Geier, V., Thomas, K., Lütke Holz, K., Athmann, M., Pude, R.,
 1338 Köpke, U., 2014. Root-length densities of various annual crops following crops with contrasting root
 1339 systems. *Soil and Tillage Research* 137, 50–57. <https://doi.org/10.1016/j.still.2013.11.005>
- 1340 Perry, C., 2007. Efficient irrigation; inefficient communication; flawed recommendations. *Irrig. and Drain.*
 1341 56, 367–378. <https://doi.org/10.1002/ird.323>
- 1342 Peterson, C.A., Nunes, P.A. de A., Martins, A.P., Bergamaschi, H., Anghinoni, I., Carvalho, P.C. de F.,
 1343 Gaudin, A.C.M., 2019. Winter grazing does not affect soybean yield despite lower soil water content in a
 1344 subtropical crop-livestock system. *Agron. Sustain. Dev.* 39, 26. [https://doi.org/10.1007/s13593-019-0573-](https://doi.org/10.1007/s13593-019-0573-3)
 1345 3
- 1346 Poeplau, C., Don, A., 2015. Carbon sequestration in agricultural soils via cultivation of cover crops – A
 1347 meta-analysis. *Agriculture, Ecosystems & Environment* 200, 33–41.
 1348 <https://doi.org/10.1016/j.agee.2014.10.024>
- 1349 Portmann, F.T., Siebert, S., Döll, P., 2010. MIRCA2000-Global monthly irrigated and rainfed crop areas
 1350 around the year 2000: A new high-resolution data set for agricultural and hydrological modeling:
 1351 MONTHLY IRRIGATED AND RAINFED CROP AREAS. *Global Biogeochem. Cycles* 24, n/a-n/a.
 1352 <https://doi.org/10.1029/2008GB003435>
- 1353 Pretty, J., Bharucha, Z.P., 2014. Sustainable intensification in agricultural systems. *Annals of Botany* 114,
 1354 1571–1596. <https://doi.org/10.1093/aob/mcu205>
- 1355 Püschel, D., Bitterlich, M., Rydlová, J., Jansa, J., 2020. Facilitation of plant water uptake by an arbuscular
 1356 mycorrhizal fungus: a Gordian knot of roots and hyphae. *Mycorrhiza* 30, 299–313.
 1357 <https://doi.org/10.1007/s00572-020-00949-9>
- 1358 Quach, M., Mele, P.M., Hayden, H.L., Marshall, A.J., Mann, L., Hu, H.-W., He, J.-Z., 2022. Proximity to
 1359 subsurface drip irrigation emitters altered soil microbial communities in two commercial processing tomato
 1360 fields. *Applied Soil Ecology* 171, 104315. <https://doi.org/10.1016/j.apsoil.2021.104315>

- 1361 Quinteiro, P., Rafael, S., Villanueva-Rey, P., Ridoutt, B., Lopes, M., Arroja, L., Dias, A.C., 2018. A
 1362 characterisation model to address the environmental impact of green water flows for water scarcity
 1363 footprints. *Science of The Total Environment* 626, 1210–1218.
 1364 <https://doi.org/10.1016/j.scitotenv.2018.01.201>
- 1365 Rawls, W.J., Pachepsky, Y.A., Ritchie, J.C., Sobecki, T.M., Bloodworth, H., 2003. Effect of soil organic
 1366 carbon on soil water retention. *Geoderma* 116, 61–76. [https://doi.org/10.1016/S0016-7061\(03\)00094-6](https://doi.org/10.1016/S0016-7061(03)00094-6)
- 1367 Renwick, L.L.R., Deen, W., Silva, L., Gilbert, M.E., Maxwell, T., Bowles, T.M., Gaudin, A.C.M., 2021.
 1368 Long-term crop rotation diversification enhances maize drought resistance through soil organic matter.
 1369 *Environ. Res. Lett.* 16, 084067. <https://doi.org/10.1088/1748-9326/ac1468>
- 1370 Richards, L.A., 1931. CAPILLARY CONDUCTION OF LIQUIDS THROUGH POROUS MEDIUMS.
 1371 *Physics* 1, 318–333. <https://doi.org/10.1063/1.1745010>
- 1372 Rillig, M.C., Mummey, D.L., 2006. Mycorrhizas and soil structure. *New Phytologist* 171, 41–53.
 1373 <https://doi.org/10.1111/j.1469-8137.2006.01750.x>
- 1374 Rockström, J., Falkenmark, M., Karlberg, L., Hoff, H., Rost, S., Gerten, D., 2009. Future water availability
 1375 for global food production: The potential of green water for increasing resilience to global change: WATER
 1376 AVAILABILITY FOR FOOD PRODUCTION. *Water Resour. Res.* 45.
 1377 <https://doi.org/10.1029/2007WR006767>
- 1378 Rost, S., Gerten, D., Bondeau, A., Lucht, W., Rohwer, J., Schaphoff, S., 2008. Agricultural green and blue
 1379 water consumption and its influence on the global water system: GLOBAL WATER USE IN
 1380 AGRICULTURE. *Water Resour. Res.* 44. <https://doi.org/10.1029/2007WR006331>
- 1381 Rubin, R.L., van Groenigen, K.J., Hungate, B.A., 2017. Plant growth promoting rhizobacteria are more
 1382 effective under drought: a meta-analysis. *Plant Soil* 416, 309–323. [https://doi.org/10.1007/s11104-017-](https://doi.org/10.1007/s11104-017-3199-8)
 1383 3199-8

- 1384 Sánchez, P.A., 2010. Tripling crop yields in tropical Africa. *Nature Geosci* 3, 299–300.
1385 <https://doi.org/10.1038/ngeo853>
- 1386 Saxton, K.E., Rawls, W.J., 2006. Soil Water Characteristic Estimates by Texture and Organic Matter for
1387 Hydrologic Solutions. *Soil Sci. Soc. Am. J.* 70, 1569–1578. <https://doi.org/10.2136/sssaj2005.0117>
- 1388 Scanlon, B.R., Zhang, Z., Reedy, R.C., Pool, D.R., Save, H., Long, D., Chen, J., Wolock, D.M., Conway,
1389 B.D., Winester, D., 2015. Hydrologic implications of GRACE satellite data in the Colorado River Basin:
1390 GRACE SATELLITE DATA IN THE COLORADO RIVER BASIN. *Water Resour. Res.* 51, 9891–9903.
1391 <https://doi.org/10.1002/2015WR018090>
- 1392 Schjønning, P., Thomsen, I.K., Petersen, S.O., Kristensen, K., Christensen, B.T., 2011. Relating soil
1393 microbial activity to water content and tillage-induced differences in soil structure. *Geoderma* 163, 256–
1394 264. <https://doi.org/10.1016/j.geoderma.2011.04.022>
- 1395 Schmidt, J.E., Gaudin, A.C.M., 2017. Toward an Integrated Root Ideotype for Irrigated Systems. *Trends in*
1396 *Plant Science* 22, 433–443. <https://doi.org/10.1016/j.tplants.2017.02.001>
- 1397 Schwartz, R.C., Baumhardt, R.L., Evett, S.R., 2010. Tillage effects on soil water redistribution and bare
1398 soil evaporation throughout a season. *Soil and Tillage Research* 110, 221–229.
1399 <https://doi.org/10.1016/j.still.2010.07.015>
- 1400 Schwen, A., Bodner, G., Scholl, P., Buchan, G.D., Loiskandl, W., 2011. Temporal dynamics of soil
1401 hydraulic properties and the water-conducting porosity under different tillage. *Soil and Tillage Research*
1402 113, 89–98. <https://doi.org/10.1016/j.still.2011.02.005>
- 1403 Sears, L., Caparelli, J., Lee, C., Pan, D., Strandberg, G., Vuu, L., Lin Lawell, C.-Y., 2018. Jevons' Paradox
1404 and Efficient Irrigation Technology. *Sustainability* 10, 1590. <https://doi.org/10.3390/su10051590>

- 1405 Siebert, S., Burke, J., Faures, J.M., Frenken, K., Hoogeveen, J., Döll, P., Portmann, F.T., 2010.
1406 Groundwater use for irrigation – a global inventory. *Hydrol. Earth Syst. Sci.* 14, 1863–1880.
1407 <https://doi.org/10.5194/hess-14-1863-2010>
- 1408 Šimůnek, J., Genuchten, M.Th., 2008. Modeling Nonequilibrium Flow and Transport Processes Using
1409 HYDRUS. *Vadose zone j.* 7, 782–797. <https://doi.org/10.2136/vzj2007.0074>
- 1410 Šimůnek, J., Genuchten, M.Th., Šejna, M., 2016. Recent Developments and Applications of the HYDRUS
1411 Computer Software Packages. *Vadose zone j.* 15, 1–25. <https://doi.org/10.2136/vzj2016.04.0033>
- 1412 Six, J., Elliott, E.T., Paustian, K., 1999. Aggregate and Soil Organic Matter Dynamics under Conventional
1413 and No-Tillage Systems. *Soil Sci. Soc. Am. j.* 63, 1350–1358. <https://doi.org/10.2136/sssaj1999.6351350x>
- 1414 Smith, C.W., 2016. Effects of Implementation of Soil Health Management Practices on Infiltration,
1415 Saturated Hydraulic Conductivity and Runoff. USDA Natural Resources Conservation Service, National
1416 Soil Survey Center.
- 1417 Smith, S.E., Read, D.J., 2010. Mycorrhizal symbiosis. Academic press.
- 1418 Soane, B.D., Ball, B.C., Arvidsson, J., Basch, G., Moreno, F., Roger-Estrade, J., 2012. No-till in northern,
1419 western and south-western Europe: A review of problems and opportunities for crop production and the
1420 environment. *Soil and Tillage Research* 118, 66–87. <https://doi.org/10.1016/j.still.2011.10.015>
- 1421 Sofo, A., Mininni, A.N., Ricciuti, P., 2020. Soil Macrofauna: A key Factor for Increasing Soil Fertility and
1422 Promoting Sustainable Soil Use in Fruit Orchard Agrosystems. *Agronomy* 10, 456.
1423 <https://doi.org/10.3390/agronomy10040456>
- 1424 Sokol, N.W., Kuebbing, Sara.E., Karlsen-Ayala, E., Bradford, M.A., 2019. Evidence for the primacy of
1425 living root inputs, not root or shoot litter, in forming soil organic carbon. *New Phytol* 221, 233–246.
1426 <https://doi.org/10.1111/nph.15361>

- 1427 Somers, E., Vanderleyden, J., Srinivasan, M., 2004. Rhizosphere Bacterial Signalling: A Love Parade
1428 Beneath Our Feet. *Critical Reviews in Microbiology* 30, 205–240.
1429 <https://doi.org/10.1080/10408410490468786>
- 1430 Sposito, G., 2013. Green Water and Global Food Security. *Vadose Zone Journal* 12, vzj2013.02.0041.
1431 <https://doi.org/10.2136/vzj2013.02.0041>
- 1432 Steffen, W., Richardson, K., Rockstrom, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter,
1433 S.R., de Vries, W., de Wit, C.A., Folke, C., Gerten, D., Heinke, J., Mace, G.M., Persson, L.M., Ramanathan,
1434 V., Reyers, B., Sorlin, S., 2015. Planetary boundaries: Guiding human development on a changing planet.
1435 *Science* 347, 1259855–1259855. <https://doi.org/10.1126/science.1259855>
- 1436 Steudle, E., 2001. The cohesion-tension mechanism and the acquisition of water by plant roots. *Annual*
1437 *review of plant biology* 52, 847–875.
- 1438 Strudley, M., Green, T., Ascoughii, J., 2008. Tillage effects on soil hydraulic properties in space and time:
1439 State of the science. *Soil and Tillage Research* 99, 4–48. <https://doi.org/10.1016/j.still.2008.01.007>
- 1440 Sun, C., Ren, L., 2014. Assessing crop yield and crop water productivity and optimizing irrigation
1441 scheduling of winter wheat and summer maize in the Haihe plain using SWAT model: CROP YIELD AND
1442 CWP AND IRRIGATION SCHEDULING IN HAIHE PLAIN. *Hydrol. Process.* 28, 2478–2498.
1443 <https://doi.org/10.1002/hyp.9759>
- 1444 Sun, M., Ren, A., Gao, Z., Wang, P., Mo, F., Xue, L., Lei, M., 2018. Long-term evaluation of tillage
1445 methods in fallow season for soil water storage, wheat yield and water use efficiency in semiarid southeast
1446 of the Loess Plateau. *Field Crops Research* 218, 24–32. <https://doi.org/10.1016/j.fcr.2017.12.021>
- 1447 Taylor, R., Zilberman, D., 2017. Diffusion of Drip Irrigation: The Case of California. *Applied Economic*
1448 *Perspectives and Policy* 39, 16–40. <https://doi.org/10.1093/aepp/ppw026>

- 1449 Thirkell, T.J., Charters, M.D., Elliott, A.J., Sait, S.M., Field, K.J., 2017. Are mycorrhizal fungi our
1450 sustainable saviours? Considerations for achieving food security. *J Ecol* 105, 921–929.
1451 <https://doi.org/10.1111/1365-2745.12788>
- 1452 Timmusk, S., Abd El-Daim, I.A., Copolovici, L., Tanilas, T., Kännaste, A., Behers, L., Nevo, E.,
1453 Seisenbaeva, G., Stenström, E., Niinemets, Ü., 2014. Drought-Tolerance of Wheat Improved by
1454 Rhizosphere Bacteria from Harsh Environments: Enhanced Biomass Production and Reduced Emissions
1455 of Stress Volatiles. *PLoS ONE* 9, e96086. <https://doi.org/10.1371/journal.pone.0096086>
- 1456 Trenberth, K.E., Dai, A., van der Schrier, G., Jones, P.D., Barichivich, J., Briffa, K.R., Sheffield, J., 2014.
1457 Global warming and changes in drought. *Nature Clim Change* 4, 17–22.
1458 <https://doi.org/10.1038/nclimate2067>
- 1459 Trout, T.J., Manning, D.T., 2019. An Economic and Biophysical Model of Deficit Irrigation. *Agronomy*
1460 *Journal* 111, 3182–3193. <https://doi.org/10.2134/agronj2019.03.0209>
- 1461 Turmel, M.-S., Speratti, A., Baudron, F., Verhulst, N., Govaerts, B., 2015. Crop residue management and
1462 soil health: A systems analysis. *Agricultural Systems* 134, 6–16. <https://doi.org/10.1016/j.agsy.2014.05.009>
- 1463 Unger, P.W., Vigil, M.F., 1998. Cover crop effects on soil water relationships. *Journal of Soil and Water*
1464 *Conservation* 53, 200–207.
- 1465 Van Gaelen, H., 2016. Evaluation of agricultural management from field to catchment scale: Development
1466 of a parsimonious agro-hydrological model.
- 1467 Van Geel, M., Verbruggen, E., De Beenhouwer, M., van Rennes, G., Lievens, B., Honnay, O., 2017. High
1468 soil phosphorus levels overrule the potential benefits of organic farming on arbuscular mycorrhizal diversity
1469 in northern vineyards. *Agriculture, Ecosystems & Environment* 248, 144–152.
1470 <https://doi.org/10.1016/j.agee.2017.07.017>

- 1471 Vardharajula, S., Zulfikar Ali, S., Grover, M., Reddy, G., Bandi, V., 2011. Drought-tolerant plant growth
1472 promoting *Bacillus* spp.: effect on growth, osmolytes, and antioxidant status of maize under drought stress.
1473 Journal of Plant Interactions 6, 1–14. <https://doi.org/10.1080/17429145.2010.535178>
- 1474 Veettil, A.V., Mishra, A.K., 2016. Water security assessment using blue and green water footprint concepts.
1475 Journal of Hydrology 542, 589–602. <https://doi.org/10.1016/j.jhydrol.2016.09.032>
- 1476 Vercambre, G., Pagès, L., Doussan, C., Habib, R., 2003. Architectural analysis and synthesis of plum tree
1477 root system in an orchard using a quantitative modelling approach. Plant and Soil 251, 1–11.
1478 <https://doi.org/10.1023/A:1022961513239>
- 1479 Vereecken, H., Schnepf, A., Hopmans, J.W., Javaux, M., Or, D., Roose, T., Vanderborght, J., Young, M.H.,
1480 Amelung, W., Aitkenhead, M., Allison, S.D., Assouline, S., Baveye, P., Berli, M., Brüggemann, N., Finke,
1481 P., Flury, M., Gaiser, T., Govers, G., Ghezzehei, T., Hallett, P., Hendricks Franssen, H.J., Heppell, J., Horn,
1482 R., Huisman, J.A., Jacques, D., Jonard, F., Kollet, S., Lafolie, F., Lamorski, K., Leitner, D., McBratney,
1483 A., Minasny, B., Montzka, C., Nowak, W., Pachepsky, Y., Padarian, J., Romano, N., Roth, K., Rothfuss,
1484 Y., Rowe, E.C., Schwen, A., Šimůnek, J., Tiktak, A., Van Dam, J., van der Zee, S.E.A.T.M., Vogel, H.J.,
1485 Vrugt, J.A., Wöhling, T., Young, I.M., 2016. Modeling Soil Processes: Review, Key Challenges, and New
1486 Perspectives. Vadose Zone Journal 15, vzj2015.09.0131. <https://doi.org/10.2136/vzj2015.09.0131>
- 1487 Villamil, M.B., Bollero, G.A., Darmody, R.G., Simmons, F.W., Bullock, D.G., 2006. No-Till
1488 Corn/Soybean Systems Including Winter Cover Crops: Effects on Soil Properties. Soil Sci. Soc. Am. J. 70,
1489 1936–1944. <https://doi.org/10.2136/sssaj2005.0350>
- 1490 Vincent, C., Rowland, D., Schaffer, B., Bassil, E., Racette, K., Zurweller, B., 2020. Primed acclimation: A
1491 physiological process offers a strategy for more resilient and irrigation-efficient crop production. Plant
1492 Science 295, 110240. <https://doi.org/10.1016/j.plantsci.2019.110240>
- 1493 von Korff, Y., Daniell, K.A., Moellenkamp, S., Bots, P., Bijlsma, R.M., 2012. Implementing Participatory
1494 Water Management: Recent Advances in Theory, Practice, and Evaluation. E&S 17, art30.
1495 <https://doi.org/10.5751/ES-04733-170130>

- 1496 Vurukonda, S.S.K.P., Vardharajula, S., Shrivastava, M., SkZ, A., 2016. Enhancement of drought stress
1497 tolerance in crops by plant growth promoting rhizobacteria. *Microbiological Research* 184, 13–24.
1498 <https://doi.org/10.1016/j.micres.2015.12.003>
- 1499 Wade, J., Beetstra, M.A., Hamilton, M.L., Culman, S.W., Margenot, A.J., 2021. Soil health
1500 conceptualization differs across key stakeholder groups in the Midwest. *Journal of Soil and Water*
1501 *Conservation* 02158. <https://doi.org/10.2489/jswc.2021.02158>
- 1502 Wade, J., Culman, S.W., Logan, J.A.R., Poffenbarger, H., Demyan, M.S., Grove, J.H., Mallarino, A.P.,
1503 McGrath, J.M., Ruark, M., West, J.R., 2020. Improved soil biological health increases corn grain yield in
1504 N fertilized systems across the Corn Belt. *Sci Rep* 10, 3917. <https://doi.org/10.1038/s41598-020-60987-3>
- 1505 Waqas, M.S., Cheema, M.J.M., Hussain, S., Ullah, M.K., Iqbal, M.M., 2021. Delayed irrigation: An
1506 approach to enhance crop water productivity and to investigate its effects on potato yield and growth
1507 parameters. *Agricultural Water Management* 245, 106576. <https://doi.org/10.1016/j.agwat.2020.106576>
- 1508 Ward, F.A., Pulido-Velazquez, M., 2008. Water conservation in irrigation can increase water use.
1509 *Proceedings of the National Academy of Sciences* 105, 18215–18220.
1510 <https://doi.org/10.1073/pnas.0805554105>
- 1511 Ward, P.R., Flower, K.C., Cordingley, N., Weeks, C., Micin, S.F., 2012. Soil water balance with cover
1512 crops and conservation agriculture in a Mediterranean climate. *Field Crops Research* 132, 33–39.
1513 <https://doi.org/10.1016/j.fcr.2011.10.017>
- 1514 Waterhouse, H., Bachand, S., Mountjoy, D., Choperena, J., Bachand, P., Dahlke, H., Horwath, W., 2020.
1515 Agricultural managed aquifer recharge—water quality factors to consider. *California Agriculture* 74, 144–
1516 154.
- 1517 Weaver, J.E., Bruner, W.E., 1928. Root Development of Vegetable Crops. *Nature* 121, 53–53.
1518 <https://doi.org/10.1038/121053a0>

- 1519 White, J.W., Jones, J.W., Porter, C., McMaster, G.S., Sommer, R., 2010. Issues of spatial and temporal
1520 scale in modeling the effects of field operations on soil properties. *Oper Res Int J* 10, 279–299.
1521 <https://doi.org/10.1007/s12351-009-0067-1>
- 1522 Williams, A., Hunter, M.C., Kammerer, M., Kane, D.A., Jordan, N.R., Mortensen, D.A., Smith, R.G.,
1523 Snapp, S., Davis, A.S., 2016. Soil Water Holding Capacity Mitigates Downside Risk and Volatility in US
1524 Rainfed Maize: Time to Invest in Soil Organic Matter? *PLoS ONE* 11, e0160974.
1525 <https://doi.org/10.1371/journal.pone.0160974>
- 1526 Williams, S.M., Weil, R.R., 2004. Crop Cover Root Channels May Alleviate Soil Compaction Effects on
1527 Soybean Crop. *Soil Sci. Soc. Am. J.* 68, 1403–1409. <https://doi.org/10.2136/sssaj2004.1403>
- 1528 Zhang, D., Chen, X., Yao, H., Lin, B., 2015. Improved calibration scheme of SWAT by separating wet and
1529 dry seasons. *Ecological Modelling* 301, 54–61. <https://doi.org/10.1016/j.ecolmodel.2015.01.018>
1530
- 1531 Zhang, J., Guan, K., Peng, B., Pan, M., Zhou, W., Jiang, C., Kimm, H., Franz, T.E., Grant, R.F., Yang, Y.,
1532 Rudnick, D.R., Heeren, D.M., Suyker, A.E., Bauerle, W.L., Miner, G.L., 2021. Sustainable irrigation based
1533 on co-regulation of soil water supply and atmospheric evaporative demand. *Nat Commun* 12, 5549.
1534 <https://doi.org/10.1038/s41467-021-25254-7>
- 1535 Zheng, W., Zeng, S., Bais, H., LaManna, J.M., Hussey, D.S., Jacobson, D.L., Jin, Y., 2018. Plant Growth-
1536 Promoting Rhizobacteria (PGPR) Reduce Evaporation and Increase Soil Water Retention. *Water Resour.*
1537 *Res.* 54, 3673–3687. <https://doi.org/10.1029/2018WR022656>
1538

1539 **Contributions**

1540 Conceptualization and organization of the article: S.E.A., H.W. and T.M.B.

1541 Figures and Tables: S.E.A. and T.M.B with input from all authors

1542 Drafted and/or revised the article: All authors.

1543 Approved the submitted version for publication: All authors.

1544 **Acknowledgments**

1545 The authors thank Dr. Carlos Bonilla, Dr. Alastair Iles, and Dr. Louise Jackson, as well as three anonymous
1546 reviewers, for providing helpful comments on the manuscript.

1547 **Funding information**


1548 S.E.A. acknowledges funding from ANID/CONICYT Doctorado Nacional Scholarship 21160742 and
1549 ANID/CONICYT/FONDECYT/Regular 1191166, Government of Chile. T.M.B. and H.W. acknowledge
1550 funding from USDA AFRI Grant # 2019-67019-29537 and NSF Coupled Natural Humans Systems Grant
1551 #1824871. T.M.B. and J.D. acknowledge funding from the California Department of Food and Agriculture
1552 Specialty Crops Grant #19-0001-016-SF. L.L.R.R. acknowledges funding from ANID Postdoctoral
1553 FONDECYT #3210036. F.B.M acknowledges funding from the USDA NIFA Hatch project 1009799.

1554 **Competing interests**

1555 The authors have declared that no competing interests exist

1556

1557 **Figures**



1558

1559

1560 **Figure 1. Annual precipitation data of California and estimated water requirements (2016).**

1561 **A:** Total annual precipitation (cm). **B:** Modeled annual ET fraction derived from green water (%) in irrigated

1562 systems. Green crop water requirements represent the amount of ET met by precipitation. Blue water

1563 requirements represent the amount of irrigation required to make up the difference between ET and the

1564 green crop water requirement, although this may overestimate blue water since in practice the full amount

1565 of water required to replace ET is sometimes not provided (e.g., with deficit irrigation). Areas

1566 corresponding to Prime Farmlands and Farmland of Statewide Importance were derived from census-based

1567 inventories as reported in Portmann et al. (2010) and the Farming Mapping and Monitoring Program (2021).

1568 Based on the dataset by Chiarelli et al. (2020) and gridMET (Abatzoglou, 2021, 2013)




Figure 2. Conceptual framework for outcomes of soil health management for water resources.

Hydrologic processes (orange circles) and plant-soil-microbe interactions below (green circles) and aboveground (pink circles) affecting blue and green water use in irrigated systems. Soil health management practices are represented in circles on the left side. Blue arrows indicate blue water flows, and green arrows indicate green water flows. Arrows with both colors represent mixed blue and green water components. Arrow size does not represent the flow magnitude. Original artwork by Elena Harley (www.elabarts.com)

Tables

Table 1. Irrigation water management strategies and how they interact with soil health management to decrease use of blue water, increase green water availability and/or increase the productive flow of water, i.e. integrate soil health-irrigation management.

Water management strategy	Interaction with soil health management	Key research questions
Delaying onset of irrigation until a benchmark level of allowable depletion has occurred (Zhang et al., 2021)	Depends on increases in green water availability via greater soil water capture and storage	What are levels of allowable soil water depletion for specific crops? How would delaying onset of irrigation affect direct seeded or transplanted annuals?
Applying fewer, deeper irrigations with reduced total water applied (Waqas et al., 2021)	Reducing evaporation and increasing rainfall infiltration by cover cropping, crop residues, mulching, and/or no-till results in less blue water needed to meet crop water demand	How can cover crops be managed across varying agro-climatic contexts to optimize impacts on soil water content for subsequent cash crops? Could this be implemented with drip irrigation systems?
Deficit irrigation or partial root-zone drying (Barrios-Masias and Jackson 2016, Lipan et al., 2019)	Greater plant-microbe interactions and root proliferation and penetration could reduce impacts on crop production	To what extent will soil health management maintain or enhance crop productivity from varying levels and approaches to deficit irrigation or partial root zone drying?
Matching irrigation strategy with root system traits (García-Tejera et al., 2018, Vincent et al., 2020)	Root proliferation in well-structured topsoil could maximize capture of shallow irrigation while deep root penetration could maximize deep stored water from higher infiltration	Is the root system architecture of crops and varieties suited to access stored soil water at different depths? How could breeding programs help match crops with patterns of water availability?

Adaptively design cropping patterns to optimize water use (Chen et al., 2022)	Planning and adaptively managing crop rotations, fallowing, and/or intercropping to facilitate complementarity in water use in space and time could enhance green water availability	How could interspecific interactions in intercropping enhance water use efficiency? Do crop rotations based on rooting depth differences increase the efficient use of stored water?
Managed groundwater recharge (Meyer et al., 2019)	Cover cropping could facilitate groundwater recharge by maintaining soil structure and reducing nutrient loading to groundwater	How does the timing of flood flow applications and cover crop establishment interact to affect nutrient retention and deep percolation? When should a cover crop be terminated to reduce non-productive water flow?
Irrigation efficiency technologies (Dumont et al., 2013)	Smaller, more frequent, and more localized irrigation could reduce other benefits from soil health improvements if only a very small portion of the soil profile is moist during the growing season	How can effective water governance policies be designed to minimize the potential for Jevon's paradox due to increases in water use efficiency at the field scale? What are the tradeoffs between irrigation technologies and soil health benefits based on water distribution in the soil profile?

1584

1585

1586

1587

Table 2. Model summary of selected models used in the comparison of both soil health management and green-blue water balance.

Model	Model type	Focal scale	‘Less drop per crop’ and ‘more crop per drop’ examples related to soil health and green-blue water balance
LPJmL	Process based	Global	- Effects of tillage in crop productivity (Lutz et al., 2019) - Mulching and conservation tillage in irrigated systems (Jägermeyr et al., 2016) - Combined effect of reservoir operation and irrigation extractions (Biemans et al., 2011)
SWAT	Physical based	Hydrologic and erosion model at the basin scale	- Importance of capillary rise (Kroes et al., 2017) - Irrigation and reservoir storage (Jouma and Dadaser-Celik, 2021) - Crop yield and irrigation optimization (Sun and Ren, 2014)
AquaCrop	Process based	Crop simulation at field scale	- Irrigation and capillary rise in crop productivity (Khan et al., 2021) - Deficit irrigation and mulching in crop productivity and water savings (Bao, 2020) - Water saving effect of mulching and drip irrigation (Nouri et al., 2019)
HYDRUS 1-D	Physical based	Water flow and solute transport through the soil profile	- HYDRUS-1D coupled with a crop growth model: impact of groundwater in productivity (M. Han et al., 2015) - Water dynamics in irrigated conservation agriculture (Bekele et al., 2020) - Water balance under no tillage and SOM incorporation (Aggarwal et al., 2017)

1591

1592 **Boxes**

1593 Box 1: Irrigation management strategies to reduce blue water inputs

1594 Deficit irrigation implies reductions in the amount of water provided to a crop compared to a fully irrigated
1595 crop based on the total crop evapotranspiration. A fully irrigated crop would receive blue water equivalent
1596 to crop evapotranspiration. Two approaches for reducing the amount of blue water applied are a controlled
1597 deficit irrigation and partial root zone drying. The former is applied during certain crop stages when drought
1598 has less impact on yield (e.g., fruit ripening in tomatoes). Partial root zone drying integrates plant
1599 physiological responses to drought (e.g., tighter regulation of transpiration rates) by wetting only part of
1600 the root system (e.g., 50%), which is alternated in each irrigation. Partial root zone drying can be applied
1601 during most of the growing season. However, not all crops or cultivars may respond well to these irrigation
1602 strategies.

1603