



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique

Elon Lindenstrauss, Amir Mohammadi and Zhiren Wang

Polynomial effective equidistribution

Volume 361 (2023), p. 507-520

Published online: 1 February 2023

<https://doi.org/10.5802/crmath.411>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 1778-3569



Dynamical systems / Systèmes dynamiques

Polynomial effective equidistribution

Elon Lindenstrauss^a, Amir Mohammadi^{*,b} and Zhiren Wang^c

^a The Einstein Institute of Mathematics, Edmond J. Safra Campus, Givat Ram, The Hebrew University of Jerusalem, Jerusalem, 91904, Israel

^b Department of Mathematics, University of California, San Diego, CA 92093, USA

^c Pennsylvania State University, Department of Mathematics, University Park, PA 16802, USA

E-mails: elon@math.huji.ac.il (E. Lindenstrauss), ammohammadi@ucsd.edu (A. Mohammadi), zhirenw@psu.edu (Z. Wang)

Abstract. We prove effective equidistribution theorems, with polynomial error rate, for orbits of the unipotent subgroups of $SL_2(\mathbb{I})$ in arithmetic quotients of $SL_2(\mathbb{C})$ and $SL_2(\mathbb{I}) \times SL_2(\mathbb{I})$.

The proof is based on the use of a Margulis function, tools from incidence geometry, and the spectral gap of the ambient space.

Résumé. Nous prouvons des théorèmes d'équidistribution effectifs, avec un taux d'erreur polynomial pour les orbites des sous-groupes unipotents de $SL_2(\mathbb{I})$ en quotients arithmétiques de $SL_2(\mathbb{C})$ et $SL_2(\mathbb{I}) \times SL_2(\mathbb{I})$.

La preuve est basée sur l'utilisation d'une fonction de Margulis, des outils de la géométrie d'incidence, et le trou spectral de l'espace ambiant.

Funding. E.L. acknowledges support by ERC 2020 grant HomDyn (grant no. 833423). A.M. acknowledges support by the NSF, grants DMS-1764246 and 2055122. Z.W. acknowledges support by the NSF grant DMS-1753042.

Manuscript received 13 April 2022, revised 27 July 2022, accepted 1 August 2022.

1. Introduction

A landmark result of Ratner [35] states that if G is a Lie group, Γ a lattice in G and if u_t is a one-parameter Ad-unipotent subgroup of G , then for *any* $x \in G/\Gamma$ the orbit $u_t.x$ is equidistributed in a periodic orbit of some subgroup $L < G$ that contains both the one parameter group u_t and the initial point x . We say an orbit $L.x$ of a group L in some space X is periodic if the stabilizer of x in L is a lattice in L , equivalently that the stabilizer of x in L is discrete and $L.x$ supports a unique L -invariant probability measure $m_{L.x}$; and $u_t.x$ is equidistributed in $L.x$ in the sense that

$$\frac{1}{T} \int_0^T f(u_t.x) dt \rightarrow \int f dm_{L.x} \quad \text{for any } f \in C_0(G/\Gamma). \quad (1)$$

* Corresponding author.

In order to prove this equidistribution result, Ratner first classified the u_t -invariant probability measures on G/Γ [33, 34]; the proof also uses the non-divergence properties of unipotent flows established by Dani and Margulis [5, 6, 29].

A key motivation behind Ratner's equidistribution theorem for one parameter unipotent flows has been to establish Raghunathan's conjecture regarding the possible orbit closures of groups generated by one-parameter unipotent groups; using the equidistribution theorem Ratner proved that if G and Γ are as above, and if $H < G$ is generated by one parameter unipotent groups, then for any $x \in G/\Gamma$ one has that $\overline{H.x} = L.x$ where $H < L < G$ and $L.x$ is periodic. Important special cases of Raghunathan's conjecture were proven earlier by Margulis and by Dani and Margulis using a different more direct approach, which in particular gave a proof of a rather strong form of the longstanding Oppenheim conjecture [7, 8, 30].

These results have had a surprisingly rich trove of applications in number theory and beyond. One drawback of this method, compared to more traditional number theoretic techniques, is that these equidistribution results were neither effective nor quantitative. Indeed, Ratner's proof relies heavily on the pointwise ergodic theorem and Lusin's theorem, both rather fundamental theorems but that do not have good effective analogues. The Dani–Margulis method is somewhat easier to make effective and a result in this direction was given by Margulis and the first named author in [26]; moreover a general result in this direction was announced by Margulis, Shah and two of us (E.L. and A.M.) with the first installment of this work appearing in [28]. However, this only gives density properties of these flows, not equidistribution, and the rate of density obtained in this way is far from optimal (polylog at best, though in most cases one only has an iterative-log type bound).

In this paper we announce a quantitative equidistribution result for orbits of a one parameter unipotent group on quotients G/Γ where G is either $\mathrm{SL}_2(\mathbb{C})$ or $\mathrm{SL}_2(\mathbb{I}) \times \mathrm{SL}_2(\mathbb{I})$ with a polynomial error rate, which is the first quantitative equidistribution statement for individual orbits of unipotent flows on quotients of semi-simple groups beyond the horospherical case. Our approach builds on the paper [27] by the first two authors, where an effective density result with a polynomial rate for orbits of a Borel subgroup of a subgroup $H \simeq \mathrm{SL}_2(\mathbb{R})$ of G was proved.

Recall that a group $N < G$ is horospheric if there is some $g \in G$ so that

$$N = \{h \in G : g^{-n} h g^n \rightarrow 1 \text{ as } n \rightarrow \infty\}.$$

For instance, the one parameter unipotent group

$$\left\{ \begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix} : r \in \mathbb{R} \right\}$$

is horospheric in $\mathrm{SL}_2(\mathbb{R})$ as are the groups

$$\left\{ \begin{pmatrix} 1 & r + is \\ 0 & 1 \end{pmatrix} : r, s \in \mathbb{R} \right\} \quad \text{and} \quad \left\{ \left(\begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} \right) : r, s \in \mathbb{R} \right\}$$

in $\mathrm{SL}_2(\mathbb{C})$ and $\mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$, respectively. The classification of invariant measures and orbit closures for horospherical flows was established prior to Ratner's work by Hedlund, Furstenberg, Dani, Veech and others, and this has been understood for some time also quantitatively since one can relate the distribution properties of individual N orbits to the ergodic theoretic properties of the action of g on G/Γ (cf. § 3 for more details).

We also mention that while our result is the first quantitative equidistribution statement for individual orbits of unipotent flows on quotients of semi-simple groups beyond the horospherical case¹ there have been some other interesting quantitative equidistribution results. When G is

¹As pointed out by Venkatesh in [42], the distinction between horospheric and non-horospheric is not completely clear cut, and indeed the results of that paper can also be recast as a nonhorospheric equidistribution problem; cf. [16] as

unipotent, an effective equidistribution result for unipotent flows on G/Γ was given by Green and Tao in [19], and was a key ingredient in a series of works by Green, Tao and Ziegler about linear equations in primes. In the case of quotients of the skew product $G = \mathrm{SL}_2(\mathbb{R}) \ltimes \mathbb{R}^2$, Strombergsson [41] has an effective equidistribution result for one parameter unipotent orbits (which are not horospheric in G , but project to a horospheric group on $\mathrm{SL}_2(\mathbb{R})$), and this has been generalized by several authors, in particular by Wooyeon Kim [24] (using a completely different argument) to $\mathrm{SL}_n(\mathbb{R}) \ltimes \mathbb{R}^n$. Moreover there is an important work of Einsiedler, Margulis and Venkatesh [11] showing that periodic orbits of semisimple subgroups H of a semisimple group G are quantitatively equidistributed in an appropriate homogeneous subspace of G/Γ if Γ is a congruence lattice and H has finite centralizer in G . Subsequently Einsiedler, Margulis, Venkatesh and the second named author by using Prasad's volume formula and a more adelic view point were able to prove such an equidistribution result for periodic orbits of maximal semisimple subgroups of G when the subgroup is allowed to vary [10] with arithmetic applications (the equidistribution of periodic orbits of semisimple groups is also closely connected to the equidistribution of Hecke points; a quantitative treatment of such equidistribution was given by Clozel, Oh and Ullmo in [4]).

In a slightly different direction, Bourgain, Furman, Mozes and the first named author [1] gave a quantitative equidistribution result with exponential rates (this is analogous to polynomial rates in our problem) for random walks by automorphisms of the torus. In this case this equidistribution result was new even without rates. There have been several extensions of this result, in particular [20] where a proximality assumption was removed. Kim in [24] used the techniques of [1] to study $\mathrm{SL}_n(\mathbb{R}) \ltimes \mathbb{R}^n$. Our work is also heavily influenced by [1].

We now proceed to describe our results and some of the ingredients of the proofs. Let

$$G = \mathrm{SL}_2(\mathbb{C}) \quad \text{or} \quad G = \mathrm{SL}_2(\mathfrak{l}) \times \mathrm{SL}_2(\mathfrak{l}).$$

Let $\Gamma \subset G$ be a lattice, and put $X = G/\Gamma$. We let m_X denote the G -invariant probability measure on X . Throughout the paper, we will denote by H a subgroup of G isomorphic to $\mathrm{SL}_2(\mathfrak{l})$, namely

$$\mathrm{SL}_2(\mathfrak{l}) \subset \mathrm{SL}_2(\mathbb{C}) \quad \text{or} \quad \{(g, g) : g \in \mathrm{SL}_2(\mathfrak{l})\} \subset \mathrm{SL}_2(\mathfrak{l}) \times \mathrm{SL}_2(\mathfrak{l}).$$

For all $t, r \in \mathbb{R}$, let a_t and u_r denote the image of

$$\begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix},$$

in H , respectively.

We fix maximal compact subgroups $\mathrm{SU}(2) \subset \mathrm{SL}_2(\mathbb{C})$ and $\mathrm{SO}(2) \times \mathrm{SO}(2) \subset \mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$. Let d be the right invariant metric on G which is defined using the Killing form and the aforementioned maximal compact subgroups. This metric induces a metric d_X on X , and natural volume forms on X and its submanifolds. We define the injectivity radius of a point $x \in X$ using this metric. In the sequel, $\|\cdot\|$ denotes the maximum norm on $\mathrm{Mat}_2(\mathbb{C})$ or $\mathrm{Mat}_2(\mathfrak{l}) \times \mathrm{Mat}_2(\mathfrak{l})$ with respect to the standard basis.

The following are the main results in this paper.

Theorem 1. *Assume Γ is an arithmetic lattice. For every $x_0 \in X$ and large enough R (depending explicitly on the injectivity radius of x_0), for any $T \geq R$, at least one of the following holds.*

- (1) *For every $\varphi \in C_c^\infty(X)$, we have*

$$\left| \int_0^1 \varphi(a_{\log T} u_r x_0) dr - \int \varphi dm_X \right| \leq C_1 \mathcal{S}(\varphi) R^{-\kappa_{\kappa_1}}$$

where $\mathcal{S}(\varphi)$ is a certain Sobolev norm.

well as [23]. There are also some quantitative equidistribution results for particular types of unipotent orbits, e.g. [3] by Chow and Lei Yang.

- (2) *There exists $x \in X$ such that Hx is periodic with $\text{vol}(Hx) \leq R$, and*

$$d_X(x, x_0) \leq C_1 R^A (\log T)^A T^{-1}.$$

The constants A, κ_1 , and C_1 are positive, and depend on X but not on x_0 .

Theorem 1 can be viewed as an effective version of [40, Theorem 1.4]. Combining Theorem 1 and the Dani–Margulis linearization method [9] (cf. also Shah [39]), that allows to control the amount of time a unipotent trajectory spends near invariant subvarieties of a homogeneous space, we also obtain an effective equidistribution theorem for long pieces of unipotent orbits (more precisely, we use a sharp form of the linearization method taken from [28]).

Theorem 2. *Assume Γ is an arithmetic lattice. For every $x_0 \in X$ and large enough R (depending explicitly on the injectivity radius of x_0), for any $T \geq R$, at least one of the following holds.*

- (1) *For every $\varphi \in C_c^\infty(X)$, we have*

$$\left| \frac{1}{T} \int_0^T \varphi(u_r x_0) dr - \int \varphi dm_X \right| \leq C_2 \mathcal{S}(\varphi) R^{-\kappa_2}$$

where $\mathcal{S}(\varphi)$ is a certain Sobolev norm.

- (2) *There exists $x \in G/\Gamma$ with $\text{vol}(H.x) \leq R^{A_1}$, and for every $r \in [0, T]$ there exists $g \in G$ with $\|g\| \leq R^{A_1}$ so that*

$$d_X(u_s x_0, gH.x) \leq C_2 R^{A_1} \left(\frac{|s-r|}{T} \right)^{1/A_2} \quad \text{for all } s, r \in [0, T].$$

- (3) *For every $r \in [0, T]$ and $t \in [\log R, \log T]$, the injectivity radius at $a_{-t} u_r x_0$ is at most $R^{A_1} e^{-t}$.*

The constants A_1, A_2, κ_2 , and C_2 are positive, and depend on X but not on x_0 .

The assumption in Theorem 1 that Γ is arithmetic may be relaxed. Let us say Γ has *algebraic entries* if the following is satisfied: there is a number field F , a semisimple F -group $\mathbf{G}$ of adjoint type, and a place v of F so that $F_v = \mathbb{R}$ and $\mathbf{G}(F_v)$ and G are locally isomorphic — in which case there is a surjective homomorphism from G onto the connected component of the identity in $\mathbf{G}(F_v)$ — and the image of Γ in $\mathbf{G}(F_v)$ (possibly after conjugation) is contained in $\mathbf{G}(F)$. Every arithmetic lattice has algebraic entries, but there are lattices with algebraic entries that are not arithmetic.

Note that the condition that Γ has *algebraic entries* is automatically satisfied if Γ is an irreducible lattice in $\text{SL}_2(\mathbb{R}) \times \text{SL}_2(\mathbb{R})$ or if $G = \text{SL}(2, \mathbb{C})$. Indeed, by arithmeticity theorems of Selberg and Margulis, irreducible lattices in $\text{SL}_2(\mathbb{R}) \times \text{SL}_2(\mathbb{R})$ are arithmetic [31, Chapter IX]. Moreover, by local rigidity, lattices in $\text{SL}_2(\mathbb{C})$ always have algebraic entries [18, Theorem 0.11] (see also [37, 43, 44]).

Theorem 3. *Assume Γ is a lattice which has algebraic entries. For every $0 < \delta < 1$, every $x_0 \in X$ and large enough T (depending explicitly on δ and the injectivity radius of x_0) at least one of the following holds.*

- (1) *For every $\varphi \in C_c^\infty(X)$, we have*

$$\left| \int_0^1 \varphi(a_{\log T} u_r x_0) dr - \int \varphi dm_X \right| \leq C_1 \mathcal{S}(\varphi) T^{-\delta^2 \kappa_1}$$

where $\mathcal{S}(\varphi)$ is a certain Sobolev norm.

- (2) *There exists $x \in X$ with*

$$d_X(x, x_0) \leq C_1 T^{-1/A},$$

satisfying the following: there are elements γ_1 and γ_2 in $\text{Stab}_H(x)$ with $\|\gamma_i\| \leq T^\delta$ for $i = 1, 2$ so that the group generated by $\{\gamma_1, \gamma_2\}$ is Zariski dense in H .

The constants A , κ_1 , and C_1 are positive, and depend on X but not on δ and x_0 .

The obstacle to effective equidistribution in Theorem 1 is much cleaner and simpler than in Theorem 2. This is not an artifact of the proof but a reflection of reality; a unipotent orbit may fail to equidistribute at the expected rate without it staying near a single period orbit of some subgroup $\{u_t\} < L < G$: one must allow a slow drift of the periodic orbit in the direction of the centralizer of u_t . Unlike the work of Shah in [40], where (in particular) a non-effective version of Theorem 1 is proved relying on Ratner's measure classification theorem for unipotent flows, our proof goes the other way, first establishing Theorem 1, and then reducing Theorem 2 from it using a linearization and non-divergence argument.

An extremely interesting analogue to unipotent flows on homogeneous spaces is given by the action of $\mathrm{SL}_2(\mathbb{R})$ and its subgroups on strata of abelian differentials. Let $g \geq 1$, and let $\alpha = (\alpha_1, \dots, \alpha_n)$ be a partition of $2g - 2$. Let $\mathcal{H}(\alpha)$ be the corresponding stratum of abelian differentials, i.e., the space of pairs (M, ω) where M is a compact Riemann surface with genus g and ω is a holomorphic 1-form on M whose zeroes have multiplicities $\alpha_1, \dots, \alpha_n$. The form ω defines a canonical flat metric on M with conical singularities and a natural area form. Let $\mathcal{H}_1(\alpha)$ be the space of unit area surfaces in $\mathcal{H}(\alpha)$. The space $\mathcal{H}(\alpha)$ admits a natural action of $\mathrm{SL}_2(\mathbb{R})$; this action preserves the unit area hyperboloid $\mathcal{H}_1(\alpha)$.

A celebrated theorem of Eskin and Mirzakhani [13] shows that any P -invariant ergodic measure is $\mathrm{SL}_2(\mathbb{R})$ -invariant and is supported on an affine invariant manifold, where P denotes the group of upper triangular matrices in $\mathrm{SL}_2(\mathbb{R})$. We shall refer to these measures as *affine invariant measures*. Moreover if we define, for any interval $I \subset \mathbb{R}$ and $x \in \mathcal{H}_1(\alpha)$, the probability measure μ_I^x on $\mathcal{H}_1(\alpha)$ by

$$\mu_I^x = |I|^{-1} \int_I \delta_{u_s x} ds,$$

then Eskin, Mirzakhani and the second named author [14] showed that for any $x \in \mathcal{H}_1(\alpha)$ the limit

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_{t=0}^T a_t \mu_{[0,1]}^x dt \quad \text{exists in weak}^* \text{ sense} \quad (2)$$

and is equal to an $(\mathrm{SL}(2, \mathbb{R})$ -invariant) affine invariant probability measure with x in its support. On the other hand, there are several results, in particular by Chaika, Smillie and B. Weiss in [2], that show that an analogue of Ratner's equidistribution theorem (or our Theorem 2) fails to hold in this setting, for instance for some x the sequence of measure $\mu_{[0,T]}^x$ may fail to converge as $T \rightarrow \infty$, or may converge to a non-ergodic measure. We believe the following strengthening of (2), which we have learned has already been conjectured by Forni in [17, Conjecture 1.4], should however hold:

Conjecture 4. *Let $\mathcal{H}_1(\alpha)$ be the space of unit area surfaces in stratum of abelian differentials on a genus g surface whose zeros have multiplicities given by $\alpha = (\alpha_1, \dots, \alpha_n)$, and let $x \in \mathcal{H}_1(\alpha)$. Then $\lim_{t \rightarrow \infty} a_t \mu_{[0,1]}^x$ exists in the weak* sense and is equal to an affine invariant measure with x in its support.*

Of course, once one establishes that $\lim_{t \rightarrow \infty} a_t \mu_{[0,1]}^x$ exists, the rest follows from [14].

2. Some preliminaries

We discuss the proofs of Theorems 1 and 3. As mentioned above, Theorem 2 is proved by combining Theorem 1 and the linearization techniques; in this announcement we focus on the proof of the former results. We note that the idea of using equidistribution of expanding translates of a fixed piece of a U orbit of the type $\{a_t u_s \cdot x : 0 \leq s \leq 1\}$ to deduce equidistribution of a large segment of a non-translated U orbit $\{u_s \cdot x : 0 \leq s \leq T\}$ is quite classical.

Let $U \subset N$ denote the group of upper triangular unipotent matrices in $H \subset G$, respectively, and let $A = \{a_t : t \in \mathbb{R}\} \subset H$. More explicitly, if $G = \mathrm{SL}_2(\mathbb{C})$, then

$$N = \left\{ n(r, s) = \begin{pmatrix} 1 & r + is \\ 0 & 1 \end{pmatrix} : (r, s) \in \mathbb{R}^2 \right\}$$

and $U = \{n(r, 0) : r \in \mathbb{R}\}$; note that $n(r, 0) = u_r$ for $r \in \mathbb{R}$. Let

$$V = \{n(0, s) = v_s : s \in \mathbb{R}\};$$

if $G = \mathrm{SL}_2(\mathbb{R}) \times \mathrm{SL}_2(\mathbb{R})$, then

$$N = \left\{ n(r, s) = \begin{pmatrix} 1 & r + s \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix} : (r, s) \in \mathbb{R}^2 \right\}$$

and $U = \{n(r, 0) : r \in \mathbb{R}\}$. As before, $n(r, 0) = u_r$ for $r \in \mathbb{R}$. Let

$$V = \{n(0, s) = v_s : s \in \mathbb{R}\}.$$

In both cases, we have $N = UV$. Let us denote the transpose of U by U^- and its elements by u_r^- .

Let $\mathfrak{g} = \mathrm{Lie}(G)$, that is, $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$ or $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{R}) \oplus \mathfrak{sl}_2(\mathbb{R})$. Let $\mathfrak{r} = i\mathfrak{sl}_2(\mathbb{R})$ if $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$ and $\mathfrak{r} = \mathfrak{sl}_2(\mathbb{R}) \oplus \{0\}$ if $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{R}) \oplus \mathfrak{sl}_2(\mathbb{R})$. In either case $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{r}$ where $\mathfrak{h} = \mathrm{Lie}(H) \simeq \mathfrak{sl}_2(\mathbb{R})$, and both $\mathfrak{h}$ and $\mathfrak{r}$ are $\mathrm{Ad}(H)$ -invariant.

We fix a norm on $\mathfrak{h}$ by taking the maximum norm where the coordinates are given by $\mathrm{Lie}(U)$, $\mathrm{Lie}(U^-)$ and $\mathrm{Lie}(A)$; similarly fix a norm on $\mathfrak{r}$. By taking maximum of these two norms, we obtain a norm on $\mathfrak{g}$. All these norms will be denoted by $\|\cdot\|$.

For all $\beta > 0$, we define $B_\beta^G := \exp(B_{\mathfrak{h}}(0, \beta)) \cdot \exp(B_{\mathfrak{r}}(0, \beta))$ where $B_\bullet(0, \beta)$ denotes the ball of radius β in $\bullet$ with respect to $\|\cdot\|$.

We also define

$$B_\beta^H := \{u_s^- : |s| \leq \beta\} \cdot \{a_t : |t| \leq \beta\} \cdot \{u_r : |r| \leq \beta\}$$

for all $0 < \beta < 1$.

For the sake of simplicity of the exposition here, let us assume X is compact. Let $0 < \eta_0 < 1$ be so that the map $g \mapsto gx$ is injective on $B_{100\eta_0}^G$ for all $x \in X$.

3. From large dimension to equidistribution

We begin with an equidistribution theorem which is of independent interest. In the proof of Theorem 1, this proposition will be applied to each of the sets obtained in the bootstrap phase, see Proposition 8.

Let us recall the following quantitative decay of correlations for the ambient space X : There exists $0 < \kappa_0 \leq 1$ so that

$$\left| \int \varphi(gx)\psi(x) dm_X - \int \varphi dm_X \int \psi dm_X \right| \ll \mathcal{S}(\varphi)\mathcal{S}(\psi)e^{-\kappa_0 d(e, g)} \quad (3)$$

for all $\varphi, \psi \in C_c^\infty(X) + \mathbb{C} \cdot 1$, where m_X is the G -invariant probability measure on X and d is our fixed right G -invariant metric on G . See, e.g., [25, § 2.4] and references there for (3); we note that κ_0 is absolute if Γ is a congruence subgroup. This is known in much greater generality, but the cases relevant to our paper are due to Selberg and Jacquet-Langlands [21, 38].

The quantitative decay of correlation can be used to establish quantitative results regarding the equidistribution of translates of pieces of an N -orbit. Specifically we employ the results in [25], but there is rich literature around the subject; a more complete list of such papers can be found in [27, § 1].

Now let $\xi : [0, 1] \rightarrow \mathfrak{r}$ be a smooth nonconstant curve. Then using the quantitative results regarding equidistribution of translates of pieces of an N -orbit such as [25], one can show that for every $x \in X$,

$$a_\tau \left\{ u_r \exp(\xi(s)) \cdot x : r, s \in [0, 1] \right\}$$

is equidistributed in X as $\tau \rightarrow \infty$ (with a rate which is polynomial in $e^{-\tau}$). The key point in the deduction of this equidistribution result from the equidistribution of shifted N orbits is that conjugation by a_τ moves $u_r \exp(\xi(s))$ to the direction of N , hence the above average essentially reduces to an average on a N orbit.

Roughly speaking, the following proposition states that one may replace the curve $\{\xi(s) : s \in [0, 1]\}$ with a measure on $\mathfrak{r}$ so long as the measure has dimension $\geq 1 - \theta$, for an appropriate choice of θ depending on κ_0 .

The precise formulation is the following.

Proposition 5. *For any $\theta > 0$ and $c > 0$ there is a κ_3 so that the following holds: Let $0 < \rho < 10^{-6}$, and let $F \subset B_{\mathfrak{r}}(0, \rho)$ be a finite set satisfying*

$$\frac{\#(F \cap B_{\mathfrak{r}}(0, b))}{\#F} \leq \rho^{-c} (b/\rho)^{1-\theta} \quad \text{for all } b \geq \rho$$

where $\rho < \rho^{20}$.

Then for all $x \in X$ and all $2 \log(1/\rho) \leq \tau \leq \frac{1}{10} \log(1/\rho)$, we have

$$\left| \int_0^1 \frac{1}{\#F} \sum_{w \in F} \varphi(a_\tau u_r \exp(w)x) dr - \int \varphi dm_X \right| \ll_X \mathcal{S}(\varphi) \max\left((\rho/\rho)^{\kappa_3}, \rho^{-2c} e^{2\tau\theta} \rho^{\kappa_0^2/M}\right),$$

where $\mathcal{S}(\varphi)$ is a certain Sobolev norm and $M \geq 100$ an absolute constant.

The proof of this proposition is significantly more delicate than that of the “toy version” of a shifted curve, and relies on an adaptation of a projection theorem due to Käenmäki, Orponen, and Venieri [22], based on the works of Wolff [45], Schlag [36], and [46], in conjunction with a sparse equidistribution argument due to Venkatesh [42]. These elements also played a crucial role in previous work by E.L. and A.M. [27] regarding quantitative density for the action of AU on the spaces we consider here.

The goal in the remaining steps is to show that unless Theorem 1 (2) holds, we can find a subset $J \subset [0, 1]$, where $[0, 1] \setminus J$ has an exponentially small measure, and up to an exponentially small error the uniform measure on $\{a_{A \log T} u_r x_0 : r \in J\}$ can be decomposed into counting measures on sets each of which satisfies the conditions in Proposition 5.

4. Inheritance of the Diophantine property

If part (2) in Theorem 1 holds, we are done. Let us, therefore, assume the alternative, which gives some Diophantine condition on the point x_0 in terms of its distance to nearby periodic H -orbits. The first step in the proof is to improve this Diophantine condition, perhaps not at x_0 , but at some (indeed all except a set of exponentially small measure) point on the translation of a the U -orbit segment $\{u_r \cdot x_0 : r \in [0, 1]\}$ by a big diagonal element a_t .

Proposition 6. *There exist D_0 (absolute) and C_3 (depending on X) so that the following holds. Let $S \geq R$, and let $x_0 \in X$ be so that*

$$d_X(x, x_0) \geq (\log S)^{D_0} S^{-1}$$

for all x with $\text{vol}(Hx) \leq R$. Then for all $s \geq \log S$ we have the following

$$\left| \left\{ r \in [0, 1] : \begin{array}{l} \text{There is } x \text{ with } \text{vol}(Hx) \leq R \\ \text{so that } d_X(x, a_s u_r x_0) \leq C_3^{-1} R^{-D_0} \end{array} \right\} \right| \leq C_3 R^{-1}.$$

In the proof of Proposition 6 we consider each periodic orbit individually, and then use the fact that the number of periodic H -orbits with volume $\leq R$ in X is $\ll R^6$, see e.g. [32, § 10] to conclude. The desired result for an individual orbit can be proved using Margulis functions for periodic H -orbits similar to those which were used in [27, § 9], see also [14, Proposition 2.13].

It is worth mentioning that even though [28, Theorem 1.4] concerns long pieces of U -orbits and Proposition 6 concerns translates of pieces of U -orbits, similar tools are applicable here as well. In particular, a version of Proposition 6 can also be proved using the methods of [28].

5. Closing lemma

Let $t > 0$ be a large parameter and fix some $e^{-0.01t} < \beta = e^{-\kappa t} < \eta_0$; in our application, κ will be chosen to be $\ll 1/D_0$ where the implied constant depends on X and D_0 is as in Proposition 6.

For every $\tau \geq 0$, put

$$E_\tau = B_\beta^H \cdot a_\tau \cdot \{u_r : r \in [0, 1]\} \subset H.$$

If $y \in X$ is so that the map $h \mapsto hy$ is injective on E_τ , then $\mu_{E_\tau \cdot y}$ denotes the pushforward of the normalized Haar measure on E_τ to $E_\tau \cdot y \subset X$.

Let $\tau \geq 0$ and $y \in X$. For every $z \in E_\tau \cdot y$, put

$$I_\tau(z) := \left\{ w \in \mathfrak{r} : 0 < \|w\| < \eta_0 \quad \text{and} \quad \exp(w)z \in E_\tau \cdot y \right\};$$

this is a finite subset of $\mathfrak{r}$ since E_τ is bounded — we will define $I_\mathcal{E}(h, z)$ for all $h \in H$ and more general sets $\mathcal{E}$ in the bootstrap phase below.

Let $0 < \alpha < 1$. Define the function $f_\tau : E_\tau \cdot y \rightarrow [1, \infty)$ as follows

$$f_\tau(z) = \begin{cases} \sum_{w \in I_\tau(z)} \|w\|^{-\alpha} & \text{if } I_\tau(z) \neq \emptyset \\ \eta_0^{-\alpha} & \text{otherwise} \end{cases}.$$

The following proposition supplies an initial dimension which we will bootstrap in the next phase. Roughly speaking, it asserts that points in $E_{8t} \cdot x_0$ (possibly after removing an exponentially small set of exceptions) are *separated transversal to H* , unless x_0 is extremely close to a periodic H orbit.

Proposition 7. *Assume Γ is arithmetic. There exists D_1 (which depends on Γ explicitly) satisfying the following. Let $D \geq D_1$ and $x_1 \in X$. Then for all large enough t at least one of the following holds.*

(1) *There is a subset $I(x_1) \subset [0, 1]$ with $|[0, 1] \setminus I(x_1)| \ll_X \beta$ such that for all $r \in I(x_1)$ we have the following*

- (a) $h \mapsto h \cdot a_{8t} u_r x_1$ is injective on E_t .
- (b) For all $z \in E_t \cdot a_{8t} u_r x_1$, we have

$$f_t(z) \leq e^{Dt}.$$

(2) *There is $x \in X$ such that Hx is periodic with*

$$\text{vol}(Hx) \leq e^{D_1 t} \quad \text{and} \quad d_X(x, x_1) \leq e^{(-D+D_1)t}.$$

This proposition is where the arithmeticity assumption on Γ is used. The proof is similar to the proof of [27, Proposition 6.1]. If we replace the assumption that Γ is arithmetic with the weaker requirement that Γ has algebraic entries, we get a version of this proposition where part (2) is replaced with the following.

(2') *There is $x \in X$ with*

$$d_X(x, x_1) \leq e^{(-D+D_1)t},$$

satisfying the following: there are elements γ_1 and γ_2 in $\text{Stab}_H(x)$ with $\|\gamma_i\| \leq e^{D_1 t}$ for $i = 1, 2$ so that the group generated by $\{\gamma_1, \gamma_2\}$ is Zariski dense in H .

6. Improving the dimension

Proposition 7 shows that up to an exponentially small error, the set $\{a_{g_t} u_r x_0 : r \in [0, 1]\}$ has a *small* positive dimension *transversal* to H at controlled scales. The objective in this step is to show that by applying elements of the form $a_\ell u_r$, for a fixed ℓ and a random $r \in [0, 1]$, we can inductively improve this dimension *transversal* to H at controlled scales to α (which will be chosen to be smaller than but quite close to 1). This is achieved by investigating the evolution of a certain Margulis function (cf. the survey [15], though the Margulis function we use here is somewhat intricate).

To state the main result which is Proposition 8, we need some notation. Let R be as in Theorem 1, and set $t = \frac{1}{D_1} \log R$ with D_1 as in Proposition 7. We will assume R is large enough so that the conclusion of Proposition 7 holds with this t .

Let

$$E = B^H_\beta \cdot \{u_r : |r| \leq \eta_0\}.$$

where $\beta = e^{-\kappa t}$ for some small $\kappa > 0$. (More explicitly, we will fix some $0 < \varepsilon \leq 10^{-8}$ which will depend on κ_0 , and let $\kappa = 10^{-6} \varepsilon / D$ where $D = D_1(D_0 + 1)$.)

It will be more convenient to *approximate* translations $\{a_s u_r x_0 : r \in [0, 1]\}$ with sets which are a disjoint union of local E -orbits as we now define. Let $F \subset B_\tau(0, \beta)$ be a finite set with $\#F \geq e^{t/2}$, and let $y \in X$. For every $w \in F$, let $E_w \subset E$ be a Borel set so that $m_H(E_w) \geq \beta^4$ and $m_H(E_w \Delta (B^H_{\beta^{10}} \cdot E_w)) \leq \beta m_H(E_w)$, where m_H denotes a fixed Haar measure on H . Put

$$\mathcal{E} = \bigcup E_w \cdot \{\exp(w)y : w \in F\}. \quad (4)$$

We equip $\mathcal{E}$ with the probability measure $\mu_{\mathcal{E}} = \frac{1}{\sum_w m_H(E_w)} \sum_w \mu_{w,y}$ where $\mu_{w,y}$ denotes the push-forward of $m_H|_{E_w}$ to $E_w \cdot \exp(w)y$ for every $w \in F$.

Let θ be a small constant depending on the decay of matrix coefficients in G/Γ (the exact value we shall use is $\theta = \kappa_0^2 / 10^4 M$, where κ_0 is as (3) and M as in Proposition 5). Let

$$\alpha = 1 - \theta \quad \text{and} \quad \sqrt{\varepsilon} = \theta.$$

Let $\ell = 0.01 \varepsilon t$, and let ν_ℓ be the probability measure on H defined by

$$\nu_\ell(\varphi) = \int_0^1 \varphi(a_\ell u_r) dr \quad \text{for all } \varphi \in C_c(H);$$

let $\nu_\ell^{(n)} = \nu_\ell \star \cdots \star \nu_\ell$ denote the n -fold convolution of ν_ℓ for all $n \in \mathbb{N}$.

The following is the main statement.

Proposition 8. *Let $x_1 \in X$, and assume that Proposition 7(2) does not hold for D , x_1 , and t . Let $r_1 \in I(x_1)$, where $I(x_1)$ is as in Proposition 7(1), and put $x_2 = a_{8t} u_{r_1} x_1$. Let*

$$J := [d_1 - 10^4 \varepsilon^{-1/2}, d_1] \cap \mathbb{N},$$

where $d_1 = 100 \lceil \frac{4D-3}{2\varepsilon} \rceil$. For every $d \in J$, there is a collection

$$\Xi_d = \{\mathcal{E}_{d,i} : 1 \leq i \leq N_d\}$$

of sets defined as in (4) for some $F_{d,i} \subset B_\tau(0, \beta)$, so that both of the following hold:

- (1) Put $\varrho = e^{-\sqrt{\varepsilon}t}$. Let $d \in J$, $1 \leq i \leq N_d$, and let $w_0 \in B_\tau(0, \beta)$. Then for every $w \in B_\tau(w_0, \varrho)$ and all $b \geq e^{-t/2}$, we have

$$\frac{\#(B_\tau(w, b) \cap B_\tau(w_0, \varrho) \cap F_{d,i})}{\#(B(w_0, \varrho) \cap F_{d,i})} \leq e^{\varepsilon t} (b/\varrho)^\alpha. \quad (5)$$

(2) For all $s \leq t$ and all $r \in [0, 2]$, we have

$$\left| \int \varphi(a_s u_r z) dv_\ell^{(d_1)} \star \mu_{E_t, x_2}(z) - \sum_{d,i} c_{d,i} \int \varphi(a_s u_r z) dv_\ell^{(d_1-d)} \star \mu_{\mathcal{E}_{d,i}}(z) \right| \ll \text{Lip}(\varphi) \beta^{(2)} \quad (6)$$

where $\varphi \in C_c^\infty(X)$, $c_{d,i} \geq 0$ and $\sum_{d,i} c_{d,i} = 1 - O(\beta^{(2)})$, $\text{Lip}(\varphi)$ is the Lipschitz norm of φ , and κ_4 and the implied constants depend on X .

Roughly speaking, the proposition states that up to an exponentially small error, $v_\ell^{(d_1)} \star \mu_{E_t, x_1}$ may be decomposed as

$$\sum_{d,i} c_{d,i} v_\ell^{(d_1-d)} \star \mu_{\mathcal{E}_{d,i}} \quad \text{where} \quad \sum_{d,i} c_{d,i} = 1 - O(\beta^{(2)})$$

(see (6)) and for all $d \in J$ and $1 \leq i \leq N_d$ the dimension of $\mathcal{E}_{d,i}$ transversal to H at controlled scales is $\geq \alpha$ (see (5)).

Combining Proposition 8 with the previous discussion, we may complete the proof of Theorem 1. A brief outline of this deduction follows: Let x_0 be as in the statement and suppose that part (2) in Theorem 1 does not hold. We first apply Proposition 6 with $s = \log T - C \log R$ for an appropriate large constant C to improve the (weak) Diophantine property of x_0 provided by the failure of part (2) in Theorem 1 to the stronger Diophantine property,

$$d_X(x, x_1) \gg R^{-D_0} \quad \text{for all } x \text{ with } \text{vol}(Hx) \leq R \quad (7)$$

for most points x_1 on $\{a_{\log T - C \log R} u_r x_0 : r \in [0, 1]\}$. Thus, in order to show that $\int_0^1 \varphi(a_{\log T} u_r x_0) dr$ is within $R^{-\star}$ of $\int \varphi$ it is enough to show the same for $\int_0^1 \varphi(a_{C \log R} u_r x_1) dr$ for x_1 satisfying the stronger Diophantine property (7).

The remaining time, i.e., $C \log R$, will be divided into *three phases*:

Phase I

Recall that $t = \frac{1}{D_1} \log R$. We apply Proposition 7 with the point x_1 . Then for every $r_1 \in I(x_1)$, the conclusion of part (1) in that proposition holds for $x_2 = a_{8t} u_{r_1} x_1$. That is, $h \mapsto hx_2$ is injective over E_t and the transverse dimension of E_t, x_2 is $\geq 1/D$ for all

$$x_2 \in \{a_{8t} u_{r_1} x_1 : r_1 \in I(x_1)\} \quad (8)$$

where $D = D_0 D_1 + D_1$. Therefore, in order to show that $\int_0^1 \varphi(a_{C \log R} u_r x_1) dr$ is within $R^{-\star}$ of $\int \varphi$, it is enough to show a similar estimate for $\int_0^1 \varphi(a_{C \log R - 8t} u_r x_2) dr$ for all x_2 as in (8).

Phase II

Let $s = 2\sqrt{\varepsilon}t$ (note that this is much larger than $\ell = 0.01\varepsilon t$). Then

$$\int_0^1 \varphi(a_{s+d_1\ell+t} u_r x_2) dr$$

is within $R^{-\star}$ of

$$\int_0^1 \int \varphi(a_s u_r z) dv_\ell^{(d_1)} \star \mu_{E_t, x_2}(z) dr.$$

We now use Proposition 8 to improve the small transversal dimension from $1/D$ to α . More precisely, Proposition 8 shows that

$$\int_0^1 \int \varphi(a_s u_r z) dv_\ell^{(d_1)} \star \mu_{E_t, x_2}(z) dr$$

is within $R^{-\star}$ of a convex combination of integrals of the form

$$\int_0^1 \int \varphi(a_s u_r z) dv_\ell^{(n)} \star \mu_{\mathcal{E}}(z) dr$$

where $0 \leq n = d_1 - d \leq 10^4 \varepsilon^{-1/2}$ and $\mathcal{E} = \mathcal{E}_{d,i}$ has dimension at least α transversal to H at controlled scales, see (5).

Phase III

It now suffices to show that $\int_0^1 \int \varphi(a_s u_r z) dv_\ell^{(n)} \star \mu_{\mathcal{E}}(z) dr$ is within $R^{-\star}$ of $\int \varphi$ for all $\mathcal{E}$ and n as above. We will use Proposition 5 to show this. First note that

$$\int_0^1 \int \varphi(a_s u_r z) dv_\ell^{(n)} \star \mu_{\mathcal{E}}(z) dr$$

is within $R^{-\star}$ of

$$\int \int_0^1 \varphi(a_{s+n\ell} u_r z) dr d\mu_{\mathcal{E}}(z).$$

Moreover, we have

$$2\sqrt{\varepsilon}t \leq s + n\ell \leq 2\sqrt{\varepsilon}t + \frac{10^4 \ell}{\sqrt{\varepsilon}} = 102\sqrt{\varepsilon}t;$$

in view of our choice of θ the right most term in the above series of inequalities is $\leq \frac{\kappa_0^2}{4M\theta} \sqrt{\varepsilon}t$. Thus, Proposition 5, applied with $\theta = 1 - \alpha$, $c = 2\varepsilon$, $\rho = e^{-\sqrt{\varepsilon}t}$, $\rho = e^{-t/2}$, and $\tau = s + n\ell$, gives

$$\left| \iint \varphi(a_{s+n\ell} u_r z) d\mu_{\mathcal{E}}(z) dr - \int \varphi dm_X \right| \ll \mathcal{S}(\varphi) e^{-\star t} = \mathcal{S}(\varphi) R^{-\star} \quad (9)$$

where the implied constants depend on X .

Note that the total time required for these three phases is $s + d_1 \ell + 9t$ which in view of the choices of s , ℓ and t is indeed a (large) constant times $\log R$. Theorem 1 follows.

In the setting of Theorem 3, we cannot utilize Proposition 6 combined with Proposition 7 as we did above; cf. the weaker conclusion in (2') following Proposition 7. Thus, we only use the three phases above (with $t = \star \delta \log T$) to improve the small dimension, namely the parameter δ in Theorem 3, to α . Thus the number of steps required is $\gg 1/\delta$ which forces $\kappa \ll \delta$. Hence, we only obtain the rate $T^{-\star \delta^2}$ in part (1) of Theorem 3.

The proof of Propositions 8 relies on the evolution of the Margulis function $\hat{f}_{\mathcal{E},\rho,\mathbf{K}}$ defined below. For every $(h, z) \in H \times \mathcal{E}$, put

$$I_{\mathcal{E},\rho}(h, z) := \left\{ w \in \mathfrak{r} : 0 < \|w\| < \rho \text{ and } \exp(w)hz \in h\mathcal{E} \right\}.$$

Since E is bounded, $I_{\mathcal{E},\rho}(h, z)$ is a finite set for all $(h, z) \in H \times \mathcal{E}$.

For every $\mathbf{K} \geq 0$, define the modified and localized Margulis function $\hat{f}_{\mathcal{E},\rho,\mathbf{K}} : H \times \mathcal{E} \rightarrow [1, \infty)$ as follows²: if $\#I_{\mathcal{E},\rho}(h, z) \leq \mathbf{K}$, put

$$\hat{f}_{\mathcal{E},\rho,\mathbf{K}}(h, z) = \rho^{-\alpha},$$

and if $\#I_{\mathcal{E},\rho}(h, z) > \mathbf{K}$, put

$$\hat{f}_{\mathcal{E},\rho,\mathbf{K}}(h, z) = \min \left\{ \sum_{w \in I} \|w\|^{-\alpha} : I \subset I_{\mathcal{E},\rho}(h, z) \text{ and } \#(I_{\mathcal{E},\rho}(h, z) \setminus I) = \mathbf{K} \right\}.$$

We begin the outline of the proof of Propositions 8 with the following observation: the set E_{t,x_1} gives rise to sets $\mathcal{E}$ which are defined as in (4) and since we assume that Proposition 7 (2) does not hold these satisfy $\hat{f}_{\mathcal{E},\rho,0}(e, z) \leq e^{Dt}$ for all $z \in \mathcal{E}$.

Let $\mathcal{E}$ be one of these sets, then up to an exponentially small error, $\nu_\ell \star \mu_{\mathcal{E}}$ may be decomposed into $\sum c'_j \mu_{\mathcal{E}'_j}$ where $c'_j \geq 0$ and $\sum c'_j = 1 - O(\beta^\star)$. This can be seen by first decomposing $\nu_\ell \star \mu_{\mathcal{E}}$ into a combination of measures supported on subsets of X which are exponentially thin in the

²If Γ is nonuniform, this Margulis function needs to be modified accordingly; but in this section we limit ourselves to the compact case.

direction of U^- (note that $a_\ell E.y$ will be exponentially thin in the direction of U^-), and then smearing these measures with B_β^H . Continuing inductively, we obtain the decomposition (6).

The fact that the energy estimate (5) is also satisfied for the terms appearing in (6) is at the heart of the proposition. The proof of this fact is based on the following bootstrap step: for all but an exponentially small set of $r \in [0, 1]$ and all but an exponentially small set of $z \in \mathcal{E}$,

$$\widehat{f}_{\mathcal{E}, \varrho, \beta^{-1}}(a_\ell u_r, z) \leq e^{-\ell/2} e^{Dt} + 4e^\ell \varrho^{-\alpha} \psi_{\mathcal{E}, \varrho}(a_\ell u_r, z). \quad (10)$$

where $\psi_{\mathcal{E}, \varrho}(h, z) := \max\{\#I_{\mathcal{E}, \varrho}(h, z), 1\}$.

The proof of (10) relies on the aforementioned works of Wolff, Schlag, and Zahl [36, 45, 46], see also the discussion below.

The set $a_\ell u_r \mathcal{E}$ can now be used to construct sets $\mathcal{E}'$ (defined as in (4)), and in view of (10), the estimate for $\widehat{f}_{\mathcal{E}', \varrho, \beta^{-1}}$ is *improved*. Continuing inductively and using $Dt - 0.5d_1 \ell \leq 3t/4$, after $d \leq d_1$ steps, we have

$$\widehat{f}_{\mathcal{E}', \varrho, d\beta^{-1}}(e, z) \leq e^{\varepsilon t} \varrho^{-\alpha} \psi_{\mathcal{E}', \varrho}(e, z), \quad (11)$$

which implies the dimension estimate (5) for the set $F' \subset B_\varepsilon(0, \beta)$ which is used in the definition of $\mathcal{E}'$, see (4).

We emphasize that our inductive scheme produces sets $\mathcal{E}'$ at every step $0 < d \leq d_1$ with an improved bound on $\widehat{f}_{\mathcal{E}', \varrho, d\beta^{-1}}$, however, it does not guarantee that (11) is only satisfied for $d \in J$. On the other hand, and as it was discussed above, the fact that our *stopping times* d satisfy $d_1 - d \leq 10^4 \varepsilon^{-1/2}$ is essential for us when we apply Proposition 5. We remedy this issue as follows: if (5) holds for some $\mathcal{E}'$ defined at step $d < d_1 - 10^4 \varepsilon^{-1/2}$, then we use the above inductive scheme to show that starting with $\mathcal{E}'$, the process *again* terminates (i.e., (11) is satisfied) in at most $10^4 \varepsilon^{-1/2}$ -many further steps.

On a related note, it should also be mentioned that ideally one would like to replace the interval of possible choices of $d \in J$ with the singleton $\{d_1\}$, i.e., to show that after exactly d_1 steps, one obtains sets which satisfy (5). Indeed, such statement can be obtained if one is content with restricting to $\alpha < 1/2$ — this can be achieved using estimates analogues to [12, Lemma 5.1] where one replaces the integral over $[0, 1]$ with integrals over *much smaller* intervals and by conditioning the random walk.

However, it is essential for us to work with $\alpha = 1 - \theta$ where $\theta > 0$ is rather small. For such choices of α , there are vectors $w \in \mathfrak{r}$ where the growth of $\|a_t u_r w\|$ is too slow. Indeed, in general, one can only guarantee that

$$\int_0^1 \|a_t u_r w\|^{-\alpha} dr \ll e^{-\theta t} \|w\|^{-\alpha}. \quad (12)$$

Using this general fact (which is an exercise in linear algebra) as an input, one can prove a version of Proposition 8 where

$$\left[d_1 - \frac{10^4}{\sqrt{\varepsilon}}, d_1 \right] \quad \text{is replaced by} \quad \left[d_1 - \frac{36}{\theta \sqrt{\varepsilon}}, d_1 \right],$$

in particular, the length of the interval cannot be made smaller than $(\theta \sqrt{\varepsilon})^{-1}$.

As it was discussed above, this improvement is pivotal to our analysis, and it is made possible by bringing to bear [36, 45, 46] in this step of the argument as well. Indeed, the *poor* rate in (12) is closely related to the existence of double zeroes for the map $r \mapsto (a_t u_r w)_{12}$ (the $(1, 2)$ -entry of $a_t u_r w$); we thus use [36, 45, 46], to control tangencies between the two parabolas $\{(u_r w_1)_{12} : r \in [0, 1]\}$ and $\{(u_r w_2)_{12} : r \in [0, 1]\}$ for most pairs $w_1, w_2 \in F$ and most $r \in [0, 1]$. This yields an improved version of (12) which we use crucially.

References

- [1] J. Bourgain, A. Furman, E. Lindenstrauss, S. Mozes, “Stationary measures and equidistribution for orbits of non-abelian semigroups on the torus”, *J. Am. Math. Soc.* **24** (2011), no. 1, p. 231-280.
- [2] J. Chaika, J. Smillie, B. Weiss, “Tremors and horocycle dynamics on the moduli space of translation surfaces”, <https://arxiv.org/abs/2004.04027>, 2020.
- [3] S. Chow, L. Yang, “An effective Ratner equidistribution theorem for multiplicative Diophantine approximation on planar lines”, <https://arxiv.org/abs/1902.06081>, 2019.
- [4] L. Clozel, H. Oh, E. Ullmo, “Hecke operators and equidistribution of Hecke points”, *Invent. Math.* **144** (2001), no. 2, p. 327-351.
- [5] S. G. Dani, “On orbits of unipotent flows on homogeneous spaces”, *Ergodic Theory Dyn. Syst.* **4** (1984), no. 1, p. 25-34.
- [6] ———, “On orbits of unipotent flows on homogeneous spaces. II”, *Ergodic Theory Dyn. Syst.* **6** (1986), no. 2, p. 167-182.
- [7] S. G. Dani, G. Margulis, “Values of quadratic forms at primitive integral points”, *Invent. Math.* **98** (1989), no. 2, p. 405-424.
- [8] ———, “Orbit closures of generic unipotent flows on homogeneous spaces of $SL(3, \mathbb{R})$ ”, *Math. Ann.* **286** (1990), no. 1-3, p. 101-128.
- [9] ———, “Asymptotic behaviour of trajectories of unipotent flows on homogeneous spaces”, *Proc. Indian Acad. Sci., Math. Sci.* **101** (1991), no. 1, p. 1-17.
- [10] M. Einsiedler, G. Margulis, A. Mohammadi, A. Venkatesh, “Effective equidistribution and property (τ) ”, *J. Am. Math. Soc.* **33** (2020), no. 1, p. 223-289.
- [11] M. Einsiedler, G. Margulis, A. Venkatesh, “Effective equidistribution for closed orbits of semisimple groups on homogeneous spaces”, *Invent. Math.* **177** (2009), no. 1, p. 137-212.
- [12] A. Eskin, G. Margulis, S. Mozes, “Upper bounds and asymptotics in a quantitative version of the Oppenheim conjecture”, *Ann. Math.* **147** (1998), no. 1, p. 93-141.
- [13] A. Eskin, M. Mirzakhani, “Invariant and stationary measures for the $SL(2, \mathbb{R})$ action on moduli space”, *Publ. Math., Inst. Hautes Étud. Sci.* **127** (2018), p. 95-324.
- [14] A. Eskin, M. Mirzakhani, A. Mohammadi, “Isolation, equidistribution, and orbit closures for the $SL(2, \mathbb{R})$ action on moduli space”, *Ann. Math.* **182** (2015), no. 2, p. 673-721.
- [15] A. Eskin, S. Mozes, “Margulis Functions and Their Applications”, in *Dynamics, Geometry, Number Theory: the impact of Margulis on modern mathematics* (D. Fisher, D. Kleinbock, G. Soifer, eds.), University of Chicago Press, 2022.
- [16] L. Flaminio, G. Forni, J. Tanis, “Effective equidistribution of twisted horocycle flows and horocycle maps”, *Geom. Funct. Anal.* **26** (2016), no. 5, p. 1359-1448.
- [17] G. Forni, “Limits of geodesic push-forwards of horocycle invariant measures”, *Ergodic Theory Dyn. Syst.* **41** (2021), no. 9, p. 2782-2804.
- [18] H. Garland, M. S. Raghunathan, “Fundamental Domains for Lattices in $\mathbb{R}$ -rank 1 Semisimple Lie Groups”, *Ann. Math.* **92** (1970), no. 2, p. 279-326.
- [19] B. Green, T. Tao, “The quantitative behaviour of polynomial orbits on nilmanifolds”, *Ann. Math.* **175** (2012), no. 2, p. 465-540.
- [20] W. He, N. de Saxcé, “Linear random walks on the torus”, <https://arxiv.org/abs/1910.13421v1>, 2019.
- [21] H. Jacquet, R. P. Langlands, *Automorphic forms on $GL(2)$* , Lecture Notes in Mathematics, vol. 114, Springer, 1970.
- [22] A. Käenmäki, T. Orponen, L. Venieri, “A Marstrand-type restricted projection theorem in $\mathbb{R}^3$ ”, <https://arxiv.org/abs/1708.04859v1>, 2017.
- [23] A. Katz, “Quantitative disjointness of nilflows from horospherical flows”, <https://arxiv.org/abs/1910.04675>, 2019.
- [24] W. Kim, “Effective equidistribution of expanding translates in the space of affine lattices”, <https://arxiv.org/abs/2110.00706>, 2021.
- [25] D. Y. Kleinbock, G. Margulis, “Bounded orbits of nonquasiunipotent flows on homogeneous spaces”, in *Sinai’s Moscow Seminar on Dynamical Systems*, American Mathematical Society Translations, Series 2, vol. 171, American Mathematical Society, 1996, p. 141-172.
- [26] E. Lindenstrauss, G. Margulis, “Effective estimates on indefinite ternary forms”, *Isr. J. Math.* **203** (2014), no. 1, p. 445-499.
- [27] E. Lindenstrauss, A. Mohammadi, “Polynomial effective density in quotients of $\mathbb{H}^3$ and $\mathbb{H}^2 \times \mathbb{H}^2$ ”, <https://arxiv.org/abs/2112.14562v1>, 2021.
- [28] E. Lindenstrauss, A. Mohammadi, G. Margulis, N. A. Shah, “Quantitative behavior of unipotent flows and an effective avoidance principle”, <https://arxiv.org/abs/1904.00290v1>, 2019.
- [29] G. Margulis, “The action of unipotent groups in a lattice space”, *Mat. Sb., N. Ser.* **86(128)** (1971), p. 552-556.
- [30] ———, “Indefinite quadratic forms and unipotent flows on homogeneous spaces”, in *Dynamical systems and ergodic theory (Warsaw, 1986)*, Banach Center Publications, vol. 23, Polish Academy of Sciences, 1989, p. 399-409.

- [31] ———, *Discrete subgroups of semisimple Lie groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 17, Springer, 1991.
- [32] A. Mohammadi, H. Oh, “Isolations of geodesic planes in the frame bundle of a hyperbolic 3-manifold”, <https://arxiv.org/abs/2002.06579v1>, 2020.
- [33] M. Ratner, “On measure rigidity of unipotent subgroups of semisimple groups”, *Acta Math.* **165** (1990), no. 3-4, p. 229-309.
- [34] ———, “On Raghunathan’s measure conjecture”, *Ann. Math.* **134** (1991), no. 3, p. 545-607.
- [35] ———, “Raghunathan’s topological conjecture and distributions of unipotent flows”, *Duke Math. J.* **63** (1991), no. 1, p. 235-280.
- [36] W. Schlag, “On continuum incidence problems related to harmonic analysis”, *J. Funct. Anal.* **201** (2003), no. 2, p. 480-521.
- [37] A. Selberg, “On discontinuous groups in higher-dimensional symmetric spaces”, in *Contributions to function theory (internat. Colloq. Function Theory, Bombay, 1960)*, Tata Institute of Fundamental Research, 1960, p. 147-164.
- [38] ———, “On the estimation of Fourier coefficients of modular forms”, in *Proc. Sympos. Pure Math., Vol. VIII*, American Mathematical Society, 1965, p. 1-15.
- [39] N. A. Shah, “Uniformly distributed orbits of certain flows on homogeneous spaces”, *Math. Ann.* **289** (1991), no. 2, p. 315-334.
- [40] ———, “Limit distributions of expanding translates of certain orbits on homogeneous spaces”, *Proc. Indian Acad. Sci., Math. Sci.* **106** (1996), no. 2, p. 105-125.
- [41] A. Strömbergsson, “An effective Ratner equidistribution result for $SL(2, \mathbb{R}) \ltimes \mathbb{R}^2$ ”, *Duke Math. J.* **164** (2015), no. 5, p. 843-902.
- [42] A. Venkatesh, “Sparse equidistribution problems, period bounds and subconvexity”, *Ann. Math.* **172** (2010), no. 2, p. 989-1094.
- [43] A. Weil, “On Discrete Subgroups of Lie Groups”, *Ann. Math.* **72** (1960), no. 2, p. 369-384.
- [44] ———, “Remarks on the Cohomology of Groups”, *Ann. Math.* **80** (1964), no. 1, p. 149-157.
- [45] T. H. Wolff, “Local smoothing type estimates on L^p for large p ”, *Geom. Funct. Anal.* **10** (2000), no. 5, p. 1237-1288.
- [46] J. Zahl, “ L^3 estimates for an algebraic variable coefficient Wolff circular maximal function”, *Rev. Mat. Iberoam.* **28** (2012), no. 4, p. 1061-1090.