

VARIATION OF CANONICAL HEIGHT AND EQUIDISTRIBUTION

LAURA DEMARCO AND NIKI MYRTO MAVRAKI

ABSTRACT. Let $\pi : E \rightarrow B$ be an elliptic surface defined over a number field K , where B is a smooth projective curve, and let $P : B \rightarrow E$ be a section defined over K with canonical height $\hat{h}_E(P) \neq 0$. In this article, we show that the function $t \mapsto \hat{h}_{E_t}(P_t)$ on $B(\overline{K})$ is the height induced from an adelically metrized line bundle with continuous semipositive metrics on B . The proof builds on work of Silverman and results from complex dynamical systems. Applying arithmetic equidistribution theorems (of Chambert-Loir, Thuillier, and Yuan), we obtain the equidistribution of points $t \in B(\overline{K})$ where P_t is torsion, and we give an explicit description of the limiting distribution on $B(\mathbb{C})$. Finally, combined with results of Masser and Zannier, we show that – for any non-special section P of a family of abelian varieties $A \rightarrow B$ that split as a product of elliptic curves – there is a positive lower bound on the height $\hat{h}_{A_t}(P_t)$, after excluding finitely many points $t \in B$, thus addressing a conjecture of Zhang from 1998.

1. INTRODUCTION

Suppose $E \rightarrow B$ is an elliptic surface defined over a number field K , so B is a smooth projective curve and all but finitely many fibers E_t , $t \in B(\overline{K})$, are smooth elliptic curves. We let $\hat{h}_E$ denote the Néron-Tate canonical height of E viewed as an elliptic curve over the function field $k = K(B)$; we let $\hat{h}_{E_t}$ denote the canonical height on a smooth fiber $E_t(\overline{K})$.

Suppose that $P : B \rightarrow E$ is a section defined over K for which $\hat{h}_E(P) \neq 0$, so, in particular, the points P_t on the fiber are not torsion in E_t for all t . Tate showed that the function

$$t \mapsto \hat{h}_{E_t}(P_t)$$

Email: demarco@northwestern.edu, Department of Mathematics, Northwestern University, Evanston, IL 60208-2730, USA.

Email: myrtomav@northwestern.edu, Department of Mathematics, Northwestern University, Evanston, IL 60208-2730, USA.

defines a Weil height on $B(\overline{K})$, up to a bounded error [Ta]. More precisely, there exists a divisor $D_P \in \text{Pic}(B) \otimes \mathbb{Q}$ of degree equal to $\hat{h}_E(P)$ so that

$$(1.1) \quad \hat{h}_{E_t}(P_t) = h_{D_P}(t) + O(1),$$

where h_{D_P} is a Weil height on $B(\overline{K})$ associated to D_P . In a series of three articles [Si1, Si3, Si4], Silverman refined statement (1.1) by analyzing the Néron decomposition of the canonical height on the fibers

$$\hat{h}_{E_t}(P_t) = \frac{1}{[K : \mathbb{Q}]} \sum_{v \in M_K} n_v \hat{\lambda}_{E_t, v}(P_t)$$

where M_K denotes the set of places of the number field K , and $n_v = [K_v : \mathbb{Q}_v]$ are the integers appearing in the product formula $\prod_{v \in M_K} |x|_v^{n_v} = 1$ for all $x \in K^*$.

In this article, we explain how Silverman's conclusions about the local functions $\hat{\lambda}_{E_t, v}(P_t)$ are precisely the input needed to show that $t \mapsto \hat{h}_{E_t}(P_t)$ is a “good” height function on the base curve B , from the point of view of equidistribution. Combining his work with methods from complex dynamics, as in [DWY], and the inequalities of Zhang on successive minima [Zh2, Zh1], we prove:

Theorem 1.1. *Let K be a number field and $k = K(B)$ for a smooth projective curve B defined over K . Fix an elliptic surface $E \rightarrow B$ defined over K and a point $P \in E(k)$ satisfying $\hat{h}_E(P) \neq 0$. Then*

$$t \mapsto \hat{h}_{E_t}(P_t),$$

defined for t with smooth fibers, extends to a height function on all of $B(\overline{K})$ induced from an adelically metrized ample line bundle $\overline{\mathcal{L}}_P$, with continuous semipositive metrics, satisfying

$$h_{\overline{\mathcal{L}}_P}(B) := c_1(\overline{\mathcal{L}}_P)^2 / (2c_1(\mathcal{L}_P)) = 0.$$

Theorem 1.1 implies that our height function on B satisfies the hypotheses of the equidistribution theorems of Chambert-Loir, Thuillier, and Yuan for points of small height on curves [CL1, Th, Y], and we deduce the following:

Corollary 1.2. *Let K be a number field and $k = K(B)$ for a smooth projective curve B defined over K . Fix an elliptic surface $E \rightarrow B$ defined over K and a point $P \in E(k)$ satisfying $\hat{h}_E(P) \neq 0$. There is a collection of probability measures $\mu_P = \{\mu_{P, v} : v \in M_K\}$ on the Berkovich analytifications B_v^{an} such that for any infinite, non-repeating sequence $t_n \in$*

$B(\overline{K})$ with

$$\hat{h}_{E_{t_n}}(P_{t_n}) \rightarrow 0 \text{ as } n \rightarrow \infty,$$

the discrete measures

$$\frac{1}{|\mathrm{Gal}(\overline{K}/K) \cdot t_n|} \sum_{t \in \mathrm{Gal}(\overline{K}/K) \cdot t_n} \delta_t$$

converge weakly on B_v^{an} to the measure $\mu_{P,v}$ at each place v of K .

Remark 1.3. The measures $\mu_{P,v}$ of Corollary 1.2 are not difficult to describe, at least at the archimedean places. At each archimedean place v , there is a canonical positive $(1, 1)$ -current T_v on the surface $E(\mathbb{C})$ (with continuous potentials away from the singular fibers) which restricts to the Haar measure on each smooth fiber $E_t(\mathbb{C})$. The measure $\mu_{P,v}$ on $B(\mathbb{C})$ is just the pull-back of this current by the section P . Moreover, at every place, the measure $\mu_{P,v}$ is the Laplacian of the local height function $\hat{\lambda}_{E_t,v}(P_t)$, away from its singularities. We give more details about (and a dynamical perspective on) the construction of the current T_v in Section 3.

As a consequence of Theorem 1.1, and combined with the work of Masser and Zannier [MZ1, MZ2, MZ3], we obtain the so-called Bogomolov extension of their theorems. Fix integer $m \geq 2$, and suppose that $E_i \rightarrow B$ is an elliptic surface over a curve B , defined over $\overline{\mathbb{Q}}$, for $i = 1, \dots, m$. We consider sections P of the fiber product $A = E_1 \times_B \dots \times_B E_m \rightarrow B$ defined over $\overline{\mathbb{Q}}$. We say that a section $P = (P_1, P_2, \dots, P_m)$ is *special* if the following two conditions are satisfied.

- (1) For each $i = 1, \dots, m$, either P_i is torsion on E_i or $\hat{h}_{E_i}(P_i) \neq 0$.
- (2) For any pair $i, j \in \{1, \dots, m\}$ such that neither P_i nor P_j is torsion, there are an isogeny $\phi : E_i \rightarrow E_j$ and nonzero group endomorphisms a, b of E_j so that $a \circ \phi(P_i) = b(P_j)$.

Note that the first condition is satisfied for every section P of A if we assume that each E_i is non-isotrivial, because then $P_i \in E_i(k)$ is torsion if and only if $\hat{h}_{E_i}(P_i) = 0$.

If a family of abelian varieties $A \rightarrow B$ is isogenous to a fiber product of elliptic surfaces (after performing a base change $B' \rightarrow B$ if needed), we say that a *section of A is special* if it is special on the fiber product.

It is not hard to see that a special section P will specialize to a torsion point P_t in the fiber A_t for infinitely many $t \in B(\overline{K})$. For a proof see [Za, Chapter 3] or, for a dynamical proof, see [De1]. The converse statement is also true, but it is much more difficult: Masser and

Zannier proved that if P_t is torsion in A_t for infinitely many $t \in B(\overline{\mathbb{Q}})$, then the section P must be special [MZ2, MZ3]. We use Theorem 1.1 to extend these results of Masser-Zannier from points of height 0 to points of small height:

Theorem 1.4. *Let B be a quasiprojective smooth algebraic curve defined over $\overline{\mathbb{Q}}$. Suppose $A \rightarrow B$ is a family of abelian varieties of relative dimension $m \geq 2$ defined over $\overline{\mathbb{Q}}$ which is isogenous to a fibered product of $m \geq 2$ elliptic surfaces. Let $\mathcal{L}$ be a line bundle on A which restricts to an ample and symmetric line bundle on each fiber A_t , and let $\hat{h}_t$ be the induced Néron-Tate canonical height on A_t , for each $t \in B(\overline{\mathbb{Q}})$. For each non-special section $P : B \rightarrow A$ defined over $\overline{\mathbb{Q}}$, there is a constant $c = c(\mathcal{L}, P) > 0$ so that*

$$\{t \in B(\overline{\mathbb{Q}}) : \hat{h}_t(P_t) < c\}$$

is finite.

If $A \rightarrow B$ is isotrivial, then Theorem 1.4 is a special case of the Bogomolov Conjecture, proved by Ullmo and Zhang [Zh3, U]. A key ingredient in their proofs is the equidistribution theorem of Szpiro, Ullmo, and Zhang [SUZ]. In his 1998 ICM lecture notes [Zh4], Zhang presented a conjecture about geometrically simple families of abelian varieties, which stated, in its most basic form:

Conjecture 1.5 (Zhang). *Let B be a quasiprojective smooth algebraic curve defined over $\overline{\mathbb{Q}}$. Suppose $A \rightarrow B$ is a non-isotrivial family of abelian varieties with fiber dimension > 1 , defined over $\overline{\mathbb{Q}}$ with a simple generic fiber. Let $\mathcal{L}$ be a line bundle on A which restricts to an ample and symmetric line bundle on each fiber A_t , and let $\hat{h}_t$ be the induced Néron-Tate canonical height on A_t , for each $t \in B(\overline{\mathbb{Q}})$. For each non-torsion section $P : B \rightarrow A$ defined over $\overline{\mathbb{Q}}$, there is a constant $c = c(\mathcal{L}, P) > 0$ so that*

$$\{t \in B(\overline{\mathbb{Q}}) : \hat{h}_t(P_t) < c\}$$

is finite.

When the dimension of the fibers A_t is equal to 2, the finiteness of $\{t \in B(\overline{\mathbb{Q}}) : \hat{h}_t(P_t) = 0\}$ for sections as in Conjecture 1.5 was established recently by Masser and Zannier in [MZ4]. It is well known that the conclusion of Conjecture 1.5 can fail to hold if A is not simple and certainly fails if it is a family of elliptic curves, as mentioned above. However, the results of Masser and Zannier in their earlier work [MZ2, MZ3] suggested a formulation of Zhang's conjecture for the non-simple case when A splits as a product of elliptic curves where the

“non-torsion” hypothesis on P should be replaced by “non-special”; this is what we proved in our Theorem 1.4.

Remark 1.6. Theorem 1.1, Corollary 1.2, and Theorem 1.4 were obtained in the special case of the Legendre family $E_t = \{y^2 = x(x-1)(x-t)\}$ over $B = \mathbb{P}^1$ and the abelian variety $A_t = E_t \times E_t$, for sections P with x -coordinates in $\overline{\mathbb{Q}}(t)$ in [DWY], using methods from complex dynamical systems, without appealing to Silverman and Tate’s results on the height function. Moreover, restricting further to sections P with constant x -coordinate (in $\mathbb{P}^1(\overline{\mathbb{Q}})$), Theorem 1.4 was obtained without relying on the theorems of Masser and Zannier and gave an alternate proof of their result. This includes the special case treated by Masser and Zannier in their article [MZ1]. For sections with constant x -coordinate, the hypothesis on P (that $\hat{h}_E(P) \neq 0$) is equivalent to asking that $x(P) \neq 0, 1, \infty$ [DWY, Proposition 1.4].

Comments and acknowledgements. This project was motivated, in part, by experiments to visualize Silverman’s results on the variation of canonical height [Si1, Si3, Si4] in terms of the measures $\mu_{P,v}$ at archimedean places, and to examine their dependence on P . In particular, the measure detects the failure of the local height function $\hat{\lambda}_{E_t,v}(P_t)$ to be harmonic; compare with the comments on non-analyticity preceding Theorem I.0.3 of [Si1]. The images appearing in Section 6 were first presented at the conference in honor of Silverman’s birthday, August 2015.

We thank Charles Favre, Dragos Ghioca, Robert Rumely, Joseph Silverman, and Amaury Thuillier for helpful suggestions. Our research was supported by the National Science Foundation and the Simons Foundation.

2. SILVERMAN’S WORK

2.1. Preliminaries. Let $\mathcal{F}$ be a product formula field of characteristic 0, so there exists a family $M_{\mathcal{F}}$ of non-trivial absolute values on $\mathcal{F}$ and a collection of positive integers n_v for $v \in M_{\mathcal{F}}$ so that

$$\prod_{v \in M_{\mathcal{F}}} |x|_v^{n_v} = 1$$

for all $x \in \mathcal{F}^*$. Let $E/\mathcal{F}$ be an elliptic curve with origin O , expressed in Weierstrass form as

$$E = \{y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6\}$$

with discriminant Δ . Denote by

$$\hat{h}_E : E(\overline{\mathcal{F}}) \rightarrow [0, \infty)$$

the Néron-Tate canonical height function; it can be defined by

$$\hat{h}_E(P) = \frac{1}{2} \lim_{n \rightarrow \infty} \frac{h(x([n]P))}{n^2}$$

where h is the naive Weil height on $\mathbb{P}^1$ and $x : E \rightarrow \mathbb{P}^1$ is the degree 2 projection to the x -coordinate.

For each $v \in M_{\mathcal{F}}$, we let $\mathcal{F}_v$ denote the completion of $\mathcal{F}$ with respect to $|\cdot|_v$ and $\mathbb{C}_v$ denote the completion of the algebraic closure of $\mathcal{F}_v$ with respect to $|\cdot|_v$. For each v , we fix an embedding of $\overline{\mathcal{F}}$ into $\mathbb{C}_v$. The canonical height has a decomposition into local heights, as

$$\hat{h}_E(P) = \frac{1}{|\text{Gal}(\overline{\mathcal{F}}/\mathcal{F}) \cdot P|} \sum_{Q \in \text{Gal}(\overline{\mathcal{F}}/\mathcal{F}) \cdot P} \sum_{v \in M_{\mathcal{F}}} n_v \hat{\lambda}_{E,v}(Q)$$

for $P \in E(\overline{\mathcal{F}}) \setminus \{O\}$, with the local heights $\hat{\lambda}_{E,v}$ characterized by three properties [Si2, Chapter 6, Theorem 1.1, page 455]:

- (1) $\hat{\lambda}_{E,v}$ is continuous on $E(\mathbb{C}_v) \setminus \{O\}$ and bounded on the complement of any v -adic neighborhood of O ;
- (2) the limit of $\hat{\lambda}_{E,v}(P) - \frac{1}{2} \log |x(P)|_v$ exists as $P \rightarrow O$ in $E(\mathbb{C}_v)$; and
- (3) for all $P = (x, y) \in E(\mathbb{C}_v)$ with $[2]P \neq O$,

$$\hat{\lambda}_{E,v}([2]P) = 4\hat{\lambda}_{E,v}(P) - \log |2y + a_1x + a_3|_v + \frac{1}{4} \log |\Delta|_v.$$

2.2. Variation of canonical height: the set up. Now let K be a number field and $E \rightarrow B$ an elliptic surface defined over a number field K with zero section $O : B \rightarrow E$. Let $P : B \rightarrow E$ be a non-zero section defined over K , and assume that

$$\hat{h}_E(P) \neq 0$$

when viewing P as a point on the elliptic curve E defined over $k = \overline{K}(B)$. For each $t \in B(\overline{K})$ such that the fiber E_t is non-singular, we have a point $P_t \in E_t(\overline{K})$. We will investigate the function

$$t \mapsto \hat{h}_{E_t}(P_t)$$

which is well defined at all but finitely many $t \in B(\overline{K})$. Furthermore, via the embedding of $\overline{K}$ into $\mathbb{C}_v$ for each place $v \in M_K$, we may view $E \rightarrow B$ as defined over $\mathbb{C}_v$ and consider the Néron local heights $\hat{\lambda}_{E_t, v}(P_t)$ on the non-singular fibers E_t as functions of $t \in B(\mathbb{C}_v)$.

Let $D_E(P)$ be the $\mathbb{Q}$ -divisor on B defined by

$$(2.1) \quad D_E(P) = \sum_{\gamma \in B(\overline{K})} \hat{\lambda}_{E, \text{ord}_\gamma}(P) \cdot (\gamma).$$

Here, $\hat{\lambda}_{E, \text{ord}_\gamma}(P)$ is the local canonical height of the point P on the elliptic curve E over $k = \overline{K}(B)$ at the place ord_γ , for each $\gamma \in B(\overline{K})$. The degree of $D_E(P)$ is equal to $\hat{h}_E(P)$. It follows from the definitions that $\text{supp } D_E(P)$ is a subset of the finite set

$$\{t \in B(\overline{K}) : E_t \text{ is singular}\} \cup \{t \in B(\overline{K}) : P_t = O_t\}.$$

By enlarging K , we may assume that the support of $D_E(P)$ is contained in $B(K)$.

Remark 2.1. That $D_E(P)$ is a $\mathbb{Q}$ -divisor is standard, following from the fact that the numbers $\hat{\lambda}_{E, \text{ord}_\gamma}(P)$ can be viewed as arithmetic intersection numbers on a Néron local model. See [Si2, Chapter III, Theorem 9.3] for a proof that $\hat{h}_E(P) \in \mathbb{Q}$; see [CS, Section 6 and p. 203] and [La1, Chapter 11 Theorem 5.1] for proofs that each local function $\hat{\lambda}_{E, v}$ also takes values in $\mathbb{Q}$; see [DG, Theorem 1.1] for a dynamical proof.

2.3. Variation of canonical height: quasi triviality. Let $h_{D_E(P)}$ be an analytic Weil height on $B(\overline{K})$ as defined in [Si4, §3 Example 1(a)]. That is, we let g be the genus of B , and for each point $\gamma \in B(K)$, we choose an element ξ_γ of $K(B)$ which has a pole of order $2g + 1$ at γ and no other poles. For each non-archimedean place v of K , set

$$\lambda_{D_E(P), v}(t) = \frac{1}{2g + 1} \sum_{\gamma \in B(K)} \hat{\lambda}_{E, \text{ord}_\gamma}(P) \log^+ |\xi_\gamma(t)|_v$$

for all $t \in B(\mathbb{C}_v) \setminus \text{supp } D_E(P)$. For archimedean places v , the local height is defined by

$$\lambda_{D_E(P), v}(t) = \frac{1}{2(2g + 1)} \sum_{\gamma \in B(K)} \hat{\lambda}_{E, \text{ord}_\gamma}(P) \log (1 + |\xi_\gamma(t)|_v^2).$$

We set

$$h_{D_E(P)}(t) = \frac{1}{|\text{Gal}(\overline{K}/K) \cdot t|} \sum_{s \in \text{Gal}(\overline{K}/K) \cdot t} \sum_{v \in M_K} \lambda_{D_E(P), v}(s)$$

for all $t \in B(\overline{K})$. For fixed choices of ξ_γ , we will call the associated height $h_{D_E(P)}$ our “reference height” for the divisor $D_E(P)$. Silverman proved:

Theorem 2.2. [Si4, Theorem III.4.1] *For any choice of reference height $h_{D_E(P)}$, there is a finite set S of places so that*

$$\hat{\lambda}_{E_t,v}(P_t) = \lambda_{D_E(P),v}(t)$$

for all $t \in B(\overline{K}) \setminus \text{supp } D_E(P)$ and all $v \in M_K \setminus S$.

2.4. Variation of canonical height: continuity. Fix a point $t_0 \in B(\overline{K})$ and a uniformizer $u \in \overline{K}(B)$ for t_0 , and consider the function

$$(2.2) \quad V_{P,t_0,v}(t) := \hat{\lambda}_{E_t,v}(P_t) + \hat{\lambda}_{E,\text{ord}_{t_0}}(P) \log |u(t)|_v,$$

which is not *a priori* defined at t_0 . Theorem 2.2 implies that

$$V_{P,t_0,v} \equiv 0$$

for all but finitely many places v in a v -adic neighborhood of t_0 . Silverman also proved the following:

Theorem 2.3. [Si3, Theorem II.0.1] *Fix $t_0 \in B(\overline{K})$ and a uniformizer u at t_0 . For all $v \in M_K$, there exists a neighborhood $U \subset B(\mathbb{C}_v)$ containing t_0 so that the function $V_{P,t_0,v}$ of (2.2) extends to a continuous function on U .*

3. A DYNAMICAL PERSPECTIVE

Recall that the Néron-Tate height $\hat{h}_E$ and its local counterparts $\hat{\lambda}_{E,v}$ can be defined dynamically. Letting E be an elliptic curve defined over a number field K , the multiplication-by-2 endomorphism ϕ on E descends to a rational function of degree 4 on $\mathbb{P}^1$, via the standard quotient π identifying a point P with its additive inverse:

$$(3.1) \quad \begin{array}{ccc} E & \xrightarrow{\phi} & E \\ \pi \downarrow & & \downarrow \pi \\ \mathbb{P}^1 & \xrightarrow{f_\phi} & \mathbb{P}^1 \end{array}$$

An elementary, but key, observation is that a point is torsion on E if and only if its quotient in $\mathbb{P}^1$ is preperiodic for f_ϕ . The height $\hat{h}_E$ on $E(\overline{K})$ satisfies

$$\hat{h}_E(P) = \frac{1}{2} \lim_{n \rightarrow \infty} \frac{1}{4^n} h(f_\phi^n(\pi P))$$

where h is the standard logarithmic Weil height on $\mathbb{P}^1(\overline{K})$. Now let $E \rightarrow B$ be an elliptic surface defined over a number field K , and let $P : B \rightarrow E$ be a section, also defined over K . In this section, we use this dynamical perspective to give a proof of the subharmonicity of the local height functions $t \mapsto \hat{\lambda}_{E_t, v}(P_t)$ and the extensions $V_{P, t_0, v}$ of (2.2). We will present this fact as an immediate consequence of now-standard complex-dynamical convergence arguments, at least when the fiber E_t is smooth and the local height $\hat{\lambda}_{E_t, v}(P_t)$ is finite. Near singular fibers, we utilize the maximum principle and standard results on removable singularities for subharmonic functions. The same reasoning applies in both archimedean and non-archimedean settings.

In §3.3 we provide the background to justify the explicit description of the distributions $\mu_{P, v}$ from Corollary 1.2 at the archimedean places v of K , as mentioned in Remark 1.3.

3.1. Canonical height and escape rates. As in §2.1, we let E be an elliptic curve in Weierstrass form, defined over a product-formula field $\mathcal{F}$ of characteristic 0. We define a rational function $f = \phi/\psi$ on $\mathbb{P}^1$ by the formula

$$f(x(P)) = x([2]P)$$

for all $P \in E(\overline{\mathcal{F}})$. Here $x(P)$ is the x -coordinate for a point P ; this function x plays the role of π in (3.1). In coordinates, we have $\phi(x) = x^4 - b_4x^2 - 2b_6x - b_8$ and $\psi(x) = 4x^3 + b_2x^2 + 2b_4x + b_6 = (2y + a_1x + a_3)^2$ for $P = (x, y)$.

By a *lift* of f , we mean any homogeneous polynomial map F on $\mathbb{A}^2$, defined over $\mathcal{F}$, so that $\tau \circ F = f \circ \tau$, where $\tau : \mathbb{A}^2 \setminus \{(0, 0)\} \rightarrow \mathbb{P}^1$ is the tautological projection. A *lift* of a point $x \in \mathbb{P}^1$ is a choice of $X \in \mathbb{A}^2 \setminus \{(0, 0)\}$ so that $\tau(X) = x$.

The *standard lift* of f will be the map $F : \mathbb{A}^2 \rightarrow \mathbb{A}^2$ defined by

$$(3.2) \quad F(z, w) = (w^4\phi(z/w), w^4\psi(z/w)).$$

For each $v \in \mathcal{M}_{\mathcal{F}}$, the v -adic *escape rate* is defined by

$$\mathcal{G}_{F, v}(z, w) = \lim_{n \rightarrow \infty} \frac{\log \|\mathbf{F}^n(z, w)\|_v}{4^n}$$

where

$$\|(z, w)\|_v = \max\{|z|_v, |w|_v\}.$$

Any other lift of f is of the form cF for some $c \in \mathcal{F}^*$; observe that

$$\mathcal{G}_{cF, v} = \mathcal{G}_{F, v} + \frac{1}{3} \log |c|_v.$$

Note that

$$\mathcal{G}_{F,v}(\alpha x, \alpha y) = \mathcal{G}_{F,v}(x, y) + \log |\alpha|_v$$

for any choice of lift F . Furthermore, $\mathcal{G}_{F,v}$ is continuous on $\mathbb{C}_v^2 \setminus \{(0, 0)\}$, as proved in the archimedean case by [HP, FS]. For non-archimedean absolute values v , $\mathcal{G}_{F,v}$ extends continuously to the product of Berkovich affine lines $\mathbb{A}_v^{1,an} \times \mathbb{A}_v^{1,an} \setminus \{(0, 0)\}$ [BR, Chapter 10].

Proposition 3.1. *For the standard lift F of f , and for each place v of $\mathcal{F}$, the local canonical height function satisfies*

$$\hat{\lambda}_{E,v}(P) = \frac{1}{2}\mathcal{G}_{F,v}(x, y) - \frac{1}{2}\log |y|_v - \frac{1}{12}\log |\Delta|_v$$

where $x(P) = (x : y)$.

Proof. The proof is immediate from the properties of $\mathcal{G}_{F,v}$ by checking the three characterizing conditions for $\hat{\lambda}_{E,v}$. $\square$

3.2. Variation of canonical height: subharmonicity. Now let K be a number field and $E \rightarrow B$ an elliptic surface defined over K with zero section $O : B \rightarrow E$. Let $k = \overline{K}(B)$; viewing E as an elliptic curve defined over k , we also fix a point $P \in E(k)$. Recall the function $V_{P,t_0,v}(t)$ defined in (2.2).

Theorem 3.2. *For every $t_0 \in B(\overline{K})$ and uniformizer u in k at t_0 , the function*

$$V_{P,t_0,v}(t) := \hat{\lambda}_{E_t,v}(P_t) + \hat{\lambda}_{E,\text{ord}_{t_0}}(P) \log |u(t)|_v,$$

extends to a continuous and subharmonic function on a neighborhood of t_0 in the Berkovich analytification B_v^{an} .

The continuity was already established in Theorem 2.3, though it was not explicitly stated for the Berkovich space. The argument below takes care of that. We begin with a lemma.

Lemma 3.3. *Fix $\alpha \in k^*$ and $t_0 \in B(\overline{K})$. Let $u \in k$ be a uniformizer at t_0 . For each place v of K , the function*

$$t \mapsto \log |\alpha_t|_v - (\text{ord}_{t_0} \alpha) \log |u(t)|_v$$

is harmonic in a neighborhood of t_0 in the Berkovich analytification B_v^{an} .

Proof. This is Silverman's [Si3, Lemma II.1.1(c)] plus a removable singularities lemma for harmonic functions. See also [BR, Proposition 7.19] for the extension of a harmonic function to a disk in the Berkovich space B_v^{an} . $\square$

Fix $P \in E(k)$. Let F and X be lifts of f and $x(P)$ to k^2 , respectively. Iterating F , we set

$$(A_n, B_n) := F^n(X) \in k^2$$

and observe that

$$(3.3) \quad \mathcal{G}_{F, \text{ord}_{t_0}}(X) = - \lim_{n \rightarrow \infty} \frac{\min\{\text{ord}_{t_0} A_n, \text{ord}_{t_0} B_n\}}{4^n}$$

from the definition of the escape rate. We let F_t and X_t denote the specializations of F and X at a point $t \in B(\overline{K})$; they are well defined for all but finitely many t . Observe that if F is the standard lift for E then so is F_t for all t .

Proposition 3.4. *Fix $P \in E(k)$, $t_0 \in B(\overline{K})$, and $v \in M_K$. For any choice of lifts F of f and X of $x(P)$, the function*

$$G_P(t; v) := \mathcal{G}_{F_t, v}(X_t) + \mathcal{G}_{F, t_0}(X) \log |u(t)|_v$$

extends to a continuous and subharmonic function in a neighborhood of t_0 in B_v^{an} .

Proof. First observe that the conclusion does not depend on the choices of F and X . Indeed,

$$\begin{aligned} \mathcal{G}_{c_t F_t, v}(\alpha_t X_t) + \mathcal{G}_{c F, t_0}(\alpha X) \log |u(t)|_v &= \mathcal{G}_{F_t, v}(X_t) + \mathcal{G}_{F, t_0}(X) \log |u(t)|_v \\ &\quad + \frac{1}{3} (\log |c_t|_v - (\text{ord}_{t_0} c) \log |u(t)|_v) \\ &\quad + \log |\alpha_t|_v - (\text{ord}_{t_0} \alpha) \log |u(t)|_v \end{aligned}$$

for any $c, \alpha \in k^*$. So by Lemma 3.3 the function $G_P(t; v)$ is continuous and subharmonic for one choice if and only if it is continuous and subharmonic for all choices.

Let F be the standard lift of f . Suppose that $P = O$. Since $F(1, 0) = (1, 0)$, we compute that

$$G_O(t; v) = \mathcal{G}_{F_t, v}(1, 0) + \mathcal{G}_{F, t_0}(1, 0) \log |u(t)|_v \equiv 0.$$

Now suppose that $P \neq O$. Fix $t_0 \in B(\overline{K})$ and local uniformizer u at t_0 . Choose a lift F of f so that the coefficients of F have no poles at t_0 , with $F_{t_0} \neq (0, 0)$. Choose lift X of $x(P)$ so that X_t is well defined for all t near t_0 and $X_{t_0} \neq (0, 0)$. As above, we write

$$F^n(X) = (A_n, B_n)$$

and put

$$a_n = \min\{\text{ord}_{t_0} A_n, \text{ord}_{t_0} B_n\}$$

so that $a_n \geq 0$ for all n and $a_0 = 0$. Set

$$F_n(t) = F_t^n(X_t)/u(t)^{a_n}.$$

For each place v of K , we set

$$h_{n,v}(t) = \frac{\log \|F_n(t)\|_v}{4^n}.$$

By construction, the limit of $h_{n,v}$ (for t near t_0 with $t \neq t_0$) is exactly the function G_P for these choices. In fact, for t in a small neighborhood of t_0 , but with $t \neq t_0$, the function f_t on $\mathbb{P}^1$ is a well-defined rational function of degree 4; so the specialization of the homogeneous polynomial map F_t satisfies $F_t^{-1}\{(0,0)\} = \{(0,0)\}$. Furthermore, as the coefficients of F_t depend analytically on t , the functions $h_{n,v}$ converge locally uniformly to the function G_P away from $t = t_0$. This can be seen with a standard telescoping sum argument, used often in complex dynamics, as in [BH, Proposition 1.2]. In particular, G_P is continuous on a punctured neighborhood of t_0 .

At the archimedean places v , and for each n , the function $h_{n,v}$ is clearly continuous and subharmonic in a neighborhood of t_0 . At non-archimedean places v , this definition extends to a Berkovich disk around t_0 , setting

$$h_{n,v}(t) = \frac{1}{4^n} \max\{\log[A_n(T)/T^{a_n}]_t, \log[B_n(T)/T^{a_n}]_t\}$$

where $[\cdot]_t$ is the seminorm on $K[[T]]$ associated to the point t . Each of these functions $h_{n,v}$ is continuous and subharmonic for t in a Berkovich disk around t_0 . (Compare [BR] Example 8.7, Proposition 8.26(D), and equation (10.9).)

Lemma 3.5. *For all v , and by shrinking the radius r if necessary, the functions $h_{n,v}$ are uniformly bounded from above on the (Berkovich) disk D_r .*

Proof. As observed above, the functions $h_{n,v}$ converge locally uniformly away from $t = t_0$ to the continuous function $G_P(t)$. Choose a small radius r , and let

$$M_v = \sup_n \max_{|t|_v=r} h_{n,v}(t)$$

which is finite by the convergence. Because the functions are subharmonic, the Maximum Principle implies that $h_{n,v}(t) \leq M_v$ throughout the disk of radius r , for all n . For the non-archimedean places, there is also a Maximum Principle on the Berkovich disk, where the role of the circle of radius r is played by the Type II point associated to the disk of radius r (see [BR] Proposition 8.14). $\square$

We can now complete the proof of Proposition 3.4. As each $h_{n,v}$ is subharmonic, and the functions are uniformly bounded from above on the disk by Lemma 3.5, we know that the (upper-semicontinuous regularization of the) limsup of these functions is subharmonic. See [BR] Proposition 8.26(F) for a proof in the non-archimedean case. $\square$

Proof of Theorem 3.2. Subharmonicity now follows from Proposition 3.1, Lemma 3.3, and Proposition 3.4. The continuity at each archimedean place follows from Theorem 2.3. The continuity at each non-archimedean place is a combination of the continuity on the punctured Berkovich disk (as in the proof of Proposition 3.4) and the continuity on Type I (classical) points given in Theorem 2.3. $\square$

3.3. The measures on the base. Here we provide more details about the description of the measures appearing in the statement of Corollary 1.2, as discussed in Remark 1.3.

Fix an archimedean place v and any point $t_0 \in B(\overline{K})$. Choosing a uniformizer u at t_0 , recall the definition of $V_{P,t_0,v}$ from (2.2). We define

$$(3.4) \quad \mu_{P,v} := dd^c V_{P,t_0,v}(t)$$

on a neighborhood of t_0 in B_v^{an} ; note that this is independent of the choice of u . Note that $\mu_{P,v}$ can be expressed as

$$\mu_{P,v} = dd^c \hat{\lambda}_{E_t,v}(P_t)$$

for t outside of the finitely many points in the support of the divisor $D_E(P)$ or where the fiber E_t is singular. Note, further, that $\mu_{P,v}$ assigns no mass to any individual point t_0 , because the potentials are bounded by Theorem 3.2. The details on the metric and the equidistribution theorem in Section 4 will show that these are exactly the measures that arise as the distribution of the points of small height in Corollary 1.2.

It is well known that the local height function on a smooth elliptic curve is a potential for the Haar measure. That is, for fixed t we have

$$dd^c \hat{\lambda}_{E_t,v}(\cdot) = \omega_t - \delta_o$$

where ω_t is the normalized Haar measure on E_t and δ_o is a delta-mass supported at the origin of E_t ; see, e.g., [La2, Theorem II.5.1]. We present an alternative proof of this fact related to dynamics as part of Proposition 3.6, as a consequence of Proposition 3.1.

Proposition 3.6. *Let $E \rightarrow B$ be an elliptic surface and $P : B \rightarrow E$ a section, both defined over a number field K . Let $\mathcal{S} \subset E$ be the union of the finitely many singular fibers in E .*

For each archimedean place v of K , there is a positive, closed $(1,1)$ current T_v on $E \setminus \mathcal{S}$ with locally continuous potentials so that $T_v|_{E_t}$ is the Haar measure on each smooth fiber, and P^*T_v is equal to the measure $\mu_{P,v}$.

Remark 3.7. As T_v has continuous potentials, the restriction $T_v|_{E_t}$ and the pullback P^*T_v are well defined. That is, we have $T_v|_{E_t} = dd^c(g|_{E_t})$ where g is a locally defined potential of T_v , and $P^*T_v = dd^c(g \circ P)$ locally on B . The measure $\mu_{P,v}$ has no atoms, so it is determined by T_v along the image of P in $E \setminus \mathcal{S}$.

Proof of Proposition 3.6. Let us fix any small neighborhood U in the base curve $B(\mathbb{C})$ so that all fibers E_t are smooth for $t \in U$. Let f_t be the map on $\mathbb{P}^1$ defined in §3.1; by shrinking U if necessary, we can find lifts F_t of f_t that are holomorphic in $t \in U$. From [HP, FS] (or the proof of [BH, Proposition 1.2]), we know that the escape rate

$$\mathcal{G}_{F_t,v}(z, w) = \lim_{n \rightarrow \infty} \frac{\log \|F_t^n(z, w)\|_v}{4^n}$$

is continuous and plurisubharmonic as a function of $(t, z, w) \in U \times (\mathbb{C}^2 \setminus \{(0, 0)\})$. The current

$$dd^c \mathcal{G}_{F_t,v}(z, w)$$

projects to a closed and positive $(1,1)$ -current $\hat{T}_v$ on the complex surface $U \times \mathbb{P}^1$, with locally continuous potentials. This current $\hat{T}_v$ has the property that, restricted to each fiber $\mathbb{P}^1$, its total mass is 1; and the induced measure on the fiber is the measure of maximal entropy for the rational map f_t [Ly, HP].

The restriction $E|_U$ of the elliptic surface E over U maps with degree 2 to the complex surface $U \times \mathbb{P}^1$ by the projection π of (3.1). The current $\hat{T}_v$ can be pulled back to E as $\frac{1}{2} dd^c(g \circ \pi)$ where g is a locally-defined continuous potential for $\hat{T}_v$. Covering the base of $E \setminus \mathcal{S}$ by sets of the form U , the local definitions glue to form the closed, positive $(1,1)$ -current T_v on $E \setminus \mathcal{S}$.

If $P : B \rightarrow E$ is a section defined over the number field K , then P^*T_v has potential given locally by

$$\frac{1}{2} g \circ \pi \circ P = \frac{1}{2} \mathcal{G}_{F_t,v}(X_t)$$

for any lift X_t of $\pi(P_t) \in \mathbb{P}^1$. Proposition 3.1 yields that P^*T_v must coincide with the measure $\mu_{P,v}$ defined in (3.4).

Finally, to conclude that $T_v|_{E_t}$ is equal to the normalized Haar measure ω_t , we may use the well-known dynamical fact that for each fixed t in the base, the measure ω_t projects by π to $\mathbb{P}^1$ to the unique measure of maximal entropy for the map f_t ; see, e.g., [Mi, §7]. $\square$

4. THE ADELIC METRIC AND EQUIDISTRIBUTION

In this section we give the proofs of Theorem 1.1 and Corollary 1.2.

We first outline the proofs. Let $E \rightarrow B$ be an elliptic surface defined over a number field K with zero section $O : B \rightarrow E$, and let $P : B \rightarrow E$ be a section also defined over K so that $\hat{h}_E(P) \neq 0$. Recall from §2.2 that we introduced a $\mathbb{Q}$ -divisor

$$D_E(P) = \sum_{\gamma \in B(\overline{K})} \hat{\lambda}_{E, \text{ord}_\gamma}(P) \cdot (\gamma)$$

on B . By enlarging K , we may assume that $\text{supp } D_E(P)$ lies in $B(K)$. We will define an adelic metric on the ample line bundle $\mathcal{L}_P$ associated to the $\mathbb{Q}$ -divisor $D_E(P)$, inducing a height function $h_{\overline{\mathcal{L}}_P}$ such that

$$h_{\overline{\mathcal{L}}_P}(t) = \hat{h}_{E_t}(P_t) \text{ for all but finitely many } t \in B(\overline{K})$$

and

$$h_{\overline{\mathcal{L}}_P}(t) \geq 0 \text{ for all } t \in B(\overline{K}).$$

Applying Silverman's results on the variation of canonical height, Theorems 2.2 and 2.3, we will deduce that the metric is continuous and adelic. From Theorem 3.2, we will conclude that the metric is also semipositive in the sense of Zhang [Zh1]. We will use Zhang's inequalities [Zh2] to deduce that the height of the underlying curve $h_{\overline{\mathcal{L}}_P}(B) := c_1(\overline{\mathcal{L}}_P)^2 / (2c_1(\mathcal{L}_P))$ may be computed as

$$h_{\overline{\mathcal{L}}_P}(B) = 0.$$

Consequently, we will be able to apply the equidistribution results of Chambert-Loir, Thuillier, and Yuan [CL1, Th, Y] to complete our proofs.

4.1. The metric and its properties. Let $m \in \mathbb{N}$ be such that

$$D = m \cdot D_E(P)$$

is an integral divisor. Let $\mathcal{L}_m$ be the associated line bundle on B . Note that $\deg(\mathcal{L}_m) = m \hat{h}_E(P) > 0$ so $\mathcal{L}_m$ is ample; by replacing m with a multiple, we may assume that $\mathcal{L}_m$ is very ample.

Fix a place v of K . Let U be an open subset of B_v^{an} . Each section $s \in \mathcal{L}_m(U)$ is identified with a meromorphic function f on U satisfying

$$(f) \geq -D.$$

We set

$$\|s(t)\|_v = \begin{cases} e^{-m \hat{\lambda}_{E_t, v}(P_t)} |f(t)|_v & \text{if } f(t) \text{ is finite and nonzero} \\ 0 & \text{if } \text{ord}_t f > -m \hat{\lambda}_{E, \text{ord}_t}(P) \\ e^{-m V_{P, t, v}(t)} & \text{otherwise,} \end{cases}$$

taking a locally-defined uniformizer $u = f^{1/\text{ord}_t f}$ at t in the definition of $V_{P, t, v}$ from (2.2).

Theorem 4.1. *The metric $\|\cdot\| = \{\|\cdot\|_v\}_{v \in M_K}$ on $\mathcal{L}_m$ is continuous, semipositive, and adelic.*

Proof. The continuity and semipositivity follows from Theorem 3.2. (In [CL2], semipositivity of a continuously metrized line bundle on a curve is defined terms of subharmonicity of potentials for the curvature form at each archimedean place, and as a uniform limit of “smooth semipositive” metrics at each non-archimedean place. In [Th], it is established that subharmonicity of potentials is a sufficient notion at all places, and he proves in [Th, Theorem 4.3.3] that this notion of semipositivity coincides with that of Zhang [Zh1]. See also [FG, Lemma 3.11, Theorem 3.12] where this same argument is applied in a dynamical context.) The adelic condition follows from Theorem 2.2. $\square$

4.2. The associated height function. A height function on $B(\overline{K})$ is defined by setting

$$(4.1) \quad h_P(t) := \frac{1}{m} \frac{1}{|\text{Gal}(\overline{K}/K) \cdot t|} \sum_{s \in \text{Gal}(\overline{K}/K) \cdot t} \sum_{v \in M_K} -n_v \log \|\phi(s)\|_v$$

where ϕ is any global section of $\mathcal{L}_m$ which is nonvanishing along the Galois orbit of t , and $\|\cdot\|_v$ is the metric of §4.1. Recall that $\text{supp } D_E(P) \subset B(K)$; we may assume that our sections ϕ are defined over K , and the product formula guarantees our height is independent of the choice of ϕ .

Our next goal is to prove the following two important facts about this height function h_P .

Proposition 4.2. *The height function h_P satisfies*

$$h_P(t) = \hat{h}_{E_t}(P_t)$$

for all $t \in B(\overline{K})$ such that the fiber E_t is smooth.

Proposition 4.3. *The height function h_P satisfies*

$$h_P(t) \geq 0$$

for all $t \in B(\overline{K})$.

Proof of Proposition 4.2. First fix $t \in B(\overline{K}) \setminus \text{supp } D_E(P)$ with smooth fiber E_t . Choose a section ϕ defined over K that does not vanish along the Galois orbit of t , and let f be the associated meromorphic function on B . Then f takes finite and nonzero values along the Galois orbit of t . We have,

$$\begin{aligned} h_P(t) &= \frac{1}{m} \frac{1}{|\text{Gal}(\overline{K}/K) \cdot t|} \sum_{s \in \text{Gal}(\overline{K}/K) \cdot t} \sum_{v \in M_K} n_v (m \hat{\lambda}_{E_s, v}(P_s) - \log |f(s)|_v) \\ &= \frac{1}{m} \frac{1}{|\text{Gal}(\overline{K}/K) \cdot t|} \sum_{s \in \text{Gal}(\overline{K}/K) \cdot t} \sum_{v \in M_K} m n_v \hat{\lambda}_{E_s, v}(P_s) \\ &= \hat{h}_{E_t}(P_t). \end{aligned}$$

where the second equality follows from the product formula.

For $t_0 \in \text{supp } D_E(P)$ such that E_{t_0} is smooth, it is necessarily the case that $P_{t_0} = O_{t_0}$ (as observed in §2.2), and therefore $\hat{h}_{E_{t_0}}(P_{t_0}) = 0$. To compute $h_P(t_0)$, observe that $t_0 \in B(K)$ so its Galois orbit is trivial; fixing a uniformizer $u \in K(B)$ at t_0 , we have

$$h_P(t_0) = \sum_{v \in M_K} n_v V_{P, t_0, v}(t_0)$$

where $V_{P, t_0, v}$ is the function of (2.2) associated to the uniformizer u .

We can compute $h_P(t_0)$ using the dynamical interpretation of the local heights, described in Section 3.1. Fix a Weierstrass equation for E in a neighborhood of t_0 and write $P = (x_P, y_P)$. The assumption that $P_{t_0} = O_{t_0}$ is equivalent to $\text{ord}_{t_0} x_P < 0$. After possibly shrinking U , write $x_P = (u)^{\text{ord}_{t_0}(x_P)} A_0$ for the chosen uniformizer u and a function $A_0 \in K(B)$ that does not vanish in U . We choose a lift X of x_P on U defined as $X = (A_0, B_0)$, where $B_0 := (u)^{-\text{ord}_{t_0}(x_P)}$. Notice that A_0 and B_0 are regular at t_0 . Let F be the standard lift in these coordinates, defined in (3.2); it satisfies $F_{t_0}(1, 0) = (1, 0)$, and we have $\mathcal{G}_{F, \text{ord}_{t_0}}(A_0, B_0) = 0$. Since $\text{ord}_{t_0} \Delta_E = 0$, Proposition 3.1 implies that

$$V_{P, t_0, v}(t) = \frac{1}{2} \mathcal{G}_{F, v}(A_0(t), B_0(t)) - \frac{1}{12} \log |\Delta_E(t)|_v$$

for all $t \in U$. Therefore,

$$\begin{aligned}
 V_{P,t_0,v}(t_0) &= \frac{1}{2} \mathcal{G}_{F_{t_0},v}(A_0(t_0), 0) - \frac{1}{12} \log |\Delta_E(t_0)|_v \\
 &= \frac{1}{2} \lim_{n \rightarrow \infty} \frac{1}{4^n} \log \|F_{t_0}^n(A_0(t_0), 0)\|_v - \frac{1}{12} \log |\Delta_E(t_0)|_v \\
 &= \frac{1}{2} \lim_{n \rightarrow \infty} \frac{1}{4^n} \log |A_0(t_0)^{4^n}|_v - \frac{1}{12} \log |\Delta_E(t_0)|_v \\
 &= \frac{1}{2} \log |A_0(t_0)|_v - \frac{1}{12} \log |\Delta_E(t_0)|_v.
 \end{aligned}$$

The product formula now yields that $h_P(t_0) = 0$, as claimed. $\square$

To prove Proposition 4.3, we first reduce to the case that the elliptic surface $E \rightarrow B$ has semi-stable reduction; that is, all of its fibers are either smooth or have multiplicative reduction. The next lemma describes how the height associated with the divisor $D_E(P)$ behaves under base extensions of the elliptic surface $E \rightarrow B$. It is adapted from [Si3, Reduction Lemma II.2.1]. We include it here for completeness.

Lemma 4.4. *Let $\mu : B' \rightarrow B$ be a finite map of smooth projective curves, let $E' \rightarrow B'$ be a minimal model for $E \times_B B'$, and let $P' : B' \rightarrow E'$ be the extension of the section P . For each $t_0 \in B(\overline{K})$ and $t'_0 \in \mu^{-1}(\{t_0\}) \subset B'(\mathbb{C}_v)$, there is a neighborhood U of t'_0 in $B'(\mathbb{C}_v)$ and a regular non-vanishing function f on U such that*

$$V_{P,t_0,v}(\mu(t')) - V_{P',t'_0,v}(t') = \log |f(t')|_v$$

on $U \setminus \{t'_0\}$. In particular,

$$V_{P,t_0,v}(t_0) - V_{P',t'_0,v}(t'_0) = \log |f(t'_0)|_v.$$

Proof. Let u be a uniformizer at t_0 , u' a uniformizer at t'_0 and $n = \text{ord}_{t'_0}(\mu^*u)$. Since local heights are invariant under base extension we have

$$(4.2) \quad \hat{\lambda}_{E', \text{ord}_{t'_0}}(P') = n \hat{\lambda}_{E, \text{ord}_{t_0}}(P).$$

Notice that for all t' in a punctured neighborhood of t'_0 the fibers $E'_{t'}$ are smooth. Hence the map $E' \rightarrow E$ gives an isomorphism between the fibers $E'_{t'} \rightarrow E_{\mu(t')}$. Under this isomorphism $P'_{t'} \in E'_{t'}$ is mapped to $P_{\mu(t')} \in E_{\mu(t')}$. Invoking now the uniqueness of the Néron local heights, we have

$$(4.3) \quad \hat{\lambda}_{E_{\mu(t')}, v}(P_{\mu(t')}) = \hat{\lambda}_{E'_{t'}, v}(P'_{t'}).$$

Combining (4.2) and (4.3) we get that for t' in a punctured neighborhood of t'_0 ,

$$V_{P,t_0,v}(\mu(t')) = V_{P',t'_0,v}(t') + \hat{\lambda}_{E,\text{ord}_{t_0}}(P) \log \left| \frac{u(\mu(t'))}{u'^n(t')} \right|_v.$$

The definition of n yields that the function $f(t') = \left(\frac{u(\mu(t'))}{u'^n(t')} \right)^{\hat{\lambda}_{E,\text{ord}_{t_0}}(P)}$ is regular and non-vanishing at t'_0 . The first part of the lemma follows.

Finally, Theorem 2.3 allows us to conclude that

$$V_{P,t_0,v}(\mu(t'_0)) - V_{P',t'_0,v}(t'_0) = \log |f(t'_0)|_v$$

at the point t'_0 , as claimed. $\square$

The following lemma will allow us to prove Proposition 4.3 in the case that a fiber has multiplicative reduction. The proof is lengthy, but it is merely a collection of computations using the explicit formulas for the local height functions, as in [Si2, Theorem VI.3.4, VI.4.2].

Lemma 4.5. *Let $E \rightarrow B$ be an elliptic surface and let $P : B \rightarrow E$ be a non-zero section defined over K . Then there exists a finite extension L of the number field K so that, for each $t_0 \in B(\overline{K})$ such that E_{t_0} has multiplicative reduction, there exists an $x(t_0) \in L^*$ so that*

$$V_{P,t_0,v}(t_0) = \log |x(t_0)|_v$$

at all places v of L .

Proof. We let

$$(4.4) \quad E : y^2 = x^3 + ax + b,$$

be a minimal Weierstrass equation for E over an affine subset $W \subset B$ defined over K with $t_0 \in W$. Here $a, b \in K(B)$ are regular functions at t_0 . Using this Weierstrass equation we write

$$P = (x_P, y_P),$$

where $x_P, y_P \in K(B)$. Since $E \rightarrow B$ has multiplicative reduction over $t_0 \in B(K)$, we have

$$(4.5) \quad N := \text{ord}_{t_0} \Delta_E \geq 1 \text{ and } \text{ord}_{t_0} a = 0;$$

see [Si5, Proposition VII.5.1]. Let v be a place of K (archimedean or non-archimedean). We denote by j_E the j -invariant of $E \rightarrow W$, given by

$$j_E(t) = 1728 \frac{(4a(t))^3}{\Delta_E(t)}.$$

Notice that equation (4.5) yields that j_E has a pole at t_0 . Hence, we can find a v -adic open neighborhood U of t_0 and an analytic map

$$\psi : U \rightarrow \{q \in \mathbb{C}_v : |q|_v < 1\},$$

such that the following holds: If j is the modular j -invariant [Si2, Chapter V], then we have

$$j_E(t) = j(\psi(t)) \text{ and } \text{ord}_{t_0} \psi = N.$$

The function $\psi(t)$ is given as

$$(4.6) \quad \psi(t) = \frac{1}{j_E(t)} + \frac{744}{j_E^2(t)} + \frac{750420}{j_E^3(t)} + \dots \in \mathbb{Z}[(j_E(t))^{-1}].$$

In the following, we choose a uniformizer $u \in K(B)$ at t_0 , and we identify ψ with its expression $\psi(t) \in \mathbb{C}_v[[u]]$ and write

$$(4.7) \quad \psi(t) = \beta u(t)^N + u(t)^{N+1} f(t), \text{ for } t \in U \setminus \{t_0\}.$$

Equation (4.6) yields that $\beta \in K \setminus \{0\}$ and $f(t) \in K[[u]]$. Following the proof of [Si3, Section 6] and after possibly shrinking U we have isomorphisms

$$(4.8) \quad E_t(\mathbb{C}_v) \xrightarrow{\sim} \mathbb{C}_v^* / \psi(t)^{\mathbb{Z}} \xrightarrow{\sim} C_{\psi(t)},$$

where $C_{\psi(t)}$ is the elliptic curve given by $y^2 = 4x^3 - g_2(\psi(t))x - g_3(\psi(t))$ for $t \in U \setminus \{t_0\}$. Under these isomorphisms, we have

$$P_t \mapsto w(t) \mapsto (\wp(w(t), \psi(t)), \wp'(w(t), \psi(t))).$$

Here g_2, g_3 are the modular invariants, given by their usual q -series

$$g_2(q) = \frac{1}{12} \left(1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n} \right), \quad g_3(q) = \frac{1}{216} \left(-1 + 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n} \right)$$

and $\wp$ is the Weierstrass $\wp$ -function given by

$$(4.9) \quad \wp(w, q) = \frac{1}{12} + \sum_{n \in \mathbb{Z}} \frac{q^n w}{(1 - q^n w)^2} - 2 \sum_{n=1}^{\infty} \frac{n q^n}{1 - q^n}, \quad \wp'(w, q) = \sum_{n \in \mathbb{Z}} \frac{q^n w (1 + q^n w)}{(1 - q^n w)^3}.$$

In view of [Si3, Lemma II.6.2], after possibly replacing P by $-P$, we may assume that $w : U \rightarrow \mathbb{C}_v$ is an analytic map satisfying

$$(4.10) \quad 0 \leq m := \text{ord}_{t_0} w \leq \frac{1}{2} \text{ord}_{t_0} \psi.$$

In the following we identify w with its series in $\mathbb{C}_v[[u]]$ and write

$$(4.11) \quad w(t) = \alpha u^m(t) + u^{m+1}(t)g(t),$$

where $\alpha \in \mathbb{C}_v$ and $g(t) \in \mathbb{C}_v[[u]]$.

We claim that $w(t) \in \overline{K}[[u]]$. To see this, notice that from [Si5, Chapter III.1] we have that for $t \in U$

$$(\wp(w(t), \psi(t)), \wp'(w(t), \psi(t))) = (\nu^{-2}(t)x_P(t), 2\nu^{-3}(t)y_P(t)),$$

where

$$\nu(t)^{12} = \frac{\Delta_E(t)}{\Delta(\psi(t))}.$$

In the equation above Δ denotes the modular discriminant given by

$$\Delta(q) = g_2(q)^3 - 27g_3(q)^3.$$

Since the functions ψ, Δ_E and Δ are defined over K , we have that $Y(t) := 2\nu^{-3}(t)y_P(t)$ is also defined over K . Since $Y(t) = \wp'(w(t), \psi(t)) \in K[[u]]$ and $\psi(t) \in K[[u]]$ we get that $w(t) \in K[[u]]$.

Therefore, there are non-zero constants $\alpha, \beta, \gamma \in \overline{K} \setminus \{0\}$, non-negative integers $k, m \in \mathbb{N}$ and functions $f(t), g(t), h(t) \in \overline{K}[[u]]$ such that for all $t \in U$

$$(4.12) \quad \psi(t) = \beta u^N(t) + f(t)u^{N+1}(t), \quad w(t) = \alpha u^m(t) + g(t)u^{m+1}(t), \quad 1 - w(t) = \gamma u^k(t) + h(t)u^{k+1}(t).$$

Next, we aim to express $x(t_0)$ (as in the statement of the lemma) in terms of $\alpha, \beta, \gamma \in \overline{K}$.

Using the isomorphisms in 4.8, the uniqueness of the local canonical heights and the explicit formulas for the local canonical heights [Si2, Theorem VI.3.4, VI.4.2], we get

$$(4.13) \quad \hat{\lambda}_{E_t, v}(P_t) = \hat{\lambda}(w(t), \psi(t)) = -\frac{1}{2}B_2 \left(\frac{\log |w(t)|_v}{\log |\psi(t)|_v} \right) \log |\psi(t)|_v - \log |1 - w(t)|_v$$

$$(4.14) \quad - \sum_{n \geq 1} \log |(1 - \psi(t)^n w(t))(1 - \psi(t)^n w(t)^{-1})|_v,$$

where $B_2(s) = s^2 - s + 1/6$ is the second Bernoulli polynomial.

Since $\text{ord}_{t_0} \psi = N \geq 1$ and using (4.10), we get

$$(4.15) \quad \lim_{t \xrightarrow{v} t_0} \sum_{n \geq 1} \log |(1 - \psi(t)^n w(t))(1 - \psi(t)^n w(t)^{-1})|_v = 0.$$

In what follows, for $F(t) \in \mathbb{C}_v[[u]]$ we write

$$F(t) := o_v(1), \text{ if } \lim_{t \rightarrow t_0} F(t) = 0.$$

In view of [Si1, Lemma I.5.1], we have

$$\begin{aligned} (4.16) \quad B_2 \left(\frac{\log |w(t)|_v}{\log |\psi(t)|_v} \right) \log |\psi(t)|_v &= \frac{\log^2 |w(t)|_v}{\log |\psi(t)|_v} - \log |w(t)|_v + \frac{1}{6} \log |\psi(t)|_v \\ &= \frac{m^2}{N} \log |u(t)|_v + \frac{m}{N^2} \log \left(\frac{|\alpha|_v^{2N}}{|\beta|_v^m} \right) - \log |\alpha|_v \\ &\quad - m \log |u(t)|_v + \frac{\log |\beta|_v}{6} + \frac{N}{6} \log |u(t)|_v + o_v(1). \end{aligned}$$

Using equations (4.15) and (4.16), equation (4.13) yields

$$\begin{aligned} (4.17) \quad \hat{\lambda}_{E,v}(P_t) + \frac{1}{2} \left(\frac{m^2}{N} - m + \frac{N}{6} + 2k \right) \log |u(t)|_v &= -\frac{1}{2} \left(\frac{m}{N^2} \log \left(\frac{|\alpha|_v^{2N}}{|\beta|_v^m} \right) - \log |\alpha|_v + \frac{\log |\beta|_v}{6} \right) \\ (4.18) \quad &\quad - \log |\gamma|_v + o_v(1). \end{aligned}$$

Finally, notice that [Si2, Theorem VI.4.2] implies

$$\hat{\lambda}_{E, \text{ord}_{t_0}}(P) = \text{ord}_{t_0}(1 - w) + \frac{1}{2} B_2 \left(\frac{\text{ord}_{t_0} w}{\text{ord}_{t_0} \psi} \right) \text{ord}_{t_0} \psi = \frac{1}{2} \left(\frac{m^2}{N} - m + \frac{N}{6} + 2k \right).$$

Therefore

$$\begin{aligned} V_{P, t_0, v}(t_0) &= \lim_{t \rightarrow t_0} V_{P, t_0, v}(t) = -\frac{1}{2} \left(\frac{m}{N^2} \log \left(\frac{|\alpha|_v^{2N}}{|\beta|_v^m} \right) - \log |\alpha|_v + \frac{\log |\beta|_v}{6} \right) - \log |\gamma|_v \\ &= \log |x(t_0)|_v, \end{aligned}$$

where $x(t_0) = \frac{\beta^{m^2/2N^2-1/2}}{\gamma \alpha^{m/N-1/2}}$ belongs to a finite extension of K , denoted by L .

□

Proof of Proposition 4.3. By [Si3, Lemma II.2.2] there is a finite map of smooth projective curves $B' \rightarrow B$ such that if $E' \rightarrow B'$ is a minimal model for $E \times_B B'$, then E' has semi-stable reduction over the singular fibers of $E \rightarrow B$. Moreover, we may choose B' so that everything is defined over K . Thus, by Lemma 4.4 and using the product formula, we may assume that the singular fibers of our elliptic surface $E \rightarrow B$ have multiplicative reduction.

For all $t \in B(\overline{K})$ for which E_t is smooth, we know from Proposition 4.2 that $h_P(t) = \hat{h}_{E_t}(P_t)$. The canonical height is always non-negative, so we may conclude that $h_P(t) \geq 0$ for all such t .

Assume now that $t_0 \in B(\overline{K})$ has a fiber with multiplicative reduction. Enlarging the number field K if necessary we may assume that $t_0 \in B(K)$ and that its corresponding $x(t_0)$ defined in the statement of Lemma 4.5 is in K^* . Then, on using the product formula, Lemma 4.5 implies that $h_P(t_0) = 0$. This completes the proof. $\square$

4.3. Proofs of Theorem 1.1 and Corollary 1.2.

Proof of Theorem 1.1. Let $\mathcal{L}_P$ be the line bundle on B induced from the divisor $D_E(P)$. From Theorem 4.1, we know that its m -th tensor power can be equipped with a continuous, adelic, semipositive metric, so that the corresponding height function is (a multiple of) the canonical height $\hat{h}_{E_t}(P_t)$ on the smooth fibers. Thus, by pulling back the metric to $\mathcal{L}_P$, we obtain a continuous, semipositive, adelic metric on $\mathcal{L}_P$ inducing a height function $h_{\overline{\mathcal{L}}_P}$ which is precisely the function h_P defined in (4.1).

It remains to show that this height h_P satisfies $h_P(B) = 0$. This is a consequence of Propositions 4.2 and 4.3 and Zhang's inequalities on successive minima [Zh2]. Indeed, from Proposition 4.3, we know that $h_P(t) \geq 0$ for all $t \in B(\overline{K})$. In addition, since $\hat{h}_E(P) \neq 0$, we know that there are infinitely many $t \in B(\overline{K})$ for which

$$\hat{h}_{E_t}(P_t) = 0.$$

(For a complex-dynamical proof, see [De1, Proposition 1.5, Proposition 2.3].) Therefore, from Proposition 4.2, we may deduce that the essential minimum of h_P on B is equal to 0. Finally, from [Zh2, Theorem 1.10], we may conclude that $h_P(B) = 0$. $\square$

Proof of Corollary 1.2. When combined with the equidistribution theorems of Yuan and Thullier [Y, Th], we immediately obtain the corollary from Theorem 1.1. The measures $\mu_{P,v}$ are the curvature distributions associated to the metrics $\|\cdot\|_v$ at each place v . From the definition of the metric in §4.1, we see that they are given locally by

$$\mu_{P,v} = dd^c V_{P,t_0,v}(t)$$

in a v -adic neighborhood of any point $t_0 \in B(\overline{K})$, and for any choice of uniformizer u at t_0 . $\square$

5. PROOF OF THEOREM 1.4.

5.1. Reduction to the case of a fiber product of elliptic surfaces. We first show that, to prove the theorem, it suffices to prove the result for sections of the fiber product $A = E_1 \times_B \cdots \times_B E_m$ of $m \geq 2$ elliptic surfaces $E_i \rightarrow B$ over the same base, and to assume that the line bundle $\mathcal{L}$ is generated by the divisor

$$\{O_{E_1}\} \times E_2 \times \cdots \times E_m + E_1 \times \{O_{E_2}\} \times \cdots \times E_m + \cdots + E_1 \times E_2 \times \cdots \times \{O_{E_m}\}.$$

Let B be a quasiprojective smooth algebraic curve defined over $\overline{\mathbb{Q}}$. Suppose $A \rightarrow B$ is family of abelian varieties defined over $\overline{\mathbb{Q}}$ that is isogenous to a fibered product of $m \geq 2$ elliptic curves. That is, there is a branched cover $B' \rightarrow B$ and $m \geq 2$ elliptic surfaces $E_i \rightarrow B'$ that give rise to an isogeny

$$E_1 \times_{B'} \cdots \times_{B'} E_m \rightarrow A$$

over B' . Now let $\mathcal{L}$ be a line bundle on A which restricts to an ample and symmetric line bundle on each fiber A_t for $t \in B$. Then the line bundle $\mathcal{L}$ pulls back to a line bundle $\mathcal{L}'$ on $E_1 \times_{B'} \cdots \times_{B'} E_m$, and it again restricts to an ample and symmetric line bundle on each fiber over $t \in B'$.

Now suppose that we have a section $P : B \rightarrow A$. The section P pulls back to a section $P' : B' \rightarrow A$, and this in turn pulls back to a (possibly multi-valued) section of $E_1 \times_{B'} \cdots \times_{B'} E_m$. If multi-valued, we can perform a base change again, passing to a branched cover $B'' \rightarrow B'$, so that the induced section $P'' : B'' \rightarrow E_1 \times_{B''} \cdots \times_{B''} E_m$ is well defined. By definition, the assumption that P is non-special on A means that it is non-special as a section of $E_1 \times_{B''} \cdots \times_{B''} E_m$.

Finally, we observe that the conclusion of Theorem 1.4 does not depend on the choice of line bundle. (We thank Joe Silverman for his help with this argument.) Recall that, on any abelian variety A defined over $\overline{\mathbb{Q}}$, the notion of a “small sequence” of points is independent of the choice of ample and symmetric line bundle. That is, if we take two ample and symmetric divisors D_1 and D_2 , then we know that there exists an integer $m_1 > 0$ so that $m_1 D_1 - D_2$ is ample; similarly there exists $m_2 > 0$ so that $m_2 D_2 - D_1$ is ample. It follows from properties of the Weil height machine that the heights h_{D_1} and h_{D_2} will then satisfy

$$\frac{1}{m_1} h_{D_2} + C_1 \leq h_{D_1} \leq m_2 h_{D_2} + C_2$$

for real constants C_1, C_2 . Upon passing to the canonical height, we conclude that

$$(5.1) \quad \frac{1}{m_1} \hat{h}_{D_2} \leq \hat{h}_{D_1} \leq m_2 \hat{h}_{D_2}$$

on the abelian variety. In particular, $\hat{h}_{D_1}(a_i) \rightarrow 0$ for some sequence in $A(\overline{\mathbb{Q}})$ if and only if $\hat{h}_{D_2}(a_i) \rightarrow 0$. Now suppose we have a family of abelian varieties $A \rightarrow B$. Two line bundles $\mathcal{L}_1$ and $\mathcal{L}_2$ associated to relatively ample and symmetric divisors induce canonical heights $\hat{h}_{\mathcal{L}_1, t}$ and $\hat{h}_{\mathcal{L}_2, t}$ on each fiber A_t . But recalling that ampleness persists on Zariski open sets [La3, Theorem 1.2.17], there exist positive integers m_1 and m_2 so that the line bundles $\mathcal{L}_1^{m_1} \otimes \mathcal{L}_2^{-1}$ and $\mathcal{L}_2^{m_2} \otimes \mathcal{L}_1^{-1}$ are relatively ample on a Zariski open subset of the base B . Passing to the canonical heights once again, we find that the relation (5.1) holds uniformly over B (after possibly excluding finitely many points). Therefore, for any section $P : B \rightarrow A$, there exists a positive constant $c(\mathcal{L}_1, P)$ of Theorem 1.4 for the height $\hat{h}_{\mathcal{L}_1}$ if and only if there exists such a constant $c(\mathcal{L}_2, P)$ for $\hat{h}_{\mathcal{L}_2}$.

5.2. Proof for a fiber product of elliptic curves. Fix integer $m \geq 2$, and let $E_i \rightarrow B$ for $i = 1, \dots, m$ be elliptic surfaces over the same base curve B , defined over $\overline{\mathbb{Q}}$. Let $A = E_1 \times_B \cdots \times_B E_m$, and let $\mathcal{L}$ be the line bundle on $E_1 \times_B \cdots \times_B E_m$ associated to the divisor

$$D = \{O_{E_1}\} \times E_2 \times \cdots \times E_m + E_1 \times \{O_{E_2}\} \times \cdots \times E_m + \cdots + E_1 \times E_2 \times \cdots \times \{O_{E_m}\}.$$

For all but finitely many $t \in B(\overline{\mathbb{Q}})$, the canonical height $\hat{h}_{\mathcal{L}_t}$ on the fiber A_t is easily seen to be the sum of canonical heights (see, e.g., [HS] for properties of the height functions), so that

$$\hat{h}_{\mathcal{L}_t} = \sum_{i=1}^m \hat{h}_{E_{i,t}}.$$

Now assume that $P = (P_1, \dots, P_m)$ is a section of $A \rightarrow B$. Define

$$\hat{h}_i(t) := \hat{h}_{E_{i,t}}(P_{i,t})$$

for $i = 1, \dots, m$ and for all $t \in B(\overline{\mathbb{Q}})$ where all $E_{i,t}$ are smooth elliptic curves. Suppose there exists an infinite sequence $\{t_n\} \subset B(\overline{\mathbb{Q}})$ for which

$$(5.2) \quad \hat{h}_i(t_n) \rightarrow 0 \text{ for all } i = 1, \dots, m$$

as $n \rightarrow \infty$. We will prove that for every pair (i, j) , there exists an infinite sequence $\{s_n\} \subset B(\overline{\mathbb{Q}})$ so that

$$\hat{h}_i(s_n) = \hat{h}_j(s_n) = 0$$

for all n . In this way, we reduce our problem to the main results of [MZ2, MZ3] which imply that the pair (P_i, P_j) must be a special section of $E_i \times_B E_j$. Finally, we observe that our definition of a special section $P = (P_1, P_2, \dots, P_m)$ is equivalent to the statement that every pair (P_i, P_j) is special. Therefore, for any non-special section P , we can conclude that there exists a constant $c = c(P) > 0$ so that the set

$$\{t \in B(\overline{\mathbb{Q}}) : \hat{h}_{\mathcal{L}_t}(P_t) < c\}$$

is finite.

Fix a pair (i, j) . First assume that neither E_i nor E_j is isotrivial. If P_i or P_j is torsion, then the section (P_i, P_j) is special. Otherwise, we have $\hat{h}_{E_i}(P_i) \neq 0$ and $\hat{h}_{E_j}(P_j) \neq 0$, and we may apply Theorem 1.1 to deduce that the height functions $\hat{h}_i$ and $\hat{h}_j$ are “good” on B . More precisely, we let $\overline{M}_i$ and $\overline{M}_j$ be the adelicly metrized line bundles on the base curve B associated to the height functions $\hat{h}_i$ and $\hat{h}_j$, from Theorem 1.1. They are both equipped with continuous adelic metrics of non-negative curvature. By assumption, we have

$$(5.3) \quad \hat{h}_i(t_n) \rightarrow 0 \quad \text{and} \quad \hat{h}_j(t_n) \rightarrow 0$$

as $n \rightarrow \infty$. Therefore, we may apply the observation of Chambert-Loir [CL2, Proposition 3.4.2], which builds upon on Zhang’s inequalities [Zh2], to conclude that there exist integers n_i and n_j so that $\overline{M}_i^{n_i}$ and $\overline{M}_j^{n_j}$ are isomorphic as line bundles on B and their metrics are scalar multiples of one another. It follows that the height functions $\hat{h}_i$ and $\hat{h}_j$ are the same, up to scale, and in particular they have the same zero sets. In other words, $P_{i,t}$ is a torsion point on $E_{i,t}$ if and only if $P_{j,t}$ is a torsion point on $E_{j,t}$ (for all but finitely many t in B), and there are infinitely many such parameters $t \in B(\overline{\mathbb{Q}})$.

Now suppose that E_i is isotrivial. The existence of the small sequence t_n in (5.3) implies that either $\hat{h}_{E_i}(P_i) \neq 0$ or P_i is torsion on E_i , and furthermore, if P_i is torsion, then it follows that (P_i, P_j) is a special section of $E_i \times_B E_j$. Similarly if E_j is isotrivial. In other words, the existence of the sequence t_n in (5.3) allows us to conclude that either (P_i, P_j) is a special pair, or we have that both $\hat{h}_{E_i}(P_i) \neq 0$ and $\hat{h}_{E_j}(P_j) \neq 0$. Therefore, we may proceed as above in the nonisotrivial case, applying Theorem 1.1 to deduce that the heights $\hat{h}_i$ and $\hat{h}_j$

coincide, up to scale, and in particular there are infinitely many parameters $s \in B(\overline{\mathbb{Q}})$ where

$$\hat{h}_i(s) = \hat{h}_j(s) = 0.$$

This concludes the proof of Theorem 1.4.

6. VARIATION OF CANONICAL HEIGHT, ILLUSTRATED

In this final section, we provide a few illustrations of the distributions $\mu_{P,v}$ arising in Corollary 1.2, for an archimedean place v . In Proposition 6.2, we present a complex-dynamical proof that the archimedean measures $\mu_{P,v}$ will have support equal to all of B .

6.1. Images. Given $E \rightarrow B$ and section P , we plot the parameters t where P_t is a torsion point on the fiber E_t of specified order. As proved in Corollary 1.2, the local height function at each place

$$t \mapsto \hat{\lambda}_{E_t,v}(P_t)$$

determines the distribution of the torsion parameters; it is a potential for the measure $\mu_{P,v}$ (away from the singularities). Recall that if we have two sections P and Q that are linearly related on E , then the distributions of their torsion parameters in B will be the same.

Figure 6.1, top, illustrates the example of Silverman from [Si1, Theorem I.0.3]. Here, we have

$$E_t = \{y^2 + xy/t + y/t = x^3 + 2x^2/t\}$$

with $B = \mathbb{P}^1$ and $P_t = (0, 0)$ in (x, y) -coordinates. Plotted are the torsion parameters of orders 2^n for all $n \leq 8$; that is, the points t in the base B where P_t is torsion of order 2^n on the fiber E_t . Roughly, a smaller yellow dot corresponds to higher order of torsion. Figure 6.1, bottom, is another section of the same family, where the x -coordinate of P_t is constant and equal to $-1/4$. (Strictly speaking, this second P is not a section of our given $E \rightarrow \mathbb{P}^1$, because the y -coordinate will not lie in $\overline{K}(B) \simeq \overline{\mathbb{Q}}(t)$ but in an extension; however, the property of being torsion and the determination of its order is independent of which point in the fiber we choose.) Observe the distinctly different pattern of yellow dots in the first and second pictures, especially in the left half of the two pictures, illustrating the linear independence in $E(k)$ of the two sections.

Figure 6.2 illustrates the torsion parameters for two independent sections of the Legendre family,

$$E_t = \{y^2 = x(x-1)(x-t)\}$$




FIGURE 6.1. At top, Silverman's example from [Si1, Theorem I.0.3], with $E_t = \{y^2 + xy/t + y/t = x^3 + 2x^2/t\}$ and $P_t = (0, 0)$, shown in the region $\{-2 \leq \operatorname{Re} t \leq 1, -1 \leq \operatorname{Im} t \leq 1\}$. The singular fibers occur at $t = 0, -2/27, -1$, and one sees the effects of numerical error in a small neighborhood of these parameters. At bottom, torsion parameters for section P having x -coordinate $x(P_t) = -1/4$ for all t .

over $B = \mathbb{P}^1$, studied in [MZ1]. The chosen sections are P_2 , with constant x -coordinate equal to 2, and P_5 , with constant x -coordinate equal to 5. As in Figure 6.1, we plot the torsion parameters of orders 2^n for all $n \leq 8$; generally, a smaller yellow dot signifies higher order




FIGURE 6.2. Torsion parameters for sections of the Legendre family, studied in [MZ1]; here, $B = \mathbb{P}^1$ and $E_t = \{y^2 = x(x-1)(x-t)\}$. At left, the section P_2 with $x((P_2)_t) = 2$ for all t ; at right, the section P_5 with $x((P_5)_t) = 5$ for all t . Both are shown in the region $\{-3 \leq \operatorname{Re} t \leq 5, -4 \leq \operatorname{Im} t \leq 4\}$

of torsion. It was proved in [DWY] that the limiting distributions for sections with constant x -coordinate satisfy $\mu_{P_x, \infty} = \mu_{P_{x'}, \infty}$ (at an archimedean place) if and only if $x = x'$. It was proved in [St] and [Ma] that there are no $t \in \mathbb{P}^1(\overline{K})$ for which both $(P_2)_t$ and $(P_5)_t$ are torsion on E_t . Again, observe the difference in the geometry of the yellow dots for the two independent sections.

Figure 6.3 illustrates our equidistribution result, Corollary 1.2, for the example of the Legendre family with the section P_5 . Plotted are the torsion parameters of orders 2^n with (a) $n \leq 6$, (b) $n \leq 8$, and (c) $n \leq 10$. Observe how the yellow dots fill in the “grid structure” in the base curve B , exactly as do the torsion points for one elliptic curve.

Remark 6.1. As mentioned above, the smaller yellow dots in the illustrations correspond, roughly, to higher orders of torsion. These images are produced with a standard escape-rate algorithm. We use the dynamical system f_t on $\mathbb{P}^1$, induced from multiplication by 2 on the elliptic curve E_t from Section 3, line (3.1). The coordinates on $\mathbb{P}^1$ are chosen so that ∞ is the image of the 0 of E_t . We mark t yellow if $|f_t^n(\pi(P_t))| \geq 10000$ for some $n \leq 8$.

6.2. Density of torsion parameters. In all of these examples, the yellow dots will fill in the picture as the order of torsion grows, and the support of the measures μ_{P_v} is equal to




FIGURE 6.3. Illustrating equidistribution: Torsion parameters of increasing orders for a section of the Legendre family, with $B = \mathbb{P}^1$ and $E_t = \{y^2 = x(x-1)(x-t)\}$ and P_5 as in Figure 6.2. At top, torsion parameters of orders 2^n for $n \leq 6$; bottom left, of orders 2^n for $n \leq 8$, and bottom right, of orders 2^n for $n \leq 10$.

$B(\mathbb{C})$. In fact, this will always be the case, for any (nontrivial) section of a complex elliptic surface, as our final result, Proposition 6.2, shows.

Let $E \rightarrow B$ be an elliptic surface over a projective curve B , defined over $\mathbb{C}$, and let $P : B \rightarrow E$ be a section for which $\hat{h}_E(P) \neq 0$ (over the function field $k = \mathbb{C}(B)$). Let μ_P

be the measure on B defined as in Proposition 3.6, that is as the pullback of the current T that restricts to Haar measure on each smooth fiber. In other words, μ_P is locally defined as the Laplacian of the function $G_P(t)$ introduced in Proposition 3.4, which is well defined when working over $\mathbb{C}$.

Proposition 6.2. *Let $E \rightarrow B$ be an elliptic surface over a projective curve B , defined over $\mathbb{C}$, and let $P : B \rightarrow E$ be a section for which $\hat{h}_E(P) \neq 0$ (over the function field $k = \mathbb{C}(B)$). Then the set*

$$\{t \in B : P_t \text{ is torsion on } E_t\}$$

is dense in $B(\mathbb{C})$ and

$$\text{supp } \mu_P = B(\mathbb{C}).$$

We give a complex-dynamical proof, viewing Proposition 6.2 as a consequence of the main result of [De1]. (We do not use the equidistribution result, Corollary 1.2.) An analytic proof is also presented in [Za, Notes to Chapter 3].

Proof. Let $B^* \subset B$ be a finitely-punctured Riemann surface such that the fiber E_t is smooth for all $t \in B^*$. Let $\pi_t : E_t \rightarrow \mathbb{P}^1$ be the degree-two projection and $f_t : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be the rational map induced by multiplication-by-2 on E_t , as defined in the introduction to Section 3. It is well known that the holomorphic family $\{f_t : t \in B^*\}$ is structurally stable; see, e.g., [Mc, Chapter 4]. Thus, over any simply-connected subset U of B^* , there is a holomorphic motion of the periodic points of f_t which extends uniquely to a holomorphic motion of all of $\mathbb{P}^1$, conjugating the dynamics.

The key observation is that μ_P is precisely the “bifurcation measure” of the pair (f, P) on B^* . See [De2, §2.7] and [De1] for definitions. The support of μ_P is equal to the bifurcation locus of (f, P) ; in particular, the parameters $t \in B^*$ for which $\pi_t(P_t)$ is preperiodic for f_t are dense in $\text{supp } \mu_P$. Therefore, it suffices to show that $\text{supp } \mu_P = B$.

Suppose to the contrary that there is an open disk $U \subset B^*$ for which $\mu_P(U) = 0$. Then the pair (f, P) is stable on U , and therefore $\pi_t(P_t)$ cannot be a repelling periodic point for any $t \in U$. From the uniqueness of the holomorphic motion, it follows that $t \mapsto \pi_t(P_t)$ is part of the holomorphic motion on U . By analytic continuity, then, we deduce that $\pi_t(P_t)$ must follow the motion of a point over all of B^* . This implies that the pair (f, P) is stable throughout B^* and the measure μ_P is 0. But this is absurd by the assumption that $\hat{h}_E(P) \neq 0$; see [De1, Theorem 1.1]. $\square$

REFERENCES

- [BR] M. Baker and R. Rumely. *Potential theory and dynamics on the Berkovich projective line*, volume 159 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [BH] B. Branner and J. H. Hubbard. The iteration of cubic polynomials. I. The global topology of parameter space. *Acta Math.* **160**(1988), 143–206.
- [CS] G. S. Call and J. H. Silverman. Canonical heights on varieties with morphisms. *Compositio Math.* **89**(1993), 163–205.
- [CL1] A. Chambert-Loir. Mesures et équidistribution sur les espaces de Berkovich. *J. Reine Angew. Math.* **595**(2006), 215–235.
- [CL2] A. Chambert-Loir. Heights and measures on analytic spaces. A survey of recent results, and some remarks. In *Motivic integration and its interactions with model theory and non-Archimedean geometry. Volume II*, volume 384 of *London Math. Soc. Lecture Note Ser.*, pages 1–50. Cambridge Univ. Press, Cambridge, 2011.
- [DG] L. DeMarco and D. Ghioca. Rationality of dynamical canonical height. To appear, *Ergodic Theory Dyn. Sys.*
- [De1] L. DeMarco. Bifurcations, intersections, and heights. *Algebra Number Theory* **10**(2016), 1031–1056.
- [De2] L. DeMarco. Dynamical moduli spaces and elliptic curves (KAWA Lecture Notes). To appear, *Ann. Fac. Sci. Toulouse Math.*
- [DWY] L. DeMarco, X. Wang, and H. Ye. Torsion points and the Lattès family. *Amer. J. Math.* **138**(2016), 697–732.
- [FG] C. Favre and T. Gauthier. Classification of special curves in the space of cubic polynomials. *Int. Math. Res. Not. IMRN.* **2**(2018), 362–411.
- [FS] J. E. Fornæss and N. Sibony. Complex dynamics in higher dimensions. In *Complex Potential Theory (Montreal, PQ, 1993)*, pages 131–186. Kluwer Acad. Publ., Dordrecht, 1994.
- [HS] M. Hindry and J.H. Silverman. *Diophantine geometry*, volume 201 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. An introduction.
- [HP] J. Hubbard and P. Papadopol. Superattractive fixed points in $\mathbf{C}^n$. *Indiana Univ. Math. J.* **43**(1994), 321–365.
- [La1] S. Lang. *Fundamentals of Diophantine geometry*. Springer-Verlag, New York, 1983.
- [La2] S. Lang. *Introduction to Arakelov theory*. Springer-Verlag, New York, 1988.
- [La3] R. Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 2004.
- [Ly] M. Lyubich. Entropy properties of rational endomorphisms of the Riemann sphere. *Ergodic Theory Dynamical Systems* **3**(1983), 351–385.
- [MZ1] D. Masser and U. Zannier. Torsion anomalous points and families of elliptic curves. *Amer. J. Math.* **132**(2010), 1677–1691.
- [MZ2] D. Masser and U. Zannier. Torsion points on families of squares of elliptic curves. *Math. Ann.* **352**(2012), 453–484.

- [MZ3] D. Masser and U. Zannier. Torsion points on families of products of elliptic curves. *Adv. Math.* **259**(2014), 116–133.
- [MZ4] D. Masser and U. Zannier. Torsion points on families of simple abelian surfaces and Pell’s equation over polynomial rings. *J. Eur. Math. Soc. (JEMS)* **17**(2015), 2379–2416. With an appendix by E. V. Flynn.
- [Ma] N.M. Mavraki. Impossible intersections in a Weierstrass family of elliptic curves. *J. Number Theory* **169**(2016), 21–40.
- [Mc] C. McMullen. *Complex Dynamics and Renormalization*. Princeton University Press, Princeton, NJ, 1994.
- [Mi] J. Milnor. On Lattès maps. In *Dynamics on the Riemann sphere*, pages 9–43. Eur. Math. Soc., Zürich, 2006.
- [Mu] D. Mumford. *Abelian varieties*. Oxford University Press, London 1970.
- [Ra1] M. Raynaud. Courbes sur une variété abélienne et points de torsion. *Invent. Math.* **71**(1983), 207–233.
- [Ra2] M. Raynaud. Sous-variétés d’une variété abélienne et points de torsion. In *Arithmetic and geometry, Vol. I*, volume 35 of *Progr. Math.*, pages 327–352. Birkhäuser Boston, Boston, MA, 1983.
- [Si1] J.H. Silverman. Variation of the canonical height on elliptic surfaces. I. Three examples. *J. Reine Angew. Math.* **426**(1992), 151–178.
- [Si2] J.H. Silverman. *Advanced Topics in the Arithmetic of Elliptic Curves*, volume 151 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
- [Si3] J.H. Silverman. Variation of the canonical height on elliptic surfaces. II. Local analyticity properties. *J. Number Theory* **48**(1994), 291–329.
- [Si4] J.H. Silverman. Variation of the canonical height on elliptic surfaces. III. Global boundedness properties. *J. Number Theory* **48**(1994), 330–352.
- [Si5] J.H. Silverman. *The Arithmetic of Elliptic Curves*, volume 106 of *Graduate Texts in Mathematics*. Springer, Dordrecht, second edition, 2009.
- [St] M. Stoll. Simultaneous torsion in the Legendre family. To appear, *Exper. Math.* Available online doi = 10.1080/10586458.2016.1201443.
- [SUZ] L. Szpiro, E. Ullmo, and S. Zhang. Équirépartition des petits points. *Invent. Math.* **127**(1997), 337–347.
- [Ta] J. Tate. Variation of the canonical height of a point depending on a parameter. *Amer. J. Math.* **105**(1983), 287–294.
- [Th] A. Thuillier. *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. Applications à la théorie d’Arakelov*. Thèse, Université de Rennes 1, 2005.
- [U] E. Ullmo. Positivité et discrétion des points algébriques des courbes. *Ann. of Math. (2)* **147**(1998), 167–179.
- [Y] X. Yuan. Big line bundles over arithmetic varieties. *Invent. Math.* **173**(2008), 603–649.

- [Za] U. Zannier. Some problems of unlikely intersections in arithmetic and geometry, With appendixes by David Masser. *Annals of Mathematics Studies*. Volume 181, Princeton University Press, Princeton, NJ, 2012.
- [Zh1] S.-W. Zhang. Positive line bundles on arithmetic varieties. *J. Amer. Math. Soc.* **8**(1995), 187–221.
- [Zh2] S.-W. Zhang. Small points and adelic metrics. *J. Algebraic Geom.* **4**(1995), 281–300.
- [Zh3] S. W. Zhang. Equidistribution of small points on abelian varieties. *Ann. of Math. (2)* **147**(1998), 159–165.
- [Zh4] S.-W. Zhang. Small points and Arakelov theory. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, number Extra Vol. II, pages 217–225 (electronic), 1998.