

Review

Interpreting metabolic complexity via isotope-assisted metabolic flux analysis

Bilal Moiz,¹ Ganesh Sriram,² and Alisa Morss Clyne ^{1,*,@}

Isotope-assisted metabolic flux analysis (iMFA) is a powerful method to mathematically determine the metabolic fluxome from experimental isotope labeling data and a metabolic network model. While iMFA was originally developed for industrial biotechnological applications, it is increasingly used to analyze eukaryotic cell metabolism in physiological and pathological states. In this review, we explain how iMFA estimates the intracellular fluxome, including data and network model (inputs), the optimization-based data fitting (process), and the flux map (output). We then describe how iMFA enables analysis of metabolic complexities and discovery of metabolic pathways. Our goal is to expand the use of iMFA in metabolism research, which is essential to maximizing the impact of metabolic experiments and continuing to advance iMFA and biocomputational techniques.

The metabolic fluxome: a dynamic description of cellular phenotype

A detailed analysis of cellular metabolism is essential for a complete understanding of cellular function in health and disease, and metabolism is also an important parameter for biotechnology applications. Most cell metabolism studies focus on experimentally measured metabolite concentrations. However, metabolite concentrations can change for varied reasons (e.g., increased upstream vs. decreased downstream reaction rates) [1]. Alternatively, metabolite concentrations could remain unchanged despite changes in flux (e.g., if upstream and downstream fluxes change by exactly the same amount). Therefore, we gain limited information from metabolite concentrations alone.

Metabolic fluxes, defined as the metabolic reaction rate (moles/time), provide much more information than metabolite concentration alone. The fluxome indicates the traffic of carbon and other elements among metabolites and therefore provides information on both metabolite concentrations and reaction rates. Since the reaction rates depend on upstream and downstream enzyme expression and activation, the fluxome integrates the cellular metabolome with the transcriptome, proteome, and regulome to offer a dynamic, comprehensive representation of the cell metabolic state [2,3].

Metabolic fluxes include extracellular fluxes (transport reactions that cross the cell membrane) and intracellular fluxes. While extracellular fluxes can be directly measured by tracking metabolite concentrations in the culture medium, intracellular fluxes cannot be directly measured. They, therefore, must be inferred from metabolite labeling patterns of metabolites [4]. In simple systems, metabolic fluxes can be determined by simply measuring how an isotopically labeled nutrient is converted to intermediate metabolites and secreted products. However, manual analysis is usually not possible in more complex metabolic systems.

iMFA is a mathematical technique based on optimization, which determines the metabolic fluxome by fitting metabolic labeling data onto a metabolic reaction network model [5,6]. iMFA

Highlights

Isotope-assisted metabolic flux analysis (iMFA) is a mathematical technique that estimates intracellular metabolic fluxes for complex biological systems.

iMFA software uses experimental data (extracellular fluxes, isotope labeling patterns) and a curated network model as inputs and produces a quantitative metabolic flux map as the output.

iMFA determines a set of flux values that produce the best match between the simulated and experimental mass distribution vectors using an iterative optimization process.

iMFA is essential for the metabolic analysis of complex systems and enables the discovery of new metabolic pathways.

Advances in iMFA are needed to enable analysis of dynamic metabolic states, multicellular co-cultures, and genome-scale networks. These new tools will significantly advance our understanding of metabolic complexity.

¹Department of Bioengineering, University of Maryland, College Park, MD 20742, USA

²Department of Chemical and Biomolecular Engineering, University of Maryland, College Park, MD 20742, USA

*Correspondence: aclyne@umd.edu (A.M. Clyne).
*Twitter: @Clyne_Lab



was originally developed to engineer microbial strains (metabolic engineering); however, it has more recently been used to analyze a variety of eukaryotic systems. iMFA improves our ability to understand metabolic complexity, and by doing so, can advance metabolomic and **fluxomic** analysis beyond what is currently possible.

The goal of this review paper is to inspire more researchers to use iMFA to maximize analytical insight from their metabolomic data. We first describe how iMFA works, including the inputs, mathematical optimization, and output. We then describe how iMFA enables us to analyze various aspects of metabolic complexity, including iMFA advances that are currently being developed that will enable us to apply iMFA to even more diverse, complex systems.

iMFA components

Input: network model

iMFA requires a metabolic network model (Figure 1), which describes the metabolic activities of the biological system. Metabolic reactions are categorized as **source**, **sink**, or **internal reactions** (see Glossary). These reactions are then organized into compartments (e.g., cytosol, mitochondria) where they take place. Finally, atom transitions are explicitly defined for each biochemical reaction. For example, glucose-6-phosphate loses the first carbon atom when entering the pentose phosphate pathway (PPP). This detail must be included in the metabolic network model for the simulation to fit the experimental isotope labeling patterns. The final network model is a curated table of metabolites and reactions, with their corresponding stoichiometry, compartment, and atom transitions.

Network models are typically curated by the end user via publicly available databases such as Kyoto Encyclopedia of Genes and Genomes (KEGG) [7,8], Virtual Metabolic Human [9], and MetaCyc [10,11]. To balance computational efficiency with metabolic accuracy, metabolic network models should be tailored to each biological system. For example, gluconeogenic pathways should be included in hepatocyte network models [12], while the tricarboxylic acid (TCA) cycle can be pruned from red blood cell network models since these cells lack mitochondria [13,14]. Network construction is often iterative, since the initially curated network model may not include all necessary metabolites, reactions, or compartmentalization. The iterative process is described in detail in the following iMFA output section.

Challenges remain in developing comprehensive metabolic network models. While atom transition information can be attained from the literature as well as the aforementioned databases, the available atom mapping information may be limited for specific reactions. Standardizing atom mapping information across different databases, along with developing algorithms to map atom transitions in complex organic reactions, will improve the accuracy of metabolic network models.

Input: experimental data

The crucial experimental inputs to iMFA are the isotope labeling states (isotope labeling patterns or **isotopomer** abundances) that result when cells or tissues process isotopically labeled tracers. ^{13}C -labeled tracers such as $[\text{U-}^{13}\text{C}_6]\text{glucose}$ (uniformly labeled glucose), $[1,2\text{-}^{13}\text{C}_2]\text{glucose}$ (glucose with ^{13}C in carbon atoms C-1 and C-2), or $[\text{U-}^{13}\text{C}_5]\text{glutamine}$ are frequently used to trace carbons. Isotopes of hydrogen (^2H) and nitrogen (^{15}N) can also be used to trace the respective elements [15]. Tracer selection requires thoughtful consideration as each tracer provides distinct metabolic information. For example, $[\text{U-}^{13}\text{C}_6]\text{glucose}$ elucidates overall glycolytic rate and glucose contribution to the TCA cycle but provides little information

Glossary

Anaplerotic: reactions that replenish intermediates of a cyclic metabolic pathway that are diverted toward biosynthesis, particularly in the case of TCA cycle intermediates. An example is the reaction $\text{pyruvate} + \text{CO}_2 \rightarrow \text{oxaloacetate}$, which replenishes the oxaloacetate diverted from the TCA cycle toward the synthesis of the aspartic family of amino acids.

Exchange flux: consider a pair of reversible reactions $A \leftrightarrow B$, where one reaction, say $A \rightarrow B$, has flux v_1 and the other reaction $B \rightarrow A$ has flux v_2 with both v_1 and $v_2 \geq 0$. In this case, the exchange flux is the minimum of v_1 and v_2 ($\min[v_1, v_2]$).

Fluxomics: an -omics method that measures intracellular and extracellular metabolic reaction rates.

Instationary MFA (INST-MFA): INST-MFA enables analysis of a system at metabolic but not isotopic steady state. INST-MFA requires the measurement of isotopomer distributions over time and preferably intracellular metabolite concentrations. INST-MFA is computationally more time-consuming than steady-state iMFA.

Internal reactions: reactions that both consume and produce metabolites and link source and sink reactions.

Isotopic steady state: a state in which fractional isotope enrichment is constant over time. Isotopic steady state can be difficult to achieve in mammalian cell cultures because high metabolite exchange leads to slow labeling, and changes in the cell state may cause metabolic variations. It is, therefore, difficult to maintain metabolic steady state long enough to reach isotopic steady state.

Isotopologues: metabolites that have different numbers of labeled atoms and therefore mass but are otherwise identical. Isotopologues can be differentiated by LC-MS or GC-MS which detect mass-to-charge ratios (m/z).

Isotopomers: isotopologues with the same number of labeled atoms and mass but different labeled atom positions. Distinguishing isotopomers requires position-specific information from tandem MS or NMR.

Liquid chromatography-mass spectrometry (LC-MS) and gas chromatography-mass spectrometry (GC-MS): LC-MS and GC-MS separate samples using liquid or gas chromatographic methods prior to

on carbon partitioning in upper glycolysis. $[1,2-^{13}\text{C}_2]$ glucose provides superior information on upper glycolysis and the PPP, especially at the glucose-6-phosphate dehydrogenase (G6PD) branchpoint or in distinguishing the oxidative from the nonoxidative PPP. $[\text{U}-^{13}\text{C}_5]$ glutamine quantifies reductive carboxylation, gluconeogenesis, and glutamine contributions to the TCA cycle [1,16]. The key to a successful labeling experiment is to add physiological concentrations of labels to the culture medium to maximize experimental information. The tracers can be metabolically rearranged in the cells into unique patterns that reflect the underlying metabolic pathways and fluxes. Mathematical tools have been developed to enable researchers to decide the best tracer or best combination of tracers to investigate a metabolic scenario [17,18].

Ideally, cells or tissues are incubated in this labeled medium until the attainment of **isotopic steady state**, after which a single labeling pattern measurement is made. However, if attainment of isotopic steady state is impractical or does not give flux results, the labeling pattern has to be measured transiently *en route* to isotopic steady state and analyzed using **instationary MFA (INST-MFA)**. INST-MFA incorporates intracellular metabolite pool size to estimate isotopomer distributions over time. INST-MFA is a more computationally demanding approach since the isotopomer balances are ordinary differential equations. However, many of the current software options support INST-MFA and allow users to input labeling data from different time points.

At the end of a labeling experiment, cells or tissue are quenched to arrest metabolism, typically by submerging in liquid nitrogen or a cold organic solvent–water mixture, after which metabolites are extracted for analysis. The labeling state is then quantified by mass spectrometry (MS) or nuclear magnetic resonance (NMR) imaging. If the isotope labeling is measured by MS, the fractional enrichment of each mass isotopomer is then assembled into a **mass distribution vector (MDV; Box 1)** for each metabolite of interest.

Different isotope tracers provide complementary information that can be integrated into one holistic iMFA model. For example, $[1,2-^{13}\text{C}_2]$ glucose could be used to provide information at the PPP branchpoint while $[\text{U}-^{13}\text{C}_5]$ glutamine could be used to analyze reductive carboxylation in the TCA cycle. However, when two or more tracers with the same labeled carbon source are fed simultaneously, they can mask each other's labeling patterns and cloud our ability to analyze metabolism. Instead, parallel labeling experiments, in which experiments are repeated with different labeled tracers, can also be used to target multiple metabolic pathways (e.g., labeled glucose for glycolysis and labeled glutamine for the TCA cycle). MDVs generated from parallel labeling experiments are challenging to interpret without a computational framework. The software tools discussed in this article enable the design and analysis of parallel labeling experiments.

The other essential experimental iMFA inputs are extracellular flux measurements, which constrain the intracellular fluxes by defining the absolute metabolite quantities that enter and leave the intracellular metabolic network. To determine extracellular fluxes, nutrients (e.g., glucose) and metabolic byproducts (e.g., lactate) in the cell culture medium are measured by HPLC [19–22], gas/liquid chromatography (GC/LC)-MS [23,24], enzymatic assays, or biochemistry analyzers [25]. External biomass flux can also be determined by measuring change in cell biomass (e.g., dry weight) over time. The extracellular fluxes are then calculated by dividing the change in concentration by the time over which the change occurred. Finally, the fluxes are normalized to cell number and growth rate. Confounding factors such as metabolite degradation and liquid evaporation should also be included.

detection by mass spectrometry.

LC-MS has simpler sample preparation and detects a broader metabolite range. GC-MS better detects sterols, sugars, and very-short-chain fatty acids.

Mass distribution vector (MDV): a distribution of isotopologues for a given metabolite. Each isotopologue's distribution is determined by dividing its ion count by the total metabolite ion count, which is also the sum of all isotopologues.

Metabolic steady state: a state in which metabolic parameters (fluxes and metabolite concentrations) are constant over time, resulting in no net metabolite accumulation. Cell cultures are typically considered at metabolic steady state when they are not proliferating or proliferating at a constant rate. The metabolic steady state assumption eliminates kinetic parameters to simplify iMFA calculations.


Net flux: consider a pair of reversible reactions $A \leftrightarrow B$, where one reaction, say $A \rightarrow B$, has flux v_1 and the other reaction $B \rightarrow A$ has flux v_2 with both v_1 and $v_2 \geq 0$. In this case, the net flux of the reversible reaction pair is the absolute value of the difference between v_1 and v_2 ($|v_1 - v_2|$).

Sink reactions: reactions that serve as an end output for the model. Examples include metabolite efflux (lactate secretion) or intracellular sinks (glycogen storage).

Source reactions: reactions that generate metabolites that are only consumed but not produced by other reactions in the model. Examples include glucose or amino acid extracellular uptake.

Sum of squared residuals (SSR): the sum of the magnitude of difference between model estimates and data measurements. In iMFA, residuals (deviations between measured and simulated isotopologues) are calculated for each metabolite and then summed to obtain the SSR.

Tandem mass spectrometry (tandem MS): multiple mass detectors are used in tandem to distinguish isotopologues with similar m/z ratios. After molecules are fragmented in the first MS, the metabolites are further fragmented in a second MS. These 'fragments of fragments' provide position-specific labeling information, improved sensitivity, and broader coverage.



Trends in Biochemical Sciences

Figure 1. iMFA workflow. Experimental data, including MDVs and extracellular fluxes, and a metabolic network model are inputted into the iMFA software. Each reaction in the network model contains detailed information on the compartments of source and product metabolites, as well as the specific carbon atom transitions that occur in the reaction. In some software, such as INCA and eFlux, compartments can be designated by adding a suffix after the metabolite name (e.g., cytoplasmic as '.c', extracellular as '.e', and mitochondrial as '.m'). The iMFA software then iteratively finds a set of fluxes that minimize the error between the simulated and experimental MDVs. When a good fit is achieved, the iMFA software outputs flux values and associated confidence intervals for each reaction flux. This information can be visualized as a flux map, in which arrow thickness represents the relative flux magnitude. The flux map shown here also indicates source reactions (light blue), sink reactions (dark red, broken), and internal reactions (black). Abbreviations: iMFA, isotope-assisted metabolic flux analysis; MDV, mass distribution vector.

Current limitations in experimental metabolite and mass isotopomer measurements decrease the power of iMFA. Intracellular sinks and sources (e.g., glycogen) cannot be either measured or experimentally constrained, thus giving cells unknown intracellular metabolite inputs and/or outputs. Experimental mass isotopomer data are largely restricted to a subset of metabolites, primarily those in 'core' canonical metabolic pathways such as glycolysis and the TCA cycle. Experiments also mix metabolites from different organelle pools during quenching and extraction, making it impossible to measure metabolite compartmentalization inside the cell. Organelle isolation can

directly quantify compartmentalized metabolites; however, organelle isolation is a lengthy process that can itself perturb metabolism [26].

iMFA process: data fitting by mathematical optimization via iMFA software

iMFA is a mathematical optimization process that determines the intracellular metabolic fluxes that best account for the experimental measurements, given the user-defined network model. iMFA uses metabolite isotopomer mass balance equations to relate the amount of each isotopomer to its precursors and products using the metabolic reaction rates. The biological system is assumed to be at metabolic and often at isotopic steady state, which simplifies the iMFA mathematical problem from ordinary differential equations into algebraic equations.

Practically, users enter the experimental data and network model into iMFA software such as eiFlux [27], INCA [28,29], OpenMebius [30], METRAN, and 13CFLUX2 [31]. The iMFA software generally begins the optimization process with an initial guess for each metabolic flux in the system. The initial fluxes are then used to calculate a simulated MDV for each model metabolite. The

Box 1. Isotopomers are arranged into mass distribution vectors (MDVs) for each metabolite

Stable isotope labeling experiments can be quantified by measuring the incorporation of labeled atoms into metabolites. These isotope labeling patterns are known as isotopomers. For each metabolite, there are 2^n possible unique isotopomers. Here, we provide an example of isotopomers for pyruvate, which has three carbons and can therefore have $2^3 = 8$ possible isotopomers (Figure I). Some isotopomers have the same number of labeled atoms but differ in the position of the labeled atoms. Isotopomers with the same total number of labeled atoms are known as **isotopologues**. These are designated by the mass shift or number of labeled atoms ($M + n$, where n is the number of total labeled atoms) (Figure I). The fractional or percent composition of isotopologues is referred to as an MDV. Traditional MS tools such as GC-MS or LC-MS can only discern between different isotopologues, while tandem MS/MS and NMR can distinguish isotopomers.




Figure I. Isotopomers versus isotopologues for pyruvate.

During LC-MS/GC-MS quantification, the ion count of each isotopologue is quantified. In the case of pyruvate, the unlabeled isotopologue (M+0) has a mass-to-charge ratio (m/z) of 88. Successive isotopologues each have one additional mass unit, with the heaviest isotopologue (M+3) having an m/z of 91. By employing chromatographic separation and MS, each isotopologue for each detectable metabolite can be quantified (Figure II). The ion count for each isotopologue is then normalized to the total ion count of each metabolite (the sum of ion counts for all isotopologues for the metabolite). This yields a fractional isotopologue composition, known as the MDV. Typically, the MDV must then be corrected for naturally occurring isotopes that can skew the calculated fractional isotopologue compositions. The final corrected MDV then serves as an input for iMFA software.




Figure II. Fractional isotopologue composition is normalized and corrected for natural isotope abundance to determine the MDV.

simulated MDVs are compared with the experimental MDVs. The difference between the simulated and experimental MDVs is then represented as an error term, typically the **sum of squared residuals (SSR)**. This process is repeated numerous times, and the flux values that result in the lowest SSR are returned. When the software does not achieve a satisfactory fit (SSR not low enough), the user must reformulate the network model (described in outputs). Once the model attains a satisfactory fit, techniques such as bootstrap Monte-Carlo [27,32] or parameter continuation [28,33] can be used to estimate confidence intervals for each flux in the network (described in detail in the next section).

Current iMFA software is being improved to make it more efficient and offer expanded features for biologically complex models. In most iMFA software, the state variables (isotopomers) and model parameters (fluxes) are kept distinct from each other, which requires the isotopomer abundances to be computed from scratch in each iteration. By contrast, eiFlux performs the optimization with no distinction between state variables and model parameters. Thus, during each iteration of the optimization, the system state is updated instead of being calculated from scratch. eiFlux also uses state-of-the-art optimization solvers such as the General Algebraic Modeling System (GAMS), thus facilitating efficient, robust parameter estimation and superior scalability to large metabolic networks [27].

New iMFA techniques are also being developed to analyze biological systems when **metabolic steady state** cannot be reached. Indeed, it is essential to study changes in cell metabolism during dynamic processes such as differentiation, activation, and proliferation. Dynamic ^{13}C -MFA (DMFA) is an emerging non-steady state approach; however, DMFA has experimental and computational challenges, since varying MDVs and fluxes complicates measurements and flux estimation. In a recent study, curve fitting was used to estimate non-steady state glycolytic fluxes from insulin-treated adipocytes that were metabolically sampled over time. While the DMFA model replicated the experimental metabolic trends, the model did not achieve a statistically acceptable fit [34]. Additional progress in DMFA is essential to estimate non-steady state fluxes.

iMFA is also being adapted to incorporate **tandem MS**, an emerging technique potentially with the benefits of both MS and NMR. Tandem MS breaks precursor ions into product ion fragments, which enables position-specific atom information to be acquired. Tandem MS has higher accuracy than **GC-MS** or **LC-MS** and improved detection of low abundance isotopomers compared with NMR. To incorporate tandem MS data, the mathematical iMFA framework must be expanded to integrate the fragmented isotopomer information. To date, only eiFlux enables users to input tandem MS data [27].

Finally, iMFA software is progressing to make computationally feasible genome-scale models, which can improve the simulation fit to the experimental data. Parallelized algorithms have been used to perform genome-scale INST-MFA with improved speed [35]. Two-scale MFA, which uses iMFA data to constrain flux predictions in a genome-scale model, has also been implemented [36]. However, genome-scale models with large metabolic networks and limited experimental data can produce large flux variabilities. Parsimonious iMFA, which posits that evolutionary pressure selects for minimal energy expenditure, provides one solution. Parsimonious iMFA runs a second optimization after standard iMFA and selects the flux solution that minimizes the weighted sum of all fluxes [37]. Parsimonious iMFA can integrate transcriptomic data to minimize fluxes associated with low gene expression [38]. Recently, a proof-of-concept study demonstrated that when compared with traditional iMFA, parsimonious iMFA results in improved flux predictions compared with traditional iMFA. However, the computational complexity of this approach limits its applicability to genome-scale models [37].

Output: metabolic flux map and statistical analysis

The statistical analysis of quantitative flux outputs is a key advantage of iMFA over manual tracer analysis. After an acceptable fit is achieved, iMFA generates a quantitative metabolic flux map of reaction flux estimates for all fluxes in the network, along with their respective confidence intervals. The confidence intervals for a given flux are then compared among experimental conditions, and if the confidence intervals do not overlap, then the fluxes are statistically significantly different. Statistical tests could be used to further test significance. Wide confidence intervals indicate high reaction flux uncertainty and suggest that the current experimental strategy is not suitable for flux resolution. In this case, the experimental design should be improved, for example, by selecting different tracers or by integrating data from parallel labeling experiments [39,40].

In some cases, iMFA does not find a set of metabolic fluxes that enable the simulation to fit the measured metabolite concentrations. When the SSR does not fall within the acceptable range, the user may need to change the metabolic network model (e.g., adding reaction reversibility, compartmentalization, or alternative routes). Visualization of the experimental and simulated MDVs with tools such as Escher-Trace [41] is helpful in determining which reactions should be altered. This iterative process can lead to valuable insights, especially in elucidating alternative or unique metabolic pathways.

iMFA resolves metabolic complexity

Large data sets

iMFA facilitates analysis of large data sets from parallel and instationary labeling experiments. These labeling strategies are well suited for mammalian cells, which can be prone to slow labeling due to high exchange rates between intracellular and extracellular metabolite pools [6,42]. Furthermore, the wealth of data can more accurately resolve metabolic fluxes in the context of the whole network, as well as buffer effects from measurement errors [43,44]. While these large data sets would be unwieldy to analyze manually, iMFA and INST-MFA are designed to easily incorporate extensive metabolic data.

In a recent INST-MFA study of resting and activated human platelets, the investigators used parallel labeling with mixed tracers to study both glucose and acetate metabolism. Platelets were incubated with either a mixture of [1,2-¹³C₂]glucose, [U-¹³C₆]glucose, and unlabeled glucose or [1-¹³C]acetate, [2-¹³C]acetate, and unlabeled acetate. Samples were then collected five times for each labeling experiment. The MDVs and extracellular fluxes were integrated into an INST-MFA model using INCA. Through this approach, the authors determined that activated platelets redirect glucose flux from the PPP and TCA cycle toward lactate [45].

INST-MFA has also been used to analyze the metabolic impact of iPSC differentiation into cardiomyocytes [46], neural stem cell differentiation [47], brown adipose tissue cold activation [48,49], paclitaxel treatment of cancer cells [50], and infection-induced metabolic changes [51,52]. Additional iMFA studies used parallel labeling, including combinations such as glucose/acetate [52], [1,2-¹³C₂]glucose/[U-¹³C₅]-glutamine [50,51,53–56], with and without palmitate [53]. The rigorous iMFA framework transforms complex data from these experiments into intuitive quantitative flux maps.


Cyclic pathways and reversible reactions

iMFA is especially useful in understanding labeling data from complex metabolic pathways with cyclic pathways, reversible reactions, and **exchange fluxes**. The TCA cycle is a classic example of metabolic complexity, since isotopomer labeling patterns change as metabolites pass through multiple TCA turns (Figure 2). Whereas manual MDV analysis can usually only account for one or two TCA cycle turns, iMFA can accurately model any number of TCA cycle turns (from one to infinity). Inclusion of infinite TCA cycle turns can clarify TCA cycle activity. For example, a recent study examined whether exogenous citrate was first metabolized in the cytosol or the mitochondria in hypoxic hepatocellular carcinoma (HCC) cells [57]. When MDVs from HCC labeled with [2,4-¹³C₂]citrate were manually interpreted assuming only one TCA cycle turn, TCA intermediate labeling patterns were indefinite and showed evidence of direct metabolism in both compartments. By contrast, iMFA with infinite TCA cycle turns unveiled significant cytosolic citrate metabolism, with exogenous citrate primarily entering the mitochondria as α-ketoglutarate. Thus, by considering infinite TCA cycle turns, iMFA uncovered compartmentalization of exogenous citrate metabolism.

iMFA can also account for reversible TCA cycle reactions, in particular reductive carboxylation of α-ketoglutarate via reversed isocitrate dehydrogenase (IDH) flux. In human bronchial epithelial cells treated with cigarette smoke condensate, [U-¹³C₅]glutamine labeling suggested increased glutamine enrichment of acetyl-coenzyme A (CoA) and [M+5]citrate. iMFA showed that cigarette smoke-treated cells completely reversed IDH flux, which was driven by glutamine consumption [58].

Similar to the TCA cycle, the PPP is complicated by both cyclic and reversible reactions, since fructose-6-phosphate (F6P) and glyceraldehyde-3-phosphate (G3P) are both products and substrates of the PPP (Figure 3). In a recent study, iMFA revealed that oxidative PPP flux increased over 200% during oxidative burst in neutrophils. The change was fueled by a switch to a cyclic PPP phenotype, as well as reversal of the reaction from F6P into G6P. These insights were then confirmed by knockout of nonoxidative PPP enzymes [59].

iMFA can also quantify bidirectional metabolite fluxes in reversible reactions. Most biochemical reactions are reversible or bidirectional and are defined in terms of the **net flux** (difference between forward and reverse fluxes) and the exchange flux (minimum of the forward and reverse fluxes). While the net flux indicates the magnitude and direction of the reaction, the exchange flux



Trends in Biochemical Sciences


Figure 2. Isotope-assisted metabolic flux helps quantify multiple tricarboxylic acid (TCA) cycle turns. Cyclic metabolic networks, such as the TCA cycle, generate a different set of isotopomers at each turn. This simplified TCA cycle example assumes that uniformly labeled glucose and an unlabeled oxaloacetate (OAA) are the only carbon contributors. Each color represents a different TCA cycle turn. The initial unlabeled OAA reacts with glucose-derived, fully labeled acetyl-coenzyme A (CoA) to form [M+2] citrate via citrate synthase (CS). This citrate is then metabolized into α-ketoglutarate (AKG) via isocitrate dehydrogenase (IDH). As the AKG progresses through the TCA cycle (metabolites not shown), ‘flipped’ isotopomers of OAA with either the first and second or the third and fourth carbons labeled can occur. When these two OAA isotopomers react with fully labeled acetyl-CoA to form [M+4] citrate in the second TCA cycle turn, two unique citrate and AKG isotopomers are created. With each turn of the TCA cycle, more isotopomers are created in different concentrations until, with infinite TCA cycle turns, the metabolites reach metabolic steady state.

quantifies the amount by which the label back-mixes through the reaction [60]. Similar labeling patterns between a product and substrate metabolite (e.g., alanine and pyruvate or succinate and malate) often indicate high exchange fluxes. If a product and substrate metabolite are labeled differently, this can suggest low exchange, which can then be quantified using iMFA. Alternatively, labeling dilution can occur if there is high exchange flux with an unlabeled source [28].

Unlabeled sources and anaplerosis

iMFA can elucidate how different sources contribute to intracellular metabolic fluxes due to its ability to integrate many input and output metabolites into a single analysis. We used iMFA to study how inhibition of glycolytic side branch pathways (hexosamine biosynthetic pathway, PPP, polyol pathway) impacted holistic human endothelial cell glucose metabolism. Manual MDV analysis showed that inhibitors decreased labeling in each pathway as well as in the TCA cycle, which we initially interpreted to indicate decreased metabolic activity. However, by integrating extracellular fluxes and isotopomer data into an iMFA model, we discovered that decreased TCA cycle labeling was explained by influx of unlabeled glutamine, which diluted TCA cycle labeling and actually fueled increased TCA cycle activity [25].

The role of other **anaplerotic** substrates in metabolism can also be clarified using iMFA. Pyruvate is particularly important for the TCA cycle, as it can generate both oxaloacetate (via pyruvate carboxylase) and acetyl-CoA (via pyruvate dehydrogenase). Each route generates a unique set of isotopomers, complicating reaction flux determination from MDVs [61]. A modified iMFA model of whole-body glucose production and gluconeogenesis was used to understand how loss of pyruvate carboxylase impacts metabolic homeostasis in liver-specific pyruvate carboxylase knock-out mice. iMFA showed that liver gluconeogenesis from TCA anaplerosis in the pyruvate



Trends in Biochemical Sciences

Figure 3. Isotope-assisted metabolic flux analysis (iMFA) helps analyze complexity from the cyclic and reversible pentose phosphate pathway (PPP) pathway. (A) In glycolysis, glucose is metabolized into glucose-6-phosphate (G6P), which can continue down through glycolysis or get shuttled into the PPP. The PPP is a complex pathway due to multiple reversible reactions and cyclic interactions with the glycolytic intermediates fructose-6-phosphate (F6P) and glyceraldehyde-3-phosphate (G3P). If cells are labeled with uniformly labeled glucose, forward PPP flux will generate uniformly labeled ribose-5-phosphate (R5P). However, reaction reversibility and complex atom rearrangements can generate all 32 possible R5P isotopomers (red). Even when simplified into mass distribution vectors (MDVs), the data can still be too complex for manual interpretation. (B) Using iMFA, we showed that endothelial cells treated with dehydroandrosterone to inhibit the first and rate-limiting PPP reaction show decreased PPP fluxes and net fluxes in the forward direction [21] (left). iMFA also showed that neutrophils undergoing oxidative burst have forward PPP flux reversal of G6P isomerase activity. This causes F6P to be metabolized into G6P, fueling cyclic PPP flux [53] (right).

carboxylase knockout mice was less than 15% of wild-type liver. However, loss of pyruvate carboxylase depleted NADPH and glutathione to increase liver oxidative stress and inflammation [62]. In our study of human endothelial cell metabolism, iMFA showed that more pyruvate was routed through pyruvate dehydrogenase than pyruvate carboxylase, and that pyruvate dehydrogenase fluxes changed under metabolic inhibition while pyruvate carboxylase fluxes remained relatively stable [25].

Pathway discovery


iMFA also enables the discovery of new metabolic pathways. When simulated MDVs cannot fit the experimental MDVs, the metabolic network model must be modified. This process often suggests alternative metabolic pathways to then be tested experimentally. In lung cancer cells grown as spheroids or monolayers, iMFA initially could not fit simulated fluxes to the spheroid isotopomer labeling data because citrate showed reductive carboxylation but palmitate, which is produced from citrate, did not. The metabolic network model was then modified so that citrate produced by reductive carboxylation in the cytosol entered the mitochondria, where it mixed with mitochondrial citrate before being exported back to the cytosol for palmitate synthesis. This alteration significantly improved the simulation fit. Cytosolic citrate transport into the mitochondria was then validated experimentally using citrate transporter knockout cells [63]. iMFA was also used to uncover evidence that transketolase-like protein 1 (TKTL1) cleaves xylulose-5-phosphate to yield G3P and acetyl-CoA in mammalian cells, since including the TKTL1 reaction was the only way to achieve an acceptable fit [64]. In our study on endothelial metabolism, we initially could not fit the simulation to the experimental data due to excess unlabeled pyruvate. By iteratively modifying the model, we identified a novel four-carbon source that fed into the malate-pyruvate shuttle as a potential source for these unlabeled carbons [25].

Compartmentalization and cell-cell interactions

iMFA can analytically parse data that were mixed experimentally back into separate model compartments. In a study of Chinese hamster ovary cells, the initial iMFA model produced a poor fit due to low labeled pyruvate, alanine, lactate, and glutamate levels. A second model, which included a mitochondrial compartment separate from the cytosol, successfully fit the data. This model further predicted that malic enzyme, which catalyzes conversion of malate to pyruvate, was only active in mitochondria [42].

Cell co-culture systems, which are vital to understand physiological interactions among cells, are even more complex to analyze metabolically. Co-cultures have multiple cell types each with their own extracellular and intracellular fluxes. Experimentally, co-cultures can be created with varying degrees of interaction (Figure 4). Conditioned media experiments keep the cells distinct, enabling extracellular and intracellular metabolite separation; transwell experiments allow cells to interact through the media while separating cells via a permeable membrane, enabling intracellular metabolite separation only; and direct co-culture mixes the cells and their metabolites. A combination of these experimental techniques is likely needed to describe metabolic interactions in a co-culture.

iMFA was used to analyze a co-culture of two *Escherichia coli* knockout strains without experimental separation. Each cell type was modeled *in silico* as a separate compartment with a complete metabolic reaction set. Each metabolite MDV was weighted by the fraction of each strain in the total cell population, and secreted metabolites were allowed to exchange between the two cell compartments. The co-culture two-compartment network model achieved a satisfactory fit [65]. In another study, cell-cell interactions via extracellular vesicles (EVs) were analyzed via a custom Exo-MFA model with terms for EV cargo packaging, secretion, and cargo release into



Trends in Biochemical Sciences

Figure 4. A variety of experimental co-culture methods are essential to determining metabolic interactions among different cell types. In 2D culture (top row), conditioned media from one cell type can be applied to a different cell type (A), cells can be co-cultured using a permeable insert so that they either share soluble factors through the media (left) or directly contact each other through the membrane (right, B), or cells can be directly co-cultured together in a dish (C). These same techniques can be used for 3D cell cultures (middle row), with cell spheroid media applied to an *in vitro* vessel-like network (D), cell spheroids co-cultured with a cell monolayer using a permeable membrane (E), or cell spheroids directly cultured on a vessel-like network (F). Finally, stimuli such as flow can be applied to co-cultures to examine metabolic interactions under more physiologically relevant conditions (bottom row). Flow can be applied to one set of cells using a cone and plate device, and conditioned media from the flow-adapted cells can be applied to another cell culture (G). Microfluidics can also be used to apply flow to cells in contact with another type of cells through a permeable membrane (H), or to 3D cell cultures like spheroids surrounding a perfused engineered vessel (I). Figure created with BioRender.

recipient cells [66]. The study showed that TCA fluxes increased 6 h after EV exposure, likely due to lactate, glutamine, and TCA intermediate cargo delivery. However, further work is required to improve co-culture analysis, including identifying novel labeling strategies and integrating multiple types of co-culture experiments.

In vivo flux determination

Over the past decade, iMFA has increasingly been used to interpret data from *in vivo* stable isotope labeling experiments in animals and humans [67]. Since *in vivo* iMFA is more complex (Box 2) and metabolite uptake and secretion cannot be measured *in vivo*, iMFA has primarily been used to study endogenous metabolite production. iMFA was used to investigate liver glucose production in mice that underwent short- and long-term fasting [41] and treadmill exercise [68], renal glucose production in liver-specific pyruvate carboxylase knockout mice [69], and liver ketogenesis [70]. Notably, the latter two examples required more complex, multicompartmental network models to integrate metabolites from multiple sources. In these studies, *in vivo* iMFA enabled the authors to both calculate metabolic fluxes *in vivo* and quantify how diverse metabolic pathways (TCA cycle, gluconeogenesis, glycogen) contribute to endogenous metabolite production. For more details, we direct the readers to other reviews on this topic [67,71,72].

Box 2. iMFA limitations

General iMFA limitations

Although iMFA provides a quantitative, systems-level perspective of metabolism, its value relies on the quality of acquired experimental data. When the selected tracer does not generate informative labeling patterns, the iMFA output fluxes will have low precision and large confidence intervals [16,73]. Measurement errors can also be problematic, as they skew flux estimates or impair model fit. These data-driven limitations can lead to erroneous conclusions, especially as the user may add or remove pathways and thereby generate an inaccurate metabolic model in an attempt to fit the data.

Current iMFA models only account for a limited subset of metabolic pathways (glycolysis, TCA, PPP, fatty acid metabolism). iMFA model sizes are limited by lack of information on atom transitions in some pathways, the inability to measure some metabolites, and computational power [74]. However, the exclusion of nonessential pathways can bias results.

Finally, most current iMFA approaches require systems to be at metabolic steady state and can therefore not account for metabolic changes over time, which are often essential to fully understanding biological processes. Dynamic iMFA approaches that can estimate non-steady state metabolic fluxes approaches are still in their infancy [34,75].

In vivo-specific iMFA limitations

iMFA can also be used to determine quantitative tissue-level and whole-body metabolic fluxes for *in vivo* studies. However, physiological complexity can greatly complicate iMFA. First, metabolite tracers must be carefully selected and validated to ensure they are metabolized by the tissue of interest and that the labeled tracers do not alter *in vivo* metabolic fluxes when added at concentrations sufficient to enrich downstream metabolites. Second, repeated sampling can be harmful to animals and therefore should be minimized for animal welfare. Thus, isotopic steady state may be difficult to measure for iMFA, and dynamic *in vivo* measurements are usually not possible for INST-MFA. Third, interorgan metabolite exchange makes quantifying tissue-specific metabolite uptake and secretion difficult, which means that there are no constraining extracellular fluxes for iMFA.

Labeled substrates can be introduced to animals via oral gavage [76] or liquid diet [77]. These methods can lead to more biologically relevant metabolism but may not provide the necessary information (e.g., infusion rate) for iMFA. A key breakthrough for *in vivo* iMFA was development of a minimally invasive, dual arterial-venous catheterization technique to continuously infuse tracers [43,72]. This both minimizes stress due to repeat experimental sampling and provides a known infusion rate, which is important for flux value estimations. Dual arterial-venous catheterization, along with sensitive MS tools, also enables sampling of microliter volumes, which make this approach practical in small animals [43]. Drawbacks to this method include technical complexity, cost, invasiveness, and reduced relevance for dietary studies [67].

Outstanding questions

iMFA co-culture analysis is complicated by the experimental mixing of metabolite pools from multiple cell types. What experimental and computational strategies can be used to determine cell-specific fluxomics and cell-cell metabolic interactions in co-culture models?

Currently, iMFA requires systems to be at a metabolic steady state; however, cell metabolism plays a critical role in dynamic cell processes. How can we advance iMFA approaches to analyze dynamic metabolic systems?

Genome-scale iMFA is limited by both experimental and computational approaches. For experimental approaches, how can we improve MS and NMR tools to collect additional isotopologues? For computational approaches, how can we increase computational efficiency to scale and integrate genome-scale isotopologue data?

Concluding remarks and future perspectives

iMFA generates quantitative flux maps from complex metabolomic data, enabling the end users to discover metabolic differences among cell states and generate new hypotheses that then can be tested experimentally. Many iMFA discoveries could not have been intuited from manual analysis of isotope labeling. We envision that iMFA will continue to yield new insights into metabolism, especially as larger models and dynamic approaches are developed. iMFA has great potential in elucidating the importance of cell metabolism in stem cell differentiation, rare inherited metabolic disorders, and diseases related to metabolic dysfunction such as cardiovascular disease, diabetes, and obesity. However, to achieve this potential, iMFA must continue to improve in its ability to manage metabolic complexity (see [Outstanding questions](#)). As computational and experimental techniques advance, iMFA approaches should be both increasingly accessible and increasingly powerful for metabolism research.

Acknowledgments

This research was funded by National Science Foundation CBET-2211966 to A.M.C. and G.S., National Institutes of Health 1R01HL140239 to A.M.C., NSF DGE-1632976 to B.M., NSF MCB-1517671 to G.S., and a seed grant from the Maryland Brain and Behavior Institute to A.M.C. and G.S.

Declaration of interests

No interests are declared.

References

- Jang, C. *et al.* (2018) Metabolomics and isotope tracing. *Cell* 173, 822–837
- Suthers, P.F. *et al.* (2007) Metabolic flux elucidation for large-scale models using ^{13}C labeled isotopes. *Metab. Eng.* 9, 387–405
- Nielsen, J. (2003) It is all about metabolic fluxes. *J. Bacteriol.* 185, 7031
- Buescher, J.M. *et al.* (2015) A roadmap for interpreting ^{13}C metabolite labeling patterns from cells. *Curr. Opin. Biotechnol.* 34, 189–201
- Antoniewicz, M.R. (2015) Methods and advances in metabolic flux analysis: a mini-review. *J. Ind. Microbiol. Biotechnol.* 42, 317–325
- Dai, Z. and Locasale, J.W. (2017) Understanding metabolism with flux analysis: from theory to application. *Metab. Eng.* 43, 94–102
- Karlsen, E. *et al.* (2018) Automated generation of genome-scale metabolic draft reconstructions based on KEGG. *BMC Bioinform.* 19, 467
- Kanehisa, M. *et al.* (2008) KEGG for linking genomes to life and the environment. *Nucleic Acids Res.* 36, D480–D484
- Noronha, A. *et al.* (2018) The Virtual Metabolic Human database: integrating human and gut microbiome metabolism with nutrition and disease. *Nucleic Acids Res.* 47, D614–D624
- Caspi, R. *et al.* (2020) The MetaCyc database of metabolic pathways and enzymes—a 2019 update. *Nucleic Acids Res.* 48, D455–D453
- Caspi, R. *et al.* (2017) The MetaCyc database of metabolic pathways and enzymes. *Nucleic Acids Res.* 46, D633–D639
- Orman, M.A. *et al.* (2012) Stoichiometry based steady-state hepatic flux analysis: computational and experimental aspects. *Metabolites* 2, 268–291
- Wiback, S.J. and Palsson, B.O. (2002) Extreme pathway analysis of human red blood cell metabolism. *Biophys. J.* 83, 808–818
- Zhang, Z.W. *et al.* (2011) Red blood cell extrudes nucleus and mitochondria against oxidative stress. *J. Biol. Chem.* 286, 560–565
- Lewis, C.A. *et al.* (2014) Tracing compartmentalized NADPH metabolism in the cytosol and mitochondria of mammalian cells. *Mol. Cell* 55, 253–263
- Metallo, C.M. *et al.* (2009) Evaluation of ^{13}C isotopic tracers for metabolic flux analysis in mammalian cells. *J. Biotechnol.* 144, 167–174
- Nargund, S. and Sriram, G. (2013) Designer labels for plant metabolism: statistical design of isotope labeling experiments for improved quantification of flux in complex plant metabolic networks. *Mol. Biosyst.* 9, 99–112
- Nöh, K. *et al.* (2018) A Pareto approach to resolve the conflict between information gain and experimental costs: multiple-criteria design of carbon labeling experiments. *PLoS Comput. Biol.* 14, e1006533
- Bergman, T. *et al.* (1986) Amino acid analysis by high performance liquid chromatography of phenylthiocarbonyl derivatives. In *Advanced Methods in Protein Microsequence Analysis* (Wittmann-Liebold, B. *et al.*, eds), pp. 45–55. Springer Berlin Heidelberg
- Sherwood, R.A. (1990) Amino acid measurement by high-performance liquid chromatography using electrochemical detection. *J. Neurosci. Methods* 34, 17–22
- Ling, Z. *et al.* (2016) Sensitive determination of glucose in Dulbecco's modified Eagle medium by high-performance liquid chromatography with 1-phenyl-3-methyl-5-pyrazolone derivatization: application to gluconeogenesis studies. *Biomed. Chromatogr.* 30, 601–605
- Robitaille, L. and Hoffer, L.J. (1988) Measurement of branched chain amino acids in blood plasma by high performance liquid chromatography. *Can. J. Physiol. Pharmacol.* 66, 613–617
- Zhu, W. *et al.* (2011) Quantitative profiling of tryptophan metabolites in serum, urine, and cell culture supernatants by liquid chromatography–tandem mass spectrometry. *Anal. Bioanal. Chem.* 401, 3249–3261
- Lau, S.K.P. *et al.* (2015) Identification of specific metabolites in culture supernatant of *Mycobacterium tuberculosis* using metabolomics: exploration of potential biomarkers. *Emerg. Microbes Infect.* 4, 1–10
- Moiz, B. *et al.* (2021) ^{13}C Metabolic flux analysis indicates endothelial cells attenuate metabolic perturbations by modulating TCA activity. *Metabolites* 11, 226
- Lee, W.D. *et al.* (2019) Spatial-fluxomics provides a subcellular-compartmentalized view of reductive glutamine metabolism in cancer cells. *Nat. Commun.* 10, 1351
- Lugar, D.J. and Sriram, G. (2022) Isotope-assisted metabolic flux analysis as an equality-constrained nonlinear program for improved scalability and robustness. *PLoS Comput. Biol.* 18, e1009831
- Young, J.D. (2014) INCA: a computational platform for isotopically non-stationary metabolic flux analysis. *Bioinformatics* 30, 1333–1335
- Rahim, M. *et al.* (2022) INCA 2.0: a tool for integrated, dynamic modeling of NMR- and MS-based isotopomer measurements and rigorous metabolic flux analysis. *Metab. Eng.* 69, 275–285
- Kajihata, S. *et al.* (2014) OpenMebius: an open source software for isotopically nonstationary ^{13}C -based metabolic flux analysis. *Biomed. Res. Int.* 2014, 627014
- Weitzel, M. *et al.* (2013) 13CFLUX2—high-performance software suite for ^{13}C -metabolic flux analysis. *Bioinformatics* 29, 143–145
- Nargund, S. and Sriram, G. (2014) Mathematical modeling of isotope labeling experiments for metabolic flux analysis. *Methods Mol. Biol.* 1083, 109–131
- Antoniewicz, M.R. *et al.* (2006) Determination of confidence intervals of metabolic fluxes estimated from stable isotope measurements. *Metab. Eng.* 8, 324–337
- Quek, L.E. *et al.* (2020) Dynamic ^{13}C flux analysis captures the reorganization of adipocyte glucose metabolism in response to insulin. *iScience* 23, 100855
- Zhang, Z. *et al.* (2020) Parallel isotope differential modeling for stationary ^{13}C fluxomics at the genome scale. *Biotechnol. Biofuels* 13, 103
- Ando, D. and Garcia, M.H. (2018) Two-scale ^{13}C metabolic flux analysis for metabolic engineering. *Methods Mol. Biol.* 1671, 170–179
- Foguet, C. *et al.* (2019) p ^{13}C MF: parsimonious ^{13}C metabolic flux analysis. *PLoS Comput. Biol.* 15, e1007310
- Jenior, M.L. *et al.* (2020) Transcriptome-guided parsimonious flux analysis improves predictions with metabolic networks in complex environments. *PLoS Comput. Biol.* 16, e1007099
- Wang, Y. *et al.* (2020) Metabolic flux analysis—linking isotope labeling and metabolic fluxes. *Metabolites* 10, 1–21
- Long, C.P. and Antoniewicz, M.R. (2019) High-resolution C metabolic flux analysis. *Methods* 14, 2856–2877
- Kumar, A. *et al.* (2020) Escher-Trace: a web application for pathway-based visualization of stable isotope tracing data. *BMC Bioinform.* 21, 297
- Nicolae, A. *et al.* (2014) Non-stationary ^{13}C metabolic flux analysis of Chinese hamster ovary cells in batch culture using extracellular labeling highlights metabolic reversibility and compartmentation. *BMC Syst. Biol.* 8, 50
- Hasenour, C.M. *et al.* (2015) Mass spectrometry-based microassay of ^2H and ^{13}C plasma glucose labeling to quantify liver metabolic fluxes in vivo. *Am. J. Physiol. Endocrinol. Metab.* 309, E191–E203
- Crown, S.B. *et al.* (2016) Comprehensive metabolic modeling of multiple ^{13}C -isotopomer data sets to study metabolism in perfused working hearts. *Am. J. Physiol. Heart Circ. Physiol.* 311, H881–H891
- Sake, C.L. *et al.* (2022) Isotopically nonstationary ^{13}C metabolic flux analysis in resting and activated human platelets. *Metab. Eng.* 69, 313–322
- Correia, C. *et al.* (2017) 3D aggregate culture improves metabolic maturation of human pluripotent stem cell derived cardiomyocytes. *Biotechnol. Bioeng.* 115, 630–644
- Sá, J.V. *et al.* (2017) Quantification of metabolic rearrangements during neural stem cells differentiation into astrocytes by metabolic flux analysis. *Neurochem. Res.* 42, 244–253
- Wang, Z. *et al.* (2020) Chronic cold exposure enhances glucose oxidation in brown adipose tissue. *EMBO Rep.* 21, e50085
- Held, N.M. *et al.* (2018) Pyruvate dehydrogenase complex plays a central role in brown adipocyte energy expenditure and fuel utilization during short-term beta-adrenergic activation. *Sci. Rep.* 8, 9562
- Araki, C. *et al.* (2018) Mass spectrometry-based method to study inhibitor-induced metabolic redirection in the central metabolism of cancer cells. *Mass Spectrom.* 7, A0067–A0067

51. Carinhas, N. *et al.* (2017) ^{13}C -metabolic flux analysis of human adenovirus infection: implications for viral vector production. *Biotechnol. Bioeng.* 114, 195–207
52. Mackenzie, J.S. *et al.* (2020) Bedaquiline reprograms central metabolism to reveal glycolytic vulnerability in *Mycobacterium tuberculosis*. *Nat. Commun.* 11, 6092
53. Parker, S.J. *et al.* (2017) LKB1 promotes metabolic flexibility in response to energy stress. *Metab. Eng.* 43, 208–217
54. Noguchi, S. *et al.* (2020) Direct and quantitative analysis of altered metabolic flux distributions and cellular ATP production pathway in fumarate hydratase-diminished cells. *Sci. Rep.* 10, 13065
55. Lesner, N.P. *et al.* (2020) α -ketobutyrate links alterations in cystine metabolism to glucose oxidation in mtDNA mutant cells. *Metab. Eng.* 60, 157–167
56. Odenwelder, D.C. *et al.* (2021) Induced pluripotent stem cells can utilize lactate as a metabolic substrate to support proliferation. *Biotechnol. Prog.* 37, e3090
57. Kumar, A. *et al.* (2021) NaCT/SLC13A5 facilitates citrate import and metabolism under nutrient-limited conditions. *Cell Rep.* 36, 109701
58. Rahman, S.M.J. *et al.* (2016) The airway epithelium undergoes metabolic reprogramming in individuals at high risk for lung cancer. *JCI Insight* 1, e88814
59. Britt, E.C. *et al.* (2022) Switching to the cyclic pentose phosphate pathway powers the oxidative burst in activated neutrophils. *Nat. Metab.* 4, 389
60. Wiechert, W. (2007) The thermodynamic meaning of metabolic exchange fluxes. *Biophys. J.* 93, 2255
61. Alves, T.C. *et al.* (2015) Integrated, step-wise, mass-isotopomer flux analysis of the TCA cycle. *Cell Metab.* 22, 936–947
62. Cappel, D.A. *et al.* (2019) Pyruvate-carboxylase-mediated anaplerosis promotes antioxidant capacity by sustaining TCA cycle and redox metabolism in liver. *Cell Metab.* 29, 1291–1305.e8
63. Jiang, L. *et al.* (2017) Quantitative metabolic flux analysis reveals an unconventional pathway of fatty acid synthesis in cancer cells deficient for the mitochondrial citrate transport protein. *Metab. Eng.* 43, 198–207
64. Ahn, W.S. *et al.* (2016) Evidence for transketolase-like TKTL1 flux in CHO cells based on parallel labeling experiments and ^{13}C -metabolic flux analysis. *Metab. Eng.* 37, 72
65. Gebreselassie, N.A. and Antoniewicz, M.R. (2015) ^{13}C -metabolic flux analysis of co-cultures: a novel approach. *Metab. Eng.* 31, 132–139
66. Achreja, A. *et al.* (2017) Exo-MFA – a ^{13}C metabolic flux analysis framework to dissect tumor microenvironment-secreted exosome contributions towards cancer cell metabolism. *Metab. Eng.* 43, 156–172
67. Fernández-García, J. *et al.* (2020) Stable isotopes for tracing mammalian-cell metabolism in vivo. *Trends Biochem. Sci.* 45, 185–201
68. Hughey, C.C. *et al.* (2017) Loss of hepatic AMP-activated protein kinase impedes the rate of glycogenolysis but not gluconeogenic fluxes in exercising mice. *J. Biol. Chem.* 292, 20125–20140
69. Rahim, M. *et al.* (2021) Multitissue $2\text{H}/^{13}\text{C}$ flux analysis reveals reciprocal upregulation of renal gluconeogenesis in hepatic PEPCK-C-knockout mice. *JCI Insight* 6, e149278
70. Deja, S. *et al.* (2021) In vivo estimation of ketogenesis using metabolic flux analysis-technical aspects and model interpretation. *Metabolites* 11, 279
71. Moiz, B. *et al.* (2022) Isotope-assisted metabolic flux analysis: a powerful technique to gain new insights into the human metabolome in health and disease. *Metabolites* 12, 1066
72. Bednarski, T.K. *et al.* (2021) In vivo $2\text{H}/^{13}\text{C}$ flux analysis in metabolism research. *Curr. Opin. Biotechnol.* 71, 1–8
73. Walther, J.L. *et al.* (2012) Optimization of ^{13}C isotopic tracers for metabolic flux analysis in mammalian cells. *Metab. Eng.* 14, 162–171
74. Basler, G. *et al.* (2018) Advances in metabolic flux analysis toward genome-scale profiling of higher organisms. *Biosci. Rep.* 38, BSR20170224
75. Martínez, V.S. *et al.* (2015) Dynamic metabolic flux analysis using B-splines to study the effects of temperature shift on CHO cell metabolism. *Metab. Eng. Commun.* 2, 46–57
76. Williams, H.C. *et al.* (2020) Oral gavage delivery of stable isotope tracer for in vivo metabolomics. *Metabolites* 10, 501
77. Sun, R.C. *et al.* (2017) Noninvasive liquid diet delivery of stable isotopes into mouse models for deep metabolic network tracing. *Nat. Commun.* 8, 1646