



OPEN ACCESS

EDITED AND REVIEWED BY
Elise Huchard,
UMR5554 Institut des Sciences de
l'Evolution de Montpellier
(ISEM), France

*CORRESPONDENCE
Richard Reed Veit
rrveit23@gmail.com

SPECIALTY SECTION

This article was submitted to
Behavioral and Evolutionary Ecology,
a section of the journal
Frontiers in Ecology and Evolution

RECEIVED 03 June 2022

ACCEPTED 29 June 2022

PUBLISHED 15 July 2022

CITATION

Veit RR, Manne LL, Zawadzki LC,
Alamo MA and Henry RW III (2022)
Editorial: Vagrancy, exploratory
behavior and colonization by birds:
Escape from extinction?
Front. Ecol. Evol. 10:960841.
doi: 10.3389/fevo.2022.960841

COPYRIGHT

© 2022 Veit, Manne, Zawadzki, Alamo
and Henry. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution
License \(CC BY\)](#). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Editorial: Vagrancy, exploratory behavior and colonization by birds: Escape from extinction?

Richard Reed Veit^{1,2*}, Lisa Louise Manne^{1,2},
Lucinda C. Zawadzki³, Marlen Acosta Alamo^{1,2} and
Robert William Henry III⁴

¹Biology Department, College of Staten Island, Staten Island, NY, United States, ²CUNY Graduate School, New York, NY, United States, ³Department of Zoology, University of Oxford, Oxford, United Kingdom, ⁴Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, Santa Cruz, CA, United States

KEYWORDS

vagrancy, exploratory behavior, dispersal, colonization, range expansion, intelligent dispersal


Editorial on the Research Topic

Vagrancy, Exploratory Behavior And Colonization by Birds: Escape From Extinction?

Vagrant birds, long thought to be mutant or otherwise navigationally incompetent individuals, have now been shown to be the outer fringe of a growing population; vagrancy is the mechanism by which growing populations can colonize newly available habitats (Grinnell, 1922; Baker, 1978; Veit, 2000, 2008; Zawadzki et al., 2019). Since changes habitats are changing at a greater rate and magnitude than ever before (Drinkwater et al., 2003; Poloczanska et al., 2013), there is a premium upon individual birds to explore, so that they can attain habitats that have become newly suitable due to changing climate. Our recent research (Veit, 2000; Zawadzki et al., 2019) clearly identifies a strong link between high rates of population growth and vagrancy, so that the largest numbers of vagrants accompany peak reproduction, as large numbers of young spread out in search of new territories. This is in direct contrast to previous ideas about vagrancy that characterize vagrancy as an act of desperation or navigational error.

Typically, birds disperse during the years prior to breeding, and the majority of long-distance vagrants are juveniles or at least pre-breeders. While species or populations of birds tend to orient in particular directions while dispersing, there are very large variations about the mean direction, resulting in many individual birds orienting in directions from that of the majority of the population. Similarly, probability distributions of distances dispersed by a population of birds are almost always strongly skewed, or “fat-tailed,” such that a small proportion of individuals within a population *normally* travel vastly farther than the mean distance. Thus, the suite of movement behaviors typical of birds predispose them toward vagrancy.

The papers with this Research Topic illustrate how vagrancy often results in colonization or recolonization of novel breeding sites. Furthermore, there is increasing



evidence that vagrancy is a positive response to improved conditions rather than a desperate move to escape catastrophe, and is thus carried out by individuals in top physical condition. This perspective on vagrancy contrasts with the traditional view of being a consequence of incompetent navigation. Combining these ideas, we suggest that, in order to colonize newly available habitats, new habitats that support elevated reproduction in response to abundant resources must exist to generate vagrants. To experience whether previously unoccupied areas are suitable for breeding, birds must first travel to these places, and this initial movement of discovery constitutes vagrancy by definition.

Growing populations produce more young birds, which consequently disperse in a variety of directions and distances. Thus, growing populations should produce vagrants, as predicted by Grinnell in 1922 to be “*the regular thing, to be expected.*” This concept has profound consequences for adjustment of species ranges as the world’s climate changes. For example, using as an example the recent range expansion of Tropical Kingbirds into North America and associated spread of vagrants north to Canada (Figure 1), demonstrates how range expansion is often accompanied by multiple instances of long-distance vagrancy.

Our Research Topic illustrates these concepts using examples of marine birds. Henry et al. show how exploratory behavior by albatrosses from growing populations in the central Pacific Ocean has preceded range expansion to distant islands in the highly productive California Current. This expansion led

to higher population growth rates in the new colonies. They further caution about the use of habitat modeling that does not consider conditions found in both the “destination” and well as the “original” range. In this sense, vagrancy can lead to novel distributions and changes in what Hutchinson (1957) termed, the “realized” portion of the “fundamental” niche. Zawadzki et al. demonstrate how population growth within the core portions of the range of Lesser Black-backed Gulls led to colonization of Iceland and Greenland, and substantial expansion of wintering range into North America. Acosta Alamo et al. take advantage of voluminous data on occurrence of vagrant North American Gulls in Europe and vagrant European Gulls in North America. This contrast allowed the authors to ask whether vagrancy was driven primarily by population growth of the species concerned rather than passive dispersal by prevailing winds. The results, though mixed, in general support the link between population growth within the core range and expansion to new environments. Veit et al. analyzed the spectacular population increase by Elegant Terns within their very limited range in the eastern Pacific and its association with increasing eastward vagrancy, culminating with successful colonization of western Europe. These papers taken together point to the biological importance of vagrancy and how it will likely be of increasing importance to species survival during intense climate change (Davis and Watson, 2018; Gill et al., 2019). Given the importance of range expansions or shifts accompanying climate change, future research should focus on examination of routes

followed by individuals, especially if those individual routes can be linked to local environmental conditions. We further need to ask whether physiological condition measures can be linked to distances traveled to address whether long distance dispersal is more likely driven by poor or good environmental conditions. Finally, our analyses to date suggest that growing bird populations are more likely to generate vagrants. If found to be generally true, this idea has important implications for the ability of species to find newly suitable habitat—or disperse through increasingly-fragmented habitat—as the climate, and perhaps other environmental characteristics, change.

Author contributions

RV conceived the idea and wrote the draft editorial. LM and RV constructed the graphic in **Figure 1**. LM, LZ, MA, and RH made modifications to the final version. RV, LM, LZ, MA, and RH shared in the writing and analyses. All authors contributed to the article and approved the submitted version.

Funding

This research was supported by NSF DEB 2049303 to RV and LM.

References

Baker, R. R. (1978). Demystifying vertebrate migration. *New Scientist*. 80, 526–528.

Davis, R. A., and Watson, D. M. (2018). Vagrants as vanguards of range shifts in a dynamic world. *Biol. Conserv.* 224, 238–241. doi: 10.1016/j.biocon.2018.06.006

Drinkwater, K. F., Belgrano, A., Borja, A., Conversi, A., Edwards, M., Greene, C. H., et al. (2003). The response of marine ecosystems to climate variability associated with the North Atlantic Oscillation. *Geophys. Monograph-Am. Geophys. Union.* 134, 211–234. doi: 10.1029/134GM10

eBird (2021). eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available online at: <http://www.ebird.org>. (accessed May 1, 2022).

Gill, J. A., Alves, J. A., and Gunnarsson, T. G. (2019). Mechanisms driving phenological and range change in migratory species. *Phil. Trans. R. Soc. B* 374, 20180047. doi: 10.1098/rstb.2018.0047

Grinnell, J. (1922). The role of the “accidental”. *The Auk.* 39, 373–380. doi: 10.2307/4073434

Hutchinson, G. E. (1957). Population studies - animal ecology and demography—concluding 806 remarks. *Cold Spring Harbor Symp. Quant. Biol.* 22, 415–427. doi: 10.1101/SQB.1957.022.01.039

Poloczanska, E. S., Brown, C. J., Sydeman, W. J., Kiessling, W., Schoeman, D. S., Moore, P. J., et al. (2013). Global imprint of climate change on marine life. *Nat. Clim. Change.* 3, 919–925. doi: 10.1038/nclimate1958

Pranty, B., Kratter, A. W., and Ponzo, V. (2016). Status and distribution in Florida of Tropical Kingbird (*Tyrannus melancholicus*) and Couch’s Kingbird (*Tyrannus couchii*). *Florida Field Natur.* 44, 83–105.

Veit, R. R. (2000). Vagrants as the expanding fringe of a growing population. *Auk* 117, 242–246. doi: 10.1093/auk/117.1.242

Veit, R. R. (2008). Why expect clusters of vagrants? *Bird Observer* 36, 213–217.

Zawadzki, L. C., Veit, R. R., and Manne, L. L., (2019). The influence of population growth and wind on vagrancy in a North American passerine. *Ardea* 107, 131–147. doi: 10.5253/arde.v107i2.a2

Acknowledgments

We thank Elise Huchard for helpful comments. We thank several reviewers for their helpful comments, Danny Bystrak at the USGS Bird Banding Lab for access to data, Jonas Knape and Jan Null for technical advice and Samantha Monier for graphics.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.