

Anthropogenic Change and the Process of Speciation

Murielle Ålund,^{1,17} Meredith Cenzer,^{2,17} Nicolas Bierne,³ Janette W. Boughman,⁴ José Cerca,⁵ Mattheau S. Comerford,⁶ Alessandro Culicchi,⁷ Brian Langerhans,⁸ S. Eryn McFarlane,^{9,10} Markus H. Möst,¹¹ Henry North,¹² Anna Qvarnström,¹³ Mark Ravinet,¹⁴ Richard Svanbäck,¹⁵ and Scott A. Taylor¹⁶

¹Department of Ecology and Genetics, Animal Ecology, Uppsala University, Uppsala 75236, Sweden

²Department of Ecology and Evolution, University of Chicago, Chicago, Illinois 60637, USA

³ISEM, Université de Montpellier, CNRS, IRD, Montpellier 34095, France

⁴Department of Integrative Biology, Michigan State University, East Lansing, Michigan 48824, USA

⁵CEES - Centre for Ecological and Evolutionary Synthesis, Department of Biosciences, University of Oslo, Oslo 0316, Norway

⁶Biology Department, UMass Boston, Boston, Massachusetts 02125, USA

⁷Department of Ecology and Genetics, Animal Ecology, Uppsala University, Uppsala 75236, Sweden

⁸Department of Biological Sciences, North Carolina State University, Raleigh, North Carolina 27695, USA

⁹Department of Botany, University of Wyoming, Laramie, Wyoming 82071, USA

¹⁰Department of Biology, York University, Toronto, Ontario M3J 1P3, Canada

¹¹Research Department for Limnology, University of Innsbruck, Innsbruck 6020, Austria

¹²Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, United Kingdom

¹³Department of Ecology and Genetics, Animal Ecology, Uppsala University, Uppsala 75236, Sweden

¹⁴School of Life Sciences, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom

¹⁵Department of Ecology and Genetics, Animal Ecology, Uppsala University, Uppsala 75236, Sweden

¹⁶Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Boulder, Colorado 80309, USA

Correspondence: murielle.alund@ebc.uu.se; mlcenzer@ucdavis.edu

Anthropogenic impacts on the environment alter speciation processes by affecting both geographical contexts and selection patterns on a worldwide scale. Here we review evidence of these effects. We find that human activities often generate spatial isolation between populations and thereby promote genetic divergence but also frequently cause sudden secondary contact and hybridization between diverging lineages. Human-caused environmental changes produce new ecological niches, altering selection in diverse ways that can drive diversification; but changes also often remove niches and cause extirpations. Human impacts that alter selection regimes are widespread and strong in magnitude, ranging from local changes in biotic and abiotic conditions to direct harvesting to global climate change.

¹⁷Co-first authors contributed equally to this work; all other authors listed alphabetically.

Editors: Catherine L. Peichel, Daniel I. Bolnick, Åke Brännström, Ulf Dieckmann, and Rebecca J. Safran
Additional Perspectives on Speciation available at www.cshperspectives.org

Copyright © 2023 Cold Spring Harbor Laboratory Press; all rights reserved
Advanced Online Article. Cite this article as *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a041455

M. Ålund et al.

Altered selection, and evolutionary responses to it, impacts early-stage divergence of lineages, but does not necessarily lead toward speciation and persistence of separate species. Altogether, humans both promote and hinder speciation, although new species would form very slowly relative to anthropogenic hybridization, which can be nearly instantaneous. Speculating about the future of speciation, we highlight two key conclusions: (1) Humans will have a large influence on extinction and “despeciation” dynamics in the short term and on early-stage lineage divergence, and thus potentially speciation in the longer term, and (2) long-term monitoring combined with easily dated anthropogenic changes will improve our understanding of the processes of speciation. We can use this knowledge to preserve and restore ecosystems in ways that promote (re-)diversification, increasing future opportunities of speciation and enhancing biodiversity.

Our current epoch is characterized by human domination of the global landscape, with impacts on the environment and climate increasing in scope and speed (Curtzen and Stoermer 2000; Canfield et al. 2010; Lewis and Maslin 2015). Many organisms on earth have had to contend with major human-induced perturbations for hundreds of thousands of years (Roebroeks and Villa 2011), but the magnitude, extent, and pace of these perturbations have accelerated at an alarming rate over the past ~200 years (Ellis and Ramankutty 2008; Lewis and Maslin 2015). Anthropogenic impacts on organisms have led to dramatic and widespread biodiversity loss (Barnosky et al. 2011). Human activities have, inadvertently or purposefully (e.g., Table 1), further altered geographic ranges, population dynamics, and selection regimes for a vast number of the persisting taxa (e.g., Ainouche et al. 2004; Antonovics 2006; Isaksson 2018; Kennedy et al. 2020). As a consequence of these effects on the evolutionary trajectories of many species, human influences are likely to have non-trivial effects on the process of speciation (Fig. 1).

Anthropogenic disturbances often involve environmental changes that can affect evolutionary processes associated with speciation, including both increases and decreases in RI between lineages. Geographic context and selection represent some of the most important factors that influence speciation (e.g., Coyne and Orr 2004). Human activities include global movement of organisms and modification of geographical barriers, as well as diverse changes in selection. We focus here on sexually reproducing species and RI, with a specific interest in anthropogenic im-

pacts on speciation processes (i.e., evolutionary increases and decreases in RI), rather than lineage persistence over longer timescales, which is extremely challenging to predict. Although the evolution of fully reproductively isolated species is expected to be slow, partial RI can evolve during human lifetimes in a variety of taxa (Table 1) and is arguably a more common stable end point than complete RI (Servedio and Hermisson 2019). Since both geographic isolation and episodes of secondary contact can be important contributors to speciation (e.g., Anderson and Weir 2022), it is important to understand the impact humans have on these processes.

Widespread anthropogenic impacts on evolutionary dynamics not only have the potential to alter speciation processes, but also provide a remarkable opportunity for the study of speciation. Modern anthropogenic global change is distinct from prior time periods in many ways that make it an exceptional context for uncovering how speciation processes occur, at least over relatively short timescales. More precisely, many anthropogenic activities are (1) repeated at various geographic, taxonomic, and temporal scales, meaning that the effects of these activities on speciation processes can be studied in a replicated manner, (2) strong and multifaceted, with disparate ecological changes of high magnitude often acting on diverging species simultaneously, and (3) occurring almost instantaneously in geological time and, importantly, can often be precisely dated. Human impacts on evolutionary processes are present everywhere and must therefore be considered when seeking to understand contemporary evolution and speciation. Just as the prior

**Table 1.** Examples of the impacts of different human activities on divergence and hybridization

Category	Driver	Effect on reproductive isolation	Estimated start date	Generation time	Type of barrier affected	Taxon	Citation
Physical environment	Habitat modification	Decreasing	1800s	2–5 yr	Premating	Gray wolves (<i>Canis lupus</i>) and coyotes (<i>Canis latrans</i>)	Lehman et al. 1991
	Habitat fragmentation	Increasing	1904–1914	2–13 yr	Premating (spatial)	Geoffrey's tamarin (<i>Saguinus geoffroyi</i>)	Díaz-Muñoz 2012
	Human-created structures	Increasing	1368	Annual, long-lived perennial	Premating	Grass (<i>Cleistogenes caespitosa</i> [Poaceae]); Siberian elm (<i>Ulmus pumila</i> [Ulmaceae]); blue knoll (<i>Heteropappus hispidus</i> [Asteraceae]); jujube (<i>Ziziphus jujuba</i> [Rhamnaceae]); Chinese chaste tree (<i>Vitex negundo</i> [Verbenaceae]); apricot (<i>Prunus armeniaca</i> [Rosaceae])	Su et al. 2003
		Increasing	1863–1939	2–4 wk	Premating (breeding site choice)	House mosquito (<i>Culex pipiens</i>)	Byrne and Nichols 1999; Haba and McBride 2022
	Water flow regime	Both	1800–1980	1–2 yr	Premating (spatial isolation + ecological isolation)	Freshwater sculpin (<i>Cottus</i>)	Nolte et al. 2005
Pollution	Metals/inorganics	Increasing	Mid- to late 1800s	Short-lived perennial	Premating (flowering time) + postmating (greater self-fertility)	Sweet vernal grass (<i>Anthoxanthum odoratum</i>)	Antonovics 2006
	Synthetic organic compounds	Decreasing	1988–1997	4–8 mo	Premating (chemical signaling)	Swordtail fishes (<i>Xiphophorus malinche</i> and <i>Xiphophorus birchmanni</i>)	Fisher et al. 2006
	Nutrients, suspended particles	Decreasing	1920s–1960s	~3 yr	Premating	Cichlids (Haplochromines, <i>Neochromis</i> , <i>Nyererei</i> complex, <i>Paralabidochromis</i>)	Seehausen et al. 1997

Anthropogenic Change and Speciation

Continued

**Table 1.** *Continued*

Category	Driver	Effect on reproductive isolation	Estimated start date	Generation time	Type of barrier affected	Taxon	Citation
Temperature	Climate change	Decreasing	1930s–1940s to 1971–1979	10–15 d	Unknown	Daphnia (<i>Daphnia galeata</i> and <i>Daphnia hyalina</i>)	Brede et al. 2009
		Decreasing	1995–2003	1 yr	Premating (spatial)	Southern and northern flying squirrel (<i>Glaucomys volans</i> and <i>Glaucomys sabrinus</i>)	Garroway et al. 2010
		Decreasing	1970–1982	1 yr	Premating	Red-breasted and red-naped sapsucker (<i>Sphyrapicus ruber</i> and <i>Sphyrapicus nuchalis</i>)	Billerman et al. 2019
Biotic interactions	Urban heat island	Increasing	~1900	5–15 yr	Premating (local adaptation)	Acorn ants (<i>Temnothorax curvispinosus</i>)	Diamond et al. 2017
	Altered predation/parasitism/pathogens	Decreasing	1997	1 yr	Premating (mate choice)	Small and medium tree finch (<i>Camarhynchus parvulus</i> and <i>Camarhynchus pauper</i>)	Kleindorfer et al. 2014
Species introductions, range expansions	Decreasing		1950s	1–3 yr	Premating/also postzygotic?	Benthic and limnetic threespine stickleback (<i>Gasterosteus aculeatus</i>)	McPhail 1992; Taylor et al. 2005; Gow et al. 2006
		Both	Mid-1900s	30–60 d	Premating (ecological isolation) + postmating (local adaptation)	Ancestral and derived ecotypes of red-shouldered soapberry bug (<i>Jadera haematoloma</i>)	Carroll and Boyd 1992; Carroll et al. 1998; Cenzer 2016
		Increasing	1690–1710	Short-lived perennial	Premating (ecological isolation) + postmating (sorting of intrinsic genetic incompatibilities)	Oxford ragwort (<i>Senecio squalidus</i>)	Nevado et al. 2020

Continued

**Table 1.** *Continued*

Category	Driver	Effect on reproductive isolation	Estimated start date	Generation time	Type of barrier affected	Taxon	Citation
Close association with humans	Agriculture	Decreasing	1970s	Perennial, flowers annually	Premating (spatial)	Smooth and California cordgrass (<i>Spartina alterniflora</i> and <i>Spartina foliosa</i>)	Ainouche et al. 2004
		Increasing	Late 1800s	Perennial, flowers annually	Postzygotic (hybrid sterility)	Smooth and small cordgrass (<i>S. alterniflora</i> and <i>S. maritima</i>)	Ainouche et al. 2004
		Increasing	Mid-1800s	1 yr	Premating (ecological isolation)	Apple maggot (<i>Rhagoletis pomonella</i>) apple and haw ecotypes	Filchak et al. 2000
	Food addition	Increasing	Early 20th century	1 yr	Premating (ecological + behavioral isolation)	House finch (<i>Carpodacus mexicanus</i>) house and desert ecotypes	Badyaev et al. 2008
	Domestication	Both	14–40 kya	~1–4 yr	Premating (spatial, behavioral, possibly mate choice)	Dogs and wolves (<i>Canis</i> sp.)	Bergström et al. 2022
		Both	~9 kya	~1 yr	Premating (spatial, behavioral, possibly mate choice)	Wild and domestic cats (<i>Felis</i> sp.)	Beaumont et al. 2001
		Both	~8 kya	~6 mo	Premating (spatial, behavioral, possibly mate choice)	Wild and Domestic/feral chicken (<i>Gallus</i> sp.)	Gering et al. 2017
		Both	~10 kya	~5–6 mo	Premating (spatial, behavioral, possibly mate choice)	Wild boar and domestic pigs (<i>Sus</i> sp.)	Ai et al. 2015
		Increasing	170–83 kya	1–21 d	Premating (ecological isolation), possibly postzygotic	Body and head louse (<i>Pediculus humanus corporis</i> and <i>Pediculus humanus capitis</i>)	Toups et al. 2011; for review, see Veracx and Raoult 2012

Continued

Anthropogenic Change and Speciation

**Table 1.** *Continued*

Category	Driver	Effect on reproductive isolation	Estimated start date	Generation time	Type of barrier affected	Taxon	Citation
		Increasing	11–6 kya	1 yr	Premating (spatial, behavioral, possibly mate choice)	House sparrow (<i>Passer domesticus</i>)	Ravinet et al. 2018
Genetic modification	Both	2002		Perennial	Possibly postzygotic	Creeping bentgrass (<i>Agrostis stolonifera</i>) and annual beard-grass (<i>Polypogon monspeliensis</i>)	Watrud et al. 2004; Zapiola and Mallory-Smith 2012

We broadly define reproductive isolation (RI) as increasing if studies showed higher genetic divergence or identified stronger isolating mechanisms in human-modified environments or time periods, and decreasing if gene flow or isolating mechanisms were reduced in human-modified environments or time periods. Estimated start date refers to the start of the specific human activity impacting that example (e.g., the construction of the Panama Canal occurred between 1904 and 1914, creating spatial isolation between populations of Geoffrey's tamarin). Generation times, when not provided in references, were drawn from the broader literature.




Figure 1. Schematic describing the general ways anthropogenic activities can influence the formation and loss of species by changing the geography, selection, and rates of hybridization between lineages. **Indicate indirect changes in selection due to changed geography or via hybridization. (RI) Reproductive isolation, (BDMIs) Bateson-Dobzhansky-Muller incompatibilities.

M. Ålund et al.

view of evolution as a slow, gradual process largely unobservable during human lifetimes has been dramatically transformed in the face of empirical evidence to the contrary, we similarly need to investigate speciation in the context of anthropogenic activities to better understand the modern pace of speciation.

In this perspective piece, we review the impact of anthropogenic change on geography and selection, two central factors in speciation processes. We highlight anthropogenic hybridization, which can act to dissolve previously distinct lineages but can also impose new patterns of selection on hybrids and parental species. We proceed with an outlook on the future of speciation, where we first ask whether speciation mechanisms may change in some way compared to their historical frequency, magnitude, or rate, and then, second, consider how we might take advantage of the ongoing occurrence of anthropogenic change to better understand processes of speciation. We argue future efforts should work toward preserving and restoring natural environments in a way that promotes diversification and future formation of new species. While most relevant studies on these topics to date have not focused on speciation per se, here we interpret prior results on the effect of human activities on evolution in the context of speciation. There is not (yet) a large literature on contemporary speciation. However, burgeoning studies on human-caused environmental change and contemporary evolution suggest that the processes of speciation should also be profoundly affected. Moreover, the recognition that RI and lineage dynamics (including extinction and hybridization) can occur on short timescales suggest we have much to gain by not ignoring modern speciation processes.

ALTERED GEOGRAPHY

Geography is a long-recognized, important feature of speciation that greatly influences the evolution of RI (Mayr 1963; Bush 1975; Coyne and Orr 2004; Bolnick and Fitzpatrick 2007; Nosil 2012). Geographical arrangement of organisms affects gene flow between populations and species, and the magnitude and frequency of gene

flow can critically influence several mechanisms involved in speciation. For instance, gene flow affects the probability and rate of the evolution of RI via genetic drift, similar selection, and divergent selection (Nosil and Flaxman 2011; Nosil 2012; Langerhans and Riesch 2013). Human activities have substantially altered geographical context (see examples in Table 1), and these effects are likely to continue or accelerate in the future. On the one hand, anthropogenic activities can create new barriers that separate previously continuous populations or species, thereby reducing gene flow (e.g., Su et al. 2003). On the other hand, humans increase connectivity by bringing incipient species together, increasing the likelihood of hybridization (van Hengstum et al. 2012). Globally, this can lead to shifts in evolutionary trajectories, affecting both speciation and extinction rates, as well as the risk of species merging together. In the latter process, distinct lineages can fuse together via hybridization—termed “despeciation” herein—resulting in the loss of biodiversity (Seehausen 2006; Vonlanthen et al. 2012).

There is broad evidence that human-constructed barriers and environmental change often increase population fragmentation in terrestrial and aquatic taxa (Templeton et al. 2001; Fuller et al. 2015; Dobbs et al. 2017; Johnson and Munshi-South 2017; Littleford-Colquhoun et al. 2017; Langerhans and Kern 2020). For example, common drivers of fragmentation involve roads, agricultural fields, power line easements, dams, and cities (Trombulak and Frissell 2000). Such fragmentation is widespread and has driven convergent signatures of reduced connectivity in many species by reducing movement between populations, therefore reducing opportunities for mating (e.g., Smith et al. 2009; Mathew et al. 2015; Benjamin et al. 2016; Lourenço et al. 2017; Miles et al. 2018). Reduced gene flow can facilitate the evolution of RI through the accumulation of neutral differences, divergent evolutionary responses to either similar or divergent selection, and even through one-allele mechanisms of RI such as the evolution of reduced migration. Among the oldest examples of human-induced fragmentation is the Great Wall of China, which has significantly reduced gene

Anthropogenic Change and Speciation

flow between plant populations since its construction began in 1368, apparently due to limiting both wind- and insect-dispersed pollen (Su et al. 2003). Over shorter timescales, intensive development and intervening roadways in the Los Angeles metropolitan area are associated with repeated patterns of reduced gene flow in birds, lizards, bobcats, and coyotes (Riley et al. 2006; Delaney et al. 2010).

Human barriers that split previously panmictic populations can rapidly result in population divergence in traits associated with RI, especially when fragmentation coincidentally results in strong environmental change, thereby inducing both reduced gene flow and divergent selection. For example, dams dramatically alter aquatic flow regimes and not only reduce connectivity, but often result in rapid phenotypic shifts in fishes (Haas et al. 2010; Franssen 2011; Franssen et al. 2013). Road construction across tidal creeks in The Bahamas not only fragmented populations of three Bahamian mosquitofish species (*Gambusia* spp.), but also led to diverse trait changes over ~35–50 years, including genital and body morphology, male coloration, and brain morphology (Heinen-Kay et al. 2014; Giery et al. 2015; Riesch et al. 2015; Jenkins et al. 2021). Human-induced fragmentation can also tip the balance between selection and gene flow, especially for species with limited dispersal. For instance, the construction of a parking lot dividing previously connected populations of walking stick insects (*Timema cristinae*) restricted gene flow between populations using two different host plants. This reduction in gene flow was apparently sufficient for adaptive divergence of genetically based cryptic coloration to emerge on the two hosts in <6 yr (Nosil 2009). Thus, human-caused fragmentation can play an important role in the earliest steps of speciation—we still have much to learn about the potential persistence of newly divergent lineages spawned by such anthropogenic impacts.

Humans also regularly transport organisms into new areas in which they are nonnative (Bullock et al. 2018). This can reduce intraspecific gene flow between populations in their native and introduced ranges and cause fast evolutionary changes in introduced populations, especially

when the new distribution differs in ecology or climate from their native range (Mooney and Cleland 2001; Prentis et al. 2008; Whitney and Gabler 2008; Buswell et al. 2011). For example, the common reed (*Phragmites australis*) diverged between its native European and introduced North American range over the last 150 years as a result of both geographic distance and environmental differences (Guo et al. 2018). Salmon introduced to Lake Washington evolved RI after fewer than 13 generations (Hendry et al. 2000). Selection likely plays a major role in most cases of rapid divergence related to species introductions (see below), but altered geography provides a key context for the divergence.

Other anthropogenic infrastructure and activity, such as channels, bridges, and maritime traffic, instead create artificial corridors of dispersal (Crispo et al. 2011; Miles et al. 2018, 2019) that can increase gene flow between populations and species. For instance, human-driven maritime translocation is credited for genetically localized introgression tracts detected between Pacific and Atlantic species of *Ciona* tunicates, which now co-occur along the heavily shipped corridors of the English Channel (Le Moan et al. 2021). This type of consistent, human-aided gene flow increases the success and impacts of invasive species both through creating disturbed ecosystems conducive to invasion, as in the fire ant *Solenopsis invicta* (Resasco et al. 2014), and through allowing the spread of adaptive alleles between invading populations, as in Asian long-horned beetles (Javal et al. 2019). Human-caused corridors, accidental or purposeful relocation, and climate-change induced shifts in geographical range can additionally bring species into contact with closely related species, sister taxa, or incipient lineages. This could lead to hybridization and reinforcement of RI through selection against interpopulation/interspecific mating (Butlin and Smadja 2018), but high rates of hybridization can also reduce RI by causing selection against incompatibilities and despeciation (see below; Thompson et al. 2023).

Perhaps the most common impact of human activity is the reduction of both population size and genetic diversity as a result of habitat loss (e.g., Lu et al. 2001; Wei et al. 2015), fragmenta-



M. Ålund et al.

tion (e.g., van der Valk et al. 2019), and/or degradation (e.g., Wisely et al. 2008; Sievers et al. 2018). Reduced population size not only increases the probability of extinction, but also decreases the efficacy of evolutionary responses to selection by reducing available genetic variation (Charlesworth 2009), which may be particularly critical in the context of adapting to, and persisting under, rapidly changing conditions (see next section). A small population size may, however, increase the probability of quick divergence and evolution of RI in the case of strong selection for local adaptation (Gavrilets 2003). Furthermore, population size asymmetry between species at secondary contact greatly affects the probability and dynamics of hybridization (Qvarnström et al. 2023).

To predict the lasting impacts of altered geographic context on speciation, we need to understand the persistence of both human-caused environmental changes and diverging lineages. Humans have a net negative influence on species number, as the rates of anthropogenic despeciation and extinctions outpace the rate at which new species may be formed. It is nevertheless also relevant to understand when and how humans interfere with processes of speciation over short and long timescales. Understanding how altered geographic contexts affect even transient divergence between populations or processes operating within incipient species in real time can prove quite consequential, as these can have long-lasting effects on both ongoing and future speciation processes. This may be especially important if anthropogenic activity increases the frequency of early-stage divergence followed by extinction.

ALTERED SELECTION

Most speciation events probably involve selection, and RI typically evolves faster in response to selection than other mechanisms (Schlüter 2009). Human activities that modify biotic and abiotic features are now well known for altering selection and adaptation (Brady et al. 2009; Daramont et al. 2009; Murúa et al. 2010; Hendry et al. 2017; Thompson et al. 2019). Anthropogenic drivers of selection are numerous and vary in

magnitude, geographic scale, and novelty (Isaksson 2018; Price 2022); we need only look around us to see the footprint of direct selection (e.g., domestication; Larson and Fuller 2014), the built environment (Alberti et al. 2017; Szulkin et al. 2020), pollution (Loria et al. 2019), altered ecological communities (Strauss et al. 2006), and climate change (Franks and Hoffmann 2012) to name a few. Altogether, these varied impacts on selection that characterize human activities may generate ecological opportunity for resilient taxa that persist in human-altered environments (Wellborn and Langerhans 2015; Isaksson 2018; Ravinet et al. 2018; Campbell-Staton 2020).

How do human-induced changes to selective landscapes affect speciation processes? Selection can influence the evolution of RI through three major mechanisms, and all are affected by human activities: divergent selection (ecological speciation), similar selection (mutation-order speciation), and reinforcement (selection against interpopulation mating after secondary contact) (Langerhans and Riesch 2013). In each scenario, which are not mutually exclusive, human-caused selection can alter traits that influence RI. Anthropogenic sources of divergent selection have received the most attention to date, while few studies have examined similar selection or reinforcement in this context. This is unfortunate, as these mechanisms may commonly occur, and the strongest evidence from the wild that responses to selection actually influence RI come from cases of reinforcement selection (because selection directly favors RI).

Many traits evolving in response to human activity have the potential to directly impact RI, like the timing and duration of reproduction (Alberti et al. 2017; Sirkiä et al. 2018), song frequency, timing, and composition (e.g., Reichard et al. 2020; Kunc and Schmidt 2021), mate-choice modality (de Jong et al. 2018), genital morphology (Heinen-Kay et al. 2014), and male color signals (Giery et al. 2015; Koneru and Caro 2022). Impacts of climate change on sexual selection are also likely to affect premating and postmating prezygotic and postzygotic reproductive barriers (Pilakouta and Ålund 2021; Garlovsky et al. 2023). Of course, adaptation

Anthropogenic Change and Speciation

does not imply speciation, and responses to selection can also reduce RI (e.g., Owens and Samuk 2020). Evidence is accumulating that anthropogenic impacts can indeed influence RI in both positive and negative manners, often involving selection (Table 1). With the growing evidence for human-induced selection and contemporary adaptation, it would be surprising if anthropogenic selection did not influence speciation. But currently less clear is how exactly this occurs, whether promoting or hindering RI is more common, where we should focus our efforts to understand it, and how to assess the potential for lineage persistence after early stages of divergence.

Anthropogenic activities often generate divergent selection, with anthropogenic environments representing novel niches far removed from ancestral environments (e.g., Isaksson 2018; Szulkin et al. 2020, Santangelo et al. 2022). When novel divergent selection pressures are paired with the movement of genes between formerly isolated areas, this can contribute to the rapid appearance of novel habitat specialists. For example, the common house mosquito (*Culex pipiens*) colonized the London Underground transit system about 100 years ago, leading to changes in breeding environment that, in turn, contributed to RI between surface- and subway-dwelling populations (Byrne and Nichols 1999). However, these lineages did not arise *in situ*. Instead, previously existing genotypes, likely originating from human-adapted *Culex pipiens molestus* in the Middle East, underwent spatial reorganization and have since colonized multiple subway systems (Haba and McBride 2022).

One striking example of a human-induced divergence event is that of a derived ecotype of the hawthorn fruit fly, which emerged along the Eastern coast of North America within the last 200 years (Walsh 1867). Here the anthropogenic introduction and proliferation of domesticated apples facilitated a host shift from their native host, hawthorne (Bush 1992). Because hawthorne and apple differ in the timing of their fruiting, allochronic isolation between sympatric populations was possible; and, much like subway mosquitos, the genes underlying allochronic RI already existed in another part of

the *Rhagoletis* range in Mexico (Feder et al. 2003). The movement of these genes northward, paired with multiple divergent selection pressures and larval host fidelity, contributed to both olfactory preference for the derived host and genetic differentiation between ecotypes (Feder et al. 1988, 1994; Linn et al. 2003; Egan et al. 2015). This host plant shift induced by human-mediated species introductions triggered local population divergence in the hawthorne flies and the emergence of a possible new lineage—the apple maggot fly (Tait et al. 2016).

Many anthropogenic impacts that are replicated across the globe can influence speciation by selection mechanisms in ways that partially depend on geography. If impacted areas are geographically isolated (whether naturally or through human modification), divergent selection between impacted and unimpacted populations could lead to parallel divergence and contribute to the evolution of RI. For instance, urban environments have been shown to act as parallel niche-space for the cosmopolitan legume *Trifolium repens*, wherein city ecosystems tend to share more similarities to one another than with the adjacent surrounding environment regardless of geographic location (Santangelo et al. 2022). If gene flow is facilitated between impacted areas, it could lead to only one primary divergence event, such as in cosmopolitan or commensal species. Moreover, if anthropogenic impacts induce similar selection across multiple geographically isolated populations, RI could evolve between independent cases of the same anthropogenic driver via adaptive responses to similar selection. That is, different populations may evolve different traits in response to the same selection regime (i.e., mutation-order speciation), and consequently result in elevated RI between populations, such as cases of intragenomic conflict (e.g., cytoplasmic male sterility, meiotic drive), sexual conflict, sexual (or social) selection for arbitrary traits, and alternative adaptive solutions to selection via competition, predation, parasitism, etc. (Price 2008; Schlüter 2009; Martin and Mendelson 2012; Langerhans and Riesch 2013). We currently know little about how these processes may unfold in an anthropogenic context.



M. Ålund et al.

Some anthropogenic effects are ubiquitous and homogenize environments, making niches less distinct. In the case of anthropogenic climate change, homogenizing effects on abiotic conditions have been demonstrated in coastal estuaries (Ferrarin et al. 2014), high arctic tundra (Stewart et al. 2018), coral reefs (Richardson et al. 2018), and others. These cases can reduce the likelihood of divergent selection between environments, but adaptation to these homogeneous environments could still increase RI. This could occur if natural geographic isolation among populations remains unchanged, or if human actions increase geographic isolation, as responses to similar selection could promote RI among different cases of these similar environments. However, if gene flow is moderate, or especially if human activities increase gene flow, there would be little chance for these mutation-order processes as adaptive alleles could easily spread throughout populations and prevent the accumulation of RI. Perhaps adaptation in the absence of RI is more likely in these scenarios, but more research is needed.

ALTERED HYBRIDIZATION

Hybridization can dissolve distinct evolutionary lineages (despeciation), but can also promote diversification through reinforcement and generation of new hybrid lineages (Mallet 2007; Arnold and Kunte 2017; Ottenburghs 2018; McFarlane and Pemberton 2019). The frequency of interspecific hybridization can increase due to the breakdown of geographic and ecological barriers, both heavily impacted by human activities (reviewed in Grabenstein and Taylor 2018; Grabenstein et al. 2023). These two processes likely often interact, for example, when species are transported to a new locality and when environmental change *in situ* alters hybridization frequency (Guo 2014; McFarlane and Pemberton 2019).

First, globalization has led to a drastic increase in the frequency and magnitude of direct species introductions via activities including horticulture, agriculture, aquaculture, shipping, and the pet trade (Lewis and Maslin 2015; Price 2022). Among birds, the extensive introduction

of the mallard duck (*Anas platyrhynchos*) to multiple locations worldwide led to the establishment of several hybrid populations with a variety of other duck species (Rhymer et al. 1994; Muñoz-Fuentes et al. 2007; Čížková et al. 2012; Ford et al. 2017; Lawson et al. 2021). In the sea, the Mediterranean mussel *Mytilus galloprovincialis* has been introduced worldwide by maritime traffic and aquaculture and formed anthropogenic admixture zones with its sister species *Mytilus trossulus* in California (Saarman and Pogson 2015) and Japan (Brannock et al. 2009), *Mytilus planulatus* in Australia (Popovic et al. 2020), *Mytilus edulis* in the English Channel (Simon et al. 2020), and *Mytilus platensis* in Argentina (Zbawicka et al. 2018).

Second, human-aided range expansions (e.g., via climate change) can also result in the secondary contact of previously isolated species, involving a simultaneous breakdown of geographic and ecological barriers (Lehman et al. 1991; Chunco 2014; reviewed in Taylor et al. 2014). For example, the range of southern flying squirrels (*Glaucomys volans*) has recently expanded northward into the range of northern flying squirrels (*Glaucomys sabrinus*), leading to hybridization (Garroway et al. 2010, 2011) and in some cases competitive exclusion (Wood et al. 2016). While ecological overlap between these two species in their new ranges is not entirely clear (O'Brien et al. 2021), the rapid movement of southern flying squirrels could potentially introduce novel parasites or diseases to the northern flying squirrel via hybridization, leading to population level threats in addition to competition or hybridization (Krichbaum et al. 2010; but see O'Brien et al. 2022).

Third, human-induced environmental change can alter the probability of hybridization and the fitness of hybrids by disrupting preexisting selective regimes (e.g., through eutrophication: Seehausen et al. 1997; Vonlanthen et al. 2012; Dokulil 2014; see also Box 1 for a detailed example) or by creating novel, often very strong selective regimes favoring hybridization and/or adaptive introgression (e.g., in the case of pesticide or pollutant resistance: Oziolor et al. 2019; Le Corre et al. 2020; Valencia-Montoya et al. 2020). A special case of secondary contact is

BOX 1. PATTERNS AND CONSEQUENCES OF HYBRIDIZATION DUE TO ANTHROPOGENIC CHANGE IN WHITEFISH-ADAPTIVE RADIATION IN PERI-ALPINE LAKES

Anthropogenic ecological change can result in a breakdown of selection regimes that maintain ecological and genetic divergence and, thus, cause increased gene flow and a reduction in divergence and diversity. One example for this process are the consequences of eutrophication on adaptive radiations of *Coregonus* whitefish in peri-Alpine lakes.

Since the last glaciation, whitefish have colonized peri-Alpine lakes and undergone replicated adaptive radiations into ecologically and genetically differentiated species over the last 10,000–15,000 years. The >30 species can be grouped into six different ecomorph classes across all lakes that are differentiated in morphology, ecology, and breeding behavior. This includes large-bodied benthic invertebrate feeders with few widely spaced gill-rakers, which spawn in summer in shallow littoral habitats; small-bodied zooplankton feeders with many, densely spaced gill-rakers, which spawn in deeper areas in summer or winter; and, in some lakes, profundal summer-spawning species with few gill-rakers (e.g., in Lake Thun and the now extinct Kilch in Lake Constance). The various ecomorphs have evolved independently in the different lake systems and their parallel evolution is explained by a polygenic adaptive architecture, with at least one major effect loci (the *edar* gene) explaining variation in gill-raker number, with secondary contact and allele sharing among flocks (De-Kayne et al. 2022).

Whitefish diversity declined precipitously with severe eutrophication of Swiss peri-alpine lakes, in part because eutrophication homogenized niche space and undermined the diversification process (Vonlanthen et al. 2012). Environmental gradients and thus both ecological and reproductive niche diversity within lakes were eroded by the ecological alterations from eutrophication. Consequently, premating isolation and divergent selection that formerly maintained high levels of extrinsic postzygotic RI among whitefish species were significantly weakened, leading to increased hybridization among sympatric species and despeciation (Vonlanthen et al. 2012). Eutrophication also caused anoxic conditions in deep water, leading to recruitment failure and subsequent loss of deep-spawning species (Vonlanthen et al. 2012; Frei et al. 2022, 2023). Diversity loss included losses of individual species, as well as lessened functional diversity and genetic differentiation of remaining whitefish (Vonlanthen et al. 2012).

But where humans have caused drastic ecological changes they can work to alleviate them. Thus, concerted efforts were undertaken to remediate pollution in these lakes through water treatment and reduced phosphorus inputs, which facilitated rapid recovery of many lakes to their former oligotrophic state (Smith and Schindler 2009). Following this ecological restoration, functional diversity (variation in gill-raker number) in some whitefish taxa was rapidly reinstated (Jacobs et al. 2019). Under eutrophication, hybridization included extensive introgression from species that became extinct and data suggest substantial introgression of putatively adaptive alleles that underpinned the initial diversification of whitefish in these lakes (Jacobs et al. 2019; Frei et al. 2022), likely facilitating rediversification and recolonization of the now restored deep-water habitats. These patterns highlight both the fragility of extrinsic RI when environments change, and also that hybridization may have helped to preserve variation now fueling the recovery of diversity after lakes were restored (Frei et al. 2023). However, while some functional diversity has been restored, species diversity has not recovered and deep-water species are still absent, indicating that there are constraints on what can be restored after severe disturbance.

This example of the consequences for eutrophication on peri-Alpine whitefish shows the drastic consequences of human activities on RI and functional diversity. However, it also highlights the importance of swift action and implementation of remediation measures that can restore ecological gradients and allow for at least some degree of rediversification using genetic variation that may still be present.



M. Ålund et al.

the feralization of domesticated species in which humans are directly involved in both the evolution of the domesticated lineage and in its secondary contact with a nondomesticated lineage. Examples of hybridization between wild and feral species include Scottish wild and domestic cats (Senn et al. 2019), wolf and dog (Pilot et al. 2018), chicken and junglefowl (Gering et al. 2015), pig and boar (Mary et al. 2022), and maize and teosinte (Le Corre et al. 2020).

Hybridization through anthropogenic secondary contact can, under some circumstances, generate biodiversity. A classic example of homoploid hybrid speciation is the Oxford ragwort (*Senecio squalidus*), which is derived and reproductively isolated from two parental species native to Italy that were cultivated and incidentally crossed at the Oxford botanic garden in the late seventeenth century (Nevado et al. 2020). The subsequent spread of *S. squalidus* throughout much of Great Britain was aided by the development of rail and road networks, whose verges appear to mimic the lava rock in the range of the parental species (Harris 2002). Thus, human activity is directly implicated in both the formation of a species and the creation of habitat necessary for that species to establish and spread. Similarly, the human-aided introduction of the house sparrow (*Passer domesticus*) into the Mediterranean region resulted in secondary contact and hybridization with the Spanish sparrow *Passer hispaniolensis*, leading to the formation of a reproductively isolated homoploid hybrid commensal species—the Italian sparrow (*Passer italiae*) (Elgvin et al. 2017; Runemark et al. 2018). A third example is seen in dock mussels—a distinct ecotype of *Mytilus* associated with port habitats that have arisen through human-mediated hybridization between *M. edulis* and *M. galloprovincialis* and colonized multiple ports in parallel, and now constitutes a stable admixed population that can hybridize with the local parental species (Simon et al. 2020).

In summary, one consequence of human-induced breakdowns in geographic and ecological barriers is an increase in the opportunity and frequency of hybridization, especially across very long distances, in many places and across multiple species within a food web and/or ecosystem at the same time. The outcomes of an-

thropogenic hybridization are more likely to be negative than positive. While adaptive introgression (Hamilton and Miller 2016), reinforcement of preexisting barriers, and hybrid speciation (see Peñalba et al. 2023; Zhang et al. 2023) are possible outcomes, many documented cases of anthropogenic hybridization have resulted in a breakdown of RI between species (Grabenstein and Taylor 2018). It is thus very unlikely for anthropogenic hybridization to result in a positive outcome for global biodiversity.

ALTERED SPECIES TURNOVER RATES

Are we changing the predominant mechanisms leading to speciation? Geographic isolation is well documented as an important, and perhaps even necessary, step for the evolution of complete RI (Anderson and Weir 2022). As we have seen above, with increasing globalization, humans are causing geographic barriers to collapse, but also causing population subdivisions via habitat fragmentation. We are directly or indirectly altering selection regimes, with consequences for evolutionary processes known to play important roles in speciation. Anthropogenic activities are likely also increasing the speed and frequency of both early divergence between populations (i.e., potentially leading to speciation) and the collapse of recently diverged lineages (i.e., “despeciation”). Since anthropogenic extinction is orders of magnitude faster and more taxonomically wide-reaching than anthropogenic speciation, and because fewer species in turn lowers chances to speciate, the global diversification rate will likely decline in the short term. Our impact on processes of speciation will inevitably vary across taxa, with habitat fragmentation having a strong impact locally, and global movement of species acting on a totally different scale. The consequences will thus depend on a species’ dispersal ability. Are we likely to witness an increase in populations/species found in a transient stage on the speciation continuum (e.g., showing partial RI; Servedio and Hermisson 2019), compared to what was observed before human disturbances? Are we introducing new biases in patterns of diversification across taxa? It is difficult to predict which, if any, of the currently diverging lineages will result in



Anthropogenic Change and Speciation

fully isolated species. However, incipient species showing some degree of RI do represent distinct lineages that may increase functional biodiversity, and can thus play an important role in ecosystem functioning.

Can our knowledge of past events of mass extinction and current data on the resilience of different taxa to global climate change allow us to predict how the world may look after us? On a much longer scale (millions of years), and somewhat paradoxically, the ongoing extinction crisis is likely to cause an increased speciation rate in the future. Based on evidence from the fossil record, elevated extinction rates generally trigger acceleration of speciation rates when ecological communities reestablish (Harries et al. 1996; Hallam and Wignall 1997; Foote 2023). For example, marine microfossils appear to flourish after large extinction events (Keller and Pardo 2004). While the extinction of many large vertebrates (megafauna) is a well-known example of human impact, other less visible taxa are also greatly affected by anthropogenic disturbance, such as insect habitat specialists (Wagner 2021). This is an exciting avenue for future studies.

Recent examples of ecosystem diversification (see Box 1) have shown that taking rapid action to restore habitats has a positive effect on diversity. While we will likely lose much of the biodiversity present before human activities, some of the adaptive genetic and phenotypic variation that allowed for initial diversification can be retained, thereby potentially allowing population persistence and ultimately, the formation of new taxa (Ottenburghs 2021). At the core of the diversification process is divergent selection arising from complex niche space, to which organisms adapt and diverge from one another. Therefore, conservation efforts may need to focus on preserving features that promote the diversification process per se, potentially allowing new species to emerge in the future.

LONG-TERM STUDIES OF ANTHROPOGENIC CHANGE AS A TOOL TO UNDERSTAND SPECIATION

The multitude of ways that human activities alter geographic context, create new sources of selec-

tion, and induce secondary contact, combined with the fact that many such events are replicated and can be precisely dated, constitute some very exciting opportunities for researchers to study speciation. Many cases of early divergence have been identified by studying organisms that have colonized human-modified environments in parallel with their “wilder” counterparts, which allowed identifying phenotypic and genomic differentiation in these new environments (Filchak et al. 2000; Fonseca et al. 2004; Wright et al. 2013). In some cases, high levels of RI have been documented. Observing the emergence of new species is rarely possible, but documenting past collapse of species or partially diverged lineages may be relatively easy in contemporary systems (Otto 2018), provided that known young species or locally adapted population pairs are followed over time (Rosenblum et al. 2012). Famous documented examples of such collapses are only possible because of long-term studies and include threespine sticklebacks (Taylor et al. 2006), cichlids (Seehausen et al. 1997), and soapberry bugs (Carroll and Boyd 1992; Carroll et al. 1998; Cenzer 2016), where human-induced changes in environmental conditions caused the rapid collapse of reproductive barriers. In one case, both the collapse and re-emergence of divergent whitefish species were observed over a few decades in Alpine lakes first affected by eutrophication, but then cleaned in hopes of reversing the consequences of this anthropogenic disturbance (see Box 1), thanks to long-term monitoring of these lakes.

Hybrid zones are often referred to as “natural laboratories” (Barton and Hewitt 1985; Hewitt 1988) and anthropogenic hybrid zones provide a particular powerful framework to study speciation (Grabenstein and Taylor 2018). Anthropogenic hybrid zones offer well controlled, natural experiments as they (1) are often replicated in space (e.g., Simon et al. 2020; Westram et al. 2021), (2) have firm dates for secondary contact from written records (e.g., Calfee et al. 2020; McFarlane et al. 2020), and (3) may have precise estimates of the number of introduced individuals (e.g., Ratcliffe 1987; reviewed in Grabenstein and Taylor 2018). Additionally, because anthropogenic hybrid zones are necessarily new, there is the opportunity to understand the initial break-

M. Ålund et al.

down or buildup of RI at secondary contact. This is often not possible in older hybrid zones where we study the separate question of how RI is maintained. Anthropogenic hybrid zones, particularly when studied over many decades, are thus a powerful opportunity to test theoretical predictions about speciation processes directly in the wild (Anderson and Weir 2022).

CONCLUSION

We have described here that the ubiquity of human impacts, affecting both geography and selection, often in parallel and very rapid ways, may change the predominant mechanisms of speciation. We may be approaching an era where it is not possible to study evolution without human disturbance. The best documented examples of human-induced early lineage divergence and species collapse were detected thanks to long-term monitoring. We would like to emphasize the power of studying natural systems over several decades or even centuries, and the need for more long-term thinking in our research field. This allows not only the detection and quantification of our impact on the environment, but also the identification of ways to remediate these impacts. By preserving and restoring ecosystems in ways that promote (re-)diversification, instead of focusing on the conservation of single species, we have the power to increase future opportunities of speciation, and ultimately enhance biodiversity.

AUTHOR CONTRIBUTIONS

All authors contributed to the conceptualization of the paper, writing of the original draft, and reviewing and editing of the final version. M.Å. and M.C. led and administered the project.

ACKNOWLEDGMENTS

The authors were supported by the following funding sources during the course of this project: Swedish research council for sustainable development (FORMAS) grant number 2018-01563 to M.Å. and A.Q. and 2018-01591 to R.S. and A.C., Swedish Research Council (VR) grant number 2020-04631 to M.Å. and A.Q.,

National Science Foundation IOS 1928891 to S.A.T., National Science Foundation (DEB) 1638778 to J.W.B., and the Modelscape project from the National Science Foundation grant number 2019528 supported S.E.M.

REFERENCES

*Reference is also in this subject collection.

Ai H, Fang X, Yang B, Huang Z, Chen H, Mao L, Zhang F, Zhang L, Cui L, He W, et al. 2015. Adaptation and possible ancient interspecies introgression in pigs identified by whole-genome sequencing. *Nat Genet* **47**: 217–225. doi:10.1038/ng.3199

Ainouche ML, Baumel A, Salmon A, Yannic G. 2004. Hybridization, polyploidy and speciation in *Spartina* (Poaceae). *New Phytologist* **161**: 165–172. doi:10.1046/j.1469-8137.2003.00926.x

Alberti M, Correa C, Marzluff JM, Henry AP, Palkovacs EP, Gotanda KM, Hunt VM, Apgar TM, Zhou Y. 2017. Global urban signatures of phenotypic change in animal and plant populations. *Proc Natl Acad Sci* **114**: 8951–8956. doi:10.1073/pnas.1606034114

Anderson SAS, Weir JT. 2022. The role of divergent ecological adaptation during allopatric speciation in vertebrates. *Science* **378**: 1214–1218. doi:10.1126/science.abe7719

Antonovics J. 2006. Evolution in closely adjacent plant populations X: long-term persistence of prereproductive isolation at a mine boundary. *Heredity (Edinb)* **97**: 33–37. doi:10.1038/sj.hdy.6800835

Arnold ML, Kunte K. 2017. Adaptive genetic exchange: a tangled history of admixture and evolutionary innovation. *Trends Ecol Evol* **32**: 601–611. doi:10.1016/j.tree.2017.05.007

Badyaev AV, Young RL, Oh KP, Addison C. 2008. Evolution on a local scale: developmental, functional and genetic bases of divergence in bill form and associated changes in song structure between adjacent habitats. *Evolution (NY)* **62**: 1951–1964. doi:10.1111/j.1558-5646.2008.00428.x

Barnosky AD, Matzke N, Tomaia S, Wogan GO, Swartz B, Quental TB, Marshall C, McGuire JL, Lindsey EL, Maguire KC, et al. 2011. Has the earth's sixth mass extinction already arrived? *Nature* **471**: 51–57. doi:10.1038/nature09678

Barton NH, Hewitt GM. 1985. Analysis of hybrid zones. *Annu Rev Ecol Syst* **16**: 113–148. doi:10.1146/annurev.es.16.110185.000553

Beaumont M, Barratt EM, Gottelli D, Kitchener AC, Daniels MJ, Pritchard JK, Bruford MW. 2001. Genetic diversity and introgression in the Scottish wildcat. *Mol Ecol* **10**: 319–336. doi:10.1046/j.1365-294x.2001.01196.x

Benjamin A, May B, O'Brien J, Finger AJ. 2016. Conservation genetics of an urban desert fish, the Arroyo chub. *Trans Am Fish Soc* **145**: 277–286. doi:10.1080/00028487.2015.1121925

Bergström A, Stanton DWG, Taron UH, Frantz L, Sinding MS, Ersmark E, Pfrengle S, Cassatt-Johnstone M, Lebrasseur O, Girdland-Flink L, et al. 2022. Grey wolf genomic history reveals a dual ancestry of dogs. *Nature* **607**: 313–320. doi:10.1038/s41586-022-04824-9



Anthropogenic Change and Speciation

Billerman SM, Murphy MA, Carling MD. 2016. Changing climate mediates sapsucker (Aves: *Sphyrapicus*) hybrid zone movement. *Ecol Evol* **6**: 7976–7990. doi:10.1002/ece3.2507

Bolnick DI, Fitzpatrick BM. 2007. Sympatric speciation: models and empirical evidence. *Annu Rev Ecol Evol Syst* **38**: 459–487. doi:10.1146/annurev.ecolsys.38.091206.095804

Brady S. 2009. Eco-evolutionary consequences of road adjacency and road salt on the wood frog, *Rana sylvatica*. *Nat Prec* doi:10.1038/npre.2009.3674.1

Brannock PM, WetHEY DS, Hilbush TJ. 2009. Extensive hybridization with minimal introgression in *Mytilus gallo-provincialis* and *M. trossulus* in Hokkaido, Japan. *Mar Ecol Prog Ser* **383**: 161–171. doi:10.3354/meps07995

Brede N, Sandrock C, Straile D, Spaak P, Jankowski T, Streit B, Schwenk K. 2009. The impact of human-made ecological changes on the genetic architecture of *Daphnia* species. *Proc Natl Acad Sci* **106**: 4758–4763. doi:10.1073/pnas.0807187106

Bullock JM, Bonte D, Pufal G, da Silva Carvalho C, Chapman DS, García C, García D, Matthysen E, Delgado MM. 2018. Human-mediated dispersal and the rewiring of spatial networks. *Trends Ecol Evol* **33**: 958–970. doi:10.1016/j.tree.2018.09.008

Bush GL. 1975. Modes of animal speciation. *Annu Rev Ecol Syst* **6**: 339–364. doi:10.1146/annurev.es.06.110175.002011

Bush G. 1992. Host race formation and sympatric speciation in *Rhagoletis* fruit flies (Diptera: Tephritidae). *Psyche (Stuttg)* **99**: 335–357. doi:10.1155/1992/67676

Buswell JM, Moles AT, Hartley S. 2011. Is rapid evolution common in introduced plant species? *J Ecol* **99**: 214–224. doi:10.1111/j.1365-2745.2010.01759.x

Butlin RK, Smadja CM. 2018. Coupling, reinforcement, and speciation. *Am Nat* **191**: 155–172. doi:10.1086/695136

Byrne K, Nichols R. 1999. *Culex pipiens* in London Underground tunnels: differentiation between surface and subterranean populations. *Heredity (Edinb)* **82**: 7–15. doi:10.1038/sj.hdy.6884120

Calfee E, Agra MN, Palacio MA, Ramírez SR, Coop G. 2020. Selection and hybridization shaped the rapid spread of African honey bee ancestry in the Americas. *PLoS Genet* **16**: e1009038. doi:10.1371/journal.pgen.1009038

Campbell-Staton SC, Winchell KM, Rochette NM, Fredette J, Maayan I, Schweier RM, Catchen J. 2020. Parallel selection on thermal physiology facilitates repeated adaptation of city lizards to urban heat islands. *Nat Ecol Evol* **4**: 652–658. doi:10.1038/s41559-020-1131-8

Canfield DE, Glazer AN, Falkowski PG. 2010. The evolution and future of Earth's nitrogen cycle. *Science* **330**: 192–196. doi:10.1126/science.1186120

Carroll SP, Boyd C. 1992. Host race radiation in the soapberry bug: natural history with the history. *Evolution (NY)* **46**: 1052–1069. doi:10.1111/j.1558-5646.1992.tb00619.x

Carroll SP, Klassen SP, Dingle H. 1998. Rapidly evolving adaptations to host ecology and nutrition in the soapberry bug. *Evol Ecol* **12**: 955–968. doi:10.1023/A:1006568206413

Cenzer ML. 2016. Adaptation to an invasive host is driving the loss of a native ecotype. *Evolution (NY)* **70**: 2296–2307. doi:10.1111/evo.13023

Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nat Rev Genet* **10**: 195–205. doi:10.1038/nrg2526

Chunco AJ. 2014. Hybridization in a warmer world. *Ecol Evol* **4**: 2019–2031. doi:10.1002/ece3.1052

Cižková D, Javůrková V, Champagnon J, Kreisinger J. 2012. Duck's not dead: does restocking with captive bred individuals affect the genetic integrity of wild mallard (*Anas platyrhynchos*) population? *Biol Conserv* **152**: 231–240. doi:10.1016/j.biocon.2012.04.008

Coyne JA, Orr HA. 2004. *Speciation*. Sinauer, Sunderland, MA.

Crispo E, Moore JS, Lee-Yaw JA, Gray SM, Haller BC. 2011. Broken barriers: human-induced changes to gene flow and introgression in animals. *Bioessays* **33**: 508–518. doi:10.1002/bies.201000154

Curtzen PJ, Stoermer EF. 2000. The “Anthropocene.” *Global Change Newsletter* **41**: 17–18.

Darimont CT, Carlson SM, Kinnison MT, Paquet PC, Reimchen TE, Wilmers CC. 2009. Human predators outpace other agents of trait change in the wild. *Proc Natl Acad Sci* **106**: 952–954. doi:10.1073/pnas.0809235106

De Jong K, Amorim MCP, Fonseca PJ, Heubel KU. 2018. Noise affects multimodal communication during courtship in a marine fish. *Front Ecol Evol* **6**: 113. doi:10.3389/fevo.2018.00113

De-Kayne R, Selz OM, Marques DA, Frei D, Seehausen O, Feulner PDG. 2022. Genomic architecture of adaptive radiation and hybridization in Alpine whitefish. *Nat Commun* **13**: 4479. doi:10.1038/s41467-022-32181-8

Delaney KS, Riley SPD, Fisher RN. 2010. A rapid, strong, and convergent genetic response to urban habitat fragmentation in four divergent and widespread vertebrates. *PLoS ONE* **5**: e12767. doi:10.1371/journal.pone.0012767

Diamond SE, Chick L, Perez A, Strickler SA, Martin RA. 2017. Rapid evolution of ant thermal tolerance across an urban-rural temperature cline. *Biol J Linn Soc* **121**: 248–257. doi:10.1093/biolinnean/blw047

Díaz-Muñoz SL. 2012. Role of recent and old riverine barriers in fine-scale population genetic structure of Geoffroy's tamarin (*Saguinus geoffroyi*) in the Panama Canal watershed. *Evol Evol* **2**: 298–309. doi:10.1002/ece3.79

Dobbs C, Nitschke C, Kendal D. 2017. Assessing the drivers shaping global patterns of urban vegetation landscape structure. *Sci Total Environ* **592**: 171–177. doi:10.1016/j.scitotenv.2017.03.058

Dokulil M. 2014. Impact of climate warming on European inland waters. *Inland Waters* **4**: 27–40. doi:10.5268/IW-4.1.705

Egan SP, Ragland GJ, Assour L, Powell THQ, Hood GR, Emrich S, Nosil P, Feder JL. 2015. Experimental evidence of genome-wide impact of ecological selection during early stages of speciation-with-gene-flow. *Ecol Lett* **18**: 817–825. doi:10.1111/ele.12460

Elgyvin TO, Trier CN, Tørresen OK, Hagen IJ, Lien S, Nederbragt AJ, Ravinet M, Jensen H, Sætre GP. 2017. The ge-

nomic mosaicism of hybrid speciation. *Sci Adv* **3**: e1602996. doi:10.1126/sciadv.1602996

Ellis EC, Ramankutty N. 2008. Putting people in the map: anthropogenic biomes of the world. *Front Ecol Environ* **6**: 439–447. doi:10.1890/070062

Feder J, Chilcote C, Bush G. 1988. Genetic differentiation between sympatric host races of the apple maggot fly *Rhagoletis pomonella*. *Nature* **336**: 61–64. doi:10.1038/336061a0

Feder JL, Opp SB, Wlazlo B, Reynolds K, Go W, Spisak S. 1994. Host fidelity is an effective premating barrier between sympatric races of the apple maggot fly. *Proc Natl Acad Sci* **91**: 7990–7994. doi:10.1073/pnas.91.17.7990

Feder JL, Berlocher SH, Roethele JB, Dambroski H, Smith JJ, Perry WL, Gavrilovic V, Filchak KE, Rull J, Aluja M. 2003. Allopatric genetic origins for sympatric host-plant shifts and race formation in *Rhagoletis*. *Proc Natl Acad Sci* **100**: 10314–10319. doi:10.1073/pnas.1730757100

Ferrarin C, Bajo M, Bellafiore D, Cucco A, De Pascalis F, Ghezzo M, Umgieser G. 2014. Toward homogenization of mediterranean lagoons and their loss of hydrodiversity. *Geophys Res Lett* **41**: 5935–5941. doi:10.1002/2014GL060843

Filchak KE, Roethele JB, Feder JL. 2000. Natural selection and sympatric divergence in the apple maggot *Rhagoletis pomonella*. *Nature* **407**: 739–742. doi:10.1038/35037578

Fisher HS, Wong BM, Rosenthal Gil G. 2006. Alteration of the chemical environment disrupts communication in a freshwater fish. *Proc Biol Sci* **273**: 1187–1193. doi:10.1098/rspb.2005.3406

Fonseca DM, Keyghobadi N, Malcolm CA, Mehmet C, Schaffner F, Mogi M, Fleischer RC, Wilkerson RC. 2004. Emerging vectors in the *Culex pipiens* complex. *Science* **303**: 1535–1538. doi:10.1126/science.1094247

Foote M. 2023. Diversity-dependent diversification in the history of marine animals. *Am Nat* **201**: 680–693. doi:10.1086/723626

Ford RJ, Selman W, Taylor SS. 2017. Hybridization between mottled ducks (*Anas fulvigula maculosa*) and mallards (*A. platyrhynchos*) in the western Gulf coast region. *Condor* **119**: 683–696. doi:10.1650/CONDOR-17-18.1

Franks SJ, Hoffmann AA. 2012. Genetics of climate change adaptation. *Ann Rev Genet* **46**: 185–208. doi:10.1146/annurev-genet-110711-155511

Franssen NR. 2011. Anthropogenic habitat alteration induces rapid morphological divergence in a native stream fish. *Evol Appl* **4**: 791–804. doi:10.1111/j.1752-4571.2011.00200.x

Franssen NR, Stewart LK, Schaefer JF. 2013. Morphological divergence and flow-induced phenotypic plasticity in a native fish from anthropogenically altered stream habitats. *Ecol Evol* **3**: 4648–4657. doi:10.1002/eece3.842

Frei D, De-Kayne R, Selz OM, Seehausen O, Feulner PGD. 2022. Genomic variation from an extinct species is retained in the extant radiation following speciation reversal. *Nat Ecol Evol* **6**: 461–468. doi:10.1038/s41559-022-01665-7

Frei D, Reichlin P, Seehausen O, Feulner PGD. 2023. Introgression from extinct species facilitates adaptation to its vacated niche. *Mol Ecol* **32**: 841–853. doi:10.1111/mec.16791

Fuller MR, Doyle MW, Strayer DL. 2015. Causes and consequences of habitat fragmentation in river networks. *Ann NY Acad Sci* **1355**: 31–51. doi:10.1111/nyas.12853

* Garlovsky MD, Whittington E, Albrech T, Arenas-Castro H, Castillo DM, Keais GL, Larson EL, Moyle LC, Plakke M, Reifová R, et al. 2023. Synthesis and scope of the role of postmating prezygotic isolation in speciation. *Cold Spring Harb Perspect Biol* doi:10.1101/cshperspect.a041429

Garroway CJ, Bowman J, Cascaden TJ, Holloway GL, Mahan CG, Malcolm JR, Steele MA, Turner G, Wilson PJ. 2010. Climate change induced hybridization in flying squirrels. *Glob Chang Biol* **16**: 113–121. doi:10.1111/j.1365-2486.2009.01948.x

Garroway CJ, Bowman J, Holloway GL, Malcolm JR, Wilson PJ. 2011. The genetic signature of rapid range expansion by flying squirrels in response to contemporary climate warming. *Glob Chang Biol* **17**: 1760–1769. doi:10.1111/j.1365-2486.2010.02384.x

Gavrilets S. 2003. Perspective: models of speciation: what have we learned in 40 years? *Evolution (NY)* **10**: 2197–2215.

Gering E, Johnsson M, Willis P, Getty T, Wright D. 2015. Mixed ancestry and admixture in Kauai's feral chickens: invasion of domestic genes into ancient red Junglefowl reservoirs. *Mol Ecol* **24**: 2112–2124. doi:10.1111/mec.13096

Giery ST, Layman CA, Langerhans RB. 2015. Anthropogenic ecosystem fragmentation drives shared and unique patterns of sexual signal divergence among three species of Bahamian mosquitofish. *Evol Appl* **8**: 679–691. doi:10.1111/eva.12275

Gow JL, Peichel CL, Taylor EB. 2006. Contrasting hybridization rates between sympatric three-spined sticklebacks highlight the fragility of reproductive barriers between evolutionarily young species. *Mol Ecol* **15**: 739–752. doi:10.1111/j.1365-294X.2006.02825.x

Grabenstein KC, Taylor SA. 2018. Breaking barriers: causes, consequences, and experimental utility of human-mediated hybridization. *Trends Ecol Evol* **33**: 198–212. doi:10.1016/j.tree.2017.12.008

Grabenstein KC, Otter KA, Burg TM, Taylor SA. 2023. Hybridization between closely related songbirds is related to human habitat disturbance. *Glob Chang Biol* **29**: 955–968. doi:10.1111/gcb.16476

Grant PR, Grant BR. 2019. Hybridization increases population variation during adaptive radiation. *Proc Natl Acad Sci* **116**: 23216–23224. doi:10.1073/pnas.1913534116

Guo Q. 2014. Plant hybridization: the role of human disturbance and biological invasion. *Divers Distrib* **20**: 1345–1354. doi:10.1111/ddi.12245

Guo WY, Lambertini C, Pyšek P, Meyerson LA, Brix H. 2018. Living in two worlds: evolutionary mechanisms act differently in the native and introduced ranges of an invasive plant. *Ecol Evol* **8**: 2440–2452. doi:10.1002/ece3.3869

Haas TC, Blum MJ, Heins DC. 2010. Morphological responses of a stream fish to water impoundment. *Biol Lett* **6**: 803–806. doi:10.1098/rsbl.2010.0401

Anthropogenic Change and Speciation

Haba Y, McBride L. 2022. Origin and status of *Culex pipiens* mosquito ecotypes. *Curr Biol* **32**: R237–R246. doi:10.1016/j.cub.2022.01.062

Hallam A, Wignall PB. 1997. *Mass extinctions and their aftermath*. Oxford University Press, Oxford.

Hamilton JA, Miller JM. 2016. Adaptive introgression as a resource for management and genetic conservation in a changing climate. *Conserv Biol* **30**: 33–41. doi:10.1111/cobi.12574

Harries PJ, Kauffman EG, Hansen TA. 1996. Models for biotic survival following mass extinction. *Geol Soc Lon Spec Publ* **102**: 41–60. doi:10.1144/GSL.SP.1996.001.01.03

Harris SA. 2002. Introduction of the Oxford ragwort *Senecio squolidus* L. (Asteraceae) into the United Kingdom. *Watsonia* **24**: 31–42.

Heinen-Kay JL, Noel HG, Layman CA, Langerhans RB. 2014. Human-caused habitat fragmentation can drive rapid divergence of male genitalia. *Evol Appl* **7**: 1252–1267. doi:10.1111/eva.12223

Hendry AP, Wenburg JK, Bentzen P, Volk EC, Quinn TP. 2000. Rapid evolution of reproductive isolation in the wild: evidence from introduced salmon. *Science* **290**: 516–518. doi:10.1126/science.290.5491.516

Hendry AP, Gotanda M, Svensson EI. 2017. Human influences on evolution, and the ecological and societal consequences. *Phil Trans R Soc Lond B Biol Sci* **372**: 20160028–20160028. doi:10.1098/rstb.2016.0028

Hewitt GM. 1988. Hybrid zones—natural laboratories for evolutionary studies. *Trends Ecol Evol* **3**: 158–167. doi:10.1016/0169-5347(88)90033-X

Isaksson C. 2018. Impact of urbanization on birds. In *Bird species. Fascinating life sciences* (ed. Tietze D). Springer, Cham, Switzerland.

Jacobs A, Carruthers M, Eckmann R, Yohannes E, Adams CE, Behrmann-Godel J, Elmer KR. 2019. Rapid niche expansion by selection on functional genomic variation after ecosystem recovery. *Nat Ecol Evol* **3**: 77–86. doi:10.1038/s41559-018-0742-9

Javal M, Lombaert E, Tsykun T, Courtin C, Kerdelhué C, Prospero S, Roques A, Roux G. 2019. Deciphering the world-wide invasion of the Asian long-horned beetle: a recurrent invasion process from the native area together with a bridgehead effect. *Mol Ecol* **28**: 951–967. doi:10.1111/mec.15030

Jenkins MR, Cummings JM, Cabe AR, Hulthén K, Peterson MN, Langerhans RB. 2021. Natural and anthropogenic sources of habitat variation influence exploration behaviour, stress response, and brain morphology in a coastal fish. *J Anim Ecol* **90**: 2446–2461. doi:10.1111/1365-2656.13557

Johnson MTJ, Munshi-South J. 2017. Evolution of life in urban environments. *Science* **358**: eaam8327. doi:10.1126/science.aam8327

Keller G, Pardo A. 2004. Disaster opportunists Guembelitridae: index for environmental catastrophes. *Mar Micropaleontol* **53**: 83–116. doi:10.1016/j.marmicro.2004.04.012

Kennedy CM, Oakleaf JR, Theobald DM, Baruch-Mordo S, Kiesecker J. 2020. *Global human modification of terrestrial systems. Palisades*. NASA Socioeconomic Data and Applications Center (SEDAC), New York.

Kleindorfer S, O'Connor JA, Dudaniec RY, Myers SA, Robertson J, Sulloway FJ. 2014. Species collapse via hybridization in Darwin's tree finches. *Am Nat* **183**: 325–341. doi:10.1086/674899

Koneru M, Care T. 2022. Animal coloration in the Anthropocene. *Front Ecol Evol* **10**: 857317. doi:10.3389/fevo.2022.857317

Krichbaum K, Mahan CG, Steele MA, Turner G, Hudson PJ. 2010. The potential role of *Strongyloides robustus* on parasite-mediated competition between two species of flying squirrels (*Glaucomys*). *J Wildl Dis* **46**: 229–235. doi:10.7589/0090-3558-46.1.229

Kunc HP, Schmidt R. 2021. Species sensitivities to a global pollutant: a meta-analysis on acoustic signals in response to anthropogenic noise. *Glob Chang Biol* **27**: 675–688. doi:10.1111/gcb.15428

Langerhans RB, Kern EMA. 2020. Urbanization and evolution in aquatic environments. In *Urban evolutionary biology* (ed. Szulkin M). Oxford University Press, Oxford.

Langerhans RB, Riesch R. 2013. Speciation by selection: a framework for understanding ecology's role in speciation. *Curr Zool* **59**: 31–52. doi:10.1093/czoolo/59.1.31

Larson G, Fuller DQ. 2014. The evolution of animal domestication. *Annu Rev Ecol Evol Syst* **45**: 115–136. doi:10.1146/annurev-ecolsys-110512-135813

Lawson DM, Williams CK, Lavretsky P, Howell DL, Fuller JC. 2021. Mallard–black duck hybridization and population genetic structure in North Carolina. *J Wildl Manage* **85**: 1243–1255. doi:10.1002/jwmg.22085

Le Corre V, Siol M, Vigouroux Y, Tenaillon MI, Delye C. 2020. Adaptive introgression from maize has facilitated the establishment of teosinte as a noxious weed in Europe. *Proc Natl Acad Sci* **117**: 25618–25627. doi:10.1073/pnas.2006633117

Lehman N, Eisenhawer A, Hansen K, Mech LD, Peterson RO, Gogan PJP, Wayne RK. 1991. Introgression of coyote mitochondrial DNA into sympatric North American gray wolf populations. *Evolution (NY)* **45**: 104–119. doi:10.2307/2409486

Le Moan AL, Roby C, Fraisse C, Daguin-Thiébaut C, Bierne N, Viard F. 2021. An introgression breakthrough left by an anthropogenic contact between two ascidians. *Mol Ecol* **30**: 6718–6732. doi:10.1111/mec.16189

Lewis SL, Maslin MA. 2015. Defining the anthropocene. *Nature* **519**: 171–180. doi:10.1038/nature14258

Linn C Jr, Feder JL, Nojima S, Dambroski HR, Berlocher SH, Roelofs W. 2003. Fruit odor discrimination and sympatric host race formation in *Rhagoletis*. *Proc Natl Acad Sci* **100**: 11490–11493. doi:10.1073/pnas.1635049100

Littleford-Colquhoun BL, Clemente C, Whiting MJ, Ortiz-Barrientos D, Frère CH. 2017. Archipelagos of the Anthropocene: rapid and extensive differentiation of native terrestrial vertebrates in a single metropolis. *Mol Ecol* **26**: 2466–2481. doi:10.1111/mec.14042

Loria A, Cristescu ME, Gonzalez A. 2019. Mixed evidence for adaptation to environmental pollution. *Evol Appl* **12**: 1259–1273. doi:10.1111/eva.12782

Lourenço A, Álvarez D, Wang JJ, Velo-Antón G. 2017. Trapped within the city: integrating demography, time

M. Ålund et al.

since isolation and population-specific traits to assess the genetic effects of urbanization. *Mol Ecol* **26**: 1498–1514. doi:10.1111/mec.14019

Lu Z, Johnson WE, Menotti-Raymond M, Yuhki N, Martenson JS, Mainka S, Shi-Qiang H, Zhihe Z, Li G, Pan W, et al. 2001. Patterns of genetic diversity in remaining giant panda populations. *Conserv Biol* **15**: 1596–1607. doi:10.1046/j.1523-1739.2001.00086.x

Mallet J. 2007. Hybrid speciation. *Nature* **446**: 279–283. doi:10.1038/nature05706

Martin MD, Mendelson TC. 2012. Signal divergence is correlated with genetic distance and not environmental differences in darters (Percidae: *Etheostoma*). *Evol Biol* **39**: 231–241. doi:10.1007/s11692-012-9179-2

Mary N, Iannuccelli N, Petit G, Bonnet N, Pinton A, Barasc H, Faure A, Calgaro A, Grosbois V, Servin B, et al. 2022. Genome-wide analysis of hybridization in wild boar populations reveals adaptive introgression from domestic pig. *Evol Appl* **15**: 1115–1128. doi:10.1111/eva.13432

Mather A, Hancox D, Riginos C. 2015. Urban development explains reduced genetic diversity in a narrow range endemic freshwater fish. *Conserv Genet* **16**: 625–634. doi:10.1007/s10592-014-0688-7

Mayr E. 1963. *Animal species and evolution*. Harvard University Press, Cambridge, MA.

McFarlane SE, Pemberton JM. 2019. Detecting the true extent of introgression during anthropogenic hybridization. *Trends in Evol Evol* **34**: 315–326. doi:10.1016/j.tree.2018.12.013

McFarlane SE, Hunter DC, Senn HV, Smith SL, Holland R, Huisman J, Pemberton JM. 2020. Increased genetic marker density reveals high levels of admixture between red deer and introduced Japanese sika in Kintyre, Scotland. *Evol Appl* **13**: 432–441. doi:10.1111/eva.12880

McPhail JD. 1992. Ecology and evolution of sympatric sticklebacks (*Gasterosteus*): evidence for a species-pair in Paxton Lake, Texada Island, British Columbia. *Can J Zool* **70**: 361–369. doi:10.1139/z92-054

Miles LS, Johnson JC, Dyer RJ, Verrelli BC. 2018. Urbanization as a facilitator of gene flow in a human health pest. *Mol Ecol* **27**: 3219–3230. doi:10.1111/mec.14783

Miles LS, Rivkin LR, Johnson MTJ, Munshi-South J, Verrelli BC. 2019. Gene flow and genetic drift in urban environments. *Mol Ecol* **28**: 4138–4151. doi:10.1111/mec.15221

Mooney HA, Cleland EE. 2001. The evolutionary impact of invasive species. *Proc Natl Acad Sci* **98**: 5446–5451. doi:10.1073/pnas.091093398

Muñoz-Fuentes V, Vilà C, Green AJ, Negro JJ, Sorenson MD. 2007. Hybridization between white-headed ducks and introduced ruddy ducks in Spain. *Mol Ecol* **16**: 629–638. doi:10.1111/j.1365-294X.2006.03170.x

Murúa M, Espinoza C, Bustamante R, Marín VH, Medel R. 2010. Does human-induced habitat transformation modify pollinator-mediated selection? A case study in *Viola portalesia* (Violaceae). *Oecologia* **163**: 153–162. doi:10.1007/s00442-010-1587-3

Nevado B, Harris S, Beaumont A, Hiscock S. 2020. Rapid homoploid hybrid speciation in British gardens: the origin of Oxford ragwort (*Senecio squalidus*). *Mol Ecol* **29**: 4221–4233. doi:10.1111/mec.15630

Nolte AW, Freyhof J, Stemshorn KC, Tautz D. 2005. An invasive lineage of sculpins, *Cottus* sp. (Pisces, Teleostei) in the Rhine with new habitat adaptations has originated from hybridization between old phylogeographic groups. *Proc R Soc Lond B Biol Sci* **272**: 2379–2387.

Nosil P. 2009. Adaptive population divergence in cryptic color-pattern following a reduction in gene flow. *Evolution (NY)* **63**: 1902–1912. doi:10.1111/j.1558-5646.2009.00671.x

Nosil P. 2012. *Ecological speciation*, Oxford Series in Ecology and Evolution. Oxford University Press, Oxford.

Nosil P, Flaxman SM. 2011. Conditions for mutation-order speciation. *Proc Biol Sci* **278**: 399–407. doi:10.1098/rspb.2010.1215

O'Brien PP, Bowman J, Coombs AB, Newar SL, Garroway CJ. 2021. Winter nest trees of sympatric northern (*Glaucomys sabrinus*) and southern (*Glaucomys volans*) flying squirrels: a test of reinforcement in a hybrid zone. *Can J Zool* **99**: 859–866. doi:10.1139/cjz-2021-0086

O'Brien PP, Bowman J, Newar SL, Garroway CJ. 2022. Testing the parasite-mediated competition hypothesis between sympatric northern and southern flying squirrels. *Int J Parasitol* **17**: 83–90.

Ottenburghs J. 2018. Exploring the hybrid speciation continuum in birds. *Evol Evol* **8**: 13027–13034. doi:10.1002/ece3.4558

Ottenburghs J. 2021. The genic view of hybridization in the Anthropocene. *Evol Appl* **14**: 2342–2360. doi:10.1111/eva.13223

Otto SP. 2018. Adaptation, speciation and extinction in the Anthropocene. *Proc Biol Sci* **285**: 20182047. doi:10.1098/rspb.2018.2047

Owens GL, Samuk K. 2020. Adaptive introgression during environmental change can weaken reproductive isolation. *Nat Clim Chang* **10**: 58–62. doi:10.1038/s41558-019-0628-0

Oziolor EM, Reid NM, Yair S, Lee KM, Guberman VerPloeg S, Bruns PC, Shaw JR, Whitehead A, Matson CW. 2019. Adaptive introgression enables evolutionary rescue from extreme environmental pollution. *Science* **364**: 455–457. doi:10.1126/science.aav4155

* Peñalba JV, Runemark A, Meier J, Singh P, Wogan G, Sánchez-Guillen R, Mallet J, Rometsch SJ, Menon M, Seehausen O, et al. 2023. The role of hybridization in species formation. *Cold Spring Harb Perspect Biol* doi:10.1101/cshperspect.a041445

Pilakouta N, Ålund M. 2021. Sexual selection and environmental change: what do we know and what comes next? *Curr Zool* **67**: 293–298. doi:10.1093/cz/zoab021

Pilot M, Greco C, von Holdt BM, Randi E, Jędrzejewski W, Sidorovich VE, Konopiński MK, Ostrander EA, Wayne RK. 2018. Widespread, long-term admixture between grey wolves and domestic dogs across Eurasia and its implications for the conservation status of hybrids. *Evol Appl* **11**: 662–680. doi:10.1111/eva.12595

Popovic I, Matias AMA, Bierne N, Riginos C. 2020. Twin introductions by independent invader mussel lineages are both associated with recent admixture with a native con-

Anthropogenic Change and Speciation

gener in Australia. *Evol Appl* **13**: 515–532. doi:10.1111/eva.12857

Prentis PJ, Wilson JR, Dormontt EE, Richardson DM, Lowe AJ. 2008. Adaptive evolution in invasive species. *Trends Plant Sci* **13**: 288–294. doi:10.1016/j.tplants.2008.03.004

Price T. 2008. *Speciation in birds*. Roberts and Company, Greenwood Village, CO.

Price T. 2022. *Ecology of a changed world*. Oxford Academic, New York.

Qvarnström A, Veen T, Husby A, Ålund M, Weissling FJ. 2023. Assortative mating in an ecological context: effects of mate choice errors and relative species abundance on the frequency and asymmetry of hybridization. *Am Nat* **201**: 125–137. doi:10.1086/722156

Ratcliffe PR. 1987. Distribution and current status of sika deer, *Cervus nippon*, in Great Britain. *Mamm Rev* **17**: 39–58. doi:10.1111/j.1365-2907.1987.tb00047.x

Ravinet M, Elgvin TO, Trier C, Aliabadian M, Gavrillov A, Sætre GP. 2018. Signatures of human-commensalism in the house sparrow genome. *Proc Biol Sci* **285**: 20181246. doi:10.1098/rspb.2018.1246

Reichard DG, Atwell JW, Pandit MM, Cardoso GC, Price TD, Ketterson ED. 2020. Urban birdsongs: higher minimum song frequency of an urban colonist persists in a common garden experiment. *Anim Behav* **170**: 33–41. doi:10.1016/j.anbehav.2020.10.007

Resasco J, Haddad NM, Orrock JL, Shoemaker D, Brudvig LA, Damschen EI, Tewksbury JJ, Levey DJ. 2014. Landscape corridors can increase invasion by an exotic species and reduce diversity of native species. *Ecology* **95**: 2033–2039. doi:10.1890/14-0169.1

Rhymer JM, Williams MJ, Braun MJ. 1994. Mitochondrial analysis of gene flow between New Zealand mallards (*Anas platyrhynchos*) and grey ducks (*A. superciliosa*). *Auk* **111**: 970–978. doi:10.2307/4088829

Richardson LE, Graham NAJ, Pratchett MS, Eurich JG, Hoey AS. 2018. Mass coral bleaching causes biotic homogenization of reef fish assemblages. *Glob Change Biol* **24**: 3117–3129. doi:10.1111/gcb.14119

Riesch R, Easter T, Layman CA, Langerhans RB. 2015. Rapid human-induced divergence of life-history strategies in Bahamian livebearing fishes (family Poeciliidae). *J Anim Ecol* **84**: 1732–1743. doi:10.1111/1365-2656.12425

Riley SP, Pollinger JP, Sauvajot RM, York EC, Bromley C, Fuller TK, Wayne RK. 2006. FAST-TRACK: a southern California freeway is a physical and social barrier to gene flow in carnivores. *Mol Ecol* **15**: 1733–1741. doi:10.1111/j.1365-294X.2006.02907.x

Roebroeks W, Villa P. 2011. On the earliest evidence for habitual use of fire in Europe. *Proc Natl Acad Sci* **108**: 5209–5214. doi:10.1073/pnas.1018116108

Rosenblum EB, Sarver BAJ, Brown JW, Des Roches S, Hardwick KM, Hether TD, Eastman JM, Pennell MW, Harmon LJ. 2012. Goldilocks meets Santa Rosalia: an ephemeral speciation model explains patterns of diversification across time scales. *Evol Biol* **39**: 255–261. doi:10.1007/s11692-012-9171-x

Runemark A, Trier CN, Eroukhmanoff F, Hermansen JS, Matschiner M, Ravinet M, Elgvin TO, Sætre GP. 2018. Variation and constraints in hybrid genome formation. *Nat Ecol Evol* **2**: 549–556. doi:10.1038/s41559-017-0437-7

Saarman NP, Pogson GH. 2015. Introgression between invasive and native blue mussels (genus *Mytilus*) in the central California hybrid zone. *Mol Ecol* **24**: 4723–4738. doi:10.1111/mec.13340

Santangelo JS, Ness RW, Cohan B, Fitzpatrick CR, Innes SG, Koch S, Miles LS, Munim S, Peres-Neto PR, Prashad C, et al. 2022. Global urban environmental change drives adaptation in white clover. *Science* **375**: 1275–1281. doi:10.1126/science.abk0989

Schlüter D. 2009. Evidence for ecological speciation and its alternative. *Science* **323**: 737–741. DOI:10.1126/science.1160006

Seehausen O. 2006. Conservation: losing biodiversity by reverse speciation. *Curr Biol* **16**: R336–R337. doi:10.1016/j.cub.2006.03.080

Seehausen O, Van Alphen JJM, Witte F. 1997. Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* **277**: 1808–1811. doi:10.1126/science.277.5333.1808

Senn HV, Ghazali M, Kaden J, Barclay D, Harrower B, Campbell RD, Macdonald DW, Kitchener AC. 2019. Distinguishing the victim from the threat: SNP-based methods reveal the extent of introgressive hybridization between wildcats and domestic cats in Scotland and inform future in situ and ex situ management options for species restoration. *Evol Appl* **12**: 399–414. doi:10.1111/eva.12720

Servedio MR, Hermisson J. 2020. The evolution of partial reproductive isolation as an adaptive optimum. *Evolution (NY)* **74**: 4–14. doi:10.1111/evo.13880

Sievers M, Hale R, Parris KM, Sweare SE. 2018. Impacts of human-induced environmental change in wetlands on aquatic animals. *Biol Rev* **93**: 529–554. doi:10.1111/brv.12358

Simon A, Arbiol C, Nielsen EE, Couteau J, Sussarellu R, Burgeot T, Bernard I, Coolen JWP, Lamy JB, et al. 2020. Replicated anthropogenic hybridisations reveal parallel patterns of admixture in marine mussels. *Evol Appl* **13**: 575–599. doi:10.1111/eva.12879

Sirkia PM, McFarlane SE, Jones W, Wheatcroft D, Ålund M, Rybinski J, Qvarnström A. 2018. Climate-driven build-up of temporal isolation within a recently formed avian hybrid zone. *Evolution (NY)* **72**: 363–374. doi:10.1111/evo.13404

Smith VH, Schindler DW. 2009. Eutrophication science: where do we go from here? *Trends Ecol Evol* **24**: 201–207. doi:10.1016/j.tree.2008.11.009

Smith RF, Alexander LC, Lamp WO. 2009. Dispersal by terrestrial stages of stream insects in urban watersheds: a synthesis of current knowledge. *J North Am Benthol Soc* **28**: 1022–1037. doi:10.1899/08-176.1

Stewart L, Simonsen CE, Svenning JC, Schmidt NM, Pellissier L. 2018. Forecasted homogenization of high Arctic vegetation communities under climate change. *J Biogeogr* **45**: 2576–2587. doi:10.1111/jbi.13434

Strauss SY, Lau JA, Carroll SP. 2006. Evolutionary responses of natives to introduced species: what do introductions tell us about natural communities? *Ecol Lett* **9**: 357–374. doi:10.1111/j.1461-0248.2005.00874.x

M. Ålund et al.

Su H, Qu LJ, He K, Zhang Z, Wang J, Chen Z, Gu H. 2003. The Great Wall of China: a physical barrier to gene flow? *Heredity (Edinb)* **90**: 212–219. doi:10.1038/sj.hdy.6800237

Szulkin M, Munshi-South J, Charmantier A. 2020. *Urban evolutionary biology*. Oxford University Press, Oxford.

Tait C, Batra S, Ramaswamy SS, Feder JL, Olsson SB. 2016. Sensory specificity and speciation: a potential neuronal pathway for host fruit odour discrimination in *Rhagoletis pomonella*. *Proc R Soc Lond B Biol Sci* **283**: 20162101.

Taylor EB, Boughman JW, Groenenboom M, Sniatynski M, Schlüter D, Gow JL. 2006. Speciation in reverse: morphological and genetic evidence of the collapse of a three-spined stickleback (*Gasterosteus aculeatus*) species pair. *Mol Ecol* **15**: 343–355. doi:10.1111/j.1365-294X.2005.02794.x

Taylor SA, Curry RL, White TA, Ferretti V, Lovette I. 2014. Spatiotemporally consistent genomic signatures of reproductive isolation in a moving hybrid zone. *Evolution (NY)* **68**: 3066–3081. doi:10.1111/evo.12510

Templeton AR, Robertson RJ, Brisson J, Strasburg J. 2001. Disrupting evolutionary processes: the effect of habitat fragmentation on collared lizards in the Missouri Ozarks. *Proc Natl Acad Sci* **98**: 5426–5432. doi:10.1073/pnas.091093098

Thompson TQ, Bellinger MR, O'Rourke SM, Prince, DJ, Stevenson, AE, Rodrigues, AT, Sloat, MR, Speller, CF, Yang, DY, Butler, VL, et al. 2019. Anthropogenic habitat alteration leads to rapid loss of adaptive variation and restoration potential in wild salmon populations. *Proc Natl Acad Sci* **116**: 177–186. doi:10.1073/pnas.1811559115

* Thompson KA, Brandvain Y, Coughlan JM, Delmore KE, Justen H, Linnen CR, Ortiz-Barrientos D, Rushworth CA, Schneemann H, Schumer M, Stelkens R. 2023. The ecology of hybrid incompatibilities. *Cold Spring Harb Perspect Biol* doi:10.1101/cshperspect.a041440

Toups M, Kitchen A, Light JE, Reed DL. 2011. Origin of clothing lice indicates early clothing use by anatomically modern humans in Africa. *Mol Biol Evol* **28**: 29–32. doi:10.1093/molbev/msq234

Trombulak SC, Frissell CA. 2000. Review of ecological effects of roads on terrestrial and aquatic communities. *Conserv Biol* **14**: 18–30. doi:10.1046/j.1523-1739.2000.99084.x

Valencia-Montoya WA, Elfekih S, North HL, Meier JI, Warren IA, Tay WT, Gordon KHJ, Specht A, Paula-Moraes SV, Rane R, et al. 2020. Adaptive introgression across semipermeable species boundaries between local *Helicoverpa zea* and invasive *Helicoverpa armigera* moths. *Mol Biol Evol* **37**: 2568–2583. doi:10.1093/molbev/msaa108

van der Valk T, Díez-del-Molino D, Marques-Bonet T, Guenschanski K, Dalén L. 2019. Historical genomes reveal the genomic consequences of recent population decline in Eastern gorillas. *Curr Biol* **19**: 165–170.e6. doi:10.1016/j.cub.2018.11.055

van Hengstum T, Lachmuth S, Oostermeijer JGB, den Nijs HCM, Meirmans PG, van Tienderen PH. 2012. Human-induced hybridization among congeneric endemic plants on Tenerife, Canary Islands. *Plant Syst Evol* **298**: 1119–1131. doi:10.1007/s00606-012-0624-6

Veracx A, Raoult D. 2012. Biology and genetics of human head and body lice. *Trends Parasitol* **28**: 563–571. doi:10.1016/j.pt.2012.09.003

Vonlanthen P, Bittner D, Hudson AG, Young KA, Müller R, Lundsgaard-Hansen B, Roy D, Di Piazza S, Largiader CR, Seehausen O. 2012. Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature* **482**: 357–362. doi:10.1038/nature10824

Wagner DL, Grames EM, Forister ML, Berenbaum MR, Stoeck D. 2021. Insect decline in the Anthropocene: death by a thousand cuts. *Proc Natl Acad Sci* **118**: e2023989118. doi:10.1073/pnas.2023989118

Walsh BJ. 1867. The apple-worm and the apple maggot. *J Horticulture* **2**: 338–343.

Watrud LS, Lee EH, Fairbrother A, Burdick C, Reichman JR, Bollman M, Storm M, King G, Van de Water PK. 2004. Evidence for landscape-level, pollen-mediated gene flow from genetically modified creeping bentgrass with CP4 EPSPS as a marker. *Proc Natl Acad Sci* **101**: 14533–14538. doi:10.1073/pnas.0405154101

Wei F, Swaisgood R, Hu Y, Nie Y, Yan L, Zhang Z, Qi D, Zhu L. 2015. Progress in the ecology and conservation of giant pandas. *Conserv Biol* **29**: 1497–1507. doi:10.1111/cobi.12582

Wellborn GA, Langerhans RB. 2015. Ecological opportunity and the adaptive diversification of lineages. *Evol Evol* **5**: 176–195. doi:10.1002/ee.3.1347

Westram AM, Faria R, Johannesson K, Butlin R. 2021. Using replicate hybrid zones to understand the genomic basis of adaptive divergence. *Mol Ecol* **30**: 3797–3814. doi:10.1111/mec.15861

Whitney KD, Gabler CA. 2008. Rapid evolution in introduced species, “invasive traits” and recipient communities: challenges for predicting invasive potential. *Divers Distrib* **14**: 569–580. doi:10.1111/j.1472-4642.2008.00473.x

Wisely SM, Santymire RM, Livieri TM, Muetting S, Howard J. 2008. Genotypic and phenotypic consequences of reintroduction history in the black-footed ferret (*Mustela nigripes*). *Conserv Genet* **9**: 389–399. doi:10.1007/s10592-007-9351-x

Wood CM, Witham JW, Hunter ML Jr. 2016. Climate-driven range shifts are stochastic processes at a local level: two flying squirrel species in Maine. *Ecosphere* **7**: e01240.

Wright KM, Lloyd D, Lowry DB, Macnair MR, Willis JH. 2013. Indirect evolution of hybrid lethality due to linkage with selected locus in *Mimulus guttatus*. *PLoS Biol* **11**: e1001497. doi:10.1371/journal.pbio.1001497

Zapiola ML, Mallory-Smith CA. 2012. Crossing the divide: gene flow produces intergeneric hybrid in feral transgenic creeping bentgrass population. *Mol Ecol* **21**: 4672–4680. doi:10.1111/j.1365-294X.2012.05627.x

Zbawicka M, Trucco MI, Wenne R. 2018. Single nucleotide polymorphisms in native South American Atlantic coast populations of smooth shelled mussels: hybridization with invasive European *Mytilus galloprovincialis*. *Genet Sel Evol* **50**: 1–14. doi:10.1186/s12711-018-0376-z

* Zhang L, Nonaka E, Egan S. 2023. Context-dependent reproductive isolation and its role in (un) speciation. *Cold Spring Harb Perspect Biol* doi:10.1101/cshperspect.a041430



Anthropogenic Change and the Process of Speciation

Murielle Ålund, Meredith Cenzer, Nicolas Bierne, Janette W. Boughman, José Cerca, Mattheau S. Comerford, Alessandro Culicchi, Brian Langerhans, S. Eryn McFarlane, Markus H. Möst, Henry North, Anna Qvarnström, Mark Ravinet, Richard Svanbäck and Scott A. Taylor

Cold Spring Harb Perspect Biol published online October 3, 2023

Subject Collection [Speciation](#)

**Mechanisms of Intrinsic Postzygotic Isolation:
From Traditional Genic and Chromosomal Views
to Genomic and Epigenetic Perspectives**

Radka Reifová, S. Lorena Ament-Velásquez, Yann Bourgeois, et al.

**Quantitative Analyses of Coupling in Hybrid
Zones**

Thomas J. Firneno, Jr., Georgy Semenov, Erik B. Dopman, et al.

**Favored Races in the Struggle for Life: Racism
and the Speciation Concept**

Joseph L. Graves, Jr.

**Anthropogenic Change and the Process of
Speciation**

Murielle Ålund, Meredith Cenzer, Nicolas Bierne, et al.

**The Impact of Chromosomal Rearrangements in
Speciation: From Micro- to Macroevolution**

Kay Lucek, Mabel D. Giménez, Mathieu Joron, et al.

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

