

Stable manifolds of biholomorphisms in \mathbb{C}^n asymptotic to formal curves

Lorena López-Hernanz¹ | Javier Ribón² |
 Fernando Sanz-Sánchez³ | Liz Vivas⁴

¹Departamento de Física y Matemáticas,
Universidad de Alcalá, Madrid, Spain

²Instituto de Matemática e Estatística,
Universidade Federal Fluminense,
Niteroi, Brazil

³Departamento de Álgebra, Análisis
Matemático, Geometría y Topología,
Universidad de Valladolid, Valladolid,
Spain

⁴Department of Mathematics, The Ohio
State University, Columbus, Ohio, USA

Correspondence

Liz Vivas, Department of Mathematics,
The Ohio State University, 231 West 18th
Ave, Columbus, OH 43214, USA.
Email: vivas@math.osu.edu

Funding information

Ministerio de Ciencia e Innovación; NSF,
Grant/Award Number: 1800777

Abstract

Given a germ of biholomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ with a formal invariant curve Γ such that the multiplier of the restricted formal diffeomorphism $F|_{\Gamma}$ is a root of unity or satisfies $|F|_{\Gamma}'(0)| < 1$, we prove that either Γ is contained in the set of periodic points of F or there exists a finite family of stable manifolds of F where all the orbits are asymptotic to Γ and whose union eventually contains every orbit asymptotic to Γ . This result generalizes to the case where Γ is a formal periodic curve.

MSC (2020)

32M25, 37C25 (primary)

1 | INTRODUCTION

In this paper, we generalize to arbitrary dimension n the main result proved by López-Hernanz et al. in [22] for $n = 2$. Namely, we consider a germ of biholomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ with a formal invariant curve Γ at the origin. Assuming that the multiplier λ of the restricted (formal) diffeomorphism $F|_{\Gamma}$ is a root of unity or satisfies $|\lambda| < 1$, we prove that either Γ is contained in the set of periodic points of F or there exists a finite family of stable manifolds of F whose union consists of and contains eventually any orbit of F asymptotic to Γ , that is, having flat contact with Γ . Note that the condition on λ corresponds to the necessary condition for the existence of stable orbits of the one-dimensional dynamics of $F|_{\Gamma}$ when Γ is convergent (see Pérez-Marco [29, 30]). It does not depend on the rest of the multipliers of F .

In the two-dimensional case, the stable manifolds obtained in [22] are either one-dimensional (saddle behavior) or open sets (node behavior). In dimension $n \geq 3$, we also obtain stable manifolds of intermediate dimension $1 < s < n$ that share some properties with both the saddle and the node cases of dimension two. Although the theorem is very similar, the proof is not a straightforward generalization of the two-dimensional one. We need to introduce new techniques that we discuss below in this introduction.

Let us first describe more precisely the statement of the main theorem.

A *stable set* of F is a subset $B \subset V$ of an open neighborhood V of 0 where F is defined which is invariant, that is, $F(B) \subset B$, and such that the orbit of each point of B converges to 0. If B is an analytic, locally closed submanifold of V then we say that B is a *stable manifold* of F (in V). Let us remark that, as in [22], our definition of stable manifold is more general than the classical one, since the stable manifolds considered in this paper do not contain the origin in general, even if by definition their closures contain it.

A formal (irreducible) curve Γ at $0 \in \mathbb{C}^n$ is a prime ideal Γ of $\mathbb{C}[[x_1, \dots, x_n]]$ such that $\mathbb{C}[[x_1, \dots, x_n]]/\Gamma$ has dimension 1. We say that Γ is *invariant* by F if $\Gamma \circ F = \Gamma$. In this case, we can consider the *restriction* $F|_{\Gamma}$, which is a formal diffeomorphism in one variable (see Section 2 for details). A non-trivial (positive) orbit O of F is *asymptotic* to a formal curve Γ if O converges to the origin and, for any finite composition σ of blow-ups of points, the lifted orbit $\sigma^{-1}(O)$ has a limit equal to the point on the transform of Γ by σ . If this is the case then Γ is necessarily invariant for F (see Section 2).

Our main result is the following:

Theorem 1. *Consider $F \in \text{Diff}(\mathbb{C}^n, 0)$ and let Γ be a formal invariant curve of F . Assume that the multiplier $\lambda = (F|_{\Gamma})'(0)$ is a root of unity or satisfies $|\lambda| < 1$. Then we have one of the following two possibilities:*

- (i) *The curve Γ is contained in the set of fixed points of some non-trivial iterate of F , or*
- (ii) *$F|_{\Gamma}$ is not periodic and there exist orbits of F asymptotic to Γ .*

In the latter case, in any sufficiently small open neighborhood V of 0 there exists a non-empty finite family of pairwise disjoint stable manifolds $S_1, \dots, S_r \subset V$ of F of pure positive dimension and with finitely many connected components such that the orbit of every point in $S_1 \cup \dots \cup S_r$ is asymptotic to Γ and such that any orbit of F asymptotic to Γ is eventually contained in $S_1 \cup \dots \cup S_r$.

The stable manifolds S_1, \dots, S_r provide a base of asymptotic convergence along Γ Ueda [38]. We can be more precise in the hyperbolic case:

Proposition 2. *Consider $F \in \text{Diff}(\mathbb{C}^n, 0)$ and let Γ be an invariant formal curve of F . Assume that the multiplier $\lambda = (F|_{\Gamma})'(0)$ satisfies $|\lambda| < 1$. Then Γ is a germ of an analytic curve at the origin and a representative of Γ is a stable manifold of F that eventually contains any orbit of F asymptotic to Γ .*

The result can be stated more generally for a formal *periodic* curve Γ (i.e., Γ is invariant for some iterate F^s of F). More precisely, we apply Theorem 1 to F^s and Γ in order to obtain stable manifolds for F^s , which induce stable manifolds for F by simple arguments that can be found in [22].

It is worth mentioning that whereas a planar diffeomorphism $F \in \text{Diff}(\mathbb{C}^2, 0)$ always has a formal periodic curve (Ribón [32], see also Corollary 4.21), this is no longer true for dimension $n \geq 3$ by an example of a holomorphic vector field of Gómez-Mont and Luengo [16], whose flow is treated by Abate and Tovena in [1]. As a consequence of the results in Section 4, we will obtain

in Section 4.6 a condition that guarantees the existence of formal invariant curves in dimension three, inspired by a result of Cerveau and Lins for vector fields [11].

Notice that in any dimension there are linear examples of biholomorphisms F with an invariant axis for which the multiplier λ either satisfies $|\lambda| > 1$ or is *irrationally neutral* (i.e., $|\lambda| = 1$ and it is not a root of unity) and Theorem 1 does not hold. Thus, the hypothesis concerning λ in Theorem 1 is necessary. In fact, for $n = 1$, as we mentioned above, if there are positive orbits of F converging to the origin then λ satisfies the hypothesis of Theorem 1.

Let us describe the structure of the proof of Theorem 1. After recalling in Section 2 the main definitions and properties concerning formal curves, blow-ups, and asymptotic orbits, in Section 3 we study the case where $F|_{\Gamma}$ is hyperbolic attracting and we prove Proposition 2. In this case, the result is a consequence of the classical stable manifold and Hartman–Grobman theorems for diffeomorphisms and the proof goes as in the two-dimensional case [22].

The case where the multiplier λ is a root of unity is the core of the paper. We assume that $F|_{\Gamma}$ is not periodic. One of the main ingredients in this case is a suitable normal form for the pair (F, Γ) , that we call *Ramis–Sibuya form*. Namely, there exist coordinates (x, y) at $0 \in \mathbb{C}^n$ such that Γ is non-singular and transverse to $x = 0$ and F is written as

$$F(x, y) = (x - x^{q+1} + bx^{2q+1} + O(x^{2q+2}), \exp(D(x) + x^q C)y + O(x^{q+1})),$$

where $q \geq 1$, $b \in \mathbb{C}$, $D(x)$ is a diagonal polynomial matrix of degree at most $q - 1$ and C is a constant matrix such that $[D(x), C] = 0$. Ramis–Sibuya form is inspired by a classical result on normal forms of systems of linear ordinary differential equations (ODEs) with formal meromorphic coefficients due to Tjurittin [37]. Such normal forms are also used for non-linear systems by Braaksma [6] and Ramis and Sibuya [31] in order to prove multisummability of the formal solutions of the system (when the coefficients are convergent). In Section 5.1 we define the analogous Ramis–Sibuya form for a pair (X, Γ) , where X is a formal vector field and Γ is a formal invariant curve of X , and in Section 5.3 we prove that any pair (X, Γ) can be reduced to Ramis–Sibuya form by means of a finite number of blow-ups with smooth centers and ramifications, all of them adapted to Γ . This result is essentially a consequence of Tjurittin’s theorem once we associate to (X, Γ) a system of $n - 1$ formal meromorphic ODEs after some initial punctual blow-ups.

Then, concerning the reduction of a pair (F, Γ) to Ramis–Sibuya form, we use a result by Binyamini [5] that guarantees that an adequate iterate F^m of F is the time-1 flow of a formal vector field X , a so-called *infinitesimal generator*, which will allow to obtain a reduction of (F^m, Γ) to Ramis–Sibuya form from the corresponding one for (X, Γ) . In order for this construction to work the condition $F = \exp(X)$ is not sufficient. We need the biholomorphism and the vector field to share some additional geometrical properties; for instance, the invariant curve Γ must be also invariant for X . We devote Section 4 to showing the geometrical nature of the correspondence between local biholomorphisms and infinitesimal generators. Moreover, we determine whether X is a geometrical infinitesimal generator of $\exp(X)$ (Theorem 4.15). The condition depends only on the eigenvalues of the linear part $D_0 X$ of X at the origin.

Using these results, in Section 5.4 we accomplish the reduction of the pair (F^m, Γ) . The fact that we are replacing F by an iterate presents no problem for the proof of Theorem 1: if S is a stable manifold of F^m composed of orbits asymptotic to the invariant curve Γ , then $\cup_{k=0}^{m-1} F^k(S)$ is also stable for F . Moreover, the blow-ups and ramifications considered in the reduction preserve the property of asymptoticity of the orbits to the formal curve. Thus, for the proof of Theorem 1, we may assume that (F, Γ) is already in Ramis–Sibuya form.

Finally, in Section 6 we prove Theorem 1 when (F, Γ) is in Ramis–Sibuya form. The family of stable manifolds in the statement is associated to the family of the q attracting directions of the restricted formal diffeomorphism $F|_{\Gamma}(x) = x - x^{q+1} + O(x^{2q+1})$. In fact, if ℓ is such an attracting direction, the Ramis–Sibuya form allows to separate the \mathbf{y} -variables in two groups, depending only on the restriction of the polynomial matrix $D(x) + x^q C$ to ℓ . Dynamically, each of these groups corresponds either to a saddle or to a node behavior along the orbits that converge to the origin tangentially to ℓ . The dimension of the stable manifold associated to ℓ will then be equal to the number of node variables plus one. The proof of the existence of those stable manifolds is a generalization of the corresponding one in [22], inspired by Hakim’s construction in [19] (see also [2] by Arizzi and Raissy). In particular, they are obtained as fixed points of an adequate continuous map. Instead of using Banach fixed point theorem as in [18, 19, 23] or [22], we use Schauder fixed point theorem, which simplifies significantly the technical computations of the proof. The lack of uniqueness in Schauder’s theorem will be compensated by an easy alternative argument.

2 | FORMAL INVARIANT CURVES, ASYMPTOTIC ORBITS, AND BLOW-UPS

In this section we introduce the main definitions and properties concerning a formal invariant curve Γ of a formal vector field X or a biholomorphism F and the behavior of both F and Γ under punctual blow-ups. The content of this section is just a generalization to higher dimension of what can already be found in [22] for dimension two. We include it for the sake of completeness and to fix notations.

First, let us consider invariance by formal vector fields. An (*irreducible*) *formal curve* at $0 \in \mathbb{C}^n$ is a prime ideal Γ of the ring $\hat{\mathcal{O}}_n = \mathbb{C}[[x_1, \dots, x_n]]$ such that the quotient ring $\hat{\mathcal{O}}_n/\Gamma$ has dimension one. It is determined by a formal parameterization $\gamma(s) = (\gamma_1(s), \dots, \gamma_n(s)) \in (s\mathbb{C}[[s]])^n \setminus \{0\}$ so that $g(\gamma(s)) \equiv 0$ if and only if $g \in \Gamma$. Let $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ be a singular formal vector field. More precisely, once we choose coordinates $\mathbf{x} = (x_1, \dots, x_n)$, we write X as

$$X = a_1(\mathbf{x}) \frac{\partial}{\partial x_1} + a_2(\mathbf{x}) \frac{\partial}{\partial x_2} + \dots + a_n(\mathbf{x}) \frac{\partial}{\partial x_n},$$

where $a_j(\mathbf{x}) = X(x_j) \in \hat{\mathcal{O}}_n$ satisfies $a_j(0) = 0$. The *multiplicity* of X , denoted by $\nu(X)$, is the minimum of the orders of the series a_j , which is independent of the chosen coordinates. The *singular locus* of X is the ideal $\text{Sing}(X) \subset \hat{\mathcal{O}}_n$ generated by the series a_1, \dots, a_n .

Recall that a formal curve Γ is invariant for X if $X(\Gamma) \subset \Gamma$. In terms of a parameterization $\gamma(s)$ of Γ , invariance is equivalent to the existence of $h_{\gamma}(s) \in \mathbb{C}[[s]]$ such that

$$X|_{\gamma(s)} = (a_1(\gamma(s)), \dots, a_n(\gamma(s))) = h_{\gamma}(s)\gamma'(s). \quad (1)$$

Notice that $h_{\gamma}(s) \equiv 0$ if and only if $\text{Sing}(X) \subset \Gamma$ and thus this property is independent of the parameterization (we say that Γ is *contained* in the singular locus of X). When Γ is invariant, we define the *restriction* of X to Γ as the one-dimensional formal vector field

$$X|_{\Gamma} = h_{\gamma}(s) \frac{\partial}{\partial s},$$

where γ is an irreducible parameterization of Γ and $h_{\gamma}(s)$ is defined by Equation (1). Actually $X|_{\Gamma}$ can be defined intrinsically since $X(\Gamma) \subset \Gamma$ implies that X defines a derivation of the ring of

formal functions $\hat{\mathcal{O}}_n/\Gamma$ of Γ . The multiplier $\lambda_\Gamma = h'_\gamma(0) \in \mathbb{C}$ is called the *inner eigenvalue* of the pair (X, Γ) . The *tangent eigenvalue* of (X, Γ) , denoted by $\lambda(\Gamma)$, is the eigenvalue of the differential D_0X corresponding to the tangent direction of Γ . These eigenvalues are related by $\nu\lambda_\Gamma = \lambda(\Gamma)$, where ν is the multiplicity of Γ at 0.

A formal curve Γ is *invariant* for $F \in \text{Diff}(\mathbb{C}^n, 0)$ if $h \circ F \in \Gamma$ for any $h \in \Gamma$. Moreover, given a parameterization $\gamma(s)$ of Γ , the invariance of Γ is equivalent to the existence of a series $\theta(s) \in \mathbb{C}[[s]]$ with $\theta(0) = 0$ and $\theta'(0) \neq 0$ such that $F \circ \gamma(s) = \gamma \circ \theta(s)$. This series $\theta(s)$ can be seen as a formal diffeomorphism in one variable, that is, $\theta(s) \in \widehat{\text{Diff}}(\mathbb{C}, 0)$. Its class of formal conjugacy is independent of the chosen parameterization $\gamma(s)$ and any representative of this class is called the *restriction* of F to Γ and denoted by $F|_\Gamma$.

If Γ is invariant for F , the multiplier $\lambda_\Gamma = (F|_\Gamma)'(0) \in \mathbb{C}^*$ does not depend on $\theta(s)$ and is called the *inner eigenvalue* of the pair (F, Γ) . Notice that this number is preserved under reparameterizations. On the other hand, the *tangent eigenvalue* $\lambda(\Gamma)$ of (F, Γ) is the eigenvalue of D_0F corresponding to the tangent direction of Γ . It is easy to check that

$$(\lambda_\Gamma)^\nu = \lambda(\Gamma), \quad (2)$$

where ν is the multiplicity of Γ at 0. In particular, we have $\lambda_\Gamma = \lambda(\Gamma)$ when Γ is non-singular.

Definition 2.1. Let Γ be a formal invariant curve of $F \in \text{Diff}(\mathbb{C}^n, 0)$ and let λ_Γ be the inner eigenvalue. We say that Γ is *hyperbolic attracting* if $|\lambda_\Gamma| < 1$, and that Γ is *rationally neutral* if λ_Γ is a root of unity.

Consider a germ of biholomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$. Denote by $\pi : \widetilde{\mathbb{C}^n} \rightarrow \mathbb{C}^n$ the blow-up of \mathbb{C}^n at the origin and by $E = \pi^{-1}(0)$ the exceptional divisor. The transformed biholomorphism $\tilde{F} = \pi^{-1} \circ F \circ \pi$ extends to an injective holomorphic map in a neighborhood of E in $\widetilde{\mathbb{C}^n}$ such that $\tilde{F}(E) = E$. We have moreover that $\tilde{F}|_E$ is the projectivization of the linear map D_0F in the identification $E \simeq \mathbb{P}_{\mathbb{C}}^{n-1}$ and hence fixed points $p \in E$ for \tilde{F} correspond to invariant lines of D_0F . Such a point p is a *first infinitely near fixed point* of F and the germ F_p of \tilde{F} at p is the *transform* of F at p . Blowing-up repeatedly, we define sequences $\{p_k\}_{k \geq 0}$ of *infinitely near fixed points* of F and corresponding transforms F_{p_k} , where $p_0 = 0$.

A formal curve Γ is also determined by its sequence of *iterated tangents* $\{q_k\}_{k \geq 0}$, defined by: $q_0 = 0$ and, for $k \geq 1$, if $\pi_{q_{k-1}}$ is the blow-up at q_{k-1} , the point $q_k \in \pi_{q_{k-1}}^{-1}(q_{k-1})$ corresponds to the tangent line of the strict transform of Γ at q_{k-1} . The formal curve Γ is invariant for F if and only if the sequence of iterated tangents of Γ is a sequence of infinitely near fixed points of F (see [22]).

Note that the inner eigenvalue is invariant under blow-up and hence the condition of Γ being hyperbolic attracting or rationally neutral is stable under blow-ups.

Given a formal curve Γ at 0 $\in \mathbb{C}^n$, a stable non-trivial orbit $O = \{a_k = F^k(a_0)\}$ of a diffeomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ is *asymptotic* to Γ if, being $\{q_k\}$ the sequence of iterated tangents of Γ , the following holds: if $\pi_1 : M_1 \rightarrow \mathbb{C}^n$ is the blow-up at the origin then $\lim_{k \rightarrow \infty} \pi_1^{-1}(a_k) = q_1$; if $\pi_2 : M_2 \rightarrow M_1$ is the blow-up at q_1 then $\lim_{k \rightarrow \infty} \pi_2^{-1} \circ \pi_1^{-1}(a_k) = q_2$; and so on. Notice that if such an orbit exists then Γ is invariant for F , since in this case any iterated tangent q_k of Γ must be an infinitely near fixed point of F .

We remark that our definition of asymptoticity to a formal curve Γ corresponds to the standard one of having Γ as “asymptotic expansion.” For instance, if Γ is non-singular and we consider

a parameterization of the form $\gamma(s) = (s, \mathbf{h}(s)) \in \mathbb{C}[[s]]^n$ in some coordinates $(x, \mathbf{y}) \in \mathbb{C} \times \mathbb{C}^{n-1}$, with $\mathbf{h}(s) = \sum_{j=1}^{\infty} \mathbf{h}_j s^j$, then a non-trivial orbit $O = \{(x_k, \mathbf{y}_k)\}$ satisfies $\lim_{k \rightarrow \infty} \pi_1^{-1}(x_k, \mathbf{y}_k) = q_1$ if and only if $\lim_{k \rightarrow \infty} \mathbf{y}_k / x_k = \mathbf{h}_1$. Then we obtain $\lim_{k \rightarrow \infty} \pi_2^{-1} \circ \pi_1^{-1}(x_k, \mathbf{y}_k) = q_2$ if and only if $\lim_{k \rightarrow \infty} \frac{\mathbf{y}_k / x_k - \mathbf{h}_1}{x_k} = \mathbf{h}_2$ and so on. Therefore O is asymptotic to Γ if and only if for any $N \in \mathbb{N}$ we have

$$\lim_{k \rightarrow \infty} \frac{\mathbf{y}_k - J_N \mathbf{h}(x_k)}{x_k^{N+1}} = \mathbf{h}_{N+1},$$

where J_N denotes the N -jet. This implies $\|\mathbf{y}_k - J_N \mathbf{h}(x_k)\| \leq (||\mathbf{h}_{N+1}|| + 1)|x_k|^{N+1}$ for some $k \geq k_0(N)$.

3 | HYPERBOLIC ATTRACTING CASE

In this section we prove Theorem 1 in the case where the formal curve is hyperbolic attracting. More precisely, we prove Proposition 2 in the introduction that we recall next for convenience.

Proposition 3.1. *Consider $F \in \text{Diff}(\mathbb{C}^n, 0)$ and let Γ be an invariant formal curve of F . Assume that Γ is hyperbolic attracting. Then Γ is a germ of an analytic curve at the origin and a representative of Γ is a stable manifold of F that eventually contains any orbit of F asymptotic to Γ .*

Proof. Let $\{q_k\}_{k \geq 0}$ be the sequence of iterated tangents of Γ . Notice that it suffices to prove the statement for F_{q_k} and Γ_k at any point q_k , where F_{q_k} is the transform of F at q_k and Γ_k is the strict transform of Γ at q_k . Notice that the curve Γ has a Puiseux parameterization by the local parameterization theorem (cf. [17, Volume II, Section D, Theorem 10]) and hence Γ can be desingularized by a sequence of punctual blow-ups. Thus, we can assume that Γ is non-singular. Let $\lambda = \lambda(\Gamma)$ be the tangent eigenvalue of (F, Γ) , which coincides with the inner eigenvalue λ_Γ since Γ is non-singular. Set $\text{spec}(D_0 F) = \{\lambda, \mu_2, \dots, \mu_n\}$. An easy computation shows that the eigenvalues of the linear part of F_{q_1} at q_1 are given by $\{\lambda, \mu_2/\lambda, \dots, \mu_n/\lambda\}$. Moreover, λ is still the tangent eigenvalue of the pair (F_{q_1}, Γ_1) since the inner eigenvalue is preserved under blow-up. Repeating this argument, it follows that, for each k , the eigenvalues of the linear part of F_{q_k} at q_k are $\{\lambda, \mu_2/\lambda^k, \dots, \mu_n/\lambda^k\}$. Now, assume that k is large enough so that $|\lambda| < 1 < |\mu_j|/|\lambda^k|$ for any $j = 2, \dots, n$. Then, by the Stable Manifold Theorem, we obtain that Γ_k is the stable manifold of F_{q_k} at q_k , hence an analytic curve. Moreover, using Hartman–Grobman theorem, we have that the unique orbits of F_{q_k} that converge to q_k are those which are eventually contained in Γ_k . \square

Remark 3.2. From the theory of one-dimensional dynamics, we have that the hyperbolic attracting case is the only one for which there is a stable set whose germ is an analytic curve at the origin (cf. [22]).

4 | INFINITESIMAL GENERATOR OF A BIHOLOMORPHISM

In this section, we recover a result due to Binyamini [5] that guarantees that for any local biholomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ there exists a formal vector field X such that the time-1 flow $\exp(X)$ of

the vector field is a non-trivial iterate F^m of F . We prove that this construction is “geometrically significant” in the sense that the geometrical properties of F^m and X are related. For example, the fixed point set of F^m coincides with the singular set of X . Moreover, the invariance of analytic sets is preserved by this correspondence between diffeomorphisms and formal vector fields; indeed, F^m preserves a germ of analytic set, or a formal analytic set, if and only if X does. In particular, if Γ is a formal invariant curve that is periodic for $F \in \text{Diff}(\mathbb{C}^n, 0)$, we will see that Γ is invariant by X . This property will be crucial in Section 5 to obtain a reduction of the pair (F^m, Γ) to Ramis–Sibuya form from the corresponding one for (X, Γ) .

4.1 | Preliminaries

The strategy to obtain a vector field X such that $\exp(X) = F^m$ for some m is a generalization to the context of diffeomorphisms of the correspondence between the connected component of the identity of a finite dimensional algebraic group and the Lie algebra of the group. This generalization is possible since the group $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$ of formal diffeomorphisms, despite being infinite dimensional, can be interpreted as a projective limit of finite dimensional algebraic groups. This approach allows to define the algebraic closure $\overline{\langle F \rangle}$ of the group $\langle F \rangle$ generated by F , its connected component of the identity and its associated Lie algebra. The infinitesimal generators of iterates of F are chosen in this Lie algebra. First, we introduce these ideas; further details can be found in [24, 35] and [34].

Consider the normal subgroup N_k of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$ defined by

$$N_k = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : x_j \circ F - x_j \in \mathfrak{m}^{k+1} \forall 1 \leq j \leq n\},$$

where \mathfrak{m} is the maximal ideal of the ring $\hat{\mathcal{O}}_n = \mathbb{C}[[x_1, \dots, x_n]]$ of formal power series. It is the subgroup of formal diffeomorphisms that have order of contact at least $k+1$ with the identity map. We denote by D_k the group $\widehat{\text{Diff}}(\mathbb{C}^n, 0)/N_k$ of k -jets of formal diffeomorphisms. Given $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$, we can uniquely associate to F the element

$$\begin{aligned} F_k &: \mathfrak{m}/\mathfrak{m}^{k+1} \rightarrow \mathfrak{m}/\mathfrak{m}^{k+1} \\ f + \mathfrak{m}^{k+1} &\mapsto f \circ F + \mathfrak{m}^{k+1} \end{aligned} \tag{3}$$

of the linear group $\text{GL}(\mathfrak{m}/\mathfrak{m}^{k+1})$ which only depends on the class of F in D_k . In this way we can interpret D_k as a subgroup of $\text{GL}(\mathfrak{m}/\mathfrak{m}^{k+1})$. Moreover, it is a (finite dimensional) algebraic matrix group since $\{F_k : F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)\}$ coincides with the group of automorphisms of the \mathbb{C} -algebra $\mathfrak{m}/\mathfrak{m}^{k+1}$ (cf. [35, Lemma 2.1]). The Lie algebra L_k of $\{F_k : F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)\}$ is the Lie algebra of derivations of the \mathbb{C} -algebra $\mathfrak{m}/\mathfrak{m}^{k+1}$ for any $k \geq 1$. Moreover, L_k can be identified with $\hat{\mathfrak{X}}(\mathbb{C}^n, 0)/K_k$ where $\hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ is the complex Lie algebra of singular formal vector fields (i.e., derivations of the \mathbb{C} -algebra \mathfrak{m}) and $K_k = \{X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0) : X(\mathfrak{m}) \subset \mathfrak{m}^{k+1}\}$.

The natural projections $\pi_{k,l} : D_k \rightarrow D_l$ and $(d\pi_{k,l})_{\text{id}} : L_k \rightarrow L_l$ when $k \geq l$ define inverse systems and the group $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$ (respectively the Lie algebra $\hat{\mathfrak{X}}(\mathbb{C}^n, 0)$) can be identified with the projective limit of the groups D_k (respectively the Lie algebras L_k) for $k \geq 1$. We denote by $\pi_k : \widehat{\text{Diff}}(\mathbb{C}^n, 0) \rightarrow D_k$ and $d\pi_k : \hat{\mathfrak{X}}(\mathbb{C}^n, 0) \rightarrow L_k$ the natural maps that send the projective limits onto their factors.

Definition 4.1. Given a subgroup G of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$, we denote by G_k the Zariski closure of $\pi_k(G)$ and by $G_{k,0}$ the connected component of the identity of G_k for $k \geq 1$. Then we define

$$\overline{G} = \varprojlim G_k = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : \pi_k(F) \in G_k \ \forall k \in \mathbb{N}\}$$

and

$$\overline{G}_0 = \varprojlim G_{k,0} = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : \pi_k(F) \in G_{k,0} \ \forall k \in \mathbb{N}\}$$

as the *Zariski closure* (or *pro-algebraic closure*) of G and its *connected component of the identity*, respectively. Given a subgroup G of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$, we say that it is *pro-algebraic* if $\overline{G} = G$. We define the *Lie algebra* \mathfrak{g} of \overline{G} as

$$\mathfrak{g} = \varprojlim \mathfrak{g}_k = \{X \in \widehat{\mathfrak{X}}(\mathbb{C}^n, 0) : d\pi_k(X) \in \mathfrak{g}_k \ \forall k \in \mathbb{N}\},$$

where \mathfrak{g}_k is the Lie algebra of G_k for $k \geq 1$.

Remark 4.2 [24, Proposition 2]. The Lie algebra \mathfrak{g} of \overline{G} satisfies

$$\mathfrak{g} = \{X \in \widehat{\mathfrak{X}}(\mathbb{C}^n, 0) : \exp(tX) \in \overline{G} \ \forall t \in \mathbb{C}\},$$

where $\exp(tX)$ is the time- t flow of X , that is, the formal diffeomorphism that satisfies, for any $g \in \hat{\mathcal{O}}_n$,

$$g \circ \exp(tX) = \sum_{j=0}^{\infty} \frac{t^j X^j(g)}{j!},$$

where $X^0(g) = g$ and $X^j(g) = X(X^{j-1}(g))$ for $j \geq 1$.

In the two following results we summarize several properties of the finite dimensional setting that generalize to the infinite dimensional one and provide a criterion that allows to identify pro-algebraic groups of formal diffeomorphisms.

Proposition 4.3. *Let G be a subgroup of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$. We have*

- (i) \overline{G}_0 is a finite index normal pro-algebraic subgroup of \overline{G} [35, Proposition 2.3 and Remark 2.9].
- (ii) Any finite index subgroup of \overline{G} is pro-algebraic and contains \overline{G}_0 [34, Lemmas 2.3 and 2.1].
- (iii) \overline{G}_0 is generated by $\exp(\mathfrak{g})$ [24, Proposition 2]. In particular $\exp(tX)$ belongs to \overline{G}_0 for any $t \in \mathbb{C}$ and any $X \in \mathfrak{g}$.

Proposition 4.4 [35, Lemma 2.4]. *Assume that H_k is an algebraic subgroup of D_k and $\pi_{k,l}(H_k) \subset H_l$ for any $k \geq l \geq 1$. Then $\varprojlim H_k$ is pro-algebraic.*

Let us remark that a pro-algebraic subgroup G of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$ can be expressed in more than one way in the form $\varprojlim H_k$ with the conditions of Proposition 4.4. Indeed, if $G = \varprojlim H_k$ then $G_k = \overline{\pi_k(G)}$ is included in H_k but they do not coincide in general.

4.2 | Construction of an infinitesimal generator

In this section we define infinitesimal generators of formal diffeomorphism and show that for any $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ there exists an index m such that F^m has an infinitesimal generator (Binyamini [5]). Before doing so, let us study the relation between the Lie algebra of $\langle F \rangle$ and the group $\langle F \rangle_0$.

Lemma 4.5. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$. Then $\overline{\langle F \rangle}_0 = \exp(\mathfrak{g})$ where \mathfrak{g} is the Lie algebra of $\overline{\langle F \rangle}$.*

Proof. Analogously as for linear algebraic groups, since $\langle F \rangle$ is abelian, its Zariski closure $\overline{\langle F \rangle}$ is also abelian [24, Lemma 1] and then \mathfrak{g} is an abelian Lie algebra [24, Proposition 3]. Since $\overline{\langle F \rangle}_0$ is generated by $\exp(\mathfrak{g})$ by Proposition 4.3, we have $\exp(\mathfrak{g}) \subset \overline{\langle F \rangle}_0$. Moreover, any element L of $\overline{\langle F \rangle}_0$ is of the form

$$L = \exp(X_1) \circ \dots \circ \exp(X_m) = \exp(X_1 + \dots + X_m)$$

where $X_1, \dots, X_m \in \mathfrak{g}$. The last equality holds since \mathfrak{g} is abelian. \square

Definition 4.6. Given a formal diffeomorphism $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$, a formal vector field $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ is an *infinitesimal generator* of F if X belongs to the Lie algebra of $\overline{\langle F \rangle}$ and $F = \exp(X)$.

Theorem 4.7 [5, Corollary 7]. *Consider $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$. There exists $m \in \mathbb{N}$ such that F^m has an infinitesimal generator.*

Proof. Since $\overline{\langle F \rangle}_0$ is a finite index normal subgroup of $\overline{\langle F \rangle}$ by Proposition 4.3, there exists $m \in \mathbb{N}$ such that $F^m \in \overline{\langle F \rangle}_0$. The result is a consequence of Lemma 4.5. \square

The existence of infinitesimal generator is well known for unipotent formal diffeomorphisms, see, for example, [15] or [25] for the one-dimensional case.

Lemma 4.8 [21, Theorem 3.17]. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ be a formal diffeomorphism whose linear part is unipotent, that is, $\text{spec}(D_0 F) = \{1\}$. Then there exists a unique nilpotent $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ (i.e., a formal vector field X with $\text{spec}(D_0 X) = \{0\}$) such that $F = \exp(X)$.*

Remark 4.9. The formal vector field X in Lemma 4.8 belongs to the Lie algebra of $\overline{\langle F \rangle}$ (see [24, Lemma 1]), that is, X is an infinitesimal generator of F .

4.3 | The index of embeddability

Given a formal diffeomorphism $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$, we define the *index of embeddability in a flow* of F as the minimum of the indexes $m \in \mathbb{N}$ such that F^m has an infinitesimal generator. We denote it by $m(F)$, or simply by m when F is implicit. Observe that if $\text{spec}(D_0 F) = \{1\}$ then $m(F) = 1$, by Remark 4.9. Note also that, by Lemma 4.5, $m(F)$ is the minimum m such that $F^m \in \overline{\langle F \rangle}_0$. The following remark allows to calculate the index of embeddability of F and its iterates.

Remark 4.10. Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ and $r \in \mathbb{Z}^*$. Observe that each coset $F^j \overline{\langle F^r \rangle}$ is a pro-algebraic set, that is, a projective limit of algebraic sets, since $\overline{\langle F^r \rangle}$ is pro-algebraic. Moreover, since $\overline{\langle F \rangle}$ is abelian, we have that the set $\cup_{j=0}^{r-1} F^j \overline{\langle F^r \rangle}$ is a pro-algebraic group and then $\overline{\langle F \rangle} = \cup_{j=0}^{r-1} F^j \overline{\langle F^r \rangle}$.

Therefore, $\overline{\langle F^r \rangle}$ is a finite index subgroup of $\overline{\langle F \rangle}$ and $\overline{\langle F \rangle} / \overline{\langle F^r \rangle}$ is a cyclic group generated by the class of F . Analogously $\overline{\langle F \rangle} / \overline{\langle F^r \rangle}_0$ is cyclic, we just need to replace $\cup_{j=0}^{r-1} F^j \overline{\langle F^r \rangle}$ with $\cup_{j=0}^{s-1} F^j \overline{\langle F^r \rangle}_0$ above where s satisfies $F^s \in \overline{\langle F^r \rangle}_0$. In particular, we obtain $m(F) = |\overline{\langle F \rangle} : \overline{\langle F \rangle}_0|$.

It is clear that $\overline{\langle F^r \rangle}_0 \subset \overline{\langle F \rangle}_0$. Since $\overline{\langle F^r \rangle}$ is a finite index subgroup of $\overline{\langle F \rangle}$ and $\overline{\langle F^r \rangle}_0$ is a finite index subgroup of $\overline{\langle F^r \rangle}$ (Proposition 4.3), we deduce that $\overline{\langle F^r \rangle}_0$ is a finite index subgroup of $\overline{\langle F \rangle}$. This implies $\overline{\langle F \rangle}_0 \subset \overline{\langle F^r \rangle}_0$ by Proposition 4.3 and hence $\overline{\langle F \rangle}_0 = \overline{\langle F^r \rangle}_0$.

By definition $m(F^r)$ is the smallest positive integer m' such that $F^{rm'} \in \overline{\langle F^r \rangle}_0 = \overline{\langle F \rangle}_0$. In particular, we obtain $m(F^r) = m(F) / \gcd(m(F), r)$ and F^r has an infinitesimal generator if and only if $m(F)$ divides r .

On the other hand, the construction of $\overline{\langle F \rangle}$ implies that the group $D_0 \overline{\langle F \rangle}$ of linear parts of the elements of $\overline{\langle F \rangle}$ is equal to $\overline{\langle D_0 F \rangle}$, where the Zariski closure of the group $\langle D_0 F \rangle$ is considered in $\text{GL}(n, \mathbb{C})$, and we have

$$|\overline{\langle F \rangle} : \overline{\langle F \rangle}_0| = |\overline{\langle D_0 F \rangle} : \overline{\langle D_0 F \rangle}_0|,$$

(see [35, Proposition 2.3]). As a consequence, the value of $m(F) = |\overline{\langle D_0 F \rangle} : \overline{\langle D_0 F \rangle}_0|$ depends only on $D_0 F$. We recall the computation of $|\overline{\langle D_0 F \rangle} : \overline{\langle D_0 F \rangle}_0|$ for the sake of completeness in order to see that it depends only on the eigenvalues of $D_0 F$.

Given a matrix $A \in \text{GL}(n, \mathbb{C})$, it admits a unique multiplicative Jordan decomposition of the form $A = A_s A_u = A_u A_s$ where $A_s, A_u \in \text{GL}(n, \mathbb{C})$, A_s is diagonalizable and A_u is unipotent, that is, $\text{spec}(A_u) = \{1\}$. Since $\overline{\langle A \rangle}$ is isomorphic to $\overline{\langle A_s \rangle} \times \overline{\langle A_u \rangle}$ and $\overline{\langle A_u \rangle}$ is always connected, we get

$$\overline{\langle A \rangle} = \overline{\langle A \rangle}_0 \Leftrightarrow \overline{\langle A_s \rangle} = \overline{\langle A_s \rangle}_0.$$

Up to a linear change of coordinates, we can suppose $A_s = \text{diag}(\lambda_1, \dots, \lambda_n)$, where $\text{spec}(A) = \{\lambda_1, \dots, \lambda_n\}$. Denote $G_\lambda = \langle \text{diag}(\lambda_1, \dots, \lambda_n) \rangle$. Consider the characters $\chi_{(m_1, \dots, m_n)} : (\mathbb{C}^*)^n \rightarrow \mathbb{C}^*$ of the torus $(\mathbb{C}^*)^n$ given by $\chi_{(m_1, \dots, m_n)}(\mu_1, \dots, \mu_n) = \mu_1^{m_1} \dots \mu_n^{m_n}$ for $(m_1, \dots, m_n) \in \mathbb{Z}^n$. There exists a one-to-one correspondence between algebraic subgroups of $(\mathbb{C}^*)^n$ and subgroups of characters (see [27, Theorem 3.2.3.5]). Indeed given a subgroup H of the group of characters, the set $G = \cap_{\chi \in H} \text{Ker}(\chi)$ is an algebraic subgroup of $(\mathbb{C}^*)^n$ such that $H = \{\chi : G \subset \text{Ker}(\chi)\}$. As a consequence of the definition of the correspondence, we deduce

$$\overline{G}_\lambda = \{\text{diag}(\mu_1, \dots, \mu_n) : \mu_1^{m_1} \dots \mu_n^{m_n} = 1 \ \forall (m_1, \dots, m_n) \in S_\lambda\}$$

where

$$S_\lambda = \{(m_1, \dots, m_n) \in \mathbb{Z}^n : \lambda_1^{m_1} \dots \lambda_n^{m_n} = 1\} \quad (4)$$

is the set of resonances of $D_0 F$.

The Lie algebra $\mathfrak{g}_\lambda = \{\text{diag}(v_1, \dots, v_n) : \text{diag}(e^{v_1}, \dots, e^{v_n}) \in \overline{G}_\lambda\}$ of \overline{G}_λ is given by

$$\mathfrak{g}_\lambda = \{\text{diag}(v_1, \dots, v_n) : m_1 v_1 + \dots + m_n v_n = 0 \ \forall (m_1, \dots, m_n) \in S_\lambda\}. \quad (5)$$

Let S'_λ be the intersection of \mathbb{Z}^n with the \mathbb{Q} -vector space generated by S_λ . Notice that we can replace S_λ with S'_λ in Equation (5). Since $\overline{G}_{\lambda,0} = \exp(\mathfrak{g}_\lambda)$, we have

$$\overline{G}_{\lambda,0} = \{\text{diag}(\mu_1, \dots, \mu_n) : \mu_1^{m_1} \dots \mu_n^{m_n} = 1 \ \forall (m_1, \dots, m_n) \in S'_\lambda\}.$$

By construction, we get

$$S'_\lambda = \{(m_1, \dots, m_n) \in \mathbb{Z}^n : \lambda_1^{m_1} \dots \lambda_n^{m_n} \text{ is a root of unity}\}.$$

Moreover, $R_\lambda = \{\lambda_1^{m_1} \dots \lambda_n^{m_n} : (m_1, \dots, m_n) \in S'_\lambda\}$ is a finite subgroup of \mathbb{S}^1 , S_λ is a finite index subgroup of S'_λ , and

$$|\overline{G}_\lambda : \overline{G}_{\lambda,0}| = |S'_\lambda : S_\lambda| = |R_\lambda|.$$

Hence, we obtain $m(F) = |S'_\lambda : S_\lambda| = |R_\lambda|$ if $\text{spec}(D_0 F) = \{\lambda_1, \dots, \lambda_n\}$.

Remark 4.11. Consider $F \in \text{Diff}(\mathbb{C}^n, 0)$ and let m be the index of embeddability of F . Since the group R_λ associated to F^m is the trivial group, every eigenvalue of $D_0 F^m$ that is a root of unity is indeed equal to 1.

Remark 4.12. Assume that $\text{spec}(D_0 F) = \{\lambda_1, \dots, \lambda_n\}$ and $S_\lambda = \{0\}$, that is, the eigenvalues have no resonances. Then $m(F) = 1$.

4.4 | Characterization of the infinitesimal generator

It is not clear, from Definition 4.6, whether a formal vector field X such that $F = \exp(X)$ is an infinitesimal generator of F . Let us show that infinitesimal generators are determined by their linear parts.

Proposition 4.13. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$. Then*

- (i) $X \in \widehat{\mathfrak{X}}(\mathbb{C}^n, 0)$ is an infinitesimal generator of F if and only if $F = \exp(X)$ and $D_0 X$ is in the Lie algebra of the matrix group $\overline{\langle D_0 F \rangle}$.
- (ii) There is a bijective correspondence $X \mapsto D_0 X$ between infinitesimal generators of F and matrices M in the Lie algebra of $\overline{\langle D_0 F \rangle}$ such that $\exp(M) = D_0 F$.

Proof. We denote by G the group $\langle F \rangle$ and by \mathfrak{g} the Lie algebra of \overline{G} , which is abelian by [24, Proposition 3]. Let us remark that G_1 (see Definition 4.1) can be identified with $\overline{\langle D_0 F \rangle}$ and hence \mathfrak{g}_1 can be identified with the Lie algebra of $\overline{\langle D_0 F \rangle}$.

Consider $M \in \text{GL}(n, \mathbb{C})$ such that $M \in \mathfrak{g}_1$ and $\exp(M) = D_0 F$. Let us show that there exists $X \in \mathfrak{g}$ such that $\exp(X) = F$ and $D_0 X = M$. Since $\pi_{k,l} : \pi_k(G) \rightarrow \pi_l(G)$ is surjective, so is $\pi_{k,l} :$

$G_k \rightarrow G_l$ and hence $(d\pi_{k,l})_{\text{Id}} : \mathfrak{g}_k \rightarrow \mathfrak{g}_l$ is also surjective for all $k \geq l$. As a consequence, the map $d\pi_k : \mathfrak{g} \rightarrow \mathfrak{g}_k$ is surjective for any $k \geq 1$. Thus there exists $Y \in \mathfrak{g}$ such that $D_0 Y = M$. The formal diffeomorphism $F' = F \circ \exp(-Y)$ is clearly tangent to the identity and belongs to \overline{G} . There exists a unique formal vector field Z with vanishing linear part such that $F' = \exp(Z)$ by Lemma 4.8. Notice that Z belongs to the Lie algebra of $\langle F' \rangle$ by Remark 4.9 and hence to \mathfrak{g} since $\langle F' \rangle \subset \langle F \rangle$. Since \mathfrak{g} is abelian, it follows that

$$F = \exp(Z) \circ \exp(Y) = \exp(Y + Z)$$

where $Y + Z \in \mathfrak{g}$ and $D_0(Y + Z) = D_0 Y = M$. Therefore the correspondence defined in (ii) is surjective.

We claim that if $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$, $Y \in \mathfrak{g}$, $D_0 X = D_0 Y$, and $\exp(X) = \exp(Y) = F$ then $X = Y$. The claim implies that the correspondence in (ii) is injective.

Assume the claim is proven. The necessary condition in property (i) is clear. Let us show the sufficient condition in (i). Since the correspondence in (ii) is surjective, there exists $Y \in \mathfrak{g}$ such that $D_0 Y = D_0 X$ and $F = \exp(Y)$. Now the claim implies $X = Y$ and hence X belongs to \mathfrak{g} and is an infinitesimal generator of F .

Let us show the claim. Since $F = \exp(X)$, we get $F_* X = X$. We define

$$H_k = \{L \in D_k : J_k(L_* X) = J_k X\}.$$

It is an algebraic subgroup of D_k since the condition $J_k(L_* X) = J_k X$ can be expressed as a finite number of algebraic equations in the coefficients of the Taylor series expansion of L of degree less than or equal to k for any $k \geq 1$. Clearly, $\pi_{k,l}(H_k) \subset H_l$ is satisfied for all $k \geq l \geq 1$. We have then that the group

$$H = \varprojlim H_k = \{L \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : L_* X = X\}$$

is a pro-algebraic subgroup of $\widehat{\text{Diff}}(\mathbb{C}^n, 0)$ by Proposition 4.4. Notice that G is a subgroup of H and thus \overline{G} is also a subgroup of H . Since $Y \in \mathfrak{g}$, we obtain $\exp(tY) \in \overline{G} \subset H$ by Proposition 4.3 and hence $\exp(tY)_* X = X$ for any $t \in \mathbb{C}$. This implies $[X, Y] = 0$. We have

$$\text{Id} = F \circ F^{-1} = \exp(X) \circ \exp(-Y) = \exp(X - Y)$$

where we used $[X, Y] = 0$ in the last equality. Since $D_0(X - Y) = 0$, the vector field $X - Y$ is the unique nilpotent vector field whose exponential is the identity map, that is, $X - Y = 0$. \square

Definition 4.14. Let $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ with $\text{spec}(D_0 X) = \{\mu_1, \dots, \mu_n\}$. We say that X is *not weakly resonant* if there is no $(m_1, \dots, m_n) \in \mathbb{Z}^n$ such that $\sum_{j=1}^n m_j \mu_j \in 2\pi i \mathbb{Q}^*$.

Similar, but slightly different, conditions of absence of weak resonances have appeared in the literature when trying to solve the equation $\exp(X) = F$ where F and $D_0 X$ are fixed ([33, 41]). Next, we characterize the formal vector fields that are infinitesimal generators.

Theorem 4.15. Let $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ with $\text{spec}(D_0 X) = \{v_1, \dots, v_n\}$. Then X is an infinitesimal generator of $\exp(X)$ if and only if X is not weakly resonant.

Proof. Let $F = \exp(X)$. The formal vector field X is an infinitesimal generator of F if and only if D_0X is in the Lie algebra of $\overline{\langle D_0F \rangle}$ by Proposition 4.13. Let $D_0X = S + N$ be the additive Jordan decomposition of D_0X as a sum of a semisimple and a nilpotent linear operators that commute. Assume that S is diagonal up to a change of basis. Notice that $D_0F = \exp(S)\exp(N)$ is the multiplicative Jordan decomposition of D_0F . Denote by \mathfrak{g} , \mathfrak{g}_S , and \mathfrak{g}_N the Lie algebras of $\langle D_0F \rangle$, $\langle \exp(S) \rangle$, and $\langle \exp(N) \rangle$, respectively. It is well known that $\mathfrak{g} = \mathfrak{g}_S \oplus \mathfrak{g}_N$ and \mathfrak{g}_N is the complex vector space generated by N [27, Sections 3.2.2, 3.2.4 and 3.3.7]. Moreover, since all the elements of \mathfrak{g}_S (respectively \mathfrak{g}_N) are semisimple (respectively nilpotent), \mathfrak{g} is abelian, and the additive Jordan decomposition is unique, we deduce that \mathfrak{g}_S (respectively \mathfrak{g}_N) is the set of semisimple (respectively nilpotent) elements of \mathfrak{g} . As a consequence, D_0X is in the Lie algebra of $\langle D_0F \rangle$ if and only if S is in the Lie algebra of $\langle \exp(S) \rangle$. Notice that

$$S = \text{diag}(v_1, \dots, v_n) \text{ and } \exp(S) = \text{diag}(\lambda_1, \dots, \lambda_n),$$

where $\lambda_j = e^{v_j}$ for $1 \leq j \leq n$. Since $\lambda_1^{m_1} \dots \lambda_n^{m_n} = e^{m_1 v_1 + \dots + m_n v_n}$, we deduce

$$S_\lambda = \{(m_1, \dots, m_n) \in \mathbb{Z}^n : m_1 v_1 + \dots + m_n v_n \in 2\pi i \mathbb{Z}\}$$

where S_λ is defined in Equation (4). Recall that $\mathfrak{g}_S = \mathfrak{g}_\lambda$ is given by Equation (5). It is clear that if X is not weakly resonant then $S \in \mathfrak{g}_S$.

Suppose $S \in \mathfrak{g}_S$. Let (m_1, \dots, m_n) with $\sum_{j=1}^n m_j v_j \in 2\pi i \mathbb{Q}$. Up to multiplication by a non-zero integer, we can assume $\sum_{j=1}^n m_j v_j \in 2\pi i \mathbb{Z}$. This implies $\lambda_1^{m_1} \dots \lambda_n^{m_n} = 1$ and hence $(m_1, \dots, m_n) \in S_\lambda$. We deduce that $\sum_{j=1}^n m_j v_j = 0$ by the description of \mathfrak{g}_S . \square

In the following proposition we prove that the infinitesimal generator is unique only in the unipotent case. This is the reason justifying why infinitesimal generators have been considered exclusively for unipotent diffeomorphisms in the literature.

Proposition 4.16. *Consider $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ such that $m(F) = 1$. The infinitesimal generator of F is unique if and only if F is unipotent. Otherwise, F has infinitely many infinitesimal generators.*

Proof. The uniqueness of the infinitesimal generator is equivalent to the uniqueness of the infinitesimal generator of $\text{diag}(\lambda_1, \dots, \lambda_n)$ in the group of diagonal matrices where $\text{spec}(D_0F) = \{\lambda_1, \dots, \lambda_n\}$. Let us recall that the conditions $m(F) = 1$, $\overline{\langle F \rangle} = \overline{\langle F \rangle}_0$ and $S_\lambda = S'_\lambda$ are equivalent (see Section 4.3). Given an infinitesimal generator $\text{diag}(\mu_1, \dots, \mu_n)$ of $\text{diag}(\lambda_1, \dots, \lambda_n)$, the infinitesimal generators of $\text{diag}(\lambda_1, \dots, \lambda_n)$ are the matrices of the form $\text{diag}(\mu_1 + 2\pi i k_1, \dots, \mu_n + 2\pi i k_n)$ where k_1, \dots, k_n are integer numbers such that $m_1 k_1 + \dots + m_n k_n = 0$ for any $(m_1, \dots, m_n) \in S'_\lambda$. The system of equations has either a unique solution, if the rank of S'_λ is equal to n , or infinitely many solutions otherwise. Notice that the rank of S'_λ is equal to n if and only if $S'_\lambda = \mathbb{Z}^n$ and this condition is equivalent to $\lambda_1 = \dots = \lambda_n = 1$ since $S_\lambda = S'_\lambda$. As a consequence F has a unique infinitesimal generator if F is unipotent and infinitely many infinitesimal generators otherwise. \square

4.5 | Geometrical properties of the infinitesimal generator

In this section, we prove that if a diffeomorphism F has an infinitesimal generator X , then F and X have the same formal analytic invariant sets (ideals). Recall that an ideal $I \subset \hat{\mathcal{O}}_n$ is invariant for a

diffeomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ if $I \circ F \subset I$, and is invariant for a formal vector field $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ if $X(I) \subset I$.

Proposition 4.17. *Given an ideal $I \subset \hat{\mathcal{O}}_n$, the set*

$$\mathcal{I}_I = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : I \circ F \subset I\}$$

is a pro-algebraic group. Moreover, it satisfies $\mathcal{I}_I = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : I \circ F = I\}$.

Proof. In order to show that \mathcal{I}_I is pro-algebraic, we will use Proposition 4.4. The ideal I is of the form $I = (f_1, \dots, f_m)$ since $\mathbb{C}[[x_1, \dots, x_n]]$ is noetherian. We define

$$H_k = \{F \in D_k : f_j \circ F \in I + \mathfrak{m}^{k+1} \ \forall 1 \leq j \leq m\}.$$

It is clear that $\pi_{k,l}(H_k) \subset H_l$ for any $k \geq l \geq 1$. The inclusion $\mathcal{I}_I \subset \varprojlim H_k$ is obvious. Any element F of D_k can be interpreted as the k th jet of a local diffeomorphism. Hence, the power series $f_j \circ F$ is of the form $\sum_{i_1, \dots, i_n} a_{i_1, \dots, i_n}^j x_1^{i_1} \dots x_n^{i_n}$ where every a_{i_1, \dots, i_n}^j is a polynomial in the coefficients of the Taylor series expansion of F at the origin. The condition $f_j \circ F \in I + \mathfrak{m}^{k+1}$ is satisfied if and only if $\sum_{i_1 + \dots + i_n \leq k} a_{i_1, \dots, i_n}^j x_1^{i_1} \dots x_n^{i_n}$ belongs to the complex vector space V_k generated by the polynomials of the form

$$J_k(x_1^{i_1} \dots x_n^{i_n} f_j) \text{ where } i_1 + \dots + i_n \leq k \text{ and } 1 \leq j \leq m.$$

The property $J_k(f_j \circ F) \in V_k$ is equivalent to a linear system of equations on the coefficients a_{i_1, \dots, i_n}^j with $i_1 + \dots + i_n \leq k$ by elementary linear algebra. Therefore the condition $J_k(f_j \circ F) \in V_k$ is equivalent to a system of polynomial equations in the coefficients of F . Hence H_k is an algebraic subset of D_k .

We denote by I_k the natural projection of I in $\mathfrak{m}/\mathfrak{m}^{k+1}$. The definition of H_k implies

$$H_k = \{F \in D_k : F(I_k) \subset I_k\},$$

(see Equation (3)). Since F defines an element of $\text{GL}(\mathfrak{m}/\mathfrak{m}^{k+1})$ and $\mathfrak{m}/\mathfrak{m}^{k+1}$ is finite dimensional, it follows that

$$H_k = \{F \in D_k : F(I_k) = I_k\}.$$

As a consequence H_k is a subgroup of D_k and hence H_k is an algebraic subgroup of D_k for any $k \geq 1$.

Finally, let us show $\mathcal{I}_I = \varprojlim H_k$. By definition, we have

$$\varprojlim H_k = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : I \circ F + \mathfrak{m}^{k+1} = I + \mathfrak{m}^{k+1} \ \forall k \geq 1\}.$$

Since $\cap_{k=1}^{\infty} (J + \mathfrak{m}^{k+1}) = J$ for any proper ideal J of $\mathbb{C}[[x_1, \dots, x_n]]$ by Krull's intersection theorem, we deduce $\varprojlim H_k = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : I \circ F = I\}$ and hence $\varprojlim H_k \subset \mathcal{I}_I$. Since $\mathcal{I}_I \subset \varprojlim H_k$, we obtain $\varprojlim H_k = \mathcal{I}_I$. Therefore \mathcal{I}_I is a pro-algebraic group by Proposition 4.4. \square

Proposition 4.18. *Given a formal curve Γ , the group*

$$\mathcal{I}'_{\Gamma} = \{F \in \mathcal{I}_{\Gamma} : (F|_{\Gamma})'(0) = 1\}$$

is pro-algebraic.

Proof. Let $\gamma(t) = (\gamma_1(t), \dots, \gamma_n(t))$ be an irreducible parameterization of Γ . We denote $J_{\nu}\gamma(t) = (\mu_1 t^{\nu}, \dots, \mu_n t^{\nu})$ where ν is the multiplicity of Γ . We define the auxiliary group

$$\mathcal{J} = \{F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0) : (\mu_1, \dots, \mu_n) \in \ker(D_0 F - \text{Id})\}$$

that is clearly pro-algebraic. It is straightforward to check out the inclusion $\mathcal{I}'_{\Gamma} \subset \mathcal{J}$. Since the intersection of pro-algebraic groups is pro-algebraic [35, Remark 2.7], we obtain that $\hat{\mathcal{I}}_{\Gamma} = \mathcal{I}_{\Gamma} \cap \mathcal{J}$ is a pro-algebraic group containing \mathcal{I}'_{Γ} . In order to show that \mathcal{I}'_{Γ} is pro-algebraic, consider the morphism of groups

$$\begin{aligned} A &: \hat{\mathcal{I}}_{\Gamma} & \rightarrow & \mathbb{C}^* \\ F &\mapsto (F|_{\Gamma})'(0). \end{aligned}$$

Notice that the tangent value $\lambda(\Gamma)$ is equal to 1 for any element of $\hat{\mathcal{I}}_{\Gamma}$. Using the relation between the tangent and the inner eigenvalues given in Equation (2), we deduce that the image of A is contained in the group of roots of unity of order ν . Therefore \mathcal{I}'_{Γ} is a finite index subgroup of $\hat{\mathcal{I}}_{\Gamma}$ and hence pro-algebraic by Proposition 4.3. \square

The next results are consequences of the above proposition and the general properties of pro-algebraic groups.

Proposition 4.19. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ and let $I \subset \hat{\mathcal{O}}_n$ be an ideal. Suppose that there exists $r \in \mathbb{Z}^*$ such that I is invariant for F^r . Then $\overline{\langle F \rangle}_0 \subset \mathcal{I}_I$. Moreover, if $I = \Gamma$ is a formal curve and $(F^r|_{\Gamma})'(0)$ is a root of unity then $\overline{\langle F \rangle}_0 \subset \mathcal{I}'_{\Gamma}$.*

Proof. Since \mathcal{I}_I is pro-algebraic, we obtain $\overline{\langle F^r \rangle} \subset \mathcal{I}_I$. Since $\overline{\langle F \rangle}_0 = \overline{\langle F^r \rangle}_0$ by Remark 4.10, we get $\overline{\langle F \rangle}_0 \subset \mathcal{I}_I$. If $I = \Gamma$ is a formal curve and $(F^r|_{\Gamma})'(0)$ is a root of unity then we can replace r with a multiple to obtain $(F^r|_{\Gamma})'(0) = 1$. The same proof shows $\overline{\langle F \rangle}_0 \subset \mathcal{I}'_{\Gamma}$. \square

Proposition 4.20. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ and let m be the index of embeddability of F . Let X be an infinitesimal generator of F^m . Given an ideal I of $\hat{\mathcal{O}}_n$, the following properties are equivalent:*

- (1) *I is invariant for X ;*
- (2) *I is invariant for F^m ;*
- (3) *I is invariant for a non-trivial iterate of F .*

Proof. The implications $(1) \Rightarrow (2)$ and $(2) \Rightarrow (3)$ are clear. Assume that (3) holds. We have $\exp(tX) \in \overline{\langle F \rangle}_0$ for any $t \in \mathbb{C}$ by Proposition 4.3. Since $\overline{\langle F \rangle}_0 \subset \mathcal{I}_I$ by Proposition 4.19, we obtain $\exp(tX) \in \mathcal{I}_I$ for any $t \in \mathbb{C}$. Thus I is invariant for X . \square

As a consequence of Proposition 4.20, we recover the following result of Ribón:

Corollary 4.21 [32]. *Let $F \in \text{Diff}(\mathbb{C}^2, 0)$ and let $m = m(F)$ be its index of embeddability. Then there exists a formal m -periodic curve of F .*

Proof. The diffeomorphism F^m has an infinitesimal generator X . The formal version of Camacho–Sad’s theorem [7] provides a formal invariant curve Γ that is invariant by X . Thus Γ is invariant by F^m . \square

Proposition 4.22. *Let $F \in \widehat{\text{Diff}}(\mathbb{C}^n, 0)$ and let Γ be a formal curve. Suppose that there exists $r \in \mathbb{Z}^*$ such that Γ is invariant for F^r and $(F^r|_{\Gamma})'(0)$ is a root of unity. Let X be an infinitesimal generator of F^m , where m is the index of embeddability of F . Then the inner eigenvalue of (X, Γ) is equal to 0. In particular, the tangent line of Γ is contained in the kernel of the linear part D_0X of X at the origin.*

Proof. Proposition 4.19 implies $\overline{\langle F \rangle}_0 \subset \mathcal{I}'_{\Gamma}$ and hence $\exp(tX) \in \mathcal{I}'_{\Gamma}$ for any $t \in \mathbb{C}$, that is, the inner eigenvalue of the pair $(\exp(tX), \Gamma)$ is equal to 1 for any t . Let $\gamma(s)$ be an irreducible parameterization of Γ . We have that $\exp(tX)(\gamma(s))$ is of the form $\gamma \circ \phi_t(s)$ where $\phi_t(0) = 0$ and $\phi'_t(0) = 1$ for any $t \in \mathbb{C}$. Since $X(\gamma(s)) = \frac{\partial \gamma \circ \phi_t(s)}{\partial t}|_{t=0}$, it follows that the multiplicity of the right-hand side is at least $\nu(\gamma) + 1$ and thus $\nu(X|_{\Gamma}) = \nu(X(\gamma(s))) - \nu(\gamma'(s)) \geq 2$. Since $\nu(X|_{\Gamma}) > \nu(\gamma(s))$, any non-zero tangent vector v of Γ at 0 is in the kernel of D_0X . \square

4.6 | Examples of diffeomorphisms possessing a formal invariant curve

Corollary 4.21 provides a formal periodic curve for any $F \in \text{Diff}(\mathbb{C}^2, 0)$. This is no longer true for dimension greater than 2. More precisely, there exist nilpotent analytic vector fields $X \in \mathfrak{X}(\mathbb{C}^3, 0)$ with no formal invariant curve by a theorem of Gómez-Mont and Luengo [16]. Then the diffeomorphism $\exp(X)$ has no formal periodic curve by Proposition 4.20.

In this section we apply our results about infinitesimal generators to obtain conditions that guarantee the existence of a formal periodic curve in dimension $n = 3$.

A formal codimension 1 foliation \mathcal{F}_{ω} in $(\mathbb{C}^3, 0)$ is determined by a non-zero 1-form

$$\omega = a_1 dx_1 + a_2 dx_2 + a_3 dx_3, \quad a_1, a_2, a_3 \in \hat{\mathcal{O}}_3,$$

satisfying the integrability condition $\omega \wedge d\omega = 0$. Two 1-forms ω and ω' define the same foliation if there exists $f \in \hat{\mathcal{K}}_3 \setminus \{0\}$ such that $\omega = f\omega'$ where $\hat{\mathcal{K}}_3$ is the field of fractions of $\hat{\mathcal{O}}_3$. We say that \mathcal{F}_{ω} has a formal integrating factor if there exists $f \in \hat{\mathcal{K}}_3 \setminus \{0\}$ such that $d(\frac{\omega}{f}) = 0$.

Proposition 4.23. *Let $F \in \text{Diff}(\mathbb{C}^3, 0)$. Suppose that either*

- (1) *there exists a foliation \mathcal{F}_{ω} with no formal integrating factor such that $F^* \omega \wedge \omega = 0$ or*
- (2) *there exists $g \in \hat{\mathcal{O}}_3 \setminus \mathbb{C}$ such that $g \circ F = g$.*

Then F^m has a formal invariant curve, where m is the index of embeddability of F .

The two cases are of different nature. Namely, in case (1) we are requiring that F preserves a foliation with “poor” integrability properties whereas in case (2) we are asking F to preserve the “fibers” of g .

Lemma 4.24. *Suppose that the hypotheses of Proposition 4.23 are satisfied. Set $\omega = dg$ in case (2). Then $\omega(X) = 0$ for any infinitesimal generator X of F^m .*

Proof. Assume that we are in case (1). Analogously as in Proposition 4.17, we can show that the group

$$\mathcal{I}_\omega = \{L \in \widehat{\text{Diff}}(\mathbb{C}^3, 0) : L^* \omega \wedge \omega = 0\}$$

is of the form $\varprojlim H_k$, where

$$H_k = \{L \in D_k : J_k(L^* \omega) = J_k(h_k \omega) \text{ for some } h_k \in \hat{\mathcal{O}}_3\}$$

is an algebraic subgroup of D_k and $\pi_{k,l}(H_k) \subset H_l$ for all $k \geq l \geq 1$. Therefore \mathcal{I}_ω is pro-algebraic by Proposition 4.4. We deduce that $\overline{\langle F \rangle}$ is contained in \mathcal{I}_ω and hence $\exp(tX)^* \omega \wedge \omega = 0$ for any infinitesimal generator X of F^m and any $t \in \mathbb{C}$ by Proposition 4.3. As a consequence the Lie derivative $L_X \omega$ satisfies $L_X \omega \wedge \omega = 0$. This implies either that $\omega(X) = 0$ or that $\omega(X)$ is a formal integrating factor of ω (see [12, Chapitre III, Proposition 1.3]). Since the latter possibility is excluded by hypothesis, we obtain $\omega(X) = 0$.

Assume that we are in case (2). The group

$$\mathcal{I}_g = \{L \in \widehat{\text{Diff}}(\mathbb{C}^3, 0) : g \circ L = g\}$$

is of the form $\varprojlim H_k$, where $H_k = \{L \in D_k : J_k(g \circ L) = J_k(g)\}$ is an algebraic subgroup of D_k and $\pi_{k,l}(H_k) \subset H_l$ for all $k \geq l \geq 1$. Arguing as in the previous case, we obtain $g \circ \exp(tX) = g$ for any infinitesimal generator X of F^m and any $t \in \mathbb{C}$. We get $\omega(X) = dg(X) = X(g) = 0$. \square

Proposition 4.23 is an immediate consequence of Lemma 4.24 and next result.

Proposition 4.25. *Let $X \in \hat{\mathfrak{X}}(\mathbb{C}^3, 0)$. Consider a formal codimension 1 foliation \mathcal{F}_ω such that $\omega(X) = 0$. Then X has a formal invariant curve.*

This result is due to Cerveau and Lins [11, Proposition 3] for holomorphic foliations and vector fields. We just adapt their proof to the formal setting.

Proof. If α is a formal differential form or a formal vector field, we denote by $\text{Sing}(\alpha)$ the ideal of its coefficients in $\hat{\mathcal{O}}_3$. We can suppose that the coefficients of ω have no common factor up to divide ω by the gcd of such coefficients. In other words we have $\text{codim}(\text{Sing}(\omega)) \geq 2$. We can assume $\dim(\text{Sing}(X)) = 0$, otherwise the result is trivial. Moreover, this implies $\omega(0) = 0$ since otherwise the foliation \mathcal{F}_ω is equal to \mathcal{F}_{dx} up to a formal change of coordinates and $dx(X) = 0$ implies $X = b(x, y, z)\partial_y + c(x, y, z)\partial_z$ and then $\dim(\text{Sing}(X)) \geq 1$.

Denote $\eta = i_X(dx \wedge dy \wedge dz)$. The property $\omega(X) = 0$ is equivalent to $\omega \wedge \eta = 0$. We claim $\text{codim}(\text{Sing}(\omega)) \neq 3$, otherwise we can apply the de Rham–Saito lemma [36] to show that the

2-form η is of the form $\omega \wedge \theta$ (the result works both in the formal and analytic settings). We have

$$\text{Sing}(\omega \wedge \theta) = \text{Sing}(\eta) = \text{Sing}(X).$$

Hence $\dim(\text{Sing}(\omega \wedge \theta)) = 0$. There exists $k \in \mathbb{N}$ such that if ω', θ' are formal 1-forms such that $J_k \omega = J_k \omega'$ and $J_k \theta = J_k \theta'$ then $\dim(\text{Sing}(\omega' \wedge \theta')) = 0$. In particular, we get $\dim(\text{Sing}(J_k \omega \wedge J_k \theta)) = 0$. This provides a contradiction since it is known that the codimension of the singular set of the exterior product of two germs of holomorphic 1-form has codimension less than or equal to 2 if it is singular at $(0,0,0)$ (see [26, Lemma 3.1.2]).

We deduce $\text{codim}(\text{Sing}(\omega)) = 2$ and hence $\text{Sing}(\omega)$ is a formal curve. Let us remark that since $\omega(X) = 0$, the curve $\text{Sing}(\omega)$ is invariant by X . \square

5 | REDUCTION TO RAMIS–SIBUYA FORM

In this section, we show that a pair (F, Γ) , where F is a diffeomorphism and Γ is a rationally neutral formal invariant curve of F not contained in the set of fixed points of a non-trivial iterate of F , can be reduced, up to iterating F , to a pair $(\tilde{F}, \tilde{\Gamma})$ in Ramis–Sibuya form. First, we perform such a reduction in the context of formal vector fields. Next, we use the results in Section 4 to adapt the reduction to diffeomorphisms.

5.1 | Ramis–Sibuya form for formal vector fields

Definition 5.1. Let X be a singular formal vector field at $0 \in \mathbb{C}^n$ and let Γ be a formal invariant curve of X . We say that the pair (X, Γ) is in *Ramis–Sibuya form* (RS-form for short) if Γ is non-singular and there exist analytic coordinates (x, \mathbf{y}) at $0 \in \mathbb{C}^n$ for which Γ is transversal to the hyperplane $x = 0$ and such that X is written as

$$X = x^{q+1}(\lambda + bx^{\max(1,q)} + x^{q+1}A(x, \mathbf{y}))\frac{\partial}{\partial x} + ((D(x) + x^qC)\mathbf{y} + x^{q+1}B(x, \mathbf{y}))\frac{\partial}{\partial \mathbf{y}}, \quad (6)$$

where $q \geq 0$, $\lambda \in \mathbb{C}^*$, $b \in \mathbb{C}$, $A(x, \mathbf{y}) \in \mathbb{C}[[x, \mathbf{y}]]$, $B(x, \mathbf{y}) \in \mathbb{C}[[x, \mathbf{y}]]^{n-1}$ and

- (i) $D(x)$ is a diagonal matrix of polynomials of degree at most $q - 1$ (equal to 0 if $q = 0$) and C is a constant matrix,
- (ii) $D(x) + x^qC \neq 0$,
- (iii) $D(x)$ commutes with C .

The polynomial vector field $\lambda x^{q+1} \frac{\partial}{\partial x} + (D(x) + x^qC)\mathbf{y} \frac{\partial}{\partial \mathbf{y}}$ is called the *principal part* of (X, Γ) in the coordinates (x, \mathbf{y}) .

Notice that $q + 1$ is the multiplicity of the restricted vector field $X|_{\Gamma}$ and thus the integer $q = q(X, \Gamma)$ is well defined for the pair (X, Γ) and is independent of the coordinates. On the other hand, if the multiplicity of X is $\nu(X) = \nu + 1$ then $\nu \leq q$ and ν coincides with the order at $x = 0$ of the polynomial matrix $D(x) + x^qC$. Thus, the number $p = p(X, \Gamma) = q - \nu \geq 0$, called the *Poincaré rank* of the pair (X, Γ) , is also independent of the coordinates.

Remark 5.2. Assume that (X, Γ) is in RS-form, written as (6) in coordinates (x, \mathbf{y}) .

- (a) Γ is not contained in the singular locus of the vector field X .
- (b) Let $l \geq 1$ be the order of contact of Γ with the x -axis, that is, Γ admits a parameterization $(s, \bar{\gamma}(s)) \in \mathbb{C}[[s]]^n$ where the minimum order of the components of $\bar{\gamma}(s)$ is equal to l . Then the invariance condition implies that the order in x of any component of the vector $X(\mathbf{y})(x, 0) \in \mathbb{C}[[x]]^{n-1}$ is at least $l + \nu$.
- (c) If $q \geq 1$ then, after a change of variables of the form $\bar{x} = ax$ where $a^q = -\lambda$, we may assume that $\lambda = -1$.
- (d) Denote by $Q_1(x), \dots, Q_l(x)$ the different polynomials in the diagonal of the matrix $D(x)$ and, up to reordering the \mathbf{y} -variables, write

$$D(x) = \text{diag}(Q_1(x)I_{n_1}, \dots, Q_l(x)I_{n_l}).$$

The property $[D(x), C] = 0$ implies that C is block-diagonal $C = \text{diag}(C^1, \dots, C^l)$ where C^j has size n_j . After a linear change of variables of the form $\bar{\mathbf{y}} = P\mathbf{y}$, we may assume that the blocks of the matrix C are in Jordan canonical form.

Let us justify our choice of the terminology in Definition 5.1. After dividing the vector field in (6) by x^ν times a unit, we can associate it to a system of $n - 1$ formal ODEs

$$x^{p+1}\mathbf{y}' = (\bar{D}(x) + x^p \bar{C} + O(x^{p+1}))\mathbf{y} + O(x^{p+1}),$$

where $\bar{D}(x) = D(x)/\lambda x^\nu$ and \bar{C} is a constant matrix. Such a system has a singular point at $x = 0$ with *Poincaré rank* equal to p (unless possibly for $q = p = 0$ if $\bar{C} = 0$). Moreover, the properties assumed for the polynomial matrix $D(x) + x^p C$ are essentially those considered in the work of Ramis and Sibuya [31], devoted to proving *multisummability* of the formal solution $\mathbf{y} = \bar{\gamma}(x)$, where $(x, \bar{\gamma}(x))$ is a parameterization of Γ , in the case where the coefficients of the system are convergent.

5.2 | Blow-ups and ramifications along an invariant curve

Let $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ be a singular vector field and let Γ be a formal invariant curve of X not contained in the singular locus of X .

A germ of holomorphic map $\phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ will be called a *permissible transformation* for the pair (X, Γ) if it is of one of the following types:

1. The germ of a holomorphic diffeomorphism.
2. Let Z be a germ of non-singular analytic submanifold at $0 \in \mathbb{C}^n$ which is invariant for X (meaning that $X(g) \in I(Z)$ for any $g \in I(Z)$, where $I(Z)$ denotes the ideal of holomorphic germs vanishing on Z) and such that the tangent line of Γ is transversal to Z . Let $\pi_Z : M \rightarrow U$ be the blow-up with center Z and let $p \in \pi_Z^{-1}(0)$ be the point corresponding to the tangent of Γ . Then there is an analytic chart τ of M at p so that ϕ is the germ of $\pi_Z \tau^{-1}$ at $0 \in \mathbb{C}^n$. We will say that Z is a *permissible center* and that ϕ is a *permissible blow-up*.
3. The curve Γ is non-singular, there are analytic coordinates $\mathbf{z} = (z_1, \dots, z_n)$ at $0 \in \mathbb{C}^n$ such that $Z = \{z_1 = 0\}$ is invariant for X and transversal to Γ and ϕ is the map $\phi(\mathbf{z}) = (z_1^l, z_2, \dots, z_n)$ for some $l \in \mathbb{N}_{>0}$. We will say that ϕ is a *permissible l-ramification* (with center Z).

In the last two cases, the non-singular hypersurface $E_\phi = \phi^{-1}(Z)$ is called the *exceptional divisor* of ϕ . For convenience, $E_\phi = \{0\}$ in the case where ϕ is a diffeomorphism. Notice that a permissible transformation ϕ is a local diffeomorphism at every point in the complement of E_ϕ .

The following result is quite well known (see for instance [8] for the three-dimensional case). We include a proof for the sake of completeness.

Proposition 5.3. *Let $\phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ be a permissible transformation for (X, Γ) . There exist a unique formal curve $\tilde{\Gamma}$ at $0 \in \mathbb{C}^n$ such that $\phi^*\Gamma \subset \tilde{\Gamma}$ (where $\phi^*\Gamma = \{g \circ \phi : g \in \Gamma\}$) and a unique formal vector field \tilde{X} at $0 \in \mathbb{C}^n$ such that $\phi_*\tilde{X} = X$. Moreover, \tilde{X} is singular and has $\tilde{\Gamma}$ as an invariant curve. In addition, the multiplicities of the restrictions satisfy $\nu(\tilde{X}|_{\tilde{\Gamma}}) \geq \nu(X|_\Gamma)$. We will call \tilde{X} and $\tilde{\Gamma}$ the transforms of X and Γ by ϕ , respectively.*

Proof. The case where ϕ is a germ of a diffeomorphism is clear.

Suppose that ϕ is a permissible blow-up with center Z . Consider analytic coordinates $\mathbf{z} = (z_1, z_2, \dots, z_n)$ so that the tangent line of Γ is tangent to the z_1 -axis and $Z = \{z_1 = z_2 = \dots = z_t = 0\}$ where $t = \text{codim } Z$ (thus $I(Z)$ is generated by z_1, \dots, z_t). We may write $\phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ as

$$\phi(\mathbf{z}) = (z_1, z_1 z_2, \dots, z_1 z_t, z_{t+1}, \dots, z_n). \quad (7)$$

Let $\gamma(s) = (\gamma_1(s), \dots, \gamma_n(s)) \in \mathbb{C}[[s]]^n$ be an irreducible parameterization of Γ in the coordinates \mathbf{z} . Then $\nu(\Gamma) = \nu(\gamma_1(s)) < \nu(\gamma_j(s))$ for $j = 2, \dots, n$, where ν denotes the order in s . Also,

$$\tilde{\gamma}(s) = \left(\gamma_1(s), \frac{\gamma_2(s)}{\gamma_1(s)}, \dots, \frac{\gamma_t(s)}{\gamma_1(s)}, \gamma_{t+1}(s), \dots, \gamma_n(s) \right) \in \mathbb{C}[[s]]^n \quad (8)$$

is a parameterization of a formal curve $\tilde{\Gamma}$ which satisfies $\phi^*\Gamma \subset \tilde{\Gamma}$. The uniqueness of $\tilde{\Gamma}$ can be seen as follows: if $\tilde{\gamma}(s) = (\tilde{\gamma}_1(s), \tilde{\gamma}_2(s), \dots, \tilde{\gamma}_n(s))$ is a parameterization of another formal curve $\tilde{\Gamma}$ satisfying $\phi^*\Gamma \subset \tilde{\Gamma}$ then we will have that $\phi \circ \tilde{\gamma}(s)$ is another parameterization of Γ and necessarily $\phi \circ \tilde{\gamma}(s) = \gamma(s)$ where $\sigma(s) \in \mathbb{C}[[s]]$. Using the expression of ϕ and Equation (8) one shows that $\tilde{\gamma}(s) = \gamma(s)$ and thus $\tilde{\Gamma} = \Gamma$.

Write $X = \sum_{i=1}^n a_i(\mathbf{z}) \frac{\partial}{\partial z_i}$. Since Γ is invariant and not contained in the singular locus of X , we have that the vector $X|_{\gamma(s)} \in \mathbb{C}[[s]]^n$ is a non-zero multiple of $\gamma'(s)$ and hence $\nu(a_1(\gamma(s))) < \nu(a_j(\gamma(s)))$ for $j = 2, \dots, n$. So $a_j(\mathbf{z})$ cannot contain a monomial of the form cz_1 , with $c \neq 0$, for $j = 2, \dots, n$. On the other hand, the condition of Z being invariant implies that, for $j = 1, \dots, t$, $a_j(\mathbf{z}) \in I(Z)$ and hence $a_j(\phi(\mathbf{z}))$ is divisible by z_1 . Using these two properties, the vector field $\tilde{X} = \sum_{i=1}^n \tilde{a}_i(\mathbf{z}) \frac{\partial}{\partial z_i}$ defined by

$$\begin{cases} \tilde{a}_j(\mathbf{z}) = \frac{a_j(\phi(\mathbf{z})) - z_j a_1(\phi(\mathbf{z}))}{z_1}, & \text{for } j = 2, \dots, t; \\ [7pt] \tilde{a}_j(\mathbf{z}) = a_j(\phi(\mathbf{z})), & \text{for } j \in \{1, t+1, \dots, n\}, \end{cases} \quad (9)$$

is formal and singular at 0, and it is the unique that satisfies $\phi_*\tilde{X} = X$. Since Γ is invariant for X , we get $X|_{\gamma(s)} = h(s)\gamma'(s)$ for some $h(s) \in \mathbb{C}[[s]]$ and one obtains that $\tilde{X}|_{\tilde{\gamma}(s)} = h(s)\tilde{\gamma}'(s)$, proving that $\tilde{\Gamma}$ is invariant for X and that $\nu(\tilde{X}|_{\tilde{\Gamma}}) = \nu(X|_\Gamma)$.

Assume now that ϕ is a permissible l -ramification, written in some coordinates \mathbf{z} as $\phi(\mathbf{z}) = (z_1^l, z_2, \dots, z_n)$. Consider a parameterization of Γ of the form $\gamma(s) = (s, \gamma_2(s), \dots, \gamma_n(s))$ in these coordinates (recall that, from the definition of permissible ramification, Γ is non-singular). Then

$$\tilde{\gamma}(s) = (s, \gamma_2(s^l), \dots, \gamma_n(s^l)) \in \mathbb{C}[[s]]^n \quad (10)$$

is a parameterization of a formal curve $\tilde{\Gamma}$ satisfying $\phi^*\Gamma \subset \tilde{\Gamma}$. Uniqueness of $\tilde{\Gamma}$ comes from the property of Γ being non-singular: Γ is generated by the series $z_j - \gamma_j(z_1)$ for $j = 2, \dots, n$ and thus, if $\bar{\Gamma}$ is a formal curve such that $\phi^*\Gamma \subset \bar{\Gamma}$, then $\tilde{\gamma}(s)$ is a parameterization of $\bar{\Gamma}$.

On the other hand, being $Z = \{z_1 = 0\}$ invariant for X , if we write $X = \sum_{i=1}^n a_i(\mathbf{z}) \frac{\partial}{\partial z_i}$ then we have $a_1(\mathbf{z}) = z_1 \bar{a}_1(\mathbf{z})$, where $\bar{a}_1(\mathbf{z})$ is a formal series. The (singular) formal vector field \tilde{X} defined by

$$\tilde{X} = \frac{z_1 \bar{a}_1(\phi(\mathbf{z}))}{l} \frac{\partial}{\partial z_1} + \sum_{i=2}^n a_i(\phi(\mathbf{z})) \frac{\partial}{\partial z_i}$$

satisfies $\phi_* \tilde{X} = X$. Since Γ is invariant and not contained in the singular locus of X , $X|_{\gamma(s)} = h(s)\gamma'(s)$ for some $h \in \mathbb{C}[[s]]$ with $\nu(h) \geq 1$. We obtain that $\tilde{\Gamma}$ is invariant for \tilde{X} and

$$\nu(\tilde{X}|_{\tilde{\Gamma}}) = \nu(X|_{\Gamma}) + (\nu(X|_{\Gamma}) - 1)(l - 1) \geq \nu(X|_{\Gamma})$$

as a consequence of $\tilde{X}|_{\tilde{\gamma}(s)} = l^{-1}s^{1-l}h(s^l)\tilde{\gamma}'(s)$. □

It is worth to notice that Proposition 5.3 remains true, except for the uniqueness of the curve $\tilde{\Gamma}$ satisfying $\phi^*\Gamma \subset \tilde{\Gamma}$, if the condition of Γ being non-singular in the definition of permissible ramification is removed (consider, for example, the curve $\Gamma = (y^2 - x^3)$ at $(\mathbb{C}^2, 0)$ and $\phi(x, y) = (x^2, y)$ where we can choose $\tilde{\Gamma} = (y - x^3)$ or $\tilde{\Gamma} = (y + x^3)$).

Remark 5.4. Observe that the expression of the transform of a vector field by a permissible transformation is *finitely determined* in the following sense. Let ϕ be a permissible transformation for (X, Γ) with center Z . Then, for any $N \in \mathbb{N}$, there exists $N' \in \mathbb{N}$ such that, if Y is another formal vector field for which Z is invariant and $J_{N'}Y = J_{N'}X$ then $J_N\tilde{Y} = J_N\tilde{X}$, where \tilde{X}, \tilde{Y} are the transforms of X, Y by ϕ , respectively. Although we do not require Y to have Γ as an invariant curve, the transform \tilde{Y} is well defined in Proposition 5.3 once we have that the center Z of ϕ is invariant for Y .

5.3 | Reduction of a vector field to Ramis–Sibuya form

Let X be a formal singular vector field at $(\mathbb{C}^n, 0)$ and let Γ be a formal invariant curve of X not contained in the singular locus of X . In this section we show that the pair (X, Γ) can be reduced to Ramis–Sibuya form by permissible transformations.

A sequence of permissible transformations for (X, Γ) is a composition

$$\Phi = \phi_l \circ \phi_{l-1} \circ \dots \circ \phi_1 : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$$

such that ϕ_1 is a permissible transformation for (X, Γ) and, for $j = 1, \dots, l-1$, (X_j, Γ_j) is the transform of (X_{j-1}, Γ_{j-1}) by ϕ_j and ϕ_{j+1} is a permissible transformation for (X_j, Γ_j) . The last pair (X_l, Γ_l) will be called the *transform* of (X, Γ) by Φ . We also define the *total divisor* of Φ as the set $E_\Phi = (\phi_l \circ \phi_{l-1} \circ \dots \circ \phi_2)^{-1}(E_{\phi_1})$, which is a normal crossing divisor at $0 \in \mathbb{C}^n$.

Theorem 5.5. *Let X be a formal singular vector field at $0 \in \mathbb{C}^n$ and let Γ be an invariant formal curve of X not contained in the singular locus of X . Then there exists a sequence Φ of permissible transformations for (X, Γ) such that the transform of (X, Γ) by Φ is in Ramis–Sibuya form.*

A composition Φ as in the statement will be called a *reduction* of (X, Γ) to RS-form.

Theorem 5.5 is not a completely new result in the theory of reduction of singularities of vector fields or in the theory of systems of meromorphic ODEs with irregular singularity. It contains in particular a result of “local uniformization” of X (i.e., reduction to non-nilpotent linear part) along the valuation corresponding to Γ (see Cano et al. [8, 10] or Panazzolo [28] for more information). The particular expression of a vector field in RS-form, that requires more than a non-nilpotent linear part, is obtained, once we associate to X a system of $n-1$ meromorphic ODEs after some initial blow-ups, from classical results in the theory of ODEs, generically known as *Turrittin’s Theorem*: see Turrittin [37], Wasow [40], Balser [3], or Barkatou [4] (for linear systems with formal coefficients), and Cano et al. [9] (for related statements for three-dimensional real vector fields). Since we could not find a statement with the precise terms of Theorem 5.5 needed for our purposes, we devote the rest of this section to provide a self-contained proof.

Let us describe the situation after a punctual blow-up. Let (x, y) be coordinates such that $\{x = 0\}$ is transversal to (the tangent line of) Γ and let $\phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ be the blow-up of \mathbb{C}^n centered at the origin (which is permissible for (X, Γ)). There is a constant vector $\xi \in \mathbb{C}^{n-1}$ so that ϕ is written as

$$\phi(x, y) = (x, x(y - \xi)). \quad (11)$$

We obtain the following properties:

- (a1) The transform of X by ϕ is written as $\tilde{X} = x^{\nu(X)-1}X'$ where $E_\phi = \{x = 0\}$ and X' is a formal singular vector field. Thus $\nu(\tilde{X}) \geq \nu(X)$ and the origin is again a permissible center for the transform $(\tilde{X}, \tilde{\Gamma})$ of (X, Γ) .
- (a2) When $\xi = 0$, the exponent of x increases at least a unit in any monomial of the coefficient $\tilde{X}(x)$ with positive degree in the y -variables and in any monomial of the components of $\tilde{X}(y)$ with degree at least two in the y -variables, whereas the order of $\tilde{X}(y)(x, 0)$ decreases in a unit.

From pre-RS form to RS form

To prove Theorem 5.5, it will be sufficient to prove that there exists a sequence Ψ of permissible transformations for (X, Γ) such that the transform $\tilde{X} = \Psi^*X$ is written in some coordinates (x, y) as

$$\tilde{X} = x^{q+1}u(x, y) \frac{\partial}{\partial x} + (B_0(x) + (D(x) + x^q C)y + O(x^{q+1}y)) \frac{\partial}{\partial y}, \quad (12)$$

where $u(0,0) \neq 0$, $B_0(0) = 0$ and the transformed curve $\tilde{\Gamma} = \Psi^*\Gamma$, together with $q, D(x), C$ satisfy the conditions of Definition 5.1.

Let us see how we can obtain RS-form from expression (12). Analogously as in Remark 5.2(b), if $\gamma(x) = (x, \tilde{\gamma}(x))$ is a parameterization of $\tilde{\Gamma}$ and $\nu(\tilde{\gamma}(x)) \geq l$ then $\nu(B_0(x)) \geq l$. Thus, by a change of variables of the form $\mathbf{y} = \tilde{\mathbf{y}} + J_{2q+2}\tilde{\gamma}(x)$ we may assume that $\nu(B_0(x)) \geq 2q+3$ and the first $2q+2$ iterated tangents of $\tilde{\Gamma}$ and of $\{\mathbf{y} = 0\}$ coincide. Taking into account the properties (a1) and (a2) above about the effect of a permissible punctual blow-up, we have that the composition Φ of the blow-ups at the first $q+1$ iterated tangents of $\tilde{\Gamma}$ is written as $\Phi(x, \mathbf{y}) = (x, x^{q+1}\mathbf{y})$ and the transform $\Phi^*\tilde{X}$ is written as in (12) with the extra hypothesis

$$u(x, \mathbf{y}) = u(x, 0) + O(x^{q+1}\mathbf{y}), \quad \nu(B_0(x)) \geq q+2.$$

Notice that the matrix $D(x) + x^q C$ has changed into $D(x) + x^q(C - (q+1)u(0,0)I_{n-1})$. If this matrix vanishes, we consider $\Phi(x, \mathbf{y}) = (x, x^{q+2}\mathbf{y})$ and $\Phi^*\tilde{X}$ is in the form (12) in which $D(x) + x^q C$ changes into $-x^q u(0,0)I_{n-1}$ and $\nu(B_0(x)) \geq q+1$. It remains to show that we can obtain $u(x, 0) = u(0, 0) + bx^{\max(1,q)} + O(x^{q+1})$. The series $u(x, 0)$ is already in the required form for $q = 0$. For the case $q \geq 1$, it suffices to consider a polynomial change of coordinates of the form $x = x + P(x)$, with $P(x) = a_2x^2 + \dots + a_qx^q$. This is the consequence of a classical result for one-dimensional vector fields: if $Y = x^{q+1}v(x)\partial_x$ is a vector field with $v(x) = v_0 + v_1x + \dots$, $v_0 \neq 0$ and $q \geq 1$, we can kill all coefficients v_1, \dots, v_{q-1} with a polynomial change of variables, tangent to identity and of degree at most q .

A pair $(\tilde{X}, \tilde{\Gamma})$ in the form (12) will be called a pair in *pre-RS-form*. In the rest of this section, we prove, to finish Theorem 5.5, that any pair (X, Γ) can be reduced to pre-RS-form by means of a finite composition of permissible transformations.

Reduction to pre-RS form

First, performing the blow-ups at the infinitely near points of Γ and by resolution of singularities of curves (see [39]), we can assume that Γ is non-singular. Moreover, using property (a1) above, there is a system of coordinates (x, \mathbf{y}) for which Γ is transversal to $\{x = 0\}$ and such that $X = x^e\tilde{X}$, where \tilde{X} is not divisible by x and $e \geq \nu(X) - 1$ (in particular $\{x = 0\}$ is contained in the singular locus of X if $\nu(X) \geq 2$). Let $\gamma(x) = (x, \tilde{\gamma}(x)) \in \mathbb{C}[[x]]^n$ be a parameterization of Γ in these coordinates and write

$$\tilde{X} = a(x, \mathbf{y}) \frac{\partial}{\partial x} + \mathbf{b}(x, \mathbf{y}) \frac{\partial}{\partial \mathbf{y}},$$

where $a(x, \mathbf{y}) \in \mathbb{C}[[x, \mathbf{y}]]$ and $\mathbf{b}(x, \mathbf{y}) \in \mathbb{C}[[x, \mathbf{y}]]^{n-1}$. Since Γ is invariant and not contained in the singular locus of X we obtain that $a(\gamma(x)) \neq 0$.

Case \tilde{X} not singular

We analyze first the case where \tilde{X} is not singular at the origin. In this case, we have $e \geq 1$ and, since Γ is the unique formal solution of \tilde{X} at 0 and it is transversal to $\{x = 0\}$, we must have $a(0) = \lambda \neq 0$. We may assume also that Γ is tangent to $\{\mathbf{y} = 0\}$. After a new blow-up at the origin, and taking

coordinates as in (11), the transform of X is written as

$$\tilde{X} = x^{e-1} \left[x(\lambda + O(x)) \frac{\partial}{\partial x} + (-\lambda y + O(x)) \frac{\partial}{\partial y} \right],$$

which is in pre-RS-form (12) with $q = e - 1 \geq 0$ and Poincaré rank $p = 0$.

Case \bar{X} singular

Assume now that \bar{X} is a singular formal vector field. Let r be the order of vector field $\bar{X}|_{\Gamma}$, it is equal to the order of the series $a(\gamma(x))$. Notice that $1 \leq r < \infty$. As in Remark 5.2(b), we can assume, up to a polynomial change of variables of the form $\phi_1(x, y) = (x, y + J_N \bar{y}(x))$, that the order in x of any component of $b(x, 0)$ is at least $2r + 2$. Let ϕ be the composition of the permissible blow-ups with center at the first $r + 1$ iterated tangents of Γ , written as $\phi(x, y) = (x, x^{r+1}y)$. Taking into account the effect, stated in property (a2), of a blow-up in the order with respect to x of the different monomials of the coefficients of X and the invariance of $\bar{X}|_{\Gamma}$ under blow-ups, we conclude that, after the transformation ϕ , the vector field X may be written as

$$X = x^e \left[x^r u(x, y) \frac{\partial}{\partial x} + (c(x) + A(x)y + x^r \Theta(x, y)) \frac{\partial}{\partial y} \right] \quad (13)$$

where $e \geq 0, r \geq 1, u(0, 0) \neq 0, \nu(c(x)) \geq r + 1, A(x) \in \mathcal{M}_{n-1}(\mathbb{C}[[x]])$ and $\Theta \in \mathbb{C}[[x, y]]^{n-1}$ has order at least 2 in the y -variables. Moreover, we may assume also that $A(0) \neq 0$: if $\nu(A(x)) \geq r$ then $x^{-(e+r)}X$ is non-singular, a case already treated above; otherwise, if $\nu(A(x)) < r$ we may rewrite X as in (13) replacing e by $e + \nu(A(x))$ and r by $r - \nu(A(x))$ so that the new matrix $A(x)$ satisfies $A(0) \neq 0$.

Put $r = s + 1$ with $s \geq 0$. Notice that if $s = 0$ then X is already in the required pre-RS-form (12) with $q = e$ and Poincaré rank $p = 0$. We assume that $s \geq 1$. To the vector field X in (13) we can associate the system of $n - 1$ formal meromorphic ODEs

$$x^{s+1} \mathbf{y}' = u(x, y)^{-1} (c(x) + A(x)y + x^r \Theta(x, y)). \quad (14)$$

We will use the following classical result, that we state more or less as it appears in the book of Wasow [40].

Theorem 5.6 (Turrittin). *Consider an m -dimensional system of formal linear ODEs*

$$x^{s+1} \mathbf{w}' = \Lambda(x) \mathbf{w}, \quad \Lambda(x) \in \mathcal{M}_m(\mathbb{C}[[x]]),$$

and assume that $s \geq 1$ and $\Lambda(0) \neq 0$. Then, after a finite number of transformations of the following types

- *Polynomial regular transformation*

$$L_{P(x)}(x, \mathbf{w}) = (x, P(x)\mathbf{w}), \quad P(x) \in \mathcal{M}_m(\mathbb{C}[x]) \text{ with } P(0) \text{ invertible.}$$

- *Shearing transformation*

$$S_{(k_1, \dots, k_m)}(x, \mathbf{w}) = (x, \text{diag}(x^{k_1}, \dots, x^{k_m})\mathbf{w}), \quad k_j \in \mathbb{N}_{\geq 0}.$$

- *Ramification*

$$R_\alpha(x, \mathbf{w}) = (x^\alpha, \mathbf{w}), \quad \alpha \in \mathbb{N}_{>0}.$$

the system transforms into a system

$$x^{p+1}\mathbf{w}' = (\bar{D}(x) + x^p\bar{C} + O(x^{p+1}))\mathbf{w},$$

where either $p = 0$ and $\bar{C} \neq 0$ or $p \geq 1$, $\bar{D}(x)$ is a diagonal matrix of polynomials of degree at most $p - 1$ commuting with \bar{C} and $\bar{D}(0) \neq 0$.

Polynomial regular transformations, shearing transformations, and ramifications, as defined in Turrin's Theorem, will be called *T-transformations*. Except for the ramifications, they are particular examples of *gauge* transformations of the system.

Remark 5.7. Note that a polynomial regular transformation does not change the number s (the Poincaré rank of the system) and that a ramification R_α multiplies it by α . The effect of a shearing transformation $S_{(k_1, \dots, k_m)}$ on the Poincaré rank depends on the parameters k_1, \dots, k_m (and on the orders of the entries of the system). Looking carefully at the proof of Theorem 5.6 (see, e.g., [40, Section 19] or, alternatively, the proof in [4]), we may observe that each shearing transformation in the process to prove Theorem 5.6 is chosen so that its application never makes the Poincaré rank increase strictly.

We resume the proof of Theorem 5.5. Assume that X is written as in (13), that Γ is non-singular and transversal to $x = 0$ and let $\gamma(s) = (s, \bar{\gamma}(s))$ be a parameterization of Γ . Consider the formal change of variables $\mathbf{y} = \hat{\mathbf{y}} + \bar{\gamma}(x)$, for which $\Gamma = \{\hat{\mathbf{y}} = 0\}$, and write X in the variables $(x, \hat{\mathbf{y}})$ as

$$x^e \left[x^{s+1} u(x, \hat{\mathbf{y}} + \bar{\gamma}(x)) \frac{\partial}{\partial x} + (\hat{A}(x)\hat{\mathbf{y}} + x^{s+1}\hat{\Theta}(x, \hat{\mathbf{y}})) \frac{\partial}{\partial \hat{\mathbf{y}}} \right]$$

where $\hat{A}(0) \neq 0$ and $\hat{\Theta}(x, \hat{\mathbf{y}}) = O(\|\hat{\mathbf{y}}\|^2)$. The system (14) associated to the vector field X becomes

$$x^{s+1}\hat{\mathbf{y}}' = u(x, \hat{\mathbf{y}} + \bar{\gamma}(x))^{-1}(\hat{A}(x)\hat{\mathbf{y}} + x^{s+1}\hat{\Theta}(x, \hat{\mathbf{y}})).$$

Apply Theorem 5.6 to the linear system

$$x^{s+1}\mathbf{w}' = u(x, \bar{\gamma}(x))^{-1}\hat{A}(x)\mathbf{w}, \quad (15)$$

associated to the formal vector field

$$Y = x^e \left[x^{s+1} u(x, \bar{\gamma}(x)) \frac{\partial}{\partial x} + \hat{A}(x)\mathbf{w} \frac{\partial}{\partial \mathbf{w}} \right].$$

We get a composition Ψ of T -transformations converting (15) into a system with the prescribed properties stated in Theorem 5.6. In terms of the associated vector field Y , if we write $\Psi(x, \mathbf{w}) = (x^\beta, \Psi_2(x, \mathbf{w}))$, where β is the product of the orders of the ramifications involved in the process and Ψ_2 is polynomial in x and linear in \mathbf{w} , we get

$$\Psi^*Y = \beta^{-1}u(x^\beta, \bar{\gamma}(x^\beta))x^\nu \left[x^{p+1} \frac{\partial}{\partial x} + \left(\bar{D}(x) + x^p \bar{C} + O(x^{p+1}) \right) \mathbf{w} \frac{\partial}{\partial \mathbf{w}} \right] \quad (16)$$

where $\nu \geq 0$ and p , $\bar{D}(x)$ and \bar{C} satisfy the properties stated in Theorem 5.6. In fact, we have $p + \nu = \beta(e + s)$.

Lemma 5.8. Ψ is a sequence of permissible transformations for $(Y, \{\mathbf{w} = 0\})$.

Proof. This is clear for polynomial regular transformations and for ramifications. On the other hand, a shearing transformation can be viewed as a composition of blow-ups. More precisely, consider $\phi = S_{(k_2, \dots, k_n)}$ where $k_j > 0$ if $2 \leq j \leq t$ and $k_j = 0$ otherwise. The expression of the shearing transformation is $\phi(x, \mathbf{w}) = (x, x^{k_2}w_2, \dots, x^{k_t}w_t, w_{t+1}, \dots, w_n)$. Put

$$\bar{Y} = x^{-e}Y = a_1(x, \mathbf{w}) \frac{\partial}{\partial x} + \sum_{i=2}^n a_i(x, \mathbf{w}) \frac{\partial}{\partial w_i}.$$

We obtain $\phi^*\bar{Y} = \tilde{a}_1(x, \mathbf{w}) \frac{\partial}{\partial x} + \sum_{i=2}^n \tilde{a}_i(x, \mathbf{w}) \frac{\partial}{\partial w_i}$ defined by

$$\begin{cases} \tilde{a}_j(x, \mathbf{w}) = \frac{a_j(\phi(x, \mathbf{w})) - k_j x^{k_j-1} w_j a_1(\phi(x, \mathbf{w}))}{x^{k_j}}, & \text{for } j = 2, \dots, t; \\ [7pt] \tilde{a}_j(x, \mathbf{w}) = a_j(\phi(x, \mathbf{w})), & \text{for } j \in \{1, t+1, \dots, n\}. \end{cases}$$

Since the Poincaré rank does not increase by the shearing transformations in Tjurittin's process (see Remark 5.7), the pull-back $\phi^*\bar{Y}$ has coefficients in $\mathbb{C}[[x, \mathbf{w}]]$ (with no poles). We deduce that a_1, \dots, a_t belong to the ideal (x, w_2, \dots, w_t) . Therefore $Z = \{x = w_2 = \dots = w_t = 0\}$ is invariant by \bar{Y} and the blow-up of Z is a permissible transformation. Then we consider the blow-up of $\{x = 0\} \cap \cap_{k_j \geq 2} \{w_j = 0\}$. Analogously as above, it is a permissible transformation. By repeating this process with centers of the form $\{x = 0\} \cap \cap_{k_j \geq d} \{w_j = 0\}$ for $1 \leq d \leq \max(k_1, \dots, k_t)$, we get that any shearing transformation is a sequence of permissible transformations for $(\bar{Y}, \{\mathbf{w} = 0\})$, and hence for $(Y, \{\mathbf{w} = 0\})$. \square

Notice that the pair $(\Psi^*Y, \{\mathbf{w} = 0\})$, where Ψ^*Y is given in (16), is in pre-RS-form. Let us show how to reduce (X, Γ) to pre-RS-form from this property. For any $m \geq 1$, consider the diffeomorphism $\phi_m : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ given by $\phi_m(x, \mathbf{y}) = (x, \mathbf{y} - J_{2m}\bar{\gamma}(x))$. The transform $X_m = \phi_m^*X$ is written as

$$X_m = x^e \left[x^{s+1}u(x, \mathbf{y} + J_{2m}\bar{\gamma}(x)) \frac{\partial}{\partial x} + (c_m(x) + A_m(x)\mathbf{y} + x^{s+1}\Theta_m(x, \mathbf{y})) \frac{\partial}{\partial \mathbf{y}} \right],$$

where $A_m(0) \neq 0$, $\Theta_m(x, \mathbf{y}) = O(\|\mathbf{y}\|^2)$ and $c_m(x), A_m(x), \Theta_m(x, \mathbf{y})$ converge, respectively to $0, \hat{A}(x), \hat{\Theta}(x, \mathbf{y})$ in the Krull topology associated to the ideal (x) (also called the (x) -adic topology)

when $m \rightarrow \infty$. Moreover, the transform $\Gamma_m = \phi_m^* \Gamma$ has a parameterization $(x, \bar{y}(x) - J_{2m} \bar{y}(x))$ that converges to $(x, 0)$ in the (x) -adic topology. Therefore, we get $\nu(c_m(x)) \geq 2m + 1$ (see Remark 5.2). Consider the map $\psi_m(x, y) = (x, x^m y)$, a composition of punctual permissible blow-ups for (X_m, Γ_m) . Let (X'_m, Γ'_m) be the transform of (X, Γ) by $\psi_m \circ \phi_m$.

Taking into account formula (9) and property (a2) above, we get that the limit of $X''_m = X'_m + mW$, where $W = x^{e+s} u(x, \bar{y}(x)) y \frac{\partial}{\partial y}$, in the (x) -adic topology is equal to Y when $m \rightarrow \infty$. Moreover, the parameterization $(x, (\bar{y}(x) - J_{2m} \bar{y}(x))/x^m)$ of $(\psi_m \circ \phi_m)^* \Gamma$ converges to $(x, 0)$ in the (x) -adic topology when $m \rightarrow \infty$. It is straightforward to check out that, since Ψ is a sequence of permissible transformations for $(Y, \{\mathbf{w} = 0\})$, there exists a neighborhood U of Y in the (x) -adic topology such that $\Psi^* Z$ is a formal vector field for any $Z \in U$ and the map $Z \mapsto \Psi^* Z$ is continuous in U where we consider the (x) -adic topology in both the source and the target.

In order to finish the proof of Theorem 5.5, it suffices to prove that if m is sufficiently big then Ψ reduces (X'_m, Γ'_m) to pre-RS-form: the map $\Psi \circ \psi_m \circ \phi_m$ will then reduce (X, Γ) to pre-RS-form. Since $\lim_{m \rightarrow \infty} X''_m = Y$, Ψ^* is continuous at Y and $\Psi^*(W) = \tau(x) \mathbf{w} \frac{\partial}{\partial \mathbf{w}}$ for $\tau(x) = x^{\beta(e+s)} u(x^\beta, \bar{y}(x^\beta))$, we deduce that Ψ is a permissible transformation for (X'_m, Γ'_m) . The continuity of Ψ^* at Y implies that the pair $(\Psi^* X'_m, \Psi^* \Gamma'_m)$ is in pre-RS-form where the matrix $x^\nu (\bar{D}(x) + x^p \bar{C})$ in Equation (16) is replaced by

$$x^\nu \bar{D}(x) + x^{p+\nu} (\bar{C} - mcI_{n-1})$$

where $J_{p+\nu} \tau(x) = cx^{p+\nu}$ (recall that $\tau(x)$ has order $\beta(e+s) = p+\nu$). Indeed, the above matrix satisfies the conditions in Definition 5.1 for $m \gg 1$. Theorem 5.5 is finished.

5.4 | Reduction of a biholomorphism to Ramis–Sibuya form

Consider a biholomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ having a formal invariant curve Γ . In this section we use Theorem 5.5 and the results in Section 4 to obtain, up to iteration of F , a reduction of the pair (F, Γ) to a form analogous to the Ramis–Sibuya form in Definition 5.1.

First, we need to adapt Proposition 5.3 to the context of flows.

Proposition 5.9. *Consider $F \in \text{Diff}(\mathbb{C}^n, 0)$ with a formal invariant curve Γ and assume that there exists a formal vector field $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ such that $F = \exp X$ and Γ is invariant for X . Let $\phi : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n, 0)$ be a permissible transformation for (X, Γ) and let $(\tilde{X}, \tilde{\Gamma})$ be the transform of (X, Γ) by ϕ . Then the diffeomorphism $\tilde{F} = \exp \tilde{X}$ is analytic, satisfies $\phi \circ \tilde{F} = F \circ \phi$ and has $\tilde{\Gamma}$ as invariant curve. We say that ϕ is a permissible transformation for (F, Γ) and that the pair $(\tilde{F}, \tilde{\Gamma})$ is the transform of (F, Γ) by ϕ .*

Proof. If ϕ is a germ of a diffeomorphism, the result is clear.

Assume that ϕ is a permissible blow-up with center Z . Consider analytic coordinates $\mathbf{z} = (z_1, z_2, \dots, z_n)$ such that Γ is tangent to the z_1 -axis, $Z = \{z_1 = z_2 = \dots = z_t = 0\}$ and $\phi(\mathbf{z}) = (z_1, z_1 z_2, \dots, z_1 z_t, z_{t+1}, \dots, z_n)$. The condition $\phi \circ \tilde{F} = F \circ \phi$ can be written as

$$z_j \circ \tilde{F}(\mathbf{z}) = \begin{cases} \frac{z_j \circ F(\phi(\mathbf{z}))}{z_1 \circ F(\phi(\mathbf{z}))}, & \text{if } j = 2, \dots, t; \\ z_j \circ F(\phi(\mathbf{z})), & \text{if } j \in \{1, t+1, \dots, n\}. \end{cases}$$

Since the tangent line of Γ is invariant for D_0F , the series $z_1 \circ F(\mathbf{z})$ has a monomial of the form az_1 , with $a \neq 0$. Moreover, since Z is invariant for F , we have $z_j \circ F(\mathbf{z}) \in (z_1, \dots, z_t)$ for $j = 1, \dots, t$. Therefore $z_1 \circ F(\phi(\mathbf{z})) = z_1(a + A(\mathbf{z}))$ with $A(0) = 0$ and $z_j \circ F(\phi(\mathbf{z}))$ is divisible by z_1 for $j = 2, \dots, t$. We conclude that $\tilde{F} \in \text{Diff}(\mathbb{C}^n, 0)$. By construction, \tilde{F} is the unique formal diffeomorphism such that $\phi \circ \tilde{F} = F \circ \phi$. We have also that $\tilde{F} = \exp \tilde{X}$ since $\tilde{X}^j(g \circ \phi) = X^j(g) \circ \phi$ for any $g \in \hat{\mathcal{O}}_n$ and any $j \geq 1$, and thus $\exp \tilde{X}$ also satisfies formally $\phi \circ \exp \tilde{X} = F \circ \phi$. Notice finally that $\tilde{\Gamma}$ is invariant for \tilde{X} by Proposition 5.3 and hence $\tilde{\Gamma}$ is also invariant for $\tilde{F} = \exp \tilde{X}$.

Assume now that ϕ is a permissible l -ramification, written in some coordinates \mathbf{z} as $\phi(\mathbf{z}) = (z_1^l, z_2, \dots, z_n)$, that is, $Z = \{z_1 = 0\}$ is the center of ϕ . As in the case of permissible blow-ups, we have that the formal diffeomorphism $\tilde{F} = \exp \tilde{X}$ satisfies $\phi \circ \tilde{F} = F \circ \phi$. This identity means

$$(z_1 \circ \tilde{F}(\mathbf{z}))^l = z_1 \circ F(\phi(\mathbf{z})); \quad z_j \circ \tilde{F}(\mathbf{z}) = z_j \circ F(\phi(\mathbf{z})), \quad j = 2, \dots, n.$$

On the other hand, since Z is invariant for F , we have $z_1 \circ F(\mathbf{z}) = z_1(a + A(\mathbf{z}))$, with $a \neq 0$ and $A(0) = 0$. We conclude that $\tilde{F} \in \text{Diff}(\mathbb{C}^n, 0)$. The proof of the invariance of $\tilde{\Gamma}$ by \tilde{F} is the same as in the previous case. \square

Remark 5.10. Assume that X is an infinitesimal generator of F in the sense of Definition 4.6. Then, the transformed vector field \tilde{X} in Proposition 5.9 is an infinitesimal generator of \tilde{F} , since being not weakly resonant is invariant by T -transformations. More precisely, given $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$ with $\text{spec}(D_0X) = \{\mu_1, \dots, \mu_n\}$, consider

$$R(X) = \{m_1\mu_1 + \dots + m_n\mu_n : m_1, \dots, m_n \in \mathbb{Q}\}.$$

It is easy to verify that $R(X) = R(\tilde{X})$ for regular transformations, ramifications, and blow-ups. It also holds for shearing transformations since they are compositions of blow-ups.

The main result in this section is the following.

Theorem 5.11 (Reduction of a diffeomorphism to Ramis–Sibuya form). *Let $F \in \text{Diff}(\mathbb{C}^n, 0)$ be a germ of a diffeomorphism having a formal invariant curve Γ . Assume that Γ is rationally neutral and not contained in the set of fixed points of any non-trivial iterate of F . Let m be the index of embeddability of F . Then there exists a finite composition Φ of permissible transformations for (F^m, Γ) and some coordinates (x, \mathbf{y}) at $0 \in \mathbb{C} \times \mathbb{C}^{n-1}$ so that, if $(\tilde{F}^m, \tilde{\Gamma})$ is the transform of (F^m, Γ) by Φ , then $\tilde{\Gamma}$ is non-singular and transversal to $\{x = 0\}$ and \tilde{F}^m is written as:*

$$\begin{cases} x \circ \tilde{F}^m(x, \mathbf{y}) = x - x^{q+1} + bx^{2q+1} + O(x^{2q+2}) \\ \mathbf{y} \circ \tilde{F}^m(x, \mathbf{y}) = \exp(D(x) + x^q C)\mathbf{y} + O(x^{q+1}), \end{cases} \quad (17)$$

where $q \geq 1$, $b \in \mathbb{C}$, $D(x)$ is a diagonal matrix of polynomials of degree at most $q - 1$, C is a constant matrix, $D(x) + x^q C \neq 0$, $[D(x), C] = 0$ and the order of contact of \tilde{F}^m with the identity coincides with the order of the matrix $D(x) + x^q C$ plus one. In this case, we say that the pair $(\tilde{F}^m, \tilde{\Gamma})$ is in Ramis–Sibuya form.

Proof. By Theorem 4.7, the iterate F^m has an infinitesimal generator $X \in \hat{\mathfrak{X}}(\mathbb{C}^n, 0)$. By Proposition 4.22, Γ is invariant for X and $\nu(X|_\Gamma) \geq 2$ (notice that Γ is rationally neutral for any iterate of

F). Moreover, Γ is not contained in the singular locus of X since, otherwise, Γ would be contained in the set of fixed points of $F^m = \exp X$. By Theorem 5.5, there exists a composition Φ of finitely many permissible transformations for (X, Γ) such that the transform $(\tilde{X}, \tilde{\Gamma})$ of (X, Γ) by Φ is in RS-form. Fix some coordinates (x, y) such that $(\tilde{X}, \tilde{\Gamma})$ is written as in Equation (6):

$$\tilde{X} = (\lambda x^{q+1} + bx^{2q+1} + O(x^{2q+2})) \frac{\partial}{\partial x} + ((D(x) + x^q C)y + O(x^{q+1})) \frac{\partial}{\partial y}.$$

Notice that $q \geq 1$ since $\nu(\tilde{X}|_{\tilde{\Gamma}}) \geq 2$ by Proposition 5.3. In particular, by Remark 5.2 we may assume that $\lambda = -1$. We conclude that the transform $\tilde{F}^m = \exp \tilde{X}$ of F by Φ is written as in Equation (17) with the required properties $q \geq 1$, $D(x) + x^q C \not\equiv 0$, $D(x)$ diagonal of degree at most $q - 1$ and $[D(x), C] = 0$. Let ν be the order of $D(x) + x^q C$. It remains to prove that the vector $y \circ \tilde{F}^m - y \in \mathbb{C}\{x, y\}^{n-1}$ has order $\nu + 1$. If \tilde{F}^m is not tangent to the identity then $D_y \tilde{F}^m(0) = \exp((D(x) + x^q C))|_{x=0} \neq I_{n-1}$, which implies $\nu = 0$ and the property holds. Suppose that \tilde{F}^m is tangent to the identity. Then, by Remarks 4.9 and 5.10, \tilde{X} is the unique nilpotent vector field such that $\tilde{F}^m = \exp \tilde{X}$ and we know that $\nu(\tilde{X}) \geq 2$ since $I = D_0 \tilde{F}^m = \exp(D_0 \tilde{X})$ and $D_0 \tilde{X}$ is nilpotent. Using the formula for the exponential, we conclude that $y \circ \tilde{F}^m - y$ has order equal to $\nu(\tilde{X}) = \nu + 1$, as wanted. \square

To finish this section, we show that stable manifolds of F , as well as asymptotic orbits to Γ , are preserved under a permissible transformation for (F, Γ) . Together with Theorem 5.11, this allows us to assume that the pair (F, Γ) is in Ramis–Sibuya form in order to prove Theorem 1 in the (remaining) case where Γ is rationally neutral and not contained in the set of fixed points of any iterate of F . Recall that to obtain Ramis–Sibuya form, we have first considered an iterate of F , then performed some punctual blow-ups along Γ and, once the transform of Γ is non-singular, some other permissible blow-ups with a center with a dimension possibly greater than 0 and ramifications. Then, for our purposes, it is sufficient to consider the statement in the following way.

Proposition 5.12. *Let ϕ be a permissible transformation for (F, Γ) with center Z and exceptional divisor $E_\phi = \phi^{-1}(Z)$. Assume that Γ is non-singular if the center Z has positive dimension. Consider a representative $\phi : \tilde{V} \rightarrow V$ such that F is defined in V and Z is an analytic smooth subvariety of V . Let $(\tilde{F}, \tilde{\Gamma})$ be the transform of (F, Γ) by ϕ . We have*

- (i) *If $\tilde{S} \subset \tilde{V}$ is a stable manifold of \tilde{F} in \tilde{V} such that $\tilde{S} \cap E_\phi = \emptyset$ then $S = \phi(\tilde{S})$ is a stable manifold of F in V . Moreover, if $\tilde{O} \subset \tilde{S}$ is a \tilde{F} -orbit asymptotic to $\tilde{\Gamma}$ then $O = \phi(\tilde{O})$ is a F -orbit asymptotic to Γ .*
- (ii) *If $S \subset V$ is a stable manifold of F such that $S \cap Z = \emptyset$ and every F -orbit in S is tangent to Γ , then $\tilde{S} = \phi^{-1}(S)$ is a stable manifold of \tilde{F} . Moreover, if $O \subset S$ is a F -orbit asymptotic to Γ then $\tilde{O} = \phi^{-1}(O)$ is a \tilde{F} -orbit asymptotic to $\tilde{\Gamma}$.*

Proof. The two assertions concerning the stable manifolds are consequences of the fact that $\phi \circ \tilde{F} = F \circ \phi$, together with the fact that ϕ is an isomorphism outside the divisor E_ϕ . The assertions concerning the asymptoticity of the orbits are immediate from the definition in the case where ϕ is the blow-up at 0. In the other cases, we take coordinates z such that Γ is parameterized by $\gamma(s) = (s, \gamma_2(s), \dots, \gamma_n(s))$ and such that ϕ is either a ramification with respect to $Z = \{z_1 = 0\}$ or is written as in (7) in the case of a blow-up. Using the corresponding formulas (10) or (8) for a

ramification of the transformed curve $\tilde{\Gamma}$ (again non-singular), the result is a consequence of the characterization of asymptoticity of orbits to a non-singular curve in terms of a parameterization of the curve (see Section 2). \square

6 | EXISTENCE OF STABLE MANIFOLDS

Consider a diffeomorphism $F \in \text{Diff}(\mathbb{C}^n, 0)$ and a formal non-singular invariant curve Γ such that the pair (F, Γ) is in Ramis–Sibuya form, that is, there exist coordinates $(x, \mathbf{y}) = (x, y_2, \dots, y_n)$ at $0 \in \mathbb{C}^n$ such that Γ is transverse to $x = 0$ and such that F is written as

$$\begin{aligned} x \circ F(x, \mathbf{y}) &= x - x^{q+1} + bx^{2q+1} + O(x^{2q+2}) \\ \mathbf{y} \circ F(x, \mathbf{y}) &= \exp(D(x) + x^q C) \mathbf{y} + O(x^{q+1}), \end{aligned}$$

where $q \geq 1$, $b \in \mathbb{C}$ and $D(x)$ and C satisfy the properties of Theorem 5.11. Denote by $k + 1$ the order of contact of F with the identity, which coincides with the order of $D(x) + x^q C$ plus one. Note that $0 \leq k \leq q$, and put $p = q - k \geq 0$.

We define the *attracting directions* of (F, Γ) as the $q = k + p$ half-lines $\{x \in \xi \mathbb{R}^+\}$, where $\xi^{k+p} = 1$. Observe that, when Γ is convergent, these directions are the limits of the secant real lines passing through the origin and points in an orbit of the restricted diffeomorphism $F|_{\Gamma} \in \text{Diff}(\mathbb{C}, 0)$, converging to 0. We classify the attracting directions of (F, Γ) as follows. Write $D(x) + x^q C = x^k(\bar{D}(x) + x^p C)$, where $\bar{D}(x) = 0$ in case $p = 0$. In case $p \geq 1$, set

$$\bar{D}(x) = \text{diag}(d_2(x), \dots, d_n(x))$$

and, for any $2 \leq j \leq n$, write $d_j(x) = A_{j, \nu_j} x^{\nu_j} + A_{j, \nu_j+1} x^{\nu_j+1} + \dots + A_{j, p-1} x^{p-1}$ if $d_j(x) \neq 0$, where ν_j is the order of d_j at 0. Given an attracting direction $\ell = \xi \mathbb{R}^+$ and $j \in \{2, \dots, n\}$, we say that ℓ is a *node direction* for (F, Γ) in the variable y_j if $p \geq 1$, $d_j(x) \neq 0$ and

$$\left(\text{Re} \left(\xi^{k+\nu_j} A_{j, \nu_j} \right), \text{Re} \left(\xi^{k+\nu_j+1} A_{j, \nu_j+1} \right), \dots, \text{Re} \left(\xi^{k+p-1} A_{j, p-1} \right) \right) < 0$$

in the lexicographic order; otherwise, we say that it is a *saddle direction* for (F, Γ) in the variable y_j . Note that, if $p = 0$, any attracting direction is a saddle direction in every variable.

The rest of this section is devoted to complete the proof of Theorem 1. After the results in Section 3 and Theorem 5.11, it suffices to show the following theorem.

Theorem 6.1. *Consider a pair (F, Γ) in Ramis–Sibuya form and let ℓ be an attracting direction of (F, Γ) . Let $s - 1 \geq 0$ be the number of variables for which ℓ is a node direction. Then, there exists a stable manifold S_{ℓ} of F of dimension s in which every orbit is asymptotic to Γ and tangent to ℓ . More precisely, there exist a connected and simply connected domain $S \subset \mathbb{C}^s$ with $0 \in \partial S$ and a holomorphic map $\varphi : S \rightarrow \mathbb{C}^{n-s}$ such that, up to reordering the variables, the set*

$$S_{\ell} = \{(x, \mathbf{w}, \varphi(x, \mathbf{w})) \in \mathbb{C} \times \mathbb{C}^{s-1} \times \mathbb{C}^{n-s} : (x, \mathbf{w}) \in S\}$$

satisfies the following properties:

- (i) S_ℓ is a stable manifold of F .
- (ii) Every orbit $\{(x_j, \mathbf{y}_j)\} \subset S_\ell$ is asymptotic to Γ and $\{x_j\}$ is tangent to ℓ .
- (iii) If $\{(x_j, \mathbf{y}_j)\} \subset \mathbb{C} \times \mathbb{C}^{n-1}$ is an orbit of F asymptotic to Γ such that $\{x_j\}$ has ℓ as tangent direction, then $(x_j, \mathbf{y}_j) \in S_\ell$ for all j sufficiently big.

Choice of coordinates

Up to a linear change of coordinates in the x -variable, we may assume that $\ell = \mathbb{R}^+$. We can also assume, without loss of generality, that ℓ is a node direction in the variables y_2, \dots, y_s and a saddle direction in the variables y_{s+1}, \dots, y_n and that C is in Jordan normal form (see Remark 5.2).

Observe that we can increase the order of contact of Γ with the x -axis by considering a polynomial change of variables of the form $(x, \mathbf{y}) \mapsto (x, \mathbf{y} - J_N \bar{\gamma}(x))$ where $\gamma(x) = (x, \bar{\gamma}(x))$ is a parameterization of Γ . Moreover, the matrices $D(x)$ and C that appear in the expression of $\mathbf{y} \circ F$ are preserved by such transformations. Note also that after a permissible punctual blow-up the transformed pair $(\tilde{F}, \tilde{\Gamma})$ is again in Ramis–Sibuya form in usual coordinates $(x, \bar{\mathbf{y}})$ with $\mathbf{y} = x\bar{\mathbf{y}}$ as in Section 5.3. Moreover, the matrix $D(x)$ is invariant by blow-up whereas C is replaced by $C + I_{n-1}$. Consequently, the saddle or node character of $\ell = \mathbb{R}^+$ in each variable does not change and, by Proposition 5.12, it suffices to prove Theorem 6.1 in the new coordinates $(x, \bar{\mathbf{y}})$. Therefore, taking N sufficiently big and up to several punctual admissible blow-ups, if we put $(x, \mathbf{y}) = (x, \mathbf{w}, \mathbf{z}) \in \mathbb{C} \times \mathbb{C}^{s-1} \times \mathbb{C}^{n-s}$ we can write F as

$$x \circ F(x, \mathbf{y}) = f(x, \mathbf{y}) = x - x^{k+p+1} + bx^{2k+2p+1} + O(x^{2k+2p+2})$$

$$\mathbf{w} \circ F(x, \mathbf{y}) = \bar{F}_1(x, \mathbf{y}) = \exp\left(x^k\left(\bar{D}_1(x) + x^p C_1\right)\right) \mathbf{w} + O(x^{k+p+1})$$

$$\mathbf{z} \circ F(x, \mathbf{y}) = \bar{F}_2(x, \mathbf{y}) = \exp\left(x^k\left(\bar{D}_2(x) + x^p C_2\right)\right) \mathbf{z} + O(x^{k+p+1}),$$

where $b \in \mathbb{C}$, $\bar{D}_1(x), \bar{D}_2(x), C_1, C_2$ are the corresponding blocks of $\bar{D}(x)$ and C (note that this decomposition is guaranteed by the commutativity of $D(x)$ and C , see Remark 5.2) and every eigenvalue of C_2 has positive real part.

In fact, we will use coordinates for which Γ has an arbitrarily big order of contact with the x -axis. Fix $m \in \mathbb{N}$, with $m \geq p+2$, and let $\gamma(x) = (x, \bar{\gamma}(x))$ be a parameterization of Γ . Consider the polynomial change of variables $\mathbf{y} \mapsto \mathbf{y}^m = \mathbf{y} - J_{p+m-1} \bar{\gamma}(x)$. In these coordinates, the order of contact of Γ with the x -axis is at least $p+m$, and the invariance of Γ implies that the order of $\mathbf{y}^m \circ F(x, 0)$ is at least $k+p+m$. Therefore, if we set $(x, \mathbf{y}^m) = (x, \mathbf{w}^m, \mathbf{z}^m) \in \mathbb{C} \times \mathbb{C}^{s-1} \times \mathbb{C}^{n-s}$ we have

$$f(x, \mathbf{y}^m) = x - x^{k+p+1} + bx^{2k+2p+1} + O(x^{2k+2p+2})$$

$$\bar{F}_1(x, \mathbf{y}^m) = \exp\left(x^k\left(\bar{D}_1(x) + x^p C_1\right)\right) \mathbf{w}^m + O(x^{k+p+1} \|\mathbf{y}^m\|, x^{k+p+m})$$

$$\bar{F}_2(x, \mathbf{y}^m) = \exp\left(x^k\left(\bar{D}_2(x) + x^p C_2\right)\right) \mathbf{z}^m + O(x^{k+p+1} \|\mathbf{y}^m\|, x^{k+p+m}).$$

Write, as above, $\bar{D}(x) = \text{diag}(d_2(x), \dots, d_n(x))$, where $d_j(x)$ is a polynomial of degree at most $p-1$ for all $2 \leq j \leq n$, and set $C_2 = \text{diag}(A_{s+1,p}, \dots, A_{n,p}) + N_2$, where $A_{j,p} \in \mathbb{C}$ for all $s+1 \leq j \leq n$ and N_2 is a nilpotent matrix.

For any $2 \leq j \leq n$, write

$$e_j(x) := d_j(x) + A_{j,p}x^p = A_{j,\nu_j}x^{\nu_j} + A_{j,\nu_j+1}x^{\nu_j+1} + \cdots + A_{j,p}x^p,$$

where ν_j is the order of e_j at 0 and $A_{j,p} = 0$ for $2 \leq j \leq s$. We set $k + \mu_j$ as the order of $x^k e_j(x) - i \operatorname{Im}[(x^k e_j(x))(0)]$. Note that $\mu_j = \nu_j$ if $k + \nu_j \geq 1$ or if $k + \nu_j = 0$ and $\operatorname{Re}(A_{j,0}) \neq 0$. Therefore, we have that

$$\operatorname{Re}(x^k e_j(x)) = \operatorname{Re}\left(A_{j,\mu_j}x^{k+\mu_j} + A_{j,\mu_j+1}x^{k+\mu_j+1} + \cdots + A_{j,p}x^{k+p}\right).$$

We define the *first asymptotic significant order* $r_j = r_j(\ell)$ of $\ell = \mathbb{R}^+$ in the variable y_j as $r_j = l_j - \mu_j$, where $\mu_j \leq l_j \leq p$ is the first index such that $\operatorname{Re}(A_{j,l_j}) \neq 0$ (note that l_j is well defined for all j because of the previous condition on the eigenvalues of C_2).

Note that r_j does not depend on m , and that $r_j < k + p$ for all j : the inequality is clear if $p = 0$ or $k + \mu_j \geq 1$; otherwise, $r_j = 0$ so it also holds.

Put $r = \max\{r_2, \dots, r_n\}$. For $d, e, \varepsilon > 0$, we define the set $R_{d,e,\varepsilon}$ as

$$R_{d,e,\varepsilon} = \{x \in \mathbb{C} : |x| < \varepsilon, \operatorname{Re} x > 0, -d(\operatorname{Re} x)^{r+1} < \operatorname{Im} x < e(\operatorname{Re} x)^{r+1}\}.$$

Lemma 6.2. Set $t = \max\{r_2 + \mu_2, \dots, r_s + \mu_s\} < p$. There exists a constant $c > 0$ such that, if $d, e, \varepsilon > 0$ are sufficiently small, then for any $x \in R_{d,e,\varepsilon}$ we have

- (i) $\operatorname{Re}(x^k d_j(x)) \leq -c|x|^{k+t}$ for any $2 \leq j \leq s$.
- (ii) $\operatorname{Re}(x^k d_j(x) + x^{k+p} A_{j,p}) \geq c|x|^{k+p}$ for any $s+1 \leq j \leq n$.

Proof. Let us prove (i), the proof of (ii) is analogous. Fix $2 \leq j \leq s$. If $r_j = 0$, we have that

$$\operatorname{Re}(x^k d_j(x)) \leq \operatorname{Re}(A_{j,\mu_j}x^{k+\mu_j})/2 \leq -c_j|x|^{k+\mu_j}$$

if d, e, ε are sufficiently small, where $-c_j = \operatorname{Re}(A_{j,\mu_j})/3$. If $r_j \geq 1$, we have that $k + \mu_j \geq 1$ and we use the same argument of [22, Lemma 5.9], that we include for the sake of completeness. We assume that $\operatorname{Im}(A_{j,\mu_j}) > 0$; the other case is analogous. Using indeterminate coefficients we can see that there exists a diffeomorphism $\rho(x) = x + \sum_{l \geq 2} \rho_l x^l$ such that

$$A_{j,\mu_j}x^{k+\mu_j} + \cdots + A_{j,p-1}x^{k+p-1} = A_{j,\mu_j}\rho(x)^{k+\mu_j},$$

with $\rho_l \in \mathbb{R}$ if $2 \leq l \leq r_j$ and $\operatorname{Im}(\rho_{r_j+1}) > 0$. Hence, to prove the inequality in (i), it suffices to show that $\operatorname{Re}(A_{j,\mu_j}x^{k+\mu_j}) \leq -c_j|x|^{k+r_j+\mu_j}$ for some $c_j > 0$ and for all $x \in \rho(R_{d,e,\varepsilon})$. It is easy to show, using the fact that $\rho_l \in \mathbb{R}$ for $2 \leq l \leq r_j$, that for any $a \in \mathbb{R}$, the image under ρ of the curve

$$\operatorname{Im} x = a(\operatorname{Re} x)^{r_j+1}$$

is a curve of the form

$$C_a : \operatorname{Im} x = (a + \operatorname{Im}(\rho_{r_j+1}))(\operatorname{Re} x)^{r_j+1} + \dots$$

Then, since $r \geq r_j$, we obtain that set $\rho(R_{d,e,\varepsilon})$ is contained, if ε is small enough, in a domain enclosed by two curves of the type C_e and C_{-d} . If d is sufficiently small, then $-d + \text{Im}(\rho_{r_j+1}) > 0$, so $d'|x|^{r_j} < \arg x < \pi/(2(k + \mu_j))$ for some $d' > 0$ and for all $x \in \rho(R_{d,e,\varepsilon})$, if d, e, ε are small enough. Then, we have

$$\begin{aligned} \text{Re}(A_{j,\mu_j} x^{k+\mu_j}) &= -\text{Im}(A_{j,\mu_j})|x|^{k+\mu_j} \sin((k + \mu_j)\arg x) \\ &\leq -\text{Im}(A_{j,\mu_j})|x|^{k+\mu_j} \sin((k + \mu_j)d'|x|^{r_j}) \end{aligned}$$

so $\text{Re}(A_{j,\mu_j} x^{k+\mu_j}) \leq -c_j|x|^{k+r_j+\mu_j}$, with $c_j = \text{Im}(A_{j,\mu_j})(k + \mu_j)d'/2$, if $d, e, \varepsilon > 0$ are small enough. This proves (i). \square

Up to a linear change of coordinates $\mathbf{z}^m \mapsto P\mathbf{z}^m$, we can assume that the non-zero terms of the nilpotent part N_2 of the matrix C_2 are all equal to $c/2$, where $c > 0$ is the constant appearing in Lemma 6.2.

Existence of the stable manifold

We prove here that for every $m \geq p + 2$ there exists a stable manifold S_m of dimension s given by a graph $\mathbf{z}^m = \varphi_m(x, \mathbf{w}^m)$ over a domain of the form

$$S_{d,e,\varepsilon}^m = \{(x, \mathbf{w}^m) \in \mathbb{C} \times \mathbb{C}^{s-1} : x \in R_{d,e,\varepsilon}, \|\mathbf{w}^m\| < |x|^{m-1}\}$$

where $d, e, \varepsilon > 0$. As we will see, these stable manifolds are essentially the same for different values of m . In the proof, we can see that the contact of S_m with Γ increases with m . This will be key in the proof of asymptoticity of the orbits inside each S_m .

We consider the vector spaces $C(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s})$ and $\mathcal{O}(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s})$ of continuous and holomorphic maps, respectively, from $S_{d,e,\varepsilon}^m$ to \mathbb{C}^{n-s} with the compact-open topology. Recall that since $S_{d,e,\varepsilon}^m$ is a locally compact second countable space and \mathbb{C}^{n-s} is a complete metric space, $C(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s})$ is completely metrizable [13, p. 272].

We will use the following result, which is an application of Schauder–Tychonoff theorem and is stated in [20, p. 15]. We include a proof for the sake of completeness.

Proposition 6.3. *Given a continuous function $L : S_{d,e,\varepsilon}^m \rightarrow \mathbb{R}_{\geq 0}$, the set*

$$\mathcal{H}_L = \left\{ \varphi \in \mathcal{O}(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s}) : \|\varphi(x)\| \leq L(x) \text{ for all } (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m \right\}$$

has the fixed point property; that is, every continuous map $T : \mathcal{H}_L \rightarrow \mathcal{H}_L$ has a fixed point.

Proof. The space $C(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s})$ is locally convex for the compact-open topology and the set \mathcal{H}_L is clearly convex and closed. Moreover, by Montel theorem \mathcal{H}_L is sequentially compact, and hence compact since $C(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s})$ is metrizable. By Schauder–Tychonoff theorem (see [14]), every compact convex subset of a locally convex linear topological space has the fixed point property and this ends the proof. \square

We define

$$\mathcal{H}_{d,e,\varepsilon}^m = \left\{ \varphi \in \mathcal{O}(S_{d,e,\varepsilon}^m, \mathbb{C}^{n-s}) : \|\varphi(x, \mathbf{w}^m)\| \leq |x|^{m-1} \text{ for all } (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m \right\}.$$

The stable manifold S_m will be given by the graph of a fixed point φ_m of a convenient continuous map $T : \mathcal{H}_{d,e,\varepsilon}^m \rightarrow \mathcal{H}_{d,e,\varepsilon}^m$.

Given $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$, we denote

$$f_\varphi(x, \mathbf{w}^m) = f(x, \mathbf{w}^m, \varphi(x, \mathbf{w}^m)), \quad \bar{F}_{1,\varphi}(x, \mathbf{w}^m) = \bar{F}_1(x, \mathbf{w}^m, \varphi(x, \mathbf{w}^m)).$$

Proposition 6.4. *If $d, e, \varepsilon > 0$ are sufficiently small, then for all $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ and $(x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m$ we have that*

$$(f_\varphi(x, \mathbf{w}^m), \bar{F}_{1,\varphi}(x, \mathbf{w}^m)) \in S_{d,e,\varepsilon}^m.$$

Proof. Since $r < k + p$, we can argue as in [22, Lemma 5.5], and we obtain that

$$f_\varphi(S_{d,e,\varepsilon}^m) \subset R_{d,e,\varepsilon}$$

if d, e, ε are sufficiently small. Now, if $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ we have by Lemma 6.2 that

$$\begin{aligned} \left\| \bar{F}_{1,\varphi}(x, \mathbf{w}^m) \right\| &= \frac{\left\| \mathbf{w}^m \left(\exp(x^k \bar{D}_1(x)) + O(x^{k+p}) \right) + O(x^{k+p+m}) \right\|}{\left| f_\varphi(x, \mathbf{w}^m) \right|^{m-1}} \\ &\leq \frac{\|\mathbf{w}^m\| \left\| \exp(x^k \bar{D}_1(x)) + O(x^{k+p}) \right\| |1 + O(x^{k+p})| + \|O(x^{k+p+1})\|}{|x|^{m-1}} \\ &\leq \frac{\|\mathbf{w}^m\|}{|x|^{m-1}} (1 - c|x|^{k+t} + \|O(x^{k+t+1})\|) |1 + O(x^{k+p})| + \|O(x^{k+p+1})\| \\ &< 1 - c|x|^{k+t} + \|O(x^{k+t+1})\| < 1 \end{aligned}$$

for all $(x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m$, if d, e, ε are sufficiently small. \square

We consider $0 < \varepsilon < 1$ and fix $d, e > 0$ small enough so that Lemma 6.2 and Proposition 6.4 hold. Given $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ and $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$, we denote

$$(x_j, \mathbf{w}_j^m) = \left(f_\varphi(x_{j-1}, \mathbf{w}_{j-1}^m), \bar{F}_{1,\varphi}(x_{j-1}, \mathbf{w}_{j-1}^m) \right), \quad j \geq 1.$$

As in the classical one-dimensional case, there exists a constant $K \geq 1$ such that

$$\lim_{j \rightarrow \infty} (k+p)jx_j^{k+p} = 1 \quad \text{and} \quad |x_j|^{k+p} \leq K \frac{|x_0|^{k+p}}{1 + (k+p)j|x_0|^{k+p}} \quad (18)$$

for all $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$ and all $j \in \mathbb{N}$, so in particular $(x_j, \mathbf{w}_j^m) \rightarrow 0$ when $j \rightarrow \infty$. Therefore, $\varphi_m \in \mathcal{H}_{d,e,\varepsilon}^m$ is a solution of the equation

$$\varphi(f_\varphi(x, \mathbf{w}^m), \bar{F}_{1,\varphi}(x, \mathbf{w}^m)) = \bar{F}_2(x, \mathbf{w}^m, \varphi(x, \mathbf{w}^m)) \quad (19)$$

if and only if the set

$$\mathcal{S}_m = \left\{ (x, \mathbf{w}^m, \varphi_m(x, \mathbf{w}^m)) : (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m \right\}$$

is a stable manifold of F .

Set $\rho = 0$ when $k \geq 1$, and $\rho = b - (p+1)/2$ when $k = 0$ (recall that b is the coefficient of x^{2p+1} in $f(x, 0)$). We define

$$E(x) = \exp \left(- \int \frac{\bar{D}_2(x) + x^p C_2}{x^{p+1}(1 - \rho x^p)} dx \right),$$

where the integral is a notation for a primitive of $x^{-(p+1)}(1 - \rho x^p)^{-1}(\bar{D}_2(x) + x^p C_2)$.

Lemma 6.5. *For any $(x, \mathbf{y}^m, \mathbf{z}^m) \in S_{d,e,\varepsilon}^m \times \{\mathbf{z}^m \in \mathbb{C}^{n-s} : \|\mathbf{z}^m\| \leq |x|^{m-1}\}$ with ε sufficiently small, we have*

$$E(x)E(f(x, \mathbf{w}^m, \mathbf{z}^m))^{-1} = \exp(-x^k(\bar{D}_2(x) + x^p C_2)) + O(x^{k+p+1}).$$

Proof. We argue as in [23, Lemma 3.7]. Observe that, since $\bar{D}_2(x)$ is diagonal and commutes with C_2 , $E(x)$ is a fundamental solution of the linear system $x^{p+1}Y' = -B(x)Y$, where $B(x) = (\bar{D}_2(x) + x^p C_2)(1 - \rho x^p)^{-1}$. Put $\Omega(x, z) = E(x + x^{p+1}z)$. If we fix x and consider Ω as a function of z then it satisfies the (regular) system

$$\frac{\partial \Omega}{\partial z} = \frac{-B(x + x^{p+1}z)}{(1 + x^p z)^{p+1}} \Omega(x, z).$$

On the other hand, we have $\Omega(x, 0) = E(x)$ and thus

$$\Omega(x, z) = \exp \left(- \int_0^z \frac{B(x + x^{p+1}u)}{(1 + x^p u)^{p+1}} du \right) E(x)$$

(using again that $\bar{D}_2(x)$ is diagonal and commutes with C_2). Hence

$$E(x)E(x + x^{p+1}z)^{-1} = E(x)\Omega(x, z)^{-1} = \exp \left(\int_0^z \frac{B(x + x^{p+1}u)}{(1 + x^p u)^{p+1}} du \right).$$

The integrand in the equation above is an analytic function of (x, u) and may be written as

$$\frac{B(x + x^{p+1}u)}{(1 + x^p u)^{p+1}} = B(x) - (p+1)x^p B(x)u + O(x^{p+1}u, x^{2p}u^2).$$

Integrating, we obtain, for any z sufficiently small,

$$E(x)E(x + x^{p+1}z)^{-1} = \exp\left(B(x)\left(z - \frac{p+1}{2}x^p z^2\right)\right) + x^{p+1}z^2\Lambda(x, z) + x^{2p}z^3\Theta(x, z),$$

where Λ and Θ are analytic at the origin. The result follows using the expression of $f(x, \mathbf{y}^m)$ and taking into account that $(x, \mathbf{y}^m) \in S_{d,e,\varepsilon}^m \times \{\|\mathbf{z}^m\| \leq |x|^{m-1}\}$, and thus $\|\mathbf{y}^m\| \leq |x|^{p+1}$, since $m \geq p+2$. \square

Lemma 6.6. *If $\varepsilon > 0$ is small enough and, given $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$, we put $x_j = f_\varphi(x_{j-1}, \mathbf{w}_{j-1}^m)$ and $\mathbf{w}_j^m = \bar{F}_{1,\varphi}(x_{j-1}, \mathbf{w}_{j-1}^m)$ for any $j \geq 1$, then:*

(i) *For any real number $l > k + p$ there exists a constant $K_l > 0$ such that for any $(x_0, \mathbf{w}_0) \in S_{d,e,\varepsilon}^m$ and any $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ we have*

$$\sum_{j \geq 0} |x_j|^l \leq K_l |x_0|^{l-k-p}.$$

(ii) *For any $(x_0, \mathbf{w}_0) \in S_{d,e,\varepsilon}^m$ and any $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$, we have $\|E(x_0)E(x_j)^{-1}\| \leq 1$ for every $j \geq 0$.*

Proof. Part (i) follows from Equation (18), as in [18, Corollary 4.3]. To prove part (ii), observe that by Lemma 6.5

$$E(x_0)E(x_1)^{-1} = \exp\left(-x_0^k(\bar{D}_2(x_0) + x_0^p C_2)\right) + \theta_\varphi(x_0, \mathbf{w}_0^m),$$

where $\|\theta_\varphi(x_0, \mathbf{w}_0^m)\| \leq M_1 |x_0|^{k+p+1}$ for any $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$ and any $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$, with some $M_1 > 0$ independent of φ . We have that

$$\exp\left(-x^k(\bar{D}_2(x) + x^p C_2)\right) = \mathcal{D} \exp(-x^{k+p} N_2) = \mathcal{D}[I - x^{k+p} N_2 + O(x^{k+p+1})],$$

where

$$\mathcal{D} = \text{diag}(\exp(-x^k d_{s+1}(x) - x^{k+p} A_{s+1,p}), \dots, \exp(-x^k d_n(x) - x^{k+p} A_{n,p})).$$

Then, using Lemma 6.2 and the fact that all the non-zero terms of N_2 are equal to $c/2$, we obtain

$$\|E(x_0)E(x_1)^{-1}\| \leq 1 - (c - c/2)|x_0|^{k+p} + M_2 |x_0|^{k+p+1} \leq 1$$

for all $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$ if $\varepsilon > 0$ is sufficiently small. We obtain the result writing $E(x_0)E(x_j)^{-1} = \prod_{l=0}^{j-1} E(x_l)E(x_{l+1})^{-1}$. \square

Define

$$H(x, \mathbf{w}^m, \mathbf{z}^m) = \mathbf{z}^m - E(x)E(f(x, \mathbf{w}^m, \mathbf{z}^m))^{-1} \bar{F}_2(x, \mathbf{w}^m, \mathbf{z}^m).$$

Using Lemma 6.5 we get

$$H(x, \mathbf{w}^m, \mathbf{z}^m) = O(x^{k+p+1} \|\mathbf{y}^m\|, x^{k+p+m}). \quad (20)$$

Proposition 6.7. *If $\varepsilon > 0$ is sufficiently small and we denote $x_j = f_\varphi(x_{j-1}, \mathbf{w}_{j-1}^m)$ and $\mathbf{w}_j^m = \bar{F}_{1,\varphi}(x_{j-1}, \mathbf{w}_{j-1}^m)$ for any $j \geq 1$, where $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$ and $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$, then the series*

$$T\varphi(x_0, \mathbf{w}_0^m) = \sum_{j \geq 0} E(x_0) E(x_j)^{-1} H(x_j, \mathbf{w}_j^m, \varphi(x_j, \mathbf{w}_j^m))$$

is normally convergent and defines a map $T : \varphi \mapsto T\varphi$ from $\mathcal{H}_{d,e,\varepsilon}^m$ to itself which is continuous for the topology of the uniform convergence. Moreover, $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ is a fixed point of T if and only if the set $\{(x, \mathbf{w}^m, \varphi(x, \mathbf{w}^m)) : (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m\}$ is a stable manifold of F .

Proof. If $\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$ and $(x_0, \mathbf{w}_0^m) \in S_{d,e,\varepsilon}^m$, then by Equation (20) we have that $\|H(x_j, \mathbf{w}_j^m, \varphi(x_j, \mathbf{w}_j^m))\| \leq M|x_j|^{k+p+m}$ for some $M > 0$, so by Lemma 6.6 we get

$$\|T\varphi(x_0, \mathbf{w}_0^m)\| \leq M \sum_{j \geq 0} |x_j|^{k+p+m}$$

and the series is normally convergent by Lemma 6.6. Moreover we have that $\|T\varphi(x, \mathbf{w}^m)\| \leq MK_{k+p+m}|x|^m \leq |x|^{m-1}$ if $\varepsilon > 0$ is sufficiently small, so $T\varphi \in \mathcal{H}_{d,e,\varepsilon}^m$. Continuity of T follows from the uniform convergence of the series with respect to φ . Finally, we rewrite

$$\begin{aligned} T\varphi(x_0, \mathbf{w}_0^m) &= E(x_0) \sum_{j \geq 0} \left[E(x_j)^{-1} \varphi(x_j, \mathbf{w}_j^m) - E(x_{j+1})^{-1} \bar{F}_2(x_j, \mathbf{w}_j^m, \varphi(x_j, \mathbf{w}_j^m)) \right] \\ &= \varphi(x_0, \mathbf{w}_0^m) - E(x_0) E(x_1)^{-1} \left[\bar{F}_2(x_0, \mathbf{w}_0^m, \varphi(x_0, \mathbf{w}_0^m)) - T\varphi(x_1, \mathbf{w}_1^m) \right]. \end{aligned}$$

From these two equalities it follows that φ is a fixed point of T if and only if φ satisfies the invariance Equation (19), that is, if and only if the set $\{(x, \mathbf{w}^m, \varphi(x, \mathbf{w}^m)) : (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m\}$ is a stable manifold of F . \square

By Proposition 6.3, T has a fixed point $\varphi_m \in \mathcal{H}_{d,e,\varepsilon}^m$. Hence, by Proposition 6.7, the set

$$S_m = \{(x, \mathbf{w}^m, \varphi_m(x, \mathbf{w}^m)) : (x, \mathbf{w}^m) \in S_{d,e,\varepsilon}^m\}$$

is a stable manifold of F .

Stable manifold as a base of asymptotic convergence

Let us show that every orbit $\{(x_j, \mathbf{y}_j^m)\}$ of F which is asymptotic to Γ and such that $\{x_j\}$ has \mathbb{R}^+ as tangent direction is eventually contained in S_m . Since the order of contact of Γ with the x -axis is at least $p+m$, any orbit $\{(x_j, \mathbf{y}_j^m)\}$ asymptotic to Γ satisfies $\|\mathbf{y}_j^m\| < |x_j|^{p+m-1}$ if j is sufficiently large. Therefore, the result is a consequence of the following lemma.

Lemma 6.8. Let $\{(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m)\}$ be a stable orbit of F such that $\{x_j\}$ has \mathbb{R}^+ as tangent direction and such that $\|\mathbf{w}_j^m\| < |x_j|^{m-1}$ for all j sufficiently big. Then $(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m) \in S_m$ for all j sufficiently big.

Proof. Since $\{x_j\}$ has \mathbb{R}^+ as tangent direction, we obtain, arguing exactly as in [22, Lemma 5.8], that $x_j \in R_{d,e,\varepsilon}$ if j is sufficiently big and hence $(x_j, \mathbf{w}_j^m) \in S_{d,e,\varepsilon}^m$ for all $j \geq j_0$. Consider the change of coordinates $\mathbf{z}^m \mapsto \mathbf{z}^m - \varphi_m(x, \mathbf{w}^m)$, valid on $S_{d,e,\varepsilon}^m \times \mathbb{C}^{n-s}$. In the new coordinates the stable manifold S_m is given by $\mathbf{z}^m = 0$ and hence F is written as

$$f(x, \mathbf{y}^m) = x - x^{k+p+1} + bx^{2k+2p+1} + O(x^{2k+2p+2})$$

$$\bar{F}_1(x, \mathbf{y}^m) = \exp\left(x^k\left(\bar{D}_1(x) + x^p C_1\right)\right) \mathbf{w}^m + O(x^{k+p+1} \|\mathbf{y}^m\|, x^{k+p+m})$$

$$\bar{F}_2(x, \mathbf{y}^m) = \exp\left(x^k\left(\bar{D}_2(x) + x^p C_2\right)\right) \mathbf{z}^m + O(x^{k+p+1} \|\mathbf{z}^m\|).$$

By Lemma 6.2 we obtain

$$\|\bar{F}_2(x_j, \mathbf{y}_j^m)\| \geq \left(1 + c|x_j|^{k+p} + O(x_j^{k+p+1})\right) \|\mathbf{z}_j^m\| \geq \|\mathbf{z}_j^m\|$$

for all $j \geq j_0$, so we conclude that if $\mathbf{z}_{j_0}^m \neq 0$ the orbit $\{(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m)\}$ cannot converge to the origin. Therefore, $(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m) \in S_m$ for any $j \geq j_0$. \square

Remark 6.9. Note that Lemma 6.8 also implies that φ_m is actually the unique fixed point of T in $\mathcal{H}_{d,e,\varepsilon}^m$.

Asymptoticity of the orbits

To finish the proof of Theorem 6.1 it only remains to prove that every orbit in S_m is asymptotic to Γ . Observe that, since the order of contact of Γ with the x -axis is at least $p+m$ and the order of contact of S_m with the x -axis is at least $m-1$, the order of contact of S_m with Γ is at least $m-1$. We will show that every orbit $\{(x_j, \mathbf{y}_j^m)\} \subset S_m$, which has order of contact at least $m-1$ with Γ , is eventually contained in S_{m+1} , and therefore its order of contact with Γ is at least m . Applying this argument recursively, we conclude that every orbit in S_m is asymptotic to Γ .

Lemma 6.10. Fix $\varepsilon, d, e > 0$ sufficiently small. Let $\{(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m)\}$ be a stable orbit of F such that $x_j \in R_{d,e,\varepsilon}$ and $\|\mathbf{z}_j^m\| < |x_j|^{m-1}$ for all j . Then $\|\mathbf{w}_j^m\| < \frac{1}{2}|x_j|^m$ for all j sufficiently large.

Proof. By Lemma 6.2 we have

$$\begin{aligned} \|\mathbf{w}_{j+1}^m\| &= \frac{\left\|\mathbf{w}_j^m \left(\exp(x_j^k \bar{D}_1(x_j)) + O(x_j^{k+p}) \right) + O(x_j^{k+p+m}) \right\|}{|x_{j+1}|^m} \\ &\leq \frac{\|\mathbf{w}_j^m\| \left(1 - c|x_j|^{k+p} + O(x_j^{k+p+1}) \right) + \|O(x_j^{k+p})\|}{|x_j|^{m+1}} \end{aligned}$$

for all j . This implies, since $t < p$, that if $\|\mathbf{w}_j^m\| < \frac{1}{2}|x_j|^m$ then $\|\mathbf{w}_{j+1}^m\| < \frac{1}{2}|x_{j+1}|^m$. Therefore, to prove the lemma it suffices to show that $\|\mathbf{w}_j^m\| < \frac{1}{2}|x_j|^m$ for some j . Suppose this is not the case, so $\|\mathbf{w}_j^m\| \geq \frac{1}{2}|x_j|^m$ for all $j \geq 0$. Then

$$\frac{\|\mathbf{w}_{j+1}^m\|}{|x_{j+1}|^m} \leq \frac{\|\mathbf{w}_j^m\|}{|x_j|^m} \left(1 - c|x_j|^{k+t} + O(x_j^{k+t+1})\right)$$

for all $j \geq 0$, so we obtain that

$$\frac{\|\mathbf{w}_{j+1}^m\|}{|x_{j+1}|^m} \leq \frac{\|\mathbf{w}_0^m\|}{|x_0|^m} \prod_{l=0}^j (1 - c|x_l|^{k+t} + O(x_l^{k+t+1})).$$

Since $\lim_{j \rightarrow \infty} (k+p)jx_j^{k+p} = 1$ and $t < p$, the product above converges to 0 when $j \rightarrow \infty$, contradicting the fact that $\|\mathbf{w}_j^m\| \geq \frac{1}{2}|x_j|^m$ for all j . \square

Consider an orbit $\{(x_j, \mathbf{w}_j^m, \mathbf{z}_j^m)\} \subset S_m$ and consider the coordinates $(x, \mathbf{y}^{m+1}) = (x, \mathbf{w}^{m+1}, \mathbf{z}^{m+1})$ satisfying $\mathbf{y}^{m+1} = \mathbf{y}^m - (J_{p+m}\bar{\gamma}(x) - J_{p+m-1}\bar{\gamma}(x))$, where $\gamma(s) = (s, \bar{\gamma}(s))$ is a parameterization of Γ . By Lemma 6.10, $\|\mathbf{w}_j^m\| < \frac{1}{2}|x_j|^m$ for all j sufficiently large, so $\|\mathbf{w}_j^{m+1}\| < \frac{1}{2}|x_j|^m + M|x_j|^{p+m}$ for some $M > 0$ and for all j sufficiently large. Then, we get that $\|\mathbf{w}_j^{m+1}\| < |x_j|^m$ for all j sufficiently large, since we can assume that $p \geq 1$ (otherwise the variables \mathbf{w}^m do not appear). Therefore, by Lemma 6.8, $(x_j, \mathbf{w}_j^{m+1}, \mathbf{z}_j^{m+1}) \in S_{m+1}$ if j is big enough. This shows that every orbit in S_m is asymptotic to Γ .

This ends the proof of Theorem 6.1.

ACKNOWLEDGEMENTS

The first, second, and third authors were partially supported by [Ministerio de Ciencia e Innovación](#), Spain, process MTM2016-77642-C2-1-P and PID2019-105621GB-I00. Fourth author was partially supported by [NSF](#)-1800777.

JOURNAL INFORMATION

The *Proceedings of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

REFERENCES

1. M. Abate and F. Tovena, *Parabolic curves in \mathbb{C}^3* , Abstr. Appl. Anal. **1** (2003), no. 5, 275–294.
2. M. Arizzi and J. Raissy, *On Écalle-Hakim's theorems in holomorphic dynamics*, Frontiers in complex dynamics, Princeton Mathematical Series 51, Princeton University Press, Princeton, NJ, 2014, pp. 387–449.
3. W. Balser, *Formal power series and linear systems of meromorphic ordinary differential equations*, Universitext, Springer, New York, 2000.
4. M. A. Barkatou, *An algorithm to compute the exponential part of a formal fundamental matrix solution of a linear differential system*, Appl. Algebra Engrg. Comm. Comput. **8** (1997), no. 1, 1–23.

5. G. Binyamini, *Finiteness properties of formal Lie group actions*, Transform. Groups **20** (2015), no. 4, 939–952.
6. B. Braaksma, *Multisummability of formal power series solutions of nonlinear meromorphic differential equations*, Ann. Inst. Fourier (Grenoble) **42** (1992), no. 3, 517–540.
7. C. Camacho and P. Sad, *Invariant varieties through singularities of holomorphic vector fields*, Ann. of Math. **115** (1982), 579–595.
8. F. Cano, R. Moussu, and J.-P. Rolin, *Non-oscillating integral curves and valuations*, J. Reine Angew. Math. **582** (2005), 107–141.
9. F. Cano, R. Moussu, and F. Sanz, *Pinceaux de courbes intégrales d'un champ de vecteurs analytique*, Astérisque **297** (2004), 1–34.
10. F. Cano, C. Roche, and M. Spivakovsky, *Reduction of singularities of three-dimensional line foliations*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **108** (2014), no. 1, 221–258.
11. D. Cerveau and A. N. Lins, *Codimension two holomorphic foliations*, J. Differential Geom. **113** (2019), no. 3, 385–416.
12. D. Cerveau and J.-F. Mattei, *Formes intégrables holomorphes singulières*, Astérisque, vol. 97, Société Mathématique de France, Paris, 1982, 193 pp.
13. J. Dugundji, *Topology*. Series in Advanced Mathematics. Allyn and Bacon, Boston, MA, 1966.
14. N. Dunford and J. T. Schwartz, *Linear Operators. I. General Theory*, Pure and Applied Mathematics, Vol. 7. Interscience Publishers, Inc., New York; Interscience Publishers, Ltd., London, 1958.
15. J. Écalle, *Théorie itérative: introduction à la théorie des invariants holomorphes*, J. Math. Pures Appl. (9) **54** (1975), 183–258.
16. X. Gómez-Mont and I. Luengo, *Germs of holomorphic vector fields in \mathbb{C}^3 without a separatrix*, Invent. Math. **109** (1992), no. 2, 211–219.
17. R. C. Gunning, *Introduction to holomorphic functions of several variables*, Vol. 2, Wadsworth, AMS, Florence, KY, 1990.
18. M. Hakim, *Analytic transformations of $(\mathbb{C}^p, 0)$ tangent to the identity*, Duke Math. J. **92** (1998), no. 2, 403–428.
19. M. Hakim, *Transformations tangent to the identity. Stable pieces of manifolds*, Prépublication d'Orsay numéro 30, Orsay, 1997.
20. M. Hukuhara, T. Kimura, and T. Matuda, *Equations différentielles ordinaires du premier ordre dans le champ complexe*. The Mathematical Society of Japan, Tokyo, 1961.
21. Y. Ilyashenko and S. Yakovenko, *Lectures on analytic differential equations*, Grad. Stud. Math., vol. 86, AMS, Providence, RI, 2008.
22. L. López-Hernanz, J. Raissy, J. Ribón, and F. Sanz Sánchez, *Stable manifolds of two-dimensional biholomorphisms asymptotic to formal curves*, Int. Math. Res. Not. IMRN (2021), no. 17, 12847–12887.
23. L. López-Hernanz and F. Sanz Sánchez, *Parabolic curves of diffeomorphisms asymptotic to formal invariant curves*, J. Reine Angew. Math. **739** (2018), 277–296.
24. M. Martelo and J. Ribón, *Derived length of solvable groups of local diffeomorphisms*, Math. Ann. **358** (2014), no. 3–4, 701–728.
25. J. Martinet and J.-P. Ramis, *Classification analytique des équations différentielles non linéaires résonnantes du premier ordre*, Ann. Sci. École Norm. Sup. (4) **16** (1983), no. 4, 571–621 (1984).
26. A. S. de Medeiros, *Singular foliations and differential p -forms*, Ann. Fac. Sci. Toulouse Math. (6) **9** (2000), no. 3, 451–466.
27. A. L. Onishchik and E. B. Vinberg, *Lie groups and algebraic groups*, Springer, Berlin, 1990.
28. D. Panazzolo, *Resolution of singularities of real-analytic vector fields in dimension three*, Acta Math. **197** (2006), no. 2, 167–289.
29. R. Pérez-Marco, *Sur une question de Dulac et Fatou*, C. R. Acad. Sci. Paris **321** s. I (1995), 1045–1048.
30. R. Pérez-Marco, *Solution to Briot and Bouquet problem on singularities of differential equations*, arXiv:1802.03630v4.
31. J. P. Ramis and Y. Sibuya, *A new proof of multisummability of formal solutions of non linear meromorphic differential equations*, Ann. Inst. Fourier **33** (1994), 811–848.
32. J. Ribón, *Families of diffeomorphisms without periodic curves*, Michigan Math. J. **53** (2005), no. 2, 243–256.
33. J. Ribón, *Embedding smooth and formal diffeomorphisms through the Jordan-Chevalley decomposition*, J. Differ. Equ. **253** (2012), no. 12, 3211–3231.
34. J. Ribón, *Finite dimensional groups of local diffeomorphisms*, Israel J. Math. **227** (2018), no. 1, 289–329.

35. J. Ribón, *The solvable length of groups of local diffeomorphisms*, J. Reine Angew. Math. **752** (2019), 105–139.
36. K. Saito, *On a generalization of de Rham lemma*, Ann. Inst. Fourier (Grenoble) **26** (1976), no. 2, vii, 165–170.
37. H. L. Turrittin, *Convergent solutions of ordinary linear homogeneous differential equations in the neighborhood of an irregular singular point*, Acta Math. **93** (1955), 27–66.
38. T. Ueda, *Local structure of analytic transformations of two complex variables, I*, J. Math. Kyoto Univ. **26-2** (1986), 233–261.
39. R. J. Walker, *Algebraic curves*, Dover Publications, Inc., New York, 1950.
40. W. Wasow, *Asymptotic expansions for ordinary differential equations*, Interscience, New York, 1965 (re-edited Dover Publications Inc. 1987).
41. X. Zhang, *The embedding flows of C^∞ hyperbolic diffeomorphisms*, J. Differ. Equ. **250** (2011), no. 5, 2283–2298.