



Mathematical aspects of the Kubo formula for electrical conductivity with dissipation

Alexander B. Watson¹ · Dionisios Margetis² · Mitchell Luskin¹

Received: 7 April 2023 / Revised: 13 August 2023 / Accepted: 22 August 2023

© The JJIAM Publishing Committee and Springer Nature Japan KK, part of Springer Nature 2023

Abstract

In this expository article, we present a systematic formal derivation of the Kubo formula for the linear-response current due to a time-harmonic electric field applied to non-interacting, spinless charged particles in a finite volume in the quantum setting. We model dissipation in a transparent way by assuming a sequence of scattering events occurring at random-time intervals modeled by a Poisson distribution. By taking the large-volume limit, we derive special cases of the formula for free electrons, continuum and tight-binding periodic systems, and the nearest-neighbor tight-binding model of graphene. We present the analogous formalism with dissipation to derive the Drude conductivity of classical free particles.

Keywords Kubo formula · Linear-response · Electrical conductivity · Dissipation · Trace formula · von Neumann equation

Mathematics Subject Classification 78A25 · 78A45 · 81S05

✉ Mitchell Luskin
luskin@umn.edu

Alexander B. Watson
abwatson@umn.edu

Dionisios Margetis
diom@umd.edu

¹ Mathematics Department, University of Minnesota Twin Cities, 206 Church St SE, Minneapolis, MN 55455, USA

² Department of Mathematics, and Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742, USA

1 Introduction

1.1 Motivation

The partial differential equations modeling solid mechanics, fluid mechanics, and electromagnetics include material moduli (heat, magnetic, and electric susceptibility, transport coefficients, etc.) that describe the macroscopic response to a small perturbation. A general approach to formulating the linear response of a physical system at equilibrium was proposed by Ryobo Kubo in the seminal papers [1, 2].

In this article, we give a systematic formal derivation of the Kubo formula for the electrical conductivity of non-interacting, spinless charged particles, e.g., spinless¹ electrons in a material. This Kubo formula plays a key role in explanations of many important phenomena such as the quantum Hall effect [1–19].²

The specific form of Kubo's formula we derive is general enough to apply to a wide variety of models of interest. However, we also show, in detail, how this form reduces to well-known simplified forms in common special cases. Our goal is to stimulate studies of this subject in the context of modern materials applications. We believe that this research direction can inspire exciting questions in mathematical modeling and numerical analysis.

1.2 General Kubo formula

We start by presenting the Kubo formula we derive without details, to communicate the main ideas. Consider a system of negatively-charged, non-interacting, spinless quantum particles in d dimensions, that are initially at equilibrium. Applying a time-harmonic electric field $\mathbf{E}(\omega)e^{-i\omega t}$ to this system will excite an electrical current density at the same temporal frequency $\mathbf{J}(\omega)e^{-i\omega t}$. Formally, we can expand the current density in powers of the field as

$$\mathbf{J}(\omega) = \delta_{\omega 0} \mathbf{J}_{\text{eq}} + \sigma(\omega) \mathbf{E}(\omega) + O(E^2), \quad (1.1)$$

where ω denotes the temporal frequency of the field, and $\mathbf{J}_{\text{eq}}$ denotes the current density in equilibrium. The Kubo formula is an expression for the linear coefficient in this expansion, σ , known as the electrical conductivity.

To state the formula, consider, more specifically, N non-interacting spinless particles confined to a subset $\Omega \subset \mathbb{R}^d$ with volume $|\Omega|$. In the quantum setting, this system can be described by the density matrix ρ , acting on a single-particle Hilbert space $\mathcal{H}$, evolving according to the von Neumann equation with Hamiltonian H . We

¹ Although electrons have spin, many phenomena of practical interest can be captured by treating electrons as spinless and then taking into account spin degeneracy at the end of the calculation.

² For further discussion of these references, see Sect. 1.5.

assume that, at equilibrium, ρ equals the Fermi–Dirac distribution function $\Phi(H)$. Then, the Kubo formula can be formulated as [11]

$$\sigma_{lm}(\omega) = -\frac{e^2}{\hbar^2} \tilde{\text{Tr}} \left\{ (\partial_l H) (\mathcal{L}_H - i\omega + \Gamma)^{-1} \partial_m \Phi(H) \right\}, \quad 1 \leq l, m \leq d. \quad (1.2)$$

In the above, $-e, \hbar, \Gamma$ denote the electron charge ($e > 0$), reduced Planck’s constant, and inverse mean scattering time, respectively; $\tilde{\text{Tr}}$ denotes the trace density in $\mathcal{H}$, defined by the trace divided by the volume $|\Omega|$; $\partial_l := -i[X_l, \cdot]$ denotes derivation³ with respect to the components, X_l , where $1 \leq l \leq d$, of the position operator X ; and $\mathcal{L}_H := \frac{i}{\hbar}[H, \cdot]$ denotes the Liouvillian of H , where $[A, B] := AB - BA$ is the commutator of the operators A, B . The notation ∂_l for the derivation is natural, since $-i[X_l, \cdot]$ acts as $\frac{\partial}{\partial k_l}$ in the Fourier and Bloch domains where $\hbar k_l$ is the l -th component of the momentum vector.

It is important to emphasize the generality of formula (1.2). It applies equally well to continuum Schrödinger Hamiltonians as to tight-binding Hamiltonians, and does not require periodicity. However, we will show that it contains many standard forms of the Kubo formula as special cases. For example, the Drude conductivity for free particles (1.3), and the Kubo formula for periodic systems including both Drude and “interband” contributions (1.9).

1.3 Formalism

The first systematic derivation of a linear relation between electric field and current in metals was introduced by Drude [21, 22]. Drude’s model gave qualitative agreement with some experimental results, but also gave incorrect predictions for the specific heat [8], for example, since it considers electrons to be classical particles. The emergence of quantum mechanics led to early improvements to the Drude model, most simply by replacing the Boltzmann–Maxwell statistics by Fermi–Dirac statistics [23]. More systematic quantum theories that utilize the band structure of crystals were proposed by Bloch [24] and others have continued to be actively developed in the physics literature.

To the best of our knowledge, the Kubo formula in the form (1.2), with $\omega = 0$, was first introduced and derived by Bellissard, van Elst, and Schulz-Baldes. The formula played a key role in their investigation of the quantum Hall effect [25], because it allowed for a formulation of the Hall conductivity even in the presence of magnetic fields and random disorder breaking translational symmetry. In particular, they were able to prove that, under appropriate conditions, the large volume limit $|\Omega| \rightarrow \infty$ of (1.2) converges to $\frac{e^2}{2\pi\hbar}$ times an integer. The formula (1.2) with $\omega \neq 0$ was then introduced and rigorously justified in [11]. The works [25] and [11] use the C^* -algebra formalism, which is well-suited for studying quantum systems without translational symmetry. Other discussions of Kubo’s formula across the physics and

³ Here we use the mathematical notion of derivation as an operation which generalizes the ordinary derivative. Derivations are linear operations which satisfy Leibniz’s law; see, e.g., [20].

mathematics literature include, for example, [4–10, 12–16, 26–28]. We discuss these works in more detail in Sect. 1.5.

The first goal of this work is to systematically derive (1.1)–(1.2), following the main ideas of [25]–[11], but avoiding the C^* -algebra formalism, and without striving for mathematical rigor. We hope that this presentation will make the core ideas of these works accessible to a wider array of practitioners across mathematics, physics, and engineering.

The main ideas of this calculation can be summarized as follows. The current density is initially defined as the trace of $\mathbf{j}\rho$, where $\mathbf{j} := \nabla H$ is the current density observable, where $\nabla := (\partial_i)_{1 \leq i \leq d}$. In the absence of dissipation, ρ evolves according to the von Neumann equation, a differential equation with time-periodic coefficients, which models the effect of the applied field $\mathbf{E}$ together with the unperturbed system Hamiltonian H . Dissipation is, then, modeled by a sequence of scattering events which return ρ to its equilibrium distribution. These events occur at random time intervals, whose lengths are modeled by a Poisson distribution with rate Γ . With these ingredients in place, the current density $\mathbf{J}$ appearing in (1.1) is defined as an average as the number of scattering events tends to infinity (in an appropriate sense); see (2.8) and (2.11). The fact that scattering and the parameter Γ are introduced at the level of the von Neumann equation makes our treatment of dissipation transparent. Equations (1.1)–(1.2) are calculated by a careful analysis of the propagator of the von Neumann equation for ρ between scattering events, for which we find a simple representation; see (3.12). We provide details of our modeling assumptions and results in Sect. 2, before giving the detailed derivation in Sect. 3.

1.4 Simplified forms of the Kubo formula

The second goal of the present work is to clarify how (1.2) unifies various well-known simplified forms of the Kubo formula, including those for free particles and for particles in a periodic potential. We provide detailed derivations of these forms starting from the trace formula (1.2) by taking a large volume limit $|\Omega| \rightarrow \infty$. Although we have not found these details in the literature, we are likely not the first to carry out such calculations. These calculations are provided in Sect. 4.

In a similar vein, we also show that the formalism discussed in Sect. 1.3 for deriving (1.1)–(1.2) has a natural classical analog. For example, we can replace the density matrix ρ by a classical phase space density, and its evolution under the von Neumann equation by evolution under the classical Liouville equation. We apply this classical formalism to the simplest case, of free particles, in order to derive the Drude conductivity, in Sect. 5. To the best of our knowledge, this calculation is original to this work. We expect it is possible to derive our classical formalism from the quantum one via a semiclassical limit, but we do not attempt this here.

For free particles, treated either quantum-mechanically via a large-volume limit of (1.2), or classically, we derive (1.1), with σ given by the isotropic Drude conductivity [8, 21, 22]

$$\sigma(\omega) = \frac{e^2 \tilde{N}}{m(\Gamma - i\omega)}, \quad \tilde{N} = \frac{N}{|\Omega|}. \quad (1.3)$$

Here, $\tilde{N}, m$ denote the electron density and mass, respectively. We derive (1.3) from (1.2) in Sect. 4.1.

For particles in a periodic potential, modeled either by a continuum Hamiltonian (Sect. 4.2) or tight-binding model (Sect. 4.3), we derive (1.1), with

$$\sigma(\omega) = \sigma^D(\omega) + \sigma^R(\omega). \quad (1.4)$$

The first term in (1.4), σ^D , known as the Drude conductivity, generalizes (1.3) as

$$\sigma_{lm}^D(\omega) = \frac{e^2 \tilde{N} (m^{\text{eff}})^{-1}_{lm}}{\Gamma - i\omega}.$$

Here, m^{eff} is a possibly anisotropic effective mass tensor, most conveniently defined through the entries of its inverse as

$$(m^{\text{eff}})^{-1}_{lm} := -\frac{1}{\hbar^2 (2\pi)^d \tilde{N}} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \times \frac{d\Phi}{dE}(E_n(\mathbf{k})) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle. \quad (1.5)$$

Here, $1 \leq l, m \leq d$, the symbol Γ^* denotes a unit cell of the reciprocal lattice (Brillouin zone), and $H(\mathbf{k})$, χ_n , and E_n denote the Bloch Hamiltonian, periodic Bloch functions, and Bloch band functions, respectively.

If every band of $H(\mathbf{k})$ is simple (separated from all other bands by gaps), formula (1.5) can be written in two more suggestive forms. The first one,

$$(m^{\text{eff}})^{-1}_{lm} = -\frac{1}{\hbar^2 (2\pi)^d \tilde{N}} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} \frac{d\Phi}{dE}(E_n(\mathbf{k})) \frac{\partial E_n}{\partial k_l}(\mathbf{k}) \frac{\partial E_n}{\partial k_m}(\mathbf{k}) d\mathbf{k}, \quad (1.6)$$

emphasizes derivatives of the Bloch bands, which equal the group velocity of wave-packets formed by superposing Bloch functions; see, e.g., [29–32]. The second one, obtained from (1.6) via integration by parts,

$$(m^{\text{eff}})^{-1}_{lm} = \frac{1}{\hbar^2 (2\pi)^d \tilde{N}} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} \Phi(E_n(\mathbf{k})) \frac{\partial^2 E_n}{\partial k_l \partial k_m}(\mathbf{k}) d\mathbf{k}, \quad (1.7)$$

emphasizes second derivatives of the Bloch bands, which are related to the “effective mass” of wave-packets formed by superposing Bloch functions; see, e.g., [33–35]. Expression (1.7) reduces straightforwardly to the diagonal matrix $\frac{1}{m}$ when there is just one band, $E(\mathbf{k}) = \frac{\hbar^2}{2m} |\mathbf{k}|^2$, so that $\tilde{N} = \frac{1}{(2\pi)^d} \int_{\Gamma^*} \Phi(E(\mathbf{k})) d\mathbf{k}$.

The second term in (1.4), σ^R , has no classical analog. It is known as the regular, or interband, conductivity

$$\sigma_{lm}^R(\omega) := -\frac{e^2}{\hbar^2(2\pi)^d} \sum_{n \neq n' \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \quad (1.8)$$

$$\times \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}) \right\rangle \left\langle \chi_{n'}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) - i\omega + \Gamma \right)}.$$

This term is responsible for materials' optical properties; see, e.g., [10]. An explicit asymptotic formula for the interband conductivity of the one-dimensional Su-Schrieffer-Heeger model [36], which demonstrates the importance of interband “resonances”, was derived in [19].

A convenient expression of the Kubo formula for periodic systems, which takes into account both the Drude and regular contributions, is [3, 8]

$$\sigma_{lm}(\omega) = -\frac{e^2}{\hbar^2(2\pi)^d} \sum_{n, n' \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \quad (1.9)$$

$$\times \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}) \right\rangle \left\langle \chi_{n'}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) - i\omega + \Gamma \right)}.$$

Similar reductions of the electrical conductivity are also possible for certain aperiodic systems under ergodicity assumptions; see Sect. 1.5. In the specific case of the nearest-neighbor tight-binding model of graphene, analytical formulas are available for $H(\mathbf{k})$, E_n , and χ_n , so that the Drude and regular conductivities can be expressed through explicit formulas. We review this case in Sect. 4.4.

1.5 Related work

Discussions of Kubo's formula in the physics literature, in addition to Kubo's original papers [1, 2], can be found in [4–10]. Mathematically rigorous approaches to the Kubo formula, other than [11, 25], include [12–16, 18, 26–28]. We do not review these works in detail since, other than [11, 25], they do not model dissipation. Note that some of these works, e.g., [15, 16, 18, 27, 28], treat the full many-body Schrödinger equation with interactions.

As discussed in Sect. 1.3, large volume limits of (1.2) exist even for some aperiodic systems under ergodicity assumptions [11, 13, 14, 25]. We do not address this case in the present work, except to mention that another case where such limits exist can be found in models of incommensurate twisted bilayers; see, e.g., [17, 37, 38].

1.6 Structure of the paper

We present our precise modeling assumptions, especially the treatment of dissipation, and our results in the quantum case, in Sect. 2. We then derive the general quantum Kubo formula in Sect. 3. We derive simplifications of the Kubo formula in the large volume limit for free particles, particles in periodic potentials modeled by continuum and tight-binding models, and the nearest-neighbor tight-binding model of graphene, in Sect. 4. We derive the classical Drude conductivity, via an analogous formalism, in Sect. 5. We give a conclusion in Sect. 6.

2 Assumptions and main results

In this section, we introduce our formalism for deriving the general Kubo formula (1.1)–(1.2), and state our result precisely. In particular, we introduce the Fermi–Dirac distribution, von Neumann equation for the evolution of the density matrix, and our treatment of dissipation via random scattering events with rate Γ . Details of the calculations which lead to the Kubo formula (2.12) will be given in Sect. 3.

Equilibrium density matrix We consider the density matrix ρ of a system of non-interacting electrons in a region $\Omega \subset \mathbb{R}^d$, with volume $|\Omega|$. We denote the effective single-particle Hamiltonian in the absence of applied fields by H , and the single-particle Hilbert space by $\mathcal{H}$. The equilibrium density matrix is, then, given by the Fermi–Dirac distribution function

$$\Phi(H) := \{e^{\beta(H-\mu)} + 1\}^{-1}. \quad (2.1)$$

Here, $\beta := (k_B T)^{-1}$ is inverse temperature T scaled by Boltzmann's constant k_B , and μ is the chemical potential. The number of particles in the region Ω , N , and the particle density, $\tilde{N}$, are given by

$$N := \text{Tr } \Phi(H), \quad \tilde{N} := \frac{N}{|\Omega|},$$

where Tr denotes the trace over $\mathcal{H}$.

Evolution of density matrix under applied field When a (possibly time-dependent) electric field $\mathbf{E}$ is applied for $t \geq 0$, and in the absence of dissipation (see below), the density matrix evolves according to the von Neumann equation:

$$\frac{d\rho}{dt} = -\mathcal{L}_{H_E}\rho = -\mathcal{L}_H\rho + \frac{e\mathbf{E}}{\hbar} \cdot \nabla\rho, \quad t \geq 0, \quad \rho(0) = \Phi(H). \quad (2.2)$$

Here, H_E denotes the Hamiltonian of the system subject to the applied electric field,

$$H_E := H + e\mathbf{E} \cdot \mathbf{X}.$$

In addition, we denote the Liouvillian operators of H and H_E by

$$\mathcal{L}_H := \frac{i}{\hbar}[H, \cdot], \quad \mathcal{L}_{H_E} := \frac{i}{\hbar}[H_E, \cdot] = \mathcal{L}_H - \frac{e}{\hbar}\mathbf{E} \cdot \nabla;$$

and introduce notation for the derivation

$$\nabla := -i[X, \cdot], \quad (\partial_m)_{1 \leq m \leq d} := (-i[X_m, \cdot])_{1 \leq m \leq d}, \quad (2.3)$$

where $X = (X_m)_{1 \leq m \leq d}$ denotes the position operator.

Modeling of dissipation We introduce dissipation as follows. We assume that scattering events occur at a random sequence of times $0 = t_0 < t_1 < \dots$, such that the differences $\tau_n := t_{n+1} - t_n > 0$ are modeled by a Poisson distribution with rate $\Gamma > 0$, so that

$$\mathbb{P}(\tau_n \leq \nu) = \int_0^\nu \Gamma e^{-\Gamma \tau_n} d\tau_n. \quad (2.4)$$

The average time between scattering events is thus Γ^{-1} . After each scattering event, we assume that the system returns to equilibrium, before evolving again according to (2.2) in-between scattering events. Hence, we assume that ρ evolves according to (for $n \in \mathbb{N}_{\geq 0}$)

$$\frac{d\rho}{dt} = -\mathcal{L}_{H_E}\rho = -\mathcal{L}_H\rho + \frac{e\mathbf{E}}{\hbar} \cdot \nabla \rho, \quad t_n \leq t < t_{n+1}, \quad \rho(t_n) = \Phi(H). \quad (2.5)$$

Remark 2.1 The electron scattering can be modeled more accurately using a scattering kernel. The replacement of such a scattering kernel formalism by the above process involving the (scalar) rate Γ is known as the relaxation time approximation [8, 11].

Current operator We invoke procedures of averaging, both over time and over scattering times modeled by (2.4), of the current density expectation $\mathbf{J}$. This expectation is defined in terms of the trace density $\tilde{\text{Tr}} := \frac{1}{|\Omega|}\text{Tr}$, where Tr is the trace over $\mathcal{H}$, and the current density operator $\mathbf{j}$. The related expression is

$$\mathbf{J} := \tilde{\text{Tr}}\{\mathbf{j}\rho\} = \frac{1}{|\Omega|}\text{Tr}\{\mathbf{j}\rho\}, \quad \mathbf{j} := -\frac{e}{\hbar}\nabla H. \quad (2.6)$$

Thus, the current density in equilibrium is

$$\mathbf{J}_{\text{eq}} := -\frac{e}{\hbar}\tilde{\text{Tr}}\{(\nabla H)\Phi(H)\}. \quad (2.7)$$

The equilibrium current (2.7) vanishes for free particles because of the invariance (even-ness) of the free dispersion relation under $\mathbf{k} \mapsto -\mathbf{k}$. The equilibrium current vanishes, similarly, for particles in a periodic potential, as long as the Bloch bands $E_n(\mathbf{k})$ are even under $\mathbf{k} \mapsto -\mathbf{k}$. A sufficient condition for this to hold is realness of the Hamiltonian $\bar{H} = H$ (often called “time-reversal symmetry” condition). This holds for tight-binding models with real coefficients, or continuum Hamiltonians $-\Delta + V$,

with V real, and hence holds for essentially all models of materials (e.g. graphene, twisted bilayer graphene) in the absence of an applied magnetic field. Alternatively, it suffices to have invariance of the Hamiltonian H under $\mathbf{x} \mapsto -\mathbf{x}$ (often called “parity symmetry”). For a discussion, see, for example, [39].

2.1 Result for constant applied field $\mathbf{E}$

We now make our result precise for the case of time-independent (constant) $\mathbf{E}$. Consider the long-time average of the current density expectation $\mathbf{J}$ (2.6), averaged over scattering times modeled by (2.4)

$$\langle \mathbf{J} \rangle := \lim_{n \rightarrow \infty} \left\langle \frac{1}{t_n} \int_0^{t_n} \mathbf{J}(t') dt' \right\rangle_{\Gamma}. \quad (2.8)$$

More precisely, $\langle \cdot \rangle_{\Gamma}$ denotes taking the expectation over the Poisson distributions defining each $\tau_n := t_{n+1} - t_n$ (2.4). Then, write the vectors $\langle \mathbf{J} \rangle$, $\mathbf{J}_{\text{eq}}$, and $\mathbf{E}$, in terms of their components as

$$\langle \mathbf{J} \rangle = (\langle J_l \rangle)_{1 \leq l \leq d}, \quad \mathbf{J}_{\text{eq}} = (J_{\text{eq},l})_{1 \leq l \leq d}, \quad \mathbf{E} = (E_m)_{1 \leq m \leq d}.$$

Our main result for constant $\mathbf{E}$ is the following expression for $\langle \mathbf{J} \rangle$ in terms of $\mathbf{E}$, known as the Kubo formula:

$$\begin{aligned} \langle J_l \rangle &= J_{\text{eq},l} + \sum_{m=1}^d \sigma_{lm} E_m + O(E^2), \quad 1 \leq l \leq d, \\ \sigma_{lm} &:= -\frac{e^2}{\hbar^2} \tilde{\text{Tr}} \left\{ (\partial_l H) (\mathcal{L}_H + \Gamma)^{-1} \partial_m \Phi(H) \right\}, \quad 1 \leq l, m \leq d. \end{aligned} \quad (2.9)$$

Note that (2.9) agrees with (1.2) when $\omega = 0$. We present a systematic formal derivation of (2.9) in Sect. 3.1.

2.2 Result for time-harmonic applied field $\mathbf{E}$

To state our result precisely for the case of time-dependent $\mathbf{E}$ is somewhat more involved. Let $\mathbf{E}$ be time-harmonic with frequency $\omega \in \mathbb{R}$, so that

$$\mathbf{E}(t; \theta) := \frac{1}{2\pi} e^{-i(\omega t + \theta)} \mathbf{E}(\omega), \quad (2.10)$$

where $\mathbf{E}(\omega)$ is a fixed vector, and $\theta \in [-\pi, \pi)$ is a parameter which we will average over to ease calculations; see (2.11). It seems possible that our results hold without this averaging step, especially since it is not necessary in the classical case (see (5.8)), but we do not investigate this in this work. We define $\rho(t; \theta)$ and $\mathbf{J}(t; \theta)$ by (2.5) and (2.6), with $\mathbf{E}$ given by (2.10).

We consider the following quantity

$$\langle \mathbf{J} \rangle(\omega) := \lim_{n \rightarrow \infty} \left\langle \frac{1}{t_n} \int_0^{t_n} \int_{-\pi}^{\pi} e^{i(\omega t + \theta)} \mathbf{J}(t; \theta) d\theta dt \right\rangle_{\Gamma}. \quad (2.11)$$

Note that we average over the phase θ of the applied field (2.10). This simplifies the calculation considerably; see (3.13). Note that this is unnecessary in the classical case, where the Liouville equation (the classical counterpart of the von Neumann equation used here) can be solved explicitly in simple closed form even for time-dependent fields; see (5.6).

We introduce notation for the vector components of $\langle \mathbf{J} \rangle$ and $\mathbf{E}$ as

$$\langle \mathbf{J} \rangle(\omega) = (\langle J_l \rangle(\omega))_{1 \leq l \leq d}, \quad \mathbf{E}(\omega) = (E_m(\omega))_{1 \leq m \leq d}.$$

Our main result for time-dependent $\mathbf{E}$ is given by the following expressions:

$$\begin{aligned} \langle J_l \rangle(\omega) &= \sum_{m=1}^d \sigma_{lm}(\omega) E_m(\omega) + O(E^2), \quad 1 \leq l \leq d, \\ \sigma_{lm}(\omega) &:= -\frac{e^2}{\hbar^2} \tilde{\text{Tr}} \left\{ (\partial_l H) (\mathcal{L}_H - i\omega + \Gamma)^{-1} \partial_m \Phi(H) \right\}, \quad 1 \leq l, m \leq d. \end{aligned} \quad (2.12)$$

We present a systematic formal derivation of (2.12) in Sect. 3.2. In Sect. 4, we study the limit of (2.12) as $|\Omega| \rightarrow \infty$. Therefore, we derive important reductions of (2.12) in special cases of the Hamiltonian H , namely, for free particles and particles in a periodic potential.

3 Derivation of general Kubo formula

3.1 Derivation for constant applied field $\mathbf{E}$

We start by re-writing the current density expectation averaged over time and scattering events (2.8) using a Tauberian theorem [40] as

$$\langle \mathbf{J} \rangle = \lim_{n \rightarrow \infty} \left\langle \frac{1}{t_n} \int_0^{t_n} \mathbf{J}(t') dt' \right\rangle_{\Gamma} = \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \delta \int_0^{t_n} e^{-\delta t'} \mathbf{J}(t') dt' \right\rangle_{\Gamma}.$$

We can now split the infinite integral into integrals over intervals between scattering events

$$\langle \mathbf{J} \rangle = \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \delta \sum_{m=0}^{n-1} \left\langle \int_{t_m}^{t_{m+1}} e^{-\delta t'} \mathbf{J}(t') dt' \right\rangle_{\Gamma}.$$

Substituting the formula for $\mathbf{J}$ (2.6) and exchanging the orders of trace and time-integral and trace and summation over n , we have

$$\langle J \rangle = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \delta \tilde{\text{Tr}} \left\{ (\nabla H) \sum_{m=0}^{n-1} \left\langle \int_{t_m}^{t_{m+1}} e^{-\delta t'} \rho(t') dt' \right\rangle_{\Gamma} \right\}. \quad (3.1)$$

We now write ρ in terms of the propagator for (2.5)

$$\rho(t') = e^{-(t'-t_n)\mathcal{L}_{H_E}} \Phi(H).$$

We can then compute the integral $\int_{t_m}^{t_{m+1}} e^{-\delta t'} \rho(t') dt' = \int_{t_m}^{t_{m+1}} e^{-\delta t'} e^{-(t'-t_m)\mathcal{L}_{H_E}} \Phi(H) dt' = (\mathcal{L}_{H_E} + \delta)^{-1} e^{-\delta t_m} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_m}) \Phi(H),$

where $\tau_m := t_{m+1} - t_m$. Substituting this back into (3.1) we have

$$\langle J \rangle = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \delta \times \tilde{\text{Tr}} \left\{ (\nabla H) (\mathcal{L}_{H_E} + \delta)^{-1} \sum_{m=0}^{n-1} \left\langle e^{-\delta t_m} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_m}) \right\rangle_{\Gamma} \Phi(H) \right\}. \quad (3.2)$$

We now perform the averaging. We will do so in detail for the $m = 0$ and $m = 1$ terms; the pattern for $m \geq 2$ will then be clear. The $m = 0$ term is (recall that $t_0 = 0$)

$$\begin{aligned} \left\langle e^{-\delta t_0} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_0}) \right\rangle_{\Gamma} &= \int_0^{\infty} \Gamma e^{-\Gamma \tau_0} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_0}) d\tau_0 \\ &= I - \Gamma (\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}. \end{aligned}$$

The $m = 1$ term is

$$\left\langle e^{-\delta t_1} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_1}) \right\rangle_{\Gamma} = \left\langle e^{-\delta \tau_0} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_1}) \right\rangle_{\Gamma},$$

which we have to average over both τ_0 and τ_1 . Since the variables are independent, we can simply average over each variable in turn. The average over τ_1 obviously gives the same factor as before, and the average over τ_0 gives

$$\int_0^{\infty} \Gamma e^{-\Gamma \tau_0} e^{-\delta \tau_0} d\tau_0 = \frac{\Gamma}{\Gamma + \delta}.$$

Hence, the $m = 1$ term in the sum is

$$\left\langle e^{-\delta t_1} (I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_1}) \right\rangle_{\Gamma} = \left(\frac{\Gamma}{\Gamma + \delta} \right) (I - \Gamma (\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}).$$

Continuing in this way, using the fact that

$$e^{-\delta t_m} = \prod_{j=0}^{m-1} e^{-\delta \tau_j},$$

we find that the m th term in the series is

$$\left(\frac{\Gamma}{\Gamma + \delta} \right)^m (I - \Gamma (\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}).$$

We can sum the series, obtaining

$$\sum_{m=0}^{\infty} \left\langle e^{-\delta t_m} \left(I - e^{-(\mathcal{L}_{H_E} + \delta)\tau_m} \right) \right\rangle_{\Gamma} = \frac{\Gamma + \delta}{\delta} (I - \Gamma(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}). \quad (3.3)$$

Substituting (3.3) into (3.2) we have

$$\langle J \rangle = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} (\Gamma + \delta) \tilde{\text{Tr}} \{ (\nabla H)(\mathcal{L}_{H_E} + \delta)^{-1} (I - \Gamma(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}) \Phi(H) \}. \quad (3.4)$$

The second resolvent identity (see, for example, Proposition 1.9 of [41]) essentially states that $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$ for invertible operators A and B . Applying this identity with $A = \mathcal{L}_{H_E} + \delta + \Gamma$ and $B = \mathcal{L}_{H_E} + \delta$, we have

$$\begin{aligned} (\mathcal{L}_{H_E} + \delta + \Gamma)^{-1} &= (\mathcal{L}_{H_E} + \delta)^{-1} - \Gamma(\mathcal{L}_{H_E} + \delta)^{-1}(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1} \\ &= (\mathcal{L}_{H_E} + \delta)^{-1} (I - \Gamma(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1}), \end{aligned}$$

so that (3.4) simplifies to

$$\langle J \rangle = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} (\Gamma + \delta) \tilde{\text{Tr}} \{ (\nabla H)(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1} \Phi(H) \}.$$

Exchanging the order of taking the trace and taking the limit $\delta \downarrow 0$ we arrive at

$$\langle J \rangle = -\frac{e}{\hbar} \tilde{\text{Tr}} \left\{ \lim_{\delta \downarrow 0} (\Gamma + \delta) (\nabla H)(\mathcal{L}_{H_E} + \delta + \Gamma)^{-1} \Phi(H) \right\},$$

where it is trivial to take the limit, resulting in

$$\langle J \rangle = -\frac{e\Gamma}{\hbar} \tilde{\text{Tr}} \{ (\nabla H)(\mathcal{L}_{H_E} + \Gamma)^{-1} \Phi(H) \}. \quad (3.5)$$

We now note that

$$\begin{aligned} (\mathcal{L}_{H_E} + \Gamma)^{-1} &= (\mathcal{L}_H + \Gamma)^{-1} + ((\mathcal{L}_{H_E} + \Gamma)^{-1} - (\mathcal{L}_H + \Gamma)^{-1}) \\ &= (\mathcal{L}_H + \Gamma)^{-1} + (\mathcal{L}_{H_E} + \Gamma)^{-1} \left(\frac{e}{\hbar} \mathbf{E} \cdot \nabla \right) (\mathcal{L}_H + \Gamma)^{-1}, \end{aligned} \quad (3.6)$$

where the second equality follows from

$$\mathcal{L}_{H_E} = \mathcal{L}_H - \frac{e}{\hbar} \mathbf{E} \cdot \nabla \quad (3.7)$$

and the second resolvent identity [41]. Substituting (3.6) into (3.5) separates (3.5) into equilibrium (corresponding to the first term on the right-hand side of (3.6)) and non-equilibrium (corresponding to the second term) contributions to the conductivity. Hence, noting that $(\mathcal{L}_H + \Gamma)^{-1} \Phi(H) = \Gamma^{-1} \Phi(H)$, we simplify (3.5) to

$$\langle J \rangle = J_{\text{eq}} - \frac{e\Gamma}{\hbar} \tilde{\text{Tr}} \left\{ (\nabla H)(\mathcal{L}_{H_E} + \Gamma)^{-1} \left(\frac{e}{\hbar} \mathbf{E} \cdot \nabla \right) (\mathcal{L}_H + \Gamma)^{-1} \Phi(H) \right\},$$

which further simplifies to

$$\langle J \rangle = J_{\text{eq}} - \frac{e^2}{\hbar^2} \tilde{\text{Tr}} \{ (\nabla H) (\mathcal{L}_{H_E} + \Gamma)^{-1} (E \cdot \nabla) \Phi(H) \}. \quad (3.8)$$

It is worth emphasizing that the calculation up to this point has been exact, in the sense that we have not made any expansion in powers of E . We now make such an approximation. Using (3.7) and the second resolvent identity once more, we obtain

$$(\mathcal{L}_{H_E} + \Gamma)^{-1} = (\mathcal{L}_H + \Gamma)^{-1} + O(E).$$

Substituting this into (3.8), and writing everything out in components, we obtain (2.9).

3.2 Derivation for time-harmonic applied field E

The first task is to find a convenient representation of the propagator for the von Neumann equation (2.5) when E is given by (2.10) and thus time-dependent. It is convenient to introduce notation

$$\mathcal{E}(\theta') := \frac{1}{2\pi} e^{-i\theta'} E(\omega), \quad (3.9)$$

so that $E(t; \theta) = \mathcal{E}(\omega t + \theta)$ and the von Neumann equation is

$$\partial_t \rho(t, \theta) = \left(-\mathcal{L}_H + \frac{e}{\hbar} \mathcal{E}(\omega t + \theta) \cdot \nabla \right) \rho(t, \theta), \quad \rho(t_n, \theta) = \Phi(H). \quad (3.10)$$

We introduce a change of variables $t' = t$ and $\theta' = \theta + \omega t$ to (3.10) and a new function P such that $\rho(t; \theta) = P(t', \theta')$ which evolves by the equivalent PDE

$$\begin{aligned} \partial_{t'} P(t', \theta') &= \left(-\mathcal{L}_H + \frac{e}{\hbar} \mathcal{E}(\theta') \cdot \nabla - \omega \partial_{\theta'} \right) P(t', \theta'), \\ P(t_n, \theta') &= \Phi(H). \end{aligned} \quad (3.11)$$

For each θ' , this equation has constant coefficients in t' , so we can write its solution as

$$P(t', \theta') = e^{-(t' - t_n) \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta') \cdot \nabla + \omega \partial_{\theta'} \right)} \Phi(H). \quad (3.12)$$

We note that the von Neumann equation (2.5) is the characteristic equation for (3.11) and that (t, θ) are the characteristic variables.

We now again use a Tauberian theorem to write

$$\langle J \rangle(\omega) = \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \delta \int_0^{t_n} \int_{-\pi}^{\pi} e^{-\delta t} e^{i(\omega t + \theta)} J(t; \theta) d\theta dt \right\rangle_{\Gamma}.$$

Splitting up the time integral and substituting the forms of J and ρ we have

$$\langle J \rangle(\omega) = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \tilde{\text{Tr}} \left\{ (\nabla H) \left\langle \delta \sum_{m=0}^{n-1} \int_{t_m}^{t_{m+1}} \int_{-\pi}^{\pi} e^{-\delta t} e^{i(\omega t + \theta)} P(t, \theta + \omega t) d\theta dt \right\rangle_{\Gamma} \right\}.$$

Changing variable in the θ integral to $\theta' = \theta + \omega t$ we find

$$\langle J \rangle(\omega) = -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \tilde{\text{Tr}} \left\{ (\nabla H) \left\langle \delta \sum_{m=0}^{n-1} \int_{t_m}^{t_{m+1}} \int_{-\pi}^{\pi} e^{-\delta t} e^{i\theta} P(t, \theta) d\theta dt \right\rangle_{\Gamma} \right\}. \quad (3.13)$$

Substituting the form of P we have

$$\begin{aligned} \langle J \rangle(\omega) = & -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \tilde{\text{Tr}} \{ (\nabla H) \delta \\ & \times \left\langle \sum_{m=0}^{n-1} \int_{t_m}^{t_{m+1}} \int_{-\pi}^{\pi} e^{-\delta t} e^{i\theta} e^{-(t-t_m) \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta) \cdot \nabla + \omega \partial_{\theta} \right)} \Phi(H) d\theta dt \right\rangle_{\Gamma} \}, \end{aligned}$$

which we can re-write as

$$\begin{aligned} \langle J \rangle(\omega) = & -\frac{e}{\hbar} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \tilde{\text{Tr}} \{ (\nabla H) \delta \\ & \times \left\langle \sum_{m=0}^{n-1} \int_{t_m}^{t_{m+1}} \int_{-\pi}^{\pi} e^{-\delta t_m} e^{i\theta} e^{-(t-t_m) \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta) \cdot \nabla + \omega \partial_{\theta} + \delta \right)} \Phi(H) d\theta dt \right\rangle_{\Gamma} \}. \end{aligned}$$

Exchanging the orders of the δ and n limits with the trace, integral over θ , and average over the Poisson process, we find

$$\begin{aligned} \langle J \rangle(\omega) = & -\frac{e}{\hbar} \tilde{\text{Tr}} \{ (\nabla H) \\ & \times \int_{-\pi}^{\pi} e^{i\theta} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \delta \left\langle \sum_{m=0}^{n-1} e^{-\delta t_m} \int_{t_m}^{t_{m+1}} e^{-(t-t_m) \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta) \cdot \nabla + \omega \partial_{\theta} + \delta \right)} \Phi(H) dt \right\rangle_{\Gamma} d\theta \}. \end{aligned}$$

We are now back to the setting of the DC case. By the same steps: performing the t integrals, averaging over the Poisson process, summing up the series, and then manipulating using the second resolvent identity [41], we arrive at

$$\begin{aligned} \langle J \rangle(\omega) = & -\frac{e}{\hbar} \tilde{\text{Tr}} \{ (\nabla H) \\ & \times \int_{-\pi}^{\pi} e^{i\theta} \lim_{\delta \downarrow 0} (\Gamma + \delta) \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta) \cdot \nabla + \omega \partial_{\theta} + \delta + \Gamma \right)^{-1} \Phi(H) d\theta \}. \end{aligned}$$

Taking the limit then yields

$$\langle J \rangle(\omega) = -\frac{e\Gamma}{\hbar} \tilde{\text{Tr}} \left\{ (\nabla H) \int_{-\pi}^{\pi} e^{i\theta} \left(\mathcal{L}_H - \frac{e}{\hbar} \mathcal{E}(\theta) \cdot \nabla + \omega \partial_{\theta} + \Gamma \right)^{-1} \Phi(H) d\theta \right\}.$$

Just as in the DC case, we can use the second resolvent identity to isolate the non-equilibrium part, (the equilibrium part contributes 0 for $\omega \neq 0$ because of the integral over θ), and then using $(\mathcal{L}_H + \omega \partial_\theta + \Gamma)^{-1} \Phi(H) = \Gamma^{-1} \Phi(H)$, we arrive at

$$\langle J \rangle(\omega) = -\frac{e^2}{\hbar^2} \tilde{\text{Tr}} \left\{ (\nabla H) \int_{-\pi}^{\pi} e^{i\theta} (\mathcal{L}_{H_\varepsilon} + \omega \partial_\theta + \Gamma)^{-1} \mathcal{E}(\theta) d\theta \cdot \nabla \Phi(H) \right\}.$$

Again, we emphasize that the calculation is, up to this point, exact with respect to E . Just as in the DC case, we now use the second resolvent identity to take the limit $\mathcal{E} \rightarrow 0$ in the resolvent. We can then insert the formula for $\mathcal{E}$ (3.9) and evaluate the integral over θ to obtain

$$\langle J \rangle(\omega) = -\frac{e^2}{\hbar^2} \tilde{\text{Tr}} \left\{ (\nabla H) (\mathcal{L}_H - i\omega + \Gamma)^{-1} E(\omega) \cdot \nabla \Phi(H) \right\} + O(E^2),$$

from which we obtain (2.12).

4 Applications of the Kubo formula

4.1 Free particles

We consider the special case of free particles, taking Ω to be a d -dimensional box with sides of length L , so that $|\Omega| = L^d$. The Hilbert space, $\mathcal{H}$, and Hamiltonian, H , are

$$\mathcal{H} = L^2 \left(\left[-\frac{L}{2}, \frac{L}{2} \right]^d \right), \quad H = -\frac{\hbar^2}{2m} \Delta_x.$$

It is now straightforward to compute the current operator (recall (2.3) and (2.6))

$$j = -\frac{ei}{\hbar} \left[X, \frac{\hbar^2 \Delta_x}{2m} \right] = -\frac{e\hbar}{m} (-i\nabla_x). \quad (4.1)$$

Note that we adopt the notation of putting a subscript x to denote ordinary differential operators in order to distinguish from our notation for derivations (2.3). A natural orthonormal basis for $\mathcal{H}$ is given by the Fourier modes

$$\left\{ \frac{e^{ik \cdot x}}{L^{\frac{d}{2}}} : k \in \frac{2\pi}{L} \mathbb{Z}^d \right\}.$$

The particle number N is given by the trace of the Fermi–Dirac distribution over $\mathcal{H}$, which we can evaluate as

$$N := \text{Tr} \left\{ e^{\beta(H-\mu)} + 1 \right\}^{-1} = \sum_{k \in \frac{2\pi}{L} \mathbb{Z}^d} \left\{ e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} + 1 \right\}^{-1}.$$

Taking the limit $L \rightarrow \infty$, the sum converges to an integral, and we derive the relationship between particle density $\tilde{N}$ and chemical potential μ

$$\tilde{N} := \lim_{L \rightarrow \infty} \frac{N}{L^d} = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \left\{ e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} + 1 \right\}^{-1} d\mathbf{k}. \quad (4.2)$$

It is straightforward to verify that the current density vanishes in equilibrium:

$$\frac{1}{L^d} \text{Tr} \{ j \Phi(H) \} = -\frac{e\hbar}{mL^d} \sum_{\mathbf{k} \in \frac{2\pi}{L} \mathbb{Z}^d} \mathbf{k} \left\{ e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} + 1 \right\}^{-1} = 0,$$

because of even-ness of the Fermi–Dirac distribution with respect to $\mathbf{k}$.

Evaluating the trace in the Kubo formula (2.12), we derive

$$\begin{aligned} \sigma_{lm}(\omega) &= -\frac{e^2}{\hbar^2 L^d} \\ &\times \sum_{\mathbf{k} \in \frac{2\pi}{L} \mathbb{Z}^d} \left(\frac{\hbar^2}{m} k_l \right) (\Gamma - i\omega)^{-1} \left(-\frac{\beta \hbar^2 k_m}{m} \right) e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} \left\{ e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} + 1 \right\}^{-2}. \end{aligned}$$

Anti-symmetry in k implies immediately that $\sigma_{lm}(\omega) = 0$ unless $l = m$, so we derive, after taking $L \rightarrow \infty$,

$$\sigma_{lm}(\omega) = -\delta_{lm} \frac{e^2}{m(2\pi)^d(\Gamma - i\omega)} \int_{\mathbb{R}^d} k_l \frac{\partial}{\partial k_l} \left\{ e^{\beta \left(\frac{\hbar^2 |k|^2}{2m} - \mu \right)} + 1 \right\}^{-1} d\mathbf{k}.$$

Integrating by parts, we arrive at (recall (4.2))

$$\sigma_{lm}(\omega) = \delta_{lm} \frac{e^2}{m(\Gamma - i\omega)} \tilde{N}, \quad \tilde{N} = \frac{N}{L^d},$$

which is consistent with the literature; see, for example, [8].

4.2 Particles in continuum periodic potential

We now consider the case of particles in a periodic potential.

Let A denote a real and invertible $d \times d$ matrix. Then, we can introduce the Bravais lattice and real space unit cell

$$\Lambda := \{ \mathbf{R} = A\mathbf{m} : \mathbf{m} \in \mathbb{Z}^d \}, \quad \Gamma := A \left[-\frac{1}{2}, \frac{1}{2} \right]^d,$$

and the reciprocal lattice and momentum space unit cell (Brillouin zone)

$$\Lambda^* := \{ \mathbf{G} = B\mathbf{n} : \mathbf{n} \in \mathbb{Z}^d \}, \quad \Gamma^* := B \left[-\frac{1}{2}, \frac{1}{2} \right]^d, \quad B := 2\pi A^{-T}.$$

We then set

$$\Omega = \Gamma_L := A \left[-\frac{L}{2}, \frac{L}{2} \right)^d,$$

so that $|\Omega| = |\Gamma|L^d$, where we restrict L to the positive even integers. We consider the Hilbert space, $\mathcal{H}$, and Hamiltonian, H , given by

$$\mathcal{H} := L^2 \left(A \left[-\frac{L}{2}, \frac{L}{2} \right)^d \right), \quad H := -\frac{\hbar^2}{2m} \Delta_x + V(x),$$

where we assume $V \in C^\infty(\mathbb{R}^d, \mathbb{R})$, and

$$V(x + \mathbf{R}) = V(x), \quad x \in \mathbb{R}^d, \mathbf{R} \in \Lambda.$$

The current operator is, again, given by (4.1).

It is natural to introduce the basis of Bloch functions, defined as follows. Consider, for $\mathbf{k} \in \Gamma^*$, the eigenvalue problem defined by

$$H\phi = E\phi, \quad \phi(x + \mathbf{R}; \mathbf{k}) = e^{i\mathbf{k} \cdot \mathbf{R}} \phi(x; \mathbf{k}), \quad x \in \Gamma, \mathbf{R} \in \Lambda,$$

for each $\mathbf{k} \in \Gamma^*$. The operator H has compact resolvent for each $\mathbf{k}$, and hence a sequence of discrete eigenvalues which can be labelled with multiplicity

$$E_1(\mathbf{k}) \leq E_2(\mathbf{k}) \leq \dots \leq E_n(\mathbf{k}) \leq \dots,$$

with associated eigenfunctions $\phi_n(x; \mathbf{k})$. To be consistent with the free case, we assume that these functions are normalized in $L^2(\Gamma)$, i.e.,

$$\int_{\Gamma} |\phi_n(x; \mathbf{k})|^2 dx = 1, \quad n \in \mathbb{N}_{>0}, \mathbf{k} \in \Gamma^*.$$

Introduce the discretized Brillouin zone

$$\Gamma_L^* := \left\{ \mathbf{k} = \frac{B\mathbf{n}}{L} : \mathbf{n} \in \left\{ -\frac{L}{2}, \dots, \frac{L}{2} - 1 \right\}^d \right\},$$

then the set of Bloch functions

$$\left\{ \frac{\phi_n(\cdot; \mathbf{k})}{L^{\frac{d}{2}}} : \mathbf{k} \in \Gamma_L^*, n \in \mathbb{N}_{>0} \right\} \quad (4.3)$$

forms an orthonormal basis of $L^2 \left(A \left[-\frac{L}{2}, \frac{L}{2} \right)^d \right)$. It is straightforward to check that each Bloch function can be decomposed as

$$\phi_n(x; \mathbf{k}) = e^{i\mathbf{k} \cdot x} \chi_n(x; \mathbf{k}),$$

where the χ 's satisfy the eigenvalue problem

$$\begin{aligned} H(\mathbf{k})\chi_n &= E\chi_n, \quad H(\mathbf{k}) := \frac{\hbar^2}{2m}(\mathbf{k} - i\nabla_{\mathbf{x}})^2 + V(\mathbf{x}) \\ \chi_n(\mathbf{x} + \mathbf{R}; \mathbf{k}) &= \chi_n(\mathbf{x}; \mathbf{k}), \quad \mathbf{x} \in \Gamma, \mathbf{R} \in \Lambda, \end{aligned} \quad (4.4)$$

for each $\mathbf{k} \in \Gamma^*$.

It is now natural to evaluate the trace in the basis (4.3) as

$$N = \text{Tr} \left\{ e^{\beta(H-\mu)} + 1 \right\}^{-1} = \sum_{n \in \mathbb{N}_{>0}} \sum_{\mathbf{k} \in \Gamma_L^*} \left\{ e^{\beta(E_n(\mathbf{k})-\mu)} + 1 \right\}^{-1},$$

where the sum over n converges because of the Weyl asymptotics [42], which state that there exist positive constants N, c_1, c_2 such that

$$c_1 n^{\frac{2}{d}} \leq E_n \leq c_2 n^{\frac{2}{d}}, \quad \forall n \geq N.$$

Taking the limit $L \rightarrow \infty$, the sum converges to an integral and we obtain

$$\tilde{N} := \lim_{L \rightarrow \infty} \frac{N}{L^d |\Gamma|} = \frac{1}{(2\pi)^d} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} \left\{ e^{\beta(E_n(\mathbf{k})-\mu)} + 1 \right\}^{-1} d\mathbf{k}.$$

We now want to evaluate the trace formula with respect to the normalized Bloch basis (4.3). We start by considering the matrix elements

$$\begin{aligned} & \frac{1}{L^d} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_l H) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} \\ &= \frac{\hbar^2}{mL^d} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} \left(-i \frac{\partial}{\partial x_l} \right) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x}. \end{aligned}$$

Expanding the Bloch functions we have

$$= \frac{\hbar^2}{mL^d} \int_{\Gamma_L} e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{x}} \overline{\chi_n(\mathbf{x}; \mathbf{k})} \left(k_l - i \frac{\partial}{\partial x_l} \right) \chi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x}. \quad (4.5)$$

Introducing the truncated lattice

$$\Lambda_L := \left\{ \mathbf{R} = A\mathbf{m} : \mathbf{m} \in \left\{ -\frac{L}{2}, \dots, \frac{L}{2} - 1 \right\}^d \right\},$$

we can further simplify (4.5) as

$$= \frac{\hbar^2}{mL^d} \sum_{\mathbf{R} \in \Lambda_L} e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{R}} \int_{\Gamma} e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{x}} \overline{\chi_n(\mathbf{x}; \mathbf{k})} \left(k_l - i \frac{\partial}{\partial x_l} \right) \chi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x}.$$

Evaluating the sum we find that

$$\frac{1}{L^d} \sum_{R \in \Lambda_L} e^{i(k' - k) \cdot R} = \delta_{kk'}, \quad \delta_{kk'} := \begin{cases} 1 & k = k', \\ 0 & k \neq k'. \end{cases}$$

We thus have that

$$\frac{1}{L^d} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_l H) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} = \delta_{kk'} \frac{\hbar^2}{m} \int_{\Gamma} \overline{\chi_n(\mathbf{x}; \mathbf{k})} \left(k_l - i \frac{\partial}{\partial x_l} \right) \chi_{n'}(\mathbf{x}; \mathbf{k}) d\mathbf{x},$$

which can also be written succinctly in terms of $H(\mathbf{k})$ (4.4) as

$$\frac{1}{L^d} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_l H) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} = \delta_{kk'} \int_{\Gamma} \overline{\chi_n(\mathbf{x}; \mathbf{k})} \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{x}; \mathbf{k}) d\mathbf{x}. \quad (4.6)$$

We now simplify

$$\begin{aligned} & \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\mathcal{L}_H - i\omega + \Gamma)^{-1} (\partial_m \Phi(H)) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} \\ &= \frac{1}{\frac{i}{\hbar} (E_n(\mathbf{k}) - E_{n'}(\mathbf{k}')) - i\omega + \Gamma} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_m \Phi(H)) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x}, \end{aligned}$$

which can be further simplified as

$$\begin{aligned} & \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_m \Phi(H)) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} \\ &= -i \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (x_m \Phi(H) - \Phi(H) x_m) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} \\ &= \frac{\Phi(E_{n'}(\mathbf{k}')) - \Phi(E_n(\mathbf{k}))}{E_{n'}(\mathbf{k}') - E_n(\mathbf{k})} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\partial_m H) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x}. \end{aligned}$$

The above expression requires interpretation when $\mathbf{k} = \mathbf{k}'$ and $n = n'$, or in the presence of degeneracies; see (4.8). Using (4.6), we then have that

$$\begin{aligned} & \frac{1}{L^d} \int_{\Gamma_L} \overline{\phi_n(\mathbf{x}; \mathbf{k})} (\mathcal{L}_H - i\omega + \Gamma)^{-1} (\partial_m \Phi(H)) \phi_{n'}(\mathbf{x}; \mathbf{k}') d\mathbf{x} \\ &= \delta_{kk'} \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \int_{\Gamma} \overline{\chi_n(\mathbf{x}; \mathbf{k})} \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_{n'}(\mathbf{x}; \mathbf{k}) d\mathbf{x}}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_n(\mathbf{k}) - E_{n'}(\mathbf{k})) - i\omega + \Gamma \right)}. \end{aligned}$$

We can now evaluate the trace per unit volume as

$$\sigma_{lm}(\omega) = -\frac{e^2}{\hbar^2 |\Gamma| L^d} \sum_{n,n' \in \mathbb{N}_{>0}} \sum_{\mathbf{k} \in \Gamma^*} \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}) \right\rangle \left\langle \chi_{n'}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) - i\omega + \Gamma \right)}.$$

Taking the limit $L \rightarrow \infty$, we obtain

$$\sigma_{lm}(\omega) = -\frac{e^2}{\hbar^2 (2\pi)^d} \sum_{n,n' \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}) \right\rangle \left\langle \chi_{n'}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) - i\omega + \Gamma \right)}, \quad (4.7)$$

which is consistent with the literature; see, for example, Chapter 13 of [8]. It is often convenient to separate the $n = n'$ and $n \neq n'$ terms in this sum. The $n = n'$ terms, known as the Drude conductivity, are

$$\sigma_{lm}^D(\omega) = -\frac{e^2}{\hbar^2 (2\pi)^d (\Gamma - i\omega)} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \times \frac{d\Phi}{dE}(E_n(\mathbf{k})) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle. \quad (4.8)$$

If every band of $H(\mathbf{k})$ is simple, the Drude conductivity can be simplified further as

$$\sigma_{lm}^D(\omega) = -\frac{e^2}{\hbar^2 (2\pi)^d (\Gamma - i\omega)} \sum_{n \in \mathbb{N}_{>0}} \int_{\Gamma^*} \frac{d\Phi}{dE}(E_n(\mathbf{k})) \frac{\partial E_n(\mathbf{k})}{\partial k_l} \frac{\partial E_n(\mathbf{k})}{\partial k_m} d\mathbf{k}.$$

The remaining terms are then known as the regular, or interband, conductivity

$$\sigma_{lm}^R(\omega) = -\frac{e^2}{\hbar^2 (2\pi)^d} \sum_{n \neq n' \in \mathbb{N}_{>0}} \int_{\Gamma^*} d\mathbf{k} \frac{(\Phi(E_{n'}(\mathbf{k})) - \Phi(E_n(\mathbf{k}))) \left\langle \chi_n(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}) \right\rangle \left\langle \chi_{n'}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_m}(\mathbf{k}) \right) \chi_n(\mathbf{k}) \right\rangle}{(E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) \left(\frac{i}{\hbar} (E_{n'}(\mathbf{k}) - E_n(\mathbf{k})) - i\omega + \Gamma \right)}.$$

4.3 Particles in periodic potential in the tight-binding limit

We now consider the case of periodic tight-binding models. Since tight-binding models approximate continuum models, it should be possible to derive the Kubo formula for tight-binding models directly from (4.7). The idea would be to approximate the inner

products of Bloch functions in (4.7) by linear combinations of inner products of atomic orbitals, following, e.g., [8, 43–48]. Here we provide a direct derivation, starting from a discrete model.

Let $\Lambda, \Gamma, A, \Lambda^*, \Gamma^*, B$ be as in Sect. 4.2. We initially consider the infinite Hilbert space

$$\mathcal{H}_{\text{inf}} := \ell^2(\Lambda; \mathbb{C}^N), \quad \mathcal{H}_{\text{inf}} \ni \psi = (\psi_R)_{R \in \Lambda}, \quad \psi_R = (\psi_R^\alpha)_{1 \leq \alpha \leq N},$$

and local Hamiltonians satisfying

$$(H\psi)_R = \sum_{R' \in \Lambda} H_{R,R'} \psi_{R'}, \quad |H_{R,R'}| \leq C e^{-\gamma|R-R'|}.$$

We assume further that H is periodic, so that

$$H_{R+v,R'+v} = H_{R,R'}, \quad R, R', v \in \Lambda.$$

Recall Λ_L and Γ_L^* , the truncated lattice and discretized Brillouin zone. We can, then, consider the restriction of H to the truncated Hilbert space

$$\mathcal{H} := \ell^2(\Lambda_L; \mathbb{C}^N), \quad \mathcal{H} \ni \psi = (\psi_R)_{R \in \Lambda_L}, \quad \psi_R = (\psi_R^\alpha)_{1 \leq \alpha \leq N},$$

where we identify points in Λ related by vectors in $L\Lambda$. A basis of $\mathcal{H}$ is provided by the discrete Bloch functions

$$\left\{ \frac{e^{ik \cdot R} \chi_n(k)}{L^{\frac{d}{2}}} : k \in \Gamma_L^*, 1 \leq n \leq N \right\}, \quad (4.9)$$

where the χ s are eigenvectors of the Bloch Hamiltonian

$$H(k) := \sum_{R' \in \Lambda} H_{0R'} e^{ik \cdot R'}.$$

Evaluating the trace of the Fermi–Dirac distribution with respect to the basis (4.9) to derive N and $\tilde{N}$ proceeds exactly as in Sect. 4.2, except that the sum over n is now a finite sum from 1 to N .

To evaluate the trace formula, we again start by evaluating the matrix elements

$$\begin{aligned} & \frac{1}{L^d} \sum_{R \in \Lambda_L} \overline{e^{ik \cdot R} \chi_n(k)} \sum_{R' \in \Lambda_L} (\partial_l H)_{RR'} e^{ik' \cdot R'} \chi_{n'}(k') \\ &= \frac{1}{L^d} \sum_{R \in \Lambda_L} e^{-ik \cdot R} \overline{\chi_n(k)} \sum_{R' \in \Lambda_L} i(R' - R)_l H_{RR'} e^{ik' \cdot R'} \chi_{n'}(k'). \end{aligned}$$

Changing variables in the sum over R' to R'' , where $R' = R + R''$, we find

$$\begin{aligned}
&= \frac{1}{L^d} \sum_{\mathbf{R} \in \Lambda_L} e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{R}} \overline{\chi_n(\mathbf{k})} \sum_{\mathbf{R}' \in \Lambda_L} iR'_l H_{\mathbf{R}\mathbf{R}+\mathbf{R}'} e^{i\mathbf{k}' \cdot \mathbf{R}'} \chi_{n'}(\mathbf{k}') \\
&= \frac{1}{L^d} \sum_{\mathbf{R} \in \Lambda_L} e^{i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{R}} \overline{\chi_n(\mathbf{k})} \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}').
\end{aligned}$$

It is now straightforward to see that

$$\frac{1}{L^d} \sum_{\mathbf{R} \in \Lambda_L} \overline{e^{i\mathbf{k} \cdot \mathbf{R}} \chi_n(\mathbf{k})} \sum_{\mathbf{R}' \in \Lambda_L} (\partial_l H)_{\mathbf{R}\mathbf{R}'} e^{i\mathbf{k}' \cdot \mathbf{R}'} \chi_{n'}(\mathbf{k}') = \delta_{\mathbf{k}\mathbf{k}'} \overline{\chi_n(\mathbf{k})} \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{n'}(\mathbf{k}).$$

Similar manipulations show that (4.7) also holds for tight-binding models, with the modifications as above.

4.4 Nearest-neighbor tight-binding model of graphene

In this section we consider the nearest-neighbor tight-binding model of graphene; for further discussion of this model, see, e.g., [46, 49]. We will show that the conductivity of this model can be written as a two-dimensional integral with explicit integrand; see (4.10)–(4.11). In a previous work [19], we used an analogous integral representation for the conductivity to derive asymptotic formulas for the interband conductivity of the one-dimensional SSH model [36].

The graphene Bravais lattice vectors are

$$\mathbf{a}_1 := \frac{a}{2} \begin{pmatrix} 1, \sqrt{3} \end{pmatrix}^\top, \quad \mathbf{a}_2 := \frac{a}{2} \begin{pmatrix} -1, \sqrt{3} \end{pmatrix}^\top, \quad \mathbf{A} := (\mathbf{a}_1, \mathbf{a}_2),$$

where $a > 0$ is the lattice constant. Within the $\mathbf{R}$ th fundamental cell of the lattice, there are two atoms, at positions $\mathbf{R} + \boldsymbol{\tau}^A$ and $\mathbf{R} + \boldsymbol{\tau}^B$, which we will take as

$$\boldsymbol{\tau}^A := \mathbf{0}, \quad \boldsymbol{\tau}^B := (0, d)^\top, \quad d := \frac{a}{\sqrt{3}}.$$

The reciprocal lattice vectors are

$$\mathbf{b}_1 := \frac{4\pi}{3d} \begin{pmatrix} \frac{\sqrt{3}}{2}, \frac{1}{2} \end{pmatrix}^\top, \quad \mathbf{b}_2 := \frac{4\pi}{3d} \begin{pmatrix} -\frac{\sqrt{3}}{2}, \frac{1}{2} \end{pmatrix}^\top, \quad \mathbf{B} := (\mathbf{b}_1, \mathbf{b}_2).$$

We consider one orbital per atom, so that there are $N = 2$ orbitals per unit cell. In the $L \rightarrow \infty$ limit, wave-functions of electrons in graphene are elements of $\ell^2(\Lambda; \mathbb{C}^2)$, written as $\psi = (\psi_{\mathbf{R}})_{\mathbf{R} \in \Lambda} = (\psi_{\mathbf{R}}^A, \psi_{\mathbf{R}}^B)_{\mathbf{R} \in \Lambda}^\top$, where $|\psi_{\mathbf{R}}^\sigma|^2$ represents the electron density on sublattice $\sigma \in \{A, B\}$ in the $\mathbf{R}$ th cell. The graphene tight-binding Hamiltonian with nearest-neighbor hopping acts as

$$(H\psi)_R = -t \begin{pmatrix} \psi_R^B + \psi_{R-a_1}^B + \psi_{R-a_2}^B \\ \psi_R^A + \psi_{R+a_1}^A + \psi_{R+a_2}^A \end{pmatrix}, \quad R \in \Lambda,$$

where $t > 0$ is the nearest-neighbor hopping energy.

The Bloch Hamiltonian is given explicitly by

$$H(\mathbf{k}) := -t \begin{pmatrix} 0 & F(\mathbf{k}) \\ \overline{F(\mathbf{k})} & 0 \end{pmatrix}, \quad F(\mathbf{k}) := 1 + e^{-ik \cdot a_1} + e^{-ik \cdot a_2},$$

and can be explicitly diagonalized, with eigenpairs

$$E_{\pm}(\mathbf{k}) := \pm t |F(\mathbf{k})|, \quad \chi_{\pm}(\mathbf{k}) := \frac{1}{\sqrt{2}} \begin{pmatrix} 1, \mp \frac{\overline{F(\mathbf{k})}}{|F(\mathbf{k})|} \end{pmatrix}^{\top}.$$

We can now evaluate the diagonal matrix elements

$$\left\langle \chi_{\pm}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{\pm}(\mathbf{k}) \right\rangle = \pm t \frac{\operatorname{Re} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_l}(\mathbf{k}) \right)}{|F(\mathbf{k})|},$$

and off-diagonal matrix elements

$$\left\langle \chi_{\mp}(\mathbf{k}) \left| \left(\frac{\partial H}{\partial k_l}(\mathbf{k}) \right) \chi_{\pm}(\mathbf{k}) \right\rangle = \pm t i \frac{\operatorname{Im} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_l}(\mathbf{k}) \right)}{|F(\mathbf{k})|},$$

from which follow formulas for the Drude conductivity

$$\begin{aligned} \sigma_{lm}^D(\omega) &= -\frac{e^2 t^2}{\hbar^2 (2\pi)^2 (\Gamma - i\omega)} \sum_{s \in \pm} \int_{\Gamma^*} d\mathbf{k} \\ &\times \frac{d\Phi}{dE}(s|F(\mathbf{k})|) \left(\frac{\operatorname{Re} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_l}(\mathbf{k}) \right)}{|F(\mathbf{k})|} \right) \left(\frac{\operatorname{Re} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_m}(\mathbf{k}) \right)}{|F(\mathbf{k})|} \right), \end{aligned} \quad (4.10)$$

and regular conductivity

$$\begin{aligned} \sigma_{lm}^R(\omega) &= -\frac{e^2 t^2}{\hbar^2 (2\pi)^2} \sum_{s, s' \in \pm, s \neq s'} \int_{\Gamma^*} d\mathbf{k} \\ &\times \frac{(\Phi(s'|F(\mathbf{k})|) - \Phi(s|F(\mathbf{k})|))}{(s'|F(\mathbf{k})| - s|F(\mathbf{k})|)(s'|F(\mathbf{k})| - s|F(\mathbf{k})| - i\omega + \Gamma)} \\ &\times \left(\frac{\operatorname{Im} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_l}(\mathbf{k}) \right)}{|F(\mathbf{k})|} \right) \left(\frac{\operatorname{Im} \left(\overline{F(\mathbf{k})} \frac{\partial F}{\partial k_m}(\mathbf{k}) \right)}{|F(\mathbf{k})|} \right). \end{aligned} \quad (4.11)$$

A similar calculation can be carried out for the Haldane model [50] (for reviews, see [51–53]). Remarkably, for parameter ranges such that the model has a bandgap, direct calculation from (1.8) shows that this model exhibits quantized transverse conductivity at zero temperature, dissipation, and frequency (this phenomenon was first observed in the context of the quantum Hall effect [3]). This calculation lies beyond our present scope.

5 Derivation of classical Drude conductivity

5.1 Assumptions and result

In this section, we show how the formalism introduced in Sect. 2 can be adapted to classical systems to derive the classical Drude conductivity (1.3). The main difference between the present derivation and standard derivations (see, for example, Chapter 1 of [8]) is that we work with the phase space density, which is the natural analog of the quantum density matrix. Working with the phase space density allows us to provide a derivation which more closely parallels the derivation of the quantum Kubo formula (1.2) than standard approaches. We expect that the present classical formalism would emerge naturally from the quantum formalism via a semiclassical limit, with, for example, the time evolution of the density matrix by the von Neumann equation being replaced by the time evolution of the phase space density by the Liouville evolution.

For simplicity, we consider non-interacting, negatively charged, classical particles, with classical Hamiltonian

$$H_0(\mathbf{p}) := \frac{p^2}{2m}.$$

We restrict the system to a d -dimensional box, with sides of length L , and periodic boundary conditions. We work with the phase space density function $\rho(\mathbf{p}, \mathbf{q}, t)$, where $\mathbf{p}$ and $\mathbf{q}$ denote d -dimensional particle momentum and position, respectively.

Equilibrium phase space density At $t = 0$, we assume the system to be at equilibrium, with equilibrium phase space density given by the Maxwellian

$$\rho(\mathbf{p}, \mathbf{q}, 0) = \Phi_M(\mathbf{p}) := \tilde{N} \left(\frac{\beta}{2\pi m} \right)^{\frac{d}{2}} e^{-\beta H_0(\mathbf{p})}. \quad (5.1)$$

Here, $\beta := (kT)^{-1}$ is the inverse temperature T scaled by Boltzmann's constant k , m denotes the particle mass, and $\tilde{N}$ is the particle density. To see that $\tilde{N}$ represents the particle density, we can verify that it equals the total particle number N normalized by the volume L^d , since

$$\frac{N}{L^d} = \tilde{N}, \text{ where } N := \int_{\left[-\frac{L}{2}, \frac{L}{2}\right]^d} \int_{\mathbb{R}^d} \Phi_M(\mathbf{p}) \, d\mathbf{p} \, d\mathbf{q}. \quad (5.2)$$

Remark 5.1 We take the Maxwellian (5.1) as the equilibrium distribution in this section because our intent is to show that the main ideas of Sects. 2 and 3 have close analogs which are entirely classical. If our goal was to give an accurate model of electrons in a real material, it would make sense to replace (5.1) by the Fermi–Dirac distribution (2.1). This approach is known as the Sommerfeld theory; see, e.g., [8].

Evolution of phase space density under applied field For $t > 0$, and in the absence of dissipation (see below), we assume that the particle density evolves according to the von Neumann equation

$$\frac{\partial \rho}{\partial t} + \left(\frac{\partial \rho}{\partial \mathbf{q}} \cdot \frac{\partial H_E}{\partial \mathbf{p}} + \frac{\partial \rho}{\partial \mathbf{p}} \cdot \frac{\partial H_E}{\partial \mathbf{q}} \right) = 0, \quad \rho(\mathbf{p}, \mathbf{q}, 0) = \Phi_M(\mathbf{p}),$$

where H_E is the classical Hamiltonian perturbed by an electric field

$$H_E(\mathbf{p}, \mathbf{q}, t) := H_0(\mathbf{p}) + e\mathbf{E}(t) \cdot \mathbf{q}.$$

Evaluating the partial derivatives of H_E we obtain

$$\frac{\partial \rho}{\partial t} + \left(\frac{\partial \rho}{\partial \mathbf{q}} \cdot \frac{\mathbf{p}}{m} - e \frac{\partial \rho}{\partial \mathbf{p}} \cdot \mathbf{E}(t) \right) = 0, \quad \rho(\mathbf{p}, \mathbf{q}, 0) = \Phi_M(\mathbf{p}). \quad (5.3)$$

The solution of the initial value problem (5.3) is explicit, given by

$$\rho(\mathbf{p}, t) = \Phi_M \left(\mathbf{p} + e \int_0^t \mathbf{E}(t') \, dt' \right).$$

In particular, the particle density $\tilde{N}$ (5.2) is conserved by the evolution.

Modeling of dissipation We introduce dissipation as follows. We assume that scattering events occur at a random sequence of times $0 = t_0 < t_1 < \dots$, such that the differences $\tau_n := t_{n+1} - t_n > 0$ are modeled by a Poisson distribution with rate $\Gamma > 0$, so that

$$\mathbb{P}(\tau_n \leq \nu) = \int_0^\nu \Gamma e^{-\Gamma \tau_n} \, d\tau_n. \quad (5.4)$$

The average time between scattering events is thus $\frac{1}{\Gamma}$. After each scattering event, we assume that the system returns to equilibrium, before evolving again according to (5.3) in between scattering events, i.e., we assume that ρ evolves according to

$$\frac{\partial \rho}{\partial t} + \left(\frac{\partial \rho}{\partial \mathbf{q}} \cdot \frac{\mathbf{p}}{m} - e \frac{\partial \rho}{\partial \mathbf{p}} \cdot \mathbf{E}(t) \right) = 0, \quad \rho(\mathbf{p}, \mathbf{q}, t_n) = \Phi_M(\mathbf{p}), \quad (5.5)$$

for each interval $t_n \leq t < t_{n+1}$, $n \in \mathbb{N}_{\geq 0}$. The solution of (5.5) is again explicit, given by

$$\rho(\mathbf{p}, t) = \Phi_M \left(\mathbf{p} + e \int_{t_n}^t \mathbf{E}(t') dt' \right). \quad (5.6)$$

Current observable Our results involve long time averages, averaged over scattering events, of the current density. The current density is defined by

$$\mathbf{J}(t) := -\frac{e}{mL^d} \int_{[-\frac{L}{2}, \frac{L}{2}]^d} \int_{\mathbb{R}^d} \mathbf{p} \rho(\mathbf{p}, \mathbf{q}, t) d\mathbf{p} d\mathbf{q}.$$

It is straightforward to verify that the current density vanishes at equilibrium, i.e.,

$$-\frac{e}{mL^d} \int_{[-\frac{L}{2}, \frac{L}{2}]^d} \int_{\mathbb{R}^d} \mathbf{p} \Phi_M(\mathbf{p}) d\mathbf{p} d\mathbf{q} = 0,$$

since $\Phi_M(-\mathbf{p}) = \Phi_M(\mathbf{p})$. In between scattering events, i.e., within each interval $t_n \leq t < t_{n+1}$, using the explicit formula (5.6), we have

$$\mathbf{J}(t) = \frac{e^2 \tilde{N}}{m} \int_{t_n}^t \mathbf{E}(t') dt'. \quad (5.7)$$

Result For simplicity, we now assume that the applied field is time-harmonic with frequency $\omega \in \mathbb{R}$, so that

$$\mathbf{E}(t) = e^{-i\omega t} \mathbf{E}(\omega). \quad (5.8)$$

Note that we could allow for a phase $e^{-i\theta}$ in the applied field. Since we do not average over phases in the classical derivation, the result would be to multiply the current by a phase without any other change. In particular, the conductivity would be unaffected.

We then consider the time-averaged current density at frequency ω , averaged over scattering times distributed according to (5.4), as the number of scattering events goes to infinity

$$\langle \mathbf{J} \rangle := \lim_{n \rightarrow \infty} \left\langle \frac{1}{t_n} \int_0^{t_n} e^{i\omega t'} \mathbf{J}(t') dt' \right\rangle_{\Gamma}, \quad (5.9)$$

where $\langle \cdot \rangle_{\Gamma}$ denotes the average over $\tau_0, \dots, \tau_{n-1}$, each distributed according to (5.4). We assume that this limit can be calculated, through a Tauberian theorem, by the sequence of limits

$$\langle \mathbf{J} \rangle(\omega) := \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \int_0^{t_n} e^{(i\omega - \delta)t'} \mathbf{J}(t') dt' \right\rangle_{\Gamma}.$$

Our result is then the following formula, known as the Drude conductivity

$$\langle J \rangle(\omega) = \frac{e^2 \tilde{N}}{m(\Gamma - i\omega)} E(\omega). \quad (5.10)$$

5.2 Derivation of classical Drude conductivity

We first consider the case $\omega = 0$, writing $\langle J \rangle := \langle J \rangle(0)$ for simplicity. We start by writing (5.9) using a Tauberian theorem as

$$\begin{aligned} \langle J \rangle &= \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta \int_{t_m}^{t_{m+1}} e^{-\delta t'} J(t') dt' \right\rangle_{\Gamma} \\ &= \frac{e^2 \tilde{N}}{m} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta \int_{t_m}^{t_{m+1}} e^{-\delta t'} (t' - t_m) dt' \right\rangle_{\Gamma} E \\ &= \frac{e^2 \tilde{N}}{m} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta e^{-\delta t_m} \int_0^{\tau_m} e^{-\delta t'} t' dt' \right\rangle_{\Gamma} E, \end{aligned}$$

using the explicit formula for J between scattering events (5.7). Let us focus on the $m = 0$ term

$$\delta \int_0^{\tau_0} e^{-\delta t'} t' dt' = \frac{1}{\delta} (1 - (1 + \tau_0 \delta) e^{-\delta \tau_0}).$$

Averaging over τ_0 we find

$$\frac{1}{\delta} \left(1 - \int_0^{\infty} \Gamma (1 + \tau_0 \delta) e^{-(\Gamma + \delta) \tau_0} d\tau_0 \right) = \frac{1}{\delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} - \frac{\Gamma \delta}{(\Gamma + \delta)^2} \right) = \frac{1}{\Gamma + \delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} \right).$$

The $n = 1$ term is

$$\frac{1}{\delta} e^{-\delta \tau_0} (1 - (1 + \tau_1 \delta) e^{-\delta \tau_1}).$$

Averaging this term over τ_0 and τ_1 we derive

$$\frac{1}{\Gamma + \delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} \right) \left(\frac{\Gamma}{\Gamma + \delta} \right).$$

More generally, noting that

$$e^{-\delta t_m} = \begin{cases} 1 & m = 0, \\ \prod_{l=0}^{m-1} e^{-\delta \tau_l} & m \geq 1, \end{cases} \quad (5.11)$$

we derive

$$\langle J \rangle = \frac{e^2 \tilde{N}}{m} \lim_{\delta \downarrow 0} \frac{1}{\Gamma + \delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} \right) \lim_{n \rightarrow \infty} \sum_{m=0}^{n-1} \left(\frac{\Gamma}{\Gamma + \delta} \right)^m.$$

Summing the series and taking the limit $n \rightarrow \infty$ we find

$$\langle J \rangle = \frac{e^2 \tilde{N}}{m} \lim_{\delta \downarrow 0} \frac{1}{\delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} \right).$$

Equation (5.10) with $\omega = 0$ then follows by noting that

$$\frac{1}{\delta} \left(1 - \frac{\Gamma}{\Gamma + \delta} \right) = \frac{1}{\delta} \left(1 - \frac{1}{1 + \frac{\delta}{\Gamma}} \right) = \frac{1}{\Gamma} + O(\delta).$$

We now consider the case with $\omega \neq 0$. We follow the exact same steps as above

$$\begin{aligned} \langle J \rangle(\omega) &= \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta \int_{t_m}^{t_{m+1}} e^{(i\omega - \delta)t'} J(t') dt' \right\rangle_{\Gamma} \\ &= \frac{e^2 \tilde{N}}{m(-i\omega)} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta \int_{t_m}^{t_{m+1}} e^{(i\omega - \delta)t'} (e^{-i\omega t'} - e^{-i\omega t_m}) dt' \right\rangle_{\Gamma} E(\omega) \\ &= \frac{e^2 \tilde{N}}{m(-i\omega)} \lim_{\delta \downarrow 0} \lim_{n \rightarrow \infty} \left\langle \sum_{m=0}^{n-1} \delta e^{-\delta t_m} \int_0^{\tau_m} e^{-\delta t'} (1 - e^{i\omega t'}) dt' \right\rangle_{\Gamma} E(\omega). \end{aligned}$$

The $m = 0$ term yields

$$\begin{aligned} \int_0^{\tau_0} e^{-\delta t} (1 - e^{i\omega t}) dt &= \left[\frac{e^{-\delta t}}{-\delta} - \frac{e^{(i\omega - \delta)t}}{i\omega - \delta} \right]_0^{\tau_0} \\ &= \left(\frac{1}{\delta} - \frac{e^{-\delta \tau_0}}{\delta} \right) - \left(\frac{e^{(i\omega - \delta)\tau_0}}{i\omega - \delta} - \frac{1}{i\omega - \delta} \right). \end{aligned}$$

Averaging over τ_0 we find

$$\begin{aligned} \frac{1}{\delta} - \frac{\Gamma}{\delta(\delta + \Gamma)} - \frac{\Gamma}{(i\omega - \delta)(\delta + \Gamma - i\omega)} + \frac{1}{i\omega - \delta} \\ = \frac{1}{\delta + \Gamma} - \frac{1}{\delta + \Gamma - i\omega}. \end{aligned}$$

Again using (5.11), the m th term in the sum is

$$\left(\frac{\Gamma}{\Gamma + \delta} \right)^m \left(\frac{1}{\delta + \Gamma} - \frac{1}{\delta + \Gamma - i\omega} \right),$$

so we arrive at

$$\langle J \rangle(\omega) = \frac{e^2 \tilde{N}}{m(-i\omega)} \lim_{\delta \downarrow 0} \delta \left(\frac{1}{\delta + \Gamma} - \frac{1}{\delta + \Gamma - i\omega} \right) \lim_{n \rightarrow \infty} \sum_{m=0}^{n-1} \left(\frac{\Gamma}{\Gamma + \delta} \right)^m.$$

Summing the series and taking the limit $n \rightarrow \infty$ we arrive at

$$\begin{aligned}\langle J \rangle(\omega) &= \frac{e^2 \tilde{N}}{m(-i\omega)} \lim_{\delta \downarrow 0} (\Gamma + \delta) \left(\frac{1}{\delta + \Gamma} - \frac{1}{\delta + \Gamma - i\omega} \right) \\ &= \frac{e^2 \tilde{N} \Gamma}{m(-i\omega)} \left(\frac{1}{\Gamma} - \frac{1}{\Gamma - i\omega} \right),\end{aligned}$$

which implies equation (5.10) with $\omega \neq 0$.

6 Conclusion

In this paper, we have provided a self-contained, systematic, formal derivation of a widely applicable form of the Kubo formula for the linear response conductivity, under assumptions sufficient for understanding many electronic properties of materials. We have also shown how many commonly-used forms of the Kubo formula, especially those used in the study of crystalline (periodic atomic structure) materials, arise as special cases of the general formula we derive. We hope that this work will stimulate further research into the mathematical modeling of materials' electronic properties.

For example, the Kubo formula is useful in the prediction of the asymptotic behavior of the optical (regular) conductivity as a function of frequency in the limit of small energy bandgap, when a symmetry of the electronic Hamiltonian is broken (see, e.g., [19]). Another interesting direction of research is the accurate numerical calculation of the conductivity in twisted multi-layer graphene, for various twist angles [17, 37, 38].

Acknowledgements The research of the first and third authors (ABW and ML) was supported in part by NSF DMREF Award No. 1922165 and Simons Targeted Grant Award No. 896630. The second author (DM) is grateful to the School of Mathematics of the University of Minnesota for hosting him as an Ordway Distinguished Visitor in the spring of 2022, when part of this work was completed. The authors wish to thank Tony Low and Tobias Stauber for useful discussions.

Data availability No datasets were generated or analysed during the current study.

References

1. Kubo, R.: Statistical-mechanical theory of irreversible processes. I. general theory and simple applications to magnetic and conduction problems. *J. Phys. Soc. Jpn.* **12**, 570–586 (1957). <https://doi.org/10.1143/JPSJ.12.570>
2. Kubo, R., Yokota, M., Nakajima, S.: Statistical-mechanical theory of irreversible processes. II. response to thermal disturbance. *J. Phys. Soc. Jpn.* **12**, 1203–1211 (1957). <https://doi.org/10.1143/JPSJ.12.1203>
3. Thouless, D.J., Kohmoto, M., Nightingale, M.P., den Nijs, M.: Quantized Hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.* **49**, 405–408 (1982)
4. Louie, S.G., Cohen, M.L.: *Conceptual Foundations of Materials: A Standard Model for Ground- and Excited-State Properties*, vol. 2. Elsevier Science, Oxford (2006)
5. Fradkin, E.: *Field Theories of Condensed Matter Physics*, 2nd edn. Cambridge University Press, Cambridge (2013). <https://doi.org/10.1017/CBO9781139015509>

6. Tong, D.: Lectures on Kinetic Theory, (2012). Online lecture notes. <https://www.damtp.cam.ac.uk/user/tong/kinetic.html>
7. Tong, D.: Quantum Hall Effect, (2016). Online book. <https://www.damtp.cam.ac.uk/user/tong/qhe.html>
8. Ashcroft, N.W., Mermin, N.D.: Solid State Physics. Saunders College, Rochester (1976)
9. Kaxiras, E., Joannopoulos, J.D.: Quantum Theory of Materials. Cambridge University Press, Cambridge (2019). <https://doi.org/10.1017/9781139030809>
10. Dresselhaus, M., Dresselhaus, G., Cronin, S., Gomes Souza Filho, A.: Solid State Properties. Springer, Berlin (2018)
11. Schulz-Baldes, H., Bellissard, J.: A kinetic theory for quantum transport in aperiodic media. J. Stat. Phys. **91**, 991–1026 (1998). <https://doi.org/10.1023/a:1023084017398>
12. Lein, M., De Nittis, G.: Linear Response Theory. Springer, Berlin (2017)
13. Bouclet, J.M., Germinet, F., Klein, A., Schenker, J.H.: Linear response theory for magnetic Schrödinger operators in disordered media. J. Funct. Anal. **226**, 301–372 (2005). <https://doi.org/10.1016/j.jfa.2005.02.002>
14. Klein, A., Lenoble, O., Müller, P.: On Mott's formula for the ac-conductivity in the Anderson model. Ann. Math. **166**, 549–577 (2007)
15. Teufel, S.: Non-equilibrium almost-stationary states and linear response for gapped quantum systems. Commun. Math. Phys. **373**, 621–653 (2020). <https://doi.org/10.1007/s00220-019-03407-6>
16. Henheik, J., Teufel, S.: Justifying Kubo' formula for gapped systems at zero temperature: a brief review and some new results. Rev. Math. Phys. **33**, 1–25 (2021). <https://doi.org/10.1142/S0129055X20600041>
17. Cancès, E., Cazeaux, P., Luskin, M.: Generalized Kubo formulas for the transport properties of incommensurate 2D atomic heterostructures. J. Math. Phys. (2017). <https://doi.org/10.1063/1.4984041>
18. Bru, J.-B., de Siqueira Pedra, W.: Lieb-Robinson Bounds for Multi-Commutators and Applications to Response Theory. Springer, Berlin (2017). <https://doi.org/10.1007/978-3-319-45784-0>
19. Margetis, D., Watson, A.B., Luskin, M.: On the Su-Schrieffer-Heeger model of electron transport: Low-temperature optical conductivity by the Mellin transform. Stud. Appl. Math. **151**(2), 555–584 (2023). <https://doi.org/10.1111/sapm.12604>
20. Bourbaki, N.: Algebra I. Springer, Berlin (1989)
21. Drude, P.: Zur elektronentheorie der metalle; II. teil. galvanomagnetische und thermomagnetische effecte. Ann. Phys. **308**(11), 369–402 (1900). <https://doi.org/10.1002/andp.19003081102>
22. Drude, P.: Zur elektronentheorie der metalle. Ann. Phys. **306**(3), 566–613 (1900). <https://doi.org/10.1002/andp.19003060312>
23. Sommerfeld, A.: Zur elektronentheorie der metalle auf grund der Fermischen statistik. Z. Phys. **47**(1–2), 1–324360 (1928)
24. Bloch, F.: Über die quantenmechanik der elektronen in kristallgittern. Z. Phys. **52**(7–8), 555–600 (1928)
25. Bellissard, J., Elst, A.V., Schulz-Baldes, H.: The noncommutative geometry of the quantum hall effect. J. Math. Phys. **35**, 5373–5451 (1994). <https://doi.org/10.1063/1.530758>
26. Elgart, A., Schlein, B.: Adiabatic charge transport and the Kubo formula for Landau-type Hamiltonians. Commun. Pure Appl. Math. **57**, 590–615 (2004). <https://doi.org/10.1002/cpa.20013>
27. Bachmann, S., Bols, A., Roeck, W.D., Fraas, M.: Quantization of conductance in gapped interacting systems. Ann. Henri Poincaré **19**, 695–708 (2018). <https://doi.org/10.1007/s00023-018-0651-0>
28. Bachmann, S., Roeck, W.D., Fraas, M.: The adiabatic theorem and linear response theory for extended quantum systems. Commun. Math. Phys. **361**, 997–1027 (2018). <https://doi.org/10.1007/s00220-018-3117-9>
29. Panati, G., Spohn, H., Teufel, S.: Effective dynamics for Bloch electrons: Peierls substitution and beyond. Commun. Math. Phys. **242**, 547–578 (2003)
30. Panati, G., Spohn, H., Teufel, S.: Motion of electrons in adiabatically perturbed periodic structures. In: Mielke, A. (ed.) Analysis, Modeling and Simulation of Multiscale Problems, pp. 595–617. Springer, Berlin (2006)
31. Lu, J.F., Yang, X.: Asymptotic analysis of quantum dynamics in crystals: the Bloch-Wigner transform, Bloch dynamics and Berry phase. Acta Math. Appl. Sin. **29**, 465–476 (2013). <https://doi.org/10.1007/s10255-011-0095-5>

32. Watson, A.B., Lu, J., Weinstein, M.I.: Wavepackets in inhomogeneous periodic media: effective particle-field dynamics and Berry curvature. *J. Math. Phys.* (2017). <https://doi.org/10.1063/1.4976200>
33. Poupaud, F., Ringhofer, C.: Semi-classical limits in a crystal with exterior potentials and effective mass theorems. *Commun. Partial Differ. Equ.* **21**, 1897–1918 (1996)
34. Allaire, G., Piatnitski, A.: Homogenization of the Schrödinger equation and effective mass theorems. *Commun. Math. Phys.* **258**, 1–22 (2005)
35. Sparber, C.: Effective mass theorems for nonlinear Schrödinger equations. *SIAM J. Appl. Math.* **66**, 820–842 (2006)
36. Su, W.P., Schrieffer, J.R., Heeger, A.J.: Solitons in polyacetylene. *Phys. Rev. Lett.* (1979). <https://doi.org/10.1103/PhysRevLett.42.1698>
37. Massatt, D., Carr, S., Lusk, M.: Efficient computation of Kubo conductivity for incommensurate 2D heterostructures. *Eur. Phys. J. B* (2020). <https://doi.org/10.1140/epjb/e2020-100518-7>
38. Etter, S., Massatt, D., Lusk, M., Ortner, C.: Modeling and computation of Kubo conductivity for 2D incommensurate bilayers. *Multiscale Model. Simul.* **18**, 1525–1564 (2020). <https://doi.org/10.1137/19M1273499>
39. Fefferman, C.L., Weinstein, M.I.: Honeycomb lattice potentials and Dirac points. *J. Am. Math. Soc.* **25**, 1169–1220 (2012). <https://doi.org/10.1090/s0894-0347-2012-00745-0>
40. Korevaar, J.: *Tauberian Theory*. Springer, Berlin (2004). <https://doi.org/10.1007/978-3-662-10225-1>
41. Hislop, P.D., Sigal, I.M.: *Introduction to Spectral Theory*, vol. 113. Springer, Berlin (1996)
42. Maciej, Z.: *Semiclassical Analysis*. Graduate Studies in Mathematics, vol. 138. AMS, Washington, D.C. (2012). <https://doi.org/10.1090/gsm/138>
43. Fefferman, C.L., Shapiro, J., Weinstein, M.I.: Lower bound on quantum tunneling for strong magnetic fields (2020). [arXiv:2006.08025](https://arxiv.org/abs/2006.08025)
44. Shapiro, J., Weinstein, M.I.: Tight-binding reduction and topological equivalence in strong magnetic fields (2020). [arXiv:2010.12097](https://arxiv.org/abs/2010.12097)
45. Shapiro, J., Weinstein, M.I.: Is the continuum SSH model topological? (2021). [arXiv:2107.09146](https://arxiv.org/abs/2107.09146)
46. Fefferman, C.L., Lee-Thorp, J.P., Weinstein, M.I.: Honeycomb Schrödinger operators in the strong binding regime. *Commun. Pure Appl. Math.* **71**, 1178–1270 (2018). <https://doi.org/10.1002/cpa.21735>
47. Helffer, B., Sjöstrand, J.: Multiple wells in the semi-classical limit I. *Commun. Partial Differ. Equ.* **9**, 337–408 (1984). <https://doi.org/10.1080/03605308408820335>
48. Helffer, B., Sjöstrand, J.: Equation de Schrödinger avec champ magnétique et équation de Harper. In: Holden, H., Jensen, A. (eds.) *Schrödinger Operators*. Lecture Notes in Physics, vol. 345, pp. 118–197. Springer, Berlin (1989). https://doi.org/10.1007/3-540-51783-9_19
49. Neto, A.H.C., Guinea, F., Peres, N.M.R., Novoselov, K.S., Geim, A.K.: The electronic properties of graphene. *Rev. Mod. Phys.* **81**, 109–162 (2009)
50. Haldane, F.D.M.: Model for a quantum hall effect without landau levels: condensed-matter realization of the parity anomaly. *Phys. Rev. Lett.* **61**, 2015–2018 (1988). <https://doi.org/10.1103/PhysRevLett.61.2015>
51. Fruchart, M., Carpentier, D.: An introduction to topological insulators. *C R Phys.* **14**, 779–815 (2013)
52. Marcelli, G., Monaco, D., Moscolari, M., Panati, G.: The Haldane model and its localization dichotomy. *Rend. Mat. Appl.* **39**(7), 307–327 (2018)
53. Colbrook, M.J., Horning, A., Thicke, K., Watson, A.B.: Computing spectral properties of topological insulators without artificial truncation or supercell approximation. *IMA J. Appl. Math.* **88**(1), 1–42 (2023). <https://academic.oup.com/imamat/article-pdf/88/1/1/49724502/hxad002.pdf>. <https://doi.org/10.1093/imamat/hxad002>