# First Measurement of Antideuteron Number Fluctuations at Energies Available at the Large Hadron Collider

S. Acharya *et al.*[*]
(ALICE Collaboration)

(Received 5 May 2022; revised 1 July 2022; accepted 15 September 2022; published 25 July 2023)

The first measurement of event-by-event antideuteron number fluctuations in high energy heavy-ion collisions is presented. The measurements are carried out at midrapidity ($|\eta| < 0.8$) as a function of collision centrality in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV using the ALICE detector. A significant negative correlation between the produced antiprotons and antideuterons is observed in all collision centralities. The results are compared with a state-of-the-art coalescence calculation. While it describes the ratio of higher order cumulants of the antideuteron multiplicity distribution, it fails to describe quantitatively the magnitude of the correlation between antiproton and antideuteron production. On the other hand, thermal-statistical model calculations describe all the measured observables within uncertainties only for correlation volumes that are different with respect to those describing proton yields and a similar measurement of net-proton number fluctuations.

DOI: 10.1103/PhysRevLett.131.041901

The production of nuclei and antinuclei in heavy-ion collisions has been extensively studied in the last two decades. Nevertheless, this wealth of results is still not able to clarify the mechanism behind nuclei and antinuclei formation in heavy-ion collisions. Indeed, the two best fitting models, the coalescence [1–3] and the statistical hadronization models (SHM) [4,5], give very similar predictions for the production rates of nuclei and antinuclei in heavy-ion collisions. This similarity calls for new observables to decisively discriminate between these two approaches.

The SHM describes the system as a hadron-resonance gas in thermal equilibrium at hadron emission, hence it predicts particle yields starting from the volume ($V$) and the temperature of the system at chemical freeze-out ($T_{\rm chem}$). The grand canonical ensemble (GCE) formulation of the SHM fits the measured production yields of light hadrons and nuclei in central Pb-Pb collisions at center-of-mass energy ($\sqrt{s_{\rm NN}}$) of 2.76 TeV with $T_{\rm chem} = 156.5$ MeV [6]. The coalescence model uses a different approach to explain the production of nuclei: the size of the nucleon-emitting source, accessible through the analysis of femtoscopic correlations [7], the momentum distribution of the nucleons, as well as the nuclear wave function, are inputs that determine the formation probability of bound states [3,8]. While using statistical hadronization it is possible to compute directly the absolute yields of particles, in the hadron coalescence model the yield of bound states can be computed only relative to the production of its components and as a function of system size.

In a recent model study [9], it is shown that the higher order cumulants of the deuteron yield distribution and correlation between proton ($p$) and deuteron ($d$) production can be used to distinguish between coalescence and SHM. Higher order cumulants $\kappa_m$ of the multiplicity distribution for $m < 4$ and the Pearson correlation coefficient ($\rho_{\rm ab}$) between different identified particles a and b can be expressed as

$$\kappa_1 = \langle n \rangle, \tag{1}$$

$$\kappa_m = \langle (n - \langle n \rangle)^m \rangle, \tag{2}$$

$$\rho_{\rm ab} = \langle (n_{\rm a} - \langle n_{\rm a} \rangle)(n_{\rm b} - \langle n_{\rm b} \rangle) \rangle / \sqrt{\kappa_{2{\rm a}} \kappa_{2{\rm b}}}, \tag{3}$$

where $n$, $\langle n \rangle$, and $m$ are the event-by-event particle numbers, event average of particle numbers, and order of the cumulants, respectively. The $\langle n_{\rm a} \rangle$ ($\langle n_{\rm b} \rangle$) and $\kappa_{2a}$ ($\kappa_{2b}$) are the first and second order cumulants of the multiplicity distribution of particle $a$ ($b$). In the GCE formulation of the SHM, the event-by-event deuteron multiplicity distribution is expected to follow the Poisson distribution [10]. Therefore various ratios between cumulants of different order of the deuteron multiplicity distribution such as $\kappa_2/\kappa_1$, $\kappa_3/\kappa_2$ are equal to unity in the GCE SHM. In a simple coalescence scenario, if deuterons are produced by the coalescence of thermally produced protons and

[*]Full author list given at the end of the Letter.



 

neutrons, then the event-by-event deuteron distribution is expected to deviate from the Poisson baseline [9]. By definition, the coalescence model also introduces a negative correlation between the measured proton and deuteron numbers in the absence of any initial correlation between proton and neutron. On the other hand, one does not expect any correlation between the measured $p$ and $d$ in the GCE SHM as the baryon productions from a thermal source are independent from each other. However, in the canonical ensemble (CE) formulation of the SHM, particle production is constrained by the conservation of the net baryon numbers on an event-by-event basis, which can also introduce a negative correlation between measured proton and deuteron in SHM and a deviation of cumulant ratios from the Poisson baseline [10,11].

In this Letter, the first measurements of the $\kappa_2/\kappa_1$ ratio of antideuteron (antiparticles are used throughout the analysis to avoid the contamination from secondary deuterons coming from spallation processes in the beam pipe) multiplicity distribution and correlation ($\rho_{\bar{p}\,\bar{d}}$) between measured antideuterons ($\bar{d}$) and antiprotons ($\bar{p}$) are presented. Measurements are compared with predictions from the SHM and coalescence model in order to shed light on the deuteron synthesis mechanism. The results presented in this Letter are obtained using data collected during the 2015 Pb-Pb LHC run at $\sqrt{s_{\rm NN}} = 5.02$ TeV.

The ALICE detector and its performance are described in detail in Refs. [12,13]. Collision events are selected by using the information from the V0C and V0A scintillator arrays [14], located on both sides of the interaction point, covering the pseudorapidity intervals $-3.7 < \eta < -1.6$ and $2.8 < \eta < 5.1$, respectively. Events are selected with a minimum-bias (MB) trigger which requires at least one hit in both the V0A and the V0C detectors. In addition, only events with the primary vertex position within 10 cm along the beam axis to the nominal interaction point are selected to benefit from the full acceptance of the detector. Furthermore, to ensure the best possible performance of the detector and proper normalisation of the results, events with more than one reconstructed primary interaction vertex (pile-up events) are rejected. In total, about $100 \times 10^6$ MB events are selected for analysis. Furthermore, the selected events are divided into centrality classes based on the measured amplitude distribution in the V0A and V0C counters as described in Ref. [15]. Central Pb-Pb collisions (head-on collisions) are obtained from the top 10% of the amplitude distribution corresponding to hadronic interactions and peripheral Pb-Pb collisions are obtained from the 70%–80% region of the same distribution.

The charged-particle tracks are reconstructed in the ALICE central barrel with the inner tracking system (ITS) [13] and the time projection chamber (TPC) [16], which are located within a solenoid that provides a homogeneous magnetic field of up to 0.5 T in the direction of the beam axis. These two subsystems provide full azimuthal coverage for charged-particle trajectories in the pseudorapidity interval $|\eta| < 0.8$. The transverse momentum range is restricted to $0.4 < p_T < 1.8$ GeV/$c$ to select the $\bar{p}$ and $\bar{d}$ with high purity. Moreover, to guarantee a track-momentum resolution of 2% in the relevant $p_T$ range and an energy loss ($dE/dx$) resolution in the TPC of 5%, the selected tracks are required to have at least 70 out of a maximum possible 159 reconstructed space points in the TPC, and at least one hit in the two innermost layers of the ITS. This selection also assures a resolution better than 300 μm [13] on the distance of the closest approach to the primary vertex in the plane perpendicular ($\mathrm{DCA}_{xy}$) and parallel ($\mathrm{DCA}_z$) to the beam axis for the selected tracks. In addition, the $\chi^2$ per space point in the TPC and the ITS from the track fit are required to be less than 4 and 36, respectively. Daughter tracks from reconstructed secondary weak-decay kink topologies were rejected and a suppression of the weak-decay particles are obtained by selecting tracks with $|\mathrm{DCA}_z|$ and $|\mathrm{DCA}_{xy}|$ less than 1.0 and 0.1 cm, respectively.

The $\bar{d}$ and $\bar{p}$ are identified via the specific energy loss $dE/dx$ in the gas volume of the TPC and the flight time of a particle from the primary vertex of the collision to the time-of-flight (TOF) detector. The $n(\sigma_i^{\rm TPC})$ variable represents the particle identification (PID) response in the TPC expressed in terms of the deviation between the measured and the expected $dE/dx$ for a particle species $i$, normalized by the detector resolution $\sigma$. The expected $dE/dx$ is computed with a parameterised Bethe-Bloch function [13]. The $\bar{p}$ and $\bar{d}$ are identified using $-2 < |n(\sigma_i^{\rm TPC})| < 4$ in the range $0.4 < p_T < 0.6$ GeV/$c$ and $0.8 < p_T < 1.0$ GeV/$c$, respectively. Particle identification on a track-by-track basis using the TPC is limited to low momenta. Therefore, to identify $\bar{d}$ ($\bar{p}$) in the range $1.0 < p_T < 1.8$ GeV/$c$ ($0.6 < p_T < 0.9$ GeV/$c$), an additional selection of $3.0 < m^2 < 4.2\,\mathrm{GeV}^2/c^4$ ($0.6 < m^2 < 1.2\,\mathrm{GeV}^2/c^4$) using the Time-of-Flight (TOF) [17] detector is applied, where the square of the particle mass, $m^2$, is obtained by combining the information of the flight time with the trajectory length of the particle. The selection of $\bar{d}$ is restricted to the range $0.8 < p_T < 1.8$ GeV/$c$ in order to keep the overall $\bar{d}$ purity above 90%. The $\bar{p}$ selection is restricted to exactly half of the $p_T$ range of $\bar{d}$ according to the coalescence mechanism. This selection results in a purity of the selected $\bar{p}$ sample above 95%. The impurity in $\bar{d}$ selection can lead to an autocorrelation with the selected $\bar{p}$ and affect the $\rho_{\bar{p}\,\bar{d}}$. The effect is negligible in our measurement as the $\bar{d}$ and $\bar{p}$ are mostly selected in separated $p_T$ regions and in the common $p_T$ interval the $\bar{d}$ purity is ∼99%. Selected $\bar{d}$ and $\bar{p}$ numbers in each event are further used to obtain the higher order cumulants and correlation.

Measured cumulants are corrected for the $\bar{d}$ and $\bar{p}$ efficiencies assuming a binomial response of the detectors.

The binomial-based method of efficiency correction [18] is a two-step method. First, the efficiency of $\bar{d}$ and $\bar{p}$ reconstruction in the ALICE detector is obtained using a simulation based on GEANT4, which correctly describes the interaction of $\bar{p}$ and $\bar{d}$ with the material of the detectors [19]. Then, the cumulants and correlation coefficient are corrected for the reconstruction efficiencies using analytic expressions as discussed in Ref. [18]. Typical reconstruction efficiencies of both $\bar{p}$ and $\bar{d}$ in the studied $p_T$ ranges are about 70% and 25% in the TPC and TOF, respectively. The efficiency-corrected cumulants and correlation are further corrected for the centrality bin width effect [20] to suppress the initial volume fluctuations which arise from the initial state (size and shape) fluctuations.

The statistical uncertainties on the efficiency corrected $\kappa_2/\kappa_1$ ratio and $\rho_{\bar{p}\,\bar{d}}$ are obtained by the subsample method [21]. The systematic uncertainties on the observables are estimated by varying the track selection and PID criteria. The systematic uncertainties due to track selection include the variation of the selection criteria on $\mathrm{DCA}_{xy}$, $\mathrm{DCA}_z$, the number of reconstructed space points in the TPC, and the quality of the track fit from their nominal values. The systematic uncertainties due to PID are calculated by varying the default $n(\sigma^{\mathrm{TPC_i}})$ and $m^2$ criteria. Systematic uncertainties due to each of these sources are considered as uncorrelated and the total systematic uncertainty on the observables is obtained by adding all the contributions in quadrature.

The resulting ratio of the second to first order cumulant for $\bar{d}$ is shown in Fig. 1 for different centrality classes. The data is found to be consistent with unity within uncertainties as expected from a Poisson distribution and does not exhibit a significant centrality dependence. Measurements are also compared with estimations from the CE version of the SHM [22] for two different correlation volumes ($V_c$) for baryon number conservation, $V_c = 4.8\ dV/dy$ (orange band in figures) and $V_c = 1.6\ dV/dy$ (green band in figures). The choice of two different $V_c$ is discussed below. In the SHM model the temperature is fixed to T = 155 MeV [5], the volume fitted to the published pion, kaon, and proton yields at midrapidity [23], and the net-baryon number set to 0. Measurements are found to be consistent with the SHM model for both of the $V_c$. In contrast to the corresponding ratio for $p$ and $\bar{p}$ [24,25], no strong dependence on the $V_c$ is seen due to the fact that only a small fraction of the total antibaryon number is carried by $\bar{d}$ [10,26]. Remarkably, the data differs from the calculations of the coalescence model, which predicts a deviation larger than 1% from the Poisson baseline as explained in Ref. [9]. Two shaded bands are shown for the coalescence model: the purple one assumes full correlation among protons and neutrons produced in the collision (model A), while the blue one assumes completely independent proton and neutron production fluctuations (model B). On the other hand, a state of art model




FIG. 1. Second order to first order cumulant ratio of the $\bar{d}$ multiplicity distribution as a function of collision centrality in Pb-Pb collisions at $\sqrt{s_{\mathrm{NN}}} = 5.02$ TeV. Statistical and systematic uncertainties are shown by the bars and boxes, respectively. Measured cumulant ratios are compared with estimations from the CE version of the SHM, from a simple coalescence model and from a MUSIC + UrQMD + Coalescence simulation. The width of the SHM model and MUSIC + UrQMD + Coalescence bands corresponds to the statistical uncertainty of the model estimation, whereas the width of the bands for the coalescence model corresponds to the uncertainty coming from the variation of the coalescence parameters.

calculation coupling coalescence to a hydrodynamical model with hadronic interactions in the final state (MUSIC + UrQMD + COAL) [27] predicts $\kappa_2/\kappa_1$ ratio ~1, in agreement with the experimental data (note that these predictions were updated after acceptance of this Letter). As discussed in [27], the main difference between the coalescence predictions in Fig. 1 and the MUSIC + UrQMD + COAL calculation is due to the different method of implementing baryon number conservation.

Figure 2 shows $\rho_{\bar{p}\,\bar{d}}$ as a function of the collision centrality. A small negative correlation of $O(0.1\%)$ is observed, i.e., in events with at least one $\bar{d}$, there are $O(0.1\%)$ less $\bar{p}$ observed than in an average event. A negative correlation as observed in data is expected by the coalescence model (shown by the blue band in Fig. 2) where $\bar{p}$ and $\bar{n}$ from two independent sources coalesce to produce $\bar{d}$. The same behavior is observed for the MUSIC + UrQMD + COAL calculation. It has to be noted that models based on fully correlated proton and neutron fluctuations (Model A in Ref. [9]) predict values of $\rho$ around 6% and are ruled out by data. On the other hand, the measured negative correlation between $\bar{p}$ and $\bar{d}$ is also expected by the CE version of the SHM which introduces a negative correlation between $\bar{p}$ and $\bar{d}$ through the conservation of a fixed net-baryon number. The predicted




FIG. 2. Pearson correlation between the measured $\bar{p}$ and $\bar{d}$ as a function of collision centrality in Pb-Pb collisions at $\sqrt{s_{\mathrm{NN}}} = 5.02$ TeV. Bars and boxes represent statistical and systematic uncertainties, respectively. Measured correlations are compared with estimations from the CE version of the SHM for two different baryon number conservation volumes, from coalescence model and from MUSIC + UrQMD + COAL.

correlation in the SHM increases with decreasing correlation volume $V_{\mathrm{c}}$ for baryon number conservation which is used in the following for a determination of $V_{\mathrm{c}}$. In order to determine the correlation volume for the baryon quantum number, a $\chi^2$ minimization is performed by varying the $V_{\mathrm{c}}$ parameter in the SHM model and comparing the result to the measured correlation as a function of centrality. The $V_{\mathrm{c}}$ interval probed in this case spans from 1 to 5 units of rapidity, and the value that describes best the measurement is $V_{\mathrm{c}} = 1.6 \pm 0.3\ dV/dy$ with a fit probability of 85%. The SHM configuration with $V_{\mathrm{c}} = 4.8\ dV/dy$ that correctly describes the net-proton number fluctuations in central Pb-Pb collisions [26,28] is compatible within uncertainties with the measured $\rho_{\bar{p}\,\bar{d}}$ only in central collisions. Conversely, this configuration is excluded with a $4\sigma$ confidence level when compared with the measurements in all centrality classes.

Several consistency checks such as the correlation between $\bar{p}$ and $\bar{d}$ from different events, the correlation between antibaryon ($\bar{d}$) and baryon ($p$) were performed for a better understanding of the observed correlation. The correlation between $\bar{p}$ and $\bar{d}$ from mixed events is served as a null hypothesis test of the measurements and the obtained results are consistent with zero as expected. However, a positive correlation is observed between antibaryon and baryon. This positive correlation is expected due to baryon number conservation [10], whereas in simple coalescence model no correlation between baryon and antibaryon is expected as $\bar{d}$ is not produced from the coalescence of $p$.

Figure 3 shows the same Pearson correlation coefficient in three centrality intervals as a function of the $\eta$ acceptance of $\bar{p}$ and $\bar{d}$ selection. The observed anticorrelation is increasing with acceptance, and the effect is more pronounced for peripheral collisions. Simple coalescence calculations do not capture this trend. On the other hand, this measurement should motivate further calculations with more refined coalescence models. The decreasing trend seen in the SHM with $V_c = 1.6\ dV/dy$ describes the experimental data. In the CE version of SHM model, anticorrelation between antibaryons depends on the fraction of antibaryon number in the acceptance out of the total conserved antibaryon numbers [10,11,25,28]. Therefore, the increased negative correlation magnitude with increasing acceptance can be understood as a consequence of baryon number conservation.

In summary, the measurement of $\bar{d}$ production fluctuation is a valuable tool to challenge the nucleosynthesis




FIG. 3. Dependence of $\bar{p}$–$\bar{d}$ correlation on pseudorapidity acceptance of $\bar{p}$ and $\bar{d}$ selection in Pb-Pb collisions at $\sqrt{s_{\mathrm{NN}}} = 5.02$ TeV for three different centrality classes. Measurements are compared with calculations from the CE version of the SHM, coalescence model and MUSIC + UrQMD + COAL.

models used for hadronic collisions. Simple coalescence models, as well as state-of-the-art MUSIC + UrQMD + COAL calculations, fail to fit simultaneously the measurement of the cumulant ratios and the correlation coefficient $\rho_{\bar{p}\,\bar{d}}$. These models show a great sensitivity to the initial correlation between the proton and the neutron production, hence further theoretical developments might improve the comparison with the measurement. In recent studies, state-of-the-art CE SHM models are describing simultaneously proton yields and net-proton fluctuation measurements finding large $V_c \approx 3$–$5 dV/dy$ [26,28,29]. Surprisingly, deuteron production measurements [5] as well as the fluctuation measurements presented here indicate a significantly smaller correlation volume for the baryon number. Under the assumption that $V_c$ is independent of collision centrality, the value $V_c = 1.6 \pm 0.3\ dV/dy$ is obtained. This discrepancy might indicate a different production mechanism for light flavored hadrons and light nuclei. However, more sophisticated approaches including partial chemical equilibrium [30] or the implementation of the interaction of hadrons through phase shift [31,32] could help in resolving this conundrum. The results of this Letter present a severe challenge to the current understanding of nuclei production in heavy-ion collisions at the LHC energies.

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundaçáo de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research—Natural Sciences, the VILLUM FONDEN and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency—BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Academico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Education and Science, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and University Politehnica of Bucharest, Romania; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA), Thailand Science Research and Innovation (TSRI) and National Science, Research and Innovation Fund (NSRF), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TENMAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America

(NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), USA. In addition, individual groups or members have received support from Marie Skłodowska Curie, Strong 2020—Horizon 2020 (Grants No. 824093, No. 896850), European Union; Academy of Finland (Center of Excellence in Quark Matter) (Grants No. 346327, No. 346328), Finland; Programa de Apoyos para la Superación del Personal Académico, UNAM, Mexico.

[1] S. Mrowczynski, Deuteron formation mechanism, J. Phys. G **13**, 1089 (1987).
[2] R. Scheibl and U. W. Heinz, Coalescence and flow in ultrarelativistic heavy ion collisions, Phys. Rev. C **59**, 1585 (1999).
[3] K.-J. Sun, C. M. Ko, and B. Dönigus, Suppression of light nuclei production in collisions of small systems at the Large Hadron Collider, Phys. Lett. B **792**, 132 (2019).
[4] A. Andronic, P. Braun-Munzinger, J. Stachel, and H. Stocker, Production of light nuclei, hypernuclei and their antiparticles in relativistic nuclear collisions, Phys. Lett. B **697**, 203 (2011).
[5] V. Vovchenko, B. Dönigus, and H. Stoecker, Multi plicity dependence of light nuclei production at LHC energies in the canonical statistical model, Phys. Lett. B **785**, 171 (2018).
[6] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, Decoding the phase structure of QCD via particle production at high energy, Nature (London) **561**, 321 (2018).
[7] S. Acharya *et al.* (ALICE Collaboration), Search for a common baryon source in high-multiplicity pp collisions at the LHC, Phys. Lett. B **811**, 135849 (2020).
[8] K. Blum, Kenny Chun Yu Ng, R. Sato, and M. Takimoto, Cosmic rays, antihelium, and an old navy spotlight, Phys. Rev. D **96**, 103021 (2017).
[9] Z. Fecková, J. Steinheimer, B. Tomášik, and M. Bleicher, Formation of deuterons by coalescence: Consequences for deuteron number fluctuations, Phys. Rev. C **93**, 054906 (2016).
[10] P. Braun-Munzinger, B. Friman, K. Redlich, A. Rustamov, and J. Stachel, Relativistic nuclear collisions: Establishing a non-critical baseline for fluctuation measurements, Nucl. Phys. **A1008**, 122141 (2021).
[11] M. Barej and A. Bzdak, Factorial cumulants from global baryon number conservation, Phys. Rev. C **102**, 064908 (2020).
[12] K. Aamodt *et al.* (ALICE Collaboration), The ALICE experiment at the CERN LHC, J. Instrum. **3**, S08002 (2008).
[13] B. B. Abelev *et al.* (ALICE Collaboration), Performance of the ALICE Experiment at the CERN LHC, Int. J. Mod. Phys. A **29**, 1430044 (2014).
[14] E. Abbas *et al.* (ALICE Collaboration), Performance of the ALICE VZERO system, J. Instrum. **8**, P10016 (2013).
[15] B. Abelev *et al.* (ALICE Collaboration), Centrality determination of Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE, Phys. Rev. C **88**, 044909 (2013).
[16] J. Alme *et al.*, The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events, Nucl. Instrum. Methods Phys. Res., Sect. A **622**, 316 (2010).
[17] A. Akindinov *et al.*, Performance of the ALICE time-of-flight detector at the LHC, Eur. Phys. J. Plus **128**, 44 (2013).
[18] T. Nonaka, M. Kitazawa, and S. I. Esumi, More efficient formulas for efficiency correction of cumulants and effect of using averaged efficiency, Phys. Rev. C **95**, 064912 (2017); **103**, 029901(E) (2021).
[19] S. Acharya *et al.* (ALICE Collaboration), Measurement of the Low-Energy Antideuteron Inelastic Cross Section, Phys. Rev. Lett. **125**, 162001 (2020).
[20] X. Luo, J. Xu, B. Mohanty, and N. Xu, Volume fluctuation and auto-correlation effects in the moment analysis of net-proton multiplicity distributions in heavy-ion collisions, J. Phys. G **40**, 105104 (2013).
[21] S. Acharya *et al.* (ALICE Collaboration), Relative particle yield fluctuations in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV, Eur. Phys. J. C **79**, 236 (2019).
[22] V. Vovchenko and H. Stoecker, Thermal-FIST: A package for heavy-ion collisions and hadronic equation of state, Comput. Phys. Commun. **244**, 295 (2019).
[23] S. Acharya *et al.* (ALICE Collaboration), Production of charged pions, kaons, and (anti-)protons in Pb-Pb and inelastic $pp$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. C **101**, 044907 (2020).
[24] M. Abdallah *et al.* (STAR Collaboration), Cumulants and correlation functions of net-proton, proton, and antiproton multiplicity distributions in Au + Au collisions at energies available at the BNL Relativistic Heavy Ion Collider, Phys. Rev. C **104**, 024902 (2021).
[25] V. Vovchenko, V. Koch, and C. Shen, Proton number cumulants and correlation functions in Au-Au collisions at sNN = 7.7–200 GeV from hydrodynamics, Phys. Rev. C **105**, 014904 (2022).
[26] S. Acharya *et al.* (ALICE Collaboration), Global baryon number conservation encoded in net-proton fluctuations measured in Pb-Pb collisions at $\sqrt{s_{\rm NN}} = 2.76$ TeV, Phys. Lett. B **807**, 135564 (2020).
[27] K.-J. Sun and C. M. Ko, Event-by-event anti-deuteron multiplicity fluctuation in Pb + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B **840**, 137864 (2023).
[28] V. Vovchenko and V. Koch, Particlization of an interacting hadron resonance gas with global conservation laws for event-by-event fluctuations in heavy-ion collisions, Phys. Rev. C **103**, 044903 (2021).
[29] V. Vovchenko, B. Dönigus, and H. Stoecker, Canonical statistical model analysis of p-p, p-Pb, and Pb-Pb collisions at energies available at the CERN Large Hadron Collider, Phys. Rev. C **100**, 054906 (2019).
[30] T. Neidig, K. Gallmeister, C. Greiner, M. Bleicher, and V. Vovchenko, Towards solving the puzzle of high temperature light (anti)-nuclei production in ultra-relativistic heavy ion collisions, Phys. Lett. B **827**, 136891 (2022).

[31] A. Andronic, P. Braun-Munzinger, B. Friman, P. M. Lo, K. Redlich, and J. Stachel, The thermal proton yield anomaly in Pb-Pb collisions at the LHC and its resolution, Phys. Lett. B **792**, 304 (2019).

[32] J. Cleymans, P. M. Lo, K. Redlich, and N. Sharma, Multiplicity dependence of (multi)strange baryons in the canonical ensemble with phase shift corrections, Phys. Rev. C **103**, 014904 (2021).

S. Acharya,[123,130] D. Adamová,[85] A. Adler,[68] G. Aglieri Rinella,[32] M. Agnello,[29] N. Agrawal,[49] Z. Ahammed,[130] S. Ahmad,[15] S. U. Ahn,[69] I. Ahuja,[36] A. Akindinov,[138] M. Al-Turany,[97] D. Aleksandrov,[138] B. Alessandro,[54] H. M. Alfanda,[6] R. Alfaro Molina,[65] B. Ali,[15] Y. Ali,[13] A. Alici,[25] N. Alizadehvandchali,[112] A. Alkin,[32] J. Alme,[20] G. Alocco,[50] T. Alt,[62] I. Altsybeev,[138] M. N. Anaam,[6] C. Andrei,[44] A. Andronic,[133] V. Anguelov,[94] F. Antinori,[52] P. Antonioli,[49] C. Anuj,[15] N. Apadula,[73] L. Aphecetche,[102] H. Appelshäuser,[62] S. Arcelli,[25] R. Arnaldi,[54] I. C. Arsene,[19] M. Arslandok,[135] A. Augustinus,[32] R. Averbeck,[97] S. Aziz,[71] M. D. Azmi,[15] A. Badalà,[51] Y. W. Baek,[39] X. Bai,[97] R. Bailhache,[62] Y. Bailung,[46] R. Bala,[90] A. Balbino,[29] A. Baldisseri,[126] B. Balis,[2] D. Banerjee,[4] Z. Banoo,[90] R. Barbera,[26] L. Barioglio,[95] M. Barlou,[77] G. G. Barnaföldi,[134] L. S. Barnby,[84] V. Barret,[123] L. Barreto,[108] C. Bartels,[115] K. Barth,[32] E. Bartsch,[62] F. Baruffaldi,[27] N. Bastid,[123] S. Basu,[74] G. Batigne,[102] D. Battistini,[95] B. Batyunya,[139] D. Bauri,[45] J. L. Bazo Alba,[100] I. G. Bearden,[82] C. Beattie,[135] P. Becht,[97] D. Behera,[46] I. Belikov,[125] A. D. C. Bell Hechavarria,[133] R. Bellwied,[112] S. Belokurova,[138] V. Belyaev,[138] G. Bencedi,[63,134] S. Beole,[24] A. Bercuci,[44] Y. Berdnikov,[138] A. Berdnikova,[94] L. Bergmann,[94] M. G. Besoiu,[61] L. Betev,[32] P. P. Bhaduri,[130] A. Bhasin,[90] I. R. Bhat,[90] M. A. Bhat,[4] B. Bhattacharjee,[40] L. Bianchi,[24] N. Bianchi,[47] J. Bielčík,[35] J. Bielčíková,[85] J. Biernat,[105] A. Bilandzic,[95] G. Biro,[134] S. Biswas,[4] J. T. Blair,[106] D. Blau,[138] M. B. Blidaru,[97] N. Bluhme,[37] C. Blume,[62] G. Boca,[21,53] F. Bock,[86] T. Bodova,[20] A. Bogdanov,[138] S. Boi,[22] J. Bok,[56] L. Boldizsár,[134] A. Bolozdynya,[138] M. Bombara,[36] P. M. Bond,[32] G. Bonomi,[53,129] H. Borel,[126] A. Borissov,[138] H. Bossi,[135] E. Botta,[24] L. Bratrud,[62] P. Braun-Munzinger,[97] M. Bregant,[108] M. Broz,[35] G. E. Bruno,[96,31] M. D. Buckland,[115] D. Budnikov,[138] H. Buesching,[62] S. Bufalino,[29] O. Bugnon,[102] P. Buhler,[101] Z. Buthelezi,[66,119] J. B. Butt,[13] A. Bylinkin,[114] S. A. Bysiak,[105] M. Cai,[6,27] H. Caines,[135] A. Caliva,[97] E. Calvo Villar,[100] J. M. M. Camacho,[107] R. S. Camacho,[43] P. Camerini,[23] F. D. M. Canedo,[108] M. Carabas,[122] F. Carnesecchi,[25] R. Caron,[124,126] J. Castillo Castellanos,[126] F. Catalano,[29] C. Ceballos Sanchez,[139] I. Chakaberia,[73] P. Chakraborty,[45] S. Chandra,[130] S. Chapeland,[32] M. Chartier,[115] S. Chattopadhyay,[130] S. Chattopadhyay,[98] T. G. Chavez,[43] T. Cheng,[6] C. Cheshkov,[124] B. Cheynis,[124] V. Chibante Barroso,[32] D. D. Chinellato,[109] E. S. Chizzali,[95,b] S. Cho,[56] P. Chochula,[32] P. Christakoglou,[83] C. H. Christensen,[82] P. Christiansen,[74] T. Chujo,[121] M. Ciacco,[29] C. Cicalo,[50] L. Cifarelli,[25] F. Cindolo,[49] M. R. Ciupek,[97] G. Clai,[49,c] F. Colamaria,[48] J. S. Colburn,[99] D. Colella,[96,31] A. Collu,[73] M. Colocci,[32] M. Concas,[54,d] G. Conesa Balbastre,[72] Z. Conesa del Valle,[71] G. Contin,[23] J. G. Contreras,[35] M. L. Coquet,[126] T. M. Cormier,[86,a] P. Cortese,[54,128] M. R. Cosentino,[110] F. Costa,[32] S. Costanza,[21,53] P. Crochet,[123] R. Cruz-Torres,[73] E. Cuautle,[63] P. Cui,[6] L. Cunqueiro,[86] A. Dainese,[52] M. C. Danisch,[94] A. Danu,[61] P. Das,[79] P. Das,[4] S. Das,[4] S. Dash,[45] A. De Caro,[28] G. de Cataldo,[48] L. De Cilladi,[24] J. de Cuveland,[37] A. De Falco,[22] D. De Gruttola,[28] N. De Marco,[54] C. De Martin,[23] S. De Pasquale,[28] S. Deb,[46] H. F. Degenhardt,[108] K. R. Deja,[131] R. Del Grande,[95] L. Dello Stritto,[28] W. Deng,[6] P. Dhankher,[18] D. Di Bari,[31] A. Di Mauro,[32] R. A. Diaz,[139,7] T. Dietel,[111] Y. Ding,[6,124] R. Divià,[32] D. U. Dixit,[18] Ø. Djuvsland,[20] U. Dmitrieva,[138] A. Dobrin,[61] B. Dönigus,[62] A. K. Dubey,[130] J. M. Dubinski,[131] A. Dubla,[97] S. Dudi,[89] P. Dupieux,[123] M. Durkac,[104] N. Dzalaiova,[12] T. M. Eder,[133] R. J. Ehlers,[86] V. N. Eikeland,[20] F. Eisenhut,[62] D. Elia,[48] B. Erazmus,[102] F. Ercolessi,[25] F. Erhardt,[88] A. Erokhin,[138] M. R. Ersdal,[20] B. Espagnon,[71] G. Eulisse,[32] D. Evans,[99] S. Evdokimov,[138] L. Fabbietti,[95] M. Faggin,[27] J. Faivre,[72] F. Fan,[6] W. Fan,[73] A. Fantoni,[47] M. Fasel,[86] P. Fecchio,[29] A. Feliciello,[54] G. Feofilov,[138] A. Fernández Téllez,[43] M. B. Ferrer,[32] A. Ferrero,[126] A. Ferretti,[24] V. J. G. Feuillard,[94] J. Figiel,[105] V. Filova,[35] D. Finogeev,[138] G. Fiorenza,[96] F. Flor,[112] A. N. Flores,[106] S. Foertsch,[66] I. Fokin,[94] S. Fokin,[138] E. Fragiacomo,[55] E. Frajna,[134] U. Fuchs,[32] N. Funicello,[28] C. Furget,[72] A. Furs,[138] J. J. Gaardhøje,[82] M. Gagliardi,[24] A. M. Gago,[100] A. Gal,[125] C. D. Galvan,[107] P. Ganoti,[77] C. Garabatos,[97] J. R. A. Garcia,[43] E. Garcia-Solis,[9] K. Garg,[102] C. Gargiulo,[32] A. Garibli,[80] K. Garner,[133]

E. F. Gauger,[106] A. Gautam,[114] M. B. Gay Ducati,[64] M. Germain,[102] S. K. Ghosh,[4] M. Giacalone,[25] P. Gianotti,[47] P. Giubellino,[54,97] P. Giubilato,[27] A. M. C. Glaenzer,[126] P. Glässel,[94] E. Glimos,[118] D. J. Q. Goh,[75] V. Gonzalez,[132] L. H. González-Trueba,[65] S. Gorbunov,[37] M. Gorgon,[2] L. Görlich,[105] S. Gotovac,[33] V. Grabski,[65] L. K. Graczykowski,[131] E. Grecka,[85] L. Greiner,[73] A. Grelli,[57] C. Grigoras,[32] V. Grigoriev,[138] S. Grigoryan,[1,139] F. Grosa,[54] J. F. Grosse-Oetringhaus,[32] R. Grosso,[97] D. Grund,[35] G. G. Guardiano,[109] R. Guernane,[72] M. Guilbaud,[102] K. Gulbrandsen,[82] T. Gunji,[120] W. Guo,[6] A. Gupta,[90] R. Gupta,[90] S. P. Guzman,[43] L. Gyulai,[134] M. K. Habib,[97] C. Hadjidakis,[71] H. Hamagaki,[75] M. Hamid,[6] Y. Han,[136] R. Hannigan,[106] M. R. Haque,[131] A. Harlenderova,[97] J. W. Harris,[135] A. Harton,[9] J. A. Hasenbichler,[32] H. Hassan,[86] D. Hatzifotiadou,[49] P. Hauer,[41] L. B. Havener,[135] S. T. Heckel,[95] E. Hellbär,[97] H. Helstrup,[34] T. Herman,[35] G. Herrera Corral,[8] F. Herrmann,[133] K. F. Hetland,[34] B. Heybeck,[62] H. Hillemanns,[32] C. Hills,[115] B. Hippolyte,[125] B. Hofman,[57] B. Hohlweger,[83] J. Honermann,[133] G. H. Hong,[136] D. Horak,[35] A. Horzyk,[2] R. Hosokawa,[14] Y. Hou,[6] P. Hristov,[32] C. Hughes,[118] P. Huhn,[62] L. M. Huhta,[113] C. V. Hulse,[71] T. J. Humanic,[87] H. Hushnud,[98] A. Hutson,[112] D. Hutter,[37] J. P. Iddon,[115] R. Ilkaev,[138] H. Ilyas,[13] M. Inaba,[121] G. M. Innocenti,[32] M. Ippolitov,[138] A. Isakov,[85] T. Isidori,[114] M. S. Islam,[98] M. Ivanov,[97] V. Ivanov,[138] V. Izucheev,[138] M. Jablonski,[2] B. Jacak,[73] N. Jacazio,[32] P. M. Jacobs,[73] S. Jadlovska,[104] J. Jadlovsky,[104] L. Jaffe,[37] C. Jahnke,[109] M. A. Janik,[131] T. Janson,[68] M. Jercic,[88] O. Jevons,[99] A. A. P. Jimenez,[63] F. Jonas,[86,133] P. G. Jones,[99] J. M. Jowett,[32,97] J. Jung,[62] M. Jung,[62] A. Junique,[32] A. Jusko,[99] M. J. Kabus,[131] J. Kaewjai,[103] P. Kalinak,[58] A. S. Kalteyer,[97] A. Kalweit,[32] V. Kaplin,[138] A. Karasu Uysal,[70] D. Karatovic,[88] O. Karavichev,[138] T. Karavicheva,[138] P. Karczmarczyk,[131] E. Karpechev,[138] V. Kashyap,[79] A. Kazantsev,[138] U. Kebschull,[68] R. Keidel,[137] D. L. D. Keijdener,[57] M. Keil,[32] B. Ketzer,[41] A. M. Khan,[6] S. Khan,[15] A. Khanzadeev,[138] Y. Kharlov,[138] A. Khatun,[15] A. Khuntia,[105] B. Kileng,[34] B. Kim,[16] C. Kim,[16] D. J. Kim,[113] E. J. Kim,[67] J. Kim,[136] J. S. Kim,[39] J. Kim,[94] J. Kim,[67] M. Kim,[94] S. Kim,[17] T. Kim,[136] S. Kirsch,[62] I. Kisel,[37] S. Kiselev,[138] A. Kisiel,[131] J. P. Kitowski,[2] J. L. Klay,[5] J. Klein,[32] S. Klein,[73] C. Klein-Bösing,[133] M. Kleiner,[62] T. Klemenz,[95] A. Kluge,[32] A. G. Knospe,[112] C. Kobdaj,[103] T. Kollegger,[97] A. Kondratyev,[139] N. Kondratyeva,[138] E. Kondratyuk,[138] J. Konig,[62] S. A. Konigstorfer,[95] P. J. Konopka,[32] G. Kornakov,[131] S. D. Koryciak,[2] A. Kotliarov,[85] O. Kovalenko,[78] V. Kovalenko,[138] M. Kowalski,[105] I. Králik,[58] A. Kravčáková,[36] L. Kreis,[97] M. Krivda,[58,99] F. Krizek,[85] K. Krizkova Gajdosova,[35] M. Kroesen,[94] M. Krüger,[62] D. M. Krupova,[35] E. Kryshen,[138] M. Krzewicki,[37] V. Kučera,[32] C. Kuhn,[125] P. G. Kuijer,[83] T. Kumaoka,[121] D. Kumar,[130] L. Kumar,[89] N. Kumar,[89] S. Kundu,[32] P. Kurashvili,[78] A. Kurepin,[138] A. B. Kurepin,[138] A. Kuryakin,[138] S. Kushpil,[85] J. Kvapil,[99] M. J. Kweon,[56] J. Y. Kwon,[56] Y. Kwon,[136] S. L. La Pointe,[37] P. La Rocca,[26] Y. S. Lai,[73] A. Lakrathok,[103] M. Lamanna,[32] R. Langoy,[117] P. Larionov,[47] E. Laudi,[32] L. Lautner,[32,95] R. Lavicka,[101] T. Lazareva,[138] R. Lea,[53,129] J. Lehrbach,[37] R. C. Lemmon,[84] I. León Monzón,[107] M. M. Lesch,[95] E. D. Lesser,[18] M. Lettrich,[95] P. Lévai,[134] X. Li,[10] X. L. Li,[6] J. Lien,[117] R. Lietava,[99] B. Lim,[16] S. H. Lim,[16] V. Lindenstruth,[37] A. Lindner,[44] C. Lippmann,[97] A. Liu,[18] D. H. Liu,[6] J. Liu,[115] I. M. Lofnes,[20] V. Loginov,[138] C. Loizides,[86] P. Loncar,[33] J. A. Lopez,[94] X. Lopez,[123] E. López Torres,[7] P. Lu,[97,116] J. R. Luhder,[133] M. Lunardon,[27] G. Luparello,[55] Y. G. Ma,[38] A. Maevskaya,[138] M. Mager,[32] T. Mahmoud,[41] A. Maire,[125] M. Malaev,[138] N. M. Malik,[90] Q. W. Malik,[19] S. K. Malik,[90] L. Malinina,[139,g] D. Mal'Kevich,[138] D. Mallick,[79] N. Mallick,[46] G. Mandaglio,[30,51] V. Manko,[138] F. Manso,[123] V. Manzari,[48] Y. Mao,[6] G. V. Margagliotti,[23] A. Margotti,[49] A. Marín,[97] C. Markert,[106] M. Marquard,[62] N. A. Martin,[94] P. Martinengo,[32] J. L. Martinez,[112] M. I. Martínez,[43] G. Martínez García,[102] S. Masciocchi,[97] M. Masera,[24] A. Masoni,[50] L. Massacrier,[71] A. Mastroserio,[48,127] A. M. Mathis,[95] O. Matonoha,[74] P. F. T. Matuoka,[108] A. Matyja,[105] C. Mayer,[105] A. L. Mazuecos,[32] F. Mazzaschi,[24] M. Mazzilli,[32] J. E. Mdhluli,[119] A. F. Mechler,[62] Y. Melikyan,[138] A. Menchaca-Rocha,[65] E. Meninno,[101,28] A. S. Menon,[112] M. Meres,[12] S. Mhlanga,[66,111] Y. Miake,[121] L. Micheletti,[54] L. C. Migliorin,[124] D. L. Mihaylov,[95] K. Mikhaylov,[138,139] A. N. Mishra,[134] D. Miśkowiec,[97] A. Modak,[4] A. P. Mohanty,[57] B. Mohanty,[79] M. Mohisin Khan,[15,e] M. A. Molander,[42] Z. Moravcova,[82] C. Mordasini,[95] D. A. Moreira De Godoy,[133] I. Morozov,[138] A. Morsch,[32] T. Mrnjavac,[32] V. Muccifora,[47] E. Mudnic,[33] S. Muhuri,[130] J. D. Mulligan,[73] A. Mulliri,[22] M. G. Munhoz,[108] R. H. Munzer,[62] H. Murakami,[120] S. Murray,[111] L. Musa,[32] J. Musinsky,[58] J. W. Myrcha,[131] B. Naik,[119] R. Nair,[78] B. K. Nandi,[45] R. Nania,[49] E. Nappi,[48] A. F. Nassirpour,[74] A. Nath,[94] C. Nattrass,[118] T. K. Nayak,[79] A. Neagu,[19] A. Negru,[122] L. Nellen,[63] S. V. Nesbo,[34]

G. Neskovic,[37] D. Nesterov,[138] B. S. Nielsen,[82] E. G. Nielsen,[82] S. Nikolaev,[138] S. Nikulin,[138] V. Nikulin,[138] F. Noferini,[49] S. Noh,[11] P. Nomokonov,[139] J. Norman,[115] N. Novitzky,[121] P. Nowakowski,[131] A. Nyanin,[138] J. Nystrand,[20] M. Ogino,[75] A. Ohlson,[74] V. A. Okorokov,[138] J. Oleniacz,[131] A. C. Oliveira Da Silva,[118] M. H. Oliver,[135] A. Onnerstad,[113] C. Oppedisano,[54] A. Ortiz Velasquez,[63] A. Oskarsson,[74] J. Otwinowski,[105] M. Oya,[92] K. Oyama,[75] Y. Pachmayer,[94] S. Padhan,[45] D. Pagano,[53,129] G. Paić,[63] A. Palasciano,[48] S. Panebianco,[126] J. Park,[56] J. E. Parkkila,[32,113] S. P. Pathak,[112] R. N. Patra,[90] B. Paul,[22] H. Pei,[6] T. Peitzmann,[57] X. Peng,[6] L. G. Pereira,[64] H. Pereira Da Costa,[126] D. Peresunko,[138] G. M. Perez,[7] S. Perrin,[126] Y. Pestov,[138] V. Petráček,[35] V. Petrov,[138] M. Petrovici,[44] R. P. Pezzi,[64] S. Piano,[55] M. Pikna,[12] P. Pillot,[102] O. Pinazza,[32,49] L. Pinsky,[112] C. Pinto,[95,26] S. Pisano,[47] M. Płoskoń,[73] M. Planinic,[88] F. Pliquett,[62] M. G. Poghosyan,[86] B. Polichtchouk,[138] S. Politano,[29] N. Poljak,[88] A. Pop,[44] S. Porteboeuf-Houssais,[123] J. Porter,[73] V. Pozdniakov,[139] S. K. Prasad,[4] S. Prasad,[46] R. Preghenella,[49] F. Prino,[54] C. A. Pruneau,[132] I. Pshenichnov,[138] M. Puccio,[32] S. Qiu,[83] L. Quaglia,[24] R. E. Quishpe,[112] S. Ragoni,[99] A. Rakotozafindrabe,[126] L. Ramello,[54,128] F. Rami,[125] S. A. R. Ramirez,[43] T. A. Rancien,[72] R. Raniwala,[91] S. Raniwala,[91] S. S. Räsänen,[42] R. Rath,[46] I. Ravasenga,[83] K. F. Read,[86,118] A. R. Redelbach,[37] K. Redlich,[78,f] A. Rehman,[20] P. Reichelt,[62] F. Reidt,[32] H. A. Reme-Ness,[34] Z. Rescakova,[36] K. Reygers,[94] A. Riabov,[138] V. Riabov,[138] R. Ricci,[28] T. Richert,[74] M. Richter,[19] W. Riegler,[32] F. Riggi,[26] C. Ristea,[61] M. Rodríguez Cahuantzi,[43] K. Røed,[19] R. Rogalev,[138] E. Rogochaya,[139] T. S. Rogoschinski,[62] D. Rohr,[32] D. Röhrich,[20] P. F. Rojas,[43] S. Rojas Torres,[35] P. S. Rokita,[131] F. Ronchetti,[47] A. Rosano,[30,51] E. D. Rosas,[63] A. Rossi,[52] A. Roy,[46] P. Roy,[98] S. Roy,[45] N. Rubini,[25] O. V. Rueda,[74] D. Ruggiano,[131] R. Rui,[23] B. Rumyantsev,[139] P. G. Russek,[2] R. Russo,[83] A. Rustamov,[80] E. Ryabinkin,[138] Y. Ryabov,[138] A. Rybicki,[105] H. Rytkonen,[113] W. Rzesa,[131] O. A. M. Saarimaki,[42] R. Sadek,[102] S. Sadovsky,[138] J. Saetre,[20] K. Šafařík,[35] S. K. Saha,[130] S. Saha,[79] B. Sahoo,[45] P. Sahoo,[45] R. Sahoo,[46] S. Sahoo,[59] D. Sahu,[46] P. K. Sahu,[59] J. Saini,[130] S. Sakai,[121] M. P. Salvan,[97] S. Sambyal,[90] T. B. Saramela,[108] D. Sarkar,[132] N. Sarkar,[130] P. Sarma,[40] V. M. Sarti,[95] M. H. P. Sas,[135] J. Schambach,[86] H. S. Scheid,[62] C. Schiaua,[44] R. Schicker,[94] A. Schmah,[94] C. Schmidt,[97] H. R. Schmidt,[93] M. O. Schmidt,[32] M. Schmidt,[93] N. V. Schmidt,[62,86] A. R. Schmier,[118] R. Schotter,[125] J. Schukraft,[32] K. Schwarz,[97] K. Schweda,[97] G. Scioli,[25] E. Scomparin,[54] J. E. Seger,[14] Y. Sekiguchi,[120] D. Sekihata,[120] I. Selyuzhenkov,[97,138] S. Senyukov,[125] J. J. Seo,[56] D. Serebryakov,[138] L. Šerkšnytė,[95] A. Sevcenco,[61] T. J. Shaba,[66] A. Shabanov,[138] A. Shabetai,[102] R. Shahoyan,[32] W. Shaikh,[98] A. Shangaraev,[138] A. Sharma,[89] D. Sharma,[45] H. Sharma,[105] M. Sharma,[90] N. Sharma,[89] S. Sharma,[90] U. Sharma,[90] A. Shatat,[71] O. Sheibani,[112] K. Shigaki,[92] M. Shimomura,[76] S. Shirinkin,[138] Q. Shou,[38] Y. Sibiriak,[138] S. Siddhanta,[50] T. Siemiarczuk,[78] T. F. Silva,[108] D. Silvermyr,[74] T. Simantathammakul,[103] G. Simonetti,[32] B. Singh,[90] B. Singh,[95] R. Singh,[79] R. Singh,[90] R. Singh,[46] V. K. Singh,[130] V. Singhal,[130] T. Sinha,[98] B. Sitar,[12] M. Sitta,[54,128] T. B. Skaali,[19] G. Skorodumovs,[94] M. Slupecki,[42] N. Smirnov,[135] R. J. M. Snellings,[57] E. H. Solheim,[19] C. Soncco,[100] J. Song,[112] A. Songmoolnak,[103] F. Soramel,[27] S. Sorensen,[118] R. Spijkers,[83] I. Sputowska,[105] J. Staa,[74] J. Stachel,[94] I. Stan,[61] P. J. Steffanic,[118] S. F. Stiefelmaier,[94] D. Stocco,[102] I. Storehaug,[19] M. M. Storetvedt,[34] P. Stratmann,[133] S. Strazzi,[25] C. P. Stylianidis,[83] A. A. P. Suaide,[108] C. Suire,[71] M. Sukhanov,[138] M. Suljic,[32] V. Sumberia,[90] S. Sumowidagdo,[81] S. Swain,[59] A. Szabo,[12] I. Szarka,[12] U. Tabassam,[13] S. F. Taghavi,[95] G. Taillepied,[97,123] J. Takahashi,[109] G. J. Tambave,[20] S. Tang,[6,123] Z. Tang,[116] J. D. Tapia Takaki,[114] N. Tapus,[122] L. A. Tarasovicova,[133] M. G. Tarzila,[44] A. Tauro,[32] G. Tejeda Muñoz,[43] A. Telesca,[32] L. Terlizzi,[24] C. Terrevoli,[112] G. Tersimonov,[3] S. Thakur,[130] D. Thomas,[106] R. Tieulent,[124] A. Tikhonov,[138] A. R. Timmins,[112] M. Tkacik,[104] T. Tkacik,[104] A. Toia,[62] N. Topilskaya,[138] M. Toppi,[47] F. Torales-Acosta,[18] T. Tork,[71] A. G. Torres Ramos,[31] A. Trifiró,[30,51] A. S. Triolo,[30,51] S. Tripathy,[49] T. Tripathy,[45] S. Trogolo,[32] V. Trubnikov,[3] W. H. Trzaska,[113] T. P. Trzcinski,[131] A. Tumkin,[138] R. Turrisi,[52] T. S. Tveter,[19] K. Ullaland,[20] B. Ulukutlu,[95] A. Uras,[124] M. Urioni,[53,129] G. L. Usai,[22] M. Vala,[36] N. Valle,[21] S. Vallero,[54] L. V. R. van Doremalen,[57] M. van Leeuwen,[83] C. A. van Veen,[94] R. J. G. van Weelden,[83] P. Vande Vyvre,[32] D. Varga,[134] Z. Varga,[134] M. Varga-Kofarago,[134] M. Vasileiou,[77] A. Vasiliev,[138] O. Vázquez Doce,[95] V. Vechernin,[138] E. Vercellin,[24] S. Vergara Limón,[43] L. Vermunt,[57] R. Vértesi,[134] M. Verweij,[57] L. Vickovic,[33] Z. Vilakazi,[119] O. Villalobos Baillie,[99] G. Vino,[48] A. Vinogradov,[138] T. Virgili,[28] V. Vislavicius,[82] A. Vodopyanov,[139] B. Volkel,[32] M. A. Völkl,[94] K. Voloshin,[138] S. A. Voloshin,[132] G. Volpe,[31] B. von Haller,[32] I. Vorobyev,[95] N. Vozniuk,[138] J. Vrláková,[36] B. Wagner,[20] C. Wang,[38] D. Wang,[38] M. Weber,[101]

A. Wegrzynek,[32] F. T. Weiglhofer,[37] S. C. Wenzel,[32] J. P. Wessels,[133] S. L. Weyhmiller,[135] J. Wiechula,[62] J. Wikne,[19] G. Wilk,[78] J. Wilkinson,[97] G. A. Willems,[133] B. Windelband,[94] M. Winn,[126] J. R. Wright,[106] W. Wu,[38] Y. Wu,[116] R. Xu,[6] A. K. Yadav,[130] S. Yalcin,[70] Y. Yamaguchi,[92] K. Yamakawa,[92] S. Yang,[20] S. Yano,[92] Z. Yin,[6] I.-K. Yoo,[16] J. H. Yoon,[56] S. Yuan,[20] A. Yuncu,[94] V. Zaccolo,[23] C. Zampolli,[32] H. J. C. Zanoli,[57] F. Zanone,[94] N. Zardoshti,[32,99] A. Zarochentsev,[138] P. Závada,[60] N. Zaviyalov,[138] M. Zhalov,[138] B. Zhang,[6] S. Zhang,[38] X. Zhang,[6] Y. Zhang,[116] M. Zhao,[10] V. Zherebchevskii,[138] Y. Zhi,[10] N. Zhigareva,[138] D. Zhou,[6] Y. Zhou,[82] J. Zhu,[6,97] Y. Zhu,[6] G. Zinovjev,[3,a] and N. Zurlo[53,129]

(ALICE Collaboration)

[1]*A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia*
[2]*AGH University of Science and Technology, Cracow, Poland*
[3]*Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine*
[4]*Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India*
[5]*California Polytechnic State University, San Luis Obispo, California, USA*
[6]*Central China Normal University, Wuhan, China*
[7]*Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba*
[8]*Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico*
[9]*Chicago State University, Chicago, Illinois, USA*
[10]*China Institute of Atomic Energy, Beijing, China*
[11]*Chungbuk National University, Cheongju, Republic of Korea*
[12]*Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic*
[13]*COMSATS University Islamabad, Islamabad, Pakistan*
[14]*Creighton University, Omaha, Nebraska, USA*
[15]*Department of Physics, Aligarh Muslim University, Aligarh, India*
[16]*Department of Physics, Pusan National University, Pusan, Republic of Korea*
[17]*Department of Physics, Sejong University, Seoul, Republic of Korea*
[18]*Department of Physics, University of California, Berkeley, California, USA*
[19]*Department of Physics, University of Oslo, Oslo, Norway*
[20]*Department of Physics and Technology, University of Bergen, Bergen, Norway*
[21]*Dipartimento di Fisica, Università di Pavia, Pavia, Italy*
[22]*Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy*
[23]*Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy*
[24]*Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy*
[25]*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy*
[26]*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy*
[27]*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy*
[28]*Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy*
[29]*Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy*
[30]*Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy*
[31]*Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy*
[32]*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
[33]*Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia*
[34]*Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway*
[35]*Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic*
[36]*Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic*
[37]*Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany*
[38]*Fudan University, Shanghai, China*
[39]*Gangneung-Wonju National University, Gangneung, Republic of Korea*
[40]*Gauhati University, Department of Physics, Guwahati, India*
[41]*Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany*
[42]*Helsinki Institute of Physics (HIP), Helsinki, Finland*
[43]*High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico*
[44]*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania*
[45]*Indian Institute of Technology Bombay (IIT), Mumbai, India*
[46]*Indian Institute of Technology Indore, Indore, India*
[47]*INFN, Laboratori Nazionali di Frascati, Frascati, Italy*

[48]INFN, Sezione di Bari, Bari, Italy
[49]INFN, Sezione di Bologna, Bologna, Italy
[50]INFN, Sezione di Cagliari, Cagliari, Italy
[51]INFN, Sezione di Catania, Catania, Italy
[52]INFN, Sezione di Padova, Padova, Italy
[53]INFN, Sezione di Pavia, Pavia, Italy
[54]INFN, Sezione di Torino, Turin, Italy
[55]INFN, Sezione di Trieste, Trieste, Italy
[56]Inha University, Incheon, Republic of Korea
[57]Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands
[58]Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic
[59]Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
[60]Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
[61]Institute of Space Science (ISS), Bucharest, Romania
[62]Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
[63]Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
[64]Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
[65]Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
[66]iThemba LABS, National Research Foundation, Somerset West, South Africa
[67]Jeonbuk National University, Jeonju, Republic of Korea
[68]Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
[69]Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
[70]KTO Karatay University, Konya, Turkey
[71]Laboratoire de Physique des 2 Infinis, Irène Joliot-Curie, Orsay, France
[72]Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
[73]Lawrence Berkeley National Laboratory, Berkeley, California, USA
[74]Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
[75]Nagasaki Institute of Applied Science, Nagasaki, Japan
[76]Nara Women's University (NWU), Nara, Japan
[77]National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
[78]National Centre for Nuclear Research, Warsaw, Poland
[79]National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
[80]National Nuclear Research Center, Baku, Azerbaijan
[81]National Research and Innovation Agency - BRIN, Jakarta, Indonesia
[82]Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
[83]Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
[84]Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
[85]Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic
[86]Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA
[87]Ohio State University, Columbus, Ohio, USA
[88]Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
[89]Physics Department, Panjab University, Chandigarh, India
[90]Physics Department, University of Jammu, Jammu, India
[91]Physics Department, University of Rajasthan, Jaipur, India
[92]Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (SKCM2), Hiroshima University, Hiroshima, Japan
[93]Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tubingen, Germany
[94]Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
[95]Physik Department, Technische Universität München, Munich, Germany
[96]Politecnico di Bari and Sezione INFN, Bari, Italy
[97]Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
[98]Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
[99]School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
[100]Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
[101]Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
[102]SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France
[103]Suranaree University of Technology, Nakhon Ratchasima, Thailand
[104]Technical University of Košice, Košice, Slovak Republic
[105]The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland

[106]The University of Texas at Austin, Austin, Texas, USA
[107]Universidad Autónoma de Sinaloa, Culiacan, Mexico
[108]Universidade de São Paulo (USP), Sao Paulo, Brazil
[109]Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
[110]Universidade Federal do ABC, Santo Andre, Brazil
[111]University of Cape Town, Cape Town, South Africa
[112]University of Houston, Houston, Texas, USA
[113]University of Jyväskylä, Jyväskylä, Finland
[114]University of Kansas, Lawrence, Kansas, USA
[115]University of Liverpool, Liverpool, United Kingdom
[116]University of Science and Technology of China, Hefei, China
[117]University of South-Eastern Norway, Kongsberg, Norway
[118]University of Tennessee, Knoxville, Tennessee, USA
[119]University of the Witwatersrand, Johannesburg, South Africa
[120]University of Tokyo, Tokyo, Japan
[121]University of Tsukuba, Tsukuba, Japan
[122]University Politehnica of Bucharest, Bucharest, Roma
[123]Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
[124]Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
[125]Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
[126]Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
[127]Università degli Studi di Foggia, Foggia, Italy
[128]Università del Piemonte Orientale, Vercelli, Italy
[129]Università di Brescia, Brescia, Italy
[130]Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
[131]Warsaw University of Technology, Warsaw, Poland
[132]Wayne State University, Detroit, Michigan, USA
[133]Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany
[134]Wigner Research Centre for Physics, Budapest, Hungary
[135]Yale University, New Haven, Connecticut, USA
[136]Yonsei University, Seoul, Republic of Korea
[137]Zentrum für Technologie und Transfer (ZTT), Worms, Germany
[138]Affiliated with an institute covered by a cooperation agreement with CERN
[139]Affiliated with an international laboratory covered by a cooperation agreement with CERN

[a]Deceased.
[b]Also at Max-Planck-Institut fur Physik, Munich, Germany.
[c]Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy.
[d]Also at Dipartimento DET del Politecnico di Torino, Turin, Italy.
[e]Also at Department of Applied Physics, Aligarh Muslim University, Aligarh, India.
[f]Also at Institute of Theoretical Physics, University of Wroclaw, Poland.
[g]Also at An institution covered by a cooperation agreement with CERN.