

Historical genomes elucidate European settlement and the African diaspora in Delaware

Highlights

- Genomic data indicate eight individuals of European and three of African ancestry
- Individuals were interred in separate burial groupings based on genomic ancestry
- Paternally related parent and child of African descent were identified
- Maternally related grandparent, parent, and child of European descent were found

Authors

Raquel E. Fleskes, Douglas W. Owsley,
Karin S. Bruwelheide,
Kathryn G. Barca, Daniel R. Griffith,
Graciela S. Cabana,
Theodore G. Schurr

Correspondence

raquel.fleskes@uconn.edu (R.E.F.),
owsleyd@si.edu (D.W.O.),
tgschurr@sas.upenn.edu (T.G.S.)

In brief

Fleskes et al. conduct ancient genomic analysis of a colonial period archaeological site in Delaware (USA). They investigate genetic ancestry of eleven burials and identify genetic kinship among individuals of both European and African descent. This study enlarges our understanding of frontier settlement and the trans-Atlantic slave trade.

Report

Historical genomes elucidate European settlement and the African diaspora in Delaware

Raquel E. Fleskes,^{1,2,7,9,*} Douglas W. Owsley,^{3,*} Karin S. Bruwelheide,³ Kathryn G. Barca,³ Daniel R. Griffith,⁴ Graciela S. Cabana,^{5,6} and Theodore G. Schurr^{2,8,*}

¹Department of Anthropology, University of Connecticut, Storrs, CT 06269, USA

²Department of Anthropology, University of Pennsylvania, Philadelphia, PA 19104, USA

³Department of Anthropology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560, USA

⁴Archaeological Society of Delaware, Inc, Dover, DE 19903, USA

⁵Department of Anthropology, University of Tennessee, Knoxville, TN 37996, USA

⁶Molecular Anthropology Laboratories, University of Tennessee, Knoxville, TN 37996, USA

⁷Twitter: [@dna_historical](#)

⁸Senior author

⁹Lead contact

*Correspondence: raquel.fleskes@uconn.edu (R.E.F.), owsleyd@si.edu (D.W.O.), tgschurr@sas.upenn.edu (T.G.S.)

<https://doi.org/10.1016/j.cub.2023.04.069>

SUMMARY

The 17th-century colonization of North America brought thousands of Europeans to Indigenous lands in the Delaware region, which comprises the eastern boundary of the Chesapeake Bay in what is now the Mid-Atlantic region of the United States.¹ The demographic features of these initial colonial migrations are not uniformly characterized, with Europeans and European-Americans migrating to the Delaware area from other countries and neighboring colonies as single persons or in family units of free persons, indentured servants, or tenant farmers.² European colonizers also instituted a system of racialized slavery through which they forcibly transported thousands of Africans to the Chesapeake region. Historical information about African-descended individuals in the Delaware region is limited, with a population estimate of less than 500 persons by 1700 CE.^{3,4} To shed light on the population histories of this period, we analyzed low-coverage genomes of 11 individuals from the Avery's Rest archaeological site (circa 1675–1725 CE), located in Delaware. Previous osteological and mitochondrial DNA (mtDNA) sequence analyses showed a southern group of eight individuals of European maternal descent, buried 15–20 feet from a northern group of three individuals of African maternal descent.⁵ Autosomal results further illuminate genomic similarities to Northwestern European reference populations or West and West-Central African reference populations, respectively. We also identify three generations of maternal kin of European ancestry and a paternal parent-offspring relationship between an adult and child of African ancestry. These findings expand our understanding of the origins and familial relationships in late 17th and early 18th century North America.

RESULTS

Genomic sequencing results

Human genomic and mitochondrial DNA (mtDNA) were extracted from the petrous pyramid of the temporal bone for each of the 11 Avery's Rest individuals. The DNA extracts underwent whole-genome enrichment, generating between 0.17 and 0.64 \times genomic coverage (Table 1; Data S1B). Estimates of present-day DNA sequence contamination were calculated, with X chromosome contamination⁶ for chromosomal males below 4.5% and present-day mtDNA contamination¹⁰ ranging between 0.73% and 15.38% (Table 1; Data S1G). Four individuals (AR03, AR07, AR09, and AR10) exhibited >5% mtDNA contamination. Consequently, we restricted sequence reads to include only those with damage using PMDtools (PMD > 3) for all subsequent population analyses (PCA, ADMIXTURE, F3 statistics, and kinship analyses) to determine if there were any

discrepancies in our findings (Figure S1; Data S2Q–S2V). Although the results were affected by the decrease in the number of useable sequence reads, they were not substantially different from initial estimates.

The overall percentage of damage at the first base position ranged between 5.16% and 21.33%, consistent with partial USER treatment during library preparation. Damage patterns were initially characterized using MapDamage,¹¹ which identified increased G-to-A substitution rates at the 3' end, but decreased C-to-T rates at the 5' end. This asymmetrical pattern of damage is likely due to the use of Phusion High-Fidelity DNA Polymerase (NEB) during library amplification, which cannot read through uracil bases but preserves signatures of damage to CpG methyl groups.^{11–13} PMDtools⁷ was run using the “platypus” option and detected elevated levels of damage to CpG methyl groups at both 5' and 3' ends, suggesting damage attributable to DNA degradation (Figure S1; Data S1G).

Table 1. Archaeological, osteological, and sequencing results summary for the Avery's Rest individuals

ID	Burial location	Age ^a (years)	Coverage	Mapped reads (Human Origins panel)		Contamination ^b (%)	Damage ^c (%)	Sex ^d	mtDNA coverage	mtDNA haplogroup	NRY haplogroup
AR01	south	30–40	0.56x	11,797	4.80%	7.22%	M	177.20x	T2a1a8	R1b1a-Y16765	
AR02	south	4–6 months	0.58x	15,882	0.73%	12.33%	M	176.83x	H1af1a	R1b1a-S378	
AR03	south	35–45	0.55x	12,533	10.91%	7.09%	M	189.75x	W	R1b1a-Z30233	
AR04	south	40–50	0.50x	11,997	3.64%	7.36%	M	174.25x	U5b2a1a1a	R1b1a-S1746	
AR05	south	35–45	0.45x	10,482	1.59%	6.85%	M	168.14x	H11a	I2a1b-FGC3617	
AR06	south	50–60	0.43x	11,225	1.65%	7.69%	F	157.10x	H1af1a	–	
AR07	south	35–45	0.17x	4,919	15.38%	8.14%	F	142.61x	H1af1a	–	
AR08	south	30–40	0.32x	8,351	4.58%	8.11%	M	167.85x	J1c2p	I1a2a-BY211	
AR09	north	32–42	0.25x	7,275	7.18%	10.30%	M	157.72x	L3e3b	E1b1a-CTS2447	
AR10	north	27–37	0.37x	10,198	7.18%	21.33%	M	170.05x	L0a1a2	E1b1a-Z5974	
AR11	north	4.5–5.5	0.64x	14,898	4.58%	5.16%	M	186.98x	L3d2	E1b1a-Z5974	

See also [Figures S1](#) and [S2](#), [Table S1](#), and [Data S1B–S1G](#).

^aEstimation based on osteological assessments in Fleskes et al.⁵

^bmtDNA contamination estimates using contamMix⁶

^cDamage estimation using PMDtools, showing percentage of damaged reads >3⁷

^dHere, we estimate sex using Skoglund et al.⁸ based on generated data related to chromosomal sex. Note that this program only estimates XX or XY states and does not account for other chromosomal sex karyotypes. While the authors contextualize sex using binary assignments of X and Y chromosomes, we recognize that sex is a far more complicated multidimensional variable.⁹

Principal component analysis

To ascertain the genomic ancestries of the Avery's Rest individuals, we called single nucleotide polymorphisms (SNPs) against transversion sites on the Affymetrix Human Origins panel ([Table S1](#)). SmartPCA¹⁴ results reinforced previous findings that showed a clear separation of the 11 individuals into two groups based on continental ancestry, with eight individuals in the southern burial group clustering in PCA space with European and Middle Eastern reference populations and the three individuals in the northern group projecting with African reference populations ([Figure 1A](#)). To verify this outcome, SmartPCA¹⁴ was also run only with damaged reads (PMD > 3),⁷ with the results showing a similar data projection ([Figure S2](#)).

For the eight individuals projecting near European populations (AR01–AR08), we investigated their genomic similarity with only European reference populations in the Human Origins panel ([Table S1](#)). The SmartPCA¹⁴ results showed that they projected close to reference populations in northwestern Europe ([Figure 1B](#)). To obtain a more detailed assessment of European ancestry, an additional SmartPCA¹⁴ was run using only European populations on the Human Origins panel. Individuals with European ancestry clustered closest to British and French populations ([Figure S3A](#)).

The genomic similarities of the three individuals with African ancestry were further evaluated using a custom panel of 78 populations and 3,558 individuals ([Table S1](#)). SmartPCA¹⁴ results indicated that these individuals cluster with different regional African reference populations ([Figure 2B](#)). Individual AR09 projected toward the middle of the PCA, with admixed African descendant populations from Barbados. Individual AR10 fell within the projected range of West-Central African Bantu-speaking populations, while AR11 clustered between West and East African populations.

To obtain better resolution of their African ancestries, we ran SmartPCA¹⁴ using data from West, West-Central, and Southern

African reference populations. Overall, the three individuals showed similarity to west and west-central biogeographic groups along the second principal component. AR09 projected close to West African populations, while AR10 clustered with West-Central African populations. In contrast, AR11 did not closely align with any specific population ([Figure S3B](#)). However, this approach assumes that present-day African reference populations are a comprehensive representation of the diversity that existed during the colonial period.¹⁵ This is problematic as populations are not static entities, population distributions in continental Africa were disrupted by the trans-Atlantic slave trade, and African reference populations are underrepresented in public databases.¹⁶

Inter- and intra-continental admixture

To investigate inter-continental population admixture for the individuals at Avery's Rest, their genomic data were analyzed using ADMIXTURE¹⁷ for K = 3 to K = 8 against transversion sites in the Human Origins panel. The results showed no major continental admixture for any of these individuals ([Figure 1C](#)). We investigated admixture patterns for the African-descended individuals by running ADMIXTURE¹⁷ with the custom African panel and the 1000 Genomes Project European-American (CEU), Han Chinese (CHS), Mexican (MXL), and Peruvian (PEL) populations, from K = 4 to K = 12 ([Figure 2C](#)). The resulting admixture profile for AR10 was similar to those for reference populations from West-Central Africa, namely Cameroon and Gabon. By contrast, AR09 and AR11 did not display admixture profiles similar to any contemporary African populations in the panel, although the predominant cluster in AR09 appeared to be highly represented in West African reference populations. The admixture profile of AR11 also contained a combination of all African clusters and did not align closely with any regional group. This pattern suggests that either this individual had




Figure 1. Genomic ancestries of Avery's Rest individuals against the global Human Origins panel

(A) PCA results showing separation of individuals with European and African ancestry components. (B) PCA results for the eight European-affiliated Avery's Rest individuals, plotted against European populations in the Human Origins panel. (C) ADMIXTURE results at $K = 8$ for the Avery's Rest series.

See also Figures S2 and S3 and Data S1C.

genetic contributions from several different African populations or the primary source population was not represented in the reference panel.

We then ran outgroup F3 statistics using popstats¹⁸ for all Avery's Rest individuals against populations in the Human Origins reference panel to ascertain patterns of shared genetic drift. We specified F3(AR, Target Population; Mbuti) against all populations in the Human Origins reference panel, whereby Mbuti represents an equal outgroup to West African and European populations.^{15,19} The results mirrored findings from the SmartPCA¹⁴ and ADMIXTURE¹⁷ analyses. There were higher F3 values with Northern or Western European populations for individuals AR01 through AR08 and higher F3 values for Bantu-speaking populations in Africa for individuals AR09, AR10, and AR11 (Data S2A–S2K; Figure S4). All values contained significant Z scores (>3).

To obtain a finer delineation of the admixture profiles of the three African-affiliated individuals, we ran outgroup F3 statistics using the arrangement F3(AR, Target Population; Mbuti).¹⁸ The results showed higher F3 values with significant Z scores (>3) across western, southern, and central Africa and did not reveal strong associations with any one population (Data S2L–S2N; Figure S4). In addition, we ran admixture F3 statistics with AR09 and AR11 to test whether they were admixed with any of the African reference populations, where F3(AR, Test Population; Target Population), but found no evidence for it (Data S2O and S2P).

Chromosomal sex estimation

Chromosomal sex was estimated for all individuals from filtered reads.⁸ The results showed that two females and nine males

were buried at Avery's Rest, verifying osteological assessments of sex for the adults⁵ and clarifying that the infant of European ancestry (AR02) and child of African ancestry (AR11) were chromosomal males. The two females of European ancestry were interred in the southern group (Table 1; Data S1F).

mtDNA and Y chromosome diversity

To investigate the maternal and paternal lineages of these individuals, we characterized their mitogenome and male-specific Y chromosome (MSY) sequences. The eight individuals in the southern burial group displayed mtDNA haplogroups

found broadly across western Europe (H, J, U, T, and W), including the United Kingdom (Table 1; Data S1E).^{20,21} Three of these individuals (two adult females [AR06 and AR07] and the infant male [AR02]) share haplogroup H1af1a, indicating a strong likelihood of maternal relatedness. The three individuals in the northern group had mtDNAs from distinctly African haplogroups, namely, L0a1a2, L3d2, and L3e3b. Of these, persons carrying L3e3 and L0a1a2 mtDNAs are present in Central Africa,^{22,23} and those with L3d mtDNAs are found at high frequencies among West African populations, although the sublineage L3d2 has also been observed in East African populations.^{24,25} L0a1 and L3e3 have also been found in other historical African-descended individuals from South Carolina (United States) and St. Helena (South Africa),^{26,27,28} and all three mtDNA haplogroups have been reported in contemporary African-descended populations in the Americas.^{29–32}

MSY haplogroups for chromosomal males (AR01, AR02, AR03, AR04, AR05, AR08, AR09, AR10, and AR11) were predicted from genomic sequence data using Yleaf³³ and showed similar patterns of ancestral distributions (Table 1; Data S1D). The southern burial group possessed a diverse set of European paternal lineages, with MSY sublineages of haplogroups I1, I2, and R1b1a being identified. I1a is commonly found in populations in northern Europe,³⁴ I2 in western and eastern Europe,³⁵ and R1b1a throughout western Europe.³⁶ By contrast, the three individuals in the northern burial group had Y chromosomes belonging to the African haplogroup E1b1a. This MSY haplogroup is observed in west, central, and southern African populations,³⁷ as well as in African-descended populations in the Americas.^{36,38,39} In addition, the adult male (AR10) and child




Figure 2. African ancestry profiles for AR09, AR10, and AR11 against a custom African reference panel

(A) Map of Africa with the countries used in the reference panel colored according to the accompanying PCA plot. The subpanel displays the Lesser Antilles Islands, for which the island of Barbados is highlighted.

(B) PCA plot of the three Avery's Rest individuals with African ancestry.

(C) ADMIXTURE profiles of the three individuals with African ancestry at $K = 9$.

See also [Figures S2 and S3](#), [Tables S1 and S2](#), and [Data S1](#).

(AR11) shared the same terminal marker in the E1b1a haplotype. While not conclusive evidence of a paternal genetic relationship, this finding does not exclude the possibility.

Genetic kinship

To explore the degree of genetic relatedness among Avery's Rest individuals, we assessed autosomal kinship using two primary approaches. We estimated first-degree relationships using average pair-wise distance with READ⁴⁰ and observed three first-degree relationship pairs (AR06 and AR07; AR07 and AR02; AR10 and AR11) (Figure 3; [Data S3A](#) and [S3B](#)). These pairs shared either the same mtDNA (H1a1af1 for AR02, AR06, and AR07) or MSY (E1b1a1-Z5974 for AR10 and AR11) haplogroup. For the individuals with African ancestry, the fact that they did not share the same mtDNA haplogroup ruled out a full-sibling relationship within the context of first-degree relatedness with identical MSY haplogroups. Thus, this evidence supports a paternal parent-offspring relationship between the African-descended adult male (AR10) and child (AR11).

We further characterized the first-degree relationship for only the individuals with European ancestry (AR02, AR06, and AR07) using NgsRelate.⁴¹ This program requires input allele frequencies generated from contemporary populations, which could not be accurately approximated for the African individuals.

For the European individuals, we used the British (GBR) and Northern European (CEU) populations from the 1000 Genomes Project to estimate allele frequencies for the intersecting SNPs.⁴² The R0, R1, and KING-coefficient statistics for the AR06-AR07 and AR02-AR07 pairs fell within the range of parent-offspring relationships, as reported by Waples and colleagues (Figure 3; [Data S3C](#)).⁴³ In addition, NgsRelate identified a second-degree relationship between AR06 and AR02. Since these individuals share the same mtDNA haplogroup, it is likely that AR06, AR07, and AR02 had an inter-generational maternal relationship, specifically that of grandmother, mother, and son.

DISCUSSION

Our findings at the Avery's Rest site illustrate how the entangled web of colonialism spans the larger Atlantic world and provide important insights into early colonial settlement, kinship, and the trans-Atlantic slave trade in Delaware at the end of the 17th century. This case study provides the first genomic evidence of the demographic makeup of European colonial migrants and their occupation in Delaware during the 17th century, which had previously only been estimated from archival records and archaeological evidence. Our results show the European-descended individuals were most genetically similar to populations

Current Biology Report




Figure 3. Kinship estimation for the Avery's Rest individuals

(A) Burial locations at the Avery's Rest site. Individuals AR01–AR08 are located in the southern burial cluster; individuals AR09–AR11 are located in the northern cluster. The space between the dots represents a 5-foot scale. Note that due to the scale of the map, burials AR01 and AR03 appear to overlap but are buried parallel to each other.

(B) READ results for the African individuals, with error bars showing one standard error for average pairwise distance.

(C and D) (C) NgsRelate results for the 2DSFS method, displaying R1 and the KING-Coefficient, and (D) R1 and the R0 values for the European-affiliated individuals.

(E) Constructed pedigree chart showing relationships among Avery's Rest individuals. Squares indicate males and circles represent females, as estimated in the [Chromosomal sex estimation](#) section of the [STAR Methods](#).

See also Figure S2 and Data S1D–S1F and S3A–S3D.

from Great Britain and France, which correlates with historical records showing British colonial settlement in Delaware following Dutch occupation of Indigenous lands.¹ Five non-genetically related males were buried along with three consecutive generations of matrilineal kin, representing a grandmother, mother, and son. This demographic profile supports historiographical findings of a blended or mosaic organizational structure for European colonists at early colonial Chesapeake and Mid-Atlantic settlements, in which unrelated males and small groups of biological kin formed social and labor units.²

Besides providing insight into the demography of colonial Delaware, the identification of individuals of European and African ancestry at Avery's Rest reflects mortuary practices in North American frontier settings. These individuals were interred in closely spaced, but separate, burial clusters divided by continental-level ancestry. Their interment location further suggests that interactions between European and African persons may have been more integrated than suggested by mortuary practices in later antebellum cemeteries, where burials were often interred in more separated locations based on race.^{44–46} The distinct clustering of the burial ground by continental ancestry nevertheless suggests that frontier lifeways

from the beginning were governed by settler-colonial practices of racialized enslavement.^{3,47}

The presence of African individuals at Avery's Rest is itself significant, as few enslaved Africans are known to have entered the Delaware region between 1665 and 1713 CE.^{1,4} However, two adult African individuals at Avery's Rest were recorded as part of an estate re-evaluation case petitioned on 7 October 1690 CE concerning debts to the estate of John Avery after his death in September 1682 CE.⁴⁸ The stated reason for re-evaluation was that the two African males had died shortly before the initial 1683 CE evaluation was conducted.⁴⁸ It is possible, although uncertain, that the two adult African individuals found at Avery's Rest are those referenced in the record. Previous osteological analysis provided information on the cause of death for AR10.⁵ Their cranium displays evidence of a facial fracture to the right zygomatic arch and maxilla caused by blunt force trauma at or near the time of death. The location of the fracture on the cranium is consistent with interpersonal violence.

The identification of a patrilineal parent-offspring relationship between adult AR10 and child AR11 represents the earliest demonstration of biological kinship among African-descended individuals using ancient DNA at a North American colonial site. The

side-by-side interment of AR10 and AR11 suggests that biological kinship may have dictated burial placement of the son (AR11), whose death likely followed the father's (AR10). Burial placement suggests that AR09, who was buried adjacent to the fence line similar to most of the other burials (Figure 3), was interred prior to AR10 and AR11. The father-son relationship is notable as families could be, and often were, separated at the will of their enslavers, reflecting the structural violence embedded in the institution of slavery.⁴⁹ This discovery is especially critical for understanding histories of African-descended individuals during the early colonial period, for which evidence of biological relatedness in the archival record is limited.

In addition, this study offers important insights into the demographic histories of African individuals forcibly enslaved and brought to peripheral locations in the colonial Atlantic world. For the father-son pair (AR10-AR11), our results indicate that the father (AR10) had ancestry tracing to west-central Africa, whereas his son (AR11) did not show any clear association with a specific African reference population. This finding may suggest that the mother of AR11, who was not found at the site, originated from a different population in Africa. The other unrelated adult African male (AR09) likely had ancestral origins in west Africa. These results align with other ancient genomic analyses of African-descended individuals in North America and the Caribbean^{50,51} and contemporary African Americans^{16,52,53} and broadly reflect the larger patterns of west and west-central African origins in the trans-Atlantic slave trade.⁵⁴ Furthermore, the genomic data suggest that the networks embedded in the trans-Atlantic slave trade were consistent between early core and peripheral locales in English colonial North America.

STAR METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
 - Lead contact
 - Materials availability
 - Data and code availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Archaeological site
 - Ethics statement
- **METHOD DETAILS**
 - Contamination control
 - DNA sample collection
 - DNA extraction and library preparation
 - Enrichment and sequencing
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - Sequence Data Processing and Authentication
 - Reference panels and Variant calling
 - Uniparental markers
 - PCA and admixture analyses
 - F3 statistics
 - Chromosomal sex estimation
 - Kinship estimation
 - Effects of contamination

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cub.2023.04.069>.

ACKNOWLEDGMENTS

The authors thank the Avery Family representatives, property owners Mr. and Mrs. Waymon Harmon, the Delaware Division of Historical and Cultural Affairs, and the Archaeological Society of Delaware, Inc., for their support of this project. We also thank the Smithsonian Institution Collections Advisory Committee for granting permission to undertake this follow-up study. With respect to genomic analyses, we thank Dr. Brenna Henn for access to the UC Davis HCLP Computing Server, Dr. Jonathan Schug at the Next Generation Sequencing Core at the University of Pennsylvania, Joe May from the Division of Biology Sequencing Facility, Veronica Brown at the Genomics Core (UTK-Knoxville), and Dr. C.E.G. Amorim (Cal State Northridge) and Dr. Iain Mathieson (University of Pennsylvania) for computational advice. We would also like to thank the reviewers, Dr. Barbara Health (UTK-Knoxville), and Dr. Morgan Hoke (University of North Carolina, Chapel Hill) for their comments on previous drafts of this paper. This project was funded by a National Science Foundation Doctoral Dissertation Research Improvement Grant (BCS-1825583, R.E.F. and T.G.S.), National Science Foundation SBE Postdoctoral Research Fellowship (SPRF-FR 2105384, R.E.F.), National Geographic Society grant (HJ-056ER-17, R.E.F.), Dissertation Completion Fellowship from the University of Pennsylvania (R.E.F.), and the Penn Museum of Anthropology and Archaeology (R.E.F.). The Rice Family Endowment for Forensic Anthropology provided critical support for Smithsonian Skeletal Biology Program contributions.

AUTHOR CONTRIBUTIONS

R.E.F., T.G.S., and D.W.O. designed research; R.E.F., D.W.O., K.S.B., K.G.B., and D.R.G. performed research; R.E.F. analyzed data; R.E.F., T.G.S., D.W.O., G.S.C., K.S.B., K.G.B., and D.R.G. wrote the paper; and T.G.S., D.W.O., and G.S.C. supervised research.

DECLARATION OF INTERESTS

The authors declare no competing interests.

INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

Received: September 16, 2022

Revised: February 1, 2023

Accepted: April 24, 2023

Published: May 18, 2023

REFERENCES

1. Munroe, J. (2006). *History of Delaware* (University of Delaware Press).
2. Tomlins, C. (2010). *Freedom Bound: Law, Labor, and Civic Identity in Colonizing English America, 1580–1865* (Cambridge University Press).
3. Essah, P. (1996). *A House Divided: Slavery and Emancipation in Delaware, 1638–1865* (University of Virginia Press).
4. Williams, W. (1996). *Slavery and Freedom in Delaware* (Scholarly Resources), pp. 1639–1865.
5. Fleskes, R.E., Bruwelheide, K.S., West, F.L., Owsley, D.W., Griffith, D.R., Barca, K.G., Cabana, G.S., and Schurr, T.G. (2019). Ancient DNA and bioarchaeological perspectives on European and African diversity and relationships on the colonial Delaware frontier. *Am. J. Phys. Anthropol.* 170, 232–245.
6. Moreno-Mayar, J.V., Korneliussen, T.S., Dalal, J., Renaud, G., Albrechtsen, A., Nielsen, R., and Malaspina, A.-S. (2020). A likelihood method for estimating present-day human contamination in ancient

Current Biology

Report



male samples using low-depth X-chromosome data. *Bioinformatics* 36, 828–841.

7. Skoglund, P., Northoff, B.H., Shunkov, M.V., Derevianko, A.P., Pääbo, S., Krause, J., and Jakobsson, M. (2014). Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. USA* 111, 2229–2234.
8. Skoglund, P., Storå, J., Götherström, A., Jakobsson, M., Götherström, A., and Jakobsson, M. (2013). Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* 40, 4477–4482.
9. Richardson, S.S. (2022). Sex contextualism. *Philosophy, Theory, and Practice in Biology* 14, 1–17.
10. Renaud, G., Slon, V., Duggan, A.T., and Kelso, J. (2015). Schmutzi: estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* 16, 224.
11. Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P.L.F., and Orlando, L. (2013). MapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinforma. Appl. Note* 29, 1682–1684.
12. Hangjø, K., and Orlando, L. (2019). Ancient epigenomics. In *Paleogenomics: Genome-Scale analysis of Ancient DNA*, C. Lindqvist, and O.P. Rajora, eds. (Springer International Publishing), pp. 75–111.
13. Cai, D., Zhu, S., Gong, M., Zhang, N., Wen, J., Liang, Q., Sun, W., Shao, X., Guo, Y., Cai, Y., et al. (2022). Radiocarbon and genomic evidence for the survival of *Equus Sussemionus* until the late Holocene. *eLife* 11, e73346. <https://doi.org/10.7554/eLife.73346>.
14. Patterson, N., Price, A.L., and Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genet.* 2, e190.
15. Barquera, R., Lamnidis, T.C., Lankapalli, A.K., Kocher, A., Hernández-Zaragoza, D.I., Nelson, E.A., Zamora-Herrera, A.C., Ramallo, P., Bernal-Felipe, N., Immel, A., et al. (2020). Origin and health status of first-generation Africans from early colonial Mexico. *Curr. Biol.* 30, 2078–2091.e11.
16. Jackson, F.L.C. (2021). So many Nigerians: why is Nigeria overrepresented as the ancestral genetic homeland of legacy African North Americans? *Am. J. Hum. Genet.* 108, 202–208.
17. Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* 19, 1655–1664.
18. Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., and Reich, D. (2012). Ancient admixture in human history. *Genetics* 192, 1065–1093.
19. Skoglund, P., Mallick, S., Bortolini, M.C., Chennagiri, N., Hünemeier, T., Petzl-Erler, M.L., Salzano, F.M., Patterson, N., and Reich, D. (2015). Genetic evidence for two founding populations of the Americas. *Nature* 525, 104–108.
20. Richards, M. (2003). The Neolithic invasion of Europe. *Annu. Rev. Anthropol.* 32, 135–162.
21. Simoni, L., Calafell, F., Pettener, D., Bertranpetti, J., and Barbujani, G. (2000). Geographic patterns of mtDNA diversity in Europe. *Am. J. Hum. Genet.* 66, 262–278.
22. Soares, P., Alshamali, F., Pereira, J.B., Fernandes, V., Silva, N.M., Afonso, C., Costa, M.D., Musilová, E., MacAulay, V., Richards, M.B., et al. (2012). The expansion of mtDNA haplogroup L3 within and out of Africa. *Mol. Biol. Evol.* 29, 915–927.
23. Silva, M., Alshamali, F., Silva, P., Carrilho, C., Mandlate, F., Jesus Trovoada, M., Černý, V., Pereira, L., and Soares, P. (2015). 60,000 years of interactions between Central and Eastern Africa documented by major African mitochondrial haplogroup L2. *Sci. Rep.* 5, 12526–13.
24. Kivisild, T., Reidla, M., Metspalu, E., Rosa, A., Brehm, A., Pennarun, E., Parik, J., Geberhiwot, T., Usanga, E., and Villem, R. (2004). Ethiopian mitochondrial DNA heritage: tracking gene flow across and around the gate of tears. *Am. J. Hum. Genet.* 75, 752–770.
25. Černý, V., Salas, A., Hájek, M., Zaloudková, M., and Brdička, R. (2007). A bidirectional corridor in the Sahel-Sudan belt and the distinctive features of the Chad Basin populations: a history revealed by the mitochondrial DNA genome. *Ann. Hum. Genet.* 71, 433–452.
26. Sandoval-Velasco, M., Jagadeesan, A., Ávila-Arcos, M.C., Gopalakrishnan, S., Ramos-Madrigal, J., Moreno-Mayar, J.V., Renaud, G., Cruz-Dávalos, D.I., Johannesson, E., Watson, J., et al. (2019). The genetic origins of Saint Helena's liberated Africans. Preprint at bioRxiv. <https://doi.org/10.1101/787515>.
27. Fleskes, R.E., Ofunniyi, A.A., Gilmore, J.K., Poplin, E., Abel, S.M., Bueschgen, W.D., Juarez, C., Butler, N., Mishoe, G., Oubré, L., et al. (2021). Ancestry, health, and lived experiences of enslaved Africans in 18th century Charleston: an osteobiographical analysis. *Am. J. Phys. Anthropol.* 175, 3–24.
28. Fleskes, R., Cabana, G.S., Gilmore, J.K., Juarez, C., Karcher, E., Oubré, L., Mishoe, G., Ofunniyi, A.A., and Schurr, T.G. (2023). Community-engaged ancient DNA project reveals diverse origins of 18th-century African descendants in Charleston, South Carolina. *Proc. Natl. Acad. Sci. USA* 120, 32201620120.
29. Ely, B., Wilson, J.L., Jackson, F., and Jackson, B.A. (2006). African-American mitochondrial DNAs often match mtDNAs found in multiple African ethnic groups. *BMC Biol.* 4, 1–14.
30. Salas, A., Richards, M., Lareu, M.V., Scozzari, R., Coppa, A., Torroni, A., Macaulay, V., and Carracedo, A. (2004). The African Diaspora: mitochondrial DNA and the Atlantic slave trade. *Am. J. Hum. Genet.* 74, 454–465.
31. Wilson, J.L., Saint-Louis, V., Auguste, J.O., and Jackson, B.A. (2012). Forensic analysis of mtDNA haplotypes from two rural communities in Haiti reflects their population history. *J. Forensic Sci.* 57, 1457–1466.
32. Carvalho, B.M., Bortolini, M.C., Santos, S.E., and Ribeiro-dos-Santos, Á.K.C. (2008). Mitochondrial DNA mapping of social-biological interactions in Brazilian Amazonian African-descendant populations. *Genet. Mol. Biol.* 31, 12–22.
33. Ralf, A., Montiel González, D., Zhong, K., and Kayser, M. (2018). Yleaf: software for human Y-chromosomal haplogroup inference from next-generation sequencing data. *Mol. Biol. Evol.* 35, 1291–1294.
34. Roots, S., Magri, C., Kivisild, T., Benazzi, G., Help, H., Bermisheva, M., Kutuev, I., Barać, L., Peričić, M., Balanovsky, O., et al. (2004). Phylogeography of Y-chromosome haplogroup I reveals distinct domains of prehistoric gene flow in Europe. *Am. J. Hum. Genet.* 75, 128–137.
35. Kivisild, T. (2017). The study of human Y chromosome variation through ancient DNA. *Hum. Genet.* 136, 529–546.
36. Resque, R., Gusmão, L., Geppert, M., Roewer, L., Palha, T., Alvarez, L., Ribeiro-dos-Santos, Á., and Santos, S. (2016). Male lineages in Brazil: Intercontinental admixture and stratification of the European background. *PLoS One* 11, e0152573.
37. de Filippo, C., Barbieri, C., Whitten, M., Mpoloka, S.W., Gunnarsdóttir, E.D., Bostoen, K., Nyambe, T., Beyer, K., Schreiber, H., de Knijff, P., et al. (2011). Y-chromosomal variation in Sub-Saharan Africa: insights into the history of Niger-Congo groups. *Mol. Biol. Evol.* 28, 1255–1269.
38. Lao, O., Vallone, P.M., Coble, M.D., Diegoli, T.M., Van Oyen, M., Van Der Gaag, K.J., Pijpe, J., De Knijff, P., and Kayser, M. (2010). Evaluating self-declared ancestry of U.S. Americans with autosomal, Y-chromosomal and mitochondrial DNA. *Hum. Mutat.* 31, E1875–E1893.
39. Benn Torres, J., Vilar, M.G., Torres, G.A., Gaiesti, J.B., Bharath Hernandez, R., Browne, Z.E., Stevenson, M., Walters, W., and Schurr, T.G. (2015). Genetic diversity in the Lesser Antilles and its implications for the settlement of the Caribbean basin. *PLoS One* 10, e0139192.
40. Monroy Kuhn, J.M., Jakobsson, M., and Günther, T. (2018). Estimating genetic kin relationships in prehistoric populations. *PLoS One* 13, e0195491.
41. Korneliussen, T.S., and Moltke, I. (2015). NgsRelate: a software tool for estimating pairwise relatedness from next-generation sequencing data. *Bioinformatics* 31, 4009–4011.
42. 1000 Genomes Project Consortium, Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., and Abecasis, G.R. (2015). A global reference for human genetic variation. *Nature* 526, 68–74.

43. Waples, R.K., Albrechtsen, A., and Moltke, I. (2019). Allele frequency-free inference of close familial relationships from genotypes or low-depth sequencing data. *Mol. Ecol.* **28**, 35–48.
44. Aufderheide, A.C., Angel, J.L., Kelley, J.O., Outlaw, A.C., Outlaw, M.A., Rapp, G., and Wittmers, L.E., Jr. (1985). Lead in bone III. Prediction of social correlates from skeletal lead content in four Colonial American populations (Catoctin Furnace, College Landing, Governor's Land, and Irene Mound). *Am. J. Phys. Anthropol.* **66**, 353–361.
45. Handler, J.S. (1996). A prone burial from a plantation slave cemetery in Barbados, West Indies: possible evidence for an African-type wotch or other negatively viewed person. *Hist. Archaeol.* **30**, 76–86.
46. Jamieson, R.W. (1995). Material culture and social death: African-American burial practices. *Hist. Archaeol.* **29**, 39–58.
47. Gall, M.J., Heinrich, A., Grossman-Bailey, I., Hayden, P.A., and McKnight, J. (2020). The place beyond the fence: slavery and cultural invention on a Delaware tenant farm. *Hist. Archaeol.* **54**, 305–333.
48. Horle, C. (1991). Records of the courts of Sussex County, Delaware, 1677–1710 (University of Pennsylvania Press).
49. Berry, D.R. (2017). The Price for Their Pound of Flesh: The Value of the Enslaved, from Womb to Grave, in the Building of a Nation (Beacon Press).
50. Schroeder, H., Ávila-Arcos, M.C., Malaspina, A.-S., Poznik, G.D., Sandoval-Velasco, M., Carpenter, M.L., Moreno-Mayar, J.V., Sikora, M., Johnson, P.L.F., Allentoft, M.E., et al. (2015). Genome-wide ancestry of 17th-century enslaved Africans from the Caribbean. *Proc. Natl. Acad. Sci. USA* **112**, 3660–3673.
51. Harney, É., Olalde, I., Bruwelheide, K., Barca, K.G., Curry, R., Comer, E., Rohland, N., Owsley, D., and Reich, D. (2022). Technical report on Ancient DNA analysis of 27 African Americans from Catoctin Furnace, Maryland. Preprint at bioRxiv. <https://doi.org/10.1101/2022.06.12.495320>.
52. Micheletti, S.J., Bryc, K., Ancona Esselmann, S.G., Freyman, W.A., Moreno, M.E., Poznik, G.D., Shastri, A.J., 23andMe Research Team, Beleza, S., Mountain, J.L., et al. (2020). Genetic consequences of the Trans-Atlantic slave trade in the Americas. *Am. J. Hum. Genet.* **107**, 265–277.
53. Jackson, F., and Borgelin, L.F.J. (2010). How genetics can provide detail into the transatlantic African diaspora. In *The African Diaspora and the Disciplines*, T. Olanian, and J.H. Sweet, eds. (Indiana University Press), pp. 53–74.
54. Eltis, D., and Richardson, D. (2013). *Routes to Slavery: Direction, Ethnicity, and Mortality in the Transatlantic Slave Trade* (Routledge).
55. Schubert, M., Lindgreen, S., and Orlando, L. (2016). AdapterRemoval v2: rapid adapter trimming, identification, and read merging. *BMC Res. Notes* **9**, 88.
56. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760.
57. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079.
58. Okonechnikov, K., Conesa, A., and García-Alcalde, F. (2016). Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294.
59. Fu, Q., Meyer, M., Gao, X., Stenzel, U., Burbano, H.A., Kelso, J., and Pääbo, S. (2013). DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl. Acad. Sci. USA* **110**, 2223–2227.
60. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., and Sham, P.C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575.
61. Weissensteiner, H., Pacher, D., Kloss-Brandstätter, A., Forer, L., Specht, G., Bandelt, H.J., Kronenberg, F., Salas, A., and Schönherr, S. (2016). HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res.* **44**, W58–63.
62. Francis, R.M. (2017). Pophelper: an R package and web app to analyse and visualize population structure. *Mol. Ecol. Resour.* **17**, 27–32.
63. Lipatov, M., Sanjeev, K., Patro, R., and Veeramah, K. (2015). Maximum likelihood estimation of biological relatedness from low coverage sequencing data. Preprint at bioRxiv. <https://doi.org/10.1101/023374>.
64. Sellers, E. (1898). Captain John Avery: President Judge at the Whorekill in Delaware Bay, and His Descendants (Press of J. B. Lippincott).
65. Gall, M., and Hayden, P. (2021). Avery's Rest: Archival Research and Button Analysis (Delaware Division of Historical and Cultural Affairs).
66. Sirak, K.A., Fernandes, D.M., Cheronet, O., Novak, M., Gamarra, B., Balassa, T., Bernert, Z., Cséki, A., Dani, J., Gallina, J.Z., et al. (2017). A minimally-invasive method for sampling human petrous bones from the cranial base for ancient DNA analysis. *Biotechniques* **62**, 283–289.
67. Pinhasi, R., Fernandes, D.M., Sirak, K., and Cheronet, O. (2019). Isolating the human cochlea to generate bone powder for ancient DNA analysis. *Nat. Protoc.* **14**, 1194–1205.
68. Dabney, J., Knapp, M., Glocke, I., Gansauge, M.T., Weihmann, A., Nickel, B., Valdiosera, C., García, N., Pääbo, S., Arsuaga, J.L., and Meyer, M. (2013). Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. USA* **110**, 15758–15763.
69. Winters, M., Monroe, C., Barta, J.L., and Kemp, B.M. (2019). Evaluating the efficiency of primer extension capture as a method to enrich DNA extractions. *J. Forensic Sci.* **64**, 1145–1151.
70. Lazaridis, I., Nadel, D., Rollefson, G., Merrett, D.C., Rohland, N., Mallick, S., Fernandes, D., Novak, M., Gamarra, B., Sirak, K., et al. (2016). Genomic insights into the origin of farming in the ancient Near East. *Nature* **536**, 419–424.
71. Chang, C.C. (2020). Data management and summary statistics with PLINK. In *Statistical Population Genomics*, J.Y. Dutheil, ed. (Springer), pp. 49–65.
72. Pagani, L., Kivisild, T., Tarekegn, A., Ekong, R., Plaster, C., Gallego Romero, I., Ayub, Q., Mehdi, S.Q., Thomas, M.G., Luiselli, D., et al. (2012). Ethiopian genetic diversity reveals linguistic stratification and complex influences on the Ethiopian gene pool. *Am. J. Hum. Genet.* **91**, 83–96.
73. Busby, G.B.J., Hellenthal, G., Montinaro, F., Tofanelli, S., Bulayeva, K., Rudan, I., Zemunik, T., Hayward, C., Toncheva, D., Karachanak-Yankova, S., et al. (2015). The role of recent admixture in forming the contemporary West Eurasian genomic landscape. *Curr. Biol.* **25**, 2878–2826.
74. Patin, E., Lopez, M., Grollemund, R., Verdu, P., Harmant, C., Quach, H., Laval, G., Perry, G.H., Barreiro, L.B., Froment, A., et al. (2017). Dispersals and genetic adaptation of Bantu-speaking populations in Africa and North America. *Science* **356**, 543–546.
75. Schlebusch, C.M., Skoglund, P., Sjödin, P., Gatupaille, L.M., Hernandez, D., Jay, F., Li, S., De Jongh, M., Singleton, A., Blum, M.G.B., et al. (2012). Genomic variation in seven Khoe-San groups reveals adaptation and complex African history. *Science* **338**, 374–379.
76. Patin, E., Siddle, K.J., Laval, G., Quach, H., Harmant, C., Becker, N., Froment, A., Régnault, B., Lemée, L., Gravel, S., et al. (2014). The impact of agricultural emergence on the genetic history of African rainforest hunter-gatherers and agriculturalists. *Nat. Commun.* **5**, 1–10.
77. Gurdasani, D., Carstensen, T., Tekola-Ayele, F., Pagani, L., Tachmazidou, I., Hatzikotoulas, K., Karthikeyan, S., Iles, L., Pollard, M.O., Choudhury, A., et al. (2015). The African Genome Variation Project shapes medical genetics in Africa. *Nature* **517**, 327–332.
78. Neparáczki, E., Kocsy, K., Tóth, G.E., Maróti, Z., Kalmár, T., Bihari, P., Nagy, I., Pálfí, G., Molnár, E., Raskó, I., and Török, T. (2017). Revising mtDNA haplotypes of the ancient Hungarian conquerors with next generation sequencing. *PLoS One* **12**, e0174886.
79. Allen Ancient DNA Resource, version 52.2. <https://reich.hms.harvard.edu/allen-ancient-dna-resource-aadr-downloadable-genotypes-present-day-and-ancient-dna-data>.
80. Fu, Q., Li, H., Moorjani, P., Jay, F., Slepchenko, S.M., Bondarev, A.a., Johnson, P.L.F., Aximu-Petri, A., Prüfer, K., de Filippo, C., et al. (2014).

Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* 514, 445–449. <https://doi.org/10.1038/nature13810>.

81. Vai, S., Amorim, C.E.G., Lari, M., and Caramelli, D. (2020). Kinship determination in archeological contexts through DNA analysis. *Front. Ecol. Evol.* 8, 1–9.

82. Korneliussen, T.S., Albrechtsen, A., and Nielsen, R. (2014). ANGSD: analysis of next generation sequencing data. *BMC Bioinf.* 15, 356.

83. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and VCFtools. *Bioinformatics* 27, 2156–2158.

84. Campbell, M.C., and Tishkoff, S.A. (2008). African genetic diversity: implications for human demographic history, modern human origins, and complex disease mapping. *Annu. Rev. Genomics Hum. Genet.* 9, 403–433.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Biological samples		
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR01 / 7S-G-57-AVERYSREST-01
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR02 / 7S-G-57-AVERYSREST-02
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR03 / 7S-G-57-AVERYSREST-03
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR04 / 7S-G-57-AVERYSREST-04
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR05 / 7S-G-57-AVERYSREST-05
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR06 / 7S-G-57-AVERYSREST-06
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR07 / 7S-G-57-AVERYSREST-07
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR08 / 7S-G-57-AVERYSREST-08
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR09 / 7S-G-57-AVERYSREST-09
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR10 / 7S-G-57-AVERYSREST-10
Archaeological Individual	This study / The Avery's Rest Archaeological Site / The Smithsonian Institution's National Museum of Natural History	AR11 / 7S-G-57-AVERYSREST-11
Chemicals, peptides, and recombinant proteins		
DNA AWAY	Molecular BioProducts	Cat# 7010
Bleach	Clorox	N/A
Agarose	Fisher Scientific	Cat# BP1356-500
Gel Red	Phenix	Cat# RGB-4103
Tris base	Fisher Scientific	Cat# BP152-1

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Boric Acid	Sigma Aldrich	Cat# B7660-500G
0.5 M EDTA	Teknova	Cat# E0306
Tween-20	Sigma Aldrich	Cat# P9416-50ML
Proteinase K	Sigma Aldrich	Cat# P2308-25MG
Guanidine Hydrochloride	Teknova	Cat# G0316
Isopropanol	Walgreens	Cat# 715455
Sodium Acetate	Sigma Aldrich	Cat# 71196-100ML
1M Tris-HCl	Invitrogen	Cat# 15568025
NEBNext Ultra II Library Preparation Kit	New England Biosciences	Cat# E7645S
USER enzyme	New England Biosciences	Cat# M5508
Phusion High-Fidelity DNA Polymerase	New England Biosciences	Cat# M0530S
Kapa HiFi Hot Start ReadyMix	Roche Diagnostics	Cat# 07958927001
10 mM Tris-Cl, 0.05% TWEEN-20 solution	Teknova	Cat# T1485
NEBNext Multiplex Oligos for Illumina	New England Biosciences	Cat# E7600S
dNTP	New England Biosciences	Cat# N0447S
DMSO	New England Biosciences	Cat# M0530S
BSA	New England Biosciences	Cat# B9000S

Critical commercial assays

PowerUp SYBR Green Master Mix	Thermo Scientific	Cat#: A25742
MinElute PCR Purification Kit	QiaGEN	Cat# 28006
myBaits Expert Whole Genome Enrichment Kit v4	Arbor Biosciences	Cat# 302508
Agilent High Sensitivity DNA Kit	Agilent Technologies	Cat# 5067-4626
MiSeq Reagent Kit v2 (300-cycles)	Illumina	Cat# MS-102-2002
NextSeq 500/550 High Output Kit v2.5 (150 Cycles)	Illumina	Cat# 20024907
NovaSeq 6000 S2 Reagent Kit v1.5 (100 cycles)	Illumina	Cat# 20028316

Deposited data

Raw data	This study	SRA: PRJNA812783
----------	------------	------------------

Software and algorithms

AdapterRemoval v2	Schubert et al. ⁵⁵	https://github.com/MikkelSchubert/adapterremoval
BWA	Li et al. ⁵⁶	https://bio-bwa.sourceforge.net/
samtools	Li et al. ⁵⁷	https://samtools.sourceforge.net/
Qualimap 2	Okonechnikov et al. ⁵⁸	http://qualimap.conesalab.org/
MapDamage 2.0	Jónsson et al. ¹¹	https://ginolhac.github.io/mapDamage/
contamMix	Fu et al. ⁵⁹	N/A
ANGSD	Moreno-Mayar et al. ⁶	https://github.com/ANGSD/angsd
PMDtools	Skoglund et al. ⁷	https://github.com/pontussk/PMDtools
PLINK	Purcell et al. ⁶⁰	https://zzz.bwh.harvard.edu/plink/
READ	Kuhn et al. ⁴⁰	https://bitbucket.org/tguenther/read/src
NGSRelate	Korneliussen et al. ⁴¹	https://github.com/ANGSD/NgsRelate
pileupCaller	N/A	https://github.com/stschiff/sequenceTools
Haplogrep 2	Weissensteiner et al. ⁶¹	https://haplogrep.i-med.ac.at/
yleaf	Ralf et al. ³³	https://github.com/genid/Yleaf
SmartPCA	Patterson et al. ¹⁴	https://www.hspf.harvard.edu/alkes-price/software/
ADMIXTURE	Alexander et al. ¹⁷	https://dalexander.github.io/admixture/
popHelper	Francis et al. ⁶²	http://www.royfrancis.com/pophelper/articles/

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
popstats	Patterson et al. ¹⁸	https://github.com/pontussk/popstats
IcMLkin	Lipatov et al. ⁶³	https://github.com/COMBINE-lab/maximum-likelihood-relatedness-estimation
Other		
Zymo Spin V	Zymo Research	Cat# C1012-50

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Raquel Fleskes (raquel.fleskes@uconn.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Genomic sequence data have been deposited at the Sequence Read Archives and are publicly available as of the date of publication. Accession numbers are listed in the [key resources table](#). This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Archaeological site

The Avery's Rest archaeological site encapsulates an early nexus of European settler colonial migration and the African diaspora. The site is located near present-day Lewes, Delaware (United States). Archival research indicates that the site was owned by the Avery family between circa A.D. 1675 to 1725, whose members migrated from the neighboring British colony of Maryland.⁶⁴ Archaeological excavations began at the site in 2006, and eventually revealed 11 burials comprised of a southern grouping of eight individuals situated 15–20 feet from a northern cluster of three individuals.⁵ All individuals buried at Avery's Rest are oriented east/west, with the head facing east, and aligned with a fence line running north/south. Previously published osteological estimations of age and biological sex indicate the presence of an infant, child, women and men at the site between the ages of 6 months to over 40 years of age.⁵ All burials are thought to date to the Avery family occupation period or slightly later.⁶⁵

Ethics statement

Consultation with representatives of Avery family descendants, the Harmon Family, who are the current property owners of the Avery's Rest site, and the Delaware Division of Historical and Cultural Affairs were conducted. Permission was obtained from the Smithsonian Institution National Museum of Natural History Department of Anthropology Collections Advisory Committee for DNA sampling in 2019. Outreach programming and public presentations were conducted through the Delaware Division of Historical and Cultural Affairs, as well as The Delaware Archaeological Society, Inc.

METHOD DETAILS

Contamination control

All laboratory analyses took place in the ancient DNA (aDNA) clean room of the Molecular Anthropology Laboratories at the University of Tennessee-Knoxville (UTK). The aDNA facility is constructed to minimize exogenous contamination through a four-chambered design, overhead UV lights, and positive-pressured HEPA-filtered airflow. Full PPE was used during all procedures conducted in the clean room laboratory. All surfaces, equipment, and gloves were wiped down using a 10% household bleach solution and/or DNA AWAY (MBP) followed by 70% ethanol before and after each use. All disposable lab supplies and small equipment were additionally subjected to UV radiation in a UV crosslinker for 30 min before use. During sample processing, gloves were changed, and all surfaces were fully cleaned using bleach and ethanol solutions between each sample to prevent cross-contamination. Negative controls were used during extraction and library preparation. The negative extraction control consisted of a blank sample tube filled with lysis buffer and no bone powder, which was carried through the extraction and library preparation process with the 11 Avery's Rest bone samples. An additional negative control (ultra-pure water) was used during the library preparation protocol. The negative controls were subject to identical protocols and quantification measures detailed in the next two sections. All were negative.

DNA sample collection

Sampling for individuals AR01 and AR03-AR10 took place at the Smithsonian Institution's National Museum of Natural History, using the minimally invasive petrosal drilling method, given that the craniums were intact.⁶⁶ A chemical fume hood was retrofitted for DNA sampling by fully decontaminating all surfaces before and after each sample using a 10% solution of household bleach followed by 70% ethanol. All sampling materials were similarly decontaminated and placed in a UV crosslinker for 10 min on each side before use.⁶⁷ Full personal protective equipment (PPE) was worn during sampling and gloves were changed between each sample collection. Bone powder was generated using a Dremel 8200 drill with a diamond-tipped bit.

Individuals AR02 and AR011 were not represented by fully intact crania. For this reason, the temporal bones from these individuals were transported to the UTK ancient DNA laboratory to allow for greater control over the sampling environment to reduce the chance of outside contamination. Petrosal samples were drilled inside a dead air box (Bone Clone Zone) that was fully cleaned with DNA AWAY (MBP) followed by 70% ethanol between each sample. The petrous pyramid was drilled from the interior aspect of the cranium to allow for easier access.

DNA extraction and library preparation

Between 0.18-0.24 g of petrous bone powder was collected in sterile 2 mL O-ring tubes for DNA extraction as per Dabney et al. (Data S1A).⁶⁸ Sample incubation occurred in an EDTA-based buffer for 24 h at 56°C with gentle shaking. DNA was then isolated using MinElute spin columns (Qiagen). Extracts and negative controls were verified through mtDNA PCR amplification and agarose gel electrophoresis.⁵

Dual-indexed sequencing libraries were prepared using a double-stranded library preparation protocol (NEBNext Ultra II) modified for degraded, low-template DNA samples. Libraries were prepared from 25 µL of DNA extracts, and subject to 3 µL of USER enzyme to partially reduce damage while still maintaining damage signatures for authentication purposes. Following ligation, libraries were dual-indexed using NEBNext Multiplex Oligos for Illumina (NEB). A StepOne Real-Time quantitative PCR (qPCR) system (Applied Biosystems) using a SYBR-Green master mix (BioRad) with library-specific primers and standards were used to verify adaptor ligation success and determine the relative target cycle number. Following qPCR results, libraries were amplified using Phusion High-Fidelity DNA Polymerase (NEB) for 15 or 16 cycles, depending on qPCR results (Data S1A). Amplified libraries were cleaned using a MinElute spin column protocol (Qiagen), following spin speeds and duration as outlined in Winters.⁶⁹

Quantification of the amplified indexed libraries, including positive and negative controls, was achieved by via a three-step process. First, the dsDNA concentrations were estimated using a Nanodrop Fluorospectrometer 3300 (Thermo Scientific). Second, the presence of adapter dimers was checked through the electrophoresis of 3 µL of extract on a 3% agarose gel. Third, a High Sensitivity Bioanalyzer assay (Agilent Technologies) was run at the UTK Division of Biology Sequencing Facility.

Enrichment and sequencing

Libraries were enriched using the myBaits Whole Genome Enrichment Kit v4 (Arbor Biosciences). Samples were independently incubated at 60°C for 48 h, with all relevant steps executed at 60°C. On-bead libraries were amplified using the HiFi Hot Start ReadyMix (Kapa Biosystems) for 14 cycles. Enriched libraries were cleaned using a modified spin-column clean up and analyzed using a High Sensitivity Bioanalyzer assay. DNA samples were pooled and sequenced on Illumina MiSeq v2 (150PE), followed by deeper sequencing on an Illumina NextSeq 500 (75SR) and NovaSeq 6000 S2 (100SR).

QUANTIFICATION AND STATISTICAL ANALYSIS

Sequence Data Processing and Authentication

AdapterRemoval v2 was used to trim adapters and filter reads based on quality scores (>30) and sequence length (>30).⁵⁵ FASTQ files were merged by sample ID before mapping to the human reference genome (hg19) with Burrows-Wheeler Aligner (BWA) aln using “I 1000 -n 0.01 -o 2” parameters, followed by BWA samse.⁵⁶ Reads with <30 mapping quality scores, duplicates, and reads with multiple mappings were removed using samtools.⁵⁷ Coverage and average read length were calculated using Qualimap 2 (Data S1B).⁵⁸

The authentication of aDNA sequences was first evaluated by assessing their damage patterns using MapDamage 2.0.¹¹ Noting the atypical damage profile likely resulting from library preparation methods, we additionally assessed damage at CpG methylation sites using the “platypus” option in PMDtools.⁷ Overall damage rates were calculated for the first 100,000 sequencing reads for each sample using the “first” and “CpG” options in PMDtools. Contamination estimates from present-day DNA sources were calculated using contamMix⁵⁹ for mitogenome data and ANGSD⁶ for X-chromosome data (Data S1G).

Reference panels and Variant calling

Variants were called against the complete Human Origins panel, a European Human Origins panel, and a custom panel of African populations (Data S1C). The Human Origins reference panel was used to determine the extent of European and African continental ancestry and admixture.⁷⁰ Variants were filtered following the protocol from Chang (2020) using PLINK.^{71,60} Specifically, variants were filtered for SNPs (–snps-only), missingness data (geno 0.1, mind 0.1), close relatives (king-cutoff 0.177), minor allele frequency (maf 0.05), and markers that deviate from Hardy-Weinberg equilibrium (–hwe 1e-25, keep-fewhet). All rsIDs were changed to chromosome and position number for consistency between datasets. This resulted in 425,308 variants for 2,056 individuals in the Human

Origins panel (Table 1; Data S1C). To assess European continental ancestry, we retained only the European populations in the Human Origins panel and filtered as described above. This resulted in the same number of variants and 450 individuals.

To obtain a more detailed understanding of African ancestry, a representative panel of 3,553 African individuals from 22 countries were assembled from the literature (Table S1).^{42,72–77} Before merging them, individual datasets were filtered as described above and a SNP list of updated rsIDs was generated in PLINK. A list of SNPs common between all datasets was generated in R using the respective lists. Each panel was then filtered for only these common SNPs and merged in PLINK. For quality purposes, the merged dataset was filtered again, resulting in 239,562 variants (Data S1C).

For admixture analyses of the three individuals with estimated African ancestry, we added the European-American (CEU), Han Chinese (CHB), Mexican (MXL), and Peruvian (PEL) populations from Phase 3 of the 1000 Genomes Project to the custom African panel to account for inter-continental admixture.⁴² The data from the selected 1000 Genomes populations were filtered as described above, and only the SNPs found on the list of common variants in the custom African panel were kept. This filtered dataset was then merged with the custom African panel and filtered again before variant calling and admixture analysis.

1000 Genomes Project data were also used to call genotypes for kinship analyses, given the significantly larger number of variants.⁴¹ After filtering, 2,493 individuals and 6,835,358 variants in the 1000 Genomes Panel were used (Data S1C).

Pseudo-haploid genotypes for the Avery's Rest individuals were independently called for each panel using samtools mpileup and pileupCaller (<https://github.com/stschiff/sequenceTools>), removing transitions and keeping SNPs with >30 mapping and base quality.⁵⁷ Subsequently, each panel was filtered for overlapping SNPs with the Avery's Rest genotypes and merged to the study individuals for downstream analyses.

Uniparental markers

The mitogenome sequences were independently mapped against the NC_012920.1 reference sequence using the mapping parameters outlined above in Sequence Data Processing and Authentication. Coverage was calculated using Qualimap 2.⁵⁸ Haploid variants were compiled using bcftools mpileup, filtering for map and base quality scores (>30), and called using bcftools call.⁵⁷ Variants were filtered for >5 depth. Haplogroups were called using the resulting VCF files with Haplogrep 2 (Data S1E).⁶¹

The mitogenome results generated from next-generation sequencing (NGS) mostly confirmed the previous mtDNA haplogroup calls generated from control region sequences in our previous study, while also increasing their phylogenetic resolution.⁵ The mtDNA haplogroups calls from two individuals (AR08 and AR11) differed from the previous Sanger sequencing results (AR08: H1 [Sanger] to J1 [NGS]; AR11: L3i [Sanger] to L3e [NGS]) generated from metacarpal or rib bone DNA extracts, respectively. This discrepancy could have resulted for different reasons, including higher error rates in Sanger sequencing for degraded DNA samples, or increased sequencing resolution with NGS technology.⁷⁸

For individuals with Y-chromosomes, paternal haplogroups were determined using yleaf, filtering for >30 quality scores, read depth >5, and calling only known ISOGG variants (Data S1D).³³

PCA and admixture analyses

Using autosomal data, principal components were calculated using SmartPCA, specifying no outliers and using lsqproject: YES.¹⁴ For admixture analyses, we first transformed the filtered comparative reference panels to make all sites homozygous to match the ancient data using a custom script (<https://github.com/cegamorim/>). We then additionally filtered for LD using indep-pairwise parameters “50 10 0.1” in PLINK.⁶⁰ We ran the program ADMIXTURE with 100 bootstrap replications for the designated values of K.¹⁷ ADMIXTURE results were visualized using popHelper in R.⁶²

F3 statistics

F3 statistics were run with popstats using the heterozygosity correction (Figure S4; Data S2A–S2V).^{18,19} Outgroup F3 statistics were run for all individuals against the Human Origins Reference panel using (AR, Target Population; Mbuti) to ascertain fine scale regional ancestry (Figure S4A). We also formally tested for shared genetic drift for the three African-descended individuals (AR09–AR11) with the Target Population representing the populations on the African panel (Figure S4B). The Mbuti samples were sourced from the publicly available 1240K dataset on the Reich lab website.^{79,80} To formally test for inter-African admixture for individuals AR09–AR11, F3 statistics were run specifying (AR, Test Population; Target Population) for all combinations of populations represented on the African reference panel/

Chromosomal sex estimation

The chromosomal sex of each individual was estimated by calculating the ratio of Y chromosome mapped reads compared to those mapped to both the X- and Y-chromosome reads (Data S1D).⁸

Kinship estimation

We used two main approaches to assess autosomal relatedness amongst the Avery's Rest individuals. First, READ was used to obtain a basic assessment of relatedness (Data S3A and S3B). READ was developed to estimate relatedness for low coverage sequences by calculating pairwise differences from pseudo-haploid genotype calls, which is ideal for aDNA studies.⁴⁰ However, READ assumes that all individuals derive from the same population and can be biased by high numbers of related individuals. To account for this issue, we independently ran READ for the European and African individuals genotyped against the 1000 Genomes reference

panel. We ran READ using the 'max' option for trios for the African-affiliated individuals, given the sample size of three. Due to the lower coverage of the samples, we used READ to characterize first-degree relationships with confidence. Individuals with second-degree relationships could not be verified using additional methods described below, and thus were not counted.

Second, we analyzed the data with NgsRelate, which produces R0, R1, and kinship coefficient values that can distinguish between parent-offspring and sibling first-degree relationships (Data S3D).^{41,43} NgsRelate uses allele frequencies from similar contemporary populations to estimate relatedness coefficients. This is also a drawback of this method, however, since an appropriate comparative population may not be available for archaeological individuals.⁸¹ Given the diverse ancestral composition of the three African individuals at Avery's Rest, a representative population could not be approximated.

For the individuals with European ancestry at Avery's Rest, we selected the European-American (CEU) and British (GBR) 1000 Genomes populations to generate representative allele frequencies.⁴² The data from these populations had been previously filtered as previously described, and were used to create a sites file to calculate genotype likelihoods in ANGSD, specifying '-gl 2 -domajorminor 3 -minMapQ 30 -domaf 1 -minmaf 0.05 -doGlf 3'.⁸² The resulting genotype likelihood positions file was used to select the positions with which to estimate allele frequencies in the CEU and GBR populations for NgsRelate.

To add additional support for the READ and NgsRelate analyses, we ran IcMLkin, which does not require estimated allele frequencies from contemporary populations.⁶³ Instead, IcMLkin estimates genotype likelihoods in a maximum likelihood framework to estimate kinship coefficients (K0, K1, K1, Pairwise) in a given population. Genotype likelihoods were estimated using the SNPbam2vcf python script against the 1000 Genomes panel. To estimate how the number of sites analyzed impacted the results of IcMLkin, we thinned every (A) 25K, (B) 50K, and (C) 100K sites and filtered ($-\text{maf} > 0.5$) using vcftools, where K equals 1,000 sites.⁸³ This resulted in (A) 104,724, (B) 53,064, and (C) 26,732, out of a possible 6,835,358 sites, respectively (Data S3E and S3F).

Given the small number of individuals at Avery's Rest, all individuals were analyzed together in IcMLkin two separate times. The first run identified related individual pairs ($\text{PI_HAT} > 0.05$), which were then used to identify the founders for the second run using the parameter -u. At all different SNP levels, relatedness between AR09 and AR10, between AR07 and AR02, and between AR07 and AR06 was estimated. The kinship coefficient was consistently higher for AR09 and AR10 than between the European pairs (Data S3G). These estimates are likely affected by the increased genomic diversity in the African-affiliated individuals relative to the European individuals in the analysis. However, separating the individuals by ancestry resulted in too small of population numbers for valid analysis. Therefore, the interpretation of the IcMLkin results were taken with caution and used only to verify the existence of biological relationships identified in READ and NgsRelate.

Our consideration of the choice of kinship estimation relied on the nature of the data analyzed. Avery's Rest represents a small 'population' comprised of two genetically diverse groups. The small sample size and proportionally higher levels of relatedness with the groups can introduce biases into analyses such as READ and IcMLkin if they are not corrected for. The importance of representative allele frequencies for NgsRelate has been noted by others,⁸¹ and is reiterated for the African-affiliated individuals analyzed in this paper. The African-affiliated individuals at Avery's Rest likely represent a diverse group of persons with West and West-Central African ancestry. African populations are known to contain the highest global levels of genetic diversity, even within regional areas.⁸⁴ This fact, combined with low sample numbers, makes determining a representative African population to estimate allele frequencies very difficult. Therefore, we support the use of multiple approaches to estimate kinship for aDNA samples to avoid the potential pitfalls noted above for a more robust estimation of biological relatedness.

Effects of contamination

For individuals with greater than 5% mtDNA contamination (AR03, AR07, AR09, and AR10), we restricted for reads with damage (>3) using PMDtools.⁷ This reduced the number of reads for all individuals (1,505,427 for AR03; 686,121 for AR07; 1,187,443 for AR09; and 1,411,230 for AR10). The damage restricted data was then analyzed to compare against downstream analyses (PCA, ADMIXTURE, outgroup F3-statistics, and kinship analyses) to deduce if the impact of contamination impacted results. We first ran SmartPCA¹⁴ following previously outlined protocols against the Human Origins Panel and Custom African Panel, respectively, which increased noise in the PCA projection but did not produce significant discrepancies in regional ancestry assignment (Figures S2A and S2B). We then conducted ADMIXTURE, as described previously, against the Human Origins Panel ($K = 3$ to $K = 8$) and the custom African Panel ($K = 3$ to $K = 14$). We also observed slight noise in the admixture profiles but did not produce significant discrepancies (Figures S2C–S2F). We then conducted outgroup F3-statistics¹⁸ using Mbuti as the designated outgroup. The resulting F-statistic and z-scores against each reference population were slightly different (Data S2Q–S2V), which is attributed to the reduced number of SNPs used in the comparison, but did not change interpretations of the findings. Taken together, the regional characterization of either African or European genomic similarity for the Avery's Rest individuals did not change.

In relation to the possible effect of contamination on kinship designations, only one individual within each of the identified kin groups displayed higher rates of mtDNA contamination (AR07 for the European-descended kin group; AR10 for the African-descended kin group). These individuals show the same previous basal mtDNA haplogroup calls as our previous paper, which used different skeletal elements (metatarsals, carpal, etc.) for control region sequencing of the mtDNA.⁵ The results are commensurate with identical sequences using different sequencing resolutions.

Damage restricted data slightly changed kinship designations by altering the degree of relatedness between individuals (Figures S2G–S2H). We conducted READ⁴⁰ and NgsRelate⁴¹ using the damage restricted sites, following previously outlined protocols. First, NgsRelate⁴¹ was conducted for the European individuals, which still identified all three relationship pairs, but reduced the R1 and KING-relatedness coefficients (Data S3D). This had an effect of reducing the degree of relatedness between AR02–AR07 from

a first degree to a second degree relationship. Alternatively, READ⁴⁰ analysis increased pairwise relatedness estimates, but maintained the same relationship pairing. For the African-descended individuals, READ⁴⁰ analysis resulted in identifying a first degree relationship between AR09 and AR10. We attribute this to the fact that pairwise estimates can be biased by the number of comparative sites, of which both AR09 and AR10 had a significantly reduced number. Therefore, we argue that using damage restricted sites for the Avery's Rest samples result in too little coverage for reliable relatedness estimations, but still demonstrated previously identified genetic kin relationships.

To verify the Y chromosome haplogroups designations, which were important in the characterization of the parent-offspring relationship between AR10 and AR11, we also reanalyzed the Y chromosome haplogroups using Yleaf³³ (Data S1D). The effect of the damage restriction significantly reduced the number of available sites, thereby reducing the resolution in Y chromosome haplogroup calls. This has the effect of eliminating some haplogroup calls from being made entirely, as well as removing the overlapping terminal marker that was shared between AR10 and AR11. However, the same broad haplogroups designations remained. Because of this, we believe that the higher resolution data is needed for this analysis.