Journal of Caribbean Ornithology

RESEARCH ARTICLE

Vol. 36:147-161. 2023

Avian use of avocado farms and intact forest in the northern Sierra de Bahoruco, Dominican Republic

Avery Young David Aborn Steven C. Latta Jennifer Nagel Boyd



Photo: Avery Young



Journal of Caribbean Ornithology

Research Article

Vol. 36:147-161. 2023

jco.birdscaribbean.org



Avian use of avocado farms and intact forest in the northern Sierra de Bahoruco, Dominican Republic

Avery Young¹,*², David Aborn^{1,3}, Steven C. Latta⁴, Jennifer Nagel Boyd^{1,5}

Associate Editor: Paige Byerly

Cover Page: Young and mature rows of avocado trees near the town of Puerto Escondido, Independéncia, Dominican Republic. This farm, and others in the area, sit in the foothills of the Northern Sierra de Bahoruco that can be seen in the distance. Many endemic and Neotropical migrant bird species inhabit these mountains including the tropical dry forest communities in the foothills where avocado farming is an increasing concern. Photograph by Avery Young in October 2021.

Published: 23 October 2023

¹Department of Biology, Geology, and Environmental Science, University of Tennessee at Chattanooga, Chattanooga, TN 37343, USA

*2e-mail: averybeayoung@icloud.com

³e-mail: david-aborn@utc.edu

⁴Department of Conservation and Field Research, National Aviary, Pittsburgh, PA 15212, USA; e-mail: steve.latta@aviary.org

⁵e-mail: jennifer-boyd@utc.edu

Abstract

The Dominican Republic is experiencing increased forest loss due to the expansion of avocado farming. While agricultural lands can provide viable habitat for birds, knowledge of avian use of avocado farms relative to native forest is limited. We conducted surveys in fall 2021 to compare the bird communities and vegetation complexity of two avocado farm sites and two native dry forest sites in the northern Sierra de Bahoruco region of the Dominican Republic to assess avian habitat use. Overall, we found that habitat associations differed for bird species distinguished by contrasting geographic ranges. In particular, the abundance, richness, and diversity of bird species endemic to Hispaniola were significantly greater in forests than in farms, while non-endemic resident bird species had the opposite habitat associations. The abundance, richness, and diversity of Neotropical migrants exhibited fewer distinct differences between farm and forest sites. Vegetation structural complexity was reduced in avocado farms relative to forests and was positively associated with the abundance, richness, and diversity of endemic bird species but negatively associated with these community measures of non-endemic resident species. Collectively, our findings suggest that avocado farms could provide viable habitat for non-endemic resident and migrant bird species, but that endemic species could be negatively impacted by an increase in avocado farming in the region. We offer recommendations for forest preservation, farming practices, and future research that could help to inform the conservation of endemic species in the region.

Keywords

agriculture, avian conservation, Dominican Republic, endemic species, habitat use, land-use change, Neotropical migrants

Resumen

Uso de granjas de cultivo de aquacate y bosques nativos por parte de las aves en el norte de la Sierra de Bahoruco, República Dominica • La República Dominicana está experimentando una mayor pérdida de bosques nativos como consecuencia de la expansión de los cultivos de aguacate. Aunque las tierras agrícolas pueden proporcionar un hábitat propicio para algunas aves, el conocimiento sobre el uso que hacen de los cultivos de aquacate en comparación con las áreas forestales nativas es limitado. En otoño de 2021, realizamos muestreos en dos granjas de aquacate y dos bosques secos nativos en la región norte de la Sierra de Bahoruco (República Dominicana), y comparamos las comunidades de aves y la complejidad de la vegetación para evaluar el uso del hábitat que hacen las aves. En general, observamos diferencias entre las asociaciones de hábitats de las especies de aves con distribuciones geográficas distintas. En particular, la abundancia, riqueza y diversidad de especies de aves endémicas de La Española fueron significativamente mayores en los bosques que en las granjas; mientras que las especies de aves residentes no endémicas mostraron asociaciones de hábitat opuestas. La abundancia, riqueza y diversidad de las aves migratorias neotropicales exhibieron menos diferencias entre las granjas y los bosques. La complejidad estructural de la vegetación se redujo en las granjas de aquacate en relación con los bosques y tuvo una correlación positiva con la abundancia, riqueza y diversidad de aves endémicas, pero negativa con las mismas medidas comunitarias de especies residentes no endémicas. En conjunto, nuestros resultados sugieren que los cultivos de aquacate podrían proporcionar un hábitat adecuado para aves residentes no endémicas y migratorias; mientras que las especies endémicas podrían verse afectadas negativamente por un aumento de

Cite this article as:

Young, A., D. Aborn, S.C. Latta, J. Nagel Boyd. 2023. Avian use of avocado farms and intact forest in the northern Sierra de Bahoruco, Dominican Republic. Journal of Caribbean Ornithology 36:147–161. https://doi.org/10.55431/jco.2023.36.147-161

estos cultivos en la región. Por último, proporcionamos recomendaciones para la conservación de los bosques, prácticas agrícolas y futuras investigaciones que podrían contribuir a la conservación de las aves endémicas de esta región.

Palabras clave

agricultura, aves migratorias neotropicales, cambio de uso de suelo, conservación de aves, especies endémicas, República Dominicana, uso de hábitat

Résumé

Utilisation par les oiseaux des plantations d'avocatiers et des forêts non exploitées dans le nord de la Sierra de Bahoruco, en République dominicaine • En République dominicaine, la forêt indigène est en régression en raison de l'augmentation de la culture des avocatiers. Bien que les terres agricoles puissent fournir un habitat approprié à certains oiseaux, les connaissances sur l'utilisation des plantations d'avocatiers par rapport à celle des forêts indigènes sont limitées. Nous avons échantillonné deux plantations d'avocatiers et deux forêts sèches indigènes dans la Sierra de Bahoruco (République dominicaine) au cours de l'automne 2021, et nous avons comparé leurs communautés d'oiseaux et la complexité de leur végétation afin d'évaluer l'utilisation de ces deux habitats par les oiseaux. Dans l'ensemble, nous avons observé que les associations entre espèces d'oiseaux et habitats différaient en fonction de la répartition géographique des oiseaux. En particulier, l'abondance, la richesse et la diversité de l'avifaune endémique à Hispaniola étaient significativement plus fortes dans les forêts que dans les plantations, tandis que les espèces d'oiseaux sédentaires non endémiques étaient inversement associées à ces habitats. L'abondance, la richesse et la diversité des migrateurs néotropicaux présentaient moins de différences distinctes entre les milieux agricoles et forestiers. La complexité de la structure de la végétation était plus faible dans les plantations d'avocatiers que dans les forêts, et était positivement corrélée à l'abondance, à la richesse et à la diversité des espèces d'oiseaux endémiques, mais négativement corrélée à ces mesures pour les communautés d'espèces sédentaires non endémiques. Dans l'ensemble, nos résultats semblent indiquer que les plantations d'avocatiers pourraient constituer un habitat viable pour les espèces d'oiseaux sédentaires et migrateurs non endémiques, mais que les espèces endémiques pourraient être affectées négativement par une augmentation de ces cultures dans la région. Nous formulons des recommandations pour la préservation des forêts, les pratiques agricoles et les recherches futures qui pourraient contribuer à la conservation des espèces endémiques dans la région.

Mots clés

agriculture, changement d'affectation des terres, conservation de l'avifaune, espèces endémiques, migrateurs néotropicaux, République dominicaine, utilisation des habitats

Rapid environmental change is threatening biodiversity worldwide, and birds are the vertebrate taxa with the greatest number of species on the brink of extinction (Ceballos et al. 2020). Owing to their high bird species diversity and high rates of habitat loss, the tropics are now home to the highest number of bird species threatened by extinction (Sodhi et al. 2011, BirdLife International 2017). In addition to affecting year-round resident bird species, land-use change in the Neotropics can affect Nearctic-Neotropical migrants that stop over or overwinter in these habitats. A recent assessment of the status of North America's avifauna concluded that a net loss of 3 billion birds has occurred during the last 50 years, among 419 migratory species (Rosenberg et al. 2019). Declines in bird populations have been linked directly to habitat encroachment from the conversion of land for agriculture and development (BirdLife International 2017, Tilman *et αl.* 2017). More than 50% of the world's tropical forests have already been modified by human use (Laurance et al. 2014), and bird species in highly biodiverse tropical systems have been especially negatively affected by industrial agriculture practices and other human activities (Newbold et al. 2013). However, differences in farming practices, such as crop intensity and organic farming methods do impact biodiversity loss (Tuck et al. 2014).

The island of Hispaniola, comprised of the Dominican Republic and Haiti, is used by more bird species than any other island in the Caribbean, with the exception of Cuba (Gerbracht and Levesque 2019). Hispaniola provides both important overwintering and stopover habitats for Neotropical migrants (Wunderle and Latta 1998, Latta and Brown 1999, Latta et al. 2003) and

permanent habitat for resident bird species (Latta et al. 2006). Forest-dependent breeding birds in tropical ecosystems, including endemic species, are considered to be the most vulnerable to habitat disturbance (Newbold et al. 2013), and Hispaniola harbors more endemic bird species than any other island in the Caribbean except for Jamaica (Latta et al. 2003). In particular, the Sierra de Bahoruco region in the southwestern Dominican Republic is home to 30 endemic bird species (Latta 2005, Latta et al. 2006), including nine species listed as vulnerable to extinction or endangered (IUCN 2022). Deforestation associated with land conversion for agriculture is an ongoing concern for avian biodiversity in the Dominican Republic, including in the Sierra de Bahoruco (Latta 2005). Over half of the country's deforestation is a direct result of agriculture growth, including illegal farming within national protected areas in Sierra de Bahoruco (León et al. 2013, Lloyd and León 2019). In total, the Dominican Republic experienced a loss of ~300,000 ha of tree cover from 2000–2019, with the majority of forest loss attributed to land conversion for agriculture (Global Forest Watch 2021).

Classic ecological studies have revealed a generally positive association between bird species diversity and vegetation structural complexity (MacArthur and MacArthur 1961), a factor that is generally greater in forests than most agricultural systems (Flynn et al. 2009, Karp et al. 2012). Although it has been proposed that bird and plant species diversity also should be correlated, empirical support for this association has been mixed (e.g., MacArthur and MacArthur 1961, Power 1972, Nilsson 1979, Erdelen 1984, Estades 1997, Rompré et al. 2007, Mohd-Azlan et al. 2019). While avian species diversity is generally greater in

tropical forests than agricultural lands (MacGregor-Fors and Schondube 2011), certain species may more successfully use converted agricultural landscapes based on their ecological and life-history traits (Newbold *et al.* 2013, Regos *et al.* 2018), such as diet, body size, and foraging height (Bain *et al.* 2020). Migrant species in particular may be better adapted to use modified landscapes than resident species because of generally more flexible requirements on non-breeding grounds (Hutto 1980, Villaseñor and Hutto 1995, Bender *et al.* 1998, Newbold *et al.* 2013). Given that endemic species are characterized by smaller ranges and stricter habitat requirements (Rabinowitz 1981, Cronk 1997, Fordham and Brook 2010), endemic bird species are often more vulnerable to human disturbance and landscape modification than geographically widespread bird species (Harris *et al.* 2014, Martin *et al.* 2020).

Understanding associations between avian diversity and land cover can provide important conservation guidance for the Sierra de Bahoruco region and other tropical forest areas of the Caribbean experiencing deforestation for agriculture. To date, many studies of bird use of agricultural lands in the tropics have focused on coffee and cacao plantations (e.g., Wunderle and Waide 1993, Perfecto et al. 1996, Wunderle and Latta 1996, Greenberg et al. 1997a, 1997b, Johnson 2001, Tejeda-Cruz and Sutherland 2004, Johnson et al. 2006, Van Bael et al. 2007, Mac-Gregor-Fors et al. 2018). Such research in the Dominican Republic and elsewhere has shown that shade-grown coffee plantations with significant overstory canopy are more extensively used as foraging habitat for resident and migrant birds compared with full-sun coffee plantations that lack overstory canopy (Wunderle and Waide 1993, Perfecto et al. 1996, Greenberg et al. 1997a, Wunderle and Latta 1998, Tejeda-Cruz and Sutherland 2004). This habitat preference of birds could be explained by the similarity of shade-grown coffee plantations to more structurally complex forest (Arendt et al. 2020) or the amount of native trees in shade coffee plantations relative to other agricultural lands (Narango et al. 2019). But studies also have concluded that even shade-grown coffee may not be suitable habitat for highly forest-dependent bird species (Greenberg et al. 1997b) or species sensitive to disturbance (Tejeda-Cruz and Sutherland 2004).

Avocados (Persea americana) are the most rapidly expanding tropical fruit export worldwide, and global production of avocados has been projected to triple from 2010 to 2030 (OECD-FAO 2021). Given a recent increase in avocado production in the Dominican Republic, the country is now the world's second-largest exporter of avocados (FAOSTAT 2021). The mid-elevation foothills of the Sierra de Bahoruco region have been identified as ideal sites to grow avocados, and native forest in the area has been increasingly converted for avocado farming as a result (León et al. 2013). There has been more documentation of forest loss from illegal and legal avocado farming on the southern slope of the Sierra de Bahoruco, with an estimated 3,400 ha of forest loss from agricultural practices including avocado and coffee farming (León et al. 2013). The impacts of avocado farming on the northern slope has not yet been studied in similar detail, but avocado agriculture threatens these native dry forests as well (León et al. 2013) with an estimated 260 ha of avocado farms in the Puerto Escondido area (Y.M. León pers. comm.). Similar to other tree plantations such as coffee and cacao, avocado plantations can comprise or be integrated within a more complex vegetation structure than non-tree crop farms and could provide viable habitat for bird species (Mokria *et al.* 2022). However, the use of avocado farms by birds, including migrant, non-endemic resident, and endemic species, has not been previously reported in the primary literature.

To investigate avian use of avocado farms in the Dominican Republic's northern Sierra de Bahoruco region, we compared bird species richness and diversity in avocado farms and native dry forest. We considered overall, migrant, and endemic bird species in our analysis, and concurrently assessed the vegetation complexity of farms and forest to better understand factors affecting the diversity and abundance of bird species. We predicted that farms would be characterized by lower vegetation complexity and that this reduced complexity would be associated with lower bird species diversity. Given their habitat associations, we categorized bird species as forest-dependent or generalist and made associated recommendations for land-use and conservation priorities.

Methods

Study Sites

The study area is located near the town of Puerto Escondido, Independéncia Province, in the foothills of the northern Sierra de Bahoruco in the southwestern Dominican Republic. Natural forests in this area are considered subtropical dry forests (Knudson et al. 1988) with elevations ranging from 421 to 522 m. Dominant vegetation in the region includes large trees such as baitoa (*Phyllostylon brasiliensis*), candelón (*Acacia scleroxyla*), and guayacán (*Guaiacum officinale*) and understory shrubs such as escobón (*Eugenia monticola*) and grenadilla (*E. ligustrina*). The town of Puerto Escondido is situated near multiple limestone water outlets, providing water for community needs and agricultural irrigation.




Fig. 1. Map depicting the location of our small farm and La Placa and Rabo de Gato forest study sites (all in green) within the Sierra de Bahoruco region of the Dominican Republic. Dark plots within the small farm area represent active avocado farms. Our large farm study site is not depicted due to a privacy agreement with the landowner.

Table 1. Details of the study sites in the northern Sierra de Bahoruco region of the Dominican Republic in which bird communities and vegetation complexity were assessed in October–December 2021. Due to landowner restrictions, we are unable to provide location information for the large farm site.

| | Municipality, | | | | |
|-----------------------------|--|---------------------------------|---------------|-----------|-------------------|
| | Province | Coordinates | Elevation (m) | Area (ha) | Survey points (n) |
| Forests | | | | | |
| La Placa | Reserve Loma de Charco Azul, Independéncia | 18°35'50.36"N, 71°65'66.01"W | 522 | 173 | 29 |
| Rabo de Gato | Puerto Escondido, Independéncia | 18°30'80.08"N, 71°59'23.04"W | 464 | 137 | 24 |
| Avocado Farms Large Farm | | | 442 | 273 | 29 |
| Small Farms | Puerto Escondido, Independéncia | 18°32'26.94"N, 71°58'99.36"W | 421 | 41 | 20 |

We conducted field work in four study sites within the area: two native forest sites and two avocado farm sites. Our first forest site was located in Reserva Loma Charco Azul, a federally protected 17,400-ha dry forest reserve (Fig. 1), known for being the main habitat of the endemic Bay-breasted Cuckoo (Coccyzus rufigularis). The second forest site was located adjacent to a portion of a well-known birding trail called Rabo de Gato (Fig. 1). Avocado farms of two sizes were chosen for the comparison to reflect the variety of such farms in the region, including potential differences in vegetation complexity. Specifically, we chose one farm site comprised of multiple, relatively small community parcelas; this is land near town that is divided into adjacent plots, each with a single owner who primarily cultivates one crop. All parcelas cultivating avocados were included in the small farm study site, which was located within 1 km of our forest site at Rabo de Gato (Fig. 1). The second farm site was located in a large organic commercial farm characterized by groves of primarily Hass and Carla avocados. Due to restricted permissions by the landowner, we are unable to disclose the specific location of our large farm site. Both avocado farm sites border native forest, which minimized potential differences in climate, topography, and human disturbance as a result of proximity to town.

Bird Community Surveys

We used point count surveys (Wunderle and Waide 1993, Wunderle and Latta 1996, Greenberg *et al.* 1997a, Marsden *et al.* 2001, Latta *et al.* 2003, Kennedy *et al.* 2010) to assess overall, endemic, and migrant bird communities in the study sites. We designated survey points using a grid system in which all points were at least 150 m apart to reduce the possibility of double-counting individual birds. We also ensured that all points were > 25 m away from the boundary of the site so that only species within the site were detected during surveying. The size and elevation of our study sites and the number of survey points in each site are listed in Table 1.

We conducted point count surveys from 13 October to 1 December 2021 to include resident overwintering migrant species. We visited each point three times during the field season, and alternated between forest and farm point counts on consecutive days to account for differences due to the effects of weather or other short-term temporal events. We began surveys at sunrise and ended no later than four hours after sunrise, encompassing the period when birds are most active and likely to be detected. We recorded all birds seen or heard within 25 m of each point for a 10-min period. Using a relatively small point count radius ensured that we were recording bird species within the habitat type we were surveying, which was especially critical in smaller avocado farms. Birds that flushed upon arrival were included in the point count, but birds flying over without stopping in the site were not counted. We averaged the number of individual birds and species across the three survey dates at each point. To investigate habitat use by non-endemic resident, endemic, and Neotropical migrant species, in particular, we similarly calculated the mean number of individuals and bird species within each group per survey point. Endemic bird species were defined as those only found in Hispaniola and its satellite islands (Latta et al. 2006), and migrant bird species were defined as those that breed north of the Tropic of Capricorn but winter south of this location (Hagan and Johnston 1992). Migrants may be passing through to locations further south or overwintering at our site, but by mid-October almost all migrants that are present are winter residents (Wunderle and Latta 1996). We used the term 'non-endemic resident' to describe more geographically widespread, non-migratory species observed in our study sites.

To characterize the diversity of the non-endemic resident, endemic, and migrant bird communities in each study site, we calculated the frequency of each species identified as the percentage of the total bird survey points in each site in which that species was observed across all three site visits. We used the Shannon Diversity Index (Shannon 1948) to calculate the diver-

sity of the overall, non-endemic resident, migrant, and endemic bird communities in each site during each visit. We averaged the diversity scores of the three visits to calculate a mean diversity measure per point. Bird survey details, including the distance that birds were identified from survey points, method of detection, weather variables, and ambient noise are provided in our Dryad dataset (Young et al. 2023).

Vegetation Complexity Measures

We assessed canopy cover, canopy height, and canopy thickness at each bird survey point and at four additional subpoints, one in each cardinal direction 25 m from the main survey point (see Van Bael et al. 2007). Canopy cover was estimated by using a tube densiometer at each survey point and subpoint. To determine canopy height and canopy thickness, we used a digital rangefinder to measure height of the highest and lowest foliage associated with trees at each survey point and subpoint. Canopy thickness was calculated as the upper height minus the lower height at each measurement location (see Muth and Bazzaz 2002, Mas and Dietsch 2004, Lee et al. 2020). We averaged the vegetation data collected at each survey point and its subpoints to calculate the mean canopy cover, canopy height, and canopy thickness for each survey point.

To assess the diversity of vegetation in each study site, we defined the five most dominant woody species ≥ 1 m tall as the species with the highest vegetative cover by visually estimating percent cover in a 25-m radius around each bird survey point (see McLaren et al. 2019). For cultivated crops, we included horticultural varieties as 'species' in our analyses given that horticultural varieties are selected for distinct differences in traits that could potentially impact habitat preference, including leaf shape and size and fruiting phenology. To characterize the diversity of the vegetation in each study site, we calculated the frequency of each dominant plant species as the percentage of the total sur-

vey circles in each site in which that species was observed across all three site visits. Vegetation survey details are provided in our Dryad dataset (Young *et al.* 2023).

Statistical Analyses

We tested for the normality of our data distribution within groups (i.e., sites) with the Shapiro-Wilk test and for homogeneity of variances among groups with Levene's test. When the assumptions of normality and homogeneity of variances were both met, we used a one-way ANOVA to test for the effect of site on vegetation complexity (i.e., canopy height, canopy cover, canopy thickness) and bird community measures (i.e., number of individuals, species richness, species diversity) followed by least-significant difference LSD post-hoc tests when the effect was significant. When the assumption of normality was met but variances were not homogenous, we used Welch's ANOVA followed by Games-Howell post-hoc tests when the effect was significant. When the assumption of normality was not met but variances were homogenous, we used non-parametric Kruskal-Wallis tests to test for the main effect of site followed by Dunn-Bonferroni post-hoc tests when the effect was significant. In all cases, differences in mean values were considered significant if $p \le 0.05$. To examine associations between vegetation complexity and bird community measures, we performed linear regression analyses. To account for the numerous associations analyzed, we calculated and utilized corrected p-values to control the false discovery rate for the group of tests for each bird community type (i.e., non-endemic resident, endemic, migrant) in accordance with methods described by Benjamini and Hochberg (1995). Results of regression analyses were considered significant if False Discovery Rate (FDR)-corrected $p \le 0.05$. All data analyses were conducted with SPSS (version 29, IBM Corp., Armonk, NY, USA).




Fig. 2. Frequency of the 20 most commonly observed bird species at each study site expressed as the percentage of total survey points at which each species was observed. Study sites included small and large avocado farms and native dry tropical forest at La Placa and Rabo de Gato. Bird species were categorized as non-endemic residents, endemics, or Neotropical migrants as described in our methods. Abbreviations, English common names, and scientific names for each species are provided in Appendix 1.




Fig. 3. Abundance of individuals (a), species richness (b), and species diversity (c) of non-endemic residents, endemic, and Neotropical migrant bird species in small and large avocado farm sites and native dry tropical forest sites at La Placa and Rabo de Gato. Values shown are means averaged across multiple survey points in each site ± 1 SE of the mean. Within species groups (i.e., non-endemic residents, endemics, migrants), values shown below the same lowercase letter do not differ at the $p \le 0.05$ level of significance.

Results

Bird Community Comparisons

The number of bird species detected across all survey points was similar in all of our study sites (38 species in the big farm, La Placa, and Rabo de Gato; 36 species in the small farm), but the relative abundance of bird species in the forest sites exhibited more evenness than in the farm sites (Fig. 2; see Appendix 1 for bird species abbreviations and scientific names). Of the 20 most frequently observed bird species at each site, the farm sites were characterized by greater numbers of non-endemic resident species than the forest sites, while the forest sites were




Fig. 4. Frequency of the 20 most commonly observed woody plant species or varieties at each study site expressed as the percentage of total survey points at which each species was observed. Study sites included small and large avocado farms and native dry tropical forest at La Placa and Rabo de Gato. Abbreviations, local common names, and scientific names for each species are provided in Appendix 2.

characterized by greater numbers of endemic bird species than the farm sites (Fig. 2). Detection frequencies of migrants were similar in the big farm and forest sites, but migrants were less frequently detected in the small farm site (Fig. 2).

The mean number of non-endemic resident bird individuals observed per survey point was significantly greater in both farm sites than in both forest sites (Welch's ANOVA: F3,49.096 = 30.14, $p \le 0.001$; Fig. 3a). Compared with the forest sites, the farm sites also were characterized by greater non-endemic resident species richness (F3,98 = 27.69, $p \le 0.001$; Fig. 3b) and diversity (H3 = 38.14, $p \le 0.001$; Fig. 3c). Endemic bird communities exhibited opposite trends. Specifically, both farm sites were characterized by significantly lower numbers of endemic individuals (F3,98 = 36.67, $p \le 0.001$; Fig. 3a), endemic species richness (Welch's ANOVA: F3,51.963 = 52.21, $p \le 0.001$; Fig. 3b), and endemic species diversity (H3 = 64.87, $p \le 0.001$; Fig. 3c) than both forest sites. The mean abundance of migrant birds observed per




Fig. 5. Canopy cover (a), canopy height (b), and canopy thickness (c) in small and large avocado farms sites and native dry tropical forest sites at La Placa and Rabo de Gato. Values shown are means averaged across multiple survey points in each site ± 1 SE of the mean. Values shown below the same lowercase letter do not differ at the $p \le 0.05$ level of significance.

sampling point was lower in the small farm than in all of the other sites, while the mean number of migrant individuals in the big farm was less than in La Placa but not Rabo de Gato (H3 = 25.89, $p \le 0.001$; Fig. 3a). The small farm also was characterized by less migrant species richness and diversity than all of the other sites, while the big farm had lower richness and diversity of migrant species than La Placa only (F3,98 = 11.61, $p \le 0.001$ for migrant species richness; F3,98 = 7.972, $p \le 0.001$ for migrant species diversity; Figs. 3b, c).

Vegetation Complexity Comparisons

More woody plant species or varieties comprised the dominant vegetation in the forest sites (25 species in La Placa, 21 species in Rabo de Gato) than in the farm sites (8 species or

varieties in the big farm, 9 species or varieties in the small farm), and the relative abundance of woody species in the forest sites exhibited more evenness than in the farm sites (Fig. 4; see Appendix 2 for plant species and variety abbreviations and scientific names). The five most dominant woody species observed in farm sites comprised ~95% of the total woody species observed in those sites (Fig. 4). In contrast, the five most dominant woody plant species observed in forest sites comprised ~35% of the total woody species observed in those sites (Fig. 4). As expected, woody vegetation in the farm sites was largely comprised of avocado varieties (Fig. 4), which collectively comprised ≥ 90% of the total trees surveyed in each farm. Both farm sites were characterized by less vegetation structural complexity assessed as mean canopy cover (H₃ = 33.53, $p \le$ 0.001; Fig. 5a), canopy height (Welch's ANOVA: F3,45.469 = 41.27, $p \le$ 0.001; Fig. 5b), and canopy thickness (Welch's ANOVA: F3,48.926 = 35.24, $p \le 0.001$; Fig. 5c) than both forest sites.

Relationships of Vegetation Complexity and Bird Communities

The number of individuals and the richness and diversity of endemic bird species were positively correlated with all measures of vegetation structural complexity (Table 2). In contrast, the mean abundance and species richness and diversity of non-resident endemic species and Neotropical migrants were not significantly correlated with any measures of vegetation complexity (Table 2).

Discussion

Avian Use of Forests versus Farms

Our comparison of bird communities in two avocado farm sites and two forest sites in the northern Sierra de Bahoruco, Dominican Republic, suggests that the increase in avocado farms could affect bird communities in the region but that the impacts may differ among types of birds and types of farms. Overall, we found that avocado farms could provide suitable habitat for non-endemic resident, endemic, and Neotropical migratory bird species as we observed numerous individuals from multiple species of all of these groups in all of the study sites. However, further study on species and species assemblage health metrics over time would be needed to determine the true viability of this habitat. We also found that avocado farms were frequented by fewer endemic species but more non-endemic resident bird species than forests (Fig. 2), suggesting that forest habitats are preferred by endemic species.

Our observation of reduced abundances and diversity of endemic bird species in both farm sites (Fig. 3) also suggests that native forest habitat is especially important to endemic bird species in the northern Sierra de Bahoruco region. Many endemic species that we observed, such as both the Broad-billed Tody (*Todus subulatus*) and Narrow-billed Tody (*T. angustirostris*), were found almost exclusively in the forests. Distinctions in vegetation complexity between farms and forest (Figs. 4, 5) and associations between vegetation complexity and endemic bird communities (Table 2) suggest that the presence of a dense canopy is important to endemic species. Given their narrow geographical ranges, endemic species are more likely to be dependent on specific habitats (Harris and Pimm 2004). For example,

Table 2. Results of linear regression analyses of associations between measures of vegetation complexity as predictors and non-endemic resident, endemic, and Neotropical migrant bird communities as response variables. Regression that are significant have p-values ≤ 0.05. Negative slopes indicate negative correlations between the predictor and response variable, and vice versa.

| Predictor | Response Variable | Slope | R ² | <i>P</i> -value |
|------------------|-----------------------|--------|----------------|-----------------|
| Canopy Cover | Non-endemic Residents | | | |
| | Number of Individuals | -0.124 | 0.936 | 0.0990 |
| | Species Richness | -0.050 | 0.853 | 0.0760 |
| | Species Diversity | -0.014 | 0.865 | 0.0788 |
| | Endemics | | | |
| | Number of Individuals | 0.084 | 0.972 | 0.0158 |
| | Species Richness | 0.062 | 0.985 | 0.0103 |
| | Species Diversity | 0.024 | 0.972 | 0.0140 |
| | Neotropical Migrants | | | |
| | Number of Individuals | 0.038 | 0.693 | 0.7515 |
| | Species Richness | 0.027 | 0.716 | 0.7515 |
| | Species Diversity | 0.008 | 0.691 | 0.5070 |
| Canopy Height | Non-endemic Residents | | | |
| .,, - | Number of Individuals | -1.258 | 0.963 | 0.1710 |
| | Species Richness | -0.516 | 0.904 | 0.0821 |
| | Species Diversity | -0.147 | 0.909 | 0.1058 |
| | Endemics | | | |
| | Number of Individuals | 0.841 | 0.992 | 0.0120 |
| | Species Richness | 0.625 | 0.997 | 0.0180 |
| | Species Diversity | 0.239 | 0.992 | 0.0090 |
| | Neotropical Migrants | | | |
| | Number of Individuals | 0.355 | 0.617 | 0.2408 |
| | Species Richness | 0.258 | 0.638 | 0.3618 |
| | Species Diversity | 0.079 | 0.615 | 0.2160 |
| Canopy Thickness | Non-endemic Residents | | | |
| | Number of Individuals | -1.463 | 0.956 | 0.0990 |
| | Species Richness | -0.598 | 0.892 | 0.0707 |
| | Species Diversity | -0.171 | 0.893 | 0.0795 |
| | Endemics | | | |
| | Number of Individuals | 0.979 | 0.989 | 0.0108 |
| | Species Richness | 0.728 | 0.994 | 0.0135 |
| | Species Diversity | 0.278 | 0.988 | 0.0090 |
| | Neotropical Migrants | | | |
| | Number of Individuals | 0.421 | 0.637 | 0.3030 |
| | Species Richness | 0.306 | 0.657 | 0.4253 |
| | Species Diversity | 0.094 | 0.635 | 0.2610 |

the Bay-breasted Cuckoo, which we observed only in La Placa, inhabits exclusively low to mid-elevation forest in the transitional zone between dry and broadleaf forest of Hispaniola (Latta et al. 2006). Another endemic species, the Hispaniolan Emerald (Riccordia swainsonii), relies on high elevation moist habitat

where it makes nests from wet forest mosses and lichen (Latta *et al.* 2006). We suggest that many endemic species in the Sierra de Bahoruco may be less able to adapt to or utilize human-modified landscapes, such as avocado farms, relative to overwintering migrants and non-endemic resident birds.

In contrast, the greater frequency, abundance, richness, and diversity of non-endemic resident bird species in our avocado farm sites compared with the forest sites (Figs. 2, 3) suggests an ability to adapt or adjust to land-use change that reduce vegetation complexity (Figs. 4, 5; Table 2). Birds frequently observed in avocado farms included numerous common resident species in the Caribbean, such as Yellow-faced Grassquit (Tiaris olivaceus), Northern Mockingbird (Mimus polyglottos), Gray Kingbird (Tyrannus dominicensis), and Smooth-billed Ani (Crotophaga ani; Latta et al. 2006). These species are not usually observed in forests, but are more generally associated with open landscapes, such as grasslands and disturbed habitats (Latta et al. 2006). Increased environmental heterogeneity in human-disturbed landscapes, such as farmlands, can lead to higher niche availability (Lack 1969, Tews et al. 2004, Stein et al. 2014), and may be one reason for the overall high diversity of non-endemic residents in avocado farms. We suggest that the commonness of species associated with such habitats could be facilitated by the high rates of deforestation in the Caribbean in recent decades (López-Carr et al. 2022). In contrast, studies have revealed that habitat specialists are less frequent in disturbed landscapes than generalist species (Devictor et al. 2008), a pattern that we suggest is reflected by the low frequency of endemic species in avocado farms in our study.

Our surveys beginning in mid-October may have included transient individuals that were only stopping over at our sites, but all species we observed are known to overwinter on the island. Although Neotropical migrant birds were more frequently observed in forest sites compared with small avocado farms (Fig. 2), our results suggest that large farms could support an abundance of migrant individuals and a diverse migrant community comparable to that of some forest habitat (Fig. 3). However, we acknowledge that our research did not include replicate large farm or small farm sites. As such, these results could reflect local-scale patterns that are not related to farm size, and we encourage future replicate studies to help determine the influence of farm size on bird diversity. Migrants have been known to utilize disturbed habitats while they were on non-breeding grounds (Hutto 1980, Villaseñor and Hutto 1995, Bender et al. 1998, Newbold et al. 2013), but our results suggest that migrant utilization of farms differs across sites and species. For example, the Prairie Warbler (Setophaga discolor) was never observed in the small farm site. Similarly, the Ovenbird (Seiurus aurocapilla) was seen only once in the small farm site but was observed in more than half of the sampling points in the big farm and both forest sites. Given the similar complexity of vegetation of our small and large farm sites (Figs. 4, 5), differences in their migrant species and communities suggest that other factors could influence migrant habitat choice. These factors could include food availability (Martin and Karr 1986, Johnson and Sherry 2001), broader matrix complexity (Conway et al. 1995), or behaviors such as site fidelity (Johnson and Sherry 2001). Previous research on migrant use of forest habitat of varying successional stages in Mexico (Smith et al. 2006) and of different habitats in the southern United States (McClure et al. 2012) also found that the structural complexity of vegetation did not affect habitat quality for migrant species.

Conservation Implications

The conversion of forests to agriculture associated with avocado production in the Sierra de Bahoruco region, while not necessarily impactful to bird species overall, could have a negative impact on endemic species in Hispaniola, many of which are already vulnerable to extinction (Latta et al. 2003, Latta 2005, Isik 2011, León et al. 2013). However, although endemic communities were more robust in our forest sites than farm sites, some specific endemic species seemed to be more generalist in their habitat preferences. For example, both Hispaniolan Woodpecker (Melanerpes striatus) and Hispaniolan Lizard-Cuckoo (Coccyzus longirostris), which are considered common throughout the island (Latta et al. 2006), were found frequently in both forest and farm sites. Other endemic species seemed to prefer habitat that is less forested. In particular, the White-necked Crow (Corvus leucognaphalus), which has a limited geographical range and is listed as vulnerable to extinction (IUCN 2022), was observed more frequently in the small farm than in any of the other sites. Notably, we did not observe this species in avocado trees, but rather in larger remnant trees in the small farm site. This observation suggests that shade trees within farmlands may improve habitat quality for bird species in agricultural landscapes.

Caribbean islands are hotspots of endemism and should thus be considered as areas of high conservation priority (Latta 2005, Buchanan et al. 2011, Devenish-Nelson et al. 2019). Hispaniola in particular is home to 31 endemic bird species and 50 endemic subspecies (Latta et al. 2006). The IUCN Red List (IUCN 2022) shows that information about habitat requirements and population numbers is deficient or lacking altogether for many bird species endemic to Hispaniola, which limits the ability to make specific conservation recommendations. Many Hispaniolan endemics are forest-dependent specialists, and our findings demonstrate the vulnerability of endemic species to habitat modification involving deforestation (Greenberg et al. 1997b). Endemic species including the Green-tailed Warbler (Microligea palustris), Narrow-billed Tody, and Flat-billed Vireo (Vireo nanus), which are all considered to have decreasing or data-deficient populations (IUCN 2022), were observed only once or not at all in the farm sites, suggesting that these endemic species are forest-dependent and therefore vulnerable to deforestation. The Bay-breasted Cuckoo was only observed at La Placa, suggesting that the distribution of this species may be especially narrow or reliant on specific habitat requirements, such as limited human disturbance and the presence of large dry forest trees for nesting. The Hispaniolan Parrot (Amazona ventralis), considered vulnerable to extinction (IUCN 2022), was also found only in La Placa. Additional species of concern such as the Broadbilled Tody and Antillean Piculet (Nesoctites micromegas) were all found in higher frequencies in the forest sites than in avocado

When evaluating avian use of agricultural land, it is important to consider differences between farms, as we found for migrant use of the large and small avocado farm sites. Our results suggest that Neotropical migrants may be less influenced by vegetation complexity than non-endemic resident and endemic bird species, but could be influenced by other habitat-selection factors. The small farm site included in our study was characterized by a patchwork of individual parcelas that are used to grow a

variety of crops. Arboreal croplands—such as avocado, coffee, and cacao—have been shown to have higher diversity of birds than non-arboreal croplands (Estrada and Coates-Estrada 1997). We suggest that crop type variation between parcelas could have resulted in smaller arboreal crop patches within the small farm site, which could have decreased utilization by migrant species relative to the more continuous arboreal cropland of the large farm site. Our study did not consider arboreal patch sizes within overall farm sites, which may be an important factor to consider in future research of avian use of avocado farm landscapes in the region. We did observe that endemics such as Hispaniolan Oriole (Icterus dominicensis) and White-necked Crow, as well as many migrant warblers, were more likely to be found in the larger trees in the small farm site, such as palm trees, mango trees, or non-pruned large local hybrid avocado trees. This same trend was not observed in the big farm site given that there were very few non-cultivated trees in the site. Overall, the differences that we observed in migrant communities in the small and large farm sites suggest that individual agricultural practices could be important to avian use.

As agriculture continues to expand into tropical forest landscapes (FAO 2009), particularly in areas of high conservation value such as endemic hotspots (Scharlemann et al. 2004), the question remains how we can best ensure compatibility of agriculture with species conservation. A variety of agricultural practices have been shown to positively impact habitat quality for birds, such as the preservation of native trees (Douglas et al. 2013) including relict shade trees (Wunderle and Latta 1996, Carlo et al. 2004), retention of native habitat patches within cultivated landscapes (Estrada-Carmona et al. 2019), and use of live fences (Harvey et al. 2005). Many tropical bird species have been found to move freely between forest and agricultural lands to benefit from seasonal resources (Estrada and Coates-Estrada 1997), which we suggest could be facilitated by close proximity of farms to intact forest. Forests and scattered trees have been demonstrated in other agricultural areas to improve ecological services including soil fertility and the provision of habitat for native species (Bommarco et al. 2013, Reed et al. 2017, Yang et al. 2020). Both of our farm sites were adjacent to native forest, and future research of avian utilization of avocado farms at different distances from intact forest could provide more understanding of bird movement patterns and farmland habitat quality. While 'bird-friendly' avocado farming practices and proximity of farms to intact forest could enhance avian use, we do not know if these practices would help to support forest specialists, such as the many bird species endemic to Hispaniola in the northern Sierra de Bahoruco. Given the strong association of endemic species with forests demonstrated in our study, we suggest that the preservation of native dry forest in federally-protected reserves like Reserva Loma Charco Azul (La Placa) will be essential for the conservation of endemic bird species.

Acknowledgments

We thank Kate Wallace and staff at Villa Barrancoli for providing site access and lodging, Jarroll (Pilo) Urbaez for field assistance and his sharing of relevant local knowledge, and Yolanda León and Héctor (Chapa) Andujar of Grupo Jaragua for help with logistics at the onset of this project. We gratefully acknowledge

support for this research from the David S. Lee Fund for Conservation of BirdsCaribbean and the Graduate School of the University of Tennessee at Chattanooga. We are grateful to the Ministerio de Medio Ambiente y Recursos Naturales de la República Dominicana and to avocado growers for access and permission to use the study sites. We appreciate Fernando Alda-Pons for translating our abstract and keywords into Spanish. We thank the two anonymous reviewers for their helpful comments on earlier drafts of this manuscript. SL and AY conceived of the project idea; AY and DA sought funding; AY, DA, and SL designed the avian methods; AY and JB designed the vegetation methods; AY collected the data; AY and JB analyzed data and wrote the paper. All authors edited versions of the manuscript.

Literature Cited

Arendt, W.J., R.C. Tuckfield, J.C. Martinez-Sanchez, J.L. Reidy, F.R. Thompson, III, J.M. Wunderle, Jr., and J.M. Zolotoff. 2020. Avian community density and distribution patterns among Nicaraguan conventional and organic shade-coffee plantations. Agricultural Sciences 11:27–53.

Bain, G.C., M.A. MacDonald, R. Hamer, R. Gardiner, C.N. Johnson, and M.E. Jones. 2020. Changing bird communities of an agricultural landscape: declines in arboreal foragers, increases in large species. Royal Society Open Science 7:1–20.

Bender, D.J., T.A. Contreras, and L. Fahrig. 1998. Habitat loss and population decline: a meta-analysis of patch size effect. Ecology 79:517–533.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B 57:289–300.

BirdLife International. 2017. Threatened birds occur in all habitats, but the majority are found in forest. datazone.birdlife. org/sowb/casestudy/threatened-birds-occur-in-all-habitats-but-the-majority-are-found-in-forest.

Bommarco, R., D. Kleijn, and S.G. Potts. 2013. Ecological intensification: harnessing ecosystem services for food security. Trends in Ecology and Evolution 28:230–238.

Buchanan, G.M., P.F. Donald, and S.H.M. Butchart. 2011. Identifying priority areas for conservation: a global assessment for forest-dependent birds. PLoS ONE 6:e29080.

Carlo, T.A., J.A. Collazo, and M.J. Groom. 2004. Influences of fruit diversity and abundance on bird use of two shaded coffee plantations. Biotropica 36:602–614.

Ceballos, G., P.R. Ehrlich, and P.H. Raven. 2020. Vertebrates on the brink as indicators of biological annihilation and the sixth mass extinction. PNAS 117:13596–13602.

Conway, C.J., G.V.N. Powell, and J.D. Nichols. 1995. Overwinter survival of Neotropical migratory birds in early-successional and mature tropical forests. Conservation Biology 9:855–864.

Cronk, Q.C.B. 1997. Islands: stability, diversity, conservation. Biodiversity and Conservation 6:477–493.

Devenish-Nelson, E.S., D. Weidemann, J. Townsend, and H.P. Nelson. 2019. Patterns in island endemic forest-dependent bird research: the Caribbean as a case-study. Biodiversity and Conservation 28:1885–1904.

Devictor, V., R. Julliard, and F. Jiguet. 2008. Distribution of spe-

- cialist and generalist species along spatial gradients of habitat disturbance and fragmentation. Oikos 117:507–514.
- Douglas, D.J.T., D. Nalwanga, R. Katebaka, P.W. Atkinson, D.E. Pomeroy, D. Nkuutu, and J.A. Vickery. 2013. The importance of native trees for forest bird conservation in tropical farmland. Animal Conservation 17:256–264.
- Erdelen, M. 1984. Bird communities and vegetation structure: I. Correlations and comparisons of simple and diversity indices. Oecologia 61:277–284.
- Estades, C.F. 1997. Bird-habitat relationships in a vegetational gradient in the Andes of central Chile. Condor 99:719–727.
- Estrada, A., and R. Coates-Estrada. 1997. Anthropogenic landscape changes and avian diversity at Los Tuxtlas, Mexico. Biodiversity and Conservation 6:19–43.
- Estrada-Carmona, N., A. Martínez-Salinas, F.A.J. DeClerck, S. Vílchez-Mendoza, and K. Garbach. 2019. Managing the farmscape for connectivity increases conservation value for tropical bird species with different forest-dependencies. Journal of Environmental Management 250:109504.
- Flynn, D.F.B., M. Gogol-Prokurat, T. Nogeire, N. Molinari, B. Trautman Richers, B.B. Lin, N. Simpson, M.M. Mayfield, and F. DeClerck. 2009. Loss of functional diversity under land use intensification across multiple taxa. Ecology Letters 12:22–33.
- Food and Agriculture Organization of the United Nations (FAO). 2009. How to Feed the World in 2050. Meeting of Experts, Rome, Italy. fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf.
- Food and Agriculture Organization of the United Nations, Statistical Division (FAOSTAT). 2021. Production Quantities of Avocados by Country. fao.org/faostat/en/#data/QCL/visualize.
- Fordham, D.A., and B.W. Brook. 2010. Why tropical island endemics are acutely susceptible to global change. Biodiversity Conservation 19:329–342.
- Gerbracht, J., and A. Levesque. 2019. The complete checklist of the birds of the West Indies: v1.1. BirdsCaribbean Checklist Committee. birdscaribbean.org/caribbean-birds/.
- Global Forest Watch. 2021. Dominican Republic. globalforest-watch.org/dashboards/country/DOM/.
- Greenberg, R., P. Bichier, A. Cruz Angon, and R. Reitsma. 1997a. Bird populations in shade and sun coffee plantations in Central Guatemala. Conservation Biology 11:448–459.
- Greenberg, R., P. Bichier, and J. Sterling. 1997b. Bird populations in rustic and planted shade coffee plantations of Eastern Chiapas, Mexico. Biotropica 29:501–514.
- Hagan, J.M., III, and D.W. Johnston. 1992. Introduction. Pp. 1–3 *in* Ecology and Conservation of Neotropical Migrant Landbirds. Smithsonian Institution Press, Washington, D.C.
- Harris, G.M, and S.L. Pimm. 2004. Bird species tolerance of secondary forest habitats and its effects on extinction. Conservation Biology 18:1607–1616.
- Harris, J.B.C., D.P. Dadang, S.D. Gregory, B.W. Brook, D.M. Prawiradilaga, N.S. Sodhi, D. Wei, D.A. Fordham, and B. Bradley. 2014. Rapid deforestation threatens mid-elevational endemic birds but climate change is most important at higher elevations. Diversity and Distributions 20:773–785.
- Harvey, C.A., C. Villanueva, J. Villacís, M. Chacón, D. Muñoz, M. López, M. Ibrahim, R. Gómez, R. Taylor, J. Martinez, A. Navas, J. Saenz, D. Sánchez, A. Medina, S. Vilchez, B. Hérnandez, A.

- Perez, F. Ruiz, F. López, I. Lang, and F.L. Sinclair. 2005. Contribution of live fences to the ecological integrity of agricultural landscapes. Agriculture, Ecosystems and Environment 111:200–230.
- Hutto, R.L. 1980. Winter habitat distribution of migratory land birds in western Mexico, with special reference to small, foliage-gleaning insectivores. Pp. 181–204 *in* Migrant Birds in the Neotropics: Ecology, Behavior, Distribution and Conservation (A. Keast and E.S. Morton, eds.). Smithsonian Institution Press, Washington, DC.
- International Union for the Conservation of Nature (IUCN). 2022. IUCN Red List of Threatened Species, version 2021-3. iucnredlist.org
- Isik, K. 2011. Rare and endemic species: why are they prone to extinction? Turkish Journal of Botany 35:411–417.
- Johnson, M.D. 2001. Effects of shade-tree species and crop structure on the winter arthropod and bird communities in a Jamaican shade coffee plantation. Biotropica 32:133–145.
- Johnson, M.D., and T.W. Sherry. 2001. Effects of food availability on the distribution of migratory warblers in Jamaica. Journal of Animal Ecology 70:546–560.
- Johnson, M.D., T.W. Sherry, R.T. Holmes, and P.P. Marra. 2006. Assessing habitat quality for a migratory songbird wintering in natural and agricultural habitats. Conservation Biology 20:1433–1444.
- Karp, D.S., A.J. Rominger, J. Zook, J. Ranganathan, P.R. Ehrlich, and G.C. Daily. 2012. Intensive agriculture erodes ß-diversity at large scales. Ecology Letters 15:963–970.
- Kennedy, C.M., P.P. Marra, W.F. Fagan, and M.C. Need. 2010. Landscape matrix and species traits mediate responses of Neotropical forest fragmentation in Jamaica. Ecological Monographs 80:651–669.
- Knudson, D.M., W.R. Chaney, and F.A. Reynoso. 1988. Fuelwood and charcoal research in the Dominican Republic. Purdue University, Department of Forestry and Natural Resources, Lafayette. IN.
- Lack, D. 1969. The number of bird species on islands. Bird Study 16:193–209.
- Latta, S.C. 2005. Complementary areas for conserving avian diversity on Hispaniola. Animal Conservation 8:69–81.
- Latta, S.C., and C. Brown. 1999. Autumn stopover ecology of the Blackpoll Warbler (*Dendroica striata*) in thorn scrub forests of the Dominican Republic. Canadian Journal of Zoology 77:114–1156.
- Latta, S., C. Rimmer, A. Keith, J. Wiley, H. Raffaele, K. McFarland, and E. Fernandez. 2006. Species Accounts. Pp. 21–217 *in* Birds of the Dominican Republic and Haiti. Princeton University Press, Princeton, NJ.
- Latta, S.C., C.C. Rimmer, and K.P. McFarland. 2003. Winter bird communities in four habitats along an elevational gradient on Hispaniola. Condor 105:179–197.
- Laurance, W.F., S. Sayer, and K.G. Cassman. 2014. Agricultural expansion and its impacts on tropical nature. Trends in Ecology and Evolution 29:107–116.
- Lee, Y.-F., Y.-M. Kuo, W.-C. Chu, and Y.-H. Lin. 2020. Perch use by flycatching *Rhinolophus formosae* in relation to vegetation structure. Journal of Mammalogy 101:455–463.
- León, Y.M., E. Garrido, and J. Almonte. 2013. Pp. 1-36 in Moni-

- toring and Mapping Broadleaf Mountain Forests of Southern Sierra de Bahoruco, Dominican. Grupo Jaragua, Inc., Santo Domingo, Dominican Republic.
- Lloyd, J., and Y. León. 2019. Forest change within and outside protected areas in the Dominican Republic. bioRxiv 558346 (preprint).
- López-Carr, D., S.J. Ryan, and M.L Clark. 2022. Global economic and diet transitions drive Latin American and Caribbean forest change during the first decade of the century: a multi-scale analysis of socioeconomic, demographic, and environmental drivers of local forest cover change. Land 11:1–11.
- MacArthur, R.H., and J.W. MacArthur. 1961. On bird species diversity. Ecology 42:592–598.
- MacGregor-Fors, I., F. Gonzáles-García, C. Hernández-Lara, and D. Santiago-Alarcon. 2018. Where are the birds in the matrix? Avian diversity in a Neotropical landscape mosaic. Wilson Journal of Ornithology 130:81–93.
- MacGregor-Fors, I., and J.E. Schondube. 2011. Use of tropical dry forests and agricultural areas by Neotropical bird communities. Biotropica 43:365–370.
- Marsden, S.J., M. Whiffin, and M. Galetti. 2001. Bird diversity and abundance in forest fragments and *Eucalyptus* plantations around an Atlantic forest reserve, Brazil. Biodiversity and Conservation 10:737–751.
- Martin, D.A., R. Andriafanomezantsoa, S. Droge, K. Osen, E. Rakotomalala, A. Wurz, A. Andrianarimisa, and H. Kreft. 2020. Bird diversity and endemism along a land-use gradient in Madagascar: the conservation value of vanilla agroforests. Biotropica 53:179–190.
- Martin, T.E., and J.R. Karr. 1986. Patch utilization by migrating birds: resource oriented? Ornis Scandinavica 17:165–174.
- Mas, A.H., and T.V. Dietsch. 2004 Linking shade coffee certification to biodiversity conservation: butterflies and birds in Chiapas, Mexico. Ecological Applications 14:642–654.
- McClure, C.J.W., B.W. Rolek, and G.E. Hill. 2012. Predicting occupancy of wintering migratory birds: is microhabitat information necessary? Condor 114:482–490.
- McLaren, M.F., C.M. White, N.J. Van Lanen, J.J. Birek, J.M. Berven, and D.J. Hanni. 2019. Integrated Monitoring in Bird Conservation Regions (IMBCR): field protocol for spatially-balanced sampling of land bird populations. Bird Conservancy of the Rockies, Brighton, CO.
- Mohd-Azlan, J., V.A.M. Fang, S.S. Kaicheen, L. Lok, and M.J. Lawes. 2019. The diversity of understory birds in fragments and oil palm plantation, Sarawak, Borneo. Journal of Oil Palm Research 3:437–447.
- Mokria, M., A. Gebrekirstos, H. Said, K. Hagdu, N. Hagazi, W. Dubale, and A. Brauning. 2022. Volume estimation models for avocado fruit. PloS One 17:e0263564.
- Muth, C.C., and F.A. Bazzaz. 2002. Tree canopy displacement at forest gap edges. Canadian Journal of Forest Research 32:247–254.
- Narango, D., D.W. Tallamy, K.J. Snyder, and R. Rice. 2019. Canopy tree preference by insectivorous birds in shade-coffee farms: implications for migratory bird conservation. Biotropica 51:387–398.
- Newbold, T., J.P.W. Scharlemann, S.H.M. Butchart, Ç.H. Şekercioğlu, R. Alkemade, H. Booth, and D.W. Purves. 2013. Eco-

- logical traits affect the response of tropical forest bird species to land-use intensity. Proceedings of the Royal Society B 280:20122131.
- Nilsson, S.G. 1979. Density and species richness of some forest bird communities in South Sweden. Oikos 33:392–401.
- Organisation for Economic Co-operation Development (OECD) and Food and Agriculture Organization of the United Nations (FAO). 2021. Chapter 11: Other products. Pp. 225–247 in OECD-FAO Agricultural Outlook 2021–2030.
- Perfecto, I., R. Rice, R. Greenberg, and M.E. Van der Voort. 1996. Shade coffee: a disappearing refuge for biodiversity. BioScience 46:598–608.
- Power, D.M. 1972. Numbers of bird species on the California islands. Evolution 26:451–463.
- Rabinowitz, D. 1981. Seven forms of rarity. Pp. 205–217 in The Biological Aspects of Rare Plant Conservation (H. Synge, ed.). John Wiley and Sons, Hoboken, NJ.
- Reed, J., J. van Vianen, S. Foli, J. Clendenning, K. Yang, M. Mac-Donald, G. Petrokofsky, C. Padoch, and T. Sunderland. 2017. Trees for life: the ecosystem service contribution of trees to food production and livelihoods in the tropics. Foreign Policy and Economics 84:62–71.
- Regos, A., L. Imbeau, M. Desrochers, A. Leduc, M. Robert, B. Jobin, L. Brotons, and P. Drapeau. 2018. Hindcasting the impacts of land-use changes on bird communities with species distribution models of bird atlas data. Ecological Implications 28:1867–1883.
- Rompré, G., W.D. Robinson, A. Desrochers, and G. Angehr. 2007. Environmental correlates of avian diversity in lowland Panama rain forests. Journal of Biogeography 34:802–815.
- Rosenberg, K.V., A.M. Dokter, P.J. Blancher, J.R. Sauer, A.C. Smith, P.A. Smith, J.C. Stanton, A. Panjabi, L. Helft, M. Parr, and P.P. Marra. 2019. Decline of the North American avifauna. Science 366:120–124.
- Scharlemann, J.P., R.E. Green, and A. Balmford. 2004. Land-use trends in Endemic Bird Areas: global expansion of agriculture in areas of high conservation value. Global Change Biology 10:2046–2051.
- Shannon, C.E. 1948. A mathematical theory of communication. The Bell System Technical Journal 27:379–423.
- Smith, A.L., J. Salgado-Ortiz, and R.J. Robertson. 2006. Distribution patters of migrant and resident birds in successional forest of the Yucatan Peninsula, Mexico. Biotropica 33:153–170.
- Sodhi, N.S., Ç.H. Şekercioğlu, J. Barlow, and S.K. Robinson. 2011. The state of tropical bird biodiversity. Pp. 1–26 *in* Conservation of Tropical Birds. John Wiley and Sons, Oxford, UK.
- Stein, A., G. Katharina, and H. Kreft. 2014. Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. Ecology Letters 17:866–880.
- Tejeda-Cruz, C., and W.J. Sutherland. 2004. Bird responses to shade coffee production. Animal Conservation 7:169–179.
- Tews, J., U. Brose, V. Grimm, K. Tielborger, M.C. Wichmann, M. Schwager, and F. Jeltsh. 2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. Journal of Biogeography 31:79–92.
- Tilman, M.C., M. Clark, D.R. Williams, K. Kimmel, S. Polasky, and C. Packer. 2017. Future threats to biodiversity and pathways to their prevention. Nature 546:83–81.

- Tuck, S.L., C. Winqvist, F. Mota, J. Ahnström, L. Turnbull, and J. Bengtsson. 2014. Land-use intensity and the effects of organic farming on biodiversity: a hierarchical meta-analysis. Journal of Applied Ecology 51:746–755.
- Van Bael, S.A., P. Bichier, I. Ochoa, and R. Greenberg. 2007. Bird diversity in cacao farms and forest fragments of Western Panama. Biodiversity and Conservation 16:2245–2256.
- Villaseñor, J.F., and R.L. Hutto. 1995. The importance of agricultural areas for the conservation of Neotropical migratory landbirds in western Mexico. Pp. 59–80 *in* Conservation of Neotropical Migratory Birds in Mexico (M.H. Wilson and S.A. Sader, eds.). Maine Agricultural and Forest Experiment Station, Orono, ME.
- Wunderle, J.M., and S.C. Latta. 1996. Avian abundance in sun and shade coffee plantations and remnant pine forest in the Cordillera Central, Dominican Republic. Ornitologia Neotrop-

- ical 7:19-34.
- Wunderle, J.M., Jr., and S. Latta. 1998. Avian resource use in Dominican shade coffee plantations. Wilson Bulletin 110:271–281
- Wunderle, J.M., and R.B. Waide. 1993. Distribution of overwintering Nearctic migrants in the Bahamas and Greater Antilles. Condor 95:904–933.
- Yang, K., S.E. Gergel, J.-Y. Duriaux-Chavarria, and F. Baudron. 2020. Forest edges near farms enhance wheat productivity measures: a test using high spatial resolution remote sensing of smallholder farms in southern Ethiopia. Frontiers in Sustainable Food Systems 4:130.
- Young, A., D. Aborn, S. Latta, J. Boyd. 2023. Data for: Avian use of avocado farms and intact forest in the northern Sierra de Bahoruco, Dominican Republic [Dataset]. Dryad.

Appendix 1. Abbreviations for common names and scientific names of bird species that were among the 20 most frequently observed species in each of the study sites.

| Species Code | English Common Name | Scientific Name |
|--------------|-----------------------------|--------------------------|
| AMKE | American Kestrel | Falco sparverius |
| AMRE | American Redstart | Setophaga ruticilla |
| ANEU | Antillean Euphonia | Chlorophonia musica |
| ANPI | Antillean Piculet | Nesoctites micromegas |
| BANA | Bananaquit | Coereba flaveola |
| BAWW | Black-and-white Warbler | Mniotilta varia |
| ВВТО | Broad-billed Tody | Todus subulatus |
| BFGR | Black-faced Grassquit | Melanospiza bicolor |
| BLPT | Black-crowned Palm-Tanager | Phaenicophilus palmarum |
| BTBW | Black-throated Blue Warbler | Setophaga caerulescens |
| CAEG | Cattle Egret | Bubulcus ibis |
| GDO | Common Ground Dove | Columbina passerina |
| CMWA | Cape May Warbler | Setophaga tigrina |
| BVI | Flat-billed Vireo | Vireo nanus |
| ABU | Greater Antillean Bullfinch | Melopyrrha violacea |
| GRAK | Gray Kingbird | Tyrannus dominicensis |
| HILC | Hispaniolan Lizard-Cuckoo | Coccyzus longirostris |
| IIMA | Hispaniolan Mango | Anthracothorax dominicus |
| IIWO | Hispaniolan Woodpecker | Melanerpes striatus |
| IPAT | Hispaniolan Parrot | Amazona ventralis |
| MODO | Mourning Dove | Zenaida macroura |
| IBTO | Narrow-billed Tody | Todus angustirostris |
| OMO | Northern Mockingbird | Mimus polyglottos |
| IOPA | Northern Parula | Setophaga americana |
| OVEN | Ovenbird | Seiurus aurocapilla |
| LAP | Plain Pigeon | Patagioenas inornata |
| PRAW | Prairie Warbler | Setophaga discolor |
| LTH | Red-legged Thrush | Turdus plumbeus |
| BAN | Smooth-billed Ani | Crotophaga ani |
| TOF | Stolid Flycatcher | Myiarchus stolidus |
| 'EHU | Vervain Hummingbird | Mellisuga minima |
| VNCR | White-necked Crow | Corvus leucognaphalus |
| ′FGR | Yellow-faced Grassquit | Tiaris olivaceus |

Appendix 2. Abbreviations for common names and scientific names of woody plant species and varieties that were among the 20 most frequently observed species and varieties in each of the study sites.

| Species Code | Local Common Name | Scientific Name |
|--------------|---------------------|---------------------------------|
| BAYA | Bayahonda | Neltuma juliflora |
| CAAG | Carga Agua | Senna atomaria |
| CANA | Palma Cana | Sabal spp. |
| CAND | Candelón | Parasenegalia skleroxyla |
| CARL | Carla | Persea americana var. carla |
| CERE | Cereza/Acerola | Malpighia emarginata |
| CHIV | Palo de Chivo | Senna atomaria |
| CHOQ | Choquete | Persea americana var. choquette |
| 000 | Palma de Coco | Cocos nucifera |
| OPE | Copey | Clusia rosea |
| ORD | | Cordia laevigata |
| RIO | Criollo | Persea americana spp. (hybrid) |
| ROT | | Croton spp. |
| RYT | | Erythroxylum brevipes |
| SCO | Escobón | Eugenia monticola |
| GREN | Grenadilla | Eugenia ligustrina |
| iuan | Guandules | Cajanus cajan |
| IUAY | Guayacán | Guaiacum officinale |
| UAZ | Guacimo | Guazuma ulmifolia |
| iUCO | Guaconejo | Amyris elemifera |
| ASS | Hass | Persea americana var. hass |
| ORS | Horsua | Coccoloba diversifolia |
| ОВО | Jobo | Spondias mombin |
| ULA | Lula | Persea americana var. lula |
| 1ELE | Melendez | Persea americana var. melendez |
| ARA | Naranja | Citrus spp. |
| IEEM | Neem/Tree of Heaven | Ailanthus altissima |
| LIV | Olivo | Morisonia ferruginea |
| AAM | Palo Amargo | Rauvolfia nitida |
| ICR | | Picramnia pentandra |
| LAT | Plátano | Musα spp. |
| NPN | Piñi Piñi/Quina | Exostema caribaeum |
| PYA | Papaya | Carica papaya |
| AND | | Randia aculeata |
| ACA | Sacasya | Vachellia macracantha |
| AVI | · | Savia sessiliflora |
| IDA | | Sida glabra |
| INK1 | | Unknown |
| JNK2 | | Unknown |